

Some non-normal Cayley digraphs of the generalized quaternion group of certain orders

Edward Dobson

Department of Mathematics and Statistics
PO Drawer MA
Mississippi State, MS 39762, U.S.A.
dobson@math.msstate.edu

Submitted: Mar 10, 2003; Accepted: Jul 30, 2003; Published: Sep 8, 2003
MR Subject Classifications: 05C25, 20B25

Abstract

We show that an action of $SL(2, p)$, $p \geq 7$ an odd prime such that $4 \nmid (p - 1)$, has exactly two orbital digraphs Γ_1, Γ_2 , such that $\text{Aut}(\Gamma_i)$ admits a complete block system \mathcal{B} of $p + 1$ blocks of size 2, $i = 1, 2$, with the following properties: the action of $\text{Aut}(\Gamma_i)$ on the blocks of \mathcal{B} is nonsolvable, doubly-transitive, but not a symmetric group, and the subgroup of $\text{Aut}(\Gamma_i)$ that fixes each block of \mathcal{B} set-wise is semiregular of order 2. If $p = 2^k - 1 > 7$ is a Mersenne prime, these digraphs are also Cayley digraphs of the generalized quaternion group of order 2^{k+1} . In this case, these digraphs are non-normal Cayley digraphs of the generalized quaternion group of order 2^{k+1} .

There are a variety of problems on vertex-transitive digraphs where a natural approach is to proceed by induction on the number of (not necessarily distinct) prime factors of the order of the graph. For example, the Cayley isomorphism problem (see [6]) is one such problem, as well as determining the full automorphism group of a vertex-transitive digraph Γ . Many such arguments begin by finding a complete block system \mathcal{B} of $\text{Aut}(\Gamma)$. Ideally, one would then apply the induction hypothesis to the groups $\text{Aut}(\Gamma)/\mathcal{B}$ and $\text{fix}_{\text{Aut}(\Gamma)}(\mathcal{B})|_B$, where $\text{Aut}(\Gamma)/\mathcal{B}$ is the permutation group induced by the action of $\text{Aut}(\Gamma)$ on \mathcal{B} , and $\text{fix}_{\text{Aut}(\Gamma)}(\mathcal{B})|_B$ is the subgroup of $\text{Aut}(\Gamma)$ that fixes each block of \mathcal{B} set-wise, and $B \in \mathcal{B}$. Unfortunately, neither $\text{Aut}(\Gamma)/\mathcal{B}$ nor $\text{fix}_{\text{Aut}(\Gamma)}(\mathcal{B})|_B$ need be the automorphism group of a digraph. In fact, there are examples of vertex-transitive graphs where $\text{Aut}(\Gamma)/\mathcal{B}$ is a doubly-transitive nonsolvable group that is not a symmetric group (see [7]), as well as examples of vertex-transitive graphs where $\text{fix}_{\text{Aut}(\Gamma)}(\mathcal{B})|_B$ is a doubly-transitive nonsolvable group that is not a symmetric group (see [2]). (There are also examples where $\text{Aut}(\Gamma)/\mathcal{B}$ is a solvable doubly-transitive group, but in practice, this is not usually

a genuine obstacle in proceeding by induction.) The only known class of examples of vertex-transitive graphs where $\text{Aut}(\Gamma)/\mathcal{B}$ is a doubly-transitive nonsolvable group, have the property that $\text{Aut}(\Gamma)/\mathcal{B}$ is a faithful representation of $\text{Aut}(\Gamma)$ and Γ is not a Cayley graph. In this paper, we give examples of vertex-transitive digraphs that are Cayley digraphs and the action of $\text{Aut}(\Gamma)/\mathcal{B}$ on \mathcal{B} is doubly-transitive, nonsolvable, not faithful, and not a symmetric group.

1 Preliminaries

Definition 1.1 Let G be a permutation group acting on Ω . If $\omega \in \Omega$, then a *sub-orbit* of G is an orbit of $\text{Stab}_G(\omega)$.

Definition 1.2 Let G be a finite group. The *socle* of G , denoted $\text{soc}(G)$, is the product of all minimal normal subgroups of G . If G is primitive on Ω but not doubly-transitive, we say G is *simply primitive*. Let G be a transitive permutation group on a set Ω and let G act on $\Omega \times \Omega$ by $g(\alpha, \beta) = (g(\alpha), g(\beta))$. The orbits of G in $\Omega \times \Omega$ are called the *orbitals* of G . The orbit $\{(\alpha, \alpha) : \alpha \in \Omega\}$ is called the *trivial orbital*. Let Δ be an orbital of G in $\Omega \times \Omega$. Define the *orbital digraph* Δ to be the graph with vertex set Ω and edge set Δ . Each orbital of G has a *paired orbital* $\Delta' = \{(\beta, \alpha) : (\alpha, \beta) \in \Delta\}$. Define the *orbital graph* Δ to be the graph with vertex set Ω and edge set $\Delta \cup \Delta'$. Note that there is a canonical bijection from the set of orbital digraphs of G to the set of sub-orbits of G (for fixed $\omega \in \Omega$).

Definition 1.3 Let G be a transitive permutation group of degree mk that admits a complete block system \mathcal{B} of m blocks of size k . If $g \in G$, then g permutes the m blocks of \mathcal{B} and hence induces a permutation in S_m , which we denote by g/\mathcal{B} . We define $G/\mathcal{B} = \{g/\mathcal{B} : g \in G\}$. Let $\text{fix}_{\mathcal{B}}(G) = \{g \in G : g(B) = B \text{ for every } B \in \mathcal{B}\}$.

Definition 1.4 Let G be transitive group acting on Ω with r orbital digraphs $\Gamma_1, \dots, \Gamma_r$. Define the *2-closure* of G , denoted $G^{(2)}$ to be $\cap_{i=1}^r \text{Aut}(\Gamma_i)$. Note that if G is the automorphism group of a vertex-transitive digraph, then $G^{(2)} = G$.

Definition 1.5 Let Γ be a graph. Define the *complement* of Γ , denoted by $\bar{\Gamma}$, to be the graph with $V(\bar{\Gamma}) = V(\Gamma)$ and $E(\bar{\Gamma}) = \{uv : u, v \in V(\Gamma) \text{ and } uv \notin E(\Gamma)\}$.

Definition 1.6 A group G given by the defining relations

$$G = \langle h, k : h^{2^{a-1}} = k^2 = m, m^2 = 1, k^{-1}hk = h^{-1} \rangle$$

is a *generalized quaternion group*.

Let $p \geq 5$ be an odd prime. Then $\text{GL}(2, p)$ acts on the set \mathbb{F}_p^2 , where \mathbb{F}_p is the field of order p , in the usual way. This action has two orbits, namely $\{0\}$ and $\Omega = \mathbb{F}_p^2 - \{0\}$. The action of $\text{GL}(2, p)$ on Ω is imprimitive, with a complete block system \mathcal{C} of $(p^2 - 1)/(p - 1) = p + 1$ blocks of size $p - 1$, where the blocks of \mathcal{C} consist of all scalar multiples of a given

vector in Ω (these blocks are usually called *projective points*), and the action of $\text{GL}(2, p)$ on the blocks of \mathcal{C} is doubly-transitive. Furthermore, $\text{fix}_{\text{GL}(2,p)}(\mathcal{C})$ is cyclic of order $p - 1$, and consists of all scalar matrices αI (where I is the 2×2 identity matrix) in $\text{GL}(2, p)$. Note that if $m|(p - 1)$, then $\text{GL}(2, p)$ admits a complete block system \mathcal{C}_m of $(p + 1)m$ blocks of size $(p - 1)/m$, and $\text{fix}_{\text{GL}(2,p)}(\mathcal{C}_m)$ consists of all scalar matrices $\alpha^i I$, where $\alpha \in \mathbb{F}_p^*$ is of order $(p - 1)/m$ and $i \in \mathbb{Z}$. Each such block of \mathcal{C}_m consists of all scalar multiples $\alpha^i v$, where v is a vector in \mathbb{F}_p^2 and $i \in \mathbb{Z}$. Hence $\text{GL}(2, p)/\mathcal{C}_m$ admits a complete block system \mathcal{D}_m consisting of $p + 1$ blocks of size m , induced by \mathcal{C}_m . Henceforth, we set $m = 2$ so that \mathcal{C}_2 consists of $2(p + 1)$ blocks of size $(p - 1)/2$, and \mathcal{D}_2 consists of $p + 1$ blocks of size 2. Note that as $p \geq 5$, $\text{SL}(2, p)$ is doubly-transitive on the set of projective points, as if $A \in \text{GL}(2, p)$, then $\det(A)^{-1}A \in \text{SL}(2, p)$. Finally, observe that $(-1)I \in \text{SL}(2, p)$. Thus $(-1)I/\mathcal{C}_2 \in \text{fix}_{\text{SL}(2,p)/\mathcal{C}_2}(\mathcal{D}_2) \neq 1$ so that $\text{SL}(2, p)/\mathcal{C}_2$ is transitive on \mathcal{C}_2 . Additionally, as $\text{fix}_{\text{GL}(2,p)}(\mathcal{C}_2) = \{\alpha^i I : |\alpha| = (p - 1)/2, i \in \mathbb{Z}\}$, $\text{SL}(2, p)/\mathcal{C}_2 \cong \text{SL}(2, p)$. That is, $\text{SL}(2, p)/\mathcal{C}_2$ is a faithful representation of $\text{SL}(2, p)$. We will thus lose no generality by referring to an element $x/\mathcal{C}_2 \in \text{SL}(2, p)/\mathcal{C}_2$ as simply $x \in \text{SL}(2, p)$. As each projective point can be written as a union of two blocks contained in \mathcal{C}_2 , we will henceforth refer to blocks in \mathcal{C}_2 as *projective half-points*.

2 Results

We begin with a preliminary result.

Lemma 2.1 *Let $p \geq 7$ be an odd prime such that $4 \nmid (p - 1)$, and let $\text{SL}(2, p)$ act as above on the $2(p + 1)$ projective half-points. Then the following are true:*

1. $\text{SL}(2, p)$ has exactly four sub-orbits; two of size 1 and 2 of size p ,
2. $\text{SL}(2, p)$ admits exactly one non-trivial complete block system which consists of $p + 1$ blocks of size 2, namely \mathcal{D}_2 , formed by the orbits of $(-1)I$.

PROOF. By [4, Theorem 2.8.1], $|\text{SL}(2, p)| = (p^2 - 1)p$. It was established above that $\text{SL}(2, p)$ admits \mathcal{D}_2 as a complete block system of $p + 1$ blocks of size 2, and this complete block system is formed by the orbits of $(-1)I$ as $(-1)I \in \text{fix}_{\text{SL}(2,p)}(\mathcal{D}_2)$ and is semi-regular of order 2. As $\text{SL}(2, p)/\mathcal{D}_2 = \text{PSL}(2, p)$ is doubly-transitive, there are two sub-orbits of $\text{SL}(2, p)/\mathcal{D}_2$, one of size 1 and the other of size p . Now, consider $\text{Stab}_{\text{SL}(2,p)}(x)$, where x is a projective half-point. Then there exists another projective half-point y such that $x \cup y$ is a projective point z . As $\{x, y\} \in \mathcal{D}_2$ is a block of size 2 of $\text{SL}(2, p)$, we have that $\text{Stab}_{\text{SL}(2,p)}(x) = \text{Stab}_{\text{SL}(2,p)}(y)$. Thus $\text{SL}(2, p)$ has at least two singleton sub-orbits. As $\text{SL}(2, p)/\mathcal{D}_2 = \text{PSL}(2, p)$ has one singleton sub-orbit, $\text{SL}(2, p)$ has exactly two singleton sub-orbits. We conclude that every non-singleton sub-orbit of $\text{SL}(2, p)$ has order a multiple of p . As the non-singleton sub-orbits of $\text{SL}(2, p)$ have order a multiple of p , $\text{Stab}_{\text{SL}(2,p)}(x)$ has either one non-singleton orbit of size $2p$ or two non-singleton orbits of size p . As the order of a non-singleton orbit must divide $|\text{Stab}_{\text{SL}(2,p)}(x)| = p(p - 1)/2$ which is odd as

$4 \nmid (p-1)$, $\text{SL}(2, p)$ must have exactly two non-singleton sub-orbits of size p . Thus 1) follows.

Suppose that \mathcal{D} is another non-trivial complete block system of $\text{SL}(2, p)$. Let $D \in \mathcal{D}$ with v a projective half-point in D . By [3, Exercise 1.5.9], D is a union of orbits of $\text{Stab}_{\text{SL}(2, p)}(v)$, so that $|D|$ is either 2, $p+1$, $p+2$, $2p$, or $2p+1$. Furthermore, as the size of a block of a permutation group divides the degree of the permutation group, $|D| = 2$ or $p+1$. If $|D| = 2$, then D is the union of two singleton orbits of $\text{Stab}_{\text{SL}(2, p)}(v)$, in which case D consists of two projective half-points whose union is a projective point. Thus if $|D| = 2$, then $D \in \mathcal{D}_2$ and $\mathcal{D} = \mathcal{D}_2$. If $|D| = p+1$, then \mathcal{D} consists of 2 blocks of size $p+1$ and D is the union of two orbits of $\text{Stab}_{\text{SL}(2, p)}(v)$, and these orbits have size 1 and p . We conclude that $\cup D$ does not contain the projective point q that contains v .

Now, $\text{fix}_{\text{SL}(2, p)}(\mathcal{D})$ cannot be trivial, as $\text{SL}(2, p)/\mathcal{D}$ is of degree 2 while $|\text{SL}(2, p)| = (p^2 - 1)p$. Then $|\text{fix}_{\text{SL}(2, p)}(\mathcal{D})| = (p^2 - 1)p/2$ as $\text{SL}(2, p)/\mathcal{D}$ is a transitive subgroup of S_2 . Furthermore, $-I \notin \text{fix}_{\text{SL}(2, p)}(\mathcal{D})$ as no block of \mathcal{D} contains the projective point q that contains v so that $-I$ permutes the two projective half-points whose union is q . Thus $\text{fix}_{\text{SL}(2, p)}(\mathcal{D}_2) \cap \text{fix}_{\text{SL}(2, p)}(\mathcal{D}) = 1$. As $\langle -I \rangle = \text{fix}_{\text{SL}(2, p)}(\mathcal{D}_2)$ and both $\text{fix}_{\text{SL}(2, p)}(\mathcal{D}_2)$ and $\text{fix}_{\text{SL}(2, p)}(\mathcal{D})$ are normal in $\text{SL}(2, p)$, we have that $\text{SL}(2, p) = \text{fix}_{\text{SL}(2, p)}(\mathcal{D}_2) \times \text{fix}_{\text{SL}(2, p)}(\mathcal{D})$. Thus a Sylow 2-subgroup of $\text{SL}(2, p)$ can be written as a direct product of two nontrivial 2-groups, contradicting [4, Theorem 8.3]. \square

Theorem 2.2 *Let $p \geq 7$ be an odd prime such that $4 \nmid (p-1)$. Then there exist exactly two digraphs Γ_i , $i = 1, 2$ of order $2(p+1)$ such that the following properties hold:*

1. Γ_i is an orbital digraph of $\text{SL}(2, p)$ in its action on the set of projective half-points and is not a graph,
2. $\text{Aut}(\Gamma_i)$ admits a unique nontrivial complete block system \mathcal{D}_2 which consists of $p+1$ blocks of size 2,
3. $\text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2) = \langle -I \rangle$ is cyclic of order 2,
4. $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2)$ is doubly-transitive but $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) \neq A_{p+1}$.

PROOF. By Lemma 2.1, $\text{SL}(2, p)$ in its action on the half-projective points has exactly four orbital digraphs; one consisting of $p+1$ independent edges (the edges of this graph consists of all edges of the form (v, w) , where $\cup\{v, w\}$ is a projective point; thus $\cup\{v, w\}$ is a block of \mathcal{D}_2), one which consists of only self-loops (and so is trivial with automorphism group S_{2p+2} and will henceforth be ignored) and two in which each vertex has in and out degree p . The orbital digraph Γ of $\text{SL}(2, p)$ consisting of $p+1$ independent edges is then $\bar{K}_{p+1} \wr K_2$. The other orbital digraphs of $\text{SL}(2, p)$, say Γ_1 and Γ_2 , each have in-degree and out-degree p .

If either Γ_1 or Γ_2 is a graph, then assume without loss of generality that Γ_1 is a graph. Then whenever $(a, b) \in E(\Gamma_1)$ then $(b, a) \in E(\Gamma_1)$. As Γ_1 is an orbital digraph, there exists $\alpha \in \text{SL}(2, p)$ such that $\alpha(a) = b$ and $\alpha(b) = a$. Raising α to an appropriate odd

power, we may assume that α has order a power of 2, and so $\alpha \in Q$, where Q is a Sylow 2-subgroup of $\text{SL}(2, p)$. As a Sylow 2-subgroup of $\text{SL}(2, p)$ is isomorphic to a generalized quaternion by [4, Theorem 8.3], Q contains a unique subgroup of order 2 (see [4, pg. 29]), which is necessarily $\langle -I \rangle$. If α is not of order 2, then $\alpha^2(a) = a$ and $\alpha^2(b) = b$ so that α has at least two fixed points. However, $(\alpha^2)^c = -I$ for some $c \in \mathbb{Z}$ and $-I$ has no fixed points, a contradiction. Thus α has order 2 and so $\alpha = -I$. Thus $(a, b) \in \bar{K}_{p+1} \wr K_2 \neq \Gamma_1$, a contradiction. Hence 1) holds.

We now establish that 2) holds. Suppose that for $i = 1$ or 2, $\text{Aut}(\Gamma_i)$ is primitive. We may then assume without loss of generality that $\text{Aut}(\Gamma_1)$ is primitive, and as $\text{Aut}(\Gamma_1) \neq K_{2(p+1)}$, $\text{Aut}(\Gamma_1)$ is simply primitive, and, of course, $\text{SL}(2, p)^{(2)} \leq \text{Aut}(\Gamma_1)$. First observe that by [11, Theorem 4.11], $\text{SL}(2, p)^{(2)}$ admits \mathcal{D}_2 as a complete block system. Let v be a projective half-point. By Lemma 2.1, $\text{SL}(2, p)$ has four sub-orbits relative to v , two of size 1, say $\mathcal{O}_1 = \{v\}$ and $\mathcal{O}_2 = \{w\}$, and two of size p , say \mathcal{O}_3 and \mathcal{O}_4 . By [11, Theorem 5.5 (ii)] the sub-orbits of $\text{SL}(2, p)^{(2)}$ relative to v are the same as the sub-orbits of $\text{SL}(2, p)$ relative to v . Thus the neighbors of v in Γ_1 consist of all elements in one of the sub-orbits \mathcal{O}_3 or \mathcal{O}_4 . Without loss of generality, assume that this sub-orbit is \mathcal{O}_3 . As $\text{Aut}(\Gamma_1)$ is primitive, by [3, Theorem 3.2A], every non-trivial orbital digraph of $\text{Aut}(\Gamma_1)$ is connected. Then the orbital digraph of $\text{Aut}(\Gamma_1)$ that contains $v\bar{w}$ is connected, and so $\mathcal{O}_2 = \{w\}$ is not a sub-orbit of $\text{Aut}(\Gamma_1)$. Of course, $\text{Aut}(\Gamma_1) = \text{Aut}(\bar{\Gamma}_1)$ so that $\text{Aut}(\bar{\Gamma}_1)$ is primitive as well. As if $\text{Aut}(\Gamma_1)$ has exactly two sub-orbits, then $\text{Aut}(\Gamma_1)$ is doubly-transitive and hence $\Gamma_1 = K_{2(p+1)}$ which is not true, $\text{Aut}(\Gamma_1)$ has exactly three sub-orbits. Clearly \mathcal{O}_3 is a sub-orbit of $\text{Aut}(\Gamma_1)$ so that the only sub-orbits of $\text{Aut}(\Gamma_1)$ relative to v are \mathcal{O}_1 , \mathcal{O}_3 , and $\mathcal{O}_2 \cup \mathcal{O}_4$. Thus the neighbors of v in $\bar{\Gamma}_1$ are all contained in one sub-orbit of $\text{Aut}(\Gamma_1)$ relative to v . However, one of these directed edges is an edge (as $\bar{\Gamma}_1 = \Gamma_2 \cup (\bar{K}_{p+1} \wr K_2)$), and so every neighbor of v in $\bar{\Gamma}_1$ is an edge. Thus every neighbor of v in Γ_1 is an edge. However, we have already established that Γ_1 is a digraph that is not a graph, a contradiction. Whence $\text{Aut}(\Gamma_i)$, $i = 1, 2$, are not primitive, and as $\text{SL}(2, p) \leq \text{Aut}(\Gamma_i)$, we have by Lemma 2.1 that \mathcal{D}_2 is the unique complete block system of $\text{Aut}(\Gamma_i)$, $i = 1, 2$. Thus (2) holds.

If $\text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2)$ is not cyclic, then there exists $1 \neq \gamma \in \text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2)$ such that $\gamma(v) = v$ for some $v \in V(\Gamma_i)$. It is then easy to see that $\text{Aut}(\Gamma_i)$ has only three sub-orbits, two of size 1, and one of size $2p$, a contradiction. Thus (3) holds.

To establish (4), as $\text{SL}(2, p)/\mathcal{D}_2 = \text{PSL}(2, p)$ which is doubly-transitive in its action on the blocks (projective points) of \mathcal{D}_2 , we have that $\text{Aut}(\Gamma_i)/\mathcal{D}_2$ is doubly-transitive. As $\text{PSL}(2, p) \leq \text{Aut}(\Gamma_i)/\mathcal{D}_2$, by [1, Theorem 5.3] $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2)$ is a doubly-transitive non-abelian simple group acting on $p+1$ points. Thus we need only show that $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) \neq A_{p+1}$.

Assume that $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) = A_{p+1}$. Recall that as p is odd, a Sylow 2-subgroup Q of $\text{SL}(2, p)$ is a generalized quaternion group. Furthermore, the unique element of Q of order 2, namely $-I$, is contained in every Sylow 2-subgroup of $\text{SL}(2, p)$ and is semiregular. Observe that as $4 \nmid (p-1)$, $4 \mid (p+1)$. Then Q contains an element δ such that δ/\mathcal{D}_2 is a product of $(p+1)/4$ disjoint 4-cycles and $\langle \delta^4 \rangle = \text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2) = \langle -I \rangle$. Let $\delta/\mathcal{D}_2 = z_0 \dots z_{\frac{p+1}{4}-1}$ be the cycle decomposition of δ/\mathcal{D}_2 . As $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) = A_{p+1}$, there

exists $\omega \in \text{Aut}(\Gamma_i)$ such that $\omega/\mathcal{D}_2 = z_0 z_1^{-1} \dots z_{\frac{p+1}{4}-1}^{-1}$ (note that if ω/\mathcal{D}_2 is not an even permutation, then δ/\mathcal{D}_2 is not an even permutation, in which case $\text{Aut}(\Gamma_i)/\mathcal{D}_2 = S_{p+1}$ and $\omega \in \text{Aut}(\Gamma_i)$). Then $|\delta\omega/\mathcal{D}_2| = 2$ so that $(\delta\omega)^2 \in \text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2)$. Let \mathcal{O}_0 be the union of the non-singleton orbits of $\langle z_0 \rangle$, and \mathcal{O}_1 be the union of the non-singleton orbits of $\langle z_1 \rangle$ (note that as $p \geq 7$, $p+1 \geq 8$, so that $(p+1)/4 \geq 2$). Let $D \in \mathcal{D}_2$ such that $D \subset \mathcal{O}_1$. Then $\delta\omega|_D$ has order 1 or 2, so that $(\delta\omega)^2|_D = 1$. Thus if $\omega|_{\mathcal{O}_0} \in \delta|_{\mathcal{O}_0}$, then $(\delta\omega)^2 \in \text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2) = \langle -I \rangle$, $(\delta\omega)^2 \neq 1$, but $(\delta\omega)^2$ has a fixed point, a contradiction. Thus $\omega|_{\mathcal{O}_0} \notin \delta|_{\mathcal{O}_0}$. Then $H = \langle \delta, \omega \rangle|_{\mathcal{O}_0}$ has a complete block system \mathcal{E} of 4 blocks of size 2 induced by \mathcal{D}_2 . Furthermore, H/\mathcal{E} is cyclic of order 4, so that $\text{fix}_H(\mathcal{E})$ has order at least 4. Then $\text{Stab}_H(v) \neq 1$ for every $v \in \mathcal{O}_0$. In particular, \mathcal{E} consists of 4 blocks of size 2, and $\text{Stab}_H(v)$ is the identity on some block of \mathcal{E} while being transitive on some other block. As each block of \mathcal{E} is also a block of \mathcal{D}_2 , $\text{Stab}_{\text{Aut}(\Gamma)}(v)$ is transitive on some block D_v of \mathcal{D}_2 . This then implies that $\text{Stab}_{\text{Aut}(\Gamma_i)}(v)$ has three orbits, two of size one and one of size $2(p+1) - 2$, a contradiction. \square

Corollary 2.3 *Let $p = 2^k - 1 > 7$ be a Mersenne prime. Then there exist exactly two digraphs Γ_i , $i = 1, 2$ of order 2^{k+1} such that the following properties hold:*

1. Γ_i is an orbital digraph of $\text{SL}(2, p)$ in its action on the set of projective half-points and is not a graph,
2. $\text{Aut}(\Gamma_i)$ admits a unique complete block system \mathcal{D}_2 which consists of 2^k blocks of size 2,
3. $\text{fix}_{\text{Aut}(\Gamma_i)}(\mathcal{D}_2)$ is cyclic of order 2,
4. $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) = \text{PSL}(2, p)$ is doubly-transitive,
5. Γ_i is a Cayley digraph of the generalized quaternion group of order 2^{k+1} .

PROOF. In view of Theorem 2.2, we need only show that $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) = \text{PSL}(2, p)$ and that each Γ_i is a Cayley digraph of the generalized quaternion group Q of order 2^{k+1} . As $|\text{SL}(2, p)| = 2^k(2^k - 1)(2^k - 2)$, a Sylow 2-subgroup of $\text{SL}(2, p)$ has order 2^{k+1} , and as p is odd, is isomorphic to a generalized quaternion group of order 2^{k+1} . As a transitive group of prime power order q^ℓ contains a transitive Sylow q -subgroup [10, Theorem 3.4'], a Sylow 2-subgroup Q of $\text{SL}(2, p)$ is transitive and thus regular. It then follows by [9] that each Γ_i is isomorphic to a Cayley digraph of Q . Furthermore, $\text{Stab}_{\text{Aut}(\Gamma_i)/\mathcal{D}_2}(v)$ is of index 2^k in $\text{Aut}(\Gamma_i)/\mathcal{D}_2$. By [5, Theorem 1] we have that either $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2)$ is A_{2^k} or $\text{PSL}(2, p)$. As by Theorem 2.2, $\text{soc}(\text{Aut}(\Gamma_i)/\mathcal{D}_2) \neq A_{2^k}$, the result follows. \square

References

- [1] Cameron, P. J., Finite permutation groups and finite simple groups, *Bull. London Math. Soc.* **13** (1981) 1–22.

- [2] Cheng, Y., and Oxley, J., On weakly symmetric graphs of order twice a prime, *J. Comb. Theory Ser. B* **42** 1987, 196-211.
- [3] Dixon, J.D., and Mortimer, B., *Permutation Groups*, Springer-Verlag New York, Berlin, Heidelberg, Graduate Texts in Mathematics, **163**, 1996.
- [4] Gorenstein, D., *Finite Groups*, Chelsea Publishing Co., New York, 1968.
- [5] Guralnick, R. M., Subgroups of prime power index in a simple group, *J. of Algebra* **81** 1983, 304-311.
- [6] Li, C. H., On isomorphisms of finite Cayley graphs - a survey, *Disc. Math.*, **246** (2002), 301-334.
- [7] Marušič, D., and Scapellato, R., Imprimitive Representations of $SL(2, 2^k)$ *J. Comb. Theory Ser. B* **58** 1993, 46-57.
- [8] Sabidussi, G., The composition of graphs, *Duke Math J.* **26** (1959), 693-696.
- [9] Sabidussi, G. O., Vertex-transitive graphs, *Monatshefte für Math.* **68** 1964, 426-438.
- [10] Wielandt, H. (trans. by R. Bercov), *Finite Permutation Groups*, Academic Press, New York, 1964.
- [11] Wielandt, H., Permutation groups through invariant relations and invariant functions, lectures given at The Ohio State University, Columbus, Ohio, 1969.
- [12] Wielandt, H., *Mathematische Werke/Mathematical works*. Vol. 1. Group theory, edited and with a preface by Bertram Huppert and Hans Schneider, Walter de Gruyter & Co., Berlin, 1994.