

Game list colouring of graphs

M. Borowiecki

Faculty of Mathematics, Computer Science and Econometrics
University of Zielona Góra
Szafrana 4a, 65–516 Zielona Góra, Poland
m.borowiecki@wmie.uz.zgora.pl

E. Sidorowicz

Faculty of Mathematics, Computer Science and Econometrics
University of Zielona Góra
Szafrana 4a, 65–516 Zielona Góra, Poland
e.sidorowicz@wmie.uz.zgora.pl

Zs. Tuza*

Computer and Automation Institute
Hungarian Academy of Sciences, Hungary
and
Department of Computer Science
University of Pannonia, Veszprém, Hungary
tuza@sztaki.hu

Submitted: Jan 11, 2006; Accepted: Mar 15, 2007; Published: Mar 22, 2007
Mathematics Subject Classifications: 05C15, 05C75

Abstract

We consider the two-player game defined as follows. Let (G, L) be a graph G with a list assignment L on its vertices. The two players, Alice and Bob, play alternately on G , Alice having the first move. Alice's goal is to provide an L -colouring of G and Bob's goal is to prevent her from doing so. A move consists in choosing an uncoloured vertex v and assigning it a colour from the set $L(v)$. Adjacent vertices of the same colour must not occur. This game will be called *game list colouring*. The *game choice number* of G , denoted by $\text{ch}_g(G)$, is defined as the least k such that Alice has a winning strategy for any k -list assignment of G .

We characterize the class of graphs with $\text{ch}_g(G) \leq 2$ and determine the game choice number for some classes of graphs.

*Research supported in part by the Hungarian Scientific Research Fund, grant OTKA T-049613.

1 Introduction and notation

All graphs considered in this paper are simple, i.e., finite, undirected, loopless and without multiple edges. The notation $(H \leq G)$ $H \subseteq G$ means that H is (an induced) subgraph of G . We say that G contains H whenever G contains (an induced) subgraph isomorphic to H . The set of neighbours of a vertex $v \in V(G)$ is denoted by $N_G(v)$, or briefly by $N(v)$. The *degree* of v is the number $|N(v)|$ and is denoted by $d(v)$. The symbol $\Delta(G)$ denotes the maximum degree of G . The vertex-disjoint union of two graphs G and H is denoted by $G \cup H$. By P_n, C_n, S_n we denote the path, the cycle, the star with n vertices. For undefined concepts we refer the reader to [5, 10].

Let N denote the set of positive integers and 2^N denote the power set of N . Let $G = (V, E)$ be a graph. A function $f : V \rightarrow N$ is a *colouring* of G if $f(v) \neq f(u)$ whenever $vu \in E$. A *list assignment* L for G is a function $L : V \rightarrow 2^N$. If $|L(v)| = k$ for every $v \in V$ then L is a *k-list-assignment* for G . A graph G with a given list assignment L will be denoted by (G, L) .

A function $f : V \rightarrow N$ is an *L-colouring* of G if f is a colouring of G with the additional requirement that $f(v) \in L(v)$ for all $v \in V(G)$. If G admits an L -colouring, then G is said to be *L-colourable*.

The graph G is *k-choosable* if it is L -colourable for every k -list assignment. The *choice number* or the *choosability* of G , denoted by $\text{ch}(G)$ is the smallest natural number k such that G is k -choosable.

The idea of considering L -colouring and k -choosability has been introduced independently by Vizing [18] and Erdős, Rubin and Taylor [6]. In both papers the choosability version of Brooks' theorem [3] was proved. The choosability version of Gallai's theorem [8] was proved independently by Thomassen [16] and by Kostochka et al. [14]. In [2] some extensions of these two basic theorems have been proved. A characterization of 2-choosable graphs is given in [6].

We consider the two-player game defined as follows. The two players are Alice and Bob and they play alternately with Alice having the first move. Given a graph G and a set C of colours, the players take turns colouring vertices of G (one vertex per move) with colours from C . If after $|V(G)|$ moves the graph is coloured then Alice wins, otherwise Bob wins; i.e., Bob wins whenever an impasse is reached before the whole graph is coloured.

The *game chromatic number* of G , denoted by $\chi_g(G)$, is defined as the least cardinality of C for which Alice has a winning strategy. The number $\chi_g(G)$ is well-defined since we have $\chi(G) \leq \chi_g(G) \leq |V(G)|$.

The game chromatic number of a graph was first introduced by Bodlaender in [1] where its computational complexity was studied. Bodlaender also showed that $\chi_g(T) \leq 5$ holds for all trees T , and exhibited trees with $\chi_g(T) \geq 4$. In [7] Bodlaender's upper bound was improved to $\chi_g(T) \leq 4$. Kierstead and Trotter proved in [12] that $\chi_g(G) \leq 33$ for every planar graph G and that the maximum game chromatic number of an outerplanar graph is between 6 and 8. The currently known best upper bounds are 18 for planar graphs [11] and 7 for outerplanar graphs [9].

The edge-colouring version of this game was considered by Cai and Zhu [4]. Nešetřil

and Sopena [15] introduced an oriented version of this game and developed some techniques for bounding the oriented game chromatic number. The conjecture of Nešetřil and Sopena was proved by Kierstead and Tuza in [13]. Further references on graph colouring games and L -colourings can be found e.g. in [17].

In this paper we are interested in another version of this game. We combine concepts of game and list colourings in the following way. Let (G, L) be a graph G with a list assignment L . Again the two players are Alice and Bob and they play alternately with Alice having the first move. Alice's goal is to provide an L -colouring of G and Bob's goal is to prevent her from doing so. A move consists in choosing an uncoloured vertex v and assigning it a colour from the set $L(v)$. This game will be called *game list colouring*.

We say that (G, L) is *game list colourable* if Alice has a winning strategy. The *game choice number* of G , denoted by $\text{ch}_g(G)$, is defined as the least k such that Alice has a winning strategy for any k -list assignment of G . It is clear that this number is well-defined.

In Section 2 we present some basic properties of the game choice number and we investigate the relations between the game choice number, the game chromatic number and the choice number. In Section 3 we characterize the graphs whose game choice number is equal to two.

2 Preliminary results

From the definitions we immediately have the following relations:

$$\begin{aligned} \chi(G) &\leq \chi_g(G) \leq \text{ch}_g(G) \leq \Delta(G) + 1, \\ \chi(G) &\leq \text{ch}(G) \leq \text{ch}_g(G) \leq \Delta(G) + 1. \end{aligned}$$

The following result provides a sufficient condition for a graph with a list assignment L to be L -colourable.

Theorem 1 [18, 6]. *Let G be a connected graph with a list assignment L satisfying the following conditions:*

- (a) $|L(v)| \geq d(v)$ for every $v \in V(G)$,
- (b) There is $v_0 \in V(G)$ such that $|L(v_0)| \geq d(v_0) + 1$.

Then G is L -colourable.

Observation 1 *Let (G, L) be a graph with $|L(v)| \geq d(v) + 1$ for every $v \in V(G)$. It is clear that (G, L) is game list colourable, i.e., there is a winning strategy for Alice.*

A natural question arises: Can Theorem 1 be extended to the game list colouring? In general the answer is NO. To explain this, we need some notation.

Let (G, L) be a graph with $|L(v)| \geq d(v)$ for every $v \in V(G)$ and let

$$\begin{aligned} X_{(G,L)} &= \{v \in V(G) : |L(v)| = d(v)\}, \\ Y_{(G,L)} &= V - X_{(G,L)} = \{v \in V(G) : |L(v)| \geq d(v) + 1\}. \end{aligned}$$

Proposition 1 For any natural number n there is a graph (G, L) with $|Y_{(G,L)}| - |X_{(G,L)}| > n$ such that (G, L) is not game list colourable.

Proof. Let (H, L_1) be a graph of even order, such that $|L_1(v)| \geq d(v) + 1$ for all $v \in V(H)$. Let (P_4, L_2) be the path with the list assignment $L_2(v) = \{1, 2\}$ for all $v \in V(P_4)$. It is easy to see that Bob has a winning strategy on the graph $G = P_4 \cup H$. Indeed, he can force Alice to be the first one who colours a vertex of the P_4 component; and then he can make P_4 uncolourable, by assigning the opposite colour at distance 2 from the firstly coloured one. If $|V(H)| = n + 4$, then Proposition 1 follows. ■

Let $d_{(G,L)} = \min\{d(v) : v \in X_{(G,L)}\}$.

Lemma 1 Let (G, L) be a graph with $|L(v)| \geq d(v)$ for every $v \in V(G)$ and let $|X_{(G,L)}| \leq \lceil \frac{d}{2} \rceil$, where $d = d_{(G,L)}$. Then Alice has a winning strategy.

Proof. Alice colours vertices of the set $X_{(G,L)}$ one by one until it gets completely coloured. It is possible because if v is a vertex of $X_{(G,L)}$ which is coloured by Alice in the k -th move, then in the set $N(v)$ there are at most $k - 1$ vertices coloured by Alice and at most $k - 1$ vertices coloured by Bob. Since $2k - 2 \leq 2 \lceil \frac{d}{2} \rceil - 2 < d(v) = |L(v)|$, Alice can choose a colour in $L(v)$ for the vertex v . Thus, the vertices of $X_{(G,L)}$ can be coloured from their lists. It is obvious (see Observation 1) that the vertices of $Y_{(G,L)}$ have a list colouring at any move of the game. ■

For complete bipartite graphs we have $\chi_g(K_{n,n}) = 3$ ($n \geq 2$). From the next theorem it follows that for almost all complete bipartite graphs the game choice number is greater than the game chromatic number.

We will use the following notation. If Alice colours a vertex v with colour c at move k , then it will be denoted by $A_k := [f(v) = c]$. Similarly, Bob's k -th move will be denoted by B_k .

Theorem 2 Let $n = \binom{k}{\lfloor \frac{k+1}{2} \rfloor} \binom{k}{\lceil \frac{k+1}{2} \rceil}$, $k \geq 2$. Then $\text{ch}_g(K_{n,n}) \geq k + 2$.

Proof. Let us denote the vertex classes of $K_{n,n}$ by $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_n\}$. We split the $2k$ -element colour set $\{1, 2, \dots, 2k\}$ into two disjoint k -sets $C_1 = \{1, 2, \dots, k\}$ and $C_2 = \{k + 1, k + 2, \dots, 2k\}$ and, as lists for the vertices, we take the $(k + 1)$ -subsets that are split equally or almost equally between C_1 and C_2 . If $k + 1$ is even, equal split is possible, each of the selected $(k + 1)$ -subsets will contain $\frac{k+1}{2}$ elements of C_1 and $\frac{k+1}{2}$ elements of C_2 . If $k + 1$ is odd, each of the selected $(k + 1)$ -subsets will contain $\frac{k}{2} + 1$ elements of C_1 and $\frac{k}{2}$ elements of C_2 . Assign each of these $(k + 1)$ -subsets of colours to one vertex of the set X and to one vertex of the set Y .

To prove the theorem it is enough to find a winning strategy for Bob. We construct it as follows. By symmetry, assume that Alice colours in the first move a vertex of X with a colour from its list, say $A_1 := [f(x_1) = c]$ and $c \in C_i$ ($i = 1$ or $i = 2$). During the first part of the game (at most k moves of Bob), Bob will play on the set X (because Alice started to colour a vertex of X). Bob's goal is to achieve a situation where at least $\lceil \frac{k+1}{2} \rceil$

colours of C_1 and $\lfloor \frac{k+1}{2} \rfloor$ colours of C_2 appear in X . If he can do so, then in the remaining part of the game the attempted choice by Alice must fail in the set Y because the entire list of some vertex $y_j \in Y$ is excluded by the partial colouring constructed on X .

If k is odd then Bob in the first move chooses an uncoloured vertex x_j and assigns a colour from $L(x_j)$ which is *not* in C_i . If k is even then Bob chooses for x_j a colour from C_i . Having done so, an even number of non-used colours remain in both C_1 and C_2 . Then, if in any one later step Alice colours a vertex of Y with a colour $c \in C_i$ ($i = 1$ or $i = 2$) then Bob colours a vertex of X which can be coloured with a new colour of the same C_i . Otherwise, if Alice colours a vertex of X then Bob again chooses an uncoloured vertex of X and assigns any new colour different than previously used by Alice and himself. The large number of vertices implies that Bob can follow this rule until more than half of C_1 and not less than half of C_2 appear in X . Thus $(K_{n,n}, L)$ is not $(k + 1)$ -game-list-colourable. ■

Let G be the r -partite graph K_{t_1, t_2, \dots, t_r} such that $t_1 = t_2 = \dots = t_r = 2$. In [6] it has been shown that the choice number of G is equal to r . It is easy to see that for this graph we have $\chi_g(G) = \text{ch}_g(G) = \Delta(G) + 1 = 2r - 1$, because Bob can repeat the colour assigned by Alice in each step, colouring the unique vertex nonadjacent to the latest choice of Alice. From the next theorem it follows that if one of the independent sets contains at least three vertices then $\text{ch}_g(K_{t_1, t_2, \dots, t_r}) \leq \Delta(K_{t_1, t_2, \dots, t_r})$.

Theorem 3 *Let $r \geq 3$, $1 \leq i < r$, $t_1 = t_2 = \dots = t_i = 3$, $t_{i+1} = \dots = t_r = 2$. Then*

$$\text{ch}_g(K_{t_1, t_2, \dots, t_r}) \leq \Delta(K_{t_1, t_2, \dots, t_r}).$$

Proof. Let (G, L) be the r -partite graph that satisfies the assumptions of the theorem, with a t -list-assignment where $t = \Delta(K_{t_1, t_2, \dots, t_r})$. Let V_1, V_2, \dots, V_r be its independent sets and let $V_1 = \{v_1, v_2, v_3\}$. We shall describe a winning strategy for Alice.

If there is a colour c such that $c \in L(v_1) \cap L(v_2) \cap L(v_3)$ then $A_1 := [f(v_1) = c]$. In the second move Alice colours v_2 or v_3 with the colour c . Let C be the set of coloured vertices after Bob's second move. The subgraph $G - C$ with the list assignment L' such that $L'(v) = L(v) \setminus \bigcup_{u \in N(v) \cap C} f(u)$ for $v \in V(G - C)$ satisfies the assumptions of Lemma 1 and then Alice wins.

Suppose that $L(v_1) \cap L(v_2) \cap L(v_3) = \emptyset$. Then each colour of $L(v_1) \cup L(v_2) \cup L(v_3)$ is in at most two of the lists $L(v_1), L(v_2), L(v_3)$. Therefore Alice can choose a vertex v_i such that $|L(v_i) \cap L(v_j)| \leq \frac{t}{2}$ for $i \neq j$. Assume that $v_i = v_1$. Let $V' \subseteq V \setminus V_1$ be the set of vertices such that the subgraph induced by V' is the $(r-1)$ -partite graph $K_{2, 2, \dots, 2}$. For any colour c let us denote the vertex set $N(c) = \{v \in V' : c \in L(v)\}$. In the first move Alice colours the vertex v_1 with the colour c_0 such that $|N(c_0)| = \min\{|N(c)| : c \in L(v_1)\}$. Suppose that $|N(c_0)| \leq r - 1$. Let v be the vertex coloured by Bob in the first move and $B_1 := [f(v) = c]$. Then the graph $(G - v_1) - v$ with the list assignment L' such that $L'(w) = L(w) \setminus \bigcup_{u \in N(w) \cap C} f(u)$ for $w \in V((G - v_1) - v)$ satisfies the assumptions of Lemma 1. Hence Alice has a winning strategy.

Assume that $|N(c_0)| > r - 1$. Let us denote $|N(c_0)| = r + d$. We consider two cases:

Case 1. In the first move Bob colours v_2 or v_3 .

Assume that Bob colours v_2 . If $c_0 \in L(v_3)$ then Alice puts $A_2 := [f(v_3) = c_0]$. Let C be the set of coloured vertices after Bob's second move. The subgraph $G - C$ with the list assignment L' such that $L'(v) = L(v) \setminus \bigcup_{u \in N(v) \cap C} f(u)$ satisfies the assumptions of Lemma 1.

Suppose that $c_0 \notin L(v_3)$. If there is $c \in L(v_3)$ such that $|N(c) \cap N(c_0)| \leq r - 1$ then again by Lemma 1 Alice wins. We may assume that $|N(c) \cap N(c_0)| \geq r$ for $c \in L(V_3)$. We show that in this case $\sum_{v \in N(c_0)} |L(v)| > |N(c_0)|t$ which contradicts the assumption that each vertex has been assigned to a list of length t . By the choice of c_0 , $|N(c)| \geq r + d$ holds for all $c \in L(v_1)$. Then for any $c \in L(v_1)$ we have

$$\sum_{v \in N(c_0)} |L(v)| \geq |N(c) \cap N(c_0)| \geq |N(c)| - |V' \setminus N(c_0)| \geq r + d - 2(r - 1) + r + d \geq 2(d + 1)$$

and

$$\begin{aligned} \sum_{v \in N(c_0)} |L(v) \cap \bigcup_{u \in V_1} L(u)| &\geq \sum_{c \in L(v_1) \cup L(v_3)} |N(c) \cap N(c_0)| \geq \\ \sum_{c \in L(v_3)} |N(c) \cap N(c_0)| + \sum_{c \in L(v_1) \setminus L(v_3)} |N(c) \cap N(c_0)| &\geq \\ rt + |L(v_1) \setminus L(v_3)|2(d + 1) &\geq rt + 2\frac{t}{2}(d + 1) > t(r + d) = |N(c_0)|t, \end{aligned}$$

a contradiction.

Case 2. In the first move Bob colours a vertex of $V - V_1$.

If $c_0 \in L(v_2)$ or $c_0 \in L(v_3)$ then by Lemma 1 Alice has a winning strategy. Assume that Bob colours the vertex v' and $B_1 := [f(v') = c']$. If $c' \notin L(v_2)$ or $c' \notin L(v_3)$ then similarly as in Case 1 we can show that there is a colour $c \in L(v_2)$ (or $c \in L(v_3)$) such that $|N(c) \cap N(c_0)| < r - 1$. Then assume that $c' \in L(v_2)$ and $c' \in L(v_3)$ and, for all $c \in (L(v_2) \cup L(v_3)) \setminus \{c'\}$, $|N(c) \cap N(c_0)| > r - 1$ holds. Then we have

$$\begin{aligned} \sum_{v \in N(c_0)} |(L(v) \cap \bigcup_{u \in V_1} L(u)) \setminus \{c'\}| &\geq \sum_{v \in N(c_0)} |L(v) \cap \bigcup_{u \in V_1} L(u)| - |N(c_0)| \geq \\ \sum_{c \in L(v_1) \cup L(v_3)} |N(c) \cap N(c_0)| - |N(c_0)| &\geq \\ \sum_{c \in L(v_3)} |N(c) \cap N(c_0)| + \sum_{c \in L(v_1) \setminus L(v_3)} |N(c) \cap N(c_0)| - |N(c_0)| &\geq \\ rt + |L(v_1) \setminus L(v_3)|2(d + 1) - |N(c_0)| &= t(r + d) + t - (r + d). \end{aligned}$$

Recall that $r + d = |N(c_0)|$. Since $|N(c_0)| \leq 2(r - 1) < t$, we have

$$\sum_{v \in N(c_0)} |L(v) \cap \bigcup_{u \in V_1} L(u)| \geq \sum_{v \in N(c_0)} |(L(v) \cap \bigcup_{u \in V_1} L(u)) \setminus \{c'\}| > t(r + d) = |N(c_0)|t, \text{ a contradiction. } \blacksquare$$

3 A characterization of graphs with game choice number two

In this section we shall characterize the graphs for which Alice has a winning strategy in the game list colouring for any 2-list assignment.

The following proposition trivially holds.

Proposition 2 *If $\text{ch}_g(G) = 2$ then $C_{2k+1} \not\subseteq G$.*

Let us denote by H_1 and H_2 the 5-vertex graphs exhibited in Figures 1 and 3, respectively.

Lemma 2 *If $\text{ch}_g(G) = 2$ then G does not contain any graph of the set $\{H_1, P_5, K_{2,3}, H_2\}$ as an induced subgraph.*

Proof. Suppose that G contains an induced subgraph H isomorphic to H_1 . Let us denote the vertices of H as in Figure 1. Let L be the 2-list-assignment such that $L(v) = \{1, 2\}$ for $v \in V(H)$ and $L(u) = \{3, 4\}$ for $u \in V(G) - V(H)$. We will give a winning strategy for Bob. If in the first move Alice colours a vertex v of H then Bob colours the vertex w of H such that $N_H(v) \cap N_H(w) \neq \emptyset$ using another colour than Alice did. Thus a third colour is necessary to colour the common neighbour of v and w . If in the first move Alice colours a vertex v of $G - H$ then Bob chooses the vertex x_1 of H and he puts $B_1 := [f(x_1) = 1]$. After the second move of Alice, Bob colours a vertex $w \in V(H)$ with $N_H(w) \cap N_H(u) = \emptyset$ (u is a vertex coloured by Alice or Bob) using a colour different from the colour of u . Such w can be chosen, no matter which vertex has been coloured by Alice (e.g., if Alice has coloured x_5 , then Bob can consider $u = x_1$ and $w = x_2$). Then a third colour is necessary to colour the common uncoloured neighbour of w and u .




Figure 1. The graph H_1

Suppose that G contains an induced subgraph P isomorphic to P_5 . Let $V(P) = \{x_1, x_2, x_3, x_4, x_5\}$ and $E(P) = \{x_1x_2, x_2x_3, x_3x_4, x_4x_5\}$. Let L be the 2-list-assignment such that $L(v) = \{1, 2\}$ for all $v \in V(P)$ and $L(u) = \{3, 4\}$ for all $u \in V(G) - V(P)$. If Alice colours x_i with colour $j \in \{1, 2\}$, the next move of Bob is to colour x_{i-2} or x_{i+2} with $3 - j$. Otherwise, if Alice colours a vertex of $G - P$, Bob puts colour 1 on x_3 , and then the colour 2 on x_1 or x_5 in his next move.

Suppose that G contains an induced subgraph K isomorphic to $K_{2,3}$. Let us denote the vertices of K as in Figure 2. Let L be the 2-list-assignment such that $L(v) = \{4, 5\}$ for $v \in V(G) - V(K)$ and as given in Figure 2 for the vertices of K . Suppose that in the first move Alice colours a vertex v of K . Then Bob chooses the vertex w such that $N_H(v) \cap N_H(w) \neq \emptyset$. It is easy to see that Bob can colour w from its list such that the common neighbour of v and w cannot be coloured. Now, suppose that in the first move Alice colours a vertex v of $G - K$. Then Bob's move is $B_1 := [f(x_1) = 1]$. This leaves the path $x_3x_2x_4$ with the reduced lists $\{3\}, \{3, 4\}, \{4\}$, which is not list-colourable, no matter how Alice and Bob play afterward.




Figure 2. The 2-list-assignment of $K_{2,3}$




Figure 3. The 2-list-assignment of H_2

Suppose that G contains an induced subgraph H isomorphic to H_2 . Let us denote the vertices of H and its 2-list-assignment L as in Figure 3 and let $L(v) = \{3, 4\}$ for $v \in V(G) - V(H)$. Assume that in the first move Alice colours a vertex of H . We may assume that Alice colours neither x_1 nor x_3 , otherwise Bob can make a move after which the vertex x_2 cannot be coloured. Suppose that Alice chooses the vertex x_2 . If she puts $A_1 := [f(x_2) = 1]$ then Bob answers $B_1 := [f(x_4) = 3]$. After such moves the vertex x_5 cannot be coloured from its list. If $A_1 := [f(x_2) = 2]$ then $B_1 := [f(x_4) = 1]$. Thus the vertex x_3 cannot be coloured. Similarly we can show that in the first move Alice cannot choose to colour the vertex x_4 . If in the first move Alice colours x_5 with colour 1 then Bob colours the vertex x_1 with 2. If Alice puts $A_1 := [f(x_5) = 3]$ then Bob's move is $B_1 := [f(x_3) = 1]$. Hence a third colour is necessary to colour the vertex x_4 . If in the first move Alice colours a vertex of $G - H$ then Bob's move is $B_1 := [f(x_1) = 1]$. If in the second move Alice does not colour the vertex x_2 or x_3 then the second Bob's move is $B_2 := [f(x_3) = 2]$. Hence the vertex x_2 cannot be coloured, but if the second move of Alice is $A_2 := [f(x_2) = 2]$ then Bob chooses $B_2 := [f(x_4) = 1]$. If $A_2 := [f(x_3) = 1]$ then Bob chooses $B_2 := [f(x_5) = 3]$. In both cases after Bob's second move there is a vertex in $V(H)$, which cannot be coloured. ■

The next corollary is an immediate consequence of Lemma 2.

Corollary 1 *If $\text{ch}_g(G) = 2$ then each component of G is isomorphic to P_4 or C_4 , or S_k for some $k \geq 1$.*

Let in the process of the game list colouring the graph G be partially coloured. We introduce the following notation. Let C be the set of coloured vertices and U be the set of uncoloured vertices. For any vertex v we will denote $N_C(v) = N(v) \cap C$ and $N_U(v) = N(v) \cap U$. We will say that the vertex x is *safe* if x is uncoloured and $|L(x) - \bigcup_{u \in N_C(x)} f(u)| > |N_U(x)|$.

The definition immediately yields the following

Proposition 3 *Suppose that in the process of the game list colouring a graph G is partially coloured. If all uncoloured vertices of G are safe then Alice wins.*

Now we will consider a game list colouring but with the first move of Bob. The rest principles of the game are the same: Two players Alice and Bob play alternately but Bob has the first move. Alice's goal is to provide a proper list colouring of the graph and Bob's goal is to prevent her from doing so. We shall refer to this version as *B-game list colouring*. The original version, i.e. Alice moving first, will sometimes be called the *A-game*.

If Alice has a winning strategy on H in the B-game then in some situations she can use it for finding a winning strategy on a larger graph G , which contains H such that $|V(G) \setminus V(H)|$ is odd. Let k be an integer and G be a graph such that $G = G_1 \cup G_2$ and $|V(G_1)|$ is odd. If Alice has a winning strategy in the A-game on G_1 with k colours and she also has a winning strategy with k colours in the B-game on G_2 then she can obtain a winning strategy on G in the following way: First Alice marks the vertex of G_1 according to her winning strategy on G_1 . After that she plays on G_i ($i = 1$ or $i = 2$) if and only if Bob has played on G_i in his previous move. In each move she follows her winning strategy in the A-game or B-game, according as $i = 1$ or $i = 2$.

Lemma 3 *Alice has a winning strategy in the B-game list colouring on P_4 .*

Proof. Let $V(P_4) = \{x_1, x_2, x_3, x_4\}$ and $E(P_4) = \{x_1x_2, x_2x_3, x_3x_4\}$. Bob has the first move.

Suppose that Bob's move is $B_1 := [f(x_1) = a]$. Alice chooses vertex x_3 . If $L(x_2) \neq L(x_3)$ then Alice colours the vertex x_3 with the colour from $L(x_3)$ which is not in $L(x_2)$. After this move all vertices of P_4 are safe; then by Proposition 3 Alice wins. Now assume that $L(x_2) = L(x_3)$. If $a \in L(x_3)$ then $A_1 := [f(x_3) = a]$. Otherwise Alice can colour x_3 with any colour of $L(x_3)$ and after her move all vertices of P_4 are safe.

If Bob's move is $B_1 := [f(x_2) = a]$ then Alice colours x_3 using another colour than Bob did. Then all vertices of P_4 are safe. The lemma follows. ■

Since $\text{ch}_g(P_4) = 3$, we also have

Corollary 2 *If $\text{ch}_g(G) = 2$ and $P_4 \leq G$ then $|V(G) - V(P_4)|$ is odd.*

Lemma 4 *Let (C_4, L) be the 4-cycle with the 2-list-assignment L as in Figure 4 or Figure 5. Then Alice has a winning strategy on (C_4, L) in the A-game and Bob has a winning strategy on (C_4, L) in the B-game.*

Proof. First let us consider A-game, Alice has the first move and $A_1 := [f(x_2) = a]$. If Bob colours the vertex x_1 or x_4 then after his move all vertices of the graph are safe. If Bob colours the vertex x_3 using the colour b or c then Alice colours x_1 with d , thus after her move the vertex x_4 can be coloured with a .

Now assume that Bob starts. It is easy to see that Bob can win, as in the first move he can colour the vertex x_1 using colour a and then the rest is uncolourable. ■




Figure 4.




Figure 5.

Corollary 3 *If $\text{ch}_g(G) = 2$ then G does not contain two vertex-disjoint C_4 s.*

Let G be a graph with a subgraph isomorphic to C_4 and with a 2-list-assignment L . From Lemma 4 it follows that there are 2-list-assignments of C_4 such that in a winning strategy of Alice, she must start choosing a vertex of C_4 . But there are also 2-list-assignments such that Alice wins if and only if in the process of the game list colouring Bob first colours a vertex of C_4 .

Lemma 5 *Let L be a 2-list-assignment of C_4 , other than the lists in Figure 4 and Figure 5. Then Alice has a winning strategy in the B-game list colouring on (C_4, L) .*

Proof. Let $V(C_4) = \{x_1, x_2, x_3, x_4\}$, $E(C_4) = \{x_1x_2, x_2x_3, x_3x_4, x_4x_1\}$. Without loss of generality assume that Bob chooses x_1 and puts $B_1 := [f(x_1) = a]$. If $a \in L(x_3)$ then Alice colours $A_1 := [f(x_3) = a]$. Hence all vertices are safe and Alice wins. Then assume that $a \notin L(x_3)$ and $L(x_3) = \{b, c\}$. If neither $A_2 := [f(x_3) = b]$ nor $A_2 := [f(x_3) = c]$ leaves both x_2 and x_4 colourable (i.e., if Alice cannot win by colouring x_3), then their lists must be $L(x_2) = \{a, b\}$ and $L(x_4) = \{a, c\}$, or vice versa. Thus, the list assignment is isomorphic to the one in Figure 4 or 5, depending on whether the second element in $L(x_1)$ is a new colour or one of b and c . ■

Corollary 4 *If $\text{ch}_g(G) = 2$ and $C_4 \leq G$ then $|V(G) - V(C_4)|$ is odd.*

Since Alice has a winning strategy on any star forest in both the A-game and B-game list colourings, we obtain the main result of this section.

Theorem 4 *Let G be a graph with at least one edge. Then $\text{ch}_g(G) = 2$ if and only if G is a star forest or $G = kP_4 \cup tC_4 \cup H$, where $k \geq 0$, $t \in \{0, 1\}$ and H is a star forest of odd order.*

References

- [1] H.L. Bodlaender, *On the complexity of some coloring games*, Internat. J. Found. Comput. Sci. **2** (1991) 133–147.
- [2] M. Borowiecki, E. Drgas-Burchardt and P. Mihók *Generalized list colourings of graphs*, Discussiones Math. Graph Theory **15** (1995) 185–193.
- [3] R.L. Brooks, *On colouring the nodes of a network*, Proc. Cambridge Phil. Soc. **37** (1941) 194–197.
- [4] L. Cai and X. Zhu, *Game chromatic index of k -degenerate graphs*, J. Graph Theory **36** (2001) 144–155.
- [5] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Second Edition, (Wadsworth & Brooks/Cole, Monterey, 1986).
- [6] P. Erdős, A.L. Rubin and H. Taylor, *Choosability in graphs*, in: Proc. West Coast Conf. on Combin., Graph Theory and Computing, Congressus Numerantium XXVI (1979) 125–157.
- [7] U. Faigle, U. Kern, H.A. Kierstead and W.T. Trotter, *On the game chromatic number of some classes of graphs*, Ars Combinatoria **35** (1993) 143–150.
- [8] T. Gallai, *Kritische Graphen II*, Magyar Tud. Akad. Mat. Kut. Int. Közl. **8** (1963) 373–395.
- [9] D.J. Guan and X. Zhu, *Game chromatic number of outerplanar graphs*, J. Graph Theory **30** (1999) 67–70.
- [10] T.R. Jensen and B. Toft, *Graph Colouring Problems*, (Wiley, New York, 1995).
- [11] H.A. Kierstead, *A simple competitive graph coloring algorithm*, J. Combin. Theory B **78** (2000) 57–68.
- [12] H.A. Kierstead and W.T. Trotter, *Planar graph coloring with an uncooperative partner*, J. Graph Theory **18** (1994) 569–584.
- [13] H.A. Kierstead and Zs. Tuza, *Marking games and the oriented game chromatic number of partial k -trees*, Graphs and Combinatorics **19** (2003) 121–129.
- [14] A.V. Kostochka, M. Stiebitz and B. Wirth, *The colour theorems of Brooks and Gallai extended*, Discrete Math. **162** (1996) 299–303.
- [15] J. Nešetřil and E. Sopena, *On the oriented game chromatic number*, Electronic J. Combin. **8** (No. 2) (2001) #R14.
- [16] C. Thomassen, *Color-critical graphs on a fixed surface*, J. Combin. Theory B **70** (1997) 67–100.
- [17] Zs. Tuza, *Graph colorings with local constraints — A survey*, Discuss. Math. Graph Theory **17** (1997) 161–228.
- [18] V.G. Vizing, *Colouring the vertices of a graph in prescribed colours*, Diskret. Analiz **29** (1976) 3–10 (in Russian).