

# The number of 0-1-2 increasing trees as two different evaluations of the Tutte polynomial of a complete graph

C. Merino\*

Instituto de Matemáticas,  
Universidad Nacional Autónoma de México,  
Circuito Exterior, C.U. Coyoacán 04510, México D.F.  
`merino@matem.unam.mx`

Submitted: Nov 21, 2007; Accepted: Jul 11, 2008; Published: Jul 21, 2008  
Mathematics Subject Classifications: 05A19

## Abstract

If  $T_n(x, y)$  is the Tutte polynomial of the complete graph  $K_n$ , we have the equality  $T_{n+1}(1, 0) = T_n(2, 0)$ . This has an almost trivial proof with the right combinatorial interpretation of  $T_n(1, 0)$  and  $T_n(2, 0)$ . We present an algebraic proof of a result with the same flavour as the latter:  $T_{n+2}(1, -1) = T_n(2, -1)$ , where  $T_n(1, -1)$  has the combinatorial interpretation of being the number of 0-1-2 increasing trees on  $n$  vertices.

## 1 Introduction

Given a graph  $G = (V, E)$ , we define the *rank function* of  $G$ ,  $r : \mathcal{P}(E) \rightarrow \mathbb{Z}$  as  $r(A) = |V| - k(A)$  for  $A \subseteq E$ , where  $k(A)$  is the number of connected components in the graph  $(V, A)$ . The 2-variable graph polynomial  $T(G; x, y)$ , known as the *Tutte polynomial* of  $G$ , is defined as

$$T(G; x, y) = \sum_{A \subseteq E} (x - 1)^{r(E) - r(A)} (y - 1)^{|A| - r(A)}. \quad (1)$$

The Tutte polynomial of  $G$  has many interesting combinatorial interpretations when evaluated on different points  $(x, y)$  and along several algebraic curves. One that is particularly interesting is along the line  $x = 1$  which can be interpreted as the generating function of critical configuration of the sandpile model, see [8], or as the generating function of the  $G$ -parking functions, see [9]. When the graph  $G$  is the complete graph on  $n$  vertices,  $K_n$ , the latter is the classical generating function of parking functions or the inversion enumerator of labelled trees on  $n$  vertices, see [10].

In the following section we prove the main theorem of the paper:

---

\*Supported by Conacyt of México.

**Theorem 1.**  $T(K_n; 2, -1) = T(K_{n+2}; 1, -1)$ .

The last section shows how this result is related to the number of 0-1-2 increasing trees on  $n$  vertices.

## 2 $T(K_n; 2, -1)$ and $T(K_{n+2}; 1, -1)$

Let us assume that the vertices of  $K_n$  are labelled  $1, 2, \dots, n$ . For a spanning tree  $A$  of  $K_n$ , an *inversion* in  $A$  is a pair of vertices labelled  $i, j$  such that  $i > j$  and  $i$  is on the unique path from 1 to  $j$  in  $A$ . Let  $\text{inv}A$  be the number of inversions in  $A$ . The *inversion enumerator*  $J_n(y)$  is then defined as the generating function of spanning trees arranged by number of inversions, that is,

$$J_n(y) = \sum_A y^{\text{inv}A},$$

where the sum is taken over all spanning trees of  $K_n$ . Now, from [10], we obtain the exponential generating function of the inversion enumerators,

$$\sum_{n \geq 0} J_{n+1}(y)(y-1)^n \frac{t^n}{n!} = \frac{\sum_{n \geq 0} y^{\binom{n+1}{2}} \frac{t^n}{n!}}{\sum_{n \geq 0} y^{\binom{n}{2}} \frac{t^n}{n!}}. \quad (2)$$

Note that our notation differs from [10], as Stanley uses  $I_n(y)$  for  $J_{n+1}(y)$ .

Let  $T_n(x, y)$  be the Tutte polynomial of  $K_n$ . Welsh in [11] gives the following exponential generating function for  $T_n(x, y)$

$$1 + (x-1) \sum_{n \geq 1} (y-1)^n T_n(x, y) \frac{t^n}{n!} = \left( \sum_{n \geq 0} y^{\binom{n}{2}} \frac{t^n}{n!} \right)^{(x-1)(y-1)} \quad (3)$$

With these two general results it is easy to prove the following:

**Theorem 2.** For  $n \geq 0$ ,  $J_{n+2}(-1) = T_n(2, -1)$ .

*Proof.* By taking  $y = -1$  in Equation (2) we get

$$\sum_{n \geq 0} J_{n+1}(-1)(-2)^n \frac{t^n}{n!} = \frac{\sum_{n \geq 0} (-1)^{\binom{n+1}{2}} \frac{t^n}{n!}}{\sum_{n \geq 0} (-1)^{\binom{n}{2}} \frac{t^n}{n!}} = \frac{F(t)}{H(t)}.$$

Clearly,  $F(t) = H'(t)$ , where  $H'(t)$  is the derivative of  $H(t)$ . Then, by integrating both sides of the previous expression and multiplying through by  $-2$  we arrive at the equality

$$\sum_{n \geq 1} J_n(-1)(-2)^n \frac{t^n}{n!} = (-2) \ln |H(t)|.$$

The function  $H(t)$  is the exponential generating function of the sequence 1, 1, -1, -1, 1, 1, -1, -1, ..., so  $H(t) = \cos(t) + \sin(t)$ . Substituting this value on the above identity we obtain

$$\sum_{n \geq 1} J_n(-1)(-2)^n \frac{t^n}{n!} = (-2) \ln |\cos(t) + \sin(t)|. \quad (4)$$

Now, by differentiating twice both sides of equation (4) we conclude that

$$\sum_{n \geq 0} J_{n+2}(-1)(-2)^n \frac{t^n}{n!} = \frac{1}{(\cos(t) + \sin(t))^2}. \quad (5)$$

Taking  $x = 2$  and  $y = -1$  in Equation (3), we get the following identities

$$1 + \sum_{n \geq 1} (-2)^n T_n(2, -1) \frac{t^n}{n!} = \left( \sum_{n \geq 0} (-1)^{\binom{n}{2}} \frac{t^n}{n!} \right)^{-2} = \frac{1}{(\cos(t) + \sin(t))^2}. \quad (6)$$

Therefore, from Equations (5) and (6),

$$1 + \sum_{n \geq 1} T_n(2, -1) \frac{(-2)^n t^n}{n!} = \sum_{n \geq 0} J_{n+2}(-1) \frac{(-2)^n t^n}{n!}.$$

As  $T_0(2, -1) = 1$ , we obtain the result by equating the corresponding coefficients.  $\square$

It is known that  $T_n(1, y) = J_n(y)$ , see [7]. Thus, Theorem 1 follows by the previous result.

A permutation  $\sigma \in S_n$  is an *up-down permutation* if  $\sigma(1) < \sigma(2) > \sigma(3) < \dots$ . Let  $a_n$  be the number of up-down permutation in  $S_n$  for  $n \geq 1$  and set  $a_0 = 1$ . The sequence  $a_n$  has a nice exponential generating function, namely

$$\sum_{n \geq 0} a_n \frac{t^n}{n!} = \tan(t) + \sec(t).$$

The result is originally from [1] but a proof may also be found in [7]. The fact that the value  $J_{n+1}(-1)$  equals  $a_n$  is mentioned in [6] but a proof of this together with other evaluations of  $J_n(x)$  is given in [7]. As a corollary we obtain

**Corollary 3.** For  $n \geq 0$ ,  $T_n(2, -1) = a_{n+1}$  and

$$\sum_{n \geq 0} T_n(2, -1) \frac{t^n}{n!} = \sec(t)(\tan(t) + \sec(t)).$$

### 3 The Tutte polynomial and increasing trees

A spanning tree in  $K_n$  with root at 1 is said to be *increasing* whenever its vertices increase along the paths away from the root. A *0–1–2 increasing tree* is an increasing tree where all the vertices have at most 2 edges going out. A remarkable result stated in [4] and proved in [5] (see also a bijective proof in [3]) is that  $a_n$  equals the number of 0–1–2 increasing trees on  $n$  vertices. By using Corollary 3 we get

**Corollary 4.**  $T_n(2, -1)$  equals the number of 0–1–2 increasing trees on  $n + 1$  vertices.

Thus, the number of 0–1–2 increasing trees on  $n$  vertices corresponds two different evaluations of the Tutte polynomial of a complete graph, that is  $T_{n-1}(2, -1)$  and  $T_{n+1}(1, -1)$ .

A similar situation occur for the number of permutations on  $n$  letters. The quantity  $T(G; 2, 0)$  equals the number of acyclic orientations of  $G$  while  $T(G; 1, 0)$  equals the number of acyclic orientations of  $G$  with a unique predefined source, see [2]. If we use this combinatorial interpretation with  $K_n$ , clearly we get that  $T_{n+1}(1, 0) = T_n(2, 0)$ . In fact, it is easy to find the exact values,  $T_n(2, 0) = n!$  and  $T_n(1, 0) = n - 1!$ . That is, the number of permutations on  $n$  letters occurs as two different evaluations of the Tutte polynomial of a complete graph,  $T_n(2, 0)$  and  $T_{n+1}(1, 0)$ .

Increasing spanning trees correspond to spanning trees with no inversions. Thus,  $J_n(0) = T_n(1, 0)$  equals the number of increasing trees in  $K_n$ . By deleting the vertex 1 in  $K_{n+1}$  we get a bijection between increasing trees in  $K_{n+1}$  and increasing spanning forests in  $K_n$ . Here a forest is increasing if it is increasing in each component. Therefore, we get the interpretation of  $T_n(2, 0)$  as the number of increasing spanning forests in  $K_n$ .

Using the same technique we get a bijection between 0–1–2 increasing trees on  $n + 1$  vertices and 0–1–2 increasing forests on  $n$  vertices with at most 2 components. Thus we get

**Corollary 5.**  $T_n(2, -1)$  equals the number of 0–1–2 increasing forests on  $n$  vertices with at most 2 components.

There are several combinatorial interpretations for evaluations of  $T(G; x, y)$  when  $x, y \geq 0$ , and even when  $x, y \leq 0$  probably because of the relationship of the Tutte polynomial with the partition function of the Potts model of statistical mechanics. But the situation is quite different when  $y < 0 < x$  or  $x < 0 < y$ . I would like to think that Corollary 5 is just the tip of the iceberg and that more combinatorial interpretations for  $T(G; x, y)$  in these regions exist.

## References

- [1] André, D.: Développements de  $\sec x$  et de  $\tan x$ . C. R. Acad. Sc. Paris, **88**, 965–967, 1879.
- [2] Brylawski, T. and Oxley, J.: The Tutte Polynomial and its Applications. In: White, N. (ed) Matroid Applications. Cambridge University Press, Cambridge, 123–225, 1992.

- [3] Donaghey, R.: Alternating permutations and binary increasing trees. *J. Combinatorial Theory Ser. A*, **18**, 141–148, 1975.
- [4] Foata, D.: Groupes de réarrangements et nombres d'Euler. *C. R. Acad. Sci. Paris Sr. A-B*, **275**, A1147–A1150, 1972.
- [5] Foata, D. and Strehl, V.: Rearrangements of the symmetric group and enumerative properties of the tangent and secant numbers. *Math. Z.*, **137**, 257–264, 1974.
- [6] Goulden, I. P. and Jackson, D. M.: *Combinatorial Enumeration*. Wiley, Chichester 1983.
- [7] Kuznetsov, A. G., Pak, I. M. and Postnikov, A. E.: Increasing trees and alternating permutations. (Russian) *Uspekhi Mat. Nauk*, **49**, 79–110, 1994; translation in *Russian Math. Surveys*, **49**, 79–114, 1994.
- [8] Merino, C.: Chip-firing and the Tutte polynomial. *Annals of Combinatorics*, **1**, 253–259, 1997.
- [9] Plautz J. and Calderer, R.:  $G$ -parking functions and the Tutte polynomial. Preprint.
- [10] Stanley, R. P.: Hyperplane arrangements, parking functions and tree inversions. In: Sagan, B. and Stanley, R. (eds) *Mathematical Essays in Honor of Gian-Carlo Rota*. Birkhäuser, Boston, Basel, 359–375, 1998.
- [11] D. J. A. Welsh, Counting colourings and flows in random graphs. In: Miklós, D., Sos, V. T. and Szönyi, T. (eds) *Combinatorics, Paul Erdős is Eighty*. Janos Bolyai Math. Soc., Budapest, 491–505, 1996.