

# For each $\alpha > 2$ there is an infinite binary word with critical exponent $\alpha$

James D. Currie\* & Narad Rampersad†

Department of Mathematics and Statistics  
University of Winnipeg  
Winnipeg, Manitoba R3B 2E9  
CANADA

e-mail: j.currie@uwinnipeg.ca,  
n.rampersad@uwinnipeg.ca

Submitted: Feb 28, 2008; Accepted: Aug 25, 2008; Published: Aug 31, 2008

Mathematics Subject Classification: 68R15

## Abstract

The critical exponent of an infinite word  $\mathbf{w}$  is the supremum of all rational numbers  $\alpha$  such that  $\mathbf{w}$  contains an  $\alpha$ -power. We resolve an open question of Krieger and Shallit by showing that for each  $\alpha > 2$  there is an infinite binary word with critical exponent  $\alpha$ .

Keywords: Combinatorics on words, repetitions, critical exponent

## 1 Introduction

If  $\alpha$  is a rational number, a word  $w$  is an  $\alpha$ -power if there exist words  $x$  and  $x'$  and a positive integer  $n$ , with  $x'$  a prefix of  $x$ , such that  $w = x^n x'$  and  $\alpha = n + |x'|/|x|$ . We refer to  $|x|$  as a *period* of  $w$ . A word is  $\alpha$ -power-free if none of its subwords is a  $\beta$ -power with  $\beta \geq \alpha$ ; otherwise, we say the word *contains an  $\alpha$ -power*.

The *critical exponent* of an infinite word  $\mathbf{w}$  is defined as

$$\sup\{\alpha \in \mathbb{Q} \mid \mathbf{w} \text{ contains an } \alpha\text{-power}\}.$$

Critical exponents of certain classes of infinite words, such as Sturmian words [8, 10] and words generated by iterated morphisms [5, 6], have received particular attention.

Krieger and Shallit [7] proved that for every real number  $\alpha > 1$ , there is an infinite word with critical exponent  $\alpha$ . As  $\alpha$  tends to 1, the number of letters required to construct

---

\*The author's research was supported by an NSERC operating grant.

†The author is supported by an NSERC Post-doctoral Fellowship.

such words tends to infinity. However, for  $\alpha > 7/3$ , Shur [9] gave a construction over a binary alphabet. For  $\alpha > 2$ , Krieger and Shallit gave a construction over a four-letter alphabet and left it as an open problem to determine if for every real number  $\alpha \in (2, 7/3]$ , there is an infinite binary word with critical exponent  $\alpha$ . Currie, Rampersad, and Shallit [3] gave examples of such words for a dense subset of real numbers  $\alpha$  in the interval  $(2, 7/3]$ . In this note we resolve the question completely by demonstrating that for every real number  $\alpha > 2$ , there is an infinite binary word with critical exponent  $\alpha$ .

## 2 Properties of the Thue-Morse morphism

In this section we present some useful properties of the *Thue-Morse morphism*; i.e., the morphism  $\mu$  defined by  $\mu(0) = 01$  and  $\mu(1) = 10$ . Note that  $|\mu^s(0)| = |\mu^s(1)| = 2^s$  for all  $s \geq 0$ .

**Lemma 1.** *Let  $s$  be a positive integer. Let  $z$  be a subword of  $\mu^{s+1}(0) = \mu^s(01)$  with  $|z| \geq 2^s$ . Then  $z$  does not have period  $2^s$ .*

*Proof.* Write  $\mu^s(0) = a_1a_2 \dots a_{2^n}$ ,  $\mu^s(1) = b_1b_2 \dots b_{2^n}$ . One checks by induction that  $a_i = 1 - b_i$  for  $1 \leq i \leq 2^n$ , and the result follows.  $\square$

Brandenburg [1] proved the following useful theorem, which was independently rediscovered by Shur [9].

**Theorem 2 (Brandenburg; Shur).** *Let  $w$  be a binary word and let  $\alpha > 2$  be a real number. Then  $w$  is  $\alpha$ -power-free if and only if  $\mu(w)$  is  $\alpha$ -power-free.*

The following sharper version of one direction of this theorem (implicit in [4]) is also useful.

**Theorem 3.** *Suppose  $\mu(w)$  contains a subword  $u$  of period  $p$ , with  $|u|/p > 2$ . Then  $w$  contains a subword  $v$  of length  $\lceil |u|/2 \rceil$  and period  $p/2$ .*

We will also have call to use the *deletion operator*  $\delta$  which removes the first (left-most) letter of a word. For example,  $\delta(12345) = 2345$ .

## 3 A binary word with critical exponent $\alpha$

We denote by  $\mathcal{L}$  the set of factors (subwords) of words of  $\mu(\{0, 1\}^*)$ .

**Lemma 4.** *Let  $00v \in \mathcal{L}$ , and suppose that  $00v$  is  $\alpha$ -power-free for some fixed  $\alpha > 2$ . Let  $r = \lceil \alpha \rceil$ . Suppose that  $0^r v = xy$  where  $u$  contains an  $\alpha$ -power. Then  $x = \epsilon$  and  $u = 0^r$ .*

*Proof.* Suppose that  $u$  has period  $p$ . Since  $00v \in \mathcal{L}$ ,  $v$  begins with 1. Since  $00v$  is  $\alpha$ -power-free, we can write  $u = 0^s v'$ , where  $x = 0^{r-s}$  for some integer  $s$ ,  $3 \leq s \leq r$ , and  $v'$  is a prefix of  $v$ . If  $0^p$  is not a prefix of  $u$  then the prefix of  $u$  of length  $p$  contains the

subword 0001. Since  $\alpha > 2$ , this means that 0001 is a subword of  $u$  at least twice, so that 0001 is a subword of  $00v$ . This is impossible, since  $00v \in \mathcal{L}$ .

Therefore,  $0^p$  is a prefix of  $u$ , and  $u$  has the form  $0^t$  for some integer  $t \geq \alpha$ . This implies that  $u$  has  $0^r$  as a prefix, so that  $x = \epsilon$  and  $u = 0^r$ .  $\square$

**Lemma 5.** *Let  $\alpha > 2$  be given, and let  $r = \lceil \alpha \rceil$ . Let  $s, t$  be positive integers, such that  $s \geq 3$  and there are words  $x, y \in \{0, 1\}^*$  such that  $\mu^s(0) = x00y$  with  $|x| = t$ . Suppose that  $2 < r - t/2^s < \alpha$  and  $00v \in \mathcal{L}$  is  $\alpha$ -power-free. Then the following statements hold.*

1. *The word  $\delta^t \mu^s(0^r v)$  has a prefix which is a  $\beta$ -power, where  $\beta = r - t/2^s$ .*
2. *Suppose that  $00v$  contains a  $\beta$ -power of period  $p$  for some  $\beta$  and  $p$ . Then  $\delta^t \mu^s(0^r v)$  contains a  $\beta$ -power of period  $2^s p$ .*
3. *The word  $\delta^t \mu^s(0^r v)$  is  $\alpha$ -power-free.*

*Proof.* We start by observing that  $\mu^s(0^r)$  has period  $2^s$ . It follows that  $\delta^t \mu^s(0^r)$  is a word of length  $r2^s - t$  with period  $2^s$ , and hence is a  $(r2^s - t)/2^s = \beta$ -power.

Now suppose  $u$  is a  $\beta$ -power of period  $p$  in  $00v$ . Then  $\mu^s(u)$  is a  $\beta$ -power of period  $2^s p$  in  $\mu^s(00v)$ . However,  $\mu^s(0^{r-1}v)$  is a suffix of  $\delta^t \mu^s(0^r v)$ , since  $t < 2^s = |\mu^s(0)|$ . Thus  $\mu^s(u)$  is a  $\beta$ -power of period  $2^s p$  in  $\delta^t \mu^s(0^r v)$ .

Next, note that  $\mu^s(0^{r-1}v)$  does not contain any  $\kappa$ -power,  $\kappa \geq \alpha$ . Otherwise, by Theorem 3 and induction,  $0^{r-1}v$  contains a  $\kappa$ -power. This is impossible by Lemma 4.

Suppose then that  $\delta^t \mu^s(0^r v)$  contains a  $\kappa$ -power  $\hat{u}$  of period  $q$ ,  $\kappa \geq \alpha$ . Using induction and Theorem 3,  $0^r v$  contains a  $\kappa$ -power  $u$  of period  $q/2^s$ . By Lemma 4, the only possibility is  $u = 0^r$ , and  $q/2^s = 1$ . Thus  $q = 2^s$ .

Since  $00v \in \mathcal{L}$ , the first letter of  $v$  is a 1. Since  $\hat{u}$  has period  $2^s$ , by Lemma 1 no subword of  $\mu^s(01)$  of length greater than  $2^s$  occurs in  $\hat{u}$ . We conclude that either  $\hat{u}$  is a subword of  $\delta^t \mu^s(0^r)$ , or of  $\mu^s(v)$ , and hence of  $\mu^s(0^{r-1}v)$ . As this second case has been ruled out earlier, we conclude that  $|\hat{u}| \leq |\delta^t \mu^s(0^r)| = r2^s - t$ . This gives a contradiction:  $\hat{u}$  is a  $\kappa$ -power, yet  $|\hat{u}|/q \leq (r2^s - t)/2^s = \beta < \alpha$ .  $\square$

By construction,  $\delta^t \mu^s(0^r v)$  has the form  $00\hat{v}$  where  $00\hat{v} \in \mathcal{L}$ .

We are now ready to prove our main theorem:

**Theorem 6.** *Let  $\alpha > 2$  be a real number. There is a word over  $\{0, 1\}$  with critical exponent  $\alpha$ .*

*Proof.* Call a real number  $\beta < \alpha$  *obtainable* if  $\beta$  can be written  $\beta = r - t/2^s$ , where  $r, s, t$  are positive integers,  $s \geq 3$ , and the word obtained by removing a prefix of length  $t$  from  $\mu^s(0)$  begins with 00. We note that  $\mu^3(0) = 01101001$  and  $\mu^3(1) = 10010110$  are of length 8, and both contain 00 as a subword; for a given  $s \geq 3$  it follows that  $r$  and  $t$  can be chosen so that  $\beta = r - t/2^s < \alpha$  and  $|\alpha - \beta| \leq 7/2^s$ ; by choosing large enough  $s$ , an obtainable number  $\beta$  can be chosen arbitrarily close to  $\alpha$ .

Let  $\{\beta_i\}$  be a sequence of obtainable numbers converging to  $\alpha$ . For each  $i$  write  $\beta_i = r_i - t_i/2^{s_i}$ , where  $r_i, s_i, t_i$  are positive integers,  $s_i \geq 3$ , and the word obtained by

removing a prefix of length  $t_i$  from  $\mu^{s_i}(0)$  begins with  $00$ . If  $00w \in \mathcal{L}$ , denote by  $\phi_i(w)$  the word  $\delta^{t_i}\mu^{s_i}(0^r w)$ .

Consider the sequence of words

$$\begin{aligned} w_1 &= \phi_1(\epsilon) \\ w_2 &= \phi_1(\phi_2(\epsilon)) \\ w_3 &= \phi_1(\phi_2(\phi_3(\epsilon))) \\ &\vdots \\ w_n &= \phi_1(\phi_2(\phi_3(\cdots(\phi_n(\epsilon))\cdots))) \\ &\vdots \end{aligned}$$

By the third part of Lemma 5, if  $00w \in \mathcal{L}$  is  $\alpha$ -power-free, then so is  $\phi_i(w)$ . Since  $00\epsilon$  is  $\alpha$ -power-free, each  $w_i$  is therefore  $\alpha$ -power-free.

By the first and second parts of Lemma 5,  $w_n$  contains  $\beta_i$ -powers,  $i = 1, 2, \dots, n$ .

Note that  $\epsilon$  is a prefix of  $\phi_{n+1}(\epsilon)$ , so that

$$w_n = \phi_1(\phi_2(\phi_3(\cdots(\phi_n(\epsilon))\cdots)))$$

is a prefix of

$$\phi_1(\phi_2(\phi_3(\cdots(\phi_n(\phi_{n+1}(\epsilon))\cdots))) = w_{n+1}.$$

We may therefore let  $w = \lim_{n \rightarrow \infty} w_n$ .

Since every prefix of  $w$  is  $\alpha$ -power-free,  $w$  is  $\alpha$ -power-free but contains  $\beta_i$ -powers for each  $i$ . The critical exponent of  $w$  is therefore  $\alpha$ .  $\square$

The following question raised by Krieger and Shallit remains open: for  $\alpha > 1$ , if  $\alpha$ -powers are avoidable on a  $k$ -letter alphabet, does there exist an infinite word over  $k$  letters with critical exponent  $\alpha$ ? In particular, for  $\alpha > \text{RT}(k)$ , where  $\text{RT}(k)$  denotes the *repetition threshold* on  $k$  letters (see [2]), does there exist an infinite word over  $k$  letters with critical exponent  $\alpha$ ? We believe that the answer is “yes”.

## Acknowledgments

We would like to thank the anonymous referee for helpful comments and suggestions.

## References

- [1] F.-J. Brandenburg, “Uniformly growing  $k$ -th power-free homomorphisms”, *Theoret. Comput. Sci.* **23** (1983), 69–82.
- [2] A. Carpi, “On Dejean’s conjecture over large alphabets”, *Theoret. Comput. Sci.* **385** (2007), 137–151.

- [3] J.D. Currie, N. Rampersad, J. Shallit, “Binary words containing infinitely many overlaps”, *Electron. J. Combin.* **13** (2006), #R82.
- [4] J. Karhumäki, J. Shallit, “Polynomial versus exponential growth in repetition-free binary words”, *J. Combin. Theory Ser. A* **104** (2004), 335–347.
- [5] D. Krieger, “On critical exponents in fixed points of binary  $k$ -uniform morphisms”. In *Proc. STACS 2006*, LNCS 3884, Springer-Verlag, 2006, pp. 104–114.
- [6] D. Krieger, “On critical exponents in fixed points of non-erasing morphisms”, *Theoret. Comput. Sci.* **376** (2007), 70–88.
- [7] D. Krieger, J. Shallit, “Every real number greater than 1 is a critical exponent”, *Theoret. Comput. Sci.* **381** (2007), 177–182.
- [8] F. Mignosi, G. Pirillo, “Repetitions in the Fibonacci infinite word”, *RAIRO Inform. Theor. Appl.* **26** (1992), 199–204.
- [9] A. M. Shur, “The structure of the set of cube-free  $\mathbb{Z}$ -words in a two-letter alphabet” (Russian), *Izv. Ross. Akad. Nauk Ser. Mat.* **64** (2000), 201–224. English translation in *Izv. Math.* **64** (2000), 847–871.
- [10] D. Vandeth, “Sturmian words and words with a critical exponent”, *Theoret. Comput. Sci.* **242** (2000), 283–300.