

Induced trees in triangle-free graphs

JIRÍ MATOUŠEK ROBERT ŠÁMAL*

Department of Applied Mathematics and
Institute of Theoretical Computer Science (ITI)
Charles University
Malostranské nám. 25, 118 00 Praha 1
Czech Republic

Submitted: Nov 29, 2007; Accepted: Feb 24, 2008; Published: Mar 12, 2008
Mathematics Subject Classification: 05C55, 05C05

Abstract

We prove that every connected triangle-free graph on n vertices contains an induced tree on $\exp(c\sqrt{\log n})$ vertices, where c is a positive constant. The best known upper bound is $(2 + o(1))\sqrt{n}$. This partially answers questions of Erdős, Saks, and Sós and of Pultr.

1 Introduction

For a graph G , let $t(G)$ denote the maximum number of vertices of an induced subgraph of G that is a tree (i.e., connected and acyclic). There are arbitrary large graphs G with $t(G) \leq 2$, namely graphs in which every connected component is a clique. To rule out these trivial examples, we need to put some restrictions on G .

Motivated by study of forbidden configurations in Priestley spaces [1], Pultr (private communication, 2002) asked how big $t(G)$ can be if G is connected and bipartite. Formally, he was interested about asymptotic properties of the function

$$f_B(n) = \min\{t(G) : |V(G)| = n, G \text{ connected and bipartite}\}.$$

Pultr's question was the starting point of our work. However, the function $t(G)$ was studied earlier and in a more general context by Erdős, Saks, and Sós [2]. They describe the influence of the number of edges of G on $t(G)$ and, more to our point, they study how small $t(G)$ can be if $\omega(G)$ is given. They observe that $t(G) \leq 2\alpha(G)$, and this allows

*Currently on leave from Institute for Theoretical Computer Science (ITI). The paper was finished while the second author was a PIMS postdoctoral fellow at Department of Mathematics, Simon Fraser University, Burnaby, B.C. V5A 1S6, Canada.

them to use estimates for Ramsey numbers. This way, they show that for any fixed $k > 3$ there are constants c_1, c_2 such that

$$c_1 \frac{\log n}{\log \log n} \leq \min\{t(G) : |V(G)| = n, G \not\supseteq K_k\} \leq c_2 \log n.$$

For $k = 3$ the lower bound still applies, but the upper bound obtained by using Ramsey numbers was only $O(\sqrt{n} \log n)$ (nowadays this approach yields $O(\sqrt{n \log n})$, due to the improved lower bound on $R(k, 3)$, see [4]). We concentrate on this case $k = 3$, that is we put

$$f_T(n) = \min\{t(G) : |V(G)| = n, G \text{ connected and triangle-free}\}.$$

Instead of applying Ramsey theory, we approach the problem directly.

It is easy to show that $f_T(n) \leq f_B(n) = O(\sqrt{n})$. The best construction we are aware of yields $f_B(n) \leq (2 + o(1))\sqrt{n}$; see Section 2. A simple “blow-up” construction, also presented in Section 2, shows that if $f_T(n_0) < \sqrt{n_0}$ for some n_0 , then $f_T(n) = O(n^{1/2-\varepsilon})$ for a positive constant $\varepsilon > 0$, and similarly for f_B . Hence, $f_T(n)$ either is of order exactly \sqrt{n} , or it is bounded above by some power strictly smaller than $1/2$. We conjecture that the second possibility holds, and that another power of n is a lower bound.

Conjecture 1.1 *There are constants $0 < \alpha < \beta < 1/2$, and c_1, c_2 such that for all n*

$$c_1 n^\alpha \leq f_T(n) \leq f_B(n) \leq c_2 n^\beta.$$

The following lower bound is the main result of this paper.

Theorem 1.2 *There is a constant $c > 0$ such that for all n*

$$f_T(n) \geq e^c \sqrt{\log n}.$$

We finish the introduction by mentioning further results concerning $t(G)$. It is interesting to consider the problem of finding induced trees in (sparse) random graphs. Vega [3] shows that $t(G_{n,c/n}) = \Omega(n)$ a.s.; Palka and Ruciński [6] prove that $t(G_{n,c \log n/n}) = \Theta(n \log \log n / \log n)$ a.s.

Krishnan and Ochem [5] search for values of $f_T(n)$ (for small n) using a computer; they succeed to find $f_T(n)$ for $n \leq 15$. They also extend results of [2] about the decision problem: “given a connected graph G and an integer t , does G have an induced tree with t vertices?”. Not only this is NP-complete for general graphs (which is proved in [2]), but it remains NP-complete even if we restrict to bipartite graphs, or to triangle-free graphs of maximum degree 4.

2 Initial observations

Observation 2.1 $f_B(n) \leq (2 + o(1))\sqrt{n}$.

Proof: It is enough to take a path with each edge replaced by a complete bipartite graph. More precisely, we take pairwise disjoint sets V_i (for $i = -(k-1), \dots, k-1$) such that $|V_i| = k - |i|$. We let G be the graph with vertices $V = \bigcup_{|i| < k} V_i$ and all possible edges between V_i and V_{i+1} (for $i = -(k-1), \dots, k-2$).

It is clear that if an induced tree in G contains a vertex from V_i and two vertices from V_{i+1} then it contains no vertex of V_j for $j > i + 1$; similarly for $i + 1$ replaced by $i - 1$. Therefore any maximum induced tree is one of trees $T_{a,b}$ ($-(k-1) \leq a < b \leq k-1$ and $b - a > 1$): it contains all vertices from two levels, V_a and V_b and one vertex from each V_i where $a < i < b$. It is easy to compute that such tree contains $2k - 1$ vertices out of the $|V| = k^2$; this proves $f_B(k^2) \leq 2k - 1$. If $(k-1)^2 < n \leq k^2$ then we take a subgraph of G to show that $f_B(n) \leq 2k - 1 < 2\sqrt{n} + 1$. \square

Lemma 2.2 (Blow-up construction) *Let G be a connected triangle-free graph and let $W \subseteq V(G)$ be a subset of m vertices ($m \geq 3$) such that any induced tree in G contains at most t vertices of W . Then we have $f_T(n) = O(n^{\ln(t-1)/\ln(m-1)})$. The same result holds with “triangle-free” replaced by “bipartite” and with f_T replaced by f_B .*

Proof: We let $W = \{w_0, \dots, w_{m-1}\}$, and write $r = m - 1$ and $q = t - 1$ to simplify expressions. As G is triangle-free it follows that $t \geq 3$, and so $q \geq 2$.

Let $T = T_{r,l}$ be a rooted tree with $l+1$ levels (counting root as one level) in which each non-leaf vertex has r sons. Next, for each vertex v of T we take a copy G_v of G (so that distinct copies are disjoint). Whenever v is a non-leaf vertex of T and u is its i -th son, we introduce an edge between w_i in G_v and w_0 in G_u ; the resulting graph will be called $T(G)$ (see Fig. 1). Clearly this graph is triangle-free/bipartite if G was triangle-free/bipartite. Moreover, $|V(T(G))| = |V(T)| \cdot |V(G)|$ and $|V(T)| = \frac{r^{l+1}-1}{r-1} = \Theta(r^l)$ (since $l \rightarrow \infty$ and $r \geq 2$).

Let S be an induced subtree of $T(G)$ and put

$$\bar{S} = \{v \in V(T) \mid G_v \text{ contains a vertex of } S\}.$$

By construction, $S \cap G_v$ is a tree in G_v for each v . So the condition on G implies that each vertex of \bar{S} has at most t neighbors in \bar{S} . Consequently, we have (since $q \geq 2$)

$$|\bar{S}| \leq 1 + \sum_{i=1}^l (q+1)q^{i-1} \leq 1 + (q+1)\frac{q^l - 1}{q-1} = \Theta(q^l).$$

Now recall that q , r , and $|V(G)|$ are constants. For a given n , choose the smallest l such that $n \leq |V(T_{r,l}(G))|$; we have $n = \Theta(r^l)$. By the above considerations,

$$f(n) \leq f(T_{r,l}(G)) \leq |V(G)| \cdot \Theta(q^l) = \Theta(r^{l \log_r q}) = \Theta(n^{\log_r q}),$$

which finishes the proof. \square




Figure 1: Graph $T_{3,2}(G)$ from the proof of Lemma 2.2.

Corollary 2.3 *If $f_T(n_0) < \sqrt{n_0}$ for some n_0 , then $f_T(n) = O(n^{1/2-\varepsilon})$ for a positive constant $\varepsilon > 0$. (The same is true for f_B .)*

Proof: Let G be the graph on n_0 vertices for which $t(G) = t < \sqrt{n_0}$. We let $W = V(G)$ and $m = n_0$ and apply Lemma 2.2. \square

As mentioned in the introduction, Krishnan and Ochem [5] search for values of $f_T(n)$ using a computer. This was motivated by hope that Corollary 2.3 would apply. It turns out, however, that for small n Observation 2.1 gives a precise estimate even for $f_T(n)$ (e.g., $f_T(15) = 7$); therefore Corollary 2.3 does not apply.

Remark. If we consider the construction from Lemma 2.2 for $G = K_3$, $W = V(G)$, $m = 3$, and $t = 2$ we recover a result of [2] that there is a graph G containing triangles (but no K_4) such that $t(G) = O(\log n)$.

3 Lower bound for bipartite graphs

Here we prove a statement weaker than Theorem 1.2—we give a bound on $f_B(n)$ instead of $f_T(n)$. The proof is simpler than that of Theorem 1.2 and it serves as an introduction to it.

We begin with a lemma about selecting induced forests of a particular kind in a bipartite graph. We introduce some terminology. Let H be a bipartite graph with color classes A and B . We will think of A as the “top” class and B as the “bottom” class (in a drawing of G in the plane, say). We write $a = |A|$ and $b = |B|$. For a subgraph F of H we write $A(F) = V(F) \cap A$, we set $a(F) = |A(F)|$, and we define $B(F)$ and $b(F)$ similarly.

Whenever we say *forest* we actually mean an induced subgraph of H that is a forest. An *up-forest* F is a forest such that every vertex in $A(F)$ has degree (in F) precisely 1 and every vertex in $B(F)$ has degree (in F) at least 1.

A *matching* is a forest F in which all vertices have degrees (in F) exactly 1.




Figure 2: An up-forest

Lemma 3.1 *Let H be a bipartite graph with color classes A and B as above, let Δ be the maximum degree of H , and let $\eta \in (0, 1)$ be a real parameter. Let us suppose that every vertex in A is connected to at least one vertex in B . Then at least one of the following cases occurs:*

(M) *There is a matching with at least $(1 - \eta)a$ edges.*

(B) *There is an up-forest F with*

$$b(F) \geq \frac{\eta}{\Delta^3} \cdot a$$

that is 2-branching, meaning that every vertex in $B(F)$ has degree at least 2 in F .



Figure 3: An illustration of Lemma 3.1

Proof. Let $B' \subseteq B$ be the set of vertices of degree 1 in B . If $|B'| \geq (1 - \eta)a$ then, clearly, case (M) occurs, so we may assume $|B'| < (1 - \eta)a$. Let $B'' \subseteq B$ consist of all vertices of degree at least 2. Since every vertex in A has degree at least 1, $|E(H \setminus N(B'))| \geq \eta a$, and so $|B''| \geq (\eta/\Delta)a$.

Let us set $B_0 = B''$ and let F_0 be an empty graph. Supposing that a set $B_{i-1} \subseteq B''$ and an up-forest F_{i-1} have already been constructed with $B_{i-1} \neq \emptyset$, we construct B_i and F_i . We let v_i be an arbitrary vertex in B_{i-1} , and we let S_i be the star formed by v_i and all of its neighbors in A . We set $F_i = F_{i-1} \cup S_i$, we let $N_i \subseteq B$ be the neighborhood of $A(S_i)$, and we let B_i be $B_{i-1} \setminus N_i$. The construction finishes when $B_i = \emptyset$, with F_i as the resulting up-forest.

It is easy to check that this construction indeed yields an up-forest F with each degree in $B(F)$ at least 2. We have $a(S_i) \leq \Delta$ and $|N_i| \leq a(S_i)(\Delta - 1) + 1$, and so in each step, at most $|N_i| \leq \Delta(\Delta - 1) + 1 \leq \Delta^2$ vertices are removed from B_i . Having started with at least $(\eta/\Delta)a$ vertices, we can proceed for at least $(\eta/\Delta^3)a$ steps, and so the resulting up-forest is as in (B). \square

Now we prove the lower bound

$$f_B(n) \geq e^{c\sqrt{\log n}}$$

for a constant $c > 0$.

Let G be a given connected bipartite graph. We assume that $n = |V(G)|$ is sufficiently large whenever convenient. We let t be the “target size” of an induced tree in G we are looking for; namely, $t = \lceil \exp(c\sqrt{\log n}) \rceil$. If G has a vertex of degree at least $t - 1$, then we can take its star for the induced tree and we are done, so we may assume that the maximum degree satisfies $\Delta \leq t - 2$.

Let us fix an arbitrary vertex of G as a root, and let L_i be the set of vertices of G at distance precisely i from the root. All edges of G go between L_{i-1} and L_i for some i , since an edge within some L_i would close an odd cycle.

We may assume that $L_t = \emptyset$, for otherwise G contains an induced path of length t . Hence there is a k with $|L_k| \geq n/t$.

Let us fix such a k . We are going to construct sets $M_i \subseteq L_i$, $i = k, k-1, \dots$, inductively, until we first reach an i with $|M_i| = 1$ (this happens for $i = 0$ at the latest since $|L_0| = 1$). We shall let ℓ be this last i .

Suppose that nonempty sets M_k, M_{k-1}, \dots, M_i have already been constructed, in such a way that the subgraph of G induced by $M_k \cup \dots \cup M_i$ is a forest, each of whose components intersects M_i in at most one vertex. We are going to construct M_{i-1} .

Let us put $A = M_i$, $B = L_{i-1}$, and let us consider the bipartite graph H induced by $A \cup B$ in G . Every vertex of A is connected to at least one vertex in B . We set $\eta = \frac{1}{t}$ and apply Lemma 3.1. This yields an up-forest F in H as in the lemma. We define $M_{i-1} = B(F)$.

If F is a matching, i.e., case (M) occurred in the lemma, we call the step from M_i to M_{i-1} a *matching step*. In this case, we have $|M_{i-1}| \geq (1 - \frac{1}{t})|M_i|$. Otherwise, F is a 2-branching forest; then we call the step a *branching step* and we have $|M_{i-1}| \geq |M_i|/(t\Delta^3) \geq |M_i|/t^4$.

Suppose that the sets M_k, \dots, M_ℓ have been constructed, $|M_\ell| = 1$. We claim that the number b of branching steps in the construction is at least $c_1\sqrt{\log n}$ for a suitable constant $c_1 > 0$. Indeed, there are no more than t matching steps, and so $1 = |M_\ell| \geq |M_k|(1 - 1/t)^t t^{-4b} \geq (n/t)e^{-1}/2 \cdot t^{-4b} = \Omega(nt^{-4b-1})$. Thus $b = \Omega(\log n / \log t) = \Omega(\sqrt{\log n})$, since $t = \lceil \exp(c\sqrt{\log n}) \rceil$.

It is easy to see that $M_k \cup M_{k-1} \cup \dots \cup M_\ell$ induces a forest in G . We let T be the component of this forest containing the single vertex of M_ℓ . Since every vertex of M_{i-1} , $\ell < i \leq k$, has at least one neighbor in M_i , and if the step from M_i to M_{i-1} was a branching step then each vertex of M_{i-1} has at least two neighbors in M_i , it follows that T has at least $2^b = \exp(\Omega(\sqrt{\log n}))$ vertices. This finishes the proof of the lower bound $f_B(n) \geq \exp(c\sqrt{\log n})$. \square

Remark. The above proof may seem wasteful in many respects. However, the result is tight up to the value of the constant in the exponent if we insist on selecting an induced tree “growing up” (i.e., made of up-forests for some choice of root and corresponding

sets L_i). Indeed, any such induced tree in the graph G_r in Figure 4 may contain at most two of the r vertices at the topmost level of the graph. Let us put $r = \exp(c\sqrt{\log n})$ and glue copies of G_r according to the pattern of a complete r -ary tree (as in the proof of Lemma 2.2), so that the resulting graph has approximately n vertices (that is, the depth is $l = \Theta(\sqrt{\log n})$). We obtain a graph with all up-growing induced trees having size at most $2^l = \exp(O(\sqrt{\log n}))$.




Figure 4: Graph G_6 in which all “up-growing trees” contain at most two vertices of the uppermost level.

4 Lower bound for triangle-free graphs

Here we prove Theorem 1.2. The scheme of the proof is very similar to the proof of the same bound for bipartite graphs in Section 3. We continue using the definitions and notation from that proof. So we decompose the given graph into the levels L_0, L_1, \dots, L_r , $r < t$. The main difference compared to the bipartite case is that there may now be edges within the levels L_i . We will need the well-known fact that any graph on n vertices with maximum degree Δ contains an independent set of size at least $n/(\Delta + 1)$. We will also need the following simple modification.

Lemma 4.1 *Let Γ be a graph (not necessarily bipartite) on n vertices with maximum degree Δ , and let $\eta \in [0, 1]$ be a real parameter. Then at least one of the following two cases occurs:*

- (IS) Γ contains an independent set with at least $(1 - \eta)n$ vertices.
- (IM) Γ contains an induced matching with at least $\frac{\eta}{2\Delta}n$ edges.

Proof. We repeatedly select edges e_1, e_2, \dots of Γ ; having selected e_i , we delete it and all the neighbors of its endvertices from the current graph. In each step we delete at most 2Δ vertices, so we either construct an induced matching as in (IM) or reach an edgeless graph after deleting at most ηn vertices, hence yielding an induced set as in (IS). \square

Proof of Theorem 1.2. We proceed similarly as in the previous section. We suppose G is a given triangle-free graph on n vertices (and that n is big enough), we put $t = \lceil \exp(c\sqrt{\log n}) \rceil$. Again, we may assume $t \leq \Delta - 2$: G is triangle-free, so a star of a vertex is an induced tree.

As before, we begin by selecting a root vertex and constructing the at most t levels L_0, L_1, \dots . We select k such that $|L_k| \geq n/t$ and we will construct sets $M_k, M_{k-1}, \dots, M_\ell$, such that $M_i \subseteq L_i$, $|M_\ell| = 1$, in such a way that their union induces a forest in G . In the induction step, we will either construct M_{i-1} from M_i , or sometimes we will go down two levels at once, producing both M_{i-1} and M_{i-2} .

We begin by selecting M_k as an independent set in the subgraph induced by L_k . By the fact mentioned before Lemma 4.1 we may assume $|M_k| \geq |L_k|/t \geq n/t^2$.

We suppose that M_i has already been constructed so that each component of the forest induced by $M_k \cup \dots \cup M_i$ intersects L_i in at most one vertex (and, in particular, M_i is an independent set). Now we proceed as in the proof in Section 3: We let $A = M_i$, $B = L_{i-1}$, and we consider the bipartite graph H induced by $A \cup B$ in G . We apply Lemma 3.1 to H with $\eta = \frac{1}{t}$, obtaining an up-forest F . We set $M'_{i-1} = B(F)$; this is not yet the final M_i since there may be edges on M'_{i-1} .

If case (B) occurred in Lemma 3.1, we have $|M'_{i-1}| \geq |M_i|/t^4$. We let M_{i-1} be an independent set of size $|M'_i|/(\Delta + 1) \geq |M_i|/t^5$ in the subgraph induced by M'_{i-1} . We call this step a *branching step*.

If case (M) occurred in Lemma 3.1, we have $|M'_{i-1}| \geq (1 - \frac{1}{t})|M_i|$. Then we apply Lemma 4.1 with $\eta = \frac{1}{t}$ to the graph Γ induced in G by M'_{i-1} . If case (IS) applies in that lemma, we let M_{i-1} be the independent set of size at least $(1 - \frac{1}{t})|M'_{i-1}| \geq (1 - \frac{1}{t})^2|M_i|$; we call this step a *matching step*. Both the matching step and the branching step go one level down, from i to $i - 1$.

If case (IM) applies in Lemma 4.1, we define M_{i-1} as the vertex set of the induced matching from the lemma. In this case we have $|M_{i-1}| \geq (1 - \frac{1}{t})|M_i|/t^2$. Note that this M_{i-1} does not satisfy the inductive assumption (it is not an independent set). We are also going to construct M_{i-2} in the same step, thus going from i to $i - 2$. To obtain M_{i-2} , we define another auxiliary bipartite graph, which we again call H to save letters. The bottom color class B is L_{i-2} , and the top color class A is obtained by contracting the edges induced by M_{i-1} . More formally, we set $A = \{uu' \in E(G) : u, u' \in M_{i-1}\}$, $B = L_{i-2}$, and $E(H) = \{\{uu', v\} : u, u' \in A, v \in B, uv \in E(G) \text{ or } u'v \in E(G)\}$. (Note that in this definition it can not happen that both uv and $u'v$ are edges of G , as G is triangle-free.) We apply Lemma 3.1 with $\eta = \frac{1}{2}$, say, to H . In both of cases (M) and (B) we obtain an up-forest F in H with $b(F) \geq |M_{i-1}|/(32t^3)$ (we note that $|A| = \frac{1}{2}|M_{i-1}|$ and that H has maximum degree no larger than $2t$). We set $M'_{i-2} = B(F)$, and finally, we select M_{i-2} as an independent set of size at least $|M'_{i-2}|/t$ in the subgraph induced by M'_{i-2} . Since G is triangle-free, one can check that $M_k \cup \dots \cup M_{i-1} \cup M_{i-2}$ induces a forest. We have $|M_{i-2}| \geq |M_{i-1}|/32t^4 \geq |M_i| \cdot (1 - \frac{1}{t})/32t^6 \geq |M_i|/t^7$. We call this step from M_i to M_{i-2} a *double-step*.

By calculation similar to that in Section 3, we find that the number b of branching steps and double-steps together is at least $\Omega(\sqrt{\log n})$. We again claim that the component of the forest induced by $M_k \cup \dots \cup M_\ell$ containing the single vertex of M_ℓ has at least 2^b vertices. Indeed, if M_i was obtained from M_{i+1} by a branching step, then each vertex of M_i has at least two successors in M_{i+1} . If M_i was obtained from M_{i+2} by a double-step, then each vertex v of M_i has at least one successor in M_{i+1} , this is connected by an edge to

precisely one other vertex of M_{i+1} , and both of these vertices have one neighbor in M_{i+2} ; consequently v has at least two successors in M_{i+2} . Theorem 1.2 is proved. \square

Acknowledgment

We would like to thank participants of a research seminar where initial steps of this work have been made, for a stimulating environment. In particular, we are indebted to Robert Babilon, Martin Bálek, Helena Nyklová, Ondra Pangrác, and Pavel Valtr for useful discussions and observations. We also want to thank the anonymous referee for helpful comments.

References

- [1] Richard N. Ball and Aleš Pultr, *Forbidden forests in Priestley spaces*, Cah. Topol. Géom. Différ. Catég. **45** (2004), no. 1, 2–22.
- [2] Paul Erdős, Michael Saks, and Vera T. Sós, *Maximum induced trees in graphs*, J. Combin. Theory Ser. B **41** (1986), no. 1, 61–79.
- [3] Wenceslas Fernandez de la Vega, *Induced trees in sparse random graphs*, Graphs Combin. **2** (1986), no. 3, 227–231.
- [4] Jeong Han Kim, *The Ramsey number $R(3, t)$ has order of magnitude $t^2/\log t$* , Random Structures Algorithms **7** (1995), no. 3, 173–207.
- [5] Sivarama Krishnan and Pascal Ochem, *Searching for induced trees*, DocCourse Prague 2005, Project Report, Charles University, Prague 2005.
- [6] Zbigniew Palka and Andrzej Ruciński, *On the order of the largest induced tree in a random graph*, Discrete Appl. Math. **15** (1986), no. 1, 75–83.