

# ALGEBRAIC MATCHING THEORY

C. D. Godsil <sup>1</sup>

Department of Combinatorics and Optimization  
University of Waterloo  
Waterloo, Ontario  
Canada N2L 3G1  
chris@bilby.uwaterloo.ca

Submitted: July 6, 1994; Accepted: April 11, 1995.

**Abstract:** The number of vertices missed by a maximum matching in a graph  $G$  is the multiplicity of zero as a root of the matchings polynomial  $\mu(G, x)$  of  $G$ , and hence many results in matching theory can be expressed in terms of this multiplicity. Thus, if  $\text{mult}(\theta, G)$  denotes the multiplicity of  $\theta$  as a zero of  $\mu(G, x)$ , then Gallai's lemma is equivalent to the assertion that if  $\text{mult}(\theta, G \setminus u) < \text{mult}(\theta, G)$  for each vertex  $u$  of  $G$ , then  $\text{mult}(\theta, G) = 1$ .

This paper extends a number of results in matching theory to results concerning  $\text{mult}(\theta, G)$ , where  $\theta$  is not necessarily zero. If  $P$  is a path in  $G$  then  $G \setminus P$  denotes the graph got by deleting the vertices of  $P$  from  $G$ . We prove that  $\text{mult}(\theta, G \setminus P) \geq \text{mult}(\theta, G) - 1$ , and we say  $P$  is  $\theta$ -essential when equality holds. We show that if, all paths in  $G$  are  $\theta$ -essential, then  $\text{mult}(\theta, G) = 1$ . We define  $G$  to be  $\theta$ -critical if all vertices in  $G$  are  $\theta$ -essential and  $\text{mult}(\theta, G) = 1$ . We prove that if  $\text{mult}(\theta, G) = k$  then there is an induced subgraph  $H$  with exactly  $k$   $\theta$ -critical components, and the vertices in  $G \setminus H$  are covered by  $k$  disjoint paths.

AMS Classification Numbers: 05C70, 05E99

---

<sup>1</sup> Support from grant OGP0009439 of the National Sciences and Engineering Council of Canada is gratefully acknowledged.

## 1. Introduction

A  $k$ -matching in a graph  $G$  is a matching with exactly  $k$  edges and the number of  $k$ -matchings in  $G$  is denoted by  $p(G, k)$ . If  $n = |V(G)|$  we define the matchings polynomial  $\mu(G, x)$  by

$$\mu(G, x) := \sum_{k \geq 0} (-1)^k p(G, k) x^{n-2k}.$$

(Here  $p(G, 0) = 1$ .) By way of example, the matchings polynomial of the path on four vertices is  $x^4 - 3x^2 + 1$ . The matchings polynomial is related to the characteristic polynomial  $\phi(G, x)$  of  $G$ , which is defined to be the characteristic polynomial of the adjacency matrix of  $G$ . In particular  $\phi(G, x) = \mu(G, x)$  if and only if  $G$  is a forest [4: Corollary 4.2]. Also the matchings polynomial of any connected graph is a factor of the characteristic polynomial of some tree. (For this, see Theorem 2.2 below.)

Let  $\text{mult}(\theta, G)$  denote the multiplicity of  $\theta$  as a zero of  $\mu(G, x)$ . If  $\theta = 0$  then  $\text{mult}(\theta, G)$  is the number of vertices in  $G$  missed by a maximum matching. Consequently many classical results in the theory of matchings provide information related to  $\text{mult}(0, G)$ . We refer in particular to Gallai's lemma and the Edmonds-Gallai structure theorem, which we now discuss briefly.

A vertex  $u$  of  $G$  is  $\theta$ -essential if  $\text{mult}(\theta, G \setminus u) < \text{mult}(\theta, G)$ . So a vertex is 0-essential if and only if it is missed by some maximum matching of  $G$ . Gallai's lemma is the assertion that if  $G$  is connected,  $\theta = 0$  and every vertex is  $\theta$ -essential then  $\text{mult}(\theta, G) = 1$ . (A more traditional expression of this result is given in [8: §3.1].) A vertex is  $\theta$ -special if it is not  $\theta$ -essential but has a neighbour which is  $\theta$ -essential. The Edmonds-Gallai structure in large part reduces to the assertion that if  $\theta = 0$  and  $v$  is a  $\theta$ -special vertex in  $G$  then a vertex  $u$  is  $\theta$ -essential in  $G$  and if and only if it is  $\theta$ -essential in  $G \setminus v$ . (For more information, see [8: §3.2].) One aim of the present paper is to investigate the extent to which these results are true when  $\theta \neq 0$ .

There is a second source of motivation for our work. Heilman and Lieb proved that if  $G$  has a Hamilton path then all zeros of  $\mu(G, x)$  are simple. (This is an easy consequence of Corollary 2.5 below.) Since all known vertex-transitive graphs have Hamilton paths we are lead to ask whether there is a vertex-transitive graph  $G$  such that  $\mu(G, x)$  has a multiple zero. As we will see, it is easy to show that if  $\theta$  is a zero of  $\mu(G, x)$  and  $G$  is vertex-transitive then every vertex of  $G$  is  $\theta$ -essential. Hence, if we could prove Gallai's lemma for general zeros of the matchings polynomial, we would have a negative answer to this question.

## 2. Identities

The first result provides the basic properties of the matchings polynomial  $\mu(G, x)$ . We write  $u \sim v$  to denote that the vertex  $u$  is adjacent to the vertex  $v$ . For the details see, for example, [6: Theorem 1.1].

**2.1 Theorem.** *The matchings polynomial satisfies the following identities:*

- (a)  $\mu(G \cup H, x) = \mu(G, x) \mu(H, x)$ ,
- (b)  $\mu(G, x) = \mu(G \setminus e, x) - \mu(G \setminus uv, x)$  if  $e = \{u, v\}$  is an edge of  $G$ ,
- (c)  $\mu(G, x) = x \mu(G \setminus u, x) - \sum_{i \sim u} \mu(G \setminus ui, x)$ , if  $u \in V(G)$ ,
- (d)  $\frac{d}{dx} \mu(G, x) = \sum_{i \in V(G)} \mu(G \setminus i, x)$ . □

Let  $G$  be a graph with a vertex  $u$ . By  $\mathcal{P}(u)$  we denote the set of paths in  $G$  which start at  $u$ . The *path tree*  $T(G, u)$  of  $G$  relative to  $u$  has  $\mathcal{P}(u)$  as its vertex set, and two paths are adjacent if one is a maximal proper subpath of the other. Note that each path in  $\mathcal{P}(u)$  determines a path starting with  $u$  in  $T(G, u)$  and with same length. We will usually denote them by the same symbol. The following result is taken from [6: Theorem 6.1.1].

**2.2 Theorem.** *Let  $u$  be a vertex in the graph  $G$  and let  $T = T(G, u)$  be the path tree of  $G$  with respect to  $u$ . Then*

$$\frac{\mu(G \setminus u, x)}{\mu(G, x)} = \frac{\mu(T \setminus u, x)}{\mu(T, x)}$$

and, if  $G$  is connected, then  $\mu(G, x)$  divides  $\mu(T, x)$ . □

Because the matchings polynomial of a tree is equal to the characteristic polynomial of its adjacency matrix, its zeros are real; consequently Theorem 2.2 implies that the zeros of the matchings polynomial of  $G$  are real, and also that they are interlaced by the zeros of  $\mu(G \setminus u, x)$ , for any vertex  $u$ . (By interlace, we mean that, between any two zeros of  $\mu(G, x)$ , there is a zero of  $\mu(G \setminus u, x)$ . This implies in particular that the multiplicity of a zero  $\theta$  in  $\mu(G, x)$  and  $\mu(G \setminus u, x)$  can differ by at most one.) For a more extensive discussion of these matters, see [6: §6.1].

We will need a strengthening of the first claim in Theorem 2.2.

**2.3 Corollary.** *Let  $u$  be a vertex in the graph  $G$  and let  $T = T(G, u)$  be the path tree of  $G$  with respect to  $u$ . If  $P \in \mathcal{P}(u)$  then*

$$\frac{\mu(G \setminus P, x)}{\mu(G, x)} = \frac{\mu(T \setminus P, x)}{\mu(T, x)}.$$

*Proof.* We proceed by induction on the number of vertices in  $P$ . If  $P$  has only one vertex, we appeal to the theorem. Suppose then that  $P$  has at least two vertices in it, and that  $v$  is the end vertex of  $P$  other than  $u$ . Let  $Q$  be the path  $P \setminus v$  and let  $H$  denote  $G \setminus Q$ . Then

$$\frac{\mu(G \setminus P, x)}{\mu(G, x)} = \frac{\mu(G \setminus P, x)}{\mu(G \setminus Q, x)} \frac{\mu(G \setminus Q, x)}{\mu(G, x)} = \frac{\mu(T(H, v) \setminus v, x)}{\mu(T(H, v), x)} \frac{\mu(T \setminus Q, x)}{\mu(T, x)},$$

where the second equality follows by induction. Now  $T(H, v)$  is one component of  $T(G, u) \setminus Q$ , and if we delete the vertex  $v$  from this component from  $T(G, u) \setminus Q$ , the graph that results is  $T(G, u) \setminus P$ . Consequently

$$\frac{\mu(T(H, v) \setminus v, x)}{\mu(T(H, v), x)} = \frac{\mu(T \setminus P, x)}{\mu(T \setminus Q, x)}.$$

The results follows immediately from this. □

Let  $\mathcal{P}(u, v)$  denote the set of paths in  $G$  which start at  $u$  and finish at  $v$ . The following result will be one of our main tools. It is a special case of [7: Theorem 6.3].

**2.4 Lemma (Heilmann and Lieb).** *Let  $u$  and  $v$  be vertices in the graph  $G$ . Then*

$$\mu(G \setminus u, x) \mu(G \setminus v, x) - \mu(G, x) \mu(G \setminus uv, x) = \sum_{P \in \mathcal{P}(u, v)} \mu(G \setminus P, x)^2. \quad \square$$

This lemma has a number of important consequences. In [5: Section 4] it is used to show that  $\text{mult}(\theta, G)$  is a lower bound on the number of paths needed to cover the vertices of  $G$ , and that the number of distinct zeros of  $\mu(G, x)$  is an upper bound on the length of a longest path. For our immediate purposes, the following will be the most useful.

**2.5 Corollary.** *If  $P$  is a path in the graph  $G$  then  $\mu(G \setminus P, x) / \mu(G, x)$  has only simple poles. In other words, for any zero  $\theta$  of  $\mu(G, x)$  we have*

$$\text{mult}(\theta, G \setminus P) \geq \text{mult}(\theta, G) - 1.$$

*Proof.* Suppose  $k = \text{mult}(\theta, G)$ . Then, by interlacing,  $\text{mult}(\theta, G \setminus u) \geq k - 1$  for any vertex  $u$  of  $G$  and  $\text{mult}(\theta, G \setminus uv) \geq k - 2$ . Hence the multiplicity of  $\theta$  as a zero of

$$\mu(G \setminus u, x) \mu(G \setminus v, x) - \mu(G, x) \mu(G \setminus uv, x)$$

is at least  $2k - 2$ . It follows from Lemma 2.4 that  $\text{mult}(\theta, G \setminus P) \geq k - 1$  for any path  $P$  in  $\mathcal{P}(u, v)$ . □

### 3. Essential Vertices and Paths

Let  $\theta$  be a zero of  $\mu(G, x)$ . A path  $P$  of  $G$  is  $\theta$ -essential if  $\text{mult}(\theta, G \setminus P) < \text{mult}(\theta, G)$ . (We will often be concerned with the case where  $P$  is a single vertex.) A vertex is  $\theta$ -special if it is not  $\theta$ -essential and is adjacent to an  $\theta$ -essential vertex. A graph is  $\theta$ -primitive if and only if every vertex is  $\theta$ -essential and it is  $\theta$ -critical if it is  $\theta$ -primitive and  $\text{mult}(\theta, G) = 1$ . (When  $\theta$  is determined by the context we will often drop the prefix ‘ $\theta$ -’ from these expressions.) If  $\theta = 0$  then a  $\theta$ -critical graph is the same thing as a factor-critical graph.

The next result implies that a vertex-transitive graph is  $\theta$ -primitive for any zero  $\theta$  of its matchings polynomial.

**3.1 Lemma.** *Any graph has at least one essential vertex.*

*Proof.* Let  $\theta$  be a zero of  $\mu(G, x)$  with multiplicity  $k$ . Then  $\theta$  has multiplicity  $k - 1$  as a zero of  $\mu'(G, x)$ . Since

$$\mu'(G, x) = \sum_{u \in V(G)} \mu(G \setminus u, x)$$

we see that if  $\text{mult}(\theta, G \setminus u) \geq k$  for all vertices  $u$  of  $G$  then  $\theta$  must have multiplicity at least  $k$  as a zero of  $\mu'(G, x)$ .  $\square$

**3.2 Lemma.** *If  $\theta \neq 0$  then any  $\theta$ -essential vertex  $u$  has a neighbour  $v$  such that the path  $uv$  is essential.*

*Proof.* Assume  $\theta \neq 0$  and let  $u$  be a  $\theta$ -essential vertex. Since

$$\mu(G, x) = x \mu(G \setminus u, x) - \sum_{i \sim u} \mu(G \setminus ui, x)$$

we see that if  $\text{mult}(\theta, G \setminus ui) \geq \text{mult}(\theta, G)$  for all neighbours  $i$  of  $u$  then  $\text{mult}(\theta, G \setminus u) \geq \text{mult}(\theta, G)$ .  $\square$

Note that the vertex  $v$  is not essential in  $G \setminus u$ . However it follows from the next lemma that the vertex  $v$  in the above lemma must be essential in  $G$ ; accordingly if  $\theta \neq 0$  then any essential vertex must have an essential neighbour.

**3.3 Lemma.** *If  $v$  is not an essential vertex of  $G$  then no path with  $v$  as an end-vertex is essential.*

*Proof.* Assume  $k = \text{mult}(\theta, G)$ . If  $v$  is not essential then  $\text{mult}(\theta, G \setminus v) \geq k$  and so, for any vertex  $u$  not equal to  $v$ , the multiplicity of  $\theta$  as a zero of

$$\mu(G \setminus u, x) \mu(G \setminus v, x) - \mu(G, x) \mu(G \setminus uv, x)$$

is at least  $2k - 1$ . By Lemma 2.4 we deduce that it is at least  $2k$  and that  $\text{mult}(\theta, G \setminus P) \geq k$  for all paths  $P$  in  $\mathcal{P}(v)$ .  $\square$

We now need some more notation. Suppose that  $G$  is a graph and  $\theta$  is a zero of  $\mu(G, x)$  with positive multiplicity  $k$ . A vertex  $u$  of  $G$  is  $\theta$ -positive if  $\text{mult}(\theta, G \setminus u) = k + 1$  and  $\theta$ -neutral if  $\text{mult}(\theta, G \setminus u) = k$ . (The ‘negative’ vertices will still be referred to as essential.) Note that, by interlacing,  $\text{mult}(\theta, G \setminus u)$  cannot be greater than  $k + 1$ .

**3.4 Lemma.** *Let  $G$  be a graph and  $u$  a vertex in  $G$  which is not essential. Then  $u$  is positive in  $G$  if and only if some neighbour of it is essential in  $G \setminus u$ .*

*Proof.* From Theorem 2.1(c) we have

$$\mu(G, x) = x \mu(G \setminus u, x) - \sum_{i \sim u} \mu(G \setminus ui, x). \quad (3.1)$$

If  $\text{mult}(\theta, G \setminus u) = k + 1$  and  $\text{mult}(\theta, G \setminus ui) \geq k + 1$  for all neighbours  $i$  of  $u$  then it follows that  $\text{mult}(\theta, G) \geq k + 1$  and  $u$  is not positive.

On the other hand, suppose  $u$  is not essential in  $G$  and  $v$  is a neighbour of  $u$  which is essential in  $G \setminus u$ . From the previous lemma we see that the path  $uv$  is not essential and thus  $\text{mult}(\theta, G \setminus uv) \geq \text{mult}(\theta, G)$ . As  $v$  is essential in  $G \setminus u$  it follows that  $\text{mult}(\theta, G \setminus u) > \text{mult}(\theta, G)$ .  $\square$

We say that  $S$  is an *extremal* subtree of the tree  $T$  if  $S$  is a component of  $T \setminus v$  for some vertex  $v$  of  $G$ .

**3.5 Lemma.** *Let  $S$  be an extremal subtree of  $T$  that is inclusion-minimal subject to the condition that  $\text{mult}(\theta, S) \neq 0$ , and let  $v$  be the vertex of  $T$  such that  $S$  is a component of  $T \setminus v$ . Then  $v$  is  $\theta$ -positive in  $T$ .*

*Proof.* Let  $u$  be the vertex of  $S$  adjacent to  $v$  and let  $e$  be the edge  $\{u, v\}$ . Then  $T \setminus e$  has exactly two components, one of which is  $S$ . Denote the other by  $R$ .

By hypothesis  $\text{mult}(\theta, S') = 0$  for any component  $S'$  of  $S \setminus u$ , therefore  $\text{mult}(\theta, S \setminus u) = 0$  by Theorem 2.1(a) and so  $u$  is essential in  $S$ . Since  $S$  is a component of  $T \setminus v$  it follows that  $u$  is essential in  $T \setminus v$ . If we can show that  $v$  is not essential then  $v$  must be positive in  $T$ , by the previous lemma.

Suppose  $\text{mult}(\theta, T) = m$ . By interlacing  $\text{mult}(\theta, T \setminus u) \geq m - 1$  and, as

$$\text{mult}(\theta, T \setminus u) = \text{mult}(\theta, R) + \text{mult}(\theta, S \setminus u) = \text{mult}(\theta, R),$$

we find that  $\text{mult}(\theta, R) \geq m - 1$ . By parts (a) and (b) of Theorem 2.1 we have

$$\mu(T, x) = \mu(R, x) \mu(S, x) - \mu(R \setminus v, x) \mu(S \setminus u, x)$$

and so, since the multiplicity of  $\theta$  as a zero of  $\mu(R, x) \mu(S, x)$  is at least  $m$ , we deduce that the multiplicity of  $\theta$  as a zero of  $\mu(R \setminus v, x) \mu(S \setminus u, x)$  is at least  $m$ . Since  $\text{mult}(\theta, S \setminus u) = 0$ , it follows that  $\text{mult}(\theta, R \setminus v) \geq m$ . On the other hand

$$\text{mult}(\theta, T \setminus v) = \text{mult}(\theta, R \setminus v) + \text{mult}(\theta, S) = \text{mult}(\theta, R \setminus v) + 1,$$

therefore  $\text{mult}(\theta, T \setminus v) \geq m + 1$  and  $v$  is positive in  $T$ . □

**3.6 Corollary (Neumaier).** *Let  $T$  be a tree and let  $\theta$  be a zero of  $\mu(T, x)$ . The following assertions are equivalent:*

- (a)  $\text{mult}(\theta, S) = 0$  for all extremal subtrees of  $T$ ,
- (b)  $T$  is  $\theta$ -critical,
- (c)  $T$  is  $\theta$ -primitive.

*Proof.* Since  $T \setminus v$  is a disjoint union of extremal subtrees for any vertex  $v$  in  $T$ , we see that if (a) holds then  $\text{mult}(\theta, T \setminus v) = 0$  for any vertex  $v$ . Hence  $T$  is  $\theta$ -critical and therefore it is also  $\theta$ -primitive. If  $T$  is  $\theta$ -primitive then no vertex in  $T$  is  $\theta$ -positive, whence Lemma 3.5 implies that (a) holds. □

Corollary 3.6 combines Theorem 3.1 and Corollary 3.3 from [9]. Note that the equivalence of (b) and (c) when  $\theta = 0$  is Gallai's lemma for trees.

**3.7 Lemma.** *Let  $G$  be a connected graph. If  $u \in V(G)$  and all paths in  $G$  starting at  $u$  are essential then  $G$  is critical.*

*Proof.* If all paths in  $\mathcal{P}(u)$  are essential then Lemma 3.3 implies that all vertices in  $G$  are essential. Hence  $G$  is primitive, and it only remains for us to show that  $\text{mult}(\theta, G) = 1$ .

Let  $T = T(G, u)$  be the path tree of  $G$  relative to  $u$ . From Theorem 2.2 we see that a path  $P$  from  $\mathcal{P}(u)$  is essential in  $G$  if and only if it is essential in  $T$ . So our hypothesis implies that all paths in  $T$  which start at  $u$  are essential, whence Lemma 3.3 yields that all vertices in  $T$  are essential. Hence  $T$  is  $\theta$ -primitive and therefore, by Corollary 3.6,  $\theta$  is a simple zero of  $\mu(T, x)$ . Using Theorem 2.2 again we deduce that  $\text{mult}(\theta, G) = 1$ .  $\square$

**3.8 Lemma.** *If  $u$  and  $v$  are essential vertices in  $G$  and  $v$  is not essential in  $G \setminus u$  then there is a  $\theta$ -essential path in  $\mathcal{P}(u, v)$ .*

*Proof.* Assume  $\text{mult}(\theta, G) = k$ . Our hypotheses imply that  $\text{mult}(\theta, G \setminus uv) \geq k - 1$ . If no path in  $\mathcal{P}(u, v)$  is essential then, by Lemma 2.4, the multiplicity of  $\theta$  as a zero of

$$\mu(G \setminus u, x) \mu(G \setminus v, x) - \mu(G, x) \mu(G \setminus uv, x)$$

is at least  $2k$ . Since  $\theta$  has multiplicity  $2k - 1$  as a zero of  $\mu(G, x) \mu(G \setminus uv, x)$  it must also have multiplicity at least  $2k - 1$  as a zero of  $\mu(G \setminus u, x) \mu(G \setminus v, x)$ . Hence  $u$  and  $v$  cannot both be essential.  $\square$

If  $u$  and  $v$  are essential in  $G$  then  $v$  is essential in  $G \setminus u$  if and only if  $u$  is essential in  $G \setminus v$ . Thus the hypothesis of Lemma 3.8 is symmetric in  $u$  and  $v$ , despite appearances.

**3.9 Corollary.** *Let  $G$  be a tree, let  $\theta$  be a zero of  $\mu(G, x)$  and let  $u$  be a vertex in  $G$ . Then all paths in  $\mathcal{P}(u)$  are essential if and only if all vertices in  $G$  are essential.*

*Proof.* It follows from Lemma 3.3 that if all paths in  $\mathcal{P}(u)$  are essential then all vertices in  $G$  are essential. Suppose conversely that all vertices in  $G$  are essential. By Corollary 3.6 it follows that  $\text{mult}(\theta, G) = 1$ . Hence the hypotheses of Lemma 3.8 are satisfied by any two vertices in  $G$ , and so any two vertices are joined by an essential path. Since  $G$  is a tree the path joining any two vertices is unique and therefore all paths in  $\mathcal{P}(u)$  are essential.  $\square$

#### 4. Structure Theorems

We now apply the machinery we have developed in the previous section.

**4.1 Lemma (De Caen [2]).** *Let  $u$  and  $v$  be adjacent vertices in a bipartite graph. If  $u$  is 0-essential then  $v$  is 0-special.*

*Proof.* Suppose that  $u$  and  $v$  are 0-essential neighbours in the bipartite graph  $G$ . As  $uv$  is a path, using Corollary 2.5 we find that

$$\text{mult}(0, G \setminus uv) \geq \text{mult}(0, G) - 1 = \text{mult}(0, G \setminus u),$$

and therefore  $v$  is not essential in  $G \setminus u$ . It follows from Lemma 3.8 that there is a 0-essential path  $P$  in  $G$  joining  $u$  to  $v$ .

We now show that  $P$  must have even length. From this it will follow that  $P$  together with the edge  $uv$  forms an odd cycle, which is impossible. From the definition of the matchings polynomial we see that  $\text{mult}(0, H)$  and  $|V(H)|$  have the same parity for any graph  $H$ . As

$$\text{mult}(0, G \setminus P) = \text{mult}(0, G) - 1$$

we deduce that  $|V(G)|$  and  $|V(G \setminus P)|$  have different parity and therefore  $P$  has even length.  $\square$

In the above proof we showed that a 0-essential path in a graph must have even length. Consequently no edge, viewed as a path of length one, can ever be 0-essential. It follows that  $K_1$  is the only connected graph such that all paths are 0-essential. In general any graph which is minimal subject to its matchings polynomial having a particular zero  $\theta$  will have the property that all its paths are  $\theta$ -essential.

Lemma 4.1 is not hard to prove without reference to the matchings polynomial. Note that it implies that in any bipartite graph there is a vertex which is covered by every maximal matching, and consequently that a bipartite graph with at least one edge cannot be 0-primitive. As noted by de Caen [2], this leads to a very simple inductive proof of König's lemma.

Our next result is a partial analog to the Edmonds-Gallai structure theorem. See, e.g., [8: Chapter 3.2].

**4.2 Theorem.** *Let  $\theta$  be a zero of  $\mu(G, x)$  with non-zero multiplicity  $k$  and let  $a$  be a positive vertex in  $G$ . Then:*

- (a) *if  $u$  is essential in  $G$  then it is essential in  $G \setminus a$ ;*
- (b) *if  $u$  is positive in  $G$  then it is essential or positive in  $G \setminus a$ ;*
- (c) *if  $u$  is neutral in  $G$  then it is essential or neutral in  $G \setminus a$ .*

*Proof.* If  $\text{mult}(\theta, G \setminus u) = k - 1$  and  $\text{mult}(\theta, G \setminus a) = k + 1$ , it follows by interlacing that  $\text{mult}(\theta, G \setminus au) = k$ . Hence  $u$  is essential in  $G \setminus a$ . Now suppose that  $u$  is positive in  $G$ . If  $\text{mult}(\theta, G \setminus au) \geq k + 1$  then  $\theta$  has multiplicity at least  $2k + 1$  as a zero of  $p(x)$  where

$$p(x) := \mu(G \setminus u, x) \mu(G \setminus a, x) - \mu(G, x) \mu(G \setminus au, x). \quad (4.1)$$

By Lemma 2.4, the multiplicity of  $\theta$  as a zero of  $p(x)$  must be even. It follows that this multiplicity must be at least  $2k + 2$  and hence that  $\theta$  has multiplicity at least  $2k + 2$  as a zero of  $\mu(G, x) \mu(G \setminus au, x)$ . Therefore  $\text{mult}(\theta, G \setminus au) \geq k + 2$  and so, by interlacing,  $\text{mult}(\theta, G \setminus au) = k + 2$  and  $u$  is positive in  $G \setminus a$ . If  $\text{mult}(\theta, G \setminus ua) = k + 2$  and  $u$  is neutral in  $G$ , then the multiplicity of  $\theta$  as a zero of  $p(x)$  is at least  $2k + 1$  and therefore at least  $2k + 2$ , but this implies that  $\theta$  is a zero of  $\mu(G \setminus u, x) \mu(G \setminus a, x)$  with multiplicity at least  $2k + 2$ . Thus we conclude that  $u$  is neutral or essential in  $G \setminus a$ .  $\square$

We note that Theorem 4.2(a) holds even if  $a$  is only neutral. If  $a$  is neutral and  $u$  is essential in  $G$  but not in  $G \setminus a$  then  $\theta$  has multiplicity at least  $2k - 1$  as a zero of (4.1) and so must have multiplicity at least  $2k$  as a zero of  $\mu(G, x) \mu(G \setminus au, x)$ . Hence its multiplicity as a zero of  $\mu(G \setminus u, x) \mu(G \setminus a, x)$  is at least  $2k$ , which is impossible.

The following consequence of Theorem 4.2 and the previous remark was proved for trees by Neumaier. (See [9: Theorem 3.4(iii)].)

**4.3 Corollary.** *Any special vertex is positive.*

*Proof.* Suppose that  $a$  is special in  $G$ , and that  $u$  is a neighbour of  $a$  which is essential in  $G$ . By part (a) of the theorem and the remark above,  $u$  is essential in  $G \setminus a$  and therefore, by Lemma 3.4,  $a$  is positive in  $G$ .  $\square$

Lemma 3.7 implies that if  $G$  is not  $\theta$ -critical then it contains a path,  $P$  say, that is not essential. If we delete  $P$  from  $G$  then the multiplicity of  $\theta$  as a zero of  $\mu(G, x)$  cannot decrease. Hence we may successively delete ‘inessential’ paths from  $G$ , to obtain a graph  $H$  such that  $\text{mult}(\theta, H) \geq \text{mult}(\theta, G)$  and all paths in  $H$  are essential. If  $k = \text{mult}(\theta, H)$  then, by Lemma 3.7 again,  $H$  contains exactly  $k$  critical components. The following result is a sharpening of this observation, since it implies that if  $\text{mult}(\theta, G) = k$  we may produce a graph with  $k$  critical components by deleting  $k$  vertex disjoint paths from  $G$ ,

**4.4 Lemma.** *Let  $G$  be a graph, let  $\theta$  be a zero of  $\mu(G, x)$  and let  $u$  be a  $\theta$ -essential vertex of  $G$ . Suppose that there is a path in  $\mathcal{P}(u)$  which is not  $\theta$ -essential. Then there is a path  $P$  in  $G$  starting at  $u$  such that  $\text{mult}(\theta, G \setminus P) = \text{mult}(\theta, G)$  and some component  $C$  of  $G \setminus P$  is critical. All vertices of  $C$  are essential in  $G$ .*

*Proof.* Suppose that there are paths in  $\mathcal{P}(u)$  which are not essential, choose one of minimum length and call it  $P$ . Let  $v$  be the end-vertex of  $P$  other than  $u$  and let  $P'$  be the path  $P \setminus v$ . Then  $P'$  is essential, hence

$$\text{mult}(\theta, G \setminus P') = \text{mult}(\theta, G) - 1$$

and, as  $P$  is not essential,

$$\text{mult}(\theta, G \setminus P) \geq \text{mult}(\theta, G).$$

But we get  $G \setminus P$  from  $G \setminus P'$  by deleting the single vertex  $v$ , therefore  $\text{mult}(\theta, G \setminus P) = \text{mult}(\theta, G)$  and  $v$  is positive in  $G \setminus P'$ . Consequently, by Lemma 3.4, there is an essential vertex  $u_1$  adjacent to  $v$  in  $(G \setminus P') \setminus v = G \setminus P$ .

We now prove by induction on the number of vertices that, if the conditions of the lemma hold, then there is a path  $P$  and a component  $C$  of  $G \setminus P$  as claimed and, further, there is a vertex  $w$  in  $C$  adjacent to the end-vertex of  $P$  distinct from  $u$  such that all paths in  $C$  that start at  $w$  are essential in  $C$ .

Let  $H$  denote  $G \setminus P$ . If all paths in  $H$  starting at  $u_1$  are essential then, by Lemma 3.7, the component  $C$  of  $H$  that contains  $u_1$  is critical. If  $Q$  is a path in  $C$  starting at  $u_1$  then  $\text{mult}(\theta, C \setminus Q) < \text{mult}(\theta, C)$ ; this implies that the path formed by the concatenation of  $P$  and  $Q$  is essential in  $G$  and hence, by Lemma 3.3, that all vertices in  $C$  are essential in  $G$ .

Thus we may suppose that there is a path in  $H$  starting at  $u_1$  that is not essential. Because  $H$  has fewer vertices than  $G$ , we may assume inductively that there is a path  $Q$  in  $H$  starting at  $u_1$  such that  $\text{mult}(\theta, H) = \text{mult}(\theta, H \setminus Q)$  and a critical component  $C$  of  $H \setminus Q$  that contains a neighbour  $w$  of the end-vertex of  $Q$  distinct from  $u_1$ . Further all the paths in  $C$  that start at  $w$  are essential.

Let  $PQ$  denote the path formed by concatenating  $P$  and  $Q$ . Then all claims of the lemma hold for  $G$ ,  $PQ$ ,  $u$  and  $C$ . □

The two results which follow provide a strengthening of the observation that the zeros of the matchings polynomial of a graph with a Hamilton path are simple.

**4.5 Lemma.** *Suppose that  $u$  and  $v$  are adjacent vertices in  $G$  such that  $\mu(G \setminus u, x)$  and  $\mu(G \setminus uv, x)$  have no common zero. Then  $\mu(G, x)$  and  $\mu(G \setminus u, x)$  have no common zero, and therefore both polynomials have only simple zeros.*

*Proof.* Assume by way of contradiction that  $\theta$  is a common zero of  $\mu(G, x)$  and  $\mu(G \setminus u, x)$ . If  $\text{mult}(\theta, G) > 1$  then by Corollary 2.5 we see that  $\theta$  is a zero of  $\mu(G \setminus u, x)$  and  $\mu(G \setminus uv, x)$ . If  $\text{mult}(\theta, G \setminus u) > 1$  then  $\text{mult}(\theta, G \setminus uv) > 0$ , by interlacing. Hence

$$\text{mult}(\theta, G) = \text{mult}(\theta, G \setminus u) = 1$$

and so  $u$  is a neutral vertex in  $G$ . It follows from Lemma 3.4 that no neighbour of  $u$  can be essential in  $G \setminus u$  and consequently  $\text{mult}(\theta, G \setminus uv) > 0$ .  $\square$

A simple induction argument on the length of  $P$  yields the following.

**4.6 Corollary.** *Let  $H$  be an induced subgraph of  $G$  and suppose that there is a vertex  $u$  in  $H$  and a path  $P$  in  $G$  such that*

$$V(H) \cap V(P) = u, \quad V(H) \cup V(P) = V(G).$$

*If  $\mu(H, x)$  and  $\mu(H \setminus u, x)$  have no common zero then all zeros of  $G$  are simple.*  $\square$

Note that the path  $P$  in this corollary does not have to be an induced path. One consequence of it is that if a graph has a Hamilton path then the zeros of its matchings polynomial are all simple. However this result shows that there will be many other graphs with all zeros simple.

## 5. Eigenvectors

Let  $G$  be a graph with adjacency matrix  $A = A(G)$ . We view an eigenvector  $f$  of  $A$  with eigenvalue  $\theta$  as a function on  $V(G)$  such that

$$\theta f(u) = \sum_{i \sim u} f(i).$$

We denote the characteristic polynomial of  $G$  by  $\phi(G, x)$ . (It is defined to be  $\det(xI - A(G))$ .) We recall that for forests the characteristic and matchings polynomials are equal. Our first result follows from the proof of Theorem 5.2 in [3].

**5.1 Lemma.** *Let  $\theta$  be an eigenvalue of the graph  $G$  and let  $u$  be a vertex in  $G$ . Then the maximum value of  $f(u)^2$  as  $f$  ranges over the eigenvectors of  $G$  with eigenvalue  $\theta$  and norm one is equal to  $\phi(G \setminus u, \theta) / \phi'(G, \theta)$ .*  $\square$

**5.2 Corollary (Neumaier [9: Theorem 3.4]).** *Let  $T$  be a tree and let  $\theta$  be a zero of its matchings polynomial. Then a vertex  $u$  is essential if and only if there is an eigenvector  $f$  of  $T$  such that  $f(u) \neq 0$ .*  $\square$

**5.3 Theorem.** *Let  $T$  be a tree, let  $\theta$  be a zero of  $\mu(T, x)$  and let  $a$  be a vertex of  $T$  which is not essential. Then a vertex is essential in  $T \setminus a$  if and only if it is essential in  $T$ . Further, if  $a$  is positive then it has an essential neighbour.*

*Proof.* Let  $W$  be the eigenspace of  $T$  belonging to  $\theta$  and let  $W_a$  be the corresponding eigenspace of  $T \setminus a$ . Then  $W_a$  is the direct sum of the eigenspaces of the component of  $T \setminus a$  belonging to  $\theta$  and  $W$  is the subspace formed by the vectors  $f$  such that

$$\sum_{i \sim a} f(i) = 0.$$

If  $a$  is neutral then  $W = W_a$  and so  $T$  and  $T \setminus a$  have the same essential vertices. If  $a$  is positive then  $W$  is a proper subspace of  $W_a$ , whence it follows that there are vectors in  $W_a$  which are not zero on all neighbours of  $a$ . For each vector in  $W_a$  there is an eigenvector in  $W$  with the same support on  $T \setminus a$ . Hence  $a$  has an essential neighbour and any vertex which is essential in  $T \setminus a$  is also essential in  $T$ .  $\square$

Theorem 5.3 is a strengthening of a result of Neumaier [9: Corollary 3.5]. Suppose that  $T$  is a tree with exactly  $s$  special vertices and  $\text{mult}(\theta, T) = k$ . Then Theorem 5.3 together with Theorem 4.2 implies that we may successively delete the special vertices, obtaining a forest  $F$  with no special vertices and  $\text{mult}(\theta, F) = k + s$ . Hence any component of  $F$  is either  $\theta$ -critical or does not have  $\theta$  as a zero of its matchings polynomial. Therefore  $F$  has exactly  $k + s$   $\theta$ -critical components, and these components form an induced subgraph of  $T$ .

## 6. Questions

Many problems remain. Here are some.

- (1) Must a positive vertex be special when  $\theta \neq 0$ ? (If  $\theta = 0$  then all vertices which are not essential are positive.)
- (2) What can be said of the graphs where every pair of vertices are joined by at least one essential path? (Or of the graphs with a vertex  $u$  such that all vertices can be joined to  $u$  by an essential path?)
- (3) Must a  $\theta$ -primitive graph be  $\theta$ -critical?

It might be interesting to investigate the case  $\theta = 1$  in depth.

## References

- [1] N. Biggs, *Algebraic Graph Theory*. (Cambridge U. P., Cambridge) 1974.
- [2] D. de Caen, On a theorem of König on bipartite graphs, *J. Comb. Inf. System Sci.* **13** (1988) 127.
- [3] C. D. Godsil, Matchings and walks in graphs, *J. Graph Theory*, **5**, (1981) 285–297.
- [4] C. D. Godsil and I. Gutman, On the theory of the matching polynomial, *J. Graph Theory*, **5** (1981), 137–144.
- [5] C. D. Godsil, Real graph polynomials, in *Progress in Graph Theory*, edited by J. A. Bondy and U. S. R. Murty, (Academic Press, Toronto) 1984, pp. 281–293.
- [6] C. D. Godsil, *Algebraic Combinatorics*. (Chapman and Hall, New York) 1993.
- [7] O. J. Heilmann and E. H. Lieb, Theory of monomer-dimer systems, *Commun. Math. Physics*, **25** (1972), 190–232.
- [8] L. Lovász and M. D. Plummer, *Matching Theory*. Annals Discrete Math. 29, (North-Holland, Amsterdam) 1986.
- [9] A. Neumaier, The second largest eigenvalue of a tree, *Linear Algebra Appl.* **48** (1982) 9–25.