

# Parking Functions and Noncrossing Partitions

Richard P. Stanley\*  
Department of Mathematics  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
rstan@math.mit.edu

Submitted: August 12, 1996; Accepted: November 12, 1996

*Dedicated to Herb Wilf on the occasion of his sixty-fifth birthday*

## Abstract

A *parking function* is a sequence  $(a_1, \dots, a_n)$  of positive integers such that if  $b_1 \leq b_2 \leq \dots \leq b_n$  is the increasing rearrangement of  $a_1, \dots, a_n$ , then  $b_i \leq i$ . A *noncrossing partition* of the set  $[n] = \{1, 2, \dots, n\}$  is a partition  $\pi$  of the set  $[n]$  with the property that if  $a < b < c < d$  and some block  $B$  of  $\pi$  contains both  $a$  and  $c$ , while some block  $B'$  of  $\pi$  contains both  $b$  and  $d$ , then  $B = B'$ . We establish some connections between parking functions and noncrossing partitions. A generating function for the flag  $f$ -vector of the lattice  $\text{NC}_{n+1}$  of noncrossing partitions of  $[n+1]$  is shown to coincide (up to the involution  $\omega$  on symmetric function) with Haiman's parking function symmetric function. We construct an edge labeling of  $\text{NC}_{n+1}$  whose chain labels are the set of all parking functions of length  $n$ . This leads to a local action of the symmetric group  $\mathfrak{S}_n$  on  $\text{NC}_{n+1}$ .

MR primary subject number: 06A07

MR secondary subject numbers: 05A15, 05E05, 05E10, 05E25

**1. Introduction.** A *parking function* is a sequence  $(a_1, \dots, a_n)$  of positive integers such that if  $b_1 \leq b_2 \leq \dots \leq b_n$  is the increasing rearrangement of

---

\*Partially supported by NSF grant DMS-9500714.

$a_1, \dots, a_n$ , then  $b_i \leq i$ .<sup>1</sup> Parking functions were introduced by Konheim and Weiss [14] in connection with a hashing problem (though the term “hashing” was not used). See this reference for the reason (formulated in a way which now would be considered politically incorrect) for the terminology “parking function.” Parking functions were subsequently related to labelled trees and to hyperplane arrangements. For further information on these connections see [31] and the references given there. In this paper we will develop a connection between parking functions and another topic, viz., noncrossing partitions.

A *noncrossing partition* of the set  $[n] = \{1, 2, \dots, n\}$  is a partition  $\pi$  of the set  $[n]$  (as defined e.g. in [29, p. 33]) with the property that if  $a < b < c < d$  and some block  $B$  of  $\pi$  contains both  $a$  and  $c$ , while some block  $B'$  of  $\pi$  contains both  $b$  and  $d$ , then  $B = B'$ . The study of noncrossing partitions goes back at least to H. W. Becker [1], where they are called “planar rhyme schemes.” The systematic study of noncrossing partitions began with Kreweras [15] and Poupard [22]. For some further work on noncrossing partitions, see [5][21][25][28] and the references given there.

A fundamental property of the set of noncrossing partitions of  $[n]$  is that it can be given a natural partial ordering. Namely, we define  $\pi \leq \sigma$  if every block of  $\pi$  is contained in a block of  $\sigma$ . In other words,  $\pi$  is a *refinement* of  $\sigma$ . Thus the poset  $\text{NC}_n$  of all noncrossing partitions of  $[n]$  is an induced subposet of the lattice  $\Pi_n$  of all partitions of  $[n]$  [29, Example 3.10.4]. In fact,  $\text{NC}_n$  is a lattice with a number of remarkable properties. We will develop additional properties of the lattice  $\text{NC}_n$  which connect it directly with parking functions.

**2. The parking function symmetric function.** Let  $P$  be a finite graded poset of rank  $n$  with  $\hat{0}$  and  $\hat{1}$  and with rank function  $\rho$ . (See [29, Ch. 3] for poset terminology and notation used here.) Let  $S$  be a subset of  $[n-1] = \{1, 2, \dots, n-1\}$ , and define  $\alpha_P(S)$  to be the number of chains  $\hat{0} = t_0 < t_1 < \dots < t_s = \hat{1}$  of  $P$  such that  $S = \{\rho(t_1), \rho(t_2), \dots, \rho(t_{s-1})\}$ . The function  $\alpha_P$  is called the *flag  $f$ -vector* of  $P$ . For  $S \subseteq [n-1]$  further define

$$\beta_P(S) = \sum_{T \subseteq S} (-1)^{|S-T|} \alpha_P(T).$$

The function  $\beta_P$  is called the *flag  $h$ -vector* of  $P$ . Knowing  $\alpha_P$  is the same as knowing  $\beta_P$  since

$$\alpha_P(S) = \sum_{T \subseteq S} \beta_P(T).$$

For further information on flag  $f$ -vectors and  $h$ -vectors (using a different terminology), see [29, Ch. 3.12].

There is a kind of generating function for the flag  $h$ -vector which is often useful in understanding the combinatorics of  $P$ . Regarding  $n$  as fixed, let

---

<sup>1</sup>Minor variations of this definition appear in the literature, but they are equivalent to the definition given here. For instance, in [31] parking functions are obtained from the definition given here by subtracting one from each coordinate.

$S \subseteq [n-1]$  and define a formal power series  $Q_S = Q_S(x) = Q_S(x_1, x_2, \dots)$  in the (commuting) indeterminates  $x_1, x_2, \dots$  by

$$Q_S = \sum_{\substack{i_1 \leq i_2 \leq \dots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

$Q_S$  is known as *Gessel's quasisymmetric function* [10] (see also [16, §5.4][18][24, Ch. 9.4]). The functions  $Q_S$ , where  $S$  ranges over all subsets of  $[n-1]$ , are linearly independent over any field. For our ranked poset  $P$  we then define

$$F_P = \sum_{S \subseteq [n-1]} \beta_P(S) Q_S.$$

This definition (in a different but equivalent form) was first suggested by R. Ehrenborg [6, Def. 4.1] and is further investigated in [30]. One of the results of [30] (Thm. 1.4) is the following proposition (which is equivalent to a simple generalization of [29, Exercise 3.65]).

**2.1 Proposition.** *Let  $P$  be as above. If every interval  $[u, v]$  of  $P$  is rank-symmetric (i.e.,  $[u, v]$  has as many elements of rank  $i$  as of corank  $i$ ), then  $F_P$  is a symmetric function of  $x_1, x_2, \dots$ .*

We now consider the case  $P = \text{NC}_{n+1}$ . (We take  $\text{NC}_{n+1}$  rather than  $\text{NC}_n$  because  $\text{NC}_{n+1}$  has rank  $n$ .) It is well-known that every interval in  $\text{NC}_{n+1}$  is self-dual and hence rank-symmetric. (This follows from the fact that  $\text{NC}_{n+1}$  is itself self-dual [15, §3][27, Thm. 1.1] and that every interval of  $\text{NC}_{n+1}$  is a product of  $\text{NC}_i$ 's [21, §1.3].) Hence  $F_{\text{NC}_{n+1}}$  is a symmetric function, and we can ask whether it is already known. In fact,  $F_{\text{NC}_{n+1}}$  has previously appeared in connection with parking functions, as stated below in Theorem 2.3. First we provide some background information related to parking functions.

Let  $\mathcal{P}_n$  denote the set of all parking functions of length  $n$ . The symmetric group  $\mathfrak{S}_n$  acts on  $\mathcal{P}_n$  by permuting coordinates. Let  $\text{PF}_n = \text{PF}_n(x)$  denote the Frobenius characteristic of the character of this action [17, Ch. 1.7]. Thus if

$$\text{PF}_n = \sum_{\lambda \vdash n} \tau_{\lambda, n} s_\lambda$$

is the expansion of  $\text{PF}_n$  in terms of Schur functions, then  $\tau_{\lambda, n}$  is the multiplicity of the irreducible character of  $\mathfrak{S}_n$  indexed by  $\lambda$  in the action of  $\mathfrak{S}_n$  on  $\mathcal{P}_n$ . The symmetric function  $\text{PF}_n$  was first considered in the context of parking functions by Haiman [13, §§2.6 and 4.1]. Following Haiman, we will give a formula for  $\text{PF}_n$  from which its expansion in terms of various symmetric function bases is immediate. The key observation (due to Pollack [8, p. 13] and repeated in [13, p. 28]) is the following (which we state in a slightly different form than Pollack). Let  $\mathbb{Z}_{n+1}$  denote the set  $\{1, 2, \dots, n+1\}$ , with addition modulo  $n+1$ . Then every coset of the subgroup  $H$  of  $\mathbb{Z}_{n+1}^n$  generated by  $(1, 1, \dots, 1)$  contains exactly one parking function. From this it follows easily that

$$\text{PF}_n = \frac{1}{n+1} [t^n] H(t)^{n+1}, \quad (1)$$

where  $[t^n]G(t)$  denotes the coefficient of  $t^n$  in the power series  $G(t)$ , and where

$$H(t) = 1 + h_1t + h_2t^2 + \dots = \frac{1}{(1 - x_1t)(1 - x_2t)\dots},$$

the generating function for the complete symmetric functions  $h_i$ . (Throughout this paper we adhere to symmetric function terminology and notation as in Macdonald [17].)

The following proposition summarizes some of the properties of  $PF_n$  which follow easily from equation (1).

**2.2 Proposition.** (a) *We have the following expansions.*

$$PF_n = \sum_{\lambda \vdash n} (n+1)^{\ell(\lambda)-1} z_\lambda^{-1} p_\lambda \tag{2}$$

$$= \sum_{\lambda \vdash n} \frac{1}{n+1} s_\lambda(1^{n+1}) s_\lambda \tag{3}$$

$$= \sum_{\lambda \vdash n} \frac{1}{n+1} \left[ \prod_i \binom{\lambda_i + n}{n} \right] m_\lambda \tag{4}$$

$$= \sum_{\lambda \vdash n} \frac{n(n-1)\dots(n-\ell(\lambda)+2)}{m_1(\lambda)! \dots m_n(\lambda)!} h_\lambda. \tag{5}$$

$$\omega PF_n = \sum_{\lambda \vdash n} \frac{1}{n+1} \left[ \prod_i \binom{n+1}{\lambda_i} \right] m_\lambda. \tag{6}$$

Here  $s_\lambda(1^{n+1})$  denotes  $s_\lambda$  with  $n+1$  variables set equal to 1 and the others to 0, and is evaluated explicitly e.g. in [17, Example 4, p. 45]. Moreover,  $\ell(\lambda)$  is the number of parts of  $\lambda$ ;  $z_\lambda$  is as in [17, p. 24];  $m_i(\lambda)$  denotes the number of parts of  $\lambda$  equal to  $i$ ; and  $\omega$  is the standard involution [17, pp. 21–22] on symmetric functions.

(b) *We also have that*

$$\sum_{n \geq 0} PF_n t^{n+1} = (tE(-t))^{(-1)}, \tag{7}$$

where  $E(t) = \sum_{n \geq 0} e_n t^n$ ,  $e_n$  denotes the  $n$ th elementary symmetric function, and  $\langle^{-1}\rangle$  denotes compositional inverse.

**Proof.** (a) Let  $C(x, y) = \prod_{i,j} (1 - x_i y_j)^{-1}$ , the well-known ‘‘Cauchy product.’’ Then  $H(t)^{n+1}$  is obtained by setting  $n+1$  of the  $y_i$ ’s equal to  $t$  and the others equal to 0. From this all the expansions in (a) follow from (1) and well-known expansions for  $C(x, y)$  and  $\omega_x C(x, y)$  (where  $\omega_x$  denotes  $\omega$  acting on the  $x$  variables only). To give just one example (needed in the first proof of Theorem 2.3), we have

$$\omega_x C(x, y) = \sum_{\lambda} m_\lambda(x) e_\lambda(y).$$

Hence

$$\omega \text{PF}_n = \sum_{\lambda \vdash n} \frac{1}{n+1} e_\lambda(1^{n+1}) m_\lambda.$$

Equation (6) now follows from the simple fact that  $e_k(1^{n+1}) = \binom{n+1}{k}$ . We should point out that (2) appears (in a dual form) in [11, (9)], (3) appears in [13, (28)], and (5) appears (again in dual form) in [13, (82)][17, Example 24(a), p. 35]. A  $q$ -analogue of  $\text{PF}_n$  and of much of our Proposition 2.2 appears in [9].

(b) This is an immediate consequence of (1), the fact that

$$\frac{1}{H(t)} = E(-t), \quad (8)$$

and the Lagrange inversion formula, as in [13, §4.1]. See also [17, Examples 2.24–2.25, pp. 35–36].  $\square$

Let  $P$  be a Cohen-Macaulay poset with  $\hat{0}$  and  $\hat{1}$  such that every interval is rank-symmetric. Thus  $F_P$  is a symmetric function. In [30, Conj. 2.3] it was conjectured that  $F_P$  is *Schur positive*, i.e., a nonnegative linear combination of Schur functions. Equation (3) confirms this conjecture in the case  $P = \text{NC}_{n+1}$ . However, it turns out that the conjecture is in fact *false*. A counterexample is provided by the following poset  $P$ . The elements of  $P$  consist of all integer vectors  $(a_1, a_2, b_1, b_2, b_3, b_4)$  such that  $0 \leq a_1 \leq 5$ ,  $0 \leq a_2 \leq 1$ ,  $0 \leq b_1 \leq 3$ ,  $0 \leq b_2, b_3, b_4 \leq 1$ , and  $a_1 + a_2 = b_1 + b_2 + b_3 + b_4$ , ordered componentwise. It can be shown that  $P$  is lexicographically shellable and hence Cohen-Macaulay, and it is easy to see that  $P$  is locally rank-symmetric (even locally self-dual). Moreover,

$$F_P = s_6 + 7s_{51} + 6s_{42} + 2s_{33} + 18s_{411} + 10s_{321} - s_{222} + 20s_{3111} + 5s_{2211} + 8s_{21111}.$$

The symmetric functions  $\text{PF}_n$  also have an unexpected connection with the multiplication of conjugacy classes in the symmetric group (the work of Farahat-Higman [7]). For further details see [11][17, Ch. I, Example 7.25, pp. 132–134]. This connection was exploited by Goulden and Jackson [11] to compute some connection coefficients for the symmetric group.

The expansion (5) of  $\text{PF}_n$  in terms of the  $h_\lambda$ 's has a simple interpretation in terms of parking functions. Suppose that  $a = (a_1, \dots, a_n) \in \mathcal{P}_n$ . Let  $r_1, \dots, r_k$  be the positive multiplicities of the elements of the multiset  $\{a_1, \dots, a_n\}$  (so  $r_1 + \dots + r_k = n$ ). Then the action of  $\mathfrak{S}_n$  on the orbit  $\mathfrak{S}_n a$  has characteristic  $h_{r_1} \cdots h_{r_k}$ . For instance, a set of orbit representatives in the case  $n = 3$  is  $(1, 1, 1), (2, 1, 1), (3, 1, 1), (2, 2, 1)$ , and  $(3, 2, 1)$ . Hence  $\text{PF}_3 = h_3 + h_2 h_1 + h_2 h_1 + h_2 h_1 + h_1^3 = h_3 + 3h_{21} + h_{111}$ . In general it follows that the coefficient  $q_\lambda$  of  $h_\lambda$  in  $\text{PF}_n$  is equal to the number of orbits of parking functions of length  $n$  such that the terms of their elements have multiplicities  $\lambda_1, \lambda_2, \dots$  (in some order). Equation (5) then gives an explicit formula for this number. The *total* number of parking functions whose terms have multiplicities  $\lambda_1, \lambda_2, \dots$  is  $q_\lambda$  times the size of the orbit, i.e.,  $q_\lambda \binom{n}{\lambda_1, \lambda_2, \dots}$ .

We are now ready to discuss the connection between  $\text{PF}_n$  and noncrossing partitions. The basic result is the following.

**2.3 Theorem.** *For any  $n \geq 0$  we have*

$$F_{\text{NC}_{n+1}} = \omega \text{PF}_n.$$

**Proof.** Let  $\lambda = (\lambda_1, \dots, \lambda_\ell)$  be a partition of  $n$  with  $\lambda_\ell > 0$ . It is immediate from the definition of  $F_P$  in Section 1 (see [30, Prop. 1.1]) that if  $F_P$  is symmetric and  $F_P = \sum_\lambda c_\lambda m_\lambda$ , then

$$c_\lambda = \alpha_P(\lambda_1, \lambda_1 + \lambda_2, \dots, \lambda_1 + \lambda_2 + \dots + \lambda_{\ell-1}). \quad (9)$$

The proof now follows by comparing equation (6) with the evaluation of  $\alpha_{\text{NC}_{n+1}}(S)$  due to Edelman [4, Thm. 3.2].  $\square$

It follows from the above discussion that  $\text{PF}_n$  encodes in a simple way the flag  $f$ -vector and flag  $h$ -vector of  $\text{NC}_{n+1}$ , viz., (1) the coefficient of  $Q_S$  in the expansion of  $\text{PF}_n$  in terms of Gessel's quasisymmetric function is equal to  $\beta_{\text{NC}_{n+1}}(S)$ , and (2) if the elements of  $S \subseteq [n-1]$  are  $j_1 < \dots < j_r$  and if  $\lambda$  is the partition whose parts are the numbers  $j_1, j_2 - j_1, j_3 - j_2, \dots, n - j_r$ , then the coefficient of  $m_\lambda$  in the expansion of  $\text{PF}_n$  in terms of monomial symmetric functions is equal to  $\alpha_{\text{NC}_{n+1}}(S)$ . There is a further statistic on  $\text{NC}_{n+1}$  closely related to  $\text{PF}_n$ , namely, the number of noncrossing partitions of  $[n+1]$  of type  $\lambda$ , i.e., with block sizes  $\lambda_1, \lambda_2, \dots$

**2.4 Proposition.** *Let  $\lambda$  be a partition of  $n$ . The coefficient of  $h_\lambda$  in the expansion of  $\text{PF}_n$  in terms of complete symmetric functions is equal to the number  $u_\lambda$  of noncrossing partitions of type  $\lambda$ .*

**First proof.** Compare equation (5) with the explicit value of the number of noncrossing partitions of type  $\lambda$  found by Kreweras [15, Thm. 4].  $\square$

**Second proof.** Our second proof is based on the following noncrossing analogue of the exponential formula due to Speicher [28, p. 616]. (For a more general result, see [21].) Given a function  $f : \mathbb{N} \rightarrow R$  (where  $R$  is a commutative ring with identity) with  $f(0) = 1$ , define a function  $g : \mathbb{N} \rightarrow R$  by  $g(0) = 1$  and

$$g(n) = \sum_{\pi = \{B_1, \dots, B_k\} \in \text{NC}_n} f(\#B_1) \cdots f(\#B_k). \quad (10)$$

Let  $F(t) = \sum_{n \geq 0} f(n)t^n$ . Then

$$\sum_{n \geq 0} g(n)t^{n+1} = \left( \frac{t}{F(t)} \right)^{\langle -1 \rangle}. \quad (11)$$

In equation (10) take  $f(n) = h_n$ , the complete symmetric function. Then  $g(n)$  becomes  $\sum_{\lambda \vdash n} u_\lambda h_\lambda$ . But Proposition 2.2(b), together with equations (11) and (8), shows that  $g(n) = \text{PF}_n$ , and the proof follows.  $\square$

Note the curious fact that Theorem 2.3 refers to  $\text{NC}_{n+1}$ , while Proposition 2.4 refers to  $\text{NC}_n$ . Proposition 2.4, together with the definition of  $\text{PF}_n$ ,

show that the number of noncrossing partitions of type  $\lambda \vdash n$  is equal to the number of  $\mathfrak{S}_n$ -orbits of parking functions of length  $n$  and part multiplicities  $\lambda$ . It is easy to give a bijective proof of this fact (shown to me by R. Simion), which we omit.

**3. An edge labeling of the noncrossing partition lattice.** If  $P$  is a locally finite poset, then an *edge* of  $P$  is a pair  $(u, v) \in P \times P$  such that  $v$  covers  $u$  (i.e.,  $u < v$  and no element  $t$  satisfies  $u < t < v$ ). An *edge labeling* of  $P$  is a map  $\Lambda : \mathcal{E}(P) \rightarrow \mathbb{Z}$ , where  $\mathcal{E}(P)$  is the set of edges of  $P$ . Edge labelings of posets have many applications; in particular, if  $P$  has what is known as an *EL-labeling*, then  $P$  is lexicographically shellable and hence Cohen-Macaulay [2][3]. An EL-labeling of  $\text{NC}_{n+1}$  was defined by Björner [2, Example 2.9] and further exploited by Edelman and Simion [5]. Here we define a new labeling, which up to an unimportant reindexing is EL and is intimately related to parking functions.

Let  $(\pi, \sigma)$  be an edge of  $\text{NC}_{n+1}$ . Thus  $\sigma$  is obtained from  $\pi$  by merging together two blocks  $B$  and  $B'$ . Suppose that  $\min B < \min B'$ , where  $\min S$  denotes the minimum element of a finite set  $S$  of integers. Define

$$\Lambda(\pi, \sigma) = \max\{i \in B : i < B'\}, \quad (12)$$

where  $i < B'$  denotes that  $i$  is less than every element of  $B'$ . For instance, if  $B = \{2, 4, 5, 15, 17\}$  and  $B' = \{7, 10, 12, 13\}$ , then  $\Lambda(\pi, \sigma) = 5$ . Note that  $\Lambda(\pi, \sigma)$  always exists since  $\min B < B'$ .

The labeling  $\Lambda$  of the edges of  $\text{NC}_{n+1}$  extends in a natural (and well-known) way to a labeling of the maximal chains. Namely, if  $\mathfrak{m} : \hat{0} = \pi_0 < \pi_1 < \dots < \pi_n = \hat{1}$  is a maximal chain of  $\text{NC}_{n+1}$ , then set

$$\Lambda(\mathfrak{m}) = (\Lambda(\pi_0, \pi_1), \Lambda(\pi_1, \pi_2), \dots, \Lambda(\pi_{n-1}, \pi_n)).$$

**3.1 Theorem.** *The labels  $\Lambda(\mathfrak{m})$  of the maximal chains of  $\text{NC}_{n+1}$  consist of the parking functions of length  $n$ , each occurring once.*

**Proof.** If  $\Lambda(\pi_j, \pi_{j+1}) = i$ , then the block of  $\pi_{j+1}$  containing  $i$  also contains an element  $k > i$ . Hence the number of  $j$  for which  $\Lambda(\pi_j, \pi_{j+1}) = i$  cannot exceed  $n + 1 - i$ , from which it follows that  $\Lambda(\mathfrak{m})$  is a parking function.

Suppose that  $\mathfrak{m}$  and  $\mathfrak{m}'$  are maximal chains of  $\text{NC}_{n+1}$  for which  $\Lambda(\mathfrak{m}) = \Lambda(\mathfrak{m}')$ . We will prove by induction on  $n$  that  $\mathfrak{m} = \mathfrak{m}'$ . The assertion is clear for  $n = 0$ . Assume true for  $n - 1$ . Let the elements of  $\mathfrak{m}$  be  $\hat{0} = \pi_0 < \pi_1 < \dots < \pi_n = \hat{1}$ . Suppose that  $\Lambda(\mathfrak{m}) = (a_1, \dots, a_n)$ . Let  $r = \max\{a_i : 1 \leq i \leq n\}$ , and let  $s = \max\{i : a_i = r\}$ . We claim that one of the blocks of  $\pi_{s-1}$  is just the singleton set  $\{r + 1\}$ . If  $r$  and  $r + 1$  are in the same block of  $\pi_{s-1}$ , then we can't have  $\Lambda(\pi_{s-1}, \pi_s) = r$ , contradicting  $a_s = r$ . Hence  $r$  and  $r + 1$  are in different blocks of  $\pi_{s-1}$ . If the block  $B$  of  $\pi_{s-1}$  containing  $r + 1$  contained some element  $t < r$ , then by the noncrossing property and the fact that  $a_s = r$  we have that  $B$  is merged with the block  $B_1$  of  $\pi_{s-1}$  containing  $r$  to get  $\pi_s$ . But  $\min B \leq t < r \in B_1$ , contradicting  $a_s = r$ . Hence every element of  $B$  is greater than  $r$ . If  $B$  contained some element  $t > r + 1$ , then (since  $r + 1 = \min B$ ) we

would have  $a_k = r + 1$  for some  $k < r$ , contradicting maximality of  $r$ . This proves the claim.

We next claim that  $\pi_s$  is obtained from  $\pi_{s-1}$  by merging the block  $B_1$  containing  $r$  with the block  $\{r + 1\}$ . Otherwise (since  $a_s = r$ )  $\pi_s$  is obtained by merging  $B_1$  with some block  $B_2$  all of whose elements are greater than  $r + 1$ . For some  $t > s$  we must obtain  $\pi_t$  from  $\pi_{t-1}$  by merging the block  $B_3$  containing  $r + 1$  with the block  $B_4$  containing  $r$ . Now  $B_3$  can't contain an element less than  $r + 1$  by the noncrossing property of  $\pi_{s-1}$  (since  $B_4$  contains both  $r$  and an element greater than  $r + 1$ ). It follows that  $\Lambda(\pi_{t-1}, \pi_t) = r$ , contradicting the maximality of  $s$  and proving the claim.

It is now clear by induction that the chain  $\mathfrak{m}$  can be uniquely recovered from the parking function  $\Lambda(\mathfrak{m}) = (a_1, \dots, a_n)$ . Namely, let  $a'$  be the sequence obtained from  $\Lambda(\mathfrak{m})$  by removing  $a_s$ . Then  $a'$  is a parking function of length  $n - 1$ . By induction there is a unique maximal chain  $\mathfrak{m}^* : \hat{0} = \pi_0^* < \pi_1^* < \dots < \pi_{n-1}^* = \hat{1}$  of  $\text{NC}_n$  such that  $\Lambda(\mathfrak{m}^*) = a'$ . By the discussion above we can then obtain  $\mathfrak{m}$  uniquely from  $\mathfrak{m}^*$  by (1) replacing each element  $i > r$  of the ambient set  $[n]$  with  $i + 1$ , (2) adjoining a singleton block  $\{r + 1\}$  to each  $\pi_i^*$  for  $i \leq s - 1$ , (3) inserting between  $\pi_{s-1}^*$  and  $\pi_s^*$  a new element obtained from  $\pi_{s-1}^*$  by merging the block containing  $r$  with the singleton block  $\{r + 1\}$ , and (4) for  $i > s$  adjoining the element  $r + 1$  to the block of  $\pi_i^*$  containing  $r$ . Hence we have shown that if  $\Lambda(\mathfrak{m}) = \Lambda(\mathfrak{m}')$ , then  $\mathfrak{m} = \mathfrak{m}'$ . But it is known [15, Cor. 5.2][4, Cor. 3.3] that  $\text{NC}_{n+1}$  has  $(n + 1)^{n-1}$  maximal chains, which is just the number of parking functions of length  $n$  [14, Lemma 1 and §6][8]. Thus every parking function of length  $n$  occurs exactly once among the sequences  $\Lambda(\mathfrak{m})$ , and the proof is complete.  $\square$

The above proof of the injectivity of the map  $\Lambda$  from maximal chains to parking functions is reminiscent of the proof [20, p. 5] that the Prüfer code of a labelled tree determines the tree. Our proof “cheated” by using the fact that the number of maximal chains is the number of parking functions. We only gave a direct proof of the injectivity of  $\Lambda$ . However, our proof actually suffices to show also surjectivity since the argument of the above paragraph is valid for any parking function, the key point being that removing an occurrence of the largest element of a parking function preserves the property of being a parking function.

If we define a new labeling  $\Lambda^*$  of  $\text{NC}_{n+1}$  by

$$\Lambda^*(\pi, \sigma) = |\pi| - \Lambda(\pi, \sigma),$$

where  $|\pi|$  is the number of blocks of  $\pi$ , then it is easy to check (using the fact that every interval of  $\text{NC}_{n+1}$  is a product of  $\text{NC}_i$ 's) that every interval  $[\pi, \tau]$  has a unique maximal chain  $\mathfrak{m} : \pi = \pi_0 < \pi_1 < \dots < \pi_j = \tau$  such that

$$\Lambda^*(\pi_0, \pi_1) \leq \Lambda^*(\pi_1, \pi_2) \leq \dots \leq \Lambda^*(\pi_{k-1}, \pi_k).$$

In other words,  $\Lambda^*$  is an *R-labeling* in the sense of [29, Def. 3.13.1]. Moreover, this maximal chain  $\mathfrak{m}$  has the lexicographically least label  $\Lambda^*(\mathfrak{m})$  of any maximal

chain of the interval  $[\pi, \tau]$ . Thus  $\Lambda^*$  is in fact an *EL-labeling*, as defined in [2, Def. 2.2] (though there it is called just an “L-labeling.”). For the significance of the EL-labeling property, see the first paragraph of this section. Here we will just be concerned with the weaker R-labeling property.

Define the *descent set*  $D(a)$  of a parking function  $a = (a_1, \dots, a_n)$  by

$$D(a) = \{i : a_i > a_{i+1}\}.$$

From the fact that  $\Lambda^*$  is an R-labeling and [29, Thm. 3.13.2], we obtain the following proposition.

**3.2 Proposition.** (a) *Let  $S \subseteq [n - 1]$ . The number of parking functions  $a$  of length  $n$  satisfying  $D(a) = S$  is equal to  $\beta_{\text{NC}_{n+1}}([n - 1] - S)$ .*

(b) *Let  $S \subseteq [n - 1]$ . The number of parking functions  $a$  of length  $n$  satisfying  $D(a) \supseteq S$  is equal to  $\alpha_{\text{NC}_{n+1}}([n - 1] - S)$ . This number is given explicitly by [4, Thm. 3.2] or by equations (4) and (9).*

The labeling  $\Lambda$  is closely related to a bijection between the maximal chains of  $\text{NC}_{n+1}$  and labelled trees, different from the earlier bijection of Edelman [4, Cor. 3.3]. Let  $\mathbf{m} : \hat{0} = \pi_0 < \pi_1 < \dots < \pi_n = \hat{1}$  be a maximal chain of  $\text{NC}_{n+1}$ . Define a graph  $\Gamma_{\mathbf{m}}$  on the vertex set  $[n + 1]$  as follows. There will be an edge  $e_i$  for each  $1 \leq i \leq n$ . Suppose that  $\pi_i$  is obtained from  $\pi_{i-1}$  by merging blocks  $B$  and  $B'$  with  $\min B < \min B'$ . Then the vertices of  $e_i$  are defined to be  $\Lambda(\pi_{i-1}, \pi_i)$  and  $\min B'$ . It is easy to see that  $\Gamma_{\mathbf{m}}$  is a tree. Root  $\Gamma_{\mathbf{m}}$  at the vertex 1 and erase the vertex labels. If  $v_i$  is the vertex of  $e_i$  farthest from the root, then move the label  $i$  of the edge  $e_i$  from  $e_i$  to the vertex  $v_i$ . Label the root with 0 and unroot the tree. We obtain a labelled tree  $T_{\mathbf{m}}$  on  $n + 1$  vertices, and one can easily check that the map  $\mathbf{m} \mapsto T_{\mathbf{m}}$  is a bijection between maximal chains of  $\text{NC}_{n+1}$  and labelled trees on  $n + 1$  vertices.

**4. A local action of the symmetric group.** Suppose that  $P$  is a graded poset of rank  $n$  with  $\hat{0}$  and  $\hat{1}$  such that  $F_P$  is a symmetric function. If  $F_P$  is Schur positive, then it is the Frobenius characteristic of a representation of  $\mathfrak{S}_n$  whose dimension is the number of maximal chains of  $P$ . Thus we can ask whether there is some “nice” representation of  $\mathfrak{S}_n$  on the vector space  $V_P$  (over a field of characteristic zero) whose basis is the set of maximal chains of  $P$ . This question was discussed in [30, §5]. A “nice” representation should somehow reflect the poset structure. With this motivation, an action of  $\mathfrak{S}_n$  on  $V_P$  is defined to be *local* [30, §5] if for every adjacent transposition  $\sigma_i = (i, i + 1)$  and every maximal chain

$$\mathbf{m} : \hat{0} = t_0 < t_1 < \dots < t_n = \hat{1}, \tag{13}$$

we have that  $\sigma_i(\mathbf{m})$  is a linear combination of maximal chains of the form  $t_0 < t_1 < \dots < t_{i-1} < t'_i < t_{i+1} < \dots < t_n$ , i.e., of maximal chains which agree with  $\mathbf{m}$  except possibly at  $t_i$ .

Now let  $P = \text{NC}_{n+1}$ . Every interval  $[\pi, \tau]$  of  $\text{NC}_{n+1}$  of length two contains either two or three elements in its middle level. In the latter case, there are three blocks  $B_1, B_2, B_3$  of  $\pi$  such that  $\tau$  is obtained from  $\pi$  by merging  $B_1, B_2, B_3$

into a single block. Moreover, any two of these blocks can be merged to form a noncrossing partition. Let  $\pi_{ij}$  be the noncrossing partition obtained by merging  $B_i$  and  $B_j$ , so that the middle elements of the interval  $[\pi, \tau]$  are  $\pi_{12}, \pi_{13}, \pi_{23}$ . Exactly one of these partitions  $\pi_{ij}$  will have the property that  $\Lambda(\pi, \pi_{ij}) = \Lambda(\pi_{ij}, \tau)$ , where  $\Lambda$  is defined by (12). Let us call this partition  $\pi_{ij}$  the *special* element of the interval  $[\pi, \tau]$ . Now define linear transformations  $\sigma'_i : V_{\text{NC}_{n+1}} \rightarrow V_{\text{NC}_{n+1}}$ ,  $1 \leq i \leq n - 1$  as follows. Let  $\mathfrak{m}$  be a maximal chain of  $\text{NC}_{n+1}$  with elements  $\hat{0} = \pi_0 < \pi_1 < \dots < \pi_n = \hat{1}$ .

*Case 1.* The interval  $[\pi_{i-1}, \pi_{i+1}]$  contains exactly two middle elements  $\pi_i$  and  $\pi'_i$ . Then set  $\sigma'_i(\mathfrak{m}) = \mathfrak{m}'$ , where  $\mathfrak{m}'$  is given by  $\pi_0 < \pi_1 < \dots < \pi_{i-1} < \pi'_i < \pi_{i+1} < \dots < \pi_n$ .

*Case 2.* The interval  $[\pi_{i-1}, \pi_{i+1}]$  contains exactly three middle elements, of which  $\pi_i$  is special. Then set  $\sigma'_i(\mathfrak{m}) = \mathfrak{m}$ .

*Case 3.* The interval  $[\pi_{i-1}, \pi_{i+1}]$  contains exactly three middle elements  $\pi_i, \pi'_i$ , and  $\pi''_i$ , of which  $\pi''_i$  is special. Then set  $\sigma'_i(\mathfrak{m}) = \mathfrak{m}'$ , where  $\mathfrak{m}'$  is given by  $\pi_0 < \pi_1 < \dots < \pi_{i-1} < \pi'_i < \pi_i < \pi_{i+1} < \dots < \pi_n$ .

**4.1 Proposition.** *The action of each  $\sigma'_i$  on  $V_{\text{NC}_{n+1}}$  defined above yields a local action of  $\mathfrak{S}_n$  on  $V_{\text{NC}_{n+1}}$ . Equivalently, there is a homomorphism  $\varphi : \mathfrak{S}_n \rightarrow \text{GL}(V_{\text{NC}_{n+1}})$  satisfying  $\varphi(\sigma_i) = \sigma'_i$ . The Frobenius characteristic of this action is given by  $\text{PF}_n$ .*

**Proof.** Each maximal chain  $\mathfrak{m}$  corresponds to a parking function  $\Lambda(\mathfrak{m})$  via Theorem 3.1. Thus the natural action of  $\mathfrak{S}_n$  on  $\mathcal{P}_n$  defined in Section 2 may be “transferred” to an action  $\psi$  of  $\mathfrak{S}_n$  on the set of maximal chains of  $\text{NC}_{n+1}$ . It is easy to check that  $\psi$  and  $\varphi$  agree on the  $\sigma_i$ ’s, and the proof follows.  $\square$

The action  $\varphi$  does not quite have the property mentioned at the beginning of this section that its characteristic is  $F_{\text{NC}_{n+1}}$ . By Theorem 2.3, the characteristic is actually  $\omega F_{\text{NC}_{n+1}}$ . However, we only have to multiply  $\varphi$  by the sign character (equivalently, define a new action  $\varphi'$  by  $\varphi'(\sigma_i) = -\varphi(\sigma_i)$ ) to get the desired property.

It is rather surprising that the simple “local” definition we have given of  $\varphi$  defines an action of  $\mathfrak{S}_n$ . Perhaps it would be interesting to look for some more examples. (We need to exclude trivial examples such as  $w(\mathfrak{m}) = \mathfrak{m}$  for all  $w \in \mathfrak{S}_n$  and all maximal chains  $\mathfrak{m}$ .) A few other examples appear in the next section and in [30, §5]. A further example (the posets of shuffles of  $C$ . Greene [12]) is discussed in [26] together with the rudiments of a systematic theory of such actions, but much work needs to be done for a satisfactory understanding of local  $\mathfrak{S}_n$ -actions.

**5. Generalizations.** In this section we will briefly discuss two generalizations of what appears above. All proofs are entirely analogous and will be omitted. Fix an integer  $k \in \mathbb{P}$ . A *k-divisible noncrossing partition* is a noncrossing partition  $\pi$  for which every block size is divisible by  $k$ . Thus  $\pi$  is a noncrossing partition of a set  $[kn]$  for some  $n \geq 0$ . Let  $\text{NC}_n^{(k)}$  be the poset of all  $k$ -divisible noncrossing partitions of  $[kn]$ . ( $\text{NC}_n^{(k)}$  is actually a join-semilattice of  $\text{NC}_{kn}$ . It has  $\hat{1}$  but not a  $\hat{0}$  when  $k > 1$ .) The combinatorial properties of

the poset  $\text{NC}_n^{(k)}$  were first considered by Edelman [4, §4]. If a pair  $(\pi, \sigma)$  is an edge of  $\text{NC}_n^{(k)}$  (i.e.,  $\sigma$  covers  $\pi$  in  $\text{NC}_n^{(k)}$ ), then  $(\pi, \sigma)$  is an edge of  $\text{NC}_n$ . Hence the edge-labeling  $\Lambda$  of  $\text{NC}_n$  restricts to an edge-labeling of  $\text{NC}_n^{(k)}$ .

Define a *k-parking function* of length  $n$  to be a sequence  $(a_1, \dots, a_n)$  of positive integers such that if  $b_1 \leq b_2 \leq \dots \leq b_n$  is the increasing rearrangement of  $a_1, \dots, a_n$ , then  $b_i \leq ki$ . Let  $\mathcal{P}_n^{(k)}$  denote the set of all *k-parking functions* of length  $n$ . The argument of Pollack mentioned in Section 2 that  $\#\mathcal{P}_n = (n + 1)^{n-1}$  easily extends to  $\mathcal{P}_n^{(k)}$ . Namely, let  $\mathbb{Z}_{k(n+1)}$  denote the set  $\{1, 2, \dots, k(n+1)\}$  with addition modulo  $k(n+1)$ . Then every coset of the subgroup  $H$  of  $\mathbb{Z}_{k(n+1)}^n$  generated by  $(1, 1, \dots, 1)$  contains exactly  $k$  *k-parking functions*. Hence  $\#\mathcal{P}_n^{(k)} = k(k(n+1))^{n-1}$ . The symmetric group  $\mathfrak{S}_n$  acts on  $\mathcal{P}_n^{(k)}$  by permuting coordinates, and we can consider its Frobenius characteristic  $\text{PF}_n^{(k)}$  just as we did for  $\mathcal{P}_n$ . The above generalization of Pollack's argument shows that

$$\text{PF}_n^{(k)} = \frac{1}{n+1} [t^n] H(t)^{k(n+1)}.$$

Proposition 2.2 generalizes straightforwardly to the case of  $\text{PF}_n^{(k)}$ . In particular, Proposition 2.2(b) takes the form

$$\sum_{n \geq 0} \text{PF}_n^{(k)} t^{n+1} = \left( tE(-t)^k \right)^{(-1)}.$$

Theorem 3.1 generalizes as follows.

**5.1 Theorem.** *The labels  $\Lambda(\mathfrak{m})$  of the maximal chains of  $\text{NC}_{n+1}^{(k)}$  consist of the *k-parking functions* of length  $n$ , each occurring once.*

Proposition 3.2 requires some modification when extended to *k-parking functions* because the posets  $\text{NC}_{n+1}^{(k)}$  do not have a  $\hat{0}$  when  $k > 1$ . For these posets we regard the minimal elements as having rank 0, and we define  $\alpha_{\text{NC}_{n+1}^{(k)}}(S)$  and  $\beta_{\text{NC}_{n+1}^{(k)}}(S)$  for  $S \subseteq \{0, 1, \dots, n-1\}$ . Thus for instance  $\alpha_{\text{NC}_{n+1}^{(k)}}(\emptyset) = \beta_{\text{NC}_{n+1}^{(k)}}(\emptyset) = 1$ , and  $\alpha_{\text{NC}_{n+1}^{(k)}}(0)$  is the number of minimal elements of  $\text{NC}_{n+1}^{(k)}$ . Write  $[0, n-1] = \{0, 1, \dots, n-1\}$ .

**5.2 Proposition.** (a) *Let  $S \subseteq [n-1]$ . The number of *k-parking functions*  $a$  of length  $n$  satisfying  $D(a) = S$  is equal to*

$$\beta_{\text{NC}_{n+1}^{(k)}}([n-1] - S) + \beta_{\text{NC}_{n+1}^{(k)}}([0, n-1] - S).$$

(b) *Let  $S \subseteq [n-1]$ . The number of *k-parking functions*  $a$  of length  $n$  satisfying  $D(a) \supseteq S$  is equal to  $\alpha_{\text{NC}_{n+1}^{(k)}}([0, n-1] - S)$ . This number is given explicitly by [4, Thm. 4.2].*

Note, however, that there does not seem to be a nice generalization of Theorem 2.3. The quasisymmetric function  $F_{\text{NC}_{n+1}^{(k)}}$  is not a symmetric function when  $k > 1$ , and we know of no simple connection between the flag *f*-vector

of  $\text{NC}_{n+1}^{(k)}$  and the symmetric function  $\text{PF}_n^{(k)}$ , nor between the number of  $k$ -divisible noncrossing partitions of a given type and  $\text{PF}_n^{(k)}$ .

Proposition 4.1 extends straightforwardly to  $\text{NC}_{n+1}^{(k)}$ . The natural action of  $\mathfrak{S}_n$  on  $\mathcal{P}_n^{(k)}$  is transferred *via* Theorem 5.1 to an action on  $V_{\text{NC}_{n+1}^{(k)}}$ . This action is a permutation representation on the maximal chains, and is readily seen to be local. Its characteristic is  $\text{PF}_n^{(k)}$ .

There is a different generalization of noncrossing partitions due to Reiner [23] (a special case had earlier appeared in a different guise in [19], as explained in [23]) that we have not looked at very closely. Reiner regards ordinary noncrossing partitions as corresponding to the root system  $A_n$  and constructs analogues for the root systems  $B_n$  and  $D_n$ . Actually, for every subset  $S \subseteq [n]$  he constructs a lattice  $\text{NC}_n^{BD}(S)$  interpolating between the  $B_n$  analogue (the case  $S = \emptyset$ ) and the  $D_n$  analogue (the case  $S = [n]$ ). The lattices  $\text{NC}_n^{BD}(S)$  do not always have self-dual intervals [23, Remark on p. 13], but at least every interval is rank-symmetric. Thus by Proposition 2.1, this implies that  $\text{NC}_n^{BD}(S)$  is a symmetric function. This symmetric function only depends on the cardinality  $s$  of  $S$ , so we write  $F_n^{BD}(s)$  for  $F_{\text{NC}_n^{BD}(S)}$ . We also write  $F_n^B$  for  $F_n^{BD}(0)$ .

Let  $[n]^n$  denote the set of all sequences  $(a_1, \dots, a_n)$  of positive integers with  $a_i \leq n$ . We call such a sequence a  $B_n$ -parking function, for the following reason. Let  $\text{PF}_n^B$  be the Frobenius characteristic of the action of  $\mathfrak{S}_n$  on  $[n]^n$  obtained by permuting coordinates. It follows from [23, Prop. 7] that

$$F_n^B = \omega \text{PF}_n^B.$$

Thus in analogy with Theorem 2.3 it makes sense to think of the elements of  $[n]^n$  as  $B_n$ -parking functions. Reiner's result [23, Prop. 7] makes it easy to give an analogue of Proposition 2.2. Let us simply mention the formula

$$\text{PF}_n^B = [t^n]H(t)^n,$$

from which the analogues of all parts of Proposition 2.2 follow easily. In particular, the analogue of Proposition 2.2(b) takes the form

$$\sum_{n \geq 1} \text{PF}_n^B \frac{t^n}{n} = \log \frac{(tE(-t))^{(-1)}}{t}. \quad (14)$$

Comparing Proposition 2.2 with equation (14) yields the curious result that

$$\exp \sum_{n \geq 1} \text{PF}_n^B \frac{t^n}{n} = \sum_{n \geq 0} \text{PF}_n t^n.$$

What is missing from the analogy between  $A_n$  and  $B_n$  noncrossing partitions is the analogue of Theorem 3.1, i.e., a labeling of  $\text{NC}_{n+1}^B$  such that the labels of the maximal chains are the  $B_n$ -parking functions. We have not looked at this question and recommend it as an interesting open problem.

For the general case of  $\text{NC}_n^{BD}(S)$ , it follows from [23, Thm. 11] that

$$F_n^{BD}(s) = F_n^B - s \cdot \text{PF}'_n,$$

where  $\text{PF}'_n$  is the Frobenius characteristic of the action of  $\mathfrak{S}_n$  on all sequences  $(a_1, \dots, a_n)$  of positive integers whose increasing rearrangement  $b_1 \leq \dots \leq b_n$  satisfies  $b_1 = 1$  and  $b_i \leq i - 1$  for  $2 \leq i \leq n$ . We have not considered further properties of the symmetric function  $F_n^{BD}(s)$ .

## References

- [1] H. W. Becker, Planar rhyme schemes, *Bull. Amer. Math. Soc.* **58** (1952), 39.
- [2] A. Björner, Shellable and Cohen-Macaulay partially ordered sets, *Trans. Amer. Math. Soc.* **260** (1980), 159–183.
- [3] A. Björner and M. Wachs, On lexicographically shellable posets, *Trans. Amer. Math. Soc.* **277** (1983), 323–341.
- [4] P. H. Edelman, Chain enumeration and non-crossing partitions, *Discrete Math.* **31** (1980), 171–180.
- [5] P. H. Edelman and R. Simion, Chains in the lattice of noncrossing partitions, *Discrete Math.* **126** (1994), 107–119.
- [6] R. Ehrenborg, On posets and Hopf algebras, *Advances in Math.* **119** (1996), 1–25.
- [7] H. K. Farahat and G. Higman, The centres of symmetric group rings, *Proc. Royal Soc. (A)* **250** (1959), 212–221.
- [8] D. Foata and J. Riordan, Mappings of acyclic and parking functions, *aequationes math.* **10** (1974), 10–22.
- [9] A. M. Garsia and M. Haiman, A remarkable  $q, t$ -Catalan sequence and  $q$ -Lagrange inversion, *J. Algebraic Combinatorics* **5** (1996), 191–244.
- [10] I. Gessel, Multipartite  $P$ -partitions and inner products of skew Schur functions, *Contemporary Math.* **34** (1984), 289–301.
- [11] I. P. Goulden and D. M. Jackson, Symmetric functions and Macdonald's result for top connexion coefficients in the symmetric group, *J. Algebra* **166** (1994), 364–378.
- [12] C. Greene, Posets of shuffles, *J. Combinatorial Theory (A)* **46** (1988), 191–206.
- [13] M. Haiman, Conjectures on the quotient ring by diagonal invariants, *J. Alg. Combinatorics* **3** (1994), 17–76.
- [14] A. G. Konheim and B. Weiss, An occupancy discipline and applications, *SIAM J. Applied Math.* **14** (1966), 1266–1274.

- [15] G. Kreweras, Sur les partitions non croisées d'un cycle, *Discrete Math.* **1** (1972), 333–350.
- [16] D. Krob and J.-Y. Thibon, Noncommutative symmetric functions IV: Quantum linear groups and Hecke algebras at  $q = 0$ , Institut Blaise Pascal, Laboratoire Informatique Théorique et Programmation, preprint LITP 96/12, March, 1996.
- [17] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, second ed., Oxford University Press, Oxford, 1995.
- [18] C. Malvenuto and C. Reutenauer, Duality between quasi-symmetric functions and the Solomon descent algebra, *J. Algebra* **177** (1995), 967–982.
- [19] C. Montenegro, The fixed point non-crossing partition lattices, preprint, 1993.
- [20] J. W. Moon, *Counting Labelled Trees*, Canadian Mathematical Monographs, No. 1, Canadian Mathematical Congress, 1970.
- [21] A. Nica and R. Speicher, A “Fourier transform” for multiplicative functions on non-crossing partitions, *J. Algebraic Combinatorics*, to appear.
- [22] Y. Poupard, Étude et dénombrement parallèles des partitions non croisées d'un cycle et des découpage d'un polygone convexe, *Discrete Math.* **2** (1972), 279–288.
- [23] V. Reiner, Non-crossing partitions for classical reflection groups, preprint dated March 1996. Available at the URL <ftp://s6.math.umn.edu/pub/papers/reiner/> or by anonymous ftp.
- [24] C. Reutenauer, *Free Lie Algebras*, Oxford University Press, Oxford, 1993.
- [25] R. Simion, Combinatorial statistics on non-crossing partitions, *J. Combinatorial Theory (A)* **66** (1994), 270–301.
- [26] R. Simion and R. Stanley, Flag-symmetry of the poset of shuffles and a local action of the symmetric group, in preparation.
- [27] R. Simion and D. Ullman, On the structure of the lattice of noncrossing partitions, *Discrete Math.* **98** (1991), 193–206.
- [28] R. Speicher, Multiplicative functions on the lattice of noncrossing partitions and free convolution, *Math. Ann.* **298** (1994), 611–628.
- [29] R. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth and Brooks/Cole, Pacific Grove, CA, 1986; second printing, Cambridge University Press, Cambridge, 1996 or 1997.
- [30] R. Stanley, Flag-symmetric and locally rank-symmetric partially ordered sets, *Electronic J. Combinatorics* **3**, R6 (1996), 22 pp.
- [31] R. Stanley, Hyperplane arrangements, parking functions and tree inversions, in *Festschrift in Honor of Gian-Carlo Rota*, Birkhäuser, Boston/Basel/Berlin, to appear.