

A note on the ranks of set-inclusion matrices

D. de Caen

Department of Mathematics and Statistics

Queen's University

Kingston, Ontario, Canada K7L 3N6

decaen@mast.queensu.ca

Submitted: June 11, 2001; Accepted: June 16, 2001

Abstract

A recurrence relation is derived for the rank (over most fields) of the set-inclusion matrices on a finite ground set.

Given a finite set X of say v elements, let $W = W_{t,k}(v)$ be the $(0,1)$ -matrix of inclusions for t -subsets versus k -subsets of X : $W_{T,K} = 1$ if T is contained in K , and 0 otherwise. These matrices play a significant part in several combinatorial investigations, see e.g. ([2], Thm. 2.4).

Let F be any field, and let $r_F(M)$ denote the rank of M over F .

Theorem. If $(k - t) \neq 0$ in the field F , then

$$r_F(W_{t,k}(v + 1)) = r_F(W_{t,k-1}(v)) + r_F((k - t + 1)W_{t-1,k}(v)). \quad (1)$$

Proof. The block-matrix identity

$$\begin{bmatrix} I & -A \\ 0 & I \end{bmatrix} \begin{bmatrix} AB & 0 \\ B & BC \end{bmatrix} \begin{bmatrix} I & -C \\ 0 & I \end{bmatrix} = \begin{bmatrix} 0 & -ABC \\ B & 0 \end{bmatrix}$$

implies that, over any field F ,

$$r_F\left(\begin{bmatrix} AB & 0 \\ B & BC \end{bmatrix}\right) = r_F(B) + r_F(ABC). \quad (2)$$

The set-inclusion matrix has the block-triangular decomposition

$$W_{t,k}(v + 1) = \begin{bmatrix} W_{t-1,k-1}(v) & 0 \\ W_{t,k-1}(v) & W_{t,k}(v) \end{bmatrix}, \quad (3)$$

as may be seen by fixing x in X and classifying t -sets and k -sets according to whether x belongs to them or not. Further, there is the elementary product formula

$$W_{t,k}(v)W_{k,l}(v) = \binom{l-t}{k-t} W_{t,l}(v) \quad (4)$$

whose proof is left as a straightforward exercise. Using (4), one may re-write (3) as

$$W_{tk}(v+1) = \begin{bmatrix} \frac{1}{(k-t)}W_{t-1,t}(v)W_{t,k-1}(v) & 0 \\ W_{t,k-1}(v) & W_{t,k-1}(v)W_{k-1,k}(v)\frac{1}{(k-t)} \end{bmatrix}$$

and so (2) is applicable:

$$\begin{aligned} r_F(W_{t,k}(v+1)) &= r_F(W_{t,k-1}(v)) + r_F(W_{t-1,t}(v)W_{t,k-1}(v)W_{k-1,k}(v)) \\ &= r_F(W_{t,k-1}(v)) + r_F((k-t+1)W_{t-1,k}(v)), \end{aligned}$$

which completes the proof of (1). □

Corollary Over the rational field \mathbb{Q} , $r_{\mathbb{Q}}(W_{t,k}(v)) = \binom{v}{t}$, provided $k+t \leq v$.

Proof. This is very easy using (1): note that the condition " $k+t \leq v$ " is inherited by the triples $(t, k-1, v-1)$ and $(t-1, k, v-1)$; so the result follows by induction. □

The corollary is a well known result, first proved by Gottlieb [3]. Wilson [4] has worked out the modular ranks of $W_{t,k}(v)$. Unfortunately, the condition $(k-t) \neq 0$ in the hypothesis of our theorem precludes a new proof of Wilson's theorem via our recursive formula. In the special case when the characteristic p of F is larger than k , our recursion does apply, with the same conclusion and proof as the above corollary.

In conclusion, we raise the question as to whether there is a q -analogue of formula (1), i.e., for the $(0,1)$ -inclusion matrix $W_{t,k}^{(q)}(v)$ of t -dimensional subspaces versus k -dimensional subspaces of a v -dimensional space over $GF(q)$; see [1], where the F -rank of $W_{t,k}^{(q)}(v)$ is computed when $\text{char}(F)$ does not divide q .

Acknowledgement Support has been provided by a grant from NSERC.

References

- [1] A. Frumkin and A. Yakir, "Rank of inclusion matrices and modular representation theory", Israel J. Math. 71 (1990), 309-320.
- [2] C. D. Godsil, "Tools from linear algebra", in Handbook of Combinatorics (eds., Graham, Grötschel, Lovász), MIT press 1995, pp. 1705-1748.
- [3] D. H. Gottlieb, "A class of incidence matrices", Proc. Amer. Math. Soc. 17 (1966), 1233-1237.
- [4] R. M. Wilson, "A diagonal form for the incidence matrix of t -subsets vs. k -subsets", European J. Combin. 11 (1990), 609-615.