

# A Concise Proof of the Littlewood-Richardson Rule

John R. Stembridge\*  
Department of Mathematics  
University of Michigan  
Ann Arbor, Michigan 48109–1109 USA  
jrs@umich.edu

Submitted January 2, 2002; Accepted March 15, 2002  
MR Subject Classification: 05E05

## Abstract

We give a short proof of the Littlewood-Richardson rule using a sign-reversing involution.

## Introduction.

The Littlewood-Richardson rule is one of the most important results in the theory of symmetric functions. It provides an explicit combinatorial rule for expressing either a skew Schur function, or a product of two Schur functions, as a linear combination of (non skew) Schur functions. Since Schur functions in  $n$  variables are the irreducible polynomial characters of  $GL_n(\mathbf{C})$ , the Littlewood-Richardson rule amounts to a tensor product rule for  $GL_n(\mathbf{C})$ .

The rule was first formulated in a 1934 paper by Littlewood and Richardson [LR], but the first complete proofs were not published until the 1970's. (For a historical account of the evolution of the rule and its proofs, see the recent survey paper of van Leeuwen [vL].) There are now many proofs available, such as those based on the Robinson-Schensted-Knuth correspondence, *jeu de taquin*, or the plactic monoid. In this note, we present a very simple, self-contained proof of the rule; the argument also proves at the same time the “bi-alternant” formula for Schur functions—the formula originally used by Cauchy to define Schur functions.

We obtained this proof by specializing a crystal graph argument that works in much greater generality (see Theorem 2.4 of [S]). The fact that crystal graphs (or the closely related Path Model of Littelmann) may be used to prove the Littlewood-Richardson rule, as well as tensor product rules for other semisimple Lie groups, is well-known (see [KN] or [L]), but we believe that it is not widely understood that there exist versions of these proofs that are self-contained, with no need to appeal to a general theory.

The proof we present here is not the first short proof. Alternatives include proofs by Berenstein and Zelevinsky [BZ], Remmel and Shimozono [RS], and Gasharov [G]. Furthermore, aside from the differences in language between semistandard tableaux and Gelfand patterns, the sign-reversing involution we use here is essentially a translation of the one used by Berenstein and Zelevinsky.

---

\* Work supported by NSF grant DMS-0070685.

## The Details.

Let  $\mathcal{P}$  denote the set of nonnegative integer sequences of the form  $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$  with finitely many nonzero terms; i.e., the set of partitions. We let  $\mathcal{P}_n$  denote the set of partitions with at most  $n$  nonzero terms, viewed (by truncation) as a subset of  $\mathbf{Z}^n$ .

Now regard  $n$  as fixed, and set  $\rho = (n - 1, \dots, 1, 0)$  and  $\emptyset = (0, \dots, 0) \in \mathcal{P}_n$ .

For each  $\lambda \in \mathbf{Z}^n$ , define  $x^\lambda = x_1^{\lambda_1} \cdots x_n^{\lambda_n}$  and  $a_\lambda = \det[x_i^{\lambda_j}] = \sum_{w \in S_n} \text{sgn}(w)x^{w\lambda}$ .

Given  $\mu, \nu \in \mathcal{P}$ , let  $D(\mu, \nu) = \{(i, j) \in \mathbf{Z}^2 : 1 \leq i \leq n, \nu_i < j \leq \mu_i\}$ . Assuming  $\nu \leq \mu$  (meaning  $\nu_i \leq \mu_i$  for all  $i$ ), define  $\mathcal{S}(\mu/\nu)$  to be the set of semistandard tableaux of shape  $\mu/\nu$ ; i.e., the set of mappings  $T : D(\mu, \nu) \rightarrow [n]$  with increasing columns ( $T(i, j) < T(i + 1, j)$ ) and weakly increasing rows ( $T(i, j) \leq T(i, j + 1)$ ). The weight of  $T$  is  $\omega(T) = (\omega_1(T), \dots, \omega_n(T)) \in \mathbf{Z}^n$ , where  $\omega_k(T) = |T^{-1}(k)|$  denotes the number of  $k$ 's in  $T$ . The generating series  $s_{\mu/\nu} = \sum_{T \in \mathcal{S}(\mu/\nu)} x^{\omega(T)}$  is a skew Schur function.

There is a well-known set of involutions  $\sigma_1, \dots, \sigma_{n-1}$  on  $\mathcal{S}(\mu/\nu)$ , due to Bender and Knuth [BK], with the property that  $\sigma_k$  acts by changing certain entries of  $T \in \mathcal{S}(\mu/\nu)$  from  $k$  to  $k + 1$  and vice-versa in such a way that  $\omega(\sigma_k(T)) = s_k \omega(T)$ , where  $s_k$  denotes the transposition  $(k, k + 1) \in S_n$ . The existence of these involutions proves that  $s_{\mu/\nu}$  is a symmetric function of  $x_1, \dots, x_n$ .

To explicitly describe the action of  $\sigma_k$  on  $T \in \mathcal{S}(\mu/\nu)$ , declare an entry  $k$  or  $k + 1$  to be *free* in  $T$  if there is no corresponding  $k + 1$  or  $k$  (respectively) in the same column. It is easy to check that the free entries in a given row must occur in consecutive columns; moreover, the entries in the free positions may be arbitrarily changed from  $k$  to  $k + 1$  and vice-versa without violating semistandardness as long as the free positions remain weakly increasing by row. The tableau  $\sigma_k(T)$  is obtained by reversing the numbers of free  $k$ 's and  $k + 1$ 's within each row; i.e., if there are  $a_i$  free  $k$ 's and  $b_i$  free  $k + 1$ 's in row  $i$  of  $T$ , then there should be  $b_i$  free  $k$ 's and  $a_i$  free  $k + 1$ 's in row  $i$  of  $\sigma_k(T)$ .

In the following,  $T_{\geq j}$  denotes the subtableau of  $T$  formed by the entries in columns  $j, j + 1, \dots$ , and we use similar notations such as  $T_{< j}$  and  $T_{> j}$  in the obvious way.

**Theorem.** For all  $\lambda \in \mathcal{P}_n$  and all  $\mu, \nu \in \mathcal{P}$  such that  $\nu \leq \mu$ , we have

$$a_{\lambda + \rho} s_{\mu/\nu} = \sum a_{\lambda + \omega(T) + \rho},$$

where the sum ranges over all  $T \in \mathcal{S}(\mu/\nu)$  such that  $\lambda + \omega(T_{\geq j}) \in \mathcal{P}_n$  for all  $j \geq 1$ .

*Proof.* As noted above, we know that  $s_{\mu/\nu}$  is symmetric, so for each  $w \in S_n$ , the quantities  $w(\lambda + \rho) + \omega(T)$  and  $w(\lambda + \rho + \omega(T))$  are identically distributed as  $T$  varies over  $\mathcal{S}(\mu/\nu)$ . Hence,

$$a_{\lambda + \rho} s_{\mu/\nu} = \sum_{w \in S_n} \sum_{T \in \mathcal{S}(\mu/\nu)} \text{sgn}(w)x^{w(\lambda + \rho + \omega(T))} = \sum_{T \in \mathcal{S}(\mu/\nu)} a_{\lambda + \omega(T) + \rho}. \quad (1)$$

We declare  $T$  to be a Bad Guy if  $\lambda + \omega(T_{\geq j})$  fails to be a partition for some  $j$ ; i.e.,

$$\lambda_k + \omega_k(T_{\geq j}) < \lambda_{k+1} + \omega_{k+1}(T_{\geq j})$$

for some pair  $k, j$ . Among all such pairs  $k, j$ , choose one that maximizes  $j$ , and among those, choose the smallest  $k$ . It must be the case that  $\lambda + \omega(T_{>j})$  is a partition, and since  $\omega_k(T_{\geq j}) - \omega_{k+1}(T_{\geq j})$  can change by at most one if we increment or decrement  $j$ , there must be a  $k + 1$  in column  $j$  of  $T$  (and no  $k$ ), and

$$\lambda_k + \omega_k(T_{\geq j}) + 1 = \lambda_{k+1} + \omega_{k+1}(T_{\geq j}). \quad (2)$$

Let  $T^*$  denote the tableau obtained from  $T$  by applying the Bender-Knuth involution  $\sigma_k$  to the subtableau  $T_{<j}$ , leaving the remainder of  $T$  unchanged. Since this involves changing some subset of the entries of  $T_{<j}$  from  $k$  to  $k + 1$  and vice-versa, and column  $j$  has a  $k + 1$  but no  $k$ , it is easy to see that  $T^*$  is semistandard. Furthermore,  $(T^*)_{\geq j}$  and  $T_{\geq j}$  are identical, so  $T \mapsto T^*$  is an involution on the set of Bad Guys. In comparing the contributions of  $T$  and  $T^*$  to (1), note that  $s_k \omega(T_{<j}) = \omega(T_{<j}^*)$ , whereas (2) implies that  $s_k$  fixes  $\lambda + \omega(T_{\geq j}) + \rho$ , whence  $s_k(\lambda + \omega(T) + \rho) = \lambda + \omega(T^*) + \rho$  and

$$a_{\lambda + \omega(T) + \rho} = -a_{\lambda + \omega(T^*) + \rho}.$$

The contributions of Bad Guys may therefore be canceled from (1).  $\square$

For the shape  $\mu = \mu/\emptyset$ , we have  $\omega(T_{\geq j}) \in \mathcal{P}_n$  for all  $j$  only if every entry in row  $i$  of  $T$  is  $i$ ; thus, there is a unique such  $T$ , it has weight  $\mu$ , and hence  $a_\rho s_\mu = a_{\mu+\rho}$ , or

**Corollary** (The Bi-Alternant Formula). *For all  $\mu \in \mathcal{P}_n$ , we have  $s_\mu = a_{\mu+\rho}/a_\rho$ .*

**Corollary.** *For all  $\lambda \in \mathcal{P}_n$  and all  $\mu, \nu \in \mathcal{P}$  such that  $\nu \leq \mu$ , we have*

$$s_\lambda s_{\mu/\nu} = \sum s_{\lambda + \omega(T)},$$

where the sum ranges over all  $T \in \mathcal{S}(\mu/\nu)$  such that  $\lambda + \omega(T_{\geq j}) \in \mathcal{P}_n$  for all  $j \geq 1$ .

This corollary is Zelevinsky's extension of the Littlewood-Richardson rule [Z].

Taking the specialization  $\lambda = \emptyset$ , we obtain the decomposition of  $s_{\mu/\nu}$  into Schur functions; it is simpler than the traditional formulation of the Littlewood-Richardson rule as found (e.g.) in [M], since it does not involve converting tableaux to words and imposing the "lattice permutation" condition. However, it still involves counting semistandard tableaux of shape  $\mu/\nu$  satisfying certain properties, and it is a not-too-difficult exercise to show that these two formulations count the *same* tableaux.

Via the specialization  $\nu = \emptyset$ , we obtain yet another formulation of the Littlewood-Richardson rule—in this case involving the decomposition of  $s_\lambda s_\mu$  into Schur functions.

## References.

- [BK] E. A. Bender and D. E. Knuth, Enumeration of plane partitions, *J. Combin. Theory Ser. A* **13** (1972), 40–54.
- [BZ] A. D. Berenstein and A. V. Zelevinsky, Involutions on Gelfand-Tsetlin schemes and multiplicities in skew  $GL_n$ -modules, *Soviet Math. Dokl.* **37** (1988), 799–802.
- [G] V. Gasharov, A short proof of the Littlewood-Richardson rule, *European J. Combin.* **19** (1998), 451–453.
- [KN] M. Kashiwara and T. Nakashima, Crystal graphs for representations of the  $q$ -analogue of classical Lie algebras, *J. Algebra* **165** (1994), 295–345.
- [L] P. Littelmann, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, *Invent. Math.* **116** (1994), 329–346.
- [M] I. G. Macdonald, “Symmetric Functions and Hall Polynomials,” Second Edition, Oxford Univ. Press, Oxford, 1995.
- [LR] D. E. Littlewood and A. R. Richardson, Group characters and algebra, *Phil. Trans. A* **233** (1934), 99–141.
- [RS] J. B. Remmel and M. Shimozono, A simple proof of the Littlewood-Richardson rule and applications, *Discrete Math.* **193** (1998), 257–266.
- [S] J. R. Stembridge, Combinatorial models for Weyl characters, *Advances in Math.*, to appear.
- [vL] M. A. A. van Leeuwen, The Littlewood-Richardson rule, and related combinatorics, in “Interactions of Combinatorics and Representation Theory,” *MSJ Memoirs* **11**, Math. Soc. Japan, Tokyo, 2001, pp. 95–145.
- [Z] A. V. Zelevinsky, A generalization of the Littlewood-Richardson rule and the Robinson-Schensted-Knuth correspondence, *J. Algebra* **69** (1981), 82–94.