

Small Ramsey Numbers

Stanisław P. Radziszowski
Department of Computer Science
Rochester Institute of Technology
Rochester, NY 14623, spr@cs.rit.edu

Submitted: June 11, 1994; Accepted: July 3, 1994
Revision #9: July 15, 2002

ABSTRACT: We present data which, to the best of our knowledge, includes all known nontrivial values and bounds for specific graph, hypergraph and multicolor Ramsey numbers, where the avoided graphs are complete or complete without one edge. Many results pertaining to other more studied cases are also presented. We give references to all cited bounds and values, as well as to previous similar compilations. We do not attempt complete coverage of asymptotic behavior of Ramsey numbers, but concentrate on their specific values.

Mathematical Reviews Subject Number 05C55.

Revisions

1993, February	preliminary version, RIT-TR-93-009 [Ra2]
1994, July 3	accepted to the <i>EJJC</i> , posted on the web
1994, November 7	<i>EJJC</i> revision #1
1995, August 28	<i>EJJC</i> revision #2
1996, March 25	<i>EJJC</i> revision #3
1997, July 11	<i>EJJC</i> revision #4
1998, July 9	<i>EJJC</i> revision #5
1999, July 5	<i>EJJC</i> revision #6
2000, July 25	<i>EJJC</i> revision #7
2001, July 12	<i>EJJC</i> revision #8
2002, July 15	<i>EJJC</i> revision #9

Contents

1. Scope and Notation	3
2. Classical Two Color Ramsey Numbers	4
2.1 Upper and lower bounds on $R(k, l)$ for $k \leq 10, l \leq 15$	4
2.2 Lower bounds on $R(k, l)$ for $l \geq 15$	6
3. Two Colors - Dropping One Edge from Complete Graph	7
4. General Graph Numbers in Two Colors	9
4.1 Paths	9
4.2 Cycles	9
4.3 Wheels	9
4.4 Books	10
4.5 Complete bipartite graphs	10
4.6 Triangle versus other graphs	10
4.7 Paths versus other graphs	11
4.8 Cycles versus complete graphs	11
4.9 Cycles versus other graphs	12
4.10 Stars versus other graphs	12
4.11 Books versus other graphs	13
4.12 Wheels versus other graphs	13
4.13 Trees and Forests	13
4.14 Mixed special cases	14
4.15 Mixed general cases	14
4.16 Other general results	14
5. Multicolor Graph Numbers	17
5.1 Bounds for multicolor classical numbers	17
5.2 Multicolor special cases	18
5.3 Multicolor results for complete graphs	19
5.4 Multicolor results for cycles	20
5.5 Other general multicolor results	20
6. Hypergraph Numbers	21
7. Cumulative Data and Surveys	22
7.1 Cumulative data for two colors	22
7.2 Cumulative data for three colors	23
7.3 Surveys	23
8. Concluding Remarks	24
9. References	25
A through D	25
E through G	28
H through L	32
M through R	36
S through Z	38

1. Scope and Notation

There is a vast literature on Ramsey type problems starting in 1930 with the original paper of Ramsey [Ram]. Graham, Rothschild and Spencer in their book [GRS] present an exciting development of Ramsey Theory. The subject has grown amazingly, in particular with regard to asymptotic bounds for various types of Ramsey numbers (see the survey paper [GrRö]), but the progress on evaluating the basic numbers themselves has been very unsatisfactory for a long time. In the last decade, however, considerable progress has been obtained in this area, mostly by employing computer algorithms. The few known exact values and several bounds for different numbers are scattered among many technical papers. This compilation is a fast source of references for the best results known for specific numbers. It is not supposed to serve as a source of definitions or theorems, but these can be easily accessed via the references gathered here.

Ramsey Theory studies conditions when a combinatorial object contains necessarily some smaller given objects. The role of Ramsey numbers is to quantify some of the general existential theorems in Ramsey Theory.

Let G_1, G_2, \dots, G_m be graphs or s -uniform hypergraphs (s is the number of vertices in each edge). $R(G_1, G_2, \dots, G_m; s)$ denotes the m -color **Ramsey number** for s -uniform graphs/hypergraphs, avoiding G_i in color i for $1 \leq i \leq m$. It is defined as the least integer n such that, in any coloring with m colors of the s -subsets of a set of n elements, for some i the s -subsets of color i contain a sub-(hyper)graph isomorphic to G_i (not necessarily induced). The value of $R(G_1, G_2, \dots, G_m; s)$ is fixed under permutations of the first m arguments.

If $s = 2$ (standard graphs) then s can be omitted. If G_i is a complete graph K_k , then we can write k instead of G_i , and if $G_i = G$ for all i we can use the abbreviation $R_m(G; s)$ or $R_m(G)$. For $s = 2$, $K_k - e$ denotes a K_k without one edge, and for $s = 3$, $K_k - t$ denotes a K_k without one triangle (hyperedge). P_i is a **path** on i vertices, C_i is a **cycle** of length i , and W_i is a **wheel** with $i - 1$ spokes, i.e. a graph formed by some vertex x , connected to all vertices of some cycle C_{i-1} . $K_{n,m}$ is a complete n by m bipartite graph, in particular $K_{1,n}$ is a **star** graph. The **book** graph $B_i = K_2 + K_i = K_1 + K_{1,i}$ has $i + 2$ vertices, and can be seen as i triangular pages attached to a single edge. The **fan** graph F_n is defined by $F_n = K_1 + nK_2$. For a graph G , $n(G)$ and $e(G)$ denote the number of vertices and edges, respectively. Finally, let $\chi(G)$ be the chromatic number of G , and let nG denote n disjoint copies of G .

Section 2 contains the data for the classical two color Ramsey numbers $R(k, l)$ for complete graphs, and section 3 for the two color case when the avoided graphs are complete or have the form $K_k - e$, but not both are complete. Section 4 lists the most studied two color cases for other graphs. The multicolor and hypergraph cases are gathered in sections 5 and 6, respectively. Finally, section 7 gives pointers to cumulative data and to some previous surveys, especially those containing data not subsumed by this compilation.

2. Classical Two Color Ramsey Numbers

l	3	4	5	6	7	8	9	10	11	12	13	14	15
k													
3	6	9	14	18	23	28	36	40 43	46 51	52 59	59 69	66 78	73 88
4		18	25	35 41	49 61	56 84	69 115	80 149	96 191	128 238	133 291	141 349	153 417
5			43 49	58 87	80 143	95 216	121 316	141 442	153	181	193	221	242
6				102 165	111 298	127 495	153 780	177 1171	253	262	278	292	374
7					205 540	216 1031	1713	2826	322	416	511		
8						282 1870	3583	6090			635		703
9							565 6588	580 12677					
10								798 23556					

Table I. Known nontrivial values and bounds for two color Ramsey numbers $R(k, l) = R(k, l; 2)$.

l	4	5	6	7	8	9	10	11	12	13	14	15
k												
3	GG	GG	Kéry	Ka2 GY	GR MZ	Ka2 GR	Ex5 RK2	Ka2 RK2	Ex12 Les	Piw1 RK2	Ex8 RK2	WW Les
4	GG	Ka1 MR4	Ex9 MR5	Ex3 Mac	Ex15 Mac	RK1 Mac	Piw1 Mac	Piw1 Spe2	SLL2 Spe2	XX1 Spe2	XX1 Spe2	XX1 Spe2
5		Ex4 MR5	Ex9 HZ1	CET Spe2	Piw1 Spe2	Haa Mac	Ex12 Mac	Ex12	Ex12	Ex12	Ex12	SLLL
6			Ka1 Mac	XX1 Mac	XX1 Mac	Ex12 Mac	XX1 Mac	XX1	XX1	XX1	XX1	SLLL
7				She1 Mac	XX1 Mac	HZ1	Mac	XX1	XX1	XX1		
8					BR Mac	Ea1	XX1 HZ1			XX1		XX1
9						She1 ShZ1	XX1 Ea1					
10							She1 Shi2					

References for Table I.

We split the data into the table of values and a table with corresponding references for the Table I. Known exact values appear as centered entries, lower bounds as top entries, and upper bounds as bottom entries.

The task of proving $R(3,3) \leq 6$ was the second problem in Part I of the William Lowell Putnam Mathematical Competition held in March 1953 [Bush].

The construction by Mathon [Mat] (see also sections 4.16 and 5.3.(h)), using data obtained by Shearer [She1], gives the following lower bounds for higher diagonal numbers: $R(11,11) \geq 1597$, $R(13,13) \geq 2557$, $R(14,14) \geq 2989$, $R(15,15) \geq 5485$, and $R(16,16) \geq 5605$. Similarly, $R(17,17) \geq 8917$, $R(18,18) \geq 11005$ and $R(19,19) \geq 17885$ were obtained in [LSL]. The same approach does not improve on an easy bound $R(12,12) \geq 1597 + 11 + 10$. The best known construction for this case showing $R(12,12) \geq 1637$ is given in [XX1] (for the general case see section 4.16).

All the critical graphs for the numbers $R(k,l)$ (graphs on $R(k,l) - 1$ vertices without K_k and without K_l in the complement) are known for $k = 3$ and $l = 3, 4, 5$ [Kéry], 6 [Ka2], 7 [RK3, MZ], and there are 1, 3, 1, 7 and 191 of them, respectively. All $(3,k)$ -graphs, for $k \leq 6$, were enumerated in [RK3], and all $(4,4)$ -graphs in [MR2]. There exists a unique critical graph for $R(4,4)$ [Ka2]. There are 4 such graphs known for $R(3,8)$ [RK2], 1 for $R(3,9)$ [Ka2] and 350904 for $R(4,5)$ [MR4], but there might be more of them. In [MR5] evidence is given for the conjecture that $R(5,5) = 43$ and that there exist 656 critical graphs on 42 vertices.

The claim that $R(5,5) = 50$ published on the web [Stone] is in error, and despite being shown so more than once, this incorrect value is being cited by some authors. The bound $R(3,13) \geq 60$ [XZ] cited in the 1995 version of this survey was shown to be incorrect in [Piw1]. The graphs constructed by Exoo in [Ex12, Ex15], and some others, are available electronically from <http://isu.indstate.edu/ge/RAMSEY>.

By taking a disjoint union of two critical graphs one can easily see that $R(k,p) \geq s$ and $R(k,q) \geq t$ imply $R(k,p+q-1) \geq s+t-1$. Xu and Xie [XX1] improved this construction to yield better general lower bounds, in particular $R(k,p+q-1) \geq s+t+k-3$. For example, this gives a lower bound $R(4,13) \geq 133$ with $p=2, q=12$. Only some higher lower bounds implied this way are shown. Some upper bounds implied by $R(k,l) \leq R(k-1,l) + R(k,l-1)$, or by its slight improvement with strict inequality when both terms on the right hand side are even, are marked [Ea1]. There are obvious generalizations of these inequalities for graphs other than complete.

The bound $R(6,6) \leq 166$, only 1 more than the best known [Mac], is an easy consequence of theorem 1 in [Walk] (see section 4.16) and the inequality $R(4,6) \leq 41$. T. Spencer [Spe2], Mackey [Mac], and Huang and Zhang [HZ1], using the bounds for minimum and maximum number of edges in $(4,5)$ Ramsey graphs listed in [MR3, MR5], were able to establish new upper bounds for several higher Ramsey numbers, improving all the previous long-standing results of Giraud [Gi3, Gi5, Gi6]. We have recomputed the bounds marked [HZ1] using the method from the paper [HZ1], because the bounds there relied on an overly optimistic personal communication from T. Spencer. Further refinements of this method are studied in [HZ2, ShZ1, Shi2].

For a more in depth study of triangle-free graphs in relation to the case of $R(3,k)$, for which considerable progress has been obtained in recent years, see also [AKS, Alon2, BBH1, BBH2, CPR, FL, Fra1, Fra2, Gri, Loc, KM1, RK3, RK4, She2, Stat, Yu1]. In 1995, Kim [Kim] obtained a breakthrough by proving that $R(3,k)$ has order of magnitude exactly

$\Theta(k^2/\log k)$. Good asymptotic bounds for $R(k, k)$ can be found, for example, in [Chu3, McS] (lower bound) and [Tho] (upper bound), and for many other asymptotic bounds in the general case of $R(k, l)$ consult [Spe1, GRS, GrRö, AP].

The lower bounds marked [XX1] were obtained by a method referenced in section 4.16. All other lower bounds for higher numbers listed in Table II were obtained by construction of cyclic graphs, except for the bound $R(5, 17) \geq 284$ established by Exoo [Ex15] using pq -groups.

$k \backslash l$	15	16	17	18	19	20	21	22	23
3	73 WW	79 WW	92 WWY1	98 WWY1	106 WWY1	109 WWY1	122 WWY1	125 WWY1	136 WWY1
4	153 XX1		182 LSS		198 LSZL	230 SLZL	242 SLZL	282 SL	
5	242 SLLL	278 LSS	284 Ex15		338 SLZL	380 LSS		422 LSZL	434 LSZL
6	374 SLLL	434 SLLL	548 SLLL	614 SLLL	710 SLLL	878 SLLL		1070 SLLL	
7			628 XX1	722 XX1	908 SLLL		1214 SLLL		
8	703 XX1		737 XX1	871 XX1	1054 XX1	1094 SLLL	1328 SLLL		

Table II. Known nontrivial lower bounds for higher two color Ramsey numbers $R(k, l)$, with references.

Exoo [Ex15] gives also the bounds $R(3, 27) \geq 158$ and $R(3, 31) \geq 198$. The constructions establishing $R(3, 26) \geq 150$, $R(3, 29) \geq 174$, $R(3, 31) \geq 198$ and $R(3, 32) \geq 212$ are presented in [SLL1], [SLL3], [LSS] and [LSZL], respectively. Yu [Yu2] constructed a special class of triangle-free cyclic graphs establishing several lower bounds for $R(3, k)$, for $k \geq 61$. Only two of these bounds, $R(3, 61) \geq 479$ and $R(3, 103) \geq 955$, cannot be easily improved by the inequality $R(3, 4k + 1) \geq 6R(3, k + 1) - 5$ from [CCD] and data from tables I and II. Finally, $R(5, 24) \geq 488$ was given in [SLL3], and $R(9, 17) \geq 1411$ and $R(10, 16) \geq 1189$ in [XX1].

3. Two Colors - Dropping One Edge from Complete Graph

$G \quad H$	K_3-e	K_4-e	K_5-e	K_6-e	K_7-e	K_8-e	K_9-e	$K_{10}-e$	$K_{11}-e$
K_3-e	3	5	7	9	11	13	15	17	19
K_3	5	7	11	17	21	25	31	37 38	42 47
K_4-e	5	10	13	17	28	29 38	34	41	
K_4	7	11	19	27 36	37 52				
K_5-e	7	13	22	31 39	40 66				
K_5	9	16	30 34	43 67	112				
K_6-e	9	17	31 39	45 70	59 135				
K_6	11	21	37 55	119	205				
K_7-e	11	28	40 66	59 135	251				
K_7	13	28 34	51 88	204					

Table III. Two types of Ramsey numbers $R(G, H)$, includes all known nontrivial values.

The exact values in Table III above involving K_3-e are trivial, since easily $R(K_3-e, K_k) = R(K_3-e, K_{k+1}-e) = 2k - 1$, for all $k \geq 2$. Other bounds (not shown in Table III) can be obtained by using Table I, an obvious generalization of the inequality $R(k, l) \leq R(k-1, l) + R(k, l-1)$, and by monotonicity of Ramsey numbers, in this case $R(K_{k-1}, G) \leq R(K_k-e, G) \leq R(K_k, G)$.

For the following numbers it was established that the critical graphs are unique: $R(K_3, K_l-e)$ for $l = 3$ [Tr], 6 and 7 [Ra1], $R(K_4-e, K_4-e)$ [FRS2], $R(K_5-e, K_5-e)$ [Ra3] and $R(K_4-e, K_7-e)$ [McR]. The number of $R(K_3, K_l-e)$ critical graphs for $l = 4, 5$ and 8 is 4, 2 and 9, respectively [MPR]. Also in [MPR] a bound $R(K_3, K_{12}-e) \geq 46$ is given. Wang, Wang and Yan in [WWY2] constructed cyclic graphs showing $R(K_3, K_{13}-e) \geq 54$, $R(K_3, K_{14}-e) \geq 59$ and $R(K_3, K_{15}-e) \geq 69$. The upper bounds in [HZ2] were obtained by a reasoning generalizing the bounds for classical numbers in [HZ1].

G	H	K_4-e	K_5-e	K_6-e	K_7-e	K_8-e	K_9-e	$K_{10}-e$	$K_{11}-e$
K_3		CH2	Clan	FRS1	GH	Ra1	Ra1	MPR MPR	WWY2 MPR
K_4-e		CH1	FRS2	McR	McR	Ea1 HZ2	Ex14	Ex14	
K_4		CH2	EHM1	Ex11 Ea1	Ex14 HZ2				
K_5-e		FRS2	CEHMS	Ex14 Ea1	Ex14 HZ2				
K_5		BH	Ex8 Ex8	Ea1 HZ2	HZ2				
K_6-e		McR	Ex14 Ea1	Ex14 HZ2	Ex14 HZ2				
K_6		McN	Ex14 Ea1	ShZ2	ShZ2				
K_7-e		McR	Ex14 HZ2	Ex14 HZ2	ShZ1				
K_7		Ea1 Ea1	Ex14 ShZ2	ShZ2					

References for Table III.

4. General Graph Numbers in Two Colors

This section includes data with respect to general graph results. We tried to include all nontrivial values and identities regarding exact results (or references to them), but only those out of general bounds and other results which, in our opinion, have a direct connection to the evaluation of specific numbers. If some small value cannot be found below, it may be covered by the cumulative data gathered in section 7, or be a special case of a general result listed in this section. Note that $B_1 = F_1 = C_3 = W_3 = K_3$, $B_2 = K_4 - e$, $P_3 = K_3 - e$, $W_4 = K_4$ and $C_4 = K_{2,2}$ imply other identities not mentioned explicitly.

4.1. Paths

$$R(P_n, P_m) = n + \lfloor m/2 \rfloor - 1 \quad \text{for all } n \geq m \geq 2 \quad [\text{GeGy}]$$

4.2. Cycles

$$R(C_3, C_3) = 6 \quad [\text{GG}]$$

$$R(C_4, C_4) = 6 \quad [\text{CH1}]$$

Result obtained independently in [Ros] and [FS1], new simple proof in [KáRos]:

$$R(C_n, C_m) = \begin{cases} 2n - 1 & \text{for } 3 \leq m \leq n, m \text{ odd}, (n, m) \neq (3, 3) \\ n - 1 + m/2 & \text{for } 4 \leq m \leq n, m \text{ and } n \text{ even}, (n, m) \neq (4, 4) \\ \max\{n - 1 + m/2, 2m - 1\} & \text{for } 4 \leq m < n, m \text{ even and } n \text{ odd} \end{cases}$$

Unions of cycles, $R(nC_p, mC_q)$, [MS, Den]

$$R(nC_3, mC_3) = 3n + 2m \quad \text{for } n \geq m \geq 1, n \geq 2 \quad [\text{BES}]$$

$$R(nC_4, mC_4) = 2n + 4m - 1 \quad \text{for } m \geq n \geq 1, (n, m) \neq (1, 1) \quad [\text{LiWa}]$$

4.3. Wheels

$$R(W_3, W_5) = 11 \quad [\text{Clan}]$$

$$R(W_3, W_n) = 2n - 1 \quad \text{for all } n \geq 6 \quad [\text{BE2}]$$

All critical colorings for $R(W_3, W_n)$ for all $n \geq 3$ [RaJi]

$$R(W_4, W_5) = 17 \quad [\text{He3}]$$

$$R(W_5, W_5) = 15 \quad [\text{HM2, He2}]$$

$$R(W_4, W_6) = 19, R(W_5, W_6) = 17 \quad \text{and } R(W_6, W_6) = 17,$$

and all critical colorings (2, 1 and 2) for these numbers [FM]

4.4. Books

$$R(B_1, B_n) = 2n + 3 \text{ for all } n > 1 \text{ [RS1]}$$

$$R(B_3, B_3) = 14 \text{ [RS1, HM2]}$$

$$R(B_2, B_5) = 16, R(B_3, B_5) = 17, R(B_5, B_5) = 21,$$

$$R(B_4, B_4) = 18, R(B_4, B_6) = 22, R(B_6, B_6) = 26 \text{ [RS1]}$$

$$254 \leq R(B_{37}, B_{88}) \leq 255 \text{ [Par6]}$$

in general $R(B_n, B_n) = 4n + 2$ for $4n + 1$ a prime power,

and several other general equalities and bounds for $R(B_n, B_m)$ [RS1, FRS7, Par6].

4.5. Complete bipartite graphs

$$R(K_{2,3}, K_{2,3}) = 10 \text{ [Bu4]}$$

$$R(K_{2,3}, K_{2,4}) = 12 \text{ [ExRe]}$$

$$R(K_{2,3}, K_{1,7}) = 13 \text{ [Par4]}$$

$$R(K_{2,3}, K_{3,3}) = 13 \text{ and } R(K_{3,3}, K_{3,3}) = 18 \text{ [HM3]}$$

$$R(K_{2,2}, K_{2,8}) = 15 \text{ and } R(K_{2,2}, K_{2,11}) = 18 \text{ [HM]}$$

$$R(K_{2,2}, K_{1,15}) = 20 \text{ [La2]}$$

$$R(nK_{1,3}, mK_{1,3}) = 4n + m - 1 \text{ for } n \geq m \geq 1, n \geq 2 \text{ [BES]}$$

Asymptotics for $K_{2,m}$ versus K_n [CLRZ]

$R(K_{1,n}, K_{1,m}) = n + m - \varepsilon$, where $\varepsilon = 1$ if both n and m are even and $\varepsilon = 0$ otherwise [Har1]. It is also a special case of multicolor numbers for stars obtained in [BuRo1].

$R(K_{2,n}, K_{2,n}) \leq 4n - 2$ for all $n \geq 2$, exact values 6, 10, 14, 18, 21, 26, 30, 33, 38, 42, 46, 50, 54, 57 and 62 of $R(K_{2,n}, K_{2,n})$ for $2 \leq n \leq 16$, respectively.

The first open case is $65 \leq R(K_{2,17}, K_{2,17}) \leq 66$ [EHM2].

4.6. Triangle versus other graphs

$$R(3, k) = \Theta(k^2 / \log k) \text{ [Kim]}$$

Explicit construction for $R(3, 4k + 1) \geq 6R(3, k + 1) - 5$, for all $k \geq 1$ [CCD]

Explicit triangle-free graphs with independence k on $\Omega(k^{3/2})$ vertices [Alon2, CPR]

$$R(K_3, K_7 - 2P_2) = R(K_3, K_7 - 3P_2) = 18 \text{ [SchSch2]}$$

$$R(K_3, K_3 + \bar{K}_m) = R(K_3, K_3 + \bar{C}_m) = 2m + 5 \text{ for } m \geq 212 \text{ [Zhou1]}$$

$R(K_3, G) = 2n(G) - 1$ for any connected G on at least 4 vertices and with at most $(17n(G) + 1)/15$ edges, in particular for $G = P_i$ and $G = C_i$, for all $i \geq 4$ [BEFRS1]

$$R(K_3, G) \leq 2e(G) + 1 \text{ for any graph } G \text{ without isolated vertices [Sid3, GK]}$$

$$R(K_3, G) \leq n(G) + e(G) \text{ for all } G, \text{ a conjecture [Sid2]}$$

$$R(K_3, K_n), \text{ see section 2}$$

$$R(K_3, K_n - e), \text{ see section 3}$$

$$R(K_3, G) \text{ for all connected } G \text{ up to 9 vertices, see section 7.1}$$

Since $B_1 = F_1 = C_3 = W_3 = K_3$, other sections apply. See also [AKS, BBH1, BBH2, FL, Fra1, Fra2, Gri, Loc, KM1, RK3, RK4, She2, Spe1, Stat, Yu1].

4.7. Paths versus other graphs

Paths versus stars [Par2, BEFRS2]
 Paths versus trees [FS4]
 Paths versus books [RS2]
 Paths versus cycles [FLPS, BEFRS2]
 Paths versus K_n [Par1]
 Paths versus $K_{n,m}$ [Häg]
 Paths versus W_5 and W_6 [SuBa1]
 Paths versus W_7 and W_8 [Bas]
 Paths versus wheels [BaSu, ChZZ]
 Paths and cycles versus trees [FSS1]
 Sparse graphs versus paths and cycles [BEFRS2]
 Graphs with long tails [Bu2, BG]

4.8. Cycles versus complete graphs

$R(C_4, K_3) = R(C_4, C_3) = 7$ [CS]
 $R(C_4, K_4) = 10$ [CH2]
 $R(C_4, K_5) = 14$ [Clan]
 $R(C_4, K_6) = 18$ [Ex2] [RoJa1]
 $R(C_4, K_7) = 22$ [RT] [JR1]
 $R(C_4, K_8) = 26$ [RT]
 $30 \leq R(C_4, K_9)$, $34 \leq R(C_4, K_{10})$ [RT]
 C_4 versus K_n [CLRZ]
 $R(C_5, K_3) = R(C_5, C_3) = 9$ [CS]
 $R(C_5, K_4) = 13$ [He2, JR4]
 $R(C_5, K_5) = 17$ [He2, JR4]
 $R(C_5, K_6) = 21$ [JR5]
 $R(C_5, K_7) = 25$ [Schi2]
 $R(C_6, K_4) = 16$ [JR2]
 $R(C_6, K_5) = 21$ [JR2, YHZ2, BJYHRZ]
 $R(C_6, K_6) = 26$ [Schi1]
 $R(C_7, K_5) = 25$ [YHZ2, BJYHRZ]
 Cycles versus K_n [BoEr, Spe1, FS4, EFRS2, CLRZ, Sud]

$R(C_n, K_m) = (n-1)(m-1) + 1$, for $n \geq m^2 - 2$ [BoEr], for $n > 3 = m$ [FS1], for $n \geq 4 = m$ [YHZ1], for $n \geq 5 = m$ [BJYHRZ], for $n \geq 6 = m$, and for $n \geq m \geq 7$ with $n \geq m(m-2)$ [Schi1]. Since 1976, it was conjectured to be true for all $n \geq m \geq 3$, except $n = m = 3$ [FS4, EFRS2].

4.9. Cycles versus other graphs C_4 versus stars [Par3, Par5, BEFRS5, Chen] C_4 versus trees [EFRS4, Bu6, BEFRS5, Chen] C_4 versus $K_{m,n}$ [HM] C_4 versus all graphs on six vertices [JR3] $R(C_4, B_n) = 7, 9, 11, 12, 13$ and 16 , for $2 \leq n \leq 7$, respectively [FRS6] $R(C_4, B_n) = 17, 18, 19, 20$ and 21 , for $8 \leq n \leq 12$, respectively [Tse] $R(C_4, W_n) = 10, 9, 10, 9, 11, 12, 13, 14, 16$ and 17 , for $4 \leq n \leq 13$, respectively [Tse] $R(C_4, G) \leq 2q + 1$ for any isolate-free graph G with q edges [RoJa2] $R(C_4, G) \leq p + q - 1$ for any connected graph G on p vertices and q edges [RoJa2] $R(C_5, W_6) = 13$ [ChvS] $R(C_5, K_6 - e) = 17$ [JR4] $R(C_5, B_1) = R(C_5, B_2) = 9$ [CRSPS] $R(C_5, B_3) = 10$, and in general $R(C_5, B_n) = 2n + 3$ for $n \geq 4$ [FRS8] C_5 versus all graphs on six vertices [JR4] $R(C_6, K_5 - e) = 17$ [JR2] C_6 versus all graphs on five vertices [JR2] $R(C_n, G) \leq 2q + \lfloor n/2 \rfloor - 1$, for $3 \leq n \leq 5$, for any isolate-free graph G with $q > 3$ edges.It is conjectured that it also holds for other n [RoJa2].

Cycles versus paths [FLPS, BEFRS2]

Cycles versus stars [La1, Clark, see Par6]

Cycles versus trees [FSS1]

Cycles versus books [FRS6, FRS8, Zhou1]

Cycles versus W_5 and W_6 [SuBB2]

Cycles versus wheels [Zhou2]

See also bipartite graphs for $K_{2,2} = C_4$ **4.10. Stars versus other graphs**Stars versus C_4 [Par3, Par5, Chen]Stars versus W_5 and W_6 [SuBa1]

Stars versus paths [Par2, BEFRS2]

Stars versus cycles [La1, Clark, see Par6]

Stars versus books [CRSPS, RS2]

Stars versus trees [Bu1, GV, ZZ]

Stars versus stripes [CL, Lor]

Stars versus $K_{2,n}$ [Par4]Stars versus $K_{n,m}$ [Stev, Par3]Stars versus $K_n - tK_2$ [Hua1, Hua2]Stars versus $2K_2$ [MO]

Union of two stars [Gro2]

4.11. Books versus other graphs

$R(B_3, K_4) = 14$ [He3]
 $20 \leq R(B_3, K_5) \leq 22$ [He2]
 Books versus paths [RS2]
 Books versus trees [EFRS7]
 Books versus stars [CRSPS, RS2]
 Books versus cycles [FRS6, FRS8, Zhou1, Tse]
 Books versus wheels [Zhou3]
 Books versus $K_2 + C_n$ [Zhou3]
 Books and $(K_1 + tree)$ versus K_n [LR1]

4.12. Wheels versus other graphs

$R(W_5, K_5 - e) = 17$ [He2][YH]
 $27 \leq R(W_5, K_5) \leq 29$ [He2]
 W_5 and W_6 versus stars and paths [SuBa1]
 W_5 and W_6 versus trees [BSNM]
 W_5 and W_6 versus cycles [SuBB2]
 W_7 and W_8 versus paths [Bas]
 Wheels versus paths [BaSu, ChZZ]
 Odd wheels versus star-like trees [SuBB]
 $R(W_6, C_5) = 13$ [ChvS]
 Wheels versus C_4 [Tse]
 Wheels versus cycles [Zhou2]
 Wheels versus books [Zhou3]
 Wheels versus linear forests [SuBa2]

4.13. Trees and Forests

Trees, forests [EG, GRS, FSS1, GV, CsKo]
 Trees versus C_4 [EFRS4, Bu6, Chen]
 Trees versus paths [FS4]
 Trees versus paths and cycles [FSS1]
 Trees versus stars [Bu1, GV, ZZ]
 Trees versus books [EFRS7]
 Trees versus W_5 and W_6 [BSNM]
 Star-like trees versus odd wheels [SuBB]
 Trees versus K_n [Chv]
 Trees versus $K_n + K_m$ [RS2, FSR]
 Trees versus bipartite graphs [BEFRS5, EFRS6]
 Trees versus almost complete graphs [GJ2]
 Trees versus small ($n(G) \leq 5$) connected G [FRS4]
 Linear forests, forests [BuRo2, FS3, CsKo]
 Linear forests versus wheels [SuBa2]

Forests versus K_n [Stahl]

Forests versus almost complete graphs [CGP]

4.14. Mixed special cases:

$$R(C_5 + e, K_5) = 17 \text{ [He5]}$$

$$R(W_5, K_5 - e) = 17 \text{ [He2][YH]}$$

$$20 \leq R(B_3, K_5) \leq 22 \text{ [He2]}$$

$$27 \leq R(W_5, K_5) \leq 29 \text{ [He2]}$$

$$25 \leq R(K_5 - P_3, K_5) \leq 28 \text{ [He2]}$$

$$26 \leq R(K_{2,2,2}, K_{2,2,2}), K_{2,2,2} \text{ is an octahedron [Ex8]}$$

4.15. Mixed general cases

Unicyclic graphs [Gro1, Köh, KrRod]

$K_{2,m}$ and C_{2m} versus K_n [CLRZ]

$K_{2,n}$ versus any graph [RoJa2]

nK_3 versus mK_3 , in particular $R(nK_3, nK_3) = 5n$ for $n \geq 2$ [BES]

nK_3 versus mK_4 [LorMu]

$R(nK_4, nK_4) = 7n + 4$ for large n [Bu7]

Stripes [CL, Lor]

Union of two stars [Gro2]

Double stars* [GHK]

Graphs with bridge versus K_n [Li]

Fans $F_n = K_1 + nK_2$ versus K_m [LR2]

$R(F_1, F_n) = R(K_3, F_n) = 4n + 1$ and bounds for $R(F_m, F_n)$ [GGS]

Multipartite complete graphs [BEFRS3, EFRS4, FRS3, Stev]

Multipartite complete graphs versus trees [EFRS8, BEFRSGJ]

Disconnected graphs versus any graph [GJ1]

Graphs with long tails [Bu2, BG]

Brooms⁺ [EFRS3]

4.16. Other general results

[Chv] $R(K_n, T_m) = (n-1)(m-1) + 1$ for any tree T on m vertices.

[CH2] $R(G, H) \geq (\chi(G) - 1)(c(H) - 1) + 1$, where $\chi(G)$ is the chromatic number of G , and $c(H)$ is the size of the largest connected component of H .

[Walk] $R(k, k) \leq 4R(k, k-2) + 2$.

[Spe1] Probabilistic asymptotic lower bounds for $R(k, l)$, also weaker bounds but with an explicit constructive approach in [AP].

* - A double star is a union of two stars with their centers joined by an edge.

+ - A broom is a star with a path attached to its center.

- [Mat] If the quadratic residues Paley graph Q_p of prime order $p = 4t + 1$ contains no K_k , then $R(k, k) \geq p + 1$ and $R(k + 1, k + 1) \geq 2p + 3$. More data was obtained in [She1, LSL].
- [XX1] For $2 \leq p \leq q$ and $3 \leq k$, if (k, p) -graph G and (k, q) -graph H have a common induced subgraph on m vertices without K_{k-1} , then $R(k, p + q - 1) > n(G) + n(H) + m$. In particular, we have $R(k, p + q - 1) \geq R(k, p) + R(k, q) + k - 3$ and $R(k, p + q - 1) \geq R(k, p) + R(k, q) + p - 2$.
- [BE1] $R(G, G) \geq \lfloor (4n(G) - 1) / 3 \rfloor$ for any connected G .
- [BE2] Graphs yielding $R(K_n, G) = (n - 1)(n(G) - 1) + 1$ and related results (see also [EFRS5]).
- [BES] Study of Ramsey numbers for multiple copies of graphs. See also [Bu1, LorMu].
- [Zeng] $R(nK_3, nG)$ for all isolate-free graphs G on 4 vertices.
- [Bu7] Study of Ramsey numbers for large disjoint unions of graphs, in particular $R(nK_k, nK_l) = n(k + l - 1) + R(K_{k-1}, K_{l-1}) - 2$, for n large enough. See also [Bu8].
- [Bu2] Graphs H yielding $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$, where $s(G)$ is a chromatic surplus of G , defined as the minimum number of vertices in some color class under all vertex colorings in $\chi(G)$ colors (such H 's are called G -good). This idea, initiated in [Bu2], is a basis of a number of exact results for $R(G, H)$ for large and sparse graphs H [BG, BEFRS2, BEFRS4, Bu5, FS, EFRS4, FRS3, BEFSRGJ, BF, LR4]. A survey of this area appeared in [FRS5].
- [BEFS] Bounds for the difference between consecutive Ramsey numbers.
- [Par3] Relations between some Ramsey graphs and block designs. See also [Par4].
- [Bra3] $R(G, H) > h(G, d)n(H)$ for all nonbipartite G and almost every d -regular H , for some h unbounded in d .
- [CSRT] $R(G, G) \leq c_d n(G)$ for all G , where constant c_d depends only on the maximum degree d in G . The constant was improved in [GRR1]. Tight lower and upper bounds for bipartite G [GRR2].
- [ChenS] $R(G, G) \leq c_d n$ for all d -arrangeable graphs G on n vertices, in particular with the same constant for all planar graphs. The constant c_d was improved in [Eaton]. An extension to graphs not containing a subdivision of K_d [RöTh]. Progress towards a conjecture that the same inequality holds for all d -generate graphs G [KoRö].
- [EFRS9] Study of graphs G , called *Ramsey size linear*, for which there exists a constant c_G such that for all H with no isolates $R(G, H) \leq c_G e(H)$. An overview and further results were given in [BSS].
- [LRS] $R(G, G) < 6n$ for all n -vertex graphs G , in which no two vertices of degree at least 3 are adjacent. This improves the result $R(G, G) \leq 12n$ in [Alon1].

- [Shi1] $R(Q_n, Q_n) \leq 2^{(3+\sqrt{5})n/2+o(n)}$, for the n -dimensional cube Q_n with 2^n vertices. This bound can be also derived from a theorem in [KoRö].
- [Gro1] Conjecture that $R(G, G) = 2n(G) - 1$ if G is unicyclic of odd girth. Further support for the conjecture was given in [Köh, KrRod].
- [RoJa2] $R(K_{2,k}, G) \leq kq + 1$, for $k \geq 2$, for isolate-free graphs G with $q \geq 2$ edges.
- [FSS1] Discussion of the conjecture that $R(T_1, T_2) \leq n(T_1) + n(T_2) - 2$ holds for all trees T_1, T_2 .
- [FM] $R(W_6, W_6) = 17$ and $\chi(W_6) = 4$. This gives a counterexample $G = W_6$ to the Erdős conjecture (see [GRS]) $R(G, G) \geq R(K_{\chi(G)}, K_{\chi(G)})$.
- [LR3] Bounds on $R(H + \bar{K}_n, K_n)$ for general H . Also, for fixed k and m , as $n \rightarrow \infty$, $R(K_k + \bar{K}_m, K_n) \leq (m + o(1))n^k / (\log n)^{k-1}$ [LRZ].
- [-] Special cases of multicolor results listed in section 5.5.
- [-] See also surveys listed in section 7.

5. Multicolor Graph Numbers

The only known value of a multicolor classical Ramsey number:

$$R_3(3) = R(3,3,3) = R(3,3,3; 2) = 17 \quad \text{[GG]}$$

2 critical colorings [KS, LayMa]

5.1. Bounds for multicolor classical numbers

$$51 \leq R_4(3) = R(3,3,3,3) \leq 62 \quad \text{[Chu1] [FKR]}$$

$$162 \leq R_5(3) \leq 307 \quad \text{[Ex10]; [FKR], 5.3.(a)}$$

$$538 \leq R_6(3) \leq 1838 \quad \text{[FreSw], 5.3.(a)}$$

$$1682 \leq R_7(3) \leq 12861 \quad \text{[FreSw], 5.3.(a)}$$

$$128 \leq R(4,4,4) \leq 236 \quad \text{[HI], 5.3.(a)}$$

$$578 \leq R(4,4,4,4) \quad \text{[ExRa]}$$

$$2295 \leq R(4,4,4,4,4) \quad \text{[XX2]}$$

$$415 \leq R(5,5,5) \quad \text{[ExRa]}$$

$$2501 \leq R(5,5,5,5) \quad \text{5.3.(j)}$$

$$25922 \leq R(5,5,5,5,5) \quad \text{5.3.(j)}$$

$$1070 \leq R(6,6,6) \quad \text{[Mat], 5.3.(h)}$$

$$10407 \leq R(6,6,6,6) \quad \text{[XX2]}$$

$$3211 \leq R(7,7,7) \quad \text{[Mat], 5.3.(h)}$$

$$42117 \leq R(7,7,7,7) \quad \text{[XX2]}$$

$$4930 \leq R(8,8,8) \quad \text{[XX2]}$$

$$8461 \leq R(9,9,9) \quad \text{[XX2]}$$

$$30 \leq R(3,3,4) \leq 31 \quad \text{[Ka2] [PR1, PR2]}$$

$$45 \leq R(3,3,5) \leq 57 \quad \text{[Ex2, KLR], 5.3.(a)}$$

$$60 \leq R(3,3,6) \quad \text{[Rob3, Rob4]}$$

$$79 \leq R(3,3,7) \quad \text{[Ex16]}$$

$$98 \leq R(3,3,8) \quad \text{[ZSL]}$$

$$110 \leq R(3,3,9) \quad \text{[SLZL]}$$

$$141 \leq R(3,3,10) \quad \text{[XX2], 5.3.(c)}$$

$$157 \leq R(3,3,11) \quad \text{[XX2], 5.3.(c)}$$

$$181 \leq R(3,3,12) \quad \text{[XX2], 5.3.(c)}$$

$$205 \leq R(3,3,13) \quad \text{[XX2], 5.3.(c)}$$

$$233 \leq R(3,3,14) \quad \text{5.3.(c)}$$

$55 \leq R(3,4,4) \leq 79$	[KLR], 5.3.(a)
$80 \leq R(3,4,5) \leq 160$	[Ex12], 5.3.(a)
$99 \leq R(3,4,6)$	5.3.(g)
$123 \leq R(3,5,5)$	5.3.(g)
$93 \leq R(3,3,3,4) \leq 153$	[Ex16], 5.3.(a)
$137 \leq R(3,3,3,5)$	[Rob3]
$171 \leq R(3,3,4,4)$	[Ex16]
$561 \leq R(3,3,3,11)$	[XX2]

The best published upper bound $R_4(3) \leq 64$ by Sánchez-Flores [San] improved a very old bound $R_4(3) \leq 65$ obtained by Folkman [Fo] in 1974. In [PR1] it is conjectured that $R(3,3,4) = 30$, and the results in [PR2] eliminate some cases which could give $R(3,3,4) = 31$.

Lower bounds for higher numbers can be obtained by applying various constructive inequalities from the section 5.3 below. For example, the bounds $261 \leq R(3,3,15)$, $241 \leq R(3,3,3,7)$ and $2501 \leq R(5,5,5,5)$ were not published explicitly but are implied by general constructions in 5.3.(c), [Rob3] and [Abb1]. Similarly, other lower bounds for parameters not listed here can be easily derived.

5.2. Multicolor special cases

$R_3(C_4) = 11$	[BS, see also Clap]
$R_3(C_5) = 17$	[YR1]
$R_3(C_6) = 12$	[YR2]
$R_3(C_7) = 25$	[FSS2]
$18 \leq R_4(C_4) \leq 21$	[Ex2] [Ir]
$27 \leq R_5(C_4) \leq 29$	[LaWo1]
$R(C_4, C_4, K_3) = 12$	[Schu]
$R(C_4, K_3, K_3) = 17$	[ExRe]
$13 \leq R(C_3, C_4, C_5)$	[Rao]
$R(K_{1,3}, C_4, K_4) = 16$	[KM2]
$R(K_4 - e, K_4 - e, P_3) = 11$	[Ex7]
$28 \leq R_3(K_4 - e) \leq 30$	[Ex7] [Piw2]
$R(C_4, C_4, C_4, T) = 16$ for $T = P_4$ and $T = K_{1,3}$	[ExRe]
$25 \leq R(C_3, C_3, C_4, C_4)$	[Rao]

All colorings on at least 14 vertices for the parameters (K_3, K_3, K_3) , and all colorings for $(K_4 - e, K_4 - e, P_3)$ were found in [Piw2].

5.3. Multicolor results for complete graphs

(a) $R(k_1, \dots, k_r) \leq 2 - r + \sum_{i=1}^r R(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_r)$, implicit in [GG]

Inequality in (a) is strict if the right hand side is even, and at least one of the terms in the summation is even. It is suspected that this upper bound is never tight for $r \geq 3$ and $k_i \geq 3$, except for $r = k_1 = k_2 = k_3 = 3$. However, the only known such case is $R_4(3) \leq 62$, for which (a) gives an upper bound of 66.

(b) $R_k(3) \geq 3R_{k-1}(3) + R_{k-3}(3) - 3$ [Chu1]

(c) $R(3, k_1, \dots, k_r) \geq 4R(k_1 - 1, k_2, \dots, k_r) - 3$ for $k_1 \geq 5, r \geq 2$ and $k_i \geq 3$ [XX2]

(d) $R(3, 3, 3, k_1, \dots, k_r) \geq 3R(3, 3, k_1, \dots, k_r) + R(k_1, \dots, k_r) - 3$ [Rob2]

(e) Bounds for $R_k(3)$ [AH, Fre, Chu2, ChGri, GrRö, Wan]

(f) $R(k_1, \dots, k_r) \geq S(k_1, \dots, k_r) + 2$, where $S(k_1, \dots, k_r)$ is the generalized Schur number [AH, Gi1, Gi2]. In particular, the special case $k_1 = \dots = k_r = 3$ has been widely studied [Fre, FreSw, Ex10, Rob3].

(g) $R(k_1, \dots, k_r) \geq L(k_1, \dots, k_r) + 1$, where $L(k_1, \dots, k_r)$ is the maximal order of cyclic (k_1, \dots, k_r) -coloring, which can be considered a special case of Schur partitions defining (symmetric) Schur numbers. Many lower bounds for Ramsey numbers were established by cyclic colorings, and thus the following recurrence can be used to derive lower bounds for higher parameters.

$$L(k_1, \dots, k_r, k_{r+1}) \geq (2k_{r+1} - 3)L(k_1, \dots, k_r) - k_{r+1} + 2$$
 [Gi2]

(h) $R_r(m) \geq p + 1$ and $R_r(m + 1) \geq r(p + 1) + 1$ if there exists a K_m -free cyclotomic r -class association scheme of order p [Mat].

(i) $R_r(m) \geq c_m (2m - 3)^r$, and some slight improvements of the exponent r for small values of m [AH, Gi1, Gi2, Song2].

(j) $R_r(pq + 1) > (R_r(p + 1) - 1)(R_r(q + 1) - 1)$ [Abb1]

(k) $R(p_1q_1 + 1, \dots, p_rq_r + 1) > (R(p_1 + 1, \dots, p_r + 1) - 1)(R(q_1 + 1, \dots, q_r + 1) - 1)$ [Song3]

(l) $R_{r+s}(m) > (R_r(m) - 1)(R_s(m) - 1)$ [Song2]

(m) $R(k_1, k_2, \dots, k_r) > (R(k_1, \dots, k_i) - 1)(R(k_{i+1}, \dots, k_r) - 1)$ in [Song1], see [ExRa, XX2].

(n) $R(k_1, k_2, \dots, k_r) > (k_1 + 1)(R(k_2 - k_1 + 1, k_3, \dots, k_r) - 1)$ [Rob5]

(o) Constructions in [XX2] show how to increase the right hand side in (k) and (m) by a suitable choice and composition of colorings for smaller parameters. Several of the best known constructions for specific parameters can be obtained this way.

All lower bounds in (b) through (o) above are constructive. (d) generalizes (b), (k) generalizes both (j) and (m), and (m) generalizes (l). Observe that the construction (k) with $q_1 = \dots = q_i = 1 = p_{i+1} = \dots = p_r$ is the same as (m).

5.4. Multicolor results for cycles

- $R_3(C_4) = 11$ [BS] and $R_4(C_4) \geq 18$ [Ex2].
- $R(C_n, C_n, C_n) \leq (4 + o(1))n$, with equality for odd n [Łuc]. It was conjectured in [BoEr, Erd] that for all odd $n \geq 5$ we have $R(C_n, C_n, C_n) = 4n - 3$.
- Formulas for $R(C_n, C_m, C_k)$ and $R(C_n, C_m, C_k, C_l)$ for n sufficiently large [EFRS1].
- $R_k(C_4) \leq k^2 + k + 1$ for all $k \geq 1$, $R_k(C_4) \geq k^2 - k + 2$ for all $k - 1$ which is a prime power [Ir, Chu2, ChGra], and $R_k(C_4) \geq k^2 + 2$ for odd prime power k [LaWo1]. The latter was extended to all prime power k in [Ling, LaMu].
- Bounds for $R_k(C_m)$ [GRS]
- See also section 5.2 above.

5.5. Other general multicolor results

- General bounds for $R_k(G)$ [CH3, Par6]
- Formulas for $R_k(G)$ for G being P_3 , $2K_2$ and $K_{1,3}$ for all k , and for P_4 if k is not divisible by 3 [Ir]. Wallis [Wall] showed $R_6(P_4) = 13$, which already implied $R_{3t}(P_4) = 6t + 1$, for all $t \geq 2$. Independently, the case $R_k(P_4)$ for $k \neq 3^m$ was completed by Lindström in [Lind], and later Bierbrauer proved $R_{3^m}(P_4) = 2 \cdot 3^m + 1$ for all $m \geq 1$.
- Bounds on $R_k(K_{s,t})$, in particular for $K_{2,2} = C_4$ and $K_{2,t}$ [ChGra, AFM].
- Bounds on $R_k(G)$ for unicyclic graphs G of odd girth. Some exact values for special graphs G , for $k = 3$ and $k = 4$ [KrRod].
- $tk^2 + 1 \leq R_k(K_{2,t+1}) \leq tk^2 + k + 2$, where upper bound is general, and lower bound holds when both t and k are prime powers [LaMu].
- Monotone paths and cycles [Lef].
- Formulas for $R(P_{n_1}, \dots, P_{n_k})$, except few cases [FS2].
- Formulas for $R(S_1, \dots, S_k)$, where S_i 's are arbitrary stars [BuRo1].
- Formulas for $R(S_1, \dots, S_k, K_n)$, where S_i 's are arbitrary stars [Jac].
- Formulas for $R(S_1, \dots, S_k, nP_2)$, where S_i 's are arbitrary stars [CL2].
- Formulas for $R(S_1, \dots, S_k, T)$, where S_i 's are stars and T is a tree [ZZ].
- Formulas for $R(pP_3, qP_3, rP_3)$ and $R(pP_4, qP_4, rP_4)$ [Scob].
- Cockayne and Lorimer [CL1] found the exact formula for $R(n_1P_2, \dots, n_kP_2)$, and later Lorimer [Lor] extended it to a more general case of $R(K_m, n_1P_2, \dots, n_kP_2)$. Still more general cases of the latter, with multiple copies of the complete graph and forests, were studied in [Stahl, LorSe, LorSo].
- If G is connected and $R(K_k, G) = (k-1)(n(G)-1) + 1$, in particular if G is any tree, then $R(K_{n_1}, \dots, K_{n_k}, G) = (R(K_{n_1}, \dots, K_{n_k}) - 1)(n(G) - 1) + 1$ [BE2]. A generali-

zation for connected G_1, \dots, G_n in place of G appeared in [Jac].

- Study of $R(S, G_1, \dots, G_k)$ for large sparse S [EFRS1, Bu3].
- Constructive bound $R(G_1, \dots, G_{t^{n-1}}) \geq t^n + 1$ for some families of decompositions of K_{t^n} [LaWo1, LaWo2].
- Bounds for trees $R_k(T)$ and forests $R_k(F)$ [EG, GRS, BB, GT, Bra1, Bra2, SwPr].
- See also surveys listed in section 7.

6. Hypergraph Numbers

The only known value of a classical Ramsey number for hypergraphs:

$$R(4,4;3) = 13 \quad \text{[MR1]}$$

more than 200000 critical colorings

Other hypergraph cases:

$$33 \leq R(4,5;3) \quad \text{[Ex13]}$$

$$63 \leq R(5,5;3) \quad \text{[Ea1]}$$

$$56 \leq R(4,4,4;3) \quad \text{[Ex8]}$$

$$34 \leq R(5,5;4) \quad \text{[Ex11]}$$

$$R(K_4 - t, K_4 - t; 3) = 7 \quad \text{[Ea2]}$$

$$R(K_4 - t, K_4; 3) = 8 \quad \text{[Sob, Ex1, MR1]}$$

$$14 \leq R(K_4 - t, K_5; 3) \quad \text{[Ex1]}$$

$$13 \leq R(K_4 - t, K_4 - t, K_4 - t; 3) \leq 17 \quad \text{[Ex1] [Ea1]}$$

The computer evaluation of $R(4,4;3)$ in [MR1] consisted of an improvement of the upper bound from 15 to 13, which followed an extensive theoretical study of this number in [Gi4, Is1, Sid1]. Exoo in [Ex1] announced the bounds $R(4,5;3) \geq 30$ and $R(5,5;4) \geq 27$ without presenting the constructions. The bound of $R(4,5;3) \geq 24$ was obtained by Isbell [Is2]. Shastri in [Sha] shows a weak bound $R(5,5;4) \geq 19$ (now 34 in [Ex11]), nevertheless his lemmas and those in [Ka3, Abb2, GRS, HuSo] can be used to derive other lower bounds for higher numbers.

Several lower bound constructions for 3-uniform hypergraphs were presented in [HuSo]. Study of lower bounds on $R(p,q;4)$ can be found in [Song3] and [SYL, Song4] (the latter two papers are almost identical in contents). Most lower bounds in these papers can be easily improved by using the same techniques, but starting with better constructions for small parameters listed above.

Let $H^{(r)}(s, t)$ be the complete r -partite r -uniform hypergraph with $r - 2$ parts of size 1, one part of size s , and one part of size t (for example, for $r = 2$ it is the same as $K_{s, t}$). For the multicolor numbers, Lazebnik and Mubayi [LaMu] proved that

$$tk^2 - k + 1 \leq R_k(H^{(r)}(2, t+1)) \leq tk^2 + k + r,$$

where the lower bound holds when both t and k are prime powers. For the general case of $H^{(r)}(s, t)$, more bounds are presented in [LaMu].

Lower bounds on $R_m(k; s)$ are discussed in [DLR, AW]. In [AS], it is shown that for some values of a, b the numbers $R(m, a, b; 3)$ are at least exponential in m . General lower bounds for large number of colors were given in an early paper by Hirschfeld [Hir], and some of them were later improved in [AL]. Other theoretical results on hypergraph numbers are gathered in [GrRö, GRS].

7. Cumulative Data and Surveys

7.1. Cumulative data for two colors

- [CH1] $R(G, G)$ for all graphs G without isolates on at most 4 vertices.
- [CH2] $R(G, H)$ for all graphs G and H without isolates on at most 4 vertices.
- [Clan] $R(G, H)$ for all graphs G on at most 4 vertices and H on 5 vertices, except five entries (now all solved).
- [He4] All critical colorings for $R(G, H)$, for isolate-free graphs G and H as in [Clan] above.
- [Bu4] $R(G, G)$ for all graphs G without isolates and with at most 6 edges.
- [He1] $R(G, G)$ for all graphs G without isolates and with at most 7 edges.
- [HM2] $R(G, G)$ for all graphs G on 5 vertices and with 7 or 8 edges.
- [He2] $R(G, H)$ for all graphs G and H on 5 vertices without isolates, except 7 entries (5 still open).
- [HoMe] $R(G, H)$ for $G = K_{1,3} + e$ and $G = K_4 - e$ versus all connected graphs H on 6 vertices, except $R(K_4 - e, K_6)$. The result $R(K_4 - e, K_6) = 21$ was claimed by McNamara [McN, unpublished].
- [FRS4] $R(G, T)$ for all connected graphs G on at most 5 vertices and all (except some cases) trees T .
- [FRS1] $R(K_3, G)$ for all connected graphs G on 6 vertices.
- [Jin] $R(K_3, G)$ for all connected graphs G on 7 vertices. Some errors in [Jin] were found by [SchSch1].
- [Brin] $R(K_3, G)$ for all connected graphs G on at most 8 vertices. The numbers for K_3 versus sets of graphs with fixed number of edges, on at most 8 vertices,

were presented in [KM1].

- [BBH1] $R(K_3, G)$ for all connected graphs G on 9 vertices. See also [BBH2].
- [JR3] $R(C_4, G)$ for all graphs G on at most 6 vertices.
- [JR4] $R(C_5, G)$ for all graphs G on at most 6 vertices.
- [JR2] $R(C_6, G)$ for all graphs G on at most 5 vertices.

Chvátal and Harary [CH1, CH2] formulated several simple but very useful observations how to discover values of some numbers. All five missing entries in the tables of Clancy [Clan] have been solved. Out of 7 open cases in [He2] 2 have been solved, the bounds for 2 were improved, and the status of the other 3 did not change. Section 4.14 lists 4 of them (labeled [He2]): 1 solved and 3 still open. $R(4, 5) = R(G_{19}, G_{23}) = 25$ is the second solved case. The other 2 open entries are K_5 versus K_5 (see section 2) and K_5 versus $K_5 - e$ (see section 3).

7.2. Cumulative data for three colors

- [YR3] $R_3(G)$ for all graphs G with at most 4 edges and no isolates.
- [YR1] $R_3(G)$ for all graphs G with 5 edges and no isolates, except $K_4 - e$. The case of $R_3(K_4 - e)$ remains open (see section 5.2).
- [YY] $R_3(G)$ for all graphs G with 6 edges and no isolates, except 10 cases.
- [AKM] $R(F, G, H)$ for most triples of isolate-free graphs with at most 4 vertices. Some of the missing cases completed in [KM2].

7.3. Surveys

- [Bu1] A general survey of results in Ramsey graph theory by S. A. Burr (1974)
- [Par6] A general survey of results in Ramsey graph theory by T. D. Parsons (1978)
- [Har2] Summary of progress by Frank Harary (1981)
- [ChGri] A general survey of bounds and values by F. R. K. Chung and C. M. Grinstead (1983)
- [JGT] Special volume of *the Journal of Graph Theory* (1983)
- [Rob1] Nice textbook-type review of Ramsey graph theory for newcomers (1984)
- [Bu6] What can we hope to accomplish in generalized Ramsey Theory ? (1987)
- [GrRö] Survey of asymptotic problems by R. L. Graham and V. Rödl (1987)
- [GRS] An excellent book by R. L. Graham, B. L. Rothschild and J. H. Spencer, second edition (1990)
- [FRS5] Survey of graph goodness results, i.e. conditions for the formula $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$ (1991)
- [Neš] A chapter in *Handbook of Combinatorics* (1996)

- [Caro] Survey of zero-sum Ramsey theory (1996)
- [Chu4] Among 114 open problems and conjectures of Paul Erdős, presented and commented by F. R. K. Chung, 31 are concerned directly with Ramsey numbers. 216 references are given. (1997)

The surveys by S. A. Burr [Bu1] and T. D. Parsons [Par6] contain extensive chapters on general exact results in graph Ramsey theory. F. Harary presented the state of the theory in 1981 in [Har2], where he also gathered many references including seven to other survey papers. Two decades ago, Chung and Grinstead in their survey paper [ChGri] gave less data than in this note, but included a broad discussion of different methods used in Ramsey computations in the classical case. S. A. Burr, one of the most experienced researchers in Ramsey graph theory, formulated in [Bu6] seven conjectures on Ramsey numbers for sufficiently large and sparse graphs, and reviewed the evidence for them found in the literature. Three of them have been refuted in [Bra3].

For newer extensive presentations see [GRS, GrRö, FRS5, Neš], though these focus on asymptotic theory not on the numbers themselves. Finally, this compilation could not pretend to be complete without mentioning a special 1983 volume of the Journal of Graph Theory [JGT] dedicated entirely to Ramsey theory. Besides a number of research papers, it includes historical notes and presents to us Frank P. Ramsey (1903-1930) as a person.

8. Concluding Remarks

This compilation does not include information on numerous variations of Ramsey numbers, nor related topics, like size Ramsey numbers, zero-sum Ramsey numbers, irredundant Ramsey numbers, induced Ramsey numbers, local Ramsey numbers, connected Ramsey numbers, chromatic Ramsey numbers, avoiding sets of graphs in some colors, coloring graphs other than complete, or the so called Ramsey multiplicities. Interested reader can find such information in the surveys listed in section 7 here.

The author apologizes for any omissions or other errors in reporting results belonging to the scope of this work. Suggestions for any kind of corrections or additions will be greatly appreciated and considered for inclusion in the next year revision of this survey.

Acknowledgement

I would like to thank Brendan McKay, Geoffrey Exoo and Heiko Harborth for their help in gathering data for earlier versions of this survey.

References

We mark the papers containing results obtained with the help of computer algorithms with stars. We identify two categories of such papers: marked with * involving some use of computers, where the results are easily verifiable with some computations, and those marked with **, where cpu intensive algorithms have to be implemented to replicate or verify the results. The first category contains mostly constructions done by algorithms, while the second mostly nonexistence results or claims of complete enumerations of special kinds of graphs.

- [Abb1] H.L. Abbott, *Ph. D. thesis*, University of Alberta, Edmonton, 1965.
- [Abb2] H.L. Abbott, A Theorem Concerning Higher Ramsey Numbers, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) North Holland, (1975) 25-28.
- [AH] H.L. Abbott and D. Hanson, A Problem of Schur and Its Generalizations, *Acta Arithmetica*, **20** (1972) 175-187.
- [AL] H.L. Abbott and Andy Liu, Remarks on a Paper of Hirschfeld Concerning Ramsey Numbers, *Discrete Mathematics*, **39** (1982) 327-328.
- [AS] H.L. Abbott and M.J. Smuga-Otto, Lower Bounds for Hypergraph Ramsey Numbers, *Discrete Applied Mathematics*, **61** (1995) 177-180.
- [AW] H.L. Abbott and E.R. Williams, Lower Bounds for Some Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **16** (1974) 12-17.
- [AKS] M. Ajtai, J. Komlós and E. Szemerédi, A Note on Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **29** (1980) 354-360.
- [Alon1] N. Alon, Subdivided Graphs Have Linear Ramsey Numbers, *Journal of Graph Theory*, **18** (1994) 343-347.
- [Alon2] N. Alon, Explicit Ramsey Graphs and Orthonormal Labelings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R12, **1** (1994), 8 pages.
- [AP] N. Alon and P. Pudlák, Constructive Lower Bounds for off-diagonal Ramsey Numbers, *Israel Journal of Mathematics*, **122** (2001) 243-251.
- [AKM] J. Arste, K. Klamroth and I. Mengersen, Three Color Ramsey Numbers for Small Graphs, *Utilitas Mathematica*, **49** (1996) 85-96.
- [AFM] M. Axenovich, Z. Füredi and D. Mubayi, On Generalized Ramsey Theory: the Bipartite Case, *Journal of Combinatorial Theory, Series B*, **79** (2000) 66-86.
- [BSS] P.N. Balister, R.H. Schelp and M. Simonovits, A Note on Ramsey Size-Linear Graphs, *Journal of Graph Theory*, **39** (2002) 1-5.
- [Bas] E.T. Baskoro, The Ramsey Number of Paths and Small Wheels, *Majalah Ilmiah Himpunan Matematika Indonesia*, MIHMI, **8** (2002) 13-16.
- [BaSu] E.T. Baskoro and Surahmat, The Ramsey Number of Paths with respect to Wheels, *preprint*, (2002).
- [BSNM] E.T. Baskoro, Surahmat, S.M. Nababan and M. Miller, On Ramsey Graph Numbers for Trees versus Wheels of Five or Six Vertices, *to appear in Graphs and Combinatorics*, (2002).
- [-] E.T. Baskoro, see also [SuBa1, SuBa2, SuBB1, SuBB2].
- [BS] A. Bialostocki and J. Schönheim, On Some Turán and Ramsey Numbers for C_4 , in *Graph Theory and Combinatorics* (ed. B. Bollobás), Academic Press, London, (1984) 29-33.
- [Bier] J. Bierbrauer, Ramsey Numbers for the Path with Three Edges, *European Journal of Combinatorics*, **7** (1986) 205-206.
- [BB] J. Bierbrauer and A. Brandis, On Generalized Ramsey Numbers for Trees, *Combinatorica*, **5** (1985) 95-107.
- [BJYHRZ] B. Bollobás, C.J. Jayawardene, Yang Jian Sheng, Huang Yi Ru, C.C. Rousseau, and Zhang Ke Min, On a Conjecture Involving Cycle-Complete Graph Ramsey Numbers, *Australasian Journal of*

- Combinatorics*, **22** (2000) 63-71.
- [BH] R. Bolze and H. Harborth, The Ramsey Number $r(K_4 - x, K_5)$, in *The Theory and Applications of Graphs*, (Kalamazoo, MI, 1980), John Wiley & Sons, New York, (1981) 109-116.
- [BoEr] J.A. Bondy and P. Erdős, Ramsey Numbers for Cycles in Graphs, *Journal of Combinatorial Theory*, Series B, **14** (1973) 46-54.
- [-] A. Brandis, see [BB].
- [Bra1] S. Brandt, Subtrees and Subforests in Graphs, *Journal of Combinatorial Theory*, Series B, **61** (1994) 63-70.
- [Bra2] S. Brandt, Sufficient Conditions for Graphs to Contain All Subgraphs of a Given Type, *Ph.D. thesis*, Freie Universität Berlin, 1994.
- [Bra3] S. Brandt, Expanding Graphs and Ramsey Numbers, *submitted*.
- [BBH1]** S. Brandt, G. Brinkmann and T. Harmuth, All Ramsey Numbers $r(K_3, G)$ for Connected Graphs of Order 9, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R7, **5** (1998), 20 pages.
- [BBH2]** S. Brandt, G. Brinkmann and T. Harmuth, The Generation of Maximal Triangle-Free Graphs, *Graphs and Combinatorics*, **16** (2000) 149-157.
- [Brin]** G. Brinkmann, All Ramsey Numbers $r(K_3, G)$ for Connected Graphs of Order 7 and 8, *Combinatorics, Probability and Computing*, **7** (1998) 129-140.
- [-] G. Brinkmann, see also [BBH1, BBH2].
- [-] H.J. Broersma, see [SuBB1, SuBB2].
- [BR]* J.P. Burling and S.W. Reyner, Some Lower Bounds of the Ramsey Numbers $n(k, k)$, *Journal of Combinatorial Theory*, Series B, **13** (1972) 168-169.
- [Bu1] S.A. Burr, Generalized Ramsey Theory for Graphs - a Survey, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 52-75.
- [Bu2] S.A. Burr, Ramsey Numbers Involving Graphs with Long Suspended Paths, *Journal of the London Mathematical Society* (2), **24** (1981) 405-413.
- [Bu3] S.A. Burr, Multicolor Ramsey Numbers Involving Graphs with Long Suspended Path, *Discrete Mathematics*, **40** (1982) 11-20.
- [Bu4] S.A. Burr, Diagonal Ramsey Numbers for Small Graphs, *Journal of Graph Theory*, **7** (1983) 57-69.
- [Bu5] S.A. Burr, Ramsey Numbers Involving Powers, *Ars Combinatoria*, **15** (1983) 163-168.
- [Bu6] S.A. Burr, What Can We Hope to Accomplish in Generalized Ramsey Theory?, *Discrete Mathematics*, **67** (1987) 215-225.
- [Bu7] S.A. Burr, On the Ramsey Numbers $r(G, nH)$ and $r(nG, nH)$ When n Is Large, *Discrete Mathematics*, **65** (1987) 215-229.
- [Bu8] S.A. Burr, On Ramsey Numbers for Large Disjoint Unions of Graphs, *Discrete Mathematics*, **70** (1988) 277-293.
- [BE1] S.A. Burr and P. Erdős, Extremal Ramsey Theory for Graphs, *Utilitas Mathematica*, **9** (1976) 247-258.
- [BE2] S.A. Burr and P. Erdős, Generalizations of a Ramsey-Theoretic Result of Chvátal, *Journal of Graph Theory*, **7** (1983) 39-51.
- [BEFRS1] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, An Extremal Problem in Generalized Ramsey Theory, *Ars Combinatoria*, **10** (1980) 193-203.
- [BEFRS2] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for the Pair Sparse Graph-Path or Cycle, *Transactions of the American Mathematical Society*, **269** (1982) 501-512.
- [BEFRS3] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Ramsey Numbers Involving Starlike Multipartite Graphs, *Journal of Graph Theory*, **7** (1983) 395-409.

- [BEFRS4] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Ramsey Number for the Pair Complete Bipartite Graph-Graph of Limited Degree, in *Graph Theory with Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley & Sons, New York, (1985) 163-174.
- [BEFRS5] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Some Complete Bipartite Graph-Tree Ramsey Numbers, *Annals of Discrete Mathematics*, **41** (1989) 79-89.
- [BEFRSGJ] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau, R.H. Schelp, R.J. Gould and M.S. Jacobson, Goodness of Trees for Generalized Books, *Graphs and Combinatorics*, **3** (1987) 1-6.
- [BEFS] S.A. Burr, P. Erdős, R.J. Faudree and R.H. Schelp, On the Difference between Consecutive Ramsey Numbers, *Utilitas Mathematica*, **35** (1989) 115-118.
- [BES] S.A. Burr, P. Erdős and J.H. Spencer, Ramsey Theorems for Multiple Copies of Graphs, *Transactions of the American Mathematical Society*, **209** (1975) 87-99.
- [BF] S.A. Burr and R.J. Faudree, On Graphs G for Which All Large Trees Are G -good, *Graphs and Combinatorics*, **9** (1993) 305-313.
- [BG] S.A. Burr and J.W. Grossman, Ramsey Numbers of Graphs with Long Tails, *Discrete Mathematics*, **41** (1982) 223-227.
- [BuRo1] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Stars, *Utilitas Mathematica*, **4** (1973) 217-220.
- [BuRo2] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Linear Forests, *Discrete Mathematics*, **8** (1974) 245-250.
- [Bush] L.E. Bush, The William Lowell Putnam Mathematical Competition, *American Mathematical Monthly*, **60** (1953) 539-542.
- [CET]* N.J. Calkin, P. Erdős and C.A. Tovey, New Ramsey Bounds from Cyclic Graphs of Prime Order, *SIAM Journal of Discrete Mathematics*, **10** (1997) 381-387.
- [Caro] Y. Caro, Zero-Sum Problems - A Survey, *Discrete Mathematics*, **152** (1996) 93-113.
- [CLRZ] Y. Caro, Li Yusheng, C.C. Rousseau and Zhang Yuming, Asymptotic Bounds for Some Bipartite Graph - Complete Graph Ramsey Numbers, *Discrete Mathematics*, **220** (2000) 51-56.
- [CGP] G. Chartrand, R.J. Gould and A.D. Polimeni, On Ramsey Numbers of Forests versus Nearly Complete Graphs, *Journal of Graph Theory*, **4** (1980) 233-239.
- [CRSPS] G. Chartrand, C.C. Rousseau, M.J. Stewart, A.D. Polimeni and J. Sheehan, On Star-Book Ramsey Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 203-214.
- [CS] G. Chartrand and S. Schuster, On the existence of specified cycles in complementary graphs, *Bulletin of the American Mathematical Society*, **77** (1971) 995-998.
- [Chen] Chen Guantao, A Result on C_4 -Star Ramsey Numbers, *Discrete Mathematics*, **163** (1997) 243-246.
- [ChenS] Chen Guantao and R.H. Schelp, Graphs with Linearly Bounded Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **57** (1993) 138-149.
- [ChZZ] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Paths versus Wheels, *preprint*, (2002).
- [Chu1] F.R.K. Chung, On the Ramsey Numbers $N(3,3,\dots,3; 2)$, *Discrete Mathematics*, **5** (1973) 317-321.
- [Chu2] F.R.K. Chung, On Triangular and Cyclic Ramsey Numbers with k Colors, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 236-241.
- [Chu3] F.R.K. Chung, A Note on Constructive Methods for Ramsey Numbers, *Journal of Graph Theory*, **5** (1981) 109-113.
- [Chu4] F.R.K. Chung, Open problems of Paul Erdős in Graph Theory, *Journal of Graph Theory*, **25** (1997) 3-36.
- [CCD] F.R.K. Chung, R. Cleve and P. Dagum, A Note on Constructive Lower Bounds for the Ramsey Numbers $R(3, t)$, *Journal of Combinatorial Theory, Series B*, **57** (1993) 150-155.

- [ChGra] F.R.K. Chung and R.L. Graham, On Multicolor Ramsey Numbers for Complete Bipartite Graphs, *Journal of Combinatorial Theory, Series B*, **18** (1975) 164-169.
- [ChGri] F.R.K. Chung and C.M. Grinstead, A Survey of Bounds for Classical Ramsey Numbers, *Journal of Graph Theory*, **7** (1983) 25-37.
- [Chv] V. Chvátal, Tree-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 93.
- [CH1] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, II. Small Diagonal Numbers, *Proceedings of the American Mathematical Society*, **32** (1972) 389-394.
- [CH2] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, III. Small Off-Diagonal Numbers, *Pacific Journal of Mathematics*, **41** (1972) 335-345.
- [CH3] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, I. Diagonal Numbers, *Periodica Mathematica Hungarica*, **3** (1973) 115-124.
- [CRST] V. Chvátal, V. Rödl, E. Szemerédi and W.T. Trotter Jr., The Ramsey Number of a Graph with Bounded Maximum Degree, *Journal of Combinatorial Theory, Series B*, **34** (1983) 239-243.
- [ChvS] V. Chvátal and A. Schwenk, On the Ramsey Number of the Five-Spoked Wheel, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 247-261.
- [Clan] M. Clancy, Some Small Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 89-91.
- [Clap] C. Clapham, The Ramsey Number $r(C_4, C_4, C_4)$, *Periodica Mathematica Hungarica*, **18** (1987) 317-318.
- [CEHMS] C. Clapham, G. Exoo, H. Harborth, I. Mengersen and J. Sheehan, The Ramsey Number of $K_5 - e$, *Journal of Graph Theory*, **13** (1989) 7-15.
- [Clark] L. Clark, On Cycle-Star Graph Ramsey Numbers, *Congressus Numerantium*, **50** (1985) 187-192.
- [-] R. Cleve, see [CCD].
- [CL1] E.J. Cockayne and P.J. Lorimer, The Ramsey Number for Stripes, *Journal of the Australian Mathematical Society, Series A*, **19** (1975) 252-256.
- [CL2] E.J. Cockayne and P.J. Lorimer, On Ramsey Graph Numbers for Stars and Stripes, *Canadian Mathematical Bulletin*, **18** (1975) 31-34.
- [CPR] B. Codenotti, P. Pudlák and G. Resta, Some Structural Properties of Low-Rank Matrices Related to Computational Complexity, *Theoretical Computer Science*, **235** (2000) 89-107.
- [CsKo] R. Csákány and J. Komlós, The Smallest Ramsey Numbers, *Discrete Mathematics*, **199** (1999) 193-199.
- [-] P. Dagum, see [CCD].
- [Den] T. Denley, The Ramsey Numbers for Disjoint Unions of Cycles, *Discrete Mathematics*, **149** (1996) 31-44.
- [DLR] D. Duffus, H. Lefmann and V. Rödl, Shift Graphs and Lower Bounds on Ramsey Numbers $r_k(l; r)$, *Discrete Mathematics*, **137** (1995) 177-187.
- [Ea1] Easy to obtain by simple combinatorics from other results, in particular by using graphs establishing lower bounds with smaller parameters.
- [Ea2] Unique 2-(6,3,2) design gives lower bound 7, upper bound is easy.
- [Eaton] N. Eaton, Ramsey Numbers for Sparse Graphs, *Discrete Mathematics*, **185** (1998) 63-75.
- [Erd] P. Erdős, On the Combinatorial Problems Which I Would Most Like to See Solved, *Combinatorica*, **1** (1981) 25-42.
- [EFRS1] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalized Ramsey Theory for Multiple Colors, *Journal of Combinatorial Theory, Series B*, **20** (1976) 250-264.
- [EFRS2] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Cycle-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **2** (1978) 53-64.
- [EFRS3] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for Brooms, *Congressus Numerantium*, **35** (1982) 283-293.

- [EFRS4] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Sparse Graph Ramsey Numbers, *Combinatorica*, **5** (1985) 311-318.
- [EFRS5] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Ramsey Problem of Harary on Graphs with Prescribed Size, *Discrete Mathematics*, **67** (1987) 227-233.
- [EFRS6] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Extremal Theory and Bipartite Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 103-112.
- [EFRS7] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Book-Tree Ramsey Numbers, *Scientia*, Series A: Mathematical Sciences, Valparaíso, Chile, **1** (1988) 111-117.
- [EFRS8] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Tree Graph Ramsey Numbers, in *Graph Theory and Its Applications: East and West, Proceedings of the First China-USA International Graph Theory Conference*, Annals of the New York Academy of Sciences, **576** (1989) 146-154.
- [EFRS9] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Size Linear Graphs, *Combinatorics, Probability and Computing*, **2** (1993) 389-399.
- [EG] P. Erdős and R.L. Graham, On Partition Theorems for Finite Sets, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) North Holland, (1975) 515-527.
- [-] P. Erdős, see also [BoEr, BE1, BE2, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, BES, CET].
- [Ex1]* G. Exoo, Ramsey Numbers of Hypergraphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **2** (1987) 5-11.
- [Ex2]* G. Exoo, Constructing Ramsey Graphs with a Computer, *Congressus Numerantium*, **59** (1987) 31-36.
- [Ex3]* G. Exoo, Applying Optimization Algorithm to Ramsey Problems, in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi ed.), SIAM Philadelphia, (1989) 175-179.
- [Ex4]* G. Exoo, A Lower Bound for $R(5, 5)$, *Journal of Graph Theory*, **13** (1989) 97-98.
- [Ex5]* G. Exoo, On Two Classical Ramsey Numbers of the Form $R(3, n)$, *SIAM Journal of Discrete Mathematics*, **2** (1989) 488-490.
- [Ex6]* G. Exoo, A Lower Bound for $r(K_5 - e, K_5)$, *Utilitas Mathematica*, **38** (1990) 187-188.
- [Ex7]* G. Exoo, On the Three Color Ramsey Number of $K_4 - e$, *Discrete Mathematics*, **89** (1991) 301-305.
- [Ex8]* G. Exoo, Indiana State University, *personal communication* (1992).
- [Ex9]* G. Exoo, Announcement: On the Ramsey Numbers $R(4, 6)$, $R(5, 6)$ and $R(3, 12)$, *Ars Combinatoria*, **35** (1993) 85. The construction of a graph proving $R(4, 6) \geq 35$ is presented in detail at <http://isu.indstate.edu/ge/RAMSEY> (2001).
- [Ex10]* G. Exoo, A Lower Bound for Schur Numbers and Multicolor Ramsey Numbers of K_3 , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R8, **1** (1994), 3 pages.
- [Ex11]* G. Exoo, Indiana State University, *personal communication* (1997).
- [Ex12]* G. Exoo, Some New Ramsey Colorings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R29, **5** (1998), 5 pages. The constructions are available electronically from <http://isu.indstate.edu/ge/RAMSEY>.
- [Ex13]* G. Exoo, Indiana State University, *personal communication* (1998), constructions available at <http://isu.indstate.edu/ge/RAMSEY>.
- [Ex14]* G. Exoo, Indiana State University, *New Lower Bounds for Table III*, (2000), and constructions available at <http://isu.indstate.edu/ge/RAMSEY>.
- [Ex15]* G. Exoo, Some Applications of pq -groups in Graph Theory, *preprint*, (2002), and constructions available at <http://isu.indstate.edu/ge/RAMSEY>.
- [Ex16]* G. Exoo, Indiana State University, *personal communication* (2002), constructions available at <http://isu.indstate.edu/ge/RAMSEY>.

- [EHM1] G. Exoo, H. Harborth and I. Mengersen, The Ramsey Number of K_4 versus $K_5 - e$, *Ars Combinatoria*, **25A** (1988) 277-286.
- [EHM2] G. Exoo, H. Harborth and I. Mengersen, On Ramsey Number of $K_{2,n}$, in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi, F.R.K. Chung, R.L. Graham and D.F. Hsu eds.), SIAM Philadelphia, (1989) 207-211.
- [ExRa]* G. Exoo and S.P. Radziszowski, Constructive and Asymptotic Lower Bounds on Classical Multicolor Ramsey Numbers, *manuscript*, (2002).
- [ExRe]* G. Exoo and D.F. Reynolds, Ramsey Numbers Based on C_5 -Decompositions, *Discrete Mathematics*, **71** (1988) 119-127.
- [-] G. Exoo, see also [CEHMS].
- [FLPS] R.J. Faudree, S.L. Lawrence, T.D. Parsons and R.H. Schelp, Path-Cycle Ramsey Numbers, *Discrete Mathematics*, **10** (1974) 269-277.
- [FM]** R.J. Faudree and B.D. McKay, A Conjecture of Erdős and the Ramsey Number $r(W_6)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **13** (1993) 23-31.
- [FRS1] R.J. Faudree, C.C. Rousseau and R.H. Schelp, All Triangle-Graph Ramsey Numbers for Connected Graphs of Order Six, *Journal of Graph Theory*, **4** (1980) 293-300.
- [FRS2] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Studies Related to the Ramsey Number $r(K_5 - e)$, in *Graph Theory and Its Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley and Sons, New York, (1985) 251-271.
- [FRS3] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalizations of the Tree-Complete Graph Ramsey Number, in *Graphs and Applications*, (F. Harary and J.S. Maybee eds.), John Wiley and Sons, New York, (1985) 117-126.
- [FRS4] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Small Order Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 119-127.
- [FRS5] R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Good Idea in Ramsey Theory, in *Graph Theory, Combinatorics, Algorithms, and Applications* (San Francisco, CA 1989), SIAM Philadelphia, PA (1991) 180-189.
- [FRS6] R.J. Faudree, C.C. Rousseau and J. Sheehan, More from the Good Book, in *Proceedings of the Ninth Southeastern Conference on Combinatorics, Graph Theory, and Computing*, Utilitas Mathematica Publ., *Congressus Numerantium XXI* (1978) 289-299.
- [FRS7] R.J. Faudree, C.C. Rousseau and J. Sheehan, Strongly Regular Graphs and Finite Ramsey Theory, *Linear Algebra and its Applications*, **46** (1982) 221-241.
- [FRS8] R.J. Faudree, C.C. Rousseau and J. Sheehan, Cycle-Book Ramsey Numbers, *Ars Combinatoria*, **31** (1991) 239-248.
- [FS1] R.J. Faudree and R.H. Schelp, All Ramsey Numbers for Cycles in Graphs, *Discrete Mathematics*, **8** (1974) 313-329.
- [FS2] R.J. Faudree and R.H. Schelp, Path Ramsey Numbers in Multicolorings, *Journal of Combinatorial Theory, Series B*, **19** (1975) 150-160.
- [FS3] R.J. Faudree and R.H. Schelp, Ramsey Numbers for All Linear Forests, *Discrete Mathematics*, **16** (1976) 149-155.
- [FS4] R.J. Faudree and R.H. Schelp, Some Problems in Ramsey Theory, in *Theory and Applications of Graphs*, (conference proceedings, Kalamazoo, MI 1976), Lecture Notes in Mathematics **642**, Springer, Berlin, (1978) 500-515.
- [FSR] R.J. Faudree, R.H. Schelp and C.C. Rousseau, Generalizations of a Ramsey Result of Chvátal, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 351-361.
- [FSS1] R.J. Faudree, R.H. Schelp and M. Simonovits, On Some Ramsey Type Problems Connected with Paths, Cycles and Trees, *Ars Combinatoria*, **29A** (1990) 97-106.

- [FSS2] R.J. Faudree, A. Schelten and I. Schiermeyer, The Ramsey Number $r(C_7, C_7, C_7)$, *preprint*, (2001).
- [FS] R.J. Faudree and M. Simonovits, Ramsey Problems and Their Connection to Turán-Type Extremal Problems, *Journal of Graph Theory*, **16** (1992) 25-50.
- [-] R.J. Faudree, see also [BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, BF, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9].
- [FKR]** S. Fettes, R.L. Kramer and S.P. Radziszowski, An Upper Bound of 62 on the Classical Ramsey Number $R(3,3,3,3)$, *preprint*, (2002).
- [Fo] J. Folkman, Notes on the Ramsey Number $N(3,3,3,3)$, *Journal of Combinatorial Theory, Series A*, **16** (1974) 371-379.
- [Fra1] K. Fraughnaugh Jones, Independence in Graphs with Maximum Degree Four, *Journal of Combinatorial Theory, Series B*, **37** (1984) 254-269.
- [Fra2] K. Fraughnaugh Jones, Size and Independence in Triangle-Free Graphs with Maximum Degree Three, *Journal of Graph Theory*, **14** (1990) 525-535.
- [FL] K. Fraughnaugh and S.C. Locke, Finding Independent Sets in Triangle-free Graphs, *SIAM Journal of Discrete Mathematics*, **9** (1996) 674-681.
- [Fre] H. Fredricksen, Schur Numbers and the Ramsey Numbers $N(3,3,\dots,3;2)$, *Journal of Combinatorial Theory, Series A*, **27** (1979) 376-377.
- [FreSw]* H. Fredricksen and M.M. Sweet, Symmetric Sum-Free Partitions and Lower Bounds for Schur Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R32, **7** (2000), 9 pages.
- [-] Z. Füredi, see [AFM].
- [GeGy] L. Gerencsér and A. Gyárfás, On Ramsey-Type Problems, *Annales Universitatis Scientiarum Budapestinensis, Eötvös Sect. Math.*, **10** (1967) 167-170.
- [Gi1] G. Giraud, Une généralisation des nombres et de l'inégalité de Schur, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A437-A440.
- [Gi2] G. Giraud, Minoration de certains nombres de Ramsey binaires par les nombres de Schur généralisés, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A481-A483.
- [Gi3] G. Giraud, Nouvelles majorations des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **268** (1969) A5-A7.
- [Gi4] G. Giraud, Majoration du nombre de Ramsey ternaire-bicolore en (4,4), *C.R. Acad. Sc. Paris, Séries A-B*, **269** (1969) A620-A622.
- [Gi5] G. Giraud, Une minoration du nombre de quadrangles unicolores et son application à la majoration des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **276** (1973) A1173-A1175.
- [Gi6] G. Giraud, Sur le problème de Goodman pour les quadrangles et la majoration des nombres de Ramsey, *Journal of Combinatorial Theory, Series B*, **27** (1979) 237-253.
- [-] A.M. Gleason, see [GG].
- [GK] W. Goddard and D.J. Kleitman, An upper bound for the Ramsey numbers $r(K_3, G)$, *Discrete Mathematics*, **125** (1994) 177-182.
- [GJ1] R.J. Gould and M.S. Jacobson, Bounds for the Ramsey Number of a Disconnected Graph Versus Any Graph, *Journal of Graph Theory*, **6** (1982) 413-417.
- [GJ2] R.J. Gould and M.S. Jacobson, On the Ramsey Number of Trees Versus Graphs with Large Clique Number, *Journal of Graph Theory*, **7** (1983) 71-78.
- [-] R.J. Gould, see also [BEFRSGJ, CGP].
- [GrRö] R.L. Graham and V. Rödl, Numbers in Ramsey Theory, in *Surveys in Combinatorics*, (ed. C. Whitehead), Cambridge University Press, 1987.

- [GRR1] R.L. Graham, V. Rödl and A. Ruciński, On Graphs with Linear Ramsey Numbers, *Journal of Graph Theory*, **35** (2000) 176-192.
- [GRR2] R.L. Graham, V. Rödl and A. Ruciński, On Bipartite Graphs with Linear Ramsey Numbers, Paul Erdős and his mathematics, *Combinatorica*, **21** (2001) 199-209.
- [GRS] R.L. Graham, B.L. Rothschild and J.H. Spencer, *Ramsey Theory*, John Wiley & Sons, 1990.
- [-] R.L. Graham, see also [ChGra, EG].
- [GY] J.E. Graver and J. Yackel, Some Graph Theoretic Results Associated with Ramsey's Theorem, *Journal of Combinatorial Theory*, **4** (1968) 125-175.
- [GG] R.E. Greenwood and A.M. Gleason, Combinatorial Relations and Chromatic Graphs, *Canadian Journal of Mathematics*, **7** (1955) 1-7.
- [GH] U. Grenda and H. Harborth, The Ramsey Number $r(K_3, K_7 - e)$, *Journal of Combinatorics, Information & System Sciences*, **7** (1982) 166-169.
- [Gri] J.R. Griggs, An Upper Bound on the Ramsey Numbers $R(3, k)$, *Journal of Combinatorial Theory, Series A*, **35** (1983) 145-153.
- [GR]** C. Grinstead and S. Roberts, On the Ramsey Numbers $R(3, 8)$ and $R(3, 9)$, *Journal of Combinatorial Theory, Series B*, **33** (1982) 27-51.
- [-] C. Grinstead, see also [ChGri].
- [Gro1] J.W. Grossman, Some Ramsey Numbers of Unicyclic Graphs, *Ars Combinatoria*, **8** (1979) 59-63.
- [Gro2] J.W. Grossman, The Ramsey Numbers of the Union of Two Stars, *Utilitas Mathematica*, **16** (1979) 271-279.
- [GHK] J.W. Grossman, F. Harary and M. Klawe, Generalized Ramsey Theory for Graphs, X: Double Stars, *Discrete Mathematics*, **28** (1979) 247-254.
- [-] J.W. Grossman, see also [BG].
- [GV] Guo Yubao and L. Volkmann, Tree-Ramsey Numbers, *Australasian Journal of Combinatorics*, **11** (1995) 169-175.
- [-] L. Gupta, see [GGS].
- [GGS] S.K. Gupta, L. Gupta and A. Sudan, On Ramsey Numbers for Fan-Fan Graphs, *Journal of Combinatorics, Information & System Sciences*, **22** (1997) 85-93.
- [GT] A. Gyárfás and Z. Tuza, An Upper Bound on the Ramsey Number of Trees, *Discrete Mathematics*, **66** (1987) 309-310.
- [-] A. Gyárfás, see also [GeGy].
- [Haa]* H. Haanpää, A Lower Bound for a Ramsey Number, *Congressus Numerantium*, **144** (2000) 189-191.
- [Häg] R. Häggkvist, On the Path-Complete Bipartite Ramsey Number, *Discrete Mathematics*, **75** (1989) 243-245.
- [Han]* D. Hanson, Sum-free Sets and Ramsey Numbers, *Discrete Mathematics*, **14** (1976) 57-61.
- [Har1] F. Harary, Recent Results on Generalized Ramsey Theory for Graphs, in *Graph Theory and Applications*, (Y. Alavi et al. eds.) Springer, Berlin (1972) 125-138.
- [Har2] F. Harary, Generalized Ramsey Theory I to XIII: Achievement and Avoidance Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 373-390.
- [-] F. Harary, see also [CH1, CH2, CH3, GHK].
- [HM1] H. Harborth and I. Mengersen, An Upper Bound for the Ramsey Number $r(K_5 - e)$, *Journal of Graph Theory*, **9** (1985) 483-485.
- [HM2] H. Harborth and I. Mengersen, All Ramsey Numbers for Five Vertices and Seven or Eight Edges, *Discrete Mathematics*, **73** (1988/89) 91-98.
- [HM3] H. Harborth and I. Mengersen, The Ramsey Number of $K_{3,3}$, in *Combinatorics, Graph Theory, and Applications*, Vol. **2** (Y. Alavi, G. Chartrand, O.R. Oellermann and J. Schwenk eds.), John Wiley &

- Sons, (1991) 639-644.
- [-] H. Harborth, see also [BH, CEHMS, EHM1, EHM2, GH].
- [HM] M. Harborth and I. Mengersen, Some Ramsey Numbers for Complete Bipartite Graphs, *Australasian Journal of Combinatorics*, **13** (1996) 119-128.
- [-] T. Harmuth, see [BBH1, BBH2].
- [He1] G.R.T. Hendry, Diagonal Ramsey Numbers for Graphs with Seven Edges, *Utilitas Mathematica*, **32** (1987) 11-34.
- [He2] G.R.T. Hendry, Ramsey Numbers for Graphs with Five Vertices, *Journal of Graph Theory*, **13** (1989) 245-248.
- [He3] G.R.T. Hendry, The Ramsey Numbers $r(K_2 + \bar{K}_3, K_4)$ and $r(K_1 + C_4, K_4)$, *Utilitas Mathematica*, **35** (1989) 40-54, addendum in **36** (1989) 25-32.
- [He4] G.R.T. Hendry, Critical Colorings for Clancy's Ramsey Numbers, *Utilitas Mathematica*, **41** (1992) 181-203.
- [He5] G.R.T. Hendry, Small Ramsey Numbers II. Critical Colorings for $r(C_5 + e, K_5)$, *Quaestiones Mathematica*, **17** (1994) 249-258.
- [-] G.R.T. Hendry, see also [YH].
- [HI]* R. Hill and R.W. Irving, On Group Partitions Associated with Lower Bounds for Symmetric Ramsey Numbers, *European Journal of Combinatorics*, **3** (1982) 35-50.
- [Hir] J. Hirschfeld, A Lower Bound for Ramsey's Theorem, *Discrete Mathematics*, **32** (1980) 89-91.
- [HoMe] M. Hoeth and I. Mengersen, Ramsey Numbers for Graphs of Order Four versus Connected Graphs of Order Six, *Utilitas Mathematica*, **57** (2000) 3-19.
- [HuSo] Huang Da Ming and Song En Min, Properties and Lower Bounds of the Third Order Ramsey Numbers (in Chinese), *Mathematica Applicata*, **9** (1996) 105-107.
- [Hua1] Huang Guotai, Some Generalized Ramsey Numbers (in Chinese), *Mathematica Applicata*, **1** (1988) 97-101.
- [Hua2] Huang Guotai, An Unsolved Problem of Gould and Jacobson (in Chinese), *Mathematica Applicata*, **9** (1996) 234-236.
- [HZ1] Huang Yi Ru and Zhang Ke Min, An New Upper Bound Formula for Two Color Classical Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **28** (1998) 347-350.
- [HZ2] Huang Yi Ru and Zhang Ke Min, New Upper Bounds for Ramsey Numbers, *European Journal of Combinatorics*, **19** (1998) 391-394.
- [-] Huang Yi Ru, see also [BJYHRZ, YHZ1, YHZ2].
- [Ir] R.W. Irving, Generalised Ramsey Numbers for Small Graphs, *Discrete Mathematics*, **9** (1974) 251-264.
- [-] R.W. Irving, see also [HI].
- [Is1] J.R. Isbell, $N(4,4;3) \geq 13$, *Journal of Combinatorial Theory*, **6** (1969) 210.
- [Is2] J.R. Isbell, $N(5,4;3) \geq 24$, *Journal of Combinatorial Theory*, Series A, **34** (1983) 379-380.
- [Jac] M.S. Jacobson, On the Ramsey Number for Stars and a Complete Graph, *Ars Combinatoria*, **17** (1984) 167-172.
- [-] M.S. Jacobson, see also [BEFRSGJ, GJ1, GJ2].
- [JR1] C.J. Jayawardene and C.C. Rousseau, An Upper Bound for the Ramsey Number of a Quadrilateral versus a Complete Graph on Seven Vertices, *Congressus Numerantium*, **130** (1998) 175-188.
- [JR2] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers $r(C_6, G)$ for All Graphs G of Order Less than Six, *Congressus Numerantium*, **136** (1999) 147-159.
- [JR3] C.J. Jayawardene and C.C. Rousseau, The Ramsey Numbers for a Quadrilateral vs. All Graphs on Six Vertices, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **35** (2000) 71-87.

- [JR4] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers $r(C_5, G)$ for All Graphs G of Order Six, *Ars Combinatoria*, **57** (2000) 163-173.
- [JR5] C.J. Jayawardene and C.C. Rousseau, The Ramsey Number for a Cycle of Length Five vs. a Complete Graph of Order Six, *Journal of Graph Theory*, **35** (2000) 99-108.
- [-] C.J. Jayawardene, see also [BJYHRZ, RoJa1, RoJa2].
- [Jin]** Jin Xia, Ramsey Numbers Involving a Triangle: Theory & Applications, *Technical Report RIT-TR-93-019*, MS thesis, Department of Computer Science, Rochester Institute of Technology, 1993.
- [-] Jin Xia, see also [RaJi].
- [JGT] Special volume on Ramsey theory of *Journal of Graph Theory*, Volume **7**, Number 1, (1983).
- [Ka1] J.G. Kalbfleisch, Construction of Special Edge-Chromatic Graphs, *Canadian Mathematical Bulletin*, **8** (1965) 575-584.
- [Ka2]* J.G. Kalbfleisch, Chromatic Graphs and Ramsey's Theorem, *Ph.D. thesis*, University of Waterloo, January 1966.
- [Ka3] J.G. Kalbfleisch, On Robillard's Bounds for Ramsey Numbers, *Canadian Mathematical Bulletin*, **14** (1971) 437-440.
- [KS] J.G. Kalbfleisch and R.G. Stanton, On the Maximal Triangle-Free Edge-Chromatic Graphs in Three Colors, *Journal of Combinatorial Theory*, **5** (1968) 9-20.
- [KáRos] G. Károlyi and V. Rosta, Generalized and Geometric Ramsey Numbers for Cycles, *Theoretical Computer Science*, **263** (2001) 87-98.
- [Kéry] G. Kéry, On a Theorem of Ramsey (in Hungarian), *Matematikai Lapok*, **15** (1964) 204-224.
- [Kim] J.H. Kim, The Ramsey Number $R(3, t)$ has Order of Magnitude $t^2/\log t$, *Random Structures and Algorithms*, **7** (1995) 173-207.
- [KM1] K. Klamroth and I. Mengersen, Ramsey Numbers of K_3 versus (p, q) -Graphs, *Ars Combinatoria*, **43** (1996) 107-120.
- [KM2] K. Klamroth and I. Mengersen, The Ramsey Number of $r(K_{1,3}, C_4, K_4)$, *Utilitas Mathematica*, **52** (1997) 65-81.
- [-] K. Klamroth, see also [AKM].
- [-] M. Klawe, see [GHK].
- [-] D.J. Kleitman, see [GK].
- [Köh] W. Köhler, On a Conjecture by Grossman, *Ars Combinatoria*, **23** (1987) 103-106.
- [-] J. Komlós, see [CsKo, AKS].
- [KoRö] A.V. Kostochka and V. Rödl, On Graphs with Small Ramsey Numbers, *Journal of Graph Theory*, **37** (2001) 198-204.
- [-] R.L. Kramer, see [FKR].
- [KrRod] I. Krasikov and Y. Roditty, On Some Ramsey Numbers of Unicyclic Graphs, *Bulletin of the Institute of Combinatorics and its Applications*, **33** (2001) 29-34.
- [KLR]* D.L. Kreher, Li Wei and S.P. Radziszowski, Lower Bounds for Multi-Colored Ramsey Numbers From Group Orbits, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 87-95.
- [-] D.L. Kreher, see also [RK1, RK2, RK3, RK4].
- [La1] S.L. Lawrence, Cycle-Star Ramsey Numbers, *Notices of the American Mathematical Society*, **20** (1973) Abstract A- 420.
- [La2] S.L. Lawrence, Bipartite Ramsey Theory, *Notices of the American Mathematical Society*, **20** (1973) Abstract A- 562.
- [-] S.L. Lawrence, see also [FLPS].

- [LayMa] C. Laywine and J.P. Mayberry, A Simple Construction Giving the Two Non-isomorphic Triangle-Free 3-Colored K_{16} 's, *Journal of Combinatorial Theory, Series B*, **45** (1988) 120-124.
- [LaMu] F. Lazebnik and D. Mubayi, New Lower Bounds for Ramsey Numbers of Graphs and Hypergraphs, *preprint*, (2001).
- [LaWo1] F. Lazebnik and A. Woldar, New Lower Bounds on the Multicolor Ramsey Numbers $r_k(C_4)$, *Journal of Combinatorial Theory, Series B*, **79** (2000) 172-176.
- [LaWo2] F. Lazebnik and A. Woldar, General Properties of Some Families of Graphs Defined by Systems of Equations, *to appear*.
- [Lef] H. Lefmann, Ramsey Numbers for Monotone Paths and Cycles, *Ars Combinatoria*, **35** (1993) 271-279.
- [-] H. Lefmann, see also [DLR].
- [Les]* A. Lesser, Theoretical and Computational Aspects of Ramsey Theory, *Examensarbeten i Matematik, Matematiska Institutionen, Stockholms Universitet*, **3** (2001).
- [LiWa] Li Da Yong and Wang Zhi Jian, The Ramsey Number $r(mC_4, nC_4)$ (in Chinese), *Journal of Shanghai Tiedao University*, **20** (1999) 66-70, 83.
- [-] Li Guiqing, see [LSZL, SLLL, SLL1, SLZL].
- [-] Li Qiao, see [SLLL, SLL2, SLL3].
- [-] Li Wei, see [KLR].
- [Li] Li Yusheng, Some Ramsey Numbers of Graphs with Bridge, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **25** (1997) 225-229.
- [LR1] Li Yusheng and C.C. Rousseau, On Book-Complete Graph Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **68** (1996) 36-44.
- [LR2] Li Yusheng and C.C. Rousseau, Fan-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **23** (1996) 413-420.
- [LR3] Li Yusheng and C.C. Rousseau, On the Ramsey Number $r(H + \overline{K}_n, K_n)$, *Discrete Mathematics*, **170** (1997) 265-267.
- [LR4] Li Yusheng and C.C. Rousseau, A Ramsey Goodness Result for Graphs with Many Pendant Edges, *Ars Combinatoria*, **49** (1998) 315-318.
- [LRS] Li Yusheng, C.C. Rousseau and L. Soltés, Ramsey Linear Families and Generalized Subdivided Graphs, *Discrete Mathematics*, **170** (1997) 269-275.
- [LRZ] Li Yusheng, C.C. Rousseau and Zang Wenan, Asymptotic Upper Bounds for Ramsey Functions, *Graphs and Combinatorics*, **17** (2001) 123-128.
- [-] Li Yusheng, see also [CLRZ].
- [-] Li Zhenchong, see [LSL].
- [Lind] B. Lindström, Undecided Ramsey-Numbers for Paths, *Discrete Mathematics*, **43** (1983) 111-112.
- [Ling] A.C.H. Ling, Some Applications of Combinatorial Designs to Extremal Graph Theory, *to appear*.
- [-] Andy Liu, see [AL].
- [-] Liu Yanwu, see [SYL].
- [Loc] S.C. Locke, Bipartite Density and the Independence Ratio, *Journal of Graph Theory*, **10** (1986) 47-53.
- [-] S.C. Locke, see also [FL].
- [Lor] P.J. Lorimer, The Ramsey Numbers for Stripes and One Complete Graph, *Journal of Graph Theory*, **8** (1984) 177-184.
- [LorMu] P.J. Lorimer and P.R. Mullins, Ramsey Numbers for Quadrangles and Triangles, *Journal of Combinatorial Theory, Series B*, **23** (1977) 262-265.

- [LorSe] P.J. Lorimer and R.J. Segegin, Ramsey Numbers for Multiple Copies of Complete Graphs, *Journal of Graph Theory*, **2** (1978) 89-91.
- [LorSo] P.J. Lorimer and W. Solomon, The Ramsey Numbers for Stripes and Complete Graphs 1, *Discrete Mathematics*, **104** (1992) 91-97. Corrigendum in *Discrete Mathematics*, **131** (1994) 395.
- [-] P.J. Lorimer, see also [CL1, CL2].
- [Łuc] T. Łuczak, $R(C_n, C_n, C_n) \leq (4 + o(1))n$, *Journal of Combinatorial Theory, Series B*, **75** (1999) 174-187.
- [LSL]* Luo Haipeng, Su Wenlong and Li Zhenchong, The Properties of Self-Complementary Graphs and New Lower Bounds for Diagonal Ramsey Numbers, *Australasian Journal of Combinatorics*, **25** (2002) 103-116.
- [LSS]* Luo Haipeng, Su Wenlong and Shen Yun-Qiu, New Lower Bounds of Ten Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **24** (2001) 81-90.
- [LSZL]* Luo Haipeng, Su Wenlong, Zhang Zhengyou and Li Guiqing, New Lower Bounds for Twelve Classical 2-Color Ramsey Numbers $R(k, l)$ (in Chinese), *Guangxi Sciences*, **7**, 2 (2000) 120-121.
- [-] Luo Haipeng, see also [SL, SLLL, SLL1, SLL2, SLL3, SLZL, ZSL].
- [Mac]* J. Mackey, Combinatorial Remedies, *Ph.D. Thesis*, Department of Mathematics, University of Hawaii, 1994.
- [Mat]* R. Mathon, Lower Bounds for Ramsey Numbers and Association Schemes, *Journal of Combinatorial Theory, Series B*, **42** (1987) 122-127.
- [-] J.P. Mayberry, see [LayMa].
- [McS] C. McDiarmid and A. Steger, Tidier Examples for Lower Bounds on Diagonal Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **74** (1996) 147-152.
- [MPR]** B.D. McKay, K. Piwakowski and S.P. Radziszowski, Ramsey Numbers for Triangles versus Almost-Complete Graphs, *preprint*, (2002).
- [MR1]** B.D. McKay and S.P. Radziszowski, The First Classical Ramsey Number for Hypergraphs is Computed, *Proceedings of the Second Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA'91, San Francisco, (1991) 304-308.
- [MR2]* B.D. McKay and S.P. Radziszowski, A New Upper Bound for the Ramsey Number $R(5, 5)$, *Australian Journal of Combinatorics*, **5** (1992) 13-20.
- [MR3]** B.D. McKay and S.P. Radziszowski, Linear Programming in Some Ramsey Problems, *Journal of Combinatorial Theory, Series B*, **61** (1994) 125-132.
- [MR4]** B.D. McKay and S.P. Radziszowski, $R(4, 5) = 25$, *Journal of Graph Theory*, **19** (1995) 309-322.
- [MR5]** B.D. McKay and S.P. Radziszowski, Subgraph Counting Identities and Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **69** (1997) 193-209.
- [MZ]** B.D. McKay and Zhang Ke Min, The Value of the Ramsey Number $R(3, 8)$, *Journal of Graph Theory*, **16** (1992) 99-105.
- [-] B.D. McKay, see also [FM].
- [McN]** J. McNamara, SUNY Brockport, *personal communication* (1995).
- [McR]** J. McNamara and S.P. Radziszowski, The Ramsey Numbers $R(K_4 - e, K_6 - e)$ and $R(K_4 - e, K_7 - e)$, *Congressus Numerantium*, **81** (1991) 89-96.
- [MO] I. Mengersen and J. Oeckermann, Matching-Star Ramsey Sets, *Discrete Applied Mathematics*, **95** (1999) 417-424.
- [-] I. Mengersen, see also [AKM, CEHMS, EHM1, EHM2, HoMe, HM1, HM2, HM3, HM, KM1, KM2].
- [-] M. Miller, see [BSNM].
- [MS] H. Mizuno and I. Sato, Ramsey Numbers for Unions of Some Cycles, *Discrete Mathematics*, **69** (1988) 283-294.

- [-] D. Mubayi, see [AFM, LaMu].
- [-] P.R. Mullins, see [LorMu].
- [-] S.M. Nababan, see [BSNM].
- [Neš] J. Nešetřil, Ramsey Theory, chapter 25 in *Handbook of Combinatorics*, ed. R.L. Graham, M. Grötschel and L. Lovász, The MIT-Press, Vol. II, 1996, 1331-1403.
- [-] J. Oeckermann, see [MO].
- [Par1] T.D. Parsons, The Ramsey Numbers $r(P_m, K_n)$, *Discrete Mathematics*, **6** (1973) 159-162.
- [Par2] T.D. Parsons, Path-Star Ramsey Numbers, *Journal of Combinatorial Theory*, Series B, **17** (1974) 51-58.
- [Par3] T.D. Parsons, Ramsey Graphs and Block Designs, I, *Transactions of the American Mathematical Society*, **209** (1975) 33-44.
- [Par4] T.D. Parsons, Ramsey Graphs and Block Designs, *Journal of Combinatorial Theory*, Series A, **20** (1976) 12-19.
- [Par5] T.D. Parsons, Graphs from Projective Planes, *Aequationes Mathematicae*, **14** (1976) 167-189.
- [Par6] T.D. Parsons, Ramsey Graph Theory, in *Selected Topics in Graph Theory*, (L.W. Beineke and R.J. Wilson eds.), Academic Press, (1978) 361-384.
- [-] T.D. Parsons, see also [FLPS].
- [Piw1]* K. Piwakowski, Applying Tabu Search to Determine New Ramsey Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R6, **3** (1996), 4 pages.
- [Piw2]** K. Piwakowski, A New Upper Bound for $R_3(K_4 - e)$, *Congressus Numerantium*, **128** (1997) 135-141.
- [PR1]** K. Piwakowski and S.P. Radziszowski, $30 \leq R(3,3,4) \leq 31$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **27** (1998) 135-141.
- [PR2]** K. Piwakowski and S.P. Radziszowski, Towards the Exact Value of the Ramsey Number $R(3,3,4)$, *Congressus Numerantium*, **148** (2001) 161-167.
- [-] K. Piwakowski, see also [MPR].
- [-] A.D. Polimeni, see [CGP, CRSPS].
- [-] L.M. Pretorius, see [SwPr].
- [-] P. Pudlák, see [AP, CPR].
- [Ra1]** S.P. Radziszowski, The Ramsey Numbers $R(K_3, K_8 - e)$ and $R(K_3, K_9 - e)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **8** (1990) 137-145.
- [Ra2] S.P. Radziszowski, Small Ramsey Numbers, *Technical Report RIT-TR-93-009*, Department of Computer Science, Rochester Institute of Technology (1993).
- [Ra3]** S.P. Radziszowski, On the Ramsey Number $R(K_5 - e, K_5 - e)$, *Ars Combinatoria*, **36** (1993) 225-232.
- [RaJi] S.P. Radziszowski and Jin Xia, Paths, Cycles and Wheels in Graphs without Antitriangles, *Australian Journal of Combinatorics*, **9** (1994) 221-232.
- [RK1]* S.P. Radziszowski and D.L. Kreher, Search Algorithm for Ramsey Graphs by Union of Group Orbits, *Journal of Graph Theory*, **12** (1988) 59-72.
- [RK2]** S.P. Radziszowski and D.L. Kreher, Upper Bounds for Some Ramsey Numbers $R(3, k)$, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 207-212.
- [RK3]** S.P. Radziszowski and D.L. Kreher, On $R(3, k)$ Ramsey Graphs: Theoretical and Computational Results, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 37-52.
- [RK4] S.P. Radziszowski and D.L. Kreher, Minimum Triangle-Free Graphs, *Ars Combinatoria*, **31** (1991) 65-92.
- [RT]* S.P. Radziszowski and Kung-Kuen Tse, A Computational Approach for the Ramsey Numbers $R(C_4, K_n)$, to appear.

- [-] S.P. Radziszowski, see also [ExRa, FKR, KLR, MPR, MR1, MR2, MR3, MR4, MR5, McR, PR1, PR2].
- [Ram] F.P. Ramsey, On a Problem of Formal Logic, *Proceedings of the London Mathematical Society*, **30** (1930) 264-286.
- [Rao]* S. Rao, Applying a Genetic Algorithm to Improve the Lower Bounds of Multi-Color Ramsey Numbers, *MS thesis*, Department of Computer Science, Rochester Institute of Technology, 1997.
- [-] G. Resta, see [CPR].
- [-] S.W. Reyner, see [BR].
- [-] D.F. Reynolds, see [ExRe].
- [Rob1] F.S. Roberts, *Applied Combinatorics*, Prentice-Hall, Englewood Cliffs, 1984.
- [-] J.A. Roberts, see [BuRo1, BuRo2].
- [-] S. Roberts, see [GR].
- [Rob2]* A. Robertson, New Lower Bounds for Some Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R12, **6** (1999), 6 pages.
- [Rob3]* A. Robertson, Some Results in Ramsey Theory, *Ph.D. Thesis*, Department of Mathematics, Temple University, 1999.
- [Rob4]* A. Robertson, Difference Ramsey Numbers and Issai Numbers, *Advances in Applied Mathematics*, **25** (2000) 153-162.
- [Rob5] A. Robertson, New Lower Bounds Formulas for Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #R13, **9** (2002), 6 pages.
- [-] Y. Roditty, see [KrRod].
- [RöTh] V. Rödl and R. Thomas, Arrangeability and Clique Subdivisions, in *The Mathematics of Paul Erdős II*, 236-239, Algorithms and Combinatorics **14**, Springer, Berlin, 1997.
- [-] V. Rödl, see also [CRST, DLR, GrRö, GRR1, GRR2, KoRö].
- [Ros] V. Rosta, On a Ramsey Type Problem of J.A. Bondy and P. Erdős, I & II, *Journal of Combinatorial Theory*, Series B, **15** (1973) 94-120.
- [-] V. Rosta, see also [KR].
- [-] B.L. Rothschild, see [GRS].
- [RoJa1] C.C. Rousseau and C.J. Jayawardene, The Ramsey Number for a Quadrilateral vs. a Complete Graph on Six Vertices, *Congressus Numerantium*, **123** (1997) 97-108.
- [RoJa2] C.C. Rousseau and C.J. Jayawardene, Harary's Problem for $K_{2,k}$, *unpublished manuscript*, (1999).
- [RS1] C.C. Rousseau and J. Sheehan, On Ramsey Numbers for Books, *Journal of Graph Theory*, **2** (1978) 77-87.
- [RS2] C.C. Rousseau and J. Sheehan, A Class of Ramsey Problems Involving Trees, *Journal of the London Mathematical Society* (2), **18** (1978) 392-396.
- [-] C.C. Rousseau, see also [BJYHRZ, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, CLRZ, CRSPS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FRS1, FRS2, FRS3, FRS4, FRS5, FRS6, FRS7, FRS8, FSR, JR1, JR2, JR3, JR4, JR5, LR1, LR2, LR3, LR4, LRS, LRZ].
- [-] P. Rowlinson, see [YR1, YR2, YR3].
- [-] A. Ruciński, see [GRR1, GRR2].
- [San] A. Sánchez-Flores, An Improved Bound for Ramsey Number $N(3,3,3,3;2)$, *Discrete Mathematics*, **140** (1995) 281-286.
- [-] I. Sato, see [MS].
- [-] R.H. Schelp, see [BSS, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRS5, BEFRSGJ, BEFS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FLPS, FRS1, FRS2, FRS3, FRS4,

- FRS5, FS1, FS2, FS3, FS4, FSR, FSS1, ChenS].
- [-] J. Schönheim, see [BS].
- [SchSch1]*A. Schelten and I. Schiermeyer, Ramsey Numbers $r(K_3, G)$ for Connected Graphs G of Order Seven, *Discrete Applied Mathematics*, **79** (1997) 189-200.
- [SchSch2] A. Schelten and I. Schiermeyer, Ramsey Numbers $r(K_3, G)$ for $G \cong K_7 - 2P_2$ and $G \cong K_7 - 3P_2$, *Discrete Mathematics*, **191** (1998) 191-196.
- [-] A. Schelten, see also [FSS2].
- [Schi1] I. Schiermeyer, All Cycle-Complete Graph Ramsey Numbers $r(C_m, K_6)$, *preprint*, (2001).
- [Schi2] I. Schiermeyer, *personal communication*, (2001).
- [-] I. Schiermeyer, see also [FSS2, SchSch1, SchSch2].
- [Schu] C. -U. Schulte, Ramsey-Zahlen für Bäume und Kreise, *Ph.D. thesis*, Heinrich-Heine-Universität Düsseldorf, (1992).
- [-] S. Schuster, see[CS].
- [-] A. Schwenk, see [ChvS].
- [Scob] M.W. Scobee, On the Ramsey Number $R(m_1P_3, m_2P_3, m_3P_3)$ and Related Results, ..., *MA thesis*, University of Louisville (1993).
- [-] R.J. Segegin, see [LorSe].
- [Sha] A. Shastri, Lower Bounds for Bi-Colored Quaternary Ramsey Numbers, *Discrete Mathematics*, **84** (1990) 213-216.
- [She1]* J.B. Shearer, Lower Bounds for Small Diagonal Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **42** (1986) 302-304.
- [She2] J.B. Shearer, A Note on the Independence Number of Triangle-free Graphs II, *Journal of Combinatorial Theory, Series B*, **53** (1991) 300-307.
- [-] J. Sheehan, see [CRSPS, CEHMS, FRS6, FRS7, FRS8, RS1, RS2].
- [-] Shen Yun-Qiu, see [LSS].
- [Shi1] Shi Ling Sheng, Cube Ramsey Numbers Are Polynomial, *Random Structures & Algorithms*, **19** (2001) 99--101.
- [Shi2] Shi Ling Sheng, Upper Bounds for Ramsey Numbers, *preprint*, (2002).
- [ShZ1] Shi Ling Sheng and Zhang Ke Min, An Upper Bound Formula for Ramsey Numbers, *preprint*, (2001).
- [ShZ2] Shi Ling Sheng and Zhang Ke Min, A Sequence of Formulas for Ramsey Numbers, *preprint*, (2001).
- [Sid1] A.F. Sidorenko, On Turán Numbers $T(n, 5, 4)$ and Number of Monochromatic 4-cliques in 2-colored 3-graphs (in Russian), *Voprosy Kibernetiki*, **64** (1980) 117-124.
- [Sid2] A.F. Sidorenko, An Upper Bound on the Ramsey Number $R(K_3, G)$ Depending Only on the Size of the Graph G , *Journal of Graph Theory*, **15** (1991) 15-17.
- [Sid3] A.F. Sidorenko, The Ramsey Number of an N -Edge Graph Versus Triangle Is at Most $2N + 1$, *Journal of Combinatorial Theory, Series B*, **58** (1993) 185-196.
- [-] M. Simonovits, see [BSS, FSS1, FS].
- [-] M.J. Smuga-Otto, see [AS].
- [Sob] A. Sobczyk, Euclidian Simplices and the Ramsey Number $R(4, 4; 3)$, *Technical Report #10*, Clemson University (1967).
- [-] W. Solomon, see [LorSo].
- [-] L. Soltés, see [LRS].
- [Song1] Song En Min, Study of Some Ramsey Numbers (in Chinese), a note (announcement of results without proofs), *Mathematica Applicata*, **4(2)** (1991) 6.

- [Song2] Song En Min, New Lower Bound Formulas for the Ramsey Numbers $N(k, k, \dots, k; 2)$ (in Chinese), *Mathematica Applicata*, **6** (1993) suppl., 113-116.
- [Song3] Song En Min, An Investigation of Properties of Ramsey Numbers (in Chinese), *Mathematica Applicata*, **7** (1994) 216-221.
- [Song4] Song En Min, Properties and New Lower Bounds of the Ramsey Numbers $R(p, q; 4)$ (in Chinese), *Journal of Huazhong University of Science and Technology*, **23** (1995) suppl. II, 1-4.
- [SYL] Song En Min, Ye Weiguo and Liu Yanwu, New Lower Bounds for Ramsey Number $R(p, q; 4)$, *Discrete Mathematics*, **145** (1995) 343-346.
- [-] Song En Min, see also [HuSo].
- [Spe1] J.H. Spencer, Asymptotic Lower Bounds for Ramsey Functions, *Discrete Mathematics*, **20** (1977) 69-76.
- [-] J.H. Spencer, see also [BES, GRS].
- [Spe2]* T. Spencer, University of Nebraska at Omaha, *personal communication* (1993), and, Upper Bounds for Ramsey Numbers via Linear Programming, *preprint*, (1994).
- [Stahl] S. Stahl, On the Ramsey Number $R(F, K_m)$ where F is a Forest, *Canadian Journal of Mathematics*, **27** (1975) 585-589.
- [-] R.G. Stanton, see [KS].
- [Stat] W. Staton, Some Ramsey-type Numbers and the Independence Ratio, *Transactions of the American Mathematical Society*, **256** (1979) 353-370.
- [-] A. Steger, see [McS].
- [Stev] S. Stevens, Ramsey Numbers for Stars Versus Complete Multipartite Graphs, *Congressus Numerantium*, **73** (1990) 63-71.
- [-] M.J. Stewart, see [CRSPS].
- [Stone] J.C. Stone, Utilizing a Cancellation Algorithm to Improve the Bounds of $R(5, 5)$, (1996), <http://oas.okstate.edu/ojas/jstone.htm>. This paper claims incorrectly that $R(5, 5) = 50$.
- [SL]* Su Wenlong and Luo Haipeng, Prime Order Cyclic Graphs and New Lower Bounds for Three Classical Ramsey Numbers $R(4, n)$ (in Chinese), *Journal of Mathematical Study*, **31**, 4 (1998) 442-446.
- [SLLL]* Su Wenlong, Luo Haipeng, Li Guiqing and Li Qiao, Lower Bounds of Ramsey Numbers Based on Cubic Residues, *Discrete Mathematics*, **250** (2002) 197-209.
- [SLL1]* Su Wenlong, Luo Haipeng and Li Guiqing, Two Lower Bounds of Classical 2-color Ramsey Numbers $R(3, q)$ (in Chinese), *Journal of Guangxi University for Nationalities*, **5**, 1 (1999) 1-4.
- [SLL2]* Su Wenlong, Luo Haipeng and Li Qiao, New Lower Bounds of Classical Ramsey Numbers $R(4, 12)$, $R(5, 11)$ and $R(5, 12)$, *Chinese Science Bulletin*, **43**, 6 (1998) 528.
- [SLL3]* Su Wenlong, Luo Haipeng and Li Qiao, New Lower Bounds for Seven Classical Ramsey Numbers $R(k, l)$ (in Chinese), *Journal of Systems Science and Mathematical Sciences*, **20**, 1 (2000) 55-57.
- [SLZL]* Su Wenlong, Luo Haipeng, Zhang Zhengyou and Li Guiqing, New Lower Bounds of Fifteen Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **19** (1999) 91-99.
- [-] Su Wenlong, see also [LSL, LSS, LSZL, ZSL].
- [Sud] B. Sudakov, A Note on Cycle-Complete Graph Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org/>, #N1, **9** (2002), 4 pages.
- [-] A. Sudan, see [GGS].
- [SuBa1] Surahmat and E.T. Baskoro, On the Ramsey Number of a Path or a Star versus W_4 or W_5 , *Proceedings of the 12-th Australasian Workshop on Combinatorial Algorithms*, Bandung, Indonesia, July 14-17 (2001) 174-179.
- [SuBa2] Surahmat and E.T. Baskoro, The Ramsey Number of Linear Forest versus Wheel, paper presented at the *13-th Australasian Workshop on Combinatorial Algorithms*, Fraser Island, Queensland, Australia, July 7-10, 2002.

- [SuBB1] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Star-like Trees versus Large Odd Wheels, *preprint*, (2002).
- [SuBB2] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Cycles versus Small Wheels, *preprint*, (2002).
- [-] Surahmat, see also [BaSu, BSNM].
- [SwPr] C.J. Swanepoel and L.M. Pretorius, Upper Bounds for a Ramsey Theorem for Trees, *Graphs and Combinatorics*, **10** (1994) 377-382.
- [-] M.M. Sweet, see [FreSw].
- [-] E. Szemerédi, see [AKS, CRST].
- [-] R. Thomas, see [RöTh].
- [Tho] A. Thomason, An Upper Bound for Some Ramsey Numbers, *Journal of Graph Theory*, **12** (1988) 509-517.
- [-] C.A. Tovey, see [CET].
- [Tr] Trivial results.
- [-] W.T. Trotter Jr., see [CRST].
- [Tse]* Kung-Kuen Tse, On the Ramsey Number of the Quadrilateral versus the Book and the Wheel, *to appear in the Australasian Journal of Combinatorics*, (2003).
- [-] Kung-Kuen Tse, see also [RT].
- [-] Z. Tuza, see [GT].
- [-] L. Volkmann, see [GV].
- [Walk] K. Walker, Dichromatic Graphs and Ramsey Numbers, *Journal of Combinatorial Theory*, **5** (1968) 238-243.
- [Wall] W.D. Wallis, On a Ramsey Number for Paths, *Journal of Combinatorics, Information & System Sciences*, **6** (1981) 295-296.
- [Wan] Wan Honghui, Upper Bounds for Ramsey Numbers $R(3, 3, \dots, 3)$ and Schur Numbers, *Journal of Graph Theory*, **26** (1997) 119-122.
- [-] Wang Gongben, see [WW, WWY1, WWY2].
- [WW]* Wang Qingxian and Wang Gongben, New Lower Bounds of Ramsey Numbers $r(3, q)$ (in Chinese), *Acta Scientiarum Naturalium, Universitatis Pekinensis*, **25** (1989) 117-121.
- [WWY1]* Wang Qingxian, Wang Gongben and Yan Shuda, A Search Algorithm And New Lower Bounds for Ramsey Numbers $r(3, q)$, *preprint*, (1994).
- [WWY2]* Wang Qingxian, Wang Gongben and Yan Shuda, The Ramsey Numbers $R(K_3, K_q - e)$ (in Chinese), *Beijing Daxue Xuebao Ziran Kexue Ban*, **34** (1998) 15-20.
- [-] Wang Zhi Jian, see [LiWa].
- [Wh] E.G. Whitehead, The Ramsey Number $N(3, 3, 3, 3; 2)$, *Discrete Mathematics*, **4** (1973) 389-396.
- [-] E.R. Williams, see [AW].
- [-] A. Woldar, see [LaWo1, LaWo2].
- [XZ]* Xie Jiguo and Zhang Xiaoxian, A New Lower Bound for Ramsey Number $r(3, 13)$ (in Chinese), *Journal of Lanzhou Railway Institute*, **12** (1993) 87-89.
- [-] Xie Zheng, see [XX1, XX2].
- [XX1]* Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on the Ramsey Numbers $r(k, l)$, *manuscript*, (2002).
- [XX2] Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on Multicolor Ramsey Numbers, *manuscript*, (2002).
- [-] J. Yackel, see [GY].

- [-] Yan Shuda, see [WWY1, WWY2].
- [YHZ1] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min, The Value of the Ramsey Number $R(C_n, K_4)$ is $3(n-1)+1$ ($n \geq 4$), *Australasian Journal of Combinatorics*, **20** (1999) 205-206.
- [YHZ2] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min, $R(C_6, K_5) = 21$ and $R(C_7, K_5) = 25$, *European Journal of Combinatorics*, **22** (2001) 561-567.
- [-] Yang Jian Sheng, see also [BJYHRZ].
- [YY]** Yang Yuansheng, On the Third Ramsey Numbers of Graphs with Six Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **17** (1995) 199-208.
- [YH]* Yang Yuansheng and G.R.T. Hendry, The Ramsey Number $r(K_1 + C_4, K_5 - e)$, *Journal of Graph Theory*, **19** (1995) 13-15.
- [YR1]** Yang Yuansheng and P. Rowlinson, On the Third Ramsey Numbers of Graphs with Five Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **11** (1992) 213-222.
- [YR2]* Yang Yuansheng and P. Rowlinson, On Graphs without 6-Cycles and Related Ramsey Numbers, *Utilitas Mathematica*, **44** (1993) 192-196.
- [YR3]* Yang Yuansheng and P. Rowlinson, The Third Ramsey Numbers for Graphs with at Most Four Edges, *Discrete Mathematics*, **125** (1994) 399-406.
- [-] Ye Weiguo, see [SYL].
- [Yu1]* Yu Song Nian, A Computer Assisted Number Theoretical Construction of $(3, k)$ -Ramsey Graphs, *Annales Universitatis Scientiarum Budapestinensis, Sect. Comput.*, **10** (1989) 35-44.
- [Yu2]* Yu Song Nian, Maximal Triangle-free Circulant Graphs and the Function $K(c)$ (in Chinese), *Journal of Shanghai University, Natural Science*, **2** (1996) 678-682.
- [-] Zang Wenan, see [LRZ].
- [Zeng] Zeng Wei Bin, Ramsey Numbers for Triangles and Graphs of Order Four with No Isolated Vertex, *Journal of Mathematical Research & Exposition*, **6** (1986) 27-32.
- [ZZ] Zhang Ke Min and Zhang Shu Sheng, Some Tree-Stars Ramsey Numbers, *Proceedings of the Second Asian Mathematical Conference 1995*, 287-291, World Sci. Publishing, River Edge, NJ, 1998.
- [-] Zhang Ke Min, see also [BJYHRZ, ChZZ, HZ1, HZ2, MZ, ShZ1, ShZ2, YHZ1, YHZ2].
- [-] Zhang Shu Sheng, see [ZZ].
- [-] Zhang Xiaoxian, see [XZ].
- [-] Zhang Yuming, see [CLRZ].
- [-] Zhang Yunqing, see [ChZZ].
- [ZSL] Zhang Zhengyou, Su Wenlong and Luo Haipeng, Edge Coloration of Complete Graph K_{97} and New Lower Bounds of Four Ramsey Numbers (in Chinese), *Application Research of Computers*, **18**, 6 (2001) 29-31, 57.
- [-] Zhang Zhengyou, see also [LSZL, SLZL].
- [Zhou1] Zhou Huai Lu, Some Ramsey Numbers for Graphs with Cycles (in Chinese), *Mathematica Applicata*, **6** (1993) 218.
- [Zhou2] Zhou Huai Lu, The Ramsey Number of an Odd Cycle with Respect to a Wheel (in Chinese), *Journal of Mathematics, Shuxue Zazhi* (Wuhan), **15** (1995) 119-120.
- [Zhou3] Zhou Huai Lu, On Book-Wheel Ramsey Number, *Discrete Mathematics*, **224** (2000) 239-249.

Out of 357 references gathered above, 286 appeared in 55 different periodicals, among which most articles were published in: *Discrete Mathematics* 47, *Journal of Combinatorial Theory* (old, Series A and B) 38, *Journal of Graph Theory* 37, and *Ars Combinatoria* 15. The results of 86 references depend on computer algorithms.