A Dynamic Survey of Graph Labeling

Joseph A. Gallian

Department of Mathematics and Statistics
University of Minnesota Duluth
Duluth, Minnesota 55812, U.S.A.

jgallian@d.umn.edu

Submitted: September 18, 1996; Accepted: November 14, 1997
Twenty-ninth edition, November 6, 2026

Mathematics Subject Classifications: 05C78

Abstract

A graph labeling is an assignment of integers to the vertices or edges, or both, subject to certain conditions. Graph labelings were first introduced in the mid-1960s. In the intervening years over 400 graph labeling techniques have been studied in over 4250 papers. Finding out what has been done for any particular kind of labeling and keeping up with new discoveries is difficult because of the sheer number of papers and because many of the papers have appeared in journals that are not widely available. In this survey, I have collected everything I could find on graph labeling. For the convenience of the reader, the survey includes a detailed table of contents and index. This edition has 54 additional pages and 297 new references that are identified with the reference number and the word "new" in the right margin. This edition has links to all the open access articles that I could fine. Many links that were not available in prior editions have been added.

Contents

1 Introduction **5**

2 Graceful and Harmonious Labelings **9**
2.1 Trees . . . 9
2.2 Cycle-Related Graphs . . . 16
2.3 Product Related Graphs . . . 23
2.4 Complete Graphs . . . 26
2.5 Disconnected Graphs . . . 28
2.6 Joins of Graphs . . . 32
2.7 Miscellaneous Results . . . 33
2.8 Summary . . . 44
Table 1: Summary of Graceful Results . . . 46
Table 2: Summary of Harmonious Results . . . 49

3 Variations of Graceful Labelings **53**
3.1 α-labelings . . . 53
Table 3: Summary of Results on α-labelings . . . 68
3.2 γ-Labelings . . . 69
3.3 Graceful-like Labelings . . . 70
Table 4: Summary of Results on Graceful-like labelings . . . 81
3.4 k-graceful Labelings . . . 82
3.5 Skolem-Graceful Labelings . . . 86
3.6 Odd-Graceful Labelings . . . 88
3.7 Cordial Labelings . . . 93
3.8 The Friendly Index–Balance Index . . . 113
3.9 k-equitable Labelings . . . 118
3.10 Odd-even Prime Graceful Labelings . . . 122
3.11 Hamming-graceful Labelings . . . 123
3.12 Hilbert Graceful Labelings . . . 123

4 Variations of Harmonious Labelings **124**
4.1 Sequential and Strongly c-harmonious Labelings . . . 124
4.2 (k, d)-arithmetic Labelings . . . 130
4.3 (k, d)-indexable Labelings . . . 132
4.4 Elegant Labelings . . . 133
4.5 Felicitous Labelings . . . 136
4.6 Odd Harmonious and Even Harmonious Labelings . . . 139

5 Magic-type Labelings **148**
5.1 Magic Labelings . . . 148
Table 5: Summary of Magic Labelings . . . 162
5.2 Edge-magic Total and Super Edge-magic Total Labelings . . . 163

Table 6: Summary of Edge-magic Total Labelings . . . 188
Table 7: Summary of Super Edge-magic Labelings . . . 190
5.3 Vertex-magic Total Labelings . . . 193
Table 8: Summary of Vertex-magic Total Labelings . . . 201
Table 9: Summary of Super Vertex-magic Total Labelings . . . 203
Table 10: Summary of Totally Magic Labelings . . . 203
5.4 H-Magic Labelings . . . 204
5.5 Magic Labelings of Type (a, b, c) . . . 209
Table 11: Summary of Magic Labelings of Type (a, b, c) . . . 211
5.6 Sigma Labelings/1-vertex magic labelings/Distance Magic . . . 212
5.7 Other Types of Magic Labelings . . . 218

6 Antimagic-type Labelings 231
6.1 Antimagic Labelings . . . 231
Table 12: Summary of Antimagic Labelings . . . 248
6.2 (a, d)-Antimagic Labelings . . . 249
Table 13: Summary of (a, d)-Antimagic Labelings . . . 258
6.3 (a, d)-Antimagic Total Labelings . . . 259
Table 14: Summary of (a, d)-Vertex-Antimagic Total and Super (a, d)-Vertex-Antimagic Total Labelings . . . 271
Table 15: Summary of (a, d)-Edge-Antimagic Total Labelings . . . 272
Table 16: Summary of (a, d)-Edge-Antimagic Vertex Labelings . . . 273
Table 17: Summary of (a, d)-Super-Edge-Antimagic Total Labelings . . . 274
6.4 Face Antimagic Labelings and d-antimagic Labeling of Type (1,1,1) . . . 275
Table 18: Summary of Face Antimagic Labelings . . . 279
Table 19: Summary of d-antimagic Labelings of Type (1,1,1) . . . 279
6.5 Product Antimagic Labelings . . . 280

7 Miscellaneous Labelings 282
7.1 Sum Graphs . . . 282
Table 20: Summary of Sum Graph Labelings . . . 293
7.2 Prime and Vertex Prime Labelings . . . 294
Table 21: Summary of Prime Labelings . . . 310
Table 22: Summary of Vertex Prime Labelings . . . 311
7.3 Edge-graceful Labelings . . . 312
Table 23: Summary of Edge-graceful Labelings . . . 322
7.4 Radio Labelings . . . 324
7.5 Product and Divisor Cordial Labelings . . . 331
7.6 Other Cordial Labelings . . . 349
7.7 Edge Product Cordial Labelings . . . 353
7.8 Difference Cordial Labelings . . . 355
7.9 Prime Cordial Labelings . . . 359
7.10 L-Cordial and Binary L Cordial . . . 364

7.11 Mean Edge Cordial . . . 365
7.12 Vector Basis S-Cordial Labeling Graphs . . . 366
7.13 Fuzzy quotient-3 Cordial Labelings . . . 366
7.14 Mean Labelings . . . 367
7.15 Pair Sum and Pair Mean Graphs . . . 399
7.16 Irregular Total Labelings . . . 402
7.17 Geometric Labelings . . . 421
7.18 Strongly Multiplicative Graphs . . . 422
7.19 Line-graceful Labelings . . . 423
7.20 Graceful k-colorings . . . 425
7.21 k-sequential Labelings . . . 425
7.22 IC-colorings . . . 426
7.23 Minimal k-rankings . . . 427
7.24 Set Graceful and Set Sequential Graphs . . . 428
7.25 Vertex Equitable Graphs . . . 430
7.26 Representations of Graphs modulo n . . . 433
7.27 Sequentially Additive Graphs . . . 435
7.28 Difference Graphs . . . 435
7.29 Square Sum Labelings and Square Difference Labelings . . . 436
7.30 Permutation and Combination Graphs . . . 441
7.31 Strongly *-graphs . . . 443
7.32 Triangular Sum Graphs . . . 443
7.33 Divisor Graphs . . . 444
7.34 Other Kinds of Labelings . . . 446

References **467**

Index **836**

1 Introduction

Most graph labeling methods trace their origin to one introduced by Rosa [3203] in 1967, or one given by Graham and Sloane [1327] in 1980. Rosa [3203] called a function f a *β-valuation* of a graph G with q edges if f is an injection from the vertices of G to the set $\{0, 1, \ldots, q\}$ such that, when each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edge labels are distinct. Golomb [1287] subsequently called such labelings *graceful* and this is now the popular term. Alternatively, Buratti, Rinaldi, and Traetta [729] define a graph G with q edges to be graceful if there is an injection f from the vertices of G to the set $\{0, 1, \ldots, q\}$ such that every possible difference of the vertex labels of all the edges is the set $\{1, 2, \ldots, q\}$. Rosa introduced β-valuations as well as a number of other labelings as tools for decomposing the complete graph into isomorphic subgraphs. In particular, β-valuations originated as a means of attacking the conjecture of Ringel [3175] that K_{2n+1} can be decomposed into $2n+1$ subgraphs that are all isomorphic to a given tree with n edges. Independently, Keevash and Staden [1948] in April 2020 and Montgomery, Pokrovskiy, and Sudakov [2547] in January in 2021 proved Ringel's 1963 conjecture that any tree with n edges packs $2n + 1$ times into the complete graph K_{2n+1} for large n. Keevash and Staden used an embedding algorithm in which the various subroutines are analyzed by a wide range of methods, some of which are adaptations of existing methods whereas Montgomery et al. used probabilistic methods. Although an unpublished result of Erdős says that most graphs are not graceful (see [1327]), most graphs that have some sort of regularity of structure are graceful. Sheppard [3504] has shown that there are exactly $q!$ gracefully labeled graphs with q edges. Rosa [3203] has identified essentially three reasons why a graph fails to be graceful: (1) G has "too many vertices" and "not enough edges," (2) G "has too many edges," and (3) G "has the wrong parity." The disjoint union of trees is a case where there are too many vertices for the number of edges. An infinite class of graphs that are not graceful for the second reason is given in [647]. As an example of the third condition Rosa [3203] has shown that if every vertex has even degree and the number of edges is congruent to 1 or 2 (mod 4) then the graph is not graceful. In particular, the cycles C_{4n+1} and C_{4n+2} are not graceful. Knuth [1971] has observed the more general condition that in any graceful labeling of a graph with the number of edges congruent to 1 or 2 (mod 4), the number of vertices with an odd degree and an odd label is always odd. Knuth [1971] proved by way a computer search that all cubic graphs on $4, 6, 8, 10, 12$, or 14 vertices, except $2K_4$ and $3K_4$, which was proved by Kotzig, are graceful. He conjectures that every connected cubic graph is graceful.

It has been known since 1975 [633] that rooted symmetric trees are graceful. However, the proofs that have been presented for this fact are either indirect inductive proofs or algorithmic descriptive proofs showing the many separate steps involved in labelling the vertices. In 2021 Rofa [3191] provided a graceful labeling for any given rooted symmetric tree in the form of a direct algebraic function that algebraically maps each vertex to a unique label. The function is a generalization of the way a path is canonically gracefully labeled as algorithmically described by Rosa in his 1967 classic paper [3203]. Rofa uses his function to show that a class of rooted symmetric trees that contains the class of

binomial trees has weakly an α-labeling (see Section 3.1) and it can provide a concise practical, computational way of producing graceful labelings of large rooted symmetric trees in relatively minimal time.

Acharya [39] proved that every graph can be embedded as an induced subgraph of a graceful graph and a connected graph can be embedded as an induced subgraph of a graceful connected graph. Acharya, Rao, and Arumugam [59] proved: every triangle-free graph can be embedded as an induced subgraph of a triangle-free graceful graph; every planar graph can be embedded as an induced subgraph of a planar graceful graph; and every tree can be embedded as an induced subgraph of a graceful tree. Sethuraman, Ragukumar, and Slater [3440] show that every tree can be embedded in a graceful tree (see also [3439]) and pose a related open problem toward settling the Graceful Tree Conjecture. Rao and Sahoo [3132] proved that every connected graph can be embedded as an induced subgraph of an Eulerian graceful graph thereby answering a question originally posed by Rao and mentioned by Acharya and Arumugum in [45]. As a consequence they deduce that the problems on deciding whether the chromatic number of a graph is less than or equal to k, for $k \geq 3$, and deciding whether the clique number of a graph is greater than or equal to k, for $k \geq 3$ are NP-complete even for Eulerian graceful graphs.

Sethuraman and Ragukumar [3438] provided an algorithm that generates a graceful tree from a given arbitrary tree by adding a sequence of new pendent edges to the given arbitrary tree thereby proving that every tree is a subtree of a graceful tree. They ask the question: If G is a graceful tree and v is any vertex of G of degree 1, is it true that $G - v$ is graceful? If the answer is affirmative, then those additional edges of the input arbitrary tree T introduced for constructing the graceful tree T by their algorithm could be deleted in some order so that the given arbitrary tree T becomes graceful. This would imply that the Graceful Tree Conjecture is true. These results demonstrate that there is no forbidden subgraph characterization of these particular kinds of graceful graphs.

Harmonious graphs naturally arose in the study by Graham and Sloane [1327] of modular versions of additive bases problems stemming from error-correcting codes. They defined a graph G with q edges to be *harmonious* if there is an injection f from the vertices of G to the group of integers modulo q such that when each edge xy is assigned the label $f(x) + f(y) \pmod q$, the resulting edge labels are distinct. When G is a tree, exactly one label may be used on two vertices. They proved that almost all graphs are not harmonious. Analogous to the "parity" necessity condition for graceful graphs, Graham and Sloane proved that if a harmonious graph has an even number of edges q and the degree of every vertex is divisible by 2^k then q is divisible by 2^{k+1}. Thus, for example, a book with seven pages (i.e., the Cartesian product of the complete bipartite graph $K_{1,7}$ and a path of length 1) is not harmonious. Liu and Zhang [2283] have generalized this condition as follows: if a harmonious graph with q edges has degree sequence $d_1, d_2, \ldots, d_p$ then $\gcd(d_1, d_2, \ldots d_p, q)$ divides $q(q-1)/2$. They have also proved that every graph is a subgraph of a harmonious graph. More generally, Sethuraman and Elumalai [3421] have shown that any given set of graphs $G_1, G_2, \ldots, G_t$ can be embedded in a graceful or harmonious graph. Determining whether a graph has a harmonious labeling was shown to be NP-complete by Auparajita, Dulawat, and Rathore in 2001 (see [2035]). As a

consequence of studying a more general problem, Uyesser and Pokrovskiy [3938] proved that all bounded degree trees admit a harmonious labeling. In [759] Catral developed new constructive frameworks for addressing the Graham–Sloane conjecture that all trees admit a harmonious labeling. By introducing inductive graph composition methods and probabilistic labelings, he gave harmonious labelings for broad families of trees including subdivided trees, graph joins, and iterated constructions. Moreover, he proposed refined conjectural boundaries and algorithms for partial labeling in general graphs.

In the early 1980s Bloom and Hsu [667], [668], [636], [669], [747] extended graceful labelings to directed graphs by defining a graceful labeling on a directed graph $D(V, E)$ as a one-to-one map θ from V to $\{0, 1, 2, \ldots, |E|\}$ such that $\theta(y) - \theta(x) \bmod (|E| + 1)$ is distinct for every edge xy in E. Graceful labelings of directed graphs also arose in the characterization of finite neofields by Hsu and Keedwell [1458], [1459]. Graceful labelings of directed graphs was the subject of Marr's 2007 Ph.D. dissertation [2441]. In [2441] and [2442] Marr presents results of graceful labelings of directed paths, stars, wheels, and umbrellas. Hegde and Kumudakshi [1412] use complete mappings to construct graceful labelings of two directed cycles. Siqinbate and Feng [3620] proved that the disjoint union of three copies of a directed cycle of fixed even length is graceful.

Over the past five decades in excess of 3,000 papers have spawned a bewildering array of graph labeling methods. Despite the unabated procession of papers, there are few general results on graph labelings. Indeed, the papers focus on particular classes of graphs and methods, and feature ad hoc arguments. In part because many of the papers have appeared in journals not widely available, frequently the same classes of graphs have been done by several authors and in some cases the same terminology is used for different concepts. In this article, we survey what is known about numerous graph labeling methods. The author requests that he be sent preprints and reprints as well as corrections for inclusion in the updated versions of the survey.

Earlier surveys, restricted to one or two labeling methods, include [629], [663], [1983], [1169], and [1171]. In [3563] Shivarajkumar, Sriraj, and Hegde provided a 2021 survey article on graceful labeling of digraphs. The book edited by Acharya, Arumugam, and Rosa [44] includes a variety of labeling methods that we do not discuss in this survey. In 2002 Eshghi [1050] wrote a 65 page paper providing an introduction to graceful graphs. The relationship between graceful digraphs and a variety of algebraic structures including cyclic difference sets, sequenceable groups, generalized complete mappings, near-complete mappings, and neofields is discussed in [667] and [668]. The connection between graceful labelings and perfect systems of difference sets is given in [632]. The computational complexity of the gracefulness of a graph is not known, but the complexity of finding a harmonious labeling of a graph is in the NP-class [214]. Labeled graphs serve as useful models for a broad range of applications such as: coding theory, x-ray crystallography, radar, astronomy, circuit design, communication network addressing, data base management, secret sharing schemes, cryptology, models for constraint programming over finite domains, ultrasound screens, [664], [665], [3836], [3054], [3640], [3641], [1007] [273], [272], [550], [3628], [2483] and network passwords–see [4137], [3784], [4136], [4138], [3322], [3889], [4321], [917] and [1273] for details. Applications of graph labelings to encryption and decryption

schemes are given in [242], [243], [142], [242], and [3149]. According to Wang, B. Yao, and M. Yao [4140], graph labelings are used for incorporating redundancy in disks, designing drilling machines, creating layouts for circuit boards, and configuring resistor networks. In [1456] Hsieh, Chen, Jiang, Liaw, and Shin use graph labelings in algorithms for image processing schemes for monitoring air quality. Zhang, Ye, Zhang, and Yao [4320] investigated the use of graph colorings and graph labelings for designing topological passwords that resist attacks. Sivakumar, Vidyanandini, Sreedevi, Nayak, and Bhoi [3623] demonstrated how the notion of total edge irregularity strength of complete tripartite graphs can be used in anti-theft networks. Nithya and Anitha [2691] investigated how graph labelings can be applied to the study of computer networks. In [2470] Medini, D'souza, Nayak, and Bhat apply graph labelings and the complements of graphs to create a new type of data-sharing mechanism. In [2394] Maheswari, Obaiys, Jamaludin, and Alwan discussed techniques of coding a message using the Graph Message Jumbled method on star twin graphs with super mean labelings and even felicitous labelings. Naqvi, Ashraf, Ovais, Rasheed, Asif, and Alameri [2651] proposed an application of H-irregularity strength labelings as a mathematical framework to model and optimize hybrid network topologies. Their results illustrate the way theoretical graph labeling and practical network technology interact, offering a novel solution to LAN design problems. [3149] new

Terms and notation not defined below follow that used in [785] and [1169].

2 Graceful and Harmonious Labelings

2.1 Trees

The Ringel-Kotzig conjecture (GTC) that all trees are graceful has been the focus of many papers. Kotzig [1463] has called the effort to prove it a "disease." Among the trees known to be graceful are: caterpillars [3203] (a *caterpillar* is a tree with the property that the removal of its endpoints leaves a path); trees with at most 4 end-vertices [1463], [4329] and [1781]; trees with diameter at most 5 [4329] and [1453]; symmetrical trees (i.e., a rooted tree in which every level contains vertices of the same degree) [633], [2845], [3297]; rooted trees where the roots have odd degree and the lengths of the paths from the root to the leaves differ by at most one and all the internal vertices have the same parity [746]; rooted trees with diameter D where every vertex has even degree except for one root and the leaves in level $\lfloor D/2 \rfloor$ [469]; rooted trees with diameter D where every vertex has even degree except for one root and the leaves, which are in level $\lfloor D/2 \rfloor$ [469]; rooted trees with diameter D where every vertex has even degree except for one root, the vertices in level $\lfloor D/2 \rfloor - 1$, and the leaves which are in level $\lfloor D/2 \rfloor$ [469]; the graph obtained by identifying the endpoints any number of paths of a fixed length except for the case that the length has the form $4r+1$, $r > 1$ and the number of paths is of the form $4m$ with $m > r$ [3337]; regular bamboo trees [3337] (a rooted tree consisting of branches of equal length the endpoints of which are identified with end points of stars of equal size); and olive trees [2789], [16] (a rooted tree consisting of k branches, where the ith branch is a path of length i); Bahls, Lake, and Wertheim [438] proved that spiders for which the lengths of every path from the center to a leaf differ by at most one are graceful. (A *spider* is a tree that has at most one vertex (called the *center*) of degree greater than 2.) Jampachon, Nakprasit, and Poomsa-ard [1568] provide graceful labelings for some classes of spiders. Panpa and Poomsa-ard [2761] showed that all spider graphs with at most four legs of lengths greater than one admit a graceful labeling. Panpa, Imnang, and Wasuanankul [2760] showed that every spider graph with five legs is graceful. In [3170], Rhys proved that every six-legged spider is graceful. Together with the previously known result for spiders with at most five legs, this shows that every spider with at most six legs is graceful. Huamaní, Bravo, and Poma [1462] proved that certain families of spiders are graceful. Jesintha and Sethuraman [1622] proved that all arbitrary fixed generalized banana trees are graceful. Shan and Zhong [3465] showed that spiders with legs $L_1, \ldots, L_s$ ($s \geq 1$) satisfying $|E(L_2)| \geq 2|E(L_1)|+4$ and $|E(L_{i+1})| \geq 2|E(L_i)|+2$ for $i \in 2, \ldots, s-1$ are graceful. They further gave explicit graceful labelings for spiders with one leg of arbitrary length and all others of length at most two such that the center is labeled by 0. This labeling enables the construction of larger graceful spiders by attaching paths. Saengsura and Poomsa-ard [3225] proved certain classes of spider graphs are graceful, focusing on conditions where leg lengths follow specific ratios or sums to generate graceful labelings, and extended their results to more general cases with even and increasing leg lengths. [3170] new [1622] new [3465] new [3225] new

In [2518], [2519], [2752], [2520], and [2515] Panda and Mishra and Panda, Mishra, and

Dash give graceful labelings for some new classes of trees with diameter six. Pradhan and Kumar [2998] proved that all combs $P_n \odot K_1$ with perfect matching are graceful. In [1347] Guruswamy and Varadhan gave a method for combining caterpillars in a specific way such that the resulting tree is graceful. Venkatesh and Balasubramanian [4081] also create graceful trees by recursively merging caterpillars. In 2006 Wilf and Yoshimura [4191] defined an ordering on the set of all rooted trees of a fixed number of vertices that leads to fast ranking and unranking algorithms. As an application to the graceful tree conjecture, they showed how their method can eliminate repeated isomorphism testing. They investigated graphs with at most 10 vertices. In 2022 Brankovic and Reynolds [704] published a survey of various computer search algorithms for finding graceful labeling of trees.

In 2018 Montgomery, Pokrovskiy, and Sudakov [2546] proved that every tree is almost-harmonious. That is, every n-vertex tree has an injective Γ-harmonious labeling for any Abelian group Γ of order $n + o(n)$. In 2022 Gnang [1277] posted a paper with a proof on arXiv of the Graceful Tree Conjecture. See [1278] for a newer proof by Gnang. In 2022 Gnang and Williams [1279] posted a proof on arXiv of the long standing Graham-Sloane conjecture that every tree admits a harmonious labeling.

Motivated by Horton's work in 2003 [1451], in 2010 Fang [1065] used a deterministic back-tracking algorithm to prove that all trees with at most 35 vertices are graceful. In 2011 Fang [1066] used a hybrid algorithm that involved probabilistic backtracking, tabu searching, and constraint programming satisfaction to verify that every tree with at most 31 vertices is harmonious. In [2405] Mahmoudzadeh and Eshghi treat graceful labelings of graphs as an optimization problem and apply an algorithm based on ant colony optimization metaheuristic to different classes of graphs and compare the results with those produced by other methods. In [3808] Suparta and Agus Ariawan provided two methods for expanding graceful trees from certain graceful trees. The edge search algorithm developed by Horton [1451] can solve graceful labeling for unrooted trees. However, it cannot accurately solve graceful labeling for rooted trees. In 2026 [287] Arnold and Vitor gave a flexible edge search algorithm that can quickly find a solution to the graceful labeling problem in rooted trees. In addition, it accurately finds which rooted trees are impossible to solve for. Computational experiments indicate that all such impossible roots are in trees of the same class: the scorpion. [287] new

Aldred, Širáň and Širáň [180] have proved that the number of graceful labelings of P_n grows at least as fast as $(5/3)^n$. They mention that this fact has an application to topological graph theory. One such application was provided by Goddyn, Richter, and Širáň [1281] who used graceful labelings of paths on $2s + 1$ vertices ($s \geq 2$) to obtain 2^{2s} cyclic oriented triangular embeddings of the complete graph on $12s + 7$ vertices. The Aldred, Širáň and Širáň bound was improved by Adamaszek [72] to $(2.37)^n$ with the aid of a computer. Cattell [761] has shown that when finding a graceful labeling of a path one has almost complete freedom to choose a particular label i for any given vertex v. In particular, he shows that the only cases of P_n when this cannot be done are when $n \equiv 3 \pmod 4$ or $n \equiv 1 \pmod{12}$, v is in the smaller of the two partite sets of vertices, and $i = (n-1)/2$. In [4126] Wang enumerated the nonequivalent graceful trees and obtained

a closed formula for the number.

Using an algorithm to run through all $n!$ graceful graphs on $n+1$ vertices Anick [259] proves that the average number of graceful labelings grows super exponentially. He provides a simple criterion to predict which trees have an exceptionally large number of graceful labelings and gives evidence that trees with an exceptionally small number of graceful labelings fall into two already known families of caterpillar graphs. Over the full set of graceful labelings for a given n, Anick shows that the distribution of vertex degrees associated with each label is very close to Poisson, with the exception of labels 0 and n. A graph is said to be *k-ubiquitously graceful* (also called "k-rotatable") if for every vertex there is a graceful labeling which assigns that vertex the label k. He also gives two new families of trees that are not k-ubiquitously graceful and includes questions suggested by his results. Pegg [2819] proved that a graceful graph with edges 0 to m can always be constructed with the nearest integer to $\sqrt{3m+9/4}+E$ vertices, where the excess E is a 0 or 1 value. For $m<51,\ E=0$. https://oeis.org/A326499

In [1051] and [1052] Eshghi and Azimi discuss a programming model for finding graceful labelings of large graphs. The computational results show that the models can easily solve the graceful labeling problems for large graphs. They used this method to verify that all trees with 30, 35, or 40 vertices are graceful. Stanton and Zarnke [3695] and Koh, Rogers, and Tan [1984], [1985], [1988] gave methods for combining graceful trees to yield larger graceful trees. In [4160] Wang, Yang, Hsu, and Cheng generalized the constructions of Stanton and Zarnke and Koh, Rogers, and Tan for building graceful trees from two smaller given graceful trees. Rogers in [3194] and Koh, Tan, and Rogers in [1987] provide recursive constructions to create graceful trees. Burzio and Ferrarese [731] have shown that the graph obtained from any graceful tree by subdividing every edge is also graceful and trees obtained from a graceful tree by replacing each edge with a path of fixed length are graceful.

The binomial tree B_0 consists of a single vertex. The binomial tree B_k consists of two binomial trees B_{k-1} that are linked together: the root of one is the leftmost child of the root of the other. Ragukumara and Sethuraman [3067] proved that all binomial trees are graceful. Sethuraman and Murugan [3433] introduced a new method of combining graceful trees called the recursive attachment method and showed that the recursively attached tree $T_i = T_{i-1} \oplus T^{A_{i-1}}$ is graceful for $i \geq 1$, where the base tree T_0 is a caterpillar and the attachment tree $T^{A_{i-1}}$ is any caterpillar. Here $T_{i-1} \oplus T^{A_{i-1}}$ represents a tree obtained by attaching a copy of $T^{A_{i-1}}$ at each vertex of degree at least two in T_{i-1}, for $i \geq 1$. Sethuraman and Murugan [3436] proved that any acyclic graph can be embedded in an unicyclic graceful graph.

In 1999 Broersma and Hoede [712] proved that an equivalent conjecture for the graceful tree conjecture is that all trees containing a perfect matching are strongly graceful (graceful with an extra condition also called an α-labeling–see Section 3.1). Wang, Yang, Hsu, and Cheng [4160] showed that there exist infinitely many equivalent versions of the graceful tree conjecture (GTC). They verify these equivalent conjectures of the graceful tree conjecture are true for trees of diameter at most 7.

In 1979 Bermond [629] conjectured that lobsters are graceful (a *lobster* is a tree with

the property that the removal of the endpoints leaves a caterpillar). Morgan [2549] has shown that all lobsters with perfect matchings are graceful. Krop [2043] proved that a lobster that has a perfect matching that covers all but one vertex (i.e., that has an almost perfect matching) is graceful. Ghosh [1266] used adjacency matrices to prove that three classes of lobsters are graceful. Broersma and Hoede [712] proved that if T is a tree with a perfect matching M of T such that the tree obtained from T by contracting the edges in M is caterpillar, then T is graceful. Superdock [3811] used this result to prove that all lobsters with a perfect matching are graceful. Mishra, Panda, and Dash [2515] gave a class of graceful lobsters with an even number of branches incident on the central path. They also provided graceful labelings for a family of lobsters in which one end vertex of the central path is attached to an even number of branches and the remaining vertices are attached to the combinations of branches. Mishra and Panda [2513] and [2517], Mishra and Bhattacharjee [2512], and Mishra, Rout, and Nayak [2516] gave graceful labeling for general classes of lobsters by applying component moving transformations on graceful caterpillars. More results on graceful labeling of lobsters are in [2514]. Sathiamoorthy, Natarajan, Ayyaswamy, and Janakiraman [3321] proved that the splitting graph of a caterpillar is graceful.

Mooney [2548] proved several results towards determining which commutative rings have zero-divisor graphs that are graceful or harmonious. He demonstrated infinite classes of rings that are graceful and harmonious, as well as infinite classes of rings that are neither graceful nor harmonious. He further provided the answer to this question for all of the zero-divisor graphs on up to 14 vertices provided by Redmond [3153]

[2548] new

[3153] new

In [2746] Pakpahan, Mursidah, Novitasari, and Sugeng showed that a super caterpillar constructed from several caterpillar graphs of the same size with each caterpillar having uniform pairs is graceful. Simarmata, Sandy, and Sugeng [3592] used an adjacency matrix to generalize the result of Pakpahan et al. by showing that a super caterpillar constructed from several star graphs of different sizes is graceful.

A *Skolem sequence* of order n is a sequence $s_1, s_2, \ldots, s_{2n}$ of $2n$ terms such that, for each $k \in \{1, 2, \ldots, n\}$, there exist exactly two subscripts $i(k)$ and $j(k)$ with $s_{i(k)} = s_{j(k)} = k$ and $|i(k) - j(k)| = k$. A Skolem sequence of order n exists if and only if $n \equiv 0$ or $1 \pmod 4$. Morgan [2550] has used Skolem sequences to construct classes of graceful trees. Morgan and Rees [2551] used Skolem and Hooked-Skolem sequences to generate classes of graceful lobsters.

Mishra and Panigrahi [2521] and [2757] found classes of graceful lobsters of diameter at least five. They show other classes of lobsters are graceful in [2522] and [2523]. In [3424] Sethuraman and Jesintha [3424] explores how one can generate graceful lobsters from a graceful caterpillar while in [3428] and [3429] (see also [1599]) they show how to generate graceful trees from a graceful star. More special cases of Bermond's conjecture have been done by Ng [2694], by Wang, Jin, Lu, and Zhang [4127], Abhyanker [15], and by Mishra and Panigrahi [2522]. Renuka, Balaganesan, Selvaraju [3162] proved spider trees with n legs of even length t and odd $n \geq 3$ and lobsters for which each vertex of the spine is adjacent to a path of length two are harmonious. Selvaraju, Balaganesan, Renuka, and Suresh [3348] showed that the squares of bistars and P_i^n are harmonious.

A tree in which all internal vertices have degrees $r+1$ except one, is called a *full r-ary* tree. A uniform full r-ary tree is a full r-ary tree in which all of its leaves are at the same level. A tree that is obtained from copies of a full r-ary tree by identifying each vertex of a fixed path with each vertex of the tree of degree r is called a *uniform-distant* tree. Suparta and Ariawan [3809] gave methods for constructing graceful classes of caterpillars, lobsters, and uniform trees that generalize results in [2567] and [3923].

Barrientos [514] defines a *y-tree* as a graph obtained from a path by appending an edge to a vertex of a path adjacent to an end point. He proves that graphs obtained from a y-tree T by replacing every edge e_i of T by a copy of K_{2,n_i} in such a way that the ends of e_i are merged with the two independent vertices of K_{2,n_i} after removing the edge e_i from T are graceful.

Sethuraman and Jesintha [3425], [3426], and [3427] (see also [1599]) proved that rooted trees obtained by identifying one of the end vertices adjacent to either of the penultimate vertices of any number of caterpillars having equal diameter at least 3 with the property that all the degrees of internal vertices of all such caterpillars have the same parity are graceful. They also proved that rooted trees obtained by identifying either of the penultimate vertices of any number of caterpillars having equal diameter at least 3 with the property that all the degrees of internal vertices of all such caterpillars have the same parity are graceful. In [3425], [3426], and [3427] (see also [1599] and [1639]) Sethuraman and Jesintha prove that all rooted trees in which every level contains pendent vertices and the degrees of the internal vertices in the same level are equal are graceful. Kanetkar and Sane [1877] show that trees formed by identifying one end vertex of each of six or fewer paths whose lengths determine an arithmetic progression are graceful.

Chen, Lü, and Yeh [795] define a *firecracker* as a graph obtained from the concatenation of stars by linking one leaf from each. They also define a *banana tree* as a graph obtained by connecting a vertex v to one leaf of each of any number of stars (v is not in any of the stars). They proved that firecrackers are graceful and conjecture that banana trees are graceful. Before Sethuraman and Jesintha [3431] and [3430] (see also [1599]) proved that all banana trees and extended banana trees (graphs obtained by joining a vertex to one leaf of each of any number of stars by a path of length of at least two) are graceful, various kinds of banana trees had been shown to be graceful by Bhat-Nayak and Deshmukh [642], by Murugan and Arumugam [2596], [2594] and by Vilfred [4098].

Consider a set of caterpillars, having equal diameter, in which one of the penultimate vertices has arbitrary degree and all the other internal vertices including the other penultimate vertex is of fixed even degree. Jesintha and Sethuraman [1641] call the rooted tree obtained by merging an end-vertex adjacent to the penultimate vertex of fixed even degree of each caterpillar an *arbitrarily fixed generalized banana tree.* They prove that such trees are graceful. From this it follows that all banana trees are graceful and all generalized banana trees are graceful.

Jeba Jesintha and Subashini proved the following graphs are graceful: the cycle of vertex switching of even cycles [1623]; the path union of vertex switching of odd cycles [1626]; the path union of vertex switching of even cycles in increasing order [1627]; the path union of vertex switching of odd cycles [1625] the path union of vertex switching

of even cycles [1630]; the path union of $P_m\theta S_n$ [1632]; and the cycle of $P_m\theta S_n$ [1632]. In [1629] they prove the cycle of caterpillar trees is graceful and as a corollary the cycle of comb graphs, cycle of paths, and the cycle of coconut trees are graceful. In [1631] Jeba Jasintha and Subashini proved the two quadrilateral snake graphs connected by a path, two alternate quadrilateral snake graphs connected by a path, two double quadrilateral snake graphs connected by a path, and two alternate double quadrilateral snake graphs connected by a path. In [1638] Jeba Jesintha, Subashini, and Rashmi Beula proved that the series of isomorphic copies of a star graph connected between two ladder graphs is graceful. In [1635] Jeba Jesintha, Subashini, and Sabu proved that two complete bipartite graphs connected by an arbitrary path of length n are graceful. In [1636] Jeba Jesintha, Subashini, and Sabu proved that the twig diamond graph with pendant edges is graceful. In [1637] Jeba Jesintha, Subashini, and Siddiqa proved that the path union of vertex switching of odd and even cycle graphs alternately is graceful.

Zhenbin [4332] has shown that graphs obtained by starting with any number of identical stars, appending an edge to exactly one edge from each star, then joining the vertices at which the appended edges were attached to a new vertex are graceful. He also shows that graphs obtained by starting with any two stars, appending an edge to exactly one edge from each star, then joining the vertices at which the appended edges were attached to a new vertex are graceful. In [1640] Jesintha and Sethuraman use a method of Hrnciar and Havier [1453] to generate graceful trees from a graceful star with n edges.

Aldred and McKay [179] used a computer to show that all trees with at most 26 vertices are harmonious. That caterpillars are harmonious was shown by Graham and Sloane [1327]. Ramya and Meenakshi [3128] gave graceful labelings, harmonious labelings, and Zumkeller labelings for ladders, banana trees, and firecrackers. Sarita, Sehgal, and Malik [3313] established that for distinct primes p and q, the prime index graph of the cyclic group Z_{p^n}, the finite Abelian group $Z_{p^n} \times Z_q$, and the group $Z_p \times Z_{p^n}$ are graceful.

Vietri [4088] utilized a counting technique that generalizes Rosa's graceful parity condition and provides constraints on possible graceful labelings of certain classes of trees. He expresses doubts about the validity of the graceful tree conjecture. In [4069] Vietri introduced a family of homogeneous polynomials (mod 2), one for every degree, having as many variables as the number of vertices, for any fixed graph; a so-called "graceful polynomial" that vanishes (mod 2) that may be useful for proving that the related graph is non-graceful (the degree 1 case dates back to Rosa's work). He also classified graphs whose graceful polynomials vanish for degrees 2 to 4, thereby obtaining some new non-graceful graphs.

Using a variant of the Matrix Tree Theorem, Whitty [4181] specifies an $n \times n$ matrix of indeterminates whose determinant is a multivariate polynomial that enumerates the gracefully labeled $(n+1)$-vertex trees. Whitty also gives a bijection between gracefully labeled graphs and rook placements on a chessboard on the Möbius strip. In [729] Buratti, Rinaldi, and Traetta use graceful labelings of paths to obtain a result on Hamiltonian cycle systems. Kumar, Mishra, Verma, and Srivastava [2061] gave graceful and alpha labelings for the graphs obtained by joining two isomorphic copies of graceful cycles by a path of length at least one. They also gave an algorithm for joining a path of desired length at a

vertex of graceful cycle.

In [706] Brankovic and Wanless describe applications of graceful and graceful-like labelings of trees to several well known combinatorial problems including complete graph decompositions, the Oberwolfach problem, which asks for a decomposition of K_v into copies of a given 2-regular graph F, and coloring. They also discuss the connection between α-labeling of paths and near transversals in Latin squares and show how spectral graph theory might be used to further the progress on the graceful tree conjecture.

In [730] Burgess, Danziger, and Traetta show that the Oberwolfach problem has a solution whenever F has a sufficiently large cycle which meets a given lower bound and, in addition, has a single-flip automorphism, which is an involutory automorphism acting as a reflection on exactly one of the cycles of F. Furthermore, they prove analogous results for the minimum covering version and the maximum packing version of the problem. They also show a similar result when the edges of K_v have multiplicity 2, but in this case they do not require that F be single-flip. Their approach allows them to explicitly construct solutions to the Oberwolfach Problem with well-behaved automorphisms. Their constructions use graceful labelings of 2-regular graphs with a vertex removed. They show that this class of graphs is graceful as long as the length of the path-component is sufficiently large. A much better lower bound on the length of the path is given for an α-labeling of such graphs to exist.

Arkut, Arkut, and Basak [272] and Basak [550] proposed an efficient method for managing Internet Protocol (IP) networks by using graceful labelings of the nodes of the spanning caterpillars of the autonomous sub-networks to assign labels to the links in the sub-networks. Graceful labelings of trees also have been used in multi protocol label switching (MPLS) routing platforms in IP networks [273], [3625], and [3889].

Despite the efforts of many, the graceful tree conjecture remains open even for trees with maximum degree 3. More specialized results about trees are contained in [629], [663], [1983], [2371], [740], [1780], and [3204]. In [997] Edwards and Howard provide a lengthy survey paper on graceful trees. Robeva [3188] provides an extensive survey of graceful labelings of trees in her 2011 undergraduate honors thesis at Stanford University. Alfalayleh, Brankovic, Giggins, and Islam [181] survey results related to the graceful tree conjecture as of 2004 and conclude with five open problems. Alfalayleh et al.: say "The faith in the [graceful tree] conjecture is so strong that if a tree without a graceful labeling were indeed found, then it probably would not be considered a tree." In his Princeton University senior thesis Superdock [3811] provided an extensive survey of results and techniques about graceful trees. He also obtained some specialized results about the gracefulness of spiders and trees with diameter 6. Arumugam and Bagga [275] discuss computational efforts aimed at verifying the graceful tree conjecture and we survey recent results on generating all graceful labelings of certain families of unicyclic graphs. Sethuraman and Murugan [3435] construct a graceful unicyclic graph G from every graceful tree T with $V(G) = V(T)$ such that the graceful labeling of G is derived from the graceful labeling of T.

2.2 Cycle-Related Graphs

Cycle-related graphs have been a major focus of attention. Rosa [3203] showed that the n-cycle C_n is graceful if and only if $n \equiv 0$ or 3 (mod 4) and Graham and Sloane [1327] proved that C_n is harmonious if and only if n is odd. Wheels $W_n = C_n + K_1$ are both graceful and harmonious – [1150], [1449], and [1327]. As a consequence we have that a subgraph of a graceful (harmonious) graph need not be graceful (harmonious). The *n-cone* (also called the *n-point suspension*; the 1-cone is the wheel; the 2-cone is also called a *double cone* of C_m) $C_m + \overline{K_n}$ has been shown to be graceful when $m \equiv 0$ or 3 (mod 12) by Bhat-Nayak and Selvam [648]. When n is even and m is 2, 6 or 10 (mod 12) $C_m + \overline{K_n}$ violates the parity condition for a graceful graph. Bhat-Nayak and Selvam [648] also prove that the following cones are graceful: $C_4 + \overline{K_n}$, $C_5 + \overline{K_2}$, $C_7 + \overline{K_n}$, $C_9 + \overline{K_2}$, $C_{11} + \overline{K_n}$ and $C_{19} + \overline{K_n}$. The *helm* H_n is the graph obtained from a wheel by attaching a pendent edge at each vertex of the n-cycle. Helms have been shown to be graceful [324] and harmonious [1276], [2296], [2297] (see also [2283], [3410], [2279], [914], [3089]), and [1385].

Koh, Rogers, Teo, and Yap, [1986] define a *web* graph as one obtained by joining the pendent points of a helm to form a cycle and then adding a single pendent edge to each vertex of this outer cycle. They asked whether such graphs are graceful. This was proved by Kang, Liang, Gao, and Yang [1883]. Yang has extended the notion of a web by iterating the process of adding pendent points and joining them to form a cycle and then adding pendent points to the new cycle. In his notation, $W(2, n)$ is the web graph whereas $W(t, n)$ is the *generalized web* with t n-cycles. Yang has shown that $W(3, n)$ and $W(4, n)$ are graceful (see [1883]), Abhyanker and Bhat-Nayak [17] have done $W(5, n)$ and Abhyanker [15] has done $W(t, 5)$ for $5 \leq t \leq 13$. Gnanajothi [1276] has shown that webs with odd cycles are harmonious. Seoud and Youssef [3410] define a *closed helm* as the graph obtained from a helm by joining each pendent vertex to form a cycle and a *flower* as the graph obtained from a helm by joining each pendent vertex to the central vertex of the helm. They prove that closed helms and flowers are harmonious when the cycles are odd. A *gear graph* is obtained from the wheel W_n by adding a vertex between every pair of adjacent vertices of the n-cycle. In 1984 Ma and Feng [2374] proved all gears are graceful while in a Master's thesis in 2006 Chen [796] proved all gears are harmonious. Liu [2296] has shown that if two or more vertices are inserted between every pair of vertices of the n-cycle of the wheel W_n, the resulting graph is graceful. Sethuraman and Sankar [3443] showed that the subdivisions of wheels are graceful for even values of $n \geq 4$. Liu [2294] has also proved that the graph obtained from a gear graph by attaching one or more pendent edges to each vertex between the vertices of the n-cycle is graceful. Pradhan and Kumar [2998] proved that graphs obtained by adding a pendent edge to each pendent vertex of hairy cycle $C_n \odot K_1$ are graceful if $n \equiv 0 \pmod{4m}$. They further provide a rule for determining the missing numbers in the graceful labeling of $C_n \odot K_1$ and of the graph obtained by adding pendent edges to each pendent vertex of $C_n \odot K_1$.

Kumar, Mishra, Kumar, and Kumar [2060] proved the following: $C_n \odot K_1$, $n \equiv 0$ (mod 4) possesses an alpha labeling with the missing number $3n/2$; the one-point union of C_{4n} and a path possesses an alpha labeling with an identifiable missing number; and

the graphs obtained by joining two isomorphic copies of the one-point union of C_{4n} and a path possesses an alpha labeling with identifiable missing numbers. Licona and Tey [2264] showed that for a given integer $m_1 \geq 3$, there exists $m^* > m_1$ such that for $m_2 \geq m^*$, if $m_1 + m_2 = 0, 3 \pmod 4$ (resp., $m_1 + m_2 = 1, 2 \pmod 4$), then there exists a graceful (resp. near-graceful) plane graph consisting of two nested cycles with sizes m_1 and m_2, respectively.

Abhyanker [15] has investigated various unicyclic (that is, graphs with exactly one cycle) graphs. He proved that the unicyclic graphs obtained by identifying one vertex of C_4 with the root of the olive tree with $2n$ branches and identifying an adjacent vertex on C_4 with the end point of the path P_{2n-2} are graceful. He showed that if one attaches any number of pendent edges to these unicyclic graphs at the vertex of C_4 that is adjacent to the root of the olive tree but not adjacent to the end vertex of the attached path, the resulting graphs are graceful. Likewise, Abhyanker proved that the graph obtained by deleting the branch of length 1 from an olive tree with $2n$ branches and identifying the root of the edge deleted tree with a vertex of a cycle of the form C_{2n+3} is graceful. He also has a number of results similar to these. In [436] Bagga, Fotso, Max, and Arumugam investigate the gracefulness of unicyclic graphs with pendent caterpillars at two adjacent vertices of the cycle, and pendent edges at some other vertices of the cycle. In [437] Bagga and Heinz give some properties of graceful graphs obtained by adding pendent edges at each vertex of a cycle. In [2811] Pattanaik, Behera, Bohidar, Bhatta, and Sethi proved the following graphs are graceful: C_3 with two of its vertices attached to caterpillars, graphs obtained by attaching a copy of P_n to each vertex of C_3, and the graphs obtained by identifying the center of a copy of $K_{1,n}$ with each vertex of C_3. [2811] new

Delorme, Maheo, Thuillier, Koh, and Teo [921] and Ma and Feng [2373] showed that any cycle with a chord is graceful. This was first conjectured by Bodendiek, Schumacher, and Wegner [673], who proved various special cases. In 1985 Koh and Yap [1989] generalized this by defining a *cycle with a P_k-chord* to be a cycle with the path P_k joining two nonconsecutive vertices of the cycle. They proved that these graphs are graceful when $k = 3$ and conjectured that all cycles with a P_k-chord are graceful. This was proved for $k \geq 4$ by Punnim and Pabhapote in 1987 [3056]. Chen [801] obtained the same result except for three cases which were then handled by Gao [1346]. In 2005, Sethuraman and Elumalai [3420] defined a *cycle with parallel P_k-chords* as a graph obtained from a cycle C_n $(n \geq 6)$ with consecutive vertices $v_0, v_1, \ldots, v_{n-1}$ by adding disjoint paths P_k, $(k \geq 3)$, between each pair of nonadjacent vertices $v_1, v_{n-1}, v_2, v_{n-2}, \ldots, v_i, v_{n-i}, \ldots, v_\alpha, v_\beta$ where $\alpha = \lfloor n/2 \rfloor - 1$ and $\beta = \lfloor n/2 \rfloor + 2$ if n is odd or $\beta = \lfloor n/2 \rfloor + 1$ if n is even. They proved that every cycle C_n $(n \geq 6)$ with parallel P_k-chords is graceful for $k = 3, 4, 6, 8$, and 10 and they conjecture that the cycle C_n with parallel P_k-chords is graceful for all even k. Xu [4228] proved that all cycles with a chord are harmonious except for C_6 in the case where the distance in C_6 between the endpoints of the chord is 2. The gracefulness of cycles with consecutive chords has also been investigated. For $3 \leq p \leq n - r$, let $C_n(p, r)$ denote the n-cycle with consecutive vertices $v_1, v_2, \ldots, v_n$ to which the r chords $v_1v_p, v_1v_{p+1}, \ldots, v_1v_{p+r-1}$ have been added. Koh and Punnin [1978] and Koh, Rogers, Teo, and Yap [1986] have handled the cases $r = 2, 3$ and $n - 3$ where n is the length of the

cycle. Goh and Lim [1286] then proved that all remaining cases are graceful. Moreover, Ma [2376] has shown that $C_n(p, n-p)$ is graceful when $p \equiv 0, 3 \pmod 4$ and Ma, Liu, and Liu [2377] have proved other special cases of these graphs are graceful. Ma also proved that if one adds to the graph $C_n(3, n-3)$ any number k_i of paths of length 2 from the vertex v_1 to the vertex v_i for $i = 2, \ldots, n$, the resulting graph is graceful. Chen [801] has shown that apart from four exceptional cases, a graph consisting of three independent paths joining two vertices of a cycle is graceful. This generalizes the result that a cycle plus a chord is graceful. Liu [2293] has shown that the n-cycle with consecutive vertices $v_1, v_2, \ldots, v_n$ to which the chords v_1v_k and v_1v_{k+2} $(2 \leq k \leq n-3)$ are adjoined is graceful.

For the cycle $C_n : v_1v_2v_3 \cdots v_nv_1$ and a cycle with a C_k- chord Venkatesh and Sivagurunathan [4083] let $C_{n,k}$ denote the graph obtained from C_n by adding a cycle C_k of length k between the non-adjacent vertices v_2 and v_n. They define a cycle with a parallel C_k chord as the graph obtained from a cycle C_n by adding a cycle C_k of length k between every pair of non-adjacent vertices $(v_2, v_n), (v_3, v_{n-1}), \ldots, (v_a, v_b)$ where $a = \lfloor \frac{n}{2} \rfloor$, $b = \lfloor \frac{n}{2} \rfloor + 2$, if n is even and $a = \lfloor \frac{n}{2} \rfloor$, $b = \lfloor \frac{n}{2} \rfloor + 3$, if n is odd. They proved that $C_{n,4}$ and $C^+_{n,4}$ are graceful for $n \equiv 0 \pmod 4$ and that $C^+_{n,6}$ is graceful for all odd values of $n \geq 5$.

In [915] Deb and Limaye use the notation $C(n, k)$ to denote the cycle C_n with k cords sharing a common endpoint called the *apex*. For certain choices of n and k there is a unique $C(n, k)$ graph and for other choices there is more than one graph possible. They call these *shell-type* graphs and they call the unique graph $C(n, n-3)$ a *shell*. Notice that the shell $C(n, n-3)$ is the same as the fan $F_{n-1} = P_{n-1} + K_1$. Kuppusamy and Guruswamy [2072] show that the subdivision graph of $K_{2,n}$ is graceful for $n \geq 1$ and the subdivision graph of the shell graph $C(n, n-3)$ is graceful for $n \geq 4$. Deb and Limaye define a *multiple shell* to be a collection of edge disjoint shells that have their apex in common. A multiple shell is said to be *balanced* with width w if every shell has order w w or $w+1$. Deb and Limaye [915] have conjectured that all multiple shells are harmonious, and have shown that the conjecture is true for the balanced double shells and balanced triple shells. Yang, Xu, Xi, and Qiao [4256] proved the conjecture is true for balanced quadruple shells. Liang [2252] proved the conjecture is true when each shell has the same order and the number of copies is odd.

Jeba Jesintha and Hilda [1609] define a *shell-butterfly* graph as a one-point union of two shells of any order with two pendent edges at the apex. They prove that certain shell-butterfly graphs are harmonious. Jeba Jesintha and Ezhilarasi Hilda [1607] proved butterfly graphs with one shell of order m and the other shell of order $2m+1$ are graceful and double shells in which each shell has the same order are graceful. Jeba Jesintha and Hilda [1613] define a *bow graph* as a double shell in which each shell has arbitrary order. A bow graph in which each shell has the same order is called a *uniform bow* graph. They prove that all uniform bow graphs are graceful. Jeba Jesintha and Ezhilarasi Hilda [1615] proved that shell-butterfly graphs are graceful. In [1608] Jeba Jesintha and Hilda prove k copies of $C(4, 1) \cup K_2$, and shell flowers (a double shell with shells of order m and $2m$) are graceful.

In [1386] Haviar and Kurtulík defined a *k-enriched fan graph* kF_n, for integers $k, n \geq 2$, as the graph of size $(k+1)n - 1$ obtained by connecting n copies of the star S_k of order

k to the fan F_n such that one vertex of each copy of the star S_k is identified with one vertex of the main path P_n of F_n. They proved that k-enriched fan graphs are graceful and provided characterizations of the k-enriched fan graphs among all simple graphs via Sheppard's labeling sequences [3504] introduced in the 1970s, as well as via labeling relations and graph chessboards.

Sethuraman and Dhavamani [3417] use $H(n,t)$ to denote the graph obtained from the cycle C_n by adding t consecutive chords incident with a common vertex. If the common vertex is u and v is adjacent to u, then for $k \geq 1,\ n \geq 4$, and $1 \leq t \leq n-3$, Sethuraman and Dhavamani denote by $G(n,t,k)$ the graph obtained by taking the union of k copies of $H(n,k)$ with the edge uv identified. They conjecture that every graph $G(n,t,k)$ is graceful. They prove the conjecture for the case that $t = n-3$.

For $i = 1, 2, \ldots, n$ let $v_{i,1}, v_{i,2}, \ldots, v_{i,2m}$ be the successive vertices of n copies of C_{2m}. Sekar [3337] defines a *chain of cycles* $C_{2m,n}$ as the graph obtained by identifying $v_{i,m}$ and $v_{i+1,m}$ for $i = 1, 2, \ldots, n-1$. He proves that $C_{6,2k}$ and $C_{8,n}$ are graceful for all k and all n. Barrientos [517] proved that all $C_{8,n}$, $C_{12,n}$, and $C_{6,2k}$ are graceful.

Truszczyński [3918] studied unicyclic graphs and proved several classes of such graphs are graceful. Among these are what he calls dragons. A *dragon* is formed by joining the end point of a path to a cycle (Koh, et al. [1986] call these *tadpoles*; Kim and Park [1963] call them *kites*). This work led Truszczyński to conjecture that all unicyclic graphs except C_n, where $n \equiv 1$ or 2 (mod 4), are graceful. Guo [1345] has shown that dragons are graceful when the length of the cycle is congruent to 1 or 2 (mod 4). Lu [2370] uses $C_n^{+(m,t)}$ to denote the graph obtained by identifying one vertex of C_n with one endpoint of m paths each of length t. He proves that $C_n^{+(1,t)}$ (a tadpole) is not harmonious when $a+t$ is odd and $C_n^{+(2m,t)}$ is harmonious when $n = 3$ and when $n = 2k+1$ and $t = k-1, k+1$ or $2k-1$. In his Master's thesis, Doma [967] investigates the gracefulness of various unicyclic graphs where the cycle has up to 9 vertices. Guruswamy and Varadhan [1347] proved that any acyclic graph can be embedded in a unicyclic graceful graph. Because of the immense diversity of unicyclic graphs, a proof of Truszczyński's conjecture seems out of reach in the near future. In [658] Biatch, Baggab, and Arumugam gave a survey of results related to Truszczynski's conjecture on the gracefulness of unicyclic graphs and provided a new class of graceful unicyclic graphs.

Cycles that share a common edge or a vertex have received some attention. Murugan and Arumugan [2595] have shown that books with n pentagonal pages (i.e., n copies of C_5 with an edge in common) are graceful when n is even and not graceful when n is odd. Lu [2370] uses $\Theta(C_m)^n$ to denote the graph made from n copies of C_m that share an edge (an n-page book with m-polygonal pages). He proved $\Theta(C_{2m+1})^{2n+1}$ is harmonious for all m and n; $\Theta(C_{4m+2})^{4n+1}$ and $\Theta(C_{4m})^{4n+3}$ are not harmonious for all m and n. Xu [4228] proved that $\Theta(C_m)^2$ is harmonious except when $m = 3$. ($\Theta(C_m)^2$ is isomorphic to $C_{2(m-1)}$ with a chord "in the middle.") Nurvazly and Sugeng [2728] proved that $\Theta(C_3)^n$ graphs (n copies of C_3 that share an edge) have graceful labelings.

A *kayak paddle* $KP(k,m,l)$ is the graph obtained by joining C_k and C_m by a path of length l. Litersky [2277] proves that kayak paddles have graceful labelings in the following cases: $k \equiv 0 \bmod 4$, $m \equiv 0$ or 3 (mod 4); $k \equiv m \equiv 2$ (mod 4) for $k \geq 3$; and $k \equiv 1$ (mod

4), $m \equiv 3 \pmod 4$. She conjectures that $KP(4k+4, 4m+2, l)$ with $2k < m$ is graceful when $l \leq 2m$ if l is even and when $l \leq 2m+1$ if l is odd; and $KP(10, 10, l)$ is graceful when $l \geq 12$. The cases are open: $KP(4k, 4m+1, l)$; $KP(4k, 4m+2, l)$; $KP(4k+1, 4m+1, l)$; $KP(4k+1, 4m+2, l)$; $KP(4k+2, 4m+3, l)$; $KP(4k+3, 4m+3, l)$.

Let $C_n^{(t)}$ denote the one-point union of t cycles of length n. Bermond, Brouwer, and Germa [630] and Bermond, Kotzig, and Turgeon [632]) proved that $C_3^{(t)}$ (that is, the *friendship graph* or *Dutch t-windmill*) is graceful if and only if $t \equiv 0$ or $1 \pmod 4$ while Graham and Sloane [1327] proved $C_3^{(t)}$ is harmonious if and only if $t \not\equiv 2 \pmod 4$. Kulkarni and Bagga [2054] used an AI-assisted search to determine the graceful-labeling behavior of a French windmill graph and its line graph can differ in either direction. [2054] new

Koh, Rogers, Lee, and Toh [1979] conjecture that $C_n^{(t)}$ is graceful if and only if $nt \equiv 0$ or $3 \pmod 4$. Yang and Lin [4248] have proved the conjecture for the case $n = 5$ and Yang, Xu, Xi, Li, and Haque [4254] did the case $n = 7$. Xu, Yang, Li and Xi [4232] did the case $n = 11$. Xu, Yang, Han and Li [4233] did the case $n = 13$. Qian [3063] verifies this conjecture for the case that $t = 2$ and n is even and Yang, Xu, Xi, and Li [4255] did the case $n = 9$. Figueroa-Centeno, Ichishima, and Muntaner-Batle [1090] have shown that if $m \equiv 0 \pmod 4$ then the one-point union of 2, 3, or 4 copies of C_m admits a special kind of graceful labeling called an α-labeling (see Section 3.1) and if $m \equiv 2 \pmod 4$, then the one-point union of 2 or 4 copies of C_m admits an α-labeling. Bodendiek, Schumacher, and Wegner [679] proved that the one-point union of any two cycles is graceful when the number of edges is congruent to 0 or 3 modulo 4. (The other cases violate the necessary parity condition.) Shee [3496] has proved that $C_4^{(t)}$ is graceful for all t. Seoud and Youssef [3408] have shown that the one-point union of a triangle and C_n is harmonious if and only if $n \equiv 1 \pmod 4$ and that if the one-point union of two cycles is harmonious then the number of edges is divisible by 4. The question of whether this latter condition is sufficient is open. Figueroa-Centeno, Ichishima, and Muntaner-Batle [1090] have shown that if G is harmonious then the one-point union of an odd number of copies of G using the vertex labeled 0 as the shared point is harmonious. Sethuraman and Selvaraju [3449] have shown that for a variety of choices of points, the one-point union of any number of non-isomorphic complete bipartite graphs is graceful. They raise the question of whether this is true for all choices of the common point.

Another class of cycle-related graphs is that of triangular cacti. The *block-cutpoint graph* of a graph G is a bipartite graph in which one partite set consists of the cut vertices of G, and the other has a vertex b_i for each block B_i of G. A *block* of a graph is a maximal connected subgraph that has no cut-vertex. A *triangular cactus* is a connected graph all of whose blocks are triangles. A *triangular snake* is a triangular cactus whose block-cutpoint-graph is a path (a triangular snake is obtained from a path $v_1, v_2, \ldots, v_n$ by joining v_i and v_{i+1} to a new vertex w_i for $i = 1, 2, \ldots, n-1$). Rosa [3205] conjectured that all triangular cacti with $t \equiv 0$ or $1 \pmod 4$ blocks are graceful. (The cases where $t \equiv 2$ or $3 \pmod 4$ fail to be graceful because of the parity condition.) Moulton [2559] proved the conjecture for all triangular snakes. A proof of the general case (i.e., all triangular cacti) seems hopelessly difficult. Liu and Zhang [2283] gave an incorrect proof that triangular snakes with an odd number of triangles are harmonious whereas triangular snakes with $n \equiv 2$

(mod 4) triangles are not harmonious. Xu [4229] subsequently proved that triangular snakes are harmonious if and only if the number of triangles is not congruent to 2 (mod 4).

A *double triangular snake* consists of two triangular snakes that have a common path. That is, a double triangular snake is obtained from a path $v_1, v_2, \ldots, v_n$ by joining v_i and v_{i+1} to a new vertex w_i for $i = 1, 2, \ldots, n-1$ and to a new vertex u_i for $i = 1, 2, \ldots, n-1$. Xi, Yang, and Wang [4224] proved that all double triangular snakes are harmonious.

A *hexagonal snake* is obtained from a path $p_1, p_2, p_3, \ldots, p_n$ by joining p_i, p_{i+1} to new vertices x_i and y_i respectively and adding edges $x_i y_i$ for $i = 1, 2, \ldots, n-1$ and replacing every edge with a 6-cycle; an *alternate hexagonal snake* is obtained from a path $p_1, p_2, p_3, \ldots, p_n$ by joining p_i, p_{i+1} to new vertices x_i and y_i (alternatively) and adding edges $x_i y_i$, where $1 \leq i \leq n-1$ for even n and $1 \leq i \leq n-2$ for odd n and replacing each alternate edge with a 6-cycle; a *double hexagonal snake* is obtained from two hexagonal snakes that share the n-path; a *double alternate hexagonal snake* is obtained from two alternative hexagonal snakes that share the n-path. Pattabiraman, Loganathan, and Rao [2810] provided graceful labelings for double hexagonal snakes, alternate hexagonal snakes, odd alternate hexagonal snakes, and double alternate hexagonal snakes.

For any graph G defining G-snake analogous to triangular snakes, Sekar [3337] has shown that C_n-snakes are graceful when $n \equiv 0 \pmod 4$ ($n \geq 8$) and when $n \equiv 2 \pmod 4$ and the number of C_n is even. Gnanajothi [1276, pp. 31-34] had earlier shown that quadrilateral snakes are graceful. Grace [1325] has proved that K_4-snakes are harmonious. Rosa [3205] has also considered analogously defined quadrilateral and pentagonal cacti and examined small cases. Yu, Lee, and Chin [4294] showed that Q_2-snakes and Q_3-snakes are graceful and, when the number of blocks is greater than 1, Q_2-snakes, Q_3-snakes and Q_4-snakes are harmonious.

Barrientos [508] calls a graph a *kC_n-snake* if it is a connected graph with k blocks whose block-cutpoint graph is a path and each of the k blocks is isomorphic to C_n. (When $n > 3$ and $k > 3$ there is more than one kC_n-snake.) If a kC_n-snake where the path of minimum length that contains all the cut-vertices of the graph has the property that the distance between any two consecutive cut-vertices is $\lfloor n/2 \rfloor$ it is called *linear*. Barrientos proves that kC_4-snakes are graceful and that the linear kC_6-snakes are graceful when k is even. He further proves that kC_8-snakes and kC_{12}-snakes are graceful in the cases where the distances between consecutive vertices of the path of minimum length that contains all the cut-vertices of the graph are all even and that certain cases of kC_{4n}-snakes and kC_{5n}-snakes are graceful (depending on the distances between consecutive vertices of the path of minimum length that contains all the cut-vertices of the graph).

Badr [431] defines a *linear cyclic snake* $(m, k)C_n$ as the graph consisting of k copies of C_n with two non-adjacent vertices in common where every copy has m copies of C_n and the block-cutpoint graph is not a path. He proves that the linear cyclic snakes $(m, k)C_4$-snake and $(m, k)C_8$-snake are graceful and conjectures that all the linear cyclic snakes $(m, k)C_n$-snakes are graceful for $n \equiv 0 \pmod 4$ or $n \equiv 3 \pmod 4$.

Several people have studied cycles with pendent edges attached. Frucht [1150] proved that any cycle with a pendent edge attached at each vertex (i.e., a *crown*) is graceful (see

also [1460]). If G has order n, the *corona of G with H*, $G \odot H$ is the graph obtained by taking one copy of G and n copies of H and joining the ith vertex of G with an edge to every vertex in the ith copy of H. Barrientos [513] also proved: if G is a graceful graph of order m and size $m-1$, then $G \odot nK_1$ and $G+nK_1$ are graceful; if G is a graceful graph of order p and size q with $q > p$, then $(G \cup (q+1-p)K_1) \odot nK_1$ is graceful; and all unicyclic graphs, other than a cycle, for which the deletion of any edge from the cycle results in a caterpillar are graceful. Pattanaik, Behera, Bohidar, and Biswal [2812] investigated the existence of certain kinds of graceful labelings of a class of corona-related graphs under specific conditions. [2812] new

The *rose corona* graph $M(C_n)$ is constructed from C_n with vertices $v_1, v_2, \ldots, v_n$ and n isolated vertices $w_1, w_2, \ldots, w_n$, and then connecting every two vertices v_i, v_{i+1} with w_i for $i = 1, 2, \ldots, n$, where $v_{n+1} = v_1$. In [1938] Khairunnisa, Nurvazly, and Indah gave graceful labelings of the corona product of the rose graphs $M(C_4) \odot \overline{K_r}$ for $r \geq 1$ and $M(C_5) \odot \overline{K_r}$ for $r = 1, 2, 3$. [1938] new

For a given cycle C_n with $n \equiv 0$ or $3 \pmod 4$ and a family of trees $\mathcal{T} = \{T_1, T_2, \ldots, T_n\}$, let u_i and $v_i, 1 \leq i \leq n$, be fixed vertices of C_n and T_i, respectively. Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima [1095] provide two construction methods that generate a graceful labeling of the unicyclic graphs obtained from C_n and $\mathcal{T}$ by amalgamating them at each u_i and v_i. Their results encompass all previously known results for unicyclic graphs whose cycle length is 0 or 3 (mod 4) and considerably extend the known classes of graceful unicyclic graphs. Khairunnisa and Sugeng [1939] let $A_{(m,n)}$ denote the graph obtained from C_m by connecting each two adjacent vertices with P_{n+1}. They prove that the graphs $A_{(3,1)} \odot \overline{K_r}$ are graceful.

In [510] Barrientos proved that helms (graphs obtained from a wheel by attaching one pendent edge to each vertex) are graceful. Grace [1324] showed that an odd cycle with one or more pendent edges at each vertex is harmonious and conjectured that $C_{2n} \odot K_1$, an even cycle with one pendent edge attached at each vertex, is harmonious. This conjecture has been proved by Liu and Zhang [2282], Liu [2296] and [2297], Hegde [1403], Huang [1465], and Bu [715]. Sekar [3337] has shown that the graph $C_m \odot P_n$ obtained by attaching the path P_n to each vertex of C_m is graceful. For any $n \geq 3$ and any t with $1 \leq t \leq n$, let C_n^{+t} denote the class of graphs formed by adding a single pendent edge to t vertices of a cycle of length n. Ropp [3202] proved that for every n and t the class C_n^{+t} contains a graceful graph. Gallian and Ropp [1169] conjectured that for all n and t, all members of C_n^{+t} are graceful. This was proved by Qian [3063] and by Kang, Liang, Gao, and Yang [1883]. Of course, such graphs are just a special case of the aforementioned conjecture of Truszczyński that all unicyclic graphs except C_n for $n \equiv 1$ or $2 \pmod 4$ are graceful. Sekar [3337] proved that the graph obtained by identifying an endpoint of a star with a vertex of a cycle is graceful. Lu [2370] shows that the graph obtained by identifying each vertex of an odd cycle with a vertex disjoint copy of C_{2m+1} is harmonious if and only if m is odd. Sudha [3713] proved that the graphs obtained by starting with two or more copies of C_4 and identifying a vertex of the i^{th} copy with a vertex of the $i+1^{th}$ copy and the graphs obtained by starting with two or more cycles (not necessarily of the same size) and identifying an edge from the i^{th} copy with an edge of the $i+1^{th}$ copy are graceful.

Sudha and Kanniga [3720] proved that the graphs obtained by identifying any vertex of C_m with any vertex of degree 1 of S_n where $n = \lceil (m-1)/2 \rceil$ are graceful.

For a given cycle C_n with $n \equiv 0$ or 3 (mod 4) and a family of trees $\mathcal{T} = \{T_1, T_2, \ldots, T_n\}$, let u_i and $v_i, 1 \leq i \leq n$, be fixed vertices of C_n and T_i, respectively. Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima [1095] provide two construction methods that generate a graceful labeling of the unicyclic graphs obtained from C_n and $\mathcal{T}$ by amalgamating them at each u_i and v_i. Their results encompass all previously known results for unicyclic graphs whose cycle length is 0 or 3 (mod 4) and considerably extend the known classes of graceful unicyclic graphs.

Solairaju and Chithra [3650] defined three classes of graphs obtained by connecting copies of C_4 in various ways. Denote the four consecutive vertices of ith copy of C_4 by $v_{i,1}, v_{i,2}, v_{i,3}, v_{i_4}$. They show that the graphs obtained by identifying $v_{i,4}$ with $v_{i+1,2}$ for $i = 1, 2, \ldots, n-1$ is graceful; the graphs obtained by joining $v_{i,4}$ with $v_{i+1,2}$ for $i = 1, 2, \ldots, n-1$ by an edge is graceful; and the graphs obtained by joining $v_{i,4}$ with $v_{i+1,2}$ for $i = 1, 2, \ldots, n-1$ with a path of length 2 is graceful.

Venkatesh [4077] showed that for positive integers m and n divisible by 4 the graphs obtained by appending a copy of C_n to each vertex of C_m by identifying one vertex of C_n with each vertex of C_m is graceful.

2.3 Product Related Graphs

Graphs that are Cartesian products and related graphs have been the subject of many papers. That planar grids, $P_m \times P_n$ $(m, n \geq 2)$, (many authors use $G \square H$ to denote the Cartesian product of G and H) are graceful was proved by Acharya and Gill [53] in 1978. In 1980, Maheo [2399] clarified the complicated-appearing construction of Acharya and Gill for $P_m \times P_2$ that readily extends to all grids. Liu, T. Zou, Y. Lu [2291] proved $P_m \times P_n \times P_2$ is graceful. In 1980 Graham and Sloane [1327] proved ladders, $P_m \times P_2$, are harmonious when $m > 2$ and in 1992 Jungreis and Reid [1806] showed that the grids $P_m \times P_n$ are harmonious when $(m, n) \neq (2, 2)$. In the answer to exercise 81 of Section 7.2.2.3 of Knuth's The Art of Computer Programming, Volume 4, Fascicle 7, he remarks that $K_m \times P_4$ is probably ungraceful for all sufficiently large m. Shisheng Li [3514] provided a short proof of this, which also covers $K_m \times P_n$ for every $n \geq 2$ and $K_m \times C_n$ for every $n \geq 3$. [3514] new

A few people have looked at graphs obtained from planar grids in various ways. Kathiresan [1918] has shown that graphs obtained from ladders by subdividing each step exactly once are graceful and that graphs obtained by appending an edge to each vertex of a ladder are graceful [1920]. Barrientos and Minion [542] showed that a graceful graph is obtained when every step of a ladder is subdivided an even number of times. In addition, they proved that when each edge of a ladder is subdivided exactly once, the resulting graph is graceful.

Acharya [42] has shown that certain subgraphs of grid graphs are graceful. Lee [2145] defines a *Mongolian tent* as a graph obtained from $P_m \times P_n$, n odd, by adding one extra vertex above the grid and joining every other vertex of the top row of $P_m \times P_n$ to the

new vertex. A *Mongolian village* is a graph formed by successively amalgamating copies of Mongolian tents with the same number of rows so that adjacent tents share a column. Lee proves that Mongolian tents and villages are graceful. A *Young tableau* is a subgraph of $P_m \times P_n$ obtained by retaining the first two rows of $P_m \times P_n$ and deleting vertices from the right hand end of other rows in such a way that the lengths of the successive rows form a nonincreasing sequence. Lee and Ng [2170] have proved that all Young tableaus are graceful. Lee [2145] has also defined a variation of Mongolian tents by adding an extra vertex above the top row of a Young tableau and joining every other vertex of that row to the extra vertex. He proves these graphs are graceful. In [3649] and [3648] Solairaju and Arockiasamy prove that various families of subgraphs of grids $P_m \times P_n$ are graceful. Sudha [3713] proved that certain subgraphs of the grid $P_n \times P_2$ are graceful. Knuth [1971] proved that $K_n \times P_3$ is graceful if and only if $n \leq 6$. Graf [1164] investigated graceful labelings for certain cases of $C_n \odot P_2$. Maharani, Romdhini, Abdurahim, Dwikasari. and Lestari [2385] introduced a new graph that starts with $P_2 \times P_3$ as the first step and recursively adding a new layer of $P_2 \times P_3$ in an upward staircase. [1164] new [2385] new

Prisms are graphs of the form $C_m \times P_n$. These can be viewed as grids on cylinders. In 1977 Bodendiek, Schumacher, and Wegner [673] proved that $C_m \times P_2$ is graceful when $m \equiv 0 \pmod 4$. According to the survey by Bermond [629], Gangopadhyay and Rao Hebbare did the case that m is even about the same time. In a 1979 paper, Frucht [1150] stated without proof that he had done all $C_m \times P_2$. A complete proof of all cases and some related results were given by Frucht and Gallian [1153] in 1988.

In 1992 Jungreis and Reid [1806] proved that all $C_m \times P_n$ are graceful when m and n are even or when $m \equiv 0 \pmod 4$. They also investigated the existence of a stronger form of graceful labeling called an α-labeling (see Section 3.1) for graphs of the form $P_m \times P_n$, $C_m \times P_n$, and $C_m \times C_n$ (see also [1171]).

Yang and Wang have shown that the prisms $C_{4n+2} \times P_{4m+3}$ [4253], $C_n \times P_2$ [4251], and $C_6 \times P_m$ ($m \geq 2$) (see [4253]) are graceful. Singh [3600] proved that $C_3 \times P_n$ is graceful for all n. In their 1980 paper Graham and Sloane [1327] proved that $C_m \times P_n$ is harmonious when n is odd and they used a computer to show $C_4 \times P_2$, the cube, is not harmonious. In 1992 Gallian, Prout, and Winters [1174] proved that $C_m \times P_2$ is harmonious when $m \neq 4$. In 1992, Jungreis and Reid [1806] showed that $C_4 \times P_n$ is harmonious when $n \geq 3$. Huang and Skiena [1467] have shown that $C_m \times P_n$ is graceful for all n when m is even and for all n with $3 \leq n \leq 12$ when m is odd. Abhyanker [15] proved that the graphs obtained from $C_{2m+1} \times P_5$ by adding a pendent edge to each vertex of an outer cycle is graceful.

Torus grids are graphs of the form $C_m \times C_n$ ($m > 2, n > 2$). Very little success has been achieved with these graphs. The graceful parity condition is violated for $C_m \times C_n$ when m and n are odd and the harmonious parity condition [1327, Theorem 11] is violated for $C_m \times C_n$ when $m \equiv 1, 2, 3 \pmod 4$ and n is odd. In 1992 Jungreis and Reid [1806] showed that $C_m \times C_n$ is graceful when $m \equiv 0 \pmod 4$ and n is even. A complete solution to both the graceful and harmonious torus grid problems will most likely involve a large number of cases.

There has been some work done on prism-related graphs. Gallian, Prout, and Winters [1174] proved that all prisms $C_m \times P_2$ with a single vertex deleted or single edge deleted are

graceful and harmonious. The *Möbius ladder* M_n is the graph obtained from the ladder $P_n \times P_2$ by joining the opposite end points of the two copies of P_n. In 1989 Gallian [1168] showed that all Möbius ladders are graceful and all but M_3 are harmonious. Ropp [3202] has examined two classes of prisms with pendent edges attached. He proved that all $C_m \times P_2$ with a single pendent edge at each vertex are graceful and all $C_m \times P_2$ with a single pendent edge at each vertex of one of the cycles are graceful. Ramachandran and Sekar [3107] proved that the graph obtained from the ladder L_n $(P_n \times P_2)$ by identifying one vertex of L_n with any vertex of the star S_m other than the center of S_m is graceful.

For a ladder $P_n \times P_2$ with the two paths with vertex labels $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$, a *triangular belt* is obtained by adding the edges $v_i u_{i+1}$ for $i = 1, 2, \ldots, n-1$. The *alternate triangular belt* is obtained from the ladder by adding the edges $v_{2i+1}u_{2i+2}$ for $i = 1, 2, \ldots, n-1$. For a path $P_n^{'} = u_1, u_2, \ldots, u_n$ and path $P_n^{''} = v_1, v_2, \ldots, v_n$, the graph Z-P_n is obtained from $P_n^{'}$ and $P_n^{''}$ by joining u_i to v_{i+1} for $i = 1, 2, \ldots, n-1$. Valani and Kanani [4038] proved that triangular belt graphs and alternate triangular belt graphs are edge even graceful and that Z-P_n $(n \geq 3)$ is edge even graceful.

Another class of Cartesian products that has been studied is that of books and "stacked" books. The *book* B_m is the graph $S_m \times P_2$ where S_m is the star with m edges. In 1980 Maheo [2399] proved that the books of the form B_{2m} are graceful and conjectured that the books B_{4m+1} were also graceful. (The books B_{4m+3} do not satisfy the graceful parity condition.) This conjecture was verified by Delorme [920] in 1980. Maheo [2399] also proved that $L_n \times P_2$ and $B_{2m} \times P_2$ are graceful. Both Grace [1323] and Reid (see [1173]) have given harmonious labelings for B_{2m}. The books B_{4m+3} do not satisfy the harmonious parity condition [1327, Theorem 11]. Gallian and Jungreis [1173] conjectured that the books B_{4m+1} are harmonious. Gnanajothi [1276] has verified this conjecture by showing B_{4m+1} has an even stronger form of labeling – see Section 4.1. Liang [2248] also proved the conjecture. In 1988 Gallian and Jungreis [1173] defined a *stacked book* as a graph of the form $S_m \times P_n$. They proved that the stacked books of the form $S_{2m} \times P_n$ are graceful and posed the case $S_{2m+1} \times P_n$ as an open question. The *n-cube* $K_2 \times K_2 \times \cdots \times K_2$ (n copies) was shown to be graceful by Kotzig [2015]—see also [2399]. Although Graham and Sloane [1327] used a computer in 1980 to show that the 3-cube is not harmonious (see also [2758]), Ichishima and Oshima [1512] proved that the n-cube Q_n has a stronger form of harmonious labeling called an α-labeling (see Section 3.1) for $n \geq 4$.

In 1986 Reid [3154] found a harmonious labeling for $K_4 \times P_n$. In 2003 Petrie and Smith [2830] investigated graceful labelings of graphs as an exercise in constraint programming satisfaction. They determined that $K_n \times P_2$ is graceful for $n = 3,\ 4$ and 5; $K_4 \times P_3$ is graceful; $K_4 \times C_3$ is graceful; $(C_n \cup C_n) + K_1$ (double wheel) is graceful for $n = 4$ and 5; and $(C_3 \cup C_3) + K_1$ is not graceful. That $K_3 \times K_3$ is not graceful follows from the parity condition given in the introduction. Using significantly better methods in 2010, Smith and Puget obtained the results about graceful labelings for $K_m \times K_1$, $K_m \times P_n$, and $K_m \times C_n$ given in Table 1. Their labeling for $K_5 \times P_2$ and $K_6 \times P_3$ are the unique graceful labelings for those graphs. Redl [3152] proved that $K_4 \times P_n$ is graceful for $n = 1, 2, 3, 4$, and 5 using a constraint programming approach and asked if all graphs of the form $K_4 \times P_n$ are

graceful

Vaidya, Kaneria, Srivastav, and Dani [3983] proved that $P_n \cup P_t \cup (P_r \times P_s)$ where $t < \min\{r, s\}$ and $P_n \cup P_t \cup K_{r,s}$ where $t \leq \min\{r, s\}$ and $r, s \geq 3$ are graceful. Kaneria, Vaidya, Ghodasara, and Srivastav [1873] proved $K_{mn} \cup (P_r \times P_s)$ where $m, n, r, s > 1$; $(P_r \times P_s) \cup P_t$ where $r, s > 1$ and $t \neq 2$; and $K_{mn} \cup (P_r \times P_s) \cup P_t$ where $m, n, r, s > 1$ and $t \neq 2$ are graceful. Xie, Zhao, and Yao [4226] proved that graphs of the form $C_n \odot T$ where T is a graceful tree are graceful.

The *composition* $G_1[G_2]$ is the graph having vertex set $V(G_1) \times V(G_2)$ and edge set $\{(x_1, y_1), (x_2, y_2)|\ x_1x_2 \in E(G_1)$ or $x_1 = x_2$ and $y_1y_2 \in E(G_2)\}$. The *symmetric product* $G_1 \oplus G_2$ of graphs G_1 and G_2 is the graph with vertex set $V(G_1) \times V(G_2)$ and edge set $\{(x_1, y_1), (x_2, y_2)|\ x_1x_2 \in E(G_1)$ or $y_1y_2 \in E(G_2)$ but not both$\}$. Seoud and Youssef [3409] have proved that $P_n \oplus \overline{K_2}$ is graceful when $n > 1$ and $P_n[P_2]$ is harmonious for all n. They also observe that the graphs $C_m \oplus C_n$ and $C_m[C_n]$ violate the parity conditions for graceful and harmonious graphs when m and n are odd.

2.4 Complete Graphs

The questions of the gracefulness and harmoniousness of the complete graphs K_n have been answered. In each case the answer is positive if and only if $n \leq 4$ ([1287], [3593], [1327], [636]). Both Rosa [3203] and Golomb [1287] proved that the complete bipartite graphs $K_{m,n}$ are graceful while Graham and Sloane [1327] showed they are harmonious if and only if m or $n = 1$. Aravamudhan and Murugan [270] have shown that the complete tripartite graph $K_{1,m,n}$ is both graceful and harmonious while Gnanajothi [1276, pp. 25–31] has shown that $K_{1,1,m,n}$ is both graceful and harmonious and $K_{2,m,n}$ is graceful. Some of the same results have been obtained by Seoud and Youssef [3404] who also observed that when m, n, and p are congruent to 2 (mod 4), $K_{m,n,p}$ violates the parity conditions for harmonious graphs. Beutner and Harborth [636] give graceful labelings for $K_{1,m,n}, K_{2,m,n}, K_{1,1,m,n}$ and conjecture that these and $K_{m,n}$ are the only complete multipartite graphs that are graceful. They have verified this conjecture for graphs with up to 23 vertices via computer.

Beutner and Harborth [636] also show that $K_n - e$ (K_n with an edge deleted) is graceful only if $n \leq 5$; any $K_n - 2e$ (K_n with two edges deleted) is graceful only if $n \leq 6$; and any $K_n - 3e$ is graceful only if $n \leq 6$. They also determine all graceful graphs of the form $K_n - G$ where G is $K_{1,a}$ with $a \leq n - 2$ and where G is a matching M_a with $2a \leq n$.

The *windmill* graph $K_n^{(m)}$ $(n > 3)$ consists of m copies of K_n with a vertex in common. A necessary condition for $K_n^{(m)}$ to be graceful is that $n \leq 5$ – see [1986]. Bermond [629] has conjectured that $K_4^{(m)}$ is graceful for all $m \geq 4$. The gracefulness of $K_4^{(m)}$ is equivalent to the existence of a $(12m + 1, 4, 1)$-perfect difference family, which are known to exist for $m \leq 1000$ (see [1467], [10], [4167], and [1235]). Bermond, Kotzig, and Turgeon [632] proved that $K_n^{(m)}$ is not graceful when $n = 4$ and $m = 2$ or 3, and when $m = 2$ and $n = 5$. Stones [3699] proved that $K_5^{(3)}$ and $K_5^{(4)}$ are graceful. In 1982 Hsu [1457] proved that $K_4^{(m)}$ is harmonious for all m. Graham and Sloane [1327] conjectured that $K_n^{(2)}$ is harmonious if and only if $n = 4$. They verified this conjecture for the cases that n is odd

or $n = 6$. Liu [2279] has shown that $K_n^{(2)}$ is not harmonious if $n = 2^a p_1^{a_1} \cdots p_s^{a_s}$ where $a, a_1, \ldots, a_s$ are positive integers and $p_1, \ldots, p_s$ are distinct odd primes and there is a j for which $p_j \equiv 3 \pmod 4$ and a_j is odd. He also shows that $K_n^{(3)}$ is not harmonious when $n \equiv 0 \pmod 4$ and $3n = 4^e(8k + 7)$ or $n \equiv 5 \pmod 8$. Koh, Rogers, Lee, and Toh [1979] and Rajasingh and Pushpam [3090] have shown that $K_{m,n}^{(t)}$, the one-point union of t copies of $K_{m,n}$, is graceful. Sethuraman and Selvaraju [3445] have proved that the one-point union of graphs of the form K_{2,m_i} for $i = 1, 2, \ldots, n$, where the union is taken at a vertex from the partite set with exactly 2 vertices is graceful if at most two of the m_i are equal. They conjecture that the restriction that at most two of the m_i are equal is not necessary. Sudha [3714] proved that two or more complete bipartite graphs having one bipartite vertex set in common are graceful. Mitra and Bhoumik [2525] proved that $K_{2n,2n} \odot K_2$ is graceful.

Koh, Rogers, Lee, and Toh [1986] introduced the notation $B(n, r, m)$ for the graph consisting of m copies of K_n with a K_r in common ($n \geq r$). (We note that Guo [1346] has used the notation $B(n, r, m)$ to denote the graph obtained by joining opposite endpoints of three disjoint paths of lengths n, r and m.) Bermond [629] raised the question: "For which m, n, and r is $B(n, r, m)$ graceful?" Of course, the case $r = 1$ is the same as $K_n^{(m)}$. For $r > 1$, $B(n, r, m)$ is graceful in the following cases: $n = 3$, $r = 2$, $m \geq 1$ [1980]; $n = 4$, $r = 2$, $m \geq 1$ [920]; $n = 4$, $r = 3$, $m \geq 1$ (see [629]), [1980]. Seoud and Youssef [3404] have proved $B(3, 2, m)$ and $B(4, 3, m)$ are harmonious. Liu [2278] has shown that if there is a prime p such that $p \equiv 3 \pmod 4$ and p divides both n and $n - 2$ and the highest power of p that divides n and $n - 2$ is odd, then $B(n, 2, 2)$ is not graceful. Smith and Puget [3641] has shown that up to symmetry, $B(5, 2, 2)$ has a unique graceful labeling; $B(n, 3, 2)$ is not graceful for $n = 6, 7, 8, 9$, and 10; $B(6, 3, 3)$ and $B(7, 3, 3)$ are not graceful; and $B(5, 3, 3)$ is graceful. Combining results of Bermond and Farhi [631] and Smith and Puget [3641] show that $B(n, 2, 2)$ is not graceful for $n > 5$. Lu [2370] obtained the following results: $B(m, 2, 3)$ and $B(m, 3, 3)$ are not harmonious when $m \equiv 1 \pmod 8$; $B(m, 4, 2)$ and $B(m, 5, 2)$ are not harmonious when m satisfies certain special conditions; $B(m, 1, n)$ is not harmonious when $m \equiv 5 \pmod 8$ and $n \equiv 1, 2, 3 \pmod 4$; $B(2m + 1, 2m, 2n + 1) \cong K_{2m} + \overline{K_{2n+1}}$ is not harmonious when $m \equiv 2 \pmod 4$.

More generally, Bermond and Farhi [631] have investigated the class of graphs consisting of m copies of K_n having exactly k copies of K_r in common. They proved such graphs are not graceful for n sufficiently large compared to r. Barrientos [514] proved that the graph obtained by performing the one-point union of any collection of the complete bipartite graphs $K_{m_1,n_1}, K_{m_2,n_2}, \ldots, K_{m_t,n_t}$, where each K_{m_i,n_i} appears at most twice and $\gcd(n_1, n_2, \ldots, n_t) = 1$, is graceful. Savitha and Sasikala [3324] investigated felicitous labelings for the following families of graphs: the join of a star and $K_{m,n}$, generalized wheel graphs with pendant vertices, ande triangular ladders. [3324] new

Sethuraman and Elumalai [3419] have shown that $K_{1,m,n}$ with a pendent edge attached to each vertex is graceful and Jirimutu [1786] has shown that the graph obtained by attaching a pendent edge to every vertex of $K_{m,n}$ is graceful (see also [218]). In [3432] Sethuraman and Kishore determine the graceful graphs that are the union of n copies of K_4 with i edges deleted for $1 \leq i \leq 5$ and with one edge in common. The only cases that

are not graceful are those graphs where the members of the union are C_4 for $n \equiv 3 \pmod 4$ and where the members of the union are P_2. They conjecture that these two cases are the only instances of edge induced subgraphs of the union of n copies of K_4 with one edge in common that are not graceful.

Renuka, Balaganesan, Selvaraju [3162] proved the graphs obtained by joining a vertex of $K_{1,m}$ to a vertex of $K_{1,n}$ by a path are harmonious. Sethuraman and Selvaraju [3451] have shown that union of any number of copies of K_4 with an edge deleted and one edge in common is harmonious.

Clemens, Coulibaly, Garvens, Gonnering, Lucas, and Winters [855] investigated the gracefulness of the one-point and two-point unions of graphs. They show the following graphs are graceful: the one-point union of an end vertex of P_n and K_4; the graph obtained by taking the one-point union of K_4 with one end vertex of P_n and the one-point union of the other end vertex of P_n with the central vertex of $K_{1,r}$; the graph obtained by taking the one-point union of K_4 with one end vertex of P_n and the one-point union of the other end of P_n with a vertex from the partite set of order 2 of $K_{2,r}$; the graph obtained from the graph just described by appending any number of edges to the other vertex of the partite set of order 2; the two-point union of the two vertices of the partite set of order 2 in $K_{2,r}$ and two vertices from K_4; and the graph obtained from the graph just described by appending any number of edges to one of the vertices from the partite set of order 2.

A *Golomb ruler* is a marked straightedge such that the distances between different pairs of marks on the straightedge are distinct. If the set of distances between marks is every positive integer up to and including the length of the ruler, then ruler is called a *perfect Golomb ruler*. Golomb [1287] proved that perfect Golomb rulers exist only for rulers with at most 4 marks. Beavers [606] examines the relationship between Golomb rulers and graceful graphs through a correspondence between rulers and complete graphs. He proves that K_n is graceful if and only if there is a perfect Golomb ruler with n marks and Golomb rulers are equivalent to complete subgraphs of graceful graphs.

2.5 Disconnected Graphs

There have been many papers dealing with graphs that are not connected. For any graph G the graph mG denotes the disjoint union of m copies of G. In 1975 Kotzig [2014] investigated the gracefulness of the graphs rC_s. When $rs \equiv 1$ or $2 \pmod 4$, these graphs violate the gracefulness parity condition. Kotzig proved that when $r = 3$ and $4k > 4$, then rC_{4k} has a stronger form of graceful labeling called α-labeling (see §3.1) whereas when $r \geq 2$ and $s = 3$ or 5, rC_s is not graceful. In 1984 Kotzig [2016] once again investigated the gracefulness of rC_s as well as graphs that are the disjoint union of odd cycles. For graphs of the latter kind he gives several necessary conditions. His paper concludes with an elaborate table that summarizes what was then known about the gracefulness of rC_s. M. He [1388] has shown that graphs of the form $2C_{2m}$ and graphs obtained by connecting two copies of C_{2m} with an edge are graceful. Cahit [743] has shown that rC_s is harmonious when r and s are odd and Seoud, Abdel Maqsoud, and Sheehan [3369] noted that when r or s is even, rC_s is not harmonious. Seoud, Abdel Maqsoud, and Sheehan [3369] proved

that $C_n \cup C_{n+1}$ is harmonious if and only if $n \geq 4$. They conjecture that $C_3 \cup C_{2n}$ is harmonious when $n \geq 3$. This conjecture was proved when Yang, Lu, and Zeng [4249] showed that all graphs of the form $C_{2j+1} \cup C_{2n}$ are harmonious except for $(n, j) = (2, 1)$. As a consequence of their results about super edge-magic labelings (see §5.2) Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima [1094] have that $C_n \cup C_3$ is harmonious if and only if $n \geq 6$ and n is even. Renuka, Balaganesan, Selvaraju [3162] proved that for odd n $C_n \cup P_3$ (see also [2695]) and $C_n \odot \overline{K_m} \cup P_3$ are harmonious. Ng, Alwie, Marjadi, and Sugeng [2695] proved: $C_m \cup P_1$ is harmonious if and only if $m \neq 2 \bmod 4$, $C_m \cup P_2$ $(m \geq 3)$, and they conjectured that $C_m \cup P_3$ is harmonious for all $m \geq 3$. Youssef [4275] has shown that if G is harmonious then mG is harmonious for all odd m.

In 1978 Kotzig and Turgeon [2019] proved that mK_n is graceful if and only if $m = 1$ and $n \leq 4$. Liu and Zhang [2283] have shown that mK_n is not harmonious for n odd and $m \equiv 2 \pmod 4$ and is harmonious for $n = 3$ and m odd. They conjecture that mK_3 is not harmonious when $m \equiv 0 \pmod 4$. Bu and Cao [716] give some sufficient conditions for the gracefulness of graphs of the form $K_{m,n} \cup G$ and they prove that $K_{m,n} \cup P_t$ and the disjoint union of complete bipartite graphs are graceful under some conditions.

Recall a *Skolem sequence* of order n is a sequence $s_1, s_2, \ldots, s_{2n}$ of $2n$ terms such that, for each $k \in \{1, 2, \ldots, n\}$, there exist exactly two subscripts $i(k)$ and $j(k)$ with $s_{i(k)} = s_{j(k)} = k$ and $|i(k) - j(k)| = k$. (A Skolem sequence of order n exists if and only if $n \equiv 0$ or $1 \pmod 4$). Abrham [25] has proved that any graceful 2-regular graph of order $n \equiv 0 \pmod 4$ in which all the component cycles are even or of order $n \equiv 3 \pmod 4$, with exactly one component an odd cycle, can be used to construct a Skolem sequence of order $n + 1$. Also, he showed that certain special Skolem sequences of order n can be used to generate graceful labelings on certain 2-regular graphs.

The graph H_n obtained from the cycle with consecutive vertices $u_1, u_2, \ldots, u_n$ $(n \geq 6)$ by adding the chords $u_2u_n, u_3u_{n-1}, \ldots, u_\alpha u_\beta$, where $\alpha = (n-1)/2$ for all n and $\beta = (n-1)/2+3$ if n is odd or $\beta = n/2+2$ if n is even is called the *cycle with parallel chords*. In Elumalai and Sethuraman [1025] prove the following: for odd $n \geq 5$, $H_n \cup K_{p,q}$ is graceful; for even $n \geq 6$ and $m = (n-2)/2$ or $m = n/2$ $H_n \cup K_{1,m}$ is graceful; for $n \geq 6$, $H_n \cup P_m$ is graceful, where $m = n$ or $n-2$ depending on $n \equiv 1$ or $3 \pmod 4$ or $m \equiv n-1$ or $n-3$ depending on $n \equiv 0$ or $2 \pmod 4$. Elumali and Sethuraman [1027] proved that every n-cycle $(n \geq 6)$ with parallel chords is graceful and every n-cycle with parallel P_k-chords of increasing lengths is graceful for $n = 2 \pmod 4$ with $1 \leq k \leq (\lfloor n/2 \rfloor - 1)$.

In 1985 Frucht and Salinas [1154] conjectured that $C_s \cup P_n$ is graceful if and only if $s + n \geq 6$ and proved the conjecture for the case that $s = 4$. The conjecture was proved by Traetta [3911] in 2012 who used his result to get a complete solution to the well known two-table Oberwolfach problem; that is, given odd number of people and two round tables when is it possible to arrange series of seatings so that each person sits next to each other person exactly once during the series. The t-table Oberwolfach problem $\mathrm{OP}(n_1, n_2, \ldots, n_t)$ asks to arrange a series of meals for an odd number $n = \sum n_i$ of people around t tables of sizes $n_1, n_2, \ldots, n_t$ so that each person sits next to each other exactly once. A solution to $\mathrm{OP}(n_1, n_2, \ldots, n_t)$ is a 2–factorization of K_n whose factors consists of t cycles of lengths $n_1, n_2, \ldots, n_t$. The λ–fold Oberwolfach problem $\mathrm{OP}_\lambda(n_1, n_2, \ldots, n_t)$ refers

to the case where K_n is replaced by λK_n. Traetta used his proof of the Frucht and Salinas conjecture to provide a complete solutions to both $OP(2r+1, 2s)$ and $OP(2r+1, s, s)$, except possibly for $OP(3, s, s)$. He also gave a complete solution of the general λ–fold Oberwolfach problem $OP_\lambda(r, s)$.

Seoud and Youssef [3411] have shown that $K_5 \cup K_{m,n}, K_{m,n} \cup K_{p,q}$ $(m, n, p, q \geq 2), K_{m,n} \cup K_{p,q} \cup K_{r,s}$ $(m, n, p, q, r, s \geq 2,$ $(p, q) \neq (2, 2))$, and $pK_{m,n}$ $(m, n \geq 2, (m, n) \neq (2, 2))$ are graceful. They also prove that $C_4 \cup K_{1,n}$ $(n \neq 2)$ is not graceful whereas Choudum and Kishore [825], [1968] have proved that $C_s \cup K_{1,n}$ is graceful for $s \geq 7$ and $n \geq 1$. Lee, Quach, and Wang [2185] established the gracefulness of $P_s \cup K_{1,n}$. Seoud and Wilson [3403] have shown that $C_3 \cup K_4, C_3 \cup C_3 \cup K_4$, and certain graphs of the form $C_3 \cup P_n$ and $C_3 \cup C_3 \cup P_n$ are not graceful. Abrham and Kotzig [30] proved that $C_p \cup C_q$ is graceful if and only if $p + q \equiv 0$ or 3 (mod 4). Zhou [4337] proved that $K_m \cup K_n$ $(n > 1, m > 1)$ is graceful if and only if $\{m, n\} = \{4, 2\}$ or $\{5, 2\}$. Knuth [1971] used a computer to show that $K_5 \cup K_2$ has a unique graceful labeling up to a complement. (C. Barrientos has called to my attention that $K_1 \cup K_n$ is graceful if and only if $n = 3$ or 4.) Shee [3495] has shown that graphs of the form $P_2 \cup C_{2k+1}$ $(k > 1)$, $P_3 \cup C_{2k+1}$, $P_n \cup C_3$, and $S_n \cup C_{2k+1}$ all satisfy a condition that is a bit weaker than harmonious. Bhat-Nayak and Deshmukh [643] have shown that $C_{4t} \cup K_{1,4t-1}$ and $C_{4t+3} \cup K_{1,4t+2}$ are graceful. Section 3.1 includes numerous families of disconnected graphs that have a stronger form of graceful labelings.

For $m = 2p + 3$ or $2p + 4$, Wang, Liu, and Li [4153] proved the following graphs are graceful: $W_m \cup K_{n,p}$ and $W_{m,2m+1} \cup K_{n,p}$; for $n \geq m$, $W_{m,2m+1} \cup K_{1,n}$; for $m = 2n + 5$, $W_{m,2m+1} \cup (C_3 + \overline{K_n})$. If G_p is a graceful graph with p edges, they proved $W_{2p+3} \cup G_p$ is graceful.

In [3189] Roblee and Abueida proved that a disjoint union of an odd cycle with a graph that consists of a central vertex that is adjacent to an endpoint of a certain number of fixed length paths are harmonious. They extended this result to include specifying that the central vertex in the tree be adjacent to different vertices in each of the t many s-paths. In [31] Abueida and Roblee proposed the following general definition of harmonious labeling that can be applied to the disjoint union of a cycle with multiple trees and showed how to extend these labelings to new disjoint unions. Let $G(V, E)$ be a simple graph with $|V| = n > m = |E| > 0$. They call an onto function $f : V \to Z_m$ a *harmonious* labeling of G provided that if $e_1 = u_1v_1$ and $e_2 = u_2v_2$ are distinct edges, then $f(u_1) + f(v_1) \neq f(u_2) + f(v_2)$ (mod m). If such a function exists, they say G is said to be *harmonious*. Observe that this agrees with the standard usage of the term "harmonious labeling." For $s \geq 3$ odd and $t \geq 2$ even, they say the graph $T = T_{st+1}$ on $st + 1$ vertices consisting of a central vertex adjacent to one endpoint of each of t-many paths each on s vertices, where $1 \leq i \leq t$ is a *starlike tree*. They proved that for $s \geq 3$ odd, $t \geq 2$ even, $r \geq 1$ and F a forest of r-many starlike trees with each central vertex v_i adjacent to an end of k_i-many paths each on s vertices, $C_s \cup F$ is harmonious. For a simple graph $G(V, E)$ with $|V| = n > m = |E| > 0$ Abueida and Roblee define an *equitably harmonious* labeling of G as a harmonious labeling of G such that for each $i, j \in Z_m$, we have $|f^{-1}|$ and $|f^{-1}(j)|$ differ by at most 1. They proved that for $s \geq 3$ odd and $t \geq 2$ even, $C_s \cup T_1 \cup T_2$, where T_1 and T_2 are isomorphic to $T_{st/2+1}$ is equitably harmonious.

Painter, Landers, and Mattox [2745] expanded results of Abueida and Roblee in [31] by exploring the numerical properties when adding values from cosets and subcosets in Z_m. For $s \geq 3$ odd, $t \geq 2$ even, $r \geq 1$ Abueida and Roblee [32] showed that the disjoint union $C_s \cup T_{st+1}$ is harmonious.

In considering graceful labelings of the disjoint unions of two or three stars S_e with e edges Yang and Wang [4252] permitted the vertex labels to range from 0 to $e+1$ and 0 to $e+2$, respectively. With these definitions of graceful, they proved that $S_m \cup S_n$ is graceful if and only if m or n is even and that $S_m \cup S_n \cup S_k$ is graceful if and only if at least one of m, n, or k is even $(m > 1, n > 1, k > 1)$.

Seoud and Youssef [3407] investigated the gracefulness of specific families of the form $G \cup K_{m,n}$. They obtained the following results: $C_3 \cup K_{m,n}$ is graceful if and only if $m \geq 2$ and $n \geq 2$; $C_4 \cup K_{m,n}$ is graceful if and only if $(m,n) \neq (1,1)$; $C_7 \cup K_{m,n}$ and $C_8 \cup K_{m,n}$ are graceful for all m and n; $mK_3 \cup nK_{1,r}$ is not graceful for all m, n and r; $K_i \cup K_{m,n}$ is graceful for $i \leq 4$ and $m \geq 2, n \geq 2$ except for $i = 2$ and $(m,n) = (2,2)$; $K_5 \cup K_{1,n}$ is graceful for all n; $K_6 \cup K_{1,n}$ is graceful if and only if n is not 1 or 3. Youssef [4277] completed the characterization of the graceful graphs of the form $C_n \cup K_{p,q}$ where $n \equiv 0$ or 3 (mod 4) by showing that for $n > 8$ and $n \equiv 0$ or 3 (mod 4), $C_n \cup K_{p,q}$ is graceful for all p and q (see also [512]). Note that when $n \equiv 1$ or 2 (mod 4) certain cases of $C_n \cup K_{p,q}$ violate the parity condition for gracefulness.

For $i = 1, 2, \ldots, m$ let $v_{i,1}, v_{i,2}, v_{i,3}, v_{i,4}$ be a 4-cycle. Yang and Pan [4247] define $F_{k,4}$ to be the graph obtained by identifying $v_{i,3}$ and $v_{i+1,1}$ for $i = 1, 2, \ldots, k-1$. They prove that $F_{m_1,4} \cup F_{m_2,4} \cup \cdots \cup F_{m_n,4}$ is graceful for all n. Pan and Lu [2751] have shown that $(P_2 + \overline{K_n}) \cup K_{1,m}$ and $(P_2 + \overline{K_n}) \cup T_n$ are graceful.

Barrientos [512] has shown the following graphs are graceful: $C_6 \cup K_{1,2n+1}$; $\bigcup_{i=1}^{t} K_{m_i,n_i}$ for $2 \leq m_i < n_i$; and $C_m \cup \bigcup_{i=1}^{t} K_{m_i,n_i}$ for $2 \leq m_i < n_i, m \equiv 0$ or 3 (mod 4), $m \geq 11$. In [1856] Kaneria, Makadia, and Viradia proved that the union of three grid graphs, $\bigcup_{l=1}^{3} (P_{m_l} \times P_{n_l})$, is graceful, the union of finitely many copies of $P_m \times P_n$ is graceful, and provided two new graceful labeling for $P_m \times P_n$.

Wang and Li [4151] use $St(n)$ to denote the star $K_{n,1}$, F_n to denote the fan $P_n \odot K_1$, and $F_{m,n}$ to denote the graph obtained by identifying the vertex of F_m with degree m and the vertex of F_n with degree n. They showed: for all positive integers n and p and $m \geq 2p+2$, $F_m \cup K_{n,p}$ and $F_{m,2m} \cup K_{n,p}$ are graceful; $F_m \cup St(n)$ is graceful; and $F_{m,2m} \cup St(n)$ and $F_{m,2m} \cup G_r$ are graceful. In [4157] Wang, Wang, and Li gave a sufficient condition for the gracefulness of graphs of the form $(P_3 + \overline{K_m}) \cup G$ and $(C_3 + \overline{K_m}) \cup G$. Wei, Wang, and Sun [4174] provided graceful labelings for the unions of some families of wheels related graphs and complete bipartite graphs. They also gave graceful labelings for some graphs of the form $G \cup (C_3 + \overline{K_m}) \cup S_n$ where G is wheel related. In [4295] Yu, Wang, and Song proved the following graphs are graceful: $K_{n,m} \cup (\overline{K_2} + \overline{K_n})$, $K_{n,m} \cup (P_3 + \overline{K_n})$, $K_{n,m} \cup (P_1 + P_{2n+2})$, and $K_{n,m} \cup K_{1,2n}$. They proved the gracefulness of such graphs for a variety of cases when G involves stars and paths. More technical results like these are given in [4159], [4158], and [749].

2.6 Joins of Graphs

A number of classes of graphs that are the join of graphs have been shown to be graceful or harmonious. Koh, Rogers, and Lim [1980] proved $G + H$ is graceful if G is a graceful tree and H is one of $\overline{K_n}$, $P_n \cup K_1$, or a star. Koh, Phoon, and Soh [1976] point out that some versions of this survey prior to 2017 incorrectly stated that Acharya [39] proved that if G is a connected graceful graph, then $G + \overline{K_n}$ is graceful. Redl [3152] showed that the double cone $C_n + \overline{K_2}$ is graceful for $n = 3, 4, 5, 7, 8, 9, 11$. That $C_n + \overline{K_2}$ is not graceful for $n \equiv 2 \pmod 4$ follows that Rosa's parity condition. Redl asks what other double cones are graceful. Bras, Gomes, and Selman [707] showed that double wheels $(C_n \cup C_n) + K_1$ are graceful. Koh, Phoon, and Soh [1976] prove that $K_3 + \overline{K_n}$ is graceful. Reid [3154] proved that $P_n + \overline{K_t}$ is harmonious. Sethuraman and Selvaraju [3450] and [3351] have shown that $P_n + K_2$ is harmonious. They ask whether $S_n + P_n$ or $P_m + P_n$ is harmonious. As stated in an earlier section, wheels are of the form $C_n + K_1$ and are graceful and harmonious. In 2006 Chen [796] proved that multiple wheels $nC_m + K_1$ are harmonious for all $n \not\equiv 0 \pmod 4$. She believes that the $n \not\equiv 0 \pmod 4$ case is also harmonious. Chen also proved that if H has at least one edge, $H + K_1$ is harmonious, and if n is odd, then $nH + K$ is harmonious.

For $n \geq t + 2$ and $t \geq 1$, Koh, Phoon, and Soh [1977] use $P(n, t)$ to denote the graph of order n consisting of a path of length t and $n - (t + 1)$ isolated vertices. For $n \geq 2t + 1$ and $t \geq 1$, they use $I(n, t)$ to denote the disjoint union of tK_2 and $\overline{K_{n-2t}}$. They proved: $\overline{K_p} + P(n, t)$ is graceful for all $p \geq 1, n \geq t + 2$ and $t \geq 1$; $\overline{K_p} + I(n, t)$ is graceful for all $p \geq 1$, $n \geq 2t + 1$ and $t \geq 1$; and for $s, t \in \{1, 2\}$, $P(m, s) + P(n, t)$ is graceful for all $m \geq s + 2$ and $n \geq t + 2$. In [1977] Koh, Phoon, and Soh ask "What can be said about the gracefulness of $C_m + P(n, t)$ where $n \geq t + 2$" and is "Is $P(m, s) + P(n, t)$ always graceful for all $m \geq s + 2$, $n \geq t + 2$, where $s \geq 3$ or $t \geq 3$?" In [1976] they state as problems about graceful graphs: $C_m + P_n$ $(m \geq 3,\ n \geq 3)$; $C_m + C_n$ $(m \geq 3,\ n \geq 3)$ and $K_{1,p} + P(n, t)$ and prove that $C_3 + P(n, t)$ is graceful for all $n \geq t + 2$, where $1 \leq t \leq 3$ and $C_5 + P(n, 1)$ is graceful for all $n \geq 3$.

Shee [3495] has proved $K_{m,n} + K_1$ is harmonious and observed that various cases of $K_{m,n} + K_t$ violate the harmonious parity condition in [1327]. Liu and Zhang [2283] have proved that $K_2 + K_2 + \cdots + K_2$ is harmonious. Youssef [4275] has shown that if G is harmonious then G^m is harmonious for all odd m. He asks the question of whether G is harmonious implies G^m is harmonious when $m \equiv 0 \pmod 4$. Yuan and Zhu [4297] proved that $K_{m,n} + K_2$ is graceful and harmonious. Gnanajothi [1276, pp. 80–127] obtained the following: $C_n + \overline{K_2}$ is harmonious when n is odd and not harmonious when $n \equiv 2, 4, 6 \pmod 8$; $S_n + \overline{K_t}$ is harmonious; and $P_n + \overline{K_t}$ is harmonious. Balakrishnan and Kumar [464] have proved that the join of $\overline{K}_n$ and two disjoint copies of K_2 is harmonious if and only if n is even. Ramírez-Alfonsín [3114] has proved that if G is graceful and $|V(G)| = |E(G)| = e$ and either 1 or e is not a vertex label then $G + \overline{K_t}$ is graceful for all t. Sudha and Kanniga [3717] proved that the graph $P_m + \overline{K_n}$ is graceful.

Seoud and Youssef [3409] have proved: the join of any two stars is graceful and harmonious; the join of any path and any star is graceful; and $C_n + \overline{K_t}$ is harmonious for

every t when n is odd. They also prove that if any edge is added to $K_{m,n}$ the resulting graph is harmonious if m or n is at least 2. Deng [924] has shown certain cases of $C_n + \overline{K_t}$ are harmonious. Seoud and Youssef [3406] proved: the graph obtained by appending any number of edges from the two vertices of degree $n \geq 2$ in $K_{2,n}$ is not harmonious; dragons $D_{m,n}$ (i.e., an endpoint of P_m is appended to C_n) are not harmonious when $m+n$ is odd; and the disjoint union of any dragon and any number of cycles is not harmonious when the resulting graph has odd order. Youssef [4274] has shown that if G is a graceful graph with p vertices and q edges with $p = q+1$, then $G + S_n$ is graceful.

Sethuraman and Elumalai [3423] have proved that for every graph G with p vertices and q edges the graph $G + K_1 + \overline{K_m}$ is graceful when $m \geq 2^p - p - 1 - q$. As a corollary they deduce that every graph is a vertex induced subgraph of a graceful graph. Balakrishnan and Sampathkumar [465] ask for which $m \geq 3$ is the graph $mK_2 + \overline{K_n}$ graceful for all n. Bhat-Nayak and Gokhale [647] have proved that $2K_2 + \overline{K_n}$ is not graceful. Youssef [4274] has shown that $mK_2 + \overline{K_n}$ is graceful if $m \equiv 0$ or $1 \pmod 4$ and that $mK_2 + \overline{K_n}$ is not graceful if n is odd and $m \equiv 2$ or $3 \pmod 4$. Ma [2375] proved that if G is a graceful tree then, $G + K_{1,n}$ is graceful. Amutha and Kathiresan [218] proved that the graph obtained by attaching a pendent edge to each vertex of $2K_2 + \overline{K_n}$ is graceful.

Wu [4213] proves that if G is a graceful graph with n edges and $n+1$ vertices then the join of G and $\overline{K_m}$ and the join of G and any star are graceful. Wei and Zhang [4173] proved that for $n \geq 3$ the disjoint union of $P_1 + P_n$ and a star, the disjoint union of $P_1 + P_n$ and $P_1 + P_{2n}$, and the disjoint union of $P_2 + \overline{K_n}$ and a graceful graph with n edges are graceful. More technical results on disjoint unions and joins are given in [4172], [4173], [4175], [4171], and [749].

2.7 Miscellaneous Results

It is easy to see that P_n^2 is harmonious [1324] while a proof that P_n^2 is graceful has been given by Kang, Liang, Gao, and Yang [1883]. (P_n^k, the kth *power of* P_n, is the graph obtained from P_n by adding edges that join all vertices u and v with $d(u,v) = k$.) This latter result proved a conjecture of Grace [1324]. Seoud, Abdel Maqsoud, and Sheehan [3369] proved that P_n^3 is harmonious and conjecture that P_n^k is not harmonious when $k > 3$. The same conjecture was made by Fu and Wu [1157]. However, Youssef [4284] has proved that P_8^4 is harmonious and P_n^k is harmonious when k is odd. Yuan and Zhu [4297] proved that P_n^{2k} is harmonious when $1 \leq k \leq (n-1)/2$. Selvaraju [3346] has shown that P_n^3 and the graphs obtained by joining the centers of any two stars with the end vertices of the path of length n in P_n^3 are harmonious.

Cahit [743] proves that the graphs obtained by joining p disjoint paths of a fixed length k to single vertex are harmonious when p is odd and when $k = 2$ and p is even. Gnanajothi [1276, p. 50] has shown that the graph that consists of n copies of C_6 that have exactly P_4 in common is graceful if and only if n is even. For a fixed n, let v_{i1}, v_{i2}, v_{i3} and v_{i4} $(1 \leq i \leq n)$ be consecutive vertices of n 4-cycles. Gnanajothi [1276, p. 35] also proves that the graph obtained by joining each v_{i1} to $v_{i+1,3}$ is graceful for all n and the generalized Petersen graph $P(n,k)$ is harmonious in all cases (see also [2190]). Recall

$P(n,k)$, where $n \geq 5$ and $1 \leq k \leq n$, has vertex set $\{a_0, a_1, \ldots, a_{n-1}, b_0, b_1, \ldots, b_{n-1}\}$ and edge set $\{a_i a_{i+1} \mid i = 0, 1, \ldots, n-1\} \cup \{a_i b_i \mid i = 0, 1, \ldots, n-1\} \cup \{b_i b_{i+k} \mid i = 0, 1, \ldots, n-1\}$ where all subscripts are taken modulo n [4170]. The standard Petersen graph is $P(5,2)$.) Redl [3152] has used a constraint programming approach to show that $P(n,k)$ is graceful for $n = 5, 6, 7, 8, 9$, and 10. In [4068] and [4086] Vietri proved that $P(8t,3)$ and $P(8t+4,3)$ are graceful for all t. He conjectured that the graphs $P(8t,3)$ have a stronger form of graceful labeling called an α-labeling (see §3.1). The gracefulness of the generalized Petersen graphs is an open problem. Shao, Deng, Li, and Vese [3487] provide a backtracking algorithm that finds graceful labelings for all generalized Petersen graphs $P(n,k)$ with $n \leq 75$ within several seconds. The algorithm strongly outperforms the standard backtracking algorithm.

Rao and Sahoo [3132] prove that every connected graph can be embedded as an induced subgraph in an Eulerian graceful graph. They also show that for an integer $k \geq 3$, the problems of deciding whether the chromatic number is less than or equal to k and whether the clique number is greater than or equal to k are NP-complete even for Eulerian graceful graphs. Sethuraman, Ragukumar, and Slater [3441] proved that any tree with m edges can be embedded in a graceful tree with less than $4m$ edges and in a graceful planar graph. A conjecture in the graph theory book by Chartrand and Lesniak [785, p. 266] that graceful graphs with arbitrarily large chromatic numbers do not exist was shown to be false by Acharya, Rao, and Arumugam [59] (see also Mahmoody [2402]).

In [520] Barrientos calculates the number of non-isomorphic harmoniously labeled graphs with n edges and at most n vertices. He provides harmonious labelings for certain unicyclic graphs obtained via the corona product and triangular grids obtained via edge amalgamation of copies of C_3 in such a way that each copy of a cycle shares at most two edges with other copies. Moreover, he uses the edge-switching technique on C_{4t} to generate unicyclic graphs with strongly felicitous labelings (see §4.4).

Bača and Youssef [430] investigated the existence of harmonious labelings for the corona graphs of a cycle and a graph G. They proved that if $G+K_1$ is strongly harmonious (that is, a harmonious labeling f for which the edge labels induced by $f(x)+f(y)$ for each edge xy are $1, \ldots, q$. with the 0 label on the vertex of K_1, then $C_n \odot G$ is harmonious for all odd $n \geq 3$. By combining this with existing results they have as corollaries that the following graphs are harmonious: $C_n \odot C_m$ for odd $n \geq 3$ and $m \not\equiv 2 \pmod 3$; $C_n \odot K_{s,t}$ for odd $n \geq 3$; and $C_n \odot K_{1,s,t}$ for odd $n \geq 3$.

Sethuraman and Selvaraju [3444] define a graph H to be a *supersubdivision* of a graph G, if every edge uv of G is replaced by $K_{2,m}$ (m may vary for each edge) by identifying u and v with the two vertices in $K_{2,m}$ that form the partite set with exactly two members. Sethuraman and Selvaraju prove that every supersubdivision of a path is graceful and every cycle has some supersubdivision that is graceful. They conjecture that every supersubdivision of a star is graceful and that paths and stars are the only graphs for which every supersubdivision is graceful. Barrientos [514] disproved this latter conjecture by proving that every supersubdivision of a y-trees is graceful (recall a y-tree is obtained from a path by appending an edge to a vertex of a path adjacent to an end point). Barrientos asks if paths and y-trees are the only graphs for which every supersubdivision is

graceful. This seems unlikely to be the case. The conjecture that every supersubdivision of a star is graceful was proved by Kathiresan and Amutha [1922]. In [3448] Sethuraman and Selvaraju prove that every connected graph has some supersubdivision that is graceful. They pose the question as to whether this result is valid for disconnected graphs. Barrientos and Barrientos [524] answered this question by proving that any disconnected graph has a supersubdivision that admits an α-labeling (see §3.1). They also proved that every supersubdivision of a connected graph admits an α-labeling. Sekar and Ramachandren proved that an arbitrary supersubdivision of disconnected graph is graceful [3338] and supersubdivisions of ladders are graceful [3110]. Sethuraman and Selvaraju also asked if there is any graph other than $K_{2,m}$ that can be used to replace an edge of a connected graph to obtain a supersubdivision that is graceful.

Sethuraman and Selvaraju [3444] call supersubdivision graphs of G where every edge uv of G is replaced by $K_{2,m}$ and m is fixed an *arbitrary supersubdivision* of G. Barrientos and Barrientos [524] answered the question of Sethuraman and Selvaraju by proving that any graph obtained from $K_{2,m}$ by attaching k pendent edges and n pendent edges to the vertices of its 2-element stable set can be used instead of $K_{2,m}$ to produce an arbitrary supersubdivision that admits an α-labeling (a *stable* set S consists of a set of vertices such that there is not an edge v_iv_j for all pairs v_i, v_j in S).

Kathiresan and Sumathi [1932] affirmatively answer the question posed by Sethuraman and Selvaraju in [3444] of whether there are graphs different from paths whose arbitrary supersubdivisions are graceful.

For a graph G Ambili and Singh [213] call the graph G^* a *strong supersubdivision* of G if G^* is obtained from G by replacing every edge e_i of G by a complete bipartite graph K_{r_i,s_i}. A strong supersubdivision G^* of G is said to be an *arbitrary strong supersubdivision* if G^* is obtained from G by replacing every edge e_i of G by a complete bipartite graph K_{r,s_i} (r is fixed and s_i may vary). They proved that arbitrary strong supersubdivisions of paths, cycles, and stars are graceful. They conjecture that every arbitrary strong supersubdivision of a tree is graceful and ask if it is true that for any non-trivial connected graph G, an arbitrary strong supersubdivision of G is graceful?

In [3447] Sethuraman and Selvaraju present an algorithm that permits one to start with any non-trivial connected graph and successively form supersubdivisions that have a strong form of graceful labeling called an α-labeling (see §3.1 for the definition).

Kathiresan [1919] uses the notation $P_{a,b}$ to denote the graph obtained by identifying the end points of b internally disjoint paths each of length a. He conjectures that $P_{a,b}$ is graceful except when a is odd and $b \equiv 2 \pmod 4$ and proves the conjecture for the case that a is even and b is odd. Liang and Zuo [2258] proved that the graph $P_{a,b}$ is graceful when both a and b are even. Daili, Wang and Xie [908] provided an algorithm for finding a graceful labeling of $P_{2r,2}$ and showed that a $P_{2r,2(2k+1)}$ is graceful for all positives r and k. Sekar [3337] has shown that $P_{a,b}$ is graceful when $a \neq 4r+1,\ r > 1, b = 4m$, and $m > r$. Yang (see [4250]) proved that $P_{a,b}$ is graceful when $a = 3, 5, 7$, and 9 and b is odd and when $a = 2, 4, 6$, and 8 and b is even (see [4250]). Yang, Rong, and Xu [4250] proved that $P_{a,b}$ is graceful when $a = 10, 12$, and 14 and b is even. Yan [4238] proved $P_{2r,2m}$ is graceful when r is odd. Yang showed that $P_{2r+1,2m+1}$ and $P_{2r,2m}$ ($r \leq 7$, and $r = 9$) are

graceful (see [3200]). Rong and Xiong [3200] showed that $P_{2r,b}$ is graceful for all positive integers r and b. Kathiresan also shows that the graph obtained by identifying a vertex of K_n with any noncenter vertex of the star with $2^{n-1} - n(n-1)/2$ edges is graceful.

For a family of graphs $G_1(u_1, u_2), G_2(u_2, u_3), \ldots, G_m(u_m, u_{m+1})$ where u_i and u_{i+1} are vertices in G_i Cheng, Yao, Chen, and Zhang [806] define a *graph-block chain* H_m as the graph obtained by identifying u_{i+1} of G_i with u_{i+1} of G_{i+1} for $i = 1, 2, \ldots, m$. They denote this graph by $H_m = G_1(u_1, u_2) \oplus G_2(u_2, u_3) \oplus \cdots \oplus G_m(u_m, u_{m+1})$. The case where each G_i has the form P_{a_i,b_i} they call a *path-block chain*. The vertex u_1 is called the *initial vertex* of H_m. They define a *generalized spider* S_m^* as a graph obtained by starting with an initial vertex u_0 and m path-block graphs and join u_0 with each initial vertex of each of the path-block graphs. Similarly, they define a *generalized caterpillar* T_m^* as a graph obtained by starting with m path-block chains $H_1, H_2, \ldots, H_m$ and a caterpillar T with m isolated vertices $v_1, v_2, \ldots, v_m$ and join each v_i with the initial vertex of each H_i. They prove several classes of path-block chains, generalized spiders, and generalized caterpillars are graceful.

The graph T_n with $3n$ vertices and $6n - 3$ edges is defined as follows. Start with a triangle T_1 with vertices $v_{1,1}, v_{1,2}$ and $v_{1,3}$. Then T_{i+1} consists of T_i together with three new vertices $v_{i+1,1}, v_{i+1,2}, v_{i+1,3}$ and edges $v_{i+1,1}v_{i,2}$, $v_{i+1,1}v_{i,3}$, $v_{i+1,2}v_{i,1}$, $v_{i+1,2}v_{i,3}$, $v_{i+1,3}v_{i,1}$, $v_{i+1,3}v_{i,2}$. Gnanajothi [1276] proved that T_n is graceful if and only if n is odd. Sekar [3337] proved T_n is graceful when n is odd and T_n with a pendent edge attached to the starting triangle is graceful when n is even.

In [611] and [3469] Begam, Palanivelrajan, Gunasekaran, and Hameed give graceful labelings for graphs constructed by combining theta graphs (that is, a collection of edge disjoint paths that have common endpoints) with paths and stars. Khatun and Abu Nayeem [1945] prove that the zero-divisor graph of the commutative ring of integers modulo n is graceful if $n = pq$, $4p$ or $9p$, where p and q are prime numbers. In [861] Constantine and Suwastika provided graceful labelings of zero-divisor graphs $\Gamma(p^2q)$ and $\Gamma(p^3q.)$ [861] new

The *torch* graph O_n is defined by $V(O_n) = \{v_i \mid 1 \leq i \leq n+4\}, E(O_n) = \{v_iv_{n+1} \mid 2 \leq i \leq n-2\} \cup \{v_iv_{n+3} \mid 2 \leq i \leq n-2\} \cup \{v_1v_i \mid 2 \leq i \leq n+4\} \cup \{v_{n-1}v_n, v_nv_{n+2}, v_nv_{n+4}, v_{n+1}v_{n+3}\}$. Manulang and Sugeng [2417] showed that the torch graph is graceful.

For a graph G, the *splitting graph* of G, $S'(G)$, is obtained from G by adding for each vertex v of G a new vertex v' so that v' is adjacent to every vertex that is adjacent to v. Sekar [3337] has shown that $S'(P_n)$ is graceful for all n and $S'(C_n)$ is graceful for $n \equiv 0, 1 \pmod 4$. Vaidya and Shah [4006] proved that the square graph of a bistar, the splitting graph of a bistar, and the splitting graph of a star are graceful graphs.

In [3718] Sudha and Kanniga proved that fans and the splitting graph of a star are graceful. Sudha and Kanniga [3719] proved that the following graphs are graceful: arbitrary supersubdivisions of wheels; combs ($P_n \odot K_1$); double fans ($P_n \odot \overline{K_2}$); $(P_m \cup P_n) \odot K_1$; and graphs obtained by starting with two star graphs S_m and S_n and identifying some of the pendent vertices of each. Sudha and Kanniga [3720] proved that the graphs obtained from $P_n \odot K_1$ by identifying the center of a S_n with the endpoint of a pendent edge at-

tached to the endpoint of P_n are graceful; and the graphs obtained from a fan $P_n \odot K_1$ by deleting a pendent edge attached to an endpoint of P_n are graceful. Sunda [3713] provided some results on graphs obtained by connecting copies of $K_{m,n}$ in certain ways. Sudha and Kanniga [3716] proved that the graphs obtained by joining the vertices of a path to any number isolated points are graceful. They also proved that the arbitrary supersubdivision of all the edges of helms, combs ($P_n \odot K_1$) and ladders ($P_n \times P_2$) with pendent edges at the vertices of degree 2 by a complete bipartite graphs $K_{2,m}$ are graceful.

The *duplication of an edge* $e = uv$ of a graph G is the graph G' obtained from G by adding an edge $e' = u'v'$ such that $N(u) = N(u')$ and $N(v) = N(v')$. The *duplication of a vertex* of a graph G is the graph G' obtained from G by adding a new vertex v' to G such that $N(v') = N(v)$. Kaneria, Vaidya, Ghodasara, and Srivastav [1873] proved the duplication of a vertex of a cycle, the duplication of an edge of an even cycle, and the graph obtained by joining two copies of a fixed cycle by an edge are graceful.

For a graph G and a vertex v of G, a *vertex switching* G_v is the graph obtained from G by removing all edges incident to v and adding edges joining v to every vertex not adjacent to v in G. Boxwala and Vashishta [701] show that the graph obtained by switching an arbitrary vertex of C_n ($n > 3$), the duplication of an arbitrary vertex on the rim of a wheel with an even number of vertices, and the mirror graph of a path are graceful. Jeba Jesintha and Subashini [1642] proved that the path union of vertex switching of even cycles in increasing order is graceful.

The *join sum* of complete bipartite graphs $< K_{m_1,n_1}, \ldots, K_{m_t,n_t} >$ is the graph obtained by starting with $K_{m_1,n_1}, \ldots, K_{m_t,n_t}$ and joining a vertex of each pair K_{m_i,n_i} and $K_{m_{i+1},n_{i+1}}$ to a new vertex v_i where $1 \leq i \leq k-1$. The *path union* of a graph G is the graph obtained by adding an edge from n copies $G_1, G_2, \ldots, G_n$ of G from G_i to G_{i+1} for $i = 1, \ldots, n-1$. We denote this graph by $P(n \cdot G)$. Kaneria, Makadia, and Meghpara [1852] proved the following graphs are graceful: the graph obtained by joining C_{4m} and C_{4n} by a path of arbitrary length; the path union of finite many copies of C_{4n}; and C_{4n} with twin chords. Kaneria, Makadia, Jariya, and Meghpara [1851] proved that the join sum of complete bipartite graphs, the star of complete bipartite graphs, and the path union of a complete bipartite graphs are graceful.

Given connected graphs $G_1, G_2, \ldots, G_n$, Kaneria, Makadia, and Jariya [1850] define a *cycle of graphs* $C(G_1, G_2, \ldots, G_n)$ as the graph obtained by adding an edge joining G_i to G_{i+1} for $i = 1, \ldots, n-1$ and an edge joining G_n to G_1. (The resulting graph can vary depending on which vertices of the G_is are chosen.) When the n graphs are isomorphic to G the notation $C(n \cdot G)$ is used. Kaneria et al. proved that $C(2t \cdot C_{4n})$ and $C(2t \cdot K_{n,n})$ are graceful. In [1853] and [1855] Kaneria, Makadia, and Meghpara prove that the following graphs are graceful: $C(2t \cdot K_{m,n})$; $C(C_{4n_1}, C_{4n_2}, \ldots, C_{4n_t})$ when t is even and $\sum_{i=1}^{\frac{t}{2}} n_i = \sum_{i=\frac{t}{2}}^{t} n_i$; $C(2t \cdot P_m \times P_n)$; the star of $P_m \times P_n$; and the path union of t copies of $P_m \times P_n$. Kaneria, Viradia, Jariya, and Makadia [1874] proved the cycle graph $C(t \cdot P_n)$ is graceful.

The *star of graphs* $G_1, G_2, \ldots, G_n$, denoted by $S(G_1, G_2, \ldots, G_n)$, is the graph obtained by identifying each vertex of $K_{1,n}$, except the center, with one vertex from each of $G_1, G_2, \ldots, G_n$. The case that $G_1 = G_2 = \cdots = G_n = G$ is denoted by $S(n \cdot G)$. In [1864] and [1865] Kaneria, Meghpara, and Makadia proved the following graphs are graceful:

$S(t \cdot K_{m,n})$; $S(t \cdot P_m \times P_n)$; the barycentric subdivision of $P_m \times P_n$ (that is, the graph obtained from $P_m \times P_n$ by inserting a new vertex in each edge); the graph obtained by replacing each edge of $K_{1,t}$ by P_n; the graph obtained by identifying each end point of $K_{1,n}$ with a vertex of $K_{m,n}$; and the graph obtained by identifying each end point of $K_{1,n}$ with a vertex of $P_m \times P_n$. Kanani and Kaneria [1821] proved that the following graphs are graceful: the barycentric subdivision of C_n-snakes (that is, the graph obtained from the subdivision of C_n by inserting a new vertex in each edge); the barycentric subdivision of alternate C_n-snakes; and quadrilateral snakes.

Kaneria and Makadia [1841] and [1842] proved the following graphs are graceful: $(P_m \times P_n) \cup (P_r \times P_s)$; $C_{2f+3} \cup (P_m \times P_n) \cup (P_r \times P_s)$, where $f = 2(mn + rs) - (m + n + r + s)$; the tensor product of P_n and P_3; the tensor product of P_m and P_n for odd m and n; the star of C_{4n}; the $t-$supersubdivision of $P_m \times P_n$; and the graph obtained by joining C_{4n} and a grid graph with a path. In [1863] Kaneria, Meghpara, and Makadia proved that the star of $K_{1,n}$ is a graceful tree.

The graph P_n^t is obtained by identifying one end point from each of t copies of P_n. The graph $P_n^t(G_1, G_2, \ldots, G_{tn})$ obtained by replacing each edge of P_n^t, except those adjacent to the vertex of degree t, by the graphs $G_1, G_2, \ldots, G_{tn}$ is called the *one point path union of* $G_1, G_2, \ldots, G_{tn}$. The case where $G_1 = G_2 = \cdots = G_{tn} = H$ is denoted by $P_n^t(tn \cdot H)$. In [1864] and [1865] Kaneria, Meghpara, and Makadia proved P_n^t and $P_n^t(tn \cdot K_{m,r})$ are graceful. In [1862] Kaneria and Meghpara proved P_n^t $(tn \cdot P_r \times P_s)$, $P_n^t(tn \cdot K_{1,m})$, $S(t \cdot C_{4n})$, and $P_n^t(tn \cdot C_{4m})$ are graceful.

A graph H is said to be a m-super subdivision of a simple graph G, if every edge of G is replaced by the complete bipartite graph $K_{m,m}$ with $m > 2$ in such a way that the end vertices of the edge are merged with any two vertices of the same partite set A or B of $K_{m,m}$ after removal of the edge of G. Srinivasan, Chidambaram, Devadoss, Pakkirisamy, and Krishnamoorthi [3690] proved that m-super subdivision of path and cycle are graceful.

Kanneria and Makadia [1843] define a *step grid graph* as the graph obtained by starting with paths $P_n, P_n, P_{n-1}, \ldots, P_2$ $(n \geq 3)$ arranged vertically parallel with the vertices in the paths forming horizontal rows and edges joining the vertices of the rows. In [1843] and [1844] they prove the following graphs are graceful: step grid graphs; one point union for a path of step grid graphs; cycles of step grid graphs; stars of step grid graphs; $m-$super subdivisions of the step grid graphs; open stars of step grid graphs; one point unions of paths of step grid graphs; and graphs obtained by joining C_{4m} and step grid graphs with a path of arbitrary length.

For n even [1845] Kaneria and Makadia [1845] define a *double step grid graph* of size n (denoted by DSt_n) as the graph obtained by starting with paths $P_n, P_n, P_{n-2}, P_{n-4}, \ldots, P_4, P_2$ arranged vertically parallel with the vertices in the paths forming horizontal rows and edges joining the vertices of the rows. They prove the following graphs are graceful: double step grid graphs; path unions of copies of DSt_n; cycles of $r \equiv 0,\ 3 \pmod 4$ copies of double step grid graphs; and stars of double step grid graphs.

In [1857] Kaneria, Makadia and Viradia prove the following graphs are graceful: open stars of double step grid graphs; one point union of paths of double step grid graphs

${P_n}^t(tn \cdot DSt_m)$; graphs obtained by joining C_{4m} and a double step grid graph with a path of arbitrary length; and graphs obtained by starting with a cycle ${C_m}^+$ ($m \equiv 2 \bmod 4$) with chords that form a triangle with an edge of the cycle and joining ${C_m}^+$ and a double step grid graph with a path of arbitrary length.

For even $n > 2$ Kaneria and Makadia [1846] define a *plus graph* of size n (denoted by Pl_n) as the graph obtained by starting with paths $P_2, P_4, \ldots, P_{n-2}, P_n, P_n, P_{n-2}, \ldots, P_4, P_2$ arranged vertically parallel with the vertices in the paths forming horizontal rows and edges joining the vertices of the rows. They prove plus graphs, path unions of copies of Pl_n, cycles of $r \equiv 0,\ 3 \pmod 4$ copies of Pl_n, and stars of plus graphs are graceful. In [1847] Kaneria and Makadia prove the following graphs are graceful: open stars of plus graphs; graphs obtained by joining C_{4m} and a plus graph with a path of arbitrary length; graphs obtained from cycles ${C_m}^+$ ($m \equiv 2 \pmod 4$) with twin chords that form a triangle with an edge of the cycle by joining ${C_m}^+$ and a plus graph with a path of arbitrary length.

Kaneria and Makadia [1848] define a *swastik graph* as the graph obtained from four copies of C_{4n} ($n > 1$) with vertices $V_{i,j}$ ($i = 1, 2, 3, 4,\ j = 1, 2, \ldots, 4n$) and identifying $V_{1,4t}$ and $V_{2,1}$, $V_{2,4t}$ and $V_{3,1}$, $V_{3,4t}$ and $V_{4,1}$, and $V_{4,4t}$ and $V_{1,1}$. They proved that path unions of swastik graphs of the same size, cycles of $r \equiv 0,\ 3 \pmod 4$ copies of swastik graphs of the same size, and the star of swastik graphs are graceful. In [1849] Kaneria and Makadia prove the following graphs are graceful: open stars of swastik graphs; one point unions for paths of swastik graphs; graphs obtained by joining C_{4m} and a swastik graph with a path of arbitrary length; graphs obtained from cycles C_m ($m \equiv 2 \pmod 4$) with twin chords that form a triangle with an edge by joining $C_m \odot K_1$ and a swastik graph with a path of arbitrary length.

In [1836] and [1835] Kaneria and Jariya define a *smooth graceful graph* as a bipartite graph G with q edges with the property that for all positive integers l there exists a map $g : V \longrightarrow \{0, 1, \ldots, \lfloor \frac{q-1}{2} \rfloor, \lfloor \frac{q+1}{2} \rfloor + l, \lfloor \frac{q+3}{2} \rfloor + l, \ldots, q + l\}$ such that the induced edge labeling map $g^\star : E \longrightarrow \{1 + l, 2 + l, \ldots, q + l\}$ defined by $g^\star(e) = |g(u) - g(v)|$ is a bijection. Note that by taking $l = 0$ a smooth graceful labeling is a graceful labeling. Kaneria and Jariya proved the following graphs are smooth graceful: P_n; C_{4n}; $K_{2,n}$; $P_m \times P_n$; and the graph obtained by joining a cycle C_{4m+2} with twin chords to C_{4n}. They also proved that the graph obtained by joining C_{4m} to W_n with a path is graceful. They proved that $K_{1,n}$ is semi smooth graceful, the star of $K_{1,n}$ is graceful, the path union of a smooth graceful tree is graceful, and the star of a smooth graceful tree is a graceful tree.

Kaneria, Makadia and Viradia [1858] proved the following: the star of a semi smooth graceful graph is graceful; $K_{m,n}$, $P(t \cdot H)$ are semi smooth graceful where H is a semi smooth graceful graph; step grid graphs; and the cycle graphs $C(t \cdot H)$ are smooth graceful, when $t \equiv \pmod 4$, H is a semi smooth; $C^t(m \cdot C_n)$, $P^t(k \cdot T)$, $< C_{n_1}, P_{n_2}, C_{n_3}, \ldots, P_{n_{2t}}, C_{n_{2t+1}} >$, $< K_{m_1,n_1}, P_{r_1}, K_{m_2,n_2}, P_{r_2}, \ldots, P_{r_{t-1}}, K_{m_t,n_t} >$, $< P_{n_1} \times P_{m_1}, P_{r_1}, P_{n_2} \times P_{m_2}, \ldots, P_{r_{t-1}}, P_{n_t} \times P_{m_t} >$ are graceful when T is semi smooth graceful tree.

Kaneria and Meghpara [1861] prove that $B_{m,n}$, the splitting graphs $S'(B_{m,n})$ and $S'(P_n)$ are semi smooth graceful and if graphs obtained by joining semi smooth graceful graph and $B^2_{m,n}$ by an arbitrary path is graceful.

A *komodo dragon* is formed by attaching a path to a vertex of degree 3 in a cycle with a chord and attaching star graphs to the end points of the path. A *komodo dragon with many tails* is formed by attaching many paths of length two to an endpoint of the path in a komodo dragon. In [3470] and [3472] Shahul Hameed, Palanivelrajan, Gunasekaran and Raziya Begam provide graceful labelings of various komodo dragon graphs and their extensions. In [3471] and [3473] Shahul Hameed et al. investigated the gracefulness of classes of graphs constructed by combining some subdivisions of certain theta graphs with stars.

For a bipartite graph G with partite sets X and Y let G' be a copy of G and X' and Y' be copies of X and Y. Lee and Liu [2163] define the *mirror graph*, $M(G)$, of G as the disjoint union of G and G' with additional edges joining each vertex of Y to its corresponding vertex in Y'. The case that $G = K_{m,n}$ is more simply denoted by $M(m, n)$. They proved that for many cases $M(m, n)$ has a stronger form of graceful labeling (see §3.1 for details).

The *total graph* $T(P_n)$ has vertex set $V(P_n) \cup E(P_n)$ with two vertices adjacent whenever they are neighbors in P_n. Balakrishnan, Selvam, and Yegnanarayanan [466] have proved that $T(P_n)$ is harmonious.

For any graph G with vertices $v_1, \ldots, v_n$ and a vector $\mathbf{m} = (m_1, \ldots, m_n)$ of positive integers the corresponding *replicated graph*, $R_{\mathbf{m}}(G)$, of G is defined as follows. For each v_i form a stable set S_i consisting of m_i new vertices $i = 1, 2, \ldots, n$ (a *stable* set S consists of a set of vertices such that there is not an edge v_iv_j for all pairs v_i, v_j in S); two stable sets $S_i, S_j, i \neq j$, form a complete bipartite graph if each v_iv_j is an edge in G and otherwise there are no edges between S_i and S_j. Ramírez-Alfonsín [3114] has proved that $R_{\mathbf{m}}(P_n)$ is graceful for all $\mathbf{m}$ and all $n > 1$ (see §3.4 for a stronger result) and that $R_{(m,1,\ldots,1)}(C_{4n}), R_{(2,1,\ldots,1)}(C_n)$ $(n \geq 8)$ and,$R_{(2,2,1,\ldots,1)}(C_{4n})$ $(n \geq 12)$ are graceful.

For any permutation f on $1, \ldots, n$, the *f-permutation* graph on a graph G, $P(G, f)$, consists of two disjoint copies of G, G_1 and G_2, each of which has vertices labeled $v_1, v_2, \ldots, v_n$ with n edges obtained by joining each v_i in G_1 to $v_{f(i)}$ in G_2. In 1983 Lee (see [2228]) conjectured that for all $n > 1$ and all permutations on $1, 2, \ldots, n$, the permutation graph $P(P_n, f)$ is graceful. Lee, Wang, and Kiang [2228] proved that $P(P_{2k}, f)$ is graceful when $f = (12)(34)\cdots(k, k+1)\cdots(2k-1, 2k)$. They conjectured that if G is a graceful nonbipartite graph with n vertices, then for any permutation f on $1, 2, \ldots, n$, the permutation graph $P(G, f)$ is graceful. Fan and Liang [1063] have shown that if f is a permutation in S_n where $n \geq 2(m-1)+2l$ then the permutation graph $P(P_n, f)$ is graceful if the disjoint cycle form of f is $\prod_{k=0}^{l-1}(m+2k, m+2k+1)$, and if $n \geq 2(m-1)+4l$ the permutation graph $P(P_n, f)$ is graceful the disjoint cycle form of f is $\prod_{k=0}^{l-1}(m+4k, m+4k+2)(m+4k+1, m+4k+3)$. For any integer $n \geq 5$ and some permutations f in S(n), Liang and Y. Miao, [2255] discuss gracefulness of the permutation graphs $P(P_n, f)$ if $f = (m, m+1, m+2, m+3, m+4), (m, m+2)(m+1, m+3), (m, m+1, m+2, m+4, m+3), (m, m+1, m+4, m+3, m+2), (m, m+2, m+3, m+4, m+1), (m, m+3, m+4, m+2, m+1)$ and $(m, m+4, m+3, m+2, m+1)$. In [2257] Liang, Zhang, Xu, Ye, Fan, and Ge prove the permutation graphs $P(P_n, f)$ where f is one of the permutations (12345), (2345), (234), (123456) and (23)(45) are graceful. Some families of graceful permutation graphs

are given in [2156], [2250], and [1359].

In [3192] Rofa defined generalized strongly graceful permutations and discovered two new permutations in addition to the known permutation that is obtained by replacing each vertex label $f(v)$ by $q - f(v)$. He uses these permutations to prove, by induction, that a lobster with a perfect matching that consists of the set of end edges of the lobster, is strongly graceful. He further showed that there exist strongly graceful labelings that assign the label 0 to four specific vertices of any tree belonging to this family of lobsters. He provided a tractable way for proving an equivalent form of Bermond conjecture which states that all lobsters are graceful. Two out of a total of three cases of the proposed equivalent form of Bermond's conjecture are completed leaving the third case open for refutation or completion. As an application of his results, he provided in [3193] number systems that are representable by all vertices of a rooted symmetric tree in such a way that the number representation of each vertex depends on its distance from the root vertex.

In [614] Bell provided methods to combine graceful bipartite graphs to create new graceful graphs. These methods unify and generalize some well-known results in the graceful labeling literature. She also found a new class of graceful trees.

The *power graph* of a finite group G has the elements of G as its vertex set and two distinct vertices of G are adjacent if one is a power of other. Sehgal, Takshak, Maan, and Malik [3336] proved that the power graph of $Z_2^{k-1} \times Z_4$ has a graceful labeling. In [3610] Singh, Takshak, Sehgal, and Malik investigated the gracefulness of graphs of the form Z_n^k for various cases. [3610] new

A graph (p, q)-graph $G(V, E)$ is said to be (k, d)-*hooked Skolem graceful* if there exists a bijection f from $V(G)$ to $\{1, 2, \ldots, p-1, p+1\}$ such that the induced edge labeling g_f from E to $\{k, k+d, \ldots, k+(n-1)d\}$ defined by $g_f(uv) = |f(u)-f(v)|$ for all uv in E is also bijective. Such a labeling f is called a (k, d)-*hooked Skolem graceful labeling* of G. Note that when $k = d = 1$, this notion coincides with that of hooked Skolem graceful labeling of the graph G. In [2824] Pereira, Singh, and Arumugam present some preliminary results on (k, d)-hooked Skolem graceful graphs and prove that nK_2 is $(2, 1)$-hooked Skolem graceful if and only if $n \equiv 1$ or 2 (mod 4).

Gnanajothi [1276, p. 51] calls a graph G *bigraceful* if both G and its line graph are graceful. She shows the following are bigraceful: P_m; $P_m \times P_n$; C_n if and only if $n \equiv 0, 3$ (mod 4); S_n; K_n if and only if $n \leq 3$; and B_n if and only if $n \equiv 3$ (mod 4). She also shows that $K_{m,n}$ is not bigraceful when $n \equiv 3$ (mod 4). (Gangopadhyay and Hebbare [1182] used the term "bigraceful" to mean a bipartite graceful graph.) Murugan and Arumugan [2593] have shown that graphs obtained from C_4 by attaching two disjoint paths of equal length to two adjacent vertices are bigraceful.

Several well-known isolated graphs have been examined. Graceful labelings have been found for the Petersen graph [1150], the cube [1201], the icosahedron and the dodecahedron. In [1202] [pp. 163-164] Gardner credits Ashenfelter and Chandra for showing the Platonic solids have graceful labelings.[1] Gardner stated that the icosahedron has only five fundamentally different graceful labelings, whereas in 2021 Knuth [1970] determined the correct number to be 24. Graham and Sloane [1327] showed that all of these solids except

[1]D. Knuth called this to my attention.

the cube are harmonious. Winters [4206] verified that the Grőtzsch graph (see [686, p. 118]), the Heawood graph (see [686, p. 236]), and the Herschel graph (see [686, p. 53]) are graceful. Graham and Sloane [1327] determined all harmonious graphs with at most five vertices. Seoud and Youssef [3408] did the same for graphs with six vertices.

In 2009 Zak [4301] defined the following generalization of harmonious labelings. For a graph $G(V, E)$ and a positive integer $t \geq |E|$ a function h from $V(G)$ to Z_t (the additive group of integers modulo t) is called a *t-harmonious labeling* of G if h is injective for $t \geq |V|$ or surjective for $t < |V|$, and $h(u) + h(v) \neq h(x) + h(y)$ for all distinct edges uv and xy. The smallest such t for which G has a t-harmonious labeling is called the *harmonious order of G*. Obviously, a graph $G(V, E)$with $|E| \geq |V|$ is harmonious if and only if the harmonious order of G is $|E|$. Zak determines the harmonious order of complete graphs, complete bipartite graphs, even cycles, some cases of P_n^k, and $2nK_3$. He presents some results about the harmonious order of the Cartesian products of graphs, the disjoint union of copies of a given graph, and gives an upper bound for the harmonious order of trees. He conjectures that the harmonious order of a tree of order n is $n + o(n)$. Hegde and Murthy [1415] proved Zak's conjecture [4301] using the value sets of polynomials, which partially proves the cordial tree conjecture by Hovey [1452] that all trees of order less than a prime p are p-cordial. (See Section 3.7.)

A graceful labeling of P_n is said to be an $(a, b; n)$-*graceful* labeling if one endpoint is labeled a and the other labeled b. A conjecture made in Gvozdjak's PhD Thesis [1349] on the Oberwolfach Problem in 2004 is: "An $(a, b; n)$-graceful labeling of P_n exists if and only if the integers a, b, n satisfy (1) $b - a$ has the same parity as $n(n + 1)/2$; (2) $0 < |b - a| \leq (n + 1)/2$ and (3) $n/2 \leq a + b \leq 3n/2$." In [4325] Zhang, Zhang, and Wang showed that the conjecture is true for every n whenever it is true for $n \leq 4a + 1$ and a is a fixed value. Moreover, they proved that the conjecture is true for $a = 0, 1, 2, 3, 4, 5, 6$.

For a graph with e edges Vietri [4087] generalizes the notion of a graceful labeling by allowing the vertex labels to be real numbers in the interval $[0, e]$. For a simple graph $G(V, E)$ he defines an injective map γ from V to $[0, e]$ to be a *real-graceful* labeling of G provided that $\sum 2^{\gamma(u)-\gamma(v)} + 2^{\gamma(v)-\gamma(u)} = 2^{e+1} - 2^{-e} - 1$, where the sum is taken over all edges uv. In the case that the labels are integers, he shows that a real-graceful labeling is equivalent to a graceful labeling. In contrast to the case for graceful labelings, he shows that the cycles C_{4t+1} and C_{4t+2} have real-graceful labelings. He also shows that the non-graceful graphs K_5, K_6, and K_7 have real-graceful labelings. With one exception, his real-graceful labels are integers.

The *gamma-number* (or gracefulness) of a graph G, denoted by $\gamma(G)$, is the smallest positive integer n for which there exists an injective function $f : V(G) \to \{0, 1, \ldots, n\}$ such that each $uv \in E(G)$ is labeled $|f(u) - f(v)|$ and the resulting edge labels are distinct. The *strong gamma-number* of a graph G, denoted by $\gamma_s(G)$, is defined to be the smallest positive integer n such that $\gamma(G) = n$ with the additional property that there exists an integer λ so that $\min\{f(u), f(v)\} \leq \lambda \max\{f(u), f(v)\}$ for each $uv \in E(G)$. The strong gamma-number is defined to be $+\infty$, otherwise. Ichishima and Oshima [1516] proved that if G is a bipartite graph, then $\gamma(mG) \leq m\gamma(G) + m - 1$ for any positive integer m. They also show that $\gamma_s(G) < +\infty$ and $\gamma_s(G) \leq 2\gamma(G) + 1$ for any bipartite

graph G. Moreover, they provide a sharp upper bound for $\gamma(G \cup H)$ in terms of $\gamma(G)$ and $\gamma_s(H)$ when G and H are graphs such that H is bipartite, and give formulas for the gamma-number of certain forests. In addition to these, they present strong gamma-number analogues to the gamma-number results and determine the exact values of the gamma-number and strong gamma-number for all cycles.

A graph G with m vertices and n edges, is said to be *prime graceful* if there is an injection ϕ from the vertices of G to $\{1, 2, \ldots, k\}$ where $k = \min\{2m, 2n\}$ such that $\gcd(\phi(v_i), \phi(v_j)) = 1$ and the induced injective function ϕ^* from the edges of G to $\{1, 2, \ldots, k-1\}$ defined by $\phi^*(v_iv_j) = |\phi(v_i) - \phi(v_j)|$, the resulting edge labels are distinct. In [**?**] Selvarajan and Subramoniam proved paths, cycles, stars, friendship graphs, bistars, $C_4 \cup P_n$, $K_{m,2}$, and $K_{m,2} \cup P_n$ have prime graceful labelings.

In 2020 [4306] Zeen El Deen introduced a new type of labeling of a graph as follows. For any positive integer δ an edge $\delta-$ graceful labeling of a graph $G(V, E)$ with p vertices and q edges is a bijective f from E to $\{\delta, 2\delta, \ldots, q\delta\}$ such $f[V] = \{f(u) = \sum f(uv) \bmod(k\delta)$ over all edges v incident to u and $k = \max(p, q)$ are pairwise distinct. He proved the existence of an edge $\delta-$graceful labeling, for any positive integer δ, for wheels, alternate triangular cycles, double wheels, $C_n \times P_2$, $W_n \times P_2$, gears, helms, butterflies, and friendship graphs. In [4309] Zeen El Deen and Elmahdy showed that for any positive integer δ there is an edge δ-graceful labeling for the following graphs: the splitting graphs of cycles, fans, and crowns; the shadow graphs of the paths, cycles, and fans; the middle graphs and the total graphs of paths, cycles, and crowns; twigs; and snails. In [4307] Zeen Deen demonstrated the existence of an edge δ-graceful labelings for certain snake and chain-related graphs. [4307] new

In [733] Byers and O'Mellan introduced a new concept that combines graceful and harmonious labelings as follows. A connected graph G with size m is said to be a *graceful-harmonious* labeling if there exits injection $f : V(G) \to \{0, 1, 2, \ldots, m\}$ such that when each edge uv is assigned either $|f(u) - f(v)|$ or $(f(u) + f(v)) \pmod q$. each edge receives a distinct label from the set $\{0, 1, 2, \ldots, m\}$. They prove that cycles, friendship graphs, and double cones $C_{4k} + \overline{K}_2$ admit graceful-harmonious labelings.

For a graph $G(V, E)$ without isolated vertices, Pereira, Singh, and Arumugam [2823] defined the gracefulness, $grac(G)$, of G as the smallest positive integer k for which there exists an injective function $f : V \to \{0, 1, 2, \ldots, k\}$ such that the edge induced function $g_f : E \to \{1, 2, \ldots, k\}$ defined by $g_f(uv) = |f(u) - f(v)|$ is also injective. Let $c(f) = \{1, 2, \ldots, i\}$ denote the edge labels and let $m(G) = \max\{c(f)\}$, where the maximum is taken over all injective functions $f : V \to N \cup \{0\}$ such that g_f is also injective. This measure $m(G)$ determines how close G is to being graceful. They determine $m(G)$ for certain cycles and friendship graphs.

A number of authors have investigated the gracefulness of the directed graphs obtained from copies of directed cycles $\vec{C}_m$ that have a vertex in common or have an edge in common. A digraph $D(V, E)$ is said to be *graceful* if there exists an injection $f : V(G) \to \{0, 1, \ldots, |E|\}$ such that the induced function $f' : E(G) \to \{1, 2, \ldots, |E|\}$ that is defined by $f'(u, v) = (f(v) - f(u)) \pmod{|E| + 1}$ for every directed edge uv is a bijection. The notations $n \cdot \vec{C}_m$ and $n - \vec{C}_m$ are used to denote the digraphs obtained from n copies of $\vec{C}_m$ with exactly one point in common and the digraphs obtained from n copies of $\vec{C}_m$ with

exactly one edge in common. In 1994 Du and Sun [979] proved that a necessary condition for $n - \vec{C}_m$ to be graceful is that mn is even and that $n \cdot \vec{C}_m$ is graceful when m is even. They conjectured that $n \cdot \vec{C}_m$ is graceful for any odd m and even n. This conjecture was proved by Jirimutu, Xu, Feng, and Bao in [1791]. In his 2006 PhD dissertation Jirimutu [1787] posed the problem: For any positive odd n and any odd $m \geq 14$ is digraph $n - \overrightarrow{C_m}$ graceful? Xu, Jirimutu, Wang, and Min [4230] proved that $n - \vec{C}_m$ is graceful for $m = 4, 6, 8, 10$ and even n. Feng and Jirimutu (see [4328]) conjectured that $n - \vec{C}_m$ is graceful for even n and asked about the situation for odd n. The cases where $m = 5, 7, 9, 11$, and 13 and even n were proved Zhao and Jirimutu [4327]. The cases for $m = 15, 17$, and 19 and even n were proved by Zhao et al. in [4326], and [3621]. Zhao, Siqintuya, and Jirimutu [4328] proved that a necessary condition for $n - \vec{C}_m$ to be graceful is that nm is even. In [1075] Feng discussed the gracefulness of the digraph $n - \overrightarrow{C_m}$ and proved the case when $m = 21$ for even n.

In a 1985 paper Bloom and Hsu [669] say a directed graph D with e edges has a graceful labeling θ if for each vertex v there is a vertex labeling θ that assigns each vertex a distinct integer from 0 to e such that for each directed edge (u, v) the integers $\theta(v) - \theta(u) \bmod (e + 1)$ are distinct and nonzero. They conjectured that digraphs whose underlying graphs are wheels and that have all directed edges joining the hub and the rim in the same direction and all directed edges in the same direction are graceful. This conjecture was proved in 2009 by Hegde and Shivarajkumar [1427]. Yao, Yao, and Cheng [4261] investigated the gracefulness for many orientations of undirected trees with short diameters and proved some directed trees do not have graceful labelings. Hegde and Kumudkshi [1413] established the gracefulness of the directed graph that is an orientation of the planar grid graph $P_m \times P_n$ in which each cell is a unicycle of length four. A *graceful difference labeling* of a directed graph G with vertex set V is a bijection $f : V \to \{1, \ldots, |V|\}$ such that, when each arc uv is assigned the difference label $f(v) - f(u)$, the resulting arc labels are distinct. Hertz and Picouleau [1437] conjectured that all disjoint unions of circuits have a graceful difference labeling, except in two particular cases. They provided partial results that support this conjecture. A survey of results on graceful digraphs by Feng, Xu, and Jirimutu is given in [1076]. Marr [2442] and [2441] summarizes previously known results on graceful directed graphs and presents some new results on directed paths, stars, wheels, and umbrellas.

In [3563] Shivarajkumar, Sriraj, and Hegde provided a 2021 survey results on graceful labeling of digraphs.

2.8 Summary

The results and conjectures discussed above are summarized in the tables following. The letter G after a class of graphs indicates that the graphs in that class are known to be graceful; a question mark indicates that the gracefulness of the graphs in the class is an open problem; we put a question mark after a "G" if the graphs have been conjectured to be graceful. The analogous notation with the letter H is used to indicate the status of the graphs with regard to being harmonious. The tables impart at a glimpse what has

been done and what needs to be done to close out a particular class of graphs. Of course, there is an unlimited number of graphs one could consider. One wishes for some general results that would handle several broad classes at once but the experience of many people suggests that this is unlikely to occur soon. The Graceful Tree Conjecture alone has withstood the efforts of scores of people over the past four decades. Analogous sweeping conjectures are probably true but appear hopelessly difficult to prove. I thank Don Knuth for his correspondence about the results of Smith and Puget [3641] in Table 1 regarding the gracefulness $K_m \times K$, $K_m \times P_n$, and $K_m \times C_n$.

Table 1: **Summary of Graceful Results**

Graph	*Graceful*
trees	G if ≤ 35 vertices [1065] G if symmetrical [633] G if at most 4 end-vertices [1463] G with diameter at most 5 [1453] G? Ringel-Kotzig G caterpillars [3203] G firecrackers [795] G bananas [3431], [3430] G? lobsters [629]
cycles C_n	G iff $n \equiv 0, 3 \pmod 4$ [3203]
wheels W_n	G [1150], [1449]
helms (see §2.2)	G [324]
webs (see §2.2)	G [1883]
gears (see §2.2)	G [2374]
cycles with P_k-chord (see §2.2)	G [921], [2373], [1989], [3056]
C_n with k consec. chords (see §2.2)	G if $k = 2, 3, n-3$ [1978], [1986]
unicyclic graphs	G? iff G $\neq C_n, n \equiv 1, 2 \pmod 4$ [3918]
P_n^k	G if $k = 2$ [1883]
$C_n^{(t)}$ (see §2.2)	$n = 3$ G iff $t \equiv 0, 1 \pmod 4$ [630], [632] G? if $nt \equiv 0, 3 \pmod 4$ [1979] G if $n = 6, t$ even [1979] G if $n = 4, t > 1$ [3496] G if $n = 5, t > 1$ [4248] G if $n = 7$ and $t \equiv 0, 3 \pmod 4$ [4254] G if $n = 9$ and $t \equiv 0, 3 \pmod 4$ [4255] G if $t = 2$ $n \not\equiv 1 \pmod 4$ [3063], [679] G if $n = 11$ [4232]

Continued on next page

Table 1 – *Continued from previous page*

Graph	*Graceful*
triangular snakes (see §2.2)	G iff no. blocks $\equiv 0, 1 \pmod 4$ [2559]
K_4-snakes (see §2.2)	?
quadrilateral snakes (see §2.2)	G [1276], [3063]
crowns $C_n \odot K_1$	G [1150]
$C_n \odot P_k$	G [3337]
grids $P_m \times P_n$	G [53]
prisms $C_m \times P_n$	G if $n = 2$ [1153], [4251] G if m even [1467] G if m odd and $3 \leq n \leq 12$ [1467] G if $m = 3$ [3600] G if $m = 6$ see [4253] G if $m \equiv 2 \pmod 4$, $n \equiv 3 \pmod 4$ [4253]
$K_m \times P_n$	G if $(m, n) = (4, 2), (4, 3), (4, 4), (4, 5),$ $(5, 2), (5, 3), (6, 3), (4, 6), (4, 7), (4, 8)$ not G if $(3, 3), (m, 2)$ $m = 6, 7, 8, 9, 10,$ 11,12 not G? for $(m, 2)$ with $m > 12$ [3641]
$K_m \times C_n$	G if $(m, n) = (4, 3), (3, 4), (4, 4), (4, 5),$ $(3, 6), (4, 6)$ not G for $(m, n) = (6, 3)$ [3641]
$K_m \odot K_1$	G if $m = 3, 4, 5, 6, 7, 8, 9$ not G if $m = 10, 11, 12, 13, 14, 15$ not G? if $m > 15$ [3641]
$K_{m,n} \odot K_1$	G [1786]
$K_m \cup K_n$ $(m, n > 1)$	G iff $\{m, n\} = \{4, 2\}$ or $\{5, 2\}$
$\bigcup_{i=1}^{t} K_{m_i,n_i}$	G $2 \leq m_i < n_i$ [512]
torus grids $C_m \times C_n$	G if $m \equiv 0 \pmod 4$, n even [1806]

Continued on next page

Table 1 – *Continued from previous page*

Graph	*Graceful*
	not G if m, n odd (parity condition)
vertex-deleted $C_m \times P_n$	G if $n = 2$ [1174]
edge-deleted $C_m \times P_n$	G if $n = 2$ [1174]
Möbius ladders M_n (see §2.3)	G [1168]
stacked books $S_m \times P_n$ (see §2.3)	$n = 2$, G iff $m \not\equiv 3 \pmod 4$ [2399], [920], [1173] G if m even [1173]
n-cube $K_2 \times K_2 \times \cdots \times K_2$	G [2015]
$K_n \times P_3$	G iff $n \leq 6$ [1971]
K_n	G iff $n \leq 4$ [1287], [3593]
$K_{m,n}$	G [3203], [1287]
$K_{1,m,n}$	G [270]
$K_{1,1,m,n}$	G [1276]
windmills $K_n^{(m)} (n > 3)$ (see §2.4)	G if $n = 4, m \leq 1000$ [1467], [10], [4167], [1235] G? if $n = 4, m \geq 4$ [629] G if $n = 5, m = 4,\ 5$ [3699] not G if $n = 4, m = 2, 3$ [629] not G if $(m, n) = (2, 5)$ [632] not G if $n > 5$ [1986]
$B(n, r, m)$ $r > 1$ (see §2.4)	G if $(n, r) = (3, 2), (4, 3)$ [1980], (4,2) [920] G $(n, r, m) = (5, 2, 2)$ [3641] not G for $(n, 2, 2)$ for $n > 5$ [631], [3641]
mK_n (see §2.5)	G iff $m = 1, n \leq 4$ [2019]
$C_m \cup P_n$	G iff $m + n \geq 6$ [3911]
$C_m \cup C_n$	G iff $m + n \equiv 0, 3 \pmod 4$ [30]
$C_n \cup K_{p,q}$	for $n > 8$ G iff $n \equiv 0, 3 \pmod 4$ [4277]

Continued on next page

Table 1 – *Continued from previous page*

Graph	*Graceful*
	G $C_6 \times K_{1,2n+1}$ [512] G $C_3 \times K_{m,n}$ iff $m, n \geq 2$ [3407] G $C_4 \times K_{m,n}$ iff $(m, n) \neq (1, 1)$ [3407] G $C_7 \times K_{m,n}$ [3407] G $C_8 \times K_{m,n}$ [3407]
$K_i \cup K_{m,n}$	G [512]
$\bigcup_{i=1}^{t} K_{m_i,n_i}$	G $2 \leq m_i < n_i$ [512]
$C_m \cup \bigcup_{i=1}^{t} K_{m_i,n_i}$	G $2 \leq m_i < n_i$, $m \equiv 0$ or 3 (mod 4), $m \geq 11$ [512]
$G + \overline{K_t}$	G for connected graceful G [1980]
double cones $C_n + \overline{K_2}$	G for $n = 3, 4, 5, 7, 8, 9, 11, 12$ not G for $n \equiv 2$ (mod 4) [3152]
t-point suspension $C_n + \overline{K_t}$	G if $n \equiv 0$ or 3 (mod 12) [648] not G if t is even and $n \equiv 2, 6, 10$ (mod 12) G if $n = 4, 7, 11$ or 19 [648] G if $n = 5$ or 9 and $t = 2$ [648]
P_n^2 (see §2.7)	G [2154]
Petersen $P(n, k)$ (see §2.7)	G for $n = 5, 6, 7, 8, 9, 10$ [3152], $(n, k) = (8t, 3)$ [4068]

Table 2: **Summary of Harmonious Results**

Graph	*Harmonious*
trees	H if ≤ 31 vertices [1066] H? [1327] H caterpillars [1327] ? lobsters

Continued on next page

Table 2 – *Continued from previous page*

Graph	*Harmonious*
cycles C_n	H iff n is odd [1327]
wheels W_n	H [1327]
helms (see §2.2)	H [1276], [2297]
webs (see §2.2)	H if cycle is odd
gears (see §2.2)	H [796]
cycles with P_k-chord (see §2.2)	?
C_n with k consec. chords (see §2.2)	?
unicyclic graphs	?
P_n^k	H if $k = 2$ [1324], k odd [3369], [4284] H if k is even and $k/2 \leq (n-1)/2$ [4297]
$C_n^{(t)}$ (see §2.2)	$n = 3$ H iff $t \not\equiv 2 \pmod 4$ [1327] H if $n = 4,\ t > 1$ [3496]
triangular snakes (see §2.2)	H if number of blocks is odd [4229] not H if number of blocks $\equiv 2 \pmod 4$ [4229]
K_4-snakes (see §2.2)	H [1325]
quadrilateral snakes (see §2.2)	?
crowns $C_n \odot K_1$	H [1324], [2282]
grids $P_m \times P_n$ prisms $C_m \times P_n$	H iff $(m,n) \neq (2,2)$ [1806] H if $n = 2, m \neq 4$ [1174] H if n odd [1327] H if $m = 4$ and $n \geq 3$ [1806]

Continued on next page

Table 2 – *Continued from previous page*

Graph	*Harmonious*
torus grids $C_m \times C_n$,	H if $m = 4$, $n \geq 3$ [1806] not H if $m \not\equiv 0 \pmod 4$, n odd [1806]
vertex-deleted $C_m \times P_n$	H if $n = 2$ [1174]
edge-deleted $C_m \times P_n$	H if $n = 2$ [1174]
Möbius ladders M_n (see §2.3)	H iff $n \neq 3$ [1168]
stacked books $S_m \times P_n$ (see §2.3)	$n = 2$, H if m even [1323], [3154] not H $m \equiv 3 \pmod 4$, $n = 2$, (parity condition) H if $m \equiv 1 \pmod 4$, $n = 2$ [1276]
n-cube $K_2 \times K_2 \times \cdots \times K_2$	H if and only if $n \geq 4$ [1512]
$K_4 \times P_n$	H [3154]
K_n	H iff $n \leq 4$ [1327]
$K_{m,n}$	H iff m or $n = 1$ [1327]
$K_{1,m,n}$	H [270]
$K_{1,1,m,n}$	H [1276]
windmills $K_n^{(m)}$ $(n > 3)$ (see §2.4)	H if $n = 4$ [1457] $m = 2$, H? iff $n = 4$ [1327] not H if $m = 2, n$ odd or 6 [1327] not H for some cases $m = 3$ [2279]
$B(n, r, m)$ $r > 1$ (see §2.4)	$(n, r) = (3, 2), (4, 3)$ [3404]
mK_n (see §2.5)	H $n = 3$, m odd [2283] not H for n odd and $m \equiv 2 \pmod 4$ [2283]
nG	H when G is harmonious and n odd [4275]

Continued on next page

Table 2 – *Continued from previous page*

Graph	*Harmonious*
G^n	H when G is harmonious and n odd [4275]
$C_m \cup P_n$	H $n = 1$ iff $m \neq 2 \bmod 4$ [2695] H $n = 2$ [2695] H $(m, 3)$ odd $m \geq 3$ [3162], [2695] H? $(m, 3)$ $m \geq 3$
fans $F_n = P_n + K_1$	H [1327]
$nC_m + K_1$ $n \not\equiv 0 \bmod 4$	H [796]
double fans $P_n + \overline{K_2}$	H [1327]
t-point suspension $P_n + \overline{K_t}$ of P_n	H [3154]
$S_m + K_1$	H [1276], [773]
t-point suspension $C_n + \overline{K_t}$ of C_n	H if n odd and $t = 2$ [3154], [1276] not H if $n \equiv 2, 4, 6 \pmod 8$ and $t = 2$ [1276]
Petersen $P(n, k)$ (see §2.7)	H [1276], [2190]

3 Variations of Graceful Labelings

3.1 α-labelings

In 1966 Rosa [3203] defined an *α-labeling* (or *α-valuation*) as a graceful labeling with the additional property that there exists an integer k so that for each edge xy either $f(x) \leq k < f(y)$ or $f(y) \leq k < f(x)$. (Other names for such labelings are *balanced*, *interlaced*, and *strongly graceful*.) It follows that such a k must be the smaller of the two vertex labels that yield the edge labeled 1. Also, a graph with an α-labeling is necessarily bipartite and therefore cannot contain a cycle of odd length. Wu [4216] has shown that a necessary condition for a bipartite graph with n edges and degree sequence $d_1, d_2, \ldots, d_p$ to have an α-labeling is that the $\gcd(d_1, d_2, \ldots, d_p, n)$ divides $n(n-1)/2$. Barrientos and Minion [541] proved that any tree of size n and excess ϵ is a spanning tree of a graph of size $n + \epsilon$ that admits an α-labeling.

For a path with consecutive vertices $v_1, v_2, \ldots, v_n$ a *triangular tree* is the tree obtained by identifying each v_i to an end vertex of the path P_i. Barrientos [519] proved that all triangular trees admit an α-labeling. He also presented several ways to combine this type of trees to construct new trees and unicyclic graphs that can be α-labeled.

A common theme in graph labeling papers is to build up graphs that have desired labelings from pieces with particular properties. In these situations, starting with a graph that possesses an α-labeling is a typical approach. (See [773], [1324], [795], and [1806].) Moreover, Jungreis and Reid [1806] showed how sequential labelings of graphs (see Section 4.1) can often be obtained by modifying α-labelings of the graphs.

Graphs with α-labelings have proved to be useful in the development of the theory of graph decompositions. Rosa [3203], for instance, has shown that if G is a graph with q edges and has an α-labeling, then for every natural number p, the complete graph K_{2qp+1} can be decomposed into copies of G in such a way that the automorphism group of the decomposition itself contains the cyclic group of order p. In the same vein El-Zanati and Vanden Eynden [1031] proved that if G has q edges and admits an α-labeling then $K_{qm,qn}$ can be partitioned into subgraphs isomorphic to G for all positive integers m and n. Although a proof of Ringel's conjecture that every tree has a graceful labeling has withstood many attempts, examples of trees that do not have α-labelings are easy to construct (one example is the subdivision graph of $K_{1,3}$ — see [3203]). Kotzig [2013] has shown however that almost all trees have α-labelings. Sethuraman and Ragukumar [3438] have proved that every tree is a subtree of a graph with an α-labeling.

As to which graphs have α-labelings, Rosa [3203] observed that the n-cycle has an α-labeling if and only if $n \equiv 0 \pmod 4$ whereas P_n always has an α-labeling. Other familiar graphs that have α-labelings include caterpillars [3203], the n-cube [2012], Möbius ladders M_n when n is odd (see §2.3) for the definition) [2786], B_{4n+1} (i.e., books with $4n+1$ pages) [1173], $C_{2m} \cup C_{2m}$ and $C_{4m} \cup C_{4m} \cup C_{4m}$ for all $m > 1$ [2014], $C_{4m} \cup C_{4m} \cup C_{4n}$ for all $(m, n) \neq 1, 1)$ [1053], $P_n \times Q_n$ [2399], $K_{1,2k} \times Q_n$ [2399], $C_{4m} \cup C_{4m} \cup C_{4m} \cup C_{4m}$ [2091], $C_{4m} \cup C_{4n+2} \cup C_{4r+2}, C_{4m} \cup C_{4n} \cup C_{4r}$ when $m + n \leq r$ [30], $C_{4m} \cup C_{4n} \cup C_{4r} \cup C_{4s}$ when $m \geq n+r+s$ [26], $C_{4m} \cup C_{4n} \cup C_{4r+2} \cup C_{4s+2}$ when $m \geq n+r+s+1$ [26], $((m+1)^2+1)C_4$ for

all m [4336], k^2C_4 for all k [4336], and $(k^2+k)C_4$ for all k [4336]. Abrham and Kotzig [28] have shown kC_4 has an α-labeling for $4 \leq k \leq 10$ and that if kC_4 has an α-labeling then so does $(4k+1)C_4, (5k+1)C_4$, and $(9k+1)C_4$. Eshghi [1047] proved that $3C_{4k}$ and $5C_{4k}$ have an α-labeling for all k. In [1053] Eshghi and Carter show several families of graphs of the form $C_{4n_1} \cup C_{4n_2} \cup \cdots \cup C_{4n_k}$ have α-labelings.

In [3242] Salarrezaei and Eshghi provided an integer programming model and a Tabu search algorithm to generate α-labelings of the quadratic graphs mC_{4k}) where $6 \geq m \geq 10$ and $2 \geq k \geq 10$. The computational complexity of the gracefulness of a graph is not known, but the complexity of finding a harmonious labeling of a graph is in the NP-class [214]. Research on programming models for finding graceful labelings of graphs can be found in [1046], [1054], [2086], [3243] [1051], [3152], [3641], [2404], and [3458].

In [214] Amini and Eshghi gave a new mathematical integer programming model for the graph labeling graphs of the form mC_n (some authors use the notation $Q(m,n)$). The advantages of this model are linearity and the existence of an objective function. They also gave two constraint programming models and a meta-heuristics algorithm that generate feasible graceful labeling and α-labeling for special classes of quadratic graphs. Their results include: mC_{4k} with $1 \leq 11$ and less than 1000 vertices has an α-labeling with the exception of $3C_4$; $12C_{4k}$ has α-labeling for $1 \leq k \leq 19$; and $13C_{4k}$ has α-labeling for $1 \leq k \leq 13$. In [1054] Eshghi and Salarrezaei proved that $7C_{4k}$ has an α-labeling for all k. Lakshmi and Vangipuram [2086] proved that $4C_{4k}$ is graceful.

Paterson and Stinson [2806] used α-valuations of the lexicographic product of cycles and the complement of a complete graph to construct various circular external difference families.

In [543], Barrientos and Minion investigated series-parallel operations with graphs that admit α-labelings. They provided necessary conditions on the graphs G_1 and G_2 to obtain a new α-labeled graph G through each of these operations. As consequence of the series operation, they proved that the one-point union of three or four copies of $K_{n,n}$ has an α-labeling, and that any tree with maximum degree four that can be decomposed into copies of the path of length eleven has an α-labeling when the distance between any pair of vertices of degree four is even. They also showed that any graph of order $n+1$ and size n with an α-labeling is an induced subgraph of a graph of order $n+3$ and size $2n+1$. Additionally, they presented an α-labeling for any graph of the form $K_{2,n} \times P_m$. In [521] Barrientos used vertex and edge duplications, replications of the entire graph, and k-vertex amalgamations to generate α-labeled graphs. He proved that for some families of graphs, it is possible to duplicate several vertices or edges. Using k-vertex amalgamations he obtained an α-labeling of a graph that can be decomposed into multiple copies of a given α-labeled graph as well as a robust family of irregular grids that can α-labeled.

Figueroa-Centeno, Ichishima, and Muntaner-Batle [1090] have shown that if $m \equiv 0 \pmod 4$ then the one-point union of 2, 3, or 4 copies of C_m admits an α-labeling, and if $m \equiv 2 \pmod 4$ then the one-point union of 2 or 4 copies of C_m admits an α-labeling. They conjecture that the one-point union of n copies of C_m admits an α-labeling if and only if $mn \equiv 0 \pmod 4$.

In [3592] Simarmata, Sandy, and Sugeng gave a new approach to showing graphs admit

an α labeling using an adjacency matrix. They use it to construct graceful labelings for superstars (rooted trees constructed from several star graphs by connecting the leaves from each star graph to a root vertex) and a super-rooted trees (rooted trees constructed from several rooted trees by connecting the root vertex of each rooted tree to a root vertex).

Pei-Shan Lee [2144] proved that $C_6 \times P_{2t+1}$ and gear graphs have α-labelings. He raises the question of whether $C_{4m+2} \times P_{2t+1}$ has an α-labeling for all m. Brankovic, Murch, Pond, and Rosa [703] conjectured that all trees with maximum degree three and a perfect matching have an α-labeling. In [3928] Uma and Rajasekaran gave an α-valuation for the tensor product of paths and cycles.

In his 2001 Ph. D. thesis Selvaraju [3346] investigated the one-point union of complete bipartite graphs. He proves that the one-point unions of the following forms have an α-labeling: K_{m,n_1} and K_{m,n_2}; K_{m_1,n_1}, K_{m_2,n_2}, and K_{m_3,n_3} where $m_1 \leq m_2 \leq m_3$ and $n_1 < n_2 < n_3$; $K_{m_1,n}$, $K_{m_2,n}$, and $K_{m_3,n}$ where $m_1 < m_2 < m_3 \leq 2n$.

Zhile [4336] uses $C_m(n)$ to denote the connected graph all of whose blocks are C_m and whose block-cutpoint-graph is a path. He proves that for all positive integers m and n, $C_{4m}(n)$ has an α-labeling but $C_m(n)$ does not have an α-labeling when m is odd.

Abrham and Kotzig [30] have proved that $C_m \cup C_n$ has an α-labeling if and only if both m and n are even and $m + n \equiv 0 \pmod 4$. Kotzig [2014] has also shown that $C_4 \cup C_4 \cup C_4$ does not have an α-labeling. He asked if $n = 3$ is the only integer such that the disjoint union of n copies of C_4 does not have an α-labeling. This was confirmed by Abrham and Kotzig in [29]. Eshghi [1046] proved that every 2-regular bipartite graph with 3 components has an α-labeling if and only if the number of edges is a multiple of four except for $C_4 \cup C_4 \cup C_4$. In [1049] Eshghi gives more results on the existence of α-labelings for various families of disjoint union of cycles.

Jungreis and Reid [1806] investigated the existence of α-labelings for graphs of the form $P_m \times P_n, C_m \times P_n$, and $C_m \times C_n$ (see also [1171]). Of course, the cases involving C_m with m odd are not bipartite, so there is no α-labeling. The only unresolved cases among these three families are $C_{4m+2} \times P_{2n+1}$ and $C_{4m+2} \times C_{4n+2}$. All other cases result in α-labelings.

Let $v_{1,j}, v_{2,j}, \ldots, v_{m,j}$ be the consecutive vertices of the jth copy of P_m in $P_m \times P_n$. An *elementary transformation* of $P_m \times P_n$ is the graph obtained by replacing the edge $v_{i,j}v_{i+1,j}$ by the new edge $v_{i-x,j}v_{i+1+x,j}$. A graph is said to be a *grid-like graph* if it is obtained through a sequence of elementary transformations. In [544] Barrientos and Minion proved the existence of an α-labeling for any grid-like graph. As consequence of this result, they showed that the graphs $C_{4t} \times P_n \cup P_n$ and $C_{4t} \times P_n \cup P_{t-1} \times P_n$ admit α-labelings.

Balakrishman [459] uses the notation $Q_n(G)$ to denote the graph $P_2 \times P_2 \times \cdots \times P_2 \times G$ where P_2 occurs $n-1$ times. Snevily [3644] has shown that the graphs $Q_n(C_{4m})$ and the cycles C_{4m} with the path P_n adjoined at each vertex have α-labelings. He [3645] also has shown that compositions of the form $G[\overline{K_n}]$ (see §2.3 for the definition) have an α-labeling whenever G does (see §2.3 for the definition of composition). Balakrishman and Kumar [463] have shown that all graphs of the form $Q_n(G)$ where G is $K_{3,3}, K_{4,4}$, or P_m have an α-labeling. Balakrishman [459] poses the following two problems. For which

graphs G does $Q_n(G)$ have an α-labeling? For which graphs G does $Q_n(G)$ have a graceful labeling?

Rosa [3203] has shown that $K_{m,n}$ has an α-labeling (see also [509]). In [1515] Ichishima and Oshima proved that if m, s and t are integers with $m \geq 1, s \geq 2$, and $t \geq 2$, then the graph $mK_{s,t}$ has an α-labeling if and only if $(m, s, t) \neq (3, 2, 2)$. Barrientos [509] has shown that for n even the graph obtained from the wheel W_n by attaching a pendent edge at each vertex has an α-labeling. In [516] Barrientos shows how to construct graceful graphs that are formed from the one-point union of a tree that has an α-labeling, P_2, and the cycle C_n. In some cases, P_2 is not needed. Qian [3063] has proved that quadrilateral snakes have α-labelings. Yu, Lee, and Chin [4294] showed that Q_3-and Q_3-snakes have α-labelings. Fu and Wu [1157] showed that if T is a tree that has an α-labeling with partite sets V_1 and V_2 then the graph obtained from T by joining new vertices $w_1, w_2, \ldots, w_k$ to every vertex of V_1 has an α-labeling. Similarly, they prove that the graph obtained from T by joining new vertices $w_1, w_2, \ldots, w_k$ to the vertices of V_1 and new vertices $u_1, u_2, \ldots, u_t$ to every vertex of V_2 has an α-labeling. They also prove that if one of the new vertices of either of these two graphs is replaced by a star and every vertex of the star is joined to the vertices of V_1 or the vertices of both V_1 and V_2, the resulting graphs have α-labelings. Fu and Wu [1157] further show that if T is a tree with an α-labeling and the sizes of the two partite sets of T differ by at most 1, then $T \times P_m$ has an α-labeling. Zhao, Ma, and Yao [4330] proved that a class of super lobster trees have α-labelings. Ghosh [1266] uses various methods of joining graceful graphs and graphs with α-labelings to obtain some classes of graceful lobsters. Lalitha and Tamilselvi [2089] proved that the hexagonal snake graph has an α-labeling.

Selvaraju and G. Sethurman [3351] prove that the graphs obtained from a path P_n by joining all the pairs of vertices u, v of P_n with $d(u, v) = 3$ and the graphs obtained by identifying one of vertices of degree 2 of such graphs with the center of a star and the other vertex the graph of degree 2 with the center of another star (the two stars needs need not have the same size) have α-labelings. They conjecture that the analogous graphs where 3 is replaced with any t with $2 \leq t \leq n-2$ have α-labelings.

In [1352] Hafez, El-Shanawany, and El Atik proved that if G has an α-labeling, then $G[\overline{K_r}], r \geq 2$ and $Spl_m(G)$ have α-labelings. Let the vertices of P_n $(n > 1)$ be $v_1 v_2 \cdots v_n$. Hafez et al. define $K(n) = \{K_s(n), s = 1, 2, \ldots\}$, where $K_s(n)$ is the graph obtained from P_n by joining the end vertices of the edge $v_i v_{i+1}$ to every vertex in the complete graph $K_{r_i}, i = 1, 2, \ldots, n-1$, of order r_i, such that (a) if $|V(K_{r_i}) \cap V(K_{r_{i+1}})| = t$, then $t \leq \min\{r_i, r_{i+1}\}$ and (b) K_{r_i} and K_{r_j} are disjoint when $|i - j| > 2$. They further prove that if G has an α-labeling, then the graph $G \bigtriangledown K(n)$ has an α-labeling ($G \bigtriangledown H$ has $V(G) \times V(H)$ as its vertex set and $u = (x_1, y_1)$ is adjacent to $v = (x_2, y_2)$ whenever $y_1 = y_2$ and $x_1 x_2 \in E(G)$ or $x_1 x_2 \in E(G)$ and $y_1 y_2 \in E(H)$.

Makadia, Karavadiya, and Kanerian [2409] proved that the graph obtained by merging t consecutive vertices of two cycle C_{4r} and C_{4s} has an α-labeling when $t \leq 2\min\{r, s\}$. They also proved that if G_1 has an α-labeling and G_2 is graceful then there exists a graceful labeling of the graph obtained by joining G_1 and G_2 by any path. Moreover, if both G_1 and G_2 have α-labelings then there exists an α-labeling of the graph obtained by

joining G_1 and G_2 by any path. Let $C_{n_1}, C_{n_2}, \ldots, C_{n_k}$ be a collection of cycles. In [542], Barrientos and Minion say that a graph G is the *coalescence* of these cycles if for every $2 \leq i \leq k$, the first t_i vertices of C_{n_i} are identified with the last t_i vertices of $C_{n_{i-1}}$, where $t_i \leq n_i/2$. They proved that the coalescence of these cycles admits an α-labeling when each $n_i \equiv 0(\mod 4)$.

Lee and Liu [2163] investigated the mirror graph $M(m,n)$ of $K_{m,n}$ (see §2.3 for the definition) for α-labelings. They proved: $M(m,n)$ has an α-labeling when n is odd or m is even; $M(1,n)$ has an α-labeling when $n \equiv 0 \pmod 4$; $M(m,n)$ does not have an α-labeling when m is odd and $n \equiv 2 \pmod 4$, or when $m \equiv 3 \pmod 4$ and $n \equiv 4 \pmod 8$.

Kumar, Mishra, Kumar, and Kumar [2060] proved that the following graphs have alpha labelings: $C_{4n} \odot K_1$, the graph obtained by joining any path to a vertex of C_{4n}, and graphs obtained by joining two isomorphic copies of $C_{4n} \odot K_1$.

Barrientos and Minion [533] proved that the Cartesian product of two α-trees is an α-tree when both trees admit α-labelings and their stable sets are balanced. (A *stable* set S consists of a set of vertices such that there is not an edge v_iv_j for all pairs v_i, v_j in S). In addition, they present a tree that has the property that when any number of pendent vertices are attached to the vertices of any subset of its smaller stable set the resulting graph is an α-tree. They also prove of an α-labeling of three types of graphs obtained by connecting, sequentially, any number of paths of equal size.

Barrientos [510] defines a *chain graph* as one with blocks $B_1, B_2, \ldots, B_m$ such that for every i, B_i and B_{i+1} have a common vertex in such a way that the block-cutpoint graph is a path. He shows that if $B_1, B_2, \ldots, B_m$ are blocks that have α-labelings then there exists a chain graph G with blocks $B_1, B_2, \ldots, B_m$ that has an α-labeling. He also shows that if $B_1, B_2, \ldots, B_m$ are complete bipartite graphs, then any chain graph G obtained by concatenation of these blocks has an α-labeling.

The *symmetric product* $G_1 \oplus G_2$ of G_1 and G_2 is the graph with vertex set $V(G_1) \times V(G_2)$ and edge set $\{(u_1,v_1)(u_2,v_2)\}$ where u_1u_2 is an edge in G_1 or v_1v_2 is an edge in G_2 but not both u_1u_2 is an edge in G_1 and v_1v_2 is an edge in G_2. A *snake* of length $n > 1$ is a packing of n congruent geometrical objects, called *cells*, such that the first and the last cell each has only one neighbor and all $n-2$ cells in between have exactly two neighbors. In [529] Barrientos and Minion define a *snake polyomino* as a snake with square cells. They prove that given two graphs of sizes m and n with α-labelings, the graph that results from the edge amalgamation (identification of two edges) of the edges of weight 1 and n, also has an α-labeling. They use that result to prove the existence of α-labelings of snake polyominoes and hexagonal chains. The result about snake polyominoes partially answers the question of Acharya. In [530], they prove that the third power of a caterpillar admits an α-labeling and that the symmetric product $G \oplus 2K_1$ has an α-labeling when G does. In addition they prove that $G \cup P_m$ is graceful provided that G admits an α-labeling that does not assign the integer $\lambda + 2$ as a label, where λ is its boundary value. They ask if all triangular chains are graceful.

In [535] Barrientos and Minion proved that under certain conditions, the union $C_r \cup G$ of the cycle C_r and a caterpillar G admits a graceful labeling when r is odd, and an

α-labeling when r is even. They also proved the existence of an α-labeling for any tree obtained by connecting with a path of length two the central vertices of G_i and G_{i+1}, where G_i is a caterpillar of diameter $2d$ with bipartite sets A_i and B_i such that $|A_i| = |B_i| + 1$ and A_i contains the vertices of maximum eccentricity in G_i.

Let $T_1, T_2, \ldots, T_s$ be trees. A *chain tree* obtained by identifying, for every $1 \leq i \leq s-1$, a vertex of T_i with a vertex of T_{i+1}. In [536], Barrientos and Minion prove that if every T_i admits an α-labeling, then there exists a chain tree that also admits an α-labeling. Let T be a tree of size n and v be a fixed vertex of T. The tree T_v^{+r} is obtained by connecting, with a path of length r, two copies of T, by identifying the end-point of this path with the vertices v of each copy of T. They give necessary conditions for the existence of an α-labeling for a tree T_v^{+2}, where v is any of the vertices labeled $\lambda, \lambda-1, \ldots, \lambda-\deg(v)-1$ by an α-labeling with boundary value λ that assigns the labels $\lambda+1, \lambda+2, \ldots, \lambda+\deg(v)$ to leaves of T. In addition they proved that T_v^{+4} has an α-labeling if there exists an α-labeling f of T, with boundary value λ, such that $f(v) = \lambda - 1$. In [536], Barrientos and Minion prove the following. The tree $\oplus(T_1, T_2, T_3, T_4)$ obtained by connecting to a new vertex w, the vertices labeled n in T_1 and T_3 and the vertices labeled $n/2$ in T_2 and T_4, where T_i is an α-labeled tree of even size n that has partite sets of cardinality $n/2$ and $n/2+1$. If G is a graph of order m and size n, with $m < n$, that admits an α-labeling, and H is any graceful graph of size $t-1$, then $tG \cup H$ is a graceful graph. For every $m \geq n$, $m \geq 3$, $n \geq 2$, and $t \geq 2$, $tK_{m,n} \cup L_{t-1}$ admits an α-labeling where L_{t-1} is any linear forest of size $t-1$. If G is a graph of order m and size n, with $m < n$, that admits an α-labeling, then $tG \cup L_{t-1}$ also admits an α-labeling when L_{t-1} is a linear forest of size $t-1$. As a consequence of this result they prove that $tG \cup P_t$ admits an α-labeling provided that G does.

Barrientos [540] showed that all lobsters constructed with k copies of any caterpillar of diameter four by connecting the central vertices of all pairs of consecutive copies with an edge have an α-labeling. Additionally, he proved that any chain-tree formed by caterpillars and this type of lobsters admits an α-labeling. Barrientos and Minion [544] say that a tree is *regular* when the cardinalities of its stable sets are equal or differ by one. They prove if S and T are regular trees that admit α-labelings then $S \times T$ also admits an α-labeling. They use this result to prove that $S \times T$ admits a sequential labeling (see Section 4.1) as well as a harmonious labeling. They define a *fence* as the tree obtained by connecting an internal vertex of P_{n_i} with an internal vertex of $P_{n_{i+1}}$ by a path of length l_i for every $1 \leq i \leq t$. They prove the existence of an α-labeling for any fence constructed with t copies of P_n, where $l_i = 2$. They define a 2-*link fence* as the graph obtained by connecting with an edge, two vertices of the ith copy of P_n, with the corresponding two vertices of the $(i+1)$th copy of P_n. They prove that all such graphs admit α-labelings. In [540] Barrientos says that a fence is *irregular* if two consecutive copies of P_n are connected by one or two pairs of corresponding vertices. He proved that all irregular fences have an α-labeling provided that all their Eulerian subgraphs have size divisible by four. In [545] Barrientos and Minion study subfamilies of 2-link fences, a subfamily of column-convex polyominoes, and a subfamily of irregular cyclic-snakes. They prove that under certain conditions, an α-labelings of these graphs can be transformed into harmonious labelings.

Barrientos and Minion [547] provided new families of harmoniously labeled graphs built on α-labeled tress. Among them are ${P_n}^k$, the join of G and tK_1 where G has a restrictive type of harmonious labeling and its order is different of its size by at most one, $K_{m,n} \cup K_{1,m-1}$, and $G \cup T$ where G is a unicyclic graph and T is a tree built with α-trees. They also showed that almost all trees admit harmonious labelings.

In [539] Barrientos and Minion extend the concept of vertex amalgamation as follows. The *k-vertex amalgamation* of G_1 and G_2 is the graph obtained by identifying k independent vertices of G_1 with k independent vertices of G_2. A *t-fold* of a graph G is obtained using t-copies of G, where the ith copy of G is k-vertex amalgamated with the $(i+1)$th copy of G. They prove that if G admits an α-labeling, then any t-fold of G admits an α-labeling. They consider a more general version of this construction for the case where G is a tree. They also introduce a new family of trees that admit α-labelings; in particular, they prove that any tree of diameter $2n$ formed by identifying the end-vertices of four caterpillars admits an α-labeling.

Fronček, Kingston, and Vezina [1136] generalized snake polyomino graphs by introducing *straight simple polynonial caterpillars* and proving that they also admit an alpha labeling. This implies that every straight simple polymonial caterpillar with n edges decomposes the complete graph K_{2kn+1} for any positive integer k. In [1128] Fronček introduced a similar family of graphs called *full hexagonal caterpillars* and prove that they admit an alpha labeling. This implies that every full hexagonal caterpillar with n edges decomposes the complete graph K_{2kn+1} for any positive integer k.

Golomb [1288] introduced polyominoes in 1953 in a talk to the Harvard Mathematics Club. *Polyominoes* are planar shapes made by connecting a certain number of equal-sized squares, each joined together with at least one other square along an edge.

A graph $G = (V(G), E(G))$ is *even graceful* if there exists an injection f from the set of vertices $V(G)$ to $\{0, 1, 2, 3, 4, \ldots, 2|E(G)|\}$ such that when each edge uv is assigned the label $|f(u) - f(v)|$, the resulting edge labels are $2, 4, 6, \ldots, 2|E(G)|$. Elsonbaty and Mohamed [1020] use even graceful labelings to give a new proof for necessary and sufficient conditions for the gracefulness of cycles. They extend this technique to odd graceful (in the obvious way) and super Fibonacci graceful labelings of cycle graphs (see §3.3). The *polar grid* graph $P_{m,n}$ consists of n copies of C_m numbered from the inner most cycle to the outer cycle as $C^(1)_m, \ldots, C^(n)_m$ and m copies of paths P_{n+1} intersected at the center vertex v_0 numbered as $P^(1)_{n+1}, \ldots, P^(m)_{n+1}$ In [1021] Elsonbaty and Daoud provided edge even graceful labelings for various classes of $P_m \times C_n$. El Dean [4305] obtained an edge even graceful labeling for Y-trees, double stars $B_{n,m}$, $\langle K_{1,2n} : K_{1,2m} \rangle$, $P_{2n-1} \odot \overline{K_{2m}}$, $\overline{K_2} + P_n$, the cycle $v_1, v_2, \ldots, v_{2n}$ with a chord from v_1 to v_n, $P_2 \odot C_n$, flags, and flowers. Zeen El Deen and Omar [4311] gave sufficient conditions for $K_{m,n}$ to have an edge even graceful labeling. They also provided edge even graceful labelings of the join of K_1 with stars, wheels, and sunflowers, and the join of $\overline{K_2}$ with stars and wheels. For results on Fibonacci trees see [1982]. Fox, Bonney, and Patten [1106] constructed odd graceful labelings of $C_n \times P_2$ for the cases of $n = 6k, 6k+2$ and $6k+4$. [1106] new

In [2281] Liu, Liu, Yao, Ma and Lian gave the following definitions. Let G_i be a simple, connected and bipartite graph with bipartition (X_i, Y_i), where $X_i = \{x_{i,1}, x_{i,2}, \ldots, x_{i,t}\}$ [2281] new

and $Y_i = \{y_{i,1}, y_{i,2}, \ldots, y_{i,n_i}\}$ for i in [1,m]. Identify the two vertices y_{i,n_i-k_i+j} and $y_{i+1,j}$ as one vertex for every i in $[1, k_i]$ and for each i in $[1, m-1]$ with $1 \leq k_i \leq \min\{n_i, n_{i+1}\}$. Denote the resulting graph by $\widehat{G_iG_{i+1}}(m)$, called a *dragon graph*. If $k_s \neq k_r$, for some distinct s, r in [1,m], $\widehat{G_iG_{i+1}}(m)$ is called an *irregular dragon graph*. They showed that the dragon graphs $K_{t_i,n_i}\widehat{K_{t_{i+1},n_i}}(m)$ admit odd-graceful labelings and that for any given positive integer q, there exists a dragon graph of q edges that admits an odd graceful labeling.

In [932] Devakirubanithi and Jeba Jesintha showed that twig diamond graphs without prime edge coupled with pendent edges, super subdivision of combs, and bistars are odd graceful. The *converse skew product* $G \bigtriangledown H$ has vertex set $V(G) \times V(H)$ and $u = (x_1, y_1)$ is adjacent to $v = (x_2, y_2)$ whenever $y_1 = y_2$ and $x_1x_2 \in E(G)$ or $x_1x_2 \in E(G)$ and $y_1y_2 \in E(H)$. Hafez, El-Shanawany, and El Atik [1352] proved that if G has an odd graceful labeling f such that G doesn't contain a path $P_3 = x_1x_2x_3$ where $f(x_1) < f(x_2) < f(x_3)$, then $G \bigtriangledown K(n)$ and $Spl_m(G)$ are odd graceful. (Hafez et al. defines $K(n)$ as $\{K_s(n), s = 1, 2, \ldots\}$, where $K_s(n)$ is the graph obtained from P_n by joining the end vertices of the edge v_iv_{i+1} to every vertex in the complete graph $K_{r_i}, i = 1, 2, \ldots, n-1$, of order r_i, such that (a) if $|V(K_{r_i}) \cap V(K_{r_{i+1}})| = t$, then $t \leq \min\{r_i, r_{i+1}\}$ and (b) K_{r_i} and K_{r_j} are disjoint when $|i - j| > 2$). Jesintha, Devasagayam, and Sherlin [1606] prove that the super subdivision of a star and the super subdivision of a path admit odd graceful labelings. In [4041] Velankanni, Raj, and Sujasree proved that the graphs obtained by the arbitrary supersubdivision of paths, caterpillars, and shell graphs are odd graceful. Jesintha and Hilda [1619] proved that all subdivided shell flower graphs are odd graceful. In [1620] they proved that the path union of 2 subdivided shell graphs of any order is odd graceful. Jesintha, Devakirubanithi, and Angel [1602] proved that a stalked subdivided shell graph joined by an edge is odd graceful. (A *stalked subdivided shell* graph is obtained by adding an edge at the apex vertex of the subdivided shell graph.) Abdel-Aal, NourEldeen, Badr, and Shabana [6] gave odd graceful labelings of the linear kC_4-snakes $\odot mK_1$, kC_8-snakes $\odot mK_1$, kC_{12}-snakes $\odot mK_1$, and kC_n-snakes $\odot mK_1$, where $n \equiv 0 \pmod 4$. In addition, they introduced a symmetric 3-layered encryption scheme in which the structure of an odd gracefully labeled graph provides an additional layer of obfuscation over a classical Trifid cipher.

[1619] new

[1620] new

[1602] new

[6] new

Wu ([4215] and [4217]) has given a number of methods for constructing larger graceful graphs from graceful graphs. Let $G_1, G_2, \ldots, G_p$ be disjoint connected graphs. Let w_i be in G_i for $1 \leq i \leq p$. Let w be a new vertex not in any G_i. Form a new graph $\oplus_w(G_1, G_2, \ldots, G_p)$ by adjoining to the graph $G_1 \cup G_2 \cup \cdots \cup G_p$ the edges $ww_1, ww_2, \ldots, ww_p$. In the case where each of $G_1, G_2, \ldots, G_p$ is isomorphic to a graph G that has an α-labeling and each w_i is the isomorphic image of the same vertex in G_i, Wu shows that the resulting graph is graceful. If f is an α-labeling of a graph, the integer k with the property that for any edge uv either $f(u) \leq k < f(v)$ or $f(v) \leq k < f(u)$ is called the *boundary value* or *critical number* of f. Wu [4215] has also shown that if $G_1, G_2, \ldots, G_p$ are graphs of the same order and have α-labelings where the labelings for each pair of graphs G_i and G_{p-i+1} have the same boundary value for $1 \leq i \leq n/2$, then $\oplus_w(G_1, G_2, \ldots, G_p)$ is graceful. In [4213] Wu proves that if G has n edges and $n + 1$

vertices and G has an α-labeling with boundary value λ, where $|n - 2\lambda - 1| \leq 1$, then $G \times P_m$ is graceful for all m.

Given graceful graphs H and G with at least one having an α-labeling Wu and Lu [4218] define four graph operations on H and G that when used repeatedly or in turns provide a large number of graceful graphs. In particular, if both H and G have α-labelings, then each of the graphs obtained by the four operations on H and G has an α-labeling.

Ajitha, Arumugan, and Germina [154] use a construction of Koh, Tan, and Rogers [1988] to create trees with α-labelings from smaller trees with graceful labelings. These in turn allows them to generate large classes of trees that have a type of called edge-antimagic labelings (see §6.1). Shiue and Lu [3562] prove that the graph obtained from $K_{1,k}$ by replacing each edge with a path of length 3 has an α-labeling if and only if $k \leq 4$. In [4078] Venkatesh and Bharathi recursively construct new trees starting with caterpillars that admit α-labelings.

Seoud and Helmi [3385] have shown that all gear graphs have an α-labeling, all dragons with a cycle of order $n \equiv 0 \pmod 4$ have an α-labeling, and the graphs obtained by identifying an endpoint of a star S_m ($m \geq 3$) with a vertex of C_{4n} has an α-labeling.

Mavonicolas and Michael [2459] say that trees $\langle T_1, \theta_1, w_1\rangle$ and $\langle T_2, \theta_2, w_2\rangle$ with roots w_1 and w_2 and $|V(T_1)| = |V(T_2)|$ are *gracefully consistent* if either they are identical or they have α-labelings with the same boundary value and $\theta_1(w_1) = \theta_2(w_2)$. They use this concept to show that a number of known constructions of new graceful trees using several identical copies of a given graceful rooted tree can be extended to the case where the copies are replaced by a set of pairwise gracefully consistent trees. In particular, let $\langle T, \theta, w\rangle$ and $\langle T_0, \theta_0, w_0\rangle$ be gracefully labeled trees rooted at w and w_0 respectively. They show that the following four constructions are adaptable to the case when a set of copies of $\langle T, \theta, w\rangle$ is replaced by a set of pairwise gracefully consistent trees. When $\theta(w) = |E(T)|$ the garland construction due to Koh, Rogers, and Tan [1981] gracefully labels the tree consisting of h copies of $\langle T, w\rangle$ with their roots connected to a new vertex r. In the case when $\theta(w) = |E(T)|$ and whenever $uw \in E(T)$ and $\theta(u) \neq 0$, then $vw \in E(T)$ where $\theta(u) + \theta(v) = |E(T)|$, the attachment construction of Koh, Tan and Rogers [1988] gracefully labels the tree formed by identifying the roots of h copies of $\langle T, w\rangle$. A construction given by Koh, Tan and Rogers [1988] gracefully labels the tree formed by merging each vertex of $\langle T_0, w_0\rangle$ with the root of a distinct copy of $\langle T, w\rangle$. When $\theta_0(w_0) = |E(T_0)|$, let N be the set of neighbors of w_0 and let x be the vertex of T at even distance from w with $\theta(x) = 0$ or $\theta(x) = |E(T)|$. Then a construction of Burzio and Ferrarese [731] gracefully labels the tree formed by merging each non-root vertex of T_0 with the root of a distinct copy of $\langle T, w\rangle$ so that for each $v \in N$ the edge vw_0 is replaced with a new edge xw_0 (where x is in the corresponding copy of T).

Snevily [3645] says that a graph G *eventually has an α-labeling* provided that there is a graph H, called a *host* of G, which has an α-labeling and that the edge set of H can be partitioned into subgraphs isomorphic to G. He defines the *α-labeling number of G* to be $G_\alpha = \min\{t :$ there is a host H of G with $|E(H)| = t|G|\}$. Snevily proved that even cycles have α-labeling number at most 2 and he conjectured that every bipartite graph has an α-labeling number. This conjecture was proved by El-Zanati, Fu, and Shiue [1028].

There are no known examples of a graph G with $G_\alpha > 2$. In [3645] Snevily conjectured that the α-labeling number for a tree with n edges is at most n. Shiue and Fu [3560] proved that the α-labeling number for a tree with n edges and radius r is at most $\lceil r/2 \rceil n$. They also prove that a tree with n edges and radius r decomposes K_t for some $t \leq (r+1)n^2+1$.

Ahmed and Snevily [145] investigated the claim that for every tree T there exists an α-labeling of T, or else there exists a graph H_T with an α-labeling such that H_T can be decomposed into two edge-disjoint copies of T. They proved this claim is true for the graphs $C_{m,k}$ obtained from $K_{1,m}$ by replacing each edge in $K_{1,m}$ with a path of length k.

A graph G with vertex set V and edge set E is called *super edge-graceful* if there is a bijection f from E to $\{0, \pm 1, \pm 2, \ldots, \pm(|E|-1)/2\}$ when $|E|$ is odd and from E to $\{\pm 1, \pm 2, \ldots, \pm|E|/2\}$ when $|E|$ is even such that the induced vertex labeling f^* defined by $f^*(u) = \sum f(uv)$ over all edges uv is a bijection from V to $\{0, \pm 1, \pm 2, \ldots, \pm(|V|-1)/2\}$ when $|V|$ is odd and from V to $\{\pm 1, \pm 2, \ldots, \pm|V|/2\}$ when $|V|$ is even. Clifton and Khodkar [856] proved that graphs formed by identifying the endpoint of a path P_n and a vertex of a cycle (kites) with $n \geq 5$ vertices, $n \neq 6$ are super edge-graceful. Khodkar, Nolen, and Perconti [1952] proved that all complete bipartite graphs except for $K_{2,2}, K_{2,3}$, and $K_{1,n}$ (n odd) are super edge-graceful. Khodkar [1955] and [1953] proved that all complete tripartite graphs except $K_{1,1,2}$ are super edge-graceful and that the union of vertex disjoint 3-cycles is super edge-graceful. Lee, Su, and Wei [2208] provide a family of trees of odd orders which are super edge-graceful.

For a tree T with m edges, the *α-deficit* $\alpha_{def}(T)$ equals $m - \alpha(T)$ where $\alpha(T)$ is defined as the maximum number of distinct edge labels over all bipartite labelings of T. Rosa and Siran [3206] showed that for every $m \geq 1$, $\alpha_{def}(C_{m,2}) = \lfloor m/3 \rfloor$, which implies that $(C_{m,2})_\alpha \geq 2$ for $m \geq 3$. Ahmed and Snevily [145] define the graph $C'_{m,j}$ as a comet-like tree with a central vertex of degree m where each neighbor of the central vertex is attached to j pendent vertices for $1 \leq j \leq (m-1)$. For $m \geq 3$ and $1 \leq j \leq (m-1)$ they prove: $(C'_{m,j})_\alpha \leq 2$; $(C'_{2k+1,j})_\alpha = 2$ for $1 \leq j \leq 2k$ and conjecture if $\Delta_T = (2k+1)$, then $\alpha_{def}(T) \leq k$. Ahmed and Snevily [145] prove that for every comet T (that is, graphs obtained from stars by replacing each edge by a path of some fixed length) there exists an α-labeling of T, or else there exists a graph H_T with an α-labeling such that H_T can be decomposed into two edge-disjoint copies of T. This is particularly noteworthy since comets are known to have arbitrarily large α-deficits.

Given two bipartite graphs G_1 and G_2 with partite sets H_1 and L_1 and H_2 and L_2, respectively, Snevily [3644] defines their *weak tensor product* $G_1 \overline{\bigotimes} G_2$ as the bipartite graph with vertex set $(H_1 \times H_2, L_1 \times L_2)$ and with edge $(h_1, h_2)(l_1, l_2)$ if $h_1 l_1 \in E(G_1)$ and $h_2 l_2 \in E(G_2)$. He proves that if G_1 and G_2 have α-labelings then so does $G_1 \overline{\bigotimes} G_2$. This result considerably enlarges the class of graphs known to have α-labelings. In [2307] López and Muntaner-Batle gave a generalization of Snevily's weak tensor product that allows them to significantly enlarges the classes of graphs admitting α-labelings, near α-labelings (defined later in this section), and bigraceful graphs.

The sequential join of graphs $G_1, G_2, \ldots, G_n$ is formed from $G_1 \cup G_2 \cup \cdots \cup G_n$ by adding edges joining each vertex of G_i with each vertex of G_{i+1} for $1 \leq i \leq n-1$. Lee and Wang [2217] have shown that for all $n \geq 2$ and any positive integers $a_1, a_2, \ldots, a_n$

the sequential join of the graphs $\overline{K}_{a_1}, \overline{K}_{a_2}, \ldots, \overline{K}_{a_n}$ has an α-labeling.

In [1169] Gallian and Ropp conjectured that every graph obtained by adding a single pendent edge to one or more vertices of a cycle is graceful. Qian [3063] proved this conjecture and in the case that the cycle is even he shows the graphs have an α-labeling. He further proves that for n even any graph obtained from an n-cycle by adding one or more pendent edges at some vertices has an α-labeling as long as at least one vertex has degree 3 and one vertex has degree 2.

In [2788] Pasotti introduced the following generalization of a graceful labeling. Given a graph G with $e = dm$ edges, an injective function from $V(\Gamma)$ to the set $\{0, 1, 2, \ldots, d(m+1)-1\}$ such that $\{|f(x) - f(y)| \mid [x, y] \in E(\Gamma)\} = \{1, 2, 3, \ldots, d(m+1)-1\} - \{m+1, 2(m+1), \ldots, (d-1)(m+1)\}$ is called a *d-divisible graceful labeling* of G. Note that for $d = 1$ and of $d = e$ one obtains the classical notion of a graceful labeling and of an odd-graceful labeling (see §3.6 for the definition), respectively. A d-divisible graceful labeling of a bipartite graph G with the property that the maximum value on one of the two bipartite sets is less than the minimum value on the other one is called a *d-divisible α-labeling* of G. Pasotti proved that these new concepts allow to obtain certain cyclic graph decompositions. In particular, if there exists a d-divisible graceful labeling of a graph G of size $e = dm$ then there exists a cyclic G-decomposition of $K_{\left(\frac{e}{d}+1\right)\times 2d}$ and that if there exists a d-divisible α-labeling of a graph Γ of size e then there exists a cyclic G-decomposition of $K_{\left(\frac{e}{d}+1\right)\times 2dn}$ for any integer $n \geq 1$. She also it is proved the following: paths and stars admit a d-divisible α-labeling for any admissible d; C_{4k} admits a 2-divisible α-labeling and a 4-divisible α-labeling for any $k \geq 1$; C_{2k} admits a 2-divisible labeling for any odd integer $k > 1$; and the ladder graph L_{2k} has a 2-divisible α-labeling if and only if k is even.

Pasotti [2788] generalized the notion of graceful labelings for graphs G with $e = d \cdot m$ edges by defining a *d-graceful labeling* as an injective function f from $V(G)$ to $\{0, 1, 2, \ldots, d(m+1)-1\}$ such that $\{|f(x) - f(y)| \mid xy \in E(G)\} = \{1, 2, \ldots, d(m+1)-1\} - \{m+1, 2(m+1), \ldots, (d-1)(m+1)\}$. The case $d = 1$ is a graceful labeling and the case that $d = e$ is an odd-graceful labeling. A *d-graceful* α-labeling of a bipartite graph is a d-graceful labeling with the property that the maximum value in one of the two bipartite sets is less than the minimum value on the other bipartite set. Pasotti [2788] proved that paths and stars have d-graceful α-labelings for all admissible d, ladders $P_n \times P_2$ have a 2-graceful labeling if and only if n is even, and provided partial results about cycles of even length. He showed that the existence of d-graceful labelings can be used to prove that certain complete graphs have cyclic decompositions. Benini and Pasotti [616] used d-divisible α-labelings to construct an infinite class of cyclic Γ-decompositions of the complete multipartite graphs, where Γ is a caterpillar, a hairy cycle or a cycle. Such labelings imply the existence of cyclic Γ-decompositions of certain complete multipartite graphs.

In [2787], Pasotti proved the existence of d-divisible α-labelings for $C_{4k} \times P_m$ for any integers $k \geq 1$, $m \geq 2$ for $d = 2m-1,\ 2(2m-1)$ and $4(2m-1)$. Benini and Pasotti [617] proved that the generalized Petersen graph $P_{8n,3}$ admits an α-labeling for any integer $n \geq 1$ confirming that the conjecture posed by A. Vietri in [4068] is true.

For any tree $T(V,E)$ whose vertices are properly 2-colored Rosa and Širáň [3206] define a *bipartite labeling* of T as a bijection $f : V \to \{0,1,2,\ldots,|E|\}$ for which there is a k such that whenever $f(u) \le k \le f(v)$, then u and v have different colors. They define the *α-size* of a tree T as the maximum number of distinct values of the induced edge labels $|f(u) - f(v)|, uv \in E$, taken over all bipartite labelings f of T. They prove that the α-size of any tree with n edges is at least $5(n+1)/7$ and that there exist trees whose α-size is at most $(5n+9)/6$. They conjectured that minimum of the α-sizes over all trees with n edges is asymptotically $5n/6$. This conjecture has been proved for trees of maximum degree 3 by Bonnington and Širáň [687]. For trees with n vertices and maximum degree 3 Brankovic, Rosa, and Širáň [705] have shown that the α-size is at least $\lfloor \frac{6n}{7} \rfloor - 1$. In [703] Brankovic, Murch, Pond, and Rose provide a lower bound for the α-size trees with maximum degree three and a perfect matching as a function of a lower bound for minimum order of such a tree that does not have an α-labeling. Using a computer search they showed that all such trees on less than 30 vertices have an α-labeling. This brought the lower bound for the α-size to $14n/15$, for such trees of order n. They conjecture that all trees with maximum degree three and a perfect matching have an α-labeling. Heinrich and Hell [1431] defined the *gracesize* of a graph G with n vertices as the maximum, over all bijections $f\colon V(G) \to \{1,2,\ldots,n\}$, of the number of distinct values $|f(u) - f(v)|$ over all edges uv of G. So, from Rosa and Širáň's result, the gracesize of any tree with n edges is at least $5(n+1)/7$.

In [711] Brinkmann, Crevals, Mélot, Rylands, and Steffan define the parameter α_{def} which measures how far a tree is from having an α-labeling as it counts the minimum number of errors, that is, the minimum number of edge labels that are missing from the set of all possible labels. Trees with an α-labeling have deficit 0. For a tree $T = (V,E)$ with bipartition classes V_1 and V_2 and a bipartite labeling $f : V \to \{0,\ldots,|V|-1\}$ the *edge parity* of T is $(\sum_{i=1}^{|E|} i) \bmod 2 = \frac{1}{2}(|V|-1)|V| \bmod 2$. So if f is an α-labeling this is the sum of all edge labels modulo 2; it is 0 if $|V| \equiv 0, 1 \bmod 4$ and 1 if $|V| \equiv 2, 3 \bmod 4$. The *vertex parity* is the parity of the number of vertices of odd degree with odd label.

Brinkmann et al. [711] proved: in a tree T with α-deficit 0 the edge parity and the vertex parities are equal; and for all non-negative integers k and d and $n \ge k^2 + k$, the number of trees T with n vertices, $\alpha_{\text{def}}(T) = d$ and maximum degree $n-k$ is the same. Furthermore, they provide computer results on the α-deficit of all trees with up to 26 vertices; with maximum degree 3 and up to 36 vertices, with maximum degree 4 and up to 32 vertices, and with maximum degree 5 and up to 31 vertices.

In [1174] Gallian weakened the condition for an α-labeling somewhat by defining a *weakly α-labeling* as a graceful labeling for which there is an integer k so that for each edge xy either $f(x) \le k \le f(y)$ or $f(y) \le k \le f(x)$. Unlike α-labelings, this condition allows the graph to have an odd cycle, but still places a severe restriction on the structure of the graph; namely, that the vertex with the label k must be on every odd cycle. Gallian, Prout, and Winters [1174] showed that the prisms $C_n \times P_2$ with a vertex deleted have α-labelings. The same paper reveals that $C_n \times P_2$ with an edge deleted from a cycle has an α-labeling when n is even and a weakly α-labeling when $n > 3$.

In [531] and [534] Barrientos and Minion focused on the enumeration of graphs with

graceful and α-labelings, respectively. They used an extended version of the adjacency matrix of a graph to count the number of labeled graphs. In [531] they count the number of gracefully-labeled graphs of size n and order m, for all possible values of m. In [534] they count the number of α-labeled graphs of size n and order m, for all possible values of m, as well as those α-labeled graphs of size n with boundary value λ. They also count the number of α-labeled graphs of size n, order m, and boundary value λ for all possible values of m and λ.

A special case of α-labeling called strongly graceful was introduced by Maheo [2386] in 1980. A graceful labeling f of a graph G is called *strongly graceful* if G is bipartite with two partite sets A and B of the same order s, the number of edges is $2t + s$, there is an integer k with $t - s \leq k \leq t + s - 1$ such that if $a \in A$, $f(a) \leq k$, and if $b \in B$, $f(b) > k$, and there is an involution π that is an automorphism of G such that: π exchanges A and B and the s edges $a\pi(a)$ where $a \in A$ have as labels the integers between $t + 1$ and $t + s$. Maheo's main result is that if G is strongly graceful then so is $G \times Q_n$. In particular, she proved that $(P_n \times Q_n) \times K_2$, B_{2n}, and $B_{2n} \times Q_n$ have strongly graceful labelings.

In 1999 Broersma and Hoede [712] conjectured that every tree containing a perfect matching is strongly graceful. Yao, Cheng, Yao, and Zhao [4258] proved that this conjecture is true for every tree with diameter at most 5 and provided a method for constructing strongly graceful trees.

El-Zanati and Vanden Eynden [1032] call a strongly graceful labeling a *strong α-labeling*. They show that if G has a strong α-labeling, then $G \times P_n$ has an α-labeling. They show that $K_{m,2} \times K_2$ has a strong α-labeling and that $K_{m,2} \times P_n$ has an α-labeling. They also show that if G is a bipartite graph with one more vertex than the number of edges, and if G has an α-labeling such that the cardinalities of the sets of the corresponding bipartition of the vertices differ by at most 1, then $G \times K_2$ has a strong α-labeling and $G \times P_n$ has an α-labeling. El-Zanati and Vanden Eynden [1032] also note that $K_{3,3} \times K_2$, $K_{3,4} \times K_2$, $K_{4,4} \times K_2$, and $C_{4k} \times K_2$ all have strong α-labelings. El-Zanati and Vanden Eynden proved that $K_{m,2} \times Q_n$ has a strong α-labeling and that $K_{m,2} \times P_n$ has an α-labeling for all n. They also prove that if G is a connected bipartite graph with partite sets of odd order such that in each partite set each vertex has the same degree, then $G \times K_2$ does not have a strong α-labeling. As a corollary they have that $K_{m,n} \times K_2$ does not have a strong α-labeling when m and n are odd.

An α-labeling f of a graph G is called *free* by El-Zanati and Vanden Eynden in [1033] if the critical number k (in the definition of α-labeling) is greater than 2 and if neither 1 nor $k - 1$ is used in the labeling. Their main result is that the union of graphs with free α-labelings has an α-labeling. In particular, they show that $K_{m,n}$, $m > 1$, $n > 2$, has a free α-labeling. They also show that Q_n, $n \geq 3$, and $K_{m,2} \times Q_n$, $m > 1$, $n \geq 1$, have free α-labelings. El-Zanati [personal communication] has shown that the Heawood graph has a free α-labeling.

Wannasit and El-Zanati [4169] proved that if G is a cubic bipartite graph each of whose components is either a prism, a Möbius ladder, or has order at most 14, then G admits free α-labeling. They conjecture that every bipartite cubic graph admits a free α-labeling.

In [2410] Makadia, Karavadiya, and Kaneria call a vertex v in a graph G with a graceful labeling f a *graceful center* of G if $f(v) = 0$ or $f(v) = |E(G)|$. They say a graph G is a *universal graceful* graph if for every $v \in V(G)$, v is a graceful center for G with respect to some graceful labeling of G. They call G a *universal α-graceful* graph if for every $v \in V(G)$, v is a graceful center for G with respect to some α-graceful labeling of G. They define the *ring sum* of two graphs G_1 and G_2 denoted $G_1 \oplus G_2$, as the graph with vertex set $(V(G_1) \cup V(G_2)$ and edge set $E(G_1) \cup E(G_2) - (E(G_1) \cap E(G_2))$. They proved: any graph G that admits α-labeling has at least four graceful centers; if G is a graceful graph, then $G \oplus K_{1,n}$ is graceful; if G is a universal graceful graph, then $G \oplus K_2$ is a graceful; if G_1 is graceful and G_2 has an α-labeling, then the ring sum $G_1 \oplus G_2$ with the graceful center of G_1 and the graceful center of G_2 as a common vertex is a graceful; and if G_1 and G_2 have α labelings, then the ring sum $G_1 \oplus G_2$ with the two graceful centers of G_1 and G_2 as a common vertex has an α labeling.

For connected bipartite graphs Grannell, Griggs, and Holroyd [1328] introduced a labeling that lies between α-labelings and graceful labelings. They call a vertex labeling f of a bipartite graph G with q edges and partite sets D and U *gracious* if f is a bijection from the vertex set of G to $\{0, 1, \ldots, q\}$ such that the set of edge labels induced by $f(u) - f(v)$ for every edge uv with $u \in U$ and $v \in D$ is $\{1, 2, \ldots, q\}$. Thus a gracious labeling of G with partite sets D and U is a graceful labeling in which every vertex in D has a label lower than every adjacent vertex. They verified by computer that every tree of size up to 20 has a gracious labeling. This led them to conjecture that every tree has a gracious labeling. For any $k > 1$ and any tree T Grannell et al. say that T has a *gracious k-labeling* if the vertices of T can be partitioned into sets D and U in such a way that there is a function f from the vertices of G to the integers modulo k such that the edge labels induced by $f(u) - f(v)$ where $u \in U$ and $v \in D$ have the following properties: the number of edges labeled with 0 is one less than the number of vertices labeled with 0 and for each nonzero integer t the number of edges labeled with t is the same as the number of vertices labeled with t. They prove that every nontrivial tree has a k-gracious labeling for $k = 2, 3, 4$, and 5 and that caterpillars are k-gracious for all $k \geq 2$. In [615] Bell and Cummins provided new methods for combining certain families of gracefully labeled graphs to produce new gracefully labeled graphs. If the constituent graphs have a gracious labeling, then the methods presented produce a gracious labeling. They also introduce new infinite families of gracious trees and new classes of graceful trees.

The same labeling that is called gracious by Grannell, Griggs, and Holroyd is called a *near α-labeling* by El-Zanati, Kenig, and Vanden Eynden [1030]. The latter prove that if G is a graph with n edges that has a near α-labeling then there exists a cyclic G-decomposition of K_{2nx+1} for all positive integers x and a cyclic G-decomposition of $K_{n,n}$. They further prove that if G and H have near α-labelings, then so does their weak tensor product (see earlier part of this section) with respect to the corresponding vertex partitions. They conjecture that every tree has a near α-labeling.

In [2384] Mahalingam and Rajendram introduced a new labeling called m-bonacci graceful labeling as follows. A graph G on n edges is *m-bonacci graceful* if the vertices can be labeled with distinct integers from the set of the first n m-bonacci numbers such that

the derived edge labels are the first n m-bonacci numbers. They showed that complete graphs, complete bipartite graphs, gear graphs, triangular grid graphs, and wheel graphs are not m-bonacci graceful. They gave m-bonacci graceful labeling for cycles, friendship graphs, polygonal snake graphs, and double polygonal snake graphs and proved that almost all trees are m-bonacci graceful.

For a simple, finite, connected, undirected, non-trivial graph G Sumathi and Raman [3773] introduced the notion of *arithmetic sequential graceful* as an injection $f : V(G) \to \{a, a+d, a+2d, a+3d, \ldots, 2(a+qd)\}$, where $a \geq 0$ and $d \geq 1$, with the property that $f^* : E(G) \to \{d, 2d, 3d, 4d, \ldots, qd\}$ defined by $f^*(uv) = |f(u) - f(v)|$ is a bijection. They proved that stars, double stars, and some star related graphs are arithmetic sequential graceful.

Another kind of labelings for trees was introduced by Ringel, Llado, and Serra [3177] in an approach to proving their conjecture $K_{n,n}$ is edge-decomposable into n copies of any given tree with n edges. If T is a tree with n edges and partite sets A and B, they define a labeling f from the set of vertices to $\{1, 2, \ldots, n\}$ to be a *bigraceful* labeling of T if f restricted to A is injective, f restricted to B is injective, and the edge labels given by $f(y) - f(x)$ where yx is an edge with y in B and x in A is the set $\{0, 1, 2, \ldots, n-1\}$. (Notice that this terminology conflicts with that given in Section 2.7 In particular, the Ringel, Llado, and Serra bigraceful does not imply the usual graceful.) Among the graphs that they show are bigraceful are: lobsters, trees of diameter at most 5, stars $S_{k,m}$ with k spokes of paths of length m, and complete d-ary trees for d odd. They also prove that if T is a tree then there is a vertex v and a nonnegative integer m such that the addition of m leaves to v results in a bigraceful tree. They conjecture that all trees are bigraceful.

A *pronic number* is one of the form $n(n+1)$, where n is a positive integer. Porchelvi and Devi [2995] defined a *pronic graceful* labeling of a graph G with $n \geq 2$ vertices as a bijection $f : V(G) \to \{0, 2, \ldots, n(n+1)\}$ such that the upon labeling each edge uv with $|f(u) - f(v)|$ the labels are distinct. A graph G is called a *pronic graceful* graph if it admits pronic graceful labeling. They proved that nontrivial cases of paths, cycles, wheels, stars, twigs, and generalized Peterson graphs are pronic graceful. [942] they proved that the generalized Peterson graphs $P(6,2), P(8,3), (10,2), P(10,3)$, and $P(12,5)$ are pronic graceful. In [975], Dorathy and Nagamani proved that C_n, $K_{1,n}$ $(n \geq 3)$, and P_n $(n \geq 3)$, helms, ladders, Möbius ladders, shells $C(n, n-3)$ $(n \geq 3)$, and shell butterflies admit pronic graceful labelings, whereas K_n $(n \geq 4)$ does not admit pronic graceful labelings. [975] new

Table 3 summarizes some of the main results about α-labelings; α indicates that the graphs have an α-labeling.

Table 3: **Summary of Results on α-labelings**

Graph	*α-labeling*
cycles C_n	α iff $n \equiv 0 \pmod 4$ [3203]
caterpillars	α [3203]
n-cube	α [2012]
books $B_{2n},\ B_{4n+1}$	α [2399], [1173]
Möbius ladders M_{2k+1}	α [2786]
$C_m \cup C_n$	α iff m, n are even and $m+n \equiv 0 \pmod 4$ [30]
$C_{4m} \cup C_{4m} \cup C_{4m}$ $(m > 1)$	α [2014]
$C_{4m} \cup C_{4m} \cup C_{4m} \cup C_{4m}$	α [2014]
$mK_{s,t}$ $(m \geq 1, s, t \geq 2)$	iff $(m, s, t) \neq (3, 2, 2)$ [1515]
$P_n \times Q_n$	α [2399]
$B_{2n} \times Q_n$	α [2399]
$K_{1,n} \times Q_n$	α [2399]
$K_{m,2} \times Q_n$	α [1032]
$K_{m,2} \times P_n$	α [1032]
$P_2 \times P_2 \times \cdots \times P_2 \times G$	α when $G = C_{4m},\ P_m,\ K_{3,3},$ $K_{4,4}$ [3644]
$P_2 \times P_2 \times \cdots \times P_2 \times P_m$	α [3644]
$P_2 \times P_2 \times \cdots \times P_2 \times K_{m,m}$	α [3644] when $m = 3$ or 4
$G[\overline{K_n}]$	α when G is α [3645]

3.2 γ-Labelings

In 2004 Chartrand, Erwin, VanderJagt, and Zhang [777] define a γ-labeling of a graph G of size m as a 1-1 function f from the vertices of G to $\{0, 1, 2, \ldots, m\}$ that induces an edge labeling f' defined by $f'(uv) = |f(u)-f(v)|$ for each edge uv. They define the following parameters of a γ-labeling: $\text{val}(f) = \Sigma f'(e)$ over all edges e of G; $\text{val}_{\max}(G) = \max\{\ \text{val}(f) : f \text{ is a } \gamma\text{-labeling of } G\}$, $\text{val}_{\min}(G) = \min\{\ \text{val}(f)\ :\ f \text{ is a } \gamma - \text{labeling of } G\}$. Among their results are the following:
$\text{val}_{\min}(P_n) = \text{val}_{\max}(P_n) = \lfloor (n^2-2)/2 \rfloor$; $\text{val}_{\min}(C_n) = 2(n-1)$; for even $n \geq 4$, $\text{val}_{\max}(C_n) = n(n+2)/2$; for odd $n \geq 3$, $\text{val}_{\max}(C_n) = (n-1)(n+3)/2$; for odd n, $\text{val}_{\min}(K_n) = \binom{n+1}{3}$; for odd n, $\text{val}_{\max}(K_n) = (n^2-1)(3n^2-5n+6)/24$; for even n, $\text{val}_{\max}(K_n) = n(3n^3-5n^2+6n-4)/24$; for every $n \geq 3$, $\text{val}_{\min}(K_{1,n-1}) = \binom{\lfloor \frac{n+1}{2} \rfloor}{2} + \binom{\lceil \frac{n+1}{2} \rceil}{2}$; $\text{val}_{\max}(K_{1,n-1}) = \binom{n}{2}$ for a connected graph of order n and size m, $\text{val}_{\min}(G) = m$ if and only if G is isomorphic to P_n; if G is maximal outerplanar of order $n \geq 2$, $\text{val}_{\min}(G) \geq 3n-5$ and equality occurs if and only if $G = P_n^2$; if G is a connected r-regular bipartite graph of order n and size m where $r \geq 2$, then $\text{val}_{\max}(G) = rn(2m-n+2)/4$.

In another paper on γ-labelings of trees Chartrand, Erwin, VanderJagt, and Zhang [778] prove for $p, q \geq 2$, $\text{val}_{\min}(S_{p,q})$ (that is, the graph obtained by joining the centers of $K_{1,p}$ and $K_{1,q}$ by an edge)$= (\lfloor p/2 \rfloor+1)^2+(\lfloor q/2 \rfloor+1)^2-(n_p\lfloor p/2 \rfloor+1)^2+(n_q\lfloor (q+2)/2 \rfloor+1)^2)$, where n_i is 1 if i is even and n_i is 0 if n_i is odd; $\text{val}_{\min}(S_{p,q}) = (p^2+q^2+4pq-3p-3q+2)/2$; for a connected graph G of order n at least 4, $\text{val}_{\min}(G) = n$ if and only if G is a caterpillar with maximum degree 3 and has a unique vertex of degree 3; for a tree T of order n at least 4, maximum degree Δ, and diameter d, $\text{val}_{\min}(T) \geq (8n+\Delta^2-6\Delta-4d+\delta_\Delta)/4$ where δ_Δ is 0 if Δ is even and δ_Δ is 0 if Δ is odd. They also give a characterization of all trees of order n at least 5 whose minimum value is $n+1$.

Saduakdee and Khemmani [3282] investigated connected graphs having the unique γ-min labeling. They determined the minimum value of a γ-labeling for some classes of trees and showed that they have no unique γ-min labeling.

In [728] Buratti and Del Fra solved the existence problem for cyclic k-cycle systems of the complete graph K_v with $v \equiv 1 \pmod{2k}$, and the existence problem for cyclic k-cycle systems of the complete m-partite graph $K_{m\times k}$ for m and k odd. As a particular consequence, a cyclic p-cycle system of K_v with p a prime exists for all admissible values of v but $(p, v) \neq (3, 9)$. This was previously known only for $p = 3, 5, 7$.

In [3276] Sanaka determined $\text{val}_{\max}(K_{m,n})$ and $\text{val}_{\min}(K_{m,n})$. In [727] Bunge, Chantasartraaamee, El-Zanati, and Vanden Eynden generalized γ-labelings by introducing two labelings for tripartite graphs. Graphs G that admit either of these labelings guarantee the existence of cyclic G-decompositions of K_{2nx+1} for all positive integers x. They also proved that, except for $C_3 \cup C_3$, the disjoint union of two cycles of odd length admits one of these labelings.

3.3 Graceful-like Labelings

As a means of attacking graph decomposition problems, Rosa [3203] invented another analogue of graceful labelings by permitting the vertices of a graph with q edges to assume labels from the set $\{0, 1, \ldots, q+1\}$, while the edge labels induced by the absolute value of the difference of the vertex labels are $\{1, 2, \ldots, q-1, q\}$ or $\{1, 2, \ldots, q-1, q+1\}$. He calls these *$\hat{\rho}$-labelings*. Frucht [1152] used the term *nearly graceful labeling* instead of *$\hat{\rho}$-labelings*. Frucht [1152] has shown that the following graphs have nearly graceful labelings with edge labels from $\{1, 2, \ldots, q-1, q+1\}$: $P_m \cup P_n$; $S_m \cup S_n$; $S_m \cup P_n$; $G \cup K_2$ where G is graceful; and $C_3 \cup K_2 \cup S_m$ where m is even or $m \equiv 3 \pmod{14}$. Seoud and Elsakhawi [3378] have shown that all cycles are nearly graceful. Barrientos [508] proved that C_n is nearly graceful with edge labels $1, 2, \ldots, n-1, n+1$ if and only if $n \equiv 1$ or $2 \pmod 4$. Nurvazly and Sugeng [2728] proved that $\Theta(C_3)^n$ graphs (n copies of C_3 that share an edge) have $\hat{\rho}$ labelings. Gao [1189] shows that a variation of banana trees is odd-graceful and in some cases has a nearly graceful labeling. (A graph G with q edges is *odd-graceful* if there is an injection f from $V(G)$ to $\{0, 1, 2, \ldots, 2q-1\}$ such that, when each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edge labels are $\{1, 3, 5, \ldots, 2q-1\}$).

For a graph G with p vertices and q directed edges that are assigned distinct vertex labels in $\{0, \ldots, q\}$ and distinct edge labels in $\{1, \ldots, p\}$ so that the label of the directed edge from u to v is $(f(v) - f(u)) \bmod (q+1)$ (this generalizes Rosa's ρ-valuations. Knuth [1971] has observed that there is a nice data structure for storing a graph or digraph in a computer. He calls this a *graceful data structure* labeling.

In 1988 Rosa [3205] conjectured that triangular snakes with $t \equiv 0$ or $1 \pmod 4$ blocks are graceful and those with $t \equiv 2$ or $3 \pmod 4$ blocks are nearly graceful (a parity condition ensures that the graphs in the latter case cannot be graceful). Moulton [2559] proved Rosa's conjecture while introducing the slightly stronger concept of *almost graceful* (also known as *near* graceful) by permitting the vertex labels to come from $\{0, 1, 2, \ldots, q-1, q+1\}$ while the edge labels are $1, 2, \ldots, q-1, q$, or $1, 2, \ldots, q-1, q+1$. More generally, Rosa [3205] conjectured that all triangular cacti are either graceful or almost graceful and suggested the use of Skolem sequences to label some types of triangular cacti. Dyer, Payne, Shalaby, and Wicks [993] verified the conjecture for two families of triangular cacti using Langford sequences to obtain Skolem and hooked Skolem sequences with specific subsequences. In [194] Alkasasbeh and Dyer used Skolem-type sequences [194] new
to show that (near) graceful labelings exist for windmills with C_3 and C_4 vanes, and provided infinite families of 3, 5-windmills and 3,6-windmills. They also showed that the graph obtained from the union of t 5-cycles with one vertex in common is graceful if and only if t is 0 or 3 mod 4 and near graceful when t is 1 or 2 mod 4. In [195] Alkasasbeh, [195] new
Dyer, and Howell obtained graceful labelings for the snakes kC_n and use this condition to label every such snake for $n = 1, 2, \ldots, 6$. They extended this result to cyclic snakes where the cycles lengths vary and obtained (near) graceful labelings of cyclic snakes based on cycles of lengths $n = 6, 10$, and 14, thereby completely solving the $n = 6$ case. In 2025 Letzter, Pokrovskiy, and Williams [2233] established an asymptotic version of the graceful [2233] new
tree conjecture by showing every sufficently large tree admits an almost graceful labeling.

Seoud and Elsakhawi [3378] and [3379] have shown that the following graphs are almost (near) graceful: C_n; $P_n + \overline{K_m}$; $P_n + K_{1,m}$; $K_{m,n}$; $K_{1,m,n}$; $K_{2,2,m}$; $K_{1,1,m,n}$; $P_n \times P_3$ $(n \geq 3)$; $K_5 \cup K_{1,n}$; $K_6 \cup K_{1,n}$, and ladders. In [2265] Licona and Tey showed that for a given integer $m \geq 3$, there exist infinitely many graceful (resp., almost graceful) two nested cycles graphs with a chord cycle of size m, [2265] new

For a graph G with p vertices, q edges, and $1 \leq k \leq q$, Eshghi [1048] defines a *holey α-labeling with respect to k* as an injective vertex labeling f for which $f(v) \in \{1, 2, \ldots, q+1\}$ for all v, $\{|f(u) - f(v)| \mid \text{for all edges } uv\} = \{1, 2, \ldots, k-1, k+1, \ldots, q+1\}$, and there exist an integer γ with $0 \leq \gamma \leq q$ such that $\min\{f(u), f(v)\} \leq \gamma \leq \max\{f(u), f(v)\}$. He proves the following: P_n has a holey α-labeling with respect to all k; C_n has a holey α-labeling with respect to k if and only if either $n \equiv 2 \pmod 4$, k is even, and $(n, k) \neq (10, 6)$, or $n \equiv 0 \pmod 4$ and k is odd.

Recall from Section 2.2 that a *kC_n-snake* is a connected graph with k blocks whose block-cutpoint graph is a path and each of the k blocks is isomorphic to C_n. In addition to his results on the graceful kC_n-snakes given in Section 2.2, Barrientos [512] proved that when k is odd the linear kC_6-snake is nearly graceful and that $C_m \cup K_{1,n}$ is nearly graceful when $m = 3, 4, 5$, and 6.

Yet another kind of labeling introduced by Rosa in his 1967 paper [3203] is a ρ-labeling. (Sometimes called a *rosy* labeling). A *ρ-labeling* (or ρ-valuation) of a graph is an injection from the vertices of the graph with q edges to the set $\{0, 1, \ldots, 2q\}$, where if the edge labels induced by the absolute value of the difference of the vertex labels are $a_1, a_2, \ldots, a_q$, then $a_i = i$ or $a_i = 2q + 1 - i$. Rosa [3203] proved that a cyclic decomposition of the edge set of the complete graph K_{2q+1} into subgraphs isomorphic to a given graph G with q edges exists if and only if G has a ρ-labeling. (A decomposition of K_n into copies of G is called *cyclic* if the automorphism group of the decomposition itself contains the cyclic group of order n.) It is known that every graph with at most 11 edges has a ρ-labeling and that all lobsters have a ρ-labeling (see [757]).

In [541] Barrientos and Minion proved that a tree admits a ρ–labeling when the deletion of some of its leaves results in a graceful tree. They use this result to prove the existence of ρ-labeling for several families of trees such as lobsters and those of diameter up to seven. Similarly, they showed that if T is any graceful tree of size n and k is an integer such that $2k \geq n + 1$, then any tree of size $n + 2k$ obtained attaching a path of length 2 to k distinct vertices of T has a ρ-labeling.

Donovan, El-Zanati, Vanden Eyden, and Sutinuntopas [968] prove that rC_m has a ρ-labeling (or a more restrictive labeling) when $r \leq 4$. They conjecture that every 2-regular graph has a ρ-labeling. Gannon and El-Zanati [1183] proved that for any odd $n \geq 7, rC_n$ admits ρ-labelings. The cases $n = 3$ and $n = 5$ were done in [965] and [1029]. Aguado, El-Zanati, Hake, Stob, and Yayla [87] give a ρ-labeling of $C_r \cup C_s \cup C_t$ for each of the cases where $r \equiv 0$, $s \equiv 1$, $t \equiv 1 \pmod 4$; $r \equiv 0$, $s \equiv 3$, $t \equiv 3 \pmod 4$; and $r \equiv 1$, $s \equiv 1$, $t \equiv 3 \pmod 4$; (iv) $r \equiv 1$, $s \equiv 2$, $t \equiv 3 \pmod 4$; (v) $r \equiv 3$, $s \equiv 3$, $t \equiv 3 \pmod 4$. Caro, Roditty, and Schőnheim [757] provide a construction for the adjacency matrix for every graph that has a ρ-labeling. They ask the following question: If H is a connected graph having a ρ-labeling and q edges and G is a new graph with q edges constructed

by breaking H up into disconnected parts, does G also have a ρ-labeling? Kézdy [1949] defines a *stunted tree* as one whose edges can be labeled with $e_1, e_2, \ldots, e_n$ so that e_1 and e_2 are incident and, for all $j = 3, 4, \ldots, n$, edge e_j is incident to at least one edge e_k satisfying $2k \leq j - 1$. He uses Alon's "Combinatorial Nullstellensatz" to prove that if $2n + 1$ is prime, then every stunted tree with n edges has a ρ-labeling.

Jeba Jesintha and Ezhilarasi Hilda [1616] introduced a variation of Rosa's ρ-labeling as follows. A *$\rho^\star$-labeling* of a graph G is an injection from the vertices of the graph with q edges to the set $\{0, 1, \ldots, 2q\}$, where if the edge labels induced by the absolute value of the difference of the vertex labels are $e_1, e_2, \ldots, e_q$, then $e_i = i$ or $e_i = 2q - i$. They prove that all paths and shell-butterfly graphs have a $\rho^\star$-labeling.

In [532] Barrientos and Minion proved the existence of ρ-labelings for some types of forests that considerably reduce the number of trees that need to be studied to prove Kotzig's Conjecture that states that K_{2n+1} can be cyclically decomposed into $2n + 1$ subgraphs isomorphic to a given tree with n edges. Among their results are the following. If T_1 and T_2 admit α-labelings such that one of the end-vertices of the edge of weight 1 in T_2 is a leaf, then $T_1 \cup T_2$ admits a ρ-labeling. If $G_1, G_2, \ldots, G_k$ is a collection of graphs that admit α-labelings, where G_k is a caterpillar of size at least $k-2$, then $\bigcup_{i=1}^{k} G_i$ admits a ρ-labeling. Let $\mathcal{R}$ denote the family that consists of all trees G such that G has a branch H, (i.e., $G - H$ is a tree) that is a caterpillar, where the excess of $G - H$ is at most the size of H. They prove that G admits a ρ-labeling when $G \in \mathcal{R}$.

Recall a kayak paddle $KP(k, m, l)$ is the graph obtained by joining C_k and C_m by a path of length l. Fronček and Tollefeson [1147], [1148] proved that $KP(r, s, l)$ has a ρ-labeling for all cases. As a corollary they have that the complete graph K_{2n+1} is decomposable into kayak paddles with n edges.

In [1125] Fronček generalizes the notion of an α-labeling by showing that if a graph G on n edges allows a certain type of ρ-labeling), called α_2-labeling, then for any positive integer k the complete graph K_{2nk+1} can be decomposed into copies of G.

In their investigation of cyclic decompositions of complete graphs El-Zanati, Vanden Eynden, and Punnim [1035] introduced two kinds of labelings. They say a bipartite graph G with n edges and partite sets A and B has a *θ-labeling* h if h is a one-to-one function from $V(G)$ to $\{0, 1, \ldots, 2n\}$ such that $\{|h(b) - h(a)| \; ab \in E(G), a \in A, b \in B\} = \{1, 2, \ldots, n\}$. They call h a *ρ^+-labeling* of G if h is a one-to-one function from $V(G)$ to $\{0, 1, \ldots, 2n\}$ and the integers $h(x) - h(y)$ are distinct modulo $2n + 1$ taken over all ordered pairs (x, y) where xy is an edge in G, and $h(b) > h(a)$ whenever $a \in A, b \in B$ and ab is an edge in G. Note that θ-labelings are ρ^+-labelings and ρ^+-labelings are ρ-labelings. They prove that if G is a bipartite graph with n edges and a ρ^+-labeling, then for every positive integer x there is a cyclic G-decomposition of K_{2nx+1}. They prove the following graphs have ρ^+-labelings: trees of diameter at most 5, C_{2n}, lobsters, and comets (that is, graphs obtained from stars by replacing each edge by a path of some fixed length). They also prove that the disjoint union of graphs with α-labelings have a θ-labeling and conjecture that all forests have ρ-labelings.

A *σ-labeling* of $G(V, E)$ is a one-to-one function f from V to $\{0, 1, \ldots, 2|E|\}$ such that $\{|f(u) - f(v)| \mid uv \in E(G)\} = \{1, 2, \ldots, |E|\}$. Such a labeling of G yields cyclic

G-decompositions of K_{2n+1} and of $K_{2n+2} - F$, where F is a 1-factor of K_{2n+2}. El-Zanati and Vanden Eynden (see [86]) have conjectured that every 2-regular graph with n edges has a ρ-labeling and, if $n \equiv 0$ or 3 (mod 4), then every 2-regular graph has a σ-labeling. Aguado and El-Zanati [86] have proved that the latter conjecture holds when the graph has at most three components.

Given a bipartite graph G with partite sets X and Y and graphs H_1 with p vertices and H_2 with q vertices, Fronček and Winters [1149] define the *bicomposition* of G and H_1 and H_2, $G[H_1, H_2]$, as the graph obtained from G by replacing each vertex of X by a copy of H_1, each vertex of Y by a copy of H_2, and every edge xy by a graph isomorphic to $K_{p,q}$ with the partite sets corresponding to the vertices x and y. They prove that if G is a bipartite graph with n edges and G has a θ-labeling that maps the vertex set $V = X \cup Y$ into a subset of $\{0, 1, 2, \ldots, 2n\}$, then the bicomposition $G[\overline{K_p}, \overline{K_q}]$ has a θ-labeling for every $p, q \geq 1$. As corollaries they have: if a bipartite graph G with n edges and at most $n + 1$ vertices has a gracious labeling (see §3.1), then the bicomposition graph $G[\overline{K_p}, \overline{K_q}]$ has a gracious labeling for every $p, q \geq 1$, and if a bipartite graph G with n edges has a θ-labeling, then for every $p, q \geq 1$, the bicomposition $G[\overline{K_p}, \overline{K_q}]$ decomposes the complete graph K_{2npq+1}.

In a paper published in 2009 [1034] El-Zanati and Vanden Eynden survey "Rosa-type" labelings. That is, labelings of a graph G that yield cyclic G-decompositions of K_{2n+1} or K_{2nx+1} for all natural numbers x. The 2009 survey by Fronček [1124] includes generalizations of ρ- and α-labelings that have been used for finding decompositions of complete graphs that are not covered in [1034].

Blinco, El-Zanati, and Vanden Eynden [662] call a non-bipartite graph *almost-bipartite* if the removal of some edge results in a bipartite graph. For these kinds of graphs G they call a labeling f a *γ-labeling* of G if the following conditions are met: f is a ρ-labeling; G is tripartite with vertex tripartition A, B, C with $C = \{c\}$ and $\overline{b} \in B$ such that $\{\overline{b}, c\}$ is the unique edge joining an element of B to c; if av is an edge of G with $a \in A$, then $f(a) < f(v)$; and $f(c) - f(\overline{b}) = n$. (In § 3.2 the term γ-labeling is used for a different kind of labeling.) They prove that if an almost-bipartite graph G with n edges has a γ-labeling then there is a cyclic G-decomposition of K_{2nx+1} for all x. They prove that all odd cycles with more than 3 vertices have a γ-labeling and that $C_3 \cup C_{4m}$ has a γ-labeling if and only if $m > 1$. In [726] Bunge, El-Zanati, and Vanden Eynden prove that every 2-regular almost bipartite graph other than C_3 and $C_3 \cup C_4$ have a γ-labeling.

In [662] Blinco, El-Zanati, and Vanden Eynden consider a slightly restricted ρ^+-labeling for a bipartite graph with partite sets A and B by requiring that there exists a number λ with the property that $\rho^+(a) \leq \lambda$ for all $a \in A$ and $\rho^+(b) > \lambda$ for all $b \in B$. They denote such a labeling by ρ^{++}. They use this kind of labeling to show that if G is a 2-regular graph of order n in which each component has even order then there is a cyclic G-decomposition of K_{2nx+1} for all x. They also conjecture that every bipartite graph has a ρ-labeling and every 2-regular graph has a ρ-labeling.

Dufour [981] and Eldergill [1000] have some results on the decomposition of complete graphs using labeling methods. Balakrishnan and Sampathkumar [465] showed that for each positive integer n the graph $\overline{K_n} + 2K_2$ admits a ρ-labeling. Balakrishnan [459] asks if

it is true that $\overline{K_n} + mK_2$ admits a ρ-labeling for all n and m. Fronček [1123] and Fronček and Kubesa [1139] have introduced several kinds of labelings for the purpose of proving the existence of special kinds of decompositions of complete graphs into spanning trees.

For positive integers c and d, let $K_{c\times d}$ denote the complete multipartite graph with c parts, each containing d vertices. Let G with n edges be the union of two vertex-disjoint even cycles. In [3700] Su et al. use Rosa-type graph labelings to show that there exists a cyclic G- decomposition of $K_(2n+1)\times t$, $K_{(n/2+1)\times 4t}, K_{5\times(n/2)t}$, and of K_{2nt} for every positive integer t. If $n \equiv 0 \pmod 4$,then there also exists a cyclic G-decomposition of $K_{n+1} \times 2t$, $K_{(n/4)+1} \times 8t$, $K_9 \times (n/4)t$, and of $K_{3\times nt}$ for every positive integer t.

For (p, q)-graphs with $p = q + 1$, Frucht [1152] has introduced a stronger version of almost graceful graphs by permitting as vertex labels $\{0, 1, \ldots, q-1, q+1\}$ and as edge labels $\{1, 2, \ldots, q\}$. He calls such a labeling *pseudograceful*. Frucht proved that P_n $(n \geq 3)$, combs, sparklers (i.e., graphs obtained by joining an end vertex of a path to the center of a star), $C_3 \cup P_n$ $(n \neq 3)$, and $C_4 \cup P_n$ $(n \neq 1)$ are pseudograceful whereas $K_{1,n}$ $(n \geq 3)$ is not. Kishore [1968] proved that $C_s \cup P_n$ is pseudograceful when $s \geq 5$ and $n \geq (s + 7)/2$ and that $C_s \cup S_n$ is pseudograceful when $s = 3, s = 4$, and $s \geq 7$. Seoud and Youssef [3411] and [3407] extended the definition of pseudograceful to all graphs with $p \leq q + 1$. They proved that K_m is pseudograceful if and only if $m = 1, 3$, or 4 [3407]; $K_{m,n}$ is pseudograceful when $n \geq 2$, and $P_m + \overline{K_n}$ $(m \geq 2)$ [3411] is pseudograceful. They also proved that if G is pseudograceful, then $G \cup K_{m,n}$ is graceful for $m \geq 2$ and $n \geq 2$ and $G \cup K_{m,n}$ is pseudograceful for $m \geq 2, n \geq 2$ and $(m, n) \neq (2, 2)$ [3407]. They ask if $G \cup K_{2,2}$ is pseudograceful whenever G is. Seoud and Youssef [3407] observed that if G is a pseudograceful Eulerian graph with q edges, then $q \equiv 0$ or 3 (mod 4). Youssef [4277] has shown that C_n is pseudograceful if and only if $n \equiv 0$ or 3 (mod 4), and for $n > 8$ and $n \equiv 0$ or 3 (mod 4), $C_n \cup K_{p,q}$ is pseudograceful for all $p, q \geq 2$ except $(p, q) = (2, 2)$. Youssef [4274] has shown that if H is pseudograceful and G has an α-labeling with k being the smaller vertex label of the edge labeled with 1 and if either $k + 2$ or $k - 1$ is not a vertex label of G, then $G \cup H$ is graceful. In [4278] Youssef shows that if G is (p, q) pseudograceful graph with $p = q + 1$, then $G \cup S_m$ is Skolem-graceful (see Section 3.5 for the definition). As a corollary he obtains that for all $n \geq 2$, $P_n \cup S_m$ is Skolem-graceful if and only if $n \geq 3$ or $n = 2$ and m is even.

In [4283] Youssef generalizes his results in [4274] and provides new families of disconnected graphs that have α-labelings and pseudo α-labelings. (A *pseudo* α-labeling f is an α-labeling for which there is an integer k_j with the property that for each edge xy of the graph either $f(x) \leq k_j < f(y)$ or $f(y) \leq k_j < f(x)$.)

For a graph G Ichishima, Muntaner-Batle, and Oshima [1489] defined the *beta-number* of G, $\beta(G)$, to be either the smallest positive integer n for which there exists an injective function f from the vertices of G to $\{1, 2, \ldots, n\}$ such that when each edge uv is labeled $|f(u) - f(v)|$ the resulting set of edge labels is $\{c, c + 1, \ldots, c + |E(G)| - 1\}$ for some positive integer c or $+\infty$ if there exists no such integer n. They defined the *strong beta-number* of G to be either the smallest positive integer n for which there exists an injective function f from the vertices of G to $\{1, 2, \ldots, n\}$ such that when each edge uv is labeled $|f(u) - f(v)|$ the resulting set of edge labels is $\{1, 2, \ldots, |E(G)|\}$ or $+\infty$ if there

exists no such integer n. They gave some necessary conditions for a graph to have a finite (strong) beta-number and some sufficient conditions for a graph to have a finite (strong) beta-number. They also determined formulas for the beta-numbers and strong beta-numbers of $C_n,\ 2C_n, K_n\ (n \geq 2),\ S_m \cup S_n,\ P_m \cup S_n$, and prove that nontrivial trees and forests without isolated vertices have finite strong beta-numbers. In [1481] Ichishima, López, Muntaner-Batte, and Oshima proved that if G is a bipartite graph and m is odd, then $\beta(mG) \leq m|E(G)| + m - 1$. If G has the additional property that G is a graceful nontrivial tree, then $\beta(mG) = m|V(G)| + m - 1$. They also investigate the (strong) beta-number of forests with components that are isomorphic to either paths or stars. They propose new conjectures on the (strong) beta-number of forests. In [1511] Ichishima and Oshima determine a formula for the (strong) beta-number of the linear forests $P_m \cup P_n$. As a corollary they provide a partial formula for the beta-number of the disjoint union of multiple copies of the same linear forest.

In [1519] Ichishima, Oshima, and Takahashi establish a lower bound for the beta-number of an arbitrary galaxy under certain conditions. They also introduce the notions of odd symmetric and even symmetric galaxies and determine formulas for the beta-number and gamma-number of odd symmetric galaxies. As corollaries, they provide formulas for the beta-number and gamma-number of the disjoint union of multiple copies of the same galaxy when the number of copies is odd. In addition to these, the present an upper bound for the beta-number of even symmetric galaxies and obtain partial formulas for the beta-number and gamma-number of even symmetric galaxies.

For a graph G of order p and size q and every positive integer n Ichishima, Muntaner-Batle, and Oshima [1491] proved if $\beta(G) = p - 1$, then there exists some positive integer c such that $q + np \leq \beta(G + nK_1) \leq c + q + np - 1$; if $\beta_s(G) = p - 1$, then $\beta(G + nK_1) = \beta_s(G + nK_1) = q + np$ and $G + nK_1$ is graceful; and if $q = p - 1$ and $\beta_s(G) = p - 1$, then $\beta(G + S_n) = \beta_s(G + S_n) = (n + 2)p + n - 1$. In particular, if T is a graceful tree of order p then $\beta(T + nK_1) = \beta_s(T + nK_1) = (n + 1)p - 1$. Moreover, $T + nK_1$ and $T + S_n$ are graceful.

In [1496] Ichishima, Muntaner-Batle, and Oshima establish a lower bound for the strong beta-number of an arbitrary galaxy (that is, a forest whose components are stars) under certain conditions. They also determine formulas for the (strong) beta-number and gracefulness of galaxies with three and four components. As corollaries, they provide formulas for the beta-number and gracefulness of the disjoint union of multiple copies of the same galaxies if the number of copies is odd. They pose some problems and conjecture. In [1486] Ichishima and Muntaner-Batle determined formulas for the (strong) beta-number and gracefulness of galaxies with five components. In [1500] Ichishima, Muntaner-Batle, and Oshima determined formulas for the (strong) beta-number and gamma-number of galaxies with five components. As a corollary of these results, they provide formulas for the beta-number and gamma-number of the disjoint union of multiple copies of the same galaxies if the number of copies is odd.

McTavish [2469] has investigated labelings of graphs with q edges where the vertex and edge labels are from $\{0, \ldots, q, q + 1\}$. She calls these *$\tilde{\rho}$-labelings*. Graphs that have $\tilde{\rho}$-labelings include cycles and the disjoint union of P_n or S_n with any graceful graph.

Frucht [1152] has made an observation about graceful labelings that yields nearly graceful analogs of α-labelings and weakly α-labelings in a natural way. Suppose $G(V, E)$ is a graceful graph with the vertex labeling f. For each edge xy in E, let $[f(x), f(y)]$ (where $f(x) \leq f(y)$) denote the interval of real numbers r with $f(x) \leq r \leq f(y)$. Then the intersection $\cap[f(x), f(y)]$ over all edges $xy \in E$ is a unit interval, a single point, or empty. Indeed, if f is an α-labeling of G then the intersection is a unit interval; if f is a weakly α-labeling, but not an α-labeling, then the intersection is a point; and, if f is a graceful but not a weakly α-labeling, then the intersection is empty. For nearly graceful labelings, the intersection also gives three distinct classes.

Let $G(V, E)$ be a graph without isolated vertices and with q edges. The *gracefulness* $\text{grac}(G)$ of G is the smallest positive integer k for which there exists an injective function $f : V \to \{0, 1, 2, \ldots, k\}$ such that the edge induced function $g_f : E \to \{1, 2, \ldots, k\}$ defined by $g_f(uv) = |f(u) - f(v)|$ for all edges uv is also injective. Let $c(f) = \max\{i : 1, 2, \ldots, i\}$ are edge labels$\}$ and let $m(G) = \max_f\{c(f)\}$ where the maximum is taken over all injective functions f from V to the nonnegative integers such that g_f is also injective. The measure $m(G)$ is called m-gracefulness of G. It determines how close G is to being graceful. Pereira, Singh, Arumugam [2822] prove that there are infinitely many nongraceful graphs with m-gracefulness $q-1$ and give necessary conditions for an Eulerian graph with q edges and K_p with q edges to have m-gracefulness $q-1$ and $q-2$. They prove that K_5 is the only complete graph to have m-gracefulness $q-1$. They also give an upper bound for the highest possible vertex label of K_p if $m(K_p) = q-2$.

A (p, q)-graph G is said to be a *super graceful* graph if there is a a bijective function $f : V(G) \cup E(G) \longrightarrow \{1, 2, \ldots, p+q\}$ such that $f(uv) = |f(u) - f(v)|$ for every edge $uv \in E(G)$ labeling. Perumal, Navaneethakrishnan, Nagarajan, Arockiaraj [2829] and [2828] show that the graphs P_n, C_n, $P_m \odot nK_1$, $K_{m,n}$, and $P_n \odot K_1$ minus a pendent edge at an endpoint of P_n are super graceful graphs. Lau, Shiu, and Ng [2112] study the super gracefulness of complete graphs, the disjoint union of certain star graphs, the complete tripartite graphs $K_{(1,1,n)}$, and certain families of trees. They also provide four methods of constructing new super graceful graphs. They prove all trees of order at most 7 are super graceful and conjecture that all trees are super graceful. Amutha and Uma Devi [219] proved the following graphs are super graceful: fans, double fans $DF_n = P_n + \overline{K_2}$ $(n \geq 2)$, and for $(m \geq 3, n \geq 2)$ the graphs obtained by identifying a central vertex of the star S_m with an end vertex of path in $P_n + K_1$. In [3705] Subbiah, Pandimadevi, and Chithra gave super graceful labelings of some trees.

For $k \geq 1$, Lau, Shiu, and Ng [2117] say a bijection $f : V \cup E \to [k, k+p+q-1]$ is a *k-super graceful labeling* if $f(uv) = |f(u) - f(v)|$ for every edge uv in G. A graph G is *k-super graceful* if it admits a k-super graceful labeling. This is a generalization of super graceful labeling defined by Perumal, Navaneethakrishnan, Nagarajan, Arockiaraj in [2828]. It was referred to as a k-sequential labeling by Slater in [3633], in which Slater gave necessary and sufficient conditions for a star to be k-sequential. In [2116] Lau, Shiu, and Ng investigated the existence of k-sequential labelings (which they call k-super graceful labelings) of paths, cycles, caterpillars, complete bipartite and complete tripartite graphs. In [2118] Lau, Shiu, Ng, Gao, and Schaffer investigated the k-super gracefulness

of some graphs with maximum degree 1 or 2.

In [1019] Elsonbaty and Daoud introduce a new version of gracefulness called an edge even graceful labeling of graphs. A bijective function f from the edges of a (p,q)-graph G to $\{2,4,\ldots,2q\}$ is said to be an *edge even graceful labeling* of G if the induced function f^* from the vertices to $\{0,2,\ldots,2q\}$ defined by $f^*(e)$ is the sum of $f(e)$ (mod $\max(p,q)$) is injective. They prove the following graphs have edge even graceful labelings: P_n if and only if n is odd, C_n if and only if n is odd, $K_{1,n}$ if and only if n is even, wheels, fans, friendship graphs, and double wheels $W_{n,n}$. The polar grid graph $P_{m,n}$ consists of n copies of C_m, a new vertex v_0, and m copies on P_{n+1} that share a endpoint at v_0 The graph is drawn as m concentric circles with a center at a new vertex v_0 and the m vertices of each cycle lie on a line with one endpoint at v_0 and the other endpoint at the outermost cycle in such a way that the n vertices of the copies on P_{n+1} other the v_0 intersect the vertices of cycles. Daoud [895] provided necessary and sufficient conditions for the polar grid graph to be edge even graceful.

Singh and Devaraj [3608] call a graph G with p vertices and q edges *triangular graceful* if there is an injection f from $V(G)$ to $\{0,1,2,\ldots,T_q\}$ where T_q is the qth triangular number and the labels induced on each edge uv by $|f(u)-f(v)|$ are the first q triangular numbers. They prove the following graphs are triangular graceful: paths, level 2 rooted trees, olive trees (see § 2.1 for the definition), complete n-ary trees, double stars, caterpillars, C_{4n}, C_{4n} with pendent edges, the one-point union of C_3 and P_n, and unicyclic graphs that have C_3 as the unique cycle. They prove that wheels, helms, flowers (see §2.2 for the definition) and K_n with $n \geq 3$ are not triangular graceful. They conjecture that all trees are triangular graceful. In [3455] Sethuraman and Venkatesh introduced a new method for combining graceful trees to obtain trees that have α-labelings.

Van Bussel [4044] considered two kinds of relaxations of graceful labelings as applied to trees. He called a labeling *range-relaxed graceful* it is meets the same conditions as a graceful labeling except the range of possible vertex labels and edge labels are not restricted to the number of edges of the graph (the edges are distinctly labeled but not necessarily labeled 1 to q where q is the number of edges). Similarly, he calls a labeling *vertex-relaxed graceful* if it satisfies the conditions of a graceful labeling while permitting repeated vertex labels. He proves that every tree T with q edges has a range-relaxed graceful labeling with the vertex labels in the range $0,1,\ldots,2q-d$ where d is the diameter of T and that every tree on n vertices has a vertex-relaxed graceful labeling such that the number of distinct vertex labels is strictly greater than $n/2$. In 2017 Sethuraman, Ragukumar, and Slater [3441] improved the bound on the range-relaxed graceful labeling given by Van Bussel in [4044] in 2002 for a tree T.

The *range-relaxed graceful game* is a maker-breaker game played in a simple graph G where two players, Alice and Bob, alternately assign an unused label $f(v) \in \{0,\ldots,k\}$ $(k \geq |E(G)|)$, to an unlabeled vertex $v \in V(G)$. If both ends of an edge $vw \in E(G)$ are already labeled, then the label of the edge is defined as $|f(v)-f(w)|$. Alice's goal is to end up with a vertex labeling of G where all edges of G have distinct labels, and Bob's goal is to prevent this from happening. When it is required that $k = |E(G)|$, the game is called a *graceful game*. The range-relaxed graceful game and the graceful game

were proposed by Tuza in 2017 [3922]. In [912] de Oliveira, Dantas, and Lui, considered a question about the least number of consecutive non-negative integer labels necessary for Alice to win the game on an arbitrary simple graph G and also asked if Alice can win the range-relaxed graceful game on G with the set of labels $\{0,\ldots,k+1\}$ once it is known that she can win with the set $\{0,\ldots,k\}$. They investigated the graceful game in Cartesian and corona products of graphs, and determined that Bob has a winning strategy in all investigated families independently of who starts the game. Additionally, they partially answer Tuza's questions presenting the first results in the range-relaxed graceful game and proving that Alice wins on any simple graph G with order n, size m, and maximum degree Δ, for any set of labels $\{0,\ldots,k\}$ with $k \geq (n-1)+2\Delta(m-\Delta)+(\Delta(\Delta-1))/2$. In [910] Oliveira, Artigas, Dantas, Frickes, and Luiz studied winning strategies for the graceful game introduced by Tuza in 2017 [3922] for the graph classes: paths, complete graphs, cycles, complete bipartite graphs, caterpillars, Wgererat trees, gears, webs, prisms, hypercubes, 2-powers of paths, wheels, and fan graphs.

In [528], Barrientos and Krop introduce left- and right-layered trees as trees with a specific representation and define the excess of a tree. Applying these ideas, they show a range-relaxed graceful labeling which improves the upper bound for maximum vertex label given by Van Bussel in [4044]. They also improve the bounds given by Rosa and Širáň in [3206] for the α-size and gracesize of lobsters.

Sekar [3337] calls an injective function ϕ from the vertices of a graph with q edges to $\{0,1,3,4,6,7,\ldots,3(q-1),3q-2\}$ *one modulo three graceful* if the edge labels induced by labeling each edge uv with $|\phi(u)-\phi(v)|$ is $\{1,4,7,\ldots,3q-2\}$. He proves that the following graphs are one modulo three graceful: P_m; C_n if and only if $n \equiv 0 \bmod 4$; $K_{m,n}$; $C_{2n}^{(2)}$ (the one-point union of two copies of C_{2n}); $C_n^{(t)}$ for $n = 4$ or 8 and $t > 2$; $C_6^{(t)}$ and $t \geq 4$; caterpillars; stars; lobsters; banana trees; rooted trees of height 2; ladders; the graphs obtained by identifying the endpoints of any number of copies of P_n; the graph obtained by attaching pendent edges to each endpoint of two identical stars and then identifying one endpoint from each of these graphs; the graph obtained by identifying a vertex of C_{4k+2} with an endpoint of a star; n-polygonal snakes (see §2.2) for $n \equiv 0 \pmod 4$; n-polygonal snakes for $n \equiv 2 \pmod 4$ where the number of polygons is even; crowns $C_n \odot K_1$ for n even; $C_{2n} \odot P_m$ (C_{2n} with P_m attached at each vertex of the cycle) for $m \geq 3$; chains of cycles (see §2.2) of the form $C_{4,m}, C_{6,2m}$, and $C_{8,m}$. He conjectures that every one modulo three graceful graph is graceful.

A *subdivided shell graph* is obtained by subdividing the edges in the path of the shell graph. Jeba Jesintha and Ezhilarasi Hilda [1611] proved that the subdivided uniform shell bow graphs (that is, double shells in which each shell has the same order) are one modulo three graceful.

In [3104] Ramachandran and Sekar introduced the notion of one modulo N graceful as follows. For a positive integer N a graph G with q edges is said to be *one modulo N graceful* if there is an injective function ϕ from the vertex set of G to $\{0,1,N,N+1,2N,2N+1,\ldots,(q-1)N,(q-1)N+1\}$ such that ϕ induces a bijection ϕ^* from the edge set of G to $\{1,N+1,2N+1,\ldots,(q-1)N+1\}$ where $\phi^*(uv) = |\phi(u)-\phi(v)|$. They proved the following graph are one modulo N graceful for all positive integers N:

paths, caterpillars, and stars [3104]; n-polygonal snakes, $C_n^{(t)}$, $P_{a,b}$ [3109]; the splitting graphs $S'(P_{2n})$, $S'(P_{2n+1})$, $S'(K_{1,n})$, all subdivision graphs of double triangular snakes, and all subdivision graphs of $2m$-triangular snakes [3105]; the graph $L_n \otimes S_m$ obtained from the ladder L_n ($P_n \times P_2$) by identifying one vertex of L_n with any vertex of the star S_m other than the center of S_m [3107]; arbitrary supersubdivisions of paths, disconnected paths, cycles, and stars [3106]; and regular bamboo trees and coconut trees [3108]. Ramachandran and Sekar [3110] proved the supersubdivisions of ladders are one modulo N graceful for all positive integers N. In [3111] Ramachandran and Sekar proved that the crowns, armed crowns, and chain of even cycles are one modulo N graceful for all positive integers N. Jeba Jesintha and Ezhilarasi Hilda [1610] proved the disjoint union of two subdivided shell graphs are one modulo three graceful.

In 1983 Bange and Barkauskas z [470] introduced the notion of Fibonacci graceful graphs as follows. A function $f : V(G) \to \{0, 1, 2, \ldots, F_q\}$, where G has q edges and F_q is the qth Fibonacci number, is a *Fibonacci graceful labeling* if the induced edge labeling $\overline{f}(uv) = |f(u) - f(v)|$ is a bijection to the set of the first q Fibonacci numbers. Such a graph is called *Fibonacci graceful*. They derived a number of properties of Fibonacci graceful graphs and provided some forbidden subgraphs of Fibonacci graceful graphs. Other results include: C_n is Fibonacci graceful if and only if $n = 0$ or 2 mod 3, trees with at least 7 vertices are Fibonacci graceful, and a maximal outerplanar graph with at least four vertices is Fibonacci graceful if and only if it has exactly two vertices of degree 2. Kathiresan and Amutha in [1923] prove the following: K_n is Fibonacci graceful if and only if $n \leq 3$; if an Eulerian graph with q edges is Fibonacci graceful, then $q \equiv 0$ (mod 3); paths are Fibonacci graceful; fans $P_n \odot K_1$ are Fibonacci graceful; squares of paths P_n^2 are Fibonacci graceful; and caterpillars are Fibonacci graceful. Dharman and Shanmuga Sundaram [954] proved that balloon trees, barycentric subdivisions of bistars, umbrella graphs, and double comb graphs are Fibonacci graceful graphs. In [222] Amuthavalli and Sundaram showed that female zodiac sign graphs are Fibonacci graceful antimagic graphs.

Kalyan and Kempepatil [1815] proved that trees and the graphs obtained by joining a vertex of C_{3m} and a vertex of C_{3n} by P_2 or P_3 admit Fibonacci graceful labelings. They define a function $f : V(G) \to \{0, F_1, F_2, \ldots, F_q\}$, where F_i is the ith Fibonacci number, to be a *super Fibonacci graceful* labeling if the induced labeling $\overline{f}(uv) = |f(u) - f(v)|$ is onto the set $\{F_1, F_2, \ldots, F_q\}$. They show that bistars $B_{n,n}$ are Fibonacci graceful but not super Fibonacci graceful for $n \geq 5$; cycles C_n are super Fibonacci graceful if and only if $n \equiv 0$ (mod 3); if G is Fibonacci or super Fibonacci graceful, then $G \odot K_1$ is Fibonacci graceful; if G_1 and G_2 are super Fibonacci graceful graphs in which no two adjacent vertices have the labels 1 and 2, then $G_1 \cup G_2$ is Fibonacci graceful; and if $G_1, G_2, \ldots, G_n$ are super Fibonacci graceful graphs in which no two adjacent vertices are labeled with 1 and 2, then the amalgamation of $G_1, G_2, \ldots, G_n$ obtained by identifying the vertices having labels 0 is also a super Fibonacci graceful. Karthikeyan, Arthi, Abinaya, Swathi, Madhumathi [1911] proved that friendship graphs $C_3^{(t)}$ and the graphs obtained by the one-point union of copies of K_4 with an edge deleted are super Fibonacci graceful.

Vaidya and Prajapati [3997] proved: the graphs obtained joining a vertex of C_{3m} and a vertex of C_{3n} by a path P_k are Fibonacci graceful; the graphs obtained by starting with

any number of copies of C_{3m} and joining each copy with a copy of the next by identifying the end points of a path with a vertex of each successive pair of C_{3m} (the paths need not be the same length) are Fibonacci graceful; the one point union of C_{3m} and C_{3n} is Fibonacci graceful; the one point union of k cycles C_{3m} is super Fibonacci graceful; every cycle C_n with $n \equiv 0 \pmod 3$ or $n \equiv 1 \pmod 3$ is an induced subgraph of a super Fibonacci graceful graph; and every cycle C_n with $n \equiv 2 \pmod 3$ can be embedded as a subgraph of a Fibonacci graceful graph. [3684] Sridevi, Navaneethakrishnan, and Nagarajan proved the following graphs are super Fibonacci graceful: the graphs obtained by identifying the apex of a fan with the end point of a path, the graphs obtained by identifying the apex of a fan with the vertex of maximum degree of $K_{1,n} \odot P_2$, the graphs obtained by identifying a vertex of C_{3n} with the end point of a path, the graphs obtained by identifying a vertex of C_{3n} with the center of a star, and the graphs obtained by identifying each endpoint a star with the center of $K_{1,2}$.

For a graph G with q edges an injective function f from the vertices of G to $\{0, F_1, F_2, \ldots, F_{q-1}, F_{q+1}\}$, where F_i is the ith Fibonacci number (as defined by Kathiresan and Amuth above), is said to be *almost super Fibonacci graceful* if the induced edge labeling $f * (uv) = |f(u) - f(v)|$ is a bijection onto the set $\{F_1, F_2, \ldots, F_q\}$ or $\{0, F_1, F_2, \ldots, F_{q-1}, F_{q+1}\}$. Sridevi, Navaneethakrishnan, and Nagarajan [3684] proved that paths, combs, graphs obtained by subdividing each edge of a star, and some special types of extension of cycle related graphs are almost super Fibonacci graceful labeling. Sridevi, Navaneethakrishnan, and Nagarajan [3683] showed that paths, combs, and some special types of extension of cycle-related graphs are almost super Fibonacci graceful.

For a graph G and a vertex v of G, a *vertex switching* G_v is the graph obtained from G by removing all edges incident to v and adding edges joining v to every vertex not adjacent to v in G. Vaidya and Vihol [4023] prove the following: trees are Fibonacci graceful; the graph obtained by switching of a vertex in cycle is Fibonacci graceful; wheels and helms are not Fibonacci graceful; the graph obtained by switching of a vertex in a cycle is super Fibonacci graceful except $n \geq 6$; the graph obtained by switching of a vertex in cycle C_n for $n \geq 6$ can be embedded as an induced subgraph of a super Fibonacci graceful graph; and the graph obtained by joining two copies of a fixed fan with an edge is Fibonacci graceful.

The *Perrin sequence* of numbers P_n is defined by the linear recurrence relation satisfying the conditions: $P_1 = 3, P_2 = 0, P_3 = 2$, and $P_n = P_{n-2} + P_{n-3}$, if $n \geq 4$. Letting P_i be the i^{th} term of the Perrin sequence and $P_0 = 0$, Sugumaran and Rajesh [3766] introduced the notion of Perrin graceful labeling as follows: A function f is called a *Perrin graceful* labeling of a graph G, if $f : V(G) \to \{P_0, P_1, P_2, ..., P_q\}$ is injective and the induced function $f^{(*)} : E(G) \to \{P_1, P_2, ..., P_q\}$ defined by $f^*(uv) = |f(u) - f(v)|$ is bijective. A graph that admits Perrin graceful labeling is called a *Perrin graceful* graph. In [3766] Sugumaran and Rajesh proved that the following graphs are Perrin graceful graphs: $K_{1,n}$, $B_{n,n}$, $P_n \odot K_1$, $C_n \odot K_1$, and $\langle K_{1,n}; 4rangle$.

For $n \geq 1$ the Pell numbers are defined as $p_0 = 0,\ p_1 = 1$, and $p_{n+1} = 2p_n + p_{n-1}$. For a graph G with q edges Muthuramakrishnan and Sutha [2619] introduced the concept of *Pell graceful* labeling as an injective function f from $V(G)$ to the Pell numbers $\{0, 1, 2, \ldots, p_q\}$

such that the induced edge labeling $f^*(uv) = |f(u) - f(v)|$ is a bijection onto the Pell numbers $\{p_1, p_2, \ldots, p_q\}$ They prove that paths, combs $P_n \odot K_1$ $(n \geq 3)$, and the graphs obtained by the one point union of paths of lengths $1, 2, \ldots, n$ $(n \geq 3)$ are Pell graceful.

In [709] Brešar and Klavžar define a natural extension of graceful labelings of certain tree subgraphs of hypercubes. A subgraph H of a graph G is called *isometric* if for every two vertices $u,\ v$ of H, there exists a shortest u-v path that lies in H. The isometric subgraphs of hypercubes are called *partial cubes*. Two edges $xy,\ uv$ of G are in Θ-relation if $d_G(x,u) + d_G(y,v) \neq d_G(x,v) + d_G(y,u)$. A Θ-relation is an equivalence relation that partitions $E(G)$ into Θ-classes. A Θ-*graceful labeling* of a partial cube G on n vertices is a bijection $f\colon V(G) \to \{0, 1, \ldots, n-1\}$ such that, under the induced edge labeling, all edges in each Θ-class of G have the same label and distinct Θ-classes get distinct labels. They prove that several classes of partial cubes are Θ-graceful and the Cartesian product of Θ-graceful partial cubes is Θ-graceful. They also show that if there exists a class of partial cubes that contains all trees and every member of the class admits a Θ-graceful labeling then all trees are graceful.

Cohen and Kovše showed that the graph obtained by merging two vertices of two 4-cycles is not a Θ-graceful partial cube, thereby answering in the negative a question of Brešar and Klavžar [709] who asked whether every partial cube is Θ-graceful.

Table 4 provides a summary results about graceful-like labelings adapted from [706]. "Y" indicates that all graphs in that class have the labeling; "N" indicates that not all graphs in that class have the labeling; "?" means unknown; "C" means conjectured.

Table 4: **Summary of Results on Graceful-like labelings**

Graph	α-labeling	β-labeling	σ-labeling	ρ-labeling
Cycle $C_n,\ n \equiv 0 \bmod 4$	Y [3203]	Y	Y	Y
Cycle $C_n,\ n \equiv 3 \bmod 4$	N [3203]	Y [3203]	Y	Y
Wheels	N	Y [1150], [1449]	Y	Y
Trees				
Yes, if order $\leq$	5	35 [1065]	54	
Paths	Y [3203]	Y	Y	Y
Caterpillars	Y [3203]	Y	Y	Y
Firecrackers	Y [795]	Y	Y	Y
Lobsters	N [663]	?C [629]	Y	Y [757]
Bananas	?	Y [3431], [3430]	Y	Y
Symmetrical trees	N [663]	Y [633]	Y	Y
Olive trees	?	Y [2789], [16]	Y	Y
Diameter < 8	N [4160]	Y	Y	Y
< 5 end vertices	N [663]	Y [3203]	Y	Y
Max degree 3	N [3206]	C	C	C
Max degree 3 and perfect matching	C [703]	C	C	C

3.4 k-graceful Labelings

A natural generalization of graceful graphs is the notion of k-graceful graphs introduced independently by Slater [3634] in 1982 and by Maheo and Thuillier [2387] in 1982. A graph G with q edges is *k-graceful* if there is labeling f from the vertices of G to $\{0, 1, 2, \ldots, q + k - 1\}$ such that the set of edge labels induced by the absolute value of the difference of the labels of adjacent vertices is $\{k, k + 1, \ldots, q + k - 1\}$. Obviously, 1-graceful is graceful and it is readily shown that any graph that has an α-labeling is k-graceful for all k. Graphs that are k-graceful for all k are sometimes called *arbitrarily graceful*. The result of Barrientos and Minion [529] that all snake polyominoes are α-graphs partially answers a question of Acharya [42] and supports his conjecture that if the length of every cycle of a graph is a multiple of 4, then the graph is arbitrarily graceful. In [3379] Seoud and Elsakhawi show that $P_2 \oplus \overline{K_2}$ $(n \geq 2)$ is arbitrarily graceful. Ng [2693] has shown that there are graphs that are k-graceful for all k but do not have an α-labeling.

Results of Maheo and Thuillier [2387] together with those of Slater [3634] show that: C_n is k-graceful if and only if either $n \equiv 0$ or 1 (mod 4) with k even and $k \leq (n-1)/2$, or $n \equiv 3$ (mod 4) with k odd and $k \leq (n^2-1)/2$. Maheo and Thuillier [2387] also proved that the wheel W_{2k+1} is k-graceful and conjectured that W_{2k} is k-graceful when $k \neq 3$ or $k \neq 4$. This conjecture was proved by Liang, Sun, and Xu [2256]. Kang [1881] proved that $P_m \times C_{4n}$ is k-graceful for all k. Lee and Wang [2215] showed that the graphs obtained from a nontrivial path of even length by joining every other vertex to one isolated vertex (a *lotus*), the graphs obtained from a nontrivial path of even length by joining every other vertex to two isolated vertices (a *diamond*), and the graphs obtained by arranging vertices into a finite number of rows with i vertices in the ith row and in every row the jth vertex in that row is joined to the jth vertex and $j + 1$st vertex of the next row (a *pyramid*) are k-graceful. Liang and Liu [2244] have shown that $K_{m,n}$ is k-graceful. Bu, Gao, and Zhang [719] have proved that $P_n \times P_2$ and $(P_n \times P_2) \cup (P_n \times P_2)$ are k-graceful for all k. Acharya (see [42]) has shown that a k-graceful Eulerian graph with q edges must satisfy one of the following conditions: $q \equiv 0$ (mod 4), $q \equiv 1$ (mod 4) if k is even, or $q \equiv 3$ (mod 4) if k is odd. Bu, Zhang, and He [723] have shown that an even cycle with a fixed number of pendent edges adjoined to each vertex is k-graceful. Lu, Pan, and Li [2372] have proved that $K_{1,m} \cup K_{p,q}$ is k-graceful when $k > 1$, and p and q are at least 2. Jirimutu, Bao, and Kong [1788] have shown that the graphs obtained from $K_{2,n}$ $(n \geq 2)$ and $K_{3,n}$ $(n \geq 3)$ by attaching $r \geq 2$ edges at each vertex is k-graceful for all $k \geq 2$. Seoud and Elsakhawi [3379] proved: paths and ladders are arbitrarily graceful; and for $n \geq 3$, K_n is k-graceful if and only if $k = 1$ and $n = 3$ or 4. Li, Li, and Yan [2242] proved that $K_{m,n}$ is k-graceful graph. Pradhan and Kamesh [2997] showed that the hairy cycle $C_n \odot rK_1$ ($n \equiv 3$ (mod 4), the graph obtained by adding a pendent edge to each pendent vertex of hairy cycle $C_n \odot K_1$; $n \equiv 0$ (mod 4), double graphs of path P_n, and double graphs of combs $P_n \odot K_1$ are k-graceful.

Yao, Cheng, Zhongfu, and Yao [4259] have shown: a tree of order p with maximum degree at least $p/2$ is k-graceful for some k; if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and

$d_{T_2}(u_2) \geq |T_2|/2$, then T is k-graceful for some positive k; if a tree T has two edges u_1u_2 and u_2u_3 such that the three components T_1, T_2, and T_3 of $T - \{u_1u_2, u_2u_3\}$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$, $d_{T_2}(u_2) \geq |T_2|/2$, and $d_{T_3}(u_3) \geq |T_3|/2$, then T is k-graceful for some $k > 1$; and every Skolem-graceful (see 3.5 for the definition) tree is k-graceful for all $k \geq 1$. They conjecture that every tree is k-graceful for some $k > 1$.

Several authors have investigated the k-gracefulness of various classes of subgraphs of grid graphs. Acharya [40] proved that all 2-dimensional polyminoes that are convex and Eulerian are k-graceful for all k; Lee [2145] showed that Mongolian tents and Mongolian villages are k-graceful for all k (see §2.3 for the definitions); Lee and K. C. Ng [2170] proved that all Young tableaus (see §2.3 for the definitions) are k-graceful for all k. (A special case of this is $P_n \times P_2$.) Lee and H. K. Ng [2170] subsequently generalized these results on Young tableaus to a wider class of planar graphs.

In [538] Barrientos and Minion say that two caterpillars Γ and Ω of size n are *analogous* if the stable sets of Γ have the same cardinalities as the stable sets of Ω. They prove that if Ω is an induced subgraph of a gracefully labeled graph G, such that the induced labeling is a bipartite k-labeling shifted c units, then the graph G' obtained by replacing Ω with any other caterpillar Γ analogous to Ω, is a graceful graph. This result is used to generalize several existing results that use k-graceful labelings of paths such as the subdivision of graceful trees [731], the α-labeling of the ith attachment tree [3455], the α-labelings of path-like trees [511], the α-labelings of the graphs obtained by identifying the end-vertices of b paths of length a with two new vertices, as well as the graceful labelings of the armed crowns [3337].

Duan and Qi [980] use $G_t(m_1, n_1; m_2, n_2; \ldots; m_s, n_s)$ to denote the graph composed of the s complete bipartite graphs $K_{m_1,n_1}, K_{m_2,n_2}, \ldots, K_{m_s,n_s}$ that have only t $(1 \leq t \leq \min\{m_1, m_2, \ldots, m_s\})$ common vertices but no common edge and $G(m_1, n_1; m_2, n_2)$ to denote the graph composed of the complete bipartite graphs K_{m_1,n_1}, K_{m_2,n_2} with exactly one common edge. They prove that these graphs are k-graceful graphs for all k.

Let $c, m, p_1, p_2, \ldots, p_m$ be positive integers. For $i = 1, 2, \ldots, m$, let S_i be a set of $p_i + 1$ integers and let D_i be the set of positive differences of the pairs of elements of S_i. If all these differences are distinct then the system $D_1, D_2, \ldots, D_m$ is called a *perfect system of difference sets starting at* c if the union of all the sets D_i is $c, c+1, \ldots, c-1+\sum_{i=1}^{m} \binom{p_i+1}{2}$. There is a relationship between k-graceful graphs and perfect systems of difference sets. A perfect system of difference sets starting with c describes a c-graceful labeling of a graph that is decomposable into complete subgraphs. A survey of perfect systems of difference sets is given in [24].

Acharya and Hegde [56] generalized k-graceful labelings to *(k, d)-graceful labelings* by permitting the vertex labels to belong to $\{0, 1, 2, \ldots, k + (q-1)d\}$ and requiring the set of edge labels induced by the absolute value of the difference of labels of adjacent vertices to be $\{k, k + d, k + 2d, \ldots, k + (q-1)d\}$. They also introduce an analog of α-labelings in the obvious way. Notice that a (1,1)-graceful labeling is a graceful labeling and a $(k, 1)$-graceful labeling is a k-graceful labeling. Bu and Zhang [724] have shown: $K_{m,n}$ is (k, d)-graceful for all k and d; for $n > 2, K_n$ is (k, d)-graceful if and only if $k = d$

and $n \leq 4$; if $m_i, n_i \geq 2$ and $\max\{m_i, n_i\} \geq 3$, then $K_{m_1,n_1} \cup K_{m_2,n_2} \cup \cdots \cup K_{m_r,n_r}$ is (k,d)-graceful for all k, d, and r; if G has an α-labeling, then G is (k,d)-graceful for all k and d; a k-graceful graph is a (kd,d)-graceful graph; a (kd,d)-graceful connected graph is k-graceful; and a (k,d)-graceful graph with q edges that is not bipartite must have $k \leq (q-2)d$.

Let T be a tree with adjacent vertices u_0 and v_0 and pendent vertices u and v such that the length of the path $u_0 - u$ is the same as the length of the path $v_0 - v$. Hegde and Shetty [1422] call the graph obtained from T by deleting u_0v_0 and joining u and v an *elementary parallel transformation* of T. They say that a tree T is a *T_p-tree* if it can be transformed into a path by a sequence of elementary parallel transformations. They prove that every T_p-tree is (k,d)-graceful for all k and d and every graph obtained from a T_p-tree by subdividing each edge of the tree is (k,d)-graceful for all k and d.

Yao, Cheng, Zhongfu, and Yao [4259] have shown: a tree of order p with maximum degree at least $p/2$ is (k,d)-graceful for some k and d; if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and T_2 is a caterpillar, then T is Skolem-graceful (see 3.5 for the definition); if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and $d_{T_2}(u_2) \geq |T_2|/2$, then T is (k,d)-graceful for some $k > 1$ and $d > 1$; if a tree T has two edges u_1u_2 and u_2u_3 such that the three components T_1, T_2, and T_3 of $T - \{u_1u_2, u_2u_3\}$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$, $d_{T_2}(u_2) \geq |T_2|/2$, and $d_{T_3}(u_3) \geq |T_3|/2$, then T is (k,d)-graceful for some $k > 1$ and $d > 1$; and every Skolem-graceful tree is (k,d)-graceful for $k \geq 1$ and $d > 0$. They conjecture that every tree is (k,d)-graceful for some $k > 1$ and $d > 1$.

Hegde [1406] has proved the following: if a graph is (k,d)-graceful for odd k and even d, then the graph is bipartite; if a graph is (k,d)-graceful and contains C_{2j+1} as a subgraph, then $k \leq jd(q-j-1)$; K_n is (k,d)-graceful if and only if $n \leq 4$; C_{4t} is (k,d)-graceful for all k and d; C_{4t+1} is $(2t,1)$-graceful; C_{4t+2} is $(2t-1,2)$-graceful; and C_{4t+3} is $(2t+1,1)$-graceful.

A *semismooth graceful graph* is a bipartite graph G with the property that for some fixed positive integer $t \leq q$ and all positive integers l there is an injective map $g : V \longrightarrow \{0, 1, \ldots, t-l, t+l+1, \ldots, q+l\}$ such that the induced edge labeling map $g^\star : E \longrightarrow \{1+l, 2+l, \ldots, q+l\}$ defined by $g^\star(e) = |g(u) - g(v)|$ is a bijection. Kaneria, Gohil, and Makadia [1833] prove every semismooth graceful graph is a (k,d)-graceful; graphs obtained by joining two semismooth graceful graphs with an arbitrary path is a semismooth graceful graph; and the notions of graceful labeling and odd-even graceful labelings are equivalent. (A graph G with q edges is *odd-even graceful* if there is an injection f from the vertices of G to $\{1, 3, 5, \ldots, 2q+1\}$ such that, when each edge uv is assigned the label $|f(u) - f(v)|$, the resulting edge labels are $\{2, 4, 6, \ldots, 2q\}$). Kaneria, Meghpara and Khoda [1840] prove: a smooth graceful labeling for a graph is also an α-labeling for the graph; a graph that has an α-labeling is a semismooth graceful graph; graphs that admit an α-labeling are semismooth graceful graphs; if m is even and H has an α-labeling, then the path union $P(m \cdot H)$ is a smooth graceful graph; and the path union $P(m \cdot H)$ has an α-labeling.

Nurvazly, Chasanah, and Wiranto [2729] showed that the corona of the ladder L_2 and $\overline{K_n}$ and of L_3 and $\overline{K_n}$, which they call *millipedes*, admits graceful, $\hat{\rho}$, and odd-even graceful labelings. In [3718] Sudha and Kanniga proved that tensor product of a star and P_2 is odd-even graceful. (The *tensor product* $G \otimes H$ of graphs G and H, has the vertex set $V(G) \times V(H)$ and any two vertices (u, u') and (v, v') are adjacent in $G \otimes H$ if and only if u' is adjacent with v' and u is adjacent with v.) In [4079] Venkatesh, Mahalakshmi, and Amirthavahini use $C_{n,k}$ to denote the dragon obtained by joining an end point of P_k with a vertex of C_n and $C_{n,k}^t$ to denote the graph obtained by taking one-point union of t copies of $C_{n,k}$ at the common vertex v. They proved that the graph $C_{n,k}^t$ admits a graceful labeling, an odd graceful labeling, and odd-even graceful labeling for all values of t with $n = 4, k = 1$, and that $C_{n,1}^t$ admits vertex cordial labeling for all values of n and t, except $n \equiv 2 \bmod 4$ (see Section 3.7). Nurvazly and Sugeng [2728] proved that $\Theta(C_3)^n$ graphs (n copies of C_3 that share an edge) have odd-even graceful labelings. Anitha, Selvam, and Thirusangu [261] provide k-graceful and odd-even graceful labelings for the extended duplicate graph of the kite graph. Dash, Behera, and Bhatta [902] provided odd-even graceful labelings for coconut trees, combs, certain classes of quadrilateral snakes, tristars, 4-stars, and a class of hairy cycles. [902] new

For a graph G let $G^{(1)}, G^{(2)}, \ldots, G^{(n)}$ be $n \geq 2$ copies of G. The graph obtained by joining vertices u, v of $G^{(i)}$ with same vertices of the graph $G^{(i+1)}$ by two edges, for all $i = 1, 2, \ldots, n-1$ is called the *double path union* of n copies of the graph G. Such graphs can obtained in $\frac{p(p-1)}{2}$ different ways, where $p = |V(G)|$ and are denoted by $D(n \cdot G)$. Kaneria, Teraiya and Meghpara [1869] prove the double path unions of C_{4m}, $K_{m,n}$, and P_{2m} have α-labelings.

Hegde [1404] calls a (k, d)-graceful graph (k, d)-*balanced* if it has a (k, d)-graceful labeling f with the property that there is some integer m such that for every edge uv either $f(u) \leq m$ and $f(v) > m$, or $f(u) > m$ and $f(v) \leq m$. He proves that if a graph is $(1, 1)$-balanced then it is (k, d)-graceful for all k and d and that a graph is $(1, 1)$-balanced graph if and only if it is (k, k)-balanced for all k. He conjectures that all trees are (k, d)-balanced for some values of k and d.

Slater [3637] has extended the definition of k-graceful graphs to countable infinite graphs in a natural way. He proved that all countably infinite trees, the complete graph with countably many vertices, and the countably infinite Dutch windmill is k-graceful for all k.

In [1428] Hegde and Shivarajkumar extend the idea of k-graceful labeling of undirected graphs to directed graphs as follows. A simple directed graph D with n vertices and e edges is labeled by assigning each vertex a distinct element from the set Z_{e+k} and assigning the edge xy from vertex x to vertex y the label $\theta(x, y) = \theta(y) - \theta(x) \bmod (e + k)$, where $\theta(y)$ and $\theta(x)$ are the values assigned to the vertices y and x respectively. A labeling is a *k-graceful* labeling if all $\theta(x, y)$ are distinct and belong to $\{k, k + 1, \ldots, k + e - 1\}$. If a digraph D admits a k-graceful labeling then D is called a *k-graceful digraph*. They provide some values of k for which the unidirectional cycles admit a k-graceful labeling; give a necessary and sufficient condition for the outspoken unicyclic wheel to be k-graceful; and prove that to provide a list of values of k for which the unicyclic wheel is k-graceful is

NP-complete.

More specialized results on k-graceful labelings can be found in [2145], [2170], [2173], [3634], [718], [720], and [719].

Graceful-type labelings methods have been used for cryptographical password construction for network data [4137], [4136], [4138], and [3784].

3.5 Skolem-Graceful Labelings

A number of authors have invented analogues of graceful graphs by modifying the permissible vertex labels. For instance, Lee (see [2199]) calls a graph G with p vertices and q edges *Skolem-graceful* if there is an injection from the set of vertices of G to $\{1, 2, \ldots, p\}$ such that the edge labels induced by $|f(x)-f(y)|$ for each edge xy are $1, 2, \ldots, q$. A necessary condition for a graph to be Skolem-graceful is that $p \geq q+1$. Lee and Wui [2229] have shown that a connected graph is Skolem-graceful if and only if it is a graceful tree. Yao, Cheng, Zhongfu, and Yao [4259] have shown that a tree of order p with maximum degree at least $p/2$ is Skolem-graceful. Although the disjoint union of trees cannot be graceful, they can be Skolem-graceful. Lee and Wui [2229] prove that the disjoint union of 2 or 3 stars is Skolem-graceful if and only if at least one star has even size. In [824] Choudum and Kishore show that the disjoint union of k copies of the star $K_{1,2p}$ is Skolem graceful if $k \leq 4p+1$ and the disjoint union of any number of copies of $K_{1,2}$ is Skolem graceful. For $k \geq 2$, let $St(n_1, n_2, \ldots, n_k)$ denote the disjoint union of k stars with $n_1, n_2, \ldots, n_k$ edges. Lee, Wang, and Wui [2222] showed that the 4-star $St(n_1, n_2, n_3, n_4)$ is Skolem-graceful for some special cases and conjectured that all 4-stars are Skolem-graceful. Denham, Leu, and, Liu [925] proved this conjecture. Kishore [1968] has shown that a necessary condition for $St(n_1, n_2, \ldots, n_k)$ to be Skolem graceful is that some n_i is even or $(k \equiv 0)$ or 1 (mod 4) (see also [4296] . He conjectures that each one of these conditions is sufficient. Yue, Yuan-sheng, and Xin-hong [4296] show that for k at most 5, a k-star is Skolem-graceful if at one star has even size or $k \equiv 0$ or 1 (mod 4). Choudum and Kishore [822] proved that all 5-stars are Skolem graceful.

Lee, Quach, and Wang [2185] showed that the disjoint union of the path P_n and the star of size m is Skolem-graceful if and only if $n = 2$ and m is even or $n \geq 3$ and $m \geq 1$. It follows from the work of Skolem [3626] that nP_2, the disjoint union of n copies of P_2, is Skolem-graceful if and only if $n \equiv 0$ or 1 (mod 4). Harary and Hsu [1371] studied Skolem-graceful graphs under the name *node-graceful*. Frucht [1152] has shown that $P_m \cup P_n$ is Skolem-graceful when $m + n \geq 5$. Bhat-Nayak and Deshmukh [644] have shown that $P_{n_1} \cup P_{n_2} \cup P_{n_3}$ is Skolem-graceful when $n_1 < n_2 \leq n_3,\ n_2 = t(n_1 + 2) + 1$ and n_1 is even and when $n_1 < n_2 \leq n_3,\ n_2 = t(n_1 + 3) + 1$ and n_1 is odd. They also prove that the graphs of the form $P_{n_1} \cup P_{n_2} \cup \cdots \cup P_{n_i}$ where $i \geq 4$ are Skolem-graceful under certain conditions. In [930] Deshmukh states the following results: the sum of all the edges on any cycle in a Skolem graceful graph is even; $C_5 \cup K_{1,n}$ if and only if $n = 1$ or 2; $C_6 \cup K_{1,n}$ if and only if $n = 2$ or 4.

Youssef [4274] proved that if G is Skolem-graceful, then $G + \overline{K_n}$ is graceful. In [4278] Youssef shows that for all $n \geq 2,\ P_n \cup S_m$ is Skolem-graceful if and only if $n \geq 3$ or $n = 2$

and m is even. Yao, Cheng, Zhongfu, and Yao [4259] have shown that if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and T_2 is a caterpillar or have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and $d_{T_2}(u_2) \geq |T_2|/2$, then T is Skolem-graceful.

A graph $G = (V, E)$ is said to be (k, d)-*Skolem graceful* if there exists a bijection f from $V(G)$ to $\{12, \ldots, |V|\}$ such that the induced edge labeling g_f defined by $g_f(uv) = |f(u) - f(v)|$ is a bijection from E to $\{k, k+d, \ldots, k+(q-1)d\}$ where k and d are positive integers. Such a labeling is called a (k, d)-*Skolem graceful labeling* of G. In [2821] Pereira, Singh, and Arumugam present a few basic results on (k, d)-Skolem graceful graphs and prove that nK_2 is $(2, 1)$-Skolem graceful if and only if $n \equiv 0$ or 3 (mod 4), which produces the Langford sequence $L(2, n)$.

Mendelsohn and Shalaby [2486] defined a *Skolem labeled* graph $G(V, E)$ as one for which there is a positive integer d and a function $L\colon V \to \{d, d+1, \ldots, d+m\}$, satisfying (a) there are exactly two vertices in V such that $L(v) = d+i$, $0 \leq i \leq m$; (b) the distance in G between any two vertices with the same label is the value of the label; and (c) if G' is a proper spanning subgraph of G, then L restricted to G' is not a Skolem labeled graph. Note that this definition is different from the Skolem-graceful labeling of Lee, Quach, and Wang. A *hooked Skolem sequence* of order n is a sequence $s_1, s_2, \ldots, s_{2n+1}$ such that $s_{2n} = 0$ and for each $j \in \{1, 2, \ldots, n\}$, there exists a unique $i \in \{1, 2, \ldots, 2n-1, 2n+1\}$ such that $s_i = s_{i+j} = j$. Mendelsohn [2485] established the following: any tree can be embedded in a Skolem labeled tree with $O(v)$ vertices; any graph can be embedded as an induced subgraph in a Skolem labeled graph on $O(v^3)$ vertices; for $d = 1$, there is a Skolem labeling or the minimum hooked Skolem (with as few unlabeled vertices as possible) labeling for paths and cycles; for $d = 1$, there is a minimum Skolem labeled graph containing a path or a cycle of length n as induced subgraph. In [2485] Mendelsohn and Shalaby prove that the necessary conditions in [2486] are sufficient for a Skolem or minimum hooked Skolem labeling of all trees consisting of edge-disjoint paths of the same length from some fixed vertex. Graham, Pike, and Shalaby [1326] obtained various Skolem labeling results for grid graphs. Among them are $P_1 \times P_n$ and $P_2 \times P_n$ have Skolem labelings if and only if $n \equiv 0$ or 1 mod 4; and $P_m \times P_n$ has a Skolem labeling for all m and n at least 3.

In [2838] Pike, Sanaei, and Shalaby introduce pseudo-Skolem sequences, which are similar to Skolem-type sequences in their structures and applications. They use known Skolem-type sequences to construct such sequences and discuss applications of these sequences to Skolem labelings of graphs such that H is bipartite, and give formulas for the gamma-number of rail-siding graphs and caterpillars.

In [854] Clark and Sanaei present (hooked) vertex Skolem labelings for generalized Dutch windmills whenever such labelings exist. They present a novel technique for showing that generalized Dutch windmills with more than two cycles cannot be Skolem labelled and that those composed of two cycles of lengths m and n, $n \geq m$ cannot be Skolem labelled if and only if $n - m \equiv 3$ or 5 (mod 8) and m is odd.

3.6 Odd-Graceful Labelings

Gnanajothi [1276, p. 182] defined a graph G with q edges to be *odd-graceful* if there is an injection f from $V(G)$ to $\{0, 1, 2, \ldots, 2q-1\}$ such that, when each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edge labels are $\{1, 3, 5, \ldots, 2q-1\}$. She proved that the class of odd-graceful graphs lies between the class of graphs with α-labelings and the class of bipartite graphs by showing that every graph with an α-labeling has an odd-graceful labeling and every graph with an odd cycle is not odd-graceful. She also proved the following graphs are odd-graceful: P_n; C_n if and only if n is even; $K_{m,n}$; combs $P_n \odot K_1$ (graphs obtained by joining a single pendent edge to each vertex of P_n); books; crowns $C_n \odot K_1$ (graphs obtained by joining a single pendent edge to each vertex of C_n) if and only if n is even; the disjoint union of copies of C_4; the one-point union of copies of C_4; $C_n \times K_2$ if and only if n is even; caterpillars; rooted trees of height 2; the graphs obtained from P_n ($n \geq 3$) by adding exactly two leaves at each vertex of degree 2 of P_n; the graphs obtained from $P_n \times P_2$ by deleting an edge that joins to end points of the P_n paths; the graphs obtained from a star by adjoining to each end vertex the path P_3 or by adjoining to each end vertex the path P_4. She conjectures that all trees are odd-graceful and proves the conjecture for all trees with order up to 10. Barrientos [515] has extended this to trees of order up to 12. Eldergill [1000] generalized Gnanajothi's result on stars by showing that the graphs obtained by joining one end point from each of any odd number of paths of equal length is odd-graceful. He also proved that the one-point union of any number of copies of C_6 is odd-graceful. Kathiresan [1921] has shown that ladders and graphs obtained from them by subdividing each step exactly once are odd-graceful. Barrientos [518] and [515] has proved the following graphs are odd-graceful: every forest whose components are caterpillars; every tree with diameter at most five is odd-graceful; and all disjoint unions of caterpillars. He conjectures that every bipartite graph is odd-graceful. In [2665] Neela and Selvaraj partially resolved a Barrientos's conjecture by showing that the following graphs are odd-graceful: finite unions of paths, stars, and caterpillars; finite unions of ladders; finite unions of paths, bistars and caterpillars; finite unions of graphs obtained by the one end point union of an odd number of paths of uniform length; and the coronas $K_{m;n} \odot rK_l$. Gao, Zhang, and Xu [1200] proved that $P_n \times P_m$ ($m = 2$, 3 or 4), generalized crown graphs $C_n \odot K_{1,t}$, and gears are odd graceful.

Seoud, Diab, and Elsakhawi [3376] have shown that a connected complete r-partite graph is odd-graceful if and only if $r = 2$ and that the join of any two connected graphs is not odd-graceful. Yan [4239] proved that $P_m \times P_n$ is odd-graceful labeling. Vaidya and Shah [4006] prove that the splitting graph and the shadow graph of bistar are odd-graceful. (The *shadow graph* $D_2(G)$ of a connected graph G is constructed by taking 2 copies G_1 and G_2 of G and joining each vertex u in G_1 to the neighbors of the corresponding vertex v in G_2. Li, Li, and Yan [2242] proved that $K_{m,n}$ is odd-graceful. Liu, Wang, and Lu [2292] that proved that a class of bicyclic graphs with a common edge is odd-graceful. Moussa and Badr [2558] proved that ladders and subdivisions of ladders with pendent edges are odd-graceful. Virk and Riasat [3160] showed that the union a specific type of tree with paths, stars, bistars, or ladders are odd-graceful. Riasat, Kanwal, and Javed [3171]

give odd-graceful labelings for disjoint unions of graphs consisting of generalized combs, ladders, stars, bistars, caterpillars and paths.

Sekar [3337] has shown the following graphs are odd-graceful: the graph obtained by identifying an end point of P_n with every vertex of C_m where $n \geq 3$ and m is even; $P_{a,b}$ when $a \geq 2$ and b is odd (see §2.7); $P_{2,b}$ and $b \geq 2$; $P_{4,b}$ and $b \geq 2$; $P_{a,b}$ when a and b are even and $a \geq 4$ and $b \geq 4$; $P_{4r+1,4r+2}$; $P_{4r-1,4r}$; all n-polygonal snakes with n even; $C_n^{(t)}$ (see §2.2 for the definition); graphs obtained by beginning with C_6 and repeatedly forming the one-point union with additional copies of C_6 in succession; graphs obtained by beginning with C_8 and repeatedly forming the one-point union with additional copies of C_8 in succession; graphs obtained from even cycles by identifying a vertex of the cycle with the endpoint of a star; $C_{6,n}$ and $C_{8,n}$ (see §2.7); the splitting graph of P_n (see §2.7) the splitting graph of C_n, n even; lobsters, banana trees, and regular bamboo trees (see §2.1).

Jeba Jesinthan, Devakirubanith, and Aathry [1601] proved that graphs obtained by attaching star graphs to the apex and all the path vertices of the subdivided shell graph are odd graceful.

Yao, Cheng, Zhongfu, and Yao [4259] have shown the following: if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and T_2 is a caterpillar, then T is odd-graceful; and if a tree T has a vertex of degree at least $|T|/2$, then T is odd-graceful. They conjecture that for trees the properties of being Skolem-graceful and odd-graceful are equivalent. Recall a banana tree is a graph obtained by starting with any number os stars and connecting one end-vertex from each to a new vertex. Zhenbin [4332] has shown that graphs obtained by starting with any number of stars, appending an edge to exactly one edge from each star, then joining the vertices at which the appended edges were attached to a new vertex are odd-graceful.

Solairaju and Chithra [3651] defined a graph G with q edges to be *edge-odd graceful* if there is a bijection f from the edges of the graph to $\{1, 3, 5, \ldots, 2q-1\}$ such that, when each vertex is assigned the sum of all the edges incident to it mod $2q$, the resulting vertex labels are distinct. They prove they following graphs are edge-odd-graceful: paths with at least 3 vertices; odd cycles; ladders $P_n \times P_2$ ($n \geq 3$); stars with an even number of edges; and crowns $C_n \odot K_1$. In [3652] they prove the following graphs have edge-odd graceful labelings: P_n ($n > 1$) with a pendent edge attached to each vertex (combs); the graph obtained by appending $2n + 1$ pendent edges to each endpoint of P_2 or P_3; and the graph obtained by subdividing each edge of the star $K_{1,2n}$. In [2057] Kumar and Gayathri showed the existence of edge-odd graceful labeling for $P_m \times P_n$. Jeba Jesintha and Yogalakshmi [1644] proved the Jahangir graph $J_{m,n}$ for $m = 3$ and $n \geq 3$ is edge-odd graceful and, in general, the Jahangir graph $J_{m,n}$ for $m \geq 5$ and odd $n \geq 3$ is also edge-odd graceful. Daoud [893] gave necessary and sufficient conditions for friendship graphs, wheels, helms, webs, double wheels, gears, fans, double fans, and polar grids to admit edge-odd graceful labelings. In [174] Aljohanis and Daoud provided edge-odd graceful labelings five new families of wheel-related graph and proved that S_n are not edge-odd graceful when n is odd.

Valani and Kanani [4039] proved that the corona product of a ladder and K_1, the corona product of slanting ladder and K_1, the H graph of path P_n and the corona product and K_1 are edge-odd graceful graphs. Soleha, Purwanto and D. Rahmadani [3653] showed that alternate triangular snakes and alternate quadrilateral snakes are edge-odd graceful. In [4040] Valani and Kanani proved that double alternate triangular snakes, double alternate quadrilateral snakes, and alternate pentagonal snakes are edge-odd graceful graphs. [4039] new [3653] new [4040] new

Gayathri, Duraisamy, and Tamilselvi [1211] calls a (p, q)-graph with $q \geq p$ *even edge-graceful* if there is an injection f from the set of edges to $\{1, 2, 3, \ldots, 2q\}$ such that the values of the induced mapping f^+ from the vertex set to $\{0, 1, 2, \ldots, 2q-1\}$ given by $f^+(x) = (\Sigma f(xy))(\text{mod } 2q)$ over all edges xy are distinct and even. In [1211] and [1210] Gayathri et al. prove the following: cycles are even edge-graceful if and only if the cycles are odd; even cycles with one pendent edge are even edge-graceful; wheels are even edge-graceful; gears (see §2.2 for the definition) are not even edge-graceful; fans $P_n + K_1$ are even edge-graceful; $C_4 \cup P_m$ for all m are even edge-graceful; $C_{2n+1} \cup P_{2n+1}$ are even edge-graceful; crowns $C_n \odot K_1$ are even edge-graceful; $C_n^{(m)}$ (see §2.2 for the definition) are even edge-graceful; sunflowers (see §3.7 for the definition) are even edge-graceful; triangular snakes (see §2.2 for the definition) are even edge-graceful; closed helms (see §2.2 for the definition) with the center vertex removed are even edge-graceful; graphs decomposable into two odd Hamiltonian cycles are even edge-graceful; and odd order graphs that are decomposable into three Hamiltonian cycles are even edge-graceful. Paley graphs are dense undirected graphs raised from the vertices as elements of an appropriate finite field by joining pairs of vertices that differ by a quadratic residue. Elsawy and Almohammadi [1018] proved that Paley graphs and generalized Paley graphs of prime order are edge graceful, edge-even graceful, and edge-odd graceful graphs. Meng [2487] proved that every tree of odd order with at most two vertices of degree two is edge-graceful. [1018] new [2487] new

In [1210] Gayathri and Duraisamy generalized the definition of even edge-graceful to include (p, q)-graphs with $q < p$ by changing the modulus from $2q$ the maximum of $2q$ and $2p$. With this version of the definition, they have shown that trees of even order are not even edge-graceful whereas, for odd order graphs, the following are even edge-graceful: banana trees (see §2.1 for the definition); graphs obtained joining the centers of two stars by a path; $P_n \odot K_{1,m}$; graphs obtained by identifying an endpoint from each of any number of copies of P_3 and P_2; bistars (that is, graphs obtained by joining the centers of two stars with an edge); and graphs obtained by appending the endpoint of a path to the center of a star. They define *edge-odd graceful* graphs in the analogous way and provide a few results about such graphs. Kathiresan, Muthumari, and Ramalakshmi [1929] proved that $K_n^c \vee 2K_2$ and the flower graph FL_n ($n \geq 3$) odd edge-graceful labelings. They further prove that $P_{a,b}$ (the graph obtained by identifying the end points of b internally disjoint paths each of length a) admit odd edge-graceful labelings when a and b are odd, whereas $P_{a,2}$ is not odd edge-graceful when $a \geq 2$.

In [175] Aljohani and Daoud introduced new families of planar graphs as follows. For $n \geq 3$, a *water wheel* is the graph with vertex set $\{u_0\} \cup \{u_i, v_i, w_i : 1 \leq i \leq n\}$ and edge set $\{u_0u_i, u_0v_i, u_iw_i, v_iw_i : 1 \leq i \leq n\} \cup \{u_iv_{i+1} : 1 \leq i \leq n-1\} \cup \{u_nv_1\}$ (see [175] for

a figures). For $n \geq 3$, a *triangulated water wheel* graph is obtained from the water wheel graph by adding edges u_0w_i for $1 \leq i \leq n$. For $n \geq 3$, a *closed water wheel* graph is obtained from the water wheel graph by adding edges w_iw_{i+1} for $1 \leq i \leq n-1$. They proved that these three families are edge-odd graceful.

For a graph G, Kulli and Muddebihai [2055] define the *lict graph* of G as the graph whose vertex set is the union of the edges of G and the set of cutpoints of G where two vertices are adjacent if and only if the corresponding members of G are adjacent or the corresponding members of G are incident. They define the *litact graph* of G as the graph whose vertex set is the union of the edges of G and the set of cutpoints of G where two vertices are adjacent if and only if the corresponding members of G are adjacent or the corresponding members of G are adjacent or incident. Mirajkar and Sthavarmath [2508] provided edge-odd graceful labeling for the lict graph of P_n for $n > 1$ and odd, the lict graph of the one-point of C_n and P_2, and the litact graph of P_n for $n \geq 4$.

A *subdivided shell graph* is obtained by subdividing the edges in the path of the shell graph. Let $G_1, G_2, \ldots, G_n$ be n subdivided shell graphs of any order. The graph SSG(n) is obtained by adding an edge to apexes of G_i and $G_{i+1}, i = 1, 2, \ldots, n-1$. Jeba Jesintha and Ezhilarasi Hilda [1618] that SSG(2) is odd graceful. In [1612] and [1617] Jeba Jesintha and Ezhilarasi Hilda proved that the subdivided uniform shell bow graphs (that is, double shells in which each shell has the same order) are odd graceful and shell butterfly graphs are edge-odd graceful. Daoud [894] provided necessary and sufficient conditions for $C_m \times P_n$ and $C_m \times C_n$ to be edge-odd graceful.

Gao [1190] has proved the following graphs are odd-graceful: the union of any number of paths; the union of any number of stars; the union of any number of stars and paths; $C_m \cup P_n$; $C_m \cup C_n$; and the union of any number of cycles each of which has order divisible by 4.

If f is an odd-graceful labeling of a bipartite graph G with bipartition (V_1, V_2) such that $\max\{f(u) : u \in V_1\} < \min\{f(v) : v \in V_2\}$, Zhou, Yao, Chen, and Tao [4341] say that f is a *set-ordered odd-graceful labeling* of G. They proved that every lobster is odd-graceful and adding leaves to a connected set-ordered odd-graceful graph is an odd-graceful graph.

In [3365] Seoud and Abdel-Aal determined all odd-graceful graphs of order at most 6 and proved that if G is odd-graceful then $G \cup K_{m,n}$ is odd-graceful. In [3385] Seoud and Helmi proved: if G has an odd-graceful labeling f with bipartition (V_1, V_2) such that $\max\{f(x) : f(x) \text{ is even}, x \in V_1\} < \min\{f(x) : f(x) \text{ is odd}, x \in V_2\}$, then G has an α-labeling; if G has an α-labeling, then $G \odot \overline{K_n}$ is odd-graceful; and if G_1 has an α-labeling and G_2 is odd-graceful, then $G_1 \cup G_2$ is odd-graceful. They also proved the following graphs have odd-graceful labelings: dragons obtained from an even cycle; graphs obtained from a gear graph by attaching a fixed number of pendent edges to each vertex of degree 2 on rim of the wheel of the graph; $C_{2m} \odot \overline{K_n}$; graphs obtained from an even cycle by attaching a fixed number of pendent edges to every other vertex; graphs obtained by identifying an endpoint of a star S_n ($n \geq 3$) with a vertex of an even cycle; the graphs consisting of two even cycles of the same order that share a common vertex with any number of pendent edges attached at the common vertex; and the graphs obtained by

joining two even cycles of the same order by an edge. Seoud, El Sonbaty, and Abd El Rehim [3377] proved that the conjunction $P_m \wedge P_n$ for all $n, m \geq 2$ and the conjunction $K_2 \wedge F_n$ for n even are odd-graceful. Jeba Jesintha and Ezhilarasi Hilda [1610] proved the disjoint union of two subdivided shell graphs is odd-graceful and the one vertex union of three subdivided shells are odd-graceful.

In [2555] and [2556] Moussa proved that $C_m \cup P_n$ is odd-graceful in some cases and gave algorithms to prove that for all $m \geq 2$ the graphs $P_{4r-1;m}$, $r = 1, 2, 3$ and $P_{4r+1;m}$, $r = 1, 2$ are odd-graceful. ($P_{n;m}$ is the graph obtained by identifying the endpoints of m paths each of length n). He also presented an algorithm that showed that closed spider graphs and the graphs obtained by joining one or two copies of P_m to each vertex of the path P_n are odd-graceful. Moussa and Badr [2554] proved that $C_m \odot P_n$ is odd-graceful if and only if m is even (see also [433]). Badr, Moussa, and Kathiresan [433] proved ladders are odd graceful.

Moussa [2557] defines the *tensor product,* $P_m \wedge P_n$, of P_m and P_n as the graph with vertices $v_i^j,\ i = 1, \ldots, n;\ j = 1, \ldots, m$ and edges $v_1^j v_2^{j+1}, v_2^{j+1} v_3^j, \ldots, v_{n-1}^j v_n^{j+1}$ for j odd and $v_1^j v_2^{j-1}, v_2^{j-1} v_3^j, \ldots, v_{n-1}^j v_n^{j-1}$ for j even. He proves that $P_m \wedge P_m$ is odd-graceful.

In [3] Abdel-Aal generalized the notions of shadow graphs and splitting graphs as follows. The m-shadow graph $D_m(G)$ of a connected graph G is constructed by taking m copies of $G_1, G_2, \ldots, G_m$ of G , and joining each vertex u in G_i to the neighbors of the corresponding vertex v in G_j for $1 \leq i, j \leq m$. The m-splitting graph $Spl_m(G)$ of a graph G is obtained by adding to each vertex v of G m new vertices, $v^1, v^2, \ldots, v^m$, such that $v^i,\ 1 \leq i \leq m$ is adjacent to every vertex that is adjacent to v in G_j. Thus the 2-shadow graph is the shadow graph $D_2(G)$ and the 1-splitting graph of G is the splitting graph of G. Abdel-Aal proved the following graphs are odd-graceful: $D_m(P_n), D_m(P_n \oplus \overline{K_2})$ (the symmetric product of P_n and $\overline{K_2}$), $D_m(K_{r,s}),\ Spl_m(P_n),\ Spl_m(K_{1,n})$, and $Spl_m(P_n \oplus \overline{K_2})$.

Vaidya and Bijukumar [3963] proved the following are odd-graceful: graphs obtained by joining two copies of C_n by a path; graphs that are two copies of an even cycle that share a common edge; graphs that are the splitting graph of a star; and graphs that are the tensor product of a star and P_2. Jeba Jesintha, Jaya Glory, and Elakiya Solai [1621] proved that the path unions of caterpillars are odd graceful.

Acharya, Germina, Princy, and Rao [52] proved that every bipartite graph G can be embedded in an odd-graceful graph H. The construction is done in such a way that if G is planar and odd-graceful, then so is H. Varkey and Sunoj [4050] investigate some new families of odd graceful graphs generated from various graph operations on the given graph.

In [791] Chawathe and Krishna extend the definition of odd-gracefulness to countably infinite graphs and show that all countably infinite bipartite graphs that are connected and locally finite have odd-graceful labelings.

Solairaju and Chithra [3651] defined a graph G with q edges to be *edge-odd graceful* if there is a bijection f from the edges of the graph to $\{1, 3, 5, \ldots, 2q-1\}$ such that, when each vertex is assigned the sum of all the edges incident to it mod $2q$, the resulting vertex labels are distinct. They prove they following graphs are odd-graceful: paths with at least 3 vertices; odd cycles; ladders $P_n \times P_2$ ($n \geq 3$); stars with an even number of edges;

and crowns $C_n \odot K_1$. In [3652] they prove the following graphs have edge-odd graceful labelings: P_n $(n > 1)$ with a pendent edge attached to each vertex (combs); the graph obtained by appending $2n+1$ pendent edges to each endpoint of P_2 or P_3; and the graph obtained by subdividing each edge of the star $K_{1,2n}$.

Singhun [3613] proved the following graphs have edge-odd graceful labelings: W_{2n}; $W_n \odot K_1$; and $W_n \odot K_m$, when n is odd, m is even, and n divides m. Seoud and Salim [3400] present edge-odd graceful labelings for the following families of graphs: W_n for $n \equiv 1, 2$ and 3 (mod 4); $C_n \odot \overline{K_{2m-1}}$; even helms; $P_n \odot \overline{K_{2m}}$; and $K_{2,s}$. They also provide two theorems about non edge-odd graceful graphs. Susanti, Ernanto1, and Surodjo [3814] found edge-odd graceful labelings for some classes of prism related graphs.

In [3685] Sridevi, Navaeethakrishnan, Nagarajan, and Nagarajan call a graph G with q edges *odd-even graceful* if there is an injection f from the vertices of G to $\{1, 3, 5, \ldots, 2q+1\}$ such that, when each edge uv is assigned the label $|f(u) - f(v)|$, the resulting edge labels are $\{2, 4, 6, \ldots, 2q\}$. They proved that P_n, combs $P_n \odot K_1$, stars $K_{1,n}, K_{1,2,n}, K_{m,n}$, and bistars $B_{m,n}$ are odd-even graceful.

Sudha and Babu [3715] say a graph G with q edges is *even-even graceful* if there is an injection f from the edges of G to $\{2, 4, 6, \ldots, 2q\}$ such that, the induced map f^+ from $V(G)$ to $\{0, 2, \ldots, 2k-2\}$ defined by $f^*(x) = \Sigma(f(xy) \pmod{2k}$ where $k = \max(p, q)$ is injective and each value is $f^*(x)$ is even. They proved that dumbbells, stars, $C_n \times P_2$, and $K_1 + C_n$ are even-even graceful.

Behera, Mishra, and Nayak [612] proved the following: bistars $B_{r,r}$ are even-even graceful, combs are even-even graceful, the trees obtained by joining and even number of pendent edges to the endpoint of a path are even-even graceful, the graphs obtained by identifying the center of a star and a vertex of C_3 are odd-even graceful, the graphs obtained by identifying the center of a star and a vertex of C_3 and two pendent edges at the other two vertices are odd-even graceful, and the graphs obtained by identifying the center of a star with a vertex of C_n and the endpoints of the star with the opposite vertices of C_n is odd-even graceful.

In [896] Daoud introduced vertex odd graceful labelings as follows. Let G be a graph with q edges. A function f is called a *vertex odd graceful* labeling of G if $f\ E(G) \to \{1, 2, 3, \ldots, 2q\}$ is an injection and the induced function $f^*\ V(G) \to \{1, 3, \ldots, 2q-1\}$ defined as $f^*(u) = \sum_{uv \in E(G)} f(uv) \pmod{2q}$ is also an injection. A graph that admits a vertex odd graceful labeling is called a *vertex odd graceful* graph. Necessary and sufficient conditions for prisms, tori, wheels, fans and books to be vertex odd graceful are given.

3.7 Cordial Labelings

Cahit [738] has introduced a variation of both graceful and harmonious labelings. Let f be a function from the vertices of G to $\{0, 1\}$ and for each edge xy assign the label $|f(x) - f(y)|$. Call f a *cordial labeling* of G if the number of vertices labeled 0 and the number of vertices labeled 1 differ by at most 1, and the number of edges labeled 0 and the number of edges labeled 1 differ at most by 1. Cahit [739] proved the following: every tree is cordial; K_n is cordial if and only if $n \leq 3$; $K_{m,n}$ is cordial for all m and n; the

friendship graph $C_3^{(t)}$ (i.e., the one-point union of t 3-cycles) is cordial if and only if $t \not\equiv 2 \pmod 4$; all fans are cordial; the wheel W_n is cordial if and only if $n \not\equiv 3 \pmod 4$ (see also [977]); maximal outerplanar graphs are cordial; and an Eulerian graph is not cordial if its size is congruent to 2 (mod 4). Kuo, Chang, and Kwong [2071] determine all m and n for which mK_n is cordial. Youssef [4278] proved that every Skolem-graceful graph (see 3.5 for the definition) is cordial. Liu and Zhu [2301] proved that a 3-regular graph of order n is cordial if and only if $n \not\equiv 4 \pmod 8$. In [1525] Imran, Cancan, Ali, Nadeem, Mushtaq, Aslam, Riaz, proved various comb related graphs are cordial. Kastrati, Myrvold, Panjer, and Williams [1916] and [1917] proved that a forest is cordial if and only if it does not have $4k+2$ components and every vertex has odd-degree.

In [696] Bosmia, Patel, Parmar, and Vihol provided necessary and sufficient conditions for the line graph of bistar $B_{n,p}$ to be cordial when $p = n+4m$ and $p = n+4m+2$. Bosmia and Kanani [694] proved that $L(B_{n,n})$ is cordial if and only if $n = t^2$ or $n = (t+1)^2 - 1$. [696] new [694] new

A *k-angular cactus* is a connected graph all of whose blocks are cycles with k vertices. In [739] Cahit proved that a k-angular cactus with t cycles is cordial if and only if $kt \not\equiv 2 \pmod 4$. This was improved by Kirchherr [1966] who showed any cactus whose blocks are cycles is cordial if and only if the size of the graph is not congruent to 2 (mod 4). Kirchherr [1967] also gave a characterization of cordial graphs in terms of their adjacency matrices. Ho, Lee, and Shee [1448] proved: $P_n \times C_{4m}$ is cordial for all m and all odd n; the composition G and H is cordial if G is cordial and H is cordial and has odd order and even size (see §2.3 for definition of composition); for $n \geq 4$ the composition $C_n[K_2]$ is cordial if and only if $n \not\equiv 2 \pmod 4$; the Cartesian product of two cordial graphs of even size is cordial. Ho, Lee, and Shee [1447] showed that a unicyclic graph is cordial unless it is C_{4k+2} and that the generalized Petersen graph (see §2.7 for the definition) $P(n,k)$ is cordial if and only if $n \not\equiv 2 \pmod 4$. Khan [1942] proved that a graph that consisting of a finite number of cycles of finite length joined at a common cut vertex is cordial if and only if the number of edges is not congruent to 2 mod 4.

Du [977] determines the maximal number of edges in a cordial graph of order n and gives a necessary condition for a k-regular graph to be cordial. Riskin [3178] proved that Möbius ladders M_n (see §2.3 for the definition) are cordial if and only if $n \geq 3$ and $n \not\equiv 2 \pmod 4$. (See also [3379].) Diab and Nada [962] show that $P_n \odot P_m$ is cordial; except for n and m both equal to 2 (mod 4), $C_n \odot C_m$ is cordial; and when $n \equiv 2 \pmod 4$ and m is odd, $C_n \odot C_m$ is not cordial. Nada, Elrokh, Elrayes, and Rabie [2627] showed that $P_m \odot P_3^4$ is cordial for all positive m when $n \geq 7$ and $n = 3$. In [3255] Salehi, Mukhin, and Saputro showed that Q_n is cordial for all $n > 1$.

In [1001] El-hay, Alsatami, ELrokh, and Rabie say that a *fourth order of lemniscate graph* is the graph obtained by identifying a common vertex shared by the fourth power two cycles. They provided sufficient conditions for the corona product of paths and fourth order of lemniscate graphs to be cordial. In [1002] El-hay, Alsatami, and ELrokh gave cordial labelings for the fifth power of paths, cycles, fans, wheels, lemniscates and the union of fifth power of paths and cycles graphs.

In [775] Charishma and Nageswari define a *wheel gripped star*, $G_{n,m}(W_m, S_n)$, as the one obtained by attaching the star graph S_m to every vertex in the wheel W_n, excluding

the central vertex. They proved that wheel gripped star graphs, the path union of square of cycles, and the star of square of cycles admit cordial labelings.

Seoud and Abdel Maqusoud [3367] proved that if G is a graph with n vertices and m edges and every vertex has odd degree, then G is not cordial when $m+n \equiv 2 \pmod 4$. They also prove the following: for $m \geq 2$, $C_n \times P_m$ is cordial except for the case $C_{4k+2} \times P_2$; P_n^2 is cordial for all n; P_n^3 is cordial if and only if $n \neq 4$; and P_n^4 is cordial if and only if $n \neq 4, 5$, or 6. Seoud, Diab, and Elsakhawi [3376] have proved the following graphs are cordial: $P_n + P_m$ for all m and n except $(m,n) = (2,2)$; $C_m + C_n$ if $m \not\equiv 0 \pmod 4$ and $n \neq 2 \pmod 4$; $C_n + K_{1,m}$ for $n \not\equiv 3 \pmod 4$ and odd m except $(n,m) = (3,1)$; $C_n + \overline{K_m}$ when n is odd, and when n is even and m is odd; $K_{1,m,n}$; $K_{2,2,m}$; the n-cube; books B_n if and only if $n \not\equiv 3 \pmod 4$; $B(3,2,m)$ for all m; $B(4,3,m)$ if and only if m is even; and $B(5,3,m)$ if and only if $m \not\equiv 1 \pmod 4$ (see §2.4 for the notation $B(n,r,m)$). In [3647] Solairaju and Arockiasamy prove that various families of subgraphs of grids $P_m \times P_n$ are cordial.

Diab [955], [956], and [958] proved the following graphs are cordial: $C_m + P_n$ if and only if $(m,n) \neq (3,3), (3,2)$, or (3,1); $P_m + K_{1,n}$ if and only if $(m,n) \neq (1,2)$; $P_m \cup K_{1,n}$ if and only if $(m,n) \neq (1,2)$; $C_m \cup K_{1,n}$; $C_m + \overline{K_n}$ for all m and n except $m \equiv 3 \pmod 4$ and n odd, and $m \equiv 2 \pmod 4$ and n even; $C_m \cup \overline{K_n}$ for all m and n except $m \equiv 2 \pmod 4$; $P_m + \overline{K_n}$; $P_m \cup \overline{K_n}$; $P_m^2 \cup P_n^2$ except for $(m,n) = (2,2)$ or (3,3); $P_n^2 + P_m$ except for $(m,n) = (3,1), (3,2), (2,2), (3,3)$, and (4,2); $P_n^2 \cup P_m$ except for $(n,m) = (2,2), (3,3)$, and (4,2); $P_n^2 + C_m$ if and only if $(n,m) \neq (1,3), (2,3)$, and $(3,3)$. $P_n + \overline{K_m}$; $C_n + K_{1,m}$ for all $n > 3$ and all m except $n \equiv 3 \pmod 4$; $C_n + K_{1,m}$ for $n \equiv 3 \pmod 4$ ($n \neq 3$) and even $m \geq 2$; and $C_m \times C_n$ if and only if $2mn$ is not congruent to 2 (mod 4).

In [957] Diab proved the graphs $W_n + W_m$ are cordial if and only if one of the following conditions is not satisfied: (i) $(n,m) = (3,3)$, (ii) $n = 3$ and $m \equiv 1 \pmod 4$, (iii) $n \equiv 1 \pmod 4$ and $m \equiv 3 \pmod 4$; the graphs $W_n \cup W_m$ are cordial if and only if one of the following conditions is not satisfied: (i) $n = 3$ and $m \equiv 1 \pmod 4$, (ii) $n \equiv 1 \pmod 4$ and $m \equiv 3 \pmod 4$; the graphs $W_n + P_m$ are cordial if and only if one of the following conditions is not satisfied: (i) $(n,m) = (3,1), (3,2)$ and $(3,3)$, (ii) $n \equiv 3 \pmod 4$ and $m = 1$. They also prove that $W_n \cup P_m$ and $W_n \cup C_m$ are cordial for all m and n and $W_n + C_m$ is cordial if and only if $(m,n) \neq (3,3)$ and $(3,4)$. In [959] Diab showed that the second power of C_n is cordial if and only if $n = 3$ or n is even and greater than 4. He also investigated the cordiality of the join and union of pairs of second power of cycles and graphs consisting of one second power of cycle with one cycle and one path.

In [2625] Nada, Diab, Elrokh, and Sabra proved that $P_n \odot C_m$ is cordial if and only if $\gcd(n,m) \neq 1$ or 3 (mod 4); in [2624] they proved $C_n \odot P_m$ is cordial for all $n \geq 3$ and $m \geq 1$. Nada, Elrokh, and Elshafey [2629] provided necessary and sufficient conditions for $F_n^2 = K_1 + P_n^2$, $F_n^2 + F_m^2$, and $F_n^2 + F_m^2$ to be cordial.

The generalized *Jahangir graph* $J_{m,n}$ $m > 3$, $n > 1$ is a graph on $mn + 1$ vertices, consisting of a cycle C_{mn} with one additional vertex that is adjacent to n vertices of C_{mn} at distance m to each other on C_{mn}. Gajjar and Des [1167] proved $J_{m,n}$ is cordial for all $m > 3$ and $n > 1$, except for $J_{1,4n-1}$.

Youssef [4280] has proved the following: If G and H are cordial and one has even

size, then $G \cup H$ is cordial; if G and H are cordial and both have even size, then $G + H$ is cordial; if G and H are cordial and one has even size and either one has even order, then $G + H$ is cordial; $C_m \cup C_n$ is cordial if and only if $m + n \not\equiv 2 \pmod 4$; mC_n is cordial if and only if $mn \not\equiv 2 \pmod 4$; $C_m + C_n$ is cordial if and only if $(m, n) \neq (3, 3)$ and $\{m \pmod 4, n \pmod 4\} \neq \{0, 2\}$; and if P_n^k is cordial, then $n \geq k + 1 + \sqrt{k-2}$. He conjectures that this latter condition is also sufficient. He confirms the conjecture for $k = 5, 6, 7, 8$, and 9. Elirokh and Rabie [1017] proved $P_n^4 + P_m^4$ and $P_n^4 \cup P_m^4$ are cordial for all $n, m \geq 7$, and $C_n^4 + C_m^4$, and $C_n^4 \cup C_m^4$ are cordial for all n, m except $(n, m) = (7, 7)$.

Lee and Liu [2164] have shown that the complete n-partite graph is cordial if and only if at most three of its partite sets have odd cardinality (see also [977]). Lee, Lee, and Chang [2138] prove the following graphs are cordial: the Cartesian product of an arbitrary number of paths; the Cartesian product of two cycles if and only if at least one of them is even; and the Cartesian product of an arbitrary number of cycles if at least one of them has length a multiple of 4 or at least two of them are even. Ali Al-Shamiri, Elrokh, El-Mashtawye, and Tallah [192] showed that the Cartesian product of a path and a cycle is cordial under some conditions and that the Cartesian product of two paths is cordial. Elrokh, Elmshtaye, and Abd El-hay [1014] provided necessary and sufficient conditions for cone and lemniscate graphs to be cordial.

Shee and Ho [3497] have investigated the cordiality of the one-point union of n copies of various graphs. For $C_m^{(n)}$, the one-point union of n copies of C_m, they prove:

(i) If $m \equiv 0 \pmod 4$, then $C_m^{(n)}$ is cordial for all n;

(ii) If $m \equiv 1$ or $3 \pmod 4$, then $C_m^{(n)}$ is cordial if and only if $n \not\equiv 2 \pmod 4$;

(iii) If $m \equiv 2 \pmod 4$, then $C_m^{(n)}$ is cordial if and only if n is even.

For $K_m^{(n)}$, the one-point union of n copies of K_m, Shee and Ho [3497] prove:

(i) If $m \equiv 0 \pmod 8$, then $K_m^{(n)}$ is not cordial for $n \equiv 3 \pmod 4$;

(ii) If $m \equiv 4 \pmod 8$, then $K_m^{(n)}$ is not cordial for $n \equiv 1 \pmod 4$;

(iii) If $m \equiv 5 \pmod 8$, then $K_m^{(n)}$ is not cordial for all odd n;

(iv) $K_4^{(n)}$ is cordial if and only if $n \not\equiv 1 \pmod 4$;

(v) $K_5^{(n)}$ is cordial if and only if n is even;

(vi) $K_6^{(n)}$ is cordial if and only if $n > 2$;

(vii) $K_7^{(n)}$ is cordial if and only if $n \not\equiv 2 \pmod 4$;

(viii) $K_n^{(2)}$ is cordial if and only if n has the form p^2 or $p^2 + 1$.

For $W_m^{(n)}$, the one-point union of n copies of the wheel W_m with the common vertex being the center, Shee and Ho [3497] show:

(i) If $m \equiv 0$ or $2 \pmod 4$, then $W_m^{(n)}$ is cordial for all n;

(ii) If $m \equiv 3 \pmod 4$, then $W_m^{(n)}$ is cordial if $n \not\equiv 1 \pmod 4$;

(iii) If $m \equiv 1 \pmod 4$, then $W_m^{(n)}$ is cordial if $n \not\equiv 3 \pmod 4$. For all n and all $m > 1$ Shee and Ho [3497] prove $F_m^{(n)}$, the one-point union of n copies of the fan $F_m = P_m + K_1$ with the common point of the fans being the center, is cordial (see also [2267]). The *flag* Fl_m is obtained by joining one vertex of C_m to an extra vertex called the *root*. Shee and Ho [3497] show all $Fl_m^{(n)}$, the one-point union of n copies of Fl_m with the common point

being the root, are cordial. In his 2001 Ph. D. thesis Selvaraju [3346] proves that the one-point union of any number of copies of a complete bipartite graph is cordial. Benson and Lee [621] have investigated the regular windmill graphs $K_m^{(n)}$ and determined precisely which ones are cordial for $m < 14$.

Diab and Mohammedm [961] proved the following: the join of two fans $F_n + F_m$ is cordial if and only if $n, m \geq 4$; $F_n \cup F_m$ is cordial if and only if $(n, m) \neq (1,1)$ or $(2,2)$; $F_n + P_m$ is cordial if and only if $(n, m) \neq (1,2)$, $(2,1)$, $(2,2)$, $(2,3)$, or $(3,2)$; $F_n \cup P_m$ is cordial if and only if $(n, m) \neq (1,2)$; $F_n + C_m$ is cordial if and only if $(n, m) \neq (1,3)$, $(2,3)$ or $(3,3)$; and $F_n \cup C_m$ is cordial if and only if $(n, m) \neq (2, 3)$. Hefnawy, Elsid, and Euat Tallah [1399] gave necessary and sufficient conditions for a cordial labeling of the sum of the second power of the path $P_n^2 + K_{1,m}$ and $P_n^2 \cup K_{1,m}$.

Andar, Boxwala, and Limaye [228], [229], and [232] have proved the following graphs are cordial: helms; closed helms; generalized helms obtained by taking a web (see 2.2 for the definitions) and attaching pendent vertices to all the vertices of the outermost cycle in the case that the number cycles is even; flowers (graphs obtained by joining the vertices of degree one of a helm to the central vertex); sunflower graphs (that is, graphs obtained by taking a wheel with the central vertex v_0 and the n-cycle $v_1, v_2, \ldots, v_n$ and additional vertices $w_1, w_2, \ldots, w_n$ where w_i is joined by edges to v_i, v_{i+1}, where $i+1$ is taken modulo n); multiple shells (see §2.2); and the one point unions of helms, closed helms, flowers, gears, and sunflower graphs, where in each case the central vertex is the common vertex.

In [3004], [3005], [3006], [3011], and [3007] Prajapati and Gajjar provided results about the existence of cordial labelings of graphs obtained from paths, cycles, flower graphs, sunflower graphs, flower snarks, lotus inside a circle graphs, helms, closed helms, armed helms ($W_n \oplus P_2$), and webs by the duplication of vertices and edges. In [1013] Elrokh and Elkom proved that certain classes any four-leaved rose graphs (the one-point union of four cycles of the same length) are cordial. Bansari, Jekil, and Mahadityasinh [473] investigated the cordial labeling for one of the Goldberg snarks and for some special forms of the Goldberg snark. [473] new

Du [978] proved that the disjoint union of $n \geq 2$ wheels is cordial if and only if n is even or n is odd and the number of vertices of in each cycle is not 0 (mod 4) or n is odd and the number of vertices of in each cycle is not 3 (mod 4). Prajapati and Gajjar [3003] prove $\overline{W_n}$ is not cordial if $n \not\equiv 4, 7 \pmod 8$ and $\overline{C_n}$ is not cordial if $n \not\equiv 4, 7 \pmod 8$. Sethuraman and Sankar [3443] showed that the subdivision of wheels are cordial for all $n \geq 4$.

Let $\mathcal{O}$ be the family of all cordial graphs of odd order and odd size for which there is no cordial labeling g such that $e_g(0) - e_g(1) = 1$. Barrientos and Minion [546] proved that if G is a cordial graph such that $G \notin \mathcal{O}$, then the corona $K_1 \odot G$ is cordial. They use this result to prove that $H \odot G$ is cordial when G and H are cordial and G has even order and even size or $G \notin \mathcal{O}$. In addition, $H \odot G$ is cordial when G is a cordial graph of odd order and even size and H is any graph of order m and size $n \in \{m-1, m, m+1\}$. If H is bipartite such that the difference of the cardinalities of its partite sets is at most one, and G is a cordial graph of even order and odd size that admits a cordial labeling g such that $e_g(0) - e_g(1) = 1$, then the corona $H \odot G$ is cordial. Barrientos and Minion

proved the cordiality of certain circulant graphs; they also proved that for every positive integer k, the k-splitting of a cordial graph of even size, results in a cordial graph. They provide sufficient conditions to prove that any super subdivision of a graph G is cordial. They study the cordiality of the join of two cordial graphs, proving that $G + H$ is cordial when G and H have even order and even size, or both have odd order and even size, or both graphs have odd order, odd size, and the dominating weight in both graphs is not 1, or G has even order, odd size, and the dominating weight on both graphs is not the same, or both G and H have odd order, but only one has odd size, and the dominating weight is 0. They also prove that when G is a cordial graph of odd order and even size, the one-point union of t copies of G is cordial.

In [546] Barrientos and Minion provide necessary conditions for the cordiality of coronas of cordial graphs, prove the cordiality of a family of circulant graphs, prove that any splitting graph of a cordial graph of even order and even size is cordial, determine a condition that a graph must satisfy in order that any super subdivision of it is cordial, prove the cordiality of the joint of two cordial graphs, and determine when a one-point union of a cordial graph is cordial.

For positive integers m and n divisible by 4 Venkatesh [4077] constructs graphs obtained by appending a copy of C_n to each vertex of C_m by identifying one vertex of C_n with each vertex of C_m and iterating by appending a copy of C_n to each vertex of degree 2 in the previous step. He proves that the graphs obtained by successive iterations are cordial.

Elumalai and Sethurman [1024] proved: cycles with parallel cords are cordial and n-cycles with parallel P_k-chords (see §2.2 for the definition) are cordial for any odd positive integer k at least 3 and any $n \not\equiv 2 \pmod 4$ of length at least 4. They call a graph H an *even-multiple subdivision graph* of a graph G if it is obtained from G by replacing every edge uv of G by a pair of paths of even length starting at u and ending at v. They prove that every even-multiple subdivision graph is cordial and that every graph is a subgraph of a cordial graph. In [4177] Wen proves that generalized wheels $C_n + mK_1$ are cordial when m is even and $n \not\equiv 2 \pmod 4$ and when m is odd and $n \not\equiv 3 \pmod 4$. Kuppusamy and Guruswamy [2072] show that the subdivision graph of $K_{2,n}$ is graceful for $n \geq 1$ and the subdivision graph of the shell graph $C(n, n-3)$ is graceful for $n \geq 4$.

Vaidya, Ghodasara, Srivastav, and Kaneria investigated graphs obtained by joining two identical graphs by a path. They prove: graphs obtained by joining two copies of the same cycle by a path are cordial [3974]; graphs obtained by joining two copies of the same cycle that has two chords with a common vertex with opposite ends of the chords joining two consecutive vertices of the cycle by a path are cordial [3974]; graphs obtained by joining two rim vertices of two copies of the same wheel by a path are cordial [3976]; and graphs obtained by joining two copies of the same Petersen graph by a path are cordial [3976]. They also prove that graphs obtained by replacing one vertex of a star by a fixed wheel or by replacing each vertex of a star by a fixed Petersen graph are cordial [3976]. In [4015] Vaidya, Ghodasara, Srivastav, and Kaneria investigated graphs obtained by joining two identical cycles that have a chord are cordial and the graphs obtained by starting with copies $G_1, G_2, \ldots, G_n$ of a fixed cycle with a chord that

forms a triangle with two consecutive edges of the cycle and joining each G_i to G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge that is incident with the endpoints of the chords in G_i and G_{i+1} are cordial. Vaidya, Dani, Kanani, and Vihol [3969] proved that the graphs obtained by starting with copies $G_1, G_2, \ldots, G_n$ of a fixed star and joining each center of G_i to the center of G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge are cordial.

Ghodasara, Rokad, and Jadav [1257] prove that the path union of $P_n \times P_n$ is cordial. They also prove that the graph obtained by joining two copies of $P_n \times P_n$ by a path is cordial. Ghodasara and Jadav [1251] prove: the graph obtained by joining a finite number of copies of $P_n \times P_n$ by path is cordial; the star of $P_n \times P_n$ is cordial; and the path union of the star of $P_n \times P_n$ is cordial. Rokad and Patadiya [3199] proved that the shadow graph, splitting graph, and the degree splitting graph of a star are cordial graphs. They also showed that the jewel graph and the jellyfish graph are cordial.

Ghodasara and Rokad prove [1258] the star of $K_{n,n}$ $(n \geq 2)$ is cordial, the path union of $K_{n,n}$ $(n \geq 2)$ is cordial, and the graph obtained by joining two copies of $K_{n,n}$ $(n \geq 2)$ by a path is cordial [1258]. In [1259] the same authors prove that a vertex switching of any non-apex vertex of a wheel graph, a vertex switching of any internal vertex of a flower graph, a vertex switching of any non-apex vertex of a gear graph, and a vertex switching of any non-apex vertex of a shell graph are cordial. In [1260] they proved that a barycentric subdivision of a shell graph, a barycentric subdivision of $K_{n,n}$, and a barycentric subdivision of a wheel are cordial. Ghodasara and Sonchhatra [1261] prove that the graph obtained by joining two copies of the same fan by a path is cordial. They also prove that the star of a fan is cordial and the graph obtained by joining two copies of the star of the same fan by a path is cordial [1261]. Elrokh, Nada, and El-Shafey [1016] showed that $P_k \odot F_m^2$ (F_m is the fan graph with $m+1$ vertices) is cordial for all $k \geq 1$ and $m \geq 4$.

Vaidya, Kanani, Srivastav, and Ghodasara [3984] proved: graphs obtained by subdividing every edge of a cycle with exactly two extra edges that are chords with a common endpoint and whose other end points are joined by an edge of the cycle are cordial; graphs obtained by subdividing every edge of the graph obtained by starting with C_n and adding exactly three chords that result in two 3-cycles and a cycle of length $n-3$ are cordial; graphs obtained by subdividing every edge of a Petersen graph are cordial. Sankar and Sethuramam [3301] showed that the subdivision graph $S(K_2, n)$ is graceful and cordial and the shell graph $S(C(n, n-3))$ is graceful and cordial for $n \geq 4$.

Recall the shell $C(n, n-3)$ is the cycle C_n with $n-3$ cords sharing a common endpoint. Vaidya, Dani, Kanani, and Vihol [3970] proved that the graphs obtained by starting with copies $G_1, G_2, \ldots, G_n$ of a fixed shell and joining common endpoint of the chords of G_i to the common endpoint of the chords of G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge are cordial. Vaidya, Dani, Kanani, and Vihol [3985] define $C_n(C_n)$ as the graph obtained by subdividing each edge of C_n and connecting the new n vertices to form a copy of C_n inscribed the original C_n. They prove that $C_n(C_n)$ is cordial if $n \neq 2 \pmod 4$; the graphs obtained by starting with copies $G_1, G_2, \ldots, G_k$ of $C_n(C_n)$ the graph obtained by joining a vertex of degree 2 in G_i to a vertex of degree 2 in G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge are cordial; and the graphs obtained by joining vertex of degree 2 from one copy of $C_n(C_n)$

to a vertex of degree 2 to another copy of $C_n(C_n)$ by any finite path are cordial. Vaidya and Shah [4011] and [4012] proved that following graphs are cordial: the shadow graph of the bistar $B_{n,n}$, the splitting graph of $B_{n,n}$, the degree splitting graph of $B_{n,n}$, alternate triangular snakes, alternate quadrilateral snakes, double alternate triangular snakes, and double alternate quadrilateral snakes. Prajapati, Asad, and Gajjar [3000] investigated the existence of cordial labelings of graphs obtained from duplication of certain graph elements in quadrilateral snake graphs. In [4014] Vaidya and Shah give cordial labelings of the degree splitting graph of paths, shells, helms, and gears. [3000] new

A graph $C(2n, n-2)$ is called an *alternate shell* if $C(2n, n-2)$ is obtained from the cycle C_{2n} $(v_0, v_1, v_2, \ldots, v_{2n-1})$ by adding $n-2$ chords between the vertex v_0 and the vertices v_{2i+1}, for $1 \leq i \leq n-2$. Sethuraman and Sankar [3442] proved that some graphs obtained by merging alternate shells and joining certain vertices by a path have α-labelings.

Vaidya, Srivastav, Kaneria, and Ghodasara [4016] proved that a cycle with two chords that share a common vertex and the opposite ends of which join two consecutive vertices of the cycle is cordial. For a graph G Vaidya, Ghodasara, Srivastav, and Kaneria [3975] introduced the graph G^* called the *star of* G as the graph obtained by replacing each vertex of the star $K_{1,n}$ by a copy of G and prove that ${C_n}^*$ admits cordial labeling. Vaidya and Dani [3965] proved that the graphs obtained by starting with n copies $G_1, G_2, \ldots, G_n$ of a fixed star and joining each center of G_i to the center of G_{i+1} by an edge as well as each of the centers to a new vertex x_i $(1 \leq i \leq n-1)$ by an edge admit cordial labelings. An *arbitrary supersubdivision* H of a graph G is the graph obtained from G by replacing every edge of G by $K_{2,m}$, where m may vary for each edge arbitrarily. Vaidya and Kanani [3977] proved that arbitrary supersubdivisions of paths and stars admit cordial labelings. Vaidya and Dani [3966] prove that arbitrary supersubdivisions of trees, $K_{m,n}$, and $P_m \times P_n$ are cordial. They also prove that an arbitrary supersubdivision of the graph obtained by identifying an end vertex of a path with every vertex of a cycle C_n is cordial except when n is odd, m_i $(1 \leq i \leq n)$ are odd, and m_i $(n+1 \leq i \leq mn)$ of the K_{2,m_i} are even. Recall for a graph G and a vertex v of G Vaidya, Srivastav, Kaneria, and Kanani [4017] define a *vertex switching* G_v as the graph obtained from G by removing all edges incident to v and adding edges joining v to every vertex not adjacent to v in G. They proved that the graphs obtained by the switching of a vertex in C_n admit cordial labelings. They also show that the graphs obtained by the switching of any arbitrary vertex of cycle C_n with one chord that forms a triangle with two consecutive edges of the cycle are cordial. Moreover they prove that the graphs obtained by the switching of any arbitrary vertex in cycle with two chords that share a common vertex the opposite ends of which join two consecutive vertices of the cycle are cordial.

The *middle graph* $M(G)$ of a graph G is the graph whose vertex set is $V(G) \cup E(G)$ and in which two vertices are adjacent if and only if either they are adjacent edges of G or one is a vertex of G and the other is an edge incident with it. Vaidya and Vihol [4019] prove that the middle graph $M(G)$ of an Eulerian graph is Eulerian with $|E(M(G))| = \sum_{i=1}^{n}(d(v_i)^2 + 2e)/2$. They prove that middle graphs of paths, crowns $C_n \odot K_1$, stars, and tadpoles (that is, graphs obtained by appending a path to a cycle) admit cordial labelings.

Vaidya and Dani [3968] define the *duplication of an edge $e = uv$ of a graph G by a new vertex w* as the graph G' obtained from G by adding a new vertex w and the edges wv and wu. They prove that the graphs obtained by duplication of an arbitrary edge of a cycle and a wheel admit a cordial labeling. Starting with k copies of fixed wheel W_n, $W_n^{(1)}, W_n^{(2)}, \ldots, W_n^{(k)}$, Vaidya, Dani, Kanani, and Vihol [3972] define $G =< W_n^{(1)} : W_n^{(2)} : \ldots : W_n^{(k)} >$ as the graph obtained by joining the center vertices of each of $W_n^{(i)}$ and $W_n^{(i+1)}$ to a new vertex x_i where $1 \leq i \leq k-1$. They prove that $< W_n^{(1)} : W_n^{(2)} : ... : W_n^{(k)} >$ are cordial graphs. Kaneria and Vaidya [1872] define the *index of cordiality* of G as n if the disjoint union of n copies of G is cordial but the disjoint union of fewer than n copies of G is not cordial. They obtain several results on index of cordiality of K_n. In the same paper they investigate cordial labelings of graphs obtained by replacing each vertex of $K_{1,n}$ by a graph G. Kaneria, Jariya, and Karavadiya [1834] proved that the index of cordiality for K_n is at most 6 for n at most 105; the index of cordiality for K_n is at most 4, when n can be expressed as sum of square of two integers; and it is at most 8 when a particular different condition on the edge labels are met. See also [1575].

In [232] Andar et al. define a *t-ply graph* $P_t(u, v)$ as a graph consisting of t internally disjoint paths joining vertices u and v. They prove that $P_t(u, v)$ is cordial except when it is Eulerian and the number of edges is congruent to 2 (mod 4). In [233] Andar, Boxwala, and Limaye prove that the one-point union of any number of plys with an endpoint as the common vertex is cordial if and only if it is not Eulerian and the number of edges is congruent to 2 (mod 4). They further prove that the path union of shells obtained by joining any point of one shell to any point of the next shell is cordial; graphs obtained by attaching a pendent edge to the common vertex of the cords of a shell are cordial; and cycles with one pendent edge are cordial.

For a graph G and a positive integer t, Andar, Boxwala, and Limaye [230] define the *t-uniform homeomorph* $P_t(G)$ of G as the graph obtained from G by replacing every edge of G by vertex disjoint paths of length t. They prove that if G is cordial and t is odd, then $P_t(G)$ is cordial; if $t \equiv 2 \pmod 4$ a cordial labeling of G can be extended to a cordial labeling of $P_t(G)$ if and only if the number of edges labeled 0 in G is even; and when $t \equiv 0 \pmod 4$ a cordial labeling of G can be extended to a cordial labeling of $P_t(G)$ if and only if the number of edges labeled 1 in G is even. In [231] Ander et al. prove that $P_t(K_{2n})$ is cordial for all $t \geq 2$ and that $P_t(K_{2n+1})$ is cordial if and only if $t \equiv 0 \pmod 4$ or t is odd and $n \not\equiv 2 \pmod 4$, or $t \equiv 2 \pmod 4$ and n is even.

In [233] Andar, Boxwala, and Limaya show that a cordial labeling of G can be extended to a cordial labeling of the graph obtained from G by attaching $2m$ pendent edges at each vertex of G. For a binary labeling g of the vertices of a graph G and the induced edge labels given by $g(e) = |g(u) - g(v)|$ let $v_g(j)$ denote the number of vertices labeled with j and $e_g(j)$ denote the number edges labeled with j. Let $i(G) = \min\{|e_g(0) - e_g(1)|\}$ taken over all binary labelings g of G with $|v_g(0) - v_g(1)| \leq 1$. Andar et al. also prove that a cordial labeling g of a graph G with p vertices can be extended to a cordial labeling of the graph obtained from G by attaching $2m + 1$ pendent edges at each vertex of G if and only if G does not satisfy either of the conditions: (1) G has an even number of edges and $p \equiv 2 \pmod 4$; (2) G has an odd number of edges and either $p \equiv 1 \pmod 4$

with $e_g(1) = e_g(0) + i(G)$ or $n \equiv 3 \pmod 4$ and $e_g(0) = e_g(1) + i(G)$. Andar, Boxwala, and Limaye [234] also prove: if g is a binary labeling of the n vertices of graph G with induced edge labels given by $g(e) = |g(u) - g(v)|$ then g can be extended to a cordial labeling of $G \odot \overline{K_{2m}}$ if and only if n is odd and $i(G) \equiv 2 \pmod 4$; $K_n \odot \overline{K_{2m}}$ is cordial if and only if $n \neq 4 \pmod 8$; $K_n \odot \overline{K_{2m+1}}$ is cordial if and only if $n \neq 7 \pmod 8$; if g is a binary labeling of the n vertices of graph G with induced edge labels given by $g(e) = |g(u) - g(v)|$ then g can be extended to a cordial labeling of $G \odot C_t$ if $t \neq 3$ mod 4, n is odd and $e_g(0) = e_g(1)$. For any binary labeling g of a graph G with induced edge labels given by $g(e) = |g(u) - g(v)|$ they also characterize in terms of $i(G)$ when g can be extended to graphs of the form $G \odot \overline{K_{2m+1}}$.

For graphs $G_1, G_2, \ldots, G_n$ ($n \geq 2$) that are all copies of a fixed graph G, Shee and Ho [3498] call a graph obtained by adding an edge from G_i to G_{i+1} for $i = 1, \ldots, n-1$ a *path union* of G (the resulting graph may depend on how the edges are chosen). Among their results they show the following graphs are cordial: path-unions of cycles; path-unions of any number of copies of K_m when $m = 4, 6$, or 7; path-unions of three or more copies of K_5; and path-unions of two copies of K_m if and only if $m-2, m$, or $m+2$ is a perfect square. They also show that there exist cordial path-unions of wheels, fans, unicyclic graphs, Petersen graphs, trees, and various compositions.

Lee and Liu [2164] give the following general construction for the forming of cordial graphs from smaller cordial graphs. Let H be a graph with an even number of edges and a cordial labeling such that the vertices of H can be divided into t parts $H_1, H_2, \ldots, H_t$ each consisting of an equal number of vertices labeled 0 and vertices labeled 1. Let G be any graph and $G_1, G_2, \ldots, G_t$ be any t subsets of the vertices of G. Let (G, H) be the graph that is the disjoint union of G and H augmented by edges joining every vertex in G_i to every vertex in H_i for all i. Then G is cordial if and only if (G, H) is. From this it follows that: all generalized fans $F_{m,n} = \overline{K_m} + P_n$ are cordial; the generalized bundle $B_{m,n}$ is cordial if and only if m is even or $n \not\equiv 2 \pmod 4$ ($B_{m,n}$ consists of $2n$ vertices $v_1, v_2, \ldots, v_n, u_1, u_2, \ldots, u_n$ with an edge from v_i to u_i and $2m$ vertices $x_1, x_2, \ldots, x_m, y_1, y_2, \ldots, y_m$ with x_i joined to v_i and y_i joined to u_i); if m is odd the generalized wheel $W_{m,n} = \overline{K_m} + C_n$ is cordial if and only if $n \not\equiv 3 \pmod 4$. If m is even, $W_{m,n}$ is cordial if and only if $n \not\equiv 2 \pmod 4$; a complete k-partite graph is cordial if and only if the number of parts with an odd number of vertices is at most 3.

Sethuraman and Selvaraju [3451] have shown that certain cases of the union of any number of copies of K_4 with one or more edges deleted and one edge in common are cordial. Youssef [4284] has shown that the kth power of C_n is cordial for all n when $k \equiv 2 \pmod 4$ and for all even n when $k \equiv 0 \pmod 4$. Ramanjaneyulu, Venkaiah, and Kothapalli [3112] give cordial labelings for a family of planar graphs for which each face is a 3-cycle and a family for which each face is a 4-cycle. Acharya, Germina, Princy, and Rao [52] prove that every graph G can be embedded in a cordial graph H. The construction is done in such a way that if G is planar or connected, then so is H.

Recall from §2.7 that a graph H is a *supersubdivision* of a graph G, if every edge uv of G is replaced by $K_{2,m}$ (m may vary for each edge) by identifying u and v with the two vertices in $K_{2,m}$ that form the partite set with exactly two members. Vaidya and

Kanani [3977] prove that supersubdivisions of paths and stars are cordial. They also prove that supersubdivisions of C_n are cordial provided that n and the various values for m are odd.

Raj and Koilraj [3084] proved that the splitting graphs of $P_n, C_n, K_{m,n}, W_n, nK_2$, and the graphs obtained by starting with k copies of stars $K_{1,n}^{(1)}, K_{1,n}^{(2)}, \ldots, K_{1,n}^{(k)}$ and joining the central vertex of $K_{1,n}^{(p-1)}$ and $K_{1,n}^{(p)}$ to a new vertex x_{p-1} for each $2 \leq p \leq k$ are cordial.

Seoud, El Sonbaty, and Abd El Rehim [3377] proved the following graphs are cordial: $K_{1,l,m,n}$ when mn is even; $P_m + K_{1,n}$ if n is even or n is odd and $(m \neq 2)$; the conjunction graph $P_4 \wedge C_n$ is cordial if n is even; and the join of the one-point union of two copies of C_n and K_1.

Recall $< K_{1,n_1}, \ldots, K_{1,n_t} >$ is the graph obtained by starting with the stars $K_{1,n_1}, \ldots, K_{1,n_t}$ and joining the center vertices of K_{1,n_i} and $K_{1,n_{i+1}}$ to a new vertex v_i where $1 \leq i \leq k-1$. Kaneria, Jariya, and Meghpara [1838] proved that $< K_{1,n_1}, \ldots, K_{1,n_t} >$ is cordial and every graceful graph with $|v_f(\text{odd}) - v_f(\text{even})| \leq 1$ is cordial. Kaneria, Meghpara, and Makadia [1866] proved that the cycle of complete graphs $C(t \cdot K_{m,n})$ and the cycle of wheels $C(t \cdot W_n)$ are cordial. Kaneria, Makadia, and Meghpara [1853] proved that the cycle of cycles $C(t \cdot C_n)$ is cordial for $t \geq 3$. Kaneria, Makadia, and Meghpara [1854] proved that a star of K_n and a cycle of n copies of K_n are cordial. Kaneria, Viradia, Jariya, and Makadia [1874] proved that the cycle of paths $C(t \cdot P_n)$ is cordial, product cordial (see Section 7.5), and total edge product cordial.

Jeba Jesintha and Subashini proved the following graphs are cordial: the cycle of vertex switching of cycles [1624]; the path union of vertex switching of wheels in increasing order [1633]; the path union of jelly fish graphs is cordial and cycle of jelly fish graphs [1634]; the star of fixed trees of diameter three tree [1626]; and the path union of vertex switching of cycles in increasing order [1628]. In [1643] Jeba Jesintha, Vinodhini, and Lakshmi, proved that a star glued with subdivided shell graph and super subdivision of circular ladders $(C_n \times K_2)$ admit cordial labelings. In [1600] Jeba Jesintha and Devakirubanithi call a planar graph with $n = 3i$ vertices, for $i = 1, 2, 3, \ldots$ that form a sequence of $n/3$ triangles where pairs of corresponding vertices on consecutive triangles are joined in the sequence *nested triangles*. They proved nested triangles, the shadow graphs of the nested triangles, and the double graph of the nested triangle graphs admit cordial labelings. A *blade* graph is obtained from a $a_0a_1 \cdots a_m$ by adding new edges joining a_0 to each of $a_2, a_3, \ldots a_{m-1}$ and new edges joining a_m to each of $a_2, a_3, \ldots a_{m-1}$. Devakirubanithi1, Jesintha, and Charishma [935] proved that blade graphs and their one-point unions admits cordial labelings.

Cordial labelings and variations of them for fractal graphs are given in [3320] and [3319].

Cahit [744] calls a graph *H-cordial* if it is possible to label the edges with the numbers from the set $\{1, -1\}$ in such a way that, for some k, at each vertex v the sum of the labels on the edges incident with v is either k or $-k$ and the inequalities $|v(k) - v(-k)| \leq 1$ and $|e(1) - e(-1)| \leq 1$ are also satisfied, where $v(i)$ and $e(j)$ are, respectively, the number of vertices labeled with i and the number of edges labeled with j. He calls a graph *H_n-cordial* if it is possible to label the edges with the numbers from the set $\{\pm 1, \pm 2, \ldots, \pm n\}$ in such

a way that, at each vertex v the sum of the labels on the edges incident with v is in the set $\{\pm 1, \pm 2, \ldots, \pm n\}$ and the inequalities $|v(i) - v(-i)| \leq 1$ and $|e(i) - e(-i)| \leq 1$ are also satisfied for each i with $1 \leq i \leq n$. Among Cahit's results are: $K_{n,n}$ is H-cordial if and only if $n > 2$ and n is even; and $K_{m,n}, m \neq n$, is H-cordial if and only if $n \equiv 0 \pmod 4$, m is even and $m > 2, n > 2$. Unfortunately, Ghebleh and Khoeilar [1250] have shown that other statements in Cahit's paper are incorrect. In particular, Cahit states that K_n is H-cordial if and only if $n \equiv 0 \pmod 4$; W_n is H-cordial if and only if $n \equiv 1 \pmod 4$; and K_n is H_2-cordial if and only if $n \equiv 0 \pmod 4$ whereas Ghebleh and Khoeilar instead prove that K_n is H-cordial if and only if $n \equiv 0$ or $3 \pmod 4$ and $n \neq 3$; W_n is H-cordial if and only if n is odd; K_n is H_2-cordial if $n \equiv 0$ or $3 \pmod 4$; and K_n is not H_2-cordial if $n \equiv 1 \pmod 4$. Ghebleh and Khoeilar also prove every wheel has an H_2-cordial labeling. In [1111] Freeda and Chellathurai prove that the following graphs are H_2-cordial: the join of two paths, the join of two cycles, ladders, and the tensor product $P_n \otimes P_2$. They also prove that the join of W_n and W_m where $n + m \equiv 0 \pmod 4$ is H-cordial. Cahit generalizes the notion of H-cordial labelings in [744].

A graph $G(V, E)$ is called H_k- *cordial* if it has an H-cordial labeling f such that for each edge e and each vertex v of G have the label $1 \leq |f(e)| \leq k$, $1 \leq |f(v)| \leq k$ and $|v_f(i) - v_f(-i))| \leq 1$, $|e_f(i)) - e_f(-i)| \leq 1$ for each i with $1 \leq i \leq k$. Ratilal and Parmar [3147] investigated H_k-cordial labelings of triangular snakes, double triangular snakes, triple triangular snakes, alternate triangular snakes, double alternate triangular snakes, irregular triangular snakes, quadrilateral snakes, double quadrilateral snakes, alternate quadrilateral snakes, and irregular quadrilateral snakes. Joshi and Parmar [1801] investigated the H-, H_2- and H_3-cordiality of the following snakes: triangular, double triangular, triple triangular, quadrilateral, double quadrilateral, alternate triangular, double alternate triangular, irregular triangular, quadrilateral, double quadrilateral, alternate quadrilateral, and irregular quadrilateral. Joshi and Pamar [1801] investigated H_k-cordial labeling of p-triangular, m-polygonal snakes, double m-polygonal snakes, alternate m-polygonal snakes, double alternate m-polygonal snakes, irregular m-polygonal snakes, and double irregular m-polygonal snakes.

Cahit and Yilmaz [748] call a graph *E_k-cordial* if it is possible to label the edges with the numbers from the set $\{0, 1, 2, \ldots, k-1\}$ in such a way that, at each vertex v, the sum of the labels on the edges incident with v modulo k satisfies the inequalities $|v(i) - v(j)| \leq 1$ and $|e(i) - e(j)| \leq 1$, where $v(s)$ and $e(t)$ are, respectively, the number of vertices labeled with s and the number of edges labeled with t. Cahit and Yilmaz prove the following graphs are E_3-cordial: P_n $(n \geq 3)$; stars S_n if and only if $n \not\equiv 1 \pmod 3$; K_n $(n \geq 3)$; C_n $(n \geq 3)$; friendship graphs; and fans F_n $(n \geq 3)$. They also prove that S_n $(n \geq 2)$ is E_k-cordial if and only if $n \not\equiv 1 \pmod k$ when k is odd or $n \not\equiv 1 \pmod{2k}$ when k is even and $k \neq 2$. Ni, Liu, and Lu [2669] demonstrate the E_3-cordiality of W_n, $P_m \times P_n$, $K_{m,n}$, and trees.

Bapat and Limaye [490] provide E_3-cordial labelings for: K_n $(n \geq 3)$; snakes whose blocks are all isomorphic to K_n where $n \equiv 0$ or $2 \pmod 3$; the one-point union of any number of copies of K_n where $n \equiv 0$ or $2 \pmod 3$; graphs obtained by attaching a copy of K_n where $n \equiv 0$ or $3 \pmod 3$ at each vertex of a path; and $K_m \odot K_n$. Sridharan and

Umarani [3689] proved: for odd $n > 1$ and $k \geq 2$, $P_n \odot K_1$ is E_k-cordial; for n even and $n \neq k/2$, $P_n \odot K_1$ is E_k-cordial; and certain cases of fans are E_k-cordial. Youssef [4281] gives a necessary condition for a graph to be E_k-cordial for certain k. He also gives some new families of E_k-cordial graphs and proves Lee's [2195] conjecture about the edge-gracefulness of the disjoint union of two cycles. Venkatesh, Salah, and Sethuraman [4082] proved that C_{2n+1} snakes and C_{2n+1}^{2t} are E_2-cordial. Liu, Liu, and Wu [2300] provide two necessary conditions for a graph G to be E_k-cordial and prove that every P_n $(n \geq 3)$ is E_p-cordial if p is odd. They also discuss the E_2-cordiality of a graph G under the condition that some subgraph of G has a 1-factor. Liu and Liu [2299] proved that a graph with no isolated vertex is E_2-cordial if and only if it does not have order $4n + 2$. Bapat and Limaye [491] prove that helms, one point unions of helms, and path unions of helms are E_3-cordial. Jinnah and Beena [1782] prove the graphs P_n $(n \geq 3), C_n$ where $n \neq 4 \bmod 8$, and K_n $(n \geq 3)$ are E_4-cordial graphs. They also prove that every graph of order at least 3 is a subgraph of an E_4-cordial graph.

Hovey [1452] introduced a simultaneous generalization of harmonious and cordial labelings. For any Abelian group A (under addition) and graph $G(V, E)$ he defines G to be *A-cordial* if there is a labeling of V with elements of A such that for all a and b in A when the edge ab is labeled with $f(a) + f(b)$, the number of vertices labeled with a and the number of vertices labeled b differ by at most one and the number of edges labeled with a and the number labeled with b differ by at most one. In the case where A is the cyclic group of order k, the labeling is called *k-cordial*. With this definition we have: if $G(V, E)$ is a graph with $|E| \geq |V| - 1$ then $G(V, E)$ is harmonious if and only if G is $|E|$-cordial; G is cordial if and only if G is 2-cordial.

Let A denote the multiplicative group $\{1, -1, i, -i\}$. In [700], Boxwala, Phadke, Shinde, and Mundlik defined the notion of *double uniform* $(t_1^{\ell_{-1}}, t_2^{\ell_2})$-ply graph and proved that it is group A-cordial for each path of length at least 5 by explicitly giving labelings for all possible cases that arise. [700] new

Hovey obtained the following: caterpillars are k-cordial for all k; all trees are k-cordial for $k = 3, 4$, and 5; odd cycles with pendent edges attached are k-cordial for all k; cycles are k-cordial for all odd k; for k even, C_{2mk+j} is k-cordial when $0 \leq j \leq \frac{k}{2} + 2$ and when $k < j < 2k$; $C_{(2m+1)k}$ is not k-cordial; K_m is 3-cordial; and, for k even, K_{mk} is k-cordial if and only if $m = 1$.

Hovey advances the following conjectures: all trees are k-cordial for all k; all connected graphs are 3-cordial; and C_{2mk+j} is k-cordial if and only if $j \neq k$, where k and j are even and $0 \leq j < 2k$. The last conjecture was verified by Tao [3865]. Tao's result combined with those of Hovey show that for all positive integers k the n-cycle is k-cordial with the exception that k is even and $n = 2mk + k$. Tao also proved that the crown with $2mk + j$ vertices is k-cordial unless $j = k$ is even, and for $4 \leq n \leq k$ the wheel W_n is k-cordial unless $k \equiv 5 \pmod 8$ and $n = (k + 1)/2$. In [3920] Tuczyński, Wenus, and Węsek proved a conjecture of Cichacz, Görlich, and Tuza [850] that all hypertrees are 2-cordial. They also proved that all hypertree are 3-cordial.

In [2809] Patrias and Pechenik initiated the study of classes of finite Abelian groups A for which particular graphs are A-cordial. Their results include: P_{2m} and P_{2m+1} are

not ${Z_2}^m$-cordial, all paths are A-cordial when A is an Abelian group of odd order, if A is an Abelian group of order n and P_n is A-cordial, then all paths are A-cordial, and P_{2n} is A-cordial when $A = Z_2 \times Z_k$ and $n = |A|$.

They conjecture that for a finite Abelian group A, all paths are A-cordial if and only if A has an element of order greater than 2. Cichacz [842] proved that all cycle graphs are A-cordial for any Abelian group A of odd order. In [1908] and [1909] Chidambaram, Athisayanathan, and Ponraj proved that hypercubes, books, $C_n \times K_2$, and $P_n \times K_3$ and the splitting graphs of paths, cycles, and wheels are $\{1, -1, i, -i\}$-cordial. Since the group $\{1, -1, i, -i\}$ is cyclic, this is same as 4-cordial. In [3064] Radha, Venkatesan, Vitaldas, and Perumal prove that triangular ladders, alternate triangular snakes, alternate quadrilateral snakes, and double triangular snakes admit $\{1, -1, i, -i\}$ cordial labelings.

Erickson [1041] et al. showed that the friendship graph F_n is Z_{3m}-cordial and conjectured that F_n is Z_m-cordial except when n is even and not divisible by 4 and $m = 3n/d$, where d is odd.

In [4288] Youssef and Al-Kuleab proved the following: if G is a (p_1, q_1) k-cordial graph and G is a (p_2, q_2) k-cordial graph with p_1 or $p_2 \equiv 0 \pmod{k}$ and q_1 or $q_2 \equiv 0 \pmod{k}$, then $G + H$ is k-cordial; if G is a (p_1, q_1) 4-cordial graph and G is a (p_2, q_2) 4-cordial graph with p_1 or $p_2 \not\equiv 2 \pmod{4}$ and q_1 or $q_2 \equiv 0 \pmod{k}$, then $G + H$ is 4-cordial; and $K_{m,n,p}$ is 4-cordial if and only if $(m, n, p) \mod 4 \not\equiv (0, 2, 2)$ or $(2, 2, 2)$.

In [4279] Youssef obtained the following results: C_{2k} with one pendent edge is not $(2k + 1)$-cordial for $k > 1$; K_n is 4-cordial if and only if $n \leq 6$; C_n^2 is 4-cordial if and only if $n \not\equiv 2 \pmod{4}$; and $K_{m,n}$ is 4-cordial if and only if $n \not\equiv 2 \pmod{4}$; He also provides some necessary conditions for a graph to be k-cordial. Driscol [987] proved that all trees are 7-cordial.

Modha and Kanani [2536] prove that the following graphs have a 5-cordial labeling: the shadow graph of a path and a cycle, graphs obtained by one point duplication and duplication of an edge by a vertex in cycle, and the graph obtained by the barycentric subdivision of wheel. In [2530] Modha and Kanani proved prisms, webs, flowers, and closed helms admit 5-cordial labelings. In [2531] they proved that fans are k-cordial for all k and double fans are k-cordial for all odd k and $n = (k + 1)/2$. In [2533] they proved that the following graphs are k-cordial: W_n for odd k, $n = mk + j, m \geq 0$, $1 \leq j \leq k - 1$ except for $j = (k - 1)/2$; the total graphs of paths (recall $T(P_n)$ has vertex set $V(P_n) \cup E(P_n)$ with two vertices adjacent whenever they are neighbors in P_n); the square C_n^2 for odd $k \leq n$; the path union of n copies of C_k where k is odd; and C_n with one pendent edge for odd $k \leq n$. Rathod and Kanani [3142] proved P_n^2 is k-cordial for all k and cycles with a single pendent edge are k-cordial for all even k. In [3140] Rathod and Kanani proved the middle graph, total graph, and splitting graph of a path are 4-cordial and P_n^2 and triangular snakes are 4-cordial.

Modha and Kanani [2534] prove the following graphs are k-cordial: $P_m \times C_k$, $P_m \times C_{k+1}$, $P_m \times C_{k+3}$ for all odd k and $m \geq 2$, and $P_m \times C_{2k-1}$ for all odd $k, m \geq 2$ and $m \neq tk$. Rathod and Kanani [3142] [3145] prove that following graphs are 4-cordial: the splitting graph of $K_{1,n}$; triangular books; and the one point union any number of copies of the fan f_3; braid graphs; triangular ladders; and irregular quadrilateral snakes obtained from the

path P_n with consecutive vertices $u_1, u_2, \ldots, u_n$ and new vertices $v_1, v_2, \ldots, v_{n-2}, w_1, w_2$, and edges $u_i v_i,\ w_i u_{i+2},\ v_i w_i$ for all $1 \leq i \leq n-2$. Rathod and Kanani [3144] prove wheels, fans, friendship graphs, double fans, and helms are 5-cordial. Driscoll, Krop, and Nguyen [986] proved that all trees are 6-cordial. In [1825], [1826], and [2532] Kanani and Modha prove that fans, friendship graphs, ladders, double fans, double wheels, wheels, helms, closed helms, and webs are 7-cordial graphs and wheels, fans and friendship graphs, gears, double fans, and helms are 4-cordial graphs. In [3323] Sathish Narayanan and Vijayaragavan obtained 3-divisor cordial labelings for graphs derived from paths.

Cichacz, Görlich and Tuza [850] extended the definition of k-cordial labeling for hypergraphs. They presented various sufficient conditions on a hypertree H (a connected hypergraph without cycles) to be k-cordial. From their theorems it follows that every k-uniform hypertree is k-cordial, and every hypertree with odd order or size is 2-cordial. Modha and Kanani [2535] prove the following graphs are k-cordial for all k: bistars, restricted square graphs $B^2_{n,n}$, the one-point union of C_3 and $K_{1,n}$, and $P_n \odot K_1$.

In [3447] Sethuraman and Selvaraju present an algorithm that permits one to start with any non-trivial connected graph G and successively form supersubdivisions (see §2.7 for the definition) that are cordial in the case that every edge in G is replaced by $K_{2,m}$ where m is even. Sethuraman and Selvaraju [3446] also show that the one-vertex union of any number of copies of $K_{m,n}$ is cordial and that the one-edge union of k copies of shell graphs $C(n, n-3)$ (see §2.2) is cordial for all $n \geq 4$ and all k. They conjectured that the one-point union of any number of copies of graphs of the form $C(n_i, n_i - 3)$ for various $n_i \geq 4$ is cordial. This was proved by Yue, Yuansheng, and Liping in [4299]. Riskin [3180] claimed that K_n is $(Z_2 \times Z_2)$-cordial if and only if n is at most 3 and $K_{m,n}$ is $(Z_2 \times Z_2)$-cordial if and only if $(m, n) \neq (2, 2)$. (Many authors use V_4 to denote $Z_2 \times Z_2$.) However, Pechenik and Wise [2814] report that the correct statement for $K_{m,n}$ is $K_{m,n}$ is $(Z_2 \times Z_2)$-cordial if and only if m and n are not both congruent to 2 mod 4. Seoud and Salim [3399] gave an upper bound on the number of edges of a graph that admits a $(Z_2 \oplus Z_2)$-cordial labeling in terms the number of vertices. Rathod and Kanani [3143] prove the following graphs are $(Z_2 \times Z_2)$-cordial for all n and m: $C_n \odot mK_1,\ C_n \odot K_2$, and graphs obtained by appending a single edge to one vertex of C_n. In Rathod and Kanani [3146] and [3141] proved the following graphs are $(Z_2 \times Z_2)$-cordial: alternate triangular snakes, alternate double triangular snakes, alternate triple triangular snakes, quadrilateral snakes, alternate quadrilateral snakes, double quadrilateral snakes, and double alternate quadrilateral snakes.

In [2814] Pechenik and Wise investigate $Z_2 \times Z_2$-cordiality of complete bipartite graphs, paths, cycles, ladders, prisms, and hypercubes. They proved that all complete bipartite graphs are $Z_2 \times Z_2$-cordial except $K_{m,n}$ where $m, n \equiv 2 \bmod 4$; all paths are $Z_2 \times Z_2$-cordial except P_4 and P_5; all cycles are $Z_2 \times Z_2$-cordial except C_4, C_5, C_k, where $k \equiv 2 \bmod 4$; and all ladders $P_2 \times P_k$ are $Z_2 \times Z_2$-cordial except C_4. They also introduce a generalization of A-cordiality involving digraphs and quasigroups, and show that there are infinitely many Q-cordial digraphs for every quasigroup Q. Jinnah and Nair [1783] proved that all trees except P_4 and P_5 are $Z_2 \times Z_2$-cordial and the graphs obtained by subdividing the pendent edges of $C_n \odot K_1$ are $Z_2 \times Z_2$ -cordial for all n.

Cairnie and Edwards [751] have determined the computational complexity of cordial and k-cordial labelings. They prove the conjecture of Kirchherr [1967] that deciding whether a graph admits a cordial labeling is NP-complete. As a corollary, this result implies that the same problem for k-cordial labelings is NP-complete. They remark that even the restricted problem of deciding whether connected graphs of diameter 2 have a cordial labeling is also NP-complete.

For a (p, q) graph G and a bijection f from $V(G)$ to $\{1, 2, \ldots, p\}$ Ponraj, Annathurai, and Kala [2858] introduced a new graph labeling as follows. For each edge uv assign the remainder when $f(u)$ is divided by $f(v)$ or when $f(v)$ is divided by $f(u)$ depending on whether $f(u) \geq f(v)$ or $f(v) \geq f(u)$. The function f is called a *remainder cordial* labeling of G if $|\eta_e - \eta_o| \leq 1$ where η_e and η_o respectively denote the number of edges labeled with even integers and the number of edges labeled with odd integers. A graph G with a remainder cordial labeling is called a *remainder cordial* graph. In [2858] and [2863] they proved that the following graphs are remainder cordial: paths, cycles, stars, bistars, crowns, combs, $K_{2,n}$, $S(K_{1,n})$, $S(B_{n,n})$, P_n^2, wheels, subdivisions of wheels, $K_{2,2n}$, and the graph obtained by subdividing the pendent edges of the bistar $B_{n,n}$. They also proved the following star related graphs are remainder cordial: $K_{1,n} \cup B_{n,n}$, $P_n \cup K_{1,n}$, $P_n \cup B_{n,n}$, $K_{1,n} \cup S(K_{1,n})$, $K_{1,n} \cup S(B_{n,n})$, $P_n^2 \cup K_{1,n}$, ${P_n}^2 \cup B_{n,n}$, and $S(K_{1,n}) \cup S(B_{n,n})$. They conjecture that K_n is remainder cordial if and only if $n \leq 3$. Ponraj, Annathurai, and Kala [2859] generalize remainder cordial labelings as follows. Let f be a function from $V(G)$ to $\{1, 2, \ldots, k\}$ where $2 < k \leq |V(G)|$. For each edge uv assign the remainder when $f(u)$ is divided by $f(v)$ or when $f(v)$ is divided by $f(u)$ depending on whether $f(u) \geq f(v)$ or $f(v) \geq f(u)$. The function f is called a *k-remainder cordial* labeling of G if $|v_f(i) - v_f(j)| \leq 1$, for $i, j \in \{1, \ldots, k\}$ where $v_f(x)$ denote the number of vertices labeled with x and $|\eta_e - \eta_o| \leq 1$ where η_e and η_o respectively denote the number of edges labeled with even integers and the number of edges labeled with odd integers. A graph that admits a k-remainder cordial labeling is called a *k-remainder cordial* graph. In [2859], [263], [264], and [2864] they proved the following. Every graph is a subgraph of a connected k-remainder cordial graph for $k \geq 4$. Note that when $k = 2$, the number of edges with label 0 is q so there does not exists a 2-remainder cordial labeling. They further investigate the 3-remainder cordial labeling behavior of paths, cycles, stars, combs, crowns, wheels, fans, squares of paths, subdivisions of wheels, subdivisions of stars, subdivisions of combs, armed crowns, and $K_{1,n} \odot K_2$. They further proved that W_n is 3-remainder cordial if and only if $n \equiv 1 \pmod 3$, $K_{1,n}$ is 3-remainder cordial if and only if $n \in \{1, 2, 3, 4, 5, 6, 7, 9\}$, and K_n is 3-remainder cordial if and only if $n \leq 3$. In [2860], [2861], and [2862] Ponraj, Annathurai, and Kala proved the following graphs are 4-remainder cordial: complete graphs, paths, cycles, crowns, stars, bistars, books, subdivisions of stars, subdivisions of bistars, subdivisions of jelly fish, flowers, sunflowers, lotuses inside a circle, friendship graphs, webs, triangular snakes, durer graphs, planar grids, mongolian tents, prisms, dragon graphs $C_m@P_n$ (the graph obtained by identifying an endpoint of P_n with one vertex of C_m), crossed prisms CP_{2n}, and $K_2 + mK_1$ ($m \equiv 0, 1, 3 \pmod 4$). They also investigate the 4-remainder cordial labeling of $L_n \odot mK_1$, $L_n \odot K_2$, $L_n \odot mK_1$, $P_n \odot K_1$, $P_n \odot 2K_1$, $C_n \odot K_1$, and $S(P_n \odot K_1)$.

In Bapat [481] introduces the following new labeling. A graph $G(V, E)$ has a L-*cordial* labeling if there is a bijection f from $E(G)$ to $\{1, 2, \ldots, |E|\}$ that assigns 0 to a vertex v if the largest label on the edges incident to v is even and assigns 1 to v otherwise and this assignment satisfies the condition that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1. A graph that admits an L-cordial labeling is called a L-*cordial* graph. He shows that stars, path, cycles, and triangular snakes are L-cordial graphs

In [784] Chartrand, Lee, and Zhang introduced the notion of uniform cordiality as follows. Let f be a labeling from $V(G)$ to $\{0, 1\}$ and for each edge xy define $f^*(xy) = |f(x) - f(y)|$. For $i = 0$ and 1, let $v_i(f)$ denote the number of vertices v with $f(v) = i$ and $e_i(f)$ denote the number of edges e with $f^*(e) = i$. They call a such a labeling f *friendly* if $|v_0(f) - v_1(f)| \leq 1$. A graph G for which every friendly labeling is cordial is called *uniformly cordial*. They prove that a connected graph of order $n \geq 2$ is uniformly cordial if and only if $n = 3$ and $G = K_3$, or n is even and $G = K_{1,n-1}$.

In [3178] Riskin introduced two measures of the noncordiality of a graph. He defines the *cordial edge deficiency* of a graph G as the minimum number of edges, taken over all friendly labelings of G, needed to be added to G such that the resulting graph is cordial. If a graph G has a vertex labeling f using 0 and 1 such that the edge labeling f_e given by $f_e(xy) = |f(x) - f(y)|$ has the property that the number of edges labeled 0 and the number of edges labeled 1 differ by at most 1, the *cordial vertex deficiency* defined as ∞. Riskin proved: the cordial edge deficiency of K_n $(n > 1)$ is $\lfloor \frac{n}{2} \rfloor - 1$; the cordial vertex deficiency of K_n is $j - 1$ if $n = j^2 + \delta$, when δ is $-2, 0$ or 2, and ∞ otherwise. In [3178] Riskin determines the cordial edge deficiency and cordial vertex deficiency for the cases when the Möbius ladders and wheels are not cordial. In [3179] Riskin determines the cordial edge deficiencies for complete multipartite graphs that are not cordial and obtains a upper bound for their cordial vertex deficiencies.

Recall a graph G the graph G^*, called the *star of* G, is the graph obtained by replacing each vertex G with the star $K_{1,n}$. In [1867] Kaneria, Patadiya and Teraiya introduced a balanced cordial labeling for a graph by saying that a cordial labeling f is a *vertex balanced cordial* if it satisfies the condition$v_f(0) = v_f(1)$; f is a *balanced cordial* if it satisfies the conditions $e_f(0) = e_f(1)$ and $v_f(0) = v_f(1)$. Kaneria, Teraiya, and Patadiya [1871] proved the path union $P(t \cdot C_{4n})$ is a balanced cordial if t is odd and it is vertex balanced cordial if t is even; $C(t \cdot C_{4n})$ is a balanced cordial if $t \equiv 0 \pmod 4$ and it is a vertex balanced cordial if $t \equiv 1, 3 \pmod 4$; and $C^{\star}_{4n}$ is balanced cordial. They proved $P_n \times C_{4t}$ is balanced cordial; $C_{2n} \times C_{4t}$ is balanced cordial; and $G_1 \odot G_2$ is cordial when G_1 is cordial and G_2 is a balanced cordial. Kaneria and Teraiya [1870] prove if G is a balanced cordial, then so is G^*; if G is a balanced cordial, then so is $P_{2n+1} \times G$; and if G is a balanced cordial, then so is $\overline{G}^*$.

An *integer cordial labeling* of a graph $G^*(p, q)$ is an injective map $g : V \to [\frac{-p}{2}, \ldots, \frac{p}{2}]^*$ or $[-\lfloor \frac{p}{2} \rfloor, \ldots, \lfloor \frac{p}{2} \rfloor]$ as p is even or odd, which induces an edge labeling $g : E \to \{0, 1\}$ defined by $g(uv) = 1$ if $g(u) + g(v) \geq 0$ and 0 otherwise such that the number of edges labeled 1 and the number of edges labeled 0 differ by at most 1. If a graph has integer cordial labeling it is called an *integer cordial graph*. In [1305] Gondalia and Rokad

investigated the existence of integer cordial labelings of star and bistar related graphs.

For a planar graph G with an integer cordial labeling g and the face labeling $g**$ from the faces of G defined by $g**(f) = 1$ if $g(v_1) + g(v_2) + \ldots + g(v_n) = 0$ and $g**(f) = 0$ otherwise, where $v_1, v_2, \ldots, v_n$ are the vertices of face f. Such a labeling is called a *face integer cordial* labeling of graph G if the number of faces labeled with 0 and the number of faces labeled with 1 differ by at most 1. Parameswari, Saradha Pritha, and Rajeswari [2771] proved that the lily graph, $2K_{1,n} + 2P_n$ $(n \geq 2)$, admits an integer cordial labeling and the vanessa graph, $2F_n + K_{1,n}$ $(n \geq 2)$, admits an integer cordial labeling and a face integer cordial labeling. Sheriff, Abbas, and Raj [3505] proved that wheels, fans, friendship graphs, triangular snakes, double triangular snakes, the star of cycles, the degree splitting graph of bistars are face integer cordial graphs.

For a simple connected graph $G(V, E)$, Sahaya, Maya, and Nicholas [3234] introduced the concept of *product integer cordial* labeling of an injective mapping $f : V \to \{1, 2, \ldots, |V|\}$ such that the induced edge labeling f^* on E defined $f^*(uv) = 1$ or 0 according as $f(u)f(v)$ even or odd respectively, has the property that the number of edges labeled with 1 and the number of edges labeled with 0 differ by at most 1. They proved that the following graphs admit product integer cordial labelings: paths, friendship graphs, C_n if and only if n is odd, and $K_{m,n}$ if and only if one of m and n is 1. In [3460] Shah and Parmar proved that nontrivial triangular snakes, double triangular snakes, triple triangular snakes, and alternate triangular snakes graph admits integer cordial labelings. In [3461] Shah and Parmar proved that m-triangular snakes, quadrilateral snakes, double quadrilateral snakes, m-quadrilateral snakes, pentagonal snakes, double pentagonal snakes, and m-pentagonal snakes are integer cordial graphs. They also proved that triangular snake graph, double triangular snakes, alternate triangular snakes, m-triangular snakes, quadrilateral snakes, double quadrilateral snakes, m-quadrilateral snakes, pentagonal snakes, double pentagonal snakes, and m-pentagonal snakes graphs are integer cordial graphs. In [3462] Shah and Parmar proved that alternate m-triangular snakes, quadrilateral snakes, alternate m-quadrilateral snakes, pentagonal snakes, alternate m-pentagonal snakes, irregular triangular snakes, irregular quadrilateral snakes, and irregular pentagonal snakes are integer cordial graphs.

If f is a binary vertex labeling of a graph G Lee, Liu, and Tan [2165] defined a partial edge labeling of the edges of G by $f^*(uv) = 0$ if $f(u) = f(v) = 0$ and $f^*(uv) = 1$ if $f(u) = f(v) = 1$. They let $e_0(G)$ denote the number of edges uv for which $f^*(uv) = 0$ and $e_1(G)$ denote the number of edges uv for which $f^*(uv) = 1$. They say G is *balanced* if it has a friendly labeling f such that if $|e_0(f) - e_1(f)| \leq 1$. In the case that the number of vertices labeled 0 and the number of vertices labeled 1 are equal and the number of edges labeled 0 and the number of edges labeled 1 are equal they say the labeling is *strongly balanced*. They prove: P_n is balanced for all n and is strongly balanced if n is even; $K_{m,n}$ is balanced if and only if m and n are even, m and n are odd and differ by at most 2, or exactly one of m or n is even (say $n = 2t$) and $t \equiv -1, 0, 1 \pmod{|m - n|}$; a k-regular graph with p vertices is strongly balanced if and only if p is even and is balanced if and only if p is odd and $k = 2$; and if G is any graph and H is strongly balanced, the composition $G[H]$ (see §2.3 for the definition) is strongly balanced. In [2008] Kong, Lee,

Seah, and Tang show: $C_m \times P_n$ is balanced if m and n are odd and is strongly balanced if either m or n is even; and $C_m \odot K_1$ is balanced for all $m \geq 3$ and strongly balanced if m is even. They also provide necessary and sufficient conditions for a graph to be balanced or strongly balanced. Lee, Lee, and Ng [2135] show that stars are balanced if and only if the number of edges of the star is at most 4. Kwong, Lee, Lo, and Wang [2078] define a graph G to be *uniformly balanced* if $|e_0(f) - e_1(f)| \leq 1$ for every vertex labeling f that satisfies if $|v_0(f) - v_1(f)| \leq 1$. They present several ways to construct families of uniformly balanced graphs. Kim, Lee, and Ng [1961] prove the following: for any graph G, mG is balanced for all m; for any graph G, mG is strongly balanced for all even m; if G is strongly balanced and H is balanced, then $G \cup H$ is balanced; mK_n is balanced for all m and strongly balanced if and only if $n = 3$ or mn is even; if H is balanced and G is any graph, the $G \times H$ is strongly balanced; if one of m or n is even, then $P_m[P_n]$ is balanced; if both m and n are even, then $P_m[P_n]$ is balanced; and if G is any graph and H is strongly balanced, then the tensor product $G \otimes H$ is strongly balanced. (The *tensor product* $G \otimes H$ of graphs G and H, has the vertex set $V(G) \times V(H)$ and any two vertices (u, u') and (v, v') are adjacent in $G \otimes H$ if and only if u' is adjacent with v' and u is adjacent with v.)

A graph G is *k-balanced* if there is a function f from the vertices of G to $\{0, 1, 2, \ldots, k-1\}$ such that for the induced function f^* from the edges of G to $\{0, 1, 2, \ldots, k-1\}$ defined by $f^*(uv) = |f(u) - f(v)|$ the number of vertices labeled i and the number of edges labeled j differ by at most 1 for each i and j. Seoud, El Sonbaty, and Abd El Rehim [3377] proved the following: if $|E| \geq 2k + 1$ and $|V| \leq k$ then $G(V, E)$ is not k-balanced; if $|E| \geq 3k + 1$, $(k \geq 2)$ and $3k - 1 \geq |V| \geq 2k + 1$ then $G(V, E)$ is not k-balanced; r-regular graphs with $3 \leq r \leq n - 1$ are not r-balanced; if G_1 has m vertices and G_2 has n vertices then $G_1 + G_2$ is not $(m + n)$-balanced for $m, n \geq 5$; $P_3 \times P_n$ with edge set E is $3n$-balanced and $|E|$-balanced; $L_n \times P_2$ ($L_n = P_n \times P_2$) with vertex set V and edge set E is $|V|$-balanced and k-balanced for $k \geq |E|$ but not n-balanced for $n \geq 2$; the one-point union of two copies of $K_{2,n}$ is $2n$-balanced, $|V|$-balanced, and $|E|$-balanced not is 3-balanced when $n \geq 4$. They also proved that the composition graph $P_n[P_2]$ is not n-balanced for $n \geq 3$, is not $2n$-balanced for $n \geq 5$, and is not $|E|$-balanced.

A graph whose edges are labeled with 0 and 1 so that the absolute difference in the number of edges labeled 1 and 0 is no more than one is called *edge-friendly*. We say an edge-friendly labeling induces a *partial vertex labeling* if vertices which are incident to more edges labeled 1 than 0, are labeled 1, and vertices which are incident to more edges labeled 0 than 1, are labeled 0. Vertices that are incident to an equal number of edges of both labels are called *unlabeled*. Call a procedure on a labeled graph a label switching algorithm if it consists of pairwise switches of labels. Krop, Lee, and Raridan [2044] prove that given an *edge-friendly labeling* of K_n, we show a label switching algorithm producing an edge-friendly relabeling of K_n such that all the vertices are labeled.

Pereira, Singh, and Arumugam [2825] extended the concept of additively graceful graphs to the realm of sigraphs as follows. Let $S = (V, E)$ be a (p, m, n) signed graph with edge partition $E = E^+(S) \cup E^-(S)$. An injective function $f : V \to \{0, 1, \ldots, m + \lceil (n + 1)/2 \rceil\}$ induces an edge labeling $f^* : E \to \mathbb{N}$ defined by [2825] new

$f^*(uv) = f(u) + f(v)$, if $uv \in E^-(S)$, and $f^*(uv) = |f(u) - f(v)|$, if $uv \in E^+(S)$. If $f^*\big(E^-(S)\big) = \{1, 2, \ldots, n\}$ and $f^*\big(E^+(S)\big) = \{1, 2, \ldots, m\}$, then f is called an *additively graceful labeling* of S. A signed graph admitting such a labeling is called an *additively graceful signed graph*. They obtained some necessary or sufficient conditions for additively graceful sigraphs and some results on Eulerian sigraphs, complete bipartite sigraphs, and complete sigraphs. D'Souza and J. Pereira [990] gave a complete characterizations for additively graceful signed paths and signed cycles. In [991] D'Souza and Pereira obtained a characterization of additively graceful signed stars and they establisheed existence, uniqueness, and non-existence results for several subclasses of additively graceful signed double stars. [990] new [991] new

In 2017 [484] Bapat introduced a new labeling as follows. A function f from the vertices of a graph $G(E, V)$ to $\{0, 1, 2, \ldots, |V| - 1\}$ is called an *extended vertex edge additive cordial* labeling if the induced function f^* from the edges of G to $\{0, 1\}$ defined by $f^*(uv) = f(u) + f(v) \pmod 2$ for all edges uv of G has the property that the number of edges labeled 0 and the number of edges labeled 1 differ by at most 1. Bapat [484] proved paths, stars, $K_{2,n}, K_{3,n}, K_{4,n}$, $P_n \odot C_3$, and $P_n \odot C_4$ admit extended vertex edge additive cordial labeling.

Let $G(p, q)$ a simple finite connected graph. Given a bijective function f from $E(G)$ to $\{0, 1, \ldots, q-1\}$ Bapat [485] calls a bijective function f^* from $E(G)$ to $\{0, 1, 2, \ldots, q-1\}$ an *extended edge vertex cordial (eevc) labeling* if the induced function f^* from $V(G)$ to $\{0, 1\}$ defined by $f^*(u) = \Sigma f(uv) \bmod 2$ where the sum is taken over all edges incident to u has the property that the number of vertices labeled with 0 differs from the number labeled with 1 by at most 1. He shows that P_n ($n \neq 2 \bmod 4$), C_n ($n \neq 2 \bmod 4$), $K_{1,n}$ ($n \neq 1 \bmod 4$), graphs obtained by joining the centers of two copies of $K_{1,2n+1}$ by an edge, and triangular snakes have eevc labelings.

Murali, Thirusangu, Madura Meenakshi [2569] say a graph $G = (V, E)$ is *biconditional cordial* if there is a function $f : V \to \{0, 1\}$ such that the induced edge function $f^* : E \to \{0, 1\}$ defined by $f^*(uv) = 1$ if $(u) = f(v)$ and 0 if $f(u) \neq f(v)$ and the number of vertices labeled with 0 and the number labeled with 1 differ by at most 1 and the number of edges labeled with 0 and the number labeled with 1 differ by at most 1. Kalaimathi and Balamurugan [1809] prove the existence of the biconditional cordial labeling for complete bipartite graphs, books with triangular pages, sunflower graphs and web graphs. Nedumaran, Thirusangu, and Celin Mary [2663] proved that the graph consisting of k copies of a double star admits a biconditional cordial labeling. Kalaimathi, Balamurugan, and Rao [1810] proved the existence of the biconditional cordial labelings for super subdivisions of ladders and grids $P_m \times P_n$.

Let $G(V, E)$ be a simple graph. A surjective function $f : V(G) \to \{0, 1, 2\}$ is said to be 1-*near mean cordial* if, for each edge uv, the induced map $f^* : E(G) \to \{0, 1\}$ is defined by $f^*(uv) = 0$ if $\frac{f(u)+f(v)}{2}$ is an integer, and $f^*(uv) = 1$ otherwise, such that $|e_f(0) - e_f(1)| \leq 1$, where $e_f(0)$ and $e_f(1)$ denote the number of edges labeled 0 and 1, respectively. A graph G is said to be a 1-*near mean cordial* graph if it admits a 1-near mean cordial labeling. Elrokh, Alsatami, and Hamza [1008] showed that strong products of two paths admit 1-near mean cordial labelings in all non-trivial cases. Nedumaran, Thirusangu, and Celin [1008] new

Mary [2663] proved that k copies of double stars admit l-near mean cordial labelings and biconditional cordial labelings. They also showed that $(P_2 \cup nK_1) + N_2$ is a 1-near mean cordial. Palani and Surya [2749] proved that wheels, globes, binary trees, and odd cycles with a chord are 1-near mean cordial. In [2750] they proved that paths, combs, fans, and crowns are 1-near mean cordial graphs. [2663] new [2749] new [2750] new

A *total cordial* labeling of a graph G is a cordial labeling of vertex set and edge set such that the number of vertices and edges labeled with 0 and the number of vertices and edges labeled with 1 differ by at most 1. Elrokh, Al-Shamiri, Nada, and El-hay [1011] provided necessary and sufficient conditions for the existence of cordial and total cordial labelings for the corona product of paths and the one-point union C_m^2 and C_n^2. In [1007] Elmshtaye, Elrokh, and El-hay showed that the following graphs are total cordial: the third power of double fans, DF_n^3 $(n > 4)$, the third power of generalized fans, $G_k(F_m^3)$ $(m > 4)$, and $P_n \odot D(F_m^3)(m > 4)$.

3.8 The Friendly Index–Balance Index

Recall a function f from $V(G)$ to $\{0, 1\}$ where for each edge xy, $f^*(xy) = |f(x) - f(y)|$, $v_i(f)$ is the number of vertices v with $f(v) = i$, and $e_i(f)$ is the number of edges e with $f^*(e) = i$ is called *friendly* if $|v_0(f) - v_1(f)| \leq 1$. Lee and Ng [2172] define the *friendly index set* of a graph G as FI(G)= $\{|e_0(f) - e_1(f)|$ where f runs over all friendly labelings f of $G\}$. They proved: for any graph G with q edges FI$(G) \subseteq \{0, 2, 4, \ldots, q\}$ if q is even and FI$(G) \subseteq \{1, 3, \ldots, q\}$ if q is odd; for $1 \leq m \leq n$, FI$(K_{m,n})$= $\{(m - 2i)^2 |\ 0 \leq i \leq \lfloor m/2 \rfloor\}$ if $m + n$ is even; and FI$(K_{m,n})$= $\{i(i + 1) |\ 0 \leq i \leq m\}$ if $m + n$ is odd. In [2175] Lee and Ng prove the following: FI$(C_{2n}) = \{0, 4, 8, \ldots, 2n\}$ when n is even; FI$(C_{2n}) = \{2, 6, 10, \ldots, 2n\}$ when n is odd; and FI$(C_{2n+1}) = \{1, 3, 5, \ldots, 2n-1\}$. Elumalai [1022] defines a *cycle with a full set of chords* as the graph PC_n obtained from $C_n = v_0, v_1, v_2, \ldots, v_{n-1}$ by adding the cords $v_1v_{n-1}, v_2v_{n-2}, \ldots, v_{(n-2)/2}, v_{(n+2)/2}$ when n is even and $v_1v_{n-1}, v_2v_{n-2}, \ldots, v_{(n-3)/2}, v_{(n+3)/2}$ when n is odd. Lee and Ng [2174] prove: FI$(PC_{2m+1}) = \{3m - 2, 3m - 4, 3m - 6, \ldots, 0\}$ when m is even and FI$(PC_{2m+1}) = \{3m - 2, 3m - 4, 3m - 6, \ldots, 1\}$ when m is odd; FI$(PC_4) = \{1, 3\}$; for $m \geq 3$, FI$(PC_{2m}) = \{3m - 5, 3m - 7, 3m - 9, \ldots, 1\}$ when m is even; FI$(PC_{2m}) = \{3m - 5, 3m - 7, 3m - 9, \ldots, 0\}$ when m is odd.

The *full edge-friendly index* is the set of all possible numbers of edges receiving label 1 under an edge-friendly labeling. Salehi and Lee [3250] determined the full friendly index for various classes of trees. Among their results are: for a tree with q edges that has a perfect matching, the friendly index is the odd integers from 1 to q and for $n \geq 2$, FI(P_n)= $\{n - 1 - 2i |\ 0 \leq i \lfloor (n - 1)/2 \rfloor$. Law [2131] determined the full friendly index sets of spiders and disproved a conjecture by Salehi and Lee [3250] that the friendly index set of a tree forms an arithmetic progression. In [2178] Lee, Ng, and Lau determine the friendly index sets of several classes of spiders. Gao, Sun, and Lee [1197] determined the full friendly index of $P_m \times P_n$ with the extra $mn + 1 - m - n$ edges $u_{ij} - u_{(i+1)(j+1)}$. Sun, Gao, and Lee [3780] determined the full friendly index and friendly index for the twisted product of Mőbius ladders. Sinha and Kaur [3597] determined the full edge friendly

index of stars, wheels, 2-regular graphs, and mP_n. In [3520] Shiu determined the full edge-friendly index sets of complete bipartite graphs. Salehi and McGinn [3253] obtained partial results about the friendly index set of Q_n and strengthen a conjecture about the friendly index set of Q_n made in [3255]. Teffilia1 and Devaraj [3872] found the friendly index set of the graphs obtained by identifying the central vertex of a fan with the endpoint of a path (*umbrella*), the graphs obtained by identifying the central vertex of a star with the endpoint of a path, the graphs obtained by identifying the endpoints of copies of P_2 (*globe*), the splitting graph of a star, and $P_2 + mK_1$. Lee, Low, Ng, and Wang [2167] determined the friendly index sets for various classes of disjoint unions of stars. Gao, Ruo-Yuan, Lee, Ren, and Sun [1195] determined $FFI(G)$, $FI(G)$ and $FPCI(G)$ for a class of cubic graphs G. Ji, Liu, Bai, and Wu [1778] determined the full friendly index sets of mC_n. In [1193] Gao, Shiu, Lee, and Lau determined the full edge-friendly index set of a one point union of cycles. Gao, Shiu, Lee, and Lau [1196] determined the full [1196] new edge-friendly index sets of one point union of cycles.

Lee and Ng [2174] define $PC(n, p)$ as the graph obtained from the cycle C_n with consecutive vertices $v_0, v_1, v_2, \ldots, v_{n-1}$ by adding the p cords joining v_i to v_{n-i} for $1 \leq p\lfloor n/2 \rfloor - 1$. They prove $\mathrm{FI}(PC(2m+1, p)) = \{2m+p-1, 2m+p-3, 2m+p-5, \ldots, 1\}$ if p is even and $\mathrm{FI}(PC(2m+1, p)) = \{2m+p-1, 2m+p-3, 2m+p-5, \ldots, 0\}$ if p is odd; $\mathrm{FI}(PC(2m, 1)) = \{2m-1, 2m-3, 2m-5, \ldots, 1\}$; for $m \geq 3$, and $p \geq 2$, $\mathrm{FI}(PC(2m, p)) = \{2m+p-4, 2m+p-6, 2m+p-8, \ldots, 0\}$ when p is even, and $\mathrm{FI}(PC(2m, p)) = \{2m+p-4, 2m+p-6, 2m+p-8, \ldots, 1\}$ when p is odd. More generally, they show that the integers in the friendly index of a cycle with an arbitrary nonempty set of parallel chords form an arithmetic progression with a common difference 2. Shiu and Kwong [3524] determine the friendly index of the grids $P_n \times P_2$. The maximum and minimum friendly indices for $C_m \times P_n$ were given by Shiu and Wong in [3557]. In [2280] Liu, Liu, and Ji generalized the concept of the full m index set as follows. For a friendly labeling f of a graph G and an edge uv, $f'(uv) = f(u)+f(v)$, where $e_{f'}(i)$ is the number of edges labeled with i, and $u_f(i)$ is the number of vertices labeled with i. The full friendly m index set of a graph G is the set $M(G) = \{e_{f'}(1) - e_{f'}(0) \mid |v_f(1) - v_f(0)|, 0 \leq m < |V(G)|\}$ taken over all friendly labelings f of G. Note that when $m = 0$ or $m = 1$, the full m index set of a graph is its full friendly index set. Liu, Liu, and Ji determined the full m index set of $P_2 \times P_n$ for all m.

In [2176] Lee and Ng prove: for $n \geq 2$, $\mathrm{FI}(C_{2n} \times P_2) = \{0, 4, 8, \ldots, 6n-8, 6n\}$ if n is even and $\mathrm{FI}(C_{2n} \times P_2) = \{2, 6, 10, \ldots, 6n-8, 6n\}$ if n is odd; $\mathrm{FI}(C_3 \times P_2) = \{1, 3, 5\}$; for $n \geq 2$, $\mathrm{FI}(C_{2m+1} \times P_2) = \{6n-1\} \cup \{6n-5-2k |$ where $k \geq 0$ and $6n-5-2k \geq 0\}$; $\mathrm{FI}(M_{4n})$ (here M_{4n} is the Möbius ladder with $4n$ steps) $= \{6n-4-4k |$ where $k \geq 0$ and $6n-4-4k \geq 0\}$; $\mathrm{FI}(M_{4n+2}) = \{6n+3\} \cup \{6n-5-2k |$ where $k \geq 0$ and $6n-5-2k > 0\}$. In [2079] Kwong, Lee, and Ng completely determine the friendly index of all 2-regular graphs. As a corollary, they show that $C_m \cup C_n$ is cordial if and only if $m + n = 0, 1$ or $3 \pmod 4$. Ho, Lee, and Ng [1445] determine the friendly index sets of stars and various regular windmills. In [4177] Wen determines the friendly index of generalized wheels $C_n + mK_1$ for all $m > 1$. In [3249] Salehi and De determine the friendly index sets of certain caterpillars of diameter 4 and disprove a conjecture of Lee and Ng [2175] that the friendly index sets of trees form

an arithmetic progression. The maximum and minimum friendly indices for $C_m \times P_n$ were given by Shiu and Wong in [3557]. Salehi and Bayot [3246] have determined the friendly index set of $P_m \times P_n$. In [2176] Lee and Ng determine the friendly index sets for two classes of cubic graphs, prisms and Möbius ladders. Sinha and Kaur [3597] investigate the full region index sets of friendly labelings of cycles, wheels fans, and $P_2 \times P_n$.

For positive integers $a \leq b \leq c$, Lee, Ng, and Tong [2181] define the *broken wheel* $W(a, b, c)$ with three spokes as the graph obtained from K_4 with vertices u_1, u_2, u_3, c by inserting vertices $x_{1,1}, x_{1,2}, \ldots, x_{1,a-1}$ along the edge u_1u_2, $x_{2,1}, x_{2,2}, \ldots, x_{2,b-1}$ along the edge u_2u_3, $x_{3,1}, x_{3,2}, \ldots, x_{3,c-1}$ along the edge u_3u_1. They determine the friendly index set for broken wheels with three spokes.

Lee and Ng [2174] define a *parallel chord* of C_n as an edge of the form v_iv_{n-i} $(i < n-1)$ that is not an edge of C_n. For $n \geq 6$, they call the cycle C_n with consecutive vertices $v_1, v_2, \ldots, v_n$ and the edges $v_1v_{n-1}, v_2v_{n-2}, \ldots, v_{(n-2)/2}v_{(n+2)/2}$ for n even and $v_2v_{n-1}, v_3v_{n-2}, \ldots, v_{(n-1)/2}v_{(n+3)/2}$ for n odd, C_n with a *full set of parallel chords*. They determine the friendly index of these graphs and show that for any cycle with an arbitrary non-empty set of parallel chords the numbers in its friendly index set form an arithmetic progression with common difference 2.

For a graph $G(V, E)$ and a graph H rooted at one of its vertices v, Ho, Lee, and Ng [1444] define a *root-union of* (H, v) *by* G as the graph obtained from G by replacing each vertex of G with a copy of the root vertex v of H to which is appended the rest of the structure of H. They investigate the friendly index set of the root-union of stars by cycles.

For a graph $G(V, E)$, the *total graph* $T(G)$ of G, is the graph with vertex set $V \cup E$ and edge set $E \cup \{(v, uv) |\ v \in V, uv \in E\}$. Note that the total graph of the n-star is the friendship graph and the total graph of P_n is a triangular snake. Lee and Ng [2171] use $SP(1^n, m)$ to denote the spider with one central vertex joining n isolated vertices and a path of length m. They show: $\mathrm{FI}(K_1 + 2nK_2)$ (friendship graph with $2n$ triangles) $= \{2n, 2n-4, 2n-8, \ldots, 0\}$ if n is even; $\{2n, 2n-4, 2n-8, \ldots, 2\}$ if n is odd; $\mathrm{FI}(K_1+(2n+1)K_2) = \{2n+1, 2n-1, 2n-3, \ldots, 1\}$; for n odd, $\mathrm{FI}(T(P_n)) = \{3n-7, 3n-11, 3n-15, \ldots, z\}$ where $z = 0$ if $n \equiv 1 \pmod 4$ and $z = 2$ if $n \equiv 3 \pmod 4$; for n even, $\mathrm{FI}(T(P_n)) = \{3n-7, 3n-11, 3n-15, \ldots, n+1\} \cup \{n-1, n-3, n-5, \ldots, 1\}$; for $m \leq n-1$ and $m+n$ even, $\mathrm{FI}(T(SP(1^n, m))) = \{3(m+n)-4, 3(m+n)-8, 3(m+n)-12, \ldots, (m+n) \pmod 4\}$; for $m+n$ odd, $\mathrm{FI}(T(SP(1^n, m))) = \{3(m+n)-4, 3(m+n)-8, 3(m+n)-12, \ldots, m+n+2\} \cup \{m+n, m+n-2, m+n-4, \ldots, 1\}$; for $n \geq m$ and $m+n$ even, $\mathrm{FI}(T(SP(1^n, m))) = \{|4k-3(m+n)| \ |(n-m+2)/2 \leq k \leq m+n\}$; for $n \geq m$ and $m+n$ odd, $\mathrm{FI}(T(SP(1^n, m))) = \{|4k-3(m+n)| \ |(n-m+3)/2 \leq k \leq m+n\}$.

Kwong and Lee [2075] determine the friendly index any number of copies of C_3 that share an edge in common and the friendly index any number of copies of C_4 that share an edge in common. Lau, Gao, Lee, and Sun determine the friendly index sets and the cordiality of the edge-gluing of a complete graph K_n and n copies of cycles C_3.

For a planar graph $G(V, E)$ Sinha and Kaur [3618] extended the notion of an index set of a friendly labeling to regions of a planar graph and determined the full region index sets of friendly labeling of cycles, wheels fans, and grids $P_n \times P_2$.

An edge-friendly labeling f of a graph G induces a function f^* from $V(G)$ to $\{0,1\}$ defined as the sum of all edge labels mod 2. The *edge-friendly index set*, $I_f(G)$, of f is the number of vertices of f labeled 1 minus the number of vertices labeled 0. The *edge-friendly index set* of a graph G, EFI(G), is $\{|I_f(G)|\}$ taken over all edge-friendly labelings f of G. The *full edge-friendly index set* of a graph G, FEFI(G), is $\{I_f(G)\}$ taken over all edge-friendly labelings f of G. Sinha and Kaur [3617] determined the full edge-friendly index sets of stars, 2-regular graphs, wheels, and mP_n. In [3619] Sinha and Kaur extended the notion of index set of an edge-friendly labeling to regions of a planar graph and determined the full region index set of edge-friendly labelings of cycles, wheels, fans $P_n + K_1$, double fans $P_n + \overline{K_2}$, and grids $P_m \times P_n$ $(m \geq 2, n \geq 3)$. Sinha and Kaur [3596] investigate the full edge-friendly index sets of double stars, fans generalized fans, and $P_n \times P_2$. In [3519] Shiu determined the extreme values of edge-friendly indices of complete bipartite graphs.

In [1962] Kim, Lee, and Ng define the *balance index set* of a graph G as $\{|e_0(f)-e_1(f)|\}$ where f runs over all friendly labelings f of G. Zhang, Lee, and Wen [2135] investigate the balance index sets for the disjoint union of up to four stars and Zhang, Ho, Lee, and Wen [4312] investigate the balance index sets for trees with diameter at most four. Kwong, Lee, and Sarvate [2083] determine the balance index sets for cycles with one pendent edge, flowers, and regular windmills. Lee, Ng, and Tong [2180] determine the balance index set of certain graphs obtained by starting with copies of a given cycle and successively identifying one particular vertex of one copy with a particular vertex of the next. For graphs G and H and a bijection π from G to H, Lee and Su [2201] define Perm(G, π, H) as the graph obtaining from the disjoint union of G and H by joining each v in G to $\pi(v)$ with an edge. They determine the balanced index sets of the disjoint union of cycles and the balanced index sets for graphs of the form Perm(G, π, H) where G and H are regular graphs, stars, paths, and cycles with a chord. They conjecture that the balanced index set for every graph of the form Perm(G, π, H) is an arithmetic progression. Lee, Ho, and Su [2151] investigated the balance index sets of k-level wheel graphs.

Wen [4176] determines the balance index set of the graph that is constructed by identifying the center of a star with one vertex from each of two copies of C_n and provides a necessary and sufficient for such graphs to be balanced. In [2204] Lee, Su, and Wang determine the balance index sets of the disjoint union of a variety of regular graphs of the same order. Kwong [2073] determines the balanced index sets of rooted trees of height at most 2, thereby settling the problem for trees with diameter at most 4. His method can be used to determine the balance index set of any tree. The *homeomorph* $Hom(G, p)$ of a graph G is the collection of graphs obtained from G by adding p $(p \geq 0)$ additional degree 2 vertices to its edges. For any regular graph G, Kong, Lee, and Lee [1997] studied the changes of the balance index sets of $Hom(G, p)$ with respect to the parameter p. They derived explicit formulas for their balance index sets provided new examples of uniformly balanced graphs. In [698] Bouchard, Clark, Lee, Lo, and Su investigate the balance index sets of generalized books and ear expansion graphs. In [3207] Rose and Su provided an algorithm to calculate the balance index sets of a graph.

In [3525] Shiu and Kwong made a major advance by introducing an easier approach

to find the balance index sets of a large number of families of graphs in a unified and uniform manner. They use this method to determine the balance index sets for r-regular graphs, amalgamations of r-regular graphs, complete bipartite graphs, wheels, one point unions of regular graphs, sun graphs, generalized theta graphs, m-ary trees, spiders, grids $P_m \times P_n$, and cylinders $C_m \times P_n$. They provide a formula that enables one to determine the balance index sets of many biregular graphs (that is, graphs with the property that there exist two distinct positive integers r and s such that every vertex has degree r or s).

A labeling f from the vertices of a graph G to $\{0,1\}$ is said to be *vertex-friendly* if the number of vertices labeled with 0 and the number labeled with 1 differ by at most 1. The *vertex balance index set* of G is $|e_0(f) - e_1(f)|$ taken over all vertex-friendly labelings f. Adiga, Subbaraya, Shrikanth and Sriraj [77] completely determined the vertex balance index set of K_n, $K_{m,n}$, $C_n \times P_2$, and complete binary trees.

Manico and Pedrano [2414] prove that if the number of edges in a vertex-friendly of a graph G is even, then the vertex balance index of G contains only even numbers, and if the number of edges in a vertex-friendly graph G is odd, then the vertex balance index of G contains only odd numbers. Furthermore, they provide the vertex balance index set of triangular snakes, quadrilateral snakes, double triangular snakes), and double quadrilateral snakes.

In [3524] Shiu and Kwong define the *full friendly index set* of a graph G as $\{e_0(f) - e_1(f)\}$ where f runs over all friendly labelings of G. The full friendly index for $P_2 \times P_n$ is given by Shiu and Kwong in [3524]. The full friendly index of $C_m \times C_n$ is given by Shiu and Ling in [3542]. In [3615] and [3616] Sinha and Kaur investigated the full friendly index sets complete graphs, cycles, fans, double fans, wheels, double stars, $P_3 \times P_n$, and the tensor product of P_2 and P_n. Shiu and Ho [3522] investigated the full friendly index sets of cylinder graphs $C_m \times P_2$ $(m \geq 3), C_m \times P_3$ $(m \geq 4)$, and $C_3 \times P_n$ $(n \geq 4)$. These results, together with previously proven ones, completely determine the full friendly index of all cylinder graphs. Shiu and Ho [1442] study the full friendly index set and the full product-cordial index set of odd twisted cylinders and two permutation Petersen graphs. Gao [1185] determined the full friendly index set of $P_m \times P_n$, but he used the terms "edge difference set" instead of "full friendly index set" and "direct product" instead of "Cartesian product." The *twisted cylinder* graph is the permutation graph on $4n$ $(n \geq 2)$ vertices, $P(2n; \sigma)$, where $\sigma = (1,2)(3,4)\cdots(2n-1,2n)$ (the product of n transpositions). Shiu and Lee [3540] determined the full friendly index sets of twisted cylinders.

In [818] and [2076] Chopra, Lee, and Su and Kwong and Lee introduce a dual of balance index sets as follows. For an edge labeling f using 0 and 1 they define a partial vertex labeling f^* by assigning 0 or 1 to $f^*(v)$ depending on whether there are more 0-edges or 1-edges incident to v and leaving $f^*(v)$ undefined otherwise. For $i = 0$ or 1 and a graph $G(V,E)$, let $e_f(i) = |\{uv \in E : f(uv) = i\}|$ and $v_f(i) = |\{v \in V : f^*(v) = i\}|$. They define the *edge-balance index* of G as $\mathrm{EBI}(G) = \{|v_f(0) - v_f(1)| :$ the edge labeling f satisfies $|e_f(0) - e_f(1)| \leq 1\}$. Among the graphs whose edge-balance index sets have been investigated by Lee and his colleagues are: fans and wheels [818]; generalized theta graphs [2076]; flower graphs [2077] and [2077]; stars, paths, spiders,

and double stars [2212]; $(p, p+1)$-graphs [2206]; prisms and Möbius ladders [4150]; 2-regular graphs, complete graphs [4149]; and the envelope graphs of stars, paths, and cycles [828]. (The *envelope graph* of $G(V, E)$ is the graph with vertex set $V(G) \cup E(G)$ and set $E(G) \cup \{(u, (u, v)) : U \in V, (u, v) \in E)\}$).

Lee, Kong, Wang, and Lee [1998] found the EBI($K_{m,n}$) for $m = 1, 2, 3, 4, 5$ and $m = n$. Krop, Minion, Patel, and Raridan [2046] did the case for complete bipartite graphs with both parts of odd cardinality. Dao, Hua, Ngo, and Raridan [891] determined the edge-balanced index sets for complete even bipartite graphs. Krop and Sikes [2048] determined EBI($K_{m,m-2a}$) for $1 \leq a \leq (m-3)/4$ and m odd. Hua and Raridan [1461] determine the balanced index sets of all complete bipartite graphs with a larger part of odd cardinality and a smaller part of even cardinality. In [892] Dao, Hua, and Raridan provided the edge-balance index sets for complete bipartite graphs where m is even and n is odd, thereby completely resolving the problem of finding the edge-balance index sets for all complete bipartite graphs.

For a graph G and a connected graph H with a distinguished vertex s, the L-product of G and (H, s), $G \times_L (H, s)$, is the graph obtained by taking $|V(G)|$ copies of (H, s) and identifying each vertex of G with s of a single copy of H. In [820] and [699] Chou, Galiardi, Kong, Lee, Perry, Bouchard, Clark, and Su investigated the edge-balance index sets of L-product of cycles with stars. Bouchard, Clark, and Su [697] gave the exact values of the edge-balance index sets of L-product of cycles with cycles.

Chopra, Lee, and Su [821] prove that the edge-balance index of the fan $P_3 + K_1$ is $\{0, 1, 2\}$ and edge-balance index of the fan $P_n + K_1$, $n \geq 4$, is $\{0, 1, 2, \ldots, n-2\}$. They define the broken fan graphs $BF(a, b)$ as the graph with $V(BF(a, b)) = \{c\} \cup \{v_1, \ldots, v_a\} \cup \{u_1, \ldots, u_b\}$ and $E(BF(a, b)) = \{(c, v_i) |\ i = 1, \ldots, a\} \cup \{(c, u_i) |\ 1, \ldots, b\} \cup E(P_a) \cup E(P_b)$ ($a \geq 2$ and $b \geq 2$). They prove the edge-balance index set of $BF(a, b)$ is $\{0, 1, 2, \ldots, a+b-4\}$. In [2202] Lee, Su, and Todt give the edge-balance index sets of broken wheels. See also [78] and [3907]. In [2136] Lee, Lee, and Su present a technique that determines the balance index sets of a graph from its degree sequence. In addition, they give an explicit formula giving the exact values of the balance indices of generalized friendship graphs, envelope graphs of cycles, and envelope graphs of cubic trees.

3.9 k-equitable Labelings

In 1990 Cahit [740] proposed the idea of distributing the vertex and edge labels among $\{0, 1, \ldots, k-1\}$ as evenly as possible to obtain a generalization of graceful labelings as follows. For any graph $G(V, E)$ and any positive integer k, assign vertex labels from $\{0, 1, \ldots, k-1\}$ so that when the edge labels induced by the absolute value of the difference of the vertex labels, the number of vertices labeled with i and the number of vertices labeled with j differ by at most one and the number of edges labeled with i and the number of edges labeled with j differ by at most one. Cahit has called a graph with such an assignment of labels *k-equitable*. Note that $G(V, E)$ is graceful if and only if it is $|E|+1$-equitable and $G(V, E)$ is cordial if and only if it is 2-equitable. Cahit [739] has shown the following: C_n is 3-equitable if and only if $n \not\equiv 3 \pmod 6$; the triangular

snake with n blocks is 3-equitable if and only if n is even; the friendship graph $C_3^{(n)}$ is 3-equitable if and only if n is even; an Eulerian graph with $q \equiv 3 \pmod 6$ edges is not 3-equitable; and all caterpillars are 3-equitable [739]. Cahit [739] claimed to prove that W_n is 3-equitable if and only if $n \not\equiv 3 \pmod 6$ but Youssef [4276] proved that W_n is 3-equitable for all $n \geq 4$. Youssef [4274] also proved that if G is a k-equitable Eulerian graph with q edges and $k \equiv 2$ or $3 \pmod 4$ then $q \not\equiv k \pmod{2k}$. Cahit conjectures [739] that a triangular cactus with n blocks is 3-equitable if and only if n is even. In [740] Cahit proves that every tree with fewer than five end vertices has a 3-equitable labeling. He conjectures that all trees are k-equitable [741]. In 1999 Speyer and Szaniszló [3682] proved Cahit's conjecture for $k = 3$. Coles, Huszar, Miller, and Szaniszlo [857] proved caterpillars, symmetric generalized n-stars (or symmetric spiders), and complete n-ary trees are 4-equitable. Vaidya and Shah [4005] proved that the splitting graphs of $K_{1,n}$ and the bistar $B_{n,n}$ and the shadow graph of $B_{n,n}$ are 3-equitable. Rokad [3195] found 3-equitable labelings of the ring sum of different graphs. Ghodasaram and Sonchhatra [1262] proved the following graphs admit 3-equitable labeling: wheels, helms, gears, cycles with one pendant edge, and the graphs obtained by joining two copies of a fan by a path of arbitrary length. In [904] Dayanand, Ahmed, and Akka proved that $C_n \odot K_1$ is 8-equitable.

Vaidya, Dani, Kanani, and Vihol [3969] proved that the graphs obtained by starting with copies $G_1, G_2, \ldots, G_n$ of a fixed star and joining each center of G_i to the center of G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge are 3-equitable. Recall the shell $C(n, n-3)$ is the cycle C_n with $n-3$ cords sharing a common endpoint called the *apex*. Vaidya, Dani, Kanani, and Vihol [3970] proved that the graphs obtained by starting with copies $G_1, G_2, \ldots, G_n$ of a fixed shell and joining each apex of G_i to the apex of G_{i+1} $(i = 1, 2, \ldots, n-1)$ by an edge are 3-equitable. For a graph G and vertex v of G, Vaidya, Dani, Kanani, and Vihol [3971] prove that the graphs obtained from the wheel W_n, $n \geq 5$, by duplicating (see 3.7 for the definition) any rim vertex is 3-equitable and the graphs obtained from the wheel W_n by duplicating the center is 3-equitable when n is even and not 3-equitable when n is odd and at least 5. They also show that the graphs obtained from the wheel W_n, $n \neq 5$, by duplicating every vertex is 3-equitable.

Vaidya, Srivastav, Kaneria, and Ghodasara [4016] prove that cycle with two chords that share a common vertex with opposite ends that are incident to two consecutive vertices of the cycle (twin chords) is 3-equitable. Vaidya, Ghodasara, Srivastav, and Kaneria [3975] prove that star of cycle ${C_n}^*$ is 3-equitable for all n. Vaidya and Dani [3965] proved that the graphs obtained by starting with n copies $G_1, G_2, \ldots, G_n$ of a fixed star and joining the center of G_i to the center of G_{i+1} by an edge and each center to a new vertex x_i $(1 \leq i \leq n-1)$ by an edge have 3-equitable labeling. Vaidya and Dani [3968] prove that the graphs obtained by duplication of an arbitrary edge of a cycle or a wheel have 3-equitable labelings.

The *Mycielski graph* of a graph G is obtained from G by adding to each vertex v a new vertex u that is adjacent to the neighbors of v and a adding a new vertex w that is adjacent to every u. In [3299] Sangeeta, Parthiban, Selvaraju proved the non-existence of 3-equitable labelings for non-trivial cases of the total graphs of fans, the middle graph of ladders, the degree splitting graphs of friendship graphs, and the Mycielskian graphs of

paths.

Recall $G =< W_n^{(1)} : W_n^{(2)} : \ldots : W_n^{(k)} >$ 1s the graph obtained by joining the center vertices of each of $W_n^{(i)}$ and $W_n^{(i+1)}$ to a new vertex x_i where $1 \leq i \leq k-1$. Vaidya, Dani, Kanani, and Vihol [3972] prove that $< W_n^{(1)} : W_n^{(2)} : ... : W_n^{(k)} >$ is 3-equitable. Vaidya and Vihol [4021] prove that any graph G can be embedded as an induced subgraph of a 3-equitable graph thereby ruling out any possibility of obtaining any forbidden subgraph characterization for 3-equitable graphs.

The *shadow graph* $D_2(G)$ of a connected graph G is constructed by taking two copies of G, G' and G'' and joining each vertex u' in G' to the neighbors of the corresponding vertex u'' in G''. Vaidya, Vihol, and Barasara [506] prove that the shadow graph of C_n is 3-equitable except for $n = 3$ and 5 while the shadow graph of P_n is 3-equitable except for $n = 3$. They also prove that the middle graph of P_n is 3-equitable and the middle graph of C_n is 3-equitable for n even and not 3-equitable for n odd.

Bhut-Nayak and Telang have shown that crowns $C_n \odot K_1$, are k-equitable for $k = n, \ldots, 2n-1$ [649] and $C_n \odot K_1$ is k-equitable for all n when $k = 2, 3, 4, 5$, and 6 [650].

In [3366] Seoud and Abdel Maqsoud prove: a graph with n vertices and q edges in which every vertex has odd degree is not 3-equitable if $n \equiv 0 \pmod 3$ and $q \equiv 3 \pmod 6$; all fans except $P_2 + \overline{K_1}$ are 3-equitable; all double fans $P_n + \overline{K_2}$ except $P_4 + \overline{K_2}$ are 3-equitable; P_n^2 is 3-equitable for all n except 3; $K_{1,1,n}$ is 3-equitable if and only if $n \equiv 0$ or 2 (mod 3); $K_{1,2,n},\ n \geq 2$, is 3-equitable if and only if $n \equiv 2 \pmod 3$; $K_{m,n},\ 3 \leq m \leq n$, is 3-equitable if and only if $(m, n) = (4, 4)$; and $K_{1,m,n},\ 3 \leq m \leq n$, is 3-equitable if and only if $(m, n) = (3, 4)$. They conjectured that C_n^2 is not 3-equitable for all $n \geq 3$. However, Youssef [4282] proved that C_n^2 is 3-equitable if and only if n is at least 8. Youssef [4282] also proved that $C_n + \overline{K_2}$ is 3-equitable if and only if n is even and at least 6 and determined the maximum number of edges in a 3-equitable graph as a function of the number of its vertices. For a graph with n vertices to admit a k-equitable labeling, Seoud and Salim [3399] proved that the number of edges is at most $k\lceil (n/k) \rfloor^2 + k - 1$.

Bapat and Limaye [488] have shown the following graphs are 3-equitable: helms $H_n,\ n \geq 4$; flowers (see §2.2 for the definition); the one-point union of any number of helms; the one-point union of any number of copies of K_4; K_4-snakes (see §2.2 for the definition); C_t-snakes where $t = 4$ or 6; C_5-snakes where the number of blocks is not congruent to 3 modulo 6. A *multiple shell* $\mathrm{MS}\{n_1^{t_1}, \ldots, n_r^{t_r}\}$ is a graph formed by t_i shells each of order $n_i,\ 1 \leq i \leq r$, that have a common apex. Bapat and Limaye [489] show that every multiple shell is 3-equitable and Chitre and Limaye [810] show that every multiple shell is 5-equitable. In [811] Chitre and Limaye define the *H-union* of a family of graphs $G_1, G_2, \ldots, G_t$, each having a graph H as an induced subgraph, as the graph obtained by starting with $G_1 \cup G_2 \cup \cdots \cup G_t$ and identifying all the corresponding vertices and edges of H in each of $G_1, \ldots, G_t$. In [811] and [812] they proved that the $\overline{K_n}$-union of gears and helms H_n $(n \geq 6)$ are edge-3-equitable.

Szaniszló [3854] has proved the following: P_n is k-equitable for all k; K_n is 2-equitable if and only if $n = 1, 2$, or 3; K_n is not k-equitable for $3 \leq k < n$; S_n is k-equitable for all k; $K_{2,n}$ is k-equitable if and only if $n \equiv k-1 \pmod k$, or $n \equiv 0, 1, 2, \ldots, \lfloor k/2 \rfloor - 1 \pmod k$, or $n = \lfloor k/2 \rfloor$ and k is odd. She also proves that C_n is k-equitable if and only if

k meets all of the following conditions: $n \neq k$; if $k \equiv 2, 3 \pmod 4$, then $n \neq k-1$ and $n \not\equiv k \pmod{2k}$. Coles, Huszar, Miller, and Szaniszló [857] proved that all caterpillars, symmetric generalized n-stars (or symmetric spiders), and complete n-ary trees for all are 4-equitable.

Vickrey [4067] has determined the k-equitability of complete multipartite graphs. He shows that for $m \geq 3$ and $k \geq 3$, $K_{m,n}$ is k-equitable if and only if $K_{m,n}$ is one of the following graphs: $K_{4,4}$ for $k = 3$; $K_{3,k-1}$ for all k; or $K_{m,n}$ for $k > mn$. He also shows that when k is less than or equal to the number of edges in the graph and at least 3, the only complete multipartite graphs that are k-equitable are $K_{kn+k-1,2,1}$ and $K_{kn+k-1,1,1}$. Partial results on the k-equitability of $K_{m,n}$ were obtained by Krussel [2049].

In [4289] Youssef and Al-Kuleab proved the following: C_n^3 is 3-equitable if and only if n is even and $n \geq 12$; gear graphs are k-equitable for $k = 3, 4, 5, 6$; ladders $P_n \times P_2$ are 3-equitable for all $n \geq 2$; $C_n \times P_2$ is 3-equitable if and only if $n \not\equiv 0 \pmod 6$; Möbius ladders M_n are 3-equitable if and only if $n \not\equiv 0 \pmod 6$; and the graphs obtained from $P_n \times P_2$ ($n \geq 2$) where by adding the edges $u_i v_{i+1}$ ($1 \leq i \leq n-1$) to the path vertices $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$.

As a corollary of the result of Cairnie and Edwards [751] on the computational complexity of cordially labeling graphs it follows that the problem of finding k-equitable labelings of graphs is NP-complete as well.

Seoud and Abdel Maqsoud [3367] call a graph *k-balanced* if the vertices can be labeled from $\{0, 1, \ldots, k-1\}$ so that the number of edges labeled i and the number of edges labeled j induced by the absolute value of the differences of the vertex labels differ by at most 1. They prove that P_n^2 is 3-balanced if and only if $n = 2, 3, 4$, or 6; for $k \geq 4$, P_n^2 is not k-balanced if $k \leq n-2$ or $n+1 \leq k \leq 2n-3$; for $k \geq 4$, P_n^2 is k-balanced if $k \geq 2n-2$; for $k, m, n \geq 3$, $K_{m,n}$ is k-balanced if and only if $k \geq mn$; for $m \leq n$, $K_{1,m,n}$ is k-balanced if and only if (i) $m = 1$, $n = 1$ or 2, and $k = 3$; (ii) $m = 1$ and $k = n+1$ or $n+2$; or (iii) $k \geq (m+1)(n+1)$.

In [4282] Youssef gave some necessary conditions for a graph to be k-balanced and some relations between k-equitable labelings and k-balanced labelings. Among his results are: C_n is 3-balanced for all $n \geq 3$; K_n is 3-balanced if and only if $n \leq 3$; and all trees are 2-balanced and 3-balanced. He conjectures that all trees are k-balanced ($k \geq 2$).

Bloom has used the term k-equitable to describe another kind of labeling (see [4207] and [4208]). He calls a graph *k-equitable* if the edge labels induced by the absolute value of the difference of the vertex labels have the property that every edge label occurs exactly k times. Bloom calls a graph of order n *minimally k-equitable* if the vertex labels are 1, 2,..., n and it is k-equitable. Both Bloom and Wojciechowski [4207], [4208] proved that C_n is minimally k-equitable if and only if k is a proper divisor of n. Barrientos and Hevia [526] proved that if G is k-equitable of size $q = kw$ (in the sense of Bloom), then $\delta(G) \leq w$ and $\Delta(G) \leq 2w$. Barrientos, Dejter, and Hevia [525] have shown that forests of even size are 2-equitable. They also prove that for $k = 3$ or $k = 4$ a forest of size kw is k-equitable if and only if its maximum degree is at most $2w$ and that if 3 divides $mn+1$, then the double star $S_{m,n}$ is 3-equitable if and only if $q/3 \leq m \leq \lfloor (q-1)/2 \rfloor$. ($S_{m,n}$ is P_2 with m pendent edges attached at one end and n pendent edges attached at the other end.) They

discuss the k-equitability of forests for $k \geq 5$ and characterize all caterpillars of diameter 2 that are k-equitable for all possible values of k. Acharya and Bhat-Nayak [62] have shown that coronas of the form $C_{2n} \odot K_1$ are minimally 4-equitable. In [507] Barrientos proves that the one-point union of a cycle and a path (dragon) and the disjoint union of a cycle and a path are k-equitable for all k that divide the size of the graph. Barrientos and Havia [526] have shown the following: $C_n \times K_2$ is 2-equitable when n is even; books B_n $(n \geq 3)$ are 2-equitable when n is odd; the vertex union of k-equitable graphs is k-equitable; and wheels W_n are 2-equitable when $n \not\equiv 3 \pmod 4$. They conjecture that W_n is 2-equitable when $n \equiv 3 \pmod 4$ except when $n = 3$. Their 2-equitable labelings of $C_n \times K_2$ and the n-cube utilized graceful labelings of those graphs.

M. Acharya and Bhat-Nayak [63] have proved the following: the crowns $C_{2n} \odot K_1$ are minimally 2-equitable, minimally $2n$-equitable, minimally 4-equitable, and minimally n-equitable; the crowns $C_{3n} \odot K_1$ are minimally 3-equitable, minimally $3n$-equitable, minimally n-equitable, and minimally 6-equitable; the crowns $C_{5n} \odot K_1$ are minimally 5-equitable, minimally $5n$-equitable, minimally n-equitable, and minimally 10-equitable; the crowns $C_{2n+1} \odot K_1$ are minimally $(2n + 1)$-equitable; and the graphs P_{kn+1} are k-equitable.

In [509] Barrientos calls a k-equitable labeling *optimal* if the vertex labels are consecutive integers and *complete* if the induced edge labels are $1, 2, \ldots, w$ where w is the number of distinct edge labels. Note that a graceful labeling is a complete 1-equitable labeling. Barrientos proves that $C_m \odot nK_1$ (that is, an m-cycle with n pendent edges attached at each vertex) is optimal 2-equitable when m is even; $C_3 \odot nK_1$ is complete 2-equitable when n is odd; and that $C_3 \odot nK_1$ is complete 3-equitable for all n. He also shows that $C_n \odot K_1$ is k-equitable for every proper divisor k of the size $2n$. Barrientos and Havia [526] have shown that the n-cube $(n \geq 2)$ has a complete 2-equitable labeling and that $K_{m,n}$ has a complete 2-equitable labeling when m or n is even. They conjecture that every tree of even size has an optimal 2-equitable labeling.

3.10 Odd-even Prime Graceful Labelings

In [2575] Muruganantham and Kowsalya introduced the notion of odd-even prime graceful graphs as follows. A graph $G(V, E)$ with p vertices and q edges is called *odd-even prime graceful* if there exists an injective function $\omega : V \to \{1, 3, 5, \ldots, 2q + 1\}$ such that $\gcd(\omega(v_i), \omega(v_j)) = 1$ for all distinct i and j and the induced edge function ω^* defined by $\omega^*(v_iv_j) = |\omega(v_i) - \omega(v_j)|$ has the property $E = \{2, 4, 6, \ldots, 2q\}$. They showed that paths, stars, bistars, and the graphs obtained by attaching an edge to a cycle are odd-even prime graceful. They also proved that the following graphs admit odd-even prime graceful labelings: P_n, C_n $(n \geq 4)$, bistars $B_{2n+3,2n+3}$, $K_{2,n}$ $(n \geq 4)$, and graphs obtained by attaching an edge to C_n. [2575] new

In [2003] Kowsalya and Keerthia introduced the notion of k-odd-even prime graceful graphs as follows. For $k \geq 2$, a graph $G(V, E)$ with p vertices and q edges is called k *odd-even prime graceful* if there exists an injective function $\omega : V \to \{1, 3, 5, \ldots, k(2q + 1)\}$ such that $\gcd(\omega(v_i), \omega(v_j)) = 1$ and induced edge function ω^* defined by $\omega^*(v_iv_j) =$ [2003] new

$|\omega(v_i) - \omega(v_j)|$ has the property $E = \{2, 4, 6, \ldots, k(2q)\}$. They proved that the following graphs admit k-odd-even prime graceful labeling: bistars $B_{2n+3,2n+3}$ and $B_{4n+1,4n+1}$ when $2n + 3$ and $4n + 1$ are not prime, $K_{1,n,n}, P_n$ $(n > 8), C_n$ $(n \geq 4)$, graphs obtained by attaching an edge to C_n $(n \geq 4), K_{2,n}$ $(n \geq 4)$, and friendship graphs F_n $(n \geq 2)$. They also proved that bistars $B_{2n+3,2n+3}$ and $B_{4n+1,4n+1}$ odd-even prime graceful when $2n + 3$ and $4n + 1$ are prime.

3.11 Hamming-graceful Labelings

Mollard, Payan, and Shixin [2542] introduced a generalization of graceful graphs called Hamming-graceful. A graph $G = (V, E)$ is called *Hamming-graceful* if there exists an injective labeling g from V to the set of binary $|E|$-tuples such that $\{d(g(v), g(u))|\ uv \in E\} = \{1, 2, \ldots, |E|\}$ where d is the Hamming distance. Shixin and Yu [3564] have shown that all graceful graphs are Hamming-graceful; all trees are Hamming-graceful; C_n is Hamming-graceful if and only if $n \equiv 0$ or 3 (mod 4); if K_n is Hamming-graceful, then n has the form k^2 or $k^2 + 2$; and K_n is Hamming-graceful for $n = 2, 3, 4, 6, 9, 11, 16$, and 18. They conjecture that K_n is Hamming-graceful for n of the forms k^2 and $k^2 + 2$ for $k \geq 5$.

3.12 Hilbert Graceful Labelings

The nth *Hilbert number* H_n is defined by $H_n = 4(n - 1) + 1$, where n is a positive integer. Let $G(V, E)$ be a simple, finite, connected, undirected, non-trivial graph with p vertices and q edges. Sureeh Kumar [2064] defines a *Hilbert graceful labeling* as an injective function H from $V(G)$ to a set of Hilbert numbers that induces a bijective function H^* from E to $\{1, 2, 3, 4, \ldots, q\}$, where for each edge uv, we have $H^*(uv) = |H(u) - H(v)|$. A graph with Hilbert graceful labeling is called a *Hilbert graceful* graph. He proves certain complete multipartite graphs are Hilbert graceful.

4 Variations of Harmonious Labelings

4.1 Sequential and Strongly c-harmonious Labelings

Chang, Hsu, and Rogers [773] and Grace [1323], [1324] have investigated subclasses of harmonious graphs. Chang et al. define an injective labeling f of a graph G with q vertices to be *strongly c-harmonious* if the vertex labels are from $\{0, 1, \ldots, q-1\}$ and the edge labels induced by $f(x) + f(y)$ for each edge xy are $c, \ldots, c+q-1$. Strongly 1-harmonious labelings are more simply called *strongly harmonious*. Grace called such a labeling *sequential*. In the case of a tree, Chang et al. modify the definition to permit exactly one vertex label to be assigned to two vertices whereas Grace allows the vertex labels to range from 0 to q with no vertex label being used twice. For graphs other than trees, we use the term c-sequential labelings interchangeably with strongly c-harmonious labelings. By taking the edge labels of a sequentially labeled graph with q edges modulo q, we obviously obtain a harmoniously labeled graph. It is not known if there is a graph that can be harmoniously labeled but not sequentially labeled. Grace [1324] proved that caterpillars, caterpillars with a pendent edge, odd cycles with zero or more pendent edges, trees with α-labelings, wheels W_{2n+1}, and P_n^2 are sequential. Liu and Zhang [2282] finished off the crowns $C_{2n} \odot K_1$. (The case $C_{2n+1} \odot K_1$ was a special case of Grace's results. Liu [2296] proved crowns are harmonious.)

Bača and Youssef [430] investigated the existence of harmonious labelings for the corona graphs of a cycle and a graph G. They proved that if $G+K_1$ is strongly harmonious with the 0 label on the vertex of K_1, then $C_n \odot G$ is harmonious for all odd $n \geq 3$. By combining this with existing results they have as corollaries that the following graphs are harmonious: $C_n \odot C_m$ for odd $n \geq 3$ and $m \not\equiv 2 \pmod 3$; $C_n \odot K_{s,t}$ for odd $n \geq 3$; and $C_n \odot K_{1,s,t}$ for odd $n \geq 3$. Bu [715] also proved that crowns are sequential as are all even cycles with m pendent edges attached at each vertex. Figueroa-Centeno, Ichishima, and Muntaner-Batle [1089] proved that all cycles with m pendent edges attached at each vertex are sequential. Wu [4214] has shown that caterpillars with m pendent edges attached at each vertex are sequential.

Singh [3600, 3602] proved the following: $C_n \odot K_2$ is sequential for all odd $n > 1$; $C_n \odot P_3$ is sequential for all odd n; $K_2 \odot C_n$ (each vertex of the cycle is joined by edges to the end points of a copy of K_2) is sequential for all odd n [3604]; helms H_n are sequential when n is even; and $K_{1,n} + K_2$, $K_{1,n} + \overline{K}_2$, and ladders are sequential [3606]. Santhosh [3302] has shown that $C_n \odot P_4$ is sequential for all odd $n \geq 3$. Both Grace [1323] and Reid (see [1173]) have found sequential labelings for the books B_{2n}. Jungreis and Reid [1806] have shown the following graphs are sequential: $P_m \times P_n$ $(m,n) \neq (2,2)$; $C_{4m} \times P_n$ $(m,n) \neq (1,2)$; $C_{4m+2} \times P_{2n}$; $C_{2m+1} \times P_n$; and $C_4 \times C_{2n}$ $(n > 1)$. The graphs $C_{4m+2} \times C_{2n+1}$ and $C_{2m+1} \times C_{2n+1}$ fail to satisfy a necessary parity condition given by Graham and Sloane [1327]. The remaining cases of $C_m \times P_n$ and $C_m \times C_n$ are open. Gallian, Prout, and Winters [1174] proved that all graphs $C_n \times P_2$ with a vertex or an edge deleted are sequential. Zhu and Liu [4342] give necessary and sufficient conditions for sequential graphs, provide a characterization of non-tree sequential graphs by way of

by vertex closure, and obtain characterizations of sequential trees.

Gnanajothi [1276] [pp. 68-78] has shown the following graphs are sequential: $K_{1,m,n}$; mC_n, the disjoint union of m copies of C_n if and only if m and n are odd; books with triangular pages or pentagonal pages; and books of the form B_{4n+1}, thereby answering a question and proving a conjecture of Gallian and Jungreis [1173]. Sun [3779] has also proved that B_n is sequential if and only if $n \not\equiv 3 \pmod 4$. Ichishima and Oshima [1515] pose determining whether or not $mK_{s,t}$ is sequential as a problem.

Yuan and Zhu [4297] have shown that mC_n is sequential when m and n are odd. Although Graham and Sloane [1327] proved that the Möbius ladder M_3 is not harmonious, Gallian [1168] established that all other Möbius ladders are sequential (see §2.3 for the definition of Möbius ladder). Chung, Hsu, and Rogers [773] have shown that $K_{m,n} + K_1$, which includes $S_m + K_1$, is sequential. Seoud and Youssef [3406] proved that if G is sequential and has the same number of edges as vertices, then $G + \overline{K_n}$ is sequential for all n. Recall that $\Theta(C_m)^n$ denotes the book with n m-polygonal pages. Lu [2370] proved that $\Theta(C_{2m+1})^{2n}$ is $2mn$-sequential for all n and $m = 1, 2, 3, 4$, and $\Theta(C_m)^2$ is $(m-2)$-sequential if $m \geq 3$ and $m \equiv 2, 3, 4, 7 \pmod 8$.

Zhou and Yuan [4339] have shown that for every c-sequential graph G with p vertices and q edges and any positive integer m the graph $(G + \overline{K_m}) + \overline{K_n}$ is also c-sequential when $q - p + 1 \leq m \leq q - p + c$. Zhou [4338] has shown that the analogous results hold for strongly c-harmonious graphs. Zhou and Yuan [4339] have shown that for every c-sequential graph G with p vertices and q edges and any positive integer m the graph $(G + \overline{K_m}) + \overline{K_n}$ is c-sequential when $q - p + 1 \leq m \leq q - p + c$.

Shee [2190] proved that every graph is a subgraph of a sequential graph. Acharya, Germina, Princy, and Rao [52] prove that every connected graph can be embedded in a strongly c-harmonious graph for some c. Miao and Liang [2488] use $C_n(d; i, j; P_k)$ to denote a cycle C_n with path P_k joining two nonconsecutive vertices x_i and x_j of the cycle, where d is the distance between x_i and x_j on C_n. They proved that the graph $C_n(d; i, j; P_k)$ is strongly c-harmonious when $k = 2, 3$ and integer $n \geq 6$. Lu [2369] provides three techniques for constructing larger sequential graphs from some smaller one: an attaching construction, an adjoining construction, and the join of two graphs. Using these, he obtains various families of sequential or strongly c-indexable graphs.

For $1 \leq s \leq n_3$, let $C_n(i : i_1, i_2, \ldots, i_s)$ denote an n-cycle with consecutive vertices $x_1, x_2, \ldots, x_n$ to which the s chords $x_i x_{i_1}, x_i x_{i_2}, \ldots, x_i x_{i_s}$ have been added. Liang [2252] proved a variety of graphs of the form $C_n(i : i_1, i_2, \ldots, i_s)$ are strongly c-harmonious.

Youssef [4279] observed that a strongly c-harmonious graph with q edges is c-cordial for all $c \geq q$ and a strongly k-indexable graph is k-cordial for every k. The converse of this latter result is not true.

In [1512] Ichishima and Oshima show that the hypercube Q_n $(n \geq 2)$ is sequential if and only if $n \geq 4$. They also introduce a special kind of sequential labeling of a graph G with size $2t + s$ by defining a sequential labeling f to be a *partitional labeling* if G is bipartite with partite sets X and Y of the same cardinality s such that $f(x) \leq t + s - 1$ for all $x \in X$ and $f(y) \geq t - s$ for all $y \in Y$, and there is a positive integer m such that the induced edge labels are partitioned into three sets $[m, m + t - 1], [m + t, m + t + s - 1]$,

and $[m+t+s, m+2t+s-1]$ with the properties that there is an involution π, which is an automorphism of G such that π exchanges X and Y, $x\pi(x) \in E(G)$ for all $x \in X$, and $\{f(x)+f(\pi(x))|\ x \in X\} = [m+t, m+t+s-1]$. They prove if G has a partitional labeling, then $G \times Q_n$ has a partitional labeling for every nonnegative integer n. Using this together with existing results and the fact that every graph that has a partitional labeling is sequential, harmonious, and felicitous (see §4.5) they show that the following graphs are partitional, sequential, harmonious, and felicitous: for $n \geq 4$, hypercubes Q_n; generalized books $S_{2m} \times Q_n$; and generalized ladders $P_{2m+1} \times Q_n$.

In [1513] Ichishma and Oshima proved the following: if G is a partitional graph, then $G \times K_2$ is partitional, sequential, harmonious and felicitous; if G is a connected bipartite graph with partite sets of distinct odd order such that in each partite set each vertex has the same degree, then $G \times K_2$ is not partitional; for every positive integer m, the book B_m is partitional if and only if m is even; the graph $B_{2m} \times Q_n$ is partitional if and only if $(m,n) \neq (1,1)$; the graph $K_{m,2} \times Q_n$ is partitional if and only if $(m,n) \neq (2,1)$; for every positive integer n, the graph $K_{m,3} \times Q_n$ is partitional when $m = 4, 8, 12$, or 16. As open problems they ask which m and n is $K_{m,n} \times K_2$ partitional and for which l, m and n is $K_{l,m} \times Q_n$ partitional?

Ichishma and Oshima [1513] also investigated the relationship between partitional graphs and strongly graceful graphs (see §3.1 for the definition) and partitional graphs and strongly felicitous graphs (see §4.5) for the definition). They proved the following. If G is a partitional graph, then $G \times K_2$ is partitional, sequential, harmonious and felicitous. Assume that G is a partitional graph of size $2t+s$ with partite sets X and Y of the same cardinality s, and let f be a partitional labeling of G such that $\lambda_1 = \max\{f(x) : x \in X\}$ and $\lambda_2 = \max\{f(y) : y \in Y\}$. If $\lambda_1 + 1 = m + 2t + s - \lambda_2$, where $m = \min\{f(x)+f(y) : xy \in E(G)\} = \min\{f(y) : y \in Y\}$, then G has a strong α-valuation. Assume that G is a partitional graph of size $2t+s$ with partite sets X and Y of the same cardinality s, and let f be a partitional labeling of G such that $\lambda_1 = \max\{f(x) : x \in X\}$ and $\lambda_2 = \max\{f(y) : y \in Y\}$. If $\lambda_1 + 1 = m + 2t + s - \lambda_2$, where $m = \min\{f(x)+f(y) : xy \in E(G)\} = \min\{f(y) : y \in Y\}$, then G is strongly felicitous. Assume that G is a partitional graph of size $2t+s$ with partite sets X and Y of the same cardinality s, and let f be a partitional labeling of G such that $\mu_1 = f(x_1) = \min\{f(x) : x \in X\}$ and $\mu_2 = f(y_1) = \min\{f(y) : y \in Y\}$. If $t+s = m+1$ and $\mu_1 + \mu_2 = m$, where $m = \min\{f(x)+f(y) : xy \in E(G)\}$ and $x_1y_1 \in E(G)$, then G has a strong α-valuation and strongly felicitous labeling.

Vaidya and Lekha [3994] proved the following graphs are odd sequential: P_n, C_n for $n \equiv 0 \pmod 4$, crowns $C_n \bigodot K_1$ for even n, the graph obtained by duplication of arbitrary vertex in even cycles, path unions of stars, arbitrary super subdivisions in P_n, and shadows of stars. They also introduced the concept of a *bi-odd sequential* labeling of a graph G as one for which both G and its line graph $L(G)$ admit odd sequential labeling. They proved P_n and C_n for $n \equiv \pmod 4$ are bi-odd sequential graphs and trees are bi-odd sequential if and only if they are paths. They also prove that P_4 is the only graph with the property that it and its complement are odd sequential.

Arockiaraj, Mahalakshmi, and Namasivayam [292] proved that the subdivision graphs

of the following graphs have odd sequential labelings (they call them *odd sum* labelings): triangular snakes; quadrilateral snakes; slanting ladders SL_n $(n > 1)$ (the graphs obtained from two paths $u_1u_2 \dots u_n$ and $v_1v_2 \dots v_n$ by joining each u_i with v_{i+1}); $C_p \odot K_1$, $H_n \odot K_1$, $C_m@C_n$ (the graph obtained by attaching paths P_n to C_m by identifying the endpoints of the paths with each successive pairs of vertices of C_m); $P_m \times P_n$; and graphs obtained by the duplication of a vertex of a path and the duplication of a vertex of a cycle. Arockiaraj, Mahalakshmi, and Namasivayam [294] investigate the odd sum labeling behavior of paths, combs, cycles, crowns, and ladders under duplication of an edge. In [295] they investigated the odd sum property of shadow graphs, edge duplication graphs and vertex identification graphs. In [1309] Gopi proved the following graphs are odd sum graphs: graphs H_n obtained from two copies of P_n $(n \geq 3)$ with vertices $v_1, v_2, \dots, v_n$ and $u_1, u_2, \dots, u_n$ by joining $v_{(n+1)/2}$ and $u_{(n+1)/2}$ if n is odd and $v_{n/2}$ and $u_{(n+2)/2}$ if n is even; graphs obtained from H_n by attaching a fixed number of pendent edges at each vertex, graphs obtained from P_n $(n \geq 4)$ by attaching a two pendent edges at each interior vertex; and graphs obtained from P_m $(m \geq 4)$ by identifying an endpoint of the star S_n $(n \geq 2)$ with each vertex of P_m. In [1313] Gopi and Irudaya Mary proved that slanting ladders, shadow graphs of stars and bistars and mirror graphs and duplicate vertex graphs of paths with at least four vertices are odd sum graphs. In [1308] Gopi proved that alternative quadrilateral snakes $A(D(Q_n))$ $(n \geq 4)$ are odd sum graphs.

Arockiaraj and Mahalakshmi [291] proved the following graphs have odd sequential labelings (odd sum labelings): P_n $(n > 1)$, C_n if and only if $n \equiv 0 \pmod 4$; $C_{2n} \odot K_1$; $P_n \times P_2$ $(n > 1)$; $P_m \odot K_1$ if m is even or m is odd and $n = 1$ or 2; the balloon graph $P_m(C_n)$ obtained by identifying an end point of P_m with a vertex of C_n if either $n \equiv 0 \pmod 4$ or $n \equiv 2 \pmod 4$ and $m \not\equiv 1 \pmod 3$; quadrilateral snakes Q_n; $P_m \odot C_n$ if $m > 1$ and $n \equiv 0 \pmod 4$; $P_m \odot Q_3$; bistars; $C_{2n} \times P_2$; the trees T_p^n obtained from n copies of T_p by joining an edge uu' between every pair of consecutive paths where u is a vertex in ith copy of the path and u' is the corresponding vertex in the $(i+1)$th copy of the path; *H_n-graphs* obtained by starting with two copies of P_n with vertices $v_1, v_2, \dots, v_n$ and $u_1, u_2, \dots, u_n$ and joining the vertices $v_{(n+1)/2}$ and $u_{(n+1)/2}$ if n is odd and the vertices $v_{n/2+1}$ and $u_{n/2}$ if n; and $H_n \odot mK_1$.

Arockiaraj and Mahalakshmi [293] proved the splitting graphs of following graphs have odd sequential labelings (odd sum labelings): P_n; C_n if and only if $n \equiv 0 \pmod 4$; $P_n \odot K_1$; $C_{2n} \odot K_1$; $K_{1,n}$ if and only if $n \leq 2$; $P_n \times P_2$ $(n > 1)$; slanting ladders SL_n $(n > 1)$; the quadrilateral snake Q_n; and H_n-graphs.

Among the strongly 1-harmonious (also called *strongly harmonious*) graphs are: fans F_n with $n \geq 2$ [773]; wheels W_n with $n \not\equiv 2 \pmod 3$ [773]; $K_{m,n} + K_1$ [773]; French windmills $K_4^{(t)}$ [1457], [1884]; the friendship graphs $C_3^{(n)}$ if and only if $n \equiv 0$ or $1 \pmod 4$ [1457], [1884], [4237]; $C_{4k}^{(t)}$ [3783]; and helms [3089].

Seoud, Diab, and Elsakhawi [3376] have shown that the following graphs are strongly harmonious: $K_{m,n}$ with an edge joining two vertices in the same partite set; $K_{1,m,n}$; the composition $P_n[P_2]$ (see §2.3 for the definition); $B(3, 2, m)$ and $B(4, 3, m)$ for all m (see §2.4 for the notation); P_n^2 $(n \geq 3)$; and P_n^3 $(n \geq 3)$. Seoud et al. [3376] have also proved: B_{2n} is strongly $2n$-harmonious; P_n is strongly $\lfloor n/2 \rfloor$-harmonious; ladders L_{2k+1}

are strongly $(k+1)$-harmonious; and that if G is strongly c-harmonious and has an equal number of vertices and edges, then $G + \overline{K_n}$ is also strongly c-harmonious.

Bača and Youssef [430] investigated the existence of harmonious labelings for the corona graphs of a cycle and a graph G, and for the corona graph of K_2 and a tree. They prove: if the join of a graph G of order p and K_1, $G + K_1$, is strongly harmonious with the 0 label on the vertex of K_1, then the corona of C_n with G, $C_n \odot G$, is harmonious for all odd $n \geq 3$; if T is a strongly c-harmonious tree of odd size q and $c = \frac{q+1}{2}$ then the corona of K_2 with T, $K_2 \odot T$, is also strongly c-harmonious; if a unicyclic graph G of odd size q is a strongly c-harmonious and $c = \frac{q-1}{2}$ then the corona of K_2 with G, $K_2 \odot G$, is also strongly c-harmonious.

Sethuraman and Selvaraju [3450] have proved that the graph obtained by joining two complete bipartite graphs at one edge is graceful and strongly harmonious. They ask whether these results extend to any number of complete bipartite graphs.

Seenivasan and Lourdusamy [3335] define an *absolutely harmonious labeling* f as an injection from the vertex set of a graph G with q edges to the set $\{0, 1, 2, \ldots, q-1\}$, if when each edge uv is assigned $f(u) + f(v)$, the resulting edge labels can be arranged as $a_0, a_1, a_2, \ldots, a_{q-1}$ where $a_i = q - i$ or $q + i$ for $0 \leq i \leq q - 1$. When G is a tree one of the vertex labels may be assigned to exactly two vertices. A graph that admits absolutely harmonious labeling is called an *absolutely harmonious* graph. Observe that a strongly harmonious graph is an absolutely harmonious graph. They prove the following graphs are absolutely harmonious: P_n $(n \geq 3)$, $P_n \odot \overline{K_m}$, $C_n \odot \overline{K_m}$, the banana tree obtained by joining a vertex of degree 1 of each of any number of copies of $K_{1,n}$ to an isolated vertex, ladders, triangular snakes, quadrilateral snakes, mK_4, K_n if and only if $n = 3$ or 4. They also prove that if G is an absolutely harmonious graph, then there exists a partition (V_1, V_2) of the vertex set $V(G)$, such that the number of edges connecting the vertices of V_1 to the vertices of V_2 is exactly $\lceil q/2 \rceil$ and that if every vertex of an absolutely harmonious graph with q edges is even then $q \equiv 1$ or 2. As corollaries of the latter condition, they have that C_n when $n \equiv 1$ or 2 (mod 4), $C_m \times C_n$ when m and n are odd, and mK_3, $m \geq 2$ are not absolutely harmonious.

Annathurai and Gomathi [265] introduced the notion of absolutely k-harmonious labelings as follows. A function $f(G) \to 0, 1, \ldots, k-1$ is called an *absolutely k-harmonious labeling* (ABH labeling) of G if the induced edge labeling $f(G) \to 0, 1, \ldots, k-1$ defined by $f(xy) = |f(x) + f(y) - k| \pmod k$ for $xy \in E(G)$ satisfies $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(i) - e_f(j)| \leq 1$ for all $i, j \in 0, 1, \ldots, k-1$. A graph admitting such a labeling is called an *absolutely k-harmonious graph*. They proved that the absolutely k-harmonious property is preserved under three principal graph product operations: the Cartesian product, the tensor product, and the strong product of two absolutely k-harmonious graphs, provided suitable divisibility conditions hold. They further proved that the property is preserved under graph complementation, when the order and the size of the graph are each divisible by k, and under disjoint union. Concretely, they established that grid graphs formed as Cartesian products of two paths, and complete bipartite graphs whose part sizes sum to a multiple of three, are absolutely 3-harmonious for all suitable orders. [265] new

For a graph $G(V, E)$ Gayathri and Hemalatha [1222] define an *even sequential harmo-*

nious labeling f of G as an injection from V to $\{0,1,2,\ldots,2|E|\}$ with the property that the induced mapping f^+ from E to $\{2,4,6,\ldots,2|E|\}$ defined by $f^+(uv) = f(u)+f(v)$ when $f(u)+f(v)$ is even, and $f^+(uv) = f(u)+f(v)+1$ when $f(u)+f(v)$ is odd, is an injection. They prove the following have even sequential harmonious labelings (all cases are the nontrivial ones): $P_n, P_n^+, C_n(\ n \geq 3)$, triangular snakes, quadrilateral snakes, Möbius ladders, $P_m \times P_n$ $(m \geq 2, n \geq 2)$, $K_{m,n}$; crowns $C_m \odot K_1$, graphs obtained by joining the centers of two copies of $K_{1,n}$ by a path; banana trees (see §2.1), P_n^2, closed helms (see §2.2), $C_3 \odot nK_1$ $(n \geq 2)$; $D \odot K_{1,n}$ where D is a dragon (see §2.2); $\langle K_{1,n} : m\rangle$ $(m,n \geq 2)$ (see §4.5); the wreath product $P_n * \overline{K_2}$ $(n \geq 2)$ (see §4.5); combs $P_n \odot K_1$; the one-point union of the end point of a path to a vertex of a cycle (tadpole); the one-point union of the end point of a tadpole and the center of a star; the graphs PC_n obtained from $C_n = v_0, v_1, v_2, \ldots, v_{n-1}$ by adding the cords $v_1v_{n-1}, v_2v_{n-2}, \ldots, v_{(n-2)/2}, v_{(n+2)/2}$ when n is even and $v_1v_{n-1}, v_2v_{n-2}, \ldots, v_{(n-3)/2}, v_{(n+3)/2}$ when n is odd (that is, cycles with a full set of cords); $P_m \odot nK_1$; the one-point union of a vertex of a cycle and the center of a star; graphs obtained by joining the centers of two stars with an edge; graphs obtained by joining two disjoint cycles with an edge (dumbbells); graphs consisting of two even cycles of the same order sharing a common vertex with an arbitrary number of pendent edges attached at the common vertex (butterflies).

In [1492] Ichishima, Muntaner-Batle, and Oshima define the *harmonious number*, $\eta(G)$, of a graph G with q edges as the smallest positive integer n for which there exists an injective function f from $V(G)$ to Z_{n+1} such that each uv of G is labeled $f(u)+f(v)$ (mod q) and the resulting edge labels are distinct, or $+\infty$ if there exists no such integer n. If such functions exist, they are called *harmonious numberings*. The *strong harmonious number*, $\eta_s(G)$, of a graph G is defined to be either the smallest positive integer n such that $n = \eta(G)$ with the additional property that there exists an integer λ such that $\min\{f(u), f(v)\} \leq \lambda \leq \max\{f(u), f(v)\}$ for each edge in G or $+\infty$ if there exists no such integer n. They provide a necessary condition for a graph to have a finite harmonious number and sufficient conditions for a graph to have an infinite (strong) harmonious number. In addition, they examine the relations between harmonious numbers, gamma-numbers, alpha-numbers, and super edge magic deficiencies (see §5.2). They determine the formulas for the (strong) harmonious numbers of some 2-regular graphs and all complete bipartite graphs.

A *k-odd sequential harmoniously* labeling is an injection f from the vertex set V to $\{k-1, k, k+1, \ldots, k+2q-2\}$ such that the induced mapping f^+ from the edge set E to $\{2k-1, 2k+1, 2k+3, \ldots, 2k+2q-3\}$, defined by $f^+(uv) = f(u)+f(v)$ if $f(u)+f(v)$ is even and $f^+(uv) = f(u)+f(v)+1$ if $f(u)+f(v)$ is odd, is injective. A graph G is said to be an *k-odd sequential harmonious* graph if it admits an k-odd sequential harmonious labeling. Recall that the quadrilateral snake Q_x is obtained from the path $v_1, v_2, v_3, \ldots, v_n$ and the double quadrilateral snake $D(Q_x)$ is obtained from Q_x by adding new vertices $w_1, w_2, w_3, \ldots, w_{2n-2}$ and new edges v_iw_{2i-2} for $2 \leq i \leq n$, and $w_{2i-1}w_{2i}$ and v_iw_{2i-1} for $1 \leq i \leq n-1$. Manonmani and Savithiri [2415] gave k-odd sequential harmonious labelings of double quadrilateral $2Q_x$-snakes for all x and k-odd sequential harmonious labelings of $2mQ_x$-snakes for all m and x. [2415] new

In her PhD thesis [2618] (see also [1223]) Muthuramakrishnan defined a labeling f of a graph $G(V, E)$ to be *k-even sequential harmonious* if f is an injection from V to $\{k-1, k, k+1, \ldots, k+2q-1\}$ such that the induced mapping f^+ from E to $\{2k, 2k+2, 2k+4, \ldots, 2k+2q-2\}$ defined by $f^+(uv) = f(u) + f(v)$ if $f(u) + f(v)$ is even and $f^+(uv) = f(u) + f(v) + 1$ if $f(u) + f(v)$ is odd are distinct. A graph G is called a *k-even sequential harmonious* graph if it admits a k-even sequential harmonious labeling. Among the numerous graphs that she proved to be k-even sequential harmonious are: paths, cycles, $K_{m,n}$, P_n^2 $(n \geq 3)$, crowns $C_m \odot K_1$, $C_m@P_n$ (the graph obtained by identifying an endpoint of P_n with one vertex of C_m), double triangular snakes, double quadrilateral snakes, bistars, grids $P_m \times P_n$ $(m, n \geq 2)$, $P_n[P_2]$, $C_3 \odot nK_1$ $(n \geq 2)$, flags Fl_m (the cycle C_m with one pendent edge), dumbbell graphs (two disjoint cycles joined by an edge) butterfly graphs B_n (two even cycles of the same order sharing a common vertex with an arbitrary number of pendent edges attached at the common vertex), $K_2 + nK_1$, $\overline{K_n} + 2K_2$, banana trees, sparklers $P_m@K_{1,n}$ $(m, n \geq 2)$, (graphs obtained by identifying an endpoint of P_m with the center of a star), twigs (graphs obtained from P_n $(n \geq 3)$ by attaching exactly two pendent edges at each internal vertex of P_n), festoon graphs $P_m \odot nK_1$ $(m \geq 2)$, the graphs $T_{m,n,t}$ obtained from a path P_t by appending m edges at one endpoint of P_t and n edges at the other endpoint of P_t, $L_n \odot K_1$ (L_n is the ladder $P_n \times P_2$), shadow graphs of paths, stars and bistars, and split graphs of paths and stars. Muthuramakrishnan also obtains some results on k-odd sequential harmonious labeling of graphs.

In [603] Beatress and Sarasija introduced a new harmonious-like labeling as follows. A graph $G(V, E)$ with n vertices and m edges is said to be a *square harmonious* graph if there exists an injection f from $V(G)$ to $\{1, 2, \ldots, m^2+1\}$ such that the induced mapping f^* from $E(G)$ to $\{1, 4, 9, \ldots, m^2\}$ defined by $f^*(uv) = (f(u) + f(v)) \bmod (m^2 + 1)$ is a bijection. Such a function f is called a *square harmonious labeling* of G. They prove that P_n $(n \geq 3), K_{1,n}$ $(n \geq 2)$, bistars, combs, $P_n \odot pK_1$ $(n \geq 2)$, and $C_3@pK_1$ $(p \geq 2)$ are square harmonious graphs. Lawas and Lim [2132] proved that stars have a square harmonious labelings.

In [548] Barrientos and Youssef generalize the concepts of harmonious and (k, d)-arithmetic graphs by relaxing the injectivity constraint of the corresponding labelings. They call these labelings *semi harmonious* and *semi (k, d)-arithmetic*. They showed the existence of a semi harmonious labeling for several types of cycle-related graphs and characterized the cycles that admit semi harmonious labelings. They proved that if G is semi (k, d)-arithmetic, then it is also semi (rk, rd)-arithmetic for every $r \geq 1$ and that nG is both semi (k, d)-arithmetic and semi $(k + d, d)$-arithmetic. They further showed that any graph whose components are complete bipartite graphs is semi (k, d)-arithmetic for any ordered pair (k, d) of positive integers, and if G_i is a semi (k_i, d)-arithmetic graph of size q_i for each $i = 1, 2$, then $G_1 \cup G_2$ is semi (k_i, d)-arithmetic.

4.2 (k, d)-arithmetic Labelings

Acharya and Hegde [56] have generalized sequential labelings as follows. Let G be a graph with q edges and let k and d be positive integers. A labeling f of G is said to be

(k,d)-*arithmetic* if the vertex labels are distinct nonnegative integers and the edge labels induced by $f(x)+f(y)$ for each edge xy are $k, k+d, k+2d, \ldots, k+(q-1)d$. They obtained a number of necessary conditions for various kinds of graphs to have a (k,d)-arithmetic labeling. The case where $k=1$ and $d=1$ was called *additively graceful* by Hegde [1400]. Hegde [1400] showed: K_n is additively graceful if and only if $n=2,3$, or 4; every additively graceful graph except K_2 or $K_{1,2}$ contains a triangle; and a unicyclic graph is additively graceful if and only if it is a 3-cycle or a 3-cycle with a single pendent edge attached. Jinnah and Singh [1784] noted that P_n^2 is additively graceful. Hegde [1401] proved that if G is strongly k-indexable, then G and $G+\overline{K_n}$ are (kd,d)-arithmetic. Acharya and Hegde [58] proved that K_n is (k,d)-arithmetic if and only if $n \geq 5$ (see also [721]). They also proved that a graph with an α-labeling is a (k,d)-arithmetic for all k and d. Bu and Shi [721] proved that $K_{m,n}$ is (k,d)-arithmetic when k is not of the form id for $1 \leq i \leq n-1$. For all $d \geq 1$ and all $r \geq 0$, Acharya and Hegde [56] showed the following: $K_{m,n,1}$ is $(d+2r,d)$-arithmetic; C_{4t+1} is $(2dt+2r,d)$-arithmetic; C_{4t+2} is not (k,d)-arithmetic for any values of k and d; C_{4t+3} is $((2t+1)d+2r,d)$-arithmetic; W_{4t+2} is $(2dt+2r,d)$-arithmetic; and W_{4t} is $((2t+1)d+2r,d)$-arithmetic. They conjecture that C_{4t+1} is $(2dt+2r,d)$-arithmetic for some r and that C_{4t+3} is $(2dt+d+2r,d)$-arithmetic for some r. Hegde and Shetty [1420] proved the following: the generalized web $W(t,n)$ (see §2.2 for the definition) is $((n-1)d/2,d)$-arithmetic and $((3n-1)d/2,d)$-arithmetic for odd n; the join of the generalized web $W(t,n)$ with the center removed and $\overline{K_p}$ where n is odd is $((n-1)d/2,d)$-arithmetic; every T_p-tree (see §3.2 for the definition) with q edges and every tree obtained by subdividing every edge of a T_p-tree exactly once is $(k+(q-1)d,d)$-arithmetic for all k and d. Lu, Pan, and Li [2372] proved that $K_{1,m} \cup K_{p,q}$ is (k,d)-arithmetic when $k > (q-1)d+1$ and $d>1$.

Yu [4292] proved that a necessary condition for C_{4t+1} to be (k,d)-arithmetic is that $k=2dt+r$ for some $r \geq 0$ and a necessary condition for C_{4t+3} to be (k,d)-arithmetic is that $k=(2t+1)d+2r$ for some $r \geq 0$. These conditions were conjectured by Acharya and Hegde [56]. Singh proved that the graph obtained by subdividing every edge of the ladder L_n is $(5,2)$-arithmetic [3601] and that the ladder L_n is $(n,1)$-arithmetic [3605]. He also proves that $P_m \times C_n$ is $((n-1)/2,1)$-arithmetic when n is odd [3605]. Acharya, Germina, and Anandavally [50] proved that the subdivision graph of the ladder L_n is (k,d)-arithmetic if either d does not divide k or $k=rd$ for some $r \geq 2n$ and that $P_m \times P_n$ and the subdivision graph of the ladder L_n are (k,k)-arithmetic if and only if k is at least 3. Lu, Pan, and Li [2372] proved that $S_m \cup K_{p,q}$ is (k,d)-arithmetic when $k > (q-1)d+1$ and $d>1$.

A graph is called *arithmetic* if it is (k,d)-arithmetic for some k and d. Singh and Vilfred [3611] showed that various classes of trees are arithmetic. Singh [3605] has proved that the union of an arithmetic graph and an arithmetic bipartite graph is arithmetic. He conjectures that the union of arithmetic graphs is arithmetic. He provides an example to show that the converse is not true.

Germina and Anandavally [1244] investigated embedding of graphs in arithmetic graphs. They proved: every graph can be embedded as an induced subgraph of an arithmetic graph; every bipartite graph can be embedded in a (k,d)-arithmetic graph for

all k and d such that d does not divide k; and any graph containing an odd cycle cannot be embedded as an induced subgraph of a connected (k,d)-arithmetic with $k < d$.

In [4260] Yao, Liu, and Yao give necessary and sufficient conditions for a tree to have the following mutually equivalent labelings: set-ordered odd-graceful, (k,d)-graceful, super edge-magic total, odd-elegant (see §4.4), harmonious, (k,d)-arithmetic, and edge-antimagic (see §6.1).

4.3 (k,d)-indexable Labelings

Acharya and Hegde [56] call a graph with p vertices and q edges (k,d)-*indexable* if there is an injective function from V to $\{0,1,2,\ldots,p-1\}$ such that the set of edge labels induced by adding the vertex labels is a subset of $\{k, k+d, k+2d, \ldots, k+q(d-1)\}$. When the set of edges is $\{k, k+d, k+2d, \ldots, k+q(d-1)\}$ the graph is said to be *strongly* (k,d)-*indexable*. A $(k,1)$-graph is more simply called k-*indexable* and strongly 1-indexable graphs are simply called *strongly indexable*. Notice that strongly indexable graphs are a stronger form of sequential graphs and for trees and unicyclic graphs the notions of sequential labelings and strongly k-indexable labelings coincide. Hegde and Shetty [1425] have shown that the notions of $(1,1)$-strongly indexable graphs and super edge-magic total labelings (see §5.2) are equivalent.

Zhou [4338] has shown that for every k-indexable graph G with p vertices and q edges the graph $(G + \overline{K_{q-p+k}}) + \overline{K_1}$ is strongly k-indexable. Acharaya and Hegde prove that the only nontrivial regular graphs that are strongly indexable are K_2, K_3, and $K_2 \times K_3$, and that every strongly indexable graph has exactly one nontrivial component that is either a star or has a triangle. Acharya and Hegde [56] call a graph with p vertices *indexable* if there is an injective labeling of the vertices with labels from $\{0,1,2,\ldots,p-1\}$ such that the edge labels induced by addition of the vertex labels are distinct. They conjecture that all unicyclic graphs are indexable. This conjecture was proved by Arumugam and Germina [277] who also proved that all trees are indexable. Bu and Shi [722] also proved that all trees are indexable and that all unicyclic graphs with the cycle C_3 are indexable. Hegde [1401] has shown the following: every graph can be embedded as an induced subgraph of an indexable graph; if a connected graph with p vertices and q edges ($q \geq 2$) is (k,d)-indexable, then $d \leq 2$; $P_m \times P_n$ is indexable for all m and n; if G is a connected $(1,2)$-indexable graph, then G is a tree; the minimum degree of any $(k,1)$-indexable graph with at least two vertices is at most 3; a caterpillar with partite sets of orders a and b is strongly $(1,2)$-indexable if and only if $|a-b| \leq 1$; in a connected strongly k-indexable graph with p vertices and q edges, $k \leq p-1$; and if a graph with p vertices and q edges is (k,d)-indexable, then $q \leq (2p-3-k+d)/d$. As a corollary of the latter, it follows that K_n ($n \geq 4$) and wheels are not (k,d)-indexable.

Lee and Lee [2134] provide a way to construct a (k,d)-strongly indexable graph from two given (k,d)-strongly indexable graphs. Lee and Lo [2166] show that every given (1,2)-strongly indexable spider can extend to an (1,2)-strongly indexable spider with arbitrarily many legs.

Seoud, Abd El Hamid, and Abo Shady [3364] proved the following graphs are in-

dexable: $P_m \times P_n$ $(m, n \geq 2)$; the graphs obtained from $P_n + K_1$ by inserting one vertex between every two consecutive vertices of P_n; the one-point union of any number of copies of $K_{2,n}$; and the graphs obtained by identifying a vertex of a cycle with the center of a star. They showed P_n is strongly $\lceil n/2 \rceil$-indexable; odd cycles C_n are strongly $\lceil n/2 \rceil$-indexable; $K_{(m,n)}$ $(m, n > 2)$ is indexable if and only if m or n is at most 2. For a simple indexable graph $G(V, E)$ they proved $|E| \leq 2|V| - 3$. Also, they determine all indexable graphs of order at most 6.

Hegde and Shetty [1424] also prove that if G is strongly k-indexable Eulerian graph with q edges then $q \equiv 0, 3 \pmod 4$ if k is even and $q \equiv 0, 1 \pmod 4$ if k is odd. They further showed how strongly k-indexable graphs can be used to construct polygons of equal internal angles with sides of different lengths.

Germina [1240] has proved the following: fans $P_n + K_1$ are strongly indexable if and only if $n = 1, 2, 3, 4, 5, 6$; $P_n + K_2$ is strongly indexable if and only if $n \leq 2$; the only strongly indexable complete m-partite graphs are $K_{1,n}$ and $K_{1,1,n}$; ladders $P_n \times P_2$ are $\lceil \frac{n}{2} \rceil$-strongly indexable, if n is odd; $K_n \times P_k$ is strongly indexable if and only if $n = 3$; $C_m \times P_n$ is 2-strongly indexable if m is odd and $n \geq 2$; $K_{1,n} + K_i$ is not strongly indexable for $n \geq 2$; for $G_i \cong K_{1,n},\ 1 \leq i \leq n$, the sequential join $G \cong (G_1 + G_2) \cup (G_2 + G_3) \cup \cdots \cup (G_{n-1} + G_n)$ is strongly indexable if and only if, either $i = n = 1$ or $i = 2$ and $n = 1$ or $i = 1, n = 3$; $P_1 \cup P_n$ is strongly indexable if and only if $n \leq 3$; $P_2 \cup P_n$ is not strongly indexable; $P_2 \cup P_n$ is $\lceil \frac{n+3}{2} \rceil$-strongly indexable; mC_n is k-strongly indexable if and only if m and n are odd; $K_{1,n} \cup K_{1,n+1}$ is strongly indexable; and $mK_{1,n}$ is $\lceil \frac{3m-1}{2} \rceil$-strongly indexable when m is odd.

Acharya and Germina [45] proved that every graph can be embedded in a strongly indexable graph and gave an algorithmic characterization of strongly indexable unicyclic graphs. In [47] they provide necessary conditions for an Eulerian graph to be strongly k-indexable and investigate strongly indexable (p, q)-graphs for which $q = 2p - 3$.

Hegde and Shetty [1420] proved that for n odd the generalized web graph $W(t, n)$ with the center removed is strongly $(n-1)/2$-indexable. Hegde and Shetty [1425] define a *level joined planar grid* as follows. Let u be a vertex of $P_m \times P_n$ of degree 2. For every pair of distinct vertices v and w that do not have degree 4, introduce an edge between v and w provided that the distance from u to v equals the distance from u to w. They prove that every level joined planar grid is strongly indexable. For any sequence of positive integers $(a_1, a_2, \ldots, a_n)$ Lee and Lee [2133] show how to associate a strongly indexable $(1, 1)$-graph. As a corollary, they obtain the aforementioned result Hegde and Shetty on level joined planar grids.

Section 5.2 of this survey includes a discussion of a labeling method called super edge-magic. In 2002 Hegde and Shetty [1425] showed that a graph has a strongly k-indexable labeling if and only if it has a super edge-magic labeling.

4.4 Elegant Labelings

In 1981 Chang, Hsu, and Rogers [773] defined an *elegant labeling* f of a graph G with q edges as an injective function from the vertices of G to the set $\{0, 1, \ldots, q\}$ such that

when each edge xy is assigned the label $f(x)+f(y) \pmod{(q+1)}$ the resulting edge labels are distinct and nonzero. An injective labeling f of a graph G with q vertices is called *strongly k-elegant* if the vertex labels are from $\{0, 1, \ldots, q\}$ and the edge labels induced by $f(x) + f(y) \pmod{(q+1)}$ for each edge xy are $k, \ldots, k+q-1$. Note that in contrast to the definition of a harmonious labeling, for an elegant labeling it is not necessary to make an exception for trees. Husna and Rahmadani [1472] showed that nontrival cases the lotus graph and its two variations are elegant graphs. [1472] new

Whereas the cycle C_n is harmonious if and only if n is odd, Chang et al. [773] proved that C_n is elegant when $n \equiv 0$ or $3 \pmod 4$ and not elegant when $n \equiv 1 \pmod 4$. Chang et al. further showed that all fans are elegant and the paths P_n are elegant for $n \not\equiv 0 \pmod 4$. Cahit [737] then showed that P_4 is the only path that is not elegant. Balakrishnan, Selvam, and Yegnanarayanan [467] have proved numerous graphs are elegant. Among them are $K_{m,n}$ and the mth-subdivision graph of $K_{1,2n}$ for all m. They prove that the bistar $B_{n,n}$ (K_2 with n pendent edges at each endpoint) is elegant if and only if n is even. They also prove that every simple graph is a subgraph of an elegant graph and that several families of graphs are not elegant. Deb and Limaye [913] have shown that triangular snakes (see §2.2 for the definition) are elegant if and only if the number of triangles is not equal to 3 (mod 4). In the case where the number of triangles is 3 (mod 4) they show the triangular snakes satisfy a weaker condition they call *semi-elegant* whereby the edge label 0 is permitted. In [914] Deb and Limaye define a graph G with q edges to be *near-elegant* if there is an injective function f from the vertices of G to the set $\{0, 1, \ldots, q\}$ such that when each edge xy is assigned the label $f(x) + f(y) \pmod{(q+1)}$ the resulting edge labels are distinct and not equal to q. Thus, in a near-elegant labeling, instead of 0 being the missing value in the edge labels, q is the missing value. Deb and Limaye show that triangular snakes where the number of triangles is 3 (mod 4) are near-elegant. For any positive integers $\alpha \le \beta \le \gamma$ where β is at least 2, the *theta graph* $\theta_{\alpha,\beta,\gamma}$ consists of three edge disjoint paths of lengths α, β, and γ having the same end points. Deb and Limaye [914] provide elegant and near-elegant labelings for some theta graphs where $\alpha = 1, 2$, or 3. Seoud and Elsakhawi [3378] have proved that the following graphs are elegant: $K_{1,m,n}$; $K_{1,1,m,n}$; $K_2 + \overline{K_m}$; $K_3 + \overline{K_m}$; and $K_{m,n}$ with an edge joining two vertices of the same partite set. Elumalai and Sethuraman [1026] proved ${P_2}^n$, $P_m^2 + \overline{K_n}$, $S_m + S_n$, $S_m + \overline{K_m}$, $C_3 \times P_m$, and even cycles C_{2n} with vertices $a_0, a_1, \ldots, a_{2n-1}, a_0$ and $2n-3$ chords $a_0a_2, a_0a_3, \ldots, a_0a_{2n-2}$ ($n \ge 2$) are elegant. Husna and Rahmadani [1472] showed that nontrival cases the lotus graph and its two variations are elegant graphs. [1472] new Zhou [4338] has shown that for every strongly k-elegant graph G with p vertices and q edges and any positive integer m the graph $(G+\overline{K_m})+\overline{K_n}$ is also strongly k-elegant when $q-p+1 \le m \le q-p+k$.

If f is a strongly k-elegant labeling of a bipartite graph G with partite sets V_1 and V_2 and $\max f(u) < \min f(v)$ for all u in V_1 and v in V_2, f is said to be a *set-ordered strongly k-elegant* labeling of G. Su, Wang, and Yao [3701] proved that if a connected (p, q)-graph admits a strongly k-elegant labeling, then $q \le 2p-3$ and if a graph is a set-ordered strongly k-elegant, then $q = p-1$. They constructed several classes of large-scale trees that have strongly k-elegant labelings through graph operations that connect edges between two

vertices or identify two vertices to form a new graph and proved that caterpillars with p vertices admits a set-ordered strongly k-elegant labelings. They further showed that a graph admits a strongly k-elegant labeling if and only if it has a super edge-magic total labeling (SEMT)–see Section 5.2.

Sethuraman and Elumalai [3423] proved that every graph is a vertex induced subgraph of a elegant graph and present an algorithm that permits one to start with any non-trivial connected graph and successively form supersubdivisions (see §2.7) that have a strong form of elegant labeling. Acharya, Germina, Princy, and Rao [52] prove that every (p, q)-graph G can be embedded in a connected elegant graph H. The construction is done in such a way that if G is planar and elegant (harmonious), then so is H.

In [3422] Sethuraman and Elumalai define a graph H to be a *$K_{1,m}$-star extension* of a graph G with p vertices and q edges at a vertex v of G where $m > p - 1 - deg(v)$ if H is obtained from G by merging the center of the star $K_{1,m}$ with v and merging $p-1-deg(v)$ pendent vertices of $K_{1,m}$ with the $p - 1 - deg(v)$ nonadjacent vertices of v in G. They prove that for every graph G with p vertices and q edges and for every vertex v of G and every $m \geq 2^{p-1} - 1 - q$, there is a $K_{1,m}$-star extension of G that is both graceful and harmonious. In the case where $m \geq 2^{p-1} - q$, they show that G has a $K_{1,m}$-star extension that is elegant. Sethuraman and Selvaraju [3451] have shown that certain cases of the union of any number of copies of K_4 with one or more edges deleted and one edge in common are elegant.

In [1039] Ephremnath and Elumlai say a graph G is a *cycle with a chord Hamiltonian path* if G is obtained from the cycle $v_0, v_1, \ldots, v_{n-1}, v_0$ $(n \geq 6)$ by adding the chords $v_1v_{n-1}, v_vv_{n-2}, \ldots, v_\alpha v_\beta$ where $\alpha = \beta = (n-2)/2$ if n is even and $\alpha = (n+3)/2$, $\beta = (n-1)/2$ if n is odd. They proved that C_n $(n \geq 6)$ with a chord Hamiltonian path is harmonious and elegant.

Gallian extended the notion of harmoniousness to arbitrary finite Abelian groups as follows. Let G be a graph with q edges and H a finite Abelian group (under addition) of order q. Define G to be *H-harmonious* if there is an injection f from the vertices of G to H such that when each edge xy is assigned the label $f(x) + f(y)$ the resulting edge labels are distinct. When G is a tree, one label may be used on exactly two vertices. Beals, Gallian, Headley, and Jungreis [598] have shown that if H is a finite Abelian group of order $n > 1$ then C_n is H-harmonious if and only if H has a non-cyclic or trivial Sylow 2-subgroup and H is not of the form $Z_2 \times Z_2 \times \cdots \times Z_2$. Thus, for example, C_{12} is not Z_{12}-harmonious but is $(Z_2 \times Z_2 \times Z_3)$-harmonious. In [999] Ehard, Glock, and Joos apply rainbow colorings to graph decompositions and harmonious labeling of graphs.

Analogously, the notion of an elegant graph can be extended to arbitrary finite Abelian groups. Let G be a graph with q edges and H a finite Abelian group (under addition) with $q + 1$ elements. We say G is *H-elegant* if there is an injection f from the vertices of G to H such that when each edge xy is assigned the label $f(x) + f(y)$ the resulting set of edge labels is the non-identity elements of H. Beals et al. [598] proved that if H is a finite Abelian group of order n with $n \neq 1$ and $n \neq 3$, then C_{n-1} is H-elegant using only the non-identity elements of H as vertex labels if and only if H has either a non-cyclic or trivial Sylow 2-subgroup. This result completed a partial characterization of elegant

cycles given by Chang, Hsu, and Rogers [773] by showing that C_n is elegant when $n \equiv 2 \pmod 4$. Mollard and Payan [2541] also proved that C_n is elegant when $n \equiv 2 \pmod 4$ and gave another proof that P_n is elegant when $n \neq 4$. In 2014 Ollis [2736] used harmonious labelings for Z_m given by Beals, Gallian, Headley, and Jungreis in [598] to construct new Latin squares of odd order.

A function f is said to be an *odd-elegant* labeling of a graph G with q edges if f is an injection from the vertices of G to the integers from 0 to $2q-1$ such that the induced mapping $f^*(uv) = f(u) + f(v) \pmod{2q}$ from the edges of G to the odd integers between 1 to $2q-1$ is a bijection. Zhou, Yao, and Chen [4340] proved that every lobster is odd-elegant. In [4135] Wang, Xu, Ma, and Zhang gave a new type of graphical passwords based on odd-elegant labeled graphs. See also [4136] and [4321].

For a graph $G(V, E)$ and an Abelian group H Valentin [4042] defines a *polychrome* labeling of G by H to be a bijection f from V to H such that the edge labels induced by $f(uv) = f(v) + f(u)$ are distinct. Valentin investigates the existence of polychrome labelings for paths and cycles for various Abelian groups.

4.5 Felicitous Labelings

Another generalization of harmonious labelings are felicitous labelings. An injective function f from the vertices of a graph G with q edges to the set $\{0, 1, \ldots, q\}$ is called *felicitous* if the edge labels induced by $f(x) + f(y) \pmod q$ for each edge xy are distinct. (Recall a harmonious labeling only allows the vertex labels $0, 1, \ldots, q-1$.) This definition first appeared in a paper by Lee, Schmeichel, and Shee in [2190] and is attributed to E. Choo. Acharya, Germina, Princy, and Rao [52] observed that every harmonious labeling of a graph is also a felicitous labeling of the graph. Balakrishnan and Kumar [464] proved the conjecture of Lee, Schmeichel, and Shee [2190] that every graph is a subgraph of a felicitous graph by showing the stronger result that every graph is a subgraph of a sequential graph. Among the graphs known to be felicitous are: C_n except when $n \equiv 2 \pmod 4$ [2190]; $K_{m,n}$ when $m, n > 1$ [2190]; $P_2 \cup C_{2n+1}$ [2190]; $P_2 \cup C_{2n}$ [3876]; $P_3 \cup C_{2n+1}$ [2190]; $S_m \cup C_{2n+1}$ [2190]; K_n if and only if $n \leq 4$ [3422]; $P_n + \overline{K_m}$ [3422]; the friendship graph $C_3^{(n)}$ for n odd [2190]; $P_n \cup C_3$ [3499]; $P_n \cup C_{n+3}$ [3876]; and the one-point union of an odd cycle and a caterpillar [3499]. Shee [3495] conjectured that $P_m \cup C_n$ is felicitous when $n > 2$ and $m > 3$. Lee, Schmeichel, and Shee [2190] ask for which m and n is the one-point union of n copies of C_m felicitous. They showed that in the case where mn is twice an odd integer the graph is not felicitous. In contrast to the situation for felicitous labelings, we remark that C_{4k} and $K_{m,n}$ where $m, n > 1$ are not harmonious and the one-point union of an odd cycle and a caterpillar is not always harmonious. Lee, Schmeichel, and Shee [2190] conjectured that the n-cube is felicitous. This conjecture was proved by Figueroa-Centeno and Ichishima in 2001 [1084]. Savitha and Sasikala [3324] investigated felicitous labelings for the following families of graphs: the join of a star and $K_{m,n}$, generalized wheels with pendant vertices, and triangular ladders. [3324] new

Balakrishnan, Selvam, and Yegnanarayanan [466] obtained numerous results on felicitous labelings. The *wreath product*, $G * H$, of graphs G and H has vertex set

$V(G) \times V(H)$ and (g_1, h_1) is adjacent to (g_2, h_2) whenever $g_1g_2 \in E(G)$ or $g_1 = g_2$ and $h_1h_2 \in E(H)$. They define $H_{n,n}$ as the graph with vertex set $\{u_1, \ldots, u_n; v_1, \ldots, v_n\}$ and edge set $\{u_iv_j |\ 1 \le i \le j \le n\}$. They let $\langle K_{1,n} : m\rangle$ denote the graph obtained by taking m disjoint copies of $K_{1,n}$, and joining a new vertex to the centers of the m copies of $K_{1,n}$. They prove the —following are felicitous: $H_{n,n}$; $P_n * \overline{K_2}$; $\langle K_{1,m} : m\rangle$; $\langle K_{1,2} : m\rangle$ when $m \not\equiv 0 \pmod 3$, or $m \equiv 3 \pmod 6$, or $m \equiv 6 \pmod{12}$; $\langle K_{1,2n} : m\rangle$ for —all m and $n \ge 2$; $\langle K_{1,2t+1} : 2n+1\rangle$ when $n \ge t$; P_n^k when $k = n-1$ and $n \not\equiv 2 \pmod 4$, or $k = 2t$ and $n \ge 3$ and $k < n-1$; the join of a star and $\overline{K_n}$; and graphs obtained by joining two end vertices or two central vertices of stars with an edge. Yegnanarayanan [4263] conjectures that the graphs obtained from an even cycle by attaching n new vertices to each vertex of the cycle is felicitous. This conjecture was verified by Figueroa-Centeno, Ichishima, and Muntaner-Batle in [1089]. In [3447] Sethuraman and Selvaraju [3451] have shown that certain cases of the union of any number of copies of K_4 with 3 edges deleted and one edge in common are felicitous. Sethuraman and Selvaraju [3447] present an algorithm that permits one to start with any non-trivial connected graph and successively form supersubdivisions (see §2.7) that have a felicitous labeling. Krisha and Dulawat [2034] give algorithms for finding graceful, harmonious, sequential, felicitous, and antimagic (see §5.7) labelings of paths. A *linear cactus* $P_m(K_n)$ is a connected graph in which all the blocks are isomorphic to a complete graph K_n and block-cutpoint is a path P_{2m-1}. Gomathi [1294] proved the follow graphs are felicitous: $P_m(K_4)$, splitting graphs of $(B_{n,n})$, planar graphs $\mathrm{Pl}_{m,n}$, and $C_{2k+1} \odot S_m$. Gomathi and Nagarajan [1296] proved the following graphs are felicitous: a vertex switching of C_n $(n \ge 4)$, a vertex switching of C_n $(n \ge 4)$ with one chord, a vertex duplication of C_n, and the square of the book $B_{n,n}$ $(n \ge 2)$. Ezhilarasi Hilda and Jeba Jesintha [1438] proved that all shell flower graphs are felicitous. Devakirubanithi, Jeba Jesintha, and Abigail [933] proved that the duplication of the apex vertex of bistars, braid graphs, and globe graphs are felicitous.

In [3722] Sudhakar, Ranjani, Swathy, and Balaji provided a technique for coding a secret messages by applying an even felicitous labeling for the union of two star graphs using a GMJ (Graph Message Jumbled) code. They include two illustrations for converting plain text into cipher text (picture coding) and a method for a felicitous labeling of a graph. In [2416] Manshath, Hariprabakaran, Veerasamy, and Balaji used a GMJ code to create a confidential message by applying an even felicitous labeling to a bistar.

In [1023] Elumalai calls the graph obtained from a shell graph $C(2n+3, 2n)$ by adding a vertex in between each pair of adjacent vertices on the cycle C_{2n}, adding an edge in apex, and adding two more chords a *stem-lotus* graph . He proved that stem-lotus graphs are graceful and felicitous.

Figueroa-Centeno, Ichishima, and Muntaner-Batle [1090] define a felicitous graph to be *strongly felicitous* if there exists an integer k so that for every edge uv, $\min\{f(u), f(v)\} \le k < \max\{f(u), f(v)\}$. For a graph with p vertices and q edges with $q \ge p-1$ they show that G is strongly felicitous if and only if G has an α-labeling (see §3.1). They also show that for graphs G_1 and G_2 with strongly felicitous labelings f_1 and f_2 the graph obtained from G_1 and G_2 by identifying the vertices u and v such that $f_1(u) = 0 = f_2(v)$ is strongly felicitous and that the one-point union of two copies of C_m where $m \ge 4$ and m

is even is strongly felicitous. As a corollary they have that the one-point union of n copies of C_m where m is even and at least 4 and $n \equiv 2 \pmod 4$ is felicitous. They conjecture that the one-point union of n copies of C_m is felicitous if and only if $mn \equiv 0, 1$, or 3 (mod 4). In [1094] Figueroa-Centeno, Ichishima, and Muntaner-Batle prove that $2C_n$ is strongly felicitous if and only if n is even and at least 4. They conjecture [1094] that mC_n is felicitous if and only if $mn \not\equiv 2 \pmod 4$ and that $C_m \cup C_n$ is felicitous if and only if $m + n \not\equiv 2 \pmod 4$.

As consequences of their results about super edge-magic labelings (see §5.2) Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima [1094] have the following corollaries: if m and n are odd with $m \geq 1$ and $n \geq 3$, then mC_n is felicitous; $3C_n$ is felicitous if and only if $n \not\equiv 2 \pmod 4$; and $C_5 \cup P_n$ is felicitous for all n.

For a graph G with q edges Shainy and Balaji [3501] call a one-to-one function f from $V(G)$ to $\{0, 1, 2, \ldots, 2q-1\}$ a *even felicitous* if the edge labels generated by $(f(r) + f(s)) \bmod (2q-1)$ for each edge are even and distinct. They proved that stars, bistars, the union two stars, and the union of three stars are even felicitous graph.

In [2413] Manickam, Marudai, and Kala prove the following graphs are felicitous: the one-point union of m copies of C_n if $mn \equiv 1, 3 \bmod 4$; the one-point union of m copies of C_4; mC_n if $mn \equiv 1, 3 \pmod 4$; and mC_4. These results partially answer questions raised by Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima in [1090] and [1094].

Chang, Hsu, and Rogers [773] have given a sequential counterpart to felicitous labelings. They call a graph with q edges *strongly c-elegant* if the vertex labels are from $\{0, 1, \ldots, q\}$ and the edge labels induced by addition are $\{c, c+1, \ldots, c+q-1\}$. (A strongly 1-elegant labeling has also been called a *consecutive* labeling.) Notice that every strongly c-elegant graph is felicitous and that strongly c-elegant is the same as $(c, 1)$-arithmetic in the case where the vertex labels are from $\{0, 1, \ldots, q\}$. Chang et al. [773] have shown: K_n is strongly 1-elegant if and only if $n = 2, 3, 4$; C_n is strongly 1-elegant if and only if $n = 3$; and a bipartite graph is strongly 1-elegant if and only if it is a star. Shee [3496] has proved that $K_{m,n}$ is strongly c-elegant for a particular value of c and obtained several more specialized results pertaining to graphs formed from complete bipartite graphs.

Seoud and Elsakhawi [3380] have shown: $K_{m,n}$ $(m \leq n)$ with an edge joining two vertices of the same partite set is strongly c-elegant for $c = 1, 3, 5, \ldots, 2n + 2$; $K_{1,m,n}$ is strongly c-elegant for $c = 1, 3, 5, \ldots, 2m$ when $m = n$, and for $c = 1, 3, 5, \ldots, m + n + 1$ when $m \neq n$; $K_{1,1,m,m}$ is strongly c-elegant for $c = 1, 3, 5, \ldots, 2m+1$; $P_n + \overline{K_m}$ is strongly $\lfloor n/2 \rfloor$-elegant; $C_m + \overline{K_n}$ is strongly c-elegant for odd m and all n for $c = (m-1)/2, (m-1)/2 + 2, \ldots, 2m$ when $(m-1)/2$ is even and for $c = (m-1)/2, (m-1)/2 + 2, \ldots, 2m - (m-1)/2$ when $(m-1)/2$ is odd; ladders L_{2k+1} $(k > 1)$ are strongly $(k+1)$-elegant; and $B(3, 2, m)$ and $B(4, 3, m)$ (see §2.4 for notation) are strongly 1-elegant and strongly 3-elegant for all m; the composition $P_n[P_2]$ (see §2.3 for the definition) is strongly c-elegant for $c = 1, 3, 5, \ldots, 5n - 6$ when n is odd and for $c = 1, 3, 5, \ldots, 5n - 5$ when n is even; P_n is strongly $\lfloor n/2 \rfloor$-elegant; P_n^2 is strongly c-elegant for $c = 1, 3, 5, \ldots, q$ where q is the number of edges of P_n^2; and P_n^3 $(n > 3)$ is strongly c-elegant for $c = 1, 3, 5, \ldots, 6k-1$ when $n = 4k$; $c = 1, 3, 5, \ldots, 6k + 1$ when $n = 4k + 1$; $c = 1, 3, 5, \ldots, 6k + 3$ when $n = 4k + 2$; $c = 1, 3, 5, \ldots, 6k + 5$ when $n = 4k + 3$.

In [537] Barrientos and Minion study a technique to transform an α-labeling of some snakes whose cells are squares into a felicitous labeling and the felicitous labeling into a harmonious labeling. They prove that all quadrilateral snakes, all snake polyominoes, and all hybrid quadrilateral snakes are both, felicitous (see §4.5) and harmonious. A *hybrid quadrilateral snake* is a snake obtained with n copies of C_4 where the ith copy of C_4 is attached to the $(i+1)$th copy via vertex amalgamation or edge amalgamation. Barrientos and Minion [537] prove that all hybrid quadrilateral snakes admit α-labelings.

4.6 Odd Harmonious and Even Harmonious Labelings

Liang and Bai [2254] introduced odd harmonious labelings by defining a function f to be an *odd harmonious* labeling of a graph G with q edges if f is an injection from the vertices of G to the integers from 0 to $2q-1$ such that the induced mapping $f^*(uv) = f(u)+f(v)$ from the edges of G to the odd integers between 1 to $2q-1$ is a bijection. A function f is said to be a *strongly odd harmonious* labeling of a graph G with q edges if f is an injection from the vertices of G to the integers from 0 to q such that the induced mapping $f^*(uv) = f(u)+f(v)$ from the edges of G to the odd integers between 1 to $2q-1$ is a bijection. Liang and Bai [2254] have shown the following: odd harmonious graphs are bipartite; if a (p,q)-graph is odd harmonious, then $2\sqrt{q} \le p \le 2q-1$; if a (p,q)-graph with degree sequence $(d_1, d_2, \ldots, d_p)$ is odd harmonious, then $\gcd(d_1, d_2, \ldots, d_p)$ divides q^2; P_n $(n>1)$ is odd harmonious and strongly odd harmonious; C_n is odd harmonious if and only if $n \equiv 0 \bmod 4$; K_n is odd harmonious if and only if $n = 2$; $K_{n_1,n_2,\ldots,n_k}$ is odd harmonious if and only if $k = 2$; K_n^t is odd harmonious if and only if $n = 2$; $P_m \times P_n$ is odd harmonious; the tadpole graph obtained by identifying the endpoint of a path with a vertex of an n-cycle is odd harmonious if $n \equiv 0 \bmod 4$; the graph obtained by appending two or more pendent edges to each vertex of C_{4n} is odd harmonious; the graph obtained by subdividing every edge of the cycle of a wheel (gear graphs) is odd harmonious; the graph obtained by appending an edge to each vertex of a strongly odd harmonious graph is odd harmonious; and caterpillars and lobsters are odd harmonious. They conjecture that every tree is odd harmonious.

Let $G(V,E)$ be a graph, and let $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ be a bijective vertex labeling. The induced edge labeling $f^* : E(G) \to \{0, 1\}$ is defined by assigning the label 1 to an edge uv whenever the labels $f(u)$ and $f(v)$ have the same parity, and the label 0 whenever they have different parity. If the difference between the number of edges labeled 0 and the number of edges labeled 1 is at most one, then f is called an *even–odd cordial* labeling. In [2655] Naresh kumar and Bharathi investigated the existence of even–odd cordial labelings for several families of zero divisor-graphs. [2655] new

Liang and Bai [2254] also showed that the kC_4-snake graph is an odd harmonious graph. Abdel-Aal [4] generalize this result by showing that the kC_n-snake with string $1, 1, \ldots, 1$ for $n \equiv 0 \pmod 4$ are odd harmonious. He also showed that the kC_4 snake with m pendent edges is odd harmonious and that all subdivisions of $2m$-triangular snakes are odd harmonious. Alyani, Firmansah, Giyarti, and Sugeng [206] constructed odd harmonious labelings for kC_n-snakes for $n = 4$ and $n = 8$ and gave odd harmonious

labelings for a variation of kC_n-snakes. An n *hair-*kC_4 *snake* is a graph obtained by attaching n leaves to vertices of degree two in a kC_4-snake graph. Mumtaz and Silaban [2566] proved that n hair-kC_4-snakes are odd harmonious. A $k(G)$ *snake graph* is a graph obtained from a path on k edges by replacing each edge by a graph isomorphic to G. Asumpta, Purwanto, and Chandra [319] showed that the snake graph $k(G)$ is odd harmonious when G is a gear graph based on W_3 and when G is $K_{2,m}$ $(m \geq 2)$. In [1102] Firmansah, Tasari, and Yuwono defined three new classes of flower graphs: lotus flower graphs, lotus flower graphs with pendant, and a variation of lotus flower graphs and proved that they are odd harmonious graphs. [1102] new

The *prism snake* graph PS_m is constructed from m copies of the cycle graph C_4 by connecting three vertices in the same position from two adjacent copies of C_4. The *generalized chain prism snake* graph $PS^+_{m,n}$ is a graph constructed from n copies of the chain prism snake graph PS^+_m by attaching each vertex and edge of two adjacent copies of PS^+_m. Winarti and Rahmadani [4209] showed that PS_m and $PS^+_{m,n}$ are odd harmonious graphs for all $m, n \in \mathbb{N}$. [4209] new

Renuka and Palanivelu [3163] proved that the one vertex union of some classes of complete bipartite graphs, the one vertex union of paths, and extended bistars and their subdivisions are odd harmonious. They further proved that the collection of paths passing through another path at a selected common mid vertex and each vertex of ladder appended by an edge are strongly odd harmonious and that the subdivision of the Cartesian product of two paths is strongly odd harmonious.

In [3383], Seoud and Hafez proved that a graph has an odd harmonious labeling if and only if it has a bigraceful labeling. They also proved that the number of distinct odd harmonious labeled graphs on m edges is $m!$ and the number of distinct strongly odd harmonious labeled graphs on m edges is $\lfloor m/2 \rfloor !$ Seoud and Hafez also proved that if G is a strongly odd harmonious graph, then the graph obtained by identifying a distinguished vertex of G to each vertex of C_{4m} is strongly odd harmoniou. They further proved that the Cartesian product of strongly odd harmonious trees is strongly odd harmonious.

Abdel-Aal also proved that a necessary condition for odd harmonious Eulerian graphs with q edges is $q \equiv 0 \pmod 4$ and that the following graphs are odd harmonious: $C_m \times P_n$ $(n \geq 2, m \equiv 0 \pmod 4)$; $C_{4m} \odot C_4$; $S_n \odot \overline{K_m}$; two copies of an even n-cycle sharing a common edge is an odd harmonious graph when $n \equiv 0 \pmod 4$; two copies of an even n-cycle sharing a common vertex is odd harmonious when $n \equiv 0 \pmod 4$; and graphs obtained from $K_{2,n}$ $(n \geq 2)$ by adding r pendent edges to one of the two vertices of degree n and s pendent edges to the other vertex of degree n. In [2565] Mumtaz, John, and Silaban proved that the grid-like graph of order $2mn+mm+n$ and size $4mn$ obtained by arranging m-rows connected rows of nC_4-snake graphs is odd harmonious.

Vaidya and Shah [4002] prove that the shadow graphs (see §3.8 for the definition) of path P_n and star K_{1n} are odd harmonious. They also show that the splitting graphs (see §2.7 for the definition)) of path P_n and star $K_{1,n}$ are odd harmonious. In [4003] Vaidya and Shah proved the following graphs are odd harmonious: the shadow graph and the splitting graph of bistar $B_{n,n}$; the arbitrary supersubdivision of paths; graphs obtained by joining two copies of cycle C_n for $n \equiv 0 (\text{mod } 4)$ by an edge; and the graphs

$H_{n,n}$, where $V(H_{n,n}) = \{v_1, v_2, \ldots, v_n, u_1, u_2, \ldots, v_n\}$ and $E(H_{n,n}) = \{v_i u_j : \ 1 \leq i \leq n, \ n - i + 1 \leq j \leq n\}$. In [4239] Yan proves that $P_m \times P_n$ is strongly odd harmonious. Koppendrayer [1999] has proved that every graph with an α-labeling is odd harmonious. Li, Li, and Yan [2242] proved that $K_{m,n}$ is odd strongly harmonious. Saputri, Sugeng, and Fronček [3311] proved that the graph obtained by joining C_n to C_k by an edge (dumbbell graph $D_{n,k,2}$) is odd harmonious for $n \equiv k \equiv 0 \pmod 4$ and $n \equiv k \equiv 2 \pmod 4$, and $C_n \times P_m$ is odd harmonious if and only if $n \equiv 0 \pmod 4$. They also observe that $C_n \odot K_1$ with $n \equiv 0 \pmod 4$ is odd harmonious. Pujiwatia, Halikinb, and Wijayac [3055] proved double stars and even cycles odd harmonious. Sugeng, Surip, and Rismayati [3740] proved that the m-shadow of C_{4n} and nontrivial gears are odd harmonious graphs.

Jeyanthi [1716] proved that the shadow and splitting graphs of $K_{2,n}$, C_{4n}, the double quadrilateral snakes $DQ(n)$ ($n \geq 2$), and the graph $H_{n,n}$ with vertex set $V(H_{n,n}) = \{v_1, v_2, \ldots, v_n, u_1, u_2, \ldots, u_n\}$ and the edge set $E(H_{n,n}) = \{v_i u_j : 1 \leq i \leq n, n - i + 1 \leq j \leq n\}$ are odd harmonious. Jeyanthi and Philo [1716] proved that the shadow graphs $D_2(K_{2,n})$ and $D_2(H_{n,n})$ are odd harmonious and the splitting of graphs of $K_{2,n}$ and $H_{n,n}$ are odd harmonious. They also showed that the shadow graph $D_2(C_n)$ is odd harmonious if $n \equiv 0 \pmod 4$, the splitting of C_n is odd harmonious if $n \equiv 0 \pmod 4$, and the double quadrilateral snake $DQ(n)$ is odd harmonious for $n \geq 2$. In [1720] Jeyanthi and Philo prove that super subdivision of cycles, ladders, $C_{4n} \oplus K_{1,m}$, and uniform fire crackers are odd harmonious graphs. Jeyanthi and Philo [1726] proved that the graph $P_{n-1}(1, 2, 3, \ldots, n)$ obtained from a path of n vertices $v_1, v_2, \ldots, v_{n-1}$ by appending a path of length $n - i$ at each v_i and certain one point unions of the end points of paths are odd harmonious.

Philo, Jeyanthi, and Davazz [3491] proved that the following graphs are odd harmonious: the path union of r copies of $K_{m,n}$; the path union of copies of complete bipartite graphs; the graphs obtained from t copies of $K_{m,n}$ by joining each one to the next one with an isolated vertex (the *join sum*) ; the one point union of t copies of complete bipartite graphs (not necessarily the same); the graphs obtained by replacing each vertex of $K_{1,t}$, except the apex vertex, with the graph $K_{m,n}$; and the one point union of isomorphic path graphs of $K_{m,n}$. Let $u_{i,1}, u_{i,2}, \ldots, u_{i,m}$ and $v_{i,1}, v_{i,2}, \ldots, v_{i,n}$ be the vertices of r copies of $K_{m,n}$. The *circle formation* [3491] of r copies of $K_{m,n}$ is the graph obtained by joining the ith copy of $K_{m,n}$ ($1 \leq i \leq r$) by joining the vertices $u_{i,m}$ to $u_{i+1,1}$ for $1 \leq i \leq r-1$ and joining the vertex $u_{r,m}$ to $u_{1,1}$. Philo, Jeyanthi, and Davazz [3491] proved that he circle formation of r copies of $K_{m,n}$ when $r = 0 (\mathrm{mod}\, 4)$ is odd harmonious.

Suptri, Sugeng, and Fronček [3311] proved that dumbbell graphs $D_{n,k,2}$ are odd harmonious if and only if $n, k = 2 (\mathrm{mod}\, 4)$. In [1722] Jeyanthi and Philo proved that the graphs obtained by attaching m pendant vertices to each vertex of paths of odd length, the splitting graph of combs, slanting ladders, graphs obtained from m copies of $\langle K_{1,n} : K_{1,n} \rangle$ by joining one leaf of ith copy of $\langle K_{1,n} : K_{1,n} \rangle$ with the center of $(i+1)$th copy of $\langle K_{1,n} : K_{1,n} \rangle$ where $1 \leq i \leq m - 1$, and H-super subdivisions of P_n and C_{4n} admit odd harmonious labeling. Moreover, they observe that all strongly odd harmonious graphs admit a mean labeling, an odd mean labeling, an odd sequential labeling, all odd sequential graphs are odd harmonious, and all odd harmonious graphs are even sequential harmonious.

They also proved the n-splitting graphs for paths, stars, and symmetric product between paths and null graphs are odd harmonious graphs for all n. Selvaraju, Balaganesan, and Renuka [3347] proved that quadrilateral snakes and k-regular caterpillars are odd harmonious. Abdel-Aal1 and Seoud [7] proved that the m-shadow graphs for paths and complete bipartite graphs are odd harmonious graphs for all m.

For a T_p-tree T with vertices $v_1, v_2, \ldots, v_n$, the graph $T\hat{\circ}P_m$ is obtained from T and n copies of P_m by identifying a pendant vertex of ith copy of P_m with vertex v_i of T. For C_n with consecutive vertices $v_1, v_2, \ldots, v_n$, the graph $C_n\hat{\circ}P_m$ is obtained from C_n and n copies of P_m by identifying a pendant vertex of ith copy of P_m with vertex v_i of C_n. Jeyanthi and Philo [1728] proved that T_p-trees, $T\hat{\circ}P_m$, $T\hat{\circ}2P_m$, regular bamboo trees, $C_n\hat{\circ}P_m$, $C_n\hat{\circ}2P_m$, and subdivided grid graphs are odd harmonious.

Recall a *subdivided shell graph* is obtained by subdividing the edges in the path of the shell graph. Let $G_1, G_2, \ldots, G_n$ be n subdivided shell graphs of any order. The graph SSG(n) is obtained by adding an edge to apexes of G_i and $G_{i+1}, i = 1, 2, \ldots, (n-1)$. Jeba Jesintha and Ezhilarasi Hilda [1618] proved that the subdivided shell graph and SSG(2) are odd harmonious.

The following definitions are taken from [1727]. The *m-shadow graph* $D_m(G)$ of a connected graph G is constructed by taking m-copies of G, G_1,G_2,G_3,...,G_m, and joining each vertex u in G_i to the neighbors of the corresponding vertex v in G_j, $1 \leq j \leq m$. The *m-splitting graph* $Spl_m(G)$ of a graph G is obtained by adding to each vertex v of G m new vertices, $v^1, v^2, \ldots, v^m$ such that v^i, $1 \leq i \leq m$, is adjacent to every vertex that is adjacent to v in G. Note that the 2-shadow graph is the shadow graph $D_2(G)$ and the 1-splitting graph is splitting graph. The *m-mirror graph* $M_m(G)$ is defined as the disjoint union of m copies of G, $G_1, G_2, \ldots, G_m$, together with additional edges joining each vertex of G_i to its corresponding vertex in G_{i+1}, $1 \leq i \leq m-1$. The graph $\overline{W_{m,n}}$ is obtained from the gear graph arising from the wheel W_n as follows: Join the vertices v_i and v_{i+2} with the new vertices v_{i+1}^j for $1 \leq j \leq m$ and $2 \leq i \leq n-2$ and join v_n and v_2 with v_{2i-1}. The graph $K_{2,n}(r, s)$ is obtained from $K_{2,n}$ $(n \geq 2)$ by adding r and s pendent edges to the two vertices of degree n. The graph $G = \langle C_n : K_{2,m} : C_r \rangle$ is obtained from $K_{2,m}$ with the partite set $\{u, v\}$ by identifying the vertex u with a vertex of C_n and the vertex v with a vertex of C_r. Let P_n be a path on n vertices denoted by $(1,1), (1,2), \ldots, (1,n)$ and with $n-1$ edges denoted by $e_1, e_2, \ldots, e_{n-1}$ where e_i is the edge joining the vertices $(1,i)$ and $(1,i+1)$. The *step ladder* graph $S(T_n)$ has $(n^2+3n-2)/2$ vertices denoted by $(1,1), (1,2), \ldots, (1,n), (2,1), (2,2), \ldots, (2,n), (3,1), (3,2), \ldots, (3,n-1), (4,1), \ldots, (4,n-2), \ldots, (n,1), (n,2)$ and n^2+n+2 edges. In any ordered pair (i,j), i denotes the row (counted from bottom to top) and j denotes the column(from left to right) in which the vertex occurs.

The *cocktail party* graph, $H_{m,n}$ $(m, n \geq 2)$, is the graph with a vertex set $V = \{v_1, v_2, \ldots, v_{mn}\}$ partitioned into n independent sets $V = \{I_1, I_2, \ldots, I_n\}$ each of size m such that $v_i v_j \in E$ for all $i, j \in \{1, 2, \ldots, mn\}$ where $i \in I_p$, $j \in I_q$, $p \neq q$.

Jeyanthi and Philo [1719] proved that the following graphs are odd harmonious: $D_m(P_n)$ for all $m, n \geq 2$; $Spl_m(P_n)$ for $m, n \geq 2$; $D_m(H_{n,n})$ for all $m \geq 2$ and $n \geq 1$; $Spl_m(H_{n,n})$ for all $m \geq 2$ and $n \geq 1$; $D_m(K_{r,s})$ for all $r, s \geq 1$; $Spl_m(K_{r,s})$ for all $m \geq 2$

and $r, s \geq 1$; $D_m(P_n \oplus \overline{K_2})$ for all $m, n \geq 2$; $Spl_m(P_n \oplus \overline{K_2})$, $m, n \geq 2$; and $Spl_m(C_n)$ if and only if $n \equiv 0 \pmod 4$.

Jeyanthi and Philo [1727] proved that following graphs are odd harmonious: $\overline{W_{m,n}}$ for $n \equiv 0 \pmod 4$, $m \geq 1$; $D_m(P_n \odot K_1)$ (the authors use the notion C_{bn} for the comb $P_n \odot K_1$) for all $m \geq 2$ and $n \geq 1$; $Spl_m(K_{2,n}(r, s))$; $\langle C_n : K_{2,m} : C_r \rangle$ for $n, r \equiv 0 \pmod 4$ and $m \geq 2$; and the graphs obtained by arranging vertices into a finite number of rows (at least 2) with i vertices in the ith row and in every row the jth vertex in that row is joined to the jth vertex and $j+1$st vertex of the next row (a *pyramid*) for $n \geq 2$. They also prove that if G is a strongly odd harmonious tree, then $M_m(G)$ is odd harmonious.

Let P_{2n} be a path of length $2n-1$ with $2n$ vertices, denoted by $(1,1), (1,2), \ldots, (1,2n)$ and with $2n-1$ edges, denoted by $e_1, e_2, \ldots, e_{2n-1}$ where e_i is the edge joining the vertices $(1,i)$ and $(1,i+1)$. On each edge e_i for $i = n+1, n+2, \ldots, 2n-1$, we erect a ladder with $2n+1-i$ steps including the edge e_i. The double sided step ladder graph $2S(T_{2\times n})$ has vertices denoted by $(1,1), (1,2), \ldots, (1,2n)$,$(2,1), (2,2), \ldots, (2,2n)$,$(3,2), (3,3), \ldots, (3,2n-1)$,$(4,3), (4,4), \ldots, (4,2n-2), \ldots, (n+1,n), (n+1,n+1)$. In any ordered pair (i,j), i denotes the row (counted from bottom to top) and j denotes the column (from left to right) in which the vertex occurs. Jeyanthi and Philo [1725] proved that the path union of t copies of $S(T_n)$, the double sided step ladder $2S(T_{2\times n})$, the path union of t copies of $2S(T_{2\times n})$, $S(t.C_{bn})$, $S(t.C_4)$, ${C_4}^t$, ${C_6}^t$, and ${C_8}^t$ are odd harmonious graphs. Jeyanthi and Philo [1721] proved that path union of r copies of $K_{m,n}$, the path union of r copies of K_{m_i,n_i}, $1 \leq i \leq r$, $K^t_{m,n}$, $K^t_{(m_1,n_1),(m_2,n_2),\ldots,(m_t,n_t)}$, the join sum of graph $\langle K_{m,n}; K_{m,n}; \ldots, K_{m,n}$ (t copies $\rangle$, $\langle K_{m_1,n_1}; K_{m_2,n_2}; \ldots, K_{m_t,n_t} \rangle$, the circle formation of r copies of $K_{m,n}$ when $r \equiv 0 \pmod 4$, $S(t.K_{m,n})$ and $P^t_n(t.n.K_{p,q})$ are odd harmonious graphs. Jeyanthi and Philo [1724] proved that the subdivided shell graphs, disjoint union of two subdivided shell graphs, subdivided shell flower graphs, and subdivided uniform shell bow graphs are odd harmonious. Jeyanthi, Philo, and Youssef [1729] proved that the path union of t copies of $P_m \times P_n$, the path union of t copies of $P_{m_i} \times P_{n_i}$ where $1 \leq i \leq t$, the vertex union of t copies of $P_m \times P_n$, the vertex union of t different copies of $P_{m_i} \times P_{n_i}$ where $1 \leq i \leq t$, the one point union of path of $P^t_n(t.n.P_m \times P_m)$, and the super subdivision of grid graph $P_m \times P_n$ are odd harmonious graphs.

Recall from Section 2.7 that for even $n > 2$ a *plus graph* of size n (denoted by Pl_n) is the graph obtained by starting with paths $P_2, P_4, \ldots, P_{n-2}, P_n, P_n, P_{n-2}, \ldots, P_4, P_2$ arranged vertically parallel with the vertices in the paths forming horizontal rows and edges joining the vertices of the rows. Jeyanthi [1718] proved that following graphs are odd harmonious: Pl_n where $n \equiv 0 \pmod 2$, $n \neq 2$; path unions of finitely many copies of Pl_n where $n \equiv 0 \pmod 2$, $n \neq 2$; open stars of plus graphs $S(t.Pl_n)$ where $n \equiv 0 \pmod 2$, $n \neq 2$ and t odd; graphs obtained by joining C_m, $m \equiv 0 \pmod 4$ and a plus graph Pl_n, $n \equiv 0 \pmod 2$, $n \neq 2$ with a path of arbitrary length; the graph obtained by replacing all vertices of $K_{1,t}$, except the apex vertex, by the path union of n copies of the graph Pl_m.

In [1720] Jeyanthi and Philo prove that super subdivision of cycles, ladders, $C_{4n} \oplus K_{1,m}$, and uniform fire crackers are odd harmonious graphs. They also proved the (m,n)-firecracker graph obtained by the concatenation of m n-stars by linking one leaf from each is odd harmonious; the arbitrary super subdivision of cycles C_m are odd harmonious; and

the super subdivision of ladders are odd harmonious. Jeyanthi and Philo [1727] proved that the m-mirror graph $M_m(G)$ $(m \geq 2)$, m-splitting graph of $K_{2,n}(r,s)$ (obtained from $K_{2,n}$, $(n > 2)$ by adding r and s $(r,s > 1)$ pendent edges to the two vertices of degree n), $\overline{W_{(m,4n)}}$ obtained from the gear graph of W_n by joining the vertices v_i and v_{i+2} with the new vertices v_{i+1}^j for $1 \leq j \leq m$ and $2 \leq i \leq n-2$ and joining v_n and v_2 with v_1^j for $1 \leq j \leq m$, $\langle C_{4n} : K_{2,m} : C_{4r} \rangle$ obtained from $K_{2,m}$ with one partite set $V_1 = \{u, v\}$ and C_r by identifying the vertex u of V_1 with a vertex of C_n and the other vertex v of V_1 with a vertex of C_r, and the pyramid graph $PY_n(n \geq 2)$ are odd harmonious graphs. They also proved that G is a strongly odd harmonious tree, then $M_m(G)$ is an odd harmonious.

The *edge comb product* of two graphs G and H is the graph formed by taking one copy of G and $|E(G)|$ copies of H, then attaching the i-th copy of H at the edge e to the i-th edge of G. In [3235] Sarasvati, Halikin, and Wijaya showed that graphs constructed by the edge comb product of P_n and C_4 and the shadow of C_4 are odd harmonious. Firmansah and Tasari [1104] gave odd harmonious labelings for a line amalgamation of double quadrilateral graphs and the graphs obtained by connecting two copies of double quadrilateral graph by an edge. Firmansah and Giyarti [1101] proved that graphs obtained by the edge amalgamation of double quadrilateral graphs are odd harmonious. Febriana and Sugeng [1074] proved that squid graphs (obtained from C_n by add pendant edges to one vertex of C_n) are odd harmonious if and only if n is even and double squid graphs (obtained from two copies of C_n that share one common vertex and adding pendant edges to the common vertex) are odd harmonious if and only $n \geq 4$ is even. In [1098], [1097], [1100] Firmansah proved that multiple net snake graphs and a variation of the double quadrilateral windmill graphs, layered graphs, and some classes of string graphs are odd harmonious graphs. Firmansah, Tasari, and Yuwono [1103] proved that the zinnia flower graph and its variations are odd harmonious graphs.

In [934] Devakirubanithi, Jeba Jesintha, and Benita define a *modified arrow graph*, $MAnm$, with dimensions m (width) and n (length) as one obtained by adding a vertex v and connecting it with the first and last superior vertices of $P_m \times P_n$ with two additional edges coming from one end. They define a *modified double arrow* graph, $MD(A_{nm}$, with dimensions m (width) and n (length) as one obtained by connecting two vertices v and w with first and last superior vertices of $P_m \times P_n$ with two additional edges coming from the two ends. They proved that modified double arrow graphs, modified arrow graphs, and the corona product of a T_p tree and a star are odd harmonious..

In [1720] Jeyanthi and Philo modified the notion of odd harmonious by defining an odd harmonious labelings as a function f to be an *odd harmonious* labeling of a graph G with q edges if f is an injection from the vertices of G to the integers from 0 to $2q-1$ such that the induced mapping $f^*(uv) = f(u) + f(v) \mod (2q)$ from the edges of G to the odd integers between 1 to $2q-1$ is a bijection. Using this definition they proved that an m-cycle and an n-cycle sharing a common vertex is an odd harmonious if and only if either both $m,\ n \equiv 0 \pmod 4$ or both $m,\ n \equiv 2 \pmod 4$ and the same holds for an m-cycle and an n-cycle sharing a common edge. They also proved that any two even cycles sharing a common vertex and a common edge are odd harmonious graphs.

Sarasija and Binthiya [3312] say a function f is an *even harmonious* labeling of a

graph G with q edges if $f : V \to \{0, 1, \ldots, 2q\}$ is injective and the induced function $f^* : E \to \{0, 2, \ldots, 2(q-1)\}$ defined as $f^*(uv) = f(u)+f(v) \pmod{2q}$ is bijective. Notice that for an even harmonious labeling of a connected graph all the vertex labels must have the same parity. Moreover, in the case of even harmonious labelings for connected graphs there is no loss of generality to assume that all the vertex labels are even integers and the duplicate vertex is 0. They proved the following graphs are even harmonious: non-trivial paths; complete bipartite graphs; odd cycles; bistars $B_{m,n}$; $K_2+\overline{K_n}$; P_n^2; and the friendship graphs F_{2n+1}. López, Muntaner-Batle and Rius-Font [2319] proved that every super edge-magic graph (see Section 5.2 for the definition of super edge-magic) with p vertices and q edges where $q \geq p-1$ has an even harmonious labeling. In [4286] Youssef provided a necessary condition for some regular graphs to be even harmonious, showed that the disjoint union of two k-sequential graphs is even $2k$-sequential under some conditions, and showed that in some cases G is k-sequential implies mG is even $2k$-sequential for all positive integer m.

Because 0 and $2q$ are equal modulo $2q$ the following restricted form of even harmonious labelings is of interest. A function f is said to be a *properly even harmonious* labeling of a graph G with q edges if f is an injection from the vertices of G to the integers from 0 to $2q-1$ and the induced function f^* from the edges of G to $\{0, 2, \ldots, 2q-2\}$ defined by $f^*(xy) = f(x)+f(y) (\mathrm{mod}\ 2q)$ is bijective. In their definition of properly even harmonious in [1175] Gallian and Schoenhard incorrectly required that the vertex labels should be the even integers from 0 to $2q-2$. For connected graphs the two definitions are equivalent but for disconnected graph they are not. They used vertex labels from 0 to $2q-1$ for their results on disconnected graphs.

A graph with a properly even harmonious labeling is said to be *properly even harmonious*. Gallian and Schoenhard [1175] say a properly even harmonious labeling of a graph with q edges is *strongly even harmonious* if it satisfies the additional condition that for any two adjacent vertices with labels u and v, $0 < u + v \leq 2q$.

Jared Bass [594] has observed that for connected graphs any harmonious labeling of a graph with q edges yields an even harmonious labeling by simply multiplying each vertex label by 2 and adding the vertex labels modulo $2q$. Thus we know that every connected harmonious graph is an even harmonious graph and every connected graph that is not a tree that has a harmonious labeling also has a properly even harmonious labeling. Conversely, a properly even harmonious labeling of a connected graph with q edges (assuming that the vertex labels are even) yields a harmonious labeling of the graph by dividing each vertex label by 2 and adding the vertex labels modulo q.

Gallian and Schoenhard [1175] proved the following: wheels W_n and helms H_n are properly even harmonious when n is odd; nP_2 is even harmonious for n odd; nP_2 is properly even harmonious if and only if n is even; K_n is even harmonious if and only if $n \leq 4$; C_{2n} is not even harmonious when n is odd; $C_n \cup P_3$ is properly even harmonious when odd $n \geq 3$; $C_4 \cup P_n$ is even harmonious when $n \geq 2$; $C_4 \cup F_n$ is even harmonious when $n \geq 2$; $S_m \cup P_n$ is even harmonious when $n \geq 2$; $K_4 \cup S_n$ is properly even harmonious; $P_m \cup P_n$ is properly even harmonious for all $m \geq 2$ and $n \geq 2$; $C_3 \cup P_n^2$ is even harmonious when $n \geq 2$; $C_4 \cup P_n^2$ is even harmonious when $n \geq 2$; the disjoint union of two or three

stars where each star has at least two edges and one has at least three edges is properly even harmonious; $P_m^2 \cup P_n$ is even harmonious for $m \geq 2$ and $2 \leq n < 4m-5$; the one-point union of two complete graphs each with at least 3 vertices is not even harmonious; $S_m \cup P_n$ is strongly even harmonious if $n \geq 2$; and $S_{n_1} \cup S_{n_2} \cup \cdots \cup S_{n_t}$ is strongly even harmonious for $n_1 \geq n_2 \geq \cdots \geq n_t$ and $t < \frac{n_1}{2} + 2$. They conjecture that $S_{n_1} \cup S_{n_2} \cup \cdots \cup S_{n_t}$ is strongly even harmonious if at least one star has more than 2 edges. They also note that $C_4, C_8, C_{12}, C_{16}, C_{20}, C_{24}$ are even harmonious and conjecture that C_{4n} is even harmonious for all n. This conjecture was proved by Youssef [4284] who also proved that if a connected even harmonious graph with q edges where q is even and each vertex has degree divisible by 2^k $(k \geq 1)$, then q is divisible by 2^{k+1}. As a corollary of the latter he gets that C_{4n+2}^2 is not even harmonious. Hall, Hillesheim, Kocina, and Schmit [1357] proved that nC_{2m+1} is properly even harmonious for all n and m.

In [1176] and [1177] Gallian and Stewart investigated properly even harmonious labelings of unions of graphs. They use ${P_m}^{+t}$ to denote the graph obtained from the path P_m by appending t edges to an endpoint; ${Cat_m}^{+t}$ to denote a caterpillar of path length m with t pendent edges; and ${C_m}^{+t}$ to denote an m-cycle with t pendent edges. They proved the following graphs are properly even harmonious: nP_m if n is even and $m \geq 2$; $P_n \cup K_{m,2}$ for n odd and $n > 1,\ m > 1$; $P_n \cup S_{m_1} \cup S_{m_2}$ for $n > 2$ and $m_1 + m_2$ is odd; $C_n \cup S_{m_1} \cup S_{m_2}$ for n odd and $m_1,\ m_2 > 3$; ${P_m}^{+t} \cup {P_n}^{+s}$; the union of any number of caterpillars; $C_m \cup {Cat_n}^{+t}$ for $m > 1$ odd, $n > 1$; $C_4 \cup {Cat_m}^{+t}$; the union of C_4 and a hairy cycle; $K_4 \cup {C_m}^{+n}$ for some cases; $W_4 \cup {C_m}^{+n}$ for some cases; $C_4 \cup (P_n + \overline{K_2})$ for $n > 1$; $K_4 \cup (P_n + \overline{K_m})$ for $n \equiv 1, 2 \pmod 4$; $C_3 \cup (P_n + \overline{K_m})$ for $n \equiv 1, 2 \pmod 4$; $W_4 \cup (P_n + \overline{K_m})$ for $n \equiv 1, 2 \pmod 4$; $W_4 \cup P_n$ for $n \equiv 1, 2 \pmod 4$; $K_4 \cup P_n$ for $n > 1$ and $n \equiv 1, 2 \pmod 4$; $K_4 \cup P_{m_1}^2 \cup P_{m_2}^2 \cup \cdots \cup P_{m_n}^2$ for $m_i > 2,\ n \geq 1$; $W_4 \cup P_{m_1}^2 \cup P_{m_2}^2 \cup \cdots \cup P_{m_n}^2$ for $m_i > 2,\ n \geq 1$; $C_m \cup P_n^2$ for $m \equiv 3 \pmod 4$ and $n > 1$; and $2P_m \cup 2P_n$. They also prove that nP_3 is even harmonious if $n > 1$ is odd and $P_{m_1}^2 \cup P_{m_2}^2 \cup \cdots \cup P_{m_n}^2$ is strongly even harmonious for $m > 2,\ n \geq 1$.

Gallian and Stewart [1178] call an injective labeling f of a graph G with q edges *even* $2a$*-sequential* if the vertex labels are from $\{0, 1, \ldots, 2q-1\}$ and the edge labels induced by $f(u) + f(v)$ for each edge uv are $2a, 2a+2, \ldots, 2a+2q-2$. When G is a tree, the allowable vertex labels are $0, 1, \ldots, 2q$. For connected a-sequential graphs, a connected $2a$-sequential graph can be obtained by multiplying all the vertex labels by 2. Notice that the vertex labels in resulting graph belong to $\{0, 2, \ldots, 2q-2\}$ (or $\{0, 2, \ldots, 2q\}$ for trees) and the edges labels are from $2a$ to $2a+2q-2$. Moreover, a connected a-sequential graph can be obtained from a connected even $2a$-sequential graph with even vertex labels by dividing all the vertex labels by 2. Likewise, a $2a$-sequential labeling of a connected graph with odd vertex labels induces an a-sequential labeling of the graph by subtracting 1 from each vertex label and dividing by 2. Thus for connected graphs, a-sequential is equivalent to $2a$-sequential. They prove that if G is even $2a$-sequential the following graphs are properly even harmonious: $G \cup P_m^2$ for $m > 2$, $G \cup P_n$ for $n > 1$, $n \equiv 1, 2 \pmod 4$, $G \cup {C_m}^{+t}$ for some cases, $G \cup {Cat_m}^{+n}$ for $m > 1$, and $G \cup W_{2n+1}$.

For n and k odd and $m, n, k, t > 1$, Mbianda and Gallian (see [2460]) proved the following graphs have properly even harmonious labelings: mP_3 for even m; $2P_m \cup 2P_n \cup S_t$;

$2P_m \cup 2P_n \cup P_k$; $2P_m \cup 2P_n \cup C_k$; $2P_m \cup 2P_n \cup C_4$; $2P_m \cup 2P_n \cup 2K_4$; $2P_m \cup 2P_n \cup 2W_4$; $2P_m \cup 2P_n \cup 2C_k$; $F_n \cup K_4$ ($F_n = P_n + K_1$ is the fan); $F_n \cup 2K_4$; $F_n \cup W_4$; $F_n \cup 2W_4$; $W_n \cup K_4$; $W_n \cup 2K_4$; $W_n \cup W_4$; $W_n \cup 2W_4$; $(C_n + K_1) \cup K_4$; $(C_n + K_1) \cup W_4$; $(C_n + K_1) \cup 2K_4$; $(C_n + K_1) \cup 2W_4$; and $(C_n + \overline{K_2}) \cup K_4$ ($(C_n + \overline{K_2})$ is the double cone). Gallian [1172] proved the following graphs have properly even harmonious labelings (in all cases $m, n > 1$): mP_n for m even; $2P_m \cup 2P_n \cup 2C_3$; $2P_m \cup 2P_n \cup 2C_4$; $2P_m \cup 2P_n \cup C_3 \cup C_4$; $F_n \cup P_4$; $F_n \cup 2P_4$; $F_n \cup C_4$; and $F_n \cup 2C_4$.

Binthiya and Sarasija [661] prove the following graphs are even harmonious: $C_n \odot mK_1$ (n odd), $P_n \odot mK_1$ ($n > 1$ odd), $C_{2n}@K_2$, P_n (n even) with $n-1$ copies of $m\overline{K_1}$, the shadow graph $D_2(K_{1,n})$, the splitting graph $spl(K_{1,n})$, and the graph obtained from the P_n (n even) with $n-1$ copies of $\overline{K_m}$ incident with first $n-1$ vertices of P_n. Vargheese and Arun [4049] prove that the triangular books, the disjoint union of two triangular book graphs, total graphs $T(P_n)$, the disjoint union of $T(P_n)$ and a triangular book, and the graph obtained by joining the centers of two disjoint copies of $K_{1,n}$ to an isolated vertex are even harmonious.

Lasim, Halikin, and Wijaya [2092] showed how to build new harmonious, odd harmonious, even harmonious labelings based on the existing such labelings.

In [602] Beatress and Sarasija introduced the notion of even-odd harmonious graphs as follows. Let G be a graph with m vertices and n edges. An injective mapping from the vertices of G to $\{1, 3, 5, \ldots, 2m-1\}$ is called an *even-odd harmonious* labeling of G if the induced edge mapping f^* from the edges of G to $\{0, 2, 4, \ldots, 2(n-1)\}$ is a bijection and $f^*(uv) = (f(u)^* + f^*(v)) \bmod 2n$ for all edges uv. They proves that the bistars, cycles with one pendent edge, crowns, $K_{1,m,n}$, ${P_n}^2$ ($n \geq 4$), and nP_2 are even-odd harmonious graphs. Kalaimathi and Balamurugan [1808] proved caterpillars, lobsters, coconut trees, spider trees, and star graphs admit even-odd harmonious labelings. Zala, Chotaliya, and Chaurasiya [4302] proved H-graphs, combs, and bistars graph have even-odd harmonious labelings. Ganeshwari and Sudhana [1181] proved that paths, cycles, stars, bistars, combs, P_n^2, $C_3@pK_1$ and $C_{2r-1}@K_1$ are even-odd average harmonious graphs.

5 Magic-type Labelings

5.1 Magic Labelings

Motivated by the notion of magic squares in number theory, magic labelings were introduced by Sedláček [3333] in 1963.[2] Responding to a problem raised by Sedláček, Stewart [3697] and [3698] studied various ways to label the edges of a graph in the mid 1960s. Stewart calls a connected graph *semi-magic* if there is a labeling of the edges with integers such that for each vertex v the sum of the labels of all edges incident with v is the same for all v. (Berge [623] used the term "regularisable" for this notion.) A semi-magic labeling where the edges are labeled with distinct positive integers is called a *magic* labeling. Stewart calls a magic labeling *supermagic* if the set of edge labels consists of consecutive positive integers. The classic concept of an $n \times n$ magic square in number theory corresponds to a supermagic labeling of $K_{n,n}$. Stewart [3697] proved the following: K_n is magic for $n = 2$ and all $n \geq 5$; $K_{n,n}$ is magic for all $n \geq 3$; fans F_n are magic if and only if n is odd and $n \geq 3$; wheels W_n are magic for $n \geq 4$; and W_n with one spoke deleted is magic for $n = 4$ and for $n \geq 6$. Stewart [3697] also proved that $K_{m,n}$ is semi-magic if and only if $m = n$. In [3698] Stewart proved that K_n is supermagic for $n \geq 5$ if and only if $n > 5$ and $n \not\equiv 0 \pmod 4$. Sedláček [3334] showed that Möbius ladders M_n (see §2.3 for the definition) are supermagic when $n \geq 3$ and n is odd and that $C_n \times P_2$ is magic, but not supermagic, when $n \geq 4$ and n is even. Shiu, Lam, and Lee [3531] have proved: the composition of C_m and $\overline{K}_n$ (see §2.3 for the definition) is supermagic when $m \geq 3$ and $n \geq 2$; the complete m-partite graph $K_{n,n,\ldots,n}$ is supermagic when $n \geq 3$, $m > 5$ and $m \not\equiv 0 \pmod 4$; and if G is an r-regular supermagic graph, then so is the composition of G and $\overline{K}_n$ for $n \geq 3$. Ho and Lee [1443] showed that the composition of K_m and $\overline{K_n}$ is supermagic for $m = 3$ or 5 and $n = 2$ or n odd. Bača, Holländer, and Lih [373] have found two families of 4-regular supermagic graphs. Shiu, Lam, and Cheng [3528] proved that for $n \geq 2$, $mK_{n,n}$ is supermagic if and only if n is even or both m and n are odd.

In 2000 Ivančo [1551] gave a characterization of all supermagic regular complete multipartite graphs. He proved that Q_n is supermagic if and only if $n = 1$ or n is even and greater than 2 and that $C_n \times C_n$ and $C_{2m} \times C_{2n}$ are supermagic. In [1551] Ivančo completely determines the supermagic graphs that are the disjoint unions of complete k-partite graphs where every partite set has the same order.

Trenklér [3913] has proved that a connected magic graph with p vertices and q edges other than P_2 exits if and only if $5p/4 < q \leq p(p-1)/2$. In [3781] Sun, Guan, and Lee give an efficient algorithm for finding a magic labeling of a graph. In [4180] Wen, Lee, and Sun show how to construct a supermagic multigraph from a given graph G by adding extra edges to G. In [1237] Geethalakshmi, Saraswathi, and Divya showed that the graphs $K_{n,n}$ $(n \neq 2)$ admit supermagic labelings. [1237] new

Sudarsana, Suryanto, Lusianti, and Putri [3711] show how magic graph labelings can be used to increase the security level of encrypted text on social media. Angel Sherin

[2] A comprehensive expository treatment of magic labelings is given by Bača, Miller, Ryan, and Semaničová-Feňovčíkováin [410].

and Maheswari [242] and Ali Ahmed and Baskar Babujee [142] used magic labeling of graph to devise encryption and decryption schemes. In [2088] Lakshmi, Sudhakar, and Sudhakar provided a cryptographic technique for data encryption and decryption using magic labelings of graphs.

In [2022] Kovář provides a general technique for constructing supermagic labelings of copies of certain kinds of regular supermagic graphs. In particular, he proves: if G is a supermagic r-regular graph ($r \geq 3$) with a proper edge r coloring, then nG is supermagic when r is even and supermagic when r and n are odd; if G is a supermagic r-regular graph with m vertices and has a proper edge r coloring and H is a supermagic s-regular graph with n vertices and has a proper edge s coloring, then $G \times H$ is supermagic when r is even or n is odd and is supermagic when s or m is odd.

Kovář, Kravčenko, Krbeček, and Silber [2024] affirmatively answered a question by Madaras about existence of supermagic graphs with arbitrarily many different degrees. Their construction provided graphs with all degrees even. They asked if there exists a supermagic graph with d different odd degrees for any positive integer d. This question was answered affirmatively by Fronček and Qiu with a construction based on the use of 3-dimensional magic rectangles.

In [976] Drajnová, Ivančo, and Semaničová proved that the maximal number of edges in a supermagic graph of order n is 8 for $n = 5$ and $\frac{n(n-1)}{2}$ for $6 \leq n \not\equiv 0 \pmod 4$, and $\frac{n(n-1)}{2} - 1$ for $8 \leq n \equiv 0 \pmod 4$. They also establish some bounds for the minimal number of edges in a supermagic graph of order n. Ivančo, and Semaničová [1561] proved that every 3-regular triangle-free supermagic graph has an edge such that the graph obtained by contracting that edge is also supermagic and the graph obtained by contracting one of the edges joining the two n-cycles of $C_n \times K_2$ ($n \geq 3$) is supermagic.

Ivančo [1553] proved: the complement of a d-regular bipartite graph of order $8k$ is supermagic if and only if d is odd; the complement of a d-regular bipartite graph of order $2n$ where n is odd and d is even is supermagic if and only if $(n, d) \neq (3, 2)$; if G_1 and G_2 are disjoint d-regular Hamiltonian graphs of odd order and $d \geq 4$ and even, then the join $G_1 \oplus G_2$ is supermagic; and if G_1 is d-regular Hamiltonian graph of odd order n, G_2 is $d-2$-regular Hamiltonian graph of order n and $4 \leq d \equiv 0 \pmod 4$, then the join $G_1 \oplus G_2$ is supermagic.

For $k \geq 2$ and graphs G and H, the graph $G \odot^k H$ defined as $(G \odot^{k-1} H) \odot H$ (where $G \odot^1 H = G \odot H$) is called the *k-multilevel corona* of G with H. Marbun and Salman [2418] proved $(W_n \odot^{k-1}) \odot C_n$ is W_n-edge magic.

A connected, undirected, simple, non-trivial graph $G(V, E)$ with p vertices and q edges an injective function $f : V(G)$ to $\{s, s + d, \ldots, s + (q + 1)d\}$ such that there is an injective function $g : E(G)$ to $\{d, 2d, \ldots, 2(q - 1)d\}$ is said to be an (s, d) *magic labeling* if $f(u) + g(uv) + f(v)$ is a constant for all edges uv. A graph G is called an (s, d) *magic graph* if it has an (s, d) magic labeling. Sumathi and Mala [3771] established the existence of (s, d) magic labelings of coconut trees, regular bamboo trees, symmetrical trees, olive treese, spiders, and the subdivisions of these families.

In [637] Bezegová and Ivančo [639] extended the notion of supermagic regular graphs by defining a graph to be *degree-magic* if the edges can be labeled with $\{1, 2, \ldots, |E(G)|\}$

such that the sum of the labels of the edges incident with any vertex v is equal to $(1 + |E(G))/\deg(v)$. They used this notion to give some constructions of supermagic graphs and proved that for any graph G there is a supermagic regular graph which contains an induced subgraph isomorphic to G. In [639] they gave a characterization of complete tripartite degree-magic graphs and in [640] they provided some bounds on the number of edges in degree-magic graphs. They say a graph G is *conservative* if it admits an orientation and a labeling of the edges by $\{1, 2, \ldots, |E(G)|\}$ such that at each vertex the sum of the labels on the incoming edges is equal to the sum of the labels on the outgoing edges. In [638] Bezegová and Ivančo introduced some constructions of degree-magic labelings for a large family of graphs using conservative graphs. Using a connection between degree-magic labelings and supermagic labelings they also constructed supermagic labelings for the disjoint union of some regular non-isomorphic graphs. Among their results are: If G is a δ-regular graph where δ is even and at least 6, and each component of G is a complete multipartite graph of even size, then G is a supermagic graph; for any δ-regular supermagic graph H, the union of disjoint graphs H and G is supermagic; if G is a δ-regular graph with $\delta \equiv 0 \pmod 8$ and each component is a circulant graph, then G is a supermagic graph; for any δ-regular supermagic graph H, the union of disjoint graphs H and G is a supermagic graph; and that the complement of the union of disjoint cycles $C_{n_1}, \ldots, C_{n_k}$ is supermagic when $k \equiv 1 \pmod 4$ and $11 \leq n_i \equiv 3 \pmod 8$ for all $i = 1, \ldots, k$. In [1544] Inpoonjal gave necessary and sufficient conditions for the existence of degree-magic labelings of graphs obtained by taking the join and composition of complete tripartite graphs.

A graph G is said to be (F, H)-sim-(super)magic if there exists a bijection f' that is simultaneously F-(super) magic and H-(super) magic. In [314], Ashari, Salman, and Simanjuntak consider (K_2, H)-sim-(super) magic graphs where H is isomorphic to three classes of graphs with varied symmetry: a cycle which is symmetric (both vertex-transitive and edge-transitive), a star which is edge-transitive but not vertex-transitive, and a path which is neither vertex-transitive nor edge-transitive. They provide forbidden subgraphs for the existence of (K_2, H)-sim-(super) magic graphs and classify classes of (K_2, H)-sim-(super) magic graphs. They also derive sufficient conditions for edge-(super) magic graphs to be (K_2, H)-sim-(super) magic and utilize such conditions to characterize some (K_2, H)-sim-(super) magic graphs.

Let G' be a copy of a simple graph G and for each vertex v_i of G let u_i be the vertex of G' corresponding with v_i. The *double graph* has vertex set $V(G) \cup V(G')$ and edge set $E(G) \cup E(G') \cup \{u_i v_j \mid u_i \in V(G);\ v_j \in V(G')\ \text{ and } u_i u_j \in E(G)\}$. Ivančo [1554] establishes sufficient conditions for generalized double graphs to be degree-magic and constructs supermagic labelings of some graphs generalizing double graphs.

Sedláček [3334] proved that graphs obtained from an odd cycle with consecutive vertices $u_1, u_2, \ldots, u_m, u_{m+1}, v_m, \ldots, v_1$ $(m \geq 2)$ by joining each u_i to v_i and v_{i+1} and u_1 to v_{m+1}, u_m to v_1 and v_1 to v_{m+1} are magic. Trenklér and Vetchý [3916] have shown that if G has order at least 5, then G^n is magic for all $n \geq 3$ and G^2 is magic if and only if G is not P_5 and G does not have a 1-factor whose every edge is incident with an end-vertex of G. Avadayappan, Jeyanthi, and Vasuki [321] have shown that k-sequential trees are

magic (see §4.1 for the definition).

Seoud and Abdel Maqsoud [3366] proved that $K_{1,m,n}$ is magic for all m and n and that P_n^2 is magic for all n. However, Serverino has reported that P_n^2 is not magic for $n = 2, 3$, and 5 [1248]. Jeurissan [1648] characterized magic connected bipartite graphs. Ivančo [1552] proved that bipartite graphs with $p \geq 8$ vertices, equal sized partite sets, and minimum degree greater than p are magic. Bača [334] characterizes the structure of magic graphs that are formed by adding edges to a bipartite graph and proves that a regular connected magic graph of degree at least 3 remains magic if an arbitrary edge is deleted. In [3647] Solairaju and Arockiasamy prove that various families of subgraphs of grids $P_m \times P_n$ are magic.

In [1545] Inpoonjai investigated the existence of supermagic labelings of the disjoint union of three copies of certain graphs G. He provded an application of supermagic graphs to balanced logistics network designs. Dayanand and Ahmed [907] investigate supermagic properties of several classes of connected and disconnected graphs. They show that there can be arbitrarily large gaps among the possible valences for certain supermagic graphs. They also prove that the disjoint union of multiple copies of a supermagic linear forest is supermagic if the number of copies is odd and that the supermagic labeling is complementary edge antimagic as well. The *broom* $B_{n,t}$ is a graph obtained by attaching $n - t$ pendent edges to an end point vertex of the path P_t. [1545] new

Marimuthu, Raja, and Raja Durga [2433] prove that $B_{n,n-1}$ is E-super vertex magic if and only if $n \geq 3$ is odd and $B_{n,t}$ is not E-super vertex magic for $n \geq 4$ and $t \geq 3$. In [2668] Nemani and Joshi defined a new class of graph called the *cartoon flower* and showed that a E-super vertex magic labeling does not exist for the class of cartoon flower graphs. They also define the wounded cartoon flower graphs and establish some sufficient conditions for the graph not to be E-super vertex magic. They give examples of some wounded cartoon flowers that admit an E-super vertex magic labeling and some others that do not. In [2065] Kumar and Marimuthu proved that semi-regular bipartite graphs are not E-super vertex-magic and gave an upper bound for the maximum degree of an E-super vertex-magic graph. They also obtained upper and lower bounds of any vertex degree d of a E-super vertex-magic graph.

A Γ-supermagic labeling of a graph $G(V, E)$ with $|E| = k$ is a bijection from E to an Abelian group Γ of order k such that the sum of labels of all incident edges of every vertex $x \in V$ is equal to the same element $\mu \in \Gamma$. An existence of a Z_{2nm}-supermagic labeling of Cartesian product of two cycles, $C_n \times C_m$ for n odd was proved recently. This along with an earlier result by Ivanč [1551] proved the existence of a Z_{2nm}-supermagic labeling of $C_m \times C_n$ for every $m, n \geq 3$ and conjectured that such labeling is possible for all $C_m \times C_n$. In [1141] Fronček and McKeown present a simple unified labeling method that Fronček [1135] used to prove this conjecture for all m, n odd that not relatively prime. In [1142] Fronček, Paananen, Sorensen proved that the Cartesian product of any two odd cycles admits a Γ-supermagic labeling for all Abelian groups Γ of order $2mn$. In [1142] they proved an analogous result for the product of two even cycles, and showed that there is a Γ-supermagic labeling for all Abelian groups Γ of order $2mn$ for all even m, n greater then 2. The case of a product of an odd and even cycle, remains widely open with three

exceptions.

Fronček, McKeown, and McKeown [1140] gave Z_{2nm}-supermagic labelings of $C_n \times C_m$ for n odd. This, together with an earlier result by Ivančo, [1551] proved that a Z_{2nm}-supermagic labeling of $C_n \times C_m$ exists for all $n, m \geq 3$. In [1141] Fronček and McKeown presented a simple unified labeling that covers all cases for Z_{2nm}-supermagic labeling of $C_n \times C_m$. Fronček [1134] provided a $(K_4 - e)$-supermagic labeling of the triangular ladder TL_{2n} $(n \geq 2)$. [1134] new

Ponnappan, Nagaraj, and Prabakaran [2847] say a vertex magic labeling f of a graph $G(V, E)$ is an *odd vertex magic* if f maps V to $\{1, 3, 5, \ldots, 2|V| - 1\}$ and E to $\{1, 2, 3, 4, \ldots, |V| + |E|\} - \{1, 3, 5, \ldots, 2|V| - 1\}$ if $|E| \geq |V| - 1)$ and otherwise f maps E to $\{2, 4, 6, \ldots, 2|E|\}$ and V to $\{1, 2, 3, 4, \ldots, |V| + |E|\} - \{2, 4, 6, \ldots, 2|E|\}$. They prove that P_n $(n \geq 3)$, C_n and mC_3 are odd vertex magic if and only if n is odd, $(3, t)-$kites are vertex magic if and only if t is even, and $C_n \odot K_1$ are not odd vertex for all n.

A triplet $[H, \phi, t]$ is called a *supermagic frame* of G if ϕ is a homomorphism of H onto G and $t : E(H) \to \{1, 2, \ldots, |E(H)|\}$ is an injective mapping such that the sum of $t(uw)$ over all $u \in \phi^{-1}(v)$ is independent of the vertex $v \in V(G)$. In 2000, Ivančo proved that if there is a supermagic frame of a graph G, then G is supermagic. Singhun, Boonklurb, and Charnsamorn [3614] construct a supermagic frame of $m \geq 2$ copies of the Cartesian product of cycles and show that m copies of the Cartesian product of cycles is supermagic.

A *prime-magic labeling* is a magic labeling for which every label is a prime. Sedláček [3334] proved that the smallest magic constant for prime-magic labeling of $K_{3,3}$ is 53 while Bača and Holländer [369] showed that the smallest magic constant for a prime-magic labeling of $K_{4,4}$ is 114. Letting σ_n be the smallest natural number such that $n\sigma_n$ is equal to the sum of n^2 distinct prime numbers we have that the smallest magic constant for a prime-magic labeling of $K_{n,n}$ is σ_n. Bača and Hollaänder [369] conjecture that for $n \geq 5$, $K_{n,n}$ has a prime-magic labeling with magic constant σ_n. They proved the conjecture for $5 \leq n \leq 17$ and confirmed the conjecture for $n = 5$, 6 and 7.

Characterizations of regular magic graphs were given by Doob [971] and necessary and sufficient conditions for a graph to be magic were given in [1648], [1777], and [928]. Some sufficient conditions for a graph to be magic are given in [969], [3912], and [2564]. Bertault, Miller, Pé-Rosés, Feria-Puron, and Vaezpour [634] provided a heuristic algorithm for finding magic labelings for specific families of graphs. The notion of magic graphs was generalized in [970] and [3275].

Let $m, n, a_1, a_2, \ldots, a_m$ be positive integers where $1 \leq a_i \leq \lfloor n/2 \rfloor$ and the a_i are distinct. The *circulant graph* $C_n(a_1, a_2, \ldots, a_m)$ is the graph with vertex set $\{v_1, v_2, \ldots, v_m\}$ and edge set $\{v_i v_{i+a_j} \mid 1 \leq i \leq n,\ 1 \leq j \leq m\}$ where addition of indices is done modulo n. In [3355] Semaničová characterizes magic circulant graphs and 3-regular supermagic circulant graphs. In particular, if $G = C_n(a_1, a_2, \ldots, a_m)$ has degree r at least 3 and $d = \gcd(a_1, n/2)$ then G is magic if and only if $r = 3$ and $n/d \equiv 2 \pmod 4$, $a_1/d \equiv 1 \pmod 2$, or $r \geq 4$ (a necessary condition for $C_n(a_1, a_2, \ldots, a_m)$ to be 3-regular is that n is even). In the 3-regular case, $C_n(a_1, n/2)$ is supermagic if and only $n/d \equiv 2 \pmod 4$, $a_1/d \equiv 1 \pmod 2$ and $d \equiv 1 \pmod 2$. Semaničová also notes that a bipartite graph that is decomposable into an even number of Hamilton cycles is supermagic. As a corollary

she obtains that $C_n(a_1, a_2, \ldots, a_{2k})$ is supermagic in the case that n is even, every a_i is odd, and $\gcd(a_{2j-1}, a_{2j}, n) = 1$ for $i = 1, 2, \ldots, 2k$ and $j = 1, 2, \ldots, k$.

Ivančo, Kovář, and Semaničová-Feňovčková [1557] characterize all pairs n and r for which an r-regular supermagic graph of order n exists. They prove that for positive integers r and n with $n \geq r + 1$ there exists an r-regular supermagic graph of order n if and only if one of the following statements holds: $r = 1$ and $n = 2$; $3 \leq r \equiv 1 \pmod 2$ and $n \equiv 2 \pmod 4$; and $4 \leq r \equiv 0 \pmod 2$ and $n > 5$. The proof of the main result is based on finding supermagic labelings of circulant graphs. The authors construct supermagic labelings of several circulant graphs.

Trenklér [3914] extended the definition of supermagic graphs to include hypergraphs and proved that the complete k-uniform n-partite hypergraph is supermagic if $n \neq 2$ or 6 and $k \geq 2$ (see also [3915]). In [3742] Sugiyama gave a generalized definition of magic graphs, for which any number of digits can be used to label a vertex and edge, and described the construction of such magic graphs and their properties. He determined the minimum and maximum magic sums for regular graphs, including polygons and polyhedrons, and provided techniques for transforming and synthesizing magic graphs using an affine transform.

For connected graphs of size at least 5, Ivančo, Lastivkova, and Semaničová [1558] provide a forbidden subgraph characterization of the line graphs that can be magic. As a corollary they obtain that the line graph of every connected graph with minimum degree at least 3 is magic. They also prove that the line graph of every bipartite regular graph of degree at least 3 is supermagic.

For any non-trivial Abelian group A under addition, a graph G is said to be *strong A-magic* if there exists a labeling f of the edges of G with non-zero elements of A such that the vertex labeling f^+ defined as $f^+(v) = \sum f(uv)$ taken over all edges uv incident at v is a constant, and the constant is same for all possible values of $|V(G)|$. Stella Arputha Mary, Navaneethakrishnan, and Nagarajan [3696] provide strong Z_4-magic labelings for various graphs and strong Z_{4p}-magic labelings for those graphs.

In [3150] Razzaq, Rizvi, and Ali introduce the concept of an *H-group magic total* labeling of a graph G over a finite Abelian group A as a bijection $\lambda : V(G) \cup E(G) \to A$ such that for any subgraph $H'(V', E')$ of G isomorphic to H, the sum $\sum_{v \in V'} \lambda(v) + \sum_{e \in E'} \lambda(e)$ is equal to magic constant k'. A graph is called *H-group magic* if it admits an H-group magic total labeling. They determine the H-group magic total labelings of fan graphs over the finite Abelian group $A \cong \mathbb{Z}_3 \times \mathbb{Z}_t$, where $t \geq 3$ and show that disjoint union of isomorphic as well as non-isomorphic copies of fan graphs are H-group magic over $A \cong \mathbb{Z}_3 \times \mathbb{Z}_t$.

For a graph $G(V, E)$ and a set of positive integers S with $|S| = |V|$, Godinho, Singh, and Arumugam [1284] say G is *S-magic* if there exists a bijection $\phi : V \to S$ such that $\sum \phi(u)$ over all $u \in N(v)$ is a constant k for all $v \in V$. They proved that if G is S-magic, then the corresponding magic constant is unique. They proved that several families of graphs are S-magic and that several families are not S-magic. They also determined the set of all S-magic constants for certain classes of graphs for different label sets S.

For a natural number h, Salehi [3244] defines a graph G to be h-magic if there is a

labeling α from the edges of G to the nonzero integers in Z_h such that for each vertex v in G the sum of all α values of edges incident to v is a constant (called the *magic sum index*) that is independent of the choice of v. If the constant is 0, G is called a *zero-sum h-magic graph*. The *null set* of graph G is the set of all natural numbers h for which G admits a zero-sum h-magic labeling. In [3244] Salehi determines the null sets for K_n, $K_{m,n}$, C_n, books, and cycles with a P_k chord. Lin and Wang [2268] determine the null sets of generalized wheels and generalized fans, and construct infinitely many examples of Z_h-magic graphs with magic sum zero and present some open problems.

In 2020, Kamatchi, Paramasivam, Prajeesh, Sabeel, and Arumugam [1817] introduced the notion group vertex magic graphs as follows. For a simple undirected graph G and an additive Abelian group A with identity 0, a mapping f from the vertices of G to the nonzero elements of A is said to be an *A-vertex magic* labeling of G if there exists a μ in A such that $w(v) = \sum f(u)$ taken over all $u \in N(v)$ is μ for all vertices v of G. If G admits such a labeling it is called an *A-vertex magic graph*. If G is an A-vertex magic for every non-trivial Abelian group A, G is called a *group vertex magic* graph. They obtained some necessary conditions for a graph to be group vertex magic and gave a characterization of trees with diameter at most 4 that are $Z_2 \times Z_2$-vertex magic. In [1910] Karthik, Sabeel, and Paramasivama provided some necessary conditions for the group vertex magicness of graphs with at least one pendant edge and for the group vertex magicness of product graphs. They also characterized group vertex magicness of trees of diameter up to 5 for all infinite Abelian groups with finitely many elements of finite order. Sabeel, Paramasivam, Prajeesh, Kamatchi, and Arumugam [3223] characterized the $Z_2 \times Z_2$-vertex magicness of any tree with diameter 5. They further characterized A-vertex magic trees of diameter at most 5 for any finite Abelian group A and proved that A-vertex magic graphs do not possess any forbidden structures. They gave a method for constructing larger A-vertex magic graphs from the existing ones. Balamoorthy, Bharanedhar, and Kamatchi [449] obtained various results about the A-vertex magicness for graphs formed using joins, tensor products, and lexicographic products of graphs. Kavitha and Stella Arputha Mary [1934] say a vertex magic labeling of V_4 ($Z_2 \times Z_2$) is a *hefty* V_4-magic if its magic number is 1. They proved regular graphs, K_n ($n \geq 3$), vertex transitive graphs, and $C_m \times C_n$ are hefty V_4-magic graphs and connected acyclic graphs are not V_4-magic graphs. Libeeshkumar [2261] provided results regarding edge induced V_4-magic labelings for the subdivision graphs of $P_n, C_n, K_n, K_{1,n}$, bi-stars, suns, flags, jelly fish graphs, triangular snakes, wheels, and combs. Mary, Krishnan, and Nagarajan [2447] proved that triangular snakes, books, the one-point union of the apexes of t fans, and the splitting graph of paths, are Z_{4p}-magic graphs. In [2259] Liao and Liu provided characterizations of unicyclic graphs with diameter at most 4 that are A-vertex magic and a characterization of bicyclic graphs of diameter 3 that are group vertex magic. In [1958] Khuluq, Krisnawati, and Hidayat determined Z_k-vertex-magic labelings of paths, complete graphs, cycles, and stars. The *prime* graph over a ring R, denoted by $PG(R)$, is a graph with $V(PG(R)) = R$ such that uv is an edge if and only if $uRv = \{0_R\}$ or $vRu = \{0_R\}$, for every vertex $u \neq v$. In [1959] Khuluq, Krisnawati, and Hidayat investigated Z_k-vertex-magic labelings of prime graphs over Z_n for $n = p, n = p^2$, and $n = pq$ for distinct primes p and q. [1959] new

In 1976 Sedláček [3334] defined a connected graph with at least two edges to be *pseudo-magic* if there exists a real-valued function on the edges with the property that distinct edges have distinct values and the sum of the values assigned to all the edges incident to any vertex is the same for all vertices. Sedláček proved that when $n \geq 4$ and n is even, the Möbius ladder M_n is not pseudo-magic and when $m \geq 3$ and m is odd, $C_m \times P_2$ is not pseudo-magic.

A *vertex magic total* labeling of a graph with p vertices and q edges is a bijection from the union of the vertex set and edge set to the consecutive integers $1, 2, \ldots, p+q$ with the property that for every vertex u, the sum of the label of u and the labels of the edges incident with u is a constant k. A vertex magic total labeling is said to be *a-vertex multiple magic* if the set of the labels of the vertices is $\{a, 2a, \ldots, na\}$ and is *b-edge multiple magic b-edge multiple magic* if the set of labels of the edges is $\{b, 2b, \ldots, mb\}$. Nagaraj, Ponnappan, and Prabakaran [2641] provide properties of a-vertex multiple magic graphs and b-edge multiple magic graphs. In [4315] Zhang and Wang verify the existence of E-super vertex magic total labeling for odd regular graphs containing a particular 3-factor. Listiana, Darmaji, and Slamin [2276] investigated the existence of vertex magic total labelings of directed sun graphs $S_n = C_n \odot K_1$ and mS_n. In [2007] Krishnappa, Srinath, and Manjunath provided a vertex magic total labeling of complete graphs as a method for public-key cryptosystem that adds one more layer of security.

A vertex magic total labeling is said to be a *V-super vertex magic* labeling if $f(V(G)) = \{1, 2, 3, \ldots, |V|\}$. A graph G is called *V-super vertex magic* if it admits a V-super vertex magic labeling. In [4113] Vimal Kumar and Vijayalakshmi establish the V-super vertex magic labelings of some classes of parity graphs (that is, for every two induced paths between the same two vertices both paths have odd length, or both have even length). In [1006] Elmorsy introduced a labeling matrix method for creating super vertex magic total labelings for cycles and complete graphs. The method constructs new super vertex magic total labelings of graphs from exiting ones.

The notion of vertex N-magic total labeling was introduced by Marimuthu and Kumar in [2431] as follows. For a simple, finite, undirected graph G a *vertex N-magic total labeling* of G is a bijection from $V(G) \cup E(G)$ onto $\{1, 2, \ldots, |V| + |E|\}$ with the property that the number of distinct degrees N is equal to the number of distinct magic constants k_i for $i \in \{1, 2, \ldots, N\}$, and the magic constants k_i must be in a strictly increasing order $k_1 < k_2 < \cdots < k_i$ for $i \in \{1, 2, \ldots, N\}$. Thay gave some properties of even vertex N-magic total labeling of non-regular graphs and even vertex N-magic total labelings of kites, products of cycles, and generalized Petersen graphs. They also gave a necessary and sufficient condition for a graph to be an even vertex magic total graph. In [2686] Nishanthini and Jeyabalan provided sharp bounds of vertex N-magic total labeling graphs in terms of their magic constants and gave a necessary condition for a vertex N-magic total labeling of trees. In [2687] they discussed applications of vertex N-magic total labelings of graphs to disease transmission networks. Nishanthini [2685] investigated the existence of odd vertex N-magic total labelings of paths, stars, suns, banana trees, and kites. He further provided practical application of vertex N-magic labeling for modeling employee skills and extracurricular activity networks, where a graph-based approach is

[2431] new

[2686] new

[2687] new

[2685] new

employed to examine the relationships and capabilities of employees within replicated corporate settings. Nishanthini and Jeyabalan [2685]investigated the existence of certain types of vertex N-magic total labelings of graphs that are a cycle where each vertex of the cycle is the center of a star and special cases of lobsters. [2685] new

In [2689] Nishanthini and Jeyabalan proved the following families admit vertex N-magic total labelings using two magic constants: P_n $(n \geq 3)$, $K_{1,n}$ $(n \geq 2)$, bi-stars $B_{n,n}$ $(n \geq 2)$, sun graphs C_n^+ $(n \geq 3)$, and wheels W_n $(n \geq 4)$. They further prove that all trees except H-trees admit vertex N-magic total labelings, fans $F_{1,n}$ $(n \geq 4)$ admit vertex N-magic total labelings using 3 magic constants, triangular snakes do not admit vertex N-magic total labelings, and H-tree graphs with three distinct degrees do not admit vertex N-magic total labelings. They conclude with applications of vertex N-magic labelings to radio station coverage. Nishanthini [2685] proved the following families of graphs admit vertex N-magic total labelings: Jahangir graphs (2 vertices connected to a common vertex, plus n pendant edges), octopus graphs (a central vertex connected to n paths of length 2), cycle-star graphs (a cycle of length 3 with n pendant edges attached to one vertex). He applied vertex N-magic labeling to chemical reaction networks to ensure stoichiometric balance and conservation of mass, charge, and energy. Furthermore, he applied this framework to water pollution analysis. [2689] new [2685] new

Kong, Lee, and Sun [2009] used the term "magic labeling" for a labeling of the edges with nonnegative integers such that for each vertex v the sum of the labels of all edges incident with v is the same for all v. In particular, the edge labels need not be distinct. They let $M(G)$ denote the set of all such labelings of G. For any L in $M(G)$, they let $s(L) = \max\{L(e) \mid e \in E\}$ and define the *magic strength* of G as $m(G) = \min\{s(L) \mid L \in M(G)\}$. To distinguish these notions from others with the same names and notation, which we will introduced in the next section for labelings from the set of vertices and edges, we call the Kong, Lee, and Sun version the *edge magic strength* and use $em(G)$ for $\min\{s(L)\colon L \text{ in } M(G)\}$ instead of $m(G)$. Kong, Lee, and Sun [2009] use $DS(k)$ to denote the graph obtained by taking two copies of $K_{1,k}$ and connecting the k pairs of corresponding leafs. They show: for $k > 1$, $em(DS(k)) = k - 1$; $em(P_k + K_1) = 1$ for $k = 1$ or 2, $em(P_k + K_1) = k$ if k is even and greater than 2, and 0 if k is odd and greater than 1; for $k \geq 3$, $em(W(k)) = k/2$ if k is even and $em(W(k)) = (k-1)/2$ if k is odd; $em(P_2 \times P_2) = 1$, $em(P_2 \times P_n) = 2$ if $n > 3$, $em(P_m \times P_n) = 3$ if m or n is even and greater than 2; $em(C_3^{(n)}) = 1$ if $n = 1$ (Dutch windmill, – see §2.4), and $em(C_3^{(n)}) = 2n - 1$ if $n > 1$. They also prove that if G and H are magic graphs then $G \times H$ is magic and $em(G \times H) = \max\{em(G), em(H)\}$ and that every connected graph is an induced subgraph of a magic graph (see also [1037] and [1087]). They conjecture that almost all connected graphs are not magic.

In [1483] Ichishima, López, Muntaner-Batle, and Takahashi introduce the parameter $l(n)$ as the minimum size of a graph G of order n for which all graphs of order n and size at least $l(n)$ have $\mu_s(G) = +\infty$, and provide lower and upper bounds for $l(G)$. Imran, Baig, and Feňovčiková, [1522] established that for $n = 0 \pmod 4$, $\mu_s(C_n \times K_2) \leq 3n/2 - 1$. Ichishima, López, Muntaner-Batle, and Takahashi, improve this bound by showing that $\mu_s(n) + 1$ when $n \geq 4$ is even. Enomoto, Lladó, Nakamigawa, and Ringel [1037] posed the

conjecture that every nontrivial tree is super edge-magic. They propose a new approach to attack this conjecture. They believe that their approach may also help to resolve the conjecture by Graham and Sloane that every nontrivial tree is harmonious [1327]. Huang, Hanif, Siddiqui, and Nadeem, [1469] showed the super edge-magicness of certain types of generalized combs and disjoint unions of generalized combs and stars. Cerioli, Fernandes, Lee, Lintzmayer [755] proved certain forests of stars admit a super edge-magic labeling and that certain forests of caterpillars admit an edge-magic labeling. Ichishima, Muntaner-Batle, Oshima, and Takahashi [1503] investigated the (super) edge-magic properties of disjoint unions of cycles and stars. The super edge-magic results they obtained yield felicitous labelings for the same class of graphs. They propose a conjecture and several open problems and settle an old problem proposed by Wallis in his book "Magic Graphs" [4120].

[1503] new

Recall a *lexicographic product* of two graphs G_1 and G_2 is a graph that arises from G_1 by replacing each vertex of G_1 by a copy of the G_2 and each edge of G_1 with $K_{n,n}$ where n is the order of G_2. Sun and Lee [3782] show that the Cartesian, conjunctive, normal, lexicographic, and disjunctive products of two magic graphs are magic and the sum of two magic graphs is magic. They also determine the edge magic strengths of the products and sums in terms of the edge magic strengths of the components graphs. In [2187] Lee, Saba, and Sun show that the edge magic strength of P_n^k is 0 when k and n are both odd.

In 1970 Kotzig and Rosa [2017] introduced the concepts of edge-magic graphs and edge-magic labelings as follows. A graph $G(V, E)$ with p vertices and q edges is *edge-magic* if there is a bijective function $f : V \cup E \to \{1, 2, \ldots, p + q\}$ and a constant k such that the $f(x) + f(xy) + f(y) = k$ for all edges xy. Such a function is called an *edge-magic labeling* of G. An edge-magic labeling f of a (p, q)-graph G with the extra property that $f(V(G)) = \{1, 2, \ldots, p\}$ is said to be a *super edge-magic labeling* of G. Super edge-magic graphs of equal order and size have been used in the study of Skolem and Langford type sequences [1482], dual shuffle primes, and the Jacobsthal numbers [2320]. Such graphs were investigated by López, Muntaner-Batle, and Prabu in [2310]. They use L to denote the loop graph and $LK_{1,n}$ to denote the graph obtained from $K_{1,n}$ by attaching a loop to the central vertex. They proved $2LK_{1,1} \cup LK_{1,n}, 2LK_{1,m} \cup LK_{1,n}, (2s + 1)LK_{1,n}$, and $LK_{1,m} \cup 2LK_{1,n} \cup (2s)LK_{1,1}$ are super edge-magic. They also proved $LK_{1,m} \cup LK_{1,n}, L \cup LK_{1,n}$, and $2L \cup LK_{1,n}$ are not super edge-magic. In [2561] Mujtahid, Salman, Hidayat, Qausar, and Zahara proved certain θ-graphs admit super edge-magic total labelings.

For a simple graph $G(V, E)$ a bijection f from $V \cup E$ to $\{1, 2, \ldots, |V| + |E|\}$ is said to be *edge-magic total labeling* of G, if there exists an integer k such that $f(u) + f(uv) + f(v) = k$ for every edge $uv \in E$. If, in addition, $f(V) = \{1, 2, \ldots, |V|\}$, f is said to be an *super edge-magic total labeling*. Deeothi [918] investigated the super edge-magic total strength of the family of unicyclic graphs having an odd cycle a varying number of pendant vertices adjacent to each vertex. In [4316] Zhang, Zhou, and Wang provided a new algorithm to determine whether a graph has a non-edge magic total labeling graph. Using their algorithm, all non-edge magic total graphs with up to 9 vertices are determined. They also proved that the joins of K_n with each of paths, cycles, and fans are non-edge-magic total labeling graphs when certain conditions are satisfied.

The following three definitions are taken form [916]. For the prism $C_n \times P_2$ denote the n consecutive vertices of one copy of C_n by $u_0, u_1, \ldots, u_{n-1}$ and the vertices a second copy by $v_0, v_1, \ldots, v_{n-1}$, with $u_i v_i$ as the n edges of P_2. A *triangulated prism,* denoted by $T\Pi_n$, is the graph obtained from $C_m \times P_2$ adding the n the edges $v_i u_{i+1}$ for $i = 0, 1, \ldots, n_1$, where $u_n = u_0$. For the wheel $C_n \times K_1$, denote the n consecutive vertices of C_n by $u_0, u_1, \ldots, u_{n-1}$ and the center by x. A *triangulated wheel*, denoted, TW_n, is the graph otained from the wheel $C_n \times K_1$ by adding n new vertces $w_0, w_1, \ldots w_{n-1}$ and the $3n$ edges $u_i w_i, u_{i+1}$, and $w_i x$. (The four vertices u_i, u_{i+1}, x, w_i form the wheel W_3 with w_i as its center.) A *double vertex wheel*, denoted DW_n, is the graph otained from the wheel $C_n \times K_1$ by adding two new vertices x_1, x_2 and the edges $u_i x_1$ and $u_i x_2$ for all i. El Deen, Hanafy, Shehda, and Omar [916] proved that triangulated prisms admit super bimagic total labeling with bimagic total labeling using the bimagic numbers $k_1 = 6r$ and $k_2 = 8r$. They also showed that triangulated wheels, the double vertex wheels graph, and the closed triangulated water wheels (see [916] for the definition) admit trimagic total labelings. In [1593] Jayasekaran and Flower proved that umbrellas, dumbells, and circular ladders admit edge trimagic total labelings. In [1594] they prove that the graphs (n, l)-dragon, Mobius ladders, and books admit edge trimagic total labelings. Edgeb-imagic total labelings and edge trimagic total labelings are said to be a *super* if the vertices in the graph have label positive integers that is smaller than edges label. Regees and Jayasekaran [3159] [3158] proved that generalized prisms and webs admit super edge trimagic total labeling. They also gave super edge trimagic total labelings of the squares of cycles and gears. They further also showed that some cases of these graphs admit super edge trimagic labelings in [3157]. [916] new [916] new [916] new [1593] new [1594] new [3159] new [3158] new [3157] new

A (p, q) graph $G(V, E)$ is called an *edge trimagic graceful* if there exists a bijection $f : V \cup E$ to $\{1, 2, 3, \ldots, p+q\}$ such that for each edge xy in E, $|f(x)–f(xy)+f(y)| = C_1, C_2$, or C_3, where C_1, C_2, and C_3 are constants. Ezhil, Regees, and Mary [1061] proved that bistars, double stars, triple stars are edge trimagic graceful graphs. [1061] new

For a connected graph G_1 with p_1 vertices and q_1 edges and a connected G_2 with p_2 vertices and q_2 edges, Baskar Babujee and Suresh [586] define $G_1 \hat{o} G_2$ to be the graph obtained by superimposing any particular vertex of G_2 on any particular vertex of G_1. They say a total edge-magic labeling f of a graph $G(V, E)$ is a *super edge edge-magic* graph, if $f(E) = \{1, 2, \ldots, |E|\}$. They proved the following: if G has a super edge edge-magic total labeling then, $G\hat{o}P_n$ and $G\hat{o}(P_n \odot K_1)$ have super edge-bimagic total labelings and if G has super edge edge-magic labeling then, $G\hat{o}K_{1,n}$ and $G + K_1$ have total edge-bimagic labelings. They also showed that if G_1 and G_2 have super edge edge-magic labelings then, $G_1 \hat{o} G_2$ has an edge-bimagic total labeling.

A *Halin graph* is a planar 3-connected graph that consists of a tree and a cycle connecting the end vertices of the tree. Let G be a (p, q)-graph in which the edges are labeled $k, k+1, \ldots, k+q-1$, where $k \geq 0$. In [2205] Lee, Su, and Wang define a graph with p vertices to be *k-edge-magic* for every vertex v the sum of the labels of the incident edges at v are constant modulo p. They investigate some classes of Halin graphs that are k-edge-magic. Lee, Su, and Wang [2207] investigated some classes of cubic graphs that are k-edge-magic and provided a counterexample to a conjecture that any cubic

graph of order $p \equiv 2 \pmod 4$ is k-edge-magic for all k. Shiu and Lau [3535] gave some necessary conditions for families of wheels with certain spokes missing to admit k-edge-magic labelings.

Lau, Alikhani, Lee, and Kocay [2096] (see also [191]) show that maximal outerplanar graphs of orders $p = 4, 5, 7$ are k-edge magic if and only if $k \equiv 2 \pmod p$ and determined all maximal outerplanar graphs that are k-edge magic for $k = 3$ and 4. They also characterize all $(p, p-h)$-graphs that are k-edge magic for $h \geq 0$ and conjecture that a maximal outerplanar graph of prime order p is k-edge magic if and only if $k \equiv 2 \pmod p$.

S. M. Lee and colleagues [2226] and [2159] call a graph G *k-magic* if there is a labeling from the edges of G to the set $\{1, 2, \ldots, k-1\}$ such that for each vertex v of G the sum of all edges incident with v is a constant independent of v. The set of all k for which G is k-magic is denoted by $\mathrm{IM}(G)$ and called the *integer-magic spectrum* of G. In [2226] Lee and Wong investigate the integer-magic spectrum of powers of paths. They prove: $\mathrm{IM}(P_4^2)$ is $\{4, 6, 8, 10, \ldots\}$; for $n > 5$, $\mathrm{IM}(P_n^2)$ is the set of all positive integers except 2; for all odd $d > 1$, $\mathrm{IM}(P_{2d}^d)$ is the set of all positive integers except 1; $\mathrm{IM}(P_4^3)$ is the set of all positive integers; for all odd $n \geq 5$, $\mathrm{IM}(P_n^3)$ is the set of all positive integers except 1 and 2; and for all even $n \geq 6$, $\mathrm{IM}(P_n^3)$ is the set of all positive integers except 2. For $k > 3$ they conjecture: $\mathrm{IM}(P_n^k)$ is the set of all positive integers when $n = k + 1$; the set of all positive integers except 1 and 2 when n and k are odd and $n \geq k$; the set of all positive integers except 1 and 2 when n and k are even and $k \geq n/2$; the set of all positive integers except 2 when n is even and k is odd and $n \geq k$; and the set of all positive integers except 2 when n and k are even and $k \leq n/2$. In [2203] Lee, Su, and Wang showed that besides the natural numbers there are two types of the integer-magic spectra of honeycomb graphs. Fu, Jhuang and Lin [1155] determine the integer-magic spectra of graphs obtained from attaching a path of length at least 2 to the end vertices of each edge of a cycle. Navas, Ajitha, and Varkey [2659] the determined the vertex integer magic spectrum of caterpillars, super caterpillars, and extended super caterpillars.

In [2159] Lee, Lee, Sun, and Wen investigated the integer-magic spectrum of various graphs such as stars, double stars (trees obtained by joining the centers of two disjoint stars $K_{1,m}$ and $K_{1,n}$ with an edge), wheels, and fans. In [3247] Salehi and Bennett report that a number of the results of Lee et al. are incorrect and provide a detailed accounting of these errors as well as determine the integer-magic spectra of caterpillars. Shiu and Low [3553] determined the integer-magic spectra and null sets of the Cartesian product of two trees.

Lee, Lee, Sun, and Wen [2159] use the notation $C_m@C_n$ to denote the graph obtained by starting with C_m and attaching paths P_n to C_m by identifying the endpoints of the paths with each successive pairs of vertices of C_m. They prove that $\mathrm{IM}(C_m@C_n)$ is the set of all positive integers if m or n is even and $\mathrm{IM}(C_m@C_n)$ is the set of all even positive integers if m and n are odd.

Lee, Valdés, and Ho [2214] investigate the integer magic spectrum for special kinds of trees. For a given tree T they define the *double tree DT* of T as the graph obtained by creating a second copy T^* of T and joining each end vertex of T to its corresponding vertex in T^*. They prove that for any tree T, $\mathrm{IM}(DT)$ contains every positive integer

with the possible exception of 2 and IM(DT) contains all positive integers if and only if the degree of every vertex that is not an end vertex is even. For a given tree T they define ADT, the *abbreviated double tree of* T, as the graph obtained from DT by identifying the end vertices of T and T^*. They prove that for every tree T, IM(ADT) contains every positive integer with the possible exceptions of 1 and 2 and IM(ADT) contains all positive integers if and only if T is a path.

Lee, Salehi, and Sun [2189] have investigated the integer-magic spectra of trees with diameter at most four. Among their findings are: if $n \geq 3$ and the prime power factorization of $n-1 = p_1^{r_1}p_2^{r_2}\cdots p_k^{r_k}$, then $\mathrm{IM}(K_{1,n}) = p_1\mathbb{N} \cup p_2\mathbb{N} \cup \cdots \cup p_k\mathbb{N}$ (here $p_i\mathbb{N}$ means all positive integer multiples of p_i); for $m, n \geq 3$, the double star IM($DS(m,m)$) (that is, stars $K_{m,1}$ and $K_{n,1}$ that have an edge in common) is the set of all natural numbers excluding all divisors of $m-2$ greater than 1; if the prime power factorization of $m-n = p_1^{r_1}p_2^{r_2}\cdots p_k^{r_k}$ and the prime power factorization of $n-2 = p_1^{s_1}p_2^{s_2}\cdots p_k^{s_k}$, (the exponents are permitted to be 0) then $\mathrm{IM}(DS(m,n)) = A_1 \cup A_2 \cup \cdots \cup A_k$ where $A_i = p_i^{1+s_i}\mathbb{N}$ if $r_i > s_i \geq 0$ and $A_i = \emptyset$ if $s_i \geq r_i \geq 0$; for $m, n \geq 3$, $\mathrm{IM}(DS(m,n)) = \emptyset$ if and only if $m-n$ divides $n-2$; if $m, n \geq 3$ and $|m-n| = 1$, then $DS(m,n)$ is not magic. Lee and Salehi [2188] give formulas for the integer-magic spectra of trees of diameter four but they are too complicated to include here. More specialized results about the integer-magic spectra of amalgamations of stars and cycles are given by Lee and Salehi in [2188].

For a graph $G(V,E)$ and a function f from the V to the positive integers, Salehi and Lee [3251] define the *functional extension of* G *by* f, as the graph H with $V(H) = \cup\{u_i|\ u \in V(G) \text{ and } i = 1, 2, \ldots, f(u)\}$ and $E(H) = \cup\{u_iu_j|\ uv \in E(G), i = 1, 2, \ldots, f(u); j = 1, 2, \ldots, f(v)\}$. They determine the integer-magic spectra for P_2, P_3, and P_4.

Hungund and Akka [1471] introduce new concepts of reverse super edge-magic labeling and reverse super edge-magic strength of a graph G as follows. A *reverse edge magic* (REM) labeling of a graph $G(V,E)$ with p vertices and q edges is a bijection $f : V(G) \cup E(G) \to \{1, 2, \ldots, p+q\}$ such that $k = f(uv) - (f(u) + f(v))$ is a constant k for any edge $uv \in E(G)$. A REM labeling f is called *reverse super edge magic* (RSEM) labeling if $f(V(G)) = \{1, 2, 3, \ldots, p\}$ and $f(E(G)) = \{p+1, p+2, p+3, \ldots, p+q\}$. They obtained the reverse super edge-magic labelings and reverse super edge-magic strength for y-trees, odd cycles, generalized Petersen graphs, and $(2m+1)C_3$. Reddy and Sharief Basha [3151] find some new classes of RSEM labeling and investigate the connection between the RSEM labeling and different classes of labelings.

In a *distance difference magic* labeling, each vertex is given a unique value from one up to the number of vertices such that for each vertex the sums of the labels of vertices in the in-neighborhood minus the sums of the labels of vertices in the out-neighborhood equals zero. In [37] Aceska, Arcila-Maya, Carlson, Marr, Parnes, Ryan, Schuerger, and Vasquez showed a connected difference distance magic oriented graph on n vertices exists for $n \geq 5$. They also constructed arbitrarily large difference distance magic oriented graphs from smaller ones using a new graph sum and exhibit a connection between linear

[37] new

algebra and this type of labeling. In [38] Aeska, Mckailyn, and Ripperger provided a method to attach oriented cycles to an existing difference distance magic oriented graph thereby creating new ones with a larger number of vertices. [38] new

Table 5 summarizes the state of knowledge about magic-type labelings. In the table, **SM** means semi-magic, **M** means magic, and **SPM** means supermagic. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The table was prepared by Petr Kovář and Tereza Kovářová.

Table 5: **Summary of Magic Labelings**

Graph	*Types*	*Notes*
K_n	M	if $n = 2,\ n \geq 5$ [3697]
	SPM	for $n \geq 5$ iff $n > 5$
		$n \not\equiv 0 \pmod 4$ [3698]
$K_{m,n}$	SM	if $n \geq 3$ [3697]
$K_{n,n}$	M	if $n \geq 3$ [3697]
fans f_n	M	iff n is odd, $n \geq 3$ [3697]
	not SM	if $n \geq 2$ [1248]
wheels W_n	M	if $n \geq 4$ [3697]
	SM	if $n = 5$ or 6 [1248]
wheels with one spoke deleted	M	if $n = 4$, $n \geq 6$ [3697]
null graph with n vertices		
Möbius ladders M_n	SPM	if $n \geq 3$, n is odd [3334]
$C_n \times P_2$	not SPM	for $n \geq 4$, n even [3334]
$C_m[\overline{K}_n]$	SPM	if $m \geq 3$, $n \geq 2$ [3531]
$K_{\underbrace{n, n, \ldots, n}_{p}}$	SPM	$n \geq 3$, $p > 5$ and
		$p \not\equiv 0 \pmod 4$ [3531]
composition of r-regular SPM graph and $\overline{K}_n$	SPM	if $n \geq 3$ [3531]
$K_k[\overline{K}_n]$	SPM	if $k = 3$ or 5, $n = 2$ or n odd [1443]
$mK_{n,n}$	SPM	for $n \geq 2$ iff n is even or both n and m are odd [3528]
Q_n	SPM	iff $n = 1$ or $n > 2$ even [1551]

Continued on next page

Table 5 – *Continued from previous page*

Graph	*Types*	*Notes*
$C_m \times C_n$	SPM	$m = n$ or m and n are even [1551]
$C_m \times C_n$	SPM?	for all m and n [1551]
connected (p, q)-graph other than P_2	M	iff $5p/4 < q \leq p(p-1)/2$ [3913]
G^i	M	$\|G\| \geq 5$, $i \geq 3$ [3916]
G^2	M	$G \neq P_5$ and G does not have a 1-factor whose every edge is incident with an end-vertex of G [3916]
$K_{1,m,n}$	M	for all m, n [3366]
P_n^2	M	for all n except 2, 3, 5 [3366], [1248]
$G \times H$	M	iff G and H are magic [2009]

5.2 Edge-magic Total and Super Edge-magic Total Labelings

In 1970 Kotzig and Rosa [2017] defined a *magic valuation* of a graph $G(V, E)$ as a bijection f from $V \cup E$ to $\{1, 2, \ldots, |V \cup E|\}$ such that for all edges xy, $f(x)+f(y)+f(xy)$ is constant (called the *magic constant*). This notion was rediscovered by Ringel and Lladó [3176] in 1996 who called this labeling *edge-magic*. To distinguish between this usage from that of other kinds of labelings that use the word magic we will use the term *edge-magic total* labeling as introduced by Wallis [4120] in 2001. (We note that for 2-regular graphs a vertex-magic total labeling is an edge-magic total labeling and vice versa.) Kotzig and Rosa proved: $K_{m,n}$ has an edge-magic total labeling for all m and n; C_n has an edge-magic total labeling for all $n \geq 3$ (see also [1280], [3190], [627], and [1037]); and the disjoint union of n copies of P_2 has an edge-magic total labeling if and only if n is odd. They further state that K_n has an edge-magic total labeling if and only if $n = 1, 2, 3, 5$, or 6 (see [2018], [862], and [1037]) and ask whether all trees have edge-magic total labelings. Wallis, Baskoro, Miller, and Slamin [4124] enumerate every edge-magic total labeling of complete graphs. They also prove that the following graphs are edge-magic total: paths, crowns, complete bipartite graphs, and cycles with a single edge attached to one vertex. Enomoto, Llado, Nakamigana, and Ringel [1037] prove that all complete bipartite graphs are edge-magic total. They also show that wheels W_n are not edge-magic total when $n \equiv 3$

(mod 4) and conjectured that all other wheels are edge-magic total. This conjecture was proved when $n \equiv 0, 1 \pmod 4$ by Phillips, Rees, and Wallis [2836] and when $n \equiv 6 \pmod 8$ by Slamin, Bača, Lin, Miller, and Simanjuntak [3629]. Fukuchi [1161] verified all cases of the conjecture independently of the work of others. Slamin et al. further show that all fans are edge-magic total. In 2002 Lee and Kong [2154] conjectured that odd star forests are super edge magic. In 2019 Cerioli, Fernandes, Lee, Lintzmayer, Mota, and da Silva [755] proved this conjecture for odd symmetric star forests and proved that odd uniform forests of caterpillars are edge-magic. Afzal, Ather, Baig, and Maheshwari [81] analyzed pyramidion ladders and C_n-books for their super edge-magicness and gave some methods for finding new super edge-magic graphs from existing ones. In [899] Darmaji and Saputro use $G(+)P_m(+)H$ $(m \geq 2)$ to denote the graph obtained by taking one copy of the graphs G, H, and P_m, then connecting one endpoint of P_m to all vertices of G and the other endpoint of P_m to all vertices of H. For any super edge-magic total graphs G, they provide some graphs H such that $G(+)P_m(+)H$ is also super edge-magic total. They further show how to construct a super edge-magic total graph from a super edge-magic total graph by considering a super edge-magic labeling of the original graph.

In [1487] Ichishima and Muntaner-Batle study the super edge-magicness of graphs of order n with degree sequences: $4, 2, 2, \ldots, 2$. They also investigate the super edge-magic properties of certain families of graphs. One such family is graph $C(m, n)$—the one-point union of the cycles C_m and C_n. One of their results is $C(m, n)$ is not super edge-magic when $m + n$ is odd. They state that an exhaustive computer search shows that $C(3, 5)$ is not super edge-magic. However, Oshima [personal communication] [2742] pointed out that this is not the case, and showed that $C(3, 4k - 3)$ is super edge-magic for integers $k \geq 2$.

In [1901] Kanwal, Riasat, Imtiaz, Iftikhar, Javed, and Ashraf define a fork as the graph obtained by starting with three paths of length t with vertices $x_{1,j}, x_{2,j}, x_{3,j}, 1 \leq j \leq t$, a single new edge $x_{2,0}$ adjacent to $x_{2,1}$, an edge joining $x_{1,1}$ and $x_{2,1}$ and an edge joining $x_{2,1}$ and $x_{3,1}$. They gave super edge-magic total labelings and deficiencies of forks, the disjoint union of a fork with a star, a bistar, and a path, and of trees obtained by starting with two copies of P_{2t+1} and adding an edge joining the middle vertex of each path. The super edge-magic total labeling strengths of forks and the trees are also determined. Girija and Karthikeyan [1275] proved that 3 copies of the jelly fish graphs are super edge magic graphs.

Inspired by Kotzig-Rosa's notion, Enomoto, Lladó, Nakamigawa, and Ringel [1037] called a graph $G(V, E)$ with an edge-magic total labeling that has the additional property that the vertex labels are 1 to $|V|$ a *super edge-magic total* labeling (SEMT).

Kanwal and Kanwal [1900] determined super edge-magic total labelings and deficiencies for forests formed by two sided generalized combs, stars, combs, and banana trees. A two-sided generalized comb $Cb^2_{a,b}$, where b is odd, is obtained from a path P_{a+1} by attaching two paths $P_{(b+1)/2}$ to each of the vertices of degree two and one vertex of degree one of P_{a+1}. In [1897] Kanwal, Azam, and Iftikhar investigate the SEMT strength of generalized comb and the SEMT labeling and deficiency of forests composed of two components, where one of the components for each forest is a generalized comb and other component is

star, bistar, comb, or path. In [1898] Kanwal, Imtiaz, Iftikhar, Ashraf, Arshad, Irfan, and Sumbal z studied the super edge-magic deficiency of paths, caterpillars, and the disjoint union of a 2-sided generalized comb with a bistar. They also provide the super edge-magic total strength for a 2-sided generalized comb. Javed, Riasat, and Kanwal [1588] study super edge-magic total labeling and deficiencies of forests consisting of combs, generalized combs, and stars. Their results provide the evidence to support a conjecture proposed by Figueroa-Centeno, Ichishima, and Muntaner-Bartle [1092]. In [1527] Imran, Cancan, Nadeem, and Nadeem study the super edge magic deficiency of jewel graphs, shadow graphs, splitting graphs, jellyfish graphs, degree splitting graphs, and quadrilateral snake graphs. They determined the exact number of minimum isolated vertices that needs to be added to these graphs to make them super edge magic graphs. In [1509] Ichishima, Muntaner-Batle, and Takahashi generalized the notions of (super) edge-magic deficiency using the concepts of perfect (super) edge-magic graphs introduced in [2314]. López, Muntaner-Batle, Rius-Font. Ichishima et al presented some results on the valences of edge-magic and super edge-magic labelings. They provide some general results in this direction and investigated the perfect (super) edge-magic deficiency of $K_{1,n}$.

Baskoro, Sudarsana, and Cholily [593] provided some constructions of new super edge-magic graphs from some old ones by attaching 1, 2, or 3 pendent vertices and edges. In [1964] Kim introduces a new construction of new super edge-magic graphs by attaching any number pendent vertices and edges under some conditions. In [899] Darmaji and Saputro define a graph $G(+)P_m(+)H$ where $m \geq 2$ as a graph obtained by taking one copy of the graphs G and H and P_m, then connect an end point of P_m to all vertices of G and the other end P_m to all vertices of H. For any super edge-magic total graphs G, they provide some graphs H such that $G(+)P_m(+)H$ is also super edge-magic total. They further show how to construct a super edge-magic total graph from a super edge-magic total graph by considering a super edge-magic labeling of the origin graph. One such instance is $(P_{2n} \cup mK_1) + 2K_1$.

Ringel and Llado [3176] prove that a graph with p vertices and q edges is not edge-magic total if q is even and $p + q \equiv 2 \pmod 4$ and each vertex has odd degree. Ringel and Llado conjecture that trees are edge-magic total. In [582] Baskar Babujee and Rao show that the path with n vertices has an edge-magic total labeling with magic constant $(5n + 2)/2$ when n is even and $(5n + 1)/2$ when n is odd. For stars with n vertices they provide an edge-magic total labeling with magic constant $3n$. In [1052] Eshghi and Azimi discuss a zero-one integer programming model for finding edge-magic total labelings of large graphs.

Santhosh [3307] proved that for n odd and at least 3, the crown $C_n \odot P_2$ has an edge-magic total labeling with magic constant $(27n + 3)/2$ and for n odd and at least 3, $C_n \odot P_3$ has an edge-magic total labeling with magic constant $(39n + 3)/2$. Ngurah and Adiwijaya [2698] investigated whether various classes of chain graphs formed from ladders, triangular ladders, diagonal ladders, C_4, and K_4 have an edge-magic or super edge-magic labelings. Baig and Afzal [443] investigated the super edge-magicness of special classes of graphs having maximum magic constant $k = 3p$.

In [1112] Freyberg introduced a generalization of edge-magic total labeling that allows

multiple labels on the vertices or edges of a graph. He used this new labeling as a tool to construct face-magic labelings of some infinite families of graphs. He then considered the question "Given a graph G, for which $a, b, c \in \{0, 1\}$ does G admit a face-magic labeling of type (a, b, c)? He completely answered this question for two families of chained cycles, ladders and subdivided ladders, fans and subdivided fans, and wheels and subdivided wheels. See [1113] for some corrections for these results. Freyberg [1114] provided $(1, 1, 1)$-face-magic labelings for square tilings, hexagon tilings on a torus, and a special class of triangle tilings on a cylinder.

Ahmad, Baig, and Imran [121] define a *zig-zag triangle* as the graph obtained from the path $x_1, x_2, \ldots, x_n$ by adding n new vertices $y_1, y_2, \ldots, y_n$ and new edges y_1x_1, y_nx_{n-1}; x_iy_i for $1 \leq i \leq n$; $y_ix_{i-1}y_ix_{i+1}$ for $2 \leq i \leq n-1$. They define a graph Cb_n as one obtained from the path $x_1, x_2, \ldots, x_n$ adding $n-1$ new vertices $y_1, y_2, \ldots, y_{n-1}$ and new edges y_ix_{i+1} for $1 \leq i \leq n-1$. The graph Cb_n^* is obtained from the Cb_n by joining a new edge x_1y_1. They prove that zig-zag triangles, graphs that are the disjoint union of a star and a banana tree, certain disjoint unions of stars, and for $n \geq 4$, $Cb_n^* \cup Cb_{n-1}$ are super edge-magic total. Baig, Afzal, Imran, and Javaid [444] investigate the existence of super edge-magic labeling of volvox and pancyclic graphs.

The *super edge-magic deficiency* of a graph G, denoted by $\mu_s(G)$, is either the minimum nonnegative integer n such that $G \cup nK_1$ is super edge-magic or $+\infty$ if there exists no such n. Krisnawati, Ngurah, Hidayat, and Alghofari [2036] investigated the super edge-magic deficiency of forests whose components are subdivided stars or paths. Imran, Afzal, and Baig investigate the super edge-magic deficiency of volvox and dumbbell type graphs in [1520]. Kanwal, Iftikhar, and Azam [1896] found super edge magic total labelings and deficiencies of forests consisting of two components, where one of the components for each forest is a generalized comb and the other component is a star, bistar, comb, or path. They also investigated the super edge magic total strength of generalized combs. In [1464] Huang, Li, and Sun provided certain conditions for the existence of edge magic total labeling for graphs.

Let G be a graph with p vertices with $V(G) = \{v_1, v_2, \ldots, v_p\}$ and let S_m be the star with m leaves. If in G, every vertex v_i is identified to the center vertex of S_{m_i}, for some $m_i \geq 0$, $1 \leq i \leq n$, where $S_0 = K_1$, then the graph obtained is denoted by $G_{(m_1,m_2,\ldots,m_p)}$. Let $M(G) = \{(m_1, m_2, \ldots, m_p) \mid G_{(m_1,m_2,\ldots,m_p)} \text{ is a super edge-magic graph}\}$. The *star super edge-magic deficiency* $S\mu^*(G)$ is defined as

$$S\mu^*(G) = \begin{cases} \min_{(m_1,m_2,\ldots,m_p)}(m_1 + m_2 + \cdots + m_p) & \text{if } M(G) \neq \emptyset, \\ +\infty, \text{ if } M(G) = \emptyset. \end{cases}$$

In [1931] Kathiresan and Sabarimalai Madha determine the star super edge-magic deficiency of certain classes of graphs. In [2037] Krisnawati, Ngurah, Hidayat, and Alghofar showed that some subdivisions of double stars have zero (consecutively) super edge-magic deficiency.

Beardon [600] extended the notion of edge-magic total to countable infinite graphs $G(V, E)$ (that is, $V \cup E$ is countable). His main result is that a countably infinite tree that posesses an infinite simple path has a bijective edge-magic total labeling using the

integers as labels. He asks whether all countably infinite trees have an edge-magic total labeling with the integers as labels and whether the graph with the integers as vertices and an edge joining every two distinct vertices has a bijective edge-magic total labeling using the integers.

Cavenagh, Combe, and Nelson [762] investigate edge-magic total labelings of countably infinite graphs with labels from a countable Abelian group A. Their main result is that if G is a countable graph that has an infinite set of mutually disjoint edges and A is isomorphic to a countable subgroup of the real numbers under addition then for any k in A there is an edge-magic labeling of G with elements from A that has magic constant k.

Balakrishnan and Kumar [464] proved that the join of $\overline{K_n}$ and two disjoint copies of K_2 is edge-magic total if and only if $n = 3$. Yegnanarayanan [4264] has proved the following graphs have edge-magic total labelings: nP_3 where n is odd; $P_n + K_1; P_n \times C_3$ $(n \geq 2)$; the crown $C_n \odot K_1$; and $P_m \times C_3$ with n pendent vertices attached to each vertex of the outermost C_3. He conjectures that for all n, $C_n \odot \overline{K_n}$, the n-cycle with n pendent vertices attached at each vertex of the cycle, and nP_3 have edge-magic total labelings. In fact, Figueroa-Centeno, Ichishima, and Muntaner-Batle, [1094] have proved the stronger statement that for all $n \geq 3$, the corona $C_n \odot \overline{K_m}$ admits an edge-magic labeling where the set of vertex labels is $\{1, 2, \ldots, |V|\}$. (See also [2412].)

Yegnanarayanan [4264] also introduces several variations of edge-magic labelings and provides some results about them. Kotzig [4122] provides some necessary conditions for graphs with an even number of edges in which every vertex has odd degree to have an edge-magic total labeling. Craft and Tesar [862] proved that an r-regular graph with r odd and $p \equiv 4 \pmod 8$ vertices cannot be edge-magic total. Wallis [4120] proved that if G is an edge-magic total r-regular graph with p vertices and q edges where $r = 2^t s+1$ $(t > 0)$ and q is even, then 2^{t+2} divides p.

Figueroa-Centeno, Ichishima, and Muntaner-Batle [1088] have proved the following graphs are edge-magic total: $P_4 \cup nK_2$ for n odd; $P_3 \cup nK_2$; $P_5 \cup nK_2$; nP_i for n odd and $i = 3, 4, 5$; $2P_n$; $P_1 \cup P_2 \cup \cdots \cup P_n$; $mK_{1,n}$; $C_m \odot nK_1$; $K_1 \odot nK_2$ for n even; W_{2n}; $K_2 \times \overline{K}_n$, nK_3 for n odd (the case nK_3 for n even and larger than 2 is done in [2463]); binary trees, generalized Petersen graphs (see also [2700]), ladders (see also [4184]), books, fans, and odd cycles with pendent edges attached to one vertex.

In [1094] Figueroa-Centeno, Ichishima, Muntaner-Batle, and Oshima, investigate super edge-magic total labelings of graphs with two components. Among their results are: $C_3 \cup C_n$ is super edge-magic total if and only if $n \geq 6$ and n is even; $C_4 \cup C_n$ is super edge-magic total if and only if $n \geq 5$ and n is odd; $C_5 \cup C_n$ is super edge-magic total if and only if $n \geq 4$ and n is even; if m is even with $m \geq 4$ and n is odd with $n \geq m/2 + 2$, then $C_m \cup C_n$ is super edge-magic total; for $m = 6, 8$, or 10, $C_m \cup C_n$ is super edge-magic total if and only if $n \geq 3$ and n is odd; $2C_n$ is strongly felicitous if and only if $n \geq 4$ and n is even (the converse was proved by Lee, Schmeichel, and Shee in [2190]); $C_3 \cup P_n$ is super edge-magic total for $n \geq 6$; $C_4 \cup P_n$ is super edge-magic total if and only if $n \neq 3$; $C_5 \cup P_n$ is super edge-magic total for $n \geq 4$; if m is even with $m \geq 4$ and $n \geq m/2+2$ then $C_m \cup P_n$ is super edge-magic total; $P_m \cup P_n$ is super edge-magic total if and only $(m, n) \neq (2, 2)$ or(3,3); and $P_m \cup P_n$ is edge-magic total if and only $(m, n) \neq (2, 2)$. In [3184] Rizvi, Ali,

Iqbal, and Gulraze give super edge-magic total labelings of forests whose components are caterpillars and stars, forests whose components are stars and banana trees, and a new families of trees.

Enomoto, Llado, Nakamigawa, and Ringel [1037] conjecture that if G is a graph of order $n+m$ that contains K_n, then G is not edge-magic total for $n \gg m$. Wijaya and Baskoro [4184] proved that $P_m \times C_n$ is edge-magic total for odd n at least 3. Ngurah and Baskoro [2700] state that $P_2 \times C_n$ is not edge-magic total. Hegde and Shetty [1416] have shown that every T_p-tree (see §4.4 for the definition) is edge-magic total. Ngurah, Simanjuntak, and Baskoro [2710] show that certain subdivisions of the star $K_{1,3}$ have edge-magic total labelings. Ali, Hussain, Shaker, and Javaid [188] provide super edge-magic total labelings of subdivisions of stars $K_{1,p}$ for $p \geq 5$. In [2705] Ngurah, Baskoro, Tomescu gave methods for construction new (super) edge-magic total graphs from old ones by adding some new pendent edges. They also proved that $K_{1,m} \cup {P_n}^m$ is super edge-magic total. Wallis [4120] proves that a cycle with one pendent edge is edge-magic total. In [4120] Wallis poses a large number of research problems about edge-magic total graphs.

Petersen graphs that they denote by $GGP(n;\pi)$, which consists of an outer n-cycle $x_0, x_1, \ldots, x_{n-1}, x_0$, a set of n-spokes $x_i y_i$, $0 \leq i \leq n-1$, and n inner edges defined by $y_i y_{\pi(i)}, i = 0, \ldots, n-1$. Notice that, for the permutation π defined by $\pi(i) = i+k \pmod n$ we have $GGP(n;\pi) = P(n;k)$. They define a second generalization of generalized Petersen graphs, $GGP(n;\pi_2, \ldots, \pi_m)$, as the graphs with vertex sets $\cup_{j=1}^{m}\{x_i^j : i = 0, \ldots, n-1\}$, an outer n-cycle $x_0^1, x_1^1, \ldots, x_{n-1}^1, x_0^1$, and inner edges $x_i^{j-1}x_i^j$ and $x_i^j x_{\pi_j(i)}^j$, for $j = 2, \ldots, m$, and $i = 0, \ldots, n-1$. Notice that, $GGP(n;\pi_2, \ldots, \pi_m) = P_m \times C_n$, when $\pi_j(i) = i+1 \pmod n$ for every $j = 2, \ldots, m$. Among their results are the Petersen graphs are super edge-magic total; for each m with $1 < l \leq m$ and $1 \leq k \leq 2$, the graph $GGP(5;\pi_2, \ldots, \pi_m)$, where $\pi_i = \sigma_1$ for $i \neq l$ and $\pi_l = \sigma_k$, is super edge-magic total; for each $1 \leq k \leq 2$, the graph $P(5n;k+5r)$ where r is the smallest integer such that $k+5r = 1 \pmod n$ is super edge-magic total.

A w-graph, $W(n)$, has vertices $\{(c_1, c_2, b, w, d) \cup (x^1, x^2, \ldots, x^n) \cup (y^1, y^2, \ldots, y^n)\}$ and edges $\{(c_1x^1, c_1x^2, \ldots, c_1x^n) \cup (c_2y^1, c_2y^2, \ldots, c_2y^n) \cup (c_1b, c_1w) \cup (c_2w, c_2d)\}$. A w-tree, $WT(n,k)$, is a tree obtained by taking k copies of a w-graph $W(n)$ and a new vertex $\mathbf{a}$ and joining $\mathbf{a}$ with in each copy d where $n \geq 2$ and $k \geq 3$. An *extended w-tree* $Ewt(n,k,r)$ is a tree obtained by taking k copies of an extended w-graph $Ew(n,r)$ and a new vertex $\mathbf{a}$ and joining $\mathbf{a}$ with the vertex d in each of the k copies for $n \geq 2$, $k \geq 3$ and $r \geq 2$. Super edge-magic total labelings for w-trees, extended w-trees, and disjoint unions of extended w-trees are given in [1585], [1582], and [187]. Javaid, Hussain, Ali, and Shaker [1586] provided super edge-magic total labelings of subdivisions of $K_{1,4}$ and w-trees. Shaker, Rana, Zobair, and Hussain [3474] gave a super edge-magic total labeling for a subdivided star with a center of degree at least 4.

In 1988 Godbod and Slater [1280] made the following conjecture. If n is odd, $n \neq 5$, C_n has an edge magic labeling with valence k, when $(5n+3)/2 \leq k \leq (7n+3)/2$. If n is even, C_n has an edge-magic labeling with valence k when $5n/2 + 2 \leq k \leq 7n/2 + 1$. Except for small values of n, very few valences for edge-magic labelings of C_n are known.

In [2317] López, Muntaner-Batle, and Rius-Font use the $\otimes_h$-product in order to prove the following two results. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ be the unique prime factorization of an odd number n. Then C_n admits at least $1 + \sum_{i=1}^{k} \alpha_i$ edge-magic labelings with at least $1 + \sum_{i=1}^{k} \alpha_i$ mutually different valences. Let $n = 2^{\alpha} p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ be the unique prime factorization of an even number n, with $p_1 > p_2 > \cdots > p_k$. Then C_n admits at least $\sum_{i=1}^{k} \alpha_i$ edge-magic labelings with at least $\sum_{i=1}^{k} \alpha_i$ mutually different valences. If $\alpha \geq 2$ this lower bound can be improved to $1 + \sum_{i=1}^{k} \alpha_i$. In [2309] López, Muntaner-Batle, and Prabu introduce a new $\otimes_h$ labeling construction that has a wider range of applications and applies it to the magic valences of cycles and crowns.

In [3853] Swita, Rafflesia, Henni Ms, Adji, and Simanihuruk use $B[(C_a, m), (C_b, n), P_t]$ to denote the graph that consists of m cycles C_a and n cycles C_b with a common path P_t. They proved that $B[(C_7, 1), (C_3, n), P_2]$ admits an edge-magic total labeling, $B[(C_a, 1), (C_3, n), P_2]$ admits a super edge-magic total labeling for all $a \equiv 3 \bmod 4$ $(a > 3)$, and $B[(C_7, 2), (C_3, n), P_2]$ admits a super edge-magic total labeling. Swita, Butar-Butar, Pramesti, Febriyan, Puspita, Saftri, Azhari, and Simanihuruk [3852] proved that the book graph $B[(C_4, \alpha), (C_3, \beta), 2]$ admits a super edge-magic total labeling for all even α and $\beta = 1$ or 2. Using the super edge-magic total labelings of $B[(C_4, 26), (C_3, 2), P_2]$, they give an illustration how to encrypt and decrypt a secret message in cryptography by the Rivest, Shamir, and Adleman (RSA) algorithm.

In 1996 Erdős asked for $M(n)$, the maximum number of edges that an edge-magic total graph of order n can have (see [862]). In 1999 Craft and Tesar [862] gave the bound $\lfloor n^2/4 \rfloor \leq M(n) \leq \lfloor n(n-1)/2 \rfloor$. For large n this was improved by Pikhurko [2840] in 2006 to $2n^2/7 + O(n) \leq M(n) \leq (0.489 + \cdots + o(1)n^2)$.

Enomoto, Lladó, Nakamigawa, and Muntaner-Batle [1037] proved that a super edge-magic total graph $G(V, E)$ with $|V| \geq 4$ and with girth at least 4 has at most $2|V| - 5$ edges. They prove this bound is tight for graphs with girth 4 and 5 in [1037] and [1505].

In his Ph.D. thesis, Barrientos [511] introduced the following notion. Let $L_1, L_2, \ldots, L_h$ be ordered paths in the grid $P_r \times P_t$ that are maximal straight segments such that the end vertex of L_i is the beginning vertex of L_{i+1} for $i = 1, 2, \ldots, h-1$. Suppose for some i with $1 < i < h$ we have $V(L_i) = \{u_0, v_0\}$ where u_0 is the end vertex of L_{i-1} and the beginning vertex of L_i and v_0 is the end vertex of L_1 and the beginning vertex of L_{i+1}. Let $u \in V(L_{i-1}) - \{u_0\}$ and $v \in V(L_{i+1}) - \{v_0\}$. The replacement of the edge u_0v_0 by a new edge uv is called an *elementary transformation* of the path P_n. A tree is called a *path-like tree* if it can be obtained from P_n by a sequence of elementary transformations on an embedding of P_n in a 2-dimensional grid. In [397] Bača, Lin, and Muntaner-Batle proved that if $T_1, T_2, \ldots, T_m$ are path-like trees each of order $n \geq 4$ where m is odd and at least 3, then $T_1 \cup T_2, \cup \cdots \cup T_m$ has a super edge-magic labeling. In [396] Bača, Lin, Muntaner-Batle and Rius-Font proved that the number of such trees grows at least exponentially with m. As an open problem Bača, Lin, Muntaner-Batle and Rius-Font ask if graphs of the form $T_1 \cup T_2 \cup \cdots \cup T_m$ where $T_1, T_2, \ldots, T_m$ are path-like trees each of order $n \geq 2$ and m is even have a super edge-magic labeling. In [511] Barrientos proved that all path-like trees admit an α-valuation. Using Barrientos's result, it is very easy to obtain that all path-like trees are a special kind of super edge-magic by using a super edge-magic

labeling of the path P_n, and hence they are also super edge-magic. Furthermore, Figueroa-Centeno, Ichishima, and Muntaner-Batle proved that if a tree is super edge-magic, then it is also harmonious. Therefore all path-like trees are also harmonious. In [2322] López, Muntaner-Batle, and Rius-Font also use a variation of the Kronecker product of matrices in order to obtain lower bounds for the number of non isomorphic super edge-magic labeling of some types of path-like trees. As a corollary they obtain lower bounds for the number of harmonious labelings of the same type of trees. López, Muntaner-Batle, and Rius-Font [2318] proved that if $m \geq 4$ is an even integer and $n \geq 3$ is an odd divisor of m, then $C_m \cup C_n$ is super edge-magic. Lee and Kong conjecture that if n is an odd, then $St(a_1, a_2, \ldots, a_n)$ is super edge-magic, and they proved that the following graphs are super edge-magic: $St(m, n)$ $(n \equiv 0 \bmod (m+1))$, $St(1, k, n)(k = 1, 2 \text{ or } n)$, $St(2, k, n)$ $(k = 2, 3)$, $St(1, 1, k, n)$ $(k = 2, 3)$, $St(k, 2, 2, n)$ $(k = 1, 2)$. Zhenbin and Chongjin [4334] proved that $St(1, m, n), St(3, m, m+1), St(n, n+1, n+2)$ are super edge-magic, and under some conditions $St(a_1, a_2, \ldots, a_{2n+1})$, $St(a_1, a_2, \ldots, a_{4n+1})$, $St(a_1, a_2, \ldots, a_{4n+3})$ are also super edge-magic. In [1087] Figueroa-Centeno, Ichishima, and Muntaner-Batle conjectured that the cycle books $B(4, m)$ that consists of m cycles C_4 with a common path P_2 is super edge-magic total if and only if m is even or $m \equiv 5 \pmod 8$. Simanihuruk, Kusmayadi, Swita, Romala, and Damanik [3584] proved this conjecture for $m \geq 36$ and m even.

For a simple graph H we say that $G(V, E)$ admits an *H-covering* if every edge in $E(G)$ belongs to a subgraph of G that is isomorphic to H. In [2321] López, Muntaner-Batle, Rius-Font study a relationship existing among (super) magic coverings and the Kronecker product of matrices. (For a simple graph H, $G(V, E)$ admits an *H-covering* if every edge in $E(G)$ belongs to a subgraph of G that is isomorphic to H.) Their results can be applied to construct S-magic partitions. For m copies of a graph G and a fixed subgraph H of each copy the graph $I(G, H, m)$ is formed by taking of all the G_i's and identifying their subgraph H. Liang [2253] determined which $I(G, H, m)$ and which mG have G supermagic coverings. Farida, Indah, and Sudiby [1067] study magic coverings of domino graphs (graphs for which every vertex is contained in at most two maximal cliques) and edge magic labelings of domino graphs. They note that coverings can be applied to secret sharing schemes and edge magic labelings can be applied to ruler models.

Bača, Lin and Muntaner-Batle in [395] using a generalization of the Kronecker product of matrices prove that the number of non-isomorphic super edge-magic labelings of the disjoint union of m copies of the path P_n, $m \equiv 2 \pmod 4$, $m \geq 2$, $n \geq 4$, is at least $(m/2)^{(2n-2)}$.

In [2319] López, Muntaner-Batle and Rius-Font proved that every super edge-magic graph with p vertices and q edges where $q \geq p - 1$ has an even harmonious labeling (See Section 4.6.) In [2313] they stated some open problems concerning relationships among super edge-magic labelings and graceful and harmonious labelings. A *Langford sequence* of order m and defect d is a sequence $(t_1, t_2, \ldots, t_{2m})$ of $2m$ numbers such that (i) for every $k \in [d, d+m-1]$ there exist exactly two subscripts $i, j \in [1, 2m]$ with $ti = tj = k$ and (ii) the subscripts i and j satisfy the condition $|i - j| = k$. López and Muntaner-Batle [2308] provided new lower bounds on the number of distinct Langford sequences with certain properties in terms of the number of 1-regular super edge-magic labeled digraphs of a

particular order.

Lee and Lee [2158] prove the following graphs are super edge-magic: $P_{2n} + \overline{K_m}$, $(P_2 \cup nK_1) + \overline{K_2}$, graphs obtained by appending a path to the apex of a fan with at least 4 vertices (*umbrella*), and *jelly fish* graphs $J(m,n)$ obtained from a 4-cycle v_1, v_2, v_3, v_4 by joining v_1 and v_3 with an edge and appending m pendent edges to v_2 and n pendent edges to v_4. In [80] Afzel introduces two new families of graphs called carrom and jukebox graphs and proves they admit super edge-magic labelings. Carroms are generalizations of $C_n \times P_2$.

Marimuthu and Balakrishnan [2422] define a graph $G(p,q)$ to be *edge magic graceful* if there exists a bijection f from $V(G) \cup E(G)$ to $\{1, 2, \ldots, p+q\}$ such $|f(u) + f(v) - f(uv)|$ is a constant for all edges uv of G. An edge magic graceful graph is said to be *super edge magic graceful* if $V(G) = \{1, 2, \ldots, p\}$. They present some properties of super edge magic graceful graphs, prove some classes of graphs are super edge magic graceful, and prove that every super edge magic graceful graph with either $f(uv) > f(u) + f(v)$ for all edges uv or $f(uv) < f(u) + f(v)$ for all edges uv is sequential, harmonious, super edge magic and not graceful. Marimuthu, Kavitha, and Balakrishnan [2423] proved that the generalized Petersen graphs $P(n,1)$ and $P(n,(n-1)/2)$ are super edge magic graceful when n is odd.

Let $G = (V,E)$ be a (p,q)-linear forest. In [396] Bača, Lin, Muntaner-Batle, and Rius-Font call a labeling f a *strong super edge-magic* labeling of G and G a *strong super edge-magic* graph if $f : V \cup E \to \{1, 2, \ldots, p+q\}$ with the extra property that if $uv \in E, u', v' \in V(G)$ and $d_G(u,u') = d_G(v,v') < +\infty$, then we have that $f(u) + f(v) = f(u') + f(v')$. In [131] Ahmad, López, Muntaner-Batle, and Rius-Font define the concept of strong super edge-magic labeling of a graph with respect to a linear forest as follows. Let $G = (V,E)$ be a (p,q)-graph and let F be any linear forest contained in G. A *strong super edge-magic labeling of G with respect to F* is a super edge-magic labeling f of G with the extra property with if $uv \in E(F), u', v' \in V(F)$ and $d_F(u,u') = d_F(v,v') < +\infty$ then we have that $f(u) + f(v) = f(u') + f(v')$. If a graph G admits a strong super edge-magic labeling with respect to some linear forest F, they say that G is a *strong super edge-magic graph with respect to F*. They prove that if m is odd and G is an acyclic graph which is strong super edge-magic with respect to a linear forest F, then mG is strong super edge-magic with respect to $F_1 \cup F_2 \cup \cdots \cup F_m$, where $F_i \simeq F$ for $i = 1, 2, \ldots, m$ and every regular caterpillar is strong super edge-magic with respect to its spine.

Noting that for a super edge-magic labeling f of a graph G with p vertices and q edges, the magic constant k is given by the formula: $k = (\sum_{u \in V} \deg(u) f(u) + \sum_{i=p+1}^{p+q} i)/q$, López, Muntaner-Batle and Rius-Font [2314] define the set $S_G = \left\{ (\sum_{u \in V} \deg(u) g(u) + \sum_{i=p+1}^{p+q} i)/q : \text{ the function } g : V \to \{i\}_{i=1}^{p} \text{ is bijective} \right\}$. If $\lceil \min S_G \rceil \leq \lfloor \max S_G \rfloor$ then the *super edge-magic interval* of G is the set $I_G = [\lceil \min S_G \rceil, \lfloor \max S_G \rfloor] \cap \mathbb{N}$. The *super edge-magic set* of G is $\sigma_G = \{k \in I_G :$ there exists a super edge-magic labeling of G with valence $k\}$. López et al. call a graph G *perfect super edge-magic* if $I_G = \sigma_G$. They show that the family of paths P_n is a family of perfect super edge-magic graphs with $|I_{P_n}| = 1$ if n is even and $|I_{P_n}| = 2$ if n is odd and raise the question of whether there is an infinite family $F_1, F_2, \ldots$ of graphs such

that each member of the family is perfect super edge-magic and $\lim_{i\to+\infty}|I_{F_i}| = +\infty$. They show that graphs $G \cong C_{p^k} \odot \overline{K_n}$ where $p > 2$ is a prime is such a family. Takahasi, Muntaner-Batle, and Ichishima [3859] investigated the perfect (super) edge-magic deficiency of $K_{1,n}$

In [2319] López et al. define the *irregular crown* $C(n; j_1, j_2, \ldots, j_n) = (V, E)$, where $n > 2$ and $j_i \geq 0$ for all $i \in \{1, 2, \ldots, n\}$ as follows: $V = \{v_i\}_{i=1}^n \cup V_1 \cup V_2 \cup \cdots \cup V_n$, where $V_k = \{v_k^1, v_k^2, \ldots, v_k^{j_k}\}$, if $j_k \neq 0$ and $V_k = \emptyset$ if $j_k = 0$, for each $k \in \{1, 2, \ldots, n\}$ and $E = \{v_i v_{i+1}\}_{i=1}^{n-1} \cup \{v_1 v_n\} \cup (\cup_{k=1, j_k\neq 0}^n \{v_k v_k^l\}_{l=1}^{j_k})$. In particular, they denote $C_m^n \cong C(m; j_1, j_2, \ldots, j_m)$, where $j_{2i-1} = n$, for each i with $1 \leq i \leq (m+1)/2$, and $j_{2i} = 0$, for each i, $1 \leq i \leq (m-1)/2$. They prove that the graphs C_3^n and C_5^n are perfect edge-magic for all $n > 1$.

López et al. [2316] define $\mathfrak{F}^k$-family and $\mathfrak{E}^k$-family of graphs as follows. The infinite family of graphs $(F_1, F_2, \ldots)$ is an $\mathfrak{F}^k$-*family* if each element F_n admits exactly k different valences for super edge-magic labelings, and $\lim_{n\to+\infty}|I(F_n)| = +\infty$. The infinite family of graphs $(F_1, F_2, \ldots)$ is an $\mathfrak{E}^k$-*family* if each element F_n admits exactly k different valences for edge-magic labelings, and $\lim_{n\to+\infty}|J(F_n)| = +\infty$. An easy observation is that $(K_{1,2}, K_{1,3}, \ldots)$ is an $\mathfrak{F}^2$-family and an $\mathfrak{E}^3$-family. They pose the two problems: for which positive integers k is it possible to find $\mathfrak{F}^k$-families and $\mathfrak{E}^k$-families? Their main results in [2316] are that an $\mathfrak{F}^k$-family exits for each $k = 1, 2, 3$; and an $\mathfrak{E}^k$-family exits for each $k = 3, 4$ and 7.

McSorley and Trono [2468] define a relaxed version of edge-magic total labelings of a graph as follows. An *edge-magic injection* μ of a graph G is an injection μ from the set of vertices and edges of G to the natural numbers such that for every edge uv the sum $\mu(u) + \mu(v) + \mu(uv)$ is some constant k_μ. They investigate $\kappa(G)$, the smallest k_μ among all edge-magic injections of a graph G. They determine $\kappa(G)$ in the cases that G is K_2, K_3, K_5, K_6 (recall that these are the only complete graphs that have edge-magic total labelings), a path, a cycle, or certain types of trees. They also show that every graph has an edge-magic injection and give bounds for $\kappa(K_n)$.

Avadayappan, Vasuki, and Jeyanthi [322] define the *edge-magic total strength* of a graph G as the minimum of all constants over all edge-magic total labelings of G. We denote this by $emt(G)$. They use the notation $< K_{1,n} : 2 >$ for the tree obtained from the bistar $B_{n,n}$ (the graph obtained by joining the center vertices of two copies of $K_{1,n}$ with an edge) by subdividing the edge joining the two stars. They prove: $emt(P_{2n}) = 5n+1$; $emt(P_{2n+1}) = 5n+3$; $emt(< K_{1,n} : 2 >) = 4n+9$; $emt(B_{n,n}) = 5n+6$; $emt((2n+1)P_2) = 9n+6$; $emt(C_{2n+1}) = 5n+4$; $emt(C_{2n}) = 5n+2$; $emt(K_{1,n}) = 2n+4$; $emt(P_n^2) = 3n$; and $emt(K_{n,m}) \leq (m+2)(n+1)$ where $n \leq m$. Using an analogous definition for super edge-magic total strength, Swaninathan and Jeyanthi [3843], [3843], [3844] provide results about the super edge-magic strength of trees, fire crackers, unicyclic graphs, and generalized theta graphs. Basha [551] determined the reverse super edge-magic strength of banana trees.

Ngurah, Simanjuntak, and Baskoro [2710] show that certain subdivisions of the star $K_{1,3}$ have super edge-magic total labelings. In [1037] Enomoto, Lladó, Nakamigawa and Ringel conjectured that all trees have a super edge-magic total labeling. Ichishima,

Muntaner-Batle, and Rius-Font [1504] have shown that any tree of order p is contained in a tree of order at most $2p-3$ that has a super edge-magic total labeling.

In [396] Bača, Lin, Muntaner-Batle, and Rius-Font use a generalization of the Kronecker product of matrices introduced by Figueroa-Centeno, Ichishima, Muntaner-Batle, and Rius-Font [1096] to obtain an exponential lower bound for the number of non-isomorphic strong super edge-magic labelings of the graph mP_n, for m odd and any n, starting from the strong super edge-magic labeling of P_n. They prove that the number of non-isomorphic strong super edge-magic labelings of the graph mP_n, $n \geq 4$, is at least $\frac{5}{2}2^{\lfloor \frac{m}{2} \rfloor}+1$ where $m \geq 3$ is an odd positive integer. This result allows them to generate an exponential number of non-isomorphic super edge-magic labelings of the forest $F \cong \bigcup_{j=1}^{m} T_j$, where each T_j is a path-like tree of order n and m is an odd integer.

López, Muntaner-Batle, and Rius-Font [2311] introduced a generalization of super edge-magic graphs called *super edge-magic models* and prove some results about them.

Yegnanarayanan and Vaidhyanathan [4265] use the term *nice* $(1,1)$ *edge-magic labeling* for a super edge-magic total labeling. They prove: a super edge-magic total labeling f of a (p,q)-graph G satisfies $2\sum_{v\in V(G)} f(v)deg(v) \equiv 0 \bmod q$; if G is (p,q) r-regular graph $(r>1)$ with a super edge-magic total labeling then q is odd and the magic constant is $(4p+q+3)/2$; every super edge-magic total labeling has at least two vertices of degree less than 4; fans $P_n + K_1$ are edge-magic total for all n and super edge-magic total if and only if n is at most 6; books B_n are edge-magic total for all n; a super edge-magic total (p,q)-graph with $q \geq p$ is sequential; a super edge-magic total tree is sequential; and a super edge-magic total tree is cordial. These last three results had been proved earlier by Figueroa-Centenoa, Ichishima, and Muntaner-Batle [1087].

In [4264] Yegnanarayanan conjectured that the disjoint union of $2t$ copies of P_3 has a $(1,1)$ edge-magic labeling and posed the problem of determining the values of m and n such that mP_n has a $(1,1)$ edge-magic labeling. Manickam and Marudai [2412] prove the conjecture and partially settle the open problem.

Hegde and Shetty [1422] (see also [1421]) define the *maximum magic strength* of a graph G as the maximum magic constant over all edge-magic total labelings of G. We use $eMt(G)$ to denote the maximum magic strength of G. Hegde and Shetty call a graph G with p vertices *strong magic* if $eMt(G) = emt(G)$; *ideal magic* if $1 \leq eMt(G) - emt(G) \leq p$; and *weak magic* if $eMt(G) - emt(G) > p$. They prove that for an edge-magic total graph G with p vertices and q edges, $eMt(G) = 3(p+q+1) - emt(G)$. Using this result they obtain: P_n is ideal magic for $n > 2$; $K_{1,1}$ is strong magic; $K_{1,2}$ and $K_{1,3}$ are ideal magic; and $K_{1,n}$ is weak magic for $n > 3$; $B_{n,n}$ is ideal magic; $(2n+1)P_2$ is strong magic; cycles are ideal magic; and the generalized web $W(t,3)$ (see §2.2 for the definition) with the central vertex deleted is weak magic.

Santhosh [3307] has shown that for n odd and at least 3, $eMt(C_n \odot P_2) = (27n+3)/2$ and for n odd and at least 3, $(39n+3)/2 \leq eMt(C_n \odot P_2) \leq (40n+3)/2$. Moreover, he proved that for n odd and at least 3 both $C_n \odot P_2$ and $C_n \odot P_3$ are weak magic. In [815] Chopra and Lee provided a number of families of super edge-magic graphs that are weak magic.

In [2592] Murugan introduces the notions of *almost-magic labeling, relaxed-magic la-*

beling, almost-magic strength, and *relaxed-magic strength* of a graph. He determines the magic strength of Huffman trees and twigs of odd order and the almost-magic strength of nP_2 (n is even) and twigs of even order. Also, he obtains a bound on the magic strength of the path-union $P_n(m)$ and on the relaxed-magic strength of kS_n and kP_n.

Enomoto, Llado, Nakamigawa, and Ringel [1037] call an edge-magic total labeling *super edge-magic* if the set of vertex labels is $\{1, 2, \ldots, |V|\}$ (Wallis [4120] calls these labelings *strongly edge-magic*). They prove the following: C_n is super edge-magic if and only if n is odd; caterpillars are super edge-magic; $K_{m,n}$ is super edge-magic if and only if $m = 1$ or $n = 1$; and K_n is super edge-magic if and only if $n = 1, 2$, or 3. They also prove that if a graph with p vertices and q edges is super edge-magic then, $q \leq 2p-3$. In [2382] MacDougall and Wallis study super edge-magic (p, q)-graphs where $q = 2p-3$. Enomoto et al. [1037] conjecture that every tree is super edge-magic. Lee and Shan [2198] have verified this conjecture for trees with up to 17 vertices with a computer. Fukuchi, and Oshima, [1163] have shown that if T is a tree of order $n \geq 2$ such that T has diameter greater than or equal to $n-5$, then T has a super edge-magic labeling.

Various classes of banana trees that have super edge-magic total labelings have been found by Swaminathan and Jeyanthi [3843] and Hussain, Baskoro, and Slamin [1475]. In [106] Ahmad, Ali, and Baskoro [106] investigate the existence of super edge-magic labelings of subdivisions of banana trees and disjoint unions of banana trees. They pose three open problems.

Kotzig and Rosa's ([2017] and [2018]) proof that nK_2 is edge-magic total when n is odd actually shows that it is super edge-magic. Kotzig and Rosa also prove that every caterpillar is super-edge magic. Figueroa-Centeno, Ichishima, and Muntaner-Batle prove the following: if G is a bipartite or tripartite (super) edge-magic graph, then nG is (super) edge-magic when n is odd [1091]; if m is a multiple of $n+1$, then $K_{1,m} \cup K_{1,n}$ is super edge-magic [1091]; $K_{1,2} \cup K_{1,n}$ is super edge-magic if and only if n is a multiple of 3; $K_{1,m} \cup K_{1,n}$ is edge-magic if and only if mn is even [1091]; $K_{1,3} \cup K_{1,n}$ is super edge-magic if and only if n is a multiple of 4 [1091]; $P_m \cup K_{1,n}$ is super edge-magic when $m \geq 4$ [1091]; $2P_n$ is super edge-magic if and only if n is not 2 or 3; $K_{1,m} \cup 2nK_2$ is super edge-magic for all m and n [1091]; $C_3 \cup C_n$ is super edge-magic if and only if $n \geq 6$ and n is even [1094] (see also [1329]); $C_4 \cup C_n$ is super edge-magic if and only if $n \geq 5$ and n is odd [1094] (see also [1329]); $C_5 \cup C_n$ is super edge-magic if and only if $n \geq 4$ and n is even [1094]; if m is even and at least 6 and n is odd and satisfies $n \geq m/2 + 2$, then $C_m \cup C_n$ is super edge-magic [1094]; $C_4 \cup P_n$ is super edge-magic if and only if $n \neq 3$ [1094]; $C_5 \cup P_n$ is super edge-magic if $n \geq 4$ [1094]; if m is even and at least 6 and $n \geq m/2 + 2$, then $C_m \cup P_n$ is super edge-magic [1094]; and $P_m \cup P_n$ is super edge-magic if and only if $(m, n) \neq (2, 2)$ or (3,3) [1094]. They [1091] conjecture that $K_{1,m} \cup K_{1,n}$ is super edge-magic only when m is a multiple of $n+1$ and they prove that if G is a super edge-magic graph with p vertices and q edges with $p \geq 4$ and $q \geq 2p-4$, then G contains triangles. In [1094] Figueroa-Centeno et al. conjecture that $C_m \cup C_n$ is super edge-magic if and only if $m+n \geq 9$ and $m+n$ is odd.

Singgih [3598] gave super edge magic total labelings for unions of books $mB(n)$ for odd m and $m(P_2 \times P_n)$ for m and n odd. In the editions of this survey published between

2017 and 2023 it was mistakenly stated that [3598] included proofs that $r(P_m \times P_n)$ for odd r and $(m,n) \neq (2,2)$ or (3,3), $r(P_3 \times mP_n)$ for odd r; mP_n for $m \equiv 2 \pmod 4$ and $n \neq 2,3$, and mP_{4n} for $m \equiv 2 \pmod 4$, $n > 1$ admitted super edge magic total labelings.

In [1162] Fukuchi and Oshima describe a construction of super edge-magic labelings of some families of trees with diameter 4. Salman, Ngurah, and Izzati [3259] use S_n^m $(n \geq 3)$ to denote the graph obtained by inserting m vertices in every edge of the star S_n. They prove that S_n^m is super edge-magic when $m = 1$ or 2. Darmaji, Rinurwati, Wahyudi, and Saputro showed how to construction of super edge-magic total graphs by adding some vertices and edges of a given super edge-magic graph so that the new graphs are also super edge-magic total graphs.

In [2320] López, Muntaner-Batle, and Ruis-Font introduce a new construction for super edge-magic labelings of 2-regular graphs which allows loops and is related to the knight jump in the game of chess. They also study the super edge-magic properties of cycles with cords.

Muntaner-Batle calls a bipartite graph with partite sets V_1 and V_2 *special super edge-magic* if is has a super edge-magic total labeling f with the property that $f(V_1) = \{1, 2, \ldots, |V_1|\}$. He proves that a tree has a special super edge-magic labeling if and only if it has an α-labeling (see §3.1 for the definition). Figueroa-Centeno, Ichishima, Muntaner-Batle, and Rius-Font [1096] use matrices to generate edge-magic total labeling and define the concept of super edge-magic total labelings for digraphs. They prove that if G is a graph with a super edge-magic total labeling then for every natural number d there exists a natural number k such that G has a (k,d)-arithmetic labeling (see §4.2 for the definition). In [2134] Lee and Lee prove that a graph is super edge-magic if and only if it is $(k,1)$-strongly indexable (see §4.3 for the definition of (k,d)-strongly indexable graphs). They also provide a way to construct (k,d)-strongly indexable graphs from two given (k,d)-strongly indexable graphs. This allows them to obtain several existing results about super edge-magic graphs as special cases of their constructions. Acharya and Germina [45] proved that the class of strongly indexable graphs is a proper subclass of super edge-magic graphs.

In [1482] Ichishima, López, Muntaner-Batle and Rius-Font show how one can use the product $\otimes_h$ of super edge-magic 1-regular labeled digraphs and digraphs with harmonious, or sequential labelings to create new undirected graphs that have harmonious, sequential labelings or partitional labelings (see §4.1 for the definition). They define the product $\otimes_h$ as follows. Let $\overrightarrow{D} = (V,E)$ be a digraph with adjacency matrix $A(\overrightarrow{D}) = (a_{ij})$ and let $\Gamma = \{F_i\}_{i=1}^m$ be a family of m digraphs all with the same set of vertices V'. Assume that $h : E \longrightarrow \Gamma$ is any function that assigns elements of Γ to the arcs of D. Then the digraph $\overrightarrow{D} \otimes_h \Gamma$ is defined by $V(D \otimes_h \Gamma) = V \times V'$ and $((a_1,b_1),(a_2,b_2)) \in E(D \otimes_h \Gamma) \Longleftrightarrow [(a_1,a_2) \in E(D) \wedge (b_1,b_2) \in E(h(a_1,a_2))]$. An alternative way of defining the same product is through adjacency matrices, since one can obtain the adjacency matrix of $\overrightarrow{D} \otimes_h \Gamma$ as follows: if $a_{ij} = 0$ then a_{ij} is multiplied by the $p' \times p'$ 0-square matrix, where $p' = |V'|$. If $a_{ij} = 1$ then a_{ij} is multiplied by $A(h(i,j))$ where $A(h(i,j))$ is the adjacency matrix of the digraph $h(i,j)$. They prove the following. Let $\overrightarrow{D} = (V,E)$ be a harmonious (p,q)-

digraph with $p \leq q$ and let h be any function from E to the set of all super edge-magic 1-regular labeled digraphs of order n, which we denote by S_n. Then the undirected graph $und(\overrightarrow{D} \otimes_h S_n)$ is harmonious. Let $\overrightarrow{D} = (V, E)$ be a sequential digraph and let $h : E \longrightarrow S_n$ be any function. Then $und(\overrightarrow{D} \otimes_h S_n)$ is sequential. Let D be a partitional graph and let $h : E \longrightarrow S_n$ be any function, where $\overrightarrow{D} = (V, E)$ is the digraph obtained by orienting all edges from one stable set to the other one. Then $und(\overrightarrow{D} \otimes_h S_n)$ is partitional.

Let $D(V, A)$ be a digraph of order p and size q. For an integer $k \geq 1$ and for $v \in V(D)$, let $w_k(v) = \sum_{e \in E_k(v)} f(e)$, where $E_k(v)$ is the set of all arcs that are at distance at most k from v. The digraph D is said to be *E_k-regular* with regularity r if $|E_k(e)| = r$ for some integer $r \geq 1$ and for all $e \in A(D)$. A V_k-super vertex out-magic total labeling (V_k-SVIMTL) is a bijection $f : V(D) \cup A(D) \to \{1, 2, \ldots, p + q\}$ with the property that $f(V(D)) = \{1, 2, \ldots, p\}$ and there exists a positive integer M such that $f(v) + w_k(v) = M$ for each vertex in $V(D)$. A digraph that admits an V_k-SVOMTL is called an V_k-super vertex out-magic total (E_k-SVOM). Mutharasu, Kumar, and Mary Berrard [2609] characterized the digraphs that are V_k-SVOM and the unidirectional cycles and union of unidirectional cycles that are V_2-SVOM.

Let $D = (V, A)$ be a directed graph with p vertices and q arcs. For an integer $k \geq 1$ and $v \in V(D)$, let $w_k(v) = \sum_{e \in E_k(v)} f(e)$, where $E_k(v)$ is the set of all in-neighborhood arcs that are at distance at most k from v. The digraph D is said to be *E_k-regular* with regularity r, if $|E_k(e)| = r$ for some integer $r \geq 1$ and for all $e \in A(D)$, where $E_k(e) = E_k(u, v) = \{u \in V(D) : 1 \leq d(u, v) \leq k\}$. An E_k-super vertex in-magic total labeling (E_k-SVIMTL) is a bijection $f : V(D) \cup A(D) \to \{1, 2, \ldots, p + q\}$ with the property that $f(A(D)) = \{1, 2, \ldots, q\}$ and for each $v \in V(D)$, $f(v) + w_k(v) = M$ for some positive integer M. A digraph that admits an E_k-SVIMTL is called E_k-super vertex in-magic total (E_k-SVIMT). Mutharasu, Bernard, and Kumar [2604] obtained a necessary and sufficient condition for the existence of E_k-SVIMTL in digraphs and the magic constant for E_k-regular digraphs. They investigated the E_k-SVIMTL of unidirectional paths and star graphs and proved necessary and sufficient conditions are for the existence of a E_2-SVIMTL in unidirectional cycles and union of unidirectional cycles. In [2608] Mutharasu and Kumar gave a necessary and sufficient condition for the existence of E_k-SVIMTL in digraphs and the magic constant for E_k-regular digraphs. They also provided a necessary and sufficient conditions for the existence of a E_2-SVIMTL in unidirectional cycles and union of unidirectional cycles and investigated the existence of E_k-SVIMTL of unidirectional paths and stars.

In [2315] López, Muntaner-Batle and Rius-Font introduce the concept of *$\{H_i\}_{i \in I}$-super edge-magic decomposable* as follows: Let $G = (V, E)$ be any graph and let $\{H_i\}_{i \in I}$ be a set of graphs such that $G = \oplus_{i \in I} H_i$ (that is, G decomposes into the graphs in the set $\{H_i\}_{i \in I}$). Then we say that G is $\{H_i\}_{i \in I}$-super edge-magic decomposable if there is a bijection $\beta : V \to [1, |V|]$ such that for each $i \in I$ the subgraph H_i meets the following two requirements: (i) $\beta(V(H_i)) = [1, |V(H_i)|]$ and (ii) $\{\beta(a) + \beta(b) : ab \in E(H_i)\}$ is a set of consecutive integers. Such function β is called an *$\{H_i\}_{i \in I}$-super edge-magic labeling* of G.

When $H_i = H$ for every $i \in I$ we just use the notation H-super edge-magic decomposable labeling. Among their results are the following. If $G = (V, E)$ is a (p, q)-graph that is $\{H_1, H_2\}$-super edge-magic decomposable for a pair of graphs H_1 and H_2 and n is an even integer, then the cycle C_n is $(n/2)K_2$-super edge-magic decomposable if and only if $n \equiv 2 \pmod 4$. Let n be odd. Then for any super edge-magic tree T there exists a bipartite connected graph $G = G(T, n)$ such that G is (nT)-super edge-magic decomposable. Let G be a $\{H_i\}_{i\in I}$-super edge magic decomposable graph, where H_i is an acyclic digraph for each $i \in I$. Assume that $\overrightarrow{G}$ is any orientation of G and $h : E(\overrightarrow{G}) \to S_p$ is any function. Then und$(\overrightarrow{G} \otimes_h S_p)$ is $\{pH_i\}_{i\in I}$-super edge magic decomposable. In [579] Babujee and Jagadesh proved that the following graphs admit super edge-bimagic labelings: $(3, n)$-kites $(n \geq 2), P_n + N_2$ $(n \geq 3), P_3 \odot K_{1,2n}, P_2 \cup mK_1 + N_2$, and $P_2 + mK_1 (m \geq 2)$.

In [226] Amuthavalli and Sugapriya define a *reverse edge-bimagic* labeling of graph G as a bijection $\phi : V(G) \cup E(G) \to \{1, 2, 3, \ldots, |V(G)| + |E(G)|\}$ such that for each edge uw in $E(G)$ the value of $\phi(uw) - (\phi(u) + \phi(w))$ is one of two distinct constants. They proved that shadow graphs, splitting graphs, jellyfishs, and jewel graphs admit reverse super edge-bimagic labelings. Martini [2438] provided reverse edge-bimagic labelings of brooms and generalized butterflies. A reverse edge-bimagic labeling is said to be a *reverse super edge-bimagic* labeling if the labels of vertices are $1, 2, \ldots, |V(G)|$ and the labels of edges are $|V(G)| + 1, |V(G)| + 2, \ldots, |V(G)| + |E(G)|$. He investigated the existence of reverse super edge-bimagic labelings for gear graphs, dove tail graphs, and hibiscus graphs. A reverse super edge trimagic is defined analogously. In [2440] Martini and Susanti proved that dove tail graphs, dove tails with a pendant graph, and the corona product of dove tail graphs and null graphs admit reverse super edge trimagic labeling. As a corollary of the last result they have that if G is a 2-regular, (1-factor)-super edge-magic decomposable graph and $\overrightarrow{G}$ is any orientation of G and $h : E(\overrightarrow{G}) \to S_p$ is any function, then und$(\overrightarrow{G} \otimes_h S_p)$ is a 2-regular, (1-factor)-super edge-magic decomposable graph. Moreover, if we denote the 1-factor of G by F then pF is the 1-factor of und$(\overrightarrow{G} \otimes_h S_p)$. In [225] Amuthavalli and Sugapriya investigated the existence of reverse super edge-bimagic labelings of star related graphs, jellyfishs, and jewel graphs.

They pose the following two open questions: Fix $p \in \mathbb{N}$. Find the maximum $r \in \mathbb{N}$ such that there is an r-regular graph of order p which is $(p/2)K_2$-super edge-magic decomposable: and characterize the set of 2-regular graphs of order n, $n \equiv 2 \pmod 4$, such that each component has even order and admits an $(n/2)K_2$-super edge-magic decomposition. In connection to open question 1 they prove: For all $r \in \mathbb{N}$, there is $n \in \mathbb{N}$ such that there exists a k-regular bipartite graph $B(n)$, with $k > r$ and $|V(B(n))| = 2 \cdot 3^n$, such that $B(n)$ is $(3^n K_2)$-super edge-magic decomposable.

Hendy, Sugeng, Salman, and Ayunda [1436] provided a sufficient condition for $\overline{C_n}[\overline{K_m}]$ to have a $P_t[\overline{K_m}]$-magic decompositions, where $n > 3,\ m > 1$, and $t = 3,\ 4,\ n - 2$.

An H-magic labeling in an H-decomposable graph G is a bijection $f\, V(G) \cup E(G) \to \{1, 2, \ldots, p + q\}$ such that, for every copy H in the decomposition, $\sum_{v\in V(H)} f(v) + \sum_{e\in E(H)} f(e)$ is constant. The function f is said to be an H-V-super magic labeling if $f(V(G)) = \{1, 2, \ldots, p\}$. In [2597] Murugan and Chandra Kumar find the magic con-

stant for H-factorable graphs that are H-V-super magic. Also, they give a necessary and sufficient condition for an H-factorable graph to be H-V-super magic and characterize the even regular graphs with a 2-factor-V-super magic labeling. In [2598] Murugan and Chandra Kumar determined the magic constant for H-factorable graphs that are H-V-super magic and gave a necessary and sufficient condition for an even regular graph to be 2-factor-V-super magic.

For a graph $G(V, E)$ with p vertices and q edges Murugan and Kumar [2599] say that G is $H-E$-*super magic graceful* if there is a surjection μ from $V(G)\cup E(G)$ to $\{1, 2, \ldots, p+q\}$ that satisfies the conditions that $\mu(E(G)) = \{1, 2, \ldots, q\}$ and the difference of the sum of all $\mu(v)$ over all vertices v in any subgraph H' of G isomorphic to H and the sum of all $\mu(e)$ over all edges e in H' isomorphic to H is a constant M. They investigated $C_m - E$-super magic graceful labeling of generalized book graphs. In [2063] Kumar, Mohankumar, Lawrance, and Bhuvaneshwari investigated $H - E$-super magic graceful labelings of fans, graphs obtained by joining a star with one isolated vertex, books and grids. [2599] new [2063] new

A bipartite graph G with partite sets X_1 and X_2 is called *consecutively super edge-magic* if there exists a bijective function $f : V(G)\cup E(G) \to \{1, 2, \ldots, |V(G)| + |E(G)|\}$ such that $f(X_1) = \{1, 2, \ldots, |X_1|\}$, $f(X_2) = \{|X_1| + 1, |X_1| + 2, \ldots, |V(G)|\}$ and $f(u) + f(v) + f(uv)$ is a constant for each $uv \in E(G)$. In [1490] Ichishima, Muntaner-Batle, and Oshima investigated for which bipartite graphs is it possible to add a finite number of isolated vertices so that the resulting graph is consecutively super edge-magic. If it is possible for a bipartite graph G, then they say that the minimum such number $\mu_c(G)$ of isolated vertices is the *consecutively super edge-magic deficiency* of G; otherwise, it is $+\infty$. Thus, the consecutively super edge-magic deficiency of a graph G is a measure of how close G is to being consecutively super edge-magic. They also include a detailed discussion of other concepts that are closely related to the consecutively super edge-magic deficiency.

In [1493] Ichishima, Muntaner-Batle, and Oshima prove that $\alpha(G) = \mu_c(G)+|V(G)|+1$. Thus a tree has a consecutively super edge-magic if and only if it has an α-valuation. They explore the relation between super edge-magic labelings and graceful labelings of trees.

In [1506] Ichishima, Oshima, and Takamashi introduce the notion of strength sum of a non-empty graph as follows. The *strength sum* $\mathrm{strs}_f(G)$ of a numbering $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ is defined by $\mathrm{strs}(G) = \min\{\mathrm{strs}_f(G) \mid f \text{ is a numbering of } G\}$, where $\mathrm{strs}_f(G) = \sum_{uv\in E(G)}(f(u) + f(v))$. A numbering f of a graph G for which $\mathrm{strs}_f(G) = \mathrm{strs}(G)$ is called a *strength sum labeling of* G. They also discuss relations among invariants on super edge-magic graphs and their its strength sums. Ichishima, Muntaner-Batle, and Oshima [1498] proved that for every $k \in [1, n - 1]$, there exists a graph G of order n satisfying $\delta(G) = k$ and $\mathrm{str}(G) = n + k$, where $\delta(G)$ denotes the minimum degree of G.

Avadayappan, Jeyanthi, and Vasuki [321] define the *super magic strength* of a graph G as $sm(G) = \min\{s(L)\}$ where L runs over all super edge-magic labelings of G. They use the notation $< K_{1,n} : 2 >$ for the tree obtained from the bistar $B_{n,n}$ (the graph obtained by joining the center vertices of two copies of $K_{1,n}$ with an edge) by subdividing the edge

joining the two stars. They prove: $sm(P_{2n}) = 5n + 1$; $sm(P_{2n+1}) = 5n + 3$; $sm(< K_{1,n} : 2 >) = 4n + 9$; $sm(B_{n,n}) = 5n + 6$; $sm((2n + 1)P_2) = 9n + 6$; $sm(C_{2n+1}) = 5n+4$; $emt(C_{2n}) = 5n+2$; $sm(K_{1,n}) = 2n+4$; and $sm(P_n^2) = 3n$. Note that in each case the super magic strength of the graph is the same as its magic strength. In [163] Akka and Warad determined the super magic strength of paths, cycles, wheels, stars, bistars, $P_n^2, < K_{1,n} : 2 >$ (the graph obtained by joining the centers of two copies of $K_{1,n}$ by a path of length 2), and $(2n + 1)P_2$.

Santhosh and Singh [3306] proved that $C_n \odot P_2$ and $C_n \odot P_3$ are super edge-magic for all odd $n \geq 3$ and prove for odd $n \geq 3$, $sm(C_n \odot P_2) = (15n + 3)/2$ and $(20n + 3) \leq sm(C_n \odot P_3) \leq (21n + 3)/2$. Gray [1330] proves that $C_3 \cup C_n$ is super edge-magic if and only if $n \geq 6$ and $C_4 \cup C_n$ is super edge-magic if and only if $n \geq 5$. His computer search shows that $C_5 \cup 2C_3$ does not have a super edge-magic labeling.

In [4120] Wallis posed the problem of investigating the edge-magic properties of C_n with the path of length t attached to one vertex. Kim and Park [1963] call such a graph an (n, t)-*kite*. They prove that an $(n, 1)$-kite is super edge-magic if and only if n is odd and an $(n, 3)$-kite is super edge-magic if and only if n is odd and at least 5. Park, Choi, and Bae [2776] show that $(n, 2)$-kite is super edge-magic if and only if n is even. Wallis [4120] also posed the problem of determining when $K_2 \cup C_n$ is super edge-magic. In [2776] and [1963] Park et al. prove that $K_2 \cup C_n$ is super edge-magic if and only if n is even. Kim and Park [1963] show that the graph obtained by attaching a pendent edge to a vertex of degree one of a star is super-edge magic and that a super edge-magic graph with edge magic constant k and q edges satisfies $q \leq 2k/3 - 3$.

Lee and Kong [2154] use $\mathrm{St}(a_1, a_2, \ldots, a_n)$ to denote the disjoint union of the n stars $\mathrm{St}(a_1)$, $\mathrm{St}(a_2)$, ... , $\mathrm{St}(a_n)$. They prove the following graphs are super edge-magic: $\mathrm{St}(m, n)$ where $n \equiv 0 \bmod(m+1)$; $\mathrm{St}(1, 1, n)$; $\mathrm{St}(1, 2, n)$; $\mathrm{St}(1, n, n)$; $\mathrm{St}(2, 2, n)$; $\mathrm{St}(2, 3, n)$; $\mathrm{St}(1, 1, 2, n)$ $(n \geq 2)$; $\mathrm{St}(1, 1, 3, n)$; $\mathrm{St}(1, 2, 2, n)$; and $\mathrm{St}(2, 2, 2, n)$. They conjecture that $\mathrm{St}(a_1, a_2, \ldots, a_n)$ is super edge-magic when $n > 1$ is odd. Gao and Fan [1194] proved that $\mathrm{St}(1, m, n)$; $\mathrm{St}(3, m, m + 1)$; and $\mathrm{St}(n, n + 1, n + 2)$ are super edge-magic, and under certain conditions $\mathrm{St}(a_1, a_2, \ldots, a_{2n+1})$, $\mathrm{St}(a_1, a_2, \ldots, a_{4n+1})$, and $\mathrm{St}(a_1, a_2, \ldots, a_{4n+3})$ are also super edge magic.

In [2381] MacDougall and Wallis investigate the existence of super edge-magic labelings of cycles with a chord. They use C_v^t to denote the graph obtained from C_v by joining two vertices that are distance t apart in C_v. They prove: C_{4m+1}^t $(m \geq 3)$ has a super edge-magic labeling for every t except $4m - 4$ and $4m - 8$; C_{4m}^t $(m \geq 3)$ has a super edge-magic labeling when $t \equiv 2 \bmod 4$; and that C_{4m+2}^t $(m > 1)$ has a super edge-magic labeling for all odd t other than 5, and for $t = 2$ and 6. They pose the problem of what values of t does C_{2n}^t have a super edge-magic labeling.

Enomoto, Masuda, and Nakamigawa [1038] have proved that every graph can be embedded in a connected super edge-magic graph as an induced subgraph. Slamin, Bača, Lin, Miller, Simanjuntak [3629] proved that the friendship graph consisting of n triangles is super edge-magic if and only if n is 3, 4, 5, or 7. Fukuchi proved [1160] the generalized Petersen graph $P(n, 2)$ (see §2.7 for the definition) is super edge-magic if n is odd and at least 3 while Xu, Yang, Xi, Haque, and Shen [4234] showed that $P(n, 3)$ is super

edge-magic for odd n is odd and at least 5. Baskoro and Ngurah [592] showed that nP_3 is super edge-magic for $n \geq 4$ and n even.

Hegde and Shetty [1425] showed that a graph is super edge-magic if and only if it is strongly k-indexable (see §4.1 for the definition). Figueroa-Centeno, Ichishima, and Muntaner-Batle [1087] proved that a graph is super edge-magic if and only if it is strongly 1-harmonious and that every super edge-magic graph is cordial. They also proved that P_n^2 and $K_2 \times C_{2n+1}$ are super edge-magic. In [1088] Figueroa-Centeno et al. show that the following graphs are super edge-magic: $P_3 \cup kP_2$ for all k; kP_n when k is odd; $k(P_2 \cup P_n)$ when k is odd and $n = 3$ or $n = 4$; and fans F_n if and only if $n \leq 6$. They conjecture that kP_2 is not super edge-magic when k is even. This conjecture has been proved by Z. Chen [799] who showed that kP_2 is super edge-magic if and only if k is odd. Figueroa-Centeno et al. proved that the book B_n is not super edge-magic when $n \equiv 1, 3, 7 \pmod 8$ and when $n = 4$. They proved that B_n is super edge-magic for $n = 2$ and 5 and conjectured that for every $n \geq 5$, B_n is super edge-magic if and only if n is even or $n \equiv 5 \pmod 8$. Yuansheng, Yue, Xirong, and Xinhong [4298] proved this conjecture for the case that n is even. They prove that every tree with an α-labeling is super edge-magic. Yokomura (see [1037]) has shown that $P_{2m+1} \times P_2$ and $C_{2m+1} \times P_m$ are super edge-magic (see also [1087]). In [1089], Figueroa-Centeno et al. proved that if G is a (super) edge-magic 2-regular graph, then $G \odot \overline{K}_n$ is (super) edge-magic and that $C_m \odot \overline{K}_n$ is super edge-magic. Fukuchi [1159] shows how to recursively create super edge-magic trees from certain kinds of existing super edge-magic trees. Ngurah, Baskoro, and Simanjuntak [2704] provide a method for constructing new (super) edge-magic graphs from existing ones. One of their results is that if G has an edge-magic total labeling and G has order p and size p or $p-1$, then $G \odot nK_1$ has an edge-magic total labeling.

Ichishima, Muntaner-Batle, Oshima [1488] enlarged the classes of super edge-magic 2-regular graphs by presenting some constructions that generate large classes of super edge-magic 2-regular graphs from previously known super edge-magic 2-regular graphs or pseudo super edge-magic graphs. By virtue of known relationships among other classes of labelings the 2-regular graphs obtained from their constructions are also harmonious, sequential, felicitous and equitable. Their results add credence to the conjecture of Holden et al. [1450] that all 2-regular graphs of odd order with the exceptions of $C_3 \cup C_4$, $3C_3 \cup C_4$, and $2C_3 \cup C_5$ possess a strong vertex-magic total labeling, which is equivalent to super edge-magic labelings for 2-regular graphs. For a 2-regular graph G with $2m+1$ vertices that has a strong vertex-magic total labeling McQuillan and McQuillan [2464] proved that $G \cup 2mC_3$, $G \cup (2m+2)C_3$, $G \cup mC_8$ and $G \cup (m+1)C_8$ also have a strong vertex-magic total labeling.

Lee and Lee [2157] investigate the existence of total edge-magic labelings and super edge-magic labelings of unicylic graphs. They obtain a variety of positive and negative results and conjecture that all unicyclic are edge-magic total.

Shiu and Lee [3539] investigated edge labelings of multigraphs. Given a multigraph G with q edges they call a bijection from the set of edges of G to $\{1, 2, \ldots, q\}$ with the property that for each vertex v the sum of all edge labels incident to v is a constant independent of v a *supermagic* labeling of G. They use $K_2[n]$ to denote the multigraph

consisting of n edges joining 2 vertices and $mK_2[n]$ to denote the disjoint union of m copies of $K_2[n]$. They prove that for m and n at least 2, $mK_2[n]$ is supermagic if and only if n is even or if both m and n are odd.

In 1970 Kotzig and Rosa [2017] defined the *edge-magic deficiency*, $\mu(G)$, of a graph G as the minimum n such that $G \cup nK_1$ is edge-magic total. If no such n exists they define $\mu(G) = \infty$. In 1999 Figueroa-Centeno, Ichishima, and Muntaner-Batle [1093] extended this notion to *super edge-magic deficiency*, $\mu_s(G)$, is the analogous way. They prove the following: $\mu_s(nK_2) = \mu(nK_2) = n - 1 \pmod 2$; $\mu_s(C_n) = 0$ if n is odd; $\mu_s(C_n) = 1$ if $n \equiv 0 \pmod 4$; $\mu_s(C_n) = \infty$ if $n \equiv 2 \pmod 4$; $\mu_s(K_n) = \infty$ if and only if $n \geq 5$; $\mu_s(K_{m,n}) \leq (m-1)(n-1)$; $\mu_s(K_{2,n}) = n-1$; and $\mu_s(F)$ is finite for all forests F. They also prove that if a graph G has q edges with $q/2$ odd, and every vertex is even, then $\mu_s(G) = \infty$ and conjecture that $\mu_s(K_{m,n}) \leq (m-1)(n-1)$. This conjecture was proved for $m = 3, 4$, and 5 by Hegde, Shetty, and Shankaran [1426] using the notion of strongly k-indexable labelings. Baig, Baskoro, and Semaničová-Feňovčíková [445] investigated the super edge-magic deficiency of a forest consisting of stars. Ngurah investigates the (super) edge-magic deficiency of chain graphs in [2699] and Ngurah and Adiwijaya does the same in [2698].

For an (n, t)-kite graph (a path of length t attached to a vertex of an n-cycle) G Ahmad, Siddiqui, Nadeem, and Imran [138] proved the following: for odd $n \geq 5$ and even $t \geq 4, \mu_s(G) = 1$; for odd $n \geq 5,\ t \geq 5,\ t \neq 11$, and $t \equiv 3, 7 \pmod 8$, $\mu_s(G) \leq 1$; for $n \geq 10,\ n \equiv 2 \pmod 4$ and $t = 4,\ \mu_s(G) \leq 1$; and for $t = 5,\ \mu_s(G) = 1$.

In [440] Baig, Ahmad, Baskoro, and Simanjuntak provide an upper bound for the super edge-magic deficiency of a forest formed by paths, stars, combs, banana trees, and subdivisions of $K_{1,3}$. Baig, Baskoro, and Semaničová-Feňovčíková [441] investigate the super edge-magic deficiency of forests consisting of stars. Among their results are: a forest consisting of $k \geq 3$ stars has super edge-magic deficiency at most $k - 2$; for every positive integer n a forest consisting of 4 stars with exactly 1, n, n, and $n + 2$ leaves has a super edge-magic total labeling; for every positive integer n a forest consisting of 4 stars with exactly 1, $n + 5, 2n + 6$, and $n + 1$ leaves has a super edge-magic total labeling; and for every positive integers n and k a forest consisting of k identical stars has super edge-magic deficiency at most 1 when k is even and deficiency 0 when k is odd. In [129] Ahmad, Javaid, Nadeem, and Hasni investigate the super edge-magic deficiency of some families of graphs related to ladder graphs. Kanwal, Javed, and Riasat [1899] give super edge-magic total labelings and the deficiency for forests consisting of extended w-trees, combs, stars and paths. In [133] Ahmad, Nadeem, and Gupta provided bounds for the super edge-magic deficiency of some Toeplitz graphs.

The *generalized Jahangir* graph $J_{n,m}$ for $m \geq 3$ is a graph on $nm+1$ vertices, consisting of a cycle C_{nm} with one additional vertex that is adjacent to m vertices of C_{nm} at distance n to each other on C_{nm}. In [442] Baig, Imran, Javaid, and Semaničová-Feňovčiková study the super edge-magic deficiencies of the web graph $Wb_{n,m}$, the generalized Jahangir graph $J_{2,n}$, crown products $L_n \odot K_1,\ K_4 \odot nK_1$, and gave the exact value of super edge-magic deficiency for one class of lobsters.

In [1092] Figueroa-Centeno, Ichishima, and Muntaner-Batle proved that $\mu_s(P_m \cup$

$K_{1,n}) = 1$ if $m = 2$ and n is odd, or $m = 3$ and n is not congruent to 0 mod 3, whereas in all other cases $\mu_s(P_m \cup K_{1,n}) = 0$. They also proved that $\mu_s(2K_{1,n}) = 1$ when n is odd and $\mu_s(2K_{1,n}) \leq 1$ when n is even. They conjecture that $\mu_s(2K_{1,n}) = 1$ in all cases. Other results in [1092] are: $\mu_s(P_m \cup P_n) = 1$ when $(m, n) = (2, 2)$ or $(3, 3)$ and $\mu_s(P_m \cup P_n) = 0$ in all other cases; $\mu_s(K_{1,m} \cup K_{1,n}) = 0$ when mn is even and $\mu_s(K_{1,m} \cup K_{1,n}) = 1$ when mn is odd; $\mu(P_m \cup K_{1,n}) = 1$ when $m = 2$ and n is odd and $\mu(P_m \cup K_{1,n}) = 0$ in all other cases; $\mu(P_m \cup P_n) = 1$ when $(m, n) = (2, 2)$ and $\mu(P_m \cup P_n) = 0$ in all other cases; $\mu_s(2C_n) = 1$ when n is even and ∞ when n is odd; $\mu_s(3C_n) = 0$ when n is odd; $\mu_s(3C_n) = 1$ when $n \equiv 0 \pmod 4$; $\mu_s(3C_n) = \infty$ when $n \equiv 2 \pmod 4$; and $\mu_s(4C_n) = 1$ when $n \equiv 0 \pmod 4$. They conjecture the following: $\mu_s(mC_n) = 0$ when mn is odd; $\mu_s(mC_n) = 1$ when $mn \equiv 0 \pmod 4$; $\mu_s(mC_n) = \infty$ when $mn \equiv 2 \pmod 4$; $\mu_s(2K_{1,n}) = 1$; and if F is a forest with two components, then $\mu(F) \leq 1$ and $\mu_s(F) \leq 1$. Santhosh and Singh [3303] proved: for n odd at least 3, $\mu_s(K_2 \odot C_n) \leq (n-3)/2$; for $n > 1,\ 1 \leq \mu_s(P_n[P_2]) = \lceil (n-1)/2 \rceil$; and for $n \geq 1,\ 1 \leq \mu_s(P_n \times K_4) \leq n$.

Ichishima and Oshima [1515] prove the following: if a graph $G(V, E)$ has an α-labeling and no isolated vertices, then $\mu_s(G) \leq |E| - |V| + 1$; if a graph $G(V, E)$ has an α-labeling, is not sequential, and has no isolated vertices, then $\mu_s(G) = |E| - |V| + 1$; and, if m is even, then $\mu_s(mK_{1,n}) \leq 1$. As corollaries of the last result they have: $\mu_s(2K_{1,n}) = 1$; when $m \equiv 2 \pmod 4$ and n is odd, $\mu_s(mK_{1,n}) = 1$; $\mu_s(mK_{1,3}) = 0$ when $m \equiv 4 \pmod 8$ or m is odd; $\mu_s(mK_{1,3}) = 1$ when $m \equiv 2 \pmod 4$; $\mu_s(mK_{2,2}) = 1$; for $n \geq 4,\ (n-4)2^{n-2} + 3 \leq \mu_s(Q_n) \leq (n-2)2^{n-1} - 4$; and for $s \geq 2$ and $t \geq 2,\ \mu_s(mK_{s,t}) \leq m(st - s - t) + 1$. They conjecture that for $s \geq 2$ and $t \geq 2,\ \mu_s(mK_{s,t}) = m(st - s - t) + 1$ and pose as a problem determining the exact value of $\mu_s(Q_n)$.

Ichishima and Oshima [1513] determined the super edge-magic deficiency of graphs of the form $C_m \cup C_n$ for m and n even and for arbitrary n when $m = 3, 4, 5$, and 7. They state a conjecture for the super edge-magic deficiency of $C_m \cup C_n$ in the general case. Afzal and Aslam [82] investigate the super edge-magic deficiency of various disjoint unions of $K_{2,n}$ with stars, paths and disjoint union of paths. The *join product* of two graphs is their graph union with additional edges that connect all vertices of the first graph to each vertex of the second graph. In [2708] Ngurah and Simanjuntak investigate the super edge-magic deficiencies of a wheel minus an edge and join products of a path, a star, and a cycle with isolated vertices. They also show that the join product of a super edge-magic graph with isolated vertices has finite super edge-magic deficiency.

A *block* of a graph is a maximal subgraph with no cut-vertex. The *block-cut-vertex graph* of a graph G is a graph H whose vertices are the blocks and cut-vertices in G; two vertices are adjacent in H if and only if one vertex is a block in G and the other is a cut-vertex in G belonging to the block. A *chain graph* is a graph with blocks $B_1, B_2, B_3, \ldots, B_k$ such that for every i, B_i and B_{i+1} have a common vertex in such a way that the block-cut-vertex graph is a path. The chain graph with k blocks where each block is identical and isomorphic to the complete graph K_n is called the *kK_n-path*.

Ngurah, Baskoro, and Simanjuntak [2703] investigate the exact values of $\mu_s(kK_n\text{-path})$ when $n = 2$ or 4 for all values of k and when $n = 3$ for $k \equiv 0, 1, 2 \pmod 4$, and give an upper bound for $k \equiv 3 \pmod 4$. They determine the exact super edge-magic deficiencies

for fans, double fans, wheels of small order and provide upper and lower bounds for the general case as well as bounds for some complete partite graphs. They also include some open problems. Lee and Wang [2218] show that various chain graphs with blocks that are complete graphs are super edge-magic. In [128] investigate the super edge-magic deficiency of some kites and $C_n \cup K_2$.

Figueroa-Centeno and Ichishima [1085] introduce the notion of the *sequential number* $\sigma(G)$ of a graph G without isolated vertices to be either the smallest positive integer n for which it is possible to label the vertices of G with distinct elements from the set $\{0, 1, \ldots, n\}$ in such a way that each $uv \in E(G)$ is labeled $f(u) + f(v)$ and the resulting edge labels are $|E(G)|$ consecutive integers or $+\infty$ if there exists no such integer n. They prove that $\sigma(G) = \mu_s(G) + |V(G)| - 1$ for any graph G without isolated vertices, and $\sigma(K_{m,n}) = mn$, which settles the conjecture of Figueroa-Centeno, Ichishima, and Muntaner-Batle [1093] that $\mu_s(K_{m,n}) = (m-1)(n-1)$.

In [1485] Ichishima and Muntaner-Batle define the *strong sequential number* $\sigma_s(G)$ of G as the smallest positive integer n for which there exists an injective function from the vertices of G to $[0, n]$ such that when each edge uv is labeled $f(u)+f(v)$, the resulting set of edge labels is $[c, c+q-1]$ for some positive integer c and there exists an integer λ so that $\min\{f(u), f(v)\} \leq \lambda < \max\{f(u), f(v)\}$ for all edges uv. Note that for G to have finite $\sigma_s(G)$, it must be bipartite. They prove for a graph G of order p, $\sigma(G) = \mu_s(G) + p - 1$. From this it follows that the problems of determining the sequential number and super edge-magic deficiency are equivalent and that for any graph G, $\sigma(G)$ is finite if and only if $\mu_s(G)$ is finite. They also introduced the following parameter as a measure of how close a graph G is to having an α-labeling. The *alpha-number* $\alpha(G)$ of a graph G with q edges is the smallest positive integer n for which there exists an injective function $f : V(G) \to [0, n]$ such that when each edge uv is labeled $|f(u) - f(v)|$ the resulting set of edge labels is $[c, c+q-1]$ for some positive integer c, and there exists an integer λ so that $\min\{f(u), f(v)\} \leq \lambda < \max\{f(u), f(v)\}$ for each $uv \in E(G)$. If no such n exists the alpha-number of G is defined to be $+\infty$. Since a graph that admits an α-labeling is necessarily bipartite, graphs with finite $\alpha(G)$ are bipartite.

Ichishima and Muntaner-Batle [1485] prove: if every vertex of graph G has even degree and $|E(G)| \equiv 2 \pmod 4$, then $\sigma(G) = \sigma_s(G) = +\infty$; for every graph G of order p, $\sigma_s(G) = \mu_c(G) + p - 1$; and if G is a super edge-magic graph with at least one edge, then the graph $G + nK_1$ is sequential for every positive integer n. As corollaries they have: for every graph $\sigma_s(G) = \alpha(G)$; a graph G has an α-labeling if and only if $\sigma_s(G) = |E(G)|$; and if a graph G of order p and size $q \geq 1$ has a super edge-magic labeling f with $s = \min\{f(u) + f(v) \ : \ uv \in E(G)\}$, then $\sigma(G + nK_1) \leq s + q + (n-1)p - 2$; if G is a graph of order p and size $q \geq 1$ and G has a super edge-magic labeling f with $s = \min\{f(u) + f(v) : uv \in E(G)\}$, then $\mu_s(G + nK_1) \leq s + q + (n-2)(p-1) - 3$; and if G is a super edge-magic graph with at least one edge, then the graph $G + nK_1$ is harmonious and felicitous for any positive integer n.

For a graph G order p and size q Ichishima, Muntaner-Batle, and Oshima [1499] prove the following: if $q = p - 1$ and $\beta_s(G) = p - 1$, then $\beta(G \odot nK_1) = \beta_s(G \odot nK_1) = (n+1)p - 1$ for every positive integer n; if $q > p - 1$ and $\beta_s(G) = q$, then there exists

a supergraph H of G such that $\beta(H \odot nK_1) = \beta_s(H \odot nK_1) = (n+1)(q+1) - 1$ for every positive integer n; if G has a subgraph H such that $\beta_s(H) = q < p - 1$, then $\beta(H \odot nK_1) = \beta_s(H \odot nK_1) = (n+1)(q+1) - 1$ for every positive integer n; and if G has a subgraph H such that $\beta_s(H) = q < p - k(H')$, where H' is a subgraph of H without isolated vertices, then $\beta(H \odot nK_1) = \beta_s(H \odot nK_1) = (n+1)(q+1) - 1$ for every positive integer n.

As the concept of super magic strength is effectively defined only for super edge-magic graphs, Ichishima, Muntaner-Batle, and Oshima [1494] generalize it for any nonempty graph as follows. A *numbering* f of a graph G of order p is a labeling that assigns distinct elements of the set $[1, p]$ to the vertices of G, where each edge uv of G is labeled $f(u)+f(v)$. The *strength*, $\mathrm{str}_f(G)$, of a numbering $f : V(G) \to [1, p]$ of G is defined by $\mathrm{str}_f(G) = \max\{f(u) + f(v) \,|uv \in E(G)\}$, that is, $\mathrm{str}_f(G)$ is the maximum edge label of G, and the *strength*, $\mathrm{str}(G)$, of a graph G itself is $\mathrm{str}(G) = \min\{\mathrm{str}_f(G) \,|f \text{ is a numbering of } G\}$. A numbering f of a graph G for which $\mathrm{str}_f(G) = \mathrm{str}(G)$ is called a *strength labeling* of G. If G is an empty graph, then $\mathrm{str}(G)$ is undefined. For a graph G of order p they prove the following: if G has order at least 3 and contains a path of order k ($k \in [2, p-1]$) as an induced subgraph, then $\mathrm{str}(G) \leq 2p - (k-1)$; if $\Delta(G) + 2 \leq \mathrm{str}(G) \leq 2p - 1$; and if $p + m + \min\{p, \delta(G) + m\} \leq \mathrm{str}(G + mK_1) \leq \mathrm{str}(G) + 2m$ for every positive integer m. They determine the exact strength for many basic families of graphs such as paths, cycles complete graphs, ladders, books, and hypercubes. They conclude with six problems and a conjecture.

In [3860] Takahashi, Ichishima, and Muntaner-Batle provide sharp lower bounds for the strength of a graph in terms of its domination number as well as its (edge) covering and (edge) independence numbers. They also provide a necessary and sufficient condition for the strength of a graph to attain an earlier bound in terms of its subgraph structure. In addition, they establish a sharp lower bound for the domination number of a graph under certain conditions.

In [1495] Ichishima, Muntaner-Batle, and Oshima determined the strength of caterpillars and complete n-ary k-level trees. The strength $\mathrm{str}(G)$ is also given for graphs G obtained by taking the corona of certain graphs and arbitrary number of isolated vertices. They further proved if G is a graph of order p with $\delta(G) \geq 1$ and $\mathrm{str}(G) = p + \delta(G)$, then $\mathrm{str}(G \odot nK_1) = (n+1)p + 1$. for every positive integer n. In [1497] Ichishima, Muntaner-Batle, Oshima show that for every $k \in [1, n-1]$, there exists a graph G of order n satisfying $\delta(G) = k$ and $\mathrm{str}(G) = n + k$, where $\delta(G)$ denotes the minimum degree of G. In [1501] Ichishima, Muntaner-Batle, and Oshima investigate minimum degree conditions for the strength of graphs. They determine certain degree sequences of graphs that naturally arise and proof that these degree sequences determine a unique graph realization. In addition, they establish a parallel bandwidth result to the one on strength of graphs and enlarge the class of k-stable properties known so far.

The following result Ichishima, Muntaner-Batle, and Oshima in [1490] shows the connection between the alpha-number of a graph and its consecutively super edge-magic deficiency. For every graph G of order p, $\alpha(G) = \mu_c(G) + p - 1$. This result shows that the problems of determining the alpha-number and consecutively super edge-magic

deficiency are equivalent.

In [2709] Ngurah and Simanjuntak proved that if G is a cycle-free graph with minimum degree one and $\mu_s(G + K_1) = 0$ then G is either a tree or a forest. They also prove: the join product of some classes of trees and forests with an isolated vertex has zero super edge-magic deficiency; for all but one tree of order at most 6, their join product with an isolated vertex has zero super edge-magic deficiency. For trees T of order at least 7 they proved that if $\mu_s(T+K_1) = 0$, then either $2K_{1,3}$ or $K_3 \cup K_{1,3}$ is a subgraph of $T+K_1$. For the super edge-magic deficiency of the join product of a tree T of order at least 2 with $m \geq 2$ isolated vertices, they showed that $\mu_s(T + mK_1) = 0$ if and only if $T = P_2$. For a tree $T \neq P_2$, they proved $\mu_s(T + mK_1) \geq \left\lfloor \frac{(m-1)(|V(T)-2|+1)}{2} \right\rfloor$ .They also present results for the super edge-magic deficiency of some chain graphs.

Z. Chen [799] has proved: the join of K_1 with any subgraph of a star is super edge-magic; the join of two nontrivial graphs is super edge-magic if and only if at least one of them has exactly two vertices and their union has exactly one edge; and if a k-regular graph is super edge-magic, then $k \leq 3$. Chen also obtained the following: there is a connected super edge-magic graph with p vertices and q edges if and only if $p - 1 \leq q \leq 2p - 3$; there is a connected 3-regular super edge-magic graph with p vertices if and only if $p \equiv 2 \pmod 4$; and if G is a k-regular edge-magic total graph with p vertices and q edges then $(p+q)(1+p+q) \equiv 0 \pmod{2d}$ where $d = \gcd(k-1, q)$. As a corollary of the last result, Chen observes that $nK_2 + nK_2$ is not edge-magic total.

Another labeling that has been called "edge-magic" was introduced by Lee, Seah, and Tan in 1992 [2196]. They defined a graph $G = (V, E)$ to be *edge-magic* if there exists a bijection $f\colon E \to \{1, 2, \ldots, |E|\}$ such that the induced mapping $f^+\colon V \to N$ defined by $f^+(u) = \sum_{(u,v)\in E} f(u, v) \pmod{|V|}$ is a constant map. Lee (see [2184]) conjectured that a cubic graph with p vertices is edge-magic if and only if $p \equiv 2 \pmod 4$. Lee, Pigg, and Cox [2184] verified this conjecture for prisms and several other classes of cubic graphs. They also show that $C_n \times K_2$ is edge-magic if and only if n is odd. Shiu and Lee [3539] showed that the conjecture is not true for multigraphs and disconnected graphs. In [3539] Lee's conjecture was modified by restricting it to simple connected cubic graphs. A computer search by Lee, Wang, and Wen [2221] showed that the new conjecture was false for a graph of order 10. Using different methods, Shiu [3517] and Lee, Su, and Wang [2207] gave proofs that it is was false.

Lee, Seah, and Tan [2196] establish that a necessary condition for a multigraph with p vertices and q edges to be edge-magic is that p divides $q(q+1)$ and they exhibit several new classes of cubic edge-magic graphs. They also proved: $K_{n,n}$ $(n \geq 3)$ is edge-magic and K_n is edge-magic for $n \equiv 1, 2 \pmod 4$ and for $n \equiv 3 \pmod 4$ $(n \geq 7)$. Lee, Seah, and Tan further proved that following graphs are not edge-magic: all trees except P_2; all unicyclic graphs; and K_n where $n \equiv 0 \pmod 4$. Schaffer and Lee [3325] have proved that $C_m \times C_n$ is always edge-magic. Lee, Tong, and Seah [2213] have conjectured that the total graph of a (p, p)-graph is edge-magic if and only if p is odd. They prove this conjecture for cycles. Lee, Kitagaki, Young, and Kocay [2153] proved that a maximal outerplanar graph with p vertices is edge-magic if and only if $p = 6$. Shiu [3516] used matrices with special properties to prove that the composition of P_n with $\overline{K_n}$ and the composition of

P_n with $\overline{K_{kn}}$ where kn is odd and n is at least 3 have edge-magic labelings. Boonklurb, Narissayaporn, and Singhun [688] show that under some conditions the m-node k-uniform hyperpaths and m-node k-uniform hypercycles are super edge-magic.

For a (p,q)-graph a bijection f from $V(G) \cup E(G)$ to $\{1,2,\ldots,p+q\}$ such that for each edge $xy \in E(G)$ the value of $f(x)+f(xy)+f(y)$ is either k_1, k_2, or k_3 is said to be an *edge trimagic total labeling.* Regees and Jayasekaran [3159] prove that $C_m \times P_n$, the generalized web graph, and the generalized web graph without a center are super edge trimagic total graphs. In [3155] proved that the star type graphs $P_3 \odot \overline{K_n}$, $B_{m,n}$, $\langle B_{m,n} : 2\rangle$ and $\langle K_{1,n}\, 3\rangle$ admits edge trimagic total labelings and super edge trimagic total labelings.

A *triangular belt* is obtained from $P_n \times P_2$ with vertices $u_1, u_2, \ldots u_n$ and $v_1, v_2, \ldots, v_{n_1}$ by adding an edge $\{(u_k v_{k+1}) \mid k = 1,2,\ldots,n-1\}$. The braid graph is obtained from a pair of paths P_n and P'_n with vertices $u_1, u_2, \ldots u_n$ and $v_1, v_2, \ldots u_n$ obtained by joining the ith vertex of P_n and the $(i+1)$th vertex with P'_n and joining the ith vertex of P'_n and the $(i+2)$th vertex with P_n for $1 \le i \le n-2$. A *semi Jahangir* graph is connected graph with a vertex set $\{u, u_k \mid 1 \le k \le n\} \cup \{s_k \mid 1 \le k \le n-1\}$ and edge set $\{u_k s_k \mid 1 \le k \le n-1\} \cup \{s_k u_{k+1} \mid 1 \le k \le n-1\} \cup \{u_k u_i \mid 1 \le k \le n\}$ A graph obtained from a graph G replacing each edge e_i by an H-graph in such a way that the ends of e_i are merged with a pendant vertex in P_2 and pendant vertex in P'_2 is called H *super subdivision* of G is denoted by $HSS(G)$, where the H-graph is a tree on 6 vertices in which exactly two vertices have degree 3. Vaghela and Parmar [3944] prove that the H-graph of a path, alternate triangular belt graph, braid graph, semi Jahangir graph, F-trees, (obtained from a path $v_1, v_2, \ldots, v_n$ by appending an edge to v_{n-1} and v_n), H-super subdivision of a path are edge magic total graphs, and an H-graph of a path, alternate triangular belt graph, braid graphs, semi Jahangir graph, H-super subdivision of a path, F-trees, the $H \odot K_1$ graph of a path are edge trimagic total graphs.

In [3156] Regees, Anisha, and Nicholas introduced the notion of edge-bimagic harmonious graphs as follows. A graph $G(V,E)$ with p vertices and q edges is called an *edge-bimagic harmonious* graph if there exists a bijective mapping f from $V \cup E$ onto $\{1,2,\ldots,p+q\}$ such that for each edge xy the value of the $(f(x)+f(y)) \bmod q)((f(x)+f(y))(\bmod q)+f(xy))$ is a constant c_1 or c_2. If there are three such constants c_1, c_2 and c_3, G said to be an edge *trimagic harmonious* graph. They proved that ladders and double ladders are edge-bimagic harmonious graphs and that circular ladders, triangular ladders are edge magic and bimagic harmonious graphs. If an edge bimagic harmonious labeling has the extra property that the vertex labels are 1 the number of vertices it called a *super edge-bimagic harmonious* labeling . In [4308] Deen, Aboamer, and Sherbiny showed that wheels and the splitting graph of odd cycles are super edge-bimagic harmonious graphs. They also showed that sunflower graphs and double sunflower graphs are super edge trimagic harmonious graphs.

Amuthavalli and Sugapriya [225] defined a *reverse edge-trimagic* labeling on a graph $G(V,E)$ with p vertices and q edges as a one-to-one map that takes the vertices and edges onto the integers $1,2,\ldots,p+q$ with the property that for every edge e, when the sum of all vertex labels incident to e is subtracted from edge label $f(e)$, the result is one of three constants. A reverse edge-trimagic labeling is said to be a *reverse super*

edge-trimagic labeling if $f(V) = \{1, 2, \ldots, p\}$ and $f(E) = \{p+1, p+2, \ldots, p+q\}$. They investigated the reverse super edge-trimagic labeling of barycentric subdivision of bistars, degree splitting graphs of $K_{1,n} + K_{1,n}$ and $K_{1,n} \cup K_{1,n}$, and the splitting graphs of stars.

Chopra, Dios, and Lee [814] investigated the edge-magicness of joins of graphs. Among their results are: $K_{2,m}$ is edge-magic if and only if $m = 4$ or 10; the only possible edge-magic graphs of the form $K_{3,m}$ are those with $m = 3, 5, 6, 15, 33$, and 69; for any fixed m there are only finitely many n such that $K_{m,n}$ is edge-magic; for any fixed m there are only finitely many trees T such that $T + \overline{K_m}$ is edge-magic; and wheels are not edge-magic.

Lee, Ho, Tan, and Su [2152] define the *edge-magic index* of a graph G to be the smallest positive integer k such that the graph kG is edge-magic. They completely determined the edge-magic indices of graphs which are stars. In [3533] Shiu, Lam, and Lee give the edge-magic index set of the second power of a path.

For any graph G and any positive integer k the graph $G[k]$, called the *k-fold* G, is the hypergraph obtained from G by replacing each edge of G with k parallel edges. Lee, Seah, and Tan [2196] proved that for any graph G with p vertices, $G[2p]$ is edge-magic and, if p is odd, $G[p]$ is edge-magic. Shiu, Lam, and Lee [3532] show that if G is an $(n+1, n)$-multigraph, then G is edge-magic if and only if n is odd and G is isomorphic to the disjoint union of K_2 and $(n-1)/2$ copies of $K_2[2]$. They also prove that if G is a $(2m+1, 2m)$-multigraph and $k \geq 2$, then $G[k]$ is edge-magic if and only if $2m+1$ divides $k(k-1)$. For a $(2m, 2m-1)$-multigraph G and k at least 2, they show that $G[k]$ is edge-magic if $4m$ divides $(2m-1)k((2m-1)k+1)$ or if $4m$ divides $(2m+k-1)k$. In [3530] Shiu, Lam, and Lee characterize the (p, p)-multigraphs that are edge-magic as $mK_2[2]$ or the disjoint union of $mK_2[2]$ and two particular multigraphs or the disjoint union of K_2, $mK_2[2]$, and four particular multigraphs. They also show for every $(2m+1, 2m+1)$-multigraph G, $G[k]$ is edge-magic for all k at least 2. Lee, Seah, and Tan [2196] prove that the multigraph $C_n[k]$ is edge-magic for $k \geq 2$.

Tables 6 and 7 summarize what is known about edge-magic total labelings and super edge-magic total labelings. We use **SEMT** to indicate the graphs have super edge-magic total labelings and **EMT** to indicate the graphs have edge-magic total labelings. A question mark following SEMT or EMT indicates that the graph is conjectured to have the corresponding property. The tables were prepared by Petr Kovář and Tereza Kovářová.

Table 6: **Summary of Edge-magic Total Labelings**

Graph	*Types*	*Notes*
P_n	EMT	[4124]
trees	EMT?	[2018], [3176]
C_n	EMT	for $n \geq 3$ [2017], [1280], [3190], [627]
K_n	EMT	iff $n = 1, 2, 3, 4, 5$, or 6 [2018], [862], [1037] enumeration of all EMT of K_n [4124]
$K_{m,n}$	EMT	[4124], [2017]
crowns $C_n \odot K_1$	EMT	[4264], [4124]
C_n with a single edge attached to one vertex	EMT	[4124]
wheels W_n	EMT	iff $n \not\equiv 3 \pmod 4$ [1037], [1161]
fans	EMT	[3629], [1087], [1088]
(p, q)-graph	not EMT	if q even, $p + q \equiv 2 \pmod 4$ [3176]
nP_2	EMT	iff n odd [2017]
$P_n + K_1$	EMT	[4264]
r-regular graph	not EMT	r odd and $p \equiv 4 \pmod 8$ [862]
$P_3 \cup nK_2$ and $P_5 \cup nK_2$	EMT	[1087], [1088]
$P_4 \cup nK_2$	EMT	n odd [1087], [1088]
nP_i	EMT	n odd, $i = 3, 4, 5$ [4264], [1087], [1088]
nP_3	EMT?	[4264]
$2P_n$	EMT	[1087], [1088]

Continued on next page

Table 6 – *Continued from previous page*

Graph	*Types*	*Notes*
$P_1 \cup P_2 \cup \cdots \cup P_n$	EMT	[1087], [1088]
$mK_{1,n}$	EMT	[1087], [1088]
unicylic graphs	EMT?	[2157]
$K_1 \odot nK_2$	EMT	n even [1087], [1088]
$K_2 \times \overline{K}_n$	EMT	[1087], [1088]
nK_3	EMT	iff $n \neq 2$ odd [1087], [1088], [2463]
binary trees	EMT	[1087], [1088]
$P(m,n)$ (generalized Petersen graph see §2.7)	EMT	[1087], [1088], [2700]
ladders	EMT	[1087], [1088]
books	EMT	[1087], [1088]
odd cycle with pendent edges attached to one vertex	EMT	[1087], [1088]
$P_m \times C_n$	EMT	n odd $n \geq 3$ [4184]
$P_m \times P_2$	EMT	m odd $m \geq 3$ [4184]
$K_{1,m} \cup K_{1,n}$	EMT	iff mn is even [1091]
$G \odot \overline{K}_n$	EMT	if G is EMT 2-regular [1089]

Table 7: **Summary of Super Edge-magic Labelings**

Graph	*Types*	*Notes*
C_n	SEMT	iff n is odd [1037]
caterpillars	SEMT	[1037], [2017], [2018]
$K_{m,n}$	SEMT	iff $m = 1$ or $n = 1$ [1037]
K_n	SEMT	iff $n = 1, 2$ or 3 [1037]
trees	SEMT?	[1037]
nK_2	SEMT	iff n odd [799]
nG	SEMT	if G is a bipartite or tripartite SEM graph and n odd [1091]
$mB(n)$	SEMT	if m is odd [3598]
$m(P_2 \times P_n)$	SEMT	if $m,\ n$ are odd [3598]
$K_{1,m} \cup K_{1,n}$	SEMT	if m is a multiple of $n + 1$ [1091]
$K_{1,m} \cup K_{1,n}$	SEMT?	iff m is a multiple of $n + 1$ [1091]
$K_{1,2} \cup K_{1,n}$	SEMT	iff n is a multiple of 3 [1091]
$K_{1,3} \cup K_{1,n}$	SEMT	iff n is a multiple of 4 [1091]
$P_m \cup K_{1,n}$	SEMT	if $m \geq 4$ is even [1091]
$2P_n$	SEMT	iff n is not 2 or 3 [1091]
$2P_{4n}$	SEMT	for all n [1091]
mP_n	SEMT	if $m \equiv 2 \pmod 4$, $n \neq 2, 3$ [3598]
mP_{4n}	SEMT	if$m \equiv 2 \pmod 4$, $n > 1$ [3598]
$K_{1,m} \cup 2nK_{1,2}$	SEMT	for all m and n [1091]

Continued on next page

Table 7 – *Continued from previous page*

Graph	*Types*	*Notes*
$C_3 \cup C_n$	SEMT	iff $n \geq 6$ even [1094], [1329]
$C_4 \cup C_n$	SEMT	iff $n \geq 5$ odd [1094], [1329]
$C_5 \cup C_n$	SEMT	iff $n \geq 4$ even [1094]
$C_m \cup C_n$	SEMT	if $m \geq 6$ even, n odd $n \geq m/2 + 2$ [1094]
$C_m \cup C_n$	SEMT?	iff $m + n \geq 9$ and $m + n$ odd [1094]
$C_4 \cup P_n$	SEMT	iff $n \neq 3$ [1094]
$C_5 \cup P_n$	SEMT	if $n \neq 4$ [1094]
$C_m \cup P_n$	SEMT	if $m \geq 6$ even, $n \geq m/2 + 2$ [1094]
$P_m \cup P_n$	SEMT	iff $(m, n) \neq (2, 2)$ or $(3, 3)$ [1094]
corona $C_n \odot \overline{K}_m$	SEMT	$n \geq 3$ [1094]
$St(m, n)$	SEMT	$n \equiv 0 \pmod{m+1}$ [2154]
$St(1, k, n)$	SEMT	$k = 1, 2$ or n [2154]
$St(2, k, n)$	SEMT	$k = 2, 3$ [2154]
$St(1, 1, k, n)$	SEMT	$k = 2, 3$ [2154]
$St(k, 2, 2, n)$	SEMT	$k = 1, 2$ [2154]
$St(a_1, \ldots, a_n)$	SEMT?	for $n > 1$ odd [2154]
C_{4m}^t	SEMT	[2381]
C_{4m+1}^t	SEMT	[2381]
friendship graph of n triangles	SEMT	iff $n = 3, 4, 5$, or 7 [3629]

Continued on next page

Table 7 – *Continued from previous page*

Graph	*Types*	*Notes*
generalized Petersen graph $P(n,2)$ (see §2.7)	SEMT	if $n \geq 3$ odd [1159]
nP_3	SEMT	if $n \geq 4$ even [592]
P_n^2	SEMT	[1087]
$K_2 \times C_{2n+1}$	SEMT	[1087]
$P_3 \cup kP_2$	SEMT	for all k [1088]
kP_n	SEMT	if k is odd [1088]
$k(P_2 \cup P_n)$	SEMT	if k is odd and $n = 3, 4$ [1088]
fans F_n	SEMT	iff $n \leq 6$ [1088]
books B_n	SEMT	if n even [4298]
books B_n	SEMT?	if $n \equiv 5 \pmod 8$ [1088]
trees with α-labelings	SEMT	[1088]
$P_{2m+1} \times P_2$	SEMT	[1037], [1087]
$C_{2m+1} \times P_m$	SEMT	[1087]
$G \odot \overline{K}_n$	SEMT	if G is SEM 2-regular graph [1089]
$C_m \odot \overline{K}_n$	SEMT	[1089]
join of K_1 with any subgraph of a star	SEMT	[799]
if G is k-regular SEMT graph		then $k \leq 3$ [799]
G is connected (p,q)-graph	SEMT	G exists iff $p - 1 \leq q \leq 2p - 3$ [799]
G is connected 3-regular	SEMT	iff $p \equiv 2 \pmod 4$ [799]

Continued on next page

Table 7 – *Continued from previous page*

Graph	*Types*	*Notes*
graph on p vertices $nK_2 + nK_2$	 not SEMT	 [799]

5.3 Vertex-magic Total Labelings

MacDougall, Miller, Slamin, and Wallis [2378] introduced the notion of a vertex-magic total labeling in 1999. For a graph $G(V, E)$ an injective mapping f from $V \cup E$ to the set $\{1, 2, \ldots, |V| + |E|\}$ is a *vertex-magic total labeling* if there is a constant k, called the *magic constant*, such that for every vertex v, $f(v) + \sum f(vu) = k$ where the sum is over all vertices u adjacent to v (some authors use the term "vertex-magic" for this concept). They prove that the following graphs have vertex-magic total labelings: C_n; P_n $(n > 2)$; $K_{m,m}$ $(m > 1)$; $K_{m,m} - e$ $(m > 2)$; and K_n for n odd. They also prove that when $n > m + 1$, $K_{m,n}$ does not have a vertex-magic total labeling. They conjectured that $K_{m,m+1}$ has a vertex-magic total labeling for all m and that K_n has vertex-magic total labeling for all $n \geq 3$. The latter conjecture was proved by Lin and Miller [2270] for the case that n is divisible by 4 while the remaining cases were done by MacDougall, Miller, Slamin, and Wallis [2378]. McQuillan [2462] provided many vertex-magic total labelings for cycles C_{nk} for $k \geq 3$ and odd $n \geq 3$ using given vertex-magic labelings for C_k. Gray, MacDougall, and Wallis [1339] then gave a simpler proof that all complete graphs are vertex-magic total. Krishnappa, Kothapalli, and Venkaiah [2006] gave another proof that all complete graphs are vertex-magic total. Senthil Amutha and Murugesan [3362] characterized connected vertex magic total labeling graphs through their ideals in topological spaces. Among other results, Wang and Zhang [4163] settle a 2006 conjecture raised by Slamin et al., which claims the existence of the vertex magic total labeling of disjoint union of multiple copies of $C_n \odot K_1$. Vimal Kumar and Vijayalakshmi [4076] investigated vertex magic total labelings of the middle and total graphs of cycles.

In [2378] MacDougall, Miller, Slamin, and Wallis conjectured that for $n \geq 5$, K_n has a vertex-magic total labeling with magic constant h if and only if h is an integer satisfying $n^3 + 3n \leq 4h \leq n^3 + 2n^2 + n$. In [2465] McQuillan and Smith proved that this conjecture is true when n is odd. Armstrong and McQuillan [274] proved that if $n \equiv 2 \pmod 4$ $(n \geq 6)$ then K_n has a vertex-magic total labeling with magic constant h for each integer h satisfying $n^3 + 6n \leq 4h \leq n^3 + 2n^2 - 2n$. If, in addition, $n \equiv 2 \pmod 8$, then K_n has a vertex-magic total labeling with magic constant h for each integer h satisfying $n^3 + 4n \leq 4h \leq n^3 + 2n^2$. They further showed that for each odd integer $n \geq 5$, $2K_n$ has a vertex-magic total labeling with magic constant h for each integer h such that $n^3 + 5n \leq 2h \leq n^3 + 2n^2 - 3n$. If, in addition, $n \equiv 1 \pmod 4$, then $2K_n$ has a vertex-magic total labeling with magic constant h for each integer h such that $n^3 + 3n \leq 2h \leq n^3 + 2n^2 - n$.

In [2463] McQuillan and McQuillan investigate the existence of vertex-magic labelings of nC_3. They prove: for every even integer $n \geq 4$, nC_3 is vertex-magic (and therefore also edge-magic); for each even integer $n \geq 6$, nC_3 has vertex-magic total labelings with at least $2n-2$ different magic constants; if $n \equiv 2 \bmod 4$, two extra vertex-magic total labelings with the highest possible and lowest possible magic constants exist; if $n = 2 \cdot 3^k,\ k > 1,\ nC_3$ has a vertex-magic total labeling with magic constant k if and only if $(1/2)(15n+4) \leq k \leq (1/2)(21n+2)$; if n is odd, there are vertex-magic total labelings for nC_3 with $n+1$ different magic constants. In [2461] McQuillan provides a technique for constructing vertex-magic total labelings of 2-regular graphs. In particular, if m is an odd positive integer, $G = C_{n_1} \cup C_{n_2} \cup \cdots \cup C_{n_k}$ has a vertex-magic total labeling, and J is any subset of $I = \{1, 2, \ldots, k\}$ then $(\cup_{i \in J}\ mC_{n_i}) \cup (\cup_{i \in I-J}\ mC_{n_i})$ has a vertex-magic total labeling.

In [846] Cichacz, Fronček and Singgih introduced a new method to expand some known vertex magic total labelings of 2-regular graphs. The also proved that for odd values of m, if $(2r+1) \not\equiv 0 \pmod 3$ and $n \not\equiv 0 \pmod{(2r+1)}$, then $2mC_{rn} \cup mC_n$ has a vertex magic total labeling.

Lin and Miller [2270] have shown that $K_{m,m}$ is vertex-magic total for all $m > 1$ and that K_n is vertex-magic total for all $n \equiv 0 \pmod 4$. Phillips, Rees, and Wallis [2837] generalized the Lin and Miller result by proving that $K_{m,n}$ is vertex-magic total if and only if m and n differ by at most 1. Cattell [760] has shown that a necessary condition for a graph of the form $H + \overline{K_n}$ to be vertex-magic total is that the number of vertices of H is at least $n-1$. As a corollary he gets that a necessary condition for $K_{m_1,m_2,\ldots,m_r,n}$ where n is the largest size of any partite set to be vertex-magic total is that $m_1+m_2+\cdots+m_r \geq n$. He poses as an open question whether graphs that meet the conditions of the theorem are vertex-magic total. Cattell also proves that $K_{1,n,n}$ has a vertex-magic total labeling when n is odd and $K_{2,n,n}$ has a vertex-magic total labeling when $n \equiv 3 \pmod 4$. In [3076] Rahim and Slamin proved the disjoint union of coronas $C_{t_1} \odot K_1 \cup C_{t_2} \odot K_1 \cup \cdots \cup C_{t_n} \odot K_1$ has a vertex-magic total labeling with magic constant $6\sum_{k=1}^{n} t_k + 1$.

Miller, Bača, and MacDougall [2495] have proved that the generalized Petersen graphs $P(n,k)$ (see §2.7) for the definition) are vertex-magic total when n is even and $k \leq n/2-1$. They conjecture that all $P(n,k)$ are vertex-magic total when $k \leq (n-1)/2$ and all prisms $C_n \times P_2$ are vertex-magic total. Bača, Miller, and Slamin [412] proved the first of these conjectures (see also [3631] for partial results) while Slamin and Miller prove the second. Slamin, Prihandoko, Setiawan, Rosita and Shaleh [3632] constructed vertex-magic total labelings for the disjoint union of two copies of $P(n,k)$ and Silaban, Parestu, Herawati, Sugeng, and Slamin [3582] extended this to any number of copies of $P(n,k)$. More generally, they proved that for $n_j \geq 3$ and $1 \leq k_j \leq \lfloor (n_j-1)/2 \rfloor$, the union $P(n_1,k_1) \cup P(n_2,k_2) \cup \cdots \cup P(n_t,k_t)$ has a vertex-magic total labeling with vertex magic constant $10(n_1+n_2+\cdots+n_t)+2$. In the same article Silaban et al. define the union of t special circulant graphs $\cup_{j=1}^{t} C_n(1,m_j)$ as the graph with vertex set $\{v_i^j |\ 0 \leq i \leq n-1, 1 \leq j \leq t\}$ and edge set $\{v_i^j v_{i+1}^j |\ 0 \leq i \leq n-1, 1 \leq j \leq t\} \cup \{v_i^j v_{i+m_j}^j |\ 0 \leq i \leq n-1, 1 \leq j \leq t\}$. They prove that for odd n at least 5 and $m_j \in \{2, 3, \ldots, (n-1)/2\}$, the disjoint union $\cup_{j=1}^{t} C_n(1,m_j)$ has a vertex-magic total labeling with constant $8tn + (n - 10/2 + 3$.

MacDougall et al. ([2378], [2380] and [1337]) have shown: W_n has a vertex-magic total labeling if and only if $n \leq 11$; fans F_n have a vertex-magic total labelings if and only if $n \leq 10$; friendship graphs have vertex-magic total labelings if and only if the number of triangles is at most 3; $K_{m,n}$ $(m > 1)$ has a vertex-magic total labeling if and only if m and n differ by at most 1. Wallis [4120] proved: if G and H have the same order and $G \cup H$ is vertex-magic total then so is $G + H$; if the disjoint union of stars is vertex-magic total, then the average size of the stars is less than 3; if a tree has n internal vertices and more than $2n$ leaves then it does not have a vertex-magic total labeling. Wallis [4121] has shown that if G is a regular graph of even degree that has a vertex-magic total labeling then the graph consisting of an odd number of copies of G is vertex-magic total. He also proved that if G is a regular graph of odd degree (not K_1) that has a vertex-magic total labeling then the graph consisting of any number of copies of G is vertex-magic total.

Gray, MacDougall, McSorley, and Wallis [1338] investigated vertex-magic total labelings of forests. They provide sufficient conditions for the nonexistence of a vertex-magic total labeling of forests based on the maximum degree and the number of internal vertices, and leaves or the number of components. They also use Skolem sequences to prove a star forest with each component a $K_{1,2}$ has a vertex-magic total labeling.

Recall a helm H_n is obtained from a wheel W_n by attaching a pendent edge at each vertex of the n-cycle of the wheel. A *generalized helm* $H(n,t)$ is a graph obtained from a wheel W_n by attaching a path on t vertices at each vertex of the n-cycle. A *generalized web* $\mathrm{W}(n,t)$ is a graph obtained from a generalized helm $H(n,t)$ by joining the corresponding vertices of each path to form an n-cycle. Thus $\mathrm{W}(n,t)$ has $(t+1)n+1$ vertices and $2(t+1)n$ edges. A *generalized Jahangir graph* $J_{k,s}$ is a graph on $ks+1$ vertices consisting of a cycle C_{ks} and one additional vertex that is adjacent to k vertices of C_{ks} at distance s to each other on C_{ks}. Rahim, Tomescu, and Slamin [3077] prove: H_n has no vertex-magic total labeling for any $n \geq 3$; $\mathrm{W}(n,t)$ has a vertex-magic total labeling for $n = 3$ or $n = 4$ and $t = 1$, but it is not vertex-magic total for $n \geq 17t + 12$ and $t \geq 0$; and $J_{n,t+1}$ is vertex-magic total for $n = 3$ and $t = 1$, but it does not have this property for $n \geq 7t + 11$ and $t \geq 1$. Recall a flower is the graph obtained from a helm by joining each pendent vertex to the central vertex of the helm. Ahmad and Tomescu [139] proved that flower graph is vertex-magic if and only if the underlying cycle is C_3.

Fronček, Kovář, and Kovářová [1137] proved that $C_n \times C_{2m+1}$ and $K_5 \times C_{2n+1}$ are vertex-magic total. Kovář [2020] furthermore proved some general results about products of certain regular vertex-magic total graphs. In particular, if G is a $(2r+1)$-regular vertex-magic total graph that can be factored into an $(r+1)$-regular graph and an r-regular graph, then $G \times K_5$ and $G \times C_n$ for n even are vertex-magic total. He also proved that if G an r-regular vertex-magic total graph and H is a $2s$-regular supermagic graph that can be factored into two s-regular factors, then their Cartesian product $G \times H$ is vertex-magic total if either r is odd, or r is even and $|H|$ is odd.

Ivančo and Polláková [1560] consider supermagic graphs having a saturated vertex (i.e., a vertex that is adjacent to every other vertex). They characterize supermagic graphs $G + K_1$, where G is a regular graph, using a connection to vertex-magic total graphs. They prove that if G is a d-regular graph of order n then the join $G + K_1$ is

supermagic if and only if G has a VMT labeling with constant h such that $(n-d-1)$ is a divisor of the non-negative integer $(n+1)h - n((d+2)/2)(n(d+2)/2)+1)$. They also prove $K_{1,n,n}$ is supermagic if and only if $n \geq 2$; $K_{1,2,2,\ldots,2}$ is supermagic except for $K_{1,2}$; and the graph obtained from $K_{n,n}$ $(n \geq 5)$ by removing all edges in a Hamilton cycle is supermagic. They also consider circulant graphs and prove that the complement of the circulant graph $C_{2n}(1,n),\ n \geq 4$, is supermagic.

In [3060] a novel algorithm is proposed based on the calculation of vertex magic total labelling value for every node in the network. Upon receiving the message from the sender node, the receiver node will quickly detect the faulty node by comparing the vertex magic total labeling pivot value. Experimental results show that the proposed approach leads to high true fault rate detection accuracy compared to the false fault rate detection.

MacDougall, Miller, and Sugeng [2379] define a *super vertex-magic total labeling* of a graph $G(V,E)$ as a vertex-magic total labeling f of G with the additional property that $f(V) = \{1,2,\ldots,|V|\}$ and $f(E) = \{|V|+1, |V|+2, \ldots, |V|+|E|\}$ (some authors use the term "super vertex-magic" for this concept). They show that a (p,q)-graph that has a super vertex-magic total labeling with magic constant k satisfies the following conditions: $k = (p+q)(p+q+1)/v - (v+1)/2$; $k \geq (41p+21)/18$; if G is connected, $k \geq (7p-5)/2$; p divides $q(q+1)$ if p is odd, and p divides $2q(q+1)$ if p is even; if G has even order either $p \equiv 0 \pmod 8$ and $q \equiv 0$ or $3 \pmod 4$ or $p \equiv 4 \pmod 8$ and $q \equiv 1$ or $2 \pmod 4$; if G is r-regular and p and r have opposite parity then $p \equiv 0 \pmod 8$ implies $q \equiv 0 \pmod 4$ and $p \equiv 4 \pmod 8$ implies $q \equiv 2 \pmod 4$. They also show: C_n has a super vertex-magic total labeling if and only if n is odd; and no wheel, ladder, fan, friendship graph, complete bipartite graph or graph with a vertex of degree 1 has a super vertex-magic total labeling. They conjecture that no tree has a super vertex-magic total labeling and that K_{4n} has a super vertex-magic total labeling when $n > 1$. The latter conjecture was proved by Gómez in [1297]. In [1298] Gómez proved that if G is a d-regular graph that has a vertex-magic total labeling and k is a positive integer such that $(k-1)(d+1)$ is even, then kG has a super vertex-magic total labeling. As a corollary, we have that if n and k are odd or if $n \equiv 0 \pmod 4$ and $n > 4$, then kK_n has a super vertex-magic total labeling. Gómez also shows how graphs with super vertex-magic total labeling can be constructed from a given graph G with super vertex-magic total labeling by adding edges to G in various ways.

Gray and MacDougall [1336] establish the existence of vertex-magic total labelings for several infinite classes of regular graphs. Their method enables them to begin with any even-regular graph and from it construct a cubic graph possessing a vertex-magic total labeling. A feature of the construction is that it produces strong vertex-magic total labelings many even order regular graphs. The construction also extends to certain families of non-regular graphs. MacDougall has conjectured (see [2021]) that every r-regular $(r > 1)$ graph with the exception of $2K_3$ has a vertex-magic total labeling. As a corollary of a general result Kovář [2021] has shown that every $2r$-regular graph with an odd number of vertices and a Hamiltonian cycle has a vertex-magic total labeling.

Gómez and Kovář [1299] proved that a super vertex-magic total labeling of kK_n exists for n odd and any k, for $4 < n \equiv 0 \pmod 4$ and any k, and for $n = 4$ and k even. They also showed kK_{4t+2} does not admit a super vertex-magic total labeling for k odd and

provide a large number of super vertex-magic total labelings of kK_{4t+2} for any k based on a super vertex-magic total labeling of kK_{4t+1}.

Beardon [599] has shown that a necessary condition for a graph with c components, p vertices, q edges and a vertex of degree d to be vertex-magic total is $(d+2)^2 \leq (7q^2+(6c+5)q+c^2+3c)/p$. When the graph is connected this reduces to $(d+2)^2 \leq (7q^2+11q+4)/p$. As a corollary, the following are not vertex-magic total: wheels W_n when $n \geq 12$; fans F_n when $n \geq 11$; and friendship graphs $C_3^{(n)}$ when $n \geq 4$.

Beardon [601] has investigated how vertices of small degree effect vertex-magic total labelings. Let $G(p,q)$ be a graph with a vertex-magic total labeling with magic constant k and let d_0 be the minimum degree of any vertex. He proves $k \leq (1+d_0)(p+q-d_0/2)$ and $q < (1+d_0)q$. He also shows that if $G(p,q)$ is a vertex-magic graph with a vertex of degree one and t is the number of vertices of degree at least two, then $t > q/3 \geq (p-1)/3$. Beardon [601] has shown that the graph obtained by attaching a pendent edge to K_n is vertex-magic total if and only if $n = 2, 3$, or 4.

Meissner and Zwierzyński [2483] used finding vertex-magic total labelings of graphs as a way to compare the efficiency of parallel execution of a program versus sequential processing.

Swaminathan and Jeyanthi [3841] prove the following graphs are super vertex-magic total: P_n if and only if n is odd and $n \geq 3$; C_n if and only if n is odd; the star graph if and only if it is P_2; and mC_n if and only if m and n are odd. In [3842] they prove the following: no super vertex-magic total graph has two or more isolated vertices or an isolated edge; a tree with n internal edges and tn leaves is not super vertex-magic total if $t > (n+1)/n$; if Δ is the largest degree of any vertex in a tree T with p vertices and $\Delta > (-3+\sqrt{1+16p})/2$, then T is not super vertex-magic total; the graph obtained from a comb by appending a pendent edge to each vertex of degree 2 is super vertex-magic total; the graph obtained by attaching a path with t edges to a vertex of an n-cycle is super vertex-magic total if and only if $n+t$ is odd. Ali, Bača, and Bashir [182] proved that mP_3 and mP_4 have no super vertex-magic total labeling

For $n > 1$ and distinct odd integers x, y and z in [1,$n-1$] Javaid, Ismail, and Salman [1576] define the *chordal ring* of order n $CR_n(x,y,z)$, as the graph with vertex set Z_n, the additive group of integers modulo n, and edges $(i, i+x), (i, i+y), (i, i+z)$ for all even i. They prove that $CR_n(1,3,n-1)$ has a super vertex-magic total labeling when $n \equiv 0 \mod 4$ and $n \geq 8$ and conjecture that for an odd integer Δ, $3 \leq \Delta \leq n-3, n \equiv 0 \mod 4$, $CR_n(1,\Delta,n-1)$ has a super vertex-magic total labeling with magic constant $23n/4+2$.

The *Knödel graphs* $W_{\Delta,n}$ with n even and degree Δ, where $1 \leq \Delta \leq \lfloor \log_2 n \rfloor$ have vertices pairs (i,j) with $i = 1,2$ and $0 \leq j \leq n/2-1$ where for every $0 \leq j \leq n/2-1$ and there is an edge between vertex $(1,j)$ and every vertex $(2,(j+2^k-1) \bmod n/2)$, for $k = 0, 1, \ldots, \Delta-1$. Xi, Yang, Mominul, and Wong [4223] have shown that $W_{3,n}$ is super vertex-magic total when $n \equiv 0 \mod 4$.

Marimuthu and Balakrishnan [2426] called a vertex magic total labeling of $G(V,E)$ *E-super vertex magic* if $f(E(G)) = \{1,2,3,\ldots,|E(G)|\}$. The *cocktail party* graph, $H_{m,n}$ $(m,n \geq 2)$, is the graph with a vertex set $V = \{v_1, v_2, \ldots, v_{mn}\}$ partitioned into n independent sets $V = \{I_1, I_2, \ldots, I_n\}$ each of size m such that $v_i v_j \in E$ for all

$i, j \in \{1, 2, \ldots, mn\}$ where $i \in I_p,\ j \in I_q,\ p \neq q$. (The graph $H_{n,n}$ is the complement of the ladder graph and the dual graph of the n-cube.) Marimuthu and Balakrishnan [2426] gave some basic properties of such labelings and proved that $H_{m,n}$ is E-super vertex magic. Wang and Zhang [4161] show the following: Hamiltonian even regular graphs of odd order are E-super magic; even-regular graphs of odd order that contains a 2-factor consisting of an odd number of odd cycles with the same size are E-super vertex magic; graphs that can be decomposed into the sum of two spanning graphs where one is E-super magic and one is regular of even degree are E-supermagic; even-regular graphs of odd order that contain a 2-factor consisting of an odd number of odd cycles with the same size are E-super vertex magic; and circulant graphs with odd order are E-super vertex magic. Swaminathan and Jeyanthi [3841] proved that mC_n is E-super magic if and only if both m and n are odd. Kumar, Nirmala, and Lawrence [2058] investigated the existence and nonexistence of E-super vertex magic graceful labelings of paths, cycles, and unions of cycles.

Samarathunge, Athapattu, and Perera [3261] proved that an E-super vertex labeling of edges adjacent to a leaf vertex must be greater than or equal to $k - (p + q)$ and sum of the parent edge labelings must be greater than or equal to $k - (p + q)$, which k is the magic constant. This implies that an E-super vertex labeling does not been proved that E-super vertex magic labeling does not exist for the perfect binary trees. A broom $B_{n,d}$ is obtained by attaching $n - d$ pendent edges to one of the pendent vertices of P_d. Marimuthu, Suganya, Kalaivani, and Balakrishnan [2429] proved the following: $B_{n,n-2}$ is E-super vertex magic if and only if n is odd; a graph obtained by subdividing all the edges of $K_{1,n}$ is super E-super vertex magic if and only if $n \leq 4$, and $K_{1,n}$ is E-super vertex magic if and only if $n = 2$.

In [2428] Marimuthu and Kumar investigate E-super vertex magic labelings of disconnected graphs. They prove: if a graph with p vertices and q edges and even order has an E-super vertex magic labeling, then either (i) $p \equiv 0 \pmod 8$ and $q \equiv 0$ or $3 \pmod 4$, or (ii) $p \equiv 4 \pmod 8$ and $q \equiv 1$ or $2 \pmod 4$; if an r-regular graph G of order p has an E-super vertex magic labeling, then p and r have opposite parity and (i) if $p \equiv 0 \pmod 8$, then $q \equiv 0 \pmod 4$ (ii) if $p \equiv 4 \pmod 8$, then $q \equiv 2 \pmod 4$; mC_n is E-super vertex magic if and only if $P_n \cup (m-1)C_n$ is E-super vertex magic; $P_m \cup K_{1,m}$ is not E-super vertex magic; $C_m \cup P_n$ is not E-super vertex magic if both m and n have the same parity; the disjoint union of two non-isomorphic suns is not E-super vertex magic; the disjoint union of any number of isomorphic suns is not E-super vertex magic; and mP_3 is not E-super vertex magic for any integer $m > 1$. They conjecture that $K_m \cup P_m$ is E-super vertex magic if $m = 8t + 2$.

In [2606] Mutharasu and Kumar generalized the notion of super vertex-magic total labelings as follows. Let $G(V, E)$ be a graph and k be an integer with $1 \leq k \leq \text{diam}(G)$. For $e \in E(G)$, let $E_k(e)$ be the set of all vertices that are at a distance at most k from e and let $E_k(v)$ be the set of all edges that are at a distance at most k from v (u and v are at distance 1 from the edge uv). A graph G is said to be *E_k-regular with regularity* r if, for all edges e, $|E_k(e)| = r$ for some positive integer r. Note that all nontrivial graphs are E_1-regular. Let G be a simple graph with p vertices and q edges. A V-

super vertex magic labeling is a bijection $f : V(G) \cup E(G) \to \{1, 2, \ldots, p+q\}$ such that $f(V(G)) = \{1, 2, \ldots, p\}$ and for each vertex $v \in V(G)$, $f(v) + \sum_{u \in N(v)} f(uv) = M$ for some positive integer M. A *V_k-super vertex magic labeling* (V_k-SVML) is a bijection $f : V(G) \cup E(G) \to \{1, 2, \ldots, p+q\}$ with the property that $f(V(G)) = \{1, 2, \ldots, p\}$ and for each $v \in V(G)$, $f(v) + \sum_{e \in E_k(v)} f(e) = M$ for some positive integer M. A graph that admits a V_k-SVML is called *V_k-super vertex magic*. Mutharasu and Kumar gave a necessary and sufficient condition for the existence of V_k-SVML in graphs, determined the magic constant for E_k-regular graphs, and obtained results about V_2-SVML labelings for cycles, complement of cycles, prisms, and a family of circulant graphs.

Balbuena, Barker, Das, Lin, Miller, Ryan, and Slamin [462] call a vertex-magic total labeling of $G(V, E)$ a *strongly vertex-magic total labeling* if the vertex labels are $\{1, 2, \ldots, |V|\}$. They prove: the minimum degree of a strongly vertex-magic total graph is at least 2; for a strongly vertex-magic total graph G with n vertices and e edges, if $2e \geq \sqrt{10n^2 - 6n + 1}$ then the minimum degree of G is at least 3; and for a strongly vertex-magic total graph G with n vertices and e edges if $2e < \sqrt{10n^2 - 6n + 1}$ then the minimum degree of G is at most 6. They also provide strongly vertex-magic total labelings for certain families of circulant graphs. In [2461] McQuillan provides a technique for constructing vertex-magic total labelings of 2-regular graphs. In particular, if m is an odd positive integer, $G = C_{n_1} \cup C_{n_2} \cup \cdots \cup C_{n_k}$ has a strongly vertex-magic total labeling, and J is any subset of $I = \{1, 2, \ldots, k\}$ then $(\cup_{i \in J}\ mC_{n_i}) \cup (\cup_{i \in I-J}\ mC_{n_i})$ has a strongly vertex-magic total labeling.

Gray [1330] proved that if G is a graph with a spanning subgraph H that possesses a strongly vertex-magic total labeling and $G - E(H)$ is even regular, then G also possesses a strongly vertex-magic total labeling. As a corollary one has that regular Hamiltonian graphs of odd order have a strongly vertex-magic total labelings.

In a series of papers Gray and MacDougall expand on McQuillan's technique to obtain a variety of results. In [1333] Gray and MacDougall show that for any $r \geq 4$, every r-regular graph of odd order at most 17 has a strong vertex-magic total labeling. They also show that several large classes of r-regular graphs of even order, including some Hamiltonian graphs, have vertex-magic total labelings. They conjecture that every 2-regular graph of odd order possesses a strong vertex-magic total labeling if and only if it is not of the form $(2t-1)C_3 \cup C_4$ or $2tC_3 \cup C_5$. They include five open problems.

In [1335] Gray and MacDougall introduce a procedure called a *mutation* that transforms one vertex-magic totaling labeling into another one by swapping sets of edges among vertices that may result in different labeling of the same graph or a labeling of a different graph. Among their results are: a description of all possible mutations of a labeling of the path and the cycle; for all $n \geq 2$ and all i from 1 to $n-1$ the graphs obtained by identifying an end points of paths of lengths $i, i+1$, and $2n-2i-1$ have a vertex-magic total labeling; for odd n, the graph obtained by attaching a path of length $n-m$ to an m cycle, (such graphs are called *$(m; n-m)$-kites*) have strong vertex-magic total labelings for $m = 3, \ldots, n-2$; $C_{2n+1} \cup C_{4n+4}$ and $3C_{2n+1}$ have a strong vertex-magic total labeling; and for $n \geq 2$, $C_{4n} \cup C_{6n-1}$ has a strong vertex-magic total labeling. They conclude with three open problems.

Kimberley and MacDougall [1965] studied mutations that involve labelings of regular graphs into labelings of other regular graphs. They present results of extensive computations which confirm how prolific this procedure is. These computations add weight to MacDougall's conjecture that all nontrivial regular graphs are vertex-magic.

Gray and MacDougall [1334] show how to construct vertex-magic total labelings for several families of non-regular graphs, including the disjoint union of two other graphs already possessing vertex-magic total labelings. They prove that if G is a d-regular graph of order v and H a t-regular graph of order u with each having a strong vertex magic total labeling and $vd^2 + 2d + 2v + 2u = 2tvd + 2t + ut^2$ then $G \cup H$ possesses a strong vertex-magic total labeling. They also provide bounds on the minimum degree of a graph with a vertex-magic total labeling.

In [1336] Gray and MacDougall establish the existence of vertex-magic total labelings for several infinite classes of regular graphs. Their method enables them to begin with any even-regular graph and construct a cubic graph possessing a vertex-magic total labeling that produces strong vertex-magic total labelings for many even order regular graphs. The construction also extends to certain families of non-regular graphs.

In [2635] Nagaraj, Ponnappan, and Prabakaran define a vertex-magic total labeling of G to be an *even vertex magic total labeling* if the set of vertex labels is $\{2, 4, 6, \ldots, 2|V(G)|\}$. They prove the following: C_n is even vertex magic total if and only if n is odd; rC_s is even vertex magic total if and only if r and s are odd; $C_n \odot K_1$ is even vertex magic total; wheels are not even vertex magic total; fans (excluding C_3) are not even vertex magic total; kites are not even vertex magic total; and K_{4n} is not even vertex magic total. In [2638] they prove that $C_3 \cup C_{2t}$ $(t > 2)$ and $C_4 \cup C_{2t+1}$ $(t \geq 2)$ have even vertex magic total labelings. In [2637] Nagaraj, Ponnappan, and Prabakan prove that the union of any finite numbers of graphs of the form $C_n \odot K_1$ (the sizes may vary) has an even vertex magic total labeling. Saduakdee and Khemmani [3283] studied conditions for which there exists an even vertex magic labeling of the t-fold wheel $W_{n,t}$ in terms of n and t. Furthermore, those having an even vertex magic total labeling are determined.

Rahim and Slamin [3074] give the bounds for the number of vertices for Jahangir graphs, helms, webs, flower graphs and sunflower graphs when the graphs considered are not vertex-magic total. Thirusangu, Nagar, and Rajeswari [3894] show that certain Cayley digraphs of cyclic groups have vertex-magic total labelings.

Balbuena, Barker, Lin, Miller, and Sugeng [468] call vertex-magic total labeling an *a-vertex consecutive magic* labeling if the vertex labels are $\{a, a+1, \ldots, a+|V|\}$. For an a-vertex consecutive magic labeling of a graph G with p vertices and q edges they prove: if G has one isolated vertex, then $a = q$ and $(p-1)^2 + p^2 = (2q+1)^2$; if $q = p-1$, then p is odd and $a = p-1$; if $p = q$, then p is odd and if G has minimum degree 1, then $a = (p+1)/2$ or $a = p$; if G is 2-regular, then p is odd and $a = 0$ or p; and if G is r-regular, then p and r have opposite parities. They also define a *b-edge consecutive magic* labeling analogously and state some results for these labelings. Marimuthu and Kumar [2427] gave some results related to a-vertex consecutive magic graphs. In [3416] Setiawan, Sugeng, Silaban, and Riama provided sufficient conditions for a graph to admit a super edge magic

total labeling and to have a b-edge consecutive edge magic total labeling. They also gave the super edge magic total labelings for banana trees, firecrackers, and identified several classes of connected graphs that have both labelings.

Wood [4211] generalizes vertex-magic total and edge-magic total labelings by requiring only that the labels be positive integers rather than consecutive positive integers. He gives upper bounds for the minimum values of the magic constant and the largest label for complete graphs, forests, and arbitrary graphs.

Exoo, Ling, McSorley, Phillips, and Wallis [1058] call a function λ a *totally magic labeling* of a graph G if λ is both an edge-magic total and a vertex-magic total labeling of G. A graph with such a labeling is called *totally magic*. Among their results are: P_3 is the only connected totally magic graph that has a vertex of degree 1; the only totally magic graphs with a component K_1 are K_1 and $K_1 \cup P_3$; the only totally magic complete graphs are K_1 and K_3; the only totally magic complete bipartite graph is $K_{1,2}$; nK_3 is totally magic if and only if n is odd; $P_3 \cup nK_3$ is totally magic if and only if n is even. In [4123] Wallis asks: Is the graph $K_{1,m} \cup nK_3$ ever totally magic? That question was answered by Calhoun, Ferland, Lister, and Polhill [750] who proved that if $K_{1,m} \cup nK_3$ is totally magic then $m = 2$ and $K_{1,2} \cup nK_3$ is totally magic if and only if n is even.

McSorley and Wallis [2467] examine the possible totally magic labelings of a union of an odd number of triangles and determine the spectrum of possible values for the sum of the label on a vertex and the labels on its incident edges and the sum of an edge label and the labels of the endpoints of the edge for all known totally magic graphs.

Gray and MacDougall [1331] define an *order n sparse semi-magic square* to be an $n \times n$ array containing the entries $1, 2, \ldots, m$ once (for some $m < n^2$), has its remaining entries equal to 0, and whose rows and columns have a constant sum of k. They prove some basic properties of such squares and provide constructions for several infinite families of squares, including squares of all orders $n \geq 3$. Moreover, they show how such arrays can be used to construct vertex-magic total labelings for certain families of graphs.

In Tables 8, 9, and 10, **VMT** means vertex-magic total labeling, **SVMT** means super vertex magic total, and **TM** means totally magic labeling. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The tables were prepared by Petr Kovář and Tereza Kovářová and updated by J. Gallian in 2007.

Table 8: **Summary of Vertex-magic Total Labelings**

Graph	*Types*	*Notes*
C_n	VMT	[2378]
P_n	VMT	$n > 2$ [2378]
$K_{m,m} - e$	VMT	$m > 2$ [2378]

Continued on next page

Table 8 – *Continued from previous page*

Graph	*Types*	*Notes*
$K_{m,n}$	VMT	iff $\|m-n\| \leq 1$ [2837], [2378], [2380]
K_n	VMT	for n odd [2378] for $n \equiv 2 \pmod 4$,$n > 2$ [2270]
nK_3	VMT	iff $n \neq 2$ [1087], [1088], [2463]
mK_n	VMT	$m \geq 1,\ n \geq 4$ [2466]
Petersen $P(n,k)$	VMT	[412]
prisms $C_n \times P_2$	VMT	[3631]
W_n	VMT	iff $n \leq 11$ [2378], [2380]
F_n	VMT	iff $n \leq 10$ [2378], [2380]
friendship graphs	VMT	iff # of triangles ≤ 3 [2378], [2380]
$G + H$	VMT	$\|V(G)\| = \|V(H)\|$ and $G \cup H$ is VMT [4120]
unions of stars	VMT	[4120]
tree with n internal vertices and more than $2n$ leaves	not VMT	[4120]
nG	VMT	n odd, G regular of even degree, VMT [4121] G is regular of odd degree, VMT, but not K_1 [4121]
$C_n \times C_{2m+1}$	VMT	[1137]
$K_5 \times C_{2n+1}$	VMT	[1137]
$G \times C_{2n}$	VMT	G $2r+1$-regular VMT [2020]
$G \times K_5$	VMT	G $2r+1$-regular VMT [2020]
$G \times H$	VMT	G r-regular VMT, r odd or r even and $\|H\|$ odd, H $2s$-regular supermagic [2020]

Table 9: **Summary of Super Vertex-magic Total Labelings**

Graph	*Types*	*Notes*
P_n	SVMT	iff $n > 1$ is odd [3841]
C_n	SVMT	iff n is odd [3841] and [2379]
$K_{1,n}$	SVMT	iff $n = 1$ [3841]
mC_n	SVMT	iff m and n are odd [3841]
W_n	not SVMT	[2379]
ladders	not SVMT	[2379]
friendship graphs	not SVMT	[2379]
$K_{m,n}$	not SVMT	[2379]
dragons (see §2.2)	SVMT	iff order is even [3842], [3842]
Knödel graphs $W_{3,n}$	SVMT	$n \equiv 0 \pmod 4$ [4223]
graphs with min. deg. 1	not SVMT	[2379]
K_{4n}	SVMT	$n > 1$ [1297]

Table 10: **Summary of Totally Magic Labelings**

Graph	*Types*	*Notes*
P_3	TM	the only connected TM graph with vertex of deg 1 [1058]
K_n	TM	iff $n = 1, 3$ [1058]

Continued on next page

Table 10 – *Continued from previous page*

Graph	*Types*	*Notes*
$K_{m,n}$	TM	iff $K_{m,n} = K_{1,2}$ [1058]
nK_3	TM	iff n is odd [1058]
$P_3 \cup nK_3$	TM	iff n is even [1058]
$K_{1,m} \cup nK_3$	TM	iff $m = 2$ and n is even [750]

5.4 H-Magic Labelings

In 2005 Gutiérrez and Lladó [1348] introduced the notion of an H-magic labeling of a graph, which generalizes the concept of a magic valuation. Let H and $G = (V, E)$ be finite simple graphs with the property that every edge of G belongs to at least one subgraph isomorphic to H. A bijection $f\colon V \cup E \to \{1, \ldots, |V| + |E|\}$ is an *H-magic labeling* of G if there exists a positive integer $m(f)$, called the *magic sum*, such that for any subgraph $H'(V', E')$ of G isomorphic to H, the sum $\sum_{v\in V'} f(v) + \sum_{e\in E'} f(e)$ is equal to the magic sum, $m(f)$. A graph is *H-magic* if it admits an H-magic labeling. If, in addition, the H-magic labeling f has the property that $\{f(v)\}_{v\in V} = \{1, \ldots, |V|\}$, then the graph is *$H$-supermagic*. A K_2-magic labeling is also known as an edge-magic total labeling. Gutiérrez and Lladó investigate the cases where $G = K_n$ or $G = K_{m,n}$ and H is a star or a path. Among their results are: a d-regular graph is not $K_{1,h}$ for any $1 < h < d$; $K_{n,n}$ is $K_{1,n}$-magic for all n; $K_{n,n}$ is not $K_{1,n}$-supermagic for $n > 1$; for any integers $1 < r < s$, $K_{r,s}$ is $K_{1,h}$-supermagic if and only if $h = s$; P_n is P_h-supermagic for all $2 \le h \le n$; K_n is not P_h-magic for any $2 < h \le n$; C_n is P_h-magic for any $2 \le h < n$ such that $\gcd(n, h(h-1)) = 1$. They also show that by uniformly gluing copies of H along edges of another graph G, one can construct connected H-magic graphs from a given 2-connected graph H and an H-free supermagic graph G.

Lladó and Moragas [2303] studied cycle-magic graphs. They proved: wheels W_n are C_3-magic for odd n at least 5; for $r \ge 3$ and $k \ge 2$ the windmill graphs $C_r^{(k)}$ (the one-point union of k copies of C_r) are C_r-supermagic; and if G is C_4-free supermagic graph of odd size, then $G \times K_2$ is C_4-supermagic. As corollaries of the latter result, they have that for n odd, prisms $C_n \times K_2$ and books $K_{1,n} \times K_2$ are C_4-magic. They define a subdivided wheel $W_n(r, k)$ as the graph obtained from a wheel W_n by replacing each radial edge $vv_i, 1 \le i \le n$ by a vv_i-path of size $r \ge 1$, and every external edge v_iv_{i+1} by a v_iv_{i+1}-path of size $k \ge 1$. They prove that $W_n(r, k)$ is C_{2r+k}-magic for any odd $n \ne 2r/k + 1$ and that $W_n(r, 1)$ is C_{2r+1}-supermagic. They also prove that the graph obtained by joining the end points of any number of internally disjoint paths of length $p \ge 2$ is C_{2p}-supermagic. Asif, Ali, Numan, and Semaničová-Feňovčíková [316] proved that if G is C_r-(super)magic, then so is nG and that $P_m \times P_n$ $(m, n \ge 4)$ is C_4-supermagic.

In [2999] Pradipta and Salman define a *calendula graph*, denoted by $Cl_{m,n}$, as the graph constructed from C_m and m copies of C_n, $C_{n_1}, C_{n_2}, \ldots, C_{n_m}$, and grafting the i-th edge of C_m to an edge of C_{n_i} for each i. They provide some cycle-supermagic labelings of calendula graphs. A p-calendula graph $Cal_{m,p}[n]$ consists of a cycle C_m with p copies of C_n amalgamated to each edge of C_m. Fronček [1135] generalized a previous result by Pradipta and Salman [2999] on 1-calendula graphs by providing C_n-supermagic labelings of $Cal_{m,p}[n]$ for all $m, n \geq 3$, $p \geq 1$, and $m \neq n$. The case $m = n$, $p > 1$ remains open. Pradipta and Salman [2999] proved that calendula graphs are C_n-supermagic for all $m, n \geq 3$ and Öner and Ate [2737] proved the same result using a dierent technique. Öner, Hussain and Banaras [9] proved that a polygonal snake δ_m^n is C_n-supermagic for every $m, n \geq 3$. In [1115] Freyberg, Jensen, and Scott studied other types of magic labelings of calendula graphs. [1135] new [2737] new [1115] new

Chithra, Marimuthu, and Kumar [807] provided some basic results on the magic constant of graphs, on cycle-supermagic labelings of generalized splitting graphs, and proved that mC_n is cycle -supermagic for $m \geq 2$ and $n \geq 3$. In [326] Azeem provided C_8-super magic labelings of zig-zag chains, linear chains, and the disjoint union of non-isomorphic copies of both chains. Yang, Rashid, Siddiqui, and Hanif [4245] provided explicit formulas for various C_n-super magic labelings of pumpkin graphs (graphs with exactly two vertices with multiple edges joining them) and two classes of planar graphs containing 8-sided and 4-sided faces or 6-sided and 4-sided faces.

A decomposition of a graph G into isomorphic copies of a graph H is H-magic if there is a bijection f from $V(G) \cup E(G)$ onto $\{0, 1, \ldots, |V(G)| + |E(G)| - 1\}$ such that the sum of labels of edges and vertices of each copy of H in the decomposition is constant. By using the results on the sumset partition problem, Inayah, Lladó, and Moragas [1531] show that K_{2m+1} admits T-magic decompositions by any graceful tree with m edges. They address analogous problems for complete bipartite graphs and for antimagic and (a, d)-antimagic decompositions.

An edge of H-magic graph G is said to be a *good edge* if it belongs to only one subgraph isomorphic to H. For $s \geq 1$, B is the collection of good edges obtained by choosing exactly s good edges from each subgraph isomorphic to H in G. A *uniform subdivided* graph $\mathcal{G}$ of the graph G is obtained by subdividing all edges of B with $k \geq 1$ vertices. A *nonuniform subdivided* graph is obtained by subdividing the edges of $E(G) \setminus B$. Rizvi, Khalid, Ali, Miller, and Ryan [3185] prove that if a graph G is a C_n-supermagic graph then its uniform subdivided graph $\mathcal{G}$ is C_{n+sk}-(super)magic for positive integers n, s, and k. Using known results on the cycle-supermagicness they immediately obtain that uniform subdivided graphs of fans, antiprisms, triangular ladders, ladders and grids are cycle-(super)magic. They also prove that some special nonuniform subdivisions of fans and triangular ladders are cycle-supermagic.

Jeyanthi and Muthuraja [1714] established that $P_{m,n}$ is C_{2m}-supermagic for all $m, n \geq 2$ and the splitting graph of C_n is C_4-supermagic for $n \neq 4$. Nirmalasari Wijaya, Ryan, and Kalinowski [2715] show that for odd n and arbitrary k, the firecracker $F_{k,n}$ is $F_{2,n}$-supermagic, the banana tree $B_{k,n}$ is $B_{1,n}$-supermagic, and flower graphs are C_3-supermagic. Kojima [1990] proved that for two positive integers m and t with $m > t \geq 2$,

if C_m is P_t-supermagic, then C_{3m} is also P_t-supermagic and for $t = 2,\ 3,\ 4,$ or 9 and C_n is P_t-supermagic if and only if n is odd with $n > t$. Nirmalasari Wijaya, Ryan, and Kalinowski [2683] proved that every d-dimensional grid graph ($d > 2$) is Q_d-supermagic where Q_d is the d-cube. Pu, Numan, Butt, Asif, Rafique, and Shao [3053] showed that toroidal fullerenes, Klein-bottle fullerenes, and the disjoint union of toroidal and Klein-bottle fullerenes are C_6-supermagic and the subdivision of toroidal fullerenes, Klein-bottle fullerenes, and any graph homeomorphic to a toroidal fullerene or Klein-bottle fullerene are cyclic-supermagic. Ulfatimah, Roswitha, and Kusmayadi [3926] proved that a star with one or more appended edges at each end-point admits a double star $S_{2,2}$-supermagic labeling and $L_m \odot P_n$ admits supermagic labeling of the one-point union of C_3 and C_4 for $m, n \geq 2$. Martini, Roswitha, and Lestari [2437] provided a C_4-supermagic labeling of $P_n \times P_m$ and a C_3 supermagic labeling of $K_1 + \overline{K_2}$.

Öner and Erol [2738] showed that the triangular book-snake graph $S(3, n, s)$ admits a C_3-supermagic labeling. Using these labelings and supermagicness of subdivisions given by Taimur, Numan, Mumtaz, and Semaničová-Feňovčíková in [3858] and the fact that $S(m, n, s)$ can be thought of as a subdivision of $S(3, n, s)$, Öner and Erol showed the C_m-supermagicness of polygonal book-snake graph $S(m, n, s)$.

The *edge corona path graph* $G_m \diamond P_n$ is the graph obtained from one copy of the gear graph G_m and $3m$ copies of P_n, P_n^i, by joining two end vertices of $e_i \in E(G_m)$ to every vertex $v_j inV(P_n)$ in the i-th copy of G_m with $i = 1, 2, \ldots, 3m$ and $j = 1, 2, \ldots, n$. Noviati, Martini, and Indriati [2718] provided a $C_3 \diamond P_n$-supermagic labeling for $f_n \diamond P_n$ and a $P_3 \diamond P_n$-supermagic labeling for $S_n \diamond P_n$ for odd $n \geq 3$.

Rizvi, Ali, and Hussian [3183] proved: the disjoint union of two or more copies of G is C_3-supermagic when G is a fan, triangular ladder, wheel, or a generalized antiprism; the disjoint union of two or more copies of G is C_3-supermagic when G is a ladder or a book; $sF_{n+1} \cup kF_n$ is C_3-supermagic; and $sL_{n+1} \cup kL_n$ is C_4-supermagic. Khalid, Rizvi, and Ali [1940] investigated whether the disjoint union of isomorphic copies of a connected cycle-supermagic graph is cycle-supermagic or not. They also study cycle-supermagic labelings for the disjoint union of isomorphic copies of fans, ladders, triangular ladders, wheels, books, and generalized antiprisms as well as disjoint unions of non-isomorphic copies of ladders and fans. Ali, Rizvi, Semaničová-Feňovčíková [189] proved that the disjoint union of an arbitrary number of isomorphic copies of prisms $C_n \times P_m$, $m \geq 2$ and $n \geq 3$, $n \neq 4$, is C_4-supermagic. They propose an open problem to find a C_4-supermagic labeling of the graph $t(C_4 \times P_m)$ for $m \geq 2$ and $t \geq 1$.

Liang [2251] proved the following: if there exist an even integer k and $m_i \equiv 0 \pmod{k}$ for every i in [1,n], then there exist $K_{k,k}$- and C_{2k}-supermagic decompositions of $K_{m_1,\ldots,m_n}$; if k and $t_n \geq k$ are even integers, then for any positive integers $t_i \equiv 0 \pmod{k}$, i in $[1, n-1]$, there exists a C_{2k}-supermagic decomposition of $K_{t_1,\ldots,t_{n-1},t_n}$; if there exists an even integer k and $K_{m,n}$ is C_{2k}-decomposable, then there exists a C_{2k}-supermagic decomposition of $K_{m,n}$; and if G is a graph with p vertices and p edges, H is a graph with q vertices and q edges, and there is an H-supermagic decomposition of G, then there exists an H-supermagic decomposition of nG. In [4182] Wichianpaisarn and Mato gave necessary and sufficient conditions for the existence of $K_{1,n-1}$-supermagic decomposition of $K_{n,n}$ minus

a one-factor.

In [2448] Maryati, Baskoro, and Salman provided P_n-(super) magic labelings of subdivisions of stars, shrubs and banana trees. Ngurah, Salman, and Sudarsana [2706] construct C_n-(super) magic labelings for some fans and ladders. For any connected graph H, Maryati, Salman, Baskoro, and Irawati [2451] proved that the disjoint union of k isomorphic copies of a connected graph H is a H-supermagic graph if and only if $|V(H)| + |E(H)|$ is even or k is odd. In [2449] Maryati, Baskoro, Salman, and Irawati give some necessary conditions for any P_n-magic graph and provide some P_n-supermagic labelings of a cycle with some pendent edges and its subdivisions.

The m-shadow of graph G, $D_m(G)$, is a graph obtained by taking m copies of G, namely, G_1, $G_2, \ldots, G_m$, and then joining every vertex u in G_i, $i \in \{1, 2, \ldots, m-1\}$, to the neighbors of the corresponding vertex v in G_{i+1}. Agustin, Susanto, Dafik, Prihandini, Alfarisi, and Sudarsana [91] studied the H-supermagic labelings of $D_m(G)$ where G are paths and cycles.

Kojima [1990] proved the following. Let G be a C_4-free super edge-magic (p, q)-graph with the minimum degree at least one and $m \geq 2$. If q odd and $m = 2$ or $|p-q| \geq 2$, then $P_m \times G$ is C_4-supermagic; if p is odd and $m = 2$ or $|p - q| = 1$ and $m \leq 5$, then $P_m \times G$ is C_4-supermagic; if $n \geq 3$ is odd and m is even, then $P_2 \times (C_n \odot \overline{K_m})$ is C_4-supermagic; if $n \geq 3$ is odd and m is odd, then $P_2 \times (C_n \odot \overline{K_m})$ is not C_4-supermagic; if G is a caterpillar, then $P_m \times G$ is C_4-supermagic for $m \geq 2$; and $P_m \times C_n$ is C_4-supermagic for $m \geq 2$ and $n \geq 3$. The latter result solved an open problem in [2707] by Ngurah, Salman, and Susilowati. Kojma also proved that if a C_4-free bipartite $(p, p-1)$-graph G with the minimum degree at least one and partite sets U and V has a super edge-magic labeling f of G such that $f(U) = \{1, 2, \ldots, |U|\}$, then $P_m \times (2G)$ is C_4-supermagic.

Maryati, Salman, Baskoro, Ryan, and Miller [2453] define a *shackle* as a graph obtained from nontrivial connected graphs $G_1, G_2, \ldots, G_k$ $(k \geq 2)$ such that G_s and G_t have no common vertex for every s and t in $[1, k]$ with $|s - t| \geq 2$, and for every i in $[1, k-1]$, G_i and G_{i+1} share exactly one common vertex that are all distinct. They prove that shackles and amalgamations constructed from copies of a connected graph H is H-supermagic. (Recall for finite collection of graph $G_1, G_2, \ldots, G_k$ with a fixed vertex v_i from each G_i, an *amalgamation*, AmalG_i, v_i), is the graph obtained by identifying the v_i.) Ashari and Salman [307] gave sufficient conditions for (H_1, H_2)-supermagic labelings for shackles involving cycles, flowers, and prisms.

Ngurah, Salman, and Susilowati [2707] proved the following: chain graphs with identical blocks each isomorphic to C_n are C_n-supermagic; fans are C_3-supermagic; ladders and books are C_4-supermagic; $K_{1,n} + K_1$ are C_3-supermagic; grids $P_m \times P_n$ are C_4-supermagic for $m \geq 3$ and $n = 3, 4$, and 5. They pose the case that $P_m \times P_n$ are C_4-supermagic for $n > 5$ as an open problem. They also have some results on P_t-(super) magic labelings of cycles. Kathiresan, Marimuthu, and Chithra investigated the existence of C_m-supermagic labelings generalized fans, generalization of a graph obtained by joining of a star K1,n with one isolated vertex, grids, and generalized books. The results given in this article are the generalizations of some results given by Ngurah, Salman, and Susilowati in [2707].

Roswitha, Baskoro, Maryati, Kurdhi, and Susanti [3208] proved: the generalized Ja-

hangir graph $J_{k,s}$ is C_{s+2}-supermagic; $K_{2,n}$ is C_4-supermagic; and W_n for n even and $n \geq 4$ is C_3-supermagic. As an open problem they asked if $K_{m,n}$, $2 < m \leq n$, admits a C_{2m}-supermagic labeling. Roswitha and Baskoro [3209] proved that double stars, caterpillars, firecrackers, and banana trees admit star-supermagic labelings.

Maryati, Salman, and Baskoro [2452] characterized all graphs G such that the disjoint union of copies of G is G-supermagic. They also showed: the disjoint union of any paths is mP_n-supermagic for certain values of m and n; some subgraph amalgamations of graphs G are G-supermagic; and for any subgraph H of G Amal(G, H, k) is G-supermagic. Salman and Maryati [3258] proved that Amal(G, P_n, k) is G-supermagic.

Selvagopal and Jeyanthi proved: for any positive integer n, a the k-polygonal snake of length n is C_k-supermagic [3343]; for $m \geq 2$, $n = 3$, or $n > 4$, $C_n \times P_m$ is C_4-supermagic [1755]; $P_2 \times P_n$ and $P_3 \times P_n$ are C_4-supermagic for all $n \geq 2$ [1755]; the one-point union of any number of copies of a 2-connected H is H-magic [1753]; graphs obtained by taking copies $H_1, H_2, \ldots, H_n$ of a 2-connected graph H and two distinct edges e_i, e_i' from each H_i and identifying e_i' of H_i with e_{i+1} of H_{i+1} where $|V(H)| \geq 4, |E(H)| \geq 4$ and n is odd or both n and $|V(H)|+|E(H)|$ are even are H-supermagic [1753]. For simple graphs H and G the *H-supermagic strength of G* is the minimum constant value of all H-magic total labelings of G for which the vertex labels are $\{1, 2, \ldots, |V|\}$. Jeyanthi and Selvagopal [1754] found the C_n-supermagic strength of n-polygonal snakes of any length and the H-supermagic strength of a chain of an arbitrary 2-connected simple graph.

Let $H_1, H_2, \ldots, H_n$ be copies of a graph H. Let u_i and v_i be two distinct vertices of H_i for $i = 1, 2, \ldots, n$. The *chain graph* H_n of H of length n is the graph obtained by identifying the vertices u_i and v_{i+1} for $i = 1, 2, \ldots, n-1$. In [1752] Jayanthi and Selvagopal show that a chain graph of any 2-connected simple graph H is H-supermagic and if H is a 2-connected (p, q) simple graph, then H_n is H-supermagic if $p+q$ is even or $p+q+n$ is even.

The *antiprism* on $2n$ vertices has vertex set $\{x_{1,1}, \ldots, x_{1,n}, x_{2,1}, \ldots, x_{2,n}\}$ and edge set $\{x_{j,i}, x_{j,i+1}\} \cup \{x_{1,i}, x_{2,i}\} \cup \{x_{1,i}, x_{2,i-1}\}$ (subscripts are taken modulo n). Jeyanthi, Selvagopal, and Sundaram [1757] proved the following graphs are C_3-supermagic: antiprisms, fans, and graphs obtained from the ladders $P_2 \times P_n$ with the two paths $v_{1,1}, \ldots, v_{1,n}$ and $v_{2,1}, \ldots, v_{2,n}$ by adding the edges $v_{1,j}v_{2,j+1}$.

Jeyanthi and Selvagopal [1756] show that for any 2-connected simple graph H the edge amalgamation of a finite number of copies of H is H-supermagic. They also show that the graph obtained by picking one endpoint v_i from each of k copies of $K_{1,k}$ then creating a new graph by joining each v_i to a fixed new vertex v is $K_{1,k}$-supermagic.

A bijection $f : V(G) \cup E(G) \to \{1, 2, \ldots, p+q\}$ is a *total neighborhood magic* labeling on a graph G with p vertices and q edges if $\sum_{v \in N(u} f(v) + f(uv) = k$ for all $u \in V(G)$, where $N(u)$ is the set of all vertices adjacent to u. A graph that admits such a labeling is said to be *total neighborhood magic*. In [3569] Shrimali, Parmar, and Patel proved that K_n (n odd) is total neighborhood magic and $C_n \odot K_1$ is not a total neighborhood magic. In [2790] Patel, Parmar, Bosmia, Vihol, Shukla and Diwan investigated total neighborhood magic labelings for completes graph with even vertices. Parmar, Patel, Bosmia, Vihol, and Prajapati [2782] constructed the total neighborhood magic labeling of disjoint union [2790] new [2782] new

of complete graphs,

An H-magic labeling is a *H-decomposition* of a graph G if there is a bijection $f : V(G) \cup E(G)$ onto $\{1, 2, \ldots, p+q\}$ such that for every copy of H in the decomposition, the sum of $f(v) + f(e)$ over all v in $V(H)$ and e in $E(H)$ is constant. The labeling f is said to be *$H - V$-super magic* if $f(V(G)) = \{1, 2, \ldots, p\}$. Marimuthu and Kumar [2432] prove that $K_{n,n}$ $(n \geq 2)$ is H-V-super magic decomposable when H is $K_{1,n}$. Marimuthu and Kumar [2430] provide a necessary and sufficient condition for the existence of V-super vertex-magic labeling and give E-super and V-super vertex-magic total labeling of certain families of generalized Petersen graphs. They also prove that no wheel is E-super vertex-magic, C_3 is the only friendship graph that is V-super vertex-magic, and C_3 is the only friendship graph that is E-super vertex-magic.

An H-magic labeling f is said to be a *H-E-super magic* labeling if $f(E(G)) = \{1, 2, \ldots, q\}$. A graph that admits an H-E-super magic labeling is called an *H-E-super magic decomposable* graph. Subbiah and Pandimadevi [3704] study some elementary properties of H-E-super magic labelings with H an m-factor and provide a necessary and sufficient condition for an even regular graph to be H-E-super magic decomposable where H is a 2-factor. Iyappan, Loganathan, and Kumar [1563] studied H-E-supermagic labelings of fans, triangle ladders, graphs obtained by joining a star $K_{1,n}$ with one isolated vertex, books, and grids.

5.5 Magic Labelings of Type (a, b, c)

A magic-type method for labeling the vertices, edges, and faces of a planar graph was introduced by Lih [2266] in 1983. Lih defines a *magic labeling of type* (1,1,0) of a planar graph $G(V, E)$ as an injective function from $\{1, 2, \ldots, |V|+|E|\}$ to $V \cup E$ with the property that for each interior face the sum of the labels of the vertices and the edges surrounding that face is some fixed value. Similarly, Lih defines a *magic labeling of type* $(1, 1, 1)$ of a planar graph $G(V, E)$ with face set F as an injective function from $\{1, 2, \ldots, |V|+|E|+|F|\}$ to $V \cup E \cup F$ with the property that for each interior face the sum of the labels of the face and the vertices and the edges surrounding that face is some fixed value. Lih calls a labeling involving the faces of a plane graph *consecutive* if for every integer s the weights of all s-sided faces constitute a set of consecutive integers. Lih gave consecutive magic labelings of type $(1, 1, 0)$ for wheels, friendship graphs, prisms, and some members of the Platonic family. In [335] Bača shows that the cylinders $C_n \times P_m$ have magic labelings of type $(1, 1, 0)$ when $m \geq 2, n \geq 3, n \neq 4$. In [345] Bača proves that the generalized Petersen graph $P(n, k)$ (see §2.7 for the definition) has a consecutive magic labeling if and only if n is even and at least 4 and $k \leq n/2 - 1$. In [122] Ahmed and Babujee provide face bimagic labelings of type (1,1,0) for certain wheels, cylinders, and disjoint unions of m copies of prism graphs. Roopa and Shobana [3201] proved the existence of face bimagic labelings of types (1,0,1), (1,1,0) and (0,1,1) for the graphs obtained by double duplication of all vertices by edges of ladder graphs. They also proved that if a graph G is (1, 0, 1)-face bimagic, except for three sided faces, then double duplication of all vertices by edges of G is face bimagic.

Bača gave magic labelings of type $(1,1,1)$ for fans [329], ladders [329], planar bipyramids (that is, 2-point suspensions of paths) [329], grids [338], hexagonal lattices [337], Möbius ladders [332], and $P_n \times P_3$ [333]. Kathiresan and Ganesan [1925] show that the graph $P_{a,b}$ consisting of $b \geq 2$ internally disjoint paths of length $a \geq 2$ with common end points has a magic labeling of type $(1,1,1)$ when b is odd, and when $a = 2$ and $b \equiv 0 \pmod 4$. They also show that $P_{a,b}$ has a consecutive labeling of type $(1,1,1)$ when b is even and $a \neq 2$. Ali, Hussain, Ahmad, and Miller [186] study magic labeling of type $(1,1,1)$ for wheels and subdivided wheels. They prove: wheels admits a magic labeling of type and $(1,1,1)$ and $(0,1,1)$, for odd n wheels W_n n admit a magic labeling of type $(0,1,0)$, and subdivided wheels admit a magic labeling of type $(1,1,0)$. As an open problem they ask for a magic labeling of type $(1,1,0)$ for W_n and n even. Ahmad [97] proves that subdivided ladders admit magic labelings of type (1,1,1) and admit consecutive magic labelings of type (1,1,0).

Bača [331], [330], [341], [339], [333], [340] and Bača and Holländer [370] gave magic labelings of type $(1,1,1)$ and type $(1,1,0)$ for certain classes of convex polytopes. Kathiresan and Gokulakrishnan [1927] provided magic labelings of type $(1,1,1)$ for the families of planar graphs with 3-sided faces, 5-sided faces, 6-sided faces, and one external infinite face. Bača [336] also provides consecutive and magic labelings of type $(0,1,1)$ (that is, an injective function from $\{1,2,\ldots,|E|+|F|\}$ to $E \cup F$ with the property that for each interior face the sum of the labels of the face and the edges surrounding that face is some fixed value) and a consecutive labeling of type $(1,1,1)$ for a kind of planar graph with hexagonal faces. Tabraiz and Hussain [3855] provide a super magic labeling of type (1, 0, 0) for ladders and a super magic labeling of type (1, 0, 0) for subdivided ladders.

A *magic labeling of type* (1,0,0) of a planar graph G with vertex set V is an injective function from $\{1,2,\ldots,|V|\}$ to V with the property that for each interior face the sum of the labels of the vertices surrounding that face is some fixed value. Kathiresan, Muthuvel, and Nagasubbu [1930] define a *lotus inside a circle* as the graph obtained from the cycle with consecutive vertices $a_1, a_2, \ldots, a_n$ and the star with central vertex b_0 and end vertices $b_1, b_2, \ldots, b_n$ by joining each b_i to a_i and a_{i+1} ($a_{n+1} = a_1$). They prove that these graphs ($n \geq 5$) and subdivisions of ladders have consecutive labelings of type $(1,0,0)$. Devaraj [937] proves that graphs obtained by subdividing each edge of a ladder exactly the same number of times has a magic labeling of type $(1,0,0)$.

In Table 11 we use following abbreviations

M(a,b,c) magic labeling of type (a,b,c)

CM(a,b,c) consecutive magic labeling of type (a,b,c).

A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The table was prepared by Petr Kovář and Tereza Kovářová.

Table 11: **Summary of Magic Labelings of Type** (a, b, c)

Graph	*Labeling*	*Notes*
W_n	CM(1,1,0)	[2266]
friendship graphs	CM(1,1,0)	[2266]
prisms	CM(1,1,0)	[2266]
cylinders $C_n \times P_m$	M(1,1,0)	$m \geq 2$, $n \geq 3$, $n \neq 4$ [335]
fans F_n	M(1,1,1)	[329]
ladders	M(1,1,1)	[329]
planar bipyramids (see §5.3)	M(1,1,1)	[329]
grids	M(1,1,1)	[338]
hexagonal lattices	M(1,1,1)	[337]
Möbius ladders	M(1,1,1)	[332]
$P_n \times P_3$	M(1,1,1)	[333]
certain classes of convex polytopes	M(1,1,1) M(1,1,0)	[331], [341], [339], [333] [340], [370]
certain classes of planar graphs with hexagonal faces	M(0,1,1) CM(0,1,1) CM(1,1,1)	[336]
lotus inside a circle (see §5.3)	CM(1,0,0)	$n \geq 5$ [1930]
subdivisions of ladders	M(1,0,0) CM(1,0,0)	[937] [1930]

5.6 Sigma Labelings/1-vertex magic labelings/Distance Magic

In 1987 Vilfred [4100] (see also [4101]) defined a *sigma-labeling* of a graph G with n vertices as a bijection f from the vertices of G to $\{1, 2, \ldots, n\}$ such that there is a constant k with the property that, at any vertex v the sum $\sum f(u)$ taken over all neighbors u of v is k. The concept of sigma labeling was independently studied in 2003 by Miller, Rodger, and Simanjuntak in [2503] under the name 1-*vertex magic*. In a 2009 article Sugeng, Fronček, Miller, Ryan, and Walker [3727] used the term *distance magic labeling.* For convenience, we will use the term distance magic. In [4103] Vilfred and Jinnah give a number of necessary conditions for a graph to have a distance magic labeling. One of them is that if u and v are vertices of a graph with a distance labeling, then the order of the symmetric difference of $N(u)$ and $N(v)$ (neighborhoods of u and v) is not 1 or 2. This condition rules out a large class of graphs as having distance magic labelings. Rao, Singh, and Parameswaran [3133] have shown $C_m \times C_n$ has a distance magic labeling if and only if $m = n \equiv 2 \pmod 4$ and $K_m \times K_n,\ m \geq 2, n \geq 3$ does not have a distance magic labeling. In [610] Benna gives necessary and sufficient condition for $K_{m,n}$ to be a distance magic graph and proves that if G_1 and G_2 are connected graphs with minimum degree 1 and at least three vertices, then $G_1 \times G_2$ does not have a distance magic labeling. Rao, Sighn, and Parameswaran [61] prove that every graph is an induced subgraph of a regular graph that has a distance magic labeling. As open problems, Rao [3131] asks for a characterize 4-regular graphs that have distance magic labelings and which graphs of the form $C_m \times C_n,\ m = n \equiv 2 \pmod 4$ have distance magic labelings. Kovář, Fronček, and Kovářová [2023] classified all orders n for which a 4-regular distance magic graph exists and also showed that there exists a distance magic graph with $k = 2t$ for every integer $t \geq 6$. Acharaya, Rao, Signh, and Parameswaran [60] proved $P_m \times C_n$ does not have a distance magic labeling when m is at least 3 and provide necessary and sufficient conditions for $K_{m,n}$ to have a distance magic labeling. However, in 2024 [3215] Rozmana and Šparl classify all distance magic Cartesian products of two cycles, thereby correcting an error in paper cited above by Rao, Singh, and Parameswaren [60]. Rozmana and Šparl further show that each distance magic labeling of a Cartesian product of cycles is determined by a pair or quadruple of suitable sequences, thereby obtaining a complete characterization of all distance magic labelings of these graphs. They also determine a lower bound on the number of all distance magic labelings of $C_m \times C_{2m}$ for $m \geq 3$ odd. In [1395] Headley showed that the only distance magic graph of the form $K_n \times C_m$ is $K_1 \times C_4$, and that $C_4 \times K_{n,t}$ is distance magic. Necessary conditions are given for $C_4 \times K_{n,t}$ to be distance magic when $n > t$. These conditions are shown to be sufficient when n and t are both even. He concluded with some examples of distance magic graphs of the form $C_4 \times K_{n,t}$ with $n > t$ in particular constructing an infinite sequence of non-isomorphic distance magic graphs of this type.

In [2001] Kovář, Rozman, Šparl introduced the concept of a self-reverse distance magic labeling of a regular graph that allows for a more compact description of the graph and the labeling in terms of the corresponding quotient graph as follows. Let Γ be a regular distance magic graph of order n and let ℓ be a corresponding distance magic labeling,

where the labels are the elements of $\{1, 2, \ldots, n\}$. For each vertex v of Γ let v^ℓ be the unique vertex of Γ such that $\ell(v) + \ell v^\ell = 0$. The distance magic labeling ℓ of Γ is said to be *self-reverse* if for each pair of vertices u and v of Γ u and v are adjacent if the vertices u^ℓ and v^ℓ are adjacent. They showed that the members of several known infinite families of tetravalent distance magic graphs admit such labelings. They also presented a novel general construction producing a new distance magic graph from two existing ones. Using it they show that for each integer $n \geq 6$, except for the odd integers up to 19, there exists a connected tetravalent graph of order n admitting a self-reverse distance magic labeling. They determined all connected tetravalent graphs up to order 30 admitting a self-reverse distance magic labeling.

Kovár and Silber [2025] proved that an $(n-3)$-regular distance magic graph with n vertices exists if and only if $n \equiv 3 \pmod 6$ and that its structure is determined uniquely. Moreover, they reduce constructions of Fronček to a single construction and provide another sufficient condition for the existence a distance magic graph with an odd number of vertices. Fronček, Kovář, and Kovářová [1138] provide a construction for distance magic graphs arising from arbitrary regular graphs based on an application of magic rectangles. They also solve a problem posed by Shafiq, Ali, and Simanjuntak [3459]. In [2026] Kováča, Silber, Kabelíıkova—Hrušsková, and Kravčenko classify of all feasible odd orders of r-regular distance magic graphs for $r = 6, 8, 10$, and 12. Godinho and Singh [1283] investigate the distance magic labelings for neighborhood expansions of graphs and present a method for embedding regular graphs into distance magic graphs.

Among the results of Miller, Rodger, and Simanjuntak in [2503]: the only trees that have a distance magic labeling are P_1 and P_3; C_n has a distance magic labeling if and only if $n = 4$; K_n has a distance magic labeling if and only if $n = 1$; the wheel $W_n = C_n + P_1$ has a distance magic labeling if and only if $n = 4$; the complete graph $K_{n,n,\ldots,n}$ with p partite sets has a distance magic labeling if and only if n is even or both n and p are odd; an r-regular graph where n is odd does not have a distance magic labeling; and $G \times \overline{K_{2n}}$ has a distance magic labeling for any regular graph G. They also give necessary and sufficient conditions for complete tripartite graphs to have a distance magic labeling. In 2003 it was conjectured by Miller, Rodger, and Simanjuntak in [2503] and in 2016 by Kotlar in [2011] that complete multipartite graphs admit distance magic labelings. In 2024 Ebrahem, Hoory, and Kotlar [995] gave an infinite family of counterexamples to this conjecture.

For a given a graph $G = (V, E)$ and a positive integer t, the *generalized Mycielskian graph* $M_t(G)$ is the graph with vertex set $\{V \times \{0, 1, \ldots, t-1\}\} \cup \{u\}$, with edges $(x, 0)(y, 0)$ and $(x, i)(y, i+1)$ where there is an edge $xy \in E$, and an edge $(x, t-1)u$ for all $x \in V$. In [2813] Pawar and Singh gave some results on distance magic labeling of generalized Mycielskian graphs. In [3216] Rozman and Ŝparl introduced a generalization of wreath graphs, called quasi wreath graphs, and classified all distance magic graphs among them. These provide infinitely many new tetravalent distance magic graphs. Moreover, they showed that the two non-wreath graphs of orders 12 and 14 are quasi wreath graphs and the one of order 16 can be obtained from a quasi wreath graph of order 14.

Generalizing definitions given by Vilfred in [4100] and [4101], Vilfred Kamalappan

[4102] defines a labeling $f : V(G) \to \{1, 2, \ldots, n\}$ on a graph G of order $n \geq 3$ as a *k-distance magic labeling* (k-DML) if $\sum_{w \in \partial N_k(u)} f(w)$ is a constant and independent of $u \in V(G)$ where $\partial N_k(u) = \{v \in V(G) : d(u, v) = k\}$, $k \in \mathbb{N}$. A graph G is called a *k-distance magic (k-DM)* if it has a k-DML. A *long brush*, denoted by $LP_{n,m}$, is a graph with vertex set $\{u_1, u_2, \ldots, u_n,\ v_1, v_2, \ldots, v_m\}$, a path $P_n = u_1\ u_2 \cdots u_n$ and edge set $E(P_n) \cup \{u_1 v_i : i = 1 \text{ to } m\} \cup E(< v_1, v_2, \ldots, v_m >)$, $m + n \geq 3$ and $m, n \in \mathbb{N}$. Using partition techniques, he obtain families of k-DM graphs and prove that (i) For $k, n \geq 3$, $m \geq 2$ and $k, m, n \in \mathbb{N}$, $LP_{n,m}$ is k-DM if and only if $m(m-1) \leq 2n$ and $k = n$; (ii) For a given $m \geq 2$ and for every $k \in \mathbb{N}_0$, $LP_{\frac{m(m-1)}{2}+k,m}$ is a $(\frac{m(m-1)}{2} + k)$-DM graph, and (iii) For $m \geq 3$, $x \geq 2$, $LP_{1,m} = K_1(u_1) + (K_{m_1} \cup K_{m_2} \cup \cdots \cup K_{m_x})$, $1 \leq m_1 \leq m_2 \leq \cdots \leq m_x$, $m_1 + m_2 + \cdots + m_x = m$, $m_1 + m_2 \geq 3$ and $m_1, m_2, ..., m_x, x \in \mathbb{N}$, $LP_{1,m}$ is 2-DM if and only if u_1 is assigned with a suitable j and $J_{m+1} \setminus \{j\}$ is partitioned into x constant sum partites of orders $m_1, m_2, ..., m_x$, $1 \leq j \leq m + 1$.

In [1126] Fronček defined the notion of a Γ-*distance magic* graph as one that has a bijective labeling of vertices with elements of an Abelian group Γ resulting in constant sums of neighbor labels. A graph that is Γ-distance magic for an Abelian group Γ is called *group distance magic*. Cichacz and Fronček [844] showed that for an r-regular distance magic graph G on n vertices, where r is odd there does not exist an Abelian group Γ of order n having exactly one involution (i.e., an element that is its own inverse) that is Γ-distance magic. Fronček [1126] proved that $C_m \times C_n$ is a Z_{mn}-distance magic graph if and only if mn is even. He also showed that $C_{2^n} \times C_{2^n}$ has a $Z_{2^{2n}}$-distance magic labeling. In [833] Cichacz showed some Γ-distance magic labelings for $C_m \times C_n$ where $\Gamma \not\approx Z_{mn}$ and $\Gamma \not\approx Z_{2^{2n}}$. Anholcer, Cichacz, Peterin, and Tepeh [253] proved that if an r_1-regular graph G_1 is Γ_1-distance magic and an r_2-regular graph G_2 is Γ_2-distance magic, then the direct product of graphs G_1 and G_2 is $\Gamma_1 \times \Gamma_2$-distance magic. Moreover they showed that if G is an r-regular graph of order n and $m = 4$ or $m = 8$ and r is even, then $C_m \times G$ is group distance magic. They proved that $C_m \times C_n$ is Z_{mn}-distance magic if and only if $m \in \{4, 8\}$ or $n \in \{4, 8\}$ or $m, n \equiv 0 \pmod 4$. They also showed that if $m, n \not\equiv 0 \pmod 4$ then $C_m \times C_n$ is not Γ-distance magic for any Abelian group Γ of order mn. Cichacz [834] gave necessary and sufficient conditions for complete k-partite graphs of odd order p to be Z_p-distance magic. Moreover, she showed that if $p \equiv 2 \pmod 4$ and k is even, then there does not exist a group Γ of order p that has a Γ-distance labeling for a k-partite complete graph of order p. She also proved that $K_{m,n}$ is a group distance magic graph if and only if $n + m \not\equiv 2 \pmod 4$. In [835] Cichacz proved that if G is an Eulerian graph, then the lexicographic product of G and C_4 is group distance magic. In the same paper she also showed that if $m + n$ is odd, then the lexicographic product of $K_{m,n}$ and C_4 is group distance magic. In [836] Cichacz gave necessary and sufficient conditions for direct product of $K_{m,n}$ and C_4 for $m + n$ odd and for $K_{m,n} \times C_8$ to be group distance magic. In [838] Cichacz proved that for n even and $r > 1$ the Cartesian product the complete r-partite graph $K_{n,n,\ldots,n}$ and C_4 is group distance magic. Godinho and Singh [1282] obtain group distance magic labelings of C_n^r for certain classes of Abelian groups and provide necessary conditions for existence of such labelings.

An orientable Γ-distance magic labeling of a graph, was introduced by Cichacz, Frey-

berg and Fronček [843] as a generalization of group distance magic labeling for oriented graphs. They showed that an even regular circulant graph of order n is orientable Z_n-distance magic, the direct product $C_n \times C_m$ is orientable Z_{nm}-distance magic. They also considered some products of circulant graphs. Moreover they proved that if G has order $n \equiv 2 \pmod 4$ and all vertices of odd degree, then there does not exist an orientable Γ-distance magic labeling of G for any Abelian group Γ of order n. Dyrlaga and Szopa in [994] gave necessary and sufficient conditions for lexicographic product $K_m \circ \overline{K_n} \cong K_{\underbrace{m, m, \ldots, m}_{n}}$ to be orientable ζ_{mn}-distance magic. As a consequence, they provide an infinite family of odd regular graphs possessing orientable ζ_n-distance magic labeling. In [1116] and [1117] Freyberg and Keranen found orientable Z_n-distance magic labelings of the Cartesian product of cycles. In [1118] they studied Z_n-distance magic labelings for the strong product of cycles.

In [2560] Mukherjee and Pawa use $L = \{1, 2, \ldots, n_p\}$ to denote the multiset obtained by reducing $1, 2, \ldots, n$ modulo p and replacing 0's, if any, by p. They call a graph G *p-distance magic* if there is a bijective map f from the vertex set V to the multiset L such that the weight of each vertex $w(x)$ is equal to the same element $\mu \pmod p$, where $w(x) = \sum f(y)$ is taken over all $y \in N(x)$. They proved that a graph G is distance magic if and only if it is p-distance magic for all $p \geq 1$. They also proved that for $p = n$, p-distance magic labeling becomes a Z_n-distance magic labeling. In some cases, they provide ways to construct different magic constants in for Z_n-distance magic labelings. In some cases, they prove the uniqueness of the magic constant for Z_n-distance magic labeling.

Anholcer, Cichacz, Peterin, and Tepeh [251] proved that the direct product of two cycles C_m and C_n is distance magic if and only if $m = 4$ or $n = 4$, or $m, n \equiv 0 \pmod 4$ (the *direct product* of graphs G and H has the vertex set $V(G) \times V(H)$ and (g, h) is adjacent to (g', h') if g is adjacent to g' in G and h is adjacent to h' in H). Zeng, Deng, and Luo [4335] characterized group distance magic labelings of the Cartesian product of two cycles. In [837] Cichacz gave necessary and sufficient conditions for circulant graph $C_n(1, 2, \ldots, p)$ to be distance magic for p odd. In [844] Cichacz and Fronček characterized all distance magic circulant graphs $C_n(1, p)$ for p odd. Cichacz, Fronček, Krop, and Raridan [845] proved that r-partite graph $K_{n,n,\ldots,n} \times C_4$ is distance magic if and only if $r > 1$ and $n > 2$ is even. Anholcer and Cichacz [247] gave necessary and sufficient conditions for lexicographic product of an r-regular graph G and $K_{m,n}$ to be distance magic. Cichacz and Görlich [849] gave necessary and sufficient conditions for the direct product of an r-regular graph G and $K_{m,n}$ to be distance magic. Kamatchi, Ramalakshmi, and Nilavarasi [1818] proved that P_n^2 and the graphs obtained from P_n ($n \neq 5$) by joining each internal vertex v with an end vertex at even distance from v are distance antimagic. In [3568] Shrimali and Parmar discuss the existence of distance magic labelings for the product, direct product, strong product, and carona product of graphs involving C_3^t and C_4. In [839] necessary and sufficient conditions for complete tripartite graphs to be group distance magic was given by Cichacz. In [280] Arumugam, Kamatchi, and Kovář give several results on distance magic graphs and open problems.

A graph $\Gamma(V, E)$ of order n is said to be a *closed distance magic labeling* of Γ is

a bijection $f : V \to \{1, 2, \ldots, n\}$ for which there exists a positive integer r such that $\sum_{x \in N[u]} f(x) = r$ for all vertices $u \in V$, where $N[u]$ is the closed neighborhood of u. A graph is said to be *closed distance magic* if it admits a closed distance magic labeling. Fernández, Maleki, Miklavič, and Razafimahatratra [1080] classified all connected closed distance magic circulants with valency at most 5, that is, Cayley graphs $\mathrm{Cay}(Z_n; S)$, where $|S| \leq 5$ and S generates Z_n. Miklavi and Ŝparl [2491] classified the distance magic circulants of valency 6 whose order are not divisible by 12.

A finite r-regular graph G has a p-partition (resp. closed p-partition) ($p \geq 2$) if there exists a partition of the set $V(G)$ into $V_1, V_2, \ldots, V_p$ such that for every $x \in V(G)$, all $V(x) \cap V_i$ (respectively, $V[x] \cap V_1$) have the same size. In [852] Cichacz and Nikodem proved the following for finite r-regular graphs G. If G is distance magic (resp. closed distance magic) graph with a p-partition and $p(t-1)$ even then tG is also distance (resp. closed distance) magic. If G has order t and H is p-regular such that tH is distance (resp. closed distance) magic, then the lexicographic product of G and H is distance (resp. closed distance) magic. If G has order t and H is such that tH is distance magic, then the lexicographic product of G and H and the direct product of G and H are distance (resp. closed distance) magic. If H is a p-regular distance magic graph with a 2-partition, then the lexicographic product of G and H and the direct product of G and H are distance magic. They further proved that if $G = C_3$ or G is the strong product of C_n and C_m for $n = 3$ and m odd, or $m, n \equiv 3 \pmod 6$, then tG is closed distance magic if and only if t is odd. (The *strong direct product* of G and H has vertex set $V(G) \times V(H)$ and (g, h) is adjacent to (g', h') if $g = g'$ and h is adjacent to h' in H, or $h = h'$ and g is adjacent to g' in G.)

In [3368] Seoud, Maqsoud, and Aldiban determined whether or not the following families of graphs have a distance magic vertex labeling: $K_n - \{e\}$; $K_n - \{2e\}$; P_n^k; C_n^2; $K_m \times C_n$; $C_m + P_n$; $C_m + C_n$; $P_m + P_n$; $K_{1,r,s}$; $K_{1,r,m,n}$; $K_{2,r,m,n}$; $K_{m,n} + P_k$; $K_{m,n} + C_k$; $C_m + \overline{K_n}$; $P_m + \overline{K_n}$; $P_m \times P_n$; $K_{m,n} \times P_k$; $K_m \times P_n$; the splitting graph of $K_{m,n}$; $K_n + G$; $K_m + \overline{K_n}$; $K_m + C_n$; $K_m + P_n$; $K_{m,n} + K_r$; $C_m \times P_n$; $C_m \times K_{1,n}$; $C_m \times K_{n,n}$; $C_m \times K_{n,n+1}$; $K_m \times K_{n,r}$; and $K_m \times K_n$. Typically, distance magic labelings exist only a few low parameter cases.

Miklavic and Sparl [2490] provided a sufficient condition for a Hamming graph to be distance magic and as a they corollary provide an infinite number of pairs (D, q) for which the corresponding Hamming graph with words of length D and over an alphabet of size q is distance magic. They classify distance magic folded hypercubes (a graph obtained from a hypercube by identifying pairs of vertices at maximal distance) by showing that the dimension-D folded hypercube is distance magic if and only if is divisible by 4.

Anholcer, Cichacz, Fronček, Simanjuntak, and Qiu [248] proved that for n odd, there does not exist a Γ-distance magic labeling of Q_n for any Abelian group Γ of order $|V(Q_n)|$, whereas for n even there exists a Γ-distance magic labeling of Q_n for every Abelian group Γ of order $|V(Q_n)|$. They also study similar distance antimagic and Γ-distance antimagic labelings where one finds a bijection such that the sums of labels are pairwise distinct for all the vertices. They show that there exists a Γ-distance antimagic labeling of Q_n for any Abelian group Γ of order $2n$ where n is odd for these labelings and give some

relationships between Γ-closed distance magic and antimagic labelings and Γ-distance antimagic labelings.

Cichacz [840] showed there exists an infinite family of odd regular graphs possessing Γ-distance magic labeling for groups Γ with more than one involution. In [837] Cichacz using a notion of a Γ-magic rectangle set $MRS_\Gamma(a,b;c)$ showed group distance labeling for Cartesian and direct product of complete r-partite graphs. These results supported a conjecture in [844] that says that if G is a distance magic graph, then G is group distance magic.

A *directed* Γ*-distance magic* labeling of an oriented graph $\overrightarrow{G} = (V, A)$ of order n is a bijective mapping f from the vertex set of G to an Abelian group Γ of order n with the property that there exists a constant $c \in \Gamma$ such that, for every vertex $v \in V(\overrightarrow{G})$, $w(v) = \sum_{u \in N_G^{\text{in}}(v)} f(u) - \sum_{u \in N_G^{\text{out}}(v)} f(u) = c$, where $N_G^{\text{in}}(v)$ is the open in-neighborhood and $N_G^{\text{out}}(v)$ is the open out-neighborhood of vertex v, that is $N_G^{\text{in}}(v) = \{u : uv \in A\}$ and $N_G^{\text{out}}(v) = \{u : vu \in A\}$. If for a graph G there exists an orientation $\overrightarrow{G}$ such that there is a directed Γ-distance magic labeling f for $\overrightarrow{G}$ the graph G is called *orientable* Γ*-distance magic*. Freyberg and Keranen [1117] proved that $C_m \times C_n$ is orientable $\mathbb{Z}_{mn}$-distance magic for all $m, n \geq 3$.

In [252] Anholcer, Cichacz, Peterin, and Tepeh introduce the notion of balanced distance magic graphs. They say that a distance magic graph G with an even number of vertices is *balanced* if there exists a bijection f from $V(G)$ to $\{1, 2, \ldots, |V(G)|\}$ such that for every vertex w the following holds: If $u \in N(w)$ with $f(u) = i$, then there exists $v \in N(w), u \neq v$ with $f(v) = |V(G)| - i + 1$. They prove that a graph G is balanced distance magic if and only if G is regular and $V(G)$ can be partitioned in pairs $(u_i, v_i), i \in \{1, 2, \ldots, |V(G)|/2$, such that $N(u_i) = N(v_i)$ for all i. Using this characterization, the following theorems are proved: if G is a regular graph and H is a graph not isomorphic to $\overline{K_n}$ where n is odd, then $G \odot H$ is a balanced distance magic graph if and only if H is a balanced distance magic graph; $G \times H$ is balanced distance magic if and only if one of G and H is balanced distance magic and the other one is regular; and $C_m \times C_n$ is distance magic if and only if $n = 4$ or $m = 4$ or $m, n \equiv 0 \pmod 4$ and $C_m \times C_n$ is balanced distance magic if and only if $n = 4$ or $m = 4$. In [255] they prove that every balanced distance magic graph is also group distance magic; the Cartesian product of a balanced distance magic graph and a regular graph is group distance magic; the direct product of C_4 or C_8 and a regular graph is group distance magic; and they show that $C_8 \times G$ is also group distance magic for any even-regular graph G. They also prove that $C_{4s} \times C_{4t}$ is $A \times B$-distance magic for any Abelian groups A and B of order $4s$ and $4t$, respectively. Moreover, they conjecture that $C_{4m} \times C_{4n}$ is a group distance magic graph. They prove that $C_m \times C_n$ is Z_{mn}-distance magic if and only if $m \in \{4, 8\}$ or $n \in \{4, 8\}$ or both n and m are divisible by 4, and that $C_m \times C_n$ with orders not divisible by 4 is not Γ-distance magic for any Abelian group Γ of order mn.

Let $G = (V, E)$ be a graph on n vertices. A bijection f from the vertices of graph G to $\{1, 2, \ldots, |V(G)|\}$ is called a *nearly distance magic* labeling of G if there exists a positive integer k such that $\sum f(x)$ over all $x \in N(v) = k$ or $k + 1$ for all v . The constant k is

called a *magic constant* of the graph and any graph which admits such a labeling is called a *nearly distance magic* graph. Godinho, Singh, and Arumugam [1285] give several basic results on nearly distance magic graphs and compute the magic constant k in terms of the fractional total domination number of the graph.

In [2446] Marr and Simanjunak defined *D-magic labelings* for oriented graphs where D is a distance set as follows. The vertices of a graph G are distinct integers in $\{1, 2, \ldots, |V(G)|\}$ such that the sum of all the vertex labels that are a distance in D from a given vertex is the same across all vertices. They gave some results related to the magic constant, constructed a few infinite families of D-magic graphs, and examined trees, cycles, and multipartite graphs.

In [1878] Kang, Chen, Li, and Hou investigated D-magic labelings of the halved n-cube ($n \geq 2$) (that is, all binary strings of length n with even number of 1's as vertices and edges between any two strings of Hamming distance 2). They show that the folded n-cube has a $\{1\}$-magic labeling (resp., a $\{0, 1\}$-magic labeling) if and only if $n = 0 \pmod 4$ (resp., $n = 3 \pmod 4$).

A survey of results on distance magic (sigma, 1-vertex) labelings through 2009 is given in [276].

5.7 Other Types of Magic Labelings

For a graph $G = (V, E)$ naturally embedded in the torus, let $\mathcal{F}(G)$ denote the set of faces of G. Then, G is called a *C_n-face-magic toroidal* graph if there exists a bijection $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ such that for every $F \in \mathcal{F}(G)$ with $F \cong C_n$, the sum of all the vertex labels along C_n is a constant S. Let $x_v = f(v)$ for all $v \in V(G)$. Then $\{x_v \mid v \in V(G)\}$ is called a *C_n-face-magic toroidal labeling* on G. Curran, Low, and Locke [872] show that, for all $m,\ n \geq 2,\ C_m \times C_n$ admits a C_4-face-magic toroidal labeling if and only if either $m = 2$ or $n = 2$, or both m and n are even. They say that a C_4-face-magic toroidal labeling $\{x_{i,j} \mid (i, j) \in V(C_{2m} \times C_{2n})\}$ on $C_{2m} \times C_{2n}$ is *antipodal balanced* if $x_{i,j} + x_{i+m,j+n} = S/2$, for all $(i, j) \in V(C_{2m} \times C_{2n})$. They show that there exists an antipodal balanced C_4-face-magic toroidal labeling on $C_{2m} \times C_{2n}$ if and only if the parity of m and n are the same. Furthermore, when both m and n are even, an antipodal balanced C_4-face-magic toroidal labeling on $C_{2m} \times C_{2n}$ is both row-sum balanced and column-sum balanced. In addition, when $m = n$ is even, an antipodal balanced C_4-face-magic toroidal labeling on $C_{2n} \times C_{2n}$ is diagonal-sum balanced. In [868] Curran and Locke proved that there are 144 distinct C_4-face-magic labelings on the 4×4 projective grid graph $P_{4,4}$ (up to symmetries on the projective plane).

For a graph $G(V, E)$ embedded in the Klein bottle, let $F(G)$ denote the set of faces of G. A *C_4-face-magic Klein bottle labeling* on G is a bijection $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ such that for any $F \in F(G)$ with $F \cong C_4$, the sum of all the vertex labels along F is a constant. A C_4-face-magic labeling $X = \{x_{i,j} : 1 \leq i, j \leq 4\}$ on the 4×4 Klein bottle grid graph is called *horizontally* (or *vertically*) *pairwise balanced* if $x_{2i-1,j} + x_{2i,j} = 17$ for $1 \leq i \leq 2$ and $1 \leq j \leq 4$ (or $x_{i,2j-1} + x_{i,2j} = 17$ for $1 \leq i \leq 4$ and $1 \leq j \leq 2$). Myers and Curran [2622] consider the $m \times n$ grid graph, denoted by $G_{m,n}$, embedded in the Klein [2622] new

bottle in the natural way. Curran, Low, and Locke [873] showed that for $m = 4$, $G_{4,n}$ admits a C_4-face-magic Klein bottle labeling if and only if n is even. They say that a bottle labeling f on $G_{m,n}$ is *equatorially balanced* if $f(v_i)+f(v_j) = mn+1$ for all antipodal vertex pairs v_i, v_j. They showed that when n is odd, a C_4-face-magic Klein bottle labeling on $G_{4,n}$ must be equatorially balanced. They further showed that when n is odd, up to symmetries on the Klein bottle, the number of C_4-face-magic Klein bottle labelings on the $4 \times n$ Klein bottle grid graph is $\tau(n)$, where $\tau(n)$ is the number of positive divisors of n. When m is an odd integer and n is an even integer, they proved that the minimum number of distinct C_4-face-magic Klein bottle labelings f on $G_{m,n}$ (up to symmetries on the Klein bottle) is either 1 if $m = 1$, or $\tau(m)$ if $m > 1$. In [2622], Myers and Curran showed that the 4×4 Klein bottle grid graph has 192 C_4-face-magic labelings up to symmetries on the Klein bottle. They classify these labelings into two categories depending on whether a C_4-face-magic label-preserving permutation of the labeling is either horizontally pairwise balanced or vertically pairwise balanced. [873] new [2622] new

Curran [866] showed that an $m \times n$ projective grid graph admits a C_4-face-magic labeling if and only if both m and n have the same parity. When m and n are even, the C_4-face-magic value of a C_4-face-magic labeling on an $m \times n$ projective grid graph must be $2mn + 2$. Also, when m and n are odd, he proved that the C_4-face-magic value of a C_4-face-magic labeling on an $m \times n$ projective grid graph is either $2mn + 1$, $2mn + 2$, or $2mn + 3$. In [867] Curran determined a category of the C_4-face-magic labelings on $\mathcal{P}_{m,n}$ having C_4-face-magic value $2mn + 1$ or $2mn + 3$. He conjectured that these are the only C_4-face-magic labelings on $\mathcal{P}_{m,n}$ having C_4-face-magic value $2mn + 1$ or $2mn + 3$.

In 2004 Baskar Babujee [561] and [562] introduced the notion of *vertex-bimagic* labeling in which there exist two constants k_1 and k_2 such that the sums involved in a specified type of magic labeling is k_1 or k_2. Thus a vertex-bimagic total labeling with bimagic constants k_1 and k_2 is the same as a vertex-magic total labeling except for each vertex v the sum of the label of v and all edges adjacent to v may be k_1 or k_2. Murugesan and Senthil Amutha [2600] proved that the bistar $B_{n,n}$ is vertex-bimagic total labeling for $n > 2$. An edge-bimagic total labeling *edge-bimagic total* of a graph $G(V, E)$ with p vertices and q edges is a bijection f from the set of vertices and edges to such that for every edge $uv \in E$, $f(u) + f(uv) + f(v)$ is one of two constants k_1 or k_2, independent of the choice of the edge. A bimagic labeling is of interest for graphs that do not have a magic labeling of a particular type. Bimagic labelings for which the number of sums equal to k_1 and the number of sums equal to k_2 differ by at most 1 are called *equitable*. When all sums except one are the same the labeling is called *almost magic*. Although the wheel W_n does not have an edge-magic total labeling when $n \equiv 3 \pmod 4$, Marr, Phillips and Wallis [2443] showed that these wheels have both equitable bimagic and almost magic labelings. They also show that whereas nK_2 has an edge-magic total labeling if and only if n is odd, nK_2 has an edge-bimagic total labeling when n is even and although even cycles do not have super edge-magic total labelings all cycles have super edge-bimagic total labelings. They conjecture that there is a constant N such that K_n has an edge-bimagic total labeling if and only if n is at most N. They show that such an N must be at least 8. They also prove that if G has an edge-magic total labeling then $2G$ has an edge-

bimagic total equitable labeling. Amara Jothi, David, and Baskar Babujee [212] provide edge-bimagic labelings for switching of paths, cycles, stars, crowns and helms. They also examine whether operations on edge magic graphs results in edge-bimagic graphs or not.

Baskar Babujee and Babitha [566] call a graph with p vertices *1-vertex bimagic* if there is a bijective labeling f from the vertices to $\{1, 2, \ldots, p\}$ such that for each vertex u the sum of all $f(v)$ where v is adjacent to u is either a constant k_1 or a constant k_2 and $k_1 \neq k_2$. A graph with p vertices is called *odd 1-vertex bimagic* if there is a bijective labeling f from the vertices to $\{1, 3, \ldots, 2p-1\}$ such that for each vertex u the sum of all $f(v)$ where v is adjacent to u is either a constant k_1 or a constant k_2 and $k_1 \neq k_2$. A graph with p vertices is called *even 1-vertex bimagic* if there is a bijective labeling f from the vertices to $\{0, 2, \ldots, 2(p-1)\}$ such that for each vertex u the sum of all $f(v)$ where v is adjacent to u is either a constant k_1 or a constant k_2 and $k_1 \neq k_2$.

Baskar Babujee and Babitha [566] prove that a necessary condition for the existence of a 1-vertex bimagic vertex labeling f of a graph G is $\sum_{x \in V(G)} d(x)f(x) = k_1p_1 + k_2p_2$ where $d(x)$ is the degree of vertex x and p_1 and p_2 are the number of vertices with common count k_1 and k_2, respectively. Among their results are: if G has a 1-vertex bimagic vertex labeling and $G \neq C_4$, then $G+K_1$ admits a 1- vertex bimagic vertex labeling; C_n a 1-vertex bimagic if and only if $n = 4$; $K_{m,n}$ is 1-vertex bimagic; graphs obtained from P_n $(n \geq 3)$ by adding edges joining every pair of vertices an odd distance apart are 1-vertex bimagic; n-partite graphs of the form $K_{p,p,\ldots,p}$ are 1-vertex bimagic for all $p > 1$ when n is even and 1-vertex bimagic for all even p when n is odd; a regular or biregular graph admits a 1-vertex bimagic labeling if and only if it the admits an odd 1-vertex bimagic labeling and if and only it admits an even 1-vertex bimagic labeling.

In [3360] Semenyuta, Nedilko, and Nedilko introduce the notion of the equivalence of vertex labelings on a given graph. They prove the equivalence of three bimagic labelings for regular graphs and obtain a particular solution for the problem of the existence a 1-vertex bimagic vertex labeling for graphs of isomorphic $K_{n,n,m}$. They prove that the sequence of biregular graphs $K_{n(ij)} = ((K_{n-1} - M) + K_1) - (u_nu_i) - (u_nu_j)$ admits a 1-vertex bimagic vertex labeling, where u_i, u_j is any pair of nonadjacent vertices in the graph $K_{n-1} - M$, u_n is the vertex of K_1, and M is the perfect matching of the complete graph K_{n-1}. They show that if an r-regular graph G of order n is a distance magic graph, then the graph $G + G$ has a 1-vertex bimagic vertex labeling with magic constants $(n+1)(n+r)/2+n^2$ and $(n+1)(n+r)/2+nr$. They also define two new types of graphs that do not admit 1-vertex bimagic vertex labeling.

Baskar Babujee and Jagadesh [562], [570], [571], and [569] proved the following graphs have super edge-bimagic labelings: cycles of length 3 with a nontrivial path attached; $P_3 \odot K_{1,n}$ n even; $P_n + \overline{K_2}$ (n odd); $P_2 + mK_1$ $(m \geq 2)$; $2P_n$ $(n \geq 2)$; the disjoint union of two stars; $3K_{1,n}$ $(n \geq 2)$; $P_n \cup P_{n+1}$ $(n \geq 2)$; $C_3 \cup K_{1,n}$; P_n; $K_{1,n}$; $K_{1,n,n}$; the graphs obtained by joining the centers of any two stars with an edge or a path of length 2; the graphs obtained by joining the centers of two copies of $K_{1,n}$ $(n \geq 3)$ with a path of length 2 then joining the center one of copies of $K_{1,n}$ to the center of a third copy of $K_{1,n}$ with a path of length 2; combs $P_n \odot K_1$; cycles; wheels; fans; gears; K_n if and only if $n \leq 5$.

Given positive integers k and λ, Yao, Chen, Yao, and Cheng [4257] say that a total

labeling f of a connected graph $G(V,E)$ from $V \cup E$ to $\{1,2,\ldots,|V|+|E|\}$ such that $f(x) \neq f(y)$ for distinct $x, y \in V \cup E$ and $f(u)+f(v) = k+\lambda f(uv)$ for each edge uv in E is a (k,λ)-*magically total labeling* of G. They provide necessary and sufficient conditions for graphs with (k,λ))-magically total labelings to also have graceful, odd-graceful, felicitous, and (a,d)-edge antimagic total labelings (see §6.2).

In [2312] López, Muntaner-Batle, and Rius-Font give a necessary condition for a complete graph to be edge-bimagic in the case that the two constants have the same parity.

In [567] Baskar Babujee, Babitha, and Vishnupriya make the following definitions. For any natural number a, a graph $G(p,q)$ is said to be *a-additive super edge-bimagic* if there exists a bijective function f from $V(G) \cup E(G)$ to $\{a+1, a+2, \ldots, a+p+q\}$ such that for every edge uv, $f(u)+f(v)+f(uv) = k_1$ or k_2. For any natural number a, a graph $G(p,q)$ is said to be *a-multiplicative super edgebimagic* if there exists a bijective f from $V(G) \cup E(G)$ to $\{a, 2a, \ldots, (p+q)a\}$ such that for every edge uv, $f(u)+f(v)+f(uv) = k_1$ or k_2. A graph $G(p,q)$ is said to be *super edge-odd bimagic* if there exists a bijection f from $V(G) \cup E(G)$ to $\{1,3,5,\ldots,2(p+q)-1\}$ such that for every edge uv $f(u)+f(v)+f(uv) = k_1$ or k_2. If f is a super edge-bimagic labeling, then a function g from $E(G)$ to $\{0,1\}$ with the property that for every edge uv, $g(uv) = 0$ if $f(u)+f(v)+f(uv) = k_1$ and $g(uv) = 1$ if $f(u)+f(v)+f(uv) = k_2$ is called a *super edge-bimagic cordial labeling* if the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. They prove: super edge-bimagic graphs are a-additive super edge-bimagic; super edge-bimagic graphs are a-multiplicative super edge-bimagic; if G is super edge-magic, then $G+K_1$ is super edge-bimagic labeling; the union of two super edge magic graphs is super edge-bimagic; and P_n, C_{2n} and $K_{1,n}$ are super edge-bimagic cordial.

For any nontrivial Abelian group A under addition a graph G is said to be *A-magic* if there exists a labeling f of the edges of G with the nonzero elements of A such that the vertex labeling f^+ defined by $f^+(v) = \Sigma f(vu)$ over all edges vu is a constant. In [3693] and [3694] Stanley noted that Z-magic graphs can be viewed in the more general context of linear homogeneous diophantine equations. Shiu, Lam, and Sun [3534] have shown the following: the union of two edge-disjoint A-magic graphs with the same vertex set is A-magic; the Cartesian product of two A-magic graphs is A-magic; the lexicographic product of two A-magic connected graphs is A-magic; for an Abelian group A of even order a graph is A-magic if and only if the degrees of all of its vertices have the same parity; if G and H are connected and A-magic, G composed with H is A-magic; $K_{m,n}$ is A-magic when $m, n \geq 2$ and A has order at least 4; K_n with an edge deleted is A-magic when $n \geq 4$ and A has order at least 4; all generalized theta graphs (§4.4 for the definition) are A-magic when A has order at least 4; $C_n + \overline{K_m}$ is A-magic when $n \geq 3, m \geq 2$ and A has order at least 2; wheels are A-magic when A has order at least 4; flower graphs $C_m@C_n$ are A-magic when $m, n \geq 2$ and A has order at least 4 ($C_m@C_n$ is obtained from C_n by joining the end points of a path of length $m-1$ to each pair of consecutive vertices of C_n).

When the constant sum of an A-magic graph is zero the graph is called *zero-sum A-magic*. The null set $N(G)$ of a graph G is the set of all positive integers h such that G is zero-sum Z_h-magic. Akbari, Ghareghani, Khosrovshahi, and Zare [155] and Akbari,

Kano, and Zare [156] proved that the null set $N(G)$ of an r-regular graph G, $r \geq 3$, does not contain the numbers 2, 3 and 4. Akbari, Rahmati, and Zare [157] proved the following: if G is an even regular graph then G is zero-sum Z_h-magic for all h; if G is an odd r-regular graph, $r \geq 3$ and $r \neq 5$ then $N(G)$ contains all positive integers except 2 and 4; if an odd regular graph is also 2-edge connected then $N(G)$ contains all positive integers except 2; and a 2-edge connected bipartite graph is zero-sum Z_h-magic for $h \geq 6$. They also determine the null set of 2-edge connected bipartite graphs, describe the structure of some odd regular graphs, $r \geq 3$, that are not zero-sum 4-magic, and describe the structure of some 2-edge connected bipartite graphs that are not zero-sum Z_h-magic for $h = 2, 3$, and 4. They conjecture that every 5-regular graph admits a zero-sum 3-magic labeling.

In [2186] Lee, Saba, Salehi, and Sun investigate graphs that are A-magic where $A = V_4 \approx Z_2 \oplus Z_2$ is the Klein four-group. Many of theorems are special cases of the results of Shiu, Lam, and Sun [3534] given in the previous paragraph. They also prove the following are V_4-magic: a tree if and only if every vertex has odd degree; the star $K_{1,n}$ if and only if n is odd; $K_{m,n}$ for all $m, n \geq 2$; $K_n - e$ (edge deleted K_n) when $n > 3$; even cycles with k pendent edges if and only if k is even; odd cycles with k pendent edges if and only if k is odd; wheels; $C_n + \overline{K_2}$; generalized theta graphs; graphs that are copies of C_n that share a common edge; and $G + \overline{K_2}$ whenever G is V_4-magic. In [2262] Libeeshkumar and Anil Kumar discussed induced V_4-magic labelings of some cycle related graphs.

In [813] Choi, Georges, and Mauro explore Z_2^k-magic graphs in terms of even edge-coverings, graph parity, factorability, and nowhere-zero 4-flows. They prove that the minimum k such that bridgeless G is zero-sum Z_2^k-magic is equal to the minimum number of even subgraphs that cover the edges of G, known to be at most 3. They also show that bridgeless G is zero-sum Z_2^k-magic for all $k \geq 2$ if and only if G has a nowhere-zero 4-flow, and that G is zero-sum Z_2^k-magic for all $k \geq 2$ if G is Hamiltonian, bridgeless planar, or isomorphic to a bridgeless complete multipartite graph, and establish equivalent conditions for graphs of even order with bridges to be Z_2^k-magic for all $k \geq 4$. In [1238] Georges, Mauro, and Wang utilized well-known results on edge-colorings in order to construct infinite families that are V_4-magic but not Z_4-magic. In [2263] Libeeshkumar and Kumar discussed some induced V_4-magic labelings of some subdivision graphs.

Low and Roberts [2360] use the combinatorial nullstellensatz to show the existence of Z_p-magic labelings (prime $p \geq 3$) for various graphs, without having to construct the Z_p-magic labelings. They give examples illustrating the usefulness and limitations in applying the combinatorial nullstellensatz to the integer-magic labeling problem. They show various classes of graphs have Z_3-magic labelings. Baskar Babujee and Shobana [585] prove that the following graphs have Z_3-magic labelings: C_{2n}; K_n $(n \geq 4)$; $K_{m,2m}$ $(m \geq 3)$; ladders $P_n \times P_2$ $(n \geq 4)$; bistars $B_{3n-1,3n-1}$; and cyclic, dihedral, and symmetric Cayley digraphs for certain generating sets. Chalise and Low [764] used the combinatorial nullstellensatz to study Z_p-magic and Z_p-antimagic graph labeling problems. They showed that, in several cases, the coefficients of monomials admit combinatorial interpretations in terms of structural properties of the graph. Siddiqui [3578] proved that generalized prisms, generalized antiprisms, fans and friendship graphs are Z_{3k}-magic for $k \geq 1$. In [819] Chou and Lee investigated Z_3-magic graphs. [764] new

Chou and Lee [819] showed that every graph is an induced subgraph of an A-magic graph for any nontrivial Abelian group A. Thus it is impossible to find a Kuratowski type characterization of A-magic graphs. Low and Lee [2358] have shown that if a graph is A_1-magic then it is A_2-magic for any subgroup A_2 of A_1 and for any nontrivial Abelian group A every Eulerian graph of even size is A-magic. For a connected graph G, Low and Lee define $T(G)$ to be the graph obtained from G by adding a disjoint uv path of length 2 for every pair of adjacent vertices u and v. They prove that for every finite nontrivial Abelian group A the graphs $T(P_{2k})$ and $T(K_{1,2n+1})$ are A-magic. Shiu and Low [3545] show that $K_{k_1,k_2,\ldots,k_n}(k_i \geq 2)$ is A-magic, for all A where $|A| \geq 3$. In [3550] Shiu and Low analyze the A-magic property for complete n-partite graphs and composition graphs with deleted edges. Lee, Salehi and Sun [2189] have shown that for $m, n \geq 3$ the double star $DS(m, n)$ is Z-magic if and only if $m = n$.

S. M. Lee [2148] calls a graph G *fully magic* if it is A-magic for all nontrivial Abelian groups A. Low and Lee [2358] showed that if G is an Eulerian graph of even size, then G is fully magic. In [2148] Lee gives several constructions that produce infinite families of fully magic graphs and proves that every graph is an induced subgraph of a fully magic graph.

In [2074] Kwong and Lee call the set of all k for which a graph is Z_k-magic the *integer-magic spectrum* of the graph. They investigate the integer-magic spectra of the coronas of some specific graphs including paths, cycles, complete graphs, and stars. Low and Sue [2363] have obtained some results on the integer-magic spectra of tessellation graphs. Shiu and Low [3546] provide the integer-magic spectra of sun graphs. Chopra and Lee [817] determined the integer-magic spectra of all graphs consisting of any number of pairwise disjoint paths with common end vertices (that is, generalized theta graphs). Low and Lee [2358] show that Eulerian graphs of even size are A-magic for every finite nontrivial Abelian group A whereas Wen and Lee [4178] provide two families of Eularian graphs that are not A-magic for every finite nontrivial Abelian group A and eight infinite families of Eulerian graphs of odd sizes that are A-magic for every finite nontrivial Abelian group A. Low and Lee [2358] also prove that if A is an Abelian group and G and H are A-magic, then so are $G \times H$ and the lexicographic product of G and H. Low and Shiu [2362] prove: $K_{1,n} \times K_{1,n}$ has a Z_{n+1}-magic labeling with magic constant 0; if $G \times H$ is Z_2-magic, then so are G and H; if G is Z_m-magic and H is Z_n-magic, then the integer-magic spectra of $G \times H$ contains all common multiples of m and n; if n is even and $k_i \geq 3$ then the integer-magic spectra of $P_{k_1} \times P_{k_2} \times \cdots \times P_{k_n} = \{3, 4, 5, \ldots\}$. In [3548] Shiu and Low determine all positive integers k for which fans and wheels have a Z_k-magic labeling with magic constant 0. Shiu and Low [3549] determined for which $k \geq 2$ a connected bicyclic graph without a pendent has a Z_k-magic labeling.

Jeyanthi and Jeya Daisy [1675] prove that P_n^2 $(n > 4)$, C_n^2, the total graph of C_n, and the splitting graph of C_{2n} are Z_k-magic graphs. They also prove: the splitting graph of C_n is Z_k-magic when n is even and n is odd and k is even, the middle graph of C_n is Z_k-magic when n and k are odd, the $m\Delta_{2n}$-snake graph is Z_k-magic when $k > m$, the graph obtained by joining the vertices u_i and u_{i+1} of C_n by a path of length m_i for $1 \leq i \leq n-1$, and u_1 and u_n by a path of length m_n is Z_k-magic if either all $m_1, m_2, \ldots, m_n$ are even

or all are odd. In [1676] Jeyanthi and Jeya Daisy prove total graphs of the paths, flower graphs, and $C_m \times P_n$ are Z_k-magic. They also prove closed helms are Z_k-magic when $k > 4$ is even, lotuses inside a circle are Z_{4k}-magic, and graphs consisting of two cycles with a common edge are Z_k-magic when at least one cycle is even. In [1681] Jeyanthi prove the following graphs are Z_k-magic: two odd cycles connected by a path; the graph obtained by identifying a vertex of C_n with a pendent vertex of a star, m-splitting graphs of paths, and m-middle graphs of paths. They prove that if G is Z_m-magic with magic constant a then $G \odot \overline{K_m}$ is Z_m-magic.

Jeyanthi and Jeya Daisy [1674] prove that the subdivision graphs of the following families of graphs are Z_k-magic: ladders, triangular ladders, the shadow graph of paths, the total graph of paths, flowers, generalized prisms $C_m \times P_n$ for m even, $m\Delta_n$-snakes, lotuses inside a circle, the square graph of paths, gears of even cycles, closed helms of even cycles, and antiprisms A_n^m for m even.

Let G be a graph and let $G_1, G_2, \ldots, G_n$ be $n \geq 2$ copies of G. The graph obtained by replacing each endpoint vertex of $K_{1,n}$ by the graphs $G_1, G_2, \ldots, G_n$ is called the *open star* of G. Jeyanthi and Jeya Daisy [1678] proved that the open star graphs of shells, flowers, double wheels, cylinders, wheels, generalized Peterson graphs, lotuses inside a circle, and closed helms are Z_k-magic graphs. They also prove that the super subdivision of any graph is Z_k-magic.

Jeyanthi and Jeya Daisy [1679] proved that the path union of $n \geq 2$ copies of the following families of graphs are Z_k-magic: odd cycles; generalized Peterson graphs $P(r, m)$ when r is odd and $1 \leq m \leq \frac{r}{2}$; shell graphs S_r when $r > 3$; wheels W_r when $r > 3$; closed helms CH_r when (i) $r > 3$ is odd and (ii) r is even and k is even; double wheels DW_r when $r > 3$ is odd; flowers Fl_r when $r > 2$; $C_r \times P_2$ when r is odd; total graphs of paths $T(P_r)$ for all $n, r > 4$; lotuses inside a circle LC_r when $r > 3$; and $C_r \odot K_1$ for odd r.

Jeyanthi and Jeya Daisy [1680] proved that the following graphs are k-magic: shell graphs S_n when n is odd or n is even and k is even; generalized Jahangir graphs $J_{n,s}$ when n and s have the same parity or n is even, s is odd, and k is even; $(P_n + P_1) \times P_2$ when n is odd; double wheels $2C_n + K_1$; mongolian tents $M(m, n)$ when m is even; flower snark graphs; slanting ladders (that is, graphs obtained from two paths $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$ by joining each u_i with v_{i+1}, $1 \leq i \leq n-1$) when n is even; double step grid graphs; double arrow graphs obtained from $P_m \times P_n$ by joining a new vertex with the m vertices of the first copy of P_m and another new vertex with the m vertices of the last copy of P_m when m is even; semi Jahangir graphs (the connected graph with vertex set $\{p, x_i, y_k : 1 \leq i \leq n+1, 1 \leq k \leq n\}$ and the edge set $\{px_i : 1 \leq i \leq n+1\} \cup \{x_iy_i : 1 \leq i \leq n\} \cup \{y_ix_{i+1} : 1 \leq i \leq n\}$); graphs obtained by connecting double wheels DW_{n_1} and DW_{n_2} by a path when n_1 and n_2 are odd; graphs obtained by joining two copies of shell graphs by a path; and the splitting graph of a Z_k magic graph with magic constant 0.

Let G be a graph with n vertices $\{u_1, u_2, \ldots, u_n\}$ and consider n copies of G, $G_1, G_2, \ldots, G_n$, with vertex sets $V(G_i) = \{u_i^j : 1 \leq i \leq n, 1 \leq j \leq n\}$. The *cycle of a graph* G, denoted by $C(n.G)$, is obtained by identifying the vertex u_1^j of G_j with u_i of G for $1 \leq i \leq n, 1 \leq j \leq n$. Jeyanthi and Jeya Daisy [1681] prove that the following graphs are Z_k-magic: $C(n.C_r)$ except r is even, n is odd, and k is odd; generalized Peter-

son graphs $C(n.P(r,m))$ except r is even, n is odd, and k is odd; cycles of shell graphs; cycles of wheel graphs; cycles of closed helms; cycles of double wheels $C(n.DW_r)$ except r is even, n is odd, and k is odd; cycles of triangular ladder graphs; cycles of flower graphs; and cycles of lotus inside a circle graphs. Jeyanthi and Jeya Daisy [1681] also prove that if G is Z_k-magic then $C(n.G)$ is Z_k-magic if n or k are even.

Shiu and Low [3547] have introduced the notion of ring-magic as follows. Given a commutative ring R with unity, a graph G is called *R-ring-magic* if there exists a labeling f of the edges of G with the nonzero elements of R such that the vertex labeling f^+ defined by $f^+(v) = \Sigma f(vu)$ over all edges vu and vertex labeling $f^\times$ defined by $f^\times(v) = \Pi f(vu)$ over all edges vu are constant. They give some results about R-ring-magic graphs.

In [745] Cahit says that a graph $G(p,q)$ is *total magic cordial* (TMC) provided there is a mapping f from $V(G) \cup E(G)$ to $\{0,1\}$ such that $(f(a) + f(b) + f(ab)) \bmod 2$ is a constant modulo 2 for all edges $ab \in E(G)$ and $|f(0) - f(1)| \leq 1$ where $f(0)$ denotes the sum of the number of vertices labeled with 0 and the number of edges labeled with 0 and $f(1)$ denotes the sum of the number of vertices labeled with 1 and the number of edges labeled with 1. He says a graph G is *total sequential cordial* (TSC) if there is a mapping f from $V(G) \cup E(G)$ to $\{0,1\}$ such that for each edge $e = ab$ with $f(e) = |f(a) - f(b)|$ it is true that $|f(0) - f(1)| \leq 1$ where $f(0)$ denotes the sum of the number of vertices labeled with 0 and the number of edges labeled with 0 and $f(1)$ denotes the sum of the number of vertices labeled with 1 and the number of edges labeled with 1. He proves that the following graphs have a TMC labeling: $K_{m,n}$ $(m,n > 1)$, trees, cordial graphs, and K_n if and only if $n = 2, 3, 5$, or 6. He also proves that the following graphs have a TSC labeling: trees; cycles; complete bipartite graphs; friendship graphs; cordial graphs; cubic graphs other than K_4; wheels W_n $(n > 3)$; K_{4k+1} if and only if $k \geq 1$ and $\sqrt{k}$ is an integer; K_{4k+2} if and only if $\sqrt{4k+1}$ is an integer; K_{4k} if and only if $\sqrt{4k+1}$ is an integer; and K_{4k+3} if and only if $\sqrt{k+1}$ is an integer. In [1663] Jeyanthi, Angel Benseera, and Cahit prove mP_2 is TMC if $m \not\equiv 2 \pmod 4$, mP_n is TMC for all $m \geq 1$ and $n \geq 3$, and obtain partial results about TMC labelings of mK_n. Neela and Selvaraj proved that the complete tripartite graphs are TMC and complete multipartite graphs are TMC when the partite sets have even sizes

The zero divisor graph of the ring Z_n has as its vertex set all elements in Z_n that are relatively prime to n. For vertices x and y, xy is an edge if $xy = 0$. For distinct odd primes p and q, Lourdusamy, Veronisha, and Beaula [2347] proved the zero-divisor graphs of Z_{2p}, Z_{4p}, and Z_{pq} are totally magic cordial. They also showed the join of two zero-divisor graphs and the product of zero-divisor graphs of Z_m and Z_n for special cases of m and n are totally magic cordial. They further provided some results about total magic cordialness of zero-divisor graphs of the form $\overline{Z_m + Z_n}$ and $\overline{Z_m} + \overline{Z_n}$ for special values of m and n.

Parameswari and Jayalakshmi e [2767] proved that *octopus* graphs (the join of a fan F_n and $K_{1,n}$ $(n > 1)$), vanessa graphs $2F_n + K_{1,n}$, lilly graphs $2K_{1,n} + 2P_n$, and lotus graphs admit cordial totally magic labelings. Parameswari [2766] proved that the square graphs and shadow graphs of bistars admit cordial totally magic labelings. In [2769] and [2770] Parameswari and Rajeswari proved that Paley digraphs, fans, gears, and shadow graphs

of paths and stars are total magic cordial. Neela and Selvaraj [2664] showed that complete tripartite graphs and complete multipartite graphs are total magic cordial when the partite sets are of even sizes.

Vaidya and Barasara [3953] proved that every graph can be embedded as an induced subgraph of a totally magic cordial graph thereby ruling out any possibility of obtaining any forbidden subgraph characterization for totally magic cordial graphs. They also proved that every connected graph can be embedded as an induced subgraph of a totally magic cordial connected graph and every planar graph can be embedded as an induced subgraph of a totally magic cordial planner graph. Similar results were also obtained for total sequential cordial labeling.

Jeyanthi and Angel Benseera [1661] investigated the existence of TMC labelings of the one-point unions of copies of cycles, complete graphs and wheels. In [1662] Jeyanthi and Angel Benseera prove that if $G_i(p_i, q_i)$, $i = 1, 2, 3, \ldots, n$ are totally magic cordial graphs with $C = 0$ such that $p_i + q_i$, $i = 1, 2, 3, ..., n$ are even, and $|p_i - 2m_i| \leq 1$, where m_i is the number of vertices labeled with 0 in G_i, $i = 1, 2, \ldots, n$, then $G_1 + G_2 + \cdots + G_n$ is TMC; if G is an odd graph with $p + q \equiv 2$ (mod 4), then G is not TMC; fans F_n are TMC for $n \geq 2$; wheels W_n $(n \geq 3)$ are TMC if and only if $n \not\equiv 3 \pmod 4$; mW_{4t+3} is TMC if and only if m is even; mW_n is TMC if $n \not\equiv 3 \pmod 4$; $C_n + \overline{K}_{2m+1}$ is TMC if and only if $n \not\equiv 3 \pmod 4$; $C_{2n+1} \odot \overline{K}_m$ is TMC if and only if m is odd; the disjoint union of $K_{1,m}$ and $K_{1,n}$ is TMC if and only if m or n is even.

For a bijection $f : V(G) \cup E(G) \to Z_k$ such that for each edge $uv \in E(G)$, $f(u) + f(v) + f(uv)$ is constant $\pmod k$ $n_f(i)$ denotes the number vertices and edges labeled by i under f. If $|n_f(i) - n_f(j)| \leq 1$ for all $0 \leq i < j \leq k - 1$, f is called a *k-totally magic cordial* labeling of G. A graph is said to be *k-totally magic cordial* if it admits a k-totally magic cordial labeling. In [1664] Jeyanthi, Angel Benseera, and Lau provide some ways to construct new families of k-totally magic cordial (k-TMC) graphs from a known k-totally magic cordial graph. Let G (respectively, H) be a (p, q)-graph (respectively, an (n, m)-graph) that admits a k-TMC labeling f (respectively, g) with constant C such that $n_f(i)$ and $v_f(i) = \frac{p}{k}$ (or $n_g(i)$ and $v_g(i) = \frac{n}{k}$) are constants for all $0 \leq i \leq k - 1$, they show that $G + H$ also admits a k-TMC labeling with constant C. They prove the following. If G is an edge magic total graph, then G is k-TMC for $k \geq 2$; if G is an odd graph with $p + q \equiv k \pmod{2k}$ and $k \equiv 2 \pmod 4$, then G is not k-TMC; if $n \equiv 7 \pmod 8$, $K_n \odot K_1$ is not $2n$-TMC; if $n \equiv 2 \pmod 4$, $C_n \odot C_3$ is not n-TMC; if $n \equiv 1 \pmod 2$, $C_n \odot K_5$ is not $2n$-TMC; if $n \equiv 2 \pmod 4$, $C_n \times P_2$ is not n-TMC; K_n $(n \geq 3)$ is n-TMC; $K_n \odot K_1$ $(n \geq 3)$ is n-TMC; S_n is n-TMC for all $n \geq 1$; $K_{m,n}$ $(m \geq n \geq 2)$ is both m-TMC and n-TMC; W_n is n-TMC for all odd $n \geq 3$ and is 3-TMC for $n \equiv 0 \pmod 6$; mK_n $(n \geq 2)$ is n-TMC if $n \geq 3$ is odd; $K_n + K_n$ is n-TMC if $n \geq 3$ is odd; $S_n + S_n$ $(n \geq 1)$ is $(n+1)$-TMC; and if $m \geq 3$ and n is odd, $C_n \times P_m$ $(n \geq 3)$ is n-TMC. In [1666] Jeyanthi, Angel Benseera, and Lau call a graph G hypo-k-TMC if $G - \{v\}$ is k-TMC for each vertex v in $V(G)$ and establish that some families of graphs admit and do not admit hypo-k-TMC labeling.

A graph $G(V, E)$ where $V = \{v_i, 1 \leq i \leq n\}$ and $E = \{v_i v_{i+1}, 1 \leq i \leq n\}$ is 0-*edge magic* if there exists a bijection $f : V(G) \to \{1, -1\}$ such that the induced edge la-

beling defined by $f^*(uv) = f(u) + f(v)$ is 0 for all $uv \in E$. Paths, cycles, complete n-ary pseudo trees, $P_m \times C_n$ where $n \equiv 0 \pmod 2$, Q_n, the graph C_m attached to mK_1, $m \equiv 0 \pmod 2$, friendship graphs $C_n^{(m)}$, and the graph $P_m \times P_m \times P_m$ are 0-edge magic graphs [2021], [1591], [3895]. Jayapriya [1589] proved the splitting graphs $spl(P_n)$, $spl(C_n)$, $spl(K_{1,n})$, $spl(B_{m,n})$, and splitting graph of any tree admits 0-edge magic labelings. Laurejas and Pedrano [2128] determine the 0-edge magic labeling of $P_m \times P_n$, $C_m \times C_n$, and the generalized Petersen graph. They also prove that odd cycles are not 0-edge magic.

A *binary magic total* labeling of a graph G is a function $f : V(G) \cup E(G) \to \{0, 1\}$ such that $f(a) + f(b) + f(ab) \equiv C \pmod 2$ for all $ab \in E(G)$. Jeyanthi and Angel Benseera [1665] define the *totally magic cordial deficiency* of G as the minimum number of vertices taken over all binary magic total labeling of G that is necessary to add so that the resulting graph is totally magic cordial. The totally magic cordial deficiency of G is denoted by $\mu_T(G)$. They provide $\mu_T(K_n)$ for some cases.

Let G be a graph rooted at a vertex u and f_i be a binary magic total labeling of G and $f_i(u) = 0$ for $i = 1, 2, \ldots, k$ and $n_{f_i}(0) = \alpha_i$, $n_{f_i}(1) = \beta_i$ for $i = 1, 2, \ldots, k$. Jeyanthi and Angel Benseera [1665] determine the totally magic cordial deficiency of the one-point union $G^{(n)}$ of n copies of G. They show that for $n \equiv 3 \pmod 4$ the totally magic cordial deficiency of W_n, $W_n^{(4t+1)}$, $W_{4t+1}^{(n)}$ and $C_n + \overline{K}_{2m+1}$ is 1; for m odd, $\mu_T(mW_{4t+3}) = 1$; and for $n \equiv 1 \pmod 4$, $\mu_T(K_4^{(n)}) = 1$.

In 2001, Simanjuntak, Rodgers, and Miller [2503] defined a *1-vertex magic* (also known as *distance magic* labeling vertex labeling of $G(V, E)$ as a bijection from V to $\{1, 2, \ldots, |V|\}$ with the property that there is a constant k such that at any vertex v the sum $\sum f(u)$ taken over all neighbors of v is k. Among their results are: $H \times \overline{K}_{2k}$ has a 1-vertex-magic vertex labeling for any regular graph H; the symmetric complete multipartite graph with p parts, each of which contains n vertices, has a 1-vertex-magic vertex labeling if and only if whenever n is odd, p is also odd; P_n has a 1-vertex-magic vertex labeling if and only if $n = 1$ or 3; C_n has a 1-vertex-magic vertex labeling if and only if $n = 4$; K_n has a 1-vertex-magic vertex labeling if and only if $n = 1$; W_n has a 1-vertex-magic vertex labeling if and only if $n = 4$; a tree has a 1-vertex-magic vertex labeling if and only if it is P_1 or P_3; and r-regular graphs with r odd do not have a 1-vertex-magic vertex labeling.

Miller, Rogers, and Simanjuntak [2503] the complete p-partite ($p > 1$) graph $K_{n,n,\ldots,n}$ ($n > 1$) has a 1-vertex-magic vertex labeling if and only if either n is even or np is odd. Shafiq, Ali, Simanjuntak [3459] proved $mK_{n,n,\ldots,n}$ has a 1-vertex-magic vertex labeling if n is even or mnp is odd and $m \geq 1, n > 1$, $p > 1$; and $mK_{n,n,\ldots,n}$ does not have a 1-vertex-magic vertex labeling if np is odd, $p \equiv 3 \pmod 4$, and m is even.

Recall if $V(G) = \{v_1, v_2, \ldots, v_p\}$ is the vertex set of a graph G and $H_1, H_2, \ldots, H_p$ are isomorphic copies of a graph H, then $G[H]$ is the graph obtained from G by replacing each vertex v_i of G by H_i and joining every vertex in H_i to every neighbor of v_i. Shafiq, Ali, Simanjuntak [3459] proved if G is an r-regular graph ($r \geq 1$) then $G[C_n]$ has a 1-vertex-magic vertex labeling if and only if $n = 4$. They also prove that for $m \geq 1$ and $n > 1$, $mC_p[\overline{K_n}]$ has 1-vertex-magic vertex labeling if and only if either n is even or mnp is odd or n is odd and $p \equiv 3 \pmod 4$.

For a graph G Jeyanthi and Angel Benseera [1660] define a function f from $V(G) \cup E(G)$ to $\{0,1\}$ to be a *totally vertex-magic cordial labeling* (TVMC) with a constant C if $f(a)+\sum_{b\in N(a)} f(ab) \equiv C \pmod 2$ for all vertices $a \in V(G)$ and $|n_f(0)-n_f(1)| \leq 1$, where $N(a)$ is the set of vertices adjacent to the vertex a and $n_f(i)$ is the sum of the number of vertices and edges with label i. They prove the following graphs have totally vertex-magic cordial labelings: vertex-magic total graphs; trees; K_n; $K_{m,n}$ whenever $|m-n| \leq 1$; $P_n + P_2$; friendship graphs with C = 0; and flower graphs Fl_n for $n \geq 3$ with C= 0. They also proved that if G is TVMC with $C = 1$, then the graph obtained by identifying any vertex of G with any vertex of a tree is TVMC with $C = 1$; if G is a (p, q) graph with $|p - q| \leq 1$, then G is TVMC with $C = 1$; and if $G(p, q)$ is a TVMC graph with constant $C = 0$ where p is odd, then $G + \overline{K_{2m}}$ is TVMC with $C = 1$ if m is odd and with $C = 0$ if m is even.

Jeyanthi, Angel Benseera, and Immaculate Mary [1659] showed that the following graphs have totally magic cordial labelings: (p, q) graphs with $|p - q| \leq 1$; flower graphs Fl_n for $n \geq 3$; ladders; and graphs obtained by identifying a vertex of C_m with each vertex of C_n. They also proved that if $G_1(p_1, q_1)$ and $G_2(p_2, q_2)$ are two disjoint totally magic cordial graphs with $p_1 = q_1$ or $p_2 = q_2$ then $G_1 \cup G_2$ is totally magic cordial. In Theorem 10 in [745] Cahit stated that K_n is totally magic cordial if and only if $n \in \{2, 3, 5, 6\}$. Jeyanthi and Angel Benseera [1665] proved that K_n is totally magic cordial if and only if $\sqrt{4k+1}$ has an integer value when $n = 4k$; $\sqrt{k+1}$ or $\sqrt{k}$ have an integer value when $n = 4k + 1$; $\sqrt{4k+5}$ or $\sqrt{4k+1}$ have an integer value when $n = 4k + 2$; or $\sqrt{k+1}$ has an integer value when $n = 4k + 3$.

A graph G is said to have a *totally magic cordial* TMC labeling with constant C if there exists a mapping $f : V(G) \cup E(G) \to \{0, 1\}$ such that $f(a) + f(b) + f(ab) \equiv C \pmod 2$ for all $ab \in E(G)$ and $|n_f(0) - n_f(1)| \leq 1$, where $n_f(i)$ $(i = 0, 1)$ is the sum of the number of vertices and edges with label i. In [1662] Jeyanthi and Angel Benseera prove that if $G_i(p_i, q_i)$, $i = 1, 2, 3, \ldots, n$ are totally magic cordial graphs with $C = 0$ such that $p_i + q_i$, $i = 1, 2, 3, \ldots, n$ are even, and $|p_i - 2m_i| \leq 1$, where m_i is the number of vertices labeled with 0 in G_i, $i = 1, 2, \ldots, n$, then $G_1 + G_2 + \cdots + G_n$ is TMC. They also prove the following. If G be an odd graph with $p + q \equiv 2 \pmod 4$, then G is not TMC; fan graph F_n is TMC for $n \geq 2$; the wheel graph W_n $(n \geq 3)$ is TMC if and only if $n \not\equiv 3 \pmod 4$; mW_{4t+3} is TMC if and only if m is even; mW_n is TMC if $n \not\equiv 3 \pmod 4$ and $m \geq 1$; $C_n + \overline{K}_{2m+1}$ is TMC if and only if $n \not\equiv 3 \pmod 4$; $C_{2n+1} \odot \overline{K}_m$ is TMC if and only if m is odd; and the disjoint union of $K_{1,m}$ and $K_{1,n}$ is TMC if and only if m or n is even.

Balbuena, Barker, Lin, Miller, and Sugeng [468] call a vertex-magic total labeling of a graph $G(V, E)$ an *a-vertex consecutive magic labeling* if the vertex labels are $\{a + 1, a + 2, \ldots, a + |V|\}$ where $0 \leq a \leq |E|$. They prove: if a tree of order n has an a-vertex consecutive magic labeling then n is odd and $a = n - 1$; if G has an a-vertex consecutive magic labeling with n vertices and $e = n$ edges, then n is odd and if G has minimum degree 1, then $a = (n + 1)/2$ or $a = n$; if G has an a-vertex consecutive magic labeling with n vertices and e edges such that $2a \leq e$ and $2e \geq \sqrt{6}n - 1$, then the minimum degree of G is at least 2; if a 2-regular graph of order n has an a-vertex consecutive magic labeling, then n is odd and $a = 0$ or n; and if a r-regular graph of order n has an a-vertex

consecutive magic labeling, then n and r have opposite parities.

Balbuena et al. also call a vertex-magic total labeling of a graph $G(V, E)$ a *b-edge consecutive magic labeling* if the edge labels are $\{b+1, b+2, \ldots, b+|E|\}$ where $0 \leq b \leq |V|$. They prove: if G has n vertices and e edges and has a b-edge consecutive magic labeling and one isolated vertex, then $b = 0$ and $(n-1)^2 + n^2 = (2e+1)^2$; if a tree with odd order has a b-edge consecutive magic labeling then $b = 0$; if a tree with even order has a b-edge consecutive magic labeling then it is P_4; a graph with n vertices and e edges such that $e \geq 7n/4$ and $b \geq n/4$ and a b-edge consecutive magic labeling has minimum degree 2; if a 2-regular graph of order n has a b-edge consecutive magic labeling, then n is odd and $b = 0$ or $b = n$; and if a r-regular graph of order n has an b-edge consecutive magic labeling, then n and r have opposite parities.

Sugeng and Miller [3731] prove: If (V, E) has an a-vertex consecutive edge magic labeling, where $a \neq 0$ and $a \neq |E|$, then G is disconnected; if (V, E) has an a-vertex consecutive edge magic labeling, where $a \neq 0$ and $a \neq |E|$, then G cannot be the union of three trees with more than one vertex each; for each nonnegative a and each positive n, there is an a-vertex consecutive edge magic labeling with n vertices; the union of r stars and a set of $r-1$ isolated vertices has an s-vertex consecutive edge magic labeling, where s is the minimum order of the stars; for every b every caterpillar has a b-edge consecutive edge magic labeling; if a connected graph G with n vertices has a b-edge consecutive edge magic labeling where $1 \leq b \leq n-1$, then G is a tree; the union of r stars and a set of $r-1$ isolated vertices has an r-edge consecutive edge magic labeling.

Baskar Babujee, Vishnupriya, and Jagadesh [589] introduced a labeling called *a-vertex consecutive edge-bimagic total* as a graph $G(V, E)$ for which there are two positive integers k_1 and k_2 and a bijection f from $V \cup E$ to $\{1, 2, \ldots, |V| + |E|\}$ such that $f(u) + f(v) + f(uv) = k_1$ or k_2 for all edges uv and $f(V) = \{a+1, a+2, \ldots, a+|V|\}$, $0 \leq a \leq |V|$. They proved the following graphs have such labelings: P_n, $K_{1,n}$, combs, bistars $B_{m,n}$, trees obtained by adding a pendent edge to a vertex adjacent to the end point of a path, trees obtained by joining the centers of two stars with a path of length 2, trees obtained from P_5 by identifying the center of a copy $K_{1,n}$ with the two end vertices and the middle vertex. In [575] Baskar Babujee and Jagadesh proved that cycles, fans, wheels, and gear graphs have a-vertex consecutive edge-bimagic total labelings. Baskar Babujee, Jagadesh, Vishnupriya [578] study the properties of a-vertex consecutive edge-bimagic total labeling for $P_3 \odot K_{1,2n}$, $P_n + \overline{K_2}$ (n is odd and $n \geq 3$), $(P_2 \cup mK_1) + \overline{K_2}$, $(P_2 + mK_1)$ $(m \geq 2)$, C_n, fans $P_n + K_1$, double fans $P_n + 2K_1$, and graphs obtained by appending a path of length at least 2 to a vertex of C_3. Baskar Babujee and Jagadesh [576] prove the following graphs have a-vertex consecutive edge-bimagic total labelings: $2P_n$ $(n \geq 2)$, $P_n \cup P_{n+1}$ $(n \geq 2)$, $K_{2,n}$, $C_n \odot K_1$, and that $C_3 \cup K_{1,n}$ an a-vertex consecutive edge-bimagic labeling for $a = n + 3$.

Vishnupriya, Manimekalai, and Baskar Babujee [4117] define a labeling f of a graph $G(p, q)$ to be a *edge-bimagic total labeling* if there exists a bijection f from $V(G) \cup E(G) \to \{1, 2, \ldots, p+q\}$ such that for each edge $e = (u, v) \in E(G)$ we have $f(u) + f(e) + f(v) = k_1$ or k_2, where k_1 and k_2 are two constants. They provide edge-bimagic total labelings for $B_{m,n}$, $K_{1,n,n}$, and trees obtained from a path by appending an edge to one of the vertices

adjacent to an endpoint of the path. An edge-bimagic total labeling is $G(V, E)$ is called an *a-vertex consecutive edge-bimagic total labeling* if the vertex labels are $\{a+1, a+2, \ldots, a+|V|\}$ where $0 \leq a \leq |E|$. Baskar Babujee and Jagadesh [573] prove the following graphs a-vertex consecutive edge-bimagic total labelings: the trees obtained from $K_{1,n}$ by adding a new pendent edge to each of the existing n pendent vertices; the trees obtained by adding a pendent path of length 2 to each of the n pendent vertices of $K_{1,n}$; the graphs obtained by joining the centers of two copies of identical stars by a path of length 2; and the trees obtained from a path by adding new pendent edges to one pendent vertex of the path. Baskar Babujee, Vishnupriya, and Jagadesh [589] proved the following graphs have such labelings: P_n, $K_{1,n}$, combs, bistars $B_{m,n}$, trees obtained by adding a pendent edge to a vertex adjacent to the end point of a path, trees obtained by joining the centers of two stars with a path of length 2, trees obtained from P_5 by identifying the center of a copy $K_{1,n}$ with the two end vertices and the middle vertex. In [575] Baskar Babujee and Jagadesh proved that cycles, fans, wheels, and gear graphs have a-vertex consecutive edge-bimagic total labelings. Baskar Babujee, Jagadesh, Vishnupriya [578] study the properties of a-vertex consecutive edge-bimagic total labeling for $P_3 \odot K_{1,2n}$, $P_n + \overline{K_2}$ (n is odd and $n \geq 3$), $(P_2 \cup mK_1) + \overline{K_2}$, $(P_2 + mK_1)$ $(m \geq 2), C_n$, fans $P_n + K_1$, double fans $P_n + 2K_1$, and graphs obtained by appending a path of length at least 2 to a vertex of C_3.

Vishnupriya, Manimekalai, and Baskar Babujee [4117] prove that bistars, trees obtained by adding a pendent edge to a vertex adjacent to the end point of a path, and trees obtained subdividing each edge of a star have edge-bimagic total labelings. Prathap and Baskar Babujee [3036] obtain all possible edge magic total labelings and edge-bimagic total labelings for the star $K_{1,n}$. Jayasekaran1 and Flower [1595] proved that the shadow graph and the splitting graph of paths stars and cycles are edge trimagic total and super edge trimagic total.

Marr, Ochel, and Perez [2444] say a digraph D with v vertices and e directed edges has an *in-magic total labeling* if there exists a bijective function λ from $V(D) \cup E(D)$ to $\{1, 2, \ldots, v+e\}$ such that for every vertex x we have $\lambda(x) + \sum \lambda(y, x) = k$ for some integer k, where the sum is taken over all directed edges (y, x). They provide such labelings for trees and cycles and discuss some relationships between this labeling and other digraph labelings.

Let D be a digraph of order p and size q. For an integer $k \geq 1$ and $v \in V(D)$, let $w_k(v) = \sum_f(e)$ over the set of all in-arcs that are at distance at most k from v. A *V_k-super vertex in-magic labeling* (V_k-SVIML) of D is an one-to-one onto function f from $V(D) \cup A(D)$ to $\{1, 2, \ldots, p+q\}$ such that $f(V(D)) = \{1, 2, \ldots, p\}$ and for every $v \in V(D)$, $f(v) + w_k(v) = M$ for some positive integer M. A digraph that admits a V_k-SVIML is called *V_k-super vertex in-magic* (V_k-SVIM). In [2605] Mutharasu, Mary Bernard, Duraisamy kumar [2605] study some properties of V_k-SVIML in digraphs. They characterized the digraphs that are V_k-SVIM and find the magic constant for E_k-regular digraphs. They furthermore characterized the unidirectional cycles and union of unidirectional cycles which are V_2-SVIML. More results about magic total labelings of digraphs are provided in [2608], [3594], [2059], [2441] and [670].

6 Antimagic-type Labelings

6.1 Antimagic Labelings

Hartsfield and Ringel [1378] introduced antimagic graphs in 1990. A graph with q edges is called *antimagic* if its edges can be labeled with $1, 2, \ldots, q$ without repetition such that the sums of the labels of the edges incident to each vertex are distinct.[3] Among the graphs they prove are antimagic are: P_n $(n \geq 3)$, cycles, wheels, and K_n $(n \geq 3)$. T. Wang [4144] has shown that the toroidal grids $C_{n_1} \times C_{n_2} \times \cdots \times C_{n_k}$ are antimagic and, more generally, graphs of the form $G \times C_n$ are antimagic if G is an r-regular antimagic graph with $r > 1$. Cheng [805] proved that all Cartesian products or two or more regular graphs of positive degree are antimagic and that if G is j-regular and H has maximum degree at most k, minimum degree at least one (G and H need not be connected), then $G \times H$ is antimagic provided that j is odd and $j^2 - j \geq 2k$, or j is even and $j^2 > 2k$. Wang and Hsiao [4145] prove the following graphs are antimagic: $G \times P_n$ $(n > 1)$ where G is regular; $G \times K_{1,n}$ where G is regular; compositions $G[H]$ (see §2.3 for the definition) where H is d-regular with $d > 1$; and the Cartesian product of any double star (two stars with an edge joining their centers) and a regular graph. In [804] Cheng proved that $P_{n_1} \times P_{n_2} \times \cdots \times P_{n_t}$ $(t \geq 2)$ and $C_m \times P_n$ are antimagic. In [3647] Solairaju and Arockiasamy prove that various families of subgraphs of grids $P_m \times P_n$ are antimagic. Liang and Zhu [2246] proved that if G is k-regular $(k \geq 2)$, then for any graph H with $|E(H)| \geq |V(H)| - 1 \geq 1$, the Cartesian product $H \times G$ is antimagic. They also showed that if $|E(H)| \geq |V(H)| - 1$ and each connected component of H has a vertex of odd degree, or H has at least $2|V(H)| - 2$ edges, then the prism of H is antimagic. Shang [3483] showed that all spiders are antimagic. Lee, Lin, and Tsai [2142] proved that C_n^2 is antimagic and the vertex sums form a set of successive integers when n is odd. Shang, Lin, and Liaw [3486] show that a star forest containing no S_1 and at most one S_2 as components is antimagic. They also prove that if a star forest mS_2 is antimagic then $m = 1$ and $mS_2 \cup S_n$ $(n \geq 3)$ is antimagic if and only if $m \leq \min\{2n + 1, 2n - 5 + \sqrt{8n^2 - 24n + 17}/2\}$. Wang, Miao, and Li [4156] show that certain graphs with even factors are antimagic. Li [2236] gives antimagic labelings for C_n^k for $k = 2, 3$, and 4. In [4162] Wang and Zhang showed that certain classes of regular graphs of odd degree with particular type of perfect matchings are antimagic. As a by-product, they get that generalized Petersen graphs and a subclass of Cayley graphs of Z_n are antimagic. Deng and Li [923] proved that caterpillars with maximum degree 3 are antimagic and [2366] Lozano, Mora, Seara and Tey proved that every caterpillar is antimagic.

Let $\tau(G)$ to denote the maximum integer such that $G \cup tP_3$ is antimagic for all $t \leq \tau(G)$. The existence of such an integer was proved by Chang, Chen, Li, and Pan in [768]. Shang, Lin, Liaw [3486] and Li [2234] gave tight bounds of $\tau(G)$ for star forests and balanced double stars. Their general bound is also tight for many other families of graphs, including cycles C_n $(3 \leq 9)$ and $2C_3$. Moreover, they discuss properties of $\tau(G)$ and pose

[3] A comprehensive expository treatment of antimagic labelings is given by Bača, Miller, Ryan, and Semaničová-Feňovčíkováin [410].

open questions, including whether their bound is always tight. Chavez, Le, Der-Fen Liu, Shurman [789] generalized the results of Shang et al by providing an upper bound of $\tau(G)$ for all graphs without isolated vertices and P_2 as components. In [2740] Ong, Chen, and Ng investigated antimagic edge labelings for disjoint unions of paths, star related graphs, broom related trees, and 4-cycles. Beaudoire, Bentz, and Picouleau [605] proved that every join graph with at least three vertices is antimagic. [2740] new [605] new

For a graph G and a vertex v of G, the *vertex switching* graph G_v is the graph obtained from G by removing all edges incident to v and adding edges joining v to every vertex not adjacent to v in G. Vaidya and Vyas [4029] proved that the graphs obtained by the switching of a pendent vertex of a path, a vertex of a cycle, a rim vertex of a wheel, the center vertex of a helm, or a vertex of degree 2 of a fan are antimagic graphs.

Phanalasy, Miller, Rylands, and Lieby [2835] in 2011 showed that there is a relationship between completely separating systems and labeling of regular graphs. Based on this relationship they proved that some regular graphs are antimagic. Phanalasy, Miller, Iliopoulos, Pissis, and Vaezpour [2833] proved the Cartesian product of regular graphs obtained from [2835] is antimagic. Ryan, Phanalasy, Miller, and Rylands introduced the generalized web and flower graphs in [3219] and proved that these families of graphs are antimagic. Rylands, Phanalasy, Ryan, and Miller extended the concept of generalized web graphs to the single apex multi-generalized web graphs and they proved these graphs to be antimagic in [3221]. Ryan, Phanalasy, Rylands and Miller extended the concept of generalized flower to the single apex multi-(complete) generalized flower graphs and constructed antimagic labeling for this family of graphs in [3220]. For more about antimagicness of generalized web and flower graphs see [2498]. Phanalasy, Ryan, Miller and Arumugam [2834] introduced the concept of generalized pyramid graphs and they constructed antimagic labeling for these graphs. Bača, Miller, Phanalasy, and Feňovčíková proved that some join graphs and incomplete join graphs are antimagic in [407]. Moreover, in [406] they proved that the complete bipartite graph $K_{m,m}$ and complete 3-partite graph $K_{m,m,m}$ are antimagic and if G is a k-regular (connected or disconnected) graph with p vertices and $k \geq 2$, then the join of G and $(p-k)K_1$ is antimagic. Deng [922] showed that every bipartite graph with minimum degree at least 15 is antimagic. In [1572] Jane and Myom modeled a dynamic encryption with implicit key exchange for end-to-end encryption using totally antimagic labeling of complete bipartite graphs over the network that makes it impossible to successfully intercept the message.

Arumugam, Miller, Phanalasy, and Ryan [282] provided antimagic labelings for a family of generalized pyramid graphs. Daykin, Iliopoulas. Miller, and Phanalasy [909] show several families of graphs recursively defined from a sequence of graphs that are generalizations of corona graphs are antimagic. Lozano, Mora, Seara, and Tey [2364] proved that caterpillars are antimagic. In [2094] Latchoumanane and Varadhan proved that the tensor product of wheels, helms, and flowers with star are antimagic.

In [2241] Li and Yang determined the exact number of paths P_3 such that the disjoint union of a double star with that number of paths P_3 is antimagic. In addition, they provide some examples of the antimagic labelings have vertex sums 1 through the number of vertices of the graphs. [2241] new

Let G be a k-regular graph with p vertices and q edges. The *generalized sausage graph*, denoted by $S(G;m)$, is the graph obtained from $G \times P_m$ ($G \times P_1 = G$), by joining each end vertex of the P_m to a new vertex (which we call *apexes*) with an edge. In particular, when $m = 1$, each vertex of the graph G joins to two vertices with two edges. The *mixed generalized sausage graph*, denoted by $MS(G;m)$, is the graph obtained from the generalized sausage graph $S(G;m)$, $m \geq 3$, by joining each vertex of each copy of the $\lceil m/2 \rceil$ copies of G on the left hand side to the left hand side apex, except the nearest copy to the apex, and similarly for the right hand side apex. The *complete mixed generalized sausage graph*, denoted by $CMS(G;m)$ is the graph obtained from the generalized sausage graph by joining each vertex of each copy of G, except the two nearest copies of G to the apexes, to each apex with an edge, and each corresponding pair of vertices of the two nearest copies of G to the apexes with an edge. The *complete mixed generalized sausage graph* $CMS^-(G;m)$ is the graph obtained from $CMS(G;m)$ by deleting the edge connecting each corresponding pair of vertices of the two nearest copies of G to the apexes. In [2832] Phanalasy proved families of generalized sausage graphs, mixed generalized sausage graphs, and complete mixed generalized sausage graphs are antimagic.

A *split* graph is a graph that has a vertex set that can be partitioned into a clique and an independent set. Tyshkevich (see [549]) defines a canonically decomposable graph as follows. For a split graph S with a given partition of its vertex set into an independent set A and a clique B (denoted by $S(A,B)$), and an arbitrary graph H the composition $S(A,B) \circ H$ is the graph obtained by taking the disjoint union of $S(A,B)$ and H and adding to it all edges having an endpoint in each of B and $V(H)$. If G contains nonempty induced subgraphs H and S and vertex subsets A and B such that $G = S(A,B) \circ H$, then G is *canonically decomposable*; otherwise G is *canonically indecomposable*. Barrus [549] proved that every connected graph on at least 3 vertices that is split or canonically decomposable is antimagic.

Hartsfield and Ringel [1378] conjecture that every tree except P_2 is antimagic and, moreover, every connected graph except P_2 is antimagic. In 2004 Alon, Kaplan, Lev, Roditty, and Yuster [201] use probabilistic methods and analytic number theory to show that this conjecture is true for all graphs with n vertices and minimum degree $\Omega(\log n)$. They also prove that if G is a graph with $n \geq 4$ vertices and $\Delta(G) \geq n - 2$, then G is antimagic and all complete partite graphs except K_2 are antimagic. Slíva [3638] proved the conjecture for graphs with a regular dominating subgraph. In 2016 Eccles [996] improved the result of Alon et al. by proving that there exists an absolute constant d_0 such that if G is a graph with average degree at least d_0 and G contains no isolated edge and at most one isolated vertex, then G is antimagic. In [708] Branson, Sutton, and Yan demonstrated how a simple evolutionary algorithm can be used to confirm the antimagic tree conjecture for all trees up to order 25. Muthukkumar and Rajendran [2612] proved that splitting graph of an antimagic graph is antimagic and that for any antimagic tree T, the graphs obtained by connecting a vertex of maximum degree of two copies of T by an edge is antimagic. In 2026, Beaudoire, Bentz, and Picouleau [605] extended the result of Alon, Kaplan, Lev, Roditty, and Yuster to graphs with maximum degree at least $n-4$ provided [605] new

that the number of edges is at least $7n$.

Chawathe and Krishna [792] proved that every complete m-ary tree is antimagic. Yilma [4268] extended results on antimagic graphs that contain vertices of large degree by proving that a connected graph with $\Delta(G) \geq |V(G)| - 3$, $|V(G)| \geq 9$ is antimagic and that if G is a graph with $\Delta(G) = \deg(u) = |V(G)| - k$, where $k \leq |V(G)|/3$ and there exists a vertex v in G such that the union of neighborhoods of the vertices u and v forms the whole vertex set $V(G)$, then G is antimagic. Sethuraman and Shermily [3454] proved that binomial trees and Fibonacci trees are antimagic. Vasuki, Shobana, and Ahmed [4071] proved the existence of face antimagic labelings for double duplication of barycentric subdivisions of cycles and some other graphs. Vasuki ·and Shobana [4072] proved the existence of face antimagic labeling of types (0, 1, 0) and (0, 1, 1) for crown related graphs.

For connected graphs $G, H_1, H_2, \ldots, H_n$ with G having the edges $e_1, e_2, \ldots, e_n$ Nivedha and Yamini [2692] define the *generalized edge corona* of G and $H_1, H_2, \ldots, H_n$ as the graph obtained by adding all possible edges between the end vertices of $e_1, e_2, \ldots e_n$ and the vertices of $H_1, H_2, \ldots, H_n$. They prove that such graphs are antimagic under certain conditions where G has only one vertex of maximum degree three and each H_i has at least 2 vertices.

In [143] Ahmed and Babujee defined a *strong face plane graph* as one that is obtained from a plane graph by adding a new vertex to every face, in such way that the faces of the resulting graph are three-sided. If the faces of original plane graphs are three sided faces, then the number of faces increase. They investigated the existence of (super) d-antimagic labeling of type (1, 1, 1) for some strong face plane graphs. Shawkat and Ahmed [3494] investigated the existence of vertex antimagic edge labelings for strong face ladders, strong face wheels, strong face fans, strong face prisms $C_m \times P_2$, and strong face friendship graphs. Kuppan and Shobana [2068] proved the existence of face antimagic labelings for the double duplication of all vertices by the edges of gear graphs G_n for $n \geq 3$, $P_3 \times P_{2n}$ $(n \geq 2)$, $C_n \times P_2$ $(n \geq 5)$, and the double duplication of all vertices by the edges of a strong face of the triangular snake graph T_n $(n \geq 3)$. They noted that an (a, d)-face antimagic labeling for double duplication of special graphs can be used to encrypt and decrypt the messages in real time. Jesintha, Vinodhini, Lakshmi [1643] proved that triangular books and double fans graph admit antimagic labelings.

In 2013 Kamachi and Arumugam [1816] introduced the concept of distance antimagic graphs as follows. Let $G = (V, E)$ be a graph of order n. Let $f : V \to \{1, 2, \ldots, n\}$ be a bijection. The weight $w(v)$ of a vertex v with respect to the labeling f is defined by $w(v) = \sum_{u \in N(v)} f(u)$, where $N(v)$ is the open neighborhood of v. The labeling f is called a *distance antimagic labeling* if $w(v_1) \neq w(v_2)$ for any two distinct vertices v_1, v_2 in V. They proved that P_n, C_n $(n \neq 4), W_n$ $(n \neq 4)$, $rK_2 + K_1$, and $G \odot K_1$ for all graphs G admit distance antimagic labelings. Cutinho, Sudha, and Arumugam [875] proved that $K_n \times K_n$ is distance antimagic if and only if $n \neq 2$ and $K_3 \times C_n$ is distance antimagic when $n \geq 3$ is odd. They included the case $K_3 \times C_n$ when n is even as a problem. Handa, Godinho, Singh, and Arumugam [2649] gave a tool for proving the existence of distance antimagic labelings for the join and the corona of two graphs. Simanjuntak, Nadeak, [1816] new [2649] new

Yasin, Wijaya, Hinding, and Sugeng [3586] proved that $P_m \cup P_n$ $(m, n \geq 3)$, copies of P_n $(n \neq 3)$, copies of wheels W_n $(n \neq 3)$, copies of fans F_n $(n \geq 0$, copies of friendship graphs f_n $(n \geq 3$, and copies of stars are distance antimagic graphs. In [1792] John, Lubis, and Sugeng showed that for any regular and distance antimagic graph copies of the graph are also distance antimagic graph. Moreover, showed that for Herschel graph and the copies of Herschel graph also distance antimagic. [3586] new [1792] new

For a commutative ring R with unity, the zero-divisor graph of R is the set of all zero-divisors of R, where zero-divisors u and v are adjacent if $uv = 0$. In [3624] Sivakumaran, Sankar, and Prabhu proved various families of zero-divisor graphs admit distance antimagic labeling.

Fronček [1127] defines a *handicap incomplete tournament* of n teams with r rounds, HIT(n, r), as a tournament in which every team plays r other teams and the total strength of the opponents that team i plays is $\vec{S}_{n,r}(i) = t - i$ for every i and some fixed constant t. (This means that the strongest team plays strongest opponents, and the lowest ranked team plays weakest opponents.) In terms of distance magic graphs this restriction corresponds to finding a distance antimagic graph with the additional property that the sequence $w(1), w(2), \ldots, w(n)$ (where team i is again the i-th ranked team) is an increasing arithmetic progression with difference one. These graphs are called *handicap distance antimagic graphs*. A *handicap distance d-antimagic labeling* of a graph $G(V, E)$ with n vertices is a bijection $\vec{f} : V \to \{1, 2, \ldots, n\}$ with the property that $\vec{f}(x_i) = i$ and the sequence of the weights $w(x_1), w(x_2), \ldots, w(x_n)$ forms an increasing arithmetic progression with difference d. A graph G is a *handicap distance d-antimagic graph* if it admits a handicap distance d-antimagic labeling, and *handicap distance antimagic graph* when $d = 1$. In [1127] Fronček establishes a relationship between handicap incomplete tournaments and distance antimagic graphs and construct some new infinite classes of distance antimagic graphs and infinite classes of handicap incomplete round robin tournaments. Fronček and Shepanik [1145] construct r-regular handicap distance antimagic graphs of order $n \equiv 0 \pmod 8$ for all feasible values of r. Fronček [1131] proved that regular handicap distance antimagic graphs exist for every feasible odd order by proving that there exists a regular handicap graph of an odd order n if and only if $n = 9$ or $n \geq 13$. In [1130] Fronček constructed a class of regular 2-handicap distance antimagic graphs for every order $n \equiv 0 \pmod{16}$. In [1132] he proved that a k-regular 2-handicap distance antimagic graph of order $n \equiv 0 \pmod{16}$ exists if and only if $n \geq 16$ and $4 \leq k \leq n - 6$. Fronček and Shepanik [1146] constructed k-regular handicap distance antimagic graphs of order $n = 4 \pmod 8$ for all feasible values of k. In [3568] Shrimali and Parmar discuss the existence distance antimagic labelings for the product, direct product, strong product, and carona product of graphs involving ${C_3}^t$ and C_4.

Cranston [863] proved that for $k \geq 2$, every k-regular bipartite graph is antimagic. For non-bipartite regular graphs, Liang and Zhu [2247] proved that every cubic graph is antimagic. That result was generalized by Cranston, Liang, and Zhu [864], who proved that odd degree regular graphs are antimagic. Hartsfield and Ringel [1378] proved that every 2-regular graph is antimagic. Bérczi, Bernäth, and Vizer [622] use a slight modification of an argument of Cranston et al. [864] to prove that k-regular graphs are antimagic

for $k \geq 2$. The same was done by Chang, Liang, Pan, and Zhu [771] proved that every even degree regular graph is antimagic.

Tai, Chia, and Ong [3856] proved that the graphs obtained by starting with two vertices and joining them with at least $r \geq 3$ edges then subdividing the edges of this graph arbitrarily so that at most one edge (*multi-bridge graphs*) is not subdivided are antimagic. For a connected, undirected, simple graph $G(V, E)$ a bijection f from V to $\{1, 2, \ldots, |V|\}$ is called a *rainbow antimagic vertex labeling* if, for any two edges uv and $u'v'$ in the same path, $w(uv) \neq w(u'v')$, where $w(uv) = f(u) + f(v)$. The *rainbow antimagic connection number* of G is the smallest number of colors taken over all rainbow colorings induced by a rainbow antimagic labeling of G. In [3413] Septory, Utoyo, Dafik, Sulistiyono, and Agustin determined the exact value of the rainbow antimagic connection numbers of Jahangir graphs, firecrackers, $K_{2,m}$, and double stars. A rainbow antimagic labeling f of a graph G is called a *strong rainbow antimagic labeling* of G if for every two vertices u and v there exists path from u to v in which no two edges of the path have the same weight. The *strong rainbow antimagic connection number* of G is the smallest number of colors taken over all strong rainbow colorings induced by strong rainbow antimagic labelings of G. In [2275] Lestari1, Dafik, Susanto, and Wahab determined the connection number of strong rainbow antimagic coloring for Jahangir graphs, Jahangir semi graphs, friendship graphs, fans, stars, and paths.

A graph $G(V, E)$ is *k-shifted antimagic* if there exists a bijection f from E to $\{k + 1, \ldots, k + |E|\}$ such that the vertex sums of all vertices are distinct; $G(V, E)$ is *absolutely antimagic* if it is k-shifted antimagic for every integer k. In [770] Chang, Li, Daphne Liu, and Pan proved that certain spider forests (a graph where each component is a spider) are k-shifted antimagic for all $k \geq 0$. In addition, they showed that for a spider forest G with m edges, there exists a positive integer $k_0 \leq m$ such that G is k-shifted antimagic for all $k \geq k_0$ and $k \leq -(m + k_0 + 11)$. Li and Wang [2240] proved that every tree of diameter four or five, except for two cases, is k-shifted antimagic for every integer k. Chang, Chen, Li, and Pan [768] proved: forest without a component isomorphic to K_2 are k-shifted-antimagic, graphs consisting of vertices of odd degrees and containing no component isomorphic to K_2 are k-shifted-antimagic for sufficiently large k, and P_n $(n \geq 6)$ are k-shifted for all k. They also gave necessary and sufficient conditions for stars, double stars, kP_3, and $2K_{1,3}$ to be k-shifted-antimagic. Dhananjaya and Li [946] proved that P_2, P_3, P_4-free linear forests and $K_{1,2}$-free forests are k-shifted-antimagic with a few exceptions. This extended the results by Shang et al. in [3483] and [3484]. Moreover, they prove that the odd tree forests are k-shifted-antimagic for all k. Chen and Kuo [803] proved that every caterpillar of diameter at least five is absolutely antimagic. [803] new

Chang, Chen, Li, and Pan [768] established connections among various concepts proposed in the literature of antimagic labelings and extend previous results in three ways: some classes of graphs, including trees and graphs whose vertices are of odd degrees, that have not been verified to be antimagic are shown to be k-shifted-antimagic for sufficiently large k; some graphs are proved k-shifted-antimagic for all k, whereas some are proved not for some particular k; and disconnected graphs are also considered.

Beck and Jackanich [608] showed that every connected bipartite graph except P_2 with

$|E|$ edges admits an edge labeling with labels from $\{1, 2, \ldots, |E|\}$, with repetition allowed, such that the sums of the labels of the edges incident to each vertex are distinct. They call such a graph *weak antimagic*.

Wang, Liu, and Li [4154] proved: mP_3 ($m \geq 2$) is not antimagic; $P_n \cup P_n$ ($n \geq 4$) is antimagic; $S_n \cup P_n$ is antimagic; $S_n \cup P_{n+1}$ is antimagic; $C_n \cup S_m$ is antimagic for $m \geq 2\sqrt{n} + 2$; mS_n is antimagic; if G and H are graphs of the same order and $G \cup H$ is antimagic, then so is $G + H$; and if G and H are r-regular graphs of even order, then $G+H$ is antimagic. In [4155] Wang, Liu, and Li proved that if G is an n-vertex graph with minimum degree at least r and H is an m-vertex graph with maximum degree at most $2r-1$ ($m \geq n$), then $G+H$ is antimagic. Bača, Kimáková, Semaničová-Feňovčiková, and Umar [382] prove the disjoint union of multiple copies of a $(a, 1)$-(super)-tree-antimagic graph is also a $(b, 1)$-(super)-tree-antimagic for certain a and b.

For any given degree sequence pertaining to a tree, Miller, Phanalasy, Ryan, and Rylands [2500] gave a construction for two vertex antimagic edge trees with the given degree sequence and provided a construction to obtain an antimagic unicyclic graph with a given degree sequence pertaining to a unicyclic graph.

Kaplan, Lev, and Roditty [1905] prove that every non-trivial rooted tree for which every vertex that is a not a leaf has at least two children is antimagic (see [2245]) for a correction of a minor error in the proof). For a graph G with m vertices and an Abelian group A they define G to be A-*antimagic* if there is a one-to-one mapping from the edges of G to the nonzero elements of A such that the sums of the labels of the edges incident to v, taken over all vertices v of G, are distinct. For any $n \geq 2$ they show that a non-trivial rooted tree with n vertices for which every vertex that is a not a leaf has at least two children is Z_n-antimagic if and only if n is odd. They also show that these same trees are A-antimagic for elementary Abelian groups G with prime exponent congruent to 1 (mod 3). Lozano, Mora, Seara, and Tey [2368] proved that trees whose vertices of even degree induce a path are antimagic, thereby extending a 2014 result of Liang, Wong, and Zhu in [2245]. [2368] new

In [765] Chan, Low, and Shiu use $[G, A]$ to denote the class of distinct A-antimagic labelings of G. They prove: for a non-trivial Abelian group A that underlies some commutative ring R with unity, if d is a unit in R and $f \in [G, A]$, then $df \in [G, A]$; if A is an Abelian group that contains a subgroup isomorphic to B and a graph G is B-antimagic, then G is A-antimagic; P_{4m+r} and C_{4m+r} are Z_k-antimagic for $k \geq 4m+r$ and $r = 0, 1, 3$; P_{4m+2} is Z_k-antimagic for $k \geq 4m + 3$; regular Hamiltonian graphs of order $4m + r$ are Z_k-antimagic for $k \geq 4m + r$ and $r = 0, 1, 3$, and Z_k-antimagic for $k \geq 4m + 3$ and $r = 2$; for odd n, S_n is Z_k-antimagic for $k \geq n > 4$; for even n, S_n is Z_k-antimagic for $k \geq n + 2 \geq 6$ but not Z_n-antimagic or Z_{n+1}-antimagic; trees of order n with exactly one vertex of even degree are Z_k-antimagic for $k \geq n$; trees of order n with exactly two vertices of even degree are Z_k-antimagic for $k \geq n + 1$; and double stars of order are Z_k-antimagic for $k \geq n + 1$ when $n \equiv 2 \pmod 4$ and Z_k-antimagic for $k \geq n$ when $n \not\equiv 2 \pmod 4$.

Chang and Liu [774] call a graph G $\mathbb{R}$-*antimagic* if for each subset A of $\mathbb{R}$ with $|A| = |E(G)|$, there is an edge labeling such that the sums of the labels assigned to

edges incident to distinct vertices are different. They proved that wheels, cycles, and complete graphs of order at least 3 are $\mathbb{R}$-antimagic, and that Cartesian products with at least two terms are $\mathbb{R}$-antimagic when each term is a complete graph of order at least 2 or a cycle.

The *integer-antimagic spectrum* of a graph G is the set $\mathrm{IAM}(G) = \{k \geq 2 : G \text{ is } Z_k\text{-antimagic}\}$. In [2733] Odabas, Roberts, and Low proved that $\mathrm{IAM}(G) \subseteq \mathrm{IAM}(G')$, a where G' is any graph obtained by adding simple edges to G (not equal to a 3-path when $k = 3$) and that if G is disconnected and the added edges do not create a new K_3-component in G', then $\mathrm{IAM}(G) \subseteq \mathrm{IAM}(G')$. Shiu, Sun, and Low [3556] determine the integer-antimagic spectra of tadpoles and lollipops. Shiu and Low [3551] determine the integer-antimagic spectra of complete bipartite graphs and complete bipartite graphs with a deleted edge. Shiu [3521] determined the integer-antimagic spectra of disjoint unions of cycles. In [2730] Odabasi, Roberts, and Low determined the integer-antimagic spectra for all Hamiltonian graphs. Roberts and Low [3186] analyzed the integer-antimagic spectra for various classes of multi-cyclic graphs. In [2731] and [2732] Odabasi, Roberts, and Low determined the integer-antimagic spectra for a disjoint union of Hamiltonian graphs and characterized the integer-antimagic spectra for the weak join of Hamiltonian graphs. Low, Roberts, and Zheng [2361] determined the integer-antimagic spectra for cycles with a chord, paths with a chord, and wheels with a chord. [2733] new [3186] new [2731] new [2732] new [2361] new

Liang, Wong, and Zhu [2245] study trees with many degree 2 vertices with a restriction on the subgraph induced by degree 2 vertices and its complement. Denoting the set of degree 2 vertices of a tree T by $V_2(T)$ Liang, Wong, and Zhu proved that if $V_2(T)$ and $V \setminus V_2(T)$ are both independent sets, or $V_2(T)$ induces a path and every other vertex has an odd degree, then T is antimagic. In [2367] Lozano, Mora, Seara, and Tey extended this result by showing that trees whose vertices of even degree induce a path are antimagic.

In [4033] Vaidya and Vyas proved that the middle graphs, total graphs, and shadow graphs of paths and cycles are antimagic. In [2032] and [2033] Krishnaa provided some results for antimagic labelings for graphs derived from wheels and antimagic labelings of helm related graphs.

Bertault, Miller, Pé-Rosés, Feria-Puron, and Vaezpour [634] approached labeling problems as combinatorial optimization problems. They developed a general algorithm to determine whether a graph has a magic labeling, antimagic labeling, or an (a, d)-antimagic labeling (see Section 6.3). They verified that all trees with fewer than 10 vertices are super edge magic and all graphs of the form $P_2^r \times P_3^s$ with less than 50 vertices are antimagic. Kuppan, Shobana, and Cangul [2067] used an (a, d)-face antimagic labeling of a strong face of the duplication of all vertices by edges of a tree T_n $(n \geq 2)$ to encrypt and decrypt thirteen secret numbers that can be extended to the double duplication of graphs to encode and decode the numbers that can be used in applications.

In [398] Bača, MacDougall, Miller, Slamin, and Wallis survey results on antimagic, edge-magic total, and vertex-magic total labelings.

A *total labeling* of a graph G is a bijection f from $V(G) \cup E(G)$ to $\{1, 2, \ldots, |V(G)| + |E(G)|\}$. When $f(V(G)) = \{1, 2, \ldots, |V(G)|\}$, we say the total labeling is *super*. For a labeling f the associated edge-weight of an edge uv is defined by $wtf(uv) = f(uv)+f(u)+$

$f(v)$. The associated vertex-weight of a vertex v is defined by $wt_f(v) = \sum_{u \in N(v)} f(uv) + f(v)$, where $N(v)$ is the set of the neighbors of v. A labeling f is called *edge-antimagic total* (*vertex-antimagic total*) if all edge-weights (vertex-weights) are pairwise distinct. A graph that admits an edge-antimagic total (vertex-antimagic total) labeling is called an *edge-antimagic total* (*vertex-antimagic total*) graph. A labeling that is simultaneously edge-antimagic total and vertex-antimagic total is called a *totally antimagic total labeling*. A graph that admits a totally antimagic total labeling is called a *totally antimagic total* graph. Ajayi [149] proved that ladders, prisms, and generalised Petersen graphs are totally antimagic total graphs. He showed that the chain graph of totally antimagic total graphs is a totally antimagic total graph. Using one of his previous results the labeling of Bhavale [655] Bhavala [654] proved that K_n $(n \geq 3)$ is super antimagic (that is, the vertex labels are 1 through the number of vertices) and totally antimagic total. He also proved that there exists an antimagic orientation of $K_n (n \geq 3)$. Akwu and Ajayi [168] proved that ladders, prisms and generalised Pertersen graphs are totally antimagic total graphs. They further showed that the chain graph of totally antimagic total graphs is a totally antimagic total graphs. [149] new [655] new [654] new [168] new

In [3785] Sundar, Jeyabalan, Nishanthini, Swathi, and Rajchakit proved that for any graph G and $m > 1$, $G \odot mK_1$ and $G \odot mK_1$ do not admit both a vertex-magic total labeling and edge-antimagic total labeling. A labeling g is said to be *ordered* (*sharp ordered*) if $wt_g(u) \leq wt_g(v)$ $(wt_g(u) < wt_g(v))$ holds for every pair of vertices $u, v \in V(G)$ such that $g(u) < g(v)$. A graph that admits a (sharp) ordered labeling is called a (*sharp*) *ordered graph*.

A *strong edge-antimagic total labeling* of a simple graph $G(V, E)$ is a labeling in which the vertex labels are consecutive integers from 1 to $|V|$ such that the total of the labels of the vertices incident to an edge form an ascending arithmetic sequence. Prasetyo [3035] proved that $mK_{1,n}$ has a strong edge-antimagic total labeling.

In [3835] Suthakaran and Jeyabalan introduced the following notion. A *parity-weighted totally antimagic total labeling*s a totally antimagic labeling in which all vertex and edge weights are either even or odd. If all vertex weights are even (or odd) and all edge weights are odd (or even), they say that the labeling is a strong or weak (WPAT) parity-weighted totally antimagic total labeling, respectively. A *strong parity weighted totally antimagic total labeling* (SPAT) is a bijection from $V \cup E$ to $\{1, 2, \ldots, |V(G)| + |E(G)|\}$ with the condition that the vertex and edge weights are pairwise distinct and the vertex weights are required to be even, while the edge weights must be odd. They provide positive and negative results regarding the existence SPAT and WPAT labeling for friendship graphs, generalized friendship graphs, fans, and wheels.

Miller, Phanalasy, and Ryan [2497] proved that all graphs have vertex-antimagic total labelings. Bača, Miller, Phanalasy, Ryan, Semaničová-Feňovčíková, and Abildgaard Sillasen [404] prove that mK_1, mK_2, P_n $(n \geq 2)$, and C_n are sharp ordered super totally antimagic total. They prove if G is an ordered super edge-antimagic total graph then $G + K_1$ is a totally antimagic total graph. As a corollary they get that stars, friendship graphs $nK_2 + K_1$, fans, and wheels are totally antimagic total. They also prove that if G is a regular ordered super edge-antimagic total graph then $G \odot nK_1$ is totally antimagic total.

As a corollary of this result, they have double-stars $K_2 \odot nK_1$ and crowns $C_m \odot nK_1$ are totally antimagic total. They show that a union of regular totally antimagic total graphs is a totally antimagic total graph. Miller, Phanalasy, Ryan, and Rylands [2499] provide a method whereby, given any degree sequence pertaining to a tree, one can construct an antimagic tree based on this sequence. By swapping the roles of edges and vertices with respect to a labeling, they provide a method to construct an edge antimagic vertex labeling for any tree. In [3352] Selvarasu and Murugan showed that twigs and combs admit vertex and edge antimagic labelings. Liunokas, Putra, Pangaribuan, and Pasangka [2302] proved that several comb related graphs admit antimagic labelings. [2302] new

Ahmed and Baskar Baskar [141] proved that complete bipartite graphs admit a totally antimagic total labeling. The same result was proved by Akwu and Ajayi [167] who also showed that the join of a complete bipartite graph and K_1 is a totally antimagic total graph. Ahmed, Baskar Babujee, Bača, Semaničová-Feňovčíková [140] proved that complete graphs admit totally antimagic total labeling. They also considered the problem of finding total labelings for prisms and for two special classes of graphs related to paths that are simultaneously edge-magic and vertex-antimagic. Ahmad, Semaničová-Feňovčíková, Siddiqui, and Kamran [136] construct α-labelings from graceful labelings of smaller trees and transform this labeling to edge-antimagic vertex labeling of trees. Shang [3484] shows that linear forests without either of the paths P_2 or P_3 as components are antimagic. Shang [3485] proved that P_2, P_3, and P_4-free linear forests are antimagic. In [3554] Shiu and Low analyzed various antimagic properties for Cartesian products, hexagonal nets and theta graphs. Ajayi and Akwu [151] proved that ladders, prisms, and generalized Petersen graphs are totally antimagic total graphs. They also showed that the chain graph of totally antimagic total graphs is a totally antimagic total graph.

In [3849] Swathi, Kumar, Jeyabalan, and Nishanthini introduced the notion of a *perfectly antimagic total labeling* as a totally antimagic total labeling whose vertex and edge weights are also pairwise distinct. A graph with such a labeling is called a *perfectly antimagic total* graph. In [3850] Swathi and Jeyabalan investigated the possibility of the perfectly antimagic total labeling of star-like trees and gave a perfectly antimagic total labeling for the corona product of a path with nK_1 and a cycle with nK_1. They further provided a particular solution to the open problem about total antimagic labelings of caterpillars posed by Lozano et al. in [2365] and [2364].

A *Fibonacci mean antimagic* labeling of a graph G is an injective function $g : V(G) \to \{f_2, f_3, \ldots, f_{n+1}\}$, where f_n is nth Fibonacci number, and the induced a function $g^* : E(G) \to N$ defined by $g^*(uv) = \lceil (g(u) + g(v))/2 \rceil$ is injective. A graph is called *Fibonacci mean antimagic* if it admits a Fibonacci mean antimagic labeling. Thirugnanasambandam and Chitra [3890] proved the following graphs have Fibonacci mean antimagic labelings: caterpillars, bistars, triangular snakes, quadrilateral snakes, ladders, and the corona product of last three graphs and K_1. Barasara and Prajapati [497] studied antimagic labelings of double triangular snakes, alternate triangular snakes, double alternate triangular snakes, quadrilateral snakes, double quadrilateral snakes, alternate quadrilateral snakes, double alternate quadrilateral snakes.

In [1397] Hefetz, Mütze, and Schwartz investigate antimagic labelings of directed

graphs. An *antimagic* labeling of a directed graph D with n vertices and m arcs is a bijection from the set of arcs of D to the integers $\{1, \ldots, m\}$ such that all n oriented vertex sums are pairwise distinct, where an oriented vertex sum is the sum of labels of all edges entering that vertex minus the sum of labels of all edges leaving it. Hefetz et al. raise the questions "Is every orientation of any simple connected undirected graph antimagic? and "Given any undirected graph G, does there exist an orientation of G which is antimagic?" They call such an orientation an *antimagic orientation* of G. Regarding the first question, they state that, except for $K_{1,2}$ and K_3, they know of no other counterexamples. They prove that there exists an absolute constant C such that for every undirected graph on n vertices with minimum degree at least $C \log n$ every orientation is antimagic. They also show that every orientation of $S_n,\ n \neq 2$, is antimagic; every orientation of W_n is antimagic; and every orientation of $K_n,\ n \neq 3$, is antimagic. For the second question they prove: for odd r, every undirected r-regular graph has an antimagic orientation; for even r every connected undirected r-regular graph that admits a matching that covers all but at most one vertex has an antimagic orientation; and if G is a graph with $2n$ vertices that admits a perfect matching and has an independent set of size n such that every vertex in the independent set has degree at least 3, then G has an antimagic orientation. They conjecture that every connected undirected graph admits an antimagic orientation and ask if it true that every connected directed graph with at least 4 vertices is antimagic. Shan [3482] supported this conjecture by proving that every bipartite graph with no vertex of degree 0 or 2 admits an antimagic orientation and every graph with minimum degree at least 33 admits an antimagic orientation. In [4244] Yang, Carlson, Owens, Perry, Singgih, Song, Zhang, and Zhang proved that a graph G admits an antimagic orientation if $\Delta(G) \geq |G| - 3$ or $\Delta(G) = |G| - t \geq 4$ for each $t = 4, 5$.

A simple graph G is said to admit an *antimagic orientation* if there exist both an orientation on the edges of G and a bijection from $E(G)$ to $\{1, 2, \ldots, |E(G)|\}$ such that the vertex sums of vertices are pairwise distinct, where the vertex sum of a vertex is defined to be the sum of the labels of the in-edges minus that of the out-edges incident to the vertex. It was conjectured by Hefetz, Mütze, and Schwartz in [1397] in 2010 that every connected simple graph admits an antimagic orientation. In [947] Dhananjaya and Li proved that several operations on graphs such as the corona product, the lexicographic product, and the Mycielski construction, together with some conditions on the graphs yield graphs satisfying the above conjecture.

For a directed graph $D = (V, A)$ with p vertices and q arcs. Kumar, Nirmala, and Mutharasu [2062] say that D has a *V-super vertex in-antimagic total graceful labeling* (V-SVIAMTG labeling) if there is a bijection $f : V(D) \cup A(D)$ to $\{1, 2, \ldots, p + q\}$ with the property that $f(V(D)) = \{1, 2, \ldots, p\}$ and for each $v \in V(D)$, $|w(v) - f(v)|$ consists of distinct integers, where $w(v)$ is $\sum(f(u, v)$ over all u incident to v. They call D a *V-super vertex in-antimagic total graceful digraph* (V-SVIAMTG digraph) if D admits a V-SVIAMTG labeling. They prove existence of V-SVIAMTG for directed paths, cycles, coronas, fans, some cylinders, and some grids.

Motivated by the Hartsfield and Ringel on antimagic labelings of graphs, in 2010 Hefetz, Mütze, and Schwartz [1397] initiated the study of antimagic orientations of graphs,

and conjectured that every connected graph admits an antimagic orientation. The conjecture has been verified to be true for regular graphs (see [[1397], [2238], [4242]]), and biregular bipartite graphs with minimum degree at least two by Shan and Yu [3481]. Yang, Carlson, Owens, Perry, Singgih, Song, Zhang, Zhang [4244] proved that every connected graph G on $n \geq 9$ vertices with maximum degree at least $n-5$ admits an antimagic orientation. Li, Song, Wang, Yang, and Zhang [2238] proved that every 2-regular graph has an antimagic orientation and for all integers $d \geq 2$, every connected $2d$-regular graph has an antimagic orientation. Gao, Yuping and Shan [1188] proved that every lobster admits an antimagic orientation. Ferraro, Newkirk, and Shan [1083] prove that subdivided caterpillars in which all pendant edges are replaced by paths of the same length admit an antimagic orientation.

Sonntag [3676] investigated antimagic labelings of hypergraphs. He shows that certain classes of cacti, cycle, and wheel hypergraphs have antimagic labelings. Javaid and Bhatti [1580] extended some of Sonntag's results to disjoint unions of hypergraphs. In [2645] Nalliah investigated the existence of antimagic labelings of some families of digraphs using hooked Skolem sequences. Marimuthu, Raja Durga, and Durga Devi [2434] investigated the existence of super vertex in-antimagic total labelings of generalized de Bruijn digraphs. In [988] Dsouza, Saumya, and Kumudakshi investigate the existence of antimagic labelings for directed wheels and directed versions of $P_n \times K_2$.

Hefetz [1396] calls a graph with q edges *k-antimagic* if its edges can be labeled with $1, 2, \ldots, q+k$ such that the sums of the labels of the edges incident to each vertex are distinct. In particular, antimagic is the same as 0-antimagic. More generally, given a weight function ω from the vertices to the natural numbers Hefetz calls a graph with q edges *(ω, k)-antimagic* if its edges can be labeled with $1, 2, \ldots, q+k$ such that the sums of the labels of the edges incident to each vertex and the weight assigned to each vertex by ω are distinct. In particular, antimagic is the same as $(\omega, 0)$-antimagic where ω is the zero function. Using Alon's combinatorial nullstellensatz [200] as his main tool, Hefetz has proved the following: a graph with 3^m vertices and a K_3 factor is antimagic; a graph with q edges and at most one isolated vertex and no isolated edges is $(\omega, 2q-4)$-antimagic; a graph with $p > 2$ vertices that admits a 1-factor is $(p-2)$-antimagic; a graph with p vertices and maximum degree $n-k$, where $k \geq 3$ is any function of p is $(3k-7)$-antimagic and, in the case that $p \geq 6k^2$, is $(k-1)$-antimagic. Hefetz, Saluz, and Tran [1398] improved the first of Hefetz's results by showing that a graph with p^m vertices, where p is an odd prime and m is positive, and a C_p factor is antimagic.

A *vertex in-out-antimagic total labeling* of a directed graph (digraph) $D(V, A)$ with n vertices and m) arcs is a bijection from $V \cup A$ to $\{1, 2, \ldots, m+n\}$ such that all n vertex in-weights are pairwise distinct, and simultaneously, all n vertex out-weights are pairwise distinct. If the smallest labels are used on vertices, the labeling is called a *super vertex in-out-antimagic total* labeling. In [383] Bača, Kovář, Kovář, and Semaničová-Feňovčíková provided a general way for determining if dense digraphs and certain sparse digraphs admit a super vertex in-out-antimagic total labelings. They also include constructions of vertex in-out-antimagic total labelings for three large infinite classes of digraphs and conjecture that all digraphs allow such a labeling.

A graph $G = (V, E)$ is *strongly antimagic* if there is a bijective mapping $f : E \to 1, 2, \ldots, |E|$ such that for any two vertices $u \neq v$, not only $\sum_{e \in E(u)} f(e) \neq \sum_{e \in E(v)} f(e)$ and also $\sum_{e \in E(u)} f(e) < \sum_{e \in E(v)} f(e)$ whenever $\deg(u) < \deg(v)$, where $E(u)$ is the set of edges incident to u. Chang, Chin, Li, and Pan [769] proved double spiders (the trees contains exactly two vertices of degree at least 3) are strongly antimagic. They raise the following two questions. Does there exist a strongly antimagic labelings for every antimagic graph? Is there a k-antimagic graph but not $(k+1)$-antimagic?

For a graph $G(V, E)$ of order p and size q having no isolated vertices, a bijection $f : V \to \{1, 2, 3, \ldots, p\}$ is called a *local edge antimagic* labeling if for any two adjacent edges uv and vw of G, we have $w(uv) \neq w(vw)$, where the edge weight $w(uv) = f(u)+f(v)$ and $w(vw) = f(v) + f(w)$. A graph G is called a *local edge antimagic* if G has a local edge antimagic labeling. The local edge antimagic chromatic number number of G is the minimum number of colors taken over all colorings induced by local edge antimagic labelings of G. Agustin, Hasan, Dafik, Alfarisi and Prihandini [90] found the local edge antimagic chromatic numbers of paths, cycles, friendship graphs, ladders, stars, wheels, complete graphs, prisms, $C_m \odot mK_1$, and $G \odot mK_1$, where G is any graph of size at least 3. Moreover, they found a relation between local edge antimagic chromatic number and local antimagic vertex chromatic number. In [2617] Muthumanickavel and Nalliah proposed a comprehensive solution to the problem of scheduling work shifts for industrial workers that utilizes local antimagic vertex colorings. Nalliah [3101] determined the local edge antimagic chromatic numbers for a friendship graphs, wheels, fans, helms, flower graphs, and closed helms. In [1393] Hadiputra and Maryati investigated characterizations of graphs with small local edge chromatic numbers, the relationship between local edge antimagic chromatic number and edge independence number, and bounds for the local edge chromatic number for any graph. In [173] Alam, Semaničová-Feňovčíková, and Popa introduced the concept of a *graceful local antimagic* as an injection f from the vertex set of a graph G to $\{0, 1, \ldots, |E(G)|\}$ such that the induced edge labeling f^* defined as $f^*(uv) = |f(u) - f(v)|$ is a local antimagic labeling. They investigated the existence of local antimagic labeling for various families of graphs, utilizing Python-based algorithms as computational tools. In [3079] Rahmadani, Putri, Wasono, and Sandariria determined the chromatic number of local edge antimagic colorings of gears and a family of fan-related graphs. In [2450] Maryati and Hadiputra investigated the local edge antimagic coloring of join product of graphs for independent sets, paths, and cycles. [3079] new [2450] new

Sandhiya and Nalliah [3467] determined the local total antimagic chromatic number for the disjoint union of stars. In [2119] Lau, Shiu, Zhang, Premalatha, and Nalliah a gave a complete characterization of s-bridge graphs with local antimagic chromatic number 2. Shankar and Nalliah [3466] studied the local vertex antimagic chromatic number of some wheel related graphs. In [3464] they provided the local antimagic chromatic number for the corona product of wheels and null graphs. Lau, Premalatha, Shiu, and Nalliah [2106] improved results given in [2104] regarding the local antimagic chromatic number of $G \vee H$ for some circulant graph G where H is a null graph or a cycle. They also obtained the local antimagic chromatic numbers of certain wheel-related graphs. Lau, Shiu, Premalatha, and Nalliah [2111] provided three ways to construct a $(3m + 2) \times (2k + 1)$ matrix that [3467] new [2119] new [3466] new [3464] new [2106] new [2111] new

meets certain properties for $m = 1, 3$ and $k \geq 1$. Consequently, they obtained many (disconnected) graphs of size $(3m+2)(2k+1)$ with local antimagic chromatic number 3.

Let $H_1, H_2, \ldots, H_n$ be graphs on $p_1, p_2, \ldots, p_n$ vertices, respectively, where $p_1 \leq p_2 \leq \cdots \leq p_n$. Then the graph $F_n(H_1, H_2, \ldots, H_n)$ consisting of graphs H_i, $1 \leq i \leq n$, with a common vertex c is called a generalized friendship graph on $H_1, H_2, \ldots, H_n$, respectively. If $H_1 \cong H_2 \cong \cdots \cong H_n$ and $p_1 = p_2 = \cdots = p_n = p$, then the resulting graph is called the *generalized friendship graph* of H. Nalliah, Shankar, and Wang [2648] determined the [2648] new
local antimagic chromatic number of various generalized friendship graphs.

For a connected graph $G(V, E)$ with $|V| = n$ and $|E| = m$, a bijection $f : V \cup E \to \{1, 2, 3, \ldots, n+m\}$ is called a *local antimagic vertex total coloring* of G if for any two adjacent vertices u and v, $w_t(u) \neq w_t(v)$, where $w_t(u)$ is the sum of $f(e) + f(u)$ over all edges e incident to u. The *local antimagic vertex total chromatic number* is the minimum number of colors taken over all colorings induced by local antimagic vertex total labelings of G. Amalia and Masruroh [209] proved that the local antimagic vertex total chromatic [209] new
number of fan graph F_n $(n \geq 3)$ is 3. They also gave a result about the local antimagic vertex total coloring number of a family of graphs obtained by grafting copies of center of F_3 to vertices of F_n in a special way.

Ahmad, Bača, Lascsáková and Semaničová-Feňovčíková [114] call a labeling of a plane graph *d-antimagic* if for every positive integer s, the set of s-sided face weights is $W_s = \{a_s, a_s + d, a_s + 2d, \ldots, a_s + (f_s - 1)d\}$ for some positive integers as a_s and d, where f_s is the number of the s-sided faces. (They allow different sets W_s for different s). A d-antimagic labeling is called *super* if the smallest possible labels appear on the vertices. In [185] they investigated the existence of super d-antimagic labelings of type $(1, 1, 0)$ for disjoint union of plane graphs for several values of difference d. Bača, Numan, and Semaničová-Feňovčíková [415] investigated the existence of super d-antimagic labelings of generalized prisms. Hussainn and Tabraiz [1477] investigated super d-antimagic labeling of type (1,1,1) on the snakes kC_5; subdivided kC_5; and isomorphic copies of kC_5 for strings $(1, 1, \ldots, 1)$ and $(2, 2, \ldots, 2)$.

In [2235] Li, Lau, and Shiu defined an edge labeling of a connected graph $G(V, E)$ of order p and size q to be a *modulo local antimagic labeling* if it is a bijection $\pi : E \to \{1, \ldots, q\}$ such that for any pair of adjacent vertices u and v, $\pi^+(u) \neq \pi^+(v)$, where the induced vertex label $\pi^+(u) = \sum \pi(e) \pmod p$, with e ranging over all the edges incident to u. The *modulo local antimagic chromatic number* of G, denoted $\pi_{la}(G)$, is the minimum number of distinct induced vertex labels over all modulo local antimagic labelings of G. They investigated the relationship among chromatic number, local antimagic chromatic number, and modulo local antimagic chromatic number of graphs. They also determined the modulo local antimagic chromatic numbers of $K_{m,m}$, P_n, tadpoles, coconut trees and combs.

Bača, Baskoro, Jendroľ, and Miller [356] investigated various k-antimagic labelings for graphs in the shape of hexagonal honeycombs. They use H_n^m to denote the honeycomb graph with m rows, n columns, and mn 6-sided faces. They prove: for n odd H_n^m, has a 0-antimagic vertex labeling and a 2-antimagic edge labeling, and if n is odd and $mn > 1$, H_n^m has a 1-antimagic face labeling. In [3552] Shiu and Low show how to

construct k-antimagic graphs from existing graphs G with particular labeling properties by joining G to cycles and dumbbell related graphs with an edge.

Huang, Wong, and Zhu [1468] say a graph G is *weighted-k-antimagic* if for any vertex weight function w from the vertices of G to the natural numbers there is an injection f from the edges of G to $\{1, 2, \ldots, |E| + k\}$ such that for any two distinct vertices u and v, $\sum(f(e) + w(v)) \neq \sum(f(e) + w(u))$ over all edges incidence to v. They proved that if G has odd prime power order p^z and has total domination number 2 with the degree of one vertex in the total dominating set not a multiple of p, then G is weighted-1-antimagic, and if G has odd prime power order p^z, $p \neq 3$ and has maximum degree at least $|V(G)| - 3$, then G is weighted-1-antimagic. Wong and Zhu [4143] proved: graphs that have a vertex that is adjacent to all other vertices are weighted-2-antimagic; graphs with a prime number of vertices that have a Hamiltonian path are weighted-1-antimagic; and connected graphs $G \neq K_2$ on n vertices are weighted-$\lfloor 3n/2 \rfloor$-antimagic. Matamala and Zamora [2455] proved that $K_{m,n}$, $3 \leq m \leq n$, $n \geq 3$, is weighted-0-antimagic and described a polynomial time algorithm that computes a $(w, 0)$-antimagic labeling of $K_{m,n}$. They also prove the following. Let H be an arbitrary complete partite graph with $n \geq 5$ vertices not isomorphic to $K_{1,n}$. Then, any graph containing H as a spanning subgraph is weighted-0-antimagic and given a weight function w, a $(w, 0)$-antimagic labeling can be computed in polynomial time. They prove that each connected graph G on $n \geq 3$ vertices having $K_{1,n}$ as a spanning subgraph is weighted-1-antimagic unless G is isomorphic to $K_{1,n}$ and n is even.

A graph G is *weighted k-list-antimagic* if for any vertex weighting $\omega : V(G) \to \mathbb{R}$ and any list assignment $L : E(G) \to 2^{\mathbb{R}}$ with $|L(e)| \geq |E(G)| + k$, there exists an edge labeling f such that $f(e) \in L(e)$ for all $e \in E(G)$, labels of edges are pairwise distinct, and the sum of the labels on edges incident to a vertex plus the weight of that vertex is distinct from the sum at every other vertex. Berikkyzy, Brandt, Jahanbekam, Larsen, and Rorabaugh [626] proved that every graph on n vertices having no K_1 or K_2 component is weighted-$\lfloor \frac{4n}{3} \rfloor$-list-antimagic.

A *distance k-antimagic* labeling of a graph $G(V, E)$ is a bijection $\bar{}f$ from V to $\{1, 2, \ldots, |V|\}$ with the property that there exists an ordering of the vertices of G such that the sequence of the weights $w(x_1), w(x_2), \ldots, w(x_n)$ forms an arithmetic progression with difference k. When $k = 1$, then $\bar{}f$ is simply called a *distance antimagic* labeling. A *distance k-antimagic* graph is a distance k-antimagic graph that admits a distance k-antimagic labeling, and is called *distance antimagic* when $k = 1$. Cichacz, Fronček, Sugeng and Zhou in [847] gave a necessary condition for a graph with an even number of vertices to be distance antimagic with respect to an Abelian group with a unique involution. They also gave sufficient conditions for a Cayley graph on an Abelian group to be distance antimagic or magic with respect to the same group, and discussed the consequences of these results to Cayley graphs on elementary Abelian groups. In [1360] Handa, Godinho, and Singh investigate the existence of distance antimagic labelings of ladders.

For a positive integer k, define $f_k : V(G) \longrightarrow \{1 + k, 2 + k, \ldots, n + k\}$ by $f_k(x) = f(x) + k$. If $w_{f_k}(u) \neq w_{f_k}(v)$ for every pair of vertices $u, v \in V$, for any $k \geq 0$ then

f is said to be an *arbitrarily distance antimagic* labeling and the graph which admits such a labeling is said to be an *arbitrarily distance antimagic* graph. Handa, Godinho, and Singh [1361] provide arbitrarily distance antimagic labelings for rP_n, the generalized Petersen graph $P(n,k)$, $n \geq 5$, the Harary graph $H_{4,n}$ for $n \neq 6$ and prove that join of these graphs is distance antimagic.

For an arbitrary set of distances $D \subseteq \{0, 1, \ldots, diam(G)\}$, a D-weight of a vertex x in a graph G under a vertex labeling $f : V \to \{1, 2, \ldots, v\}$ is defined as $w_D(x) = \sum_{y \in N_D(x)} f(y)$, where $N_D(x) = \{y \in V \mid d(x,y) \in D\}$. A graph G is said to be *D-distance magic* if all vertices has the same D-vertex-weight, it is said to be *D-distance antimagic* if all vertices have distinct D-vertex-weights. In [3587] Simanjuntak and Wijaya gave some necessary conditions for the existence of D-distance antimagic graphs and conjectured that those conditions are sufficient. They also gave $\{1\}$-distance antimagic labelings for cycles, suns, prisms, complete graphs, wheels, fans, and friendship graphs.

In [279] Arumugam and Kamatchi introduced the notion of (a,d)-distance antimagic graphs as follows. Let G be a graph with vertex set V and $f : V \to \{1, 2, \ldots, |V|\}$ be a bijection. If for all v in G the set of sums $\sum f(u)$ taken over all neighbors u of v is the arithmetic progression $\{a, a+d, a+2d, \ldots, a+(|V|-1)d\}$, f is called an (a,d)-*distance antimagic* labeling and G is called a (a,d)-*distance antimagic graph*. Arumugam and Kamatchi [279] proved: C_n is (a,d)-distance antimagic if and only if n is odd and $d = 1$; there is no $(1,d)$-distance antimagic labeling for P_n when $n \geq 3$; a graph G is $(1,d)$-distance antimagic graph if and only if every component of G is K_2; $C_n \times K_2$ is $(n+2,1)$-distance antimagic; and the graph obtained from $C_{2n} = (v_1, v_2, \ldots, v_{2n})$ by adding the edges v_1v_{n+1} and v_iv_{2n+2-i} for $i = 2, 3, \ldots, n$ is $(2n+2,1)$-distance antimagic. In [1127] and [1129] Fronček proved that disjoint copies of the Cartesian product of two complete graphs and its complement are $(a,2)$-distance antimagic and $(a,1)$-distance antimagic. He also proved that disjoint copies of the hypercube Q_3 is $(a,1)$-distance antimagic. Semeniuta [3358] proved that the crown $P_n \odot P_1$ does not admit an $(a,1)$-distance antimagic labeling for $n \geq 2$ and $a \geq 2$ and determines the values of a for which P_n can be an $(a,1)$-distance antimagic graph. The circulant graph is also investigated. Semenyuta [3359] proved that $P_n \odot P_1$ is not an (a,d)-distance antimagic graph for all a and d and that Q_n is a $(2^n+n-1, n-2)$-distance antimagic graph. He found two types of graphs that do not allow 1-vertex bimagic vertex labeling and established a relation between the distance magic labeling of a regular graph G with 1-vertex bimagic vertex labeling $G \cup G$. In [2697] Ngurah constructed new classes of (super) (a,d)-antimagic 2-regular graphs from known super edge-magic total 2-regular graphs. Dak, and Meganingtyas, Dafik, and Slamin [2481] investigated the existence of super (a,d)-vertex antimagic labeling of directed cycles.

Patel and Vasva [2805] use $Circ(n,k)$ $n \geq 3$ to denote the graph $Circ(n, \{k\})$ defined as follows: for a subset $S \subset \{1, 2, \ldots, n\}$, the circulant graph $Circ(n,S)$ is the graph with vertex set $\{v_1, v_2, \ldots, v_n\}$ and there is an edge between vertices v_i and v_j if and only if $|i-j| \in S \bigcup \{1, n-1\}$). In [2805] they proved the existence or non-existence of (a,d)-distance antimagic labelings for the following graphs: $Circ(2n, \{1,n\})$ is $(2n+2,1)$-distance antimagic for all even n; mK_{2n} is $(n(2mn-2m+1),1)$-distance antimagic for

all m and n; $3K_{2n+1}$ is $(6n^2+n-1,1)$-distance antimagic for all n; $2K_{2n+1}$ is not (a,d)-distance antimagic for all n; helms H_n is not (a,d)-distance antimagic for any n; books $B_n = S_n \times P_2$ of order $2n+2$ are not (a,d)-distance antimagic for any n; and $K_n \odot K_1$ is not (a,d)-distance antimagic for $n > 1$.

Kamatchi, Vijayakumar, Ramalakshmi, Nilavarasi, and Arumugam [1820] prove that the hypercube is (a,d)-distance antimagic and the bistar $K_2(n,n)$ is distance antimagic. They also show that if G is a regular distance antimagic graph, then $2G$ is also distance antimagic and several families of disconnected graphs are distance antimagic graphs.

A connected graph $G = (V,E)$ with m edges is called *universal antimagic* if for each set B of m positive integers there is an bijective function $f : E \to B$ such that the function $\tilde{f} : V \to \mathbb{N}$ defined at each vertex v as the sum of all labels of edges incident to v is injective. Matamala and Zamora [2454] proved that paths, cycles, split graphs, and graphs that contains the complete bipartite graph $K_{2,n}$ as a spanning subgraph are universal antimagic. A universal antimagic graph is *weight universal antimagic* if, in addition, for any weight function w on the vertices, all $w(u)+f^u$ are distinct. Generalizing their previous result [2454], in [2456] the authors show constructively that if a graph has a complete bipartite graph $K_{m,n}$ as a spanning subgraph with $m,n \geq 3$, then it is weighted universal antimagic (hence universal antimagic). They also show that for all other values of m and n, the graph is universal antimagic.

In [3896] Thirusangu and Selvaganapathy introduced the notion of Gaussian antimagic labeling as follows. For a graph $G(p,q)$ a *Gaussian antimagic labeling* of G is a function $f : V(G) \to \{a+ib \mid a,b \in \mathbb{N},\ 1 \leq a < b \leq q\}$ such that the induced function $f^* : E(G) \to \mathbb{N}$, defined by $f^*(uv) = f(u)^2 + f(v)^2$, assigns distinct labels to all edges. A graph that admits a Gaussian antimagic labeling is called a *Gaussian antimagic graph.* They proved that paths, cycles, Y-trees, combs, triangular snakes, and chordal graphs are Gaussian antimagic. In [3899] Thirusangu and Selvaganapathy investigated the existence of Gaussian antimagic labelings for shadow graphs, split graphs, jewel graphs, jellyfish graphs, modified extended duplicate graphs of stars and paths, $K_p \odot K_t$, $W_p \odot K_t$, $S_p \odot K_t$, and $H_p \odot K_t, C_m + K_1$, Mongolian tents, Petersen graphs, coconuts, centipede trees, and square graphs of paths. In [3899] Thirusangu and Selvaganapathy investigated the existence of Gaussian antimagic labelings for sunlet graphs (that is, $C_n \odot K_1$), path unions of sunlet graphs, the one point union of two copies of sunlet graphs, the join sum of two copies of a sunlet graph sharing a common edge, alternate triangular snakes, double triangular snakes, alternate double triangular snakes, quadrilateral snakes, and alternate quadrilateral snakes. In [3264] Selvaganapathy and Thirusangu investigated the existence of Gaussian antimagic labelings for the path union of fans, double fans, globes, pagodas, ladders, and triangular ladders. In [3265] they investigated the existence of Gaussian antimagic labelings for certain spider-related graphs and n-pans (obtained by joining C_n to K_1). Selvaganapathy and Ramesh [3263] investigated the existence of Gaussian antimagic labelings for various families of trees. Selvaganapathy and Lalithkumar [3262] demonstrated the existence of Gaussian antimagic labelings for varvious families of graphs. Thirusangu and Selvaganapathy [3898] studied the existence of Gaussian antimagic labelings for middle graphs of paths, split graphs of paths, shadow graphs of paths, total

[3896] new
[3899] new
[3899] new
[3264] new
[3265] new
[3263] new
[3262] new
[3898] new

path graphs, middle cycle graphs, ladders, square combs, Splitting comb graphs, Theta graphs, the sum of two copies of Theta graphs by the path of length two, cubic graphs, and the path union of two copies of a cubic graph.

In 2019 [411] Bača, Miller, Ryan, and Semaničová-Feňovčíková published a monograph that focuses on variations of magic and antimagic type labelings and includes new results, techniques, constructions, and open problems and conjectures.

In Table 12 we use the abbreviation **A** to mean antimagic. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The table was prepared by Petr Kovář and Tereza Kovářová and updated by J. Gallian in 2014.

Table 12: **Summary of Antimagic Labelings**

Graph	*Labeling*	*Notes*
P_n	A	for $n \geq 3$ [1378]
C_n	A	[1378]
W_n	A	[1378]
K_n	A	for $n \geq 3$ [1378]
every tree except K_2	A?	[1378]
caterpillars	A	[2364]
regular graphs	A	[2247], [1378], [771]
every connected graph except K_2	A?	[1378]
$n \geq 4$ vertices $\Delta(G) \geq n-2$	A	[201]
all complete partite graphs except K_2	A	[201]
$C_m \times P_n$	A	[804]
$P_{m_1} \times P_{m_2} \times \cdots \times P_{m_k}$	A	[804]

Continued on next page

Table 12 – *Continued from previous page*

Graph	*Labeling*	*Notes*
$C_{m_1} \times C_{m_2} \times \cdots \times C_{m_k}$	A	[4144]
C_n^2	A	[2142]
mP_3 $m \geq 2$	not A	[4154]

6.2 (a, d)-Antimagic Labelings

The concept of an (a, d)-antimagic labelings was introduced by Bodendiek and Walther [674] in 1993. A connected graph $G = (V, E)$ is said to be (a, d)-*antimagic* if there exist positive integers a, d and a bijection $f\colon E \to \{1, 2, \ldots, |E|\}$ such that the induced mapping $g_f\colon V \to N$, defined by $g_f(v) = \sum\{f(uv)|\ uv \in E(G)\}$, is injective and $g_f(V) = \{a, a+d, \ldots, a+(|V|-1)d\}$. (In [2271] Lin, Miller, Simanjuntak, and Slamim called these (a, d)-*vertex-antimagic edge labelings*). Bodendick and Walther ([676] and [677]) prove the Herschel graph is not (a, d)-antimagic and obtain both positive and negative results about (a, d)-antimagic labelings for various cases of graphs called *parachutes* $P_{g,p}$. ($P_{g,p}$ is the graph obtained from the wheel W_{g+p} by deleting p consecutive spokes.) In [371] Bača and Holländer prove that necessary conditions for $C_n \times P_2$ to be (a, d)-antimagic are $d = 1,\ a = (7n+4)/2$ or $d = 3,\ a = (3n+6)/2$ when n is even, and $d = 2,\ a = (5n+5)/2$ or $d = 4,\ a = (n + 7)/2$ when n is odd. Bodendiek and Walther [675] conjectured that $C_n \times P_2$ $(n \geq 3)$ is $((7n + 4)/2, 1)$-antimagic when n is even and is $((5n + 5)/2, 2)$-antimagic when n is odd. These conjectures were verified by Bača and Holländer [371] who further proved that $C_n \times P_2$ $(n \geq 3)$ is $((3n + 6)/2, 3)$-antimagic when n is even. Bača and Holländer [371] conjecture that $C_n \times P_2$ is $((n + 7)/2, 4)$-antimagic when n is odd and at least 7. Bodendiek and Walther [675] also conjectured that $C_n \times P_2$ $(n \geq 7)$ is $((n+7)/2, 4)$-antimagic. Miller and Bača [2493] prove that the generalized Petersen graph $P(n, 2)$ is $((3n+6)/2, 3)$-antimagic for $n \equiv 0 \pmod 4$, $n \geq 8$ and conjectured that $P(n, k)$ is $((3n + 6)/2, 3)$-antimagic for even n and $2 \leq k \leq n/2 - 1$ (see §2.7 for the definition of $P(n, k)$). This conjecture was proved for $k = 3$ by Xu, Yang, Xi, and Li [4235]. Jirimutu and Wang proved that $P(n, 2)$ is $((5n + 5)/2, 2)$-antimagic for $n \equiv 3 \pmod 4$ and $n \geq 7$. Xu, Xu, Lü, Baosheng, and Nan [4231] proved that $P(n, 2)$ is $((3n + 6)/2, 2)$-antimagic for $n \equiv 2 \pmod 4$ and $n \geq 10$. Xu, Yang, Xi, and Li [4235] proved that $P(n, 3)$ is $((3n+6)/2, 3)$-antimagic for even $n \geq 10$ and for $n \equiv 0 \pmod 4$, $n \geq 8$. In [2274] Lingqi, Linna, Yuan show that $P(n, 3)$ is $(5n + 5)/2, 2)$-antimagic for odd $n \geq 7$. Feng, Hong, Yang, and Jirimutu [1077] show that $P(n, 5)$ is $(3n + 6)/2, 3)$-antimagic for even $n \geq 12$. Bao, Zhao, Yang, Feng, and Jirimutu [480] proved that $P(n, 7)$ is $(\frac{3n+6}{2}, 3)$-antimagic for even $n \geq 16$. Ivančo [1554] investigated $(a, 1)$-antimagic labelings and their connection with supermagic generalized double graphs. Bodendiek and Walther [678] proved that the following graphs are not (a, d)-antimagic: even cycles; paths of even order; stars;

$C_3^{(k)}; C_4^{(k)}$; trees of odd order at least 5 that have a vertex that is adjacent to three or more end vertices; n-ary trees with at least two layers when $d = 1$; the Petersen graph; K_4 and $K_{3,3}$. They also prove: P_{2k+1} is $(k, 1)$-antimagic; C_{2k+1} is $(k+2, 1)$-antimagic; if a tree of odd order $2k+1$ $(k > 1)$ is (a, d)-antimagic, then $d = 1$ and $a = k$; if K_{4k} $(k \geq 2)$ is (a, d)-antimagic, then d is odd and $d \leq 2k(4k-3)+1$; if K_{4k+2} is (a, d)-antimagic, then d is even and $d \leq (2k+1)(4k-1)+1$; and if K_{2k+1} $(k \geq 2)$ is (a, d)-antimagic, then $d \leq (2k+1)(k-1)$. Lin, Miller, Simanjuntak, and Slamin [2271] show that no wheel W_n $(n > 3)$ has an (a, d)-antimagic labeling. In [3817] Susanto provided super (a, d)-C_n-antimagic total labelings for various cases of mC_n. In [1928], Kathiresan and Laurence posed the problem of characterizing the super $(a, 1) - P_3$-antimagic total labelings of the stars S_n, where $n = 6, 7, 8$, and 9. This problem was completely solved in [3103] by Rajkumar, Nalliah, and Venkataraman. In [4262] Yatin1, Awanis, and Wardhana showed that for odd m, $P_n \odot P_m$ is super $(9m2n + 4mn + m - n + 3, 1) - P_2 \odot P_m$-antimagic, and for m even, $P_n \odot P_m$ is super $(9m2n + 4mn + m - 2n + 5, 3) - P_2 \odot P_m$-antimagic.

In [1562] Ivančo, and Semaničová show that a 2-regular graph is super edge-magic if and only if it is $(a, 1)$-antimagic. As a corollary we have that each of the following graphs are $(a, 1)$-antimagic: kC_n for n odd and at least 3; $k(C_3 \cup C_n)$ for n even and at least 6; $k(C_4 \cup C_n)$ for n odd and at least 5; $k(C_5 \cup C_n)$ for n even and at least 4; $k(C_m \cup C_n)$ for m even and at least 6, n odd, and $n \geq m/2 + 2$. Extending a idea of Kovář they prove if G is $(a_1, 1)$-antimagic and H is obtained from G by adding an arbitrary $2k$-factor then H is $(a_2, 1)$-antimagic for some a_2. As corollaries they observe that the following graphs are $(a, 1)$-antimagic: circulant graphs of odd order; $2r$-regular Hamiltonian graphs of odd order; and $2r$-regular graphs of odd order $n < 4r$. They further show that if G is an $(a, 1)$-antimagic r-regular graph of order n and $n - r - 1$ is a divisor of the non-negative integer $a + n(1 + r - (n+1)/2)$, then $G \oplus K_1$ is supermagic. As a corollary of this result they have if G is $(n-3)$-regular for n odd and $n \geq 7$ or $(n-7)$-regular for n odd and $n \geq 15$, then $G \oplus K_1$ is supermagic.

For a graph $G(V, E)$ a function $f : V \cup E \to \{1, 2, \ldots, |V| + |E|\}$ is said to be an (a, d)-*edge antimagic total labeling* if the set of edge weights $\{f(x) + f(xy) + f(y) : xy \in E\}$ forms an arithmetic progression with an initial term a and a common difference d. Such a labeling is called a *super (a, d)-edge antimagic total* labeling) if the smallest labels are assigned to the vertices. Such a labeling is called a *local super (a,d) edge-antimagic total* labeling if $f(E) = \{1, 2, \ldots, |E|\}$. Muthuselvi and Saratha Devi [2621] investigated the existence of the local super (a, d)-edge antimagic total labelings for certain graph classes and obtained a relationship between a local super $(a, 0)$-edge antimagic total labeling and a local super $(a, 2)$-edge antimagic total labeling of any graph. [2621] new

Bertault, Miller, Feria-Purón, and Vaezpour [634] approached labeling problems as combinatorial optimization problems. They developed a general algorithm to determine whether a graph has a magic labeling, antimagic labeling, or an (a, d)-antimagic labeling. They verified that all trees with fewer than 10 vertices are super edge magic and all graphs of the form $P_2^r \times P_3^s$ with less than 50 vertices are antimagic. Javaid, Hussain, Ali, and Dar [1585] and Javaid, Bhatti, and Hussain [1582] constructed super (a, d)-edge-antimagic total labelings for w-trees and extended w-trees (see 5.2 for the definitions) as well as super

(a,d)-edge-antimagic total labelings for disjoint union of isomorphic and non-isomorphic copies of extended w-trees. In [1583] Javaid and Bhatt defined a generalized w-tree and proved that they admit a super (a,d)-edge-antimagic total labeling. In [4152] Wang, Li, and Wang proved that some classes of graphs derived from regular or regular bipartite graphs are antimagic. A subdivided star $T(n_1,n_2,\ldots,n_r)$ is a tree obtained by inserting $n_i \geq 1,\ 1 \leq i \leq r$ with $r \geq 3$ vertices.

In [3070] Raheem, Javaid, and Baig study a super (a,d)-edge-antimagic total labelings of the subdivided stars $T(n,n+1,n_3,\ldots,n_r)$ when n is even and $T(n,n,n+1,n_4,\ldots,n_r)$ when n is odd for all possible values of d. In [3069] Raheem and Baig proved the super edge antimagicness of subdivided stars for all possible values of d. Bhatti, Tahir, and Javaid [652] give super (a,d)-edge antimagic total labelings of some wheel-like graphs. In [1581] Javaid, Bhatti, and Aslam investigated the existence of super (a,d)-edge antimagic total labelings of subdivided stars for all possible values of d but mainly $d=1$ and 3. In [283] investigate the existence of super (a,d)-edge antimagic total labelings for friendship graphs and generalized friendship graphs. Girija and Karthikeyan [1275] proved that 3 copies of the jelly fish graphs are super (a,d)-edge antimagic vertex graphs and super (a,d)-edge antimagic total graphs. The Hill cipher is a cryptographic algorithm that uses modulo arithmetic and a matrix as a key to perform encryption and decryption. In [3041] Prihandini and Adawiyah1 discussed how to use a super $(3n+5,2)$-edge antimagic total labeling to construct the Hill cipher algorithm. They stated that variations of the edge weight function and the corresponding edge label on the graph will make the constructed lock more complicated to hack.

In [83] Afzal, Javaid, Alanazi, and Alshehri investigated the super $(a,0)$ edge-antimagicness of the union of the networks of stars, paths, and copies of paths and the rooted product of C_n with $K_{2,n}$. They also provided super $(a,0)$ edge-antimagic labelings of the rooted product of cycles and planar pancyclic networks, and give super $(a,0)$ edge-antimagic labelings for a pancyclic network containing chains of C_6, and three different symmetrically designed lattices.

For graphs G and F, if every edge of G belongs to a subgraph of G isomorphic to F and there exists a total labeling λ of G such that for every subgraph F' of G that is isomorphic to F, the set $\{\Sigma\lambda(F') :\ F' \cong F,\ F' \subseteq G\}$ forms an arithmetic progression starting with a with common difference d, Lee, Tsai, and Lin [2141] say that G is (a,d)-*F-antimagic.* Furthermore, if $\lambda(V(G)) = \{1,2,\ldots,|V(G)|\}$ then G is said to be *super* (a,d)-*F-antimagic* and λ is said to be a *super* (a,d)-*F-antimagic* labeling of G. Lee, Tsai, and Lin [2141] proved that $P_m \times P_n$ $(m,n \geq 2)$ is super $(a,1)$-C_4-antimagic. In [3344] Selvagopal, Jeyanthi, Muthuraja, and Semaničová-Feňovčiková investigated the existence super (a,d)-star-antimagic labelings of a particular class of banana trees and construct a star-antimagic graph. In [1763] Jeyanthi, Selvi, and Ramya proved the existence of super (a,d)-C_n-antimagic labelings of fan graphs and ladders. Inayah [1530] proved the existence of an $(a,b)-P_4$-antimagic decomposition of a generalized Petersen graph $GP_z(n,3)$ for several values of b. Taimur, Ali, Numan, Aslam, and Anoh Yannick [3857] provided super (a,d)-C_3-antimagic total labelings for the generalized antiprism for $d=0$ and 1 and a super (a,d)-C_8-antimagic total labeling for the toroidal octagonal map for $d=1,2,\ldots,7$.

The *edge corona path graph* $G_m \diamond P_n$ is the graph obtained from one copy of the gear graph G_m and $3m$ copies of P_n, P_n^i, by joining two end vertices of $e_i \in E(G_m)$ to every vertex $v_j \in V(P_n)$ in the i-th copy of G_m with $i = 1, 2, \ldots, 3m$ and $j = 1, 2, \ldots, n$. The graph $G_m \cdot C_n$ is the graph obtained from G_m and $2m+1$ copies of C_n namely C_n^i by joined every vertex $v_i \in G_m$ to all vertices $v_i \in C_n$ for $i \in \{1, 2, \ldots, 2m+1\}$. Nistyawati and Martini [2716] proved that for every odd m, the gear edge corona path graph $G_m \diamond P_n$ is super $C_4 \diamond P_n$-antimagic and for every odd m, the gear corona cycle graph $G_m \cdot C_n$ is super $C_4 \cdot C_n$-antimagic. Roswitha, Martini, and S. A. Nugroho [3210] proved: for $n \geq 5$ the double cone $DC_n = C_n + \overline{K_2}$ is $(14+7n+(n+1)2, 1)$-C_3-antimagic and $(a, 1)$-C_3-antimagic; DC_n is (a, d)-W_n-antimagic; DC_{2n} is $(a, 1)$-W_{2n}-antimagic; and DC_{2n+1} is $(a, 2)$-W_{2n+1}-antimagic.

Yegnanarayanan [4264] introduced several variations of antimagic labelings and provides some results about them.

The *antiprism* on $2n$ vertices has vertex set $\{x_{1,1}, \ldots, x_{1,n}, x_{2,1}, \ldots, x_{2,n}\}$ and edge set $\{x_{j,i}, x_{j,i+1}\} \cup \{x_{1,i}, x_{2,i}\} \cup \{x_{1,i}, x_{2,i-1}\}$ (subscripts are taken modulo n). For $n \geq 3$ and $n \not\equiv 2 \pmod 4$ Bača [343] gives $(6n+3, 2)$-antimagic labelings and $(4n+4, 4)$-antimagic labelings for the antiprism on $2n$ vertices. He conjectures that for $n \equiv 2 \pmod 4$, $n \geq 6$, the antiprism on $2n$ vertices has a $(6n+3, 2)$-antimagic labeling and a $(4n+4, 4)$-antimagic labeling.

Nicholas, Somasundaram, and Vilfred [2713] prove the following: If $K_{m,n}$ where $m \leq n$ is (a, d)-antimagic, then d divides $((m-n)(2a+d(m+n-1)))/4 + dmn/2$; if $m+n$ is prime, then $K_{m,n}$, where $n > m > 1$, is not (a, d)-antimagic; if $K_{n,n+2}$ is (a, d)-antimagic, then d is even and $n+1 \leq d < (n+1)^2/2$; if $K_{n,n+2}$ is (a, d)-antimagic and n is odd, then a is even and d divides a; if $K_{n,n+2}$ is (a, d)-antimagic and n is even, then d divides $2a$; if $K_{n,n}$ is (a, d)-antimagic, then n and d are even and $0 < d < n^2/2$; if G has order n and is unicylic and (a, d)-antimagic, then $(a, d) = (2, 2)$ when n is even and $(a, d) = (2, 2)$ or $(a, d) = ((n+3)/2, 1)$ when n is odd; a cycle with m pendent edges attached at each vertex is (a, d)-antimagic if and only if $m = 1$; the graph obtained by joining an endpoint of P_m with one vertex of the cycle C_n is $(2, 2)$-antimagic if $m = n$ or $m = n-1$; if $m+n$ is even the graph obtained by joining an endpoint of P_m with one vertex of the cycle C_n is (a, d)-antimagic if and only if $m = n$ or $m = n-1$. They conjecture that for n odd and at least 3, $K_{n,n+2}$ is $((n+1)(n^2-1)/2, n+1)$-antimagic and they have obtained several results about (a, d)-antimagic labelings of caterpillars.

In [2365] Lozano, Mora, and Seara prove that any caterpillar of order n is $(\lfloor (n-1)/2, \rfloor - 2)$-antimagic. Furthermore, if C is a caterpillar with a spine of order s, they prove that when C has at least $\lfloor (3s+1)/2 \rfloor$ leaves or $\lfloor (s-1)/2 \rfloor$ consecutive vertices of degree at most 2 at one end of a longest path, then C is antimagic. As a consequence of a result of Wong and Zhu [4210], they also prove that if p is a prime number, any caterpillar with a spine of order p, $p-1$ or $p-2$ is 1-antimagic.

In [4104] Vilfred and Florida proved the following: the one-sided infinite path is $(1, 2)$-antimagic; P_{2n} is not (a, d)-antimagic for any a and d; P_{2n+1} is (a, d)-antimagic if and only if $(a, d) = (n, 1)$; C_{2n+1} has an $(n+2, 1)$-antimagic labeling; and that a 2-regular graph G is (a, d)-antimagic if and only if $|V(G)| = 2n+1$ and $(a, d) = (n+2, 1)$. They

also prove that for a graph with an (a,d)-antimagic labeling, q edges, minimum degree δ and maximum degree Δ, the vertex labels lie between $\delta(\delta+1)/2$ and $\Delta(2q-\Delta+1)/2$.

Chelvam, Rilwan, and Kalaimurugan [793] proved that Cayley digraph of any finite group admits a super vertex (a,d)-antimagic labeling depending on d and the size of the generating set. They provide algorithms for constructing the labelings. Thirusangu and Bhrathiraja [3892] proved the existence of super vertex (a,d)- antimagic labeling and vertex magic total labeling for Cayley digraphs arising from the groups $Z_{p^n} \oplus Z_{p^n}$ and $Z_m \oplus Z_n$.

Irfan and Semaničová-Feňovčiková [1547] provide some classes of graphs that admit a labeling that is simultaneously a super edge-magic total and a super vertex-antimagic total and give some results for fans, sun graphs, caterpillars, and prisms.

For $n > 1$ and distinct odd integers x, y and z in [1,$n-1$] Javaid, Ismail, and Salman [1576] define the *chordal ring* of order n, $CR_n(x,y,z)$, as the graph with vertex set Z_n, the additive group of integers modulo n, and edges $(i,i+x),(i,i+y),(i,i+z)$ for all even i. They prove that $CR_n(1,3,7)$ and $CR_n(1,5,n-1)$ have (a,d)-antimagic labelings when $n \equiv 0 \bmod 4$ and conjecture that for an odd integer $\Delta,\ 3 \le \Delta \le n-3, n \equiv 0 \bmod 4$, $CR_n((1,\Delta,n-1)$ has an $((7n+8)/4,1)$-antimagic labeling.

For an arbitrary set of distances $D \subseteq \{0,1,\ldots,diam(G)\}$, a D-weight of a vertex x in a graph G under a vertex labeling $f : V \to \{1,2,\ldots,v\}$ is defined as $w_D(x) = \sum_{y \in N_D(x)} f(y)$, where $N_D(x) = \{y \in V | d(x,y) \in D\}$. A graph G is said to be D-distance magic if all vertices have the same D-vertex-weight, it is said to be D-distance antimagic index$D$-distance antimagic if all vertices have distinct D-vertex-weights, and it is called $(a,d)-D$-distance antimagic if the D-vertex-weights constitute an arithmetic progression with difference d and starting value a. In [3587] Simanjuntak and Wijaya gave some necessary conditions for the existence of D-distance antimagic graphs and conjectured that those conditions are sufficient. They also gave $\{1\}$-distance antimagic labelings for cycles, suns, prisms, complete graphs, wheels, fans, and friendship graphs. Arumugam and Kamatchi [279] characterized (a,d)-distance antimagic cycles and (a,d)-distance antimagic labelings for paths and prisms. In [1127] and [1129] Fronček proved that disjoint copies of the Cartesian product of two complete graphs and its complement are $(a,2)$-distance antimagic and $(a,1)$-distance antimagic. He also proved that disjoint copies of the hypercube Q_3 is $(a,1)$-distance antimagic. In [1360] Handa, Godinho and Singh investigate the existence of distance antimagic labeling of ladders.

In [4105] Vilfred and Florida call a graph $G = (V,E)$ *odd antimagic* if there exist a bijection $f\colon E \to \{1,3,5,\ldots,2|E|-1\}$ such that the induced mapping $g_f\colon V \to N$, defined by $g_f(v) = \sum\{f(uv) |\ uv \in E(G)\}$, is injective and *odd* (a,d)*-antimagic* if there exist positive integers a, d and a bijection $f\colon E \to \{1,3,5,\ldots,2|E|-1\}$ such that the induced mapping $g_f\colon V \to N$, defined by $g_f(v) = \sum\{f(uv) |\ uv \in E(G)\}$, is injective and $g_f(V) = \{a, a+d, a+2d, \ldots, a+(|V|-1)d\}$. Although every (a,d)-antimagic graph is antimagic, C_4 has an antimagic labeling but does not have an (a,d)-antimagic labeling. They prove: P_{2n+1} is not odd (a,d)-antimagic for any a and d; C_{2n+1} has an odd $(2n+2,2)$-antimagic labeling; if a 2-regular graph G has an odd (a,d)-antimagic labeling, then $|V(G)| = 2n+1$ and $(a,d) = (2n+2,2)$; C_{2n} is odd magic; and an odd magic graph

with at least three vertices, minimum degree δ, maximum degree Δ, and $q \geq 2$ edges has all its vertex labels between δ^2 and $\Delta(2q - \Delta)$.

Combining the notions of 1-vertex-magic vertex labelings and antimagic labelings Swaminathan and Jeyanthi [3846] introduced a new labeling as follows. For a graph with p vertices a 1-1 mapping from the vertices to $\{1, 2, \ldots, p\}$ is called an (a, d)-1-*vertex-antimagic vertex labeling* if the sums of the labels of the vertices adjacent to each vertex taken over all vertices form the set $\{a, a+d, a+2d, \ldots, a+(p-1)d\}$. They give some basic properties of such labelings and provide some results for some classes of regular graphs.

For a graph $G = (V, E)$, a bijection g from $V(G) \cup E(G)$ into $\{1, 2, \ldots, |V(G)|+|E(G)|\}$ is called a (a, d)-*edge-antimagic graceful* labeling of G if the edge-weights $w(xy) = |g(x) + g(y) - g(xy)|$, $xy \in E(G)$, form an arithmetic progression starting from a and having a common difference d. An (a, d)-edge-antimagic graceful labeling is called *super* (a, d)-*edge-antimagic graceful* if $g(V(G)) = \{1, 2, \ldots, |V(G)|\}$. Marimuthu and Krishnaveni [2425] proved mC_n has a super $(0, 1)$-edge-antimagic graceful labeling for every $m \geq 2$ and $n \geq 3$; and mK_n and MP_n have a super $(a, 1)$-edge-antimagic graceful labeling for every $m \geq 2$ and $n \geq 2$. Marimuthu and Krishnaveni [2424] proved that $K_n, (n \geq 3)$ is super (a, d)-edge-antimagic graceful if and only $d \leq 1$ and that $K_{n,n}$ $(n \geq 2)$ is super (a, d)-edge-antimagic graceful if and only $d = 1$. [2424] new

In [144] Ahmed, Semaničová-Feňovčíková, Bača, Baskar Babujee, and Shobana introduced the notion of graceful antimagic graphs as follows. A graceful labeling that is simultaneously antimagic (that is, the sums of labels of all edges incident to a given vertex are pairwise distinct for different vertices) is said to be *graceful antimagic* . They obtain results about paths, stars, double stars, cycles, complete graphs, complete graphs with deleted edges, and complete bipartite graphs. They include many conjectures and open problems.

In [286] Arumugam, Premalatha, Bacă, and Semaničová-Feňovčíková introduced a new graph coloring parameter as follows. Let f be a local antimagic labeling of a connected graph G. The assignment of $w(u)$ to u for each vertex u of G induces naturally a proper vertex coloring of G, called a of G. The , denoted $\chi_{la}(G)$, is the minimum number of colors taken over all local antimagic colorings of G. Arumugam, Lee, Premalatha, Wang [281] proved that for $m > 1$, $\chi_{la}(C_3 \odot O_m) = 3m + 3$, where O_m is the null graph on m vertices, and for $n > 1$, $\chi_{la}(K_n \odot K_1) = 2n - 1$. In [2103] Lau and Nalliah corrected a mistake in [1440] for the lower bounds of the local antimagic chromatic number of the corona product of friendship and fan graphs with null graph and obtain a sharp lower bound that gives the exact local antimagic chromatic number of the corona product of friendship and null graph. In [2104] Lau, Ng, and Shiu give counterexamples to the lower bound of $\chi_{la}(G \vee O_2)$ that was obtained in [286]. A sharp lower bound of $\chi_{la}(G \vee O_n)$ and sufficient conditions for the given lower bound to be attained are obtained. Moreover, they improve a theorem and solve a problem stated in [286] and determine the local antimagic chromatic number of complete bipartite graphs. In [2116] Lau, Shiu, and Ng obtain a sharp lower bound of the local antimagic chromatic number of a graph with cut-vertices given by pendants is obtained. They also determine the exact value of the local antimagic chromatic number of many families of graphs with cut-vertices (possibly

given by pendant edges). Uma and Rajasekaran [3929] provided the vertex local antimagic chromatic number for some Knödel graphs, Fibonacci graphs, the disjoint union of Knödel graphs, the join graphs of Knödel graphs, complete graphs, cycles, complete bipartite graphs, and graphs consisting of any number n isolated vertices.

For a graph $G(V, E)$ a graph of order p without isolated vertices, a bijection $f : V \to \{1, 2, \ldots, p\}$ is called a *local distance antimagic* labeling, if the weights of every two adjacent vertices are unequal, where the weight of a vertex is defined to be the sum of labels of adjacent vertices. The *local distance antimagic chromatic number* is the minimum number of colors taken over all colorings of G induced by local distance antimagic labelings of G. In [177] Almeida and Singh studied the local distance antimagic chromatic number for the join of graphs and the lexicographic product of graphs with empty graphs. In [2115] Lau, Shiu, and Ng provide sharp upper and lower bounds of $\chi_{la}(G)$ for G with pendant vertices, and give sufficient conditions for equality. They show that there are infinitely many graphs with $k \geq \chi(G) - 1$ pendant vertices and $\chi_{la}(G) = k + 1$. They conjecture that every tree T_k, other than certain caterpillars, spiders and lobsters, with k pendant vertices has $\chi_{la}(T_k) = k+1$. Bača, Semaničová-Feňovčíková, Lai, and Wang [421] verified a conjecture by Arumugam, Lee, Premalatha, and Wang [281] that for every tree T the local antimagic chromatic number satisfies $l + 1 \leq \chi_{la}(T) \leq l + 2$, where l is the number of leaves of T, by determining the exact value for the local antimagic chromatic number of all complete full t-ary trees is $l + 1$ for odd t. In [3046] Priyadharshini and Nalliah determined the graphs whose local distance antimagic chromatic number is 2. In [73] Adawiyah, Makhfudloh, and Prihandini obtained results about the local (a, d)-antimagic chromatic number of sunflower graphs and umbrella graphs.

Lau, Shiu, and Ng [2121] obtained sharp upper and lower bounds of $\chi_{la}(G)$ for G with pendant vertices, and gave sufficient conditions for the bounds to equal. They also showed that for $k \geq 1$, there are infinitely many graphs with $k \geq \chi(G) - 1$ pendant vertices and $\chi_{la}(G) = k + 1$. In [2120] Lau, Premalatha, Shiu, and Nalliah determined the local antimagic chromatic numbers of the join product $G \vee H$, where G is a circulant graph and H is a null graph or a cycle. The local antimagic chromatic number of certain wheel related graphs are also obtained.

Lau, Shiu, and Soo [2125] showed that a d-leg spider graph has $d + 1 \leq \chi_{la} \leq d + 2$; each 3-leg spider has $\chi_{la} = 4$ if not all legs are of odd length; and that there are no 3-leg spiders with all odd leg lengths and $\chi_{la} = 5$. These results provide partial solutions to the characterization of k-pendant trees T with $\chi_{la}(T) = k + 1$ or $k + 2$. They also obtained many sufficient conditions such that both the values are attainable. They conjectured that almost all d-leg spiders of size q that satisfy $d(d + 1) \leq 2(2q - 1)$ when each leg length at least 2 has $\chi_{la} = d + 1$.

In [2123] Lau and Shiu showed the existence of non-complete regular graphs with arbitrarily large order, regularity, and local antimagic chromatic numbers. They also determined the local antimagic chromatic number of regular graphs of order at most 8. In [2124] for graphs G and H Lau and Shiu denote the lexicographic product of graphs G and H by $G[H]$. They obtained sharp upper bound of $\chi_{la}(G[O_n])$ $n \geq 3$), the lexicographic product of graph G with the null graph and determined the local antimagic chromatic

number of infinitely many (connected and disconnected) regular graphs that partially support the conjecture of the existence of an r-regular graph G of order p such that (i) $\chi_{la}(G) = \chi(G) = k$, and (ii) $\chi_{la}(G) = \chi(G) + 1 = k$ for each possible r, p, k. They further provided sufficient conditions for even regular bipartite and tripartite graphs G to have $\chi_{la}(G) = 3$.

In [2122] Lau, Shiu, Premalatha, Zhang and Nalliah obtained sufficient conditions for $\chi_{la}(G[H]) \leq \chi_{la}(G)\chi_{la}(H)$. They also gave examples of G and H such that $\chi_{la}(G[H]) = \chi(G)\chi(H)$, where $\chi(G)$ is the chromatic number of G. Premalatha, Lau, Arumugam, and Shiu [3039] gave sufficient conditions under which $\chi_{la}(G \vee H)$, where H is either a cycle or the null graph, satisfies a sharp upper bound. They further determined the value of $\chi_{la}(G \vee H)$ for many wheel-related graph G. Lau, Li, and Shiu [2102] showed the existence of infinitely many bipartite and tripartite graphs with local antimagic chromatic number 2 or 3. They also investigate the local antimagic chromatic number of some circulant graphs.

For a connected graph G with q edges a bijection $f : E \to \{1, 2, \ldots, q\}$ is called a *local antimagic labeling* if for any two adjacent vertices u and v, $w(u) \neq w(v)$, where $w(u) = \sum_{e \in E(u)} f(e)$, and $E(u)$ is the set of edges incident to u. In [286] Arumugam, Premalatha, Bača, and Semaničová-Feňovčíková proved several basic results on this new parameter and conjectured that any connected graph other than K_2 admits a local antimagic labeling. This conjecture was proved by Haslegrave [1382] using the probabilistic method, proves that the local antimagic conjecture is true. Lau [2095] proved that every graph admits a local antimagic total labeling. For any graph G, the graph $H = G \vee O_n$, $n \geq 1$, is defined by $V(H) = V(G) \cup \{v_i : 1 \leq i \leq n\}$ and $E(H) = E(G) \cup \{uv_i : u \in V(G)\}$. In [286, Theorem 2.16], it was claimed that for any G with order $m \geq 4$,

$$\chi_{la}(G) + 1 \leq \chi_{la}(G \vee O_2) \leq \begin{cases} \chi_{la}(G) + 1 & \text{if } m \text{ is even,} \\ \chi_{la}(G) + 2 & \text{if } m \text{ is odd.} \end{cases}$$

Lau and Shiu [2109] determined the local antimagic total chromatic number of graphs that are the amalgamation of complete graphs. Fachriza and Sugeng [1062] proved that barbell wheel graphs obtained by connecting the centers of two copies of wheels W_n by the path P_k ($k \geq 2$) have local antimagic total labelings and determined their local antimagic total chromatic numbers. In [4243] Yang, Bian, Yu, and Liu gave the local antimagic total chromatic numbers of some wheel-related graphs. for fans, flowers, and other wheel-related graphs. [4243] new

In [2108] Lau, Schaffer, and Shiu proved every graph is local antimagic total and provided sharp upper bound for $\chi_{lat}(G)$, the local antimagic total chromatic number of G. They also determined $\chi_{lat}(G)$, where G is a complete bipartite graph, a path, or the Cartesian product of two cycles, and $\chi_{lat}(G \vee K_1)$ and $\chi_{lat}(G)$ for a class of 2-regular graphs G. In [2110] Lau and Shiu determined the local antimagic total chromatic number the amalgamation of complete graphs. They also obtained the local antimagic (total) chromatic number of the disjoint union of complete graphs, and the join of K_1 and amalgamation of complete graphs under various conditions.

The number of distinct induced vertex labels under a local antimagic labeling f is denoted by $c(f)$, and is called the *color number* of f. The *local antimagic chromatic number* of G, denoted by $\chi_{la}(G)$, is $\min\{c(f) : f \text{ is a local antimagic labeling of } G\}$. In [1382], Haslegrave proved that the local antimagic chromatic number is well-defined for every connected graph other than K_2. Thus, for every connected graph $G \neq K_2$, $\chi_{la}(G) \geq \chi(G)$, the chromatic number of G. In [2113] Lau, Shiu, and Ng provided several sufficient conditions for $\chi_{la}(H) \leq \chi_{la}(G)$, where H is obtained from G with a certain edge deleted or added. The further determined the exact value of the local antimagic chromatic number of many cycle-related join graphs. In [2105] Lau, Premalatha, Arumugam, and Shiu determined the local antimagic chromatic number of $P_{2m} \vee O_n$, $P_{2m} \vee C_{2n-1}$, $P_{2m} \vee K_n$, $C_{2m} \vee O_{2n-1}$, $C_{2m} \vee C_{2n-1}$, $C_{2m} \vee K_{2n-1}$, $C_{2m-1} \vee K_{2n}$ and $C_{2n+1} \vee O_{2n}$ (O_n denotes the null graph of order n). In [2107] Lau, Shiu, and Ng determined the sharp lower bound of the local antimagic chromatic number of a graph with cut-vertices given by pendants. They gave antimagic chromatic numbers of one point unions of cycles, tadpole graphs, and caterpillar graphs. These results partially answer Problem 3.1 posed by Arumugam, Premalatha, Bača, and Semaničová-Feňovčíková in [286].

Let $G(V, E)$ be a simple graph and f be a bijection $f : V \cup E \to \{1, 2, \ldots, |V| + |E|\}$ where $f(V) = \{1, 2, \ldots, |V|\}$. For a vertex $x \in V$, define the weight of x, $w(x)$, as the sum of labels of all edges incident with x and the vertex label itself. Such an f is called a *super vertex local antimagic total* labeling (SLAT) if for every two adjacent vertices their weights are different. The *super vertex local antimagic total chromatic number* $\chi_{slat}(G)$ is the minimum number of colors taken over all colorings induced by super vertex local antimagic total labelings of G. Hadiputra, Sugeng, Silaban, Maryati, and Fronček classify all trees T that have $\chi_{slat}(T) = 2$, present a class of trees that have $\chi_{slat}(T) = 3$, and show that for any positive integer $n \geq 2$ there is a tree T with $\chi_{slat}(T) = n$.

In Table 13 we use the abbreviation (a, d)-**A** to mean that the graph has an (a, d)-antimagic labeling. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The table was prepared by Petr Kovář and Tereza Kovářová and updated by J. Gallian in 2008.

Table 13: **Summary of (a,d)-Antimagic Labelings**

Graph	*Labeling*	*Notes*
P_{2n}	not (a,d)-A	[678]
P_{2n+1}	iff $(n,1)$-A	[678]
C_{2n}	not (a,d)-A	[678]
C_{2n+1}	$(n+2,1)$-A	[678]
stars	not (a,d)-A	[678]
$C_3^{(k)}, C_4^{(k)}$	not (a,d)-A	[678]
$K_{3,3}$	not (a,d)-A	[678]
K_4	not (a,d)-A	[678]
Petersen graph	not (a,d)-A	[678]
W_n	not (a,d)-A	$n>3$ [2271]
antiprism on $2n$ vertices (see §6.2)	$(6n+3,2)$-A	$n\geq 3$, $n\not\equiv 2 \pmod 4$ [343]
	$(4n+4,4)$-A	$n\geq 3$, $n\not\equiv 2 \pmod 4$ [343]
	$(2n+5,6)$-A?	$n\geq 4$ [343]
	$(6n+3,2)$-A?	$n\geq 6$, $n\not\equiv 2 \pmod 4$ [343]
	$(4n+4,4)$-A?	$n\geq 6$, $n\not\equiv 2 \pmod 4$ [343]
Hershel graph (see [785])	not (a,d)-A	[674], [676]
parachutes $P_{g,p}$ (see §6.2)	(a,d)-A	for certain classes [674], [676]
prisms $C_n \times P_2$	$((7n+4)/2,1)$-A	$n\geq 3$, n even [675], [371]
	$((5n+5)/2,2)$-A	$n\geq 3$, n odd [675], [371]
	$((3n+6)/2,3)$-A	$n\geq 3$, n even [371]
	$((n+7)/2,4)$-A?	$n\geq 7$, [676], [371]
generalized Petersen graph $P(n,2)$	$((3n+6)/2,3)$-A	$n\geq 8,\ n\equiv 0 \pmod 4$ [372]

6.3 (a, d)-Antimagic Total Labelings

Bača, Bertault, MacDougall, Miller, Simanjuntak, and Slamin [361] introduced the notion of a (a, d)-vertex-antimagic total labeling in 2000. For a graph $G(V, E)$, an injective mapping f from $V \cup E$ to the set $\{1, 2, \ldots, |V| + |E|\}$ is a (a, d)-*vertex-antimagic total labeling* if the set $\{f(v) + \sum f(vu)\}$ where the sum is over all vertices u adjacent to v for all v in G is $\{a, a+d, a+2d, \ldots, a+(|V|-1)d\}$. In the case where the vertex labels are 1,2, $\ldots$, $|V|$, (a, d)-vertex-antimagic total labeling is called a *super* (a, d)*-vertex-antimagic total labeling.* Among their results are: every super-magic graph has an $(a, 1)$-vertex-antimagic total labeling; every (a, d)-antimagic graph $G(V, E)$ is $(a+|E|+1, d+1)$-vertex-antimagic total; and, for $d > 1$, every (a, d)-antimagic graph $G(V, E)$ is $(a + |V| + |E|, d - 1)$-vertex-antimagic total. They also show that paths and cycles have (a, d)-vertex-antimagic total labelings for a wide variety of a and d. In [362] Bača et al. use their results in [361] to obtain numerous (a, d)-vertex-antimagic total labelings for prisms, and generalized Petersen graphs (see §2.7 for the definition). (See also [378] and [3733] for more results on generalized Petersen graphs.)

Sugeng, Miller, Lin, and Bača [3733] prove: C_n has a super (a, d)-vertex-antimagic total labeling if and only if $d = 0$ or 2 and n is odd, or $d = 1$; P_n has a super (a, d)-vertex-antimagic total labeling if and only if $d = 2$ and $n \geq 3$ is odd, or $d = 3$ and $n \geq 3$; no even order tree has a super $(a, 1)$-vertex antimagic total labeling; no cycle with at least one tail and an even number of vertices has a super $(a, 1)$-vertex-antimagic labeling; and the star S_n, $n \geq 3$, has no super (a, d)-super antimagic labeling. As open problems they ask whether $K_{n,n}$ has a super (a, d)-vertex-antimagic total labeling and the generalized Petersen graph has a super (a, d)-vertex-antimagic total labeling for specific values a, d, and n. In [3068] Raheem proved that various subclasses of stars admit super (a, d)-edge antimagic total labelings for $d = 1, 2$, and 3. Lin, Miller, Simanjuntak, and Slamin [2271] have shown that for $n > 20$, W_n has no (a, d)-vertex-antimagic total labeling. Tezer and Cahit [3881] proved that neither P_n nor C_n has (a, d)-vertex-antimagic total labelings for $a \geq 3$ and $d \geq 6$. Kovář [2021] has shown that every $2r$-regular graph with n vertices has an $(s, 1)$-vertex antimagic total labeling for $s \in \{(rn + 1)(r + 1) + tn \mid t = 0, 1, \ldots, r\}$. Dafik, Slamin, Romdhani, and Arianti [889] studied the super (a, d)-antimagicness of generalized flower and disk brake graphs. Dafik, Slamin, Eka, and Sya'diyah [888] proved that triangular books and diamond ladder graphs admit a super (a, d)-edge-antimagic total labeling of graph for $d \in \{0, 1, 2\}$.

Several papers have been written about vertex-antimagic total labeling of graphs that are the disjoint union of suns. The sun graph S_n is $C_n \odot K_1$. Rahim and Sugeng [3075] proved that $S_{n_1} \cup S_{n_2} \cup \cdots \cup S_{n_t}$ is $(a, 0)$-vertex-antimagic total (or vertex magic total). Parestu, Silaban, and Sugeng [2774] and [2775] proved $S_{n_1} \cup S_{n_2} \cup \cdots \cup S_{n_t}$ is (a, d)-vertex-antimagic total for $d = 1, 2, 3, 4$, and 6 and particular values of a. In [3072] Rahim, Ali, Kashif, and Javaid provide (a, d)-vertex antimagic total labelings of disjoint unions of cycles, sun graphs, and disjoint unions of sun graphs. In [1037] Enomoto et al. proposed the conjecture that every tree is a super $(a, 0)$-edge-antimagic total graph. Javaid [1578] gave (a, d)-edge-antimagic total labelings for certain subclasses of subdivided

stars. Javaid [1579] gave a super (a,d)-edge-antimagic total labeling for the subdivided star $T(n,n,n+4,n+4,n_5,n_6,\ldots,n_r)$ for $d=0,1,2$, where $n_p=2^{p-4}(n+3)+1,\ 5\le p\le r$ and $n\ge 3$ is odd.

In [2701] Ngurah, Baskova, and Simanjuntak provide (a,d)-vertex-antimagic total labelings for the generalized Petersen graphs $P(n,m)$ for the cases: $n\ge 3,\ 1\le m\le \lfloor (n-1)/2\rfloor,\ (a,d)=(8n+3,2)$; odd $n\ge 5,\ m=2,\ (a,d)=((15n+5)/2,1)$; odd $n\ge 5,\ m=2,\ (a,d)=((21n+5)/2,1)$; odd $n\ge 7,\ m=3,\ (a,d)=((15n+5)/2,1)$; odd $n\ge 7,\ m=3,\ (a,d)=((21n+5)/2,1)$; odd $n\ge 9,\ m=4,\ (a,d)=((15n+5)/2,1)$; and $(a,d)=((21n+5)/2,1)$. They conjecture that for n odd and $1\le m\le \lfloor (m-1)/2\rfloor$, $P(n,m)$ has an $((21n+5)/2,1)$-vertex-antimagic labeling. In [3738] Sugeng and Silaban show: the disjoint union of any number of odd cycles of orders $n_1,n_2,\ldots,n_t$, each at least 5, has a super $(3(n_1+n_2+\cdots+n_t)+2,1)$-vertex-antimagic total labeling; for any odd positive integer t, the disjoint union of t copies of the generalized Petersen graph $P(n,1)$ has a super $(10t+2)n-\lfloor n/2\rfloor+2,1)$-vertex-antimagic total labeling; and for any odd positive integers t and n $(n\ge 3)$, the disjoint union of t copies of the generalized Petersen graph $P(n,2)$ has a super $(21tn+5)/2,1)$-vertex-antimagic total labeling. Ahmad, Ali, Bača, Kovar, and Semaničová-Feňovčíková, investigated the vertex-antimagicness of regular graphs and the existence of (super) (a,d)-vertex antimagic total labelings for regular graphs in general.

Ali, Bača, Lin, and Semaničová-Feňovčiková [185] investigated super-(a,d)-vertex antimagic total labelings of disjoint unions of regular graphs. Among their results are: if m and $(m-1)(r+1)/2$ are positive integers and G is an r-regular graph that admits a super-vertex magic total labeling, then mG has a super-$(a,2)$-vertex antimagic total labeling; if G has a 2-regular super-$(a,1)$-vertex antimagic total labeling, then mG has a super-$(m(a-2)+2,1),1)$-vertex antimagic total labeling; mC_n has a super-(a,d)-vertex antimagic total labeling if and only if either d is 0 or 2 and m and n are odd and at least 3 or $d=1$ and $n\ge 3$; and if G is an even regular Hamilton graph, then mG has a super-$(a,1)$-vertex antimagic total labeling for all positive integers m.

In [425] Bača, A. Semaničová-Feňovčíková, Wang, and Zhang investigate the existence of $(a,1)$-vertex-antimagic edge labelings for disconnected 3-regular graphs. As an extension of (a,d)-vertex-antimagic edge labeling they also introduce the concept of (a,d)-vertex-antimagic edge deficiency for measuring how close a graph is away from being an (a,d)-antimagic graph. In [285] Arumugam and Nalliah investigate the existence of a super (a,d)-edge-antimagic total labelings of disconnected graphs.

Ahmad, Ali, Bača, Kovář and Semaničová-Feňovčíková [104] provided a technique that allows one to construct several (a,r)-vertex-antimagic edge labelings for any $2r$-regular graph G of odd order provided the graph is Hamiltonian or has a 2-regular factor that has $(b,1)$-vertex-antimagic edge labeling. A similar technique allows them to construct a super (a,d)-vertex-antimagic total labeling for any $2r$-regular Hamiltonian graph of odd order with differences $d=1,2,\ldots,r$ and $d=2r+2$.

For $n\ge 2$ Dafik, Setiawani, and Azizah [883] define a *shackle* as a graph constructed from connected graphs $G_1,G_2,\ldots,G_n$, all isomorphic to G, such that G_s and G_t are disjoint when $|s-t|\ge 2$ and for every $i=1,2,\ldots,n-1,\ G_i$ and G_{i+1} share exactly one

common vertex v. In a *generalized shackle* a common subgraph is shared by each G_i and G_{i+1}. Dafik, Setiawani, and Azizah prove that the generalized shackle of a fan of order four and five admits a super (a,d)-edge antimagic total labeling for $d = 0, 1, 2$.

Sugeng and Bong [3726] show how to construct super (a,d)-vertex antimagic total labelings for the circulant graphs $C_n(1,2,3)$, for $d = 0,1,2,3,4,8$. Thirusangu, Nagar, and Rajeswari [3894] show that certain Cayley digraphs of dihedral groups have (a,d)-vertex-magic total labelings.

For a simple graph H we say that $G(V,E)$ admits an *H-covering* if every edge in $E(G)$ belongs to a subgraph of G that is isomorphic to H. Inayah, Salman, and Simanjuntak [1532] define an *(a,d)-H-antimagic total labeling* of G as a bijective function ξ from $V \cup E \to \{1,2,\ldots,|V|+|E|\}$ such that for all subgraphs H' isomorphic to H, the H-weights $w(H') = \sum_{v\in V(H')} \xi(v) + \sum_{e\in E(H')} \xi(e)$ constitute an arithmetic progression $a, a+d, a+2d, \ldots, a+(t-1)d$ where a and d are positive integers and t is the number of subgraphs of G isomorphic to H. Such a labeling ξ is called a *super (a,d)-H-antimagic total labeling*, if $\xi(V) = \{1,2,\ldots,|V|\}$. Inayah et al. study some basic properties of such labeling and give (a,d)-cycle-antimagic labelings of fans. Taimur, Numan, Mumtaz, and Semaničová-Feňovčíková [3858] proved that if a graph G is super cycle-antimagic then the subdivided graph of G also admits a super cycle-antimagic labeling and they showed that the subdivided wheel is super (a,d)-cycle-antimagic for wide range of values. Laurence and Kathiresan [2126] investigated super (a,d)-P_n-antimagic total labeling of stars. In [3042] Prihandini, Dafik, Agustin, Alfarisi, Adawiyah, and Santoso investigated the existence of super (a,d)-$P_2 \triangleright H$-antimagic total labelings of the disjoint union graph $C_n \triangle H$. Bača, Jeyanthi, Selvagopal, Muthu Raja, and Semaničová-Feňovčíková [328] proved the existence of super (a,d)-H-antimagic labelings of fan graphs and ladders for H isomorphic to a cycle.

In [3356] Semaničová-Feňovčíková, Bača, Lascsáková, Miller, and Ryan investigated the super (a,d)-C_n-antimagic total labelings of wheels and super (a,d)-P_n-antimagic total labelings of cycles and paths. Ovais, Umar, Bača, and Semaničová-Feňovčíková [2743] proved that fans admits a super (a,d)-C_k-antimagic labeling for $d = 1, 3, 2k-5, 2k-1, 3k-1, k-7, k+1, 3k-9$. They also prove that fans admits a super (a,d)-C_3-antimagic labeling for $d = 0,1,2,3,4,5,6,8$, and a super (a,d)-C_4-antimagic labeling for $d = 0,1,2,3,4,5,6,7,11$. They propose an open problem to find a super (a,d)-C_k-antimagic labeling of fans for $d \neq 1, 3, k-7, k+1, 2k-5, 2k-1, 3k-1, 3k-9$. Bača, Miller, Ryan, and Semaničová-Feňovčíková [409] study super (a,d)-H-antimagic labelings of a disjoint union of graphs for $d = |E(H)| - |V(H)|$.

For a vertex u of a graph G, $G_u[S_n]$ is the graph obtained by identifying u with the center of S_n. Then for any vertex w of S_n $G+e$, $e = uw$ is a subgraph of $G_u[S_n]$. Kathiresan and Laurence [1928] prove that the graph $G_u[S_n]$ admits a super-(a,d)-$(G+e)$-antimagic total labeling if and only if $d \in \{0,1,2,\ldots,|V(G)|+|E(G)|+2\}$. Moreover, they show that a caterpillar $S_{n_1,n_2,\ldots,n_k}$ has a super-$(a,4n^2)$-$S_{n,n}$-antimagic total labeling for $n_1 = n_2 = \cdots = n_k = n$. In [3102] Rajkumar, Nalliah, Uma Maheswari provided a partial solution to the problem of characterizing the super $(a,d) - G + e$-antimagic total labeling of the graph $G_u[S_n]$, where $n \geq 3$ and $4 \leq d \leq p+q+2$ posed in 2015 by

Kathiresan and Laurence in [1928].

Jeyanthi, Muthuraja, Semaničová-Feňovčíková, and Dharshikha proved [1715] proved that fans, triangular ladders, and middle graphs of cycles are super (a,d)-C_3-antimagic for some values of a and d. They also proved that ladder are super (a,d)-C_4-antimagic for $1 \leq d \leq 8$. Inayah, Simanjuntak, and Salman [1533] proved that there exists a super $(a,d)-H$-antimagic total labelings for shackles of a connected graph H. Nadzima and Martini [2633] determined (a,d)-H-antimagic total labeling for certain cases of $W_n \odot P_n$ with H as $C_3 \odot P_n$ and $W_n \odot C_n$ with H as $C_3 \odot C_n$.

A graph G is said to have an $(H_1, H_2, \ldots, H_k)$-*covering* if every edge in G belongs to at least one of the H_i's. Susilowati, Sania, and Estuningsih [3834] investigated such antimagic labelings for the ladders $P_n \times P_2$ with C_t-coverings for $t = 4,\ 6$, and 8 for some value of d.

Simanjuntak, Bertault, and Miller [3585] define an (a,d)-*edge-antimagic vertex labeling* for a graph $G(V,E)$ as an injective mapping f from V onto the set $\{1,2,\ldots,|V|\}$ such that the set $\{f(u)+f(v)|uv \in E\}$ is $\{a, a+d, a+2d, \ldots, a+(|E|-1)d\}$. (The equivalent notion of (a,d)-*indexable* labeling was defined by Hegde in 1989 in his Ph. D. thesis– see [1401].) Similarly, Simanjuntak et al. define an (a,d)-*edge-antimagic total labeling* for a graph $G(V,E)$ as an injective mapping f from $V \cup E$ onto the set $\{1,2,\ldots,|V|+|E|\}$ such that the set $\{f(v)+f(vu)+f(v)|uv \in E\}$ where v ranges over all of V is $\{a, a+d, a+2d, \ldots, a+(|V|-1)d\}$. Among their results are: C_{2n} has no (a,d)-edge-antimagic vertex labeling; C_{2n+1} has a $(n+2,1)$-edge-antimagic vertex labeling and a $(n+3,1)$-edge-antimagic vertex labeling; P_{2n} has a $(n+2,1)$-edge-antimagic vertex labeling; P_n has a $(3,2)$-edge-antimagic vertex labeling; C_n has $(2n+2,1)$- and $(3n+2,1)$-edge-antimagic total labelings; C_{2n} has $(4n+2,2)$- and $(4n+3,2)$-edge-antimagic total labelings; C_{2n+1} has $(3n+4,3)$- and $(3n+5,3)$-edge-antimagic total labelings; P_{2n+1} has $(3n+4,2)$-, $(3n+4,3)$-, $(2n+4,4)$-, $(5n+4,2)$-, $(3n+5,2)$-, and $(2n+6,4)$-edge-antimagic total labelings; P_{2n} has $(6n,1)$- and $(6n+2,2)$-edge-antimagic total labelings; and several parity conditions for (a,d)-edge-antimagic total labelings. They conjecture: C_{2n} has a $(2n+3,4)$- or a $(2n+4,4)$-edge-antimagic total labeling; C_{2n+1} has a $(n+4,5)$- or a $(n+5,5)$-edge-antimagic total labeling; paths have no (a,d)-edge-antimagic vertex labelings with $d > 2$; and cycles have no (a,d)-antimagic total labelings with $d > 5$. The first and last of these conjectures were proved by Zhenbin in [4333] and the last two were verified by Bača, Lin, Miller, and Simanjuntak [392] who proved that a graph with v vertices and e edges that has an (a,d)-edge-antimagic vertex labeling must satisfy $d(e-1) \leq 2v-1-a \leq 2v-4$. As a consequence, they obtain: for every path there is no (a,d)-edge-antimagic vertex labeling with $d > 2$; for every cycle there is no (a,d)-edge-antimagic vertex labeling with $d > 1$; for K_n $(n > 1)$ there is no (a,d)-edge-antimagic vertex labeling (the cases for $n = 2$ and $n = 3$ are handled individually); $K_{n,n}$ $(n > 3)$ has no (a,d)-edge-antimagic vertex labeling; for every wheel there is no (a,d)-edge-antimagic vertex labeling; for every generalized Petersen graph there is no (a,d)-edge-antimagic vertex labeling with $d > 1$. They also study the relationship between graphs with (a,d)-edge-antimagic labelings and magic and antimagic labelings. They conjecture that every tree has an $(a,1)$-edge-antimagic total labeling.

Bača and Barrientos [347] prove that if a tree T has an α-labeling and $\{A,B\}$ is the

bipartition of the vertices of T, then T also admits an $(a,1)$-edge-antimagic vertex labeling and it admits a $(3,2)$-edge-antimagic vertex labeling if and only if $||A|-|B|| \leq 1$.

In [392] Bača, Lin, Miller, and Simanjuntak prove: if P_n has an (a,d)-edge-antimagic total labeling, then $d \leq 6$; P_n has $(2n+2,1)$-, $(3n,1)$-, $(n+4,3)$-, and $(2n+2,3)$-edge-antimagic total labelings; P_{2n+1} has $(3n+4,2)$-,$(5n+4,3)$-, $(2n+4,4)$-, and $(2n+6,4)$-edge-antimagic total labelings; and P_{2n} has $(3n+3,2)$- and $(5n+1,2)$-edge-antimagic total labelings. Ngurah [2696] proved P_{2n+1} has $(4n+4,1)$-, $(6n+5,3)$-,$(4n+4,2)$-,$(4n+5,2)$-edge-antimagic total labelings and C_{2n+1} has $(4n+4,2)$- and $(4n+5,2)$-edge-antimagic total labelings. Silaban and Sugeng [3583] prove: P_n has $(n+4,4)$- and $(6,6)$-edge-antimagic total labelings; if $C_m \odot \overline{K_n}$ has an (a,d)-edge-antimagic total labeling, then $d \leq 5$; $C_m \odot \overline{K_n}$ has (a,d)-edge-antimagic total labelings for $m > 3, n > 1$ and $d = 2$ or 4; and $C_m \odot \overline{K_n}$ has no (a,d)-edge-antimagic total labelings for m and d and $n \equiv 1 \bmod 4$. They conjecture that P_n $(n \geq 3)$ has $(a,5)$-edge-antimagic total labelings. In [3741] Sugeng and Xie use adjacency methods to construct super edge magic graphs from (a,d)-edge-antimagic vertex graphs. Pushpam and Saibulla [3059] determined super (a,d)-edge antimagic total labelings for graphs derived from copies of generalized ladders, fans, generalized prisms and web graphs. Liu, Aslam, Javaid, and Raheen [2290] compute bounds of the minimum and maximum edge-weights for super (a,d)-edge-antimagic labelings on a generalized class of subdivided caterpillars. They also investigate the existence of super (a,d)-edge-antimagic total labeling for the validation of the obtained bounds.

In [429] Bača and Youssef used parity arguments to find a large number of conditions on p, q and d for which a graph with p vertices and q edges cannot have an (a,d)-edge-antimagic total labeling or vertex-antimagic total labeling. Bača and Youssef [429] made the following connection between (a,d)-edge-antimagic vertex labelings and sequential labelings: if G is a connected graph other than a tree that has an (a,d)-edge-antimagic vertex labeling, then $G + K_1$ has a sequential labeling.

In [3710] Sudarsana, Ismaimuza, Baskoro, and Assiyatun prove: for every $n \geq 2$, $P_n \cup P_{n+1}$ has a $(6n+1,1)$- and a $(4n+3,3)$-edge-antimagic total labeling, for every odd $n \geq 3$, $P_n \cup P_{n+1}$ has a $(6n,1)$- and a $(5n+1,2)$-edge-antimagic total labeling, for every $n \geq 2$, $nP_2 \cup P_n$ has a $(7n,1)$- and a $(6n+1,2)$-edge-antimagic total labeling. In [3707] the same authors show that $P_n \cup P_{n+1}$, $nP_2 \cup P_n$ $(n \geq 2)$, and $nP_2 \cup P_{n+2}$ are super edge-magic total. They also show that under certain conditions one can construct new super edge-magic total graphs from existing ones by joining a particular vertex of the existing super edge-magic total graph to every vertex in a path or every vertex of a star and by joining one extra vertex to some vertices of the existing graph. Baskoro, Sudarsana, and Cholily [593] also provide algorithms for constructing new super edge-magic total graphs from existing ones by adding pendent vertices to the existing graph. A corollary to one of their results is that the graph obtained by attaching a fixed number of pendent edges to each vertex of a path of even length is super edge-magic. Baskoro and Cholily [591] show that the graphs obtained by attaching any numbers of pendent edges to a single vertex or a fix number of pendent edges to every vertex of the following graphs are super edge-magic total graphs: odd cycles, the generalized Petersen graphs $P(n,2)$ (n odd and at least 5), and $C_n \times P_m$ (n odd, $m \geq 2$).

Arumugam and Nalliah [284] proved: the friendship graph $C_3^{(n)}$ with $n \equiv 0, 8 \pmod{12}$ has no super $(a, 2)$-edge-antimagic total labeling; $C_n^{(n)}$ with $n \equiv 2 \pmod 4$ has no super $(a, 2)$-edge-antimagic total labeling; and the generalized friendship graph $F_{2,p}$ consisting of 2 cycles of various lengths, having a common vertex, and having order p where $p \geq 5$, has a super $(2p + 2, 1)$-edge-antimagic total labeling if and only if p is odd.

An (a, d)-edge-antimagic total labeling of $G(V, E)$ is called a *super* (a, d)-*edge-antimagic total* if the vertex labels are $\{1, 2, \ldots, |V(G)|\}$ and the edge labels are $\{|V(G)| + 1, |V(G)| + 2, \ldots, |V(G)| + |E(G)|\}$. Bača, Baskoro, Simanjuntak, and Sugeng [360] prove the following: C_n has a super (a, d)-edge-antimagic total labeling if and only if either d is 0 or 2 and n is odd, or $d = 1$; for odd $n \geq 3$ and $m = 1$ or 2, the generalized Petersen graph $P(n, m)$ has a super $(11n + 3)/2, 0)$-edge-antimagic total labeling and a super $((5n + 5)/2, 2)$-edge-antimagic total labeling; for odd $n \geq 3,\ P(n, (n - 1)/2)$ has a super $((11n + 3)/2, 0)$-edge-antimagic total labeling and a super $((5n + 5)/2, 2)$-edge-antimagic total labeling. They also prove: if $P(n, m), n \geq 3,\ 1 \leq m \leq \lfloor (n - 1)/2 \rfloor$ is super (a, d)-edge-antimagic total, then $(a, d) = (4n + 2, 1)$ if n is even, and either $(a, d) = ((11n + 3)/2, 0)$, or $(a, d) = (4n + 2, 1)$, or $(a, d) = ((5n + 5)/2, 2)$, if n is odd; and for odd $n \geq 3$ and $m = 1, 2$, or $(n - 1)/2,\ P(n, m)$ has an $(a, 0)$-edge-antimagic total labeling and an $(a, 2)$-edge-antimagic total labeling. (In a personal communication MacDougall argues that "edge-magic" is a better term than "$(a, 0)$-edge-antimagic" for while the latter is technically correct, "antimagic" suggests different weights whereas "magic" emphasizes equal weights and that the edge-magic case is much more important, interesting, and fundamental rather than being just one subcase of equal value to all the others.) They conjecture that for odd $n \geq 9$ and $3 \leq m \leq (n-3)/2,\ P(n, m)$ has a $(a, 0)$-edge-antimagic total labeling and an $(a, 2)$-edge-antimagic total labeling. Ngurah and Baskoro [2700] have shown that for odd $n \geq 3,\ P(n, 1)$ and $P(n, 2)$ have $((5n + 5)/2, 2)$-edge-antimagic total labelings and when $n \geq 3$ and $1 \leq m < n/2,\ P(n, m)$ has a super $(4n + 2, 1)$-edge-antimagic total labeling. In [2701] Ngurah, Baskova, and Simanjuntak provide (a, d)-edge-antimagic total labelings for the generalized Petersen graphs $P(n, m)$ for the cases $m = 1$ or 2, odd $n \geq 3$, and $(a, d) = ((9n + 5)/2, 2)$.

In [3708] Sudarsana, Baskoro, Uttunggadewa, and Ismaimuza show how to construct new larger super (a, d)-edge-antimagic-total graphs from existing smaller ones.

In [2702] Ngurah, Baskoro, and Simanjuntak prove that mC_n $(n \geq 3)$ has an (a, d)-edge-antimagic total in the following cases: $(a, d) = (5mn/2 + 2, 1)$ where m is even; $(a, d) = (2mn+2, 2)$; $(a, d) = ((3mn+5)/2, 3)$ for m and n odd; and $(a, d) = ((mn+3), 4)$ for m and n odd; and mC_n has a super $(2mn + 2, 1)$-edge-antimagic total labeling.

Bača and Barrientos [348] have shown that mK_n has a super (a, d)-edge-antimagic total labeling if and only if $(i)\ d \in \{0, 2\},\ n \in \{2, 3\}$ and $m \geq 3$ is odd, or $(ii)\ d = 1,\ n \geq 2$ and $m \geq 2$, or $(iii)\ d \in \{3, 5\},\ n = 2$ and $m \geq 2$, or $(iv)\ d = 4,\ n = 2$, and $m \geq 3$ is odd. In [347] Bača and Barrientos proved the following: if a graph with q edges and $q+1$ vertices has an α-labeling, than it has an $(a, 1)$-edge-antimagic vertex labeling; a tree has a $(3, 2)$-edge-antimagic vertex labeling if and only if it has an α-labeling and the number of vertices in its two partite sets differ by at most 1; if a tree with at least two vertices has a super (a, d)-edge-antimagic total labeling, then d is at most 3; if a graph has an

$(a,1)$-edge-antimagic vertex labeling, then it also has a super $(a_1,0)$-edge-antimagic total labeling and a super $(a_2,2)$-edge-antimagic total labeling.

Bača and Youssef [429] proved the following: if G is a connected (a,d)-edge-antimagic vertex graph that is not a tree, then $G+K_1$ is sequential; mC_n has an (a,d)-edge-antimagic vertex labeling if and only if m and n are odd and $d=1$; an odd degree (p,q)-graph G cannot have a (a,d)-edge-antimagic total labeling if $p \equiv 2 \pmod 4$ and $q \equiv 0 \pmod 4$, or $p \equiv 0 \pmod 4$, $q \equiv 2 \pmod 4$, and d is even; a (p,q)-graph G cannot have a super (a,d)-edge-antimagic total labeling if G has odd degree, $p \equiv 2 \pmod 4$, q is even, and d is odd, or G has even degree, $q \equiv 2 \pmod 4$, and d is even; C_n has a $(2n+2,3)$- and an $(n+4,3)$-edge-antimagic total labeling; a (p,q)-graph is not super (a,d)-vertex-antimagic total if: $p \equiv 2 \pmod 4$ and d is even; $p \equiv 0 \pmod 4$, $q \equiv 2 \pmod 4$, and d is odd; $p \equiv 0 \pmod 8$ and $q \equiv 2 \pmod 4$.

In [3710] Sudarsana, Ismaimuza, Baskoro, and Assiyatun prove: for every $n \geq 2$, $P_n \cup P_{n+1}$ has super $(n+4,1)$- and $(2n+6,3)$-edge antimagic total labelings; for every odd $n \geq 3$, $P_n \cup P_{n+1}$ has super $(4n+5,1)$-,$(3n+6,2)$-, $(4n+3,1)$- and $(3n+4,2)$-edge antimagic total labelings; for every $n \geq 2$, $nP_2 \cup P_n$ has super $(6n+2,1)$- and $(5n+3,2)$-edge antimagic total labelings; and for every $n \geq 1$, $nP_2 \cup P_{n+2}$ has super $(6n+6,1)$- and $(5n+6,2)$-edge antimagic total labelings. They pose a number of open problems about constructing (a,d)-edge antimagic labelings and super (a,d)-edge antimagic labelings for the graphs $P_n \cup P_{n+1}$, $nP_2 \cup P_n$, and $nP_2 \cup P_{n+2}$ for specific values of d.

Dafik, Miller, Ryan, and Bača [885] investigated the super edge-antimagicness of the disconnected graph mC_n and mP_n. For the first case they prove that mC_n, $m \geq 2$, has a super (a,d)-edge-antimagic total labeling if and only if either d is 0 or 2 and m and n are odd and at least 3, or $d=1$, $m \geq 2$, and $n \geq 3$. For the case of the disjoint union of paths they determine all feasible values for m,n and d for mP_n to have a super (a,d)-edge-antimagic total labeling except when m is even and at least 2, $n \geq 2$, and d is 0 or 2. In [887] Dafik, Miller, Ryan, and Bača obtain a number of results about super edge-antimagicness of the disjoint union of two stars and state three open problems. Nalliah and Arumugam [2647] proved that $K_{1,6} \cup K_{1,5}$ does not have such a labeling and prove that some special cases of $K_{1,n+1} \cup K_{1,n}$ do have them.

Sudarsana, Hendra, Adiwijaya, and Setyawan [3709] show that the t-joint copies of wheel W_n have a super edge antimagic $((2n+2)t+2,1)$-total labeling for $n \geq 4$ and $t \geq 2$.

In [386] Bača, Lascsáková, and Semaničová investigated the connection between graphs with α-labelings and graphs with super (a,d)-edge-antimagic total labelings. Among their results are: If G is a graph with n vertices and $n-1$ edges ($n \geq 3$) and G has an α-labeling, then mG is super (a,d)-edge-antimagic total if either d is 0 or 2 and m is odd, or $d=1$ and n is even; if G has an α-labeling and has n vertices and $n-1$ edges with vertex bipartition sets V_1 and V_2 where $|V_1|$ and $|V_2|$ differ by at most 1, then mG is super (a,d)–edge-antimagic total for $d=1$ and $d=3$. In the same paper Bača et al. prove: caterpillars with odd order at least 3 have super $(a,1)$-edge-antimagic total labelings; if G is a caterpillar of odd order at least 3 and G has a super $(a,1)$-edge-antimagic total labeling, then mG has a super $(b,1)$-edge-antimagic total labeling for some b that is a function of a and m.

In [884] Dafik, Miller, Ryan, and Bača investigated the existence of antimagic labelings of disjoint unions of s-partite graphs. They proved: if $s \equiv 0$ or $1 \pmod 4$, $s \geq 4, m \geq 2, n \geq 1$ or mn is even , $m \geq 2,\ n \geq 1,\ s \geq 4$, then the complete s-partite graph $mK_{n,n,\ldots,n}$ has no super $(a,0)$-edge-antimagic total labeling; if $m \geq 2$ and $n \geq 1$, then $mK_{n,n,n,n}$ has no super $(a,2)$-antimagic total labeling; and for $m \geq 2$ and $n \geq 1$, $mK_{n,n,n,n}$ has an $(8mn+2,1)$-edge-antimagic total labeling. They conjecture that for $m \geq 2,\ n \geq 1$ and $s \geq 5$, the complete s-partite graph $mK_{n,n,\ldots,n}$ has a super $(a,1)$-antimagic total labeling.

In [413] Bača, Muntaner-Batle, Semaničová-Feňovčiková, and Shafiq investigate super (a,d)-edge-antimagic total labelings of disconnected graphs. Among their results are: If G is a (super) $(a,2)$-edge-antimagic total labeling and m is odd, then mG has a (super) $(a',2)$-edge-antimagic-total labeling where $a' = m(a-3) + (m+1)/2 + 2$; and if d a positive even integer and k a positive odd integer, G is a graph with all of its vertices having odd degree, and the order and size of G have opposite parity, then $2kG$ has no (a,d)-edge-antimagic total labeling. Bača and Brankovic [363] have obtained a number of results about the existence of super (a,d)-edge-antimagic totaling of disjoint unions of the form $mK_{n,n}$. In [367] Bača, Dafik, Miller, and Ryan provide (a,d)-edge-antimagic vertex labelings and super (a,d)-edge-antimagic total labelings for a variety of disjoint unions of caterpillars. Bača and Youssef [429] proved that mC_n has an (a,d)-edge-antimagic vertex labeling if and only if m and n are odd and $d = 1$. Bača, Dafik, Miller, and Ryan [368] constructed super (a,d)-edge-antimagic total labeling for graphs of the form $m(C_n \odot \overline{K}_s)$ and $mP_n \cup kC_n$ while Dafik, Miller, Ryan, and Bača [886] do the same for graphs of the form $mK_{n,n,n}$ and $K_{1,m} \cup 2sK_{1,n}$. Both papers provide a number of open problems. In [397] Bača, Lin, and Muntaner-Batle provide super (a,d)-edge-antimagic total labeling of forests in which every component is a specific kind of tree. In [384] Bača, Kovář, Semaničová-Feňovčiková, and Shafiq prove that every even regular graph and every odd regular graph with a 1-factor are super $(a,1)$-edge-antimagic total and provide some constructions of non-regular super $(a,1)$-edge-antimagic total graphs. Bača, Lin, and Semaničová-Feňovčiková [399] show: the disjoint union of m graphs with super $(a,1)$-edge antimagic total labelings have super $(m(a-2)+2,1)$-edge antimagic total labelings; the disjoint union of m graphs with super $(a,3)$-edge antimagic total labelings have super $(m(a-3)+3,3)$-edge antimagic total labelings; if G has a $(a,1)$-edge antimagic total labelings then mG has an $(b,1)$-edge antimagic total labeling for some b; and if G has a $(a,3)$-edge antimagic total labelings then mG has an $(b,3)$-edge antimagic total labeling for some b.

Bača, Miller, Ryan, and Semaničová-Feňovčíková [409] prove that if G admits a (super) (a,d)-H-antimagic labeling, where $d = |E(H)| - |V(H)|$, then mG admits a (super) (b,d)-H-antimagic labelling. By considering special H-coverings of a given H-antimagic graph G they derive many corollaries. In [3357] Semaničová-Feňovčíková, Bača, and Lascsáková provide two constructions of (super) H-antimagic graphs obtained from smaller (super) H'-antimagic graphs. Dafik, Slamin, Tana, Semaničová-Feňovčíková, and Bača [890] show a connection between a constructions of H-antimagic labelings of graph and edge-antimagic total labelings and describe how to obtain the H-antimagic graph using smaller edge-antimagic graph. Bača, Semaničová-Feňovčikovă, Umar, and Welyyanti [424] gave

sufficient conditions for $G_1 \times G_2$ to admit an H-supermagic or a super (a,d)-H-antimagic labeling but provide no examples of graphs that satisfy the given conditions.

For $t \geq 2$ and $n \geq 4$ the *Harary graph*, C_p^t, is the graph obtained by joining every two vertices of C_p that are at distance t in C_p. In [3072] Rahim, Ali, Kashif, and Javaid provide super (a,d)-edge antimagic total labelings for disjoint unions of Harary graphs and disjoint unions of cycles. In [1473] Hussain, Ali, Rahim, and Baskoro construct various (a,d)-vertex-antimagic labelings for Harary graphs and disjoint unions of identical Harary graphs. For p odd and at least 5, Balbuena, Barker, Das, Lin, Miller, Ryan, Slamin, Sugeng, and Tkac [462] give a super $((17p+5)/2)$-vertex-antimagic total labeling of C_p^t. MacDougall and Wallis [2381] have proved the following: $C_{4m+3}^t,\ m \geq 1$, has a super $(a,0)$-edge-antimagic total labeling for all possible values of t with $a = 10m+9$ or $10m+10$; $C_{4m+1}^t,\ m \geq 3$, has a super $(a,0)$-edge-antimagic total labeling for all possible values except $t = 5, 9, 4m-4$, and $4m-8$ with $a = 10m+4$ and $10m+5$; $C_{4m+1}^t,\ m \geq 1$, has a super $(10m+4,0)$-edge-antimagic total labeling for all $t \equiv 1 \pmod 4$ except $4m-3$; $C_{4m}^t,\ m > 1$, has a super $(10m+2,0)$-edge-antimagic total labeling for all $t \equiv 2 \pmod 4$; $C_{4m+2}^t,\ m > 1$, has a super $(10m+7,0)$-edge-antimagic total labeling for all odd t other than 5 and for $t = 2$ or 6. In [1474] Hussain, Baskoro, and Ali prove the following: for any $p \geq 4$ and for any $t \geq 2$, C_p^t admits a super $(2p+2,1)$-edge-antimagic total labeling; for $n \geq 4$, $k \geq 2$ and $t \geq 2$, kC_n^t admits a super $(2nk+2,1)$-edge-antimagic total labeling; and for $p \geq 5$ and $t \geq 2$, C_p^t admits a super $(8p+3,1)$-vertex-antimagic total labeling, provided if $p \neq 2t$.

Bača and Murugan [419] have proved: if $C_n^t,\ n \geq 4, 2 \leq t \leq n-2$, is super (a,d)-edge-antimagic total, then $d = 0, 1$, or 2; for $n = 2k+1 \geq 5,\ C_n^t$ has a super $(a,0)$-edge-antimagic total labeling for all possible values of t with $a = 5k+4$ or $5k+5$; for $n = 2k+1 \geq 5,\ C_n^t$ has a super $(a,2)$-edge-antimagic total labeling for all possible values of t with $a = 3k+3$ or $3k+4$; for $n \equiv 0 \pmod 4$, C_n^t has a super $(5n/2+2,0)$-edge-antimagic total labeling and a super $(3n/2+2,0)$-edge-antimagic total labeling for all $t \equiv 2 \pmod 4$; for $n = 10$ and $n \equiv 2 \pmod 4$, $n \geq 18,\ C_n^t$ has a

super $(5n/2+2,0)$-edge-antimagic total labeling and a super $(3n/2+2,0)$-edge-antimagic total labeling for all $t \equiv 3 \pmod 4$ and for $t = 2$ and 6; for odd $n \geq 5,\ C_n^t$ has a super $(2n+2,1)$-edge-antimagic total labeling for all possible values of t; for even $n \geq 6,\ C_n^t$ has a super $(2n+2,1)$-edge-antimagic total labeling for all odd $t \geq 3$; and for even $n \equiv 0 \pmod 4$, $n \geq 4,\ C_n^t$ has a super $(2n+2,1)$-edge-antimagic total labeling for all $t \equiv 2 \pmod 4$. They conjecture that there is a super $(2n+2,1)$-edge-antimagic total labeling of C_n^t for $n \equiv 0 \pmod 4$ and for $t \equiv 0 \pmod 4$ and for $n \equiv 2 \pmod 4$ and for t even.

In [393] Bača, Lin, Miller, and Youssef prove: if the friendship $C_3^{(n)}$ is super (a,d)-antimagic total, then $d < 3$; $C_3^{(n)}$ has an $(a,1)$-edge antimagic vertex labeling if and only if $n = 1, 3, 4, 5$, and 7; $C_3^{(n)}$ has a super (a,d)-edge-antimagic total labelings for $d = 0$ and 2; $C_3^{(n)}$ has a super $(a,1)$-edge-antimagic total labeling; if a fan F_n $(n \geq 2)$ has a super (a,d)-edge-antimagic total labeling, then $d < 3$; F_n has a super (a,d)-edge-antimagic total labeling if $2 \leq n \leq 6$ and $d = 0, 1$ or 2; the wheel W_n has a super (a,d)-edge-antimagic total labeling if and only if $d = 1$ and $n \not\equiv 1 \pmod 4$; $K_n,\ n \geq 3$, has a super (a,d)-edge-

antimagic total labeling if and only if either $d = 0$ and $n = 3$, or $d = 1$ and $n \geq 3$, or $d = 2$ and $n = 3$; and $K_{n,n}$ has a super (a, d)-edge antimagic total labeling if and only if $d = 1$ and $n \geq 2$.

Bača, Lin, and Muntaner-Batle [394] have shown that if a tree with at least two vertices has a super (a, d)-edge-antimagic total labeling, then d is at most three and $P_n,\ n \geq 2$, has a super (a, d)-edge-antimagic total labeling if and only if $d = 0, 1, 2$, or 3. They also characterize certain path-like graphs in a grid that have super(a, d)-edge-antimagic total labelings.

In [3732] Sugeng, Miller, and Bača prove that the ladder, $P_n \times P_2$, is super (a, d)-edge-antimagic total if n is odd and $d = 0, 1$, or 2 and $P_n \times P_2$ is super $(a, 1)$-antimagic total if n is even. They conjecture that $P_n \times P_2$ is super $(a, 0)$- and $(a, 2)$-edge-antimagic when n is even. Sugeng, Miller, and Bača [3732] prove that $C_m \times P_2$ has a super (a, d)-edge-antimagic total labeling if and only if either $d = 0, 1$ or 2 and m is odd and at least 3, or $d = 1$ and m is even and at least 4. They conjecture that if m is even, $m \geq 4,\ n \geq 3$, and $d = 0$ or 2, then $C_m \times P_n$ has a super (a, d)-edge-antimagic total labeling. In [2140] M.-J. Lee studied super $(a, 1)$-edge-antimagic properties of $m(P_4 \times P_n)$ for $m, n \geq 1$ and $m(C_n \odot \overline{K_t})$ for n even and $m, t \geq 1$. He also proved that for $n \geq 2$ the graph $P_4 \times P_n$ has a super $(8n + 2, 1)$-edge antimagic total labeling.

In [207] and [208] Azizu, Yulianti, Sy, Narwen proved that $(C_m \times P_2) \odot \overline{K_n}$ (branched-prism) for odd $m \geq 3$ and $n \geq 1$ admits an edge magic and a super (a, d)-edge antimagic total labelings for odd $m \geq 3$ and $n \geq 1$.

Sugeng, Miller, and Bača [3732] define a variation of a ladder, $\mathbb{L}_n$, as the graph obtained from $P_n \times P_2$ by joining each vertex u_i of one path to the vertex v_{i+1} of the other path for $i = 1, 2, \ldots, n-1$. They prove $\mathbb{L}_n,\ n \geq 2$, has a super (a, d)-edge-antimagic total labeling if and only if $d = 0, 1$, or 2.

In [351] Bača, Bashir, and Semaničová showed that for $n \geq 4$ and $d = 0, 1, 2, 3, 4, 5$, and 6 the antiprism A_n has a super d-antimagic labeling of type $(1, 1, 1)$. The *generalized antiprism* A_m^n is obtained from $C_m \times P_n$ by inserting the edges $\{v_{i,j+1}, v_{i+1,j}\}$ for $1 \leq i \leq m$ and $1 \leq j \leq n - 1$ where the subscripts are taken modulo m. Sugeng et al. prove that $A_m^n,\ m \geq 3,\ n \geq 2$, is super (a, d)-edge-antimagic total if and only if $d = 1$.

A *toroidal polyhex* (toroidal fullerene) is a cubic bipartite graph embedded on the torus such that each face is a hexagon. Note that the torus is a closed surface that can carry a toroidal polyhex such that all its vertices have degree 3 and all faces of the embedding are hexagons. Bača and Shabbir [426] proved the toroidal polyhex $\mathbb{H}_m^n$ with mn hexagons, $m, n \geq 2$, admits a super (a, d)-edge-antimagic total labeling if and only if $d = 1$ and $a = 4mn + 2$.

Bača, Miller, Phanalasy, and A. Semaničová-Feňovčíková [405] investigated the existence of (super) 1-antimagic labelings of type $(1, 1, 1)$ for disjoint union of plane graphs. They prove that if a plane graph $G(V, E, F)$ has a (super) 1-antimagic labeling h of type $(1, 1, 1)$ such that $h(z_{ext}) = |V(G)| + |E(G)| + |F(G)|$ where z_{ext} denotes the unique external face then, for every positive integer m, the graph mG also admits a (super) 1-antimagic labeling of type $(1, 1, 1)$; and if a plane graph $G(V, E, F)$ has 4-sided inner faces and h is a (super) d-antimagic labeling of type $(1, 1, 1)$ of G such that

$h(z_{ext}) = |V(G)| + |E(G)| + |F(G)|$ where $d = 1, 3, 5, 7, 9$ then, for every positive integer m, the graph mG also admits a (super) d-antimagic labeling of type $(1, 1, 1)$. They also give a similar result about plane graphs with inner faces that are 3-sided.

Sugeng, Miller, Slamin, and Bača [3734] proved: the star S_n has a super (a, d)-antimagic total labeling if and only if either $d = 0, 1$ or 2, or $d = 3$ and $n = 1$ or 2; if a nontrivial caterpillar has a super (a, d)-edge-antimagic total labeling, then $d \leq 3$; all caterpillars have super $(a, 0)$-, $(a, 1)$- and $(a, 2)$-edge-antimagic total labelings; all caterpillars have a super $(a, 1)$-edge-antimagic total labeling; if m and n differ by at least 2 the double star $S_{m,n}$ (that is, the graph obtained by joining the centers of $K_{1,m}$ and $K_{1,n}$ with an edge) has no $(a, 3)$-edge-antimagic total labeling.

Sugeng and Miller [3730] show how to manipulate adjacency matrices of graphs with (a, d)-edge-antimagic vertex labelings and super (a, d)-edge-antimagic total labelings to obtain new (a, d)-edge-antimagic vertex labelings and super (a, d)-edge-antimagic total labelings. Among their results are: every graph can be embedded in a connected (a, d)-edge-antimagic vertex graph; every (a, d)-edge-antimagic vertex graph has a proper (a, d)-edge-antimagic vertex subgraph; if a graph has a $(a, 1)$-edge-antimagic vertex labeling and an odd number of edges, then it has a super $(a, 1)$-edge-antimagic total labeling; every super edge magic total graph has an $(a, 1)$-edge-antimagic vertex labeling; and every graph can be embedded in a connected super (a, d)-edge-antimagic total graph.

Rahmawati, Sugeng, Silaban, Miller, and Bača [3080] construct new larger (a, d)-edge-antimagic vertex graphs from an existing (a, d)-edge-antimagic vertex graph using adjacency matrix for difference $d = 1, 2$. The results are extended for super (a, d)-edge-antimagic total graphs with differences $d = 0, 1, 2, 3$.

Ajitha, Arumugan, and Germina [154] show that $(p, p-1)$ graphs with α-labelings (see §3.1) and partite sets with sizes that differ by at most 1 have super (a, d)-edge antimagic total labelings for $d = 0, 1, 2$ and 3. They also show how to generate large classes of trees with super (a, d)-edge-antimagic total labelings from smaller graceful trees.

Bača, Lin, Miller, and Ryan [391] define a *Möbius grid*, M_n^m, as the graph with vertex set $\{x_{i,j} |\ i = 1, 2, \ldots, m+1, j = 1, 2, \ldots, n\}$ and edge set $\{x_{i,j}x_{i,j+1} |\ i = 1, 2, \ldots, m+1, j = 1, 2, \ldots, n-1\} \cup \{x_{i,j}x_{i+1,j} |\ i = 1, 2, \ldots, m,\ j = 1, 2, \ldots, n\} \cup \{x_{i,n}x_{m+2-i,1} |\ i = 1, 2, \ldots, m+1\}$. They prove that for $n \geq 2$ and $m \geq 4$, M_n^m has no d-antimagic vertex labeling with $d \geq 5$ and no d-antimagic-edge labeling with $d \geq 9$.

Ali, Bača, and Bashir, [182] investigated super (a, d)-vertex-antimagic total labelings of the disjoint unions of paths. They prove: mP_2 has a super (a, d)-vertex-antimagic total labeling if and only if m is odd and $d = 1$; mP_3, $m > 1$, has no super $(a, 3)$-vertex-antimagic total labeling; mP_3 has a super $(a, 2)$-vertex-antimagic total labeling for $m \equiv 1 \pmod 6$; and mP_4 has a super $(a, 2)$-vertex-antimagic total labeling for $m \equiv 3 \pmod 4$.

Lee, Tsai, and Lin [2143] denote the subdivision of a star S_n obtained by inserting m vertices into every edge of the star S_n by S_m^n. They proved that for $n \geq 3$, the graph kS_m^n is super (a, d)-edge antimagic total for certain values. In [1482] Ichishima, López, Muntaner-Batle and Rius-Font proved that if G is tripartite and has a (super) (a, d)-edge antimagic total labeling, then nG $(n \geq 3)$ has a (super) (a, d)-edge antimagic total labeling for $d = 1$ and for $d = 0, 2$ when n is odd.

Let $p, t_1, t_2, \ldots, t_k$ be integers such that $1 \leq t_1 < t_2 < \cdots < t_k < p$. A *Toeplitz* graph, denoted by $T_p\langle t_1 \ldots, t_k\rangle$, is a graph with vertex set $\{v_1, v_2, \ldots, v_p\}$ and edge set $\{v_i v_j : |i-j| \in \{t_1, t_2, \ldots, t_k\}\}$. Bača, Bashir, Nadeem, and Shabbir [350] give an upper bound on the difference d when a Toeplitz graph $T_p\langle t_1, t_2, \ldots t_k\rangle$ is super (a,d)-edge-antimagic total. They also construct a super $(a,1)$-edge-antimagic total labeling for an arbitrary Toeplitz graph without isolated vertices and prove that the Toeplitz graph $T_p\langle t_1\rangle$ admits a super $(a,3)$-edge-antimagic total labeling. Moreover, when p and t_1 satisfy certain conditions $T_p\langle t_1\rangle$ also admits a super (a,d)-edge-antimagic total labeling for $d=0$ and $d=2$. When $k=2$ they show the existence of a super $(a,2)$-edge-antimagic total labeling for the Toeplitz graph $T_p\langle t_1, t_1+1\rangle$.

In [216] Amudha, Jayapriya, and Gowri provide an algorithmic encryption method that employs antimagic labelings of graphs.

Pandimadevi and Subbiah [2754] show the existence and nonexistence of (a,d)-vertex antimagic total labeling for several class of digraphs and show how to construct labelings for generalized de Bruijn digraphs.

A *vertex antimagic total labeling* is a distinct assignment of positive integers to all vertices and edges of a graph such that the total weight obtained by adding the label of each vertex and the labels of all edges incident with it, is the same for every vertex in that graph. Kumar, Sathiamoorthy, and Balachandran [2066] proved the vertex antimagic total labeling of a modified fan graph obtained by merging isolated vertices from n copies of the disjoint union $K_1 \cup \overline{K_2} \cup \overline{K_m}$ to a single apex vertex. [2066] new

In [1249] Getzimah and Palani investigated vertex antimagic total labelings and edge antimagic total labelings of Z_{p+q}, cycles, stars, complete bipartite graphs, the subdivision graphs of ladders, and combs. They also investigate totally (a,d)-edge antimagic graphs, totally super vertex graphs, edge antimagic graphs, and determine the bounds for the vertices and the edges under total labelings.

For a graph G and an integer $s > 1$, Li [2239] z denotes the graph obtained from G by replacing each edge of G with P_s by G^s. He showed that G^s is antimagic when G a complete graph or a complete bipartite graph.

The book [403] by Bača and Miller has a wealth of material and open problems on super edge-antimagic labelings. In [359] Bača, Baskoro, Miller, Ryan, Simanjuntak, and Sugeng provide detailed survey of results on edge antimagic labelings and include many conjectures and open problems. In 2015 Nalliah [2646] published a list of open problems on super (a,d)-edge antimagic total labelings of graphs. In 2017 Brankovic, Jendrol', Lin, Phanalasy, Ryan, Semaničová-Feňovčíková, Slamin, and Sugeng [352] provided a survey of recent results on face-antimagic labelings. It was dedicated to the memory of Mirka Miller, who introduced the concept of face-antimagic labeling of plane graphs in 2003.

In Tables 14, 15, 16 and 17 we use the abbreviations

(a,d)-**VAT** (a,d)-vertex-antimagic total labeling

(a,d)-**SVAT** super (a,d)-vertex-antimagic total labeling

(a,d)-**EAT** (a,d)-edge-antimagic total labeling

(a,d)-**SEAT** super (a,d)-edge-antimagic total labeling

(a,d)-**EAV** (a,d)-edge-antimagic vertex labeling

A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The tables were prepared by Petr Kovář and Tereza Kovářová and updated by J. Gallian in 2008.

Table 14: **Summary of (a,d)-Vertex-Antimagic Total and Super (a,d)-Vertex-Antimagic Total Labelings**

Graph	*Labeling*	*Notes*
P_n	(a,d)-VAT	wide variety of a and d [361]
P_n	(a,d)-SVAT	iff $d=3,\ d=2, n\geq 3$ odd or $d=3,\ n\geq 3$ [3733]
C_n	(a,d)-VAT	wide variety of a and d [360]
C_n	(a,d)-SVAT	iff $d=0,2$ and n odd or $d=1$ [3733]
generalized Petersen graph $P(n,k)$	(a,d)-VAT $(a,1)$-VAT	[362] $n\geq 3,\ 1\leq k\leq n/2$ [3735]
prisms $C_n\times P_2$	(a,d)-VAT	[362]
antiprisms	(a,d)-VAT	[362]
$S_{n_1}\cup\ldots\cup S_{n_t}$	(a,d)-VAT	$d=1,2,3,4,6$ [2775], [3075]
W_n	not (a,d)-VAT	for $n>20$ [2271]
$K_{1,n}$	not (a,d)-SVAT	$n\geq 3$ [3733]

Table 15: **Summary of (a, d)-Edge-Antimagic Total Labelings**

Graph	*Labeling*	*Notes*
trees	$(a, 1)$-EAT?	[392]
P_n	not (a, d)-EAT	$d > 2$ [392]
P_{2n}	$(6n, 1)$-EAT	[3585]
	$(6n + 2, 2)$-EAT	[3585]
P_{2n+1}	$(3n + 4, 2)$-EAT	[3585]
	$(3n + 4, 3)$-EAT	[3585]
	$(2n + 4, 4)$-EAT	[3585]
	$(5n + 4, 2)$-EAT	[3585]
	$(3n + 5, 2)$-EAT	[3585]
	$(2n + 6, 4)$-EAT	[3585]
C_n	$(2n + 2, 1)$-EAT	[3585]
	$(3n + 2, 1)$-EAT	[3585]
	not (a, d)-EAT	$d > 5$ [392]
C_{2n}	$(4n + 2, 2)$-EAT	[3585]
	$(4n + 3, 2)$-EAT	[3585]
	$(2n + 3, 4)$-EAT?	[3585]
	$(2n + 4, 4)$-EAT?	[3585]
C_{2n+1}	$(3n + 4, 3)$-EAT	[3585]
	$(3n + 5, 3)$-EAT	[3585]
	$(n + 4, 5)$-EAT?	[3585]
	$(n + 5, 5)$-EAT?	[3585]
K_n	not (a, d)-EAT	$d > 5$ [392]
$K_{n,n}$	(a, d)-EAT	iff $d = 1, n \geq 2$ [393]
caterpillars	(a, d)-EAT	$d \leq 3$ [3734]
W_n	not (a, d)-EAT	$d > 4$ [392]
generalized Petersen	not (a, d)-EAT	$d > 4$ [392]
graph $P(n, k)$	$((5n + 5)/2, 2)$-EAT	$n \geq 3$ odd, $k = 1, 2$ [2700]
	super $(4n + 2, 1)$-EAT	$n \geq 3$, $1 \leq k \leq n/2$ [2700]

Table 16: **Summary of (a, d)-Edge-Antimagic Vertex Labelings**

Graph	*Labeling*	*Notes*
P_n	$(3, 2)$-EAV	[3585]
	not (a, d)-EAV	$d > 2$ [3585]
P_{2n}	$(n + 2, 1)$-EAV	[3585]
C_n	not (a, d)-EAV	$d > 1$ [392]
C_{2n}	not (a, d)-EAV	[3585]
C_{2n+1}	$(n + 2, 1)$-EAV	[3585]
	$(n + 3, 1)$-EAV	[3585]
K_n	not (a, d)-EAV	for $n > 1$ [392]
$K_{n,n}$	not (a, d)-EAV	for $n > 3$ [392]
W_n	not (a, d)-EAV	[392]
$C_3^{(n)}$ (friendship graph)	$(a, 1)$-EAV	iff $n = 1, 3, 4, 5, 7$ [393]
generalized Petersen graph $P(n, k)$	not (a, d)-EAV	$d > 1$ [392]

Table 17: **Summary of (a,d)-Super-Edge-Antimagic Total Labelings**

Graph	*Labeling*	*Notes*
$C_n \odot K_1$	(a,d)-SEAT	variety of cases [340], [419]
$P_n \times P_2$ (ladders)	(a,d)-SEAT	n odd, $d \leq 2$ [3732]
		n even, $d = 1$ [3732]
	(a,d)-SEAT?	$d = 0, 2,\ n$ even [3732]
$C_n \times P_2$	(a,d)-SEAT	iff $d \leq 3$ n odd [3732]
		or $d = 1,\ n \geq 4$ even [3732]
$C_m \times P_n$	(a,d)-SEAT?	$m \geq 4$ even,
		$n \geq 3,\ d = 0, 2$ [3732]
caterpillars	$(a,1)$-SEAT	[3734]
$C_3^{(n)}$ (friendship graphs)	(a,d)-SEAT	$d = 0, 1, 2$ [393]
F_n $(n \geq 2)$ (fans)	(a,d) SEAT	only if $d < 3$ [393]
	(a,d)-SEAT	$2 \leq n \leq 6, d = 0, 1, 2$ [393]
W_n	(a,d)-SEAT	iff $d = 1, n \not\equiv 1 \pmod 4$ [393]
K_n $(n \geq 3)$	(a,d) SEAT	iff $d = 0, n = 3$ [393]
		$d = 1, n \geq 3$ [393]
		$d = 2, n = 3$ [393]
trees	(a,d)-SEAT	only if $d \leq 3$ [394]
P_n $(n > 1)$	(a,d)-SEAT	iff $d \leq 3$ [394]
mK_n	(a,d)-SEAT	iff $d \in \{0,2\}, n \in \{2,3\}$,
		$m \geq 3$ odd [348]
		$d = 1, m, n \geq 2$ [348]
		$d = 3$ or $5, n = 2, m \geq 2$ [348]
		$d = 4, n = 2,\ m \geq 3$ odd [348]
C_n	(a,d)-SEAT	iff $d = 0$ or 2, n odd [394]
		$d = 1$ [360]
$P(m,n)$	(a,d)-SEAT	many cases [360]

6.4 Face Antimagic Labelings and d-antimagic Labeling of Type (1,1,1)

Bača [342] defines a connected plane graph G with edge set E and face set F to be (a,d)-*face antimagic* if there exist positive integers a and d and a bijection $g\colon E \to \{1,2,\ldots,|E|\}$ such that the induced mapping $\psi_g\colon F \to \{a, a+d, \ldots, a+(|F(G)|-1)d\}$, where for a face f, $\psi_g(f)$ is the sum of all $g(e)$ for all edges e surrounding f is also a bijection. In [344] Bača proves that for n even and at least 4, the prism $C_n \times P_2$ is $(6n+3,2)$-face antimagic and $(4n+4,4)$-face antimagic. He also conjectures that $C_n \times P_2$ is $(2n+5,6)$-face antimagic. In [389] Bača, Lin, and Miller investigate (a,d)-face antimagic labelings of the convex polytopes $P_{m+1} \times C_n$. They show that if these graphs are (a,d)-face antimagic then either $d=2$ and $a=3n(m+1)+3$, or $d=4$ and $a=2n(m+1)+4$, or $d=6$ and $a=n(m+1)+5$. They also prove that if n is even, $n \geq 4$ and $m \equiv 1 \pmod 4$, $m \geq 3$, then $P_{m+1} \times C_n$ has a $(3n(m+1)+3,2)$-face antimagic labeling and if n is at least 4 and even and m is at least 3 and odd, or if $n \equiv 2 \pmod 4$, $n \geq 6$ and m is even, $m \geq 4$, then $P_{m+1} \times C_n$ has a $(3n(m+1)+3,2)$-face antimagic labeling and a $(2n(m+1)+4,4)$-face antimagic labeling. They conjecture that $P_{m+1} \times C_n$ has $(3n(m+1)+3,2)$- and $(2n(m+1)+4,4)$-face antimagic labelings when $m \equiv 0 \pmod 4$, $n \geq 4$, and for m even and $m \geq 4$, that $P_{m+1} \times C_n$ has a $(n(m+1)+5,6)$-face antimagic labeling when n is even and at least 4. Bača, Baskoro, Jendroľ, and Miller [356] proved that graphs in the shape of hexagonal honeycombs with m rows, n columns, and mn 6-sided faces have d-antimagic labelings of type $(1,1,1)$ for $d=1,2,3$, and 4 when n odd and $mn>1$.

In [401] Bača and Miller define the class Q_n^m of convex polytopes with vertex set $\{y_{j,i} : i=1,2,\ldots,n; j=1,2,\ldots,m+1\}$ and edge set $\{y_{j,i}y_{j,i+1} : i=1,2,\ldots,n; j=1,2,\ldots,m+1\} \cup \{y_{j,i}y_{j+1,i} : i=1,2,\ldots,n; j=1,2,\ldots,m\} \cup \{y_{j,i+1}y_{j+1,i} : 1+1,2,\ldots,n; j=1,2,\ldots,m, j \text{ odd}\} \cup \{y_{j,i}y_{j+1,i+1} : i=1,2,\ldots,n; j=1,2,\ldots,m, j \text{ even}\}$ where $y_{j,n+1}=y_{j,1}$. They prove that for m odd, $m \geq 3, n \geq 3$, Q_n^m is $(7n(m+1)/2+2,1)$-face antimagic and when m and n are even, $m \geq 4, n \geq 4$, Q_n^m is $(7n(m+1)/2+2,1)$-face antimagic. They conjecture that when n is odd, $n \geq 3$, and m is even, then Q_n^m is $((5n(m+1)+5)/2,2)-$face antimagic and $((n(m+1)+7)/2,4)$-face antimagic. They further conjecture that when n is even, $n>4, m>1$ or n is odd, $n>3$ and m is odd, $m>1$, then Q_n^m is $(3n(m+1)/2+3,3)$-face antimagic. In [346] Bača proves that for the case $m=1$ and $n \geq 3$ the only possibilities for (a,d)-antimagic labelings for Q_n^m are $(7n+2,1)$ and $(3n+3,3)$. He provides the labelings for the first case and conjectures that they exist for the second case. Bača [342] and Bača and Miller [400] describe (a,d)-face antimagic labelings for a certain classes of convex polytopes.

In [355] Bača et al. provide a detailed survey of results on face antimagic labelings and include many conjectures and open problems.

For a plane graph G, Bača and Miller [402] call a bijection h from $V(G) \cup E(G) \cup F(G)$ to $\{1,2,\ldots,|V(G)|+|E(G)| \cup |F(G)|\}$ a *d-antimagic labeling of type* $(1,1,1)$ if for every number s the set of s-sided face weights is $W_s=\{a_s, a_s+d, a_s+2d, \ldots, a_s+(f_s-1)d\}$ for some integers a_s and d, where f_s is the number of s-sided faces (W_s varies with s). They show that the prisms $C_n \times P_2$ $(n \geq 3)$ have a 1-antimagic labeling of type $(1,1,1)$ and

that for $n \equiv 3 \pmod 4$, $C_n \times P_2$ have a d-antimagic labeling of type $(1,1,1)$ for $d = 2,3,4$, and 6. They conjecture that for all $n \geq 3$, $C_n \times P_2$ has a d-antimagic labeling of type $(1,1,1)$ for $d = 2,3,4,5$, and 6. This conjecture has been proved for the case $d = 3$ and $n \neq 4$ by Bača, Miller, and Ryan [408] (the case $d = 3$ and $n = 4$ is open). The cases for $d = 2,4,5$, and 6 were done by Lin, Slamin, Bača, and Miller [2272]. Bača, Lin, and Miller [390] prove: for $m, n > 8$, $P_m \times P_n$ has no d-antimagic edge labeling of type $(1,1,1)$ with $d \geq 9$; for $m \geq 2, n \geq 2$, and $(m,n) \neq (2,2)$, $P_m \times P_n$ has d-antimagic labelings of type $(1,1,1)$ for $d = 1,2,3,4$, and 6. They conjecture the same is true for $d = 5$. Butt, Numan, Shah, and Ali [732] prove that the generalized prisms $C_n \times P_m$ have d-antimagic face labelings of type (1,1,1) for $n \geq 5$ and $m \geq 2$.

Bača, Miller, and Ryan [408] also prove that for $n \geq 4$ the antiprism (see §6.1 for the definition) on $2n$ vertices has a d-antimagic labeling of type $(1,1,1)$ for $d = 1,2$, and 4. They conjecture the result holds for $d = 3,5$, and 6 as well. Lin, Ahmad, Miller, Sugeng, and Bača [2269] did the cases that $d = 7$ for $n \geq 3$ and $d = 12$ for $n \geq 11$. Sugeng, Miller, Lin, and Bača [3735] did the cases: $d = 7,8,9,10$ for $n \geq 5$; $d = 15$ for $n \geq 6$; $d = 18$ for $n \geq 7$; $d = 12,14,17,20,21,24,27,30,36$ for n odd and $n \geq 7$; and $d = 16,26$ for n odd and $n \geq 9$.

Baca, Numan, and Semaničová-Feňovčíková [416] investigated the problem of labeling the vertices, edges, and faces of a disjoint union of r copies $C_n \times P_m$ by the consecutive integers starting from 1 in such a way that the sum of the labels of a face and the labels of vertices and edges surrounding that face for all s-sided faces form an arithmetic progression with common difference d.

Ali, Bača, Bashir, and Semaničová-Feňovčíková [184] investigated antimagic labelings for disjoint unions of prisms and cycles. They prove: for $m \geq 2$ and $n \geq 3$, $m(C_n \times P_2)$ has no super d-antimagic labeling of type $(1,1,1)$ with $d \geq 30$; for $m \geq 2$ and $n \geq 3$, $n \neq 4$, $m(C_n \times P_2)$ has super d-antimagic labeling of type $(1,1,1)$ for $d = 0,1,2,3,4$, and 5; and for $m \geq 2$ and $n \geq 3$, mC_n has $(m(n+1)+3,3)$- and $(2mn+2,2)$-vertex-antimagic total labeling. Bača and Bashir [349] proved that for $m \geq 2$ and $n \geq 3$, $n \neq 4$, $m(C_n \times P_2)$ has super 7-antimagic labeling of type $(1,1,1)$ and for $n \geq 3$, $n \neq 4$ and $2 \leq m \leq 2n$ $m(C_n \times P_2)$ has super 6-antimagic labeling of type $(1,1,1)$.

Bača, Numan and Siddiqui [418] investigated the existence of the super d-antimagic labeling of type $(1,1,1)$ for the disjoint union of m copies of antiprism mA_n. They proved that for $m \geq 2$, $n \geq 4$, mA_n has super d-antimagic labelings of type $(1,1,1)$ for $d = 1,2,3,5,6$. Ahmad, Bača, Lascsáková, and Semaničová-Feňovčíková [114] investigated super d-antimagicness of type $(1,1,0)$ for mG in a more general sense. They prove: if there exists a super 0-antimagic labeling of type $(1,1,0)$ of a plane graph G then, for every positive integer m, the graph mG also admits a super 0-antimagic labeling of type $(1,1,0)$; if a plane graph G with 3-sided inner faces admits a super d-antimagic labeling of type $(1,1,0)$ for $d = 0,6$ then, for every positive integer m, the graph mG also admits a super d-antimagic labeling of type $(1,1,0)$; if a plane graph G with 3-sided inner faces is a tripartite graph with a super d-antimagic labeling of type $(1,1,0)$ for $d = 2,4$ then, for every positive integer m, the graph mG also admits a super d-antimagic labeling of type $(1,1,0)$; if a plane graph G with 4-sided inner faces admits a super d-antimagic labeling of

type $(1,1,0)$ for $d = 0, 4, 8$ then the disjoint union of arbitrary number of copies of G also admits a super d-antimagic labeling of type $(1,1,0)$; if a plane graph G with k-sided inner faces, $k \geq 3$, admits a super d-antimagic labeling of type $(1,1,0)$ for $d = 0, 2k$ then, for every positive integer m, the graph mG also admits a super d-antimagic labeling of type $(1,1,0)$; if a plane graph G with k-sided inner faces admits a super k-antimagic labeling of type $(1,1,0)$ for k even then, for every positive integer m, the graph mG also admits a super k-antimagic labeling of type $(1,1,0)$.

Bača, Jendraľ, Miller, and Ryan [378] prove: for n even, $n \geq 6$, the generalized Petersen graph $P(n,2)$ has a 1-antimagic labeling of type $(1,1,1)$; for n even, $n \geq 6$, $n \neq 10$, and $d = 2$ or 3, $P(n,2)$ has a d-antimagic labeling of type $(1,1,1)$; and for $n \equiv 0 \pmod 4$, $n \geq 8$ and $d = 6$ or 9, $P(n,2)$ has a d-antimagic labeling of type $(1,1,1)$. They conjecture that there is an d-antimagic labeling of type (1,1,1) for $P(n,2)$ when $n \equiv 2 \pmod 4$, $n \geq 6$, and $d = 6$ or 9.

In [365] Bača, Brankovic, and A. Semaničová-Feňovčikovă provide super d-antimagic labelings of type (1,1,1) for friendship graphs F_n $(n \geq 2)$ and several other families of planar graphs.

Bača, Brankovic, Lascsáková, Phanalasy, and Semaničová-Feňovčíková [364] provided super d-antimagic labeling of type $(1,1,0)$ for friendship graphs F_n, $n \geq 2$, for $d \in \{1,3,5,7,9,11,13\}$. Moreover, they show that for $n \equiv 1 \pmod 2$ the graph F_n also admits a super d-antimagic labeling of type $(1,1,0)$ for $d \in \{0,2,4,6,8,10\}$.

Bača, Baskoro, and Miller [357] have proved that hexagonal planar honeycomb graphs with an even number of columns have 2-antimagic and 4-antimagic labelings of type $(1,1,1)$. They conjecture that these honeycombs also have d-antimagic labelings of type $(1,1,1)$ for $d = 3$ and 5. They pose the odd number of columns case for $1 \leq d \leq 5$ as an open problem. Bača, Baskoro, and Miller [358] give d-antimagic labelings of a special class of plane graphs with 3-sided internal faces for $d = 0$, 2, and 4. Bača, Lin, Miller, and Ryan [391] prove for odd $n \geq 3$, $m \geq 1$ and $d = 0, 1, 2$ or 4, the Möbius grid M_n^m has an d-antimagic labeling of type $(1,1,1)$. Siddiqui, Numan, and Umar [3581] examined the existence of super d-antimagic labelings of type (1,1,1) for Jahangir graphs for certain differences d.

Bača, Numan, and Shabbir [417] studied the existence of super d-antimagic labelings of type $(1,1,1)$ for the toroidal polyhex $\mathbb{H}_m^n$. They labeled the edges of a 1-factor by consecutive integers and then in successive steps they labeled the edges of $2m$-cycles (respectively $2n$-cycles) in a 2-factor by consecutive integers. This technique allowed them to construct super d-antimagic labelings of type $(1,1,1)$ for $\mathbb{H}_m^n$ with $d = 1, 3, 5$. They suppose that such labelings exist also for $d = 0, 2, 4$.

Kathiresan and Ganesan [1926] define a class of plane graphs denoted by P_a^b $(a \geq 3, b \geq 2)$ as the graph obtained by starting with vertices $v_1, v_2, \ldots, v_a$ and for each $i = 1, 2 \ldots, a-1$ joining v_i and v_{i+1} with b internally disjoint paths of length $i+1$. They prove that P_a^b has d-antimagic labelings of type $(1,1,1)$ for $d = 0, 1, 2, 3, 4$, and 6. Lin and Sugen [2273] prove that P_a^b has a d-antimagic labeling of type $(1,1,1)$ for $d = 5, 7a-2, a+1, a-3, a-7, a+5, a-4, a+2, 2a-3, 2a-1, a-1, 3a-3, a+3, 2a+1, 2a+3, 3a+1, 4a-1, 4a-3, 5a-3, 3a-1, 6a-5, 6a-7, 7a-7$, and $5a-5$. Similarly, Bača, Baskoro,

and Cholily [354] define a class of plane graphs denoted by C_a^b as the graph obtained by starting with vertices $v_1, v_2, \ldots, v_a$ and for each $i = 1, 2 \ldots, a$ joining v_i and v_{i+1} with b internally disjoint paths of length $i+1$ (subscripts are taken modulo a). In [354] and [353] they prove that for $a \geq 3$ and $b \geq 2$, C_a^b has a d-antimagic labeling of type $(1, 1, 1)$ for $d = 0, 1, 2, 3, a+1, a-1, a+2$, and $a-2$.

In [366] Bača, Brankovic, and Semaničová-Feňovčiková investigated the existence of super d-antimagic labelings of type (1,1,1) for plane graphs containing a special kind of Hamilton path. They proved: if there exists a Hamilton path in a plane graph G such that for every face except the external face, the Hamilton path contains all but one of the edges surrounding that face, then G is super d-antimagic of type (1,1,1) for $d = 0, 1, 2, 3, 5$; if there exists a Hamilton path in a plane graph G such that for every face except the external face, the Hamilton path contains all but one of the edges surrounding that face and if $2(|F(G)| - 1) \leq |V(G)|$, then G is super d-antimagic of type $(1, 1, 1)$ for $d = 0, 1, 2, 3, 4, 5, 6$; if G is a plane graph with $M = \lfloor \frac{|V(G)|}{|F(G)|-1} \rfloor$ and a Hamilton path such that for every face, except the external face, the Hamilton path contains all but one of the edges surrounding that face, then for $M = 1$, G admits a super d-antimagic labeling of type (1,1,1) for $d = 0, 1, 2, 3, 5$; and for $M \geq 2$, G admits a super d-antimagic labeling of type (1,1,1) for $d = 0, 1, 2, 3, \ldots, M+4$. They also proved that $P_n \times P_2$ $(n \geq 3)$ admits a super d-antimagic labeling of type (1,1,1) for $d \in \{0, 1, 2, \ldots, 15\}$ and the graph obtained from $P_n \times P_m$ $(n \geq 2)$ by adding a new edge in every 4-sided face such that the added edges are "parallel" admits a super d-antimagic labeling of type (1,1,1) for $d \in \{0, 1, 2, \ldots, 9\}$.

In [1529] Imran, Siddiqui, and Numan examine the existence of super d-antimagic labelings of type (1,1,1) for uniform subdivision of wheel for certain differences d.

In the following tables we use the abbreviations

(a, d)-**FA** (a, d)-face antimagic labeling

d-**AT(1,1,1)** d-antimagic labeling of type $(1, 1, 1)$.

A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property. The tables were prepared by Petr Kovář and Tereza Kovářová and updated by J. Gallian in 2008.

Table 18: **Summary of Face Antimagic Labelings**

Graph	*Labeling*	*Notes*
Q_n^m (see §6.4)	$(7n(m+1)/2+2,1)$-FA	$m \geq 3$, $n \geq 3$, m odd [401]
	$(7n(m+1)/2+2,1)$-FA	$m \geq 4$, $n \geq 4$, m, n even [401]
	$((5n(m+1)+5)/2,2)$-FA?	$m \geq 2$, $n \geq 3$, m even, n odd [401]
	$((n(m+1)+7)/2,4)$-FA?	$m \geq 2$, $n \geq 3$, m even, n odd [401]
	$(3n(m+1)/2+3,3)$-FA?	$m > 1$, $n > 4$, n even [401]
	$(3n(m+1)/2+3,3)$-FA?	$m > 1$, $n > 3$, m odd, n odd [401]
$C_n \times P_2$	$(6n+3,2)$-FA	$n \geq 4$, n even [344]
	$(4n+4,4)$-FA	$n \geq 4$, n even [344]
	$(2n+5,6)$-FA?	[344]
$P_{m+1} \times C_n$	$(3n(m+1)+3,2)$-FA	$n \geq 4$, n even and [389]
		$m \geq 3$, $m \equiv 1 \pmod 4$,
	$(3n(m+1)+3,2)$-FA and	$n \geq 4$, n even and [389]
	$(2n(m+1)+4,4)$-FA	$m \geq 3$, m odd [389],
		or $n \geq 6$, $n \equiv 2 \pmod 4$ and
		$m \geq 4$, m even
	$(3n(m+1)+3,2)$-FA?	$m \geq 4$, $n \geq 4$, $m \equiv 0 \pmod 4$ [389]
	$(2n(m+1)+4,4)$-FA?	$m \geq 4$, $n \geq 4$, $m \equiv 0 \pmod 4$ [389]
	$(n(m+1)+5,6)$-FA?	$n \geq 4$, n even [389]

Table 19: **Summary of d-antimagic Labelings of Type (1,1,1)**

Graph	*Labeling*	*Notes*
$P_m \times P_n$	not d-AT(1,1,1)	$m, n, d \geq 9$, [390]
$P_m \times P_n$	d-AT(1,1,1)	$d = 1, 2, 3, 4, 6$;
		$m, n \geq 2$, $(m, n) \neq (2, 2)$ [390]
$P_m \times P_n$	5-AT(1,1,1)	$m, n \geq 2$, $(m, n) \neq (2, 2)$ [390]
$C_n \times P_2$	1-AT(1,1,1)	[402]
	d-AT(1,1,1)	$d = 2, 3, 4$ and 6 [402]
		for $n \equiv 3 \pmod 4$
	d-AT(1,1,1)	$d = 2, 4, 5, 6$ for $n \geq 3$ [2272]

Continued on next page

Table 19 – *Continued from previous page*

Graph	*Labeling*	*Notes*
	d-AT(1,1,1)	$d = 3$ for $n \geq 5$ [408]
$P_m \times P_n$	5-AT(1,1,1)?	[2272]
	not d-AT	$m, n > 8,\ d \geq 9$ [2272]
antiprism on $2n$ vertices	d-AT(1,1,1)	$d = 1, 2$ and 4 for $n \geq 4$ [408]
	d-AT(1,1,1)?	$d = 3, 5$ and 6 for $n \geq 4$ [408]
M_n^m (Möbius grids)	d-AT(1,1,1)	$n \geq 3$ odd, $d = 0, 1, 2, 4$ [391]
		$d = 7,\ n \geq 3$ [2269]
		$d = 12,\ n \geq 11$ [2269]
		$d = 7, 8, 9, 10,\ n \geq 5$ [3735]
		$d = 15,\ n \geq 6$ [3735]
		$d = 18\ n \geq 7$ [3735]
$P(n, 2)$	d-AT(1,1,1)	$d = 1;\ d = 2, 3,\ n \geq 6,$ $n \neq 10$ [378]
$P(4n, 2)$	d-AT(1,1,1)	$d = 6, 9,\ n \geq 2,\ n \neq 10$ [378]
$P(4n + 2, 2)$	d-AT(1,1,1)?	$d = 6, 9,\ n \geq 1,\ n \neq 10$ [378]
honeycomb graphs with even number of columns	d-AT(1,1,1)	$d = 2, 4$ [357]
	d-AT(1,1,1)?	$d = 3, 5$ [357]
$C_n \times P_2$	d-AT(1,1,1)	$d = 1, 2, 4, 5, 6$ [2272], [402]
$C_n \times P_2$	3-AT(1,1,1)	$n \neq 4$ [408]

6.5 Product Antimagic Labelings

Figueroa-Centeno, Ichishima, and Muntaner-Batle [1086] have introduced multiplicative analogs of magic and antimagic labelings. They define a graph G of size q to be *product magic* if there is a labeling from $E(G)$ onto $\{1, 2, \ldots, q\}$ such that, at each vertex v, the product of the labels on the edges incident with v is the same. They call a graph G of size q *product antimagic* if there is a labeling$f$ from $E(G)$ onto $\{1, 2, \ldots, q\}$ such that the products of the labels on the edges incident at each vertex v are distinct. They prove: a graph of size q is product magic if and only if $q \leq 1$ (that is, if and only if it is $K_2, \overline{K_n}$ or

$K_2 \cup \overline{K_n}$); P_n ($n \geq 4$) is product antimagic; every 2-regular graph is product antimagic; and, if G is product antimagic, then so are $G + K_1$ and $G \odot \overline{K}_n$. They conjecture that a connected graph of size q is product antimagic if and only if $q \geq 3$. Kaplan, Lev, and Roditty [1904] proved the following graphs are product antimagic: the disjoint union of cycles and paths where each path has least three edges; connected graphs with n vertices and m edges where $m \geq 4n\ln n$; graphs $G = (V, E)$ where each component has at least two edges and the minimum degree of G is at least $8\sqrt{\ln |E| \ln (\ln |E|)}$; all complete k-partite graphs except K_2 and $K_{1,2}$; and $G \odot H$ where G has no isolated vertices and H is regular. Wang and Gao [4142] show that caterpillars with at least three edges are product antimagic by an $O(m \log m)$ algorithm.

In [2842] Pikhurko characterizes all large graphs that are product antimagic graphs. More precisely, it is shown that there is an n_0 such that a graph with $n \geq n_0$ vertices is product antimagic if and only if it does not belong to any of the following four classes: graphs that have at least one isolated edge; graphs that have at least two isolated vertices; unions of vertex-disjoint of copies of $K_{1,2}$; graphs consisting of one isolated vertex; and graphs obtained by subdividing some edges of the star $K_{1,k+l}$.

In [2545] Mora and Tey proved that every connected graph of size $m \geq 3$ admits an antimagic labeling for every arithmetic sequence of m positive real numbers, if every vertex of degree at least three has at least one pendant edge incident with it. As a corollary, this shows that these graphs are antimagic. Moreover, these graphs admit also a product antimagic with these labels provided that the smallest label is at least one. [2545] new

In [1086] Figueroa-Centeno, Ichishima, and Muntaner-Batle also define a graph G with p vertices and q edges to be *product edge-magic* if there is a labeling f from $V(G) \cup E(G)$ onto $\{1, 2, \ldots, p+q\}$ such that $f(u) \cdot f(v) \cdot f(uv)$ is a constant for all edges uv and *product edge-antimagic* if there is a labeling f from $V(G) \cup E(G)$ onto $\{1, 2, \ldots, p+q\}$ such that for all edges uv the products $f(u) \cdot f(v) \cdot f(uv)$ are distinct. They prove $K_2 \cup \overline{K}_n$ is product edge-magic, a graph of size q without isolated vertices is product edge-magic if and only if $q \leq 1$ and every graph other than K_2 and $K_2 \cup \overline{K}_n$ is product edge-antimagic.

%

7 Miscellaneous Labelings

7.1 Sum Graphs

In 1990, Harary [1368] introduced the notion of a sum graph. A graph $G(V, E)$ is called a *sum graph* if there is a bijection f from V to a set of positive integers S such that $xy \in E$ if and only if $f(x) + f(y) \in S$. Since the vertex with the highest label in a sum graph cannot be adjacent to any other vertex, every sum graph must contain isolated vertices. In 1991 Harary, Hentzel, and Jacobs [1370] defined a *real sum graph* in an analogous way by allowing S to be any finite set of positive real numbers. However, they proved that every real sum graph is a sum graph. Bergstrand, Hodges, Jennings, Kuklinski, Wiener, and Harary [625] defined a *product graph* analogous to a sum graph except that 1 is not permitted to belong to S. They proved that every product graph is a sum graph and vice versa.

For a connected graph G, let $\sigma(G)$, the *sum number of* G, denote the minimum number of isolated vertices that must be added to G so that the resulting graph is a sum graph (some authors use $s(G)$ for the sum number of G). A labeling that makes G together with $\sigma(G)$ isolated points a sum graph is called an *optimal sum graph* labeling. Ellingham [1005] proved the conjecture of Harary [1368] that $\sigma(T) = 1$ for every tree $T \neq K_1$. Smyth [3642] proved that there is no graph G with e edges and $\sigma(G) = 1$ when $n^2/4 < e \leq n(n-1)/2$. Smyth [3643] conjectures that the disjoint union of graphs with sum number 1 has sum number 1. More generally, Kratochvil, Miller, and Nguyen [2029] conjecture that $\sigma(G \cup H) \leq \sigma(G) + \sigma(H) - 1$. Hao [1362] has shown that if $d_1 \leq d_2 \leq \cdots \leq d_n$ is the degree sequence of a graph G, then $\sigma(G) > \max(d_i - i)$ where the maximum is taken over all i. Bergstand et al. [624] proved that $\sigma(K_n) = 2n - 3$. Hartsfield and Smyth [1379] claimed to have proved that $\sigma(K_{m,n}) = \lceil 3m + n - 3 \rceil / 2$ when $n \geq m$ but Yan and Liu [4240] found counterexamples to this assertion when $m \neq n$. Pyatkin [3061], Liaw, Kuo, and Chang [2260], Wang, and Liu [4168], and He, Shen, Wang, Chang, Kang, and Yu [1391] have shown that for $2 \leq m \leq n$, $\sigma(K_{m,n}) = \lceil \frac{n}{p} + \frac{(p+1)(m-1)}{2} \rceil$ where $p = \lceil \sqrt{\frac{2n}{m-1} + \frac{1}{4}} - \frac{1}{2} \rceil$ is the unique integer such that $\frac{(p-1)p(m-1)}{2} < n \leq \frac{(p+1)p(m-1)}{2}$.

Miller, Ryan, Slamin, and Smyth [2505] proved that $\sigma(W_n) = \frac{n}{2} + 2$ for n even and $\sigma(W_n) = n$ for $n \geq 5$ and n odd (see also [3839]). Miller, Ryan, and Smyth [2507] prove that the complete n-partite graph on n sets of 2 nonadjacent vertices has sum number $4n - 5$ and obtain upper and lower bounds on the complete n-partite graph on n sets of m nonadjacent vertices. Fernau, Ryan, and Sugeng [1082] proved that the generalized friendship graphs $C_n^{(t)}$ (see §2.2) has sum number 2 except for C_4. Gould and Rödl [1321] investigated bounds on the number of isolated points in a sum graph. A group of six undergraduate students [1306] proved that $\sigma(K_n - \text{edge}) \leq 2n - 4$. The same group of six students also investigated the difference between the largest and smallest labels in a sum graph, which they called the *spum*. They proved spum of K_n is $4n - 6$ and the spum

of C_n is at most $4n - 10$. Kratochvil, Miller, and Nguyen [2029] have proved that every sum graph on n vertices has a sum labeling such that every label is at most 4^n. Konečný, Kučera, Novotná, Pekárek, Šimsa, and Töpfer [1996] showed that if one allows for non-injective labelings or graphs with loops then there are sum graphs without a minimal sum labeling, which partially answers the question posed by Miller, Ryan and Smyth in [2507]. At a conference in 2000 Miller [2492] posed the following two problems: Given any graph G, does there exist an optimal sum graph labeling that uses the label 1; Find a class of graphs G that have sum number of the order $|V(G)|^s$ for $s > 1$. (Such graphs were shown to exist for $s = 2$ by Gould and Rödl in [1321]). Fernau and Gajjar [1078] provided a complete characterization of the sum number of graphs of maximum degree two. As an immediate corollary they have that all graphs that are the disjoint unions of paths and cycles are sum labeled graphs.

In [3628] Slamet, Sugeng, and Miller show how one can use sum graph labelings to distribute secret information to set of people so that only authorized subsets can reconstruct the secret.

Chang [772] generalized the notion of sum graph by permitting $x = y$ in the definition of sum graph. He calls graphs that have this kind of labeling *strong sum graphs* and uses $i^*(G)$ to denote the minimum positive integer m such that $G \cup mK_1$ is a strong sum graph. Chang proves that $i^*(K_n) = \sigma(K_n)$ for $n = 2, 3$, and 4 and $i^*(K_n) > \sigma(K_n)$ for $n \geq 5$. He further shows that for $n \geq 5$, $3n^{\log_2 3} > i^*(K_n) \geq 12\lfloor n/5 \rfloor - 3$.

In [1079] Fernau and Gajjar initiate the study of sum graphs from the viewpoint of computational complexity. They show that every n-vertex, m-edge, d-degenerate graph (a graph is d-degenerate if all its subgraphs have a vertex of degree at most d) can be made a sum graph by adding at most m isolated vertices to it, such that the largest numbers used as vertex labels grows as $\Theta(n^2 d)$. As a consequence, we have that such a graph can be stored using $\Theta(m \log n)$ bits of memory. The previously best known upper bound on the numbers needed for labeling general graphs with the minimum number of isolated vertices was $\Theta(4n)$, due to Kratochvíl, Miller and Nguyen [2029] in 2001. Moreover, Fernau and Gajjar's labeling can be constructed in polynomial time. Their results show the ever-expanding database that is gradually built up can be efficiently implemented and then accessed using their sum-labeling scheme.

In 1994 Harary [1369] generalized sum graphs by permitting S to be any set of integers. He calls these graphs *integral sum graphs*. Unlike sum graphs, integral sum graphs need not have isolated vertices. Sharary [3488] has shown that C_n and W_n are integral sum graphs for all $n \neq 4$. Chen [798] proved that trees obtained from a star by extending each edge to a path and trees all of whose vertices of degree not 2 are at least distance 4 apart are integral sum graphs. He conjectures that all trees are integral sum graphs. In [798] and [800] Chen gives methods for constructing new connected integral sum graphs from given integral sum graphs by identifying vertices. Chen [800] has shown that every graph is an induced subgraph of a connected integral sum graph. Chen [800] calls a vertex of a graph *saturated* if it is adjacent to every other vertex of the graph. He proves that every integral sum graph except K_3 has at most two saturated vertices and gives the exact structure of all integral sum graphs that have exactly two saturated vertices. Chen [800]

also proves that a connected integral sum graph with $p > 1$ vertices and q edges and no saturated vertices satisfies $q \leq p(3p-2)/8 - 2$. Wu, Mao, and Le [4212] proved that mP_n are integral sum graphs. They also show that the conjecture of Harary [1369] that the sum number of C_n equals the integral sum number of C_n if and only if $n \neq 3$ or 5 is false and that for $n \neq 4$ or 6 the integral sum number of C_n is at most 1. Vilfred and Nicholas [4097] prove that graphs G of order n with $\Delta(G) = n-1$ and $|V_\Delta(G)| > 2$ are not integral sum graphs, except K_3, and that integral sum graphs G of order n with $\Delta(G) = n-1$ and $|V_\Delta(G)| = 2$ exist and are unique up to isomorphism. Chen [802] proved that if $G(V, E)$ is an integral sum other than K_3 that has vertex of degree $|V|-1$, then the edge-chromatic number of G is $|V|-1$. In [2611] Muthukkumar and Rajendran proved that all trees are integral sum graphs and that every bipartite graph is an induced subgraph of a sum graph with sum number 1. Elizeche and Tripathi [1004] characterized sum and integral sum labelings of complete graphs, symmetric complete bipartite graphs, star graphs, and deduced the spum, integral spum, and integral radius for these classes of graphs. In [2673] Nicholas, Somasundaram, and Vilfred showed that $\sigma(G) = \zeta(G)$ for $G = P_n \times P_2$ and the equality does not hold for $G = mC_n$ ($n \neq 4$). They also proved that a special class of caterpillars are integral sum graphs. [2673] new

In 2021 Singla, Tiwari, and Tripathi [3595] were the first to investigated the spum and integral spum (denoted ispum(G). They provided results on complete graphs, symmetric complete bipartite graphs, star graphs, cycles, and paths and gave sharp lower bounds for the spum and the integral spum of connected graphs. Shortly after that paper appeared, Li [2237] used some of the techniques in [3595], as well as his own, to correct multiple errors and improve many results in [3595]. The paper by Singla, Tiwari, and Tripathi and the one by Li open up new areas for research.

Following are some of the corrected and improved results. For $3 \leq n \leq 6$, $\text{spum}(P_n) = 2n-3$, and for $n \geq 7$, $2n-2 \leq \text{spum}(P_n) \leq 2n+1$ if n is odd, and $2n-2 \leq \text{spum}(P_n) \leq 2n-1$ if n is even. Li conjectures that for $n \geq 8$, we have $\text{spum}(P_n) = 2n+1$ if n is odd and $\text{spum}(P_n) = 2n-1$ if n is even. He verified the conjecture for n up to 15. In [3595] Singla, Tiwari, and Tripathi gave a flawed proof that $\text{spum}(C_n) = 2n-1$ for all $n \geq 13$. Li [2237] proved $\text{spum}(C_3) = 6$ and $\text{spum}(C_n) = 2n-1$ for $n \geq 4$.

Li [2237] improved the previous best upper bounds given for ispum(C_n) in [2484] as follows. For $n \geq 12$, we have ispum$(C_n) \leq 8(n-9)$ if n is odd and $\frac{3}{2}(3n-14)$ if n is even.

In 1990 Harary [1368] showed that $\zeta(nK_2) = 0$ for all positive integers n, but Li is the first to investigate the spum and ispum of nK_2. He proved that for all positive integers n $\sigma(nK_2) = 1$, $\text{spum}(nK_2) = 4n-2$, and ispum$(2K_2) = 4$ and ispum$(nK_2) = 4n-3$ for $n \neq 2$.

Motivated by the fact that there is no obvious relationship between ispum(G) and spum(G), due to $\zeta(G)$ possibly being strictly smaller than $\sigma(G)$, i.e., when the inequality $\zeta(G) \leq \sigma(G)$ is strict, Li introduced the following modification of spum. The *sum-diameter* of a graph G, denoted sd(G), is the minimum possible value of range(L) for a set L of positive integer labels, such that the induced sum graph of L consists of the disjoint union of G with any number of isolated vertices. The difference between the definition of sum-diameter and spum is that in spum, one is required to use the minimum

number of additional isolated vertices possible. Similarly, Li defines the *integral sum-diameter* of a graph G, denoted $\mathrm{isd}(G)$, by allowing the labels to be arbitrary distinct integers. He notes that $\mathrm{sd}(G) \leq \mathrm{spum}(G)$, as any set of labels L from the definition of $\mathrm{spum}(G)$, i.e., on $n + \sigma(G)$ vertices, is a valid labeling for the definition of $\mathrm{sd}(G)$. Similarly, $\mathrm{isd}(G) \leq \mathrm{sd}(G)$. Noting that any labeling L from the definition of $\mathrm{sd}(G)$ is a valid labeling for the definition of $\mathrm{isd}(G)$, yields $\mathrm{isd}(G) \leq \mathrm{sd}(G)$ for all graphs G.

Li observes that the following stronger result, which was initially stated for $\mathrm{spum}(G)$ in [3595] rather than $\mathrm{sd}(G)$, follows from the same proof. For graphs G of order n without any isolated vertices, with maximum and minimum vertex degree Δ and δ, respectively, $\mathrm{sd}(G) \geq 2n - (\Delta - \delta) - 2$.

Similarly, he noticed that the proof of Theorem 2.2 in [3595] does not use the assumption that exactly $\zeta(G)$ isolated vertices are present when bounding $\mathrm{ispum}(G)$, and thus the following strengthening is true. For graphs G of order n without any isolated vertices, with maximum degree Δ, we have $\mathrm{isd}(G) \geq 2n - \Delta - 3$.

Li uses the notion of Sidon sets to prove that if G is a graph with n vertices, then $\mathrm{sd}(G) \leq 64n^2 - 64n + 9$. Because $\mathrm{isd}(G) \leq \mathrm{sd}(G)$, we also have that $\mathrm{isd}(G) \leq 64n^2 - 64n + 9$. To show this upper bound is asymptotically tight, for a graph G with n vertices, he lets $f(n)$ be the maximum value of $\mathrm{sd}(G)$ over all graphs G with n vertices. Then Li proves $f(n) \geq \frac{n^2}{4} - O(n \log n) = \Omega(n^2)$.

For K_n, Li proves $\mathrm{sd}(K_n) = \mathrm{spum}(K_n) = 4n-6$ for $n \geq 2$ and $\mathrm{isd}(K_n) = \mathrm{spum}(K_n) = 4n-6$ for all $n \geq 2$ and $\mathrm{isd}(K_n) = \mathrm{ispum}(K_n) = n-1$ for $n \leq 3$ and $\mathrm{ispum}(K_n) = 4n-6$ when $n \geq 4$.

For cycles Li proves: for all $n \geq 4$, $2n-2 \leq \mathrm{sd}(C_n) \leq 2n-1$ and $2n-5 \leq \mathrm{isd}(C_n) \leq 2n-1$.

The current best bounds for $\mathrm{ispum}(P_n)$ are due to Singla, Tiwari, and Tripathi [3595]: $2n-5 \leq \mathrm{ispum}(P_n) \leq \frac{5}{2}(n-3)$ if n is odd and $2n-5 \leq \mathrm{ispum}(P_n) \leq 2n-3$ if n is even. For the sum-diameter, Li obtains a much tighter result that for $n \geq 3$, we have $2n-3 \leq \mathrm{sd}(P_n) \leq 2n-2$. An exhaustive computer search yields that $\mathrm{sd}(P_n) = 2n-3$ for $3 \leq n \leq 6$ and $\mathrm{sd}(P_n) = 2n-2$ for $7 \leq n \leq 13$. This led Li to conjecture that $\mathrm{sd}(P_n) = 2n-3$ for $3 \leq n \leq 6$ and $\mathrm{sd}(P_n) = 2n-2$ for $n \geq 7$. For $\mathrm{isd}(P_n)$ Li obtains: $2n-5 \leq \mathrm{isd}(P_n) \leq 2n-2$ if n is odd and $2n-5 \leq \mathrm{isd}(P_n) \leq 2n-3$ if n is even.

Li provides upper bounds on the sum-diameter for the disjoint union of two graphs and the join of two graphs as follows. For disjoint graphs G_1 and G_2 with no isolated vertices $\mathrm{sd}(G_1 \cup G_2) \leq 10 \max\{\mathrm{sd}(G_1), \mathrm{sd}(G_2)\} + \mathrm{sd}(G_1) + \mathrm{sd}(G_2) + 2$. If one wishes to add isolated vertices to a graph G, Li proves that if G is a graph with no isolated vertices, and N_k is the empty graph on k vertices, then $\mathrm{sd}(G \cup N_k) \leq \max\{k, 4\mathrm{sd}(G)\} + k - 5$. If one wishes to add a vertex with arbitrary edges incident to a graph G with no isolated vertices to form a new graph G' obtained by adding a vertex to G along with any desired edges incident to this new vertex, then $\mathrm{sd}(G') \leq 4\mathrm{sd}(G) - 1$. For two graphs G_1 and G_2 with no isolated vertices, where without loss of generality $\mathrm{sd}(G_1) \leq \mathrm{sd}(G_2)$, Li proves $\mathrm{sd}(G_1 + G_2) \leq 8\mathrm{sd}(G_2) + 11\mathrm{sd}(G_1) - 5$.

For a subset U of vertices from a graph $G(V, E)$ with no isolated vertices and $G[U]$ denoting the subgraph of G consisting of U and all of the edges connecting pairs of vertices

in U, Li proves that $\text{sd}(G[U]) \leq 2\text{sd}(G) - 2$. As an open question he asks is: For any induced subgraph $G[U]$ of G, is $\text{sd}(G[U]) \leq \text{sd}(G)$? For graphs obtained by deleting an edge e from a graph G with no isolated vertices (denoted by $G \setminus e$) or by contracting an edge e from G (denoted by G/e), Li proves that $\text{sd}(G \setminus e) \leq 4\text{sd}(G) - 1$ and $\text{sd}(G/e) \leq 4\text{sd}(G) - 1$. For a graph G with no isolated vertices and an edge $e = (u, v)$ between two vertices u and v of G that is not present in G, Li proves that $\text{sd}(G + e) \leq 4\text{sd}(G) - 1$.

Li concludes his paper by introducing the generalization of sum-diameter to hypergraphs, and generalizes his previous general upper and lower bounds to provide preliminary bounds on both sides for arbitrary k-uniform hypergraphs and with some remarks on areas for further research and some open questions.

He, Wang, Mi, Shen, and Yu [1389] say that a graph has a *tail* if the graph contains a path for which each interior vertex has degree 2 and an end vertex of degree at least 3. They prove that every tree with a tail of length at least 3 is an integral sum graph.

B. Xu [4227] has shown that the following are integral sum graphs: the union of any three stars; $T \cup K_{1,n}$ for all trees T; mK_3 for all m; and the union of any number of integral sum trees. Xu also proved that if $2G$ and $3G$ are integral sum graphs, then so is mG for all $m > 1$. Xu poses the question as to whether all disconnected forests are integral sum graphs. Nicholas and Somasundaram [2711] prove that all banana trees (see Section 2.1 for the definition) and the union of any number of stars are integral sum graphs.

Liaw, Kuo, and Chang [2260] proved that all caterpillars are integral sum graphs (see also [4212] and [4227] for some special cases of caterpillars). This shows that the assertion by Harary in [1369] that $K(1, 3)$ and $S(2, 2)$ are not integral sum graphs is incorrect. They also prove that all cycles except C_4 are integral sum graphs and they conjecture that every tree is an integral sum graph. Singh and Santhosh show that the crowns $C_n \odot K_1$ are integral sum graphs for $n \geq 4$ [3609] and that the subdivision graphs of $C_n \odot K_1$ are integral sum graphs for $n \geq 3$ [3308]. In 2000 Liaw, Kuo, and Chang [2260]conjectured that every tree is an integral sum graph. In 2024 [2610] Muthukkumar proved that all trees are integral sum graphs and that every bipartite graph is an induced subgraph of a sum graph with sum number 1. Wang, Li, and Wei [4133] proved that there exists a connected integral sum graph with any minimum degree and give an upper bound for the relation between the vertex number and the edge number of a connected integral sum graph with no saturated vertex.

For graphs with n vertices, Tiwari and Tripathi [3906] show that there exist sum graphs with m edges if and only if $m \leq \lfloor (n - 1^2)/4 \rfloor$ and that there exists integral sum graphs with m edges if and only if $m \leq \lceil 3(n-1)^2/8 \rceil + \lfloor (n-1)/2 \rfloor$, except for $m = \lceil 3(n-1)^2/8 \rceil + \lfloor (n-1)/2 \rfloor - 1$ when n is of the form $4k+1$. They also characterize sets of positive integers (respectively, integers) that are in bijection with sum graphs (respectively, integral sum graphs) of maximum size for a given order.

The *integral sum number*, $\zeta(G)$, of G is the minimum number of isolated vertices that must be added to G so that the resulting graph is an integral sum graph. Thus, by definition, G is a integral sum graph if and only if $\zeta(G) = 0$. Harary [1369] conjectured that $\zeta(K_n) = 2n - 3$ for $n \geq 4$. This conjecture was verified by Chen [797], by Sharary [3488], and by B. Xu [4227]. Yan and Liu proved: $\zeta(K_n - E(K_r)) = n - 1$ when $n \geq 6$, $n \equiv$

0 (mod 3) and $r = 2n/3 - 1$ [4241]; $\zeta(K_{m.m}) = 2m - 1$ for $m \geq 2$ [4241]; $\zeta(K_n \backslash -\text{edge}) = 2n - 4$ for $n \geq 4$ [4241], [4227]; if $n \geq 5$ and $n - 3 \geq r$, then $\zeta(K_n \backslash E(K_r)) \geq n - 1$ [4241]; if $\lceil 2n/3 \rceil - 1 > r \geq 2$, then $\zeta(K_n \backslash E(K_r)) \geq 2n - r - 2$ [4241]; and if $2 \leq m < n$, and $n = (i+1)(im - i + 2)/2$, then $\sigma(K_{m,n}) = \zeta(K_{m,n}) = (m-1)(i+1) + 1$ while if $(i+1)(im - i + 2)/2 < n < (i+2)[(i+1)m - i + 1]/2$, then $\sigma(K_{m,n}) = \zeta(K_{m,n}) = \lceil ((m-1)(i+1)(i+2) + 2n)/(2i+2) \rceil$ [4241]. Wang [4128] proved that $\sigma(K_{n+1} \backslash E(K_{1,r})) = \zeta(K_{n+1} \backslash E(K_{1,r})) = 2n - 2$ when $r + 1$, $2n - 3$ when $2 \leq r \leq n - 1$, and $2n - 4$ when $r = n$.

Nagamochi, Miller, and Slamin [2623] have determined upper and lower bounds on the sum number a graph. For most graphs $G(V, E)$ they show that $\sigma(G) = \Omega(|E|)$. He, Yu, Mi, Sheng, and Wang [1390] investigated $\zeta(K_n \backslash E(K_r))$ where $n \geq 5$ and $r \geq 2$. They proved that $\zeta(K_n \backslash E(K_r)) = 0$ when $r = n$ or $n - 1$;
$\zeta(K_n \backslash E(K_r)) = n - 2$ when $r = n - 2; \zeta(K_n \backslash E(K_r)) = n - 1$ when $n - 3 \geq r \geq \lceil 2n/3 \rceil - 1; \zeta(K_n \backslash E(K_r)) = 3n - 2r - 4$ when $\lceil 2n/3 \rceil - 1 > r \geq n/2; \zeta(K_n \backslash E(K_r)) = 2n - 4$ when $\lceil 2n/3 \rceil - 1 \geq n/2 > r \geq 2$. Moreover, they prove that if $n \geq 5, r \geq 2$, and $r \neq n - 1$, then $\sigma(K_n \backslash E(K_r)) = \zeta(K_n \backslash E(K_r))$.

Dou and Gao [973] prove that for $n \geq 3$, the fan $F_n = P_n + K_1$ is an integral sum graph, $\rho(F_4) = 1, \rho(F_n) = 2$ for $n \neq 4$, and $\sigma(F_4) = 2, \sigma(F_n) = 3$ for $n = 3$ or $n \geq 6$ and n even, and $\sigma(F_n) = 4$ for $n \geq 6$ and n odd.

Wang and Gao [4129] and [4130] determined the sum numbers and the integral sum numbers of the complements of paths, cycles, wheels, and fans as follows: $0 = \zeta(\overline{P_4}) < \sigma(\overline{P_4}) = 1$; $1 = \zeta(\overline{P_5}) < \sigma(\overline{P_5}) = 2$; $3 = \zeta(\overline{P_6}) < \sigma(\overline{P_6}) = 4$; $\zeta(\overline{P_n}) = \sigma(\overline{P_n}) = 0$, $n = 1, 2, 3$; $\zeta(\overline{P_n}) = \sigma(\overline{P_n}) = 2n - 7$, $n \geq 7$. $\zeta(\overline{C_n}) = \sigma(\overline{C_n}) = 2n - 7$, $n \geq 7$. $\zeta(\overline{W_n}) = \sigma(\overline{W_n}) = 2n - 8$, $n \geq 7$. $0 = \zeta(\overline{F_5}) < \sigma(\overline{F_5}) = 1$; $2 = \zeta(\overline{F_6}) < \sigma(\overline{F_6}) = 3$; $\zeta(\overline{F_n}) = \sigma(\overline{F_n}) = 0, n = 3, 4$; $\zeta(\overline{F_n}) = \sigma(\overline{F_n}) = 2n - 8$, $n \geq 7$.

Wang, Yang and Li [4134] proved: $\zeta(K_n \backslash E(C_{n-1}) = 0$ for $n = 4, 5, 6, 7$; $\zeta(K_n \backslash E(C_{n-1}) = 2n - 7$ for $n \geq 8$; $\sigma(K_4 \backslash E(C_{n-1}) = 1$; $\sigma(K_5 \backslash E(C_{n-1}) = 2$; $\sigma(K_6 \backslash E(C_{n-1}) = 5$; $\sigma(K_7 \backslash E(C_{n-1}) = 7$; $\sigma(K_n \backslash E(C_{n-1}) = 2n - 7$ for $n \geq 8$.

Wang and Li [4132] proved: a graph with $n \geq 6$ vertices and degree greater than $(n+1)/2$ is not an integral sum graph; for $n \geq 8$, $\zeta(K_n \setminus E(2P_3)) = \sigma(K_n \setminus E(2P_3)) = \epsilon(K_n \setminus E(2P_3)) = \epsilon(K_n \setminus E(2P_3)) = 2n - 7$; for $n \geq 7$, $\zeta(K_n \setminus E(K_2)) = \sigma(K_n \setminus E(K_2)) = 2n - 4$; and for $n \geq 7$ and $1 \leq r \leq \lceil \frac{n}{2} \rceil$, $\zeta(K_n \setminus E(rK_2)) = \sigma(K_n \setminus E(rK_2)) = 2n - 5$.

Chen [797] has given some properties of integral sum labelings of graphs G with $\Delta(G) < |V(G)| - 1$ whereas Nicholas, Somasundaram, and Vilfred [2713] provided some general properties of connected integral sum graphs G with $\Delta(G) = |V(G)| - 1$. They have shown that connected integral sum graphs G other than K_3 with the property that G has exactly two vertices of maximum degree are unique and that a connected integral sum graph G other than K_3 can have at most two vertices with degree $|V(G)| - 1$ (see also [4110]).

Vilfred and Florida [4107] have examined one-point unions of pairs of small complete graphs. They show that the one-point union of K_3 and K_2 and the one-point union of K_3 and K_3 are integral sum graphs whereas the one-point union of K_4 and K_2 and the one-point union of K_4 and K_3 are not integral sum graphs. In [4108] Vilfred and Florida

defined and investigated properties of maximal integral sum graphs.

Vilfred and Nicholas [4111] have shown that the following graphs are integral sum graphs: banana trees, the union of any number of stars, fans $P_n + K_1$ $(n \geq 2)$, Dutch windmills $K_3^{(m)}$, and the graph obtained by starting with any finite number of integral sum graphs $G_1, G_2, \ldots, G_n$ and any collections of n vertices with $v_i \in G_i$ and creating a graph by identifying $v_1, v_2, \ldots, v_n$. The same authors [4112] also proved that $G + v$ where G is a union of stars is an integral sum graph.

Melnikov and Pyatkin [2484] have shown that every 2-regular graph except C_4 is an integral sum graph and that for every positive integer r there exists an r-regular integral sum graph. They also show that the cube is not an integral sum graph. For any integral sum graph G, Melnikov and Pyatkin define the *integral radius of* G as the smallest natural number $r(G)$ that has all its vertex labels in the interval $[-r(G), r(G)]$. For the family of all integral sum graphs of order n they use $r(n)$ to denote maximum integral radius among all members of the family. Two questions they raise are: Is there a constant C such that $r(n) \leq C_n$ and for $n > 2$, is $r(n)$ equal to the $(n-2)$th prime?

The concepts of sum number and integral sum number have been extended to hypergraphs. Sonntag and Teichert [3678] prove that every hypertree (i.e., every connected, non-trivial, cycle-free hypergraph) has sum number 1 provided that a certain cardinality condition for the number of edges is fulfilled. In [3679] the same authors prove that for $d \geq 3$ every d-uniform hypertree is an integral sum graph and that for $n \geq d + 2$ the sum number of the complete d-uniform hypergraph on n vertices is $d(n-d)+1$. They also prove that the integral sum number for the complete d-uniform hypergraph on n vertices is 0 when $d = n$ or $n-1$ and is between $(d-1)(n-d-1)$ and $d(n-d)+1$ for $d \leq n-2$. They conjecture that for $d \leq n-2$ the sum number and the integral sum number of the complete d-uniform hypergraph are equal. Teichert [3874] proves that hypercycles have sum number 1 when each edge has cardinality at least 3 and that hyperwheels have sum number 1 under certain restrictions for the edge cardinalities. (A *hypercycle* $\mathcal{C}_n = (\mathcal{V}_n, \mathcal{E}_n)$ has $\mathcal{V}_n = \cup_{i=1}^{n}\{v_1^i, v_2^i, \ldots, v_{d_i-1}^i\}, \mathcal{E}_n = \{e_1, e_2, \ldots, e_n\}$ with $e_i = \{v_1^i, \ldots, v_{d_i}^i = v_1^{i+1}\}$ where $i+1$ is taken modulo n. A *hyperwheel* $\mathcal{W}_n = (\mathcal{V}_n', \mathcal{E}_n')$ has $\mathcal{V}_n' = \mathcal{V}_n \cup \{c\} \cup_{i=1}^{n} \{{v_2}^{n+i}, \ldots, {v_{d_{n+i}-1}}^{n+i}\}, \mathcal{E}_n' = \mathcal{E}_n \cup \{e_{n+1}, \ldots, e_{2n}\}$ with $e_{n+i} = \{{v_1}^{n+i} = c, {v_2}^{n+i}, \ldots, {v_{d_{n+i}-1}}^{n+i}, {v_{d_{n+i}}}^{n+i} = {v_1}^i\}$.)

Teichert [3873] determined an upper bound for the sum number of the d-partite complete hypergraph $K^d_{n_1,\ldots,n_d}$. In [3875] Teichert defines the *strong hypercycle* $\mathcal{C}_n^d$ to be the d-uniform hypergraph with the same vertices as C_n where any d consecutive vertices of C_n form an edge of $\mathcal{C}_n^d$. He proves that for $n \geq 2d+1 \geq 5$, $\sigma(\mathcal{C}_n^d) = d$ and for $d \geq 2$, $\sigma(\mathcal{C}_{d+1}^d) = d$. He also shows that $\sigma(\mathcal{C}_5^3) = 3$; $\sigma(\mathcal{C}_6^3) = 2$, and he conjectures that $\sigma(\mathcal{C}_n^d) < d$ for $d \geq 4$ and $d + 2 \leq n \leq 2d$.

In [2714] Nicholas and Vilfred define the *edge reduced sum number* of a graph as the minimum number of edges whose removal from the graph results in a sum graph. They show that for K_n, $n \geq 3$, this number is $(n(n-1)/2 + \lfloor n/2 \rfloor)/2$. They ask for a characterization of graphs for which the edge reduced sum number is the same as its sum number. They conjecture that an integral sum graph of order p and size q exists if and only if $q \leq 3(p^2-1)/8 - \lfloor (p-1)/4 \rfloor$ when p is odd and $q \leq 3(3p-2)/8$ when p is

even. They also define the *edge reduced integral sum number* in an analogous way and conjecture that for K_n this number is $(n-1)(n-3)/8 + \lfloor (n-1)/4 \rfloor$ when n is odd and $n(n-2)/8$ when n is even.

For certain graphs G Vilfred and Florida [4106] investigated the relationships among $\sigma(G), \zeta(G), \chi(G)$, and $\chi'(G)$ where $\chi(G)$ is the chromatic number of G and $\chi'(G)$ is the edge chromatic number of G. They prove: $\sigma(C_4) = \zeta(C_4) > \chi(C_4) = \chi'(C_4)$; for $n \geq 3$, $\zeta(C_{2n}) < \sigma(C_{2n}) = \chi(C_{2n}) = \chi'(C_{2n})$; $\zeta(C_{2n+1}) < \sigma(C_{2n+1}) < \chi(C_{2n+1}) = \chi'(C_{2n+1})$; for $n \geq 4$, $\chi'(K_n) \leq \chi(K_n) < \zeta(K_n) = \sigma(K_n)$; and for $n \geq 2$, $\chi(P_n \times P_2) < \chi'(P_n \times P_2) = \zeta(P_n \times P_2) = \sigma(P_n \times P_2)$.

Alon and Scheinermann [202] generalized sum graphs by replacing the condition $f(x) + f(y) \in S$ with $g(f(x), f(y)) \in S$ where g is an arbitrary symmetric polynomial. They called a graph with this property a *g-graph* and proved that for a given symmetric polynomial g not all graphs are g-graphs. On the other hand, for every symmetric polynomial g and every graph G there is some vertex labeling such that G together with at most $|E(G)|$ isolated vertices is a g-graph.

Boland, Laskar, Turner, and Domke [685] investigated a modular version of sum graphs. They call a graph $G(V, E)$ a *mod sum graph* (MSG) if there exists a positive integer n and an injective labeling from V to $\{1, 2, \ldots, n-1\}$ such that $xy \in E$ if and only if $(f(x)+f(y)) \pmod n = f(z)$ for some vertex z. Obviously, all sum graphs are mod sum graphs. However, not all mod sum graphs are sum graphs. Boland et al. [685] have shown the following graphs are MSG: all trees on 3 or more vertices; all cycles on 4 or more vertices; and $K_{2,n}$. They further proved that K_p $(p \geq 2)$ is not MSG (see also [1272]) and that W_4 is MSG. They conjecture that W_p is MSG for $p \geq 4$. This conjecture was refuted by Sutton, Miller, Ryan, and Slamin [3840] who proved that for $n \neq 4$, W_n is not MSG (the case where n is prime had been proved in 1994 by Ghoshal, Laskar, Pillone, and Fricke [1272]. In the same paper Sutton et al. also showed that for $n \geq 3$, $K_{n,n}$ is not MSG. Ghoshal, Laskar, Pillone, and Fricke [1272] proved that every connected graph is an induced subgraph of a connected MSG graph and any graph with n vertices and at least two vertices of degree $n-1$ is not MSG.

In [635] Beste, de Wiljes, and Kreh investigated the sum and mod sum graphs using arithmetic progressions and prime numbers and characterized the induced sum graphs and mod sum graphs.

Sutton, Miller, Ryan, and Slamin [3840] define the *mod sum number*, $\rho(G)$, of a connected graph G to be the least integer r such that $G \cup \overline{K_r}$ is MSG. Recall the cocktail party graph $H_{m,n}$, $m, n \geq 2$, as the graph with a vertex set $V = \{v_1, v_2, \ldots, v_{mn}\}$ partitioned into n independent sets $V = \{I_1, I_2, \ldots, I_n\}$ each of size m such that $v_i v_j \in E$ for all $i, j \in \{1, 2, \ldots, mn\}$ where $i \in I_p$, $j \in I_q$, $p \neq q$. The graphs $H_{m,n}$ can be used to model relational database management systems (see [3836]). Sutton and Miller [3838] prove that $H_{m,n}$ is not MSG for $n > m \geq 3$ and $\rho(K_n) = n$ for $n \geq 4$. In [3837] Sutton, Draganova, and Miller prove that for n odd and $n \geq 5$, $\rho(W_n) = n$ and when n is even, $\rho(W_n) = 2$. Wang, Zhang, Yu, and Shi [4166] proved that fan $F_n (n \geq 2)$ are not mod sum graphs and $\rho(F_n) = 2$ for even n at least 6. They also prove that $\rho(K_{n,n}) = n$ for $n \geq 3$.

Dou and Gao [974] obtained exact values for $\rho(K_{m,n})$ and $\rho(K_m - E(K_n))$ for some cases of m and n and bounds in the remaining cases. They call a graph $G(V, E)$ a *mod integral sum graph* if there exists a positive integer n and an injective labeling from V to $\{0, 1, 2, \ldots, n-1\}$ (note that 0 is included) such that $xy \in E$ if and only if $(f(x) + f(y)) \pmod n = f(z)$ for some vertex z. They define the *mod integral sum number*, $\psi(G)$, of a connected graph G to be the least integer r such that $G \cup \overline{K_r}$ is a mod integral sum graph. They prove that for $m+n \geq 3$, $\psi(K_{m,n}) = \rho(K_{m,n})$ and obtained exact values for $\psi(K_m - E(K_n))$ for some cases of m and n and bounds in the remaining cases.

Wallace [4119] has proved that $K_{m,n}$ is MSG when n is even and $n \geq 2m$ or when n is odd and $n \geq 3m-3$ and that $\rho(K_{m,n}) = m$ when $3 \leq m \leq n < 2m$. He also proves that the complete m-partite $K_{n_1,n_2,\ldots,n_m}$ is not MSG when there exist n_i and n_j such that $n_i < n_j < 2n_i$. He poses the following conjectures: $\rho(K_{m,n}) = n$ when $3m-3 > n \geq m \geq 3$; if $K_{n_1,n_2,\ldots,n_m}$ where $n_1 > n_2 > \cdots > n_m$, is not MSG, then $(m-1)n_m \leq \rho(K_{n_1,n_2,\ldots,n_m}) \leq (m-1)n_1$; if G has n vertices, then $\rho(G) \leq n$; and determining the mod sum number of a graph is NP-complete (Sutton has observed that Wallace probably meant to say 'NP-hard'). Miller [2492] has asked if it is possible for the mod sum number of a graph G be of the order $|V(G)|^2$.

In a sum graph G, a vertex w is called a *working vertex* if there is an edge uv in G such that $w = u+v$. If $G = H \cup \overline{H_r}$ has a sum labeling such that H has no working vertex the labeling is called an *exclusive sum labeling of H with respect G*. The *exclusive sum number*, $\epsilon(H)$, of a graph H is the smallest integer r such that $G \cup \overline{K_r}$ has an exclusive sum labeling. The exclusive sum number is known in the following cases (see [2496] and [2506]): for $n \geq 3$, $\epsilon(P_n) = 2$; for $n \geq 3$, $\epsilon(C_n) = 3$; for $n \geq 3$, $\epsilon(K_n) = 2n-3$; for $n \geq 4$, $\epsilon(F_n) = n$ (fan of order $n+1$); for $n \geq 4$, $\epsilon(W_n) = n$; $\epsilon(C_3^{(n)}) = 2n$ (friendship graph–see §2.2); $m \geq 2$, $n \geq 2$, $\epsilon(K_{m,n}) = m+n-1$; for $n \geq 2$, $S_n = n$ (star of order $n+1$); $\epsilon(S_{m,n}) = \max\{m, n\}$ (double star); $H_{2,n} = 4n-5$ (cocktail party graph); and $\epsilon(\text{caterpillar } G) = \Delta(G)$. Dou [972] showed that $H_{m,n}$ is not a mod sum graph for $m \geq 3$ and $n \geq 3$; $\rho(H_{m,3}) = m$ for $m \geq 3$; $H_{m,n} \cup \rho(H_{m,n})K_1$ is exclusive for $m \geq 3$ and ≥ 4; and $m(n-1) \leq \rho(H_{m,n}) \leq mn(n-1)/2$ for $m \geq 3$ and $n \geq 4$. Vilfred and Florida [4109] proved that $\epsilon(P_3 \times P_3) = 4$ and $\epsilon(P_n \times P_2) = 3$. In [1430] Hegde and Vasudeva provide an $O(n^2)$ algorithm that produces an exclusive sum labeling of a graph with n vertices given its adjacency matrix. In [3882] Thakare and Acharya provided an optimal exclusive sum labeling for prism graph D_n and prove that $\epsilon(D_n) = 5$ for $n \geq 3$.

In 2001 Kratochvil, Miller, and Nguyen proved that $\sigma(G \cup H) \leq \sigma(G) + \sigma(H) - 1$. In 2003 Miller, Ryan, Slamin, Sugeng, and Tuga [2502] posed the problem of finding the exclusive sum number of the disjoint union of graphs. In 2010 Wang and Li [4131] proved the following. Let G_1 and G_2 be graphs without isolated vertices, L_i be an exclusive sum labeling of $G_i \cup \epsilon(G_i)K_1$, and C_i be the isolated set of L_i for $i = 1$ and 2. If $\max C_1$ and $\min C_2$ are relatively prime, then $\epsilon(G_1 \cup G_2) \leq \epsilon(G_1) + \epsilon(G_2) - 1$. Wang and Li also proved the following: $\epsilon(K_{r,s}) = s+r-1$; $\epsilon(K_{r,s} - E(K_2)) = s-1$; for $s \geq r \geq 2$, $\epsilon(K_{r,s} - E(rK_2)) = s+r-3$. For $n \geq 5$ they prove: $\epsilon(K_n - E(K_n)) = 0$; $\epsilon(K_n - E(K_{n-1})) = n-1$; for $2 \leq r < n/2$, $\epsilon(K_n - E(K_r)) = 2n-4$; for $n/2 \leq r \leq n-2$, $\epsilon(K_n - E(K_r)) = 3n-2r-4$, and $\epsilon(C_n \odot K_1)$ is 3 or 4. They show that $\epsilon(C_3 \odot K_1) = 3$ and guess that for $n \geq 4$ the

class E_k of hypergraphs with a k-exclusive sum labeling is hereditary, but nontrivial to characterize even for $k = 1$. In [3057] Purcell, Ryan, Ryjáček, and Skyvová provided a complete description of the minimal forbidden induced subhypergraphs of E_1 that are 3-uniform with maximum vertex degree 2. They also show that every hypertree has a 1-exclusive sum labeling and every combinatorial design does not.

A survey of exclusive sum labelings of graphs is given by Ryan in [3217].

If $\epsilon(G) = \Delta(G)$, then G is said to be an Δ-*optimum summable graph.* An exclusive sum labeling of a graph G using $\Delta(G)$ isolates is called a Δ-*optimum exclusive sum labeling* of G. Tuga, Miller, Ryan, and Ryjáček [3921] show that some families of trees that are Δ-optimum summable and some that are not. They prove that if G is a tree that has at least one vertex that has two or more neighbors that are not leaves then $\epsilon(G) = \Delta(G)$.

Koh, Miller, Smyth, and Wang [1975] show the following: the graphs obtained by identifying one end of a q-path with a vertex of a p-cycle are 1-optimum summable, and that two of these graphs can be joined via a new edge to create a 2-optimum summable graph; generalized θ-graphs are 2-optimum summable; $\theta(p, q, r)$ which consists of a pair of vertices joined by 3 independent paths of lengths p, q and r (with a few small exceptions) are 2-optimum summable; there exists a 3-optimum summable graph of order $4l + 3$ for all $l \geq 1$; how to construct for all $k \geq 4$ a k-optimum summable graph; and if G is a k-optimum summable graph of order n, then $n \geq 2k$.

In [1577] Javaid, Khalid, Ahmad, and Imran introduce a weaker version of sum labeling of graphs as follows. Let $H = (V, E)$ be a simple, finite, undirected graph with $|V| = p$. H is a *weak sum* graph if there exists a labeling L (called a w-sum) of the vertices of V by distinct positive integers such that $(u, v) \in E$ if there exists a vertex $w \in V$ such that $L(w) = L(u) + L(v)$. (A sum graph also requires the "only if" condition). If H is a w-sum graph with the additional constraint that the labels L all fall in the range $1, \ldots, p$, then H is called a *super weak sumgraph* (sw-sumgraph). Because sumgraphs must have isolated vertices we may write $H = G + K_\delta$, where G is connected and K_δ denotes δ isolated vertices If δ is a minimum with respect to G, we say that the sumgraph (respectively, w-sumgraph, sw-sumgraph) H is δ-optimal and that G is δ-optimal summable (respectively, w-summable, sw-summable). Javaid et al. prove: paths are 1-optimal sw-summable; cycles are 2-optimal sw-summable; wheels are 3-optimal sw-summable; K_n is $(n-1)$-optimal sw-summable; and $G = K_{n_1,n_2,\ldots,n_q}$ are t-optimal sw-summable, where t is the minimum degree of any vertex in G. They also prove that for $n \geq 5$, the Cayley graph $\mathrm{Cay}(\mathbb{Z}_n, \pm 1, \pm 2)$ is 4-optimal w-summable. They conjecture that all connected graphs are δ-optimal w-summable for some δ. See also [1975] and [2502].

Grimaldi [1340] has investigated labeling the vertices of a graph $G(V, E)$ with n vertices with distinct elements of the ring Z_n so that $xy \in E$ whenever $(x + y)^{-1}$ exists in Z_n.

In his 2001 Ph. D. thesis Sutton [3836] introduced two methods of graph labelings with applications to storage and manipulation of relational database links specifically in mind. He calls a graph $G = (V_p \cup V_i, E)$ a *sum* graph* of $G_p = (V_p, E_p)$ if there is an injective labeling λ of the vertices of G with non-negative integers with the property that $uv \in E_p$ if and only if $\lambda(u) + \lambda(v) = \lambda(z)$ for some vertex $z \in G$. The *sum* number*, $\sigma^*(G_p)$, is the minimum cardinality of a set of new vertices V_i such that there exists a

sum* graph of G_p on the set of vertices $V_p \cup V_i$. A *mod sum* graph* of G_p is defined in the identical fashion except the sum $\lambda(u) + \lambda(v)$ is taken modulo n where the vertex labels of G are restricted to $\{0, 1, 2, \ldots, n-1\}$. The *mod sum* number*, $\rho^*(G_p)$, of a graph G_p is defined in the analogous way. Sum* graphs are a generalization of sum graphs and mod sum* graphs are a generalization of mod sum graphs. Sutton shows that every graph is an induced subgraph of a connected sum* graph. Sutton [3836] poses the following conjectures: $\rho(H_{m,n}) \leq mn$ for $m, n \geq 2$; $\sigma^*(G_p) \leq |V_p|$; and $\rho^*(G_p) \leq |V_p|$.

The following table summarizes what is known about sum graphs, mod sum graphs, sum* graphs, and mod sum* graphs is reproduced from Sutton's Ph. D. thesis [3836]. It was updated by J. Gallian in 2006. A question mark indicates the value is unknown. The results on sum* and mod sum* graphs are found in [3836].

Table 20: **Summary of Sum Graph Labelings**

Graph	$\sigma(G)$	$\rho(G)$	$\sigma^*(G)$	$\rho^*(G)$
$K_2 = S_1$	1	1	0	0
stars, $S_n,\ n \geq 2$	1	0	0	0
trees $T_n,\ n \geq 3$ when $T_n \neq S_n$	1	0	1	0
C_3	2	1	1	0
C_4	3	0	2	0
$C_n,\ n > 4$	2	0	2	0
W_4	4	0	2	0
$W_n,\ n \geq 5,\ n$ odd	n	n	2	0
$W_n,\ n \geq 6,\ n$ even	$\frac{n}{2} + 2$	2	2	0
fan, F_4,	2	1	1	0
fans, $F_n,\ n \geq 5,\ n$ odd	?	2	1	0
fans, $F_n,\ n \geq 6,\ n$ even	3	2	1	0
$K_n,\ n \geq 4$	$2n - 3$	n	$n - 2$	0
cocktail party graphs, $H_{2,n}$	$4n - 5$	0	?	0
$C_n^{(t)}\ (n,t) \neq (4,1)$ (see §2.2)	2	?	?	?
$K_{n,n}$	$\lceil \frac{4n-3}{2} \rceil$	$n (n \geq 3)$	?	?
$K_{m,n},\ 2nm \geq n \geq 3$	?	n	?	?
$K_{m,n}\ m \geq 3n - 3,\ n \geq 3,\ m$ odd	?	0	?	0
$K_{m,n},\ m \geq 2n,\ n \geq 3,\ m$ even	?	0	?	0
$K_{m,n},\ m < n$ $k = \lceil \sqrt{1 + (8m + n - 1)(n - 1)/2}\ \rceil$	$\lceil \frac{kn-k}{2} + \frac{m}{k-1} \rceil$	?	?	?
$K_{n,n} - E(nK_2),\ n \geq 6$	$2n - 3$	$n - 2$	?	?

7.2 Prime and Vertex Prime Labelings[4]

The notion of a prime labeling originated with Entringer and was introduced in a paper by Tout, Dabboucy, and Howalla [3910]. A graph with vertex set V is said to have a *prime labeling* if its vertices are labeled with distinct integers $1, 2, \ldots, |V|$ such that for each edge xy the labels assigned to x and y are relatively prime. Around 1980, Entringer conjectured that all trees have a prime labeling. Little progress was made on this conjecture until 2011 when Haxell, Pikhurko, Taraz [1387] proved that all large trees are prime. Also, their method allowed them to determine the smallest size of a non-prime connected order-n graph for all large n, proving a conjecture of Rao [3134] in this range. Among the classes of trees known to have prime labelings are: paths, stars, complete binary trees, spiders (i.e., trees with one vertex of degree at least 3 and with all other vertices with degree at most 2), olive trees (i.e., a rooted tree consisting of k branches such that the ith branch is a path of length i), all trees of order up to 50, palm trees (i.e., trees obtained by appending identical stars to each vertex of a path), banana trees, and binomial trees (the binomial tree B_0 of order 0 consists of a single vertex; the binomial tree B_n of order n has a root vertex whose children are the roots of the binomial trees of order $0, 1, 2, \ldots, n-1$ (see [3260], [2841], [3910], [1156], and [3187]). Tout, Dabboucy, and Howalla [3910] showed t-toe caterpillars (the internal vertices on the spine are regular in degree) are prime and that all caterpillars with maximum degree at most 5 are prime.

Seoud, Sonbaty, and Mahran [3402] provide necessary and sufficient conditions for a graph to be prime. They also give a procedure to determine whether or not a graph is prime. Other graphs with prime labelings include all cycles and the disjoint union of C_{2k} and C_n [927]. The complete graph K_n does not have a prime labeling for $n \geq 4$ and W_n is prime if and only if n is even (see [2230]). Lee, Wui, and Yeh [2230] proved that friendship graphs have prime labelings. Diefenderfer et al. [964] and [963] proved that the graph obtained by identifying a vertex of C_n with an endpoint of the star S_m where $1 \leq m \leq 9$, chains of C_n where $n = 4, 6$, or 8, $C_n \times P_2$ where $n - 1$ is prime and $n \geq 4$, generalized books $S_n \times P_m$ where $3 \leq m \leq 7$, and other families of unicylic graphs have prime vertex labelings. In [1902] Kanzariya, Chaurasia, and Rupani gave prime labelings for graphs obtained from the disjoint union of two copies of helms under various edge subdivision operations. These include subdivisions of edges incident to the apex vertex, edges of C_n, edges joining C_n to pendent vertices, and the barycentric subdivision of the helm graph. [1902] new

Seoud, Diab, and Elsakhawi [3376] have shown the following graphs are prime: fans; helms; flowers (see §2.2); stars; $K_{2,n}$; and $K_{3,n}$ unless $n = 3$ or 7. They also shown that $P_n + \overline{K_m}$ $(m \geq 3)$ is not prime. Berliner, Dean, Hook, Marr, Mbirka, and McBee give consecutive cyclic prime labelings of certain classes of ladders. Although $K_{n,n}$ does not have a prime labeling when $n > 2$, Berliner et al. give minimal prime labelings for all n-values $1 \leq n \leq 23$ and give conditions on m and n for which $K_{m,n}$ are prime. They provide specific values of n for m up to 13. Dissanayake, Abeysekara, Dhananjaya, Perera, and

[4]I am grateful to John Asplund and N. Bradley Fox for their helpful comments on the results in this section.

Ranasinghe [966] provide necessary and sufficient conditions for $K_{1,m,n}$ to have a prime labeling. In [659] Bigham, Donovan, Pack, Turley, and Wiglesworth [659] investigated the existence of prime labelings of certain snake graphs.

Kanzariya and Rupani [1903] proved that expanding the intermediate vertices with a path of a jewel graph without prime edges and jelly fish graphs without prime edges into paths results in graphs that admit prime labelings. They also showed zigzag graphs admit prime labelings. Karthik Chidambaram [1907] proved that the graphs obtained by replacing every edge of a star graph $K_{1,n}$ by C_m ($n \geq 1$ and $m \geq 4$) are prime graphs. [1903] new [1907] new

Tout, Dabboucy, and Howalla [3910] proved that $C_m \odot \overline{K_n}$ is prime for all m and n. Vaidya and Prajapati [3999] proved that the graphs obtained by duplication of a vertex by a vertex in P_n and $K_{1,n}$ are prime graphs and the graphs obtained by duplication of a vertex by an edge, duplication of an edge by a vertex, duplication of an edge by an edge in P_n, $K_{1,n}$, and C_n are prime graphs. They also proved that graph obtained by duplication of every vertex by an edge in P_n, $K_{1,n}$, and C_n are not prime graphs. Ghorbani and Kamali [1265] proved that ladders have prime labelings. In [2027] Kowsalya and Keerthika proved $C_3 \odot K_{1,n}$ and $C_4 \odot K_{1,n}$ are prime graphs. Vaithilingam and Meena [4037] and [2480]showed that certain crown related graphs as well as helms and certain helm related graphs are prime graphs. In [3078] Rahmadani, Aldiansyah, Pratiwi, Yunus, and Kusumasari proved that two classes of amalgamation of flower graphs are prime graphs. Triwahyuniti and Rahmadani [3917] proved that tensor products of P_2 and C_n and the tensor product of P_2 and P_n ($n \geq 2$) are prime graphs. [3917] new

The *prime* graph over a ring R, denoted by $PG(R)$, is a graph with $V(PG(R)) = R$ such that uv is an edge if and only if $uRv = \{0_R\}$ or $vRu = \{0_R\}$, for every vertex $u \neq v$. In [1959] Khuluq, Krisnawati, and Hidayat investigated Z_k-vertex-magic labelings of prime graphs over Z_n for $n = p$, $n = p^2$, and $n = pq$ for distinct primes p and q. [1959] new

In 2012 Ravi and Kala [3139] a introduced the notion of total prime graphs as follows. A graph $G(V, E)$ with p vertices and q edges is said to be a *total prime* graph if there is a bijection f from $V \cup E$ to $\{1, 2, 3, \ldots, p+q\}$ such that for each edge uv, $f(u)$ and $f(v)$ are relatively prime and, for each edge vertex v of degree at least 2, the greatest common divisor of the labels on the edges incident to v edges is 1. They proved that the following graphs are total prime graphs: paths, stars, bistars, even order cycles, helms, $K_{2,n}, C_3^n$, and fans. They also showed that cycles of order are not prime total.

Tharunraj and Ragukuma [3888] focused on converting non-prime graph classes to a prime graph classes by applying corona product with complete graph on one vertex. They proved the following graphs are prime: $P_n^2 \odot K_1$ ($n \geq 3$), $C_n^2 \odot K_1$ ($n \geq 4$), the corona product of the Möbius ladder M_n with K_1 for even $n \geq 6$, the corona product of graphs obtained by duplicating every vertex of P_n by an edge with K_1, the corona product of graphs obtained by duplicating every vertex of C_n by an edge with K_1, and the corona product graphs obtained by duplicating every vertex $K_{1,n}$ by an edge with K_1. For an arbitrary tree T, Tharunraj and Ragukumar [3887] developed an algorithm to construct a prime labeled tree T' such that T is a subtree of T'. They extended this algorithm for k arbitrary trees T_i, where $i = 1, 2, 3, \ldots, k$, to construct a larger prime labeled tree T' such that each T_i is a subtree of T'. They also developed a log-linear algorithm to construct a [3887] new

larger prime labeled tree T' from k prime trees T_i, where $i = 1, 2, 3, \ldots, k$, such that T_i is a subtree of T'. They proved the correctness of the proposed algorithms and obtained the time complexity of the proposed algorithms. "'

For m and n at least 3, Seoud and Youssef [3405] define $S_n^{(m)}$, the (m, n)-*gon star*, as the graph obtained from the cycle C_n by joining the two end vertices of the path P_{m-2} to every pair of consecutive vertices of the cycle such that each of the end vertices of the path is connected to exactly one vertex of the cycle. Seoud and Youssef [3405] have proved the following graphs have prime labelings: books; $S_n^{(m)}$; $P_n + \overline{K_2}$ if and only if $n = 2$ or n is odd; $C_n \odot K_1$ with a complete binary tree of order $2^k - 1$ ($k \geq 2$) attached at each pendent vertex, and that C_m-snakes are prime (see §2.2) for the definition). They also prove that every spanning subgraph of a prime graph is prime and every graph is a subgraph of a prime graph. They conjecture that all unicycle graphs have prime labelings. Diefenderfer, Hastings, Heath, Prawzinsky, Preston, White, and Whittemore [963] proved that certain families of graphs that are special cases of Seoud and Youssef's conjecture [3405] have prime labelings. Seoud and Youssef [3405] proved the following graphs are not prime: $C_m + C_n$; C_n^2 for $n \geq 4$; P_n^2 for $n = 6$ and for $n \geq 8$; and Möbius ladders M_n for n even (see §2.3 for the definition). They also give an exact formula for the maximum number of edges in a prime graph of order n and an upper bound for the chromatic number of a prime graph.

Youssef and Elsakhawi [4291] have shown: the union of stars $S_m \cup S_n$, are prime; the union of cycles and stars $C_m \cup S_n$ are prime; $K_m \cup P_n$ is prime if and only if m is at most 3 or if $m = 4$ and n is odd; $K_n \odot K_1$ is prime if and only if $n \leq 7$; $K_n \odot \overline{K_2}$ is prime if and only if $n \leq 16$; $6K_m \cup S_n$ is prime if and only if the number of primes less than or equal to $m + n + 1$ is at least m; and that the complement of every prime graph with order at least 20 is not prime. Michael and Youssef [2489] determined all self-complementary graphs that have prime labelings.

For positive integers m, k, q with $k \geq 3$ and $1 \leq q \leq \lfloor k/2 \rfloor$, the *uniform cycle snake* graph $C_{k,q}^m$ is constructed by taking a path with m edges and replacing each edge by a k-cycle by identifying two vertices at distance q in the cycle with the vertices of the original path edge.

Salmasian [3260] has shown that every tree with n vertices ($n \geq 50$) can be labeled with n integers between 1 and $4n$ such that every two adjacent vertices have relatively prime labels. Pikhurko [2841] has improved this by showing that for any $c > 0$ there is an N such that any tree of order $n > N$ can be labeled with n integers between 1 and $(1 + c)n$ such that labels of adjacent vertices are relatively prime.

Baskar Babujee and Vishnupriya [587] proved the following graphs have prime labelings: nP_2, $P_n \cup P_n \cup \cdots \cup P_n$, bistars (that is, the graphs obtained by joining the centers of two identical stars with an edge), and the graph obtained by subdividing the edge joining edge of a bistar. Baskar Babujee [564] obtained prime labelings for the graphs: $(P_m \cup nK_1) + \overline{K_2}$, $(C_m \cup nK_1) + \overline{K_2}$, $(P_m \cup C_n \cup \overline{K_r}) + \overline{K_2}$, $C_n \cup C_{n+1}$, $(2n-2)C_{2n}$ ($n > 1$), $C_n \cup mP_k$ and the graph obtained by subdividing each edge of a star once. In [574] Baskar Babujee and Jagadesh prove the following graphs have prime labelings: bistars B_m, n; $P_3 \odot K_{1,n}$; the union of $K_{1,n}$ and the graph obtained from $K_{1,n}$ by appending a

pendent edge to every pendent edge of $K_{1,n}$; and the graph obtained by identifying the center of $K_{1,n}$ with the two endpoints and the middle vertex of P_5.

In [3995] Vaidya and Prajapati prove the following graphs have prime labelings: a t-ply graph of prime order; graphs obtained by joining center vertices of wheels W_m and W_n to a new vertex w where m and n are even positive integers such that $m+n+3=p$ and p and $p-2$ are twin primes; the disjoint union of the wheel W_{2n} and a path; the graph obtained by identifying any vertex of a wheel W_{2n} with an end vertex of a path; the graph obtained from a prime graph of order n by identifying an end vertex of a path with the vertex labeled with 1 or n; the graph obtained by identifying the center vertices of any number of fans (that is, a "multiple shell"); the graph obtained by identifying the center vertices of m wheels $W_{n_1}, W_{n_2}, \ldots, W_{n_m}$ where each $n_i \geq 4$ is an even integer and each n_i is relatively prime to $2+\sum_{k=1}^{i-1} n_k$ for each $i \in \{2,3,\ldots,m\}$. Prajapati and Suther [3028] provided results about the existence of prime labelings of graphs obtained from $K_{2,n}$ by the duplication of vertices and edges. In [3001] Prajapati and Gajjar provided conditions under which the disjoint union of two graphs admit a prime labeling. They showed that $C_{2n+1} \times P_2$ is not prime, $\overline{W_n}$ is prime if and only if $3 \leq n \leq 6$, and, for a prime $p \geq 3$, $C_{p-1} \times P_2$ is prime and a wheel graph of odd order is switching invariant. In [3013] they proved that generalized Petersen graph $P(n,k)$ is prime then n must be even and k must be odd and found some classes of generalized Petersen graphs that admit prime labelings.

In [4204] Wilson and Jini define the torch O_n as the graph that has $n+4$ vertices and $2n+3$ edges with $V(O_n) = \{v-i \mid 1 \leq i \leq n+4\}$ and $E(O_n) = \{v_i v_{n+1} \mid 2 \leq i \leq n-2\} \cup \{v_i v_{n+3} \mid 2 \leq i \leq n-2\} \cup \{v_1 v_i \mid n \leq i \leq n+4\} \cup \{v_{n-1}v_n, v_n v_{n+2}, v_n v_{n+4}, v_{n+1}v_{n+3}\}$. They proved that the torch graph O_n is a prime graph.

The *Knödel graphs* $W_{\Delta,n}$ with n even and degree Δ, where $1 \leq \Delta \leq \lfloor \log_2 n \rfloor$ have vertices pairs (i,j) with $i = 1,2$ and $0 \leq j \leq n/2-1$ where for every $0 \leq j \leq n/2-1$ and there is an edge between vertex $(1,j)$ and every vertex $(2,(j+2^k-1) \bmod n/2)$, for $k = 0,1,\ldots,\Delta-1$. Haque, Lin, Yang, and Zhao [1367] have shown that $W_{3,n}$ is prime when $n \leq 130$.

In [1060] Ezhal investigated prime labelings for some bipartite and cycle related graphs and prime labelings the joint sum of paths joining of bipartite graphs and cycles. Abughazaleh and Abughneim [33] investigated the existence prime labelings of graphs obtained by connecting wheels to paths and cycles.

Samuel and Kalaivani wrote a series of papers about graphs that admit prime labelings. Among them are graphs in the context of duplication, fusion, switching in an octopus graphs [3269]; planter related graphs [3267]: vanessa related graphs [3271]; vanessa related graphs [3268]; graphs in the context of the graph operations duplication, fusion, switching in drums graphs; and graphs related to brush graph $P_n \odot P_2$. Ganesan, Bhaalamurugan, and Christy [1179] investigated prime labelings for some graphs related to jelly fish graphs, subdivision of H-graphs, path relaed graphs, lotus inside a circle graphs, and umbrella graphs.

[3269] new
[3267] new
[3271] new
[3268] new
[1179] new

In [3327] Schuchter and Wilson gave evidence for their conjecture that a generalized Petersen graph $P(n,k)$ is prime if and only if it is bipartite, which occurs for n even and

k. They show that it is true for all even n and odd k such that $n \leq 9000$ and $1 \leq k \leq \frac{n}{2}$. They conjectured that all cubic bipartite graphs with at least 8 vertices are prime and verified it for all such graphs G, connected or not, satisfying $8 \leq V(G) \leq 22$.

In the 24th edition of this survey, the following was stated. "Schroeder [3330] proved that every bipartite graph is prime except $K_{3,3}$. This result establishes that the generalized Petersen graph $P(n,k)$ is prime precisely when it is bipartite, the Knődel graph $W_{3,n}$ is prime for all even $n \geq 4$, and the union of any number of even cycles is prime." Schroeder's result about bipartite graphs was later modified [3331] to state that for $4 \leq n \leq 32$, cubic bipartite graphs on $2n$ vertices are prime. Consequently, the statements about the generalized Petersen graph, the Knődel graph, and the union of even cycles are open. In [2553] Mostafa and Ghorbani characterized Hamiltonian k-coprime graphs, which implied a conjecture of Schroeder in [3328] that all 2-regular graphs are prime if every union of even cycles is prime. They then cited Schroeder's original result to claim the 1982 conjecture by Tout, Dabboucy, Howalla [3910] that all 2-regular graphs are prime. So, that conjecture is still open as well.

Prajapati and Shah [3027] investigated the existence of prime labelings for graphs obtained by the duplication of paths, cycles, stars, and wheels. Sundaram, Ponraj, and Somasundaram [3791] investigated the prime labeling behavior of all graphs of order at most 6 and established that only one graph of order 4, one graph of order 5, and 42 graphs of order 6 are not prime.

Kansagara and Patel [1895] define the *web graph without a center* as the graph obtained by joining the n pendent vertices of $C_n \odot K_1$ to form an n-cycle and then adding a single pendent edge to each vertex of this outer cycle. They investigated prime labelings for web graphs without a center and graphs resulting from the subdivision of some specific edges in it. They also provide results on prime labeling of the union of a web graph without center with a wheel, generalized Jahangir graphs, and drum graphs.

Given a collection of graphs $G_1, \ldots, G_n$ and some fixed vertex v_i from each G_i, Lee, Wui, and Yeh [2230] define Amal$\{(G_i, v_i)\}$, the amalgamation of $\{(G_i, v_i)|\ i = 1, \ldots, n\}$, as the graph obtained by taking the union of the G_i and identifying $v_1, v_2, \ldots, v_n$. They proved *Amal*$\{(G_i, v_i)\}$ has a prime labeling when G_i are paths and when G_i are cycles. They also showed that the amalgamation of any number of copies of W_n, n odd, with a common vertex is not prime. They conjecture that for any tree T and any vertex v from T, the amalgamation of two or more copies of T with v in common is prime. They further conjecture that the amalgamation of two or more copies of W_n that share a common point is prime when n is even ($n \neq 4$). Vilfred, Somasundaram, and Nicholas [4115] have proved this conjecture for the case that $n \equiv 2 \pmod 4$ where the central vertices are identified.

Dean [905], and independently Ghorbani and Kamali [1265], showed that the ladder $P_n \times P_2$ has a prime labeling. In [874] Curran and Ollis pointed out a flaw in Dean's proof and showed that a stronger condition is needed for it to hold. They conjecture that this stronger condition is true. They also offer an alternative construction inspired by Dean's approach that shows that if the Even Goldbach Conjecture and a particular strengthening of Lemoine's Conjecture are true, then the Prime Ladder Conjecture follows.

Vilfred, Somasundaram, and Nicholas [4115] proved the following graphs are prime:

helms; $P_m \times P_n$ where n is prime, $m \leq 3$; double fans $P_n + \overline{K_2}$ if and only if n is odd; and cycles with a P_k-chord. They conjecture that $P_m \times P_n$ where $m < n$ and n is prime is prime and ladders $P_n \times P_2$ are prime. The conjecture about grids was proved by Sundaram, Ponraj, and Somasundaram [3789]. In the same article they also showed that $P_n \times P_n$ is prime when n is prime. Kanetkar [1876] proved: $P_6 \times P_6$ is prime; $P_{n+1} \times P_{n+1}$ is prime when n is a prime with $n \equiv 3$ or 9 (mod 10) and $(n+1)^2+1$ is also prime; and $P_n \times P_{n+2}$ is prime when n is an odd prime with $n \not\equiv 2 \pmod 7$. Curran [865] showed that $P_p \times P_n$ has a prime labeling for any odd prime p and any integer n such that $p < n \leq p^2$. Combining the results of Sundaram, Ponraj, and Somasundaram [3789] and Curran, we have that $P_p \times P_n$ has a prime labeling for any odd prime p and any integer n such that $1 \leq n \leq p^2$.

Seoud, El Sonbaty, and Abd El Rehim [3377] proved that for $m = p_{n+t-1} - (t+n)$ where p_i is the i^{th} prime number in the natural order $K_n \cup K_{t,m}$ is prime and graphs obtained from $K_{2,n},\ (n \geq 2)$ by adding p and q edges out from the two vertices of degree n of $K_{2,n}$ are prime. They also proved that if G is not prime, then $G \cup K_{1,n}$ is prime if $\pi(n+m+1) \geq m$ where m is the order of G and $\pi(x)$ is the number of primes less than or equal to x.

Recall that $C_n^{(k)}$ is the graph obtained from the $k \geq 2$ copies of the cycle C_n by identifying exactly one vertex of each of these k copies of C_n. Patel and Vasava [2804] proved the following: $C_n^{(j)} \cup C_m^{(k)}$ is a prime graph if and only if either n is even or m is even; $C_{2n}^{(2)} \cup C_{2m}^{(2)} \cup C_k^{(2)}$ is a prime graph for all n, m and k; $C_{2n} \cup C_{2n} \cup C_{2n} \cup C_{2n} \cup C_{2m} \cup C_k$ is a prime graph for all n, m and k; and $G = \left(\bigcup_{k=1}^{N} C_{n_k}^{(2)}\right) \cup \left(\bigcup_{j=1}^{M} C_{m_j}^{(2)}\right)$ is not a prime graph if $M \leq N-2$ They also provided conditions for which $G = C_{2n}^{(2)} \cup C_{2m+1}^{(2)} \cup C_{2k+1}^{(2)}$ is a prime graph.

For any finite collection $\{G_i, u_i v_i\}$ of graphs G_i, each with a fixed edge $u_i v_i$, Carlson [756] defines the edge amalgamation *Edgeamal*$\{(G_i, u_i v_i)\}$ as the graph obtained by taking the union of all the G_i and identifying their fixed edges. The case where all the graphs are cycles she calls *generalized books*. She proves that all generalized books are prime graphs. Moreover, she shows that graphs obtained by taking the union of cycles and identifying in each cycle the path P_n are also prime.

In [563] Baskar Babujee proves that the maximum number of edges in a simple graph with n vertices that has a prime labeling is $\sum_{k=2}^{n} \phi(k)$. He also shows that the planar graphs having n vertices and $3(n-2)$ edges (i.e., the maximum number of edges for a planar graph with n vertices) obtained from K_n $(n \geq 5)$ with vertices $v_1, v_2, \ldots, v_n$ by deleting the edges joining v_s and v_t for all s and t satisfying $3 \leq s \leq n-2$ and $s+2 \leq t \leq n$ has a prime labeling if and only if n is odd.

By showing that for every even $n \leq 2.468 \times 10^9$ there exists $1 \leq s \leq n-1$ such that both $n+s$ and $2n+s$ are prime, Schluchter, Schroeder, Cokus, Ellingson, Harris, Rarity, and Wilson [3326] prove the generalized Peterson graph $P(n,1)$ (which is isomorphic to $C_n \times P_2$) is prime for all even $4 \leq n \leq 2.468 \times 10^9$. For a fixed n they also describe a method for labeling $P(n,k)$ that is a prime labeling for multiple values of k. Using this method, they prove $P(n,k)$ is prime for all even $n \leq 50$ and odd $k < n/2$.

Yao, Cheng, Zhongfu, and Yao [4259] have shown: a tree of order p with maximum

degree at least $p/2$ is prime; a tree of order p with maximum degree at least $p/2$ has a vertex subdivision that is prime; if a tree T has an edge u_1u_2 such that the two components T_1 and T_2 of $T - u_1u_2$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$ and $d_{T_2}(u_2) \geq |T_2|/2$, then T is prime when $|T_1| + |T_2|$ is prime; if a tree T has two edges u_1u_2 and u_2u_3 such that the three components T_1, T_2, and T_3 of $T - \{u_1u_2, u_2u_3\}$ have the properties that $d_{T_1}(u_1) \geq |T_1|/2$, $d_{T_2}(u_2) \geq |T_2|/2$, and $d_{T_3}(u_3) \geq |T_3|/2$, then T is prime when $|T_1| + |T_2| + |T_3|$ is prime.

Vaidya and Prajapati [3996] define a *vertex switching* G_v of a graph G as the graph obtained by taking a vertex v of G, removing all the edges incident to v and adding edges joining v to every other vertex that is not adjacent to v in G. They say a prime graph G is *switching invariant* if for every vertex v of G, the graph G_v obtained by switching the vertex v in G is also a prime graph. They prove: P_n and $K_{1,n}$ are switching invariant; the graph obtained by switching the center of a wheel is a prime graph; and the graph obtained by switching a rim vertex of W_n is a prime graph if $n+1$ is a prime. They also prove that the graph obtained by switching a rim vertex in W_n is not a prime graph if $n+1$ is an even integer greater than 9.

Prajapati and Gajjar [3001] prove the following graphs are prime: graphs obtained from P_{m+1} and m copies of C_n by identifying each edge of P_{m+1} with an edge of a corresponding copy of C_n; graphs obtained from C_m and m copies of C_n by identifying each edge of C_m with an edge of corresponding copy of C_n; for a prime $p \geq 3$ and $p-2$ copies of C_{p+1}, the graph obtained by identifying one vertex of each copy of C_{p+1} with corresponding pendent vertex of $K_{1,p-2}$; for a prime $p \geq 3$, $C_{p-1} \times P_2$; and for a prime $p \geq 3$, the graphs obtained by joining every rim vertex of a wheel graph W_{p-1} with the corresponding vertex of C_{p-1}. They also prove that the complement of W_n is prime if and only if $3 \leq n \leq 6$; for odd $n \geq 3$ $C_n \times P_2$ is not prime; and W_{2n} is switching invariant.

Selvaraju and Moha [3350] proved that the one-point union of any number of cycles and the one-point union of any number of wheels at the center are prime graphs. Haque, Xiaohui, Yuansheng, and Pingzhong proved that the generalized Petersen graph $P(n,k)$ is prime for all even $n \leq 2500$ when $k = 1$ [1364] and for all even $n \leq 100$ when $k = 3$ [1366]. They show $P(n,3)$ is not prime for odd n and conjecture that $P(n,3)$ are prime for all even n.

In [3382] Seoud, El-Sonbaty, and Mahran discuss the primality of some corona graphs $G \odot H$ and conjecture that $K_n \odot \overline{K_m}$ is prime if and only if $n \leq \pi(nm+n)+1$, where $\pi(x)$ is the number of primes less than or equal to x. For $m \leq 20$ they give the exact values of n for which $K_n \odot \overline{K_m}$ is prime. They also show that $K_{m,n}$ is prime if and only if $\min\{m,n\} \leq \pi(m+n) - \pi((m+n)/2) + 1$.

In [2804] Patel and Vasava provided the following results about prime graphs: $W_n \cup C_m$ is a prime graph if and only if n and m both are even; $(P_n + \overline{K_2}) \cup C_m$ is a prime graph if and only if either $n = 2$, or n is odd and m is even; $C_n^{(2)} \cup C_m$ is a prime graph if and only if at least one of n and m is even; $S_n^{(m)} \cup S_k^{(j)}$ is not a prime graph if m and j are even and n and k are odd ($S_n^{(m)}$ denotes the (m,n)-gon star obtained from C_n and n copies of P_{m-2} by joining the two end vertices of P_{m-2} to each pair of consecutive vertices of the cycle such that each of the end vertices of the path is adjacent to exactly one vertex of

the cycle); $S_{2n}^{(2m)} \cup S_k^{(2m)}$ is a prime graph for all n, m, and k; $H_n \bigcup B_n$ is a prime graph for all n and m; $Circ(n,k)$ is not a prime graph when n and k both are even; and $Circ(n,k)$ is not a prime graph when n is odd.

In [2796] Patel and Kansagara prove that the following graphs are prime: $(C_n \odot K_1) \cup (C_n \odot K_1)$ for all n; $G_n \cup G_n$ for all n (G_n is the gear graph); $H_n \cup H_n$ for all n (H_n is the helm graph); $B_n \cup (K_{1,n} \times P_2)$ for all n; $H_n \cup G_n$ for all n; $(C_n \odot K_1) \cup G_n$ for all n; $(C_n \odot K_1) \cup H_n$ for all n; and $C_n(C_n) \cup C_n(C_n)$ if $n \not\equiv 1 \pmod 3$ ($C_n(C_n)$ is the graph obtained by taking barycentric subdivision of C_n and joining each newly inserted vertices of incident edges by an edge).

In [2798] Patel, Kansagava, and Vasva provided prime labeling of graphs obtained by taking barycentric subdivision of W_n, flower graphs, and $(C_n \odot K_1) \cup (C_n \odot K_1)$; and the graphs by obtained by taking subdivision of certain edges W_n, H_n, $(C_n \odot K_1)$ and $(C_n \odot K_1) \cup (C_n \odot K_1)$.

Hamm, Hamm, and Way [1358] proved that if a complete k-partite k-uniform hypergraph has enough vertices and every pod of vertices is large enough, then it does not have a prime labeling. They also proved that if a pod of vertices in a complete k-partite k-uniform is small enough, the graph it does have a prime labeling.

Klee, Lehmann, and Park [1969] we extended the notion of prime labeling to the Gaussian integers. They showed that paths, stars, spiders, graphs obtained by joining the centers of two stars with a path, and some firecrackers admit Gaussian prime labelings. In [3572] Shrimali and Singh proved the following graphs have Gaussian vertex prime labelings: books; kayaks; the one point union of k copies of W_n; the one point union of k copies of a gear graph; the one point union of k copies a graph obtained from a wheel by replacing each cycle edge by P_n; and the one point union of k copies a graph obtained from a wheel by replacing each spoke by P_n. Kavitha and Jayalalitha [1933] proved the following graphs admit Gaussian prime distance labeling: grids, paths, bipartite graphs, non-trivial Dutch windmills, non-trivial stars, double stars, and spider trees.

The Prime Ladder Conjecture states that every ladder $P_n \times P_2$ is prime. This was proved by Dean [905] in 2017. He conjectures that every integer $n \geq 50$ has a canonical partition with at most three terms and he states that this conjecture was verified by computer up to 5,000,000.

Lau, Chu, Suhadak, Foo, and Ng [2097] introduced SD-prime cordial labelings as follows. Given a finite, simple graph G with n vertices and a bijection $f : V(G) \to \{1, 2, \ldots, n\}$, for each edge uv let $S = f(u) + f(v)$ and $D = |f(u) - f(v)|$. For each edge uv define f' induced by f by assigning $f'(uv) = 1$ if $\gcd(S, D) = 1$ and $f'(uv) = 0$ otherwise. Then f' is said to be SD-*prime cordial* if $f'(uv) = 1$ for all edges uv. They provide results about paths, complete bipartite graphs, stars, double stars, wheels, fans, double fans, ladders, and grids. They conjecture that $P_m \times P_n$ is SD-prime for all $m \geq 2$ and $n \geq 2$. Lau, Shiu, Ng, and Jeyanthi [2114] give sufficient conditions for a theta graph to have an SD-prime cordial labeling, provide a way to construct new SD-prime cordial graphs from existing ones, and investigate SD-prime cordialness of some general graphs. Lourdusamy and Patrick [2336] provide a way to construct SD-prime cordial graphs from an existing graph G with an SD-prime cordial labeling by identifying a vertex of G having

a particular label with a vertex of maximum degree of a star or fan or with an endpoint of a path. In [2337] Lourdusamy and Patrick investigated SD-prime cordial labelings of subdivision graphs, splitting graphs, shadow graphs of stars and bistars, $T(P_n), T(C_n)$, the graph obtained by duplication of each vertex of a path and a cycle by an edge, Q_n, $A(T_n)$, triangular ladders, $P_n \odot K_1$, $C_n \odot K_1$, and jewel graphs. Shiu and Lau [3538] provided SD-prime labelings for some one-point unions of gear graphs.

The following definitions appear in [2951], [2938], [2939], and [2940]. A *double triangular snake* DT_n consists of two triangular snakes that have a common path; a *double quadrilateral snake* DQ_n consists of two quadrilateral snakes that have a common path; an *alternate triangular snake* $A(T_n)$ is the graph obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertex v_i (that is, every alternate edge of a path is replaced by C_3); a *snake graphs double alternate triangular snake* $DA(T_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to two new vertices v_i and w_i; an *alternate quadrilateral snake* $A(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i and w_i respectively and then joining v_i and w_i (that is, every alternate edge of a path is replaced by a cycle C_4); a *double alternate quadrilateral snake* $DA(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i, x_i and w_i and y_i respectively and then joining v_i and w_i and x_i and y_i.

Prajapati and Vantiya [3029] proved that the following snake graphs have SD-prime cordial labelings: triangular (except for $n = 3$), alternate triangular, quadrilateral, alternate quadrilateral, double triangular, double alternate triangular, double quadrilateral, and double alternate quadrilateral. Lourdusamy, Wency, and Patrick [2349] proved that the union of stars and paths, subdivision of combs, subdivision of ladders, and the graph obtained by attaching a star at one end of a path are SD-prime graphs. They proved that the union of two SD-prime cordial graphs need not be SD-prime cordial graphs. Also, they proved that given a positive integer n, there is SD-prime cordial graph with n vertices. In [3030] Prajapati and Vantiya investigated SD-prime cordial labelings of various kinds of snakes involving K_4. In [3031] they investigate the SD-prime cordial labeling of alternate k-polygonal snake graphs of type-1, type-2 and type-3.

Vaidya and Prajapati [3995] have introduced the notion of k-prime labeling. A *k-prime labeling* of a graph G is an injective function $f : V(G) \to \{k+1, k+2, k+3, \ldots, k+|V(G)|-1\}$ for some positive integer k that induces a function f^+ on the edges of G defined by $f^+(uv) = \gcd(f(u), f(v))$ such that $\gcd(f(u), f(v)) = 1$ for all edges uv. A graph that admits a k-prime labeling is called a *k-prime* graph. They prove the following are prime graphs: a tadpole (that is, a graph obtained by identifying a vertex of a cycle to an end vertex of a path); the union of a prime graph of order n and a $(n+1)$-prime graph; the graph obtained by identifying the vertex labeled with n in an n-prime graph with either of the vertices labeled with 1 or n in a prime graph of order n.

For connected graphs $G_1(p_1, q_1)$ and $G_2(p_2, q_2)$, $G_1 \hat{o} G_2$ is the graph obtained by superimposing any selected vertex of G_2 on any selected vertex of G_1. Then for any graph k-prime graph G, Vijayalashmi and Arockiamary [4094] proved the following graphs have k-prime labelings: $C_{2n} \cup C_{2n}$, $G \cup tP_m (m > 1)$, $G \cup K_{1,n}, G \hat{o} P_n, G \hat{o} K_{1,n}$ and $G \hat{o} (P_n + K_1)$.

Chaurasiya, Limda, Parmar, and Rupani [788] proved that $P_n \odot 3K_1, P_n \odot K_1$, ladders, and quadrilateral snakes admit k-prime labelings.

In [305] Teresa Arockiamary and Vijayalakshmi proved that the graphs obtained by identifying an endpoint of $r \geq 2$ copies of P_n $(n \geq 2)$ admit a vertex k-prime labelings for all positive integers k. Further, they make study on comparison between prime labelings and vertex k-prime labelings for the one point union of path graph using the C++ programming language. Arockiamary and Vijayalakshmi [301] proved that shell graphs, uniform shell butterfly graphs, and shell-flower graphs admit k-prime total labelings. In [304] Teresa Arockiamary and Vijayalakshmi introduced a generalization of a theta graph and gave conditions for which these graphs admit vertex k-prime labelings.

A dual of prime labelings has been introduced by Deretsky, Lee, and Mitchem [927]. They say a graph with edge set E has a *vertex prime labeling* if its edges can be labeled with distinct integers $1, \ldots, |E|$ such that for each vertex of degree at least 2 the greatest common divisor of the labels on its incident edges is 1. Deretsky, Lee, and Mitchem show the following graphs have vertex prime labelings: forests; all connected graphs; $C_{2k} \cup C_n$; $C_{2m} \cup C_{2n} \cup C_{2k+1}$; $C_{2m} \cup C_{2n} \cup C_{2t} \cup C_k$; and $5C_{2m}$. They further prove that a graph with exactly two components, one of which is not an odd cycle, has a vertex prime labeling and a 2-regular graph with at least two odd cycles does not have a vertex prime labeling. They conjecture that a 2-regular graph has a vertex prime labeling if and only if it does not have two odd cycles. Let $G = \bigcup_{i=1}^{t} C_{2n_i}$ and $N = \sum_{i=1}^{t} n_i$. In [690] Borosh, Hensley and Hobbs proved that there is a positive constant n_0 such that the conjecture of Deretsky et al. is true for the following cases: G is the disjoint union of at most seven cycles; G is a union of cycles all of the same even length $2n$ where $n \leq 150\,000$ or where $n \geq n_0$; $n_i \geq (\log N)^{4 \log\log\log n}$ for all $i = 1, \ldots, t$; and when each C_{2n_i} is repeated at most n_i times. They end their paper with a discussion of graphs whose components are all even cycles, and of graphs with some components that are not cycles and some components that are odd cycles. In [2129] Lavanya and Ganesa provided vertex prime labelings for graphs related to $C_n \odot K_1$.

In [481] Bapat proved the following graphs have vertex prime labelings: kayak paddles $KP(k, m, l)$; books; irregular books not necessarily with pages of the same size; triangular snakes; m-fold triangular snakes of length n obtained from a path $v_1, v_2, \ldots, v_n, v_{n+1}$ by joining v_i and v_{i+1} to new m vertices $w_1^i, w_2^i, \ldots, w_m^i$, for $i = 1, 2, \ldots, n$ giving edges $v_i w_j^i$ and $w_j^i v_{i+1)}$, for $j = 1, \ldots, m,\ i = 1, 2, \ldots, n$; m-fold petal sunflowers obtained from a cycle $v_1, v_2, \ldots, v_n$ by joining v_i and v_{i+1} to new m vertices $w_1^i, w_2^i, \ldots, w_m^i$, for $i = 1, 2, \ldots, n$ giving edges $v_i w_j^i$ and $w_j^i v_{i+1)}$ for $j = 1, \ldots, m,\ i = 1, 2, \ldots, n (v_{n+1} = v_1)$; and one-point unions of cycles not necessarily of the same length.

A bijection f from $V(G$ to $\{1, 2, \ldots, |V| + |E|\}$ is said to be a *total prime* if for each edge uv the labels assigned to u and v are relatively prime and for each vertex of degree at least 2, the labels on the incident edges are relatively prime. A graph that admits a total prime labeling is called a *total prime graph*. In [3010] Prajapati and Gajjar defined a *braided star* graph as follows. Let a_0 be the apex vertex and $a_1, a_2, \ldots, a_{n-1}, a_n$ be consecutive n rim vertices of W_n, $(n \geq 3)$. Let $b_1, b_2, b_3, \ldots, b_{2n-1}, b_{2n}$ be consecutive $2n$ vertices of C_{2n} $(n > 1)$; and let $c_1, c_2, c_3, \ldots, c_{2n-1}, c_{2n}$ be consecutive $2n$ vertices of a

second copy of C_{2n}. Join each a_i to b_{2i-1} by an edge and b_{2i} to c_{2i} by an edge. For each i, join a new vertices d_i to each c_{2i-1} and c_{2i+1} by an edge taking the subscripts modulo n. They proved that braided stars are prime, total prime, and vertex prime.

In [288] Arockiamary, Baskar Babujee, and Vijayalakshmi say a graph G has a *k-prime total labeling* if k is a positive integer and there is a bijection $f : V(G) \cup E(G) \to \{k, k+1, k+2, \ldots, k+V(G) \cup E(G)-1\}$ such that for every edge uv the labels $f(u), f(v)$, and $f(uv)$ are pairwise relatively prime. A graph that admits a k-prime total labeling is called a *k-prime total graph*. They proved that P_n is k-prime total for odd k, $K_{1,n}$ is k-prime total for odd k and $n \geq 2$, the bistar $B(m, n)$ is a $k-$prime total for odd k and $m,\ n \geq 2$, and various other tree-related graphs are k-prime total.

Jothi [1803] calls a graph G *highly vertex prime* if its edges can be labeled with distinct integers $\{1, 2, \ldots, |E|\}$ such that the labels assigned to any two adjacent edges are relatively prime. Such labeling is called a *highly vertex prime* labeling. He proves: if G is highly vertex prime then the line graph of G is prime; cycles are highly vertex prime; paths are highly vertex prime; K_n is highly vertex prime if and only if $n \leq 3$; $K_{1,n}$ is highly vertex prime if and only if $n \leq 2$; even cycles with a chord are highly vertex prime; $C_p \cup C_q$ is not highly vertex prime when both p and q are odd; and crowns $C_n \odot K_1$ are highly vertex prime.

For a finite simple graph $G(V, E)$ with n vertices and $v \in V$ let $N(v)$ denote the open neighborhood of v. Patel and Shrimali [2800] say a bijective function $f :\to \{1, 2, 3, \ldots, n\}$ is a *neighborhood-prime labeling* of G, if for every vertex $v \in V$ with $deg(v) > 1$, $gcd\{f(u) : u \in N(v)\} = 1$. A graph that admits a neighborhood-prime labeling is called a *neighborhood-prime* graph. In [2800], [2801], and [2802] they prove the following graphs have a prime-neighborhood labeling: graphs with a vertex of degree $|V| - 1$; paths; C_n if and only if $n \not\equiv 2 \pmod 4$; helms; closed helms; flowers; graphs obtained by the duplication of an arbitrary vertex of cycle or path; $G_1 + G_2$ where each of G_1 and G_2 have at least 2 vertices; $C_n \cup C_m$ is a neighborhood-prime graph if and only if $n \equiv 0 \pmod 4$ and $m \equiv 0 \pmod 4$, or $n \equiv 0 \pmod 4$ and $m \equiv 1 \pmod 2$; $W_m \cup W_n$; the union of a finite number of paths; $P_m \times P_n$; and the tensor product of two paths of the same order. They also prove that if G is neighborhood-prime graph and v is a vertex in G that is not adjacent to any pendent vertices, then the graph obtained by duplicating the vertex v is neighborhood-prime [2800].

Patel [2795] showed that the generalized Petersen graph $P(n, k)$ is neighborhood-prime when the greatest common divisor of n and k is $1, 2$, or 4 and that $P(n, 8)$ is neighborhood-prime for all n. Patel and Kansagara [2797] proved that the snake graphs of the type C_k^m are neighborhood-prime if and only if either $k \not\equiv 2 \pmod 4$ or $m \not\equiv 1 \pmod 4$ and that $C_{k,2}^m$ and $C_{k,3}^m$ are neighborhood-prime for all $m \geq 2$. Rozario Raj and Sheriff [3214] gave neighborhood-prime labelings for books with triangular and rectangle pages. Shrimali, Rathod, and Vihol [3571] proved the following graphs are neighborhood-prime: graphs obtained from the helm H_n by identifying each vertex of degree 1 with a vertex of W_n, graphs obtained from the helm H_n by identifying each vertex of degree 1 with a vertex of the fan F_n graphs obtained by identifying each pendent vertex of H_n with a vertex of outer cycle of closed helm of H_n, and graphs obtained by identifying each pendent vertex

of H_n by vertex of outer cycle of the Petersen graph.

Let $G(V, E)$ be a graph with p vertices and Q edges. Rajesh Kumar and Mathew Varkey [3100] call a bijection from $V(G) \cup E(G)$ to $\{1, 2, \ldots, p+q\}$ an *total neighborhood prime* labeling if for each vertex of degree at least two, the gcd of labeling on its neighborhood vertices is 1 and for each vertex of degree at least two, the gcd of labeling on the induced edges is 1. They proved that paths, combs, and C_{4n+2} are total neighborhood-prime graphs. Shrimali and Pandya [3565] proved that the following graphs have total neighborhood-prime labelings: combs $P_n \odot K_1$, $P_m \cup P_n$, $(P_m \odot K_1) \cup (P_n \odot K_1)$, $W_m \cup W_n$, graphs obtained from a copy of P_n and n copies $K_{1,n}$ by joining the ith vertex of P_n with an edge to the center vertex of the ith copy of $K_{1,m}$, $C_n \odot mK_1$, and subdivisions of bistars $B_{m,n}$. Shrimali, Rathod, and Vihol [3571] proved that following graphs are neighborhood-prime graphs: the graph obtained by identifying each pendent vertex of a helm H_n with a rim vertex of the wheel W_n; the graph obtained by identifying each pendent vertex of a helm H_n with a vertex of maximum degree of the fan $P_n + K_1$; and the graph obtained by identifying each pendent vertex of H_n with a vertex of outer cycle of closed helm graph $\overline{H_n}$. Rajesh Kumar [2069] proved that double stars, spiders, caterpillars, and firecrackers admit total neighborhood-prime labelings.

In [3099] Rajesh Kumar and Mathew Varkey extend the neighborhood-prime labeling concept to Gaussian integers. Using the spiral order on the Gaussian integers, they showed the following graphs have Gaussian neighborhood-prime labelings: graphs obtained by connecting the centers of two stars with a path, combs $P_n \odot K_1$, spiders, C_n where $n \neq 2 \mod 4$, C_4 ($n \geq 4$) with a cord, graphs obtained by switching of any vertex C_n, and graphs obtained by duplicating arbitrary vertex of C_n. Rajesh Kumar, Jerome, and Santhosh Kumar [3097] extended the Gaussian neighborhood-prime labeling concept to Gaussian total neighborhood-prime labeling. Among the graphs they prove to have a Gaussian neighborhood-prime labelings are: paths, stars, trees with one vertex of degree at least 3 and all other vertices having degree 1 or 2, and caterpillars. In [3573] Shrimali and Trivedi provided results depending upon the Hamiltonicity of a graph that guarantee that the graph is Gaussian neighborhood-prime graph under the spiral order and proved that generalized Petersen graphs are Gaussian neighborhood-prime graphs under the spiral order.

In [2755] Pandya and Shrimali introduce a vertex-edge neighborhood-prime labeling of graphs as follows. For a graph G an injective function $f : V(G) \cup E(G) \to \{1, 2, \ldots, |V(G)| + |E(G|\}$ is said to be a *vertex-edge neighborhood-prime labeling* if it has the following properties: If u has degree 1, then $\gcd\{f(w), f(uw)\} = 1$ taken over all vertices w adjacent to u; if u has degree greater than 1, then $\gcd\{f(w)\} = 1$ taken over all vertices w adjacent to u and $\gcd\{f(wu)\} = 1$ taken over all vertices w adjacent to u. A graph that admits vertex-edge neighborhood-prime labeling is called a *vertex-edge neighborhood-prime* graph. They give vertex-edge neighborhood-prime labelings for paths, helms, $C_n \oplus K_1$, bistars, the central edge subdivision of bistars, and subdivisions of edges of bistars. They observe that every vertex-edge neighborhood-prime graph is a total neighborhood-prime graph and that a total neighborhood-prime graph that does not have a vertex of degree 1 is vertex-edge neighborhood-prime. Shrimali and Rathod [3570]

proved that coconut trees, double coconut trees, spiders with legs of equal length, olive trees, combs, and $F(n,2)$-firecrackers have vertex-edge neighborhood-prime labelings.

Let $G(V,E)$ be a graph with p vertices and q edges. For $s \in V$, let $N_V[s] = \{s, t \in V \mid t \text{ is adjacent to } s\}$ and $N_E[s] = \{s \in V, e \in E \mid e \text{ is incident with } s\}$. A one-one mapping $\phi : V \cup E \to \{1, 2, \ldots, p+q\}$ is called a VECN *prime labeling* if, for each vertex s with $d(s) \geq 1$, where $d(s)$ is the degree of s, $\gcd(\phi(s), \phi(t)) \mid t \in N_V[s] = 1$ and $\gcd(\phi(s), \phi(e) \mid e \in N(E[s]) = 1$. A graph that admits VECN prime labeling is called a VECN *prime* graph. In [3798] Sunitha and Revathi investigate VECN prime labelings of m-fold quadrilateral snakes, m-fold alternate quadrilateral snakes, and m-fold irregular triangular snakes.

Patel and Ghodasara [2792] proved the following graphs are neighborhood-prime: one point union $C_n^{(k)}$ $(k \geq 2, n \geq 3)$ of k copies of cycle C_n, the barycentric subdivision of wheels and gears, the middle and total graph of crowns $C_n \odot K_1$ $(n \geq 3)$, the square of crowns, tadpoles $T(n,l)$ $(n \geq 3, l \geq 1)$, cycles, and umbrellas. In [919] Delman, Koilraj, and Raj gave neighborhood-prime labelings of arbitrary super subdivision of helms, tadpoles, and triangular snakes. In [2794] Pate1 and Ghodasara proved the following graphs are permutation graphs: wheels, restricted square and degree splitting graphs of bistars, arbitrary super subdivision of paths, stars, and cycles. Sonchhatra and Ghodasara [3673] proved that $P_2 + K_n$, books, cycles with one chord, cycles with twin chords, tadpoles, and lotus inside a circle graphs are permutation graphs.

Let $G(V,E)$ be a graph with p vertices and q edges. For $s \in V$, let $N_V[s] = \{s, t \in V \mid t \text{ is adjacent to } s\}$ and $N_E[s] = \{s \in V, e \in E \mid e \text{ is incident with } s\}$. A one-one mapping $\phi : V \cup E \to \{1, 2, \ldots, p+q\}$ is called a VECN *prime labeling* if, for each vertex s with $d(s) \geq 1$, where $d(s)$ is the degree of s, $\gcd(\phi(s), \phi(t)) \mid t \in N_V[s] = 1$ and $\gcd(\phi(s), \phi(e) \mid e \in N(E[s]) = 1$. A graph that admits VECN prime labeling is called a VECN *prime* graph. In [3798] Sunitha and Revathi investigate VECN prime labelings of m-fold quadrilateral snakes, m-fold alternate quadrilateral snakes, and m-fold irregular triangular snakes.

In [3025] Prajapati and Shah introduce an odd prime labeling as follows. Let $G(V,E)$ be a graph. A bijection f from V to $\{1, 3, \ldots, 2|V|-1\}$ is called an *odd prime labeling* if for each edge uv, $\gcd(f(u), f(v)) = 1$. A graph that admits odd prime labeling is called an *odd prime graph.* They prove paths, ladders, complete bipartite graphs, wheels, gears, flowers, helms, closed helms, and generalized Petersen graphs $P(n,2)$ are odd prime and conjecture that generalized Petersen graphs $P(n,k)$ and every prime graph is an odd prime graph. In [3026] they proved the following graphs are odd prime graphs: graphs obtained by duplication of a vertex of paths, stars, and wheels, and graphs obtained by duplication of an edge of cycles, stars, and wheels. Carter and Fox [758] prove that the disjoint union of cycles; cycle chains; snakes; books; triangular, pentagonal, and hexagonal stacked prisms; spiders; perfect binary trees; special cases of caterpillars; and firecrackers are odd prime. They characterize when odd prime labelings exist for powers of paths and cycles and make progress towards proving the conjecture that all prime graphs are also odd prime. Meena and Gajalakshmi [2474] proved that the corona product of $C_n \times P_2$ and $K_1, \overline{K_2}$, and $\overline{K_3}$ are odd prime graphs.

The *volcano graph* V_n has vertices $\{x_k \mid k = 1, 2, 3\} \cup \{y_1, y_2, \ldots, y_n\}$ and edges $\{x_1x_3\} \cup \{x_kx_{k+1} \mid k = 1, 2\} \cup \{x_1y_t \mid t = 1, 2, \ldots, n\}$. Komarullah, Hidayat, Krisnawati, and Wijaya [1995] showed that volcano graphs, graphs obtained by identifying a vertex of C_3 and vertex of maximum degree in a fan, $C_3 \odot \overline{K_n}$, tadpoles, and palm trees are all both prime and odd prime graphs. In [1994] they investigated prime and odd prime labelings of broom graphs and some related graphs. [1995] new [1994] new

In [1107] Fox and Mooney provided several infinite families of commutative rings that have prime labelings for their zero-divisor graphs. They also gave infinite families of commutative rings that do not have prime labelings for their zero-divisor graphs. They further determined which commutative rings will have prime labelings for their zero-divisor graphs for all rings with 14 or fewer vertices in their zero-divisor graph. In [1108] Fox and Spaeth showed that snakes, books, prisms, prime trees, certain families of windmills, and other families of graphs are total prime. [1107] new [1108] new

For a graph $G(V, E)$ with p vertices and q edges Shiu, Lau, and Lee [3536] call a bijection f from E to $\{1, 2, \ldots, q\}$ an *edge-prime labeling* if for each edge uv in E, we have $\gcd(f^+(u), f^+(v)) = 1$, where $f^+(u) = \Sigma f(uw)$ over all $uw \in E$. A graph that admits an edge-prime labeling is called an *edge-prime graph*. A bijection f from E to $\{1, 2, \ldots, q\}$ is aa *semi-edge-prime labeling* if for each edge uv in E, we have $\gcd(f^+(u), f^+(v)) = 1$ or $f^+(u) = f^+(v)$. They obtained a necessary and sufficient condition for the disjoint union of paths to be edge-prime, proved that all 2-regular graphs are edge-prime, proved that many bipartite and tripartite graphs are edge-prime (or not edge-prime), and showed that certain bipartite and tripartite graphs are semi-edge-prime graphs. In [2101] Lau, Lee, and Shiu proved that if G is a cubic graph and every component is of order 4, 6 or 8, then G is edge-prime if and only if $G \not\approx K_4$ or $nK_{3,3}$ and $n = 2$ or 3 (mod 4). They conjectured that a connected cubic graph G is not edge-prime if and only if $G \approx K_4$.

In [1566] Jagadesh and Baskar Babujee introduced an *edge vertex prime* labeling of a graph G as an injection f from $V(G) \cup E(G)$ to $\{1, 2, \ldots, |V(G)| + |E(G)|\}$ such that for every edge uv, the labels $f(u), f(v)$, and $f(uv)$ are pairwise relatively prime. A graph that admits such a labeling is called an *edge vertex prime* graph. They proved that paths, cycles, and stars are edge vertex prime. In [2780] and [2781] Parmar proved that wheels, fans, friendship graphs, and $K_{2,n}$ are edge vertex prime. Simaringa and Muthukumaran [3588] proved that following graphs have edge vertex prime labelings: triangular and rectangular books, butterfly graphs, $\overline{K_n} \cup K_{1,m}$, $K_{1,m} + K_1$, $\overline{K_m} \cup \overline{K_n}$, Jahangir graphs $J_{n,3}$ and $J_{n,4}$.

For two connected graphs $G_1(p_1, q_1)$ and $G_2(p_2, q_2)$ Baskar Babujee and Jagadesh [577] defined the graph $G_1\hat{O}G_2$ as one obtained by superimposing any vertex of G_2 on any vertex of G_1. Simaringa and S. Muthukumara [3588] proved that if $G(p, q)$ is not a path or W_4 with $p + q$ odd and G has an edge vertex prime labeling, then there exists a graph that admits edge vertex prime labeling for the following classes: $G\hat{O}W_n$, $G\hat{O}f_n$ (f_n is a fan graph), and $G\hat{O}F_n$ (F_n is a friendship graph). If $G(p, q)$ where $p + q$ even and G has an edge vertex prime labeling, they proved that there exists a graph that admits edge vertex prime labeling for the classes $G\hat{O}P_n$ and $G\hat{O}C_3$. They further showed that crowns and the union of cycles are edge vertex prime graphs.

In [3589] Simaringa and Muthukumaran investigated the existence of edge vertex prime labelings for crowns, unions of cycles, and wheel relate graphs. Shrimali and Parmar [3566] proved that the following graphs have edge vertex prime labelings: bistars $B(m,n)$, n-centipede trees, coconut trees obtained from the path P_n by appending m new pendent edges at an end vertex of P_n), double coconut trees (graphs obtained by attaching $n>1$ pendent vertices to one end of the path P_r and $m>1$ pendent vertices to the other end of path P_r), and special classes of banana trees and firecrackers. In [3567] Shrimali and Parmar showed that helms and graphs obtained by joining a fan ($n \geq 2$) and C_{2n} by sharing a common vertex are edge vertex prime graphs and that $P_{n_1} \cup P_{n_2} \cup \cdots \cup P_{n_k}$ and $K_{1,n_1} \cup K_{1,n_2} \cup \cdots \cup K_{1,n_k}$ are not an edge vertex prime graphs. They also provided a necessary condition for being edge vertex prime graph. Rilwan and Radha [3174] investigated edge vertex prime labelings and super edge vertex prime labelings of Cayley graphs and Cayley digraphs of finite groups. Baskar Babujee and Jagadesh [577] proved generalized stars $K_{1,n_1,n_2,n_3,\ldots,n_m}$ and generalized cycle stars $C_n * (P_{n_1} \cup P_{n_2} m \cup \ldots \cup P_{n_m})$ that are obtained by joining one of the pendent vertices of each of the paths with an edge to any vertex of C_n, admit edge vertex prime labelings. Simaringa and Muthukumaran [3590] proved that the following graphs are edge vertex prime graphs: $C_m \cup K_{1,n}, C_m \cup P_n, K_{2,m} \cup C_n, C_n \cup C_n$ when $n = 0$ or $2 \bmod n$, and the one point union of wheel and cycle-related graphs. They also proved that for graphs G with $|V(G)|+|E(G)|$ even $G \cup K_{1,n}$ and $G \cup P_n$ are edge vertex prime graphs.

In [4290] Youssef and Almoreed gave a new variation of the prime labeling as follows: A graph $G(V,E)$ has an *odd prime labeling* if its vertices can be labeled with distinct odd integers from 1 to $2|V(G)|-1$ such that for every edge xy in E the labels assigned to the vertices of x and y are relatively prime. A graph that admits an odd prime labeling is called an *odd prime graph*. They provided some families of odd prime graphs and give some necessary conditions for a graph to be odd prime. They conjectured that every prime graph is odd prime graph.

A *coprime* labeling of vertices of a graph G with distinct labels from the set $\{1,\ldots,m\}$ for some integer $m \geq n$ such that adjacent labels are relatively prime. The minimum value m for which G has a coprime labeling is defined as the *minimum coprime number* , denoted by $\mathrm{pr}(G)$, and a coprime labeling of G with largest label being $\mathrm{pr}(G)$ is called a *minimum coprime* labeling of G. Obviously, if G is a prime graph of order n, then $\mathrm{pr}(G) = n$. In [318] Asplund and Fox focus on the problem of determining the minimum coprime number for graphs that have been shown to not be prime. Among them are K_n ($n \geq 4$), W_{2n+1}, and the union of odd cycles. C. Lee [2137] determined the minimum coprime number of coronas of complete graphs with empty graphs, the joins of two paths, and prisms. She also proved that gears are prime, double wheels DW_n are prime if and only if n is even, and the graph that obtained by attaching P_2 to each vertex of C_n followed by attaching the star S_m at its center to each pendent vertex is prime. Komarullah, Wijaya, Hidayat, and Krisnawati [1973] provided the exact values for the minimum coprime numbers for mC_n with n odd and the exact values of the mimimum coprime numbers of mC_n for several other infinite families. They also gave exact minimum coprime numbers for mK_n and characterize prime and coprime labelings of dumbbell graphs according to the parity

[1973] new

of their parameters.

Tonapi and Paquin [3908] introduced a novel extension of coprime labeling for time-sensitive networks they call *dynamic coprime labeling* They proved that a graph admits a dynamic coprime labeling if and only if it admits a classical coprime labeling (existence equivalence). [3908] new

In [2826] Periasamy, Venugopal, and Rozario Raj introduced the kth Fibonacci prime labeling of graphs as follows. A kth *Fibonacci prime labeling of* a graph $G(V, E)$ with $|V(G)| = n$ is an injective function g from $V(G)$ to $\{f_k, f_{k+1}, \ldots, f_{k+n-1}\}$ where f_k is the kth Fibonacci number, that induces a function g^* from $E(G)$ to the non-negative integers defined by $g(uv) = \gcd(g(u), g(v)) = 1$ for all $uv \in E(G)$. A graph that admits a kth Fibonacci prime labeling and is called a kth *Fibonacci prime* graph. They proved that paths, cycles, $P_n \odot K_1$, triangular snakes, quadrilateral snakes, and tadpoles are kth Fibonacci prime graphs.

The tables following summarize the state of knowledge about prime labelings and vertex prime labelings. In the table, **P** means prime labeling exists, and **VP** means vertex prime labeling exists. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property.

Table 21: **Summary of Prime Labelings**

Graph	*Types*	*Notes*
P_n	P	[1156]
stars	P	[1156]
complete binary trees	P	[1156]
spiders	P	[1156]
trees	P?	[2230]
C_n	P	[927]
$C_n \cup C_{2m}$	P	[927]
K_n	P	iff $n \leq 3$ [2230]
W_n	P	iff n is even [3910]
helms	P	[3376]
fans	P	[3376]
flowers	P	[3376]
$K_{2,n}$	P	[3376]
$K_{3,n}$	P	$n \neq 3, 7$ [3376]
$P_n + \overline{K_m}$	not P	$n \geq 3$ [3376]
$P_n + \overline{K_2}$	P	iff $n = 2$ or n is odd [3376]
books	P	[3405]
$C_m + C_n$	not P	[3405]
C_n^2	not P	$n \geq 4$ [3405]
P_n^2	not P	$n \geq 6,\ n \neq 7$ [3405]

Continued on next page

Table 21 – *Continued from previous page*

Graph	*Types*	*Notes*
M_n (Möbius ladders)	not P	n even [3405]
$S_m \cup S_n$	P	[4291]
$C_m \cup S_n$	P	[4291]
$K_m \cup S_n$	P	iff no. of primes $\leq m+n+1$ is at least m [4291]
$K_n \odot K_1$	P	iff $n \leq 7$ [4291]
$P_m \times P_n$ (grids)	P	$m \leq 3,\ n$ prime [4115] m prime $> 2,\ p < n \leq p^2$ [865]
$C_n \odot \overline{K_i}$ (crowns)	P	[3910]
$P_n \odot \overline{K_2}$	P	iff $n \neq 2$ [4115]
C_m-snakes (see §2.2)	P	[756]
unicyclic	P?	[3376]

Table 22: **Summary of Vertex Prime Labelings**

Graph	*Types*	*Notes*
$C_m + C_n$	not P	[3405]
C_n^2	not P	$n \geq 4$ [3405]
P_n	not P	$n = 6,\ n \geq 8$ [3405]
M_{2n} (Möbius ladders)	not P	[3405]
connected graphs	VP	[927]
forests	VP	[927]

Continued on next page

Table 22 – *Continued from previous page*

Graph	*Types*	*Notes*
$C_{2m} \cup C_n$	VP	[927]
$C_{2m} \cup C_{2n} \cup C_{2k+1}$	VP	[927]
$C_{2m} \cup C_{2n} \cup C_{2t} \cup C_k$	VP	[927]
$5C_{2m}$	VP	[927]
$G \cup H$	VP	if G, H are connected and one is not an odd cycle [927]
2-regular graph G [927]	not VP	G has at least 2 odd cycles
	VP?	iff G has at most 1 odd cycle [927]

7.3 Edge-graceful Labelings

In 1985, Lo [2305] introduced the notion of edge-graceful graphs. A graph $G(V, E)$ is said to be *edge-graceful* if there exists a bijection f from E to $\{1, 2, \ldots, |E|\}$ such that the induced mapping f^+ from V to $\{0, 1, \ldots, |V|-1\}$ given by $f^+(x) = (\sum f(xy)) \pmod{|V|}$ taken over all edges xy is a bijection. Note that an edge-graceful graph is antimagic (see §6.1). A necessary condition for a graph with p vertices and q edges to be edge-graceful is that $q(q+1) \equiv p(p+1)/2 \pmod{p}$. Lee [2147] notes that this necessary condition extends to any multigraph with p vertices and q edges. It was conjectured by Lee [2147] that any connected simple (p, q)-graph with $q(q+1) \equiv p(p-1)/2 \pmod{p}$ vertices is edge-graceful. Lee, Kitagaki, Young, and Kocay [2153] prove that the conjecture is true for maximal outerplanar graphs. Lee and Murthy [2139] proved that K_n is edge-graceful if and only if $n \not\equiv 2 \pmod 4$. (An edge-graceful labeling given in [2305] for K_n for $n \not\equiv 2 \pmod 4$ is incorrect.) Lee [2147] notes that a multigraph with $p \equiv 2 \pmod 4$ vertices is not edge-graceful and conjectures that this condition is sufficient for the edge-gracefulness of connected graphs. Lee [2146] has conjectured that all trees of odd order are edge-graceful. Small [3639] has proved that spiders for which every vertex has odd degree with the property that the distance from the vertex of degree greater than 2 to each end vertex is the same are edge-graceful. Keene and Simoson [1946] proved that all spiders of odd order with exactly three end vertices are edge-graceful. Cabaniss, Low, and Mitchem [734] have shown that regular spiders of odd order are edge-graceful. For a (p, q)

connected edge-graceful graph G with $q = kp + r$, where $k$is an integer and $0 \leq r < p$. Kayathri and Amutha [1935] proved that every edge-graceful labeling f of G induces $((k+1)!)^r(k!)^{p-r}$ edge-graceful labelings of G.

Lee and Seah [2192] have shown that $K_{n,n,\ldots,n}$ is edge-graceful if and only if n is odd and the number of partite sets is either odd or a multiple of 4. Lee and Seah [2191] have also proved that C_n^k (the kth power of C_n) is edge-graceful for $k < \lfloor n/2 \rfloor$ if and only if n is odd and C_n^k is edge-graceful for $k \geq \lfloor n/2 \rfloor$ if and only if $n \not\equiv 2 \pmod 4$ (see also [734]). Lee, Seah, and Wang [2197] gave a complete characterization of edge-graceful P_n^k graphs. Shiu, Lam, and Cheng [3529] proved that the composition of the path P_3 and any null graph of odd order is edge-graceful. Uma and Mazuda Shanofer [3927] prove that $C_{2n+1} \odot \overline{K_2}$ is edge graceful and the graphs obtained by starting with C_n and, for each edge of C_n, adjoining a copy of C_n that shares an edge with the starting copy (the *flower graph* FL_n) is not edge-graceful. Rafael, Antalan, Angel, Loureynz, Gamurot, and Tagle [3065] provided edge-graceful labelings for all edge-graceful fan graphs $F_{1,n}$. Rafael, Antalan, Tagle, Angel, Loureynz, and Gamurot [3066] established that the bubble sort graph B_n is not edge-graceful when $n > 4$ is odd and that W_3 is the only edge-graceful Kulli-cycle windmill graph. [3065] new [3066] new

Lo [2305] proved that all odd cycles are edge-graceful and Wilson and Riskin [4205] proved the Cartesian product of any number of odd cycles is edge-graceful. Lee, Ma, Valdes, and Tong [2168] investigated the edge-gracefulness of grids $P_m \times P_n$. The necessity condition of Lo [2305] that a (p, q) graph must satisfy $q(q+1) \equiv 0$ or $p/2 \pmod p$ severely limits the possibilities. Lee et al. prove the following: $P_2 \times P_n$ is not edge-graceful for all $n > 1$; $P_3 \times P_n$ is edge-graceful if and only if $n = 1$ or $n = 4$; $P_4 \times P_n$ is edge-graceful if and only if $n = 3$ or $n = 4$; $P_5 \times P_n$ is edge-graceful if and only if $n = 1$; $P_{2m} \times P_{2n}$ is edge-graceful if and only if $m = n = 2$. They conjecture that for all $m, n \geq 10$ of the form $m = (2k+1)(4k+1)$, $n = (2k+1)(4k+3)$, the grids $P_m \times P_n$ are edge-graceful. Riskin and Weidman [3181] proved: if G is an edge-graceful $2r$-regular graph with p vertices and q edges and $(r, kp) = 1$, then kG is edge-graceful when k is odd; when n and k are odd, kC_n^r is edge-graceful; and if G is the Cartesian product of an odd number of odd cycles and k is odd, then kG is edge-graceful. They conjecture that the disjoint union of an odd number of copies of a $2r$-regular edge-graceful graph is edge-graceful.

Shiu, Lee, and Schaffer [3541] investigated the edge-gracefulness of multigraphs derived from paths, combs, and spiders obtained by replacing each edge by k parallel edges. Lee, Ng, Ho, and Saba [2177] construct edge-graceful multigraphs starting with paths and spiders by adding certain edges to the original graphs. Lee and Seah [2193] have also investigated edge-gracefulness of various multigraphs.

Lee and Seah (see [2147]) define a *sunflower* graph $SF(n)$ as the graph obtained by starting with an n-cycle with consecutive vertices $v_1, v_2, \ldots, v_n$ and creating new vertices $w_1, w_2, \ldots, w_n$ with w_i connected to v_i and v_{i+1} (v_{n+1} is v_1). In [2194] they prove that $SF(n)$ is edge-graceful if and only if n is even. In the same paper they prove that C_3 is the only triangular snake that is edge-graceful. Lee and Seah [2191] prove that for $k \leq n/2$, C_n^k is edge-graceful if and only if n is odd, and for $k \geq n/2$, C_n^k is edge-graceful if and only if $n \not\equiv 2 \pmod 4$. Lee, Seah, and Lo (see [2147]) have proved that for n odd,

$C_{2n} \cup C_{2n+1}, C_n \cup C_{2n+2}$, and $C_n \cup C_{4n}$ are edge-graceful. They also show that for odd k and odd n, kC_n is edge-graceful. Lee and Seah (see [2147]) prove that the generalized Petersen graph $P(n,k)$ (see Section 2.7 for the definition) is edge-graceful if and only if n is even and $k < n/2$. In particular, $P(n,1) = C_n \times P_2$ is edge-graceful if and only if n is even.

Schaffer and Lee [3325] proved that $C_m \times C_n$ $(m > 2, n > 2)$ is edge-graceful if and only if m and n are odd. They also showed that if G and H are edge-graceful regular graphs of odd order then $G \times H$ is edge-graceful and that if G and H are edge-graceful graphs where G is c-regular of odd order m and H is d-regular of odd order n, then $G \times H$ is edge-magic if $\gcd(c,n) = \gcd(d,m) = 1$. They further show that if H has odd order, is $2d$-regular and edge-graceful with $\gcd(d,m) = 1$, then $C_{2m} \times H$ is edge-magic, and if G is odd-regular, edge-graceful of even order m that is not divisible by 3, and G can be partitioned into 1-factors, then $G \times C_m$ is edge-graceful.

In 1987 Lee (see [2195]) conjectured that $C_{2m} \cup C_{2n+1}$ is edge-graceful for all m and n except for $C_4 \cup C_3$. Lee, Seah, and Lo [2195] have proved this for the case that $m = n$ and m is odd. They also prove: the disjoint union of an odd number copies of C_m is edge-graceful when m is odd; $C_n \cup C_{2n+2}$ is edge-graceful; and $C_n \cup C_{4n}$ is edge-graceful for n odd. Bu [714] gave necessary and sufficient conditions for graphs of the form $mC_n \cup P_{n-1}$ to be edge-graceful.

Kendrick and Lee (see [2147]) proved that there are only finitely many n for which $K_{m,n}$ is edge-graceful and they completely solve the problem for $m = 2$ and $m = 3$. Ho, Lee, and Seah [1446] use $S(n; a_1, a_2, \ldots, a_k)$ where n is odd and $1 \le a_1 \le a_2 \le \cdots \le a_k < n/2$ to denote the (n, nk)-multigraph with vertices $v_0, v_1, \ldots, v_{n-1}$ and edge set $\{v_i v_j |\ i \ne j, i - j \equiv a_t \pmod{n}$ for $t = 1, 2, \ldots, k\}$. They prove that all such multigraphs are edge-graceful. Lee and Pritikin (see [2147]) prove that the Möbius ladders (see §2.2 for definition) of order $4n$ are edge-graceful. Lee, Tong, and Seah [2213] have conjectured that the total graph of a (p,p)-graph is edge-graceful if and only if p is even. They have proved this conjecture for cycles. In [1957] Khodkar and Vinhage proved that there exists a super edge-graceful labeling of the total graph of $K_{1,n}$ and the total graph of C_n. Wang and Zang [4164] proved that a regular graph of odd degree is edge-graceful if it contains either a quasi-prism factor or a claw factor. In [266] Antalan, Angel, Gamurot, and Tagle investigated the edge-gracefulness of fan graphs.

For $n \ge 3$ a *water wheel* graph, denoted by WW_n, is the graph with $V(WW_n) = \{u_0\} \cup \{u_i, v_i, w_i : 1 \le i \le n\}$; and $E(WW_n) = \{u_0u_i, u_0v_i, u_iw_i, v_iw_i : 1 \le i \le n\} \cup \{u_iv_{i+1} : 1 \le i \le n-1\} \cup \{u_nv_1\}$. In [241] Angel, Antalan, Loureynz, Gamurot, and Tagle introduced two new related graphs, the right water wheel graphs (RWW_n) and left water wheel graphs (LWW_n). They proved the non edge-gracefulness of these graphs and determined all the edge-graceful splitting graphs of WW_n, RWW_n, and LWW_n. Using Python computer programs they proved that among these graph families, only $S(WW3)$, $S(RWW_3)$, and $S(LWW_3)$ are edge-graceful. [241] new

Kuang, Lee, Mitchem, and Wang [2050] have conjectured that unicyclic graphs of odd order are edge-graceful. They have verified this conjecture in the following cases: graphs obtained by identifying an endpoint of a path P_m with a vertex of C_n when $m + n$ is

even; crowns with one pendent edge deleted; graphs obtained from crowns by identifying an endpoint of P_m, m odd, with a vertex of degree 1; amalgamations of a cycle and a star obtained by identifying the center of the star with a cycle vertex where the resulting graph has odd order; graphs obtained from C_n by joining a pendent edge to $n-1$ of the cycle vertices and two pendent edges to the remaining cycle vertex.

In [4165] Wang and Zhang introduced the notion called edge-graceful deficiency, which is a parameter to measure how close a graph is away from being an edge-graceful graph. The *edge-graceful deficiency* of a graph G is the minimum value of k such that the edge labeling $fE \to \{1, 2, \ldots, q+k\}$ is edge-graceful. They proved that an odd regular graph is edge-graceful if it contains a quasi-prism factor or a claw factor and completely determine the edge-graceful deficiency of Hamiltonian regular graphs of even degree.

Gayathri and Subbiah [1231] say a graph $G(V, E)$ has a *strong edge-graceful* labeling if there is an injection f from the E to $\{1, 2, 3, \ldots, \langle 3|E|/2\rangle\}$ such that the induced mapping f^+ from V defined by $f^+(u) = (\Sigma f(uv)) \pmod{2|V|}$ taken all edges uv is an injection. They proved the following graphs have strong edge graceful labelings: $P_n(n \geq 3), C_n, K_{1,n}(n \geq 2)$, crowns $C_n \odot K_1$, and fans $P_n + K_1(n \geq 2)$. In his Ph.D. thesis [3703] Subbiah provided edge-graceful and strong edge-graceful labelings for a large variety of graphs. Among them are bistars, twigs, y-trees, spiders, flags, kites, friendship graphs, mirror of paths, flowers, sunflowers, graphs obtained by identifying a vertex of a cycle with an endpoint of a star, and $K_2 \odot C_n$, and various disjoint unions of path, cycles, and stars.

Hefetz [1396] has shown that a graph $G(V, E)$ of the form $G = H \cup f_1 \cup f_2 \cup \cdots \cup f_r$ where $H = (V, E')$ is edge-graceful and the f_i's are 2-factors is also edge-graceful and that a regular graph of even degree that has a 2-factor consisting of k cycles each of length t where k and t are odd is edge-graceful.

Bača and Holländer [372] investigated a generalization of edge-graceful labeling called (a, b)-consecutive labelings. A connected graph $G(V, E)$ is said to have an (a, b)-*consecutive labeling* where a is a nonnegative integer and b is a positive proper divisor of $|V|$, if there is a bijection from E to $\{1, 2, \ldots, |E|\}$ such that if each vertex v is assigned the sum of all edges incident to v the vertex labels are distinct and they can be partitioned into $|V|/b$ intervals
$W_j = [w_{\min} = (j-1)b + (j-1)a, w_{\min} + jb + (j-1)a - 1]$, where $1 \leq j \leq p/b$ and $w_{\min}$ is the minimum value of the vertices. They present necessary conditions for (a, b)-consecutive labelings and describe (a, b)-consecutive labelings of the generalized Petersen graphs for some values of a and b.

A graph with p vertices and q edges is said to be *k-edge-graceful* if its edges can be labeled with $k, k+1, \ldots, k+q-1$ such that the sums of the edges incident to each vertex are distinct modulo p. In [2216] Lee and Wang show that for each $k \neq 1$ there are only finitely many trees that are k-edge graceful (there are infinitely many 1-edge graceful trees). They describe completely the k-edge-graceful trees for $k = 0, 2, 3, 4$, and 5. Gayathri and Sarada Devi [1206] obtained some necessary conditions and characterizations for k-edge-gracefulness of trees. They also proved that specific families of trees are edge-graceful and k-edge-graceful and conjecture that all odd trees are k-edge-graceful.

Gayathri and Sarada Devi [943] defined a *k-even edge-graceful* labeling of a (p, q) graph $G(V, E)$ as an injection f from E to $\{2k-1, 2k, 2k+1, \ldots, 2k+2q-2\}$ such that the induced mapping f^+ of V defined by $f^+(x) = \sum f(xy) \pmod{2s}$ taken over all edges xy, are distinct and even, where $s = \max\{p, q\}$ and k is a positive integer. A graph G that admits a k-even-edge-graceful labeling is called a *k-even-edge-graceful* graph. In [943], [1207], [1208], and [1209] Gayathri and Sarada Devi investigate the k-even edge-gracefulness of a wide variety of graphs. Among them are: paths; stars; bistars; cycles with a pendent edge; cycles with a cord; crowns $C_n \odot K_1$; graphs obtained from P_n by replacing each edge by a fixed number of parallel edges; and sparklers (paths with a star appended at an endpoint of the path).

In 1991 Lee [2147] defined the *edge-graceful spectrum* of a graph G as the set of all nonnegative integers k such that G has a k-edge graceful labeling. In [2219] Lee, Wang, Ng, and Wang determine the edge-graceful spectrum of the following graphs: $G \odot K_1$ where G is an even cycle with one chord; two even cycles of the same order joined by an edge; and two even cycles of the same order sharing a common vertex with an arbitrary number of pendent edges attached at the common vertex (*butterfly graph*). Lee, Chen, and Wang [2150] have determined the edge-graceful spectra for various cases of cycles with a chord and for certain cases of graphs obtained by joining two disjoint cycles with an edge (i.e., *dumbbell graphs*). More generally, Shiu, Ling, and Low [3543] call a connected with p vertices and $p+1$ edges *bicyclic*. In particular, the family of bicyclic graphs includes the one-point union of two cycles, two cycles joined by a path and cycles with one cord. In [3544] they determine the edge-graceful spectra of bicyclic graphs that do not have pendent edges. Kang, Lee, and Wang [1882] determined the edge-graceful spectra of wheels and Wang, Hsiao, and Lee [4147] determined the edge-graceful spectra of the square of P_n for odd n. Results about the edge-graceful spectra of three types of $(p, p+1)$-graphs are given by Chen, Lee, and Wang [794]. In [4148] Wang and Lee determine the edge-graceful spectra of the one-point union of two cycles, the corona product of the one-point union of two cycles with K_1, and the cycles with one chord.

Lee, Levesque, Lo, and Schaffer [2162] investigate the edge-graceful spectra of cylinders. They prove: for odd $n \geq 3$ and $m \equiv 2 \pmod 4$, the spectra of $C_n \times P_m$ is $\emptyset$; for $m = 3$ and $m \equiv 0, 1$ or $3 \pmod 4$, the spectra of $C_4 \times P_m$ is $\emptyset$; for even $n \geq 4$, the spectra of $C_n \times P_2$ is all natural numbers; the spectra of $C_n \times P_4$ is all odd positive integers if and only if $n \equiv 3 \pmod 4$; and $C_n \times P_4$ is all even positive integers if and only if $n \equiv 1 \pmod 4$. They conjecture that $C_4 \times P_m$ is k-edge-graceful for some k if and only if $m \equiv 2 \pmod 4$. Shiu, Ling, and Low [3544] determine the edge-graceful spectra of all connected bicyclic graphs without pendent edges.

A graph $G(V, E)$ is called *super edge-graceful* if there is a bijection f from E to $\{0, \pm 1, \pm 2, \ldots, \pm(|E|-1)/2\}$ when $|E|$ is odd and from E to $\{\pm 1, \pm 2, \ldots, \pm|E|/2\}$ when $|E|$ is even such that the induced vertex labeling f^* defined by $f^*(u) = \Sigma f(uv)$ over all edges uv is a bijection from V to $\{0, \pm 1, \pm 2, \ldots, \pm(p-1)/2\}$ when p is odd and from V to $\{\pm 1, \pm 2, \ldots, \pm p/2\}$ when p is even. Lee, Wang, Nowak, and Wei [2220] proved the following: $K_{1,n}$ is super-edge-magic if and only if n is even; the double star $DS(m, n)$ (that is, the graph obtained by joining the centers of $K_{1,m}$ and $K_{1,n}$ with an edge) is

super edge-graceful if and only if m and n are both odd. They conjecture that all trees of odd order are super edge-graceful. In [2208] Lee, Su, and Wei exhibit a family of trees of odd orders which are super edge-graceful. Chung, Lee, Gao and Schaffer [830] posed the problems of characterizing the paths and trees of diameter 4 that are super edge-graceful.

In [829] Chung, Lee, and Gao prove various classes of caterpillars, combs, and amalgamations of combs and stars of even order are super edge-graceful. Lee, Sun, Wei, Wen, and Yiu [2209] proved that trees obtained by starting with the paths the P_{2n+2} or P_{2n+3} and identifying each internal vertex with an endpoint of a path of length 2 are super edge-graceful.

Shiu [3515] has shown that $C_n \times P_2$ is super-edge-graceful for all $n \geq 2$. More generally, he defines a family of graphs that includes $C_n \times P_2$ and generalized Petersen graphs as follows. For any permutation θ on n symbols without a fixed point the *θ-Petersen graph* $P(n;\theta)$ is the graph with vertex set $\{u_1, u_2, \ldots, u_n\} \cup \{v_1, v_2, \ldots, v_n\}$ and edge set $\{u_iu_{i+1}, u_iw_i, w_iw_{\theta(i)} \mid 1 \leq i \leq n\}$ where addition of subscripts is done modulo n. (The graph $P(n;\theta)$ need not be simple.) Shiu proves that $P(n;\theta)$ is super-edge-graceful for all $n \geq 2$. He also shows that certain other families of connected cubic multigraphs are super-edge-graceful and conjectures that every connected cubic of multigraph except K_4 and the graph with 2 vertices and 3 edges is super-edge-graceful.

In [3527] Shiu and Lam investigated the super-edge-gracefulness of fans and wheel-like graphs. They showed that fans F_{2n} and wheels W_{2n} are super-edge-graceful. Although F_3 and W_3 are not super-edge-graceful the general cases F_{2n+1} and W_{2n+1} are open. For a positive integer n_1 and even positive integers $n_2, n_3, \ldots, n_m$ they define an *m-level wheel* as follows. A wheel is a 1-level wheel and the cycle of the wheel is the 1-level cycle. An i-level wheel is obtained from an $(i-1)$-level wheel by appending $n_i/2$ pairs of edges from any number of vertices of the $i-1$-level cycle to n_i new vertices that form the vertices in the i-level cycle. They prove that all m-level wheels are super-edge-graceful. They also prove that for n odd $C_m \odot \overline{K_n}$ is super-edge-graceful, for odd $m \geq 3$ and even $n \geq 2$ $C_m \odot \overline{K_n}$ is edge-graceful, and for $m \geq 3$ and $n \geq 1$ $C_m \odot \overline{K_n}$ is super-edge-graceful. For a cycle C_m with consecutive vertices $v_1, v_2, \ldots, v_m$ and nonnegative integers $n_1, n_2, \ldots, n_m$ they define the graph $A(m; n_1, n_2, \ldots, n_m)$ as the graph obtained from C_m by attaching n_i edges to the vertex v_i for $1 \leq i \leq m$. They prove $A(m; n_1, n_2, \ldots, n_m)$ is super-edge-graceful if m is odd and $A(m; n_1, n_2, \ldots, n_m)$ is super-edge-graceful if m is even and all the n_i are positive and have the same parity. Chung, Lee, Gao, and Schaffer [830] provide super edge-graceful labelings for various even order paths, spiders and disjoint unions of two stars. In [827] Chung and Lee characterize spiders of even orders that are not super-edge-graceful and exhibit some spiders of even order of diameter at most four that are super-edge-graceful. They raised the question of which paths are super edge-graceful. This was answered by Cichacz, Fronček, and Xu [848] who showed that the only paths that are not super edge-graceful are P_2 and P_4. Cichacz et al. also proved that the only cycles that are not super edge-graceful are C_4 and C_6. Gao and Zhang [1199] proved that some cases of caterpillars are super edge-graceful.

In [830] Chung, Lee, Gao, and Schaffer asked for a characterize trees of diameter 4 that are super edge-graceful. Krop, Mutiso, and Raridan [2047] provide a super edge-graceful

labelings for all caterpillars and even size lobsters of diameter 4 that permit such labelings. They also provide super edge-graceful labelings for several families of odd size lobsters of diameter 4. They were unable to find general methods that describe super edge-graceful labelings for a few families of odd size lobsters of diameter 4, although they are able to show that certain lobsters in these families are super-edge graceful. They conclude with three conjectures about rooted trees of height 2 and diameter 4.

Although it is not the case that a super edge-graceful graph is edge-graceful, Lee, Chen, Yera, and Wang [2149] proved that if G is a super edge-graceful with p vertices and q edges and $q \equiv -1 \pmod{p}$ when q is even, or $q \equiv 0 \pmod{p}$ when q is odd, then G is also edge-graceful. They also prove: the graph obtained from a connected super edge-graceful unicyclic graph of even order by joining any two nonadjacent vertices by an edge is super edge-graceful; the graph obtained from a super edge-graceful graph with p vertices and $p+1$ edges by appending two edges to any vertex is super edge-graceful; and the one-point union of two identical cycles is super edge-graceful. Collins, Magnant, and Wang [860] present a stronger concept of "tight" super-edge-graceful labeling. Such a super-edge graceful labeling has an additional constraint on the edge and vertices with the largest and smallest labels. They use this concept to recursively construct tight super-edge graceful trees of any order.

Paley graphs are dense undirected graphs raised from the vertices as elements of an appropriate finite field by joining pairs of vertices that differ by a quadratic residue. In [1819], Kamaraj and Thangakani study the construction of edge even (odd) graceful labeling for Paley graphs and prove that Paley graphs of prime order are edge even (odd) graceful. Lee, Pan, and Tsai [2183] call a graph G with p vertices and q edges *vertex-graceful* if there exists a labeling $f\, V(G) \to \{1, 2, \ldots, p\}$ such that the induced labeling f^+ from $E(G)$ to Z_q defined by $f^+(uv) = f(u) + f(v) \pmod{q}$ is a bijection. Vertex-graceful graphs can be viewed the dual of edge-graceful graphs. They call a vertex-graceful graph *strong vertex-graceful* if the values of $f^+(E(G)$ are consecutive. They observe that the class of vertex-graceful graphs properly contains the super edge-magic graphs and strong vertex-graceful graphs are super edge-magic. They provide vertex-graceful and strong vertex-graceful labelings for various $(p, p+1)$-graphs of small order and their amalgamations.

Shiu and Wong [3558] proved the one-point union of an m-cycle and an n-cycle is vertex-graceful only if $m + n \equiv 0 \pmod{4}$; for $k \geq 2$, $C(3, 4k-3)$ is strong vertex-graceful; $C(2n+3, 2n+1)$ is strong vertex-graceful for $n \geq 1$; and if the one-point union of two cycles is vertex-graceful, then it is also strong vertex-graceful. In [3654] Somashekara and Veena found the number of $(n, 2n-3)$ strong vertex graceful graphs. Gao, Zhang, and Xu [1184] proved that C_n, $C_n \odot K_1$ and $C_n \odot K_{1,t}$ are vertex-graceful if n is odd; C_n is super vertex-graceful if $n \neq 4, 6$; and $C_n \odot K_1$ is super vertex-graceful if n is even. They proposed two conjectures on (super)vertex-graceful labelings. Selvaraju, Balaganesan, Renuka, and Suresh [3349] showed that $P_n \times C_m$ is vertex-graceful for $n > 1$ and odd $m \geq 5$, squares of bistars are vertex-graceful, and $P_n \times P_2$ (n odd) are vertex-graceful. They further proved that $C_j \cup C_k \cup C_l$ is vertex graceful for odd $j + k + l$ with $l \geq j + k + 5$.

As a dual to super edge-graceful graphs Lee and Wei [2223] define a graph $G(V,E)$ to be *super vertex-graceful* if there is a bijection f from V to $\{\pm 1, \pm 2, \ldots, \pm(|V|-1)/2\}$ when $|V|$ is odd and from V to $\{\pm 1, \pm 2, \ldots, \pm|V|/2\}$ when $|V|$ is even such that the induced edge labeling f^* defined by $f^+(uv) = f(u)+f(v)$ over all edges uv is a bijection from E to $\{0, \pm 1, \pm 2, \ldots, \pm(|E|-1)/2\}$ when $|E|$ is odd and from E to $\{\pm 1, \pm 2, \ldots, \pm|E|/2\}$ when $|E|$ is even. They show: for m and $n_1, n_2, \ldots, n_m$ each at least 3, $P_{n_1} \times P_{n_2} \times \cdots \times P_{n_m}$ is not super vertex-graceful; for n odd, books $K_{1,n} \times P_2$ are not super vertex-graceful; for $n \geq 3$, $P_n^2 \times P_2$ is super vertex-graceful if and only if $n = 3, 4$, or 5; and $C_m \times C_n$ is not super vertex-graceful. They conjecture that $P_n \times P_n$ is super vertex-graceful for $n \geq 3$.

In [2227] Lee and Wong generalize super edge-vertex graphs by defining a graph $G(V,E)$ to be $P(a)Q(1)$-*super vertex-graceful* if there is a bijection f from V to $\{0, \pm a, \pm(a+1), \ldots, \pm(a-1+(|V|-1)/2)\}$ when $|V|$ is odd and from V to $\{\pm a, \pm(a+1), \ldots, \pm(a-1+|V|/2)\}$ when $|V|$ is even such that the induced edge labeling f^* defined by $f^+(uv) = f(u)+f(v)$ over all edges uv is a bijection from E to $\{0, \pm 1, \pm 2, \ldots, \pm(|E|-1)/2\}$ when $|E|$ is odd and from E to $\{\pm 1, \pm 2, \ldots, \pm|E|/2\}$ when $|E|$ is even. They show various classes of unicyclic graphs are $P(a)Q(1)$-super vertex-graceful. In [2161] Lee, Leung, and Ng more simply refer to $P(1)Q(1)$-super vertex-graceful graphs as *super vertex-graceful* and show how to construct a variety of unicyclic graphs that are super vertex-graceful. They conjecture that every unicyclic graph is an induced subgraph of a super vertex-graceful unicyclic graph. Lee and Leung [2160] determine which trees of diameter at most 6 are super vertex-graceful graphs and propose two conjectures. Lee, Ng, and Sun [2179] found many classes of caterpillars that are super vertex-graceful. In [1191] Gao shows that the generalized butterfly graph B_n^t is super vertex-graceful when $t > 0$ is even, B_n^0 is super vertex-graceful when $n \equiv 0$ or 3 (mod 4), and $C_3^{(t)}$ is super vertex-graceful if and only if $t = 1, 2, 3, 5$, or 7.

In [816] Chopra and Lee define a graph $G(V,E)$ to be $Q(a)P(b)$-*super edge-graceful* if there is a bijection f from E to $\{\pm a, \pm(a+1), \ldots, \pm(a+(|E|-2)/2)\}$ when $|E|$ is even and from E to $\{0, \pm a, \pm(a+1), \ldots, \pm(a+(|E|-3)/2)\}$ when $|E|$ is odd and $f^+(u)$ is equal to the sum of $f(uv)$ over all edges uv is a bijection from V to $\{\pm b, \pm(b+1), \ldots, (|V|-2)/2\}$ when $|V|$ is even and from V to $\{0, \pm b, \pm(b+1), \ldots, \pm(|V|-3)/2\}$ when $|V|$ is odd. They say a graph is *strongly super edge-graceful* if it is $Q(a)P(b)$-super edge-graceful for all $a \geq 1$. Among their results are: a star with n pendent edges is strongly super edge-graceful if and only if n is even; wheels with n spokes are strongly super edge-graceful if and only if n is even; coronas $C_n \odot K_1$ are strongly super edge-graceful for all $n \geq 3$; and double stars $DS(m,n)$ are strongly super edge-graceful in the case that m is odd and at least 3 and n is even and at least 2 and in the case that both m and n are odd and one of them is at least 3. Lee, Song, and Valdés [2200] investigate the $Q(a)P(b)$-super edge-gracefulness of wheels W_n for $n = 3, 4, 5$, and 6.

In [2224] Lee, Wang, and Yera proved that some Eulerian graphs are super edge-graceful, but not edge-graceful, and that some are edge-graceful, but not super edge-graceful. They also showed that a Rosa-type condition for Eulerian super edge-graceful graphs does not exist and pose some conjectures, one of which was: For which n, is K_n is super edge-graceful? It was known that the complete graphs K_n for $n = 3, 5, 6, 7, 8$ are

super edge-graceful and K_4 is not super edge-graceful. Khodkar, Rasi, and Sheikholeslami, [1956] answered this question by proving that all complete graphs of order $n \geq 3$, except 4, are super edge-graceful.

Let ω be a primitive cube root of unity and G be a graph with a finite number of vertices where each vertex is labeled with $1, \omega$, or ω^2 and each edge uv is labeled with the product of the labels of u and v. Such a labeling is called a *cubic roots cordial labeling* of G if, for very pair of distinct labels x and y, the number of vertices labeled with x and y differ by at most 1 and number of edges labeled with x and y differ by at most 1. In [439] Bai and Senbagamalar proved that sunlet graphs ($C_n \odot K_1$) and friendship graphs admit cubic roots cordial labelings.

In 1997 Yilmaz and Cahit [4269] introduced a weaker version of edge-graceful called E-cordial. Let G be a graph with vertex set V and edge set E and let f a function from E to $\{0, 1\}$. Define f on V by $f(v) = \sum\{f(uv)|uv \in E\} \pmod 2$. The function f is called an *E-cordial* labeling of G if the number of vertices labeled 0 and the number of vertices labeled 1 differ by at most 1 and the number of edges labeled 0 and the number of edges labeled 1 differ by at most 1. A graph that admits an E-cordial labeling is called *E-cordial*. Yilmaz and Cahit prove the following graphs are E-cordial: trees with n vertices if and only if $n \not\equiv 2 \pmod 4$; K_n if and only if $n \not\equiv 2 \pmod 4$; $K_{m,n}$ if and only if $m + n \not\equiv 2 \pmod 4$; C_n if and only if $n \not\equiv 2 \pmod 4$; regular graphs of degree 1 on $2n$ vertices if and only if n is even; friendship graphs $C_3^{(n)}$ for all n (see §2.2 for the definition); fans F_n if and only if $n \not\equiv 1 \pmod 4$; and wheels W_n if and only if $n \not\equiv 1 \pmod 4$. They observe that graphs with $n \equiv 2 \pmod 4$ vertices can not be E-cordial. They generalized E-cordial labelings to E_k-cordial ($k > 1$) labelings by replacing $\{0, 1\}$ by $\{0, 1, 2, \ldots, k-1\}$. Of course, E_2-cordial is the same as E-cordial (see §3.7). Vaidya and Barasara [3951] proved that every graph can be embedded as an induced subgraph of an E-cordial graph thereby ruling out any possibility of obtaining any forbidden subgraph characterization for E-cordial graphs. They also proved that a connected graph can be embedded as an induced subgraph of an E-cordial connected graph and every planar graph can be embedded as an induced subgraph of an E-cordial planar graph.

In [3017] Prajapati and Patel proved the following graphs admit edge product cordial labelings: the graphs ottained by switching of any vertex with degree two that is adjacent to a vertex with degree two in triangular snakes, switching of any vertex with degree one in P_m for odd $m \geq 3$, switching of vertex with degree two in $P_m = u_1u_2 \cdots u_m$ except vertices u_2 or u_{m-1} for $m > 4$, and switching of any vertex in C_m. [3017] new

In [3226] Saibulla, Raj, Suganya, and Kareem use the term *edge-cordial* for the labeling we call E-cordial. They call an edge-cordial labeling in which the number of vertices and edges labeled with 0 and the number of vertices and edges labeled with 1 differ by at most 1. They say an edge-cordial is a *total edge-cordial* labeling. if the number of edges labeled with 0 and the number of vertices and edges labeled with 1 differ by at most 1. They investigated the existence of edge-cordial and total edge cordial labeling of the fractal graphs derived from Koch Curve and Koch Snowflake.

Liu, Liu, and Wu [2300] provide two necessary conditions for a graph to be E_k-cordial and prove that P_n ($n \geq 3$) is E_p-cordial for odd p. They also discuss the E_2-cordiality of

graphs that have a subgraph that is a 1-factor. Devaraj [936] has shown that $M(m,n)$, the mirror graph of $K(m,n)$, is E-cordial when $m+n$ is even and the generalized Petersen graph $P(n,k)$ is E-cordial when n is even. (Recall that $P(n,1)$ is $C_n \times P_2$.)

In [4026] Vaidya and Vyas prove that the following graphs are E-cordial: the mirror graphs (see §2.3 for the definition) even paths, even cycles, and the hypercube are E-cordial. In [3992] they show that the middle graph, the total graph, and the splitting graph of a path are E-cordial and the composition of P_2n with P_2. (See §2.7 for the definitions of middle, total and splitting graphs.) In [3993] Vaidya and Lekha [3993] prove the following graphs are E-cordial: the graph obtained by duplication of a vertex (see §2.7 for the definition) of a cycle; the graph obtained by duplication of an edge (see §2.7 for the definition) of a cycle; the graph obtained by joining of two copies of even cycle by an edge; the splitting graph of an even cycle; and the shadow graph (see §3.8 for the definition) of a path of even order.

Vaidya and Vyas [4027] proved the following graphs have E-cordial labelings: $K_{2n} \times P_2$; $P_{2n} \times P_2$; $W_n \times P_2$ for odd n; and $K_{1,n} \times P_2$ for odd n. Vaidya and Vyas [4028] proved that the Möbius ladders, the middle graph of C_n, and crowns $C_n \odot K_1$ are E-cordial graphs for even n while bistars $B_{n,n}$ and its square graph $B^2_{n,n}$ are E-cordial graphs for odd n. In [4030] and [4031] Vaidya and Vyas proved the following graphs are E-cordial: flowers, closed helms, double triangular snakes, gears, graphs obtained by switching of an arbitrary vertex in C_n except $n \equiv 2 \pmod 4$, switching of rim vertex in wheel W_n except $n \equiv 1 \pmod 4$, switching of an apex vertex in helms, and switching of an apex vertex in closed helms. Sugumaran and Vishnu Prakash [3758] proved that the following graphs are E-cordial: theta graphs, duplication of any vertex in theta graphs, switching of any vertex in theta graphs, the fusion of any two vertices in theta graphs, and the open star of n copies of a fixed theta graph (that is, the graph obtained by replacing each endpoint vertex of $K_{1,n}$ by copies of the theta graph).

In her PhD thesis [4046] Vanitha defines a (p,q) graph G to be *directed edge-graceful* if there exists an orientation of G and a labeling of the arcs of G with $\{1,2,\ldots,q\}$ such that the induced mapping g on V defined by $g(v) = |f^+(v) - f^-(v)| \pmod p$ is a bijection where, $f^+(v)$ is the sum of the labels of all arcs with head v and $f^-(v)$ is the sum of the labels of all arcs with tail v. She proves that a necessary condition for a graph with p vertices to be directed edge-graceful is that p is odd. Among the numerous graphs that she proved to be directed edge-graceful are: odd paths, odd cycles, fans F_{2n} $(n \geq 2)$, wheels W_{2n}, nC_3-snakes, butterfly graphs B_n (two even cycles of the same order sharing a common vertex with an arbitrary number of pendent edges attached at the common vertex), $K_{1,2n}$ $(n \geq 2)$, odd order y-trees with at least 5 vertices, flags Fl_{2n} (the cycle C_{2n} with one pendent edge), festoon graphs $P_n \odot mK_1$, the graphs $T_{m,n,t}$ obtained from a path P_t $(t \geq 2)$ by appending m edges at one endpoint of P_t and n edges at the other endpoint of P_t, C_3^n, $P_3 \cup K_{1,2n+1}$, $P_5 \cup K_{1,2n+1}$, and $K_{1,2n} \cup K_{1,2m+1}$.

In [689] Boonklurb, Ruamkaew, and Singhun use $C(c \times a)$ to denote the graph obtained by identifying a vertex of c cycles C_a to a single vertex. They provide directed edge-graceful labelings for $C(c \times a)$ in the case when $a \geq 3$ is an odd integer and $c \geq 2$ and in the case where a and c are even with $a \geq 4$ and $c \geq 2$.

The table following summarizes the state of knowledge about edge-graceful labelings. In the table **EG** means edge-graceful labeling exists. A question mark following an abbreviation indicates that the graph is conjectured to have the corresponding property.

Table 23: **Summary of Edge-graceful Labelings**

Graph	*Types*	*Notes*
K_n	EG	iff $n \not\equiv 2 \pmod 4$ [2139]
odd order trees	EG?	[2146]
$K_{n,n,\ldots,n}$ (k terms)	EG	iff n is odd or $k \not\equiv 2 \pmod 4$ [2192]
$C_n^k,\ k < \lfloor n/2 \rfloor$	EG	iff n is odd [2191]
$C_n^k,\ k \geq \lfloor n/2 \rfloor$	EG	iff $n \not\equiv 2 \pmod 4$ [2191]
$P_3[K_n]$	EG	n is odd [2191]
M_{4n} (Möbius ladders)	EG	[2147]
odd order dragons	EG	[2050]
odd order unicyclic graphs	EG?	[2050]
$P_{2m} \times P_{2n}$	EG	iff $m = n = 2$ [2168]
$C_n \cup P_2$	EG	n even [2195]
$C_{2n} \cup C_{2n+1}$	EG	n odd [2195]
$C_n \cup C_{2n+2}$	EG	[2195]
$C_n \cup C_{4n}$	EG	n odd [2195]
$C_{2m} \cup C_{2n+1}$	EG?	$(m,n) \neq (4,3)$ odd [2196]
$P(n,k)$ generalized Petersen graph	EG	n even, $k < n/2$ [2147]

Continued on next page

Table 23 – *Continued from previous page*

Graph	*Types*	*Notes*
$C_m \times C_n$	EG?	$(m,n) \neq (4,3)$ [2196]

7.4 Radio Labelings

In 2001 Chartrand, Erwin, Zhang, and Harary [781] were motivated by regulations for channel assignments of FM radio stations to introduce radio labelings of graphs. A *radio labeling* of a connected graph G is an injection c from the vertices of G to the natural numbers such that $d(u,v) + |c(u) - c(v)| \geq 1 + diam(G)$ for every two distinct vertices u and v of G. The *radio number* of c, $rn(c)$, is the maximum number assigned to any vertex of G. The *radio number* of G, $rn(G)$, is the minimum value of $rn(c)$ taken over all radio labelings c of G. Chartrand et al. and Zhang [4317] gave bounds for the radio numbers of cycles. The exact values for the radio numbers for paths and cycles were reported by Liu and Zhu [2289] as follows: for odd $n \geq 3$, $rn(P_n) = (n-1)^2/2 + 2$; for even $n \geq 4$, $rn(P_n) = n^2/2 - n + 1; rn(C_{4k}) = (k+2)(k-2)/2 + 1$; $rn(C_{4k+1}) = (k+1)(k-1)/2$; $rn(C_{4k+2}) = (k+2)(k-2)/2+1$; and $rn(C_{4k+3}) = (k+2)(k-1)/2$. However, Chartrand, Erwin, and Zhang [780] obtained different values than Liu and Zhu for P_4 and P_5. Chartrand, Erwin, and Zhang [780] proved: $rn(P_n) \leq (n-1)(n-2)/2 + n/2 + 1$ when n is even; $rn(P_n) \leq n(n-1)/2 + 1$ when n is odd; $rn(P_n) < rn(P_{n+1})$ $(n > 1)$; for a connected graph G of diameter d, $rn(G) \geq (d+1)^2/4 + 1$ when d is odd; and $rn(G) \geq d(d+2)/4 + 1$ when d is even. Karst, Langowitz, Oehrlein, and Troxell [1906] provide general lower bounds for $rc_k(C_n)$ for all cycles C_n when $k \geq \text{diam}(C_n)$ and show that these bounds are exact values when $k = \text{diam}(C_n) + 1$. In [671] Bloomfield, Liu, and Ramirez combine a lower bound approach with the cyclic group structure to determine the value of $\text{rnk}(C_n)$ for $k \geq n-3$. For $d \leq k < n-3$, they obtain the values of $\text{rnk}(C_n)$ when n and k have the same parity, and prove partial results when n and k and have different parities. These results extend the known values of $\text{rn}_d(C_n)$ and $\text{rn}_{d+1}(C_n)$ shown by Liu and Zhu [2289], and by Karst, Langowitz, Oehrlein, and Troxell in [1906], respectively. Benson, Porter, and Tomova [620] determined the radio numbers of all graphs of order n and diameter $n-2$. In [2284] Liu obtained lower bounds for the radio number of trees and the radio number of spiders (trees with at most one vertex of degree greater than 2) and characterized the graphs that achieve these bounds.

In [2286] Liu, Saha, and Das gave lower bounds for some trees whose radio number exceeds Liu's lower bound in [2284]. Sharp bounds for complete binary trees that improved results in [2243] [18] and odd paths in [2289] are given. They also extended the results of Halasza and Tuza [1356] on complete level-wise regular trees. Bantva, Vaidya, and Zhou [479] and [597] give a lower bound for the radio number of trees and a necessary and sufficient condition for their bound to be achieved. They determine the radio number for symmetric trees (that is, trees whose non-leaf vertices all have the same degree and whose leaf vertices all have the same eccentricity), banana trees, and firecracker trees. In [1991] Kola and Panigrahi provide the radio number for a class of caterpillars. In [478] Bantva and Der-Fen Liu give a lower bound for the radio number of the Cartesian product of two trees. Moreover, they present three necessary and sufficient conditions, and three sufficient conditions for the product of two trees to achieve this bound. Applying these results, they determine the radio number of the Cartesian product of two stars as well as a path and a star.

Gomathi and Venugopal [1292] gave upper and lower bounds for the radio number of triangular and rhombic honeycomb networks. In [4267] Yenoke, Selvagopal, Smitha, and Cranston determined the bounds for the radio number of an enhanced mesh. They state that their results have implications in the optimization of radio frequency assignment in wireless networks. [1292] new [4267] new

Nazeer, Khan, Kousar, and Nazeer [2660] investigated the radio number for some families of generalized caterpillar graphs. Chakraborty, Nandi, Sen, and Supraja [763] gave the exact values of $\mathrm{rc}_k(G)$ for powers of paths where the diameter of the graph is strictly less than k. Their proof provides a linear time algorithm for providing such labelings.

Chartrand, Erwin, Zhang, and Harary [781] proved: $rn(K_{n_1,n_2,\ldots,n_k}) = n_1 + n_2 + \cdots + n_k + k - 1$; if G is a connected graph of order n and diameter 2, then $n \leq rn(G) \leq 2n-2$; and for every pair of integers k and n with $n \leq k \leq 2n-2$, there exists a connected graph of order n and diameter 2 with $rn(G) = k$. They further provide a characterization of connected graphs of order n and diameter 2 with prescribed radio number.

Fernandez, Flores, Tomova, and Wyels [1081] proved $rn(K_n) = n$; $rn(W_n) = n+2$; and the radio number of the gear graph obtained from W_n by inserting a vertex between each vertex of the rim is $4n + 2$. Morris-Rivera, Tomova, Wyels, and Yeager [2552] determine the radio number of $C_n \times C_n$. In [3231] Saha and Panigrahi determined the radio number of $C_m \times C_n$ when $mn = 0 \pmod 2$. In [3508] Shigehalli and Usmanbasha provided the radio numbers for the Mycielskian graphs of paths, cycles and stars. [3231] new [3508] new

Martinez, Ortiz, Tomova, and Wyels [2445] define *generalized prisms,* denoted $Z_{n,s}$, $s \geq 1$, $n \geq s$, as the graphs with vertex set $\{(i,j) \mid i = 1,2 \text{ and } j = 1,\ldots,n\}$ and edge set $\{((i,j),(i,j \pm 1))\} \cup \{((1,i),(2,i+\sigma)) \mid \sigma = -\left\lfloor \frac{s-1}{2} \right\rfloor \ldots, 0, \ldots, \left\lfloor \frac{s}{2} \right\rfloor\}$. They determine the radio number of $Z_{n,s}$ for $s = 1, 2$ and 3. In [474] and [475] Bantva determines the radio number for three families of trees obtained by taking a graph operation on a given tree or a family of trees and the radio number for the middle graph of paths. Sun [3778] determined the radio number for the middle graph of a dandelion (the one-point union of a star and a path), which generalized Bantva's results in [475]. Zhang, Nazeer, Habib, Zia, and Ren [4314] determined the radio number for the generalized Petersen graphs $P(4k+2, 2)$ and provided a lower bound for $P(4k, 2)$. In [4061] Vasoya and Bantva gave a lower bound for the radio number for the Cartesian product of the generalized Petersen graph and a tree. They give two necessary and sufficient conditions and three other sufficient conditions to achieve the lower bound. They use these results to determine the radio number for the Cartesian product of the Petersen graph and stars.

Elrokh, Badr, Al-Shamiri, and Ramadhan [1012] provided upper bounds for the radio number of triangular snakes and double triangular snakes that improved on those given by Badr and Moussa [432] and by Saha and Panigrahi in [3230] and [3233]. DeVito, Niedzialomski, and Warren [945] determined the radio number of all diameter 3 Hamming graphs and showed that an infinite subset of them have radio graceful labelings. In [193] Alkasasbeh, · Badr, · Attiya, and · Shabana provided radio numbers of the friendship graphs $F_{3,k}, F_{4,k}, F_{5,k}$, and $F_{6,k}$. Kang, Nazeer, Nazeer, Kousar and Jung [1879] determined the radio number for a family of caterpillars In [2419] Mari and Jeyaraj determined the

radio number of the supersub-division of P_n $(n \geq 3)$ with $K_{2,m}$.

Gomathi and Venugopal [1291] determined lower and upper bounds of the radio antipodal numbers of triangular grids and the upper bound of the ladder-related graphs. In [1290] Gomathi, Venugopal, and Jose determined bounds of the radio numbers of triangular grid graphs, triangular ladders, and pagoda graphs. In [596] Basunia, Saha, and Tiwary determined the radio numbers of some special type of path related graphs. William and Kenneth [4192] also determined the radio numbers of some special type of path related graphs. In [2420] Mari and Jeyaraj determined the radio numbers of various families of graphs with diameter 3.

In [1392] Ramyal and Sooryanarayana generalized the notion of radio labeling as follows. Let M be a subset of non-negative integers and $(M, \star)$ be a monoid with the identity e. A *radio $\star$-labeling* of graph $G(V, E)$ is a mapping $f : V \to M$ such that $|f(u) - f(v)| \star d(u, v) \geq diam(G) + 1 - e$, for all $u, v \in V$. The *radio $\star$-number* $rn \star (f)$ of a radio $\star$ -labeling f of G is the maximum label assigned to a vertex of G. The radio $\star$-number of G, denoted by $rn \star (G)$, is the minimum of $rn \star (f)$ taken over all radio $\star$-labeling f of G. They completely determine $rn \star (G)$ of some transformation graphs of paths and cycles where $\star$ is the usual multiplication of integers.

Sooryanarayana and Ranghunath [3680] define a radio labeling f of a graph G to be a *consecutive radio labeling* of G if $f(V(G)) = \{1, 2, \ldots, |V(G)|\}$. They call a graph for which a consecutive radio labeling exists *radio graceful*. In her Ph.D. thesis [2674] Niedzialomski (see also [2675]) investigated the existence of radio graceful labelings of Cartesian products of graphs. Among her results are: for $n \geq 3$ and $1 \leq t \leq n - 1$ the Cartesian product of t copies of K_n is radio graceful; for $2 \leq p \leq n_2$ the Cartesian product of $p \cdot \lceil n/p \rceil$ copies of K_n is radio graceful; the Cartesian product $K_{n_1} \times K_{n_2}, \ldots, \times K_{n_s}$ is radio graceful when $n_1, n_2, \ldots, n_s$ are relatively prime; certain families of generalized Petersen graphs are radio graceful; and the Cartesian product of $t \geq 1 + n(n^2 - 1)/6$ copies of K_n is not radio graceful. Locke and Niedzialomski [2306] proved that $K_n \times P$ is radio graceful where P is the Peterson graph. Wyels and Tomova [2306] proved that $P \times P$ is radio graceful. Sooryanarayana and Raghunath [3680] determine the values of n for which C_n^3 is radio graceful. Sarkar [3236] used a recursive construction to show that for each positive integer $n \geq 9$, there exists a bipartite graph with $n+1$ vertices such that both a graph and its complement are radio graceful graphs. [3236] new

For a simple connected graph G with at least 3 vertices the *triameter* of G, denoted by $\mathrm{tr}(G)$, is the smallest positive integer M such that $d(u, v) + d(v, w) + d(w, u) \leq M$ for every triplet u, v, and w in $V(G)$. Saha and Basunia [3227] proved that a graph with diameter 2 is radio graceful if and only if it contains a Hamiltonian path. They also prove that if both a graph G and its complement $\overline{G}$ are connected and $\mathrm{tr}(G) > 9$, then $\overline{G}$ is radio graceful only if G contains a Hamiltonian path and $\mathrm{tr}(\overline{G}) = 5$ or 6.

The generalized gear graph $J_{t,n}$ is obtained from a wheel W_n by introducing t-vertices between every pair (v_i, v_{i+1}) of adjacent vertices on the n-cycle of wheel. Ali, Rahim, Ali, and Farooq [190] gave an upper bound for the radio number of generalized gear graph, which coincided with the lower bound found in and [3073]. They proved for $t < n-1$ and $n \geq 7$, $\mathrm{rn}(J_{t,n}) = (nt^2+4nt+3n+4)/2$. They pose the determination of the radio number

of $J_{t,n}$ when $n \leq 7$ and $t > n-1$ as an open problem. In [3796] Sunitha determine the radio number of double triangular snake graphs.

Saha and Panigrahi [3231] determined the radio number of the toroidal grid $C_m \times C_n$ when at least one of m and n is an even integer and gave a lower bound for the radio number when both m and n are odd integers. Liu and Xie [2287] determined the radio numbers of squares of cycles for most values of n. In [2288] Liu and Xie proved that $rn(P_n^2)$ is $\lfloor n/2 \rfloor + 2$ if $n \equiv 1 \pmod 4$ and $n \geq 9$ and $rn(P_n^2)$ is $\lfloor n/2+1 \rfloor$ otherwise. In [2285] Liu found a lower bound for the radio number of trees and characterizes the trees that achieve the bound. She also provides a lower bound for the radio number of spiders in terms of the lengths of their legs and characterizes the spiders that achieve this bound. Sweetly and Joseph [3851] prove that the radio number of the graph obtained from the wheel W_n by subdividing each edge of the rim exactly twice is $5n-3$. Marinescu-Ghemeci [2435] determined the radio number of the caterpillar obtained from a path by attaching a new terminal vertex to each non-terminal vertex of the path and the graph obtained from a star by attaching k new terminal vertices to each terminal vertex of the star. Ahmad and Marinescu-Ghemeci [132] determined the radio numbers of Mongolian tents, diamonds, fans, and double fans.

Sooryanarayana and Raghunath [3680] determined the radio number of C_n^3, for $n \leq 20$ and for $n \equiv 0$ or 2 or 4 (mod 6). Sooryanarayana, Vishu Kumar, Manjula [3681] determine the radio number of P_n^3, for $n \geq 4$. Lo and Alegria [2304] completely determine the radio number for the fourth-power of P_n for $n \geq 6$, except when $n \equiv 1 \pmod 8$. Saha and Panigrahi [3232] prove that for an n-vertex simple connected graph G, the difference between the upper and lower bounds of the radio number of G^2 is at most $\lfloor (n-1)/2 \rfloor$. They also determine the radio number for square of graphs belonging to some specific class and apply this to find the radio number for square of hypercube Q_n^2 ($n \not\equiv 0 \pmod 4$), the square of toroidal grid $T_{m,n}^2$ ($m+n \equiv 1,2,3,4,6 \pmod 8$), and the square of some generalized prism graphs. Wang, Xu, Yang, Zhang, Luo, and Wang [4139] determine the radio number of ladder graphs. Jiang [1779] completely determined the radio number of the grid graph $P_m \times P_n$ ($m, n > 2$). In [4024] Vaidya and Vihol determined upper bounds on radio numbers of cycles with chords and determined the exact radio numbers for the splitting graph and the middle graph of C_n. In [2243] Li, Mak, and Zhou determine the radio number of complete m-ary trees. Kim, Hwang, and Song [1960] determine the radio numbers of P_n with $n \geq 4$ and K_m with $m \geq 3$. Bantva [476] improved the lower bound for the radio number of graphs given by Das *et al.* in [900] and gave necessary and sufficient condition to achieve the lower bound. He also determined the radio number for Cartesian product of paths P_n and the Peterson graph P and provided a short proof for the radio number of Cartesian product of paths P_n and complete graphs K_m given by Kim [1960]. Bantva [477] determined the radio number of the Cartesian product of a path and a wheel.

A radio k-coloring of G when $k = \mathrm{diam}(G) - 1$ is called a *radio antipodal labeling*. The minimum span of a radio antipodal labeling of G is called the *radio antipodal number* of G and is denoted by $an(G)$. Khennoufa and Togni [1950] determined the radio number and the radio antipodal number of the hypercube by using a generalization of binary Gray

codes. They proved that $rn(Q_n) = (2^{n-1}-1)\lceil\frac{n+3}{2}\rceil+1$ and $an(Q_n) = (2^{n-1}-1)\lceil\frac{n}{2}\rceil+\varepsilon(n)$, with $\varepsilon(n) = 1$ if $n \equiv 0 \bmod 4$, and $\varepsilon(n) = 0$ otherwise. In [1293] Gomathi and Venugopal provided bounds for the radio antipodal number of honeycombs derived networks—triangular and rhombic honeycombs. These bounds give the optimum number of channels (bandwidth) needed for these honeycomb derived networks for effective communication without interference.

Let G be a finite, connected, simple graph with vertex set $V(G)$ and diameter D. For $u, v \in V(G)$, let $d_G(u, v)$ denote the distance between u and v in G. An *antipodal labeling* of G is a function $f(G) \to \mathbb{N}_0$ such that, for every pair of distinct vertices $u, v \in V(G)$, $|f(u) - f(v)| \geq D - d_G(u, v)$. The *span* of f is defined by $\mathrm{sp}(f) = \max|f(u) - f(v)|, v \in V(G)$. The *antipodal number* of G, denoted by $\mathrm{an}(G)$, is the minimum span over all antipodal labelings of G; that is, $\mathrm{an}(G) = \min \mathrm{sp}(f)$ is an antipodal labeling of G. In 2012 Juan and Liu [1804] proved that $\mathrm{an}(C_{4k}) \leq 2k^2 - 1$ and conjectured that $\mathrm{an}(C_{4k}) = 2k^2 - 1$ for every positive integer k. They verified their conjecture for $k \leq 5$. In 2026 Campbell [752] proved this conjecture for all positive integers k. Basunia, Das, Saha, and Tiwary [595] gave an exact formula for the antipodal number of full m-ary trees, together with an explicit construction of an optimal antipodal labeling. This paper extends earlier work on antipodal labelings of full binary trees and complements known results on radio labelings of complete m-ary trees. [752] new [595] new

In [4222] Xavier and Thivyarathi introduced the notion of radio antipodal mean labeling as follows. A *radio antipodal mean labeling* of a graph G is a function f that assigns to each vertex a non-negative integer such that for distinct vertices u and v of G, $d(u, v) < \mathrm{diam}(G)$ and $d(u, v) + \lceil(f(u) + f(v))/2\rceil \geq d$. The *radio antipodal mean number* of f, denoted by $r_{amn}(f)$, is the maximum number assigned to any vertex of G. The *radio antipodal mean number* of G, denoted by $r_{amn}(G)$, is the minimum value of $r_{amn}(f)$ taken over all radio antipodal mean labelings of G. They determined an upper bound for the antipodal mean number of paths, cycles, wheels, meshes ($P_m \times P_n$), and enhanced meshes. Jose, Prabakaran, and Gomath [1797] determined the radio antipodal mean number of triangular grids and torus grids. Yenoke, Jose, and Venugopal [4266] obtained the radio antipodal mean number of mongolian tents, triangular ladders, and Pagoda graphs.

In [2661] Nazeer, Kousar, and Nazeer give radio and radio antipodal labelings for certain circulant graphs. In [2662] Nazeer, Kousar, and Munir determined the radio number and radio antipodal number of non-bipartite cubic graphs of order 2^k. Shen, Dong, Zheng, and Guo [3500] use $C(m, t)$ to denote the caterpillar consisting of a path $x_1x_2\cdots x_m$ with t pendent edges at each inner vertex. They determine the exact value of the radio number of $C(m, t)$ for all integers $m \geq 4$ and $t \geq 2$, and explicitly construct an optimal radio labeling. They also show that the radio number and the construction of optimal radio labelings of paths are the special cases of $C(m, t)$ with $t = 2$. An edge-joint graph G is a 1-edge connected graph having an edge uv such that eccentricity of u equals the eccentricity of v and deletion of uv disconnects G.

Chavez, Liu, and Shurman [790] extended Liu's [2284] lower bound on radio numbers of trees to radio-k-numbers of trees. They established properties trees whose radio k- [790] new

number is this lower bound, and apply them to obtain shorter proofs and generalizations of some known results. They investigated the minimum integer k such that a given tree T has a radio k number equal to Liu's lower bound. They also defined the joined union operation on trees, where trees are composed by merging a vertex from each tree into a single vertex and used this operation to construct new k-lower bound trees in three ways. Finally, they posed several questions.

Niranjan and Rao Kola [2681] determined $\mathrm{rn}(P_n \odot C_m)$ when n is even and $m \geq 5$ and gave an upper bound for the same when n is odd. For $m \geq 4$ they determined the radio number of $P_n \odot P_m$ when n is even, and gave both upper and lower bounds for $\mathrm{rn}(P_n \odot P_m)$ when n is odd.

In [754] Canales, Tomova, and Wyels investigated the question of which radio numbers of graphs of order n are achievable. They proved that the achievable radio numbers of graphs of order n must lie in the interval $[n, rn(P_n)]$, and that these bounds are the best possible. They also show that for odd n, the integer $rn(P_n) - 1 = \frac{(n-1)^2}{2} + 2$ is an unachievable radio number for any graph of order n. In [3646] Sokolowsky settled the question of exactly which radio numbers are achievable for a graph of order n.

Adefokun and Ajayi [74] investigated the radio number of $K_{1,m} \times P_{2n}$ for the case $m \geq 4$. They also obtained new lower and upper bounds of the radio number for the case that $m = 3$ that improve similar their results in [150]. For a graph H, Naseem, Shabbir, and Shaker [2656] use JuH to denote the graph obtained by joining two disjoint copies of a graph H by an edge between the vertices labeled with u in H. They provided a lower bound for the radio number of such graphs and show that their lower bound is optimal (i.e., equal to the radio number) for certain a subfamily of JuH. They obtained similar results for graphs obtained by contracting the edge uv of JuH.

For any connected graph G and positive integer k Chartrand, Erwin, and Zhang, [779] define a *radio k-coloring* as an injection f from the vertices of G to the natural numbers such that $d(u, v) + |f(u) - f(v)| \geq 1 + k$ for every two distinct vertices u and v of G. Using $rc_k(f)$ to denote the maximum number assigned to any vertex of G by f, the *radio k-chromatic number* of G, $rc_k(G)$, is the minimum value of $rc_k(f)$ taken over all radio k-colorings of G. Note that $rc_1(G)$ is $\chi(G)$, the chromatic number of G, and when $k = diam(G)$, $rc_k(G)$ is $rn(G)$, the radio number of G. Chartrand, Nebesky, and Zang [787] gave upper and lower bounds for $rc_k(P_n)$ for $1 \leq k \leq n - 1$. Kchikech, Khennoufa, and Togni [1936] improved Chartrand et al.'s lower bound for $rc_k(P_n)$ and Kola and Panigrahi [1992] improved the upper bound for certain special cases of n. The exact value of $rc_{n-2}(P_n)$ for $n \geq 5$ was given by Khennoufa and Togni in [1951] and the exact value of $rc_{n-3}(P_n)$ for $n \geq 8$ was given by Kola and Panigrahi in [1992]. Kola and Panigrahi [1992] gave the exact value of $rc_{n-4}(P_n)$ when n is odd and $n \geq 11$ and an upper bound for $rc_{n-4}(P_n)$ when n is even and $n \geq 12$. In [3229] Saha and Panigrahi provided an upper and a lower bound for $rc_k(C_n^r)$ for all possible values of n, k and r and showed that these bounds are sharp for antipodal number of C_n^r for several values of n and r. Kchikech, Khennoufa, and Togni [1937] gave upper and lower bounds for $rc_k(G \times H)$ and $rc_k(Q_n)$. In [1936] the same authors proved that $rc_k(K_{1,n}) = n(k-1)+2$ and for any tree T and $k \geq 2$, $rc_k(T) \leq (n-1)(k-1)$. In [3228] Saha, Basunia, Das,

and Tiwary [3228] gave some lower bounds for the radio k-chromatic number and the radio antipodal number (when $k = \text{diam}(G) - 1$) for trees, and they computed the exact values for complete m-ary trees of height h for $k \geq 2h - 1$. Das, Lehtila, Nandi, Sen, and Supraja [901] improved the known lower bound of radio k-chromatic number for an arbitrary power of the infinite path. [901] new

In [763] Chakraborty, Nandi, Sen, and Supraja provided the exact values of radio k-coloring numbers of graphs that are a power of a path with diameter strictly less than k. Karst, Langowitz, Oehrlein, and Troxell [1906] provide general lower bounds for $rc_k(C_n)$ for all cycles C_n when $k \geq \text{diam}(C_n)$ and show that these bounds are exact values when $k = \text{diam}(C_n) + 1$.

In [736] Čada, Ekstein, Holub, and Togni [736] defined a k-labeling of a connected graph as an assignment c of non-negative integers to the vertices of the graph so that for every pair of vertices x and y, $|c(x) - c(y)| \geq k + 1 - d(x, y)$. The radio k-number of a graph is the largest number that has to be used in a k-labeling of the graph. A *distance graph* is a graph with integer vertices, where two vertices are adjacent if the absolute value of their difference is in some chosen set D. They established some lower and upper bounds for the radio k-number of graphs with distance sets $D(1, 2, \ldots, t)$, $D(1, t)$, and $D(t-1, t)$ for a positive t. Korze, Shao, and Vesel [2010] improved some lower and upper bounds for the radio k-labeling number of the same three families studied by Čada et al. and conjectured that some of these upper bounds for the radio k-labeling number of $D(1, 2, \ldots, t)$ are exact radio k-labeling numbers. They obtained their main results using theoretical constructions and the mathematical optimization methods of integer linear programming. Arockiamary and Vijayalakshmi [302] investigated the existence of vertex k-labeling of various cyclic snakes and $mC_n \odot K_1$.

For a positive integer k, Arockiamary and Vijayalakshmi [303] define a *vertex k-prime labeling* f of a graph $G(V, E)$ as a bijective function from V to $\{k, k+1, \ldots, k+|V|-1\}$ such that $\gcd(f(u), f(v) = 1$ for every edge uv. A graph G that admits vertex k-prime labeling is called a *vertex k-prime* graph. They proved that the class of planar graphs containing the maximum number of edges possible in a graph with n vertices is vertex k-prime if and only if n is odd, $G \cup K_{1,n}$ is vertex k-prime, and if G and H are k-prime labelings, then so is $G \cup H$. They also prove that K_n $(n \geq 4)$ is not vertex k-prime.

The Dd-distance was introduced by Anto Kinsley and Siva Ananthi [267] as follows. For a connected graph G, the Dd-*length* of a connected uv path is defined as $\text{Dd}(u, v) = \text{D}(u, v) + \deg(u) + \deg(v)$. Letting $D^{Dd}(u, v)$ denote the Dd-distance between u and v and $\text{diam}^{Dd}(G)$ denote the Dd-diameter of G, Viola and Nicholas [4116] define a *radio mean Dd-distance labeling* of a connected graph G as an injective map f from $V(G)$ to the positive integers such that for two distinct vertices u and v of G, $D^{Dd}(u, v) + \lceil (f(u) + f(v))/2 \rceil \geq 1 + \text{diam}^{Dd}(G)$. The *radio mean Dd-distance number* of f, $rmn^{Dd}(f)$, is the maximum label assigned to any vertex of G. The *radio mean Dd-distance number* of G, $rmn^{Dd}(G)$, is the minimum value of f of G. They determined the radio mean Dd-distance number of complete graphs, stars, bistars, subdivisions of stars, paths, fans, and friendship graphs.

In [1796] Jose and Giridaran introduced the notion a radial radio mean labeling of

graphs as follows. A *radial radio mean labeling* of a graph is a function f from $V(G)$ to the non-negative integers that satisfies the condition $d(u,v)+\lceil(f(u)+f(v))/2\rceil \geq 1+r(G)$, where u and v are vertices of G and $r(G)$ is the radius of G. The *span* of a radial radio mean labeling is the largest integer in the range of f. The *radial radio mean number* of a graph G is the minimum span taken over all radial radio mean labelings of G. They determined the radial radio mean number of Mongolian tents and diamond graphs. In [2765] Paramasivam, Yenoke, and Muralidharan investigated the upper bounds for radial radio number of chess board graphs and king's graph. Kaabar and Yenoke [1807] studied the relationship between the radio number and radial radio number for any connected graph. They provided upper bounds of the radio numbers and radial radio numbers for certain sunflower graphs. In [435] Badr, Alaybani, Elrokh, El-hay, and Almutairi determined the radio mean numbers for cycles and paths, provided an algorithm that approximates an upper bound for radio mean number of a given graph, and introduced a novel integer linear programing formulation for the radio mean problem.

The survey article by Panigrahi [2756] includes background information and further results about radio k-colorings.

7.5 Product and Divisor Cordial Labelings

Sundaram, Ponraj, and Somasundaram [3786] introduced the notion of product cordial labelings. A *product cordial labeling* of a graph G with vertex set V is a function f from V to $\{0,1\}$ such that if each edge uv is assigned the label $f(u)f(v)$, the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1, and the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. A graph with a product cordial labeling is called a *product cordial* graph. In [3786] and [3795] Sundaram, Ponraj, and Somasundaram prove the following graphs are product cordial: trees; unicyclic graphs of odd order; triangular snakes; dragons; helms; $P_m \cup P_n$; $C_m \cup P_n$; $P_m \cup K_{1,n}$; $W_m \cup F_n$ (F_n is the fan $P_n + K_1$); $K_{1,m} \cup K_{1,n}$; $W_m \cup K_{1,n}$; $W_m \cup P_n$; $W_m \cup C_n$; the total graph of P_n (the total graph of P_n has vertex set $V(P_n) \cup E(P_n)$ with two vertices adjacent whenever they are neighbors in P_n); C_n if and only if n is odd; $C_n^{(t)}$, the one-point union of t copies of C_n, provided t is even or both t and n are even; K_2+mK_1 if and only if m is odd; $C_m \cup P_n$ if and only if $m+n$ is odd; $K_{m,n} \cup P_s$ if $s > mn$; $C_{n+2} \cup K_{1,n}$; $K_n \cup K_{n,(n-1)/2}$ when n is odd; $K_n \cup K_{n-1,n/2}$ when n is even; and P_n^2 if and only if n is odd. They also prove that $K_{m,n}$ $(m,n>2)$, $P_m \times P_n$ $(m,n>2)$ and wheels are not product cordial and if a (p,q)-graph is product cordial graph, then $q \leq (p-1)(p+1)/4+1$.

The *bicyclic* graph $B[m,n]$ is the one-point union of C_m and C_n. Meena and Usharani [2478] provided product cordial labelings for some bicyclic related graphs. Among them are $B[n,n]$, the graphs obtained by adjoining the center of $K_{1,m}$ to each vertex of $B[n,n]$, $B[n,n] \odot K_2$, and $B[n,n] \odot K_3$. In [3489] Sharma and Parthiban investigted the existence of divisor cordial labelings of certain classes of planar graphs obtained from complete graphs and complete bipartite graphs using vertex duplication. [3489] new

In [3384] Seoud and Helmi obtained the following results: K_n is not product cordial

for all $n \geq 4$; C_m is product cordial if and only if m is odd; the gear graph G_m is product cordial if and only if m is odd; all web graphs are product cordial; the corona of a triangular snake with at least two triangles is product cordial; the C_4-snake is product cordial if and only if the number of 4-cycles is odd; $C_m \odot \overline{K_n}$ is product cordial; and they determine all graphs of order less than 7 that are not product cordial. Seoud and Helmi define the *conjunction* $G_1 \hat{} G_2$ of graphs G_1 and G_2 as the graph with vertex set $V(G_1) \times V(G_2)$ and edge set $\{(u_1, v_1)(u_2, v_2) |\ u_1u_2 \in E(G_1), v_1v_2 \in E(G_2)\}$. They prove: $P_m \hat{} P_n$ $(m, n \geq 2)$ and $P_m \hat{} S_n$ $(m, n \geq 2)$ are product cordial. Nada, Diab, Elrokh, and Sabra [2625] proved that $P_n \odot C_m$ is product cordial if and only if $(n, m) \neq (1, 3) \pmod 4$. Rokad [3197] proved the following graphs are product cordial: double wheels $DW_n = 2C_n + K_1$, path unions of finite number of copies of double wheels, the graphs obtained by joining two copies of double wheels by a path of arbitrary length, $DW_n \oplus K_{1,n}$, and $DF_n \oplus K_{1,n}$ $(DF_n = P_n + \overline{K}_2)$.

Vaidya and Kanani [3978] prove the following graphs are product cordial: the path union of k copies of C_n except when k is odd and n is even; the graph obtained by joining two copies of a cycle by path; the path union of an odd number copies of the shadow of a cycle (see §3.8 for the definition); and the graph obtained by joining two copies of the shadow of a cycle by a path of arbitrary length. In [3981] Vaidya and Kanani prove the following graphs are product cordial: the path union of an even number of copies of $C_n(C_n)$; the graph obtained by joining two copies of $C_n(C_n)$ by a path of arbitrary length; the path union of any number of copies of the Petersen graph; and the graph obtained by joining two copies of the Petersen graph by a path of arbitrary length.

Vaidya and Barasara [3939] prove that the following graphs are product cordial: friendship graphs; the middle graph of a path; odd cycles with one chord except when the chord joins the vertices at a diameter distance apart; and odd cycles with two chords that share a common vertex and form a triangle with an edge of the cycle and neither chord joins vertices at a diameter apart. In [3962] Vaidya and Barasara investigated the product cordial labeling of the line graph of the middle graphs of paths, triangular snakes, armed crowns, the square of paths, the splitting graphs of paths, and the total graph of paths.

In [3967] Vaidya and Dani prove the following graphs are product cordial: $< S_n^{(1)} : S_n^{(2)} : \ldots : S_n^{(k)} >$ except when k odd and n even; $< K_{1,n}^{(1)} : K_{1,n}^{(2)} : \ldots : K_{1,n}^{(k)} >$; and $< W_n^{(1)} : W_n^{(2)} : \ldots : W_n^{(k)} >$ if and only if k is even or k is odd and n is even with $k > n$. (See §3.7 for the definitions.)

Vaidya and Barasara [3946] proved the following graphs are product cordial: closed helms, web graphs, flower graphs, double triangular snakes obtained from the path P_n if and only if n is odd, and gear graphs obtained from the wheel W_n if and only if n is odd. Vaidya and Barasara [3948] proved that the graphs obtained by the duplication of an edge of a cycle, the mutual duplication of pair of edges of a cycle, and mutual duplication of pair of vertices between two copies of C_n admit product cordial labelings. Moreover, if G and G' are the graphs such that their orders or sizes differ at most by 1 then the new graph obtained by joining G and G' by a path P_k of arbitrary length admits product cordial labeling.

Vaidya and Barasara [3947] define the *duplication of a vertex* v of a graph G *by a new*

edge $u'v'$ as the graph G' obtained from G by adding the edges $u'v'$, vu' and vv' to G. They define the *duplication of an edge* uv of a graph G *by a new vertex* v' as the graph G' obtained from G by adding the edges uv' and vv' to G. They proved the following graphs have product cordial labelings: the graph obtained by duplication of an arbitrary vertex by a new edge in C_n or P_n $(n > 2)$; the graph obtained by duplication of an arbitrary edge by a new vertex in C_n $(n > 3)$ or P_n $(n > 3)$; and the graph obtained by duplicating all the vertices by edges in path P_n. They also proved that the graph obtained by duplicating all the vertices by edges in C_n $(n > 3)$ and the graph obtained by duplicating all the edges by vertices in C_n are not product cordial.

Recall (see [2951], [2938], [2939], [2940]) a *double triangular snake* DT_n consists of two triangular snakes that have a common path; a *double quadrilateral snake* DQ_n consists of two quadrilateral snakes that have a common path; an *alternate triangular snake* $A(T_n)$ is the graph obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertex v_i (that is, every alternate edge of a path is replaced by C_3); a *double alternate triangular snake* $DA(T_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to two new vertices v_i and w_i; an *alternate quadrilateral snake* $A(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i and w_i respectively and then joining v_i and w_i (that is, every alternate edge of a path is replaced by a cycle C_4); a *double alternate quadrilateral snake* $DA(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i, x_i and w_i and y_i respectively and then joining v_i and w_i and x_i and y_i.

Vaidya and Barasara [3950] prove that the shell graph S_n is product cordial for odd n and not product cordial for even n. They also show that $D_2(C_n)$; $D_2(P_n)$; C_n^2; $M(C_n)$; $S'(C_n)$; circular ladder CL_n; Möbius ladder M_n; step ladder $S(T_n)$ and $H_{n,n}$ does not admit product cordial labeling.

Vaidya and Vyas [4034] prove the following graphs are product cordial: alternate triangular snakes $A(T_n)$ except $n \equiv 3 \pmod 4$; alternate quadrilateral snakes $A(QS_n)$ except except $n \equiv 2 \pmod 4$; double alternate triangular snakes $DA(T_n)$ and double alternate quadrilateral snakes $DA(QS_n)$. Vaidya and Vyas [4035] prove the following graphs are product cordial: the splitting graph of bistar $S'(B_{n,n})$; duplicating each edge by a vertex in bistar $B_{n,n}$ and duplicating each vertex by an edge in bistar $B_{n,n}$. They also proved that $D_2(B_{n,n})$ is not product cordial.

Ghodasara and Vaghasiya [1263] prove the following graphs admit product cordial labelings: the path union of an odd number of copies of C_n with a chord except for $n = 4$, the path union of an odd number of copies of C_n with twin chords except when $n = 6$, the path union of C_n $(n > 6)$ with three cords that form two triangles and a cycle of length $n - 3$, the graph obtained by joining two copies of the same cycle that has one chord by a path, the graph obtained by joining two copies of same cycle that has twin chords by a path, and the graph obtained by joining two copies of C_n $(n \geq 7)$ with three cords that form two triangles and a cycle of length $n-3$ by a path. Ghodasara and Vaghasiya [1264] prove the following graphs are product cordial: the path union of helms, the path union of closed helms, the path union of gear graphs G_n for odd n, the graph obtained by joining two copies of the same helm by a path, the graph obtained by joining two copies of the

same closed helm by a path, and the graph obtained by joining two copies of the same gear graph by a path.

In [483] Bapat proves the following graphs are product cordial: graphs obtained by identifying an endpoint of P_n with each vertex of C_3, graphs obtained by identifying an endpoint of P_n with each vertex of C_4, graphs obtained by identifying the degree m vertex of $K_{1,m}$ with each vertex of C_3, and graphs obtained by identifying the degree m vertex of $K_{1,m}$ with each vertex of the shell $C_{n,n-3)}$ (C_n with $n-3$ chords that share a common endpoint) if and only n is even or n is odd and m is even. In [482] Bapat proves $K_5 \odot C_n$ and kayak paddles are product cordial, the one-point union of n copies of K_m is product cordial if and only if n is even, and graphs obtained by identifying one edge of K_5 with each edge of P_n is product cordial if and only if n is even.

Prajapati and Raval [3019] investigated product cordial labelings of the graphs obtained by duplication of vertices and edges of gears and graphs obtained by the vertex switching operation of gears. In [3020] Prajapati and Raval proved that the book $B_{m,n}$ is a product cordial graph if and only if m and n both are odd and $m \geq 3$. They showed that graphs obtained from books by duplicating or deleting vertices or edges are product cordial. For graphs with an even number of vertices they proved that the duplication of each of the vertices of a product cordial graph with an edge is a product cordial graph and that for graphs that have an odd number of vertices and even number of edges the duplication of each of the vertices of a product cordial graph with an edge is a product cordial graph.

Kwong, Lee, and Ng [2082] determine the product-cordial index sets of Möbius ladders and the graphs obtained by subdividing an edge of K_4 and an edge of a Möbius ladder that is not a rung and joining the two new vertices by an edge. They show that no Möbius ladder is product cordial. Gao, Sun, Zhang, Meng, and Lau [1198] provide sufficient conditions for a graph to admit (or not admit) a product cordial labeling. Gao, Lau, and Lee [1192] investigated the friendly index and product-cordial index sets of a family of cubic graphs known as Möbius-like graphs. Prajapati and Raval [3018] proved that windmills, barbells, the one point union at the apex of copies of a wheel (*generalized wheel*), and the one point union of copies of a wheel connected at one common rim vertex of the wheel are product cordial graphs. They also showed that duplicating all rim edges with a vertex and duplicating all the vertices with an edge of generalized wheels, and the graphs obtained by switching an apex vertex in a generalized wheel are product cordial graphs. Patel, Prajapati, and Kansagara [2799] proved that graphs obtained from the barbell graph by duplicating all vertices by edges and duplicating all edges by vertices in the path joining complete graphs are product cordial and the graphs obtained by switching a vertex of path in the barbell graph are product cordial.

In [3245] Salehi called the set $\{|e_f(0) - e_f(1)| : f \text{ is a friendly labeling of } G\}$ the *product-cordial set* of G. He determines the product-cordial sets for paths, cycles, wheels, complete graphs, bipartite complete graphs, double stars, and complete graphs with an edge deleted. Salehi and Mukhin [3254] say a graph G of size q is *fully product-cordial* if its product cordial set is $\{q - 2k : 0 \leq k \leq \lfloor q/2 \rfloor\}$. They proved: P_n ($n \geq 2$) is fully product-cordial; trees with a perfect matching are fully product-cordial; and $P_2 \times P_n$ is not

fully product-cordial. They determine the product-cordial sets of $P_2 \times P_n$, $P_n \times P_{2m}$, and $P_n \times P_{2m+1}$, where $m \geq n$. Because the product-cordial set is the multiplicative version of the friendly index set, Kwong, Lee, and Ng [2080] called it the *product-cordial index set* of G. They determined the exact values of the product-cordial index set of C_m and $C_m \times P_n$ and that $P_m \times P_n$ has the maximum product cordial-index $2mn-m-n$. In [2081] Kwong, Lee, and Ng determined the friendly index sets and product-cordial index sets of 2-regular graphs and the graphs obtained by identifying the centers of any number of wheels. In [3248] z Salehi, Churchman, Hill, and Jordan determine the product-cordial index sets of certain classes of trees.

In [3526] Shiu and Kwong define the *full product-cordial index* of G under f as $\mathrm{FPCI}(G) = \{i_f^*(G) \mid f \text{ is a friendly labeling of } G\}$. They provide a relation between the friendly index and the product-cordial index of a regular graph. As applications, they determine the full product-cordial index sets of C_m and $C_m \times C_n$, which was asked by Kwong, Lee, and Ng in [2080]. Shiu [3518] determined the product-cordial index sets of grids $P_m \times P_n$. Recall the *twisted cylinder* graph is the permutation graph on $4n$ ($n \geq 2$) vertices, $P(2n;\sigma)$, where $\sigma = (1,2)(3,4)\cdots(2n-1,2n)$ (the product of n transpositions). Shiu and Lee [3540] determined the full friendly index sets and the full product-cordial index sets of twisted cylinders.

Jeyanthi and Maheswari [1688] define a mapping $f : V(G) \to \{0,1,2\}$ to be a 3-*product cordial* labeling if $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(i) - e_f(j)| \leq 1$ for any $i,j \in \{0,1,2\}$, where $v_f(i)$ denotes the number of vertices labeled with $i, e_f(i)$ denotes the number of edges xy with $f(x)f(y) \equiv i \pmod 3$. A graph with a 3-product cordial labeling is called a 3-*product cordial* graph. In [1688] they prove that for a (p,q) 3-product cordial graph: $p \equiv 0 \pmod 3$ implies $q \leq \frac{p^2-3p+6}{3}$; $p \equiv 1 \pmod 3$ implies $q \leq \frac{p^2-2p+7}{3}$; and $p \equiv 2 \pmod 3$ implies $q \leq \frac{p^2-p+4}{3}$. They prove the following graphs are 3-product cordial: paths; stars; C_n if and only if $n \equiv 1,2 \pmod 3$; $C_n \cup P_n$, $C_m \odot \overline{K_n}; P_m \odot \overline{K_n}$ for $m \geq 3$ and $n \geq 1; W_n$ when $n \equiv 1 \pmod 3$; and the graph obtained by joining the centers of two identical stars to a new vertex. They also prove that K_n is not 3-product cordial for $n \geq 3$ and if G_1 is a 3-product cordial graph with $3m$ vertices and $3n$ edges and G_2 is any 3-product cordial graph, then $G_1 \cup G_2$ is a 3-product cordial graph. In [1689] they prove that ladders, $< W_n^{(1)} : W_n^{(2)} : \ldots : W_n^{(k)} >$ (see §3.7 for the definition), graphs obtained by duplicating an arbitrary edge of a wheel, graphs obtained by duplicating an arbitrary vertex of a cycle or a wheel are 3-product cordial. They also prove that the graphs obtained by from the ladders $L_n = P_n \times P_2$ ($n \geq 2$) by adding the edges u_iv_{i+1} for $1 \leq i \leq n-1$, where the consecutive vertices of two copies of P_n are $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$ and the edges are u_iv_i. They call these graphs *triangular ladders* . The graph $B_{n,n}^*$ is obtained from the bistar $B_{n,n}$ with $V(B_{n,n}) = \{u, v, u_i, v_i \mid 1 \leq i \leq n\}$ and $E(B_{n,n}) = \{uv, uu_i, vv_i, vu_i, uv_i \mid 1 \leq i \leq n\}$ by joining u with v_i and v with u_i for $1 \leq i \leq 4$. Jeyanthi and Maheswari [1695] proved: the splitting graphs $S'(K_{1,n})$ and $S'(B_{n,n})$ are 3-product cordial graphs; $B_{n,n}^*$ is a 3-product cordial graph if and only if $n \equiv 0,1 \pmod 3$; and the shadow graph $D_2(B_{n,n})$ is a 3-product cordial graph if and only if $n \equiv 0,1 \mod 3$. Jeyanthi, Maheswari, and Vijaya Laksmi [1709] prove the following: graphs obtained by switching an apex vertex in a closed helm are 3-product

cordial; W_n are 3-product cordial if and only if $n \equiv 2 \pmod 3$; double fans are 3-product cordial if and only if $n \equiv 0 \pmod 3$; books are 3-product cordial; and permutation graphs $P(K_2 + mK_1; T)$ are 3-product cordial if and only if $m \equiv 2 \pmod 3$. In [1708] Jeyanthi, Maheswari, and Vijayalaksmi investigated the 3-product cordial behavior of alternate triangular snakes, double alternate triangular snakes, and triangular snakes.

A *k-product cordial* labeling of a graph G is a map f from $V(G)$ to $\{0, 1, \ldots, k-1\}$ where k is a positive integer at most $|V(G)|$ such that when each edge uv is assigned the label $f(u)f(v) \pmod k$, the number of vertices (edges) labeled with i and the number of edges (vertices) labeled with j differ by at most 1. In [1649] and [1653] Jeya Daisy, Sabibha, Jeyanthi, and Youssef showed that the following graphs admit k-product cordial labelings: fans and double fans when $k = 4$ and 5; cones $(\overline{K_1} + C_m)$ and double cones $(\overline{K_2} + C_m)$ for $k = 5$. They also proved that double cones are not 4-product cordial. In [1653] they investigated the k-product cordial behavior of $G + K_t$, where G is a k-product cordial graph and found an upper bound of the size of k-product cordial graphs. In [1649] they investigated the k-product cordial behavior of union of graphs. In [1651] Jeya Daisy, Santrin Sabibha, Jeyanthi, and Youssef proved that the 1-cone $C_n + K_1$ and double cone $C_m + \overline{K_2}$ admit 5-product cordial labelings. They also showed that the double cone does not admit 4-product cordial labeling. In [1650] they define the *Napier bridge*, $P_n(t)$, as the graph obtained from the path P_n by joining the pairs of vertices u, v of P_n with $d(u, v) = t$. They provided necessary and significant conditions for $P_n(3), P_n(4)$ and $P_n(5)$ to have 3-product and 4-product cordial labelings. In [1652] Jeya Daisy, Sabibha, Jeyanthi and Youssef proved that fans F_n and double fans DF_n admit 4-product cordial labelings and 5-product cordial labelings. In [1654] they investigated the existence of k-product cordial labelings of direct products, Cartesian products, strong products, and lexicographic products of graphs. Jeya Daisya, Sabibhab, Jeyanthi, and Youssef [1655] showed that the splitting graph of star graphs admit k-product cordial labeling. In [3224] Sabibha, Jeya Daisy, Jeyanthi, and Youssef proved that paths admit k-product cordial labelings. Jeya Daisy, Santrin Sabibha, Jeyanthi, and Youssef [1656] investigated the k-product cordial behavior of $G + K_t$ and provided an upper bound of the size of connected k-product cordial graphs.

In [3555] Shiu, Sabibha, Jeyanthi, Daisy, and Lau introduced the notions of bijective product k-cordial labeling and bijective product square k-cordial labelings as follows. A *bijective product k-cordial* labeling for a graph $G(V, E)$ is a bijection from V to $\{1, 2, \ldots, |V|\}$ such that the induced edge labeling $f^x : E \to Z_k = \{1, 2, \ldots, k-1\}$ defined by $f^x(uv) = f(u)f(v) \pmod k$ for every edge uv satisfies the condition the number of edges labeled with i under f^x and the number of edges labeled with j under f^x differ by at most 1. A graph that admits a bijective product k-cordial labeling is called a *bijective product k-cordial* graph. A *bijective product square k-cordial* labeling for a graph $G(V, E)$ is a bijection from V to $\{1, 2, \ldots, |V|\}$ such that the induced edge labeling $f^* : E \to \{i^2 \bmod k\}$ defined as $f^*(uv) = f(u)f(v)^2$ for every edge uv satisfies the condition the number of edges labeled with i under f^* and the number of edges labeled with j under f^* differ by at most 1. A graph that admits a bijective product k-cordial labeling is called a *bijective product square k-cordial* graph.. They showed that some standard

graphs admit bijective product k-cordial labeling, where $k = 2, 3$ and that paths, cycles, flowers, helms, and gears are bijective product square 3-cordial graphs. In [3310] Santrin Sabibha, Jeya Daisy, Jeyanthi, and Youssef gave an upper limit of the size of the bijective product square k-cordial graph for $k = 3$ and 5 proved that every graph is a subgraph of a connected bijective product square k-cordial graph for all odd prime k. They also established a relation between bijective product square k-cordial and k-product cordial graphs. Bashammakh, Shiu, Sabibha, Jeyanthi, and Abdel-Aal [555] studied bijective product k-cordiality for paths and cycles, where k is an odd prime. They determined bijective product k-cordiality for paths and cycles for 3, 5, 7, 9, 11, and 13. They also established the bijective product k-cordial labeling of stars and provided the bijective product 4-cordial labeling of bistars and the splitting graphs of stars and bistars.

Sundaram and Somasundaram [3790] introduced the notion of total product cordial labelings. A *total product cordial labeling* of a graph G with vertex set V is a function f from V to $\{0, 1\}$ such that if each edge uv is assigned the label $f(u)f(v)$ the number of vertices and edges labeled with 0 and the number of vertices and edges labeled with 1 differ by at most 1. A graph with a total product cordial labeling is called a *total product cordial* graph. In [3790] and [3788] Sundaram, Ponraj, and Somasundaram prove the following graphs are total product cordial: every product cordial graph of even order or odd order and even size; trees; all cycles except C_4; $K_{n,2n-1}$; C_n with m edges appended at each vertex; fans; double fans; wheels; helms; $C_n \times P_2$; $K_{2,n}$ if and only if $n \equiv 2 \pmod 4$; $P_m \times P_n$ if and only if $(m, n) \neq (2, 2)$; $C_n + 2K_1$ if and only if n is even or $n \equiv 1 \pmod 3$; $\overline{K_n} \times 2K_2$ if n is odd, or $n \equiv 0$ or $2 \pmod 6$, or $n \equiv 2 \pmod 8$. Y.-L. Lai, the reviewer for MathSciNet, called attention to some errors in [3788]. Pedrano and Rulete [2815] determined the total product cordial labeling of $P_m \times C_n$, $C_m \times C_n$ and the generalized Petersen graph $P(m, n)$. In [2816] Pedrano and Rulete determined the total product cordial labeling of $P_m \odot C_n$, $P_m \odot P_n$, $C_m \odot P_n$, $P_m \odot F_n$, $P_m \odot W_n$, and $P_m \odot K_n$. Villar [4075] proved $P_n \odot C_m$ $(n \geq 2, m \geq 3)$ is product cordial, $P_n \odot P_m$ $(n, m \geq 2)$ is product cordial except when n and m are both even, and $P_{2n+1} \odot K_m$ $(n \geq 1, m \geq 4)$ is not product cordial. In [2817] Pedrano and Rulete provided total product cordial labelings for $P_m \times C_n$ except when $(m, n) = (1, 4), C_m \times C_n$, and the generalized Petersen graph $P(m, n)$ for $m \geq 3$ and $n \geq 3$. Ryan, Marquez, and Pedrano [3218] determined the total product cordial labeling of the snake graphs T_n, $A(T_n)$, $D(T_n)$, $DA(T_n)$, Q_n, $A(Q_n)$, $D(Q_n)$, and $DA(Q_n)$.

[2817] new

[3218] new

Gao, Sun, Zhang, Meng, and Lau [1198] proved that P_{n+1}^m is total product cordial. Ramanjaneyulu, Venkaiah, and Kothapalli [3112] give total product cordial labeling for a family of planar graphs for which each face is a 4-cycle.

Vaidya and Vihol [4018] prove the following graphs have total product cordial labelings: a split graph; the total graph of C_n; the star of C_n (recall that the *star of a graph* G is the graph obtained from G by replacing each vertex of star $K_{1,n}$ by a graph G); the friendship graph F_n; the one point union of k copies of a cycle; and the graph obtained by the switching of an arbitrary vertex in C_n.

Sundaram, Ponraj, and Somasundaram [3793] introduced the notion of EP-*cordial labeling* (extended product cordial) labeling of a graph G as a function f from the

vertices of a graph to $\{-1, 0, 1\}$ such that if each edge uv is assigned the label $f(u)f(v)$, then $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(i) - e_f(j)| \leq 1$ where $i, j \in \{-1, 0, 1\}$ and $v_f(k)$ and $e_f(k)$ denote the number of vertices and edges respectively labeled with k. An EP-*cordial graph* is one that admits an EP-cordial labeling. In [3793] Sundaram, Ponraj, and Somasundaram prove the following: every graph is an induced subgraph of an EP-cordial graph, K_n is EP-cordial if and only if $n \leq 3$; C_n is EP-cordial if and only if $n \equiv 1, 2 \pmod 3$, W_n is EP-cordial if and only if $n \equiv 1 \pmod 3$; and caterpillars are EP-cordial. They prove that all $K_{2,n}$, paths, stars and the graphs obtained by subdividing each edge of a star exactly once are EP-cordial. They also prove that if a (p, q) graph is EP-cordial, then $q \leq 1 + p/3 + p^2/3$. They conjecture that every tree is EP-cordial.

Ponraj, Sivakumar, and Sundaram [2973] introduced the notion of k-product cordial labeling of graphs. Let f be a map from $V(G)$ to $\{0, 1, 2, \ldots, k-1\}$, where $2 \leq k \leq |V|$. For each edge uv assign the label $f(u)f(v) \pmod k$. f is called a k-product cordial labeling if $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(i) - e_f(j)| \leq 1$, $i, j \in \{0, 1, 2, \ldots, k-1\}$, where $v_f(x)$ and $e_f(x)$ denote the number of vertices and edges labeled with x. A graph with a k-product cordial labeling is called a *k-product cordial graph*. Observe that 2-product cordial labeling is simply a product cordial labeling and 3-product cordial labeling is an EP-cordial labeling. In [2973] and [2980] Ponraj et al. prove the following are 4-product cordial: P_n if and only $n \leq 11$, C_n if and only if $n = 5, 6, 7, 8, 9$, or 10, K_n if and only if $n \leq 2$, $P_n \odot K_1$, $P_n \odot 2K_1$, $K_{2,n}$ if and only if $n \equiv 0, 3 \pmod 4$, W_n if and only if $n = 5$ or 9, $\overline{K_n} + 2K_2$ if and only $n \leq 2$, and the subdivision graph of $K_{1,n}$. Sivakumar [3622] proved the following coronas are 4-total product cordial: $P_n \odot K_1$, $P_n \odot 2K_1$, $S(P_n \odot K_1)$, $S(P_n \odot 2K_1)$, $S(C_n \odot K_1)$ and $S(C_n \odot 2K_1)$. Jeyanthi, Maheswari, and Vijayalakshmi [1708] investigated the 3-product cordial behavior of alternate triangular snakes, double alternate triangular snakes, and triangular snake graphs. In [1710] they establish that vertex switching graphs of wheels, gears, and degree splitting of bistars are 3-product cordial graphs.

Let f be a map from $V(G)$ to $\{0, 1, 2, \ldots, k-1\}$ where $2 \leq k \leq |V|$. For each edge uv assign the label $f(u)f(v) \pmod k$. Ponraj, Sivakumar, and Sundaram [2975] define f to be a *k-total product cordial labeling* if $|f(i) - f(j)| \leq 1$, $i, j \in \{0, 1, 2, \ldots, k-1\}$, where $f(x)$ denote the number of vertices and edges labeled with x. A graph with a k-total product cordial labeling is called a *k-total product cordial graph* . A 2-total product cordial labeling is simply a total product cordial labeling. In [2975], [2976], [2977], [2978] and [2979], Ponraj et al. proved the following graphs are 3-total product cordial: P_n, C_n if and only if $n \neq 3$ or 6, $K_{1,n}$ if and only if $n \equiv 0, 2 \pmod 3$, $P_n \odot K_1$, $P_n \odot 2K_1$, $K_2 + mK_1$ if and only if $m \equiv 2 \pmod 3$, helms, wheels, $C_n \odot 2K_1$, $C_n \odot K_2$, dragons $C_m@P_n$ (obtained by identifying an endpoint of P_n with a vertex of C_m), $C_n \odot K_1$, bistars $B_{m,n}$, and the subdivision graphs of $K_{1,n}$, $C_n \odot K_1$, $K_{2,n}$, $P_n \odot K_1$, $P_n \odot 2K_1$, $C_n \odot K_2$, wheels and helms. They also proved that every graph is a subgraph of a connected k-total product cordial graph, $B_{m,n}$ is $(n+2)$-total product cordial, and $K_{m,n}$ is $(n+2)$-total product cordial. Ahmad, Hasni, Irfan, Naseem, and Siddiqui [125] proved that $P_m \times P_n$ for $m, n \geq 2$ is 3-total edge product cordial. Sharon Philomena and Thirusangu [3492] proved that the flower graph is 3-total product cordial. Ahmada, Bača, Naseemc, and

Semaničová-Feňovčíková [116] described a method for obtaining a 3-total edge product cordial labeling of the hexagonal grid from a smaller hexagonal grid. In [99] Ahmad proved that the generalized Petersen graphs $P(n, m)$ are 3-total edge product cordial. In [325] Azaizeh, Hasni, Lau, and Ahmad proved that complete graphs, bipartite graphs and generalized friendship graphs have 3-total edge product cordial labelings. Ahmad, Ali, Bilal, Zafar, and Zahid [103] prove that webs, helms, gears, and $L_n \odot 2K_1$ (L_n is the ladder with $2n$ vertices) have 3-total edge product cordial labelings. Ivančo [1555] characterized graphs admitting a 2-total edge product cordial labeling and proved that dense graphs and regular graphs of degree $2(k-1)$ admit a k-total edge product cordial labeling. Javed and Jamil [1587] proved that the rhombic grid graphs R_1^m, R_2^m and R_3^m are 3-total edge product cordial for $m \geq 1$ and that the rhombic grid graph R_n^m is 3-total edge product cordial for $m, n \geq 1$. Tenguria and Verma [3877] proved that paths, cycles, and $K_{1,n}$ ($n = 0$ or 2 mod 3) admit 3-total edge sum cordial labelings.

Ullah, Rahmat, Numan, Anoh Yannick, and Aslam [3925] proved that the stellation of $P_n \times P_n$ (that is, the graph obtained from $P_n \times P_n$ by adding a vertex in each square of $P_n \times P_n$ and then joining this vertex to each vertex of that square) admits a 3-total edge product cordial labeling. Bala, Krithika Devi, and Thirusangu [451] proved the existence of 3-total cordial edge magic labeling, 3-total sum cordial labeling, and total product cordial labeling for the square graph of a comb. Selva and Lathamaheswari [3342] proved that paths, stars, double stars, cycles, and wheels are total 3-sum cordial graphs. Veronisha, Lourdusamy, and Beaula [4084] investigated total 3-sum cordial labelings of zero-divisor graphs obtained from various rings of integers modulo n. Jeba Jesintha, Devakirubanithi, and Monisha [1605] provided edge product cordial labelings for the duplication of prism graphs.

In [2070] Kumari and Mehra call a vertex labeling f of a graph G with 0 and 1 with the induced edge labeling f given by $f(uv) = f(u)f(v)$ a *vertex product cordial* labeling if the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1 and the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. They prove the following graphs have vertex product cordial labelings: P_n^2 if and only if n is odd, the path unions of k copies of P_n^2, $P_n \odot K_1$, helms, gears G_n for odd n, graphs obtained from C_n after switching of a vertex, $C_n \odot K_1$, $C_n \odot \bar{K}_m$, and certain banana trees.

A labeling f of a graph $G(V, E)$ from $V \cup E$ to $\{0, 1\}$ is said to be a *total product cordial* labeling of G if $f(xy) = f(x)f(y)$ for all $x, y \in V$ and $xy \in E$ and the difference of the total number vertices and edges labeled 0 and the total number vertices and edges labeled 1 differ by at most 1. Cyrile, Valdehueza, and Pedrano [881] determined the total edge product cordial labeling of the corona graphs $P_n \odot P_m$ and $P_n \odot C_m$.

In [1165] Gajjar and Raval define an *E-sum cordial labeling* of a graph G as an edge labeling $f^* : E(G) \to \{0, 1\}$, where the induced vertex labeling defined by $f(u) = \sum f(uv)$ mod 2 over all edges incident to u, has the property that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1, and the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. They introduce two new graphs insprited by origami models that they call the *north*

star graph and the *lotus star* (see article for the definitions) and investigate the existence of cordial labelings, sum cordial labelings, signed product cordial labelings, and E-sum cordial labelings for these new graph.

In [3088] Soundar Rajan and Baskar Babujee investigated the existence of signed product cordial labeling for splitting graphs of bull graphs, the splitting graphs of stars, P_n^2, $C_n \odot 3K_1$, and the graphs obtained by joining two copies of the helm H_4 by a path of arbitrary length.

In [3198] Rokad and Ghodasara introduced the notion of Fibonacci cordial labelings as follows. For a graph $G(V, E)$ an injective function $f : V(G) \to \{F_0, F_1, F_2, \ldots, F_n\}$, where $n = |V| - 1$ and F_j is the jth Fibonacci number, is said to be a *Fibonacci cordial labeling* if the induced function $f^* : E(G) \to \{0, 1\}$ defined by $f^*(uv) = (f(u) + f(v)) \mod 2$ satisfies the condition the number of edges with label 0 and the number of edges labeled with 1 differer by at most 1. A graph that admits Fibonacci cordial labeling is called *Fibonacci cordial* graph. Rokad and Ghodasara proved that the following graphs are Fibonacci cordial: Petersen graphs, wheels, shells, bistars, graphs obtained by joining the centers of two copies of a star with an edge, graphs obtained by identifying the center of $K_{1,n}$ with a vertex of C_n, graphs obtained by vertex switching of C_n, graphs obtained by joining the apexes of two copies of a fan with an edge, and graphs obtained by joining the centers of two copies of a wheel with an edge. Sulayman and Pedrano [**?**] proved that for all non-trivial cases alternate triangular snakes, quadrilateral snakes, double alternate quadrilateral snakes, and cycle quadrilateral snakes admit Fibonacci cordial labelings. Gondalia [1302] proved that the graphs obtained by vertex switching of the following graphs are Fibonacci product cordial: cycles, cycles with one chord, and cycles with twin chords where the chords form two triangles and one cycle C_{n-2}.

In [2526] Mitra and Bhoumik characterized the complete graphs that are Fibonacci cordial. They further investigated the Fibonacci cordialness of paths and cycles. In [2528] Mitra and Bhoumik investigated the existence and construction of Fibonacci cordial labelings for generalized Petersen graphs, open and closed helm graphs, and circulant graphs of small order. Mitra, Pritchard, and Bhoumik proved the following planar graphs admit Fibonacci cordial labelings: combs, jellyfish, H-graphs, certain coconut graphs, and graphs obtained by identifying an endpoint of $K_{1,n}$ ($n > 1$ odd with an endpoint of another copy of $K_{1,n}$.

In [2383] Mader and Mitra investigated the Fibonacci cordiality of $C_n \odot K_p$ for $p = 4, 5$; gave exact conditions for which the Fibonacci cordiality of $K_{n,n} \odot K_p$ for $p = 1, 2$, and 3 fails; etabished exact conditions for which the Fibonacci cordiality of $C_n \odot K_p$ fails; and investigated when $K_m \odot K_m$ and $K_n \odot K_{n,n} \odot K_p$ admit Fibonacci cordial labelings. [2383] new

Sulayman and Pedrano [3769] proved that all non-trival cases of the following graphs admit Fibonacci cordial labelings: alternate triangular snakes, quadrilateral snakes, double alternate quadrilateral snakes, and cycle quadrilateral snakes. [3769] new

The sequence T_n of tribonacci numbers is defined by the third-order linear recurrence relation ($n \geq 0$) $T_{n+3} = T_n + T_{n+1} + T_{n+2}$ where the first three tribonacci numbers are 0, 1, and 1. In [2527] Mitra and Bhouik introduced tribonacci cordial labeling as an extension of Fibonacci cordial labeling as follows. For a graph $G(V, E)$ an injective

function $f : V(G) \to \{T_0, T_1, T_2, \ldots, T_n\}$, where $n = |V| - 1$, is said to be a *tribonacci cordial labeling* if the induced function $f^* : E(G) \to \{0, 1\}$ defined by $f^*(uv) = (f(u) + f(v)) \bmod 2$ satisfies the condition the number of edges with label 0 and the number labeled with 1 differer by at most 1. A graph that admits a tribonacci cordial labeling is called a *tribonacci cordial*graph . They investigated whether paths, cycles, complete graphs, wheels, shells and generalized friendship graphs are tribonacci cordial. In [4090] Vieve, Gudin, and Pedrano determined which triangular snakes, double triangular snakes, quadrilateral snakes, double quadrilateral snake graph, and cycle quadrilateral snakes admit tribonacci cordial labelings.

In 2019 the concept of a Fibonacci product cordial labeling was introduced by Abraham and Jose [23] as follows. An injective function f from the vertices of a graph to the set $\{F_0, F_1, F_2, \ldots, F_{n+1}\}$ where F_j is the jth Fibonacci number is said to be a *Fibonacci product cordial* labeling if the induced function f^* from the edges to the set $\{0, 1\}$ defined by $f^*(uv) = (f(u)fv))(\text{mod } 2)$ has the property that the number of edges with label 0 and the number of edges with label 1 differ by at most 1. A graph which admits a Fibonacci product cordial labeling is called a *Fibonacci product cordial* graph. They proved that the following graphs are Fibonacci product cordial: paths, cycles, wheels W_n except when $n = 2 \bmod 3$, and the Petersen graph. In [1342] Gudin and Pedrano gave [1342] new tribonacci cordial labelings of triangular snakes, double triangular snakes, quadrilateral snakes, double quadrilateral snakes, and cycle quadrilateral snakes.

In [776] Charishma and Nageswari defined an L-Fibonacci cordial labeling as follows. An injective function $g : V(G) \to \{F_0, F_1, \ldots, F_n\}$, where $n = |V| - 1$ and F_j denotes the jth Fibonacci number, is said to be an *L-Fibonacci cordial labeling* of $G(V, E)$ if the induced function $g : E \to \{0, 1\}$ results in edge labeling of 0 if the maximum label among the two adjacent vertices is an even number and 1 otherwise, satisfies the condition that the number of vertices labeled with 0 and the number labeled with 1 differ by at most 1. A graph that admits an L-Fibonacci cordial labeling is called an *L-Fibonacci cordial graph* . Charishma and Nageswari showed that L-Fibonacci cordial labelings is admissible for paths, cycles, Petersen graphs, n-pan graphs, and bistars.

For a graph G Sundaram, Ponraj, and Somasundaram [3794] defined the *index of product cordiality*, $i_p(G)$, of G as the minimum of $\{|e_f(0) - e_f(1)|\}$ taken over all the 0-1 binary labelings f of G with $|v_f(i) - v_f(j)| \leq 1$ and $f(uv) = f(u)f(v)$, where $e_f(k)$ and $v_f(k)$ denote the number of edges and the number of vertices labeled with k. They established that $i_p(K_n) = \lfloor n/2 \rfloor^2$; $i_p(C_n) = 2$ if n is even; $i_p(W_n) = 2$ or 4 according as n is even or odd; $i_p(K_{2,n}) = 4$ or 2 according as n is even or odd; $i_p(K_2 + nK_1) = 3$ if n is even; $i_p(G \times P_2) \leq 2i_p(G)$; $i_p(G_1 \cup G_2) \leq i_p(G_1) + i_p(G_2) + 2 \min\{\Delta(G_1), \Delta(G_2)\}$ where G_1 and G_2 are graphs of odd order; and $i_p(G_1 \odot G_2) \leq i_p(G_1) + i_p(G_2) + 2\delta(G_2) + 3$ where G_1 and G_2 have odd order.

The notion of topological cordial labeling was introduced by Selistin and Asha in 2020 [3340] as follows. A *topological cordial labeling* of a graph $G = (V, E)$ is an injective function f from V to the set 2^X, where X is a set such that $0 < |X| < |V|$ and it forms a topology on X that is induced a function f^* from E to $\{0, 1\}$ defined by $f^*(uv) = 1$ if $|f(u) \cap f(v)| > 1$ and $f(uv) = 0$ otherwise and has the property that the number

of edges labeled with and 0 and the number of edges labeled with 1 differ by at most 1 and likewise for the vertices. A graph that satisfies the condition of the topological cordial labeling is called a . Selestin and Asha [3340] proved that the vertex switching of cycles are topological cordial graphs. Siva Prijith, Subbulakshmi, and Chandrakala [3043] proved that $P_2 \odot nK_1$, coconut trees, and various graphs obtained by joining fans and paths in special ways are topological cordial. Vertudes and Pedrano [4085] proved that the following graphs are topologial cordial: splitting of a bistars; a degree splitting bistars; firecrackers $F_{2,n}$; banana trees $BN_{2,k}$; jellyfish $J_{m,n}$, where $m-n=i$ where $i=0,1,2$ and m and n are positive integer. In [3341] Selistin and Asha proved that the Durer graph, the Herchel graph, and some constructed graphs are topological cordial graph. Siva Prijith, Subbulakshmi, and Chandrakala [3043] proved that $P_2 \odot nK_1$, coconut trees, and various graphs obtained by joining paths are topological cordial.

In [3879] Tenguria and Verma called a mapping f from $V(G)$ to $\{0,1,2\}$ such that each edge uv is labeled $(f(u)+f(v))$ mod 3, a 3-*total super sum cordial labeling* if $|f(i)-f(j)| \leq 1$ for $i,\ j \in \{0,1,2\}$, where $f(x)$ denotes the total number of vertices and edges labeled with x and for each edge $uv,\ |f(u)-f(v)| \leq 1$. A graph that has a 3-total super sum cordial labeling is called 3-*total super sum cordial* graph. They proved $P_m \cup P_n,\ C_m \cup C_n$, and $K_{1,m} \cup K_{1,n}$ are 3-total super sum cordial graphs. (These results also appeared in [3878] and [3880]).

Sridevi, Nagarajan, Nellaimurugan, and Navaneethakrishnan [3686] introduced the notion of *Fibonacci divisor cordial* labeling of a graph $G(V,E)$ as a bijection $f: V \to \{F_1, F_2, F_3, \ldots, F_p\}$, where F_i is the i^{th} Fibonacci number such that if each edge uv is assigned the label 1 if $f(u)$ divides $f(v)$ or $f(v)$ divides $f(u)$ and 0 otherwise, then the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. If a graph has a Fibonacci divisor cordial labeling, then it is called a *Fibonacci divisor cordial graph*. They proved $P_n,\ C_n,\ K_{2,n}$, and the subdivision of the bistar $B_{n,n}$ are Fibonacci divisor cordial graphs. They further proved that K_n $(n \geq 3)$ is not a Fibonacci divisor cordial graph. Ghosh and Pal [1268] proved that the graphs obtained by the switching of an arbitrary vertex of cycles and wheels, the duplication of an arbitrary vertex of cycles, the degree splitting graphs of paths, umbrella graphs, and bistars $B_{n,n}$ are also Fibonacci divisor cordial graphs. Sridevi, Palani, Navanaeethakrishnan, and Nagarajan [3687] proved that $K_{1,n}$ for $n \leq 9$ and $n=11$, W_n for $n=7,8,9,10$ and $C_m@P_n$ are Fibonacci divisor cordial graphs. They also we prove that $K_{1,n}$ for $n \geq 12$ and $n=10$, and W_n for $n=4,5,6$ and $n \geq 11$ are not Fibonacci divisor cordial graphs.

Vaidya and Vyas [4025] define the *tensor product* $G_1(T_p)G_2$ of graphs G_1 and G_2 as the graph with vertex set $V(G_1) \times V(G_2)$ and edge set
$\{(u_1,v_1)(u_2,v_2)|\ u_1u_2 \in E(G_1), v_1v_2 \in V(G_2)\}$. They proved the following graphs are product cordial: $P_m(T_p)P_n$; $C_{2m}(T_p)P_{2n}$; $C_{2m}(T_p)C_{2n}$; the graph obtained by joining two components of $P_m(T_p)P_n$ an by arbitrary path; the graph obtained by joining two components of $C_{2m}(T_p)P_{2n}$ by an arbitrary path; and the graph obtained by joining two components of $C_{2m}(T_p)C_{2n}$ by an arbitrary path.

In [2848] Ponraj introduced the notion of an $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial labeling of a graph. Let $S = \{\alpha_1, \alpha_2, \ldots, \alpha_k\}$ be a finite set of distinct integers and f be a function from a

vertex set $V(G)$ to S. For each edge uv of G assign the label $f(u)f(v)$. He calls f an $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-*cordial labeling* of G if $|v_f(\alpha_i) - v_f(\alpha_j)| \leq 1$ for all $i, j \in \{1, 2, \ldots, k\}$ and $|e_f(\alpha_i\alpha_j) - e_f(\alpha_r\alpha_s)| \leq 1$ for all $i, j, r, s \in \{1, 2, \ldots, k\}$, where $v_f(t)$ and $e_f(t)$ denote the number of vertices labeled with t and the number of edges labeled with t, respectively. A graph that admits an $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial labeling is called an $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-*cordial graph*. Note that an $(-\alpha, \alpha)$-cordial graph is simply a cordial graph and a $(0, \alpha)$-cordial graph is a product cordial graph. Ponraj proved that $K_{1,n}$ is $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial if and only if $n \leq k$ and for $\alpha_1 \neq 0,\ \alpha_2 \neq 0,\ \alpha_1 + \alpha_2 \neq 0$ proved the following: K_n is (α_1, α_2)-cordial if and only if $n \leq 2$; P_n is (α_1, α_2)-cordial; C_n is (α_1, α_2)-cordial if and only if $n > 3$; $K_{m,n}$ $(m, n > 2)$ is not (α_1, α_2)-cordial; the bistar $B_{n,n+1}$ is (α_1, α_2)-cordial; $B_{n+2,n}$ is (α_1, α_2)-cordial if and only if $n \equiv 1, 2 \pmod 3$; $B_{n+3,n}$ is (α_1, α_2)-cordial if and only if $n \equiv 0, 2 \pmod 3$; and $B_{n+r,n}, r > 3$ is not (α_1, α_2)-cordial. He also proved that if G is an (α_1, α_2)-cordial graph with p vertices and q edges, then $q \leq 3p^2/8 - p/2 + 9/8$. In [2848] Ponraj proved that combs $P_n \odot K_1$ are (α_1, α_2)-cordial; coronas $C_n \odot K_1$ are (α_1, α_2)-cordial for $n \equiv 0, 2, 4, 5 \pmod 6$; $C_3^{(t)}$ is not (α_1, α_2)-cordial; W_n is not (α_1, α_2)-cordial; and $\overline{K_n} + 2K_2$ is (α_1, α_2)-cordial if and only if $n = 2$.

In [4047] Varatharajan, Navanaeethakrishnan Nagarajan define a *divisor cordial* labeling of a graph G with vertex set V as a bijection f from V to $\{1, 2, \ldots, |V|\}$ such that an edge uv is assigned the label 1 if one $f(u)$ or $f(v)$ divides the other and 0 otherwise, then the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. If graph that has a divisor cordial labeling, it is called a *divisor cordial* graph. They proved the standard graphs such as paths, cycles, wheels, stars and some complete bipartite graphs are divisor cordial. They also proved that complete graphs are not divisor cordial. In [4048] they proved dragons, coronas, wheels, and complete binary trees are divisor cordial. For t copies $S_1, S_2, \ldots, S_t$ of an n-star $K_{1,n}$ they define $\langle S_1, S_2, \ldots, S_t\rangle$ as the graph obtained by starting with $S_1, S_2, \ldots, S_t$ and joining the central vertices of S_{k-1} and S_k to a new vertex x_{k-1}. They prove that $\langle S_1, S_2\rangle$ and $\langle S_1, S_2, S_3\rangle$ are divisor cordial. In [504] Barasara and Thakkar gave e some characterizations of various divisor cordial type labelings.

Vaidya and Shah [4007] proved that the splitting graphs of stars and bistars are divisor cordial and the shadow graphs and the squares of bistars are divisor cordial. In [4009] they proved that helms, flower graphs, and gears are divisor cordial graphs. They also proved that graphs obtained by switching of a vertex in a cycle, switching of a rim vertex in a wheel, and switching of an apex vertex in a helm admit divisor cordial labelings. Raj and Valli [3086] proved the following graphs divisor cordial: the duplication of a vertex of a cycle; graphs obtained by joining two wheels of the same size by a path of length at least 3; $G_v \odot K_1$, where G_v is a graph obtained by switching any vertex of a cycle of size at least 4; graphs obtained by joining the apex vertices of two shells of the same size to an isolated vertex; graphs obtained by joining the centers of two wheels of the same size to an isolated vertex; and a class of graphs obtained by removing certain edges from complete graphs. Bosmia and Kanani [695] proved that the graphs of the form $G \odot K_1$ where G any of the following admits a divisor cordial labeling: $K_{1,n},\ K_{2,n},\ K_{3,n}$, a wheel, a helm, a flower, a fan, a double fan, and a barycentric subdivision of a star. Bosmia

and Kanani [693] prove that the following graphs admit divisor cordial labelings: bistars, the splitting graph of bistars, the degree splitting graph of bistars, the shadow graph of bistars, the restricted square graph of bistars, the barycentric subdivision of bistars, and the corona product of a bistar with K_1. Thirusangu and Madhu [3893] proved that the extended duplicate graph of star and bistar graphs are divisor cordial graphs.

In [2537] Sugumaran and Mohan proved that the following graphs are divisor cordial graphs: degree splitting graph of $K_{1,n,n}$), the splitting graph of the graph obtained from two isolated vertices are joined by n paths of length 2, the $W-$ graph (the graph obtained from two copies of $K_{1,n}$ and identifying the last pendent vertex of the first copy with the first pendent edge of the second copy), $B(n) \odot_{u_m} K_1$, where $B(n) = 2P_n + K_1$ (*bow* graph, the Herschel graph H_s and switching of an apex vertex in the Herschel graph (see [686, p. 53]. In [3747] Sugumaran and Suresh proved that the following graphs are divisor cordial graphs: fans, Petersen graphs, $C_m \odot K_1$, friendship graphs F_n, and the switching of a end vertex in path P_n, switching of any one of the inner vertices of Petersen graph P_e.

Barasara and Thakkar [499] investigated divisor cordial labelings of armed crowns, closed helms, webs, and the one point union of t copies of C_n). In [495] they investigated divisor cordial labeling of the combs, P_n^2, C_n^2 for $n \neq 4$, and middle graphs of paths and cycles. Barasara and Thakkar [498] proved that ladders, circular ladders, Möbius ladders, total graphs of paths, and total graphs of cycles are divisor cordial graphs.

Barasara and Thakkar [500] proved that the triangular snake T_n and the quadrilateral snake Q_n are divisor cordial graphs. Moreover, the graphs $DS(P_n)$, $DS(S_n)(n \geq 5)$, $DS(C_n \odot K_1)$, $DS(P_n \odot K_1)$ $(n \geq 4)$, and the degree splitting graph of cycle C_n with one chord are divisor cordial graphs. In [501], Barasara and Thakkar showed that the C_n-quadrilateral snake is a divisor cordial graph. Also, the graphs $DS(T_n)$ $(n \geq 4)$, $DS(Q_n)$ $(n \geq 4)$, $DS(L_n)$ $(n \geq 3)$, and $DS(H_n)$ are divisor cordial graphs. Moreover, the graph obtained by duplication of an arbitrary edge by a new vertex in P_n, the graph obtained by duplication of an arbitrary vertex by a new edge in P_n, the graph obtained by duplication of an arbitrary vertex by a new vertex in P_n, and the graph obtained by duplication of an arbitrary edge by a new edge in P_n are divisor cordial graphs.

Vaidya and Barasara [3951] proved that every graph can be embedded as an induced subgraph of a product cordial graph, thereby ruling out any possibility of obtaining any forbidden subgraph characterization for product cordial graphs. They also proved that every connected graph can be embedded as an induced subgraph of a product cordial connected graph and every planar graph can be embedded as an induced subgraph of a product cordial planar graph. Similar results for total product cordial labeling were also obtained. In [3961] they prove that every graph can be embedded as an induced subgraph of a divisor cordial graph, thereby ruling out any possibility of obtaining any forbidden subgraph characterization for divisor cordial graphs. They also proved that every connected graph can be embedded as an induced subgraph of a divisor cordial connected graph and every planar graph can be embedded as an induced subgraph of a divisor cordial planar graph.

Gondalia [1300] proved that the Herschel graph, the fusion of any two adjacent vertices of degree 3 in a Herschel graph, the duplication of any vertex of degree 3 in a Herschel

graph, the switching of a central vertex in the Herschel graph, the joint sum of two copies of a Herschel graph, and the degree splitting of Herschel graph are divisor cordial graphs.

Motivated by the concept of divisor cordial labeling, Lourdusamy and Patrick [2332] introduced a new concept of divisor cordial labeling called sum divisor cordial labeling. Let $G = (V(G), E(G))$ be a simple graph and f be a bijection from $V(G)$ to $\{1, 2, \ldots, |V(G)|\}$. For each edge uv, assign the label 1 if 2 divides $f(u)+f(v)$ and the label 0 otherwise. The function f is called a *sum divisor cordial* labeling if the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. A graph which admits a sum divisor cordial labeling is called a *sum divisor cordial* . They prove that paths, combs, stars, complete bipartite, K_2+mK_1, bistars, jewels, crowns, flowers, gears, subdivisions of stars, the graph obtained from $K_{1,3}$ by attaching the center of $K_{1,n}$ at each pendent vertex of $K_{1,3}$, and the square $B_{n,n}$ are sum divisor cordial graphs. In [2340] Lourdusamy and Patrick proved that every transformed tree admits a sum divisor cordial labeling. They also investigated the sum divisor cordial labelings of the graphs obtained by identifying a vertex of graphs of paths, middle graphs of paths, and the splitting graphs of cycles. Krithika and Stanley [2042] proved that subdivided shell graphs, disjoint union of two subdivided shell graphs and uniform shell bow graphs are sum divisor cordial. In [2334] Lourdusamy and Patrick proved that the following graphs admit divisor cordial labelings: shadow graphs of stars, triangular ladders, slanting ladders, closed helms, the corona of a ladders and K_1, the corona of a triangular ladders and K_1, and the duplication of a vertex of a star or bistars $B_{n,n}$. Lakshmi and Bharathi [2087] proved that the path union and cycle of zero divisor graphs are sum divisor cordial.

Adalja and Ghodasara [68] proved that quadrilateral snake, double quadrilateral snake, alternate quadrilateral snake, double alternate quadrilateral snake are sum divisor cordial graphs. Krithika and Hilda [1439] proved that disjoint union of copies of strong face wheel graphs and prism graphs are sum divisor cordial graphs. In [2038] they proved that the disjoint union of k copies of C_n with two chords, pentagon snakes, and alternate pentagon snakes are sum divisor cordial. Adalja [67] proved that bistars, splitting graph of bistars, degree splitting graphs of bistars, shadow graphs of bistars, restricted square graphs of bistars, barycentric subdivisions of bistars, and corona products of bistars with K_1 admit sum divisor cordial labelings. In [66] Adalja proved that the graphs obtained by switching of a vertex in bistars, combs, apex vertex of helms, apex vertex of closed helms are sum divisor cordial. Adalja and Ghodasara [70] showed that cycles, cycles with one chord, cycles with a twin chord, cycles with a triangle, wheels, helms, webs, shells, flowers, and double fans admit sum divisor cordial labelings. Adalja and Patel [71] investigated that path union of two copies of a cycle, a cycle with triangle, wheels, helms, and closed helms admit sum divisor cordial labelings. Chuaephon [832] showed that $K_{m,n}$, the splitting graph of $K_{m,n}$, graphs obtained by attaching the apex vertices of two copies of a fan by an edge, and $\overline{F_m} + P_n$ admit sum divisor cordial labelings.

[67] new

[66] new

[70] new

[71] new

[832] new

In [3760], [3761], [3762], and [3763] Sugumaran and Rajesh proved that the following graphs are sum divisor cordial: swastiks, path unions of finite number of copies of swastiks, cycles of k copies of swastiks, when k is odd, jelly fish, Petersen graphs, theta graphs, the fusion of any two vertices in the cycle of swastiks, duplication of any vertex in the

cycle of swastiks, the switchings of a central vertex in swastiks, the path unions of two copies of a swastik, the star graph of the theta graphs, the Herschel graph, the fusion of any two adjacent vertices of degree 3 in Herschel graphs, the duplication of any vertex of degree 3 in the Herschel graph, the switching of central vertex in Herschel graph, the path union of two copies of the Herschel graph, H-graph H_n, when n is odd, $C_3@K_{1,n}$ (obtained by identifying the center of $K_{1,n}$ with a vertex of C_3), $< F_n^1 \triangle F_n^2 >$ (the graph obtained by joining the apex vertices of F_n^1 and F_n^2 by an edge and by joining the two apex vertices to a new vertex v') and open star of swastik graphs $S(t.Sw_n)$, when t is odd. In [3764], [3765] and [3767] Sugumaran and Rajesh proved that the following graphs are sum divisor cordial graphs: H-graph H_n, when n is even, duplication of all edges of the H-graph H_n, when n is even, $H_n \odot K_1$, $P(r.H_n)$, $C(r.H_n)$, plus graphs, umbrella graphs, path unions of odd cycles, kites, complete binary trees, drums graph (two copies of C_n that share exactly one vertex v and two copies of P_n that have an end point at v, twigs (graphs obtained from a path by attaching exactly two pendent edges to each internal vertices of the path), fire crackers of the form $P_n \odot S_n$, where n is even, and the double arrow graph DA_m^n, where $|m-n| \leq 1$ and n is even (obtained from $P_m \times P_n$ by adding two new vertices u and v such that each of the top row vertices of $P_m \times P_n$ are connected to u by an edge and the bottom row vertices of $P_m \times P_n$ are connected to v by an edge). Sugumaran and Rajesh [3746] proved that the following graphs are sum divisor cordial: $P_n + P_n$ (n is odd), $P_n@K_{1,m}$ (obtained by identifying an endpoint of P_n with the center of $K_{1,n}$), $C_n@K_{1,m}$ (n is odd), the graph obtained from W_{2n} by attaching the apex vertex of a copy of $K_{1,m}$ to each rim vertex, the graph obtained by joining the central vertices of two copies of $K_{1,n,n}$ by an edge and to a new vertex, the graph obtained by starting with C_n and, for each edge of C_n, adjoining a copy of C_n that shares an edge with the starting copy (the *flower graph* FL_n). In [69] Adalja and Ghodasara provide sum divisor cordial labelings for the graphs resulting from the duplication of graph elements in stars, cycles, and paths. In [3071] Raheem, Javaid, Numan, and Hasni investigated the existence sum divisor cordial labelings for the disjoint union of paths and subdivided stars. In [2350] Arockiams, Wencys, and Patrick investigate sum divisor cordial labelings of transformed tree related graphs.

Lourdusamy and Patrick [2339] proved P_2^n, $P_n \odot mK_1$, the subdivision graph of $P_n \odot mK_1$, $D_2(P_n)$, the total graphs of paths and cycles, the graphs obtained by duplication of each vertex of path by an edge, the shadow of cycles, the graphs obtained by duplication of each vertex of cycle by an edge, $C_4^{(t)}$, books, quadrilateral snakes, and alternate triangular snakes are sum divisor cordial graphs. Annamma and Vijitha [262] provided sum divisor cordial labelings of splitting graph of paths, the union of shadow graph of C_n and the shell graph $C_{n,n-3}$, and $2K_1 \odot (C_n \odot K_1)$. Krithika, Kowsalya, and Lavanya [2041] proved that graphs obtained from shell graph of C_n by joining vertices v_1 and v_n of C_n to a new vertex u (*ice cream graph*), circular fans with four chords, and circular fans with two chords are sum divisor cordial.

The *zero divisor graph* of a commutative ring R, $\Gamma(R)$, is the graph whose vertex set is set of zero divisors in R in which two distinct vertices u, v are adjacent if $uv = 0$. In [2348] Lourdusamy, Jenifer Wency, and Joy Beaula investigate the existence of sum

divisor cordial labelings of graphs of the form $\Gamma(Z_n), \Gamma(Z_m) + \Gamma(Z_n)$, and $\overline{\Gamma(Z_m)} + \Gamma(Z_n)$.

Murugesan [2602] introduced a *square divisor cordial* labeling. Let G be a simple graph and $f :\to \{1, 2, \ldots, |V(G)|\}$ a bijection. For each edge uv, assign the label 1 if either $(f(u))^2$ divides $f(v)$ or $(f(v))^2$ divides $f(u)$ and the label 0 otherwise. Call f a *square divisor cordial labeling* if $|e_f(0) - e_f(1)| \leq 1$. A graph with a square divisor cordial labeling is called a *square divisor cordial* graph. Murugesan proved that the following are square divisor cordial graphs: P_n $(n \leq 12)$, C_n $(3 \leq n \leq 11)$, wheels, some stars, some complete bipartite graphs, and some complete graphs. Vaidya and Shah [4013] proved that the following are square divisor cordial graphs: flowers, bistars, shadow graphs of stars, splitting graphs of stars and bistars, degree splitting graphs of paths and bistars.

Kanani and Bosmia [1822] define a *cube divisor cordial* labeling f of a simple graph G as a bijection from $V(G)$ to $\{1, 2, \ldots, |V(G)|\}$ such that, when each edge uv is assigned the label 1 if $(f(u))^3$ divides $f(v)$ or $(f(v))^3$ divides $f(u)$ and the label 0 otherwise, it holds that $|e_f(0) - e_f(1)| \leq 1$. A graph with a cube divisor cordial labeling is called a *cube divisor cordial* graph. They proved that the following graphs admit cube divisor cordial labelings: K_n if and only if $n = 1, 2, 3$; $K_{1,n}$ if and only if $n = 1, 2, 3$; $K_{2,n}$ for all n; $K_{3,n}$ if and only if $n = 1, 2$; bistars $B_{n,n}$ for all n ; and the graph obtained by joining leaves of one star of a bistar with the center of the opposite star of the bistar. Kanani and Bosmia [1822] prove: the edge deleted graph of a cube divisor cordial graph is also a cube divisor cordial graph; P_n is a cube divisor cordial graph if and only if $n = 1, 2, 3, 4, 5, 6, 8$; C_n is a cube divisor cordial graph if and only if $n = 3, 4, 5$; and wheels, flowers and fans are cube divisor cordial,

The Lucas sequence of numbers is a linear recurrence relation satisfying the conditions: $l_1 = 1, l_2 = 3$ and $l_n = l_{n-1} + l_{n-2}, n \geq 3$. Let $G = (V, E)$ be a simple graph and $f : V(G) \to \{l_1, l_2, \ldots, l_{|V(G)|}\}$ be a bijection such that each edge uv, assign the label 1 if either $f(u)$ divides $f(v)$ or $f(v)$ divides $f(u)$ and label 0 otherwise. In [3768] Sugumaran and Rajesh call such an f a *Lucas divisor cordial labeling* if $|e_f(0) - e_f(1)| \leq 1$. A graph with a Lucas divisor cordial labeling is called a *Lucas divisor cordial* graph. In [3768] Sugumaran and Rajesh proved that the following graphs are Lucas divisor cordial graphs: bistars, jelly fish, square graphs of bistars, switching of a vertex in cycles, and switching of a pendent vertex in paths.

A variation of divisor cordial labeling called vertex odd divisor cordial labeling was introduced by Muthaiyan and Pugalenthi (see [2603]) as follows. Let G be a graph with p vertices and a bijection f from $V(G)$ to $\{1, 3, 5, \ldots, 2p-1\}$ such that if each edge uv is assigned the label 1 if $f(u)$ divides $f(v)$ or $f(v)$ divides $f(u)$, and the label 0 otherwise. The function f is called a *vertex odd divisor cordial labeling* if $|e_f(0) - e_f(1)| \leq 1$. A graph with vertex odd divisor cordial labeling is called a *vertex odd divisor cordial graph.* Muthaiyan and Pugalenthi (see [2603]) proved paths, cycles, $K_{2,n}, K_{1,n} \cup K_{1,m}$, helms, flowers, $< K_{1,n}^{(1)}, K_{1,n}^{(2)} >$, the switching of the apex vertex in helms, and the splitting graph of stars are vertex odd divisor cordial graphs under some conditions. In [2603] Muthaiyan and Pugalenthi proved the following graphs have vertex odd divisor cordial labelings: wheels, the switching of a pendent vertex in paths and cycles, bistars $B_{n,n}$, the subdivision graph of $K_{1,n}, B_{n,n}^2, DS(B_{n,n})$, the splitting graph of $B_{n,n}$, and $< K_{1,n}^{(1)}, K_{1,n}^{(2)}, K_{1,n}^{(3)} >$.

Let G_1 and G_2 be two copies of any graph G that has an apex vertex. The graph obtained by joining the apex vertices of G_1 and G_2 by an edge and the graph obtained by joining the apex vertices of G_1 and G_2 by an edge and by joining the two apex vertices to a new vertex v', is denoted $G_1 \triangle G_2$. For any vertex u of $K_{m,n}$ the graph obtained by joining u to a new pendent vertex is denoted by $K_{m,n} \odot u(K_1)$. In [3751] Sugumaran and Suresh proved that the following graphs are vertex odd divisor cordial graphs: the shadow graph of $K_{1,n}$, $K_{2,n} \odot u(K_1)$, $K_{1,n} \triangle K_{1,n}$, the subdivision of the edge between the apex vertices of $B_{n,n}$, and the graph $K_{1,n} * P_{n+2}$ (the graph obtained by identifying an end vertex of P_{n+2} with the apex vertex of $K_{1,n}$). In [3750] they showed that the graphs $F_n \triangle F_n$, $K_{1,n} \triangle K_{1,n} \triangle K_{1,n}$, $K_{1,n} \triangle K_{1,n} \triangle K_{1,n} \triangle K_{1,n}$, theta graphs, and switching of a vertex in a Petersen graph are vertex odd divisor cordial graphs. In [3752] they proved that gears, switching of an apex vertex in $S(K_{1,n})$, $P_2 + mK_1$, $C(n, n-3)$, and $C(n, n-4)$ are vertex odd divisor cordial graphs. In [3748] they showed that the globe $Gl(n)$, jewels, $G * W_n$ (appending the central vertex of wheel W_n with any one of the vertices of G), $G * C(n, n-3)$, and wheels are vertex odd divisor cordial graphs.

Let $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ be an injective map. For each edge uv assign the label r where r is the remainder when $f(u)$ is divided by $f(v)$ or $f(v)$ is divided by $f(u)$ according as $f(u) \geq f(v)$ or $f(v) \geq f(u)$. The function f is called a *remainder cordial* labeling of G if $|e_f(0) - e_f(1)| \leq 1$ where $e_f(0)$ and $e_f(1)$ respectively denote the number of edges labeled with even integers and number of edges labeled with odd integers. A graph G with admits a remainder cordial labeling is called a *remainder cordial* graph. In [2863] Ponraj, Annathurai, and Kala investigated the remainder cordial behavior of $S(K_{1,n})$, $S(B_{n,n})$, $S(W_n)$ and union of some star related graphs. Ponraj, Gayathri, and Somasundaram [2885] investigated the 4-remainder cordial labeling behavior of parachutes, twigs, and kayak paddles graphs.

The *Durer* graph has vertex set $V(C_n) \cup \{u_i, v_i : 1 \leq i \leq n\}$ and edge set $E(C_n) \cup \{u_i v_i : 1 \leq i \leq n\} \cup \{u_i u_{i+2} : 1 \leq i \leq n-2\} \cup \{u_n u_2, u_{n-1} u_1\}$. In [2865] Ponraj, Annathurai, and Kalainvestigated the 4-remainder cordial labeling behavior of prisms, crossed prisms, webs, triangular snakes, $L_n \odot K_1$, Durer graphs, and dragon graphs.

A *double divisor cordial labeling* of a graph $G(V, E)$ is a bijective function ψ from V to $\{1, 2, 3, \ldots, |V|\}$ such that each edge ab is given label 1 if $2\psi(a) | \psi(b)$ or $2\psi(b) | \psi(a)$ and 0 otherwise, then the number of edges labeled 0 and the number of edges labeled 1 differ by at most 1. An graph that admits a double divisor cordial labeling is said to be a *double divisor cordial graph*. In [2785] Parthiban and Sharma proved that the following graphs admit double divisor cordial labelings: full binary trees, P_n $(n \geq 3)$, C_n, W_n (n odd), helms, fans, stars, bistars, jelly fish, and friendship graphs. They also prove that K_n $(n \geq 5)$ does not admit a double divisor cordial labeling. Sharma and Parthiban [3490] proved that the following graphs admit double divisor cordial labelings: the barycentric subdivision of stars and bistars $B(n, n)$; the splitting graph of $K_{1,n}$ $(n \geq 2)$; the splitting graph of $B(m, n)$ $(m, n \geq 2)$; $\langle K^{(1)}_{1,n}, K^{(2)}_{1,n} \rangle$; $\langle K^{(1)}_{1,n}, K^{(2)}_{1,n}, K^{(3)}_{1,n} \rangle$; the graph constructed by connecting an apex vertex of one copy of $B_{n,n}$ to a new vertex r an apex vertex of another copy of $B_{n,n}$ to another vertices s; and $K_{i,n} \odot K_1$ for $i = 1, 2$, and 3.

7.6 Other Cordial Labelings

In [502] Barasara and Thakkar introduced a new concept called power cordial labeling which is defined as follows: a *power cordial labeling* of a graph $G = (V, E)$ is a bijection $f : V(G) \to \{1, 2, \ldots, |V(G)|\}$ such that when an edge $e = uv$ is assigned the label 1 if $f(u) = (f(v))^n$ or $f(v) = (f(u))^n$, for some $n \in \mathbb{N} \cup \{0\}$ and the label 0 otherwise, the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. A graph that admits a power cordial labeling is called a *power cordial graph*. They proved that the helm H_n is a power cordial graph for $n \leq 9$ and $n = 13$ and not a power cordial graph for $n = 10, 11, 12$ and $n \geq 14$, the flower Fl_n is a power cordial graph, the gear graph G_n is a power cordial graph for $n \leq 9$ and $n = 13$ and not a power cordial graph for $n = 10, 11, 12$ and $n \geq 14$, the fan F_n is a power cordial graph, the jewel graph J_n is a power cordial graph, the graph $S(K_{1,n})$ is a power cordial graph, the graph $2K_{1,n}$ is a power cordial graph, the graph $S'(K_{1,n})$ is a power cordial graph for $n \leq 5$ and $n = 7$ and not a power cordial graph for $n = 6$ and $n \geq 8$, the graph $G = \langle K_{1,n}^{(1)}, K_{1,n}^{(2)} \rangle$ is a power cordial graph and the graph $S'(B_{n,n})$ is a power cordial graph for $n \leq 8$ and not a power cordial graph for $n > 8$. In [503] Barasara and Thakkar proved that any graph G can be embedded as an induced subgraph of a power cordial graph. This rules out the possibility of forbidden subgraph characterization for power cordial labeling. Also, they showed that for given positive integer n, there is a power cordial graph G that has n vertices. If G is a power cordial graph of even size, then $G - e$ is also a power cordial graph for all $e \in E(G)$, If G is a power cordial graph of odd size, then $G - e$ is also a power cordial graph for some $e \in E(G)$, for G be a power cordial graph of even size and $G * K_n$, then adding an edge between any two nonadjacent vertices of G is also a power cordial graph. Moreover they proved that for given graph G with size n and G has a vertex v such that $\left\lfloor \frac{n}{2} \right\rfloor \leq d(v) \leq \left\lceil \frac{n}{2} \right\rceil$ then G is a power cordial graph.

In [2971] Ponraj, Sathish Narayanan, and Ramasamy introduced a new graph labeling called parity combination cordial labeling. Let G be a (p, q)-graph. Let f be an injective map from $V(G)$ to $\{1, 2, \ldots, p\}$. For each edge xy, assign the label $\binom{x}{y}$ or $\binom{y}{x}$ according as $x > y$ or $y > x$. Call f a *parity combination cordial* labeling if f is a one to one map and $|e_f(0) - e_f(1)| \leq 1$, where $e_f(0)$ and $e_f(1)$ denote the number of edges labeled with an even number and odd number, respectively. A graph with a parity combination cordial labeling is called a *parity combination cordial graph*. They proved that the following are parity combination cordial graphs: paths, cycles, stars, triangular snakes, alternate triangular snakes, olive trees, combs, crowns, fans, umbrellas, P_n^2, helms, dragons, bistars, butterfly graphs, and graphs obtained from C_n and $K_{1,m}$ by unifying a vertex of C_n and a pendent vertex of $K_{1,m}$. They also proved that W_n admits a parity combination cordial labeling if and only if $n \geq 4$ and conjectured that for $n \geq 4$, K_n is not a parity combination cordial graph. In [2970], Ponraj, Rajpal Singh, and Sathish Narayanan proved that if G is a parity combination cordial graph, then $G \cup P_n$ is also parity combination cordial if $n \neq 2, 4$. In [3363] Seoud and Aboshady surveyed all graphs of order at most six with regard to whether they have a parity combination cordial labeling or not. They obtained an upper bound for the number of edges of any graph that satisfies a certain condition,

and described the parity combination cordial labelings for two families of graphs.

In [2630] Naduvath defines the notion of the set-cordial labeling of a graph as follows. For a non-empty set X, a function f from the vertices of a graph G to the power set of X is said to be a *set-cordial* labeling of G if $|f(v_i)| - |f(v_j)| = \pm 1$ for all edges v_iv_j of G. A graph that admits a set-cordial labeling is called a *set-cordial* graph . He proves that paths are set-cordial and that a graph G admits a set-cordial labeling if and only if G is bipartite. He defines the *glutting number* of a graph G as the minimum number of edges of G that can be removed so that the resulting graph admits a set-cordial labeling and provides the glutting number of wheels, helms, and complete graphs.

In 2011 Murugan and Selvaraj [2588] introduced the concept of V-*cordial* labeling of a graph G with vertex set V as a bijective function $\phi : V \to \{0, 1\}$ such that the induced function $\phi^* : E \to \{0, 1\}$ is defined by $\phi(uv) = 0$ if $\phi(u) = \phi(v) = 0$ and 1 otherwise with the condition that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1, and likewise for the edges. A graph that admits a V-cordial labeling is called a V-*cordial* graph. In 2015 Murugan and Mathubala introduced the concept of homo-cordial labeling as follows. A *homo-cordial* labeling of a graph G with vertex set V is a bijection $\phi : V \to \{0, 1\}$ such that the induced function $\phi^* : E \to \{0, 1\}$ given by $\phi^*(uv) = 1$, if $\phi(u) = \phi(v)$ and 0 otherwise with the condition that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1, and likewise for the edges. A graph that admits a homo-cordial labeling is called a *homo-cordial* graph. In [2589] Murugan and Vidhya say that a *hetro-cordial* labeling of a graph G with vertex set V is a bijection $\phi : V \to \{0, 1\}$ such that the induced function $\phi^* : E \to \{0, 1\}$ given by $\phi^*(uv) = 0$, if $\phi(u) = \phi(v)$ and 1, otherwise with the condition that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1, and likewise for the edges. A graph that admits a hetro-cordial labeling is called a *hetro-cordial* graph. Murugan, and Mathubala [2574] proved that odd cycles, shadow graphs of cycles, graphs obtained by identifying the end points of n paths of length 2 (globe graphs), and double triangular snakes are homo-cordial graphs. Murugan, and Selva Vidhya [2589] proved that paths, combs, fans, double fans, and ladders are hetro-cordial graphs. Murugan and Selvaraj [2588] proved that paths, fans double fans, and combs are V cordial graphs. Bala, Sundarraj, and Thirusangu [452] proved existence of a homo-cordial labeling, a hetro-cordial labeling, and a V-cordial labeling for the extended triplicate graph of a comb.

In [2672] Nicholas and Maya introduced the concept of an *integer edge cordial* labeling of a graph G with edge set E as an injective map f from E to $[-q/2, \ldots, q/2]$ or $[\lfloor -q/2 \rfloor, \ldots, \lfloor q/2 \rfloor]$ as q is even or odd, which induces a vertex labeling $f^* : V \to \{0, 1\}$ such that, a vertex u is assigned the label 1 if $\sum(f^*(e_i) \geq 0$ taken over all i, and 0 otherwise, and the number of vertices labeled with 1 and the number of vertices labeled with 0 differ by at most by 1. A graph that has integer edge cordial labeling is called an *integer edge cordial graph*. They proved that P_n $(n \geq 3)$, cycles, W_n $(n > 3)$, helms, closed helms, $K_{1,2n}$, and flower graphs are integer edge cordial graphs. They also proved that $K_{n,n}$ is not integer edge cordial and that $K_{n,n} \setminus M$ is integer edge cordial if n is even, where M is a perfect matching of $K_{n,n}$. In [2671] Nicholas and Maya proved the following graphs [2671] new

are integer cordial: cycles, K_n $(n > 3)$, wheels W_n $(n > 3)$, stars, helms, closed helms, $K_{n,n}$ if and only if n is even, and $K_{n,n} - M$, where M is a perfect matching.

For a planar graph $G(p, q)$ suppose that $g : E(G) \to [\frac{-q}{2}, \ldots, \frac{q}{2}]$ or $[-\lfloor\frac{q}{2}\rfloor, \ldots, \lfloor\frac{q}{2}\rfloor]$ as q is even or odd, is an injective map that induces a vertex labeling $g^* : V(G) \to \{0, 1\}$ defined by $g^*(v) = 1$ if the sum of all $g(e)$ over all edges e of G adjacent to v is nonnegative and $g^*(v) = 0$ otherwise, and the face labeling function g^{**} from the faces of G to $\{0, 1\}$ defined by $g^{**}(f) = 1$, if the sum of $g(e)$ over all edges e of f is nonnegative, and $g^{**}(f) = 0$ otherwise. Then g is called a *face integer edge cordial labeling* if the number of vertices labeled 0 and the number of vertices labeled 1 differ by at most 1 and the number of faces labeled 0 and the number of faces labeled 1 differ by at most 1. A graph that admits a face integer edge cordial labeling is called *face integer edge cordial*. Sheriff and Abbas [3506] gave face integer edge cordial labelings of the duplication of each vertex by an edge in nontrivial fans, the planar graph G' obtained from joining the outer vertex of the two cop

The concept of multiply divisor cordial labeling was introduced in 2019 by Gondalia and Rokad as follows. A *multiply divisor cordial labeling* of a graph G having vertex set V is a bijective h from V to $\{1, 2, \ldots, |V|\}$ such that an edge xy is assigned the label 1 if 2 divides $h(x) \cdot h(y)$ and 0 otherwise, then the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. A graph having multiply divisor cordial labeling is said to be a *multiply divisor cordial graph*. In [1303] and [1304] Gondalia Rokad proved that cycles, cycles with one chord, cycles with a twin chord, cycles with a triangle, paths, stars, jelly fish, and coconut trees are multiply divisor cordial graph. Further they proved that ring sum of star with a cycle, the ring sum of star with a cycle having one chord, the ring sum of star with a cycle having twin chords, the ring sum of star with a cycle having triangle, the ring sum of star with double fan, the ring sum of star with double wheel, and the ring sum of star with helm graph are multiply divisor cordial labeling.

In [1301] Gondalia proved the following graphs admit multiply divisor cordial labelings: graphs obtained by joining two copies of Petersen graph by a path of arbitrary length; graphs obtained by joining two copies of C_n with one chord by a path of arbitrary length; graphs obtained by joining two copies of C_n with twin chords by a path of arbitrary length; graphs obtained by joining two copies of C_n with a triangle by a path of arbitrary length; graphs obtained by joining two copies of a fan F_n by a path of arbitrary length; and graphs obtained by joining two copies of flower graph Fl_n by a path of arbitrary length.

For a graph $G(V, E)$ with p vertices and q edges Ponraj and Sutharson [2989] introduced the notion of fractional product cordial labeling of graph as follows.
Let

$$M = \begin{cases} 1, 2, \cdots, \frac{p}{2}, \frac{1}{2}, \frac{1}{3}, \cdots, \frac{2}{p+2}, & \text{if } p \text{ is even} \\ 1, 2, \cdots, \frac{p-1}{2}, \frac{1}{2}, \frac{1}{3}, \cdots, \frac{2}{p+3}, & \text{if } p \text{ is odd.} \end{cases}$$

Let $\chi : V(G) \to M$ be a bijection. For each edge xy assign the label $\chi(\lceil\chi(x)\chi(y)\rceil)$, called a *fractional product cordial labeling* if $|\Pi_\chi(0) - \Pi_\chi(1)| \leq 1$, where $\Pi_\chi(1)$ and $\Pi_\chi(0)$, respectively denotes the number of edges labeled with 1 and not labeled with 1. A graph

with a fractional product cordial labeling is called a *fractional product cordial* graph. Ponraj and Sutharson investigated the fractional product cordial labeling behavior of paths, stars, cycles, webs, dumbbells, jelly fish, jewel graphs tadpoles, snake graphs, helms, sunflowers, and subdivision of combs. In [2992] Ponraj and Sutharson investigated the fractional product cordial labeling behaviorf parachutes, twigs, kayak paddales, alternate triangular snakes, and alternate quadrilateral snakes. [2992] new

In [853] Clarisse, Prudencio, and Rulete gave the following definition. Let G be a connected graph of order n and let p be an odd prime. The bijective function $f : V(G) \to \{1, 2, \ldots, n\}$ is called a *Legendre product cordial labeling modulo p*, if there is an induced function $f_{p*} : E(G) \to \{0, 1\}$, defined by $f_{p*}(uv) = 0$ whenever $(f(u)f(v))/p) = -1$ or $f(u)f(v) = 0 \pmod p$, and $f_{p*}(uv) = 1$ whenever $(f(u)f(v))/p = 1$, that satisfies the condition that the number of edges with labels 0 and the number of edges labeled with 1 differ by at most 1. They explored the Legendre product cordial labeling modulo p of $P_{p-1}, C_{p-1}, K_{p-1}$ and several other families of graphs. [853] new

For an odd prime p and a simple connected graph $G = (V, E)$, Andoyo and Rulete [240] call a bijective function $f : V(G) \to \{1, 2, \ldots, |V|\}$ a *Legendre cordial modulo p* labeling, if the induced function $f_{p^*} : E(G) \to \{0, 1\}$ defined by $f_{p^*}(uv) = 0$ when $f(u)+f(v) = -1$ or $f(u)+f(v) = 0$, and $f_{p^*}(uv) = 1$ when $(f(u)+f(v))/p = 1$, satisfies the condition that the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. They proved that $P_p, C_p, K_{1,p}$, tadpoles $T_{p,p}$, kayak paddles $KP_{p,p,p}$, bistars $B_{p,p}$, and fans F_{p+1} admit a Legendre cordial labeling modulo p, where p is an odd prime.

In 2024 Tharani and Saradha [3883] introduced a stronger form of cordial labelings as follows. A *discrete labeling* of a graph is a cordial labeling of the vertices such that the edges receive labels depending on the incident vertex labels using EX-OR operation with the condition that for every vertex the cardinality of neighbors labeled 0 and 1 differs by at most 1. They proved that paths, combs, brooms, bistars, the Herschel graph, the Petersen graph, C_n if and only if $n = 0 \pmod 4$, $K_{m,n}$, and K_n if and only if $n \leq 3$ admit discrete labelings. In [660] Bijukumar showed that the following graphs admit discrete labelings: super subdivision of P_n by $K_{2,n}$, graphs obtained by switching of one of the pendent vertices of P_n, and shadow graphs of paths, stars, and bistars.

In [939] Devi, Bernie, Faridha, and Rajeshwaran introduced the following variation of cordial labelings. A labeling $h : V(G) \to \{1, 2\}$ with the induced label of an edge uv defined by $h^*(uv) = \lfloor (h(u)h(v) + \lfloor h(u)/h(v) \rfloor)/2 \rfloor$ labeling is said to be TPD-*edge balanced*, if the number of edges labeled with 1 and the number of edges labeled with 2 differ by at most 1. They provided a partition-based characterization theorem and gave TPD-edge balanced labelings for paths, cycles, stars and fan graphs. In addition, they discussed structural properties, extremal bounds, and potential applications to communication network and fault-tolerant systems. [939] new

Let η be a fixed positive integer. Let S be a subset of $\mathbb{Z}$, let $\star : S \times S \to \mathbb{Z}$ be a binary function, and let $\zeta_\eta : \{\xi \in \mathbb{Z} : \gcd(\xi, \eta) = 1\} \to \{0, 1\}$ be a function. For a simple connected graph G of order n, a bijective function $f : V(G) \to S$ (where $|S| = n$) is called an *arithmetic cordial labeling modulo* η under the arithmetic structure $\langle S, \zeta_\eta, \star \rangle$ if the induced function $f^*_\eta : E(G) \to \{0, 1\}$, defined by $f^*_\eta(ab) = 1$ whenever

$\gcd(f(a) \star f(b), \eta) = 1$ and $\zeta_\eta(f(a) \star f(b)) = 1$, and $f^*_\eta(ab) = 0$ otherwise, satisfies the condition $|e_{f^*_\eta}(0) - e_{f^*_\eta}(1)| \leq 1$, where $e_{f^*_\eta}(i)$ is the number of edges with label i for $i \in \{0, 1\}$.

In 2026 Andoyo, Clarisse, Prudencio, and Rulete [239] introduced the notion of arithmetic cordial labeling of graphs as follows. Let η be a fixed positive integer. Let S be a subset of $\mathbb{Z}$, $\star : S \times S \to \mathbb{Z}$ be a binary function, and $\zeta_\eta : \{\xi \in \mathbb{Z} : \gcd(\xi, \eta) = 1\} \to \{0, 1\}$ be a function. For a simple graph G of order n, a bijective function $f : V(G) \to S$ (where $|S| = n$) is called an *arithmetic cordial labeling modulo η under the arithmetic structure $\langle S, \zeta_\eta, \star\rangle$ if the induced function $f^*_\eta : E(G) \to \{0, 1\}$*, defined by $f^*_\eta(ab) = 1$ whenever $\gcd(f(a) \star f(b), \eta) = 1$ and $\zeta_\eta(f(a) \star f(b)) = 1$; otherwise, $f^*_\eta(ab) = 0$, satisfies the condition $|e_{f^*_\eta}(0) - e_{f^*_\eta}(1)| \leq 1$, where $e_{f^*_\eta}(i)$ is the number of edges with label i $(i = 0, 1)$. Andoyo et al. explored the arithmetic cordial labelings of stars, ladders, alternate cycle snakes, join graphs, corona graphs, and tensor product graphs. In [238] Andoja studied the arithmetic cordial labelings of the product operations: corona, lexicographic, Cartesian, tensor, and strong under the operation of addition. [239] new [238] new

7.7 Edge Product Cordial Labelings

Vaidya and Barasara [3951] introduced the concept of edge product cordial labeling as edge analogue of product cordial labeling. An *edge product cordial labeling* of graph G is an edge labeling function $f : E(G) \to \{0, 1\}$ that induces a vertex labeling function $f^* : V(G) \to \{0, 1\}$ defined as $f^*(u) = \prod\{f(uv) \mid uv \in E(G)\}$ such that the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1 and the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1. A graph with an edge product cordial labeling is called an *edge product cordial graph*.

In [3951], [3961], [3954], [3955], and [3958] Vaidya and Barasara proved the following graphs are edge product cordial: C_n for n odd; trees with order greater than 2; unicyclic graphs of odd order; $C_n^{(t)}$, the one point union of t copies of C_n for t even or t and n both odd; $C_n \odot K_1$; armed crowns $C_m \odot P_n$; helms; closed helms; webs; flowers; gears; shells S_n for odd n; tadpoles $C_n@P_m$ for $m + n$ even or $m + n$ odd and $m > n$ while not edge product cordial for $m + n$ odd and $m < n$; triangular snakes; for odd n, double triangular snakes DT_n, quadrilateral snakes Q_n and double quadrilateral snakes DQ_n; P_n^2 for odd n; $M(P_n)$, $T(P_n)$; $S'(P_n)$ for even n; the tensor product of P_m and P_n; and the tensor product of C_n and C_m if m and n are even. In [3959] Vaidya and Barasara investigate product and edge product cordial labelings of the degree splitting graphs of paths, shells, bistars, and gear graphs. They proved the following graphs are not edge product cordial: C_n for n even; K_n for $n \geq 4$; $K_{m,n}$ for $m, n \geq 2$; wheels; the one point union of t copies of C_n for t odd and n even; shells S_n for even n; tadpoles $C_n@P_m$ for $m + n$ odd and $m < n$; for n even double triangular snake DT_n, quadrilateral snake Q_n and double quadrilateral snake DQ_n; double fans; C_n^2 for $n > 3$; P_n^2 for even n; $D_2(C_n)$, $D_2(P_n)$; $M(C_n)$; $T(C_n)$; $S'(C_n)$; $S'(P_n)$ for odd n; $P_m \times P_n$ and $C_m \times C_n$; the tensor product of C_n and C_m if m or n odd; and $P_n[P_2]$ and $C_n[P_2]$.

Barasara [494] proved that C_n with one chord except when n is even and the chord joins vertices that are at diameter distance, C_n with twin chords except when n is even and the chords join vertices that are at diameter distance, and $P_n \odot K_1$ are edge product cordial graphs, whereas triangular ladders are not edge product cordial. In [3952], Barasara proved that C_n with one chord, C_n with twin chords, $P_n \odot K_1$, and triangular ladders are total edge product cordial graphs. Vaidya and Barasara [3960] proved that every graph can be embedded as an induced subgraph of an edge product cordial graph, thereby ruling out any possibility of obtaining any forbidden subgraph characterization for edge product cordial graphs. They also proved that every connected graph can be embedded as an induced subgraph of an edge product cordial connected graph and every planar graph can be embedded as an induced subgraph of an edge product cordial planar graph. In [493], Barasara proved that the shadow graph of W_n and the splitting graph of W_n are not edge product cordial graphs, whereas the middle graph of W_n is edge product cordial for $n > 5$, and the total graph of W_n is edge product cordial for $n > 8$. He also investigated total edge product cordial labeling of the shadow graph, the splitting graph, the middle graph, and the total graph of W_n.

Vaidya and Barasara [3961] proved that every graph can be embedded as an induced subgraph of a total edge product cordial graph thereby ruling out any possibility of obtaining any forbidden subgraph characterization for total edge product cordial graphs. They also proved that a connected graph can be embedded as an induced subgraph of a total edge product cordial connected graph and every planar graph can be embedded as an induced subgraph of a total edge product cordial planar graph.

Prajapati and Shah [3024] proved the following graphs are edge product cordial: graphs obtained from a crown by duplication of a vertex, duplication of a vertex by an edge, or duplication of an edge by a vertex; graphs obtained from a gear graph by duplication of each of the vertices of degree three by an edge; and the graph obtained from a helm by duplication of each of the pendent vertices by a new vertex. In [3015] Prajapati and Patel provided results about the existence of edge product cordial labelings closed webs, lotus inside a circle, and sunflower graphs.

Vaidya and Barasara [3956] introduced the concept of a total edge product cordial labeling as edge analogue of total product cordial labeling. An *total edge product cordial labeling* of graph G is an edge labeling function $f : E(G) \to \{0, 1\}$ that induces a vertex labeling function $f^* : V(G) \to \{0, 1\}$ defined as $f^*(u) = \prod\{f(uv) \mid uv \in E(G)\}$ such that the number of edges and vertices labeled with 0 and the number of edges and vertices labeled with 1 differ by at most 1. A graph with total edge product cordial labeling is called a *total edge product cordial graph*.

In [3956] and [3957] Vaidya and Barasara proved the following graphs are total edge product cordial: C_n for $n \neq 4$; K_n for $n > 2$; W_n; $K_{m,n}$ except $K_{1,1}$ and $K_{2,2}$; gears; $C_n^{(t)}$, the one point union of t copies of C_n; fans; double fans; C_n^2; $M(C_n)$; $D_2(C_n)$; $T(C_n)$; $S'(C_n)$; P_n^2 for $n > 2$; $M(C_n)$; $D_2(C_n)$ for $n > 2$; $T(C_n)$; $S'(C_n)$. Moreover, they prove that every edge product cordial graph of either even order or even size admits total edge product cordial labeling. Bača, Irfan, Javad, and Semaničová-Feňočová [375] investigated the existence of total edge product cordial labeling of toroidal fullerenes and for Klein-

bottle fullerenes. Prajapati and Patel [3016] proved that the one point union of t copies of a wheel with a rim vertex in common is edge product cordial if and only is t is even; all pentagonal snakes (obtained from the path by replacing every edge of a path by C_5) are edge product cordial; and a double pentagonal snakes (two pentagonal snakes that have a common path) is edge product cordial if and only is t is odd.

In [3033] Prasad and Maheswari developed a technique for coding a secret messages using sunflower graphs by subdividing edges and applying edge product cordial labeling.

7.8 Difference Cordial Labelings

Ponraj, Sathish Narayanan, and Kala [2950] introduced the notion of difference cordial labelings. A *difference cordial labeling* of a graph G is an injective function f from $V(G)$ to $\{1, \ldots, |V(G)|\}$ such that if each edge uv is assigned the label $|f(u)-f(v)|$, the number of edges labeled with 1 and the number of edges not labeled with 1 differ by at most 1. A graph with a difference cordial labeling is called a *difference cordial* graph. Jeba Jesintha, Devakirubanithi, and Jasma Pershi [1604] proved that the graphs obtained by switching a central vertex in the path is difference cordial and the crown graph with r copies of fan graph attached to it is also difference cordial. In [2393] Uma Maheswari, Margaret, Jebarani, and Balaji provide a technique for coding a message using GMJ code on the Fibonacci web graph by applying the difference cordial labeling.

The following definitions appear in [2951], [2938], [2939], and [2940]. A *double triangular snake* DT_n consists of two triangular snakes that have a common path; a *double quadrilateral snake* DQ_n consists of two quadrilateral snakes that have a common path; an *alternate triangular snake* $A(T_n)$ is the graph obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertex v_i (that is, every alternate edge of a path is replaced by C_3); a *double alternate triangular snake* $DA(T_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to two new vertices v_i and w_i; an *alternate quadrilateral snake* $A(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i and w_i respectively and then joining v_i and w_i (that is, every alternate edge of a path is replaced by a cycle C_4); a *double alternate quadrilateral snake* $DA(Q_n)$ is obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to new vertices v_i, x_i and w_i and y_i respectively and then joining v_i and w_i and x_i and y_i.

In [2939] and [2940] Ponraj and Sathish Narayanan define the *irregular triangular snake* IT_n as the graph obtained from the path $P_n : u_1, u_2, \ldots, u_n$ with vertex set $V(IT_n) = V(P_n) \cup \{v_i : 1 \leq i \leq n \leq 2\}$ and the edge set $E(IT_n) = E(P_n) \cup \{u_iv_i, v_iu_{i+2} : 1 \leq i \leq n-2\}$. The *irregular quadrilateral snake* IQ_n is obtained from the path $P_n : u_1, u_2, \ldots, u_n$ with vertex set $V(IQ_n) = V(P_n) \cup \{v_i, w_i : 1 \leq i \leq n-2\}$ and edge set $E(IQ_n) = E(P_n) \cup \{u_iv_i, w_iu_{i+2}, v_iw_i : 1 \leq i \leq n-2\}$. They proved the following graphs are difference cordial: triangular snakes T_n, quadrilateral snakes, alternate triangular snakes, alternate quadrilateral snakes, irregular triangular snakes, irregular quadrilateral snakes, double triangular snakes DT_n if and only if $n \leq 6$, double quadrilateral snakes, double alternate triangular snakes $DA(T_n)$, and double alternate quadrilateral snakes.

In [2950], [2937], [2951], and [2938] Ponraj, Sathish Narayanan, and Kala proved the following graphs have difference cordial labelings: paths; cycles; wheels; fans; gears; helms; $K_{1,n}$ if and only if $n \leq 5$; K_n if and only if $n \leq 4$; $K_{2,n}$ if and only if $n \leq 4$; $K_{3,n}$ if and only if $n \leq 4$; bistar $B_{1,n}$ if and only if $n \leq 5$; $B_{2,n}$ if and only if $n \leq 6$; $B_{3,n}$ if and only if $n \leq 5$; $DT_n \odot K_1$; $DT_n \odot 2K_1$; $DT_n \odot K_2$; $DQ_n \odot K_1$; $DQ_n \odot 2K_1$; $DQ_n \odot K_2$; $DA(T_n) \odot K_1$; $DA(T_n) \odot 2K_1$; $DA(T_n) \odot K_2$; $DA(Q_n) \odot K_1$; $DA(Q_n) \odot 2K_1$; and $DA(Q_n) \odot K_2$. They also proved: if G is a (p, q) difference cordial graph, then $q \leq 2p - 1$; if G is a r-regular graph with $r \geq 4$, then G is not difference cordial; if $m \geq 4$ and $n \geq 4$, then $K_{m,n}$ is not difference cordial; if $m + n > 8$ then the bistar $B_{m,n}$ is not difference cordial; and every graph is a subgraph of a connected difference cordial graph. If G is a book, sunflower, lotus inside a circle, or square of a path, they prove that $G \odot mK_1$ $(m = 1, 2)$ and $G \odot K_2$ is difference cordial.

In [2952], [2954], and [2953] Ponraj, Sathish Narayanan, and Kala proved that the following graphs are difference cordial: crowns $C_n \odot K_1$; combs $P_n \odot K_1$; $P_n \odot C_m$; $C_n \odot C_m$; $W_n \odot K_2$; $W_n \odot 2K_1$; $G_n \odot K_1$ where G_n is the gear graph; $G_n \odot 2K_1$; $G_n \odot K_2$; $(C_n \times P_2) \odot K_1$; $(C_n \times P_2) \odot 2K_1$; $(C_n \times P_2) \odot K_2$; $L_n \odot K_1$; $L_n \odot 2K_1$; and $L_n \odot K_2$. Ponraj, Sathish Narayanan and Kala proved that the following subdivision graphs are difference cordial: $S(T_n)$; $S(Q_n)$; $S(DT_n)$; $S(DQ_n)$; $S(A(T_n))$; $S(DA(T_n))$; $S(AQ_n)$; $S(DAQ_n)$; $S(K_{1,n})$; $S(K_{2,n})$; $S(W_n)$; $S(P_n \odot K_1)$; $S(P_n \odot 2K_1)$; $S(LC_n)$; $S(P_n^2)$; $S(K_2 + mK_1)$; subdivision graphs of sunflowers $S(SF_n)$; subdivisions graphs flowers $S(Fl_n)$; $S(B_m)$ (B_m is a book with m pages); $S(C_n \times P_2)$; $S(B_{m,n})$; subdivisions n-cubes; $S(J(m, n))$; $S(W(t, n))$; subdivisions of Young tableaus $S(Y_{n,n})$; and if $S(G)$ is difference cordial, then $S(G \odot mK_1)$ is difference cordial. For graphs G that are a tree, a unicycle, or when $|E(G)| = |V(G)| + 1$, they proved that $G \odot P_n$ and $G \odot mK_1$ $(m = 1, 2, 3)$ are difference cordial.

In [3008] Prajapati and Gajjar define a *holiday star* as follows. Let $v_1, v_2, \ldots, v_{4n-1}, v_{4n}$ be the consecutive $4n$ vertices of C_{4n} $(n \geq 3)$. Let u_0 be the central vertex and $u_1, u_2, \ldots, u_{2n-1}, u_{2n}$ be end vertices of $K_{1,2n}$. Join u_0 to v_{4i-3} by an edge; for each i from 1 to n. In [3009] they define a *Kusadama flower graph* as follows. Let v_0 be the apex vertex and $v_1, v_2, v_3, \ldots, v_{2n-1}, v_{2n}$ be $2n$ consecutive rim vertices of the wheel W_{2n} $(n \geq 3)$. Subdivide the spoke edge v_0v_{2i-1} by a vertex w_i and at each w_i join two copies of path of length 2; $P_2^\ell = v_0, u_{2i-1}, w_i$ and $P_2^r = v_0, u_{2i}, w_i$, for each $i \in [n]$. In [3008] and [3009] Prajapati and Gajjar proved that the holiday star graph and the Kusudama flower graph admit cordial, E-cordial, difference cordial, prime, vertex prime, and total prime labelings. In [3010] Prajapati and Gajjar define a *braided star graph* as follows: Let a_0 be the apex vertex and $a_1, a_2, \ldots, a_{n-1}, a_n$ be consecutive n rim vertices of W_n $(n \geq 3)$; let $b_1, b_2, b_3, \ldots, b_{2n-1}, b_{2n}$ be $2n$ consecutive vertices of the cycle C_{2n}; let $c_1, c_2, \ldots, c_{2n-1}, c_{2n}$ be consecutive $2n$ vertices of C_{2n}. Join each a_i to b_{2i-1} by an edge and b_{2i} to c_{2i} by an edge. Take a new vertex d_i and join each d_i to c_{2i-1} and c_{2i+1} by an edge for each $i \in [n]$ where subscripts are taken modulo n. Prajapati and Gajjar [3010] proved that braided star graphs are cordial, E-cordial and difference cordial. In [3012] Prajapati and Gajjar investigated the existence of cordial, E-cordial, vertex prime, and difference cordial labelings of graphs that are inspired by the origami model called an *aboreale star* (see their

paper for the definition).

Recall the splitting graph of G, $S'(G)$, is obtained from G by adding for each vertex v of G a new vertex v' so that v' is adjacent to every vertex that is adjacent to v and the shadow graph $D_2(G)$ of a connected graph G is constructed by taking two copies of G, G' and G'', and joining each vertex u' in G' to the neighbors of the corresponding vertex u' in G''.

Ponraj and Sathish Narayanan [2939], [2940] proved the following graphs are difference cordial: $S'(P_n)$; $S'(C_n)$; $S'(P_n \odot K_1)$; and $S'(K_{1,n})$ if and only if $n \leq 3$. They proved following are not difference cordial: $S'(W_n)$; $S'(K_n)$; $S'(C_n \times P_2)$; the splitting graph of a flower graph; $DS(SF_n)$; $DS(LC_n)$; $DS(Fl_n)$; $D_2(G)$ where G is a (p,q) graph with $q \geq p$; and $DS(B_{m,n})$ $(m \neq n)$ with $m+n > 8$.

Let $G(V,E)$ be a graph with $V = S_1 \cup S_2 \cup \cdots \cup S_t \cup T$ where each S_i is a set of vertices having at least two vertices and having the same degree. Panraj and Sathish Narayanan [2939], [2940] define the *degree splitting graph* of G denoted by $DS(G)$ as the graph obtained from G by adding vertices $w_1, w_2, \ldots, w_t$ and joining w_i to each vertex of S_i $(1 \leq i \leq t)$. They proved the following graphs are difference cordial: $DS(P_n)$; W_n; $DS(C_n)$; $DS(K_n)$ if and only if $n \leq 3$; $DS(K_{1,n})$ if and only if $n \leq 4$; $DS(W_n)$ if and only if $n = 3$; $DS(K_n^c + 2K_2)$ if and only if $n = 1$; $DS(K_2 + mK_1)$ if and only if $n \leq 3$; $DS(K_{n,n})$ if and only if $n \leq 2$; $DS(T_n)$ if and only if $n \leq 5$; $DS(Q_n)$ if and only if $n \leq 5$; $DS(L_n)$ if and only if $n \leq 5$; $DS(B_{n,n})$ if and only if $n \leq 2$; $DS(B_{1,n})$ if and only $n \leq 4$; $DS(B_{2,n})$ if and only $n \leq 4$; $D_2(P_n)$; $D_2(K_n)$ if and only if $n \leq 2$; and $D_2(K_{1,m})$ if and only if $m \leq 2$.

In [2941], Ponraj and Sathish Narayanan proved the following graphs are difference cordial: $T_n \odot K_1$, $T_n \odot 2K_1$, $T_n \odot K_2$, $A(T_n) \odot K_1$, $A(T_n) \odot 2K_1$ and $A(T_n) \odot K_2$ where T_n and $A(T_n)$ are triangular snake and alternate triangular snake respectively. In [2955,2956] Ponraj, Sathish Narayanan, and Kala proved the following graphs are difference cordial: $C_n \times P_2$; Mobius ladders; the n-cube; sunflower graphs; lotuses inside a circle; pyramids; books with n pentagonal pages; mongolian tents; graphs obtained from a ladder by subdividing each step exactly once; permutation graphs $P(P_{2k}, f)$ where $f = (1\,2)(3\,4)\cdots(k\;k{+}1)\cdots(2k{-}1\;2k)$; and $P(P_n, I)$, $P(C_n, I)$, $P(P_n \odot K_1, I)$, $P(P_n \odot 2K_1, I)$ where I is the identity permutation. Ponraj, Sathish Narayanan, and Kala [2955] [2956] proved the following graphs are not difference cordial: $G_1(p_1, q_1) \times G_2(p_2, q_2)$ with $q_1 \geq p_1$ and $q_2 \geq p_2$; $C_m \times C_n$; $G \times K_n$ where G connected graph and $n \geq 5$, $G + K_1$ where $|E(G) > |V(G) + 1$; $G_1 + G_2$ where G_1 and G_2 are connected and $|E(G_1)| > 1$ and $E(G_2)| > 3$; permutation graphs $P(G \times K_2, f)$ where $|E(G)| \geq |V(G)|$ and f is any permutation; $P(W_n, f)$ for any permutation f; $P(S'(G), f)$ where $S'(G)$ is the splitting graph of G, $|E(G)| \geq |V(G)|$, and f is any permutation; and $P(Fl_n, f)$ where Fl_n is a flower graph and f is any permutation. They also obtained the following necessary and sufficient conditions for difference cordiality: $K_m \times P_2$ if and only if $m \leq 3$; for a connected graph G, $G \times W_n$ if and only if $G = K_1$; books B_m if and only if $m \leq 6$; $G + G$ if and only if $|V(G)| \leq 3$ and $|E(G)| \leq 1$; $K_2 + mK_1$ if and only if $m \leq 4$; $\overline{K_n} + 2K_2$ if and only if $n \leq 2$; the double fan DF_n if and only if $n \leq 4$; the t-fold wheel $W_n + \overline{K_t}$ if and only if $t \leq 2$ and $n = 3$; cocktail party graphs $H_{n,n}$ if and only $n \leq 6$; $P(K_n, I)$ if and only if

$n \leq 3$; $P(K_2 + mK_1, I)$ if and only if $m \leq 3$; and $P(K_{m,n}, I)$ $(m, n > 1)$ if and only if $m = n = 2$ and $n = 3, 4, 5$.

Vaidya and Barasara [3953] proved that every graph can be embedded as an induced subgraph of a difference cordial graph, thereby ruling out any possibility of obtaining any forbidden subgraph characterization for difference cordial graphs. They also proved that every connected graph can be embedded as an induced subgraph of a difference cordial connected graph and every planar graph can be embedded as an induced subgraph of a difference cordial planar graph.

In [2856], Ponraj, Maria Adaickalam, and Kala introduced a new graph labeling called a k-difference cordial labeling. Let G be a (p, q)-graph and $2 \leq k \leq |V(G)|$. Let $f : V(G) \to \{1, 2, \ldots, k\}$ be a map. For each edge uv, assign the label $|f(u) - f(v)|$. They say f is a *k-difference cordial* labeling of G if $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(0) - e_f(1)| \leq 1$, where $v_f(x)$ denotes the number of vertices labeled with x, $e_f(1)$ denotes the number of edges labeled with 1, and $e_f(0)$ denotes the number of edges that are not labeled with 1. A graph with a k-difference cordial labeling is called a *k-difference cordial* graph. They proved the following: every graph is a subgraph of a connected k-difference cordial graph; if k is even, then k-copies of $K_{1,p}$ is k-difference cordial; and if $n \equiv 0 \pmod{k}$ and $k \geq 6$, then $K_{1,n}$ is not k-difference cordial. They further prove the following are 3-difference cordial graphs: paths; C_n where $n \equiv 0, 3 \pmod{4}$; $K_{m,n}$ $(m \leq n)$ and m is even; combs; double combs; quadrilateral snakes; bistars; subdivisions of a star; subdivisions of a bistar; $C_4^{(t)}$; K_n if and only if $n \in \{1, 2, 3, 4, 6, 7, 9, 10\}$; and $K_{1,n}$ if and only if $n \in \{1, 2, 3, 4, 5, 6, 7, 9\}$.

In [2851], [2852], [2853], Ponraj and Maria Adaickalam proved the following are 3-difference cordial graphs: $K_{1,n} \odot K_2$, $P_n \odot 3K_1$, $C_n \odot K_2$, mC_4, splitting graph of a star, fan, double fan, W_n where $n \equiv 0, 1 \pmod{3}$, helms, flower, sunflower graph, lotus inside a circle, closed helm, double wheel DW_n where $V(DW_n) = V(W_n) \cup \{v_i : 1 \leq i \leq n\}$ and edge set $E(DW_n) = E(W_n) \cup \{uv_i : 1 \leq i \leq n\} \cup \{v_iv_{i+1} : 1 \leq i \leq n-1\} \cup \{v_1v_n\}$, degree splitting graph of a bistar, $spl(K_{1,n}) \cup K_{1,n}$, $spl(K_{1,n}) \cup P_n$, $K_{3,n} \cup spl(K_{1,n})$, $DF_n \cup spl(K_{1,n})$, $S(K_{1,n}) \cup S(B_{n,n})$, $K_{2,n} \cup S(K_{1,n})$, $F_n \cup S(K_{1,n})$, $W_n \cup S(K_{1,n})$, $B_{n,n} \cup S(B_{n,n})$, $K_{2,n} \cup B_{n,n}$, $(C_n \odot K_1) \cup (P_n \odot K_1)$, $F_n \cup F_n$, jelly fish, $P_n \cup K_{1,n}$, $K_{1,n} \cup K_{2,n}$, $K_{1,n} \cup S(K_{1,n})$, are Let C_n be the cycle $u_1u_2 \ldots u_nu_1$. If G is (p, q) 3-difference cordial graph with $p \equiv 0 \pmod{2}$ and $q \equiv 0 \pmod{3}$, then $G \cup G$ also 3-difference cordial. Let G be the graph obtained from C_n with $V(G) = V(C_n) \cup \{v_i : 1 \leq i \leq \lceil \frac{n}{2} \rceil\}$ and $E(G) = \{u_iv_i, u_{i+1}v_i : 1 \leq i \leq n\}$. Then G is 3-difference cordial. The graph G_n with the vertex set $V(G_n) = \{u_i, v_i, w_i : 1 \leq i \leq n\}$ and $E(G_n) = \{u_iu_{i+1}, v_iv_{i+1} : 1 \leq i \leq n-1\} \cup \{u_nu_1, v_1u_1\} \cup \{u_iv_i, v_iw_i : 1 \leq i \leq n\}$ is 3-difference cordial. Let C_3 be the cycle $u_1u_2u_3u_1$. Let G be a graph obtained from C_3 with $V(G) = V(C_3) \cup \{v_i, w_i, z_i : 1 \leq i \leq n\}$ and $E(G) = E(C_3) = \{u_1v_i, u_2w_i, u_3z_i : 1 \leq i \leq n\}$. Then G is 3-difference cordial if $n \equiv 0, 2, 3 \pmod{4}$. If $n \equiv 0, 1 \pmod{3}$, then $K_{1,n} \cup K_{1,n}$ is 3-difference cordial. Ponraj, Adaickalam, and Kala [2857] proved the following graphs have 3-difference cordial labelings: $DA(T_n) \odot K_1$, $DA(T_n) \odot 2K_1$, $DA(T_n) \odot K_2$, $DA(Q_n) \odot K_1$, and $DA(Q_n) \odot 2K_1$ (T_n is a triangular snake.) In [2854] Ponraj, Adaickalam, Maria Adaickalam, and Kala investigated the 3-difference cordial labeling behavior of ladders, books, dumbbell graphs,

and umbrella graphs. Ponraj, Subbulakshmi, and Somasundarami [2988] investigated the 4-difference cordial labeling behavior of jelly fish, jewel graphs, combs, subdivisions of stars, subdivisions of bistars, and books with triangle pages.

For graphs G and H and a vertex v of G the graph $G \odot_v H$ is obtained by joining any particular vertex of H to vertex v. In [2538] Sugumaran and Mohan proved that the following graphs are difference cordial graphs: the path union of r copies of P_n^2 (that is, $P(r.P_n^2)$)–see Section 2.7 for the definition), the cycle union of r copies of C_n^2, (that is, $C(r.C_n^2)$), the open star of r copies of the square graph P_n^2 (that is, $S(r.P_n^2)$), the graph $C_n^2 \odot_{v_n} P_k$, and the graph $C_n^2 \odot_{v_n} P_k^2$. In [2539] they proved that the plus graph Pl_n, the path union of plus graph $P(r.Pl_n)$, the cycle union of plus graph $C(r.Pl_n)$, the barycentric subdivision of Pl_n, the hanging pyramid HP_{y_n} graph, and the path union of hanging pyramid $P(r.HP_{y_n})$. In [3745] they proved that switching of a pendent vertex in path P_n, switching of an apex vertex in CH_n, the graph obtained by duplication of each vertex of path P_n by an edge, the barycentric subdivision of $C_n \odot K_1$, the path union of r copies of fan $P(r.F_n)$, the cycle union of r copies of fan $C(r.F_n)$, and the open star of r copies of fan $S(r.F_n)$ are difference cordial graphs.

7.9 Prime Cordial Labelings

Sundaram, Ponraj, and Somasundaram [3787] have introduced the notion of prime cordial labelings. A *prime cordial labeling* of a graph G with vertex set V is a bijection f from V to $\{1, 2, \ldots, |V|\}$ such that if each edge uv is assigned the label 1 if $\gcd(f(u), f(v)) = 1$ and 0 if $\gcd(f(u), f(v)) > 1$, then the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. In [3787] Sundaram, Ponraj, and Somasundram prove the following graphs are prime cordial: C_n if and only if $n \geq 6$; P_n if and only if $n \neq 3$ or 5; $K_{1,n}$ (n odd); the graph obtained by subdividing each edge of $K_{1,n}$ if and only if $n \geq 3$; bistars; dragons; crowns; triangular snakes if and only if the snake has at least three triangles; ladders; $K_{1,n}$ if n is even and there exists a prime p such that $2p < n+1 < 3p$; $K_{2,n}$ if n is even and if there exists a prime p such that $3p < n+2 < 4p$; and $K_{3,n}$ if n is odd and if there exists a prime p such that $5p < n+3 < 6p$. They also prove that if G is a prime cordial graph of even size, then the graph obtained by identifying the central vertex of $K_{1,n}$ with the vertex of G labeled with 2 is prime cordial, and if G is a prime cordial graph of odd size, then the graph obtained by identifying the central vertex of $K_{1,2n}$ with the vertex of G labeled with 2 is prime cordial. They further prove that $K_{m,n}$ is not prime cordial for a number of special cases of m and n. Sundaram and Somasundaram [3790] and Youssef [4284] observed that for $n \geq 3$, K_n is not prime cordial provided that the inequality $\phi(2) + \phi(3) + \cdots + \phi(n) \geq n(n-1)/4 + 1$ is valid for $n \geq 3$ (ϕ is the Euler phi-function). This inequality was proved by Yufei Zhao [4331]. Haque, Lin, Yang, and Zhao [1365] show that with the exception of $P(4, 1)$, all generalized Petersen graphs are prime cordial. Haque, Lin, Yang, and Zhang [1363] show that the flower snark and related graphs are prime cordial. In [1267] Ghosh, Mohanty, and Pal gave an algorithmic approach to find cordial labelings of the Cartesian product of two balanced bipartite graphs. In [4060] Vasanthi1 and Ramya proved that the Heawood graph, the Pappus graph, and the

Tutte-Coxeter graph admit for prime cordial labelings. In [3756] Sugumaran and Vishnu Prakash investigated the prime cordial labeling of theta graphs in the context of the graph operations of duplication, switching, fusion, path union, and the graphs obtained by joining two copies of theta graph by a path of arbitrary length. Rayjada, Sarvaiya and Gadhiya [3149] applied prime cordial labelings for internode connection modeling that enhances the cryptographic complexity. [3149] new

A *signed product cordial* labeling of graph $G(V, E)$ is a function f from V to $\{-1, 1\}$ such that when each edge uv is assigned the label $f(u)f(v)$, the number of vertices labeled -1 and the number of vertices labeled 1 differ by at most 1, and likewise for edges. A graph that admits signed product cordial labeling is called a *signed product cordial* graph. Signed product cordial labelings for fractal graphs given by Santhi and Albert in [3304] and by Raval and Prajapati in [3138]. Ghosh and Pal [**?**] proved that $K_{m,m} \times P_n$ admits signed product cordial labelings and total signed product cordial labelings. El-hay, Elmshtaye, and Elrokh [1003] provided necessary and sufficient conditions for which there exist a signed product cordial labeling of corona product of paths and third power of lemniscate graph.

Seoud and Salim [3394] give an upper bound for the number of edges of a graph with a prime cordial labeling as a function of the number of vertices. For bipartite graphs they give a stronger bound. They prove that K_n does not have a prime cordial labeling for $2 < n < 500$ and conjecture that K_n is not prime cordial for all $n > 2$. They determine all prime cordial graphs of order at most 6. For a graph with n vertices to admit a prime cordial labeling, Seoud and Salim [3399] proved that the number of edges must be less than $n(n-1) - 6n^2/\pi^2 + 3$. As a corollary they get that K_n $(n > 2)$ is not prime cordial thereby proving their earlier conjecture.

In [1252] Ghodasara and Jena prove that the following graphs are prime cordial: C_n with one chord, C_n with twin chords (that is, two cords that form a triangle with an edge of the cycle), C_n with three cords that form two triangles and a cycle of length $n-3$ $(n \geq 7)$, the graph obtained by joining two copies of C_n with one chord by a path, and the graph obtained by joining two copies of the same cycle with twin chords by a path is prime cordial.

In [583] Baskar Babujee and Shobana proved sun graphs $C_n \odot K_1$; C_n with a path of length $n-3$ attached to a vertex; and P_n $(n \geq 6)$ with $n-3$ pendent edges attached to a pendent vertex of P_n have prime cordial labelings. Additional results on prime cordial labelings are given in [584].

In [4020] and [4022] Vaidya and Vihol prove following graphs are prime cordial: the total graph of P_n and the total graph of C_n for $n \geq 5$ (see §2.7 for the definition); $P_2[P_m]$ for all $m \geq 5$; the graph obtained by joining two copies of a fixed cycle by a path; and the graph obtained by switching of a vertex of C_n except for $n = 5$ (see §3.6 for the definition); the graph obtained by duplicating each edge by a vertex in C_n except for $n = 4$ (see §2.7 for the definition); the graph obtained by duplicating a vertex by an edge in cycle C_n (see §2.7 for the definition); the path union of any number of copies of a fixed cycle (see §3.7 for the definition); and the friendship graph F_n for $n \geq 3$. Vaidya and Shah [4001] prove following results: P_n^2 is prime cordial for $n = 6$ and $n \geq 8$; C_n^2 is prime cordial for $n \geq 10$;

the shadow graphs of $K_{1,n}$ (see §3.8 for the definition) for $n \geq 4$ and the bistar $B_{n,n}$ are prime cordial graphs.

In [176] Aljouiee provided a complete characterization of complete graphs that are prime cordial, showed that closed helms are prime cordial, and presented a new prime cordial labeling for P_n^2. He also corrected the proof of that $C_n^{@}$ is prime cordial for ≥ 10 given in [4001].

Let G_n be a simple nontrivial connected cubic graph with vertex set $V(G_n) = \{a_i, b_i, c_i, d_i : 0 \leq i \leq n-1\}$, and edge set $E(G_n) = \{a_ia_{i+1}, b_ib_{i+1}, c_ic_{i+1}, d_ia_i, d_ib_i, d_ic_i : 0 \leq i \leq n-1\}$, where the edge labels are taken modulo n. Let H_n be a graph obtained from G_n by replacing the edges $b_{n-1}b_0$ and $c_{n-1}c_0$ with $b_{n-1}c_0$ and $c_{n-1}b_0$ respectively. For odd $n \geq 5$, H_n is called a *flower snark* whereas G_n, H_3 and all H_n with even $n \geq 4$, are called the *related graphs* of a flower snark. Mominul Haque, Lin, Yang, and Zhang [2543] proved that flower snarks and related graphs are prime cordial for all $n \geq 3$.

In [4004] Vaidya and Shah prove that the following graphs are prime cordial: split graphs of $K_{1,n}$ and $B_{n,n}$; the square graph of $B_{n,n}$; the middle graph of P_n for $n \geq 4$; and W_n if and only if $n \geq 8$. Vaidya and Shah [4004] prove following graphs are prime cordial: the splitting graphs of $K_{1,n}$ and $B_{n,n}$; the square of $B_{n,n}$; the middle graph of P_n for $n \geq 4$; and wheels W_n for $n \geq 8$. Gajjar [1166] proved that the generalized web graph $W(t, n)$ $(n \geq 3)$ is prime cordial for $t = 1, 2, 3$, and 4 and that $W(t, 2p)$ is prime cordial for all t and odd primes p.

In [4008] [4010] Vaidya and Shah proved following graphs are prime cordial: gear graphs G_n for $n \geq 4$; helms; closed helms CH_n for $n \geq 5$; flower graphs Fl_n for $n \geq 4$; degree splitting graphs of P_n and the bistar $B_{n,n}$; double fans Df_n for $n = 8$ and $n \geq 10$; the graphs obtained by duplication of an arbitrary rim edge by an edge in W_n where $n \geq 6$; and the graphs obtained by duplication of an arbitrary spoke edge by an edge in wheel W_n where $n = 7$ and $n \geq 9$.

Let $G(p, q)$ with $p \geq 4$ be a prime cordial graph and $K_{2,n}$ be a bipartite graph with bipartition $V = V_1 \cup V_2$ with $V_1 = \{v_1, v_2\}$ and $V_2 = \{u_1, u_2, \ldots, u_n\}$. If G_1 is the graph obtained by identifying the vertices v_1 and v_2 of $K_{2,n}$ with the vertices of G having labels 2 and 4 respectively, Vaidya and Prajapati [3999] proved that G_1 admits a prime cordial labeling if n is even; if n, p, q are odd and with $e_f(0) = \lfloor q/2 \rfloor$; and if n is odd, p is even and q is odd with $e_f(0) = \lceil q/2 \rceil$. Prajapati and Gajjar [3002] proved the following graphs are prime cordial: $C_n \times P_2$ except for $n = 1, 2$ and 4, $C_n \times P_4$ $(n \geq 3)$, $C_3 \times P_n$ $(n > 1)$, $C_5 \times P_n$ $(n > 1)$, $C_6 \times P_n$ $(n > 1)$, $C_{2p} \times P_n$ where p is an odd prime and $n > 1$, and $C_4 \times P_n$ $(n > 2)$.

In [3743] Sugumaran and Mohan proved the following graphs are prime cordial: the *cycle butterfly* graph $B_{n,m}$ (two copies of C_n that share a common vertex with m pendent vertices attached to the common vertex), $W-$ graph (obtained by starting with the two copies of $K_{1,n}$ and merging the last pendent vertex in the first copy of $K_{1,n}$ with the initial pendent vertex in the second copy of $K_{1,n}$), H_n graph (the graph obtained from two paths $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$ by joining the vertices $u_{(n+1)/2}$ and $v_{(n+1)/2}$ if n is odd and joining $u_{n/2}$ and $v_{n/2+1}$ if n is even), and duplication of all edges of an H_n graph. In [3744] Sugumaran and Mohan proved that the following graphs are prime cordial: $H_n \odot K_1$, the

path union of r copies of H_n, the cycle union of r copies of an H_n, the open star of r copies of an H_n-graph (obtained by replacing each pendent vertex of $K_{1,n}$ by a copy of H_n.

In [3749] Sugumaran and Suresh proved that the following graphs are prime cordial graphs: the duplication of each vertex by an edge of paths, stars, jelly fish, bistars, and $C_n \odot K_1$. Sugumaran and Vishnu Prakash [3754] proved that the following graphs are prime cordial graphs: duplication of any vertex of degree 3 in theta graph , switching of any vertex of degree 3 in theta graph, fusion of any two vertices in theta graph, the path union of two copies of theta graph, and two copies of theta graph joined by a path of any length. They further proved that the theta graph is not a prime cordial labeling. In [3755] they showed that one point union of path graph $P_n^t(tn.T_\alpha)$, the open star of theta graph, and any path union of even number of theta graphs are prime cordial graphs. Also they proved [3753] the subdivision of bistar $B_{n,n}$, $P_n \bigodot K_{1,n-1}$, (that is, each i^{th} vertex of path P_n is append with the apex vertex of i^{th} copy of $K_(1, n-1)$), the disconnected graph $P_n \cup P_m$ are prime cordial graphs.

Vaidya and Barasara [3951] proved that every graph can be embedded as an induced subgraph of a prime cordial graph thereby ruling out any possibility of obtaining any forbidden subgraph characterization for prime cordial graphs. They also proved that a connected graph can be embedded as an induced subgraph of a prime cordial connected graph and every planar graph can be embedded as an induced subgraph of a prime cordial planar graph. In [1234] Gayathri, Thanjavur, Maniammai, and Sakar proved that 8-polygonal snakes containing n 8-polygons, splitting graphs of C_n for $n \geq 5$, and armed crowns $C_{2k} \odot P_m$ for all $k \geq 3$ and $m \geq 2$ admit prime cordial labelings. Barasara and Prajapati [496] showed that switching of a vertex of degree 1 in the path P_n is a prime cordial graph for $n \neq 2, 3, 4, 5, 7$ and not a prime cordial graph for $n = 2, 3, 4, 5, 7$, the armed crown AC_n is a prime cordial graph for all n, all web graphs are prime cordial, and $C_n^{(t)}$, one-point union of cycles, are prime cordial for $t \neq 2$ and $n \neq 3, 4$ and not a prime cordial graph for $t = 2$ and $n = 3, 4$. Babitha and Baskar Babujee [327] proved that if G is prime cordial, then so is G with an edge deleted, $K_{1,m} \odot \ P_n$ $(m, n > 2)$ is prime cordial, and in certain cases, the one-point union of $K_{1,m}$ and P_n and the one-point union of a prime cordial graph G and $K_{1,n}$ are prime cordial. They further provided some characterization results.

For a graph $G(V, E)$ and the group S_3 of all permutations of $\{1, 2, 3\}$ Chandra and Kala [766] define a function $g : V(G) \to S_3$ such that $xy \in E$ if $g(x)$ and $g(y)$ have relatively prime orders. Let $n_j(g)$ denote the number of vertices of G having label j under g. Then g is called a $g$roup S_3 cordial prime labeling if $|n_i(g) - n_j(g)| \leq 1$ for every $i, j \in S_3$. A graph that admits a group S_3 cordial prime labeling is called a $g$roup S_3 cordial prime cordial prime graph. Chandra and Kala [766] prove that all paths, cycles, gears, ladders, and fans are group S_3 cordial prime and characterize wheels that are group S_3 cordial prime.

In [2784] Parthiban and Sharma gave a comprehensive survey on prime cordial and divisor cordial labeling of graphs.

Vaidya and Prajapati [3998] call a graph *strongly prime cordial* if for any vertex v

there is a prime labeling f of G such that $f(v) = 1$. They prove the following: the graphs obtained by identifying any two vertices of $K_{1,n}$ are prime cordial; the graphs obtained by identifying any two vertices of P_n are prime cordial; C_n, P_n, and $K_{1,n}$ are strongly prime cordial; and W_n is a strongly prime cordial for every even integer $n \geq 4$. Prajapati and Gajjar [3014] proved that generalized prism graphs $Y_{n,2}$ is prime cordial except for $n = 1, 2$ and 4; $Y_{n,4}$ is prime cordial for $n \geq 3$; $Y_{3,n}, Y_{5,n}, Y_{6,n}$ and $Y_{2p,n}$ (for odd prime p) are prime cordial for $n > 1$; and $Y_{4,n}$ is prime cordial for $n > 2$. They also proved the following graphs are prime cordial: $C_n \times P_2$ except for $n = 1, 2$ and 4, $C_n \times P_4$ $(n \geq 3)$, $C_3 \times P_n$ $(n > 1)$, $C_5 \times P_n$ $(n > 1)$, $C_6 \times P_n$ $(n > 1)$, $C_{2p} \times P_n$ where p is an odd prime and $n > 1$, and $C_4 \times P_n$ $(n > 2)$.

In [2965] Ponraj, Singh, Kala, and Sathish Narayanan introduced a new graph labeling called k-prime cordial labeling. Let G be a (p, q)-graph and $2 \leq p \leq k$ and let $f : V(G) \to \{1, 2, \ldots, k\}$ be a map. For each edge uv, assign the label $\gcd(f(u), f(v))$. They say that f is a *k-prime cordial* labeling of G if $|v_f(i) - v_f(j)| \leq 1$ for $i, j \in \{1, 2, \ldots, k\}$ and $|e_f(0) - e_f(1)| \leq 1$, where $v_f(x)$ denotes the number of vertices labeled with x, and $e_f(1)$ and $e_f(0)$, respectively, denote the number of edges labeled with 1 and not labeled with 1. A graph with a k-prime cordial labeling is a *k-prime cordial graph*. They proved that every graph is a subgraph of a connected k-prime cordial graph; if k is even, then P_n, $n \neq 3$, is k-prime cordial; C_n, $n \neq 3$, is k-prime cordial when k is even; and the bistar $B_{n,n}$ is k-prime cordial for all even k. They studied 3-prime cordiality of paths, cycles, and olive trees. They also proved that if T is a 3-prime cordial tree, then $T \odot K_1$ is 3-prime cordial; $K_{1,n}$ is 3-prime cordial if and only if $n \leq 3$; K_n is 3-prime cordial if and only if $n < 3$; combs $P_n \odot K_1$ are 3-prime cordial; and $C_n \odot K_1$ is 3-prime cordial if and only if $n \neq 3$. They proved that $K_2 + mK_1$, $K_{2,n}$, and wheels are not 3-prime cordial graphs. In [2966] Ponraj, Singh, and Sathish Narayana proved if G is 3-prime cordial, then $G \cup P_n$ is a 3-prime cordial for $n > 12$, the splitting graph of a star is not a 3-prime cordial graph, and the jelly fish $J(m, n)$ is 3-prime cordial if $10m \geq n + 2$.

In [1354] Hailat called a k-prime cordial labeling f of a graph G a *k-prime cordial weak* labeling of G if, for distinct i and j from $\{1, 2, \ldots, k\}$, the number of vertices labeled i and the number of vertices labeled j differ by at most 2 and the number of edges labeled 0 and the number of edges labeled with 1 differ by at most 2. Hailat also introduced the concept of the n-fold of a graph, denoted $T_n(G)$, constructed by joining corresponding vertices across n copies of a base graph G. He showed show that $T_n(C_n)$ admits a 4-prime cordial weak labeling for all $n \geq 2$. [1354] new

For a 4-prime cordial graph G Ponraj and Rajpal Singh [2964] proved $G \cup P_n$ $(n \geq 5)$, $G \cup 2mK_{n,n}$, and $G \cup 2mK_{1,n}$ are 4-prime cordial. For a $(4t, q)$ 4-prime cordial graph G they prove that $G + K_1$ and $G + 2K_1$ are 4-prime cordial. Ponraj, Rajpal Singh, and Kala [2967] prove that $P_m \times P_n$ and subdivisions of wheels and helms are 4-prime cordial. They also show that if G is bipartite then $G \cup G$ is 4-prime cordial; and if G is 4-prime cordial then $G \odot K_1$ is 4-prime cordial. Ponraj, Singh, and Kala [2968] proved the following graphs are 4-prime cordial: $2m(K_{n,n})$, $2m(P_n \times P_2)$, $m(C_n \oplus K_1)$, $mB_{n,n}$, and $2W_{2n+1}$.

In [2928] and [2929] Ponraj, Rajpal Singh, and Narayanan investigated 4-prime cordial

graph labelings for complete graphs, books, flowers, mC_n, the corona of wheels, gears, double cones, helms, closed helms, butterfly graphs, and friendship graphs. In [2930,2931] Ponraj, Rajpal Singh, and Kala investigated the existence of 4-prime cordial labeling of the lotus inside a circle, sunflower graphs, $S(K_2 + mK_1)$, $S(P_n \odot K_1)$, dodecahedrons, and shadow graphs of a paths, cycles, stars, the degree splitting graphs of a bistars, jelly fish, splitting graphs of paths and stars. Ponraj and Rajpal Singh [2932] investigated the 4-prime cordial labeling of triangular and diamond snake graphs and applied the principles of prime cordiality to their structures. Ponraj, Rajpal Singh, and Kala [2933] and [2934] studied the 4-prime cordial labeling behavior of some cycle and star related graphs and they explored the concept of a 3-prime cordial labeling for specific graphs related to cycles. Ponraj, Rajpal Singh, and Kalam [2935] determined conditions for a 4-prime cordial labeling of degree splitting graphs.

In [1353] Hailat introduced the following notion. A *trigraph* of G, $T_3(G)$, of a connected graph G is constructed by taking two copies of G, G' and G'' and joining each vertex v in G to the neighbors of the corresponding vertex v' in G' and v'' in G'' and joining v' to the neighbors of the corresponding vertex v'' in G''. They showed that both $T_3(P_n)$ and $T_3(C_n)$ are 4-prime cordial graphs. [1353] new

Murugesan, Jayaraman, and Shiama (see [3757]) defined a 3-*equitable prime cordial* labeling of a graph G as a bijection f from $V(G)$ to $\{1, 2, \ldots, |V(G)|\}$ such that if an edge uv is assigned the label 1 when $\gcd(f(u), f(v)) = 1$ and $\gcd(f(u)+f(v), f(u)-f(v)) = 1$, the label 2 when $\gcd(f(u), f(v)) = 1$ and $\gcd(f(u) + f(v), f(u) - f(v)) = 2$, and the label 0 otherwise, then the number of edges labeled with i and the number of edges labeled with j differ by at most 1 for $0 \leq i \leq 2$ and $0 \leq j \leq 2$. A graph that has a 3-equitable prime cordial labeling is called a 3-*equitable prime cordial* graph. Sugumaran and Vishnu Prakash [3757] proved the following graphs are 3-equitable prime cordial graphs: bistars, combs, ladders, kites, and slanting ladders. In [3759] they showed that theta graphs, the duplication of any vertex in theta graphs, switching of any vertex in theta graphs, the fusion of any two vertices in theta graphs, path unions of two copies of theta graphs, open star graphs of copies of a fixed theta graph are 3-equitable prime cordial graphs. Seoud and Jaber [3388] proved that the butterfly $BF_{n,m}$, helms H_n, the graph $\langle W_n : W_m \rangle$ obtained by joining apex vertices of two wheels with a new vertex are prime cordial, and determine the prime cordial graphs of order 7. They also gave an algorithm to calculate the maximum number of edges in a 3-equitable prime cordial graph. Murugesan, Jayaraman, and Shiama [2601] proved the following graphs are 3-equitable prime cordial: P_n, C_n, $(n \geq 4)$, $K_{1,n}$ if and only if $n = 2 \pmod 3$, and K_n if and only if $n \leq 2$.

7.10 L-Cordial and Binary L Cordial

In [486] Bapat introduced the notion of L-cordial labaling as follows. An *L-cordial labeling* of graph $G(V, E)$ is one for which each edge is labeled with a distinct integer from 1 to $|E|$, each vertex is assigned a label 0 or 1 based on the parity of the largest edge label incident to it, and the number of vertices labeled 0 and 1 differs by at most 1. Bapat proved that [486] new

the following graph families are L-cordial: books, n copies of C_4 sharing a common edge, n copies of C_5 sharing a common edge, n triangles sharing a common vertex, and a specific type of snake graph built from C_3 paths. Arthy [306] provided L-cordiality for ladders, snakes, path-related graphs, generalized antiprisms, H-graphs and related graphs, braid graphs, and cycle-duplications. Bapat [487] proved the following graphs are L-cordial: the *flag of* C_n (obtained by fusing an edge with any vertex of C_n); *bull graphs* (cycles that have an edge appended to two adjacent vertices) crowns; shells; and graphs obtained by fusing t distinct edges at each vertex of a shell graph. [306] new [487] new

In [2678] Nila Steffi and Subhashini introduced binary L-cordial labelings and product binary L-cordial labelings as follows. Let $h : E(G) \to \{1, 2, \ldots, |E(G)|\}$ be a bijection. For each vertex u, let uv_i and uv_j be incident edges having the smallest and greatest h-values, respectively. In a *binary L-cordial* labeling, the vertex u is assigned label 1 when $h(uv_i) * h(uv_j)$ is odd and label 0 otherwise, where $*$ is a binary operation. In addition, the numbers of vertices labeled 0 and 1 must differ by at most one. A *product binary L-cordial* labeling is the special case in which the binary operation is multiplication. They established product binary L-cordiality for paths, odd cycles, even-order stars, combs, dumbbells, bull graphs based on odd cycles, and crowns. Nila Steffi and Subhashini [2679] gave binary L-cordial labelings for paths, cycles, $P_n \odot K_1$, flag graphs, and hypercubes. [2678] new [2679] new

In [2680] Nila Steffi and Subhashini introduced the notion of mean binary L-cordial labelings as follows. A graph $G(V, E)$ is said to be *mean binary L-cordial* if there is a bijection $h : E(G) \to 1, 2, \ldots, |E|$ such that for each vertex u, if $\frac{h(uv_i)+h(uv_j)}{2}$ has an odd integral part, assign the label 1, and assign the label 0 for an even integral part, where uv_i, uv_j have the smallest and greatest h-values, respectively and it also satisfies the condition $|V_h(0) - V_h(1)| \leq 1$, where $V_h(0)$ is the number of vertices with label 0 and $V_h(1)$ is the number of vertices with label 1. They showed paths, cycles, subdivisions of stars, actinia graphs (unicyclic graphs obtained from a cycle where each vertex of cycle is joined to fixed number new vertices of null graph on that number of vertices), dumbbells, and crowns and combs with an odd number of vertices are binary L-cordial graphs. [2680] new

7.11 Mean Edge Cordial

In [4193] Wilson K and Vivik J introduced the concept of mean edge cordial labeling in which a binary edge labeling function $f : E(G) \to \{0, 1\}$ induces a vertex labeling function $f^* : V(G) \to \{0, 1\}$ determined by the mean of the labels of the edges incident with each vertex. A vertex receives label 0 if the average of its incident edge labels is less than 0.5 and label 1 if the average is at least 0.5. A graph is said to be *mean edge cordial* if the numbers of edges labeled 0 and 1 differ by at most one and the numbers of vertices labeled 0 and 1 also differ by at most one. They call such a labeling *mean edge cordial*. They proved that paths, combs, fans, double fans, Z-P_n graphs, and square of paths are mean edge cordial. In [4194,4195] they proved that wheels, gears, helms, closed helms, webs, flower graphs, sunflower graphs, lotus inside a circle graphs, rose flower graphs, kusudama flower graphs, and cherry blossom graphs are edge mean cordial. In [4196,4197] they proved the following graphs admit mean edge cordial labelings: cycles with one chord, cycles with [4193] new [4194, 4195] new [4196, 4197] new

twin chords, sunlet graphs, armed crown graphs, splitting graphs of paths, shadow graphs of paths, total graphs of paths, middle graphs of paths, vertex duplication graphs of paths, and edge duplication graphs of paths. They also showed that C_n is mean edge cordial if and only if n is odd and that every graph is a subgraph of a connected mean edge cordial graph. In [4198–4200] they showed subdivision graphs of rectangular grids, rectangular grids, super subdivision graphs of rectangular grids, open ladders, triangular ladders, open triangular ladders, diagonal ladders, open diagonal ladders, Möbius ladders, and slanting ladders are mean edge cordial. Ladders, subdivision of ladders, and subdivision of combs were shown to admit mean edge cordial labeling in [4201]. [4198–4200] new [4201] new

7.12 Vector Basis S-Cordial Labeling Graphs

In [2889] Ponraj and Jeya introduced the vector basis labeling of graphs as follows. Let G be a (p, q) graph and let V be an inner product space with basis S. Denote inner product of the vectors x and y by $< x, y >$. Let $\phi : V(G) \to S$ be a function. For edge uv, assign the label $< \phi(u), \phi(v) >$. Then ϕ is called a *vector basis S-cordial* labeling of G if $|\phi_x - \phi_y| \leq 1$ and $|\gamma_i - \gamma_j| \leq 1$, where ϕ_x denotes the number of vertices labeled with the vector x and γ_i denotes the number of edges labeled with the scalar i. A graph that admits a vector basis S-cordial labeling is called a *vector basis S-cordial graph.* In [2889] Ponraj, and Jeya investigated vector basis S-cordial labelings of paths, stars, cycles, and complete graphs with different bases S. In [2890,2891], [2892,2895], and [2896] Ponraj and Jeya studied vector basis $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$ S-cordial labelings of thorn graphs, generalized friendship graphs, tadpoles, gears, $D(T_n) \odot mK_1$, $D(Q_n) \odot mK_1$, and certain trees and flower graphs. In [2893, 2894, 2899] they investigated vector basis S-cordial labelings of friendship graphs, fans, lilly graphs, Mongolian tents, and parachutes.

In [2897] Ponraj and Jeya proved the following: a graph is vector basis $\{(1,0),(0,1)\}$ S-cordial if and only if it is cordial; P_n is vector basis $\{(1,0,0),(0,1,0),(0,0,1)\}$ S-cordial and $\{(1,0,0,),(0,1,0,0),(0,0,1,0),(0,0,0,1)\}$ S-cordial if and only if $n \neq 3$; $K_{1,n}$ is vector basis $\{1,0,0),(0,1,0),(0,0,1)\}$ S–cordial if and only if $n \neq 1, 3$; C_n is vector basis $\{(1,0,0,0),(0,1,0,0),(0,0,1,0),(0,0,0,1)\}$ S-cordial if and only if $n \not\equiv 0 \pmod 4$; all stars and combs are vector basis $\{(1,0,0,0),(0,1,0,0),(0,0,1,0),(0,0,0,1)\}$ S-cordial; and K_n is vector basis $\{(1,0,0,0),(0,1,0,0),(0,0,1,0),(0,0,0,1)\}$ S-cordial if and only if $n \leq 3$. In [2898] they prove that the balloon of triangular snake graph is a vector basis $\{(1,0,0,0),(0,1,0,0),(0,0,1,0),(0,0,0,1)\}$ S-cordial. [2897] new [2898] new

7.13 Fuzzy quotient-3 Cordial Labelings

For a simple, finite and planar graph G of order p and size q. Sumathi and J. Suresh Kumar [3777] introduced the concept of fuzzy quotient-3 cordial labeling as follows. Let $\sigma : V(G) \to [0,1]$ be a function defined by $\sigma(v) = \frac{r}{10}$, $r \in Z_4 - \{0\}$. For each edge uv define $\mu : E(G) \to [0,1]$ by $\mu(uv) = \frac{1}{10}\frac{3\sigma(u)}{\sigma(v)}$ where $\sigma(u) \leq \sigma(v)$. The function σ is called

fuzzy quotient-3 cordial labeling of G if the number of vertices labeled with i and the number of vertices labeled with j differ by at most 1, the number of edges labeled with i and the number of edges labeled with j differ by at most 1 where $i \neq j$ have the form $\frac{r}{10}$, where $r \in \{1, 2, 3\}$. They proved that stars and star related graphs are fuzzy quotient-3 cordial.

7.14 Mean Labelings

Somasundaram and Ponraj [3657] introduced the notion of mean labelings of graphs. A graph G with p vertices and q edges is called a *mean graph* if there is an injective function f from the vertices of G to $\{0, 1, 2, \ldots, q\}$ such that when each edge uv is labeled with $(f(u)+f(v))/2$ if $f(u)+f(v)$ is even, and $(f(u)+f(v)+1)/2$ if $f(u)+f(v)$ is odd, then the resulting edge labels are distinct. In [3657], [3658], [3659], [3660], [2985], and [2986] they prove the following graphs are mean graphs: P_n, C_n, $K_{2,n}$, $K_2 + mK_1$, $\overline{K_n} + 2K_2$, $C_m \cup P_n$, $P_m \times P_n$, $P_m \times C_n$, $C_m \odot K_1$, $P_m \odot K_1$, triangular snakes, quadrilateral snakes, K_n if and only if $n < 3$, $K_{1,n}$ if and only if $n < 3$, bistars $B_{m,n}$ $(m > n)$ if and only if $m < n+2$, the subdivision graph of the star $K_{1,n}$ if and only if $n < 4$, the friendship graph $C_3^{(t)}$ if and only if $t < 2$, the one point union of two copies a fixed cycle, dragons (the one point union of C_m and P_n, where the chosen vertex of the path is an end vertex), the one point union of a cycle and $K_{1,n}$ for small values of n, and the arbitrary super subdivision of a path, which is obtained by replacing each edge of a path by $K_{2,m}$. They also prove that W_n is not a mean graph for $n > 3$ and enumerate all mean graphs of order less than 5.

Gayathri and Gopi [1219] prove the following are mean graphs: double triangular snakes; double quadrilateral snakes; generalized antiprisms; graphs obtained by joining the 2 vertices of $K_{2,n}$ of degree n with an edge; and graphs obtained from C_n with consecutive vertices $v_1, v_2, \ldots, v_n$ by adding the chords joining v_i and v_{n-i+2} for $2 \leq i \leq \lfloor n/2 \rfloor$. In [1217] Gayathri and Gopi gave various necessary conditions for mean labelings. Kannan, Manivannan, Loganathan, and Gyeltshen [1886] investigated the existence of mean labelings for the graphs obtained by duplicating an edge or vertex of paths, cycles, combs, and the splitting graph of paths. In [1523] Imran, Cancan, and Ali proved that various families of comb related graphs are mean graphs.

Lourdusamy and Seenivasan [2343] prove that kC_n-snakes are means graphs and every cycle has a super subdivision that is a mean graph. They define a generalized kC_n-*snake* in the same way as a C_n-snake except that the sizes of the cycle blocks can vary (see Section 2.2). They prove that generalized kC_n-snakes are mean graphs. Recall that $P_{a,b}$ denotes the graph obtained by identifying the endpoints of b internally disjoint paths each of length a. Vasuki and Nagarajan [4063] proved that the following graphs admit mean labelings: $P_{r,2m+1}$ for all r and m; $P_{r,2m}$ for all m and $2 \leq r \leq 6$; P_r^{2m+1} for all r and m; and P_r^{2m} for all m and $2 \leq r \leq 6$. Anusa, Sandhya, and Somasundaram [227] proved that triangular ladders, triangular snakes, double triangular snakes, quadrilateral snakes, and double quadrilateral snakes are mean graphs.

Lourdusamy and Seenivasan [2345] define an *edge linked cyclic snake,* $EL(kC_n)$, as the connected graph obtained from k copies of C_n $(n \geq 4)$ by identifying an edge of the

$(i+1)^{th}$ copy to an edge of the i^{th} copy for $i = 1, 2, \ldots, k-1$ in such a way that the consecutive edges so chosen are not adjacent. They proved that all $EL(kC_{2n})$ are mean graphs and some cases of $EL(C_{2n-1})$ are mean graphs. They also define a *generalized edge linked cyclic snake* in the same way but allow the cycle lengths (at least 4) to vary. They prove that certain cases of generalized edge linked cyclic snakes are mean graphs.

A *tortoise graph* ($n \geq 5$ and $n \equiv 1, 3 \pmod 4$) is obtained from a path P_n by attaching an edge between v_i and v_{n-i+1} for $1 \leq i \leq \left\lfloor \frac{n}{2} \right\rfloor$. The graph PC_n ($n \geq 5$) is obtained from $C_n = v_1v_2, v_2v_3, \ldots, v_nv_1$ by adding chords between v_i and v_{n-i+2} for $2 \leq i \leq l$, where $l = \frac{n}{2}$ if n is even and $l = \frac{n-1}{2}$ if n is odd. In [2843] Pipaliya1 and Chaurasia1 proved that tortoise graphs, PC_n graph, k-triangular snake graph, and alternate k-triangular snake graph admit mean labelings. [2843] new

Barrientos and Krop [527] proved that there exist $n!$ graphs of size n that admit mean labelings. They give two necessary conditions for the existence of a mean labeling of a graph G with m vertices and n edges: if G is a mean graph, then $n + 1 \geq m$; if G is a mean graph with n edges and maximum degree $\Delta(G)$, then $\Delta(G) \leq \frac{n+3}{2}$ when n is odd and $\Delta(G) \leq \frac{n+2}{2}$ when n is even. They proved that the disjoint union of n copies of C_3 is a mean graph and if a mean r-regular graph has n vertices, then $r < n - 2$. They established a connection between α-labelings and mean labelings by proving that every tree that admits an α-labeling is a mean graph when the size of its stable sets differ by at most one. When the tree is a caterpillar, this difference can be up to two. Barrientos and Krop call a mean labeling of a bipartite graph an *α-mean labeling* if the labels assigned to vertices of the same color have the same parity. They show that the complementary labeling of a α-mean labeling is also an α-mean labeling. They use graphs with α-mean labelings to construct new mean graphs. One construction consists of connecting a pair of corresponding vertices of two copies of an α-mean graph by an edge. The other construction identifies a pair of suitable vertices from two α-mean graphs. Barrientos and Krop also proved that every quadrilateral snake admits an α-mean labeling. They conjecture that all trees of size n and maximum degree at most $\lceil (n+1)/2 \rceil$ are mean graphs and state some open problems. In [522] Barrientos proves that all trees with up to four end-vertices except $K_{1,4}$ are mean graphs. Bailey and Barrientos [446] prove the following are mean graphs: $C_n \cup C_m,\ C_n \cup P_m,\ K_2 + nK_1,\ 2K_2 + nK_1, C_n \times K_2$.

In [446] Bailey and Barrientos study several operations with mean graphs. They prove that the coronas $G \odot K_1$ and $G \odot K_2$ are mean graphs when G is an α-mean graph. Also, if G and H are mean graphs with n vertices and $n-1$ edges and H is an α-mean graph, then $G \times H$ is a mean graph. They prove that given two mean graphs G and H, there exists a mean graph obtained by identifying an edge from G with an edge from H and uses this result to prove that the graphs R_n ($n \geq 2$) of order $2n$ and size $4n-3$ with vertex set $V(R_n) = \{v_1, v_2, \ldots, v_{2n}\}$ and edge set $E(R_n) = \{v_iv_{i+1} \mid 1 \leq i \leq n-1 \text{ and } n+1 \leq i \leq 2n-1\} \cup \{v_iv_{n+i} \mid 1 \leq i \leq n\} \cup \{v_iv_{n+i-1} \mid 2 \leq i \leq n\}$ (*rigid ladders*) are mean graphs.

Barrientos, Abdel-Aal, Minion, and Williams [523] use A_n to denote the set of all α-mean labeled graphs of size n such that the difference of the cardinalities of the bipartite sets of the vertices of the graphs is at most one. They prove that the class A_n is equivalent

to the class of α-labeled graphs of size n with bipartite sets that differ by at most one. They also prove that when $G \in A_n$, the coronas $G \odot mK_1$, $G \odot P_2$, and $G \odot P_3$ admit mean labelings.

In [3964] Vaidya and Bijukumar define two methods of creating new graphs from cycles as follows. For two copies of a cycle C_n the *mutual duplication* of a pair of vertices v_k and v'_k respectively from each copy of C_n is the new graph G such that $N(v_k) = N(v'_k)$. For two copies of a cycle C_n and an edge $e_k = v_k v_{k+1}$ from one copy of C_n with incident edges $e_{k-1} = v_{k-1}v_k$ and $e_{k+1} = v_{k+1}v_{k+2}$ and an edge $e'_m = u_m u_{m+1}$ in the second copy of C_n with incident edges $e'_{m-1} = u_{m-1}u_m$ and $e'_{m+1} = u_{m+1}u_{m+2}$, the *mutual duplication* of a pair of edges e_k and e'_m respectively from two copies of C_n is the new graph G such that $N(v_k) - v_{k+1} = N(u_m) - u_{m+1} = \{v_{k-1}, u_{m-1}\}$ and $N(v_{k+1}) - v_k = N(u_{m+1}) - u_m = \{v_{k+2}, u_{m+2}\}$. They proved that the graph obtained by mutual duplication of a pair of vertices each from each copy of a cycle and the mutual duplication of a pair of edges from each copy of a cycle are mean graphs. Moreover, they proved that the shadow graphs of the stars $K_{1,n}$ and bistars $B_{n,n}$ are mean graphs.

Vasuki and Nagarajan [4065] proved the following graphs are admit mean labelings: the splitting graphs of paths and even cycles; $C_m \odot P_n$; $C_m \odot 2P_n$; $C_n \cup C_n$; disjoint unions of any number of copies of the hypercube Q_3; and the graphs obtained from by starting with m copies of C_n and identifying one vertex of one copy of C_n with the corresponding vertex in the next copy of C_n.) Jeyanthi and Ramya [1731] define the jewel graph J_n as the graph with vertex set $\{u, x, v, y, u_i : \ 1 \le i \le n\}$ and edge set $\{ux, vx, uy, vy, xy, uu_i, vu_i : \ 1 \le i \le n\}$. They proved that the jewel graphs, jelly fish graphs, and the graph obtained by joining any number of isolated vertices to the two endpoints of P_3 are mean graphs. Ramya and Jeyanthi [3121] proved several families of graphs constructed from T_p-tree are mean graphs. Ahmad, Imran, and Semaničová-Feňovčiková [127] studied the relation between mean labelings and (a, d)-edge-antimagic vertex labelings. They show that two classes of caterpillars admit mean labelings. Revathi [3164] proved that the shadow graphs of bistars, combs, and the splitting graph of combs have mean labelings.

Recall from Section 2.7 that given connected graphs $G_1, G_2, \ldots, G_n$, Kaneria, Makadia, and Jariya [1850] define a *cycle of graphs* $C(G_1, G_2, \ldots, G_n)$ as the graph obtained by adding an edge joining G_i to G_{i+1} for $i = 1, \ldots, n-1$ and an edge joining G_n to G_1. (The resulting graph can vary depending on which vertices of the G_i are chosen.) When the n graphs are isomorphic to G the notation $C(n \cdot G)$ is used. Also recall Kanneria and Makadia [1843] define a *step grid graph* St_n as the graph obtained by starting with paths $P_n, P_n, P_{n-1}, \ldots, P_2$ $(n \ge 3)$ arranged vertically parallel with the vertices in the paths forming horizontal rows and edges joining the vertices of the rows. In [1875], [1860], and [1863], Kaneria, Viradia, and Makadia proved the following graphs are mean graphs: the path union of any number of copies of a mean graph; $C(2t \cdot P_n)$; $C(2t \cdot C_n)$; $C(2t \cdot P_n \times P_m)$; $C(2r \cdot B^2_{n,n})$ ($B^2_{n,n}$ is the square of the bistar $B_{n,n}$); $C(2r \cdot M(C_n))$ ($M(C_n)$ is the middle graph of C_n); $C(2r \cdot (P_{2n} + 2K_1))$; step grid graphs; the path union of finitely copies of the step grid graphs; cycles of step grid graphs $C(2r \cdot St_n)$; and $C(2t \cdot K_{2,m})$.

For a fixed vertex v of C_m Avadayappan and Vasuki [323] use $(P_m; C_n)$ to denote the

graph obtained from m copies of C_n and the path $P_m : u_1u_2 \cdots u_m$ by joining u_i with v of the ith copy of C_n with an edge for $1 \leq i \leq m$. They define $(P_m; Q_3)$, $(P_{2n}; S_m)$, $(P_n; S_1)$ and $(P_n; S_2)$, where v is a fixed vertex of the cube Q_3 and v is the center of the star S_k, in an analogous way. For $C_n : v_1v_2 \ldots v_nv_1$ they use $[P_m; C_n]$ to denote the graph obtained from m copies of C_n with vertices $v_{1_1}, v_{1_2}, \ldots, v_{1_n}, v_{2_1}, \ldots, v_{2_n}, \ldots, v_{m_1}, \ldots, v_{m_n}$ by joining v_{i_j} and $v_{(i+1)_j}$ with an edge, for some j and $1 \leq i \leq m-1$. They define $[P_m; Q_3]$ and $[P_m; C_m^{(2)}]$, where $C_m^{(2)}$ is the friendship graph, similarly. In [323] they prove these families are mean graphs.

Ramya, Ponraj, and Jeyanthi [3124] called a mean graph *super mean* if vertex labels and the edge labels are $\{1, 2, \ldots, p+q\}$. They prove following graphs are super mean: paths, combs, odd cycles, P_n^2, $L_n \odot K_1$, $C_m \cup P_n$ $(n \geq 2)$, the bistars $B_{n,n}$ and $B_{n+1,n}$. They also prove that unions of super mean graphs are super mean and K_n and $K_{1,n}$ are not super mean when $n > 3$. In [1736] Jeyanthi, Ramya, and Thangavelu prove the following are super mean: $nK_{1,4}$; the graphs obtained by identifying an endpoint of P_m $(m \geq 2)$ with each vertex of C_n; the graphs obtained by identifying an endpoint of two copies of P_m $(m \geq 2)$ with each vertex of C_n; the graphs obtained by identifying an endpoint of three copies of P_m $(m \geq 2)$; and the graphs obtained by identifying an endpoint of four copies of P_m $(m \geq 2)$. In [1732] Jeyanthi and Ramya prove the following graphs have super mean labelings: the graph obtained by identifying the endpoints of two or more copies of P_5; the graph obtained from C_n $(n \geq 4)$ by joining two vertices of C_n distance 2 apart with a path of length 2 or 3; Jeyanthi and Rama [1734] use $S(G)$ to denote the graph obtained from a graph G by subdividing each edge of G by inserting a vertex. They prove the following graphs have super mean labelings: $S(P_n \odot K_1), S(B_{n,n}), C_n \odot \overline{K_2}$; the graphs obtained by joining the central vertices of two copies of $K_{1,m}$ by a path P_n (denoted by $\langle B_{m,m} : P_n \rangle$); generalized antiprisms (see §6.2 for the definition), and the graphs obtained from the paths $v_1, v_2, v_3, \ldots, v_n$ by joining each v_i and v_{i+1} to two new vertices u_i and w_i (double triangular snakes).

In [2231] Leena, Sudhakar, Anuradha, Senthil, and Balaj presented two approaches for coding messages using super mean labeling of the union of four stars. Sudhakar, Leen, Anuradha, Indhumathi, Avinash, and Balaji [3721] introduced an innovative cryptographic framework based on super mean labeling of the union of five-star graphs for encoding messages using super mean labelings that represent complex multi-population disease dynamics. Maheswari, Hariprabakaran, and Balaji [2401] introduced a coding technique for converting a text message using a subsuper mean labeling on two and three star graphs to a picture coding message. For more on coding a text message using a graph labelings see [1372], [2397], and [2392].

[3721] new

Lourdusamy and Seenivasan [2344] introduced the notion of super vertex mean labeling as follows. For a (p, q)-graph and an injective function f from the edges to the set $\{1, 2, 3, \ldots, p+q\}$ that induces for each vertex v the label defined by $f^*(v) = (\text{Round} \sum_{e \in E_v} f(e))/d(v)$, where E_v denotes the set of edges in G that are incident to the vertex v, $d(v)$ is the degree of v, and $\text{Round}(x)$ is the integer nearest to x, such that the set of all edge labels and the induced vertex labels is $\{1, 2, 3, \ldots, p+q\}$ is called a *super vertex mean* labeling of G and G is called a *super vertex mean graph*. In [2328] they

investigated the all graphs of order up to 5 and regular graphs of order up to 7 for the property of being super vertex mean and proved that all linear triangular snakes are super vertex mean. Lourdusamy, George, and Seenivasan [2330] proved that all cycles except C_4 are super vertex mean and Lourdusamy and George [2329] proved that linear C_n snakes with at least 2 blocks are super vertex mean graphs for the following cases: $n = 4, 5, 6$, and 7; $n \geq 8$ even; $n \geq 9$ and $n \equiv 1 \bmod 4$; and $n \geq 11$ and $n \equiv 3 \bmod 4$. Lourdusamy and George [2327] showed that all cycles except C_4 have super vertex mean labelings. Inayah, Sudarsana, Musdalifah, and Mangesa [1534] have showed that the total graphs of paths and cycles are super mean graphs. In [3711] Sudarsana, Suryanto, Lusianti, and Putri show how super mean graph labelings can be used to increase the security level of encrypted text on social medias.

A graph G with q edges is called a *k-mean graph* if there is an injective function f from the vertices of G to $\{0, 1, 2, \ldots, k + q - 1\}$ such that when each edge uv is labeled with $(f(u) + f(v))/2$ if $f(u) + f(v)$ is even, and $(f(u) + f(v) + 1)/2$ if $f(u) + f(v)$ is odd, the resulting edge labels are $\{k, k + 1, k + 2, \ldots, k + q - 1\}$. A graph G with q edges is said to have a *restricted k-mean* labeling if there is an injective function f from the vertices of G to $\{k-1, k, k+1, \ldots, k+q-1\}$ such that when each edge uv is labeled with $\{k, k + 1, k + 2, \ldots, k + q - 1\}$, the resulting edge labels $\{k, k + 1, k + 2, \ldots, k + q - 1\}$ are distinct where k is a positive integer. A graph that admits a restricted k-mean labeling is called a *restricted k-mean* graph. Gayathri and Gopi proved some properties of k-mean labelings in [1220]. In [1221] they proved that if G_1 and G_2 are restricted k-mean graphs for all k, then $G_1 \cup G_2$ is restricted k-mean for all k, and if G_1 is a restricted k-mean graph for all $k \geq k_1$ and G_2 is a restricted k-mean graphs for all k, then $G_1 \cup G_2$ is restricted k-mean for all $k \geq k_1$.

A mean graph is called *k-super mean* if vertex labels and the edge labels are $\{k, k + 1, k + 2, \ldots, p + q + k - 1\}$. Jeyanthi, Ramya, Thangavelu [1737] give super mean labelings for $C_m \cup C_n$ and k-super mean labelings for a variety of graphs. Tamilselvi, Akilandeswari, and Suguna [3862] proved that the following graphs admit k-super mean labelings: the graph obtained by subdividing the central edge of the bistar $B_{n,n}$ $(n \geq 2)$, the subdivision graph of $B_{n,n}$, and the corona product of a triangular snake and K_1. Tamilselvi, and Akilandeswari [3861] proved that the following graphs admit k-super heronian mean labelings: $(P_n \times P_2) \odot K_1$ $(n > 1)$, subdivision graphs of triangular snakes T_n $(n > 1)$, subdivision graphs of $T_n \odot K_1$, graphs obtained from a P_n by fusing one edge of C_4 to each vertex of P_n, and graphs obtained from P_n by fusing one vertex two copies of C_6 to each vertex of P_n.

Vasuki and Nagarajan [4064] define H_n, called the H-graph of a path P_n, as the graph obtained from two copies of P_n with vertices $v_1, v_2, \ldots, v_n$ and $u_1, u_2, \ldots, u_n$ by joining the vertices $v_{(n+1)/2}$ and $u_{(n+1)/2}$ if n is odd, and the vertices $v_{\frac{n}{2}+1}$ and $u_{\frac{n}{2}}$ if n is even, and a cyclic snake mC_n as the graph obtained from m copies of C_n by identifying the vertex $v_{(k+2)_j}$ in the jth copy of the vertex $v_{1_{j+1}}$ in the $(j + 1)$th copy if $n = 2k + 1$ and identifying the vertex $v_{(k+1)_j}$ in the jth copy with the vertex $v_{1_{j+1}}$ in the $(j + 1)$th copy if $n = 2k$. They establish the super meanness of even cycles, H-graphs, the coronas of H-graphs, 2-coronas of H-graphs, coronas of cycles, mC_n-snakes $(n \neq 4)$, dragons $P_n(C_m)$

for $m \neq 4$, and $C_m \times P_n$ for $m = 3$ and 5. Vasuki, Sugirtha, and Venkateswari [4073] proved that the subdivision of the following graphs are super mean graphs: H_n, $H_n \odot K_1$, H_n with two pendent edges attached to each vertex, $C_n \odot K_1$ $(n \geq 3)$, slanting ladders, triangular snakes with a pendent edge at each vertex, and $C_m@C_n$ (the graph obtained by attaching paths P_n to C_m by identifying the endpoints of the paths with each successive pairs of vertices of C_m).

Let $G(V, E)$ be a simple graph of order p and size q. Then G is said to be a *relaxed mean graph* if it is possible to label the vertices $x \in V$ with distinct elements $f(x)$ from $\{0, 1, 2 \ldots, q-1, q+1\}$ in such a way that when each edge uv is labeled with $(f(u)+f(v))/2$ if $f(u)+f(v)$ is even and $(f(u)+f(v)+1)/2$ if $f(u)+f(v)$ is odd, then the resulting edge labels $\{1, 2, 3, \ldots, q\}$ are distinct. Such an f is called a *relaxed mean labeling* of G. Balaji, Ramesh, and Sudhaker [455] prove that the disjoint union of any path with $n-1$ edges joining the pendent vertices of distinct paths is a relaxed mean graph and $K_{1,m}$ is not a relaxed mean graph for $m \geq 5$. They also prove that the graph consisting of two stars $K_{1,m}$ and $K_{n,1}$ with an edge in common is a relaxed mean graph if and only if $|m-n| \leq 5$. Balaji and Maheswari [453] proved the following graphs are relaxed mean graphs C_n $(n > 4)$; $K_{2,n}$; triangular snakes; quadrilateral snakes; ${P_n}^2$; $(P_n \times P_2) \odot K_1$; $\overline{K_n} + 2K_2$; $K_2 + mK_1$; $C_3^{(t)}$; the union of any two trees; $P_m \times P_n$ $(m > 1, n > 1)$; and $P_m \times C_n$ $(m > 1)$. They also prove that K_n is not a relaxed mean graph. Maheswari, Ramesh, and Balaji [2398] proved the following graphs are relaxed mean graphs: P_n $(n > 5)$; bistars $B_{m,n}$ if and only if $|m - n| \leq 3$; combs; and $C_3 \cup P_n$ $(n > 1)$. They also proved that $K_{1,n}$ is not a relaxed mean graph for $n > 5$. In [454] Balaji, Ramesh, and Subramanian proved that the following graphs admit relaxed mean labelings: C_n $(n > 4)$, $K_{2,n}$, triangular snakes and quadrilateral snakes, P_m^2, $K_n^c + 2K2$, $K_2 + mK_1$, $P_m \times P_n$ $(m > 2, n > 2)$, and $P_m \times C_n$ $(m > 2, n > 3)$. They also proved that K_n is not a relaxed mean graph. Francis and Balaji proved that the wheels W_n $(n > 4)$ are relaxed mean graphs. Balajia, Ramesh, and Sudhakar [455] proved that the disjoint union of P_{n-1} and the graph obtained by joining the pendent vertices of distinct paths is a relaxed mean graph, $K_{1,m} \cup K_{1,n}$ with an edge in common is a relaxed mean graph if and only if $|m - n| \leq 5$, and that $K_{1,m}$ is not a relaxed mean graph for $m > 5$.

In [447] Balaji, Ramesh, and Subramanian say a graph $G(V, E)$ with p vertices and q edges has a *relaxed Skolem mean labeling* if there is a function f from V to $\{1, 2, \ldots, p+1\}$ such that the induced edge edge map f^* from E to $\{2, 3, \ldots, p+1\}$ given by $f^*(uv) = (f(u) + f(v))/2$ if $f(u) + f(v)$ is even and $f^*(uv) = (f(u) + f(v) + 1)/2 if f(u) + f(v)$ is odd. A graph with a relaxed Skolem mean labeling is called a relaxed Skolem mean graph. They provided necessary and sufficient conditions for graphs that the union of three, four, or five stars to be Skolem mean graphs. They also have a conjecture about the general case of t stars. Ramesh and Jovanna [3119] proved that certain 5-star graphs with partition 3, 2 are relaxed Skolem mean graphs. Results about relaxed Skolem mean labelings of unions of certain four stars are given by Samuel, Vinolin, and Ramesh in [3273]. In [1802] Jovanna proved that the 4-star graph $K_{1,\alpha_1} \cup K_{1,\alpha_2} \cup K_{1,\alpha_3} \cup K_{1,\beta}$ where $\beta - \alpha_1 - \alpha_2 - \alpha_3 = 6$ is a relaxed Skolem mean graph. Karthiyayini1 and Jovanna [1913] proved that the 4-star graph $K_{1,\alpha_1} \cup K_{1,\alpha_2} \cup K_{1,\alpha_3} \cup K_{1,\beta}$ is a relaxed Skolem mean graph when $\alpha_1 \leq \alpha_2 \leq \alpha_3 \leq \beta$ [1913] new

and $\beta - \alpha_1 - \alpha_2 - \alpha_3 = -6$.

In [298] Arockiaraj, Rajesh Kannan, and Durai Baskar introduced the F-centroidal mean labeling of graphs by defining a function f to be an *F-centroidal mean* labeling of a graph $G(V, E)$ with q edges if $f : V(G) \to \{1, 2, 3, \ldots, q+1\}$ is injective and the induced function $f^* : E(G) \to \{1, 2, 3, \ldots, q\}$ defined as $f^*(uv) = \left\lfloor \frac{2\ [f(u)^2+f(u)f(v)+f(v)^2]}{3\ [f(u)+f(v)]} \right\rfloor$ for all $uv \in E(G)$ is bijective. A graph that admits an F-centroidal mean labeling is called an *F-centroidal mean* graph. They discussed the F-centroidal meanness of the tree $P_n(X_1, X_2, \ldots X_n)$ obtained from a path on n vertices by attaching X_i pendent vertices at each i^{th} vertex of the path for $1 \leq i \leq n$, the twig graph $TW(P_n)$, the graph $P_n \circ S_m$ for $m \leq 4$, $P_m \times P_n$ for $m \leq 3$, ladders, $P_n \circ K_2$, P_a^b for $a \geq 2$ and $b \leq 3$, the middle graphs and splitting graphs of paths, the total graphs of paths, P_n^2, and $P(1, 2, \ldots, n-1)$ the graph obtained by replacing each i^{th} edge of P_n by identifying its end vertices with the vertices of the two element component of $K_{2,i}$. In the same article they introduced super F-centroidal mean graphs [298] as follows. Let G be a graph and $f : V(G) \to \{1, 2, 3, \ldots, p+q\}$ be an injection. With f^* defined as for the F-centroidal case f is called a *super F-centroidal mean* labeling if $f(V(G)) \cup \{f^*(uv) : uv \in E(G)\} = \{1, 2, 3, \ldots, p+q\}$. A graph that admits a super F-centroidal mean labeling is called a *super F-centroidal mean* graph. They proved that the following graphs are super F-centroidal mean graphs: paths, cycles, the union of any number of paths, the mirror graph of P_n, $P_n \circ S_m$, $TW(P_n)$, $P_n \cup C_m$, P_n^2, and dragons $P_n(C_m)$ the graph obtained from C_m by identifying an end vertex of P_n at a vertex of C_m.

In [1317] Gopi and Kumar proved the following graphs are F-centroidal mean graphs: the mirror graph $M(P_n)$ $(n \geq 3)$, $\mathrm{Spl}(P_n)$ $(n \geq 3)$, P_n^2 $(n \geq 3)$, $VD(G)$, the *tortoise* graph T_n $(n \geq 4)$ (obtained from a path $v_1, v_2, \ldots, v_n$ by attaching an edge between v_i and v_{n-i+1} for $1 \leq i \leq \lfloor n/2 \rfloor$. $(n = 1, 3 \pmod 4)$, and PC_n $(n \geq 5)$ (obtained from $C_n = v_1, v_2, \ldots, v_n$ by adding chords joining v_i and v_{n-i+2} for $2 \leq i \leq t$, where $t = \lfloor n/2 \rfloor$), In [1316] they proved $D(T_n)$ $(n \geq 3)$, $AD(T_n)$ $(n \geq 4)$, Q_n $(n \geq 2)$, $A(Q_n)$ $(n \geq 2)$ $AD(Q_n)$ $(n \geq 2)$, and SL_n $(n \geq 2)$ are F-centroidal mean graphs.

Arockiaraj, Kannan, Manivannan, and Durai Baskar [300] investigated the F-centroidal mean property for paths, cycles, stars, K_n, $P_n \circ S_1$, the triangular snakes, arbitrary subdivisions of $K_{1,3}$, and some line graphs.

In [457] and [458] Balaji, Ramesh, and Subramanian use the term "Skolem mean" labeling for super mean labeling. They prove: P_n is Skolem mean; $K_{1,m}$ is not Skolem mean if $m \geq 4$; $K_{1,m} \cup K_{1,n}$ is Skolem mean if and only if $|m - n| \leq 4$; $K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is Skolem mean if $|m-n| = 4+l$ for $l = 1, 2, 3, \ldots, m = 1, 2, 3, \ldots$, and $l \leq m < n$; $K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is not Skolem mean if $|m - n| > 4 + l$ for $l = 1, 2, 3, \ldots, m = 1, 2, 3, \ldots, n \geq l + m + 5$ and $l \leq m < n$; $K_{1,l} \cup K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is Skolem mean if $|m - n| = 4 + 2l$ for $l = 2, \ldots, m = 2, 3, 4 \ldots, n = 2l + m + 4$ and $l \leq m < n$; $K_{1,l} \cup K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is not Skolem mean if $|m - n| > 4 + l$ for $l = 1, 2, 3, \ldots, m = 1, 2, 3, \ldots, n \geq l + m + 5$ and $l \leq m < n$; $K_{1,l} \cup K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is not Skolem mean if $|m - n| > 4 + 2l$ for $l = 2, \ldots, m = 2, 3, 4 \ldots, n \geq 2l + m + 5$ and $l \leq m < n$; $K_{1,l} \cup K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is Skolem mean if $|m - n| = 7$ for $m = 1, 2, 3, \ldots, n = m + 7$ and $1 \leq m < n$; and $K_{1,l} \cup K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is not Skolem mean if $|m - n| > 7$ for $m = 1, 2, 3, \ldots, n \geq m + 8$

and $1 \leq m < n$. Balaji [456] proved that $K_{1,l} \cup K_{1,m} \cup K_{1,n}$ is Skolem mean if $|m-n| < 4+l$ for integers $1, m \geq 1$ and $l \leq m < n$. In [3468] Shainy and Balaji determined necessary and sufficient conditions for the disjoint union of three stars to be Skolem mean. Ramesh, Sopna, and Gnanaselvi [3120] proved that certain graphs that are the union of four star graphs admit Skolem mean labelings.

In [3032] Prakash, Gopi, and Shalini introduced the following graph labeling. A graph $G(V, E)$ with p vertices and q edges where $p < q + 1$ is said to be an *anti skolem mean graph* if it is possible to label the vertices with distinct labels from $\{0, 1, 2, \ldots, q + 1\}$ in such a way that if each edge uv is labeled with $f^*(uv) = \lceil (f(u) + f(v))/rceil$, the resulting edge labels are distinct labels from the set $\{2, 3, \ldots, q + 1\}$. They proved that quadrilateral snake related graphs have anti skolem mean labelings.

In [3502] Shendra Shainy, Hariprabakaran, Swathy, and Balaji provided a technique for coding a secret messages by applying a Skolem mean-like labeling on a graph obtained from the two stars of the form $K_{1,n}$ and $K_{1,n+1}$ that are joined by an edge from one vertex of degree 1 in $K_{1,n}$ to one vertex of degree 1 in $K_{1,n+1}$.

In [1758] Jeyanthi, Selvi, and Ramya prove that $C_m \cup C_n$, $(P_n + K_1) \cup (n-2)K_2$ $(n > 2)$, $(P_n + K_2) \cup (2n-3)K_2$ $(n \geq 2)$ and $W_n \cup (n-1)K_2$ $(n \geq 3)$ are Skolem difference mean graphs. In [1759] they show that the union of any finite number of paths, the union of any finite number of stars, $G \cup nK_2$ where G is Skolem difference mean and all the vertex labels are odd, $C_m \cup P_m$ $(m \geq 2)$, $K_{m,n} \cup (m-1)(n-1)K_2$, and $K_{1,1,n} \cup (n-1)K_2$. are Skolem difference mean graphs. Vijaya Kumar, Thulukkanam, and Thirusangu [4095] proved the exitence of Skolem difference mean labelings of duplicate paths, dupicate combs, duplicate ladders, duplicate triangular snakes, and duplicate quadrilateral snakes. [4095] new

In [1738] Jeyanthi, Ramya, and Thangavelu proved the following graphs have super mean labelings: the one point union of any two cycles, graphs obtained by joining any two cycles by an edge (dumbbell graphs), $C_{2n+1} \odot C_{2m+1}$, graphs obtained by identifying a copy of an odd cycle C_m with each vertex of C_n, the quadrilateral snake Q_n, where n is odd, and the graphs obtained from an odd cycle $u_1, u_2, \ldots, u_n$ by joining the vertices u_i and u_{i+1} by the path P_m (m is odd) for $1 \leq i \leq n-1$ and joining vertices u_n and u_1 by the path P_m. Jeyanthi, Ramya, Thangavelu, and Aditanar [1736] give super mean labelings of $C_m \cup C_n$ and T_p-trees. Vasuki and Arockiaraj [4062] proved that nC_4, $n > 1$, triangular grid graphs, the edge mC_n-snakes, and the braid graphs are super mean graphs. They further proved that the graphs obtained by identifying an edge of two cycles C_m and C_n is a super mean graph.

In [1730] Jeyanthi and Ramya define $S_{m,n}$ as the graph obtained by identifying one endpoint of each of n copies of P_m and $< S_{m,n} : P_m >$ as a graph obtained by identifying one end point of a path P_m with the vertex of degree n of a copy of $S_{m,n}$ and the other endpoint of the same path to the vertex of degree n of another copy of $S_{m,n}$. They prove the following graphs have super mean labelings: caterpillars, $< S_{m,n} : P_{m+1} >$, and the graphs obtained from P_{2m} and $2m$ copies of $K_{1,n}$ by identifying a leaf of ith copy of $K_{1,n}$ with ith vertex of P_{2m}. They further establish that if T is a T_p-tree, then $T \odot K_1$, $T \odot \overline{K_2}$, and, when T has an even number of vertices, $T \odot \overline{K_n}$ $(n \geq 3)$ are super mean graphs.

In [948] Dhanalakshmi and Parvathi define a *mean square cordial* labeling of a graph

$G(V,E)$ with p vertices and q edges as one for which there is a bijection from V to $\{0,1\}$ such that when each edge uv is assigned the label $\lceil (f(u)^2+f(v)^2)/2 \rceil$ the condition that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1 and the number of edges labeled with 0 and the number of edges labeled 0 and the number of edges labeled with 1 differ by at most 1 is satisfied. In [948] Dhanalakshmi and Parvathi proved that helms, closed helms, gears, sunlet graphs, fans, and $C_p \odot nK_1$ admit mean square cordial labelings. In [949] Dhanalakshmi and Parvathi showed that the paths, combs, n-centipedes, and stars admit mean square cordial labelings. They also proved that the induced subgraph obtained by the upper approximation of any subgraph H of the above acyclic graphs admits a mean square cordial labeling. In [950] Dhanalakshmi and Parvathi prove that bistars, subdivisions of stars, coconut trees, banana trees, the one-point union of C_n and $K_{1,n}$, and $P_m \odot K_{1,n}$ admit mean square cordial labelings. Dhanalakshmim [951] proved that shell-butterfly graphs with shell orders $m > 2$ and $n > 2$ admit mean square cordial labelings. Dhanalakshmi and Thirunavukkarasu [952] z proved pentagonal snakes, subdivision of a pentagonal snakes, double pentagonal snakes, and alternate pentagonal snakes admit mean square cordial labelings.

In [3292] Sandhya, Somasundaram, and Anusa say a graph with q edges is a *root square mean* graph if it is possible to label the vertices with distinct elements from $\{1,2,\ldots,q+1\}$ in such a way that when each edge uv is labeled with $\lfloor \sqrt{(f(u)^2+f(v)^2)/2} \rfloor$ or $\lceil \sqrt{(f(u)^2+f(v)^2)/2} \rceil$, the resulting edge labels are distinct. They prove that paths, cycles, combs, ladders, triangular snakes, quadrilateral snakes, $K_{1,n)}$ $(1 \leq 6)$, and K_n for $n = 1, 2$ and 3 are root square mean graphs. In [3281] they proved that the following graphs admit root square mean labelings: double triangular snakes, alternate double triangular snakes, double quadrilateral snakes, alternate double quadrilateral snakes, and polygonal chains. Sandhya, Somasundaram, and Anusa [3277] showed that the following graphs are root square mean: cycles union triangular snakes, crowns union triangular snakes, the corona product of cycles and double triangular snakes, crowns union double triangular snakes, cycles union quadrilateral snakes, crowns union quadrilateral snakes, the corona product of cycles and double triangular snakes, crowns union double quadrilateral snakes, Sandhya, Somasundaram, and Anusa [3279] provided square mean labelings for some degree splitting graphs. In [3280]they showed that the following graphs are root square mean: $C_m \cup (P_n \odot K_1), (C_m \odot K_1) \cup (P_n \odot K_1), (C_m \odot \overline{K_2}) \cup (P_n \odot K_1), C_m \cup (P_n \times P_2), (C_m \odot K_1) \cup (P_n \odot P_2), C_m \cup (P_n \odot K_3), (C_m \odot K_1) \cup (P_n \odot K_3), (C_m \odot K_3) \cup P_n$ and $(C_m \odot K_3) \cup (P_n \odot K_3)$. Sandhya, Somasundaram, and Anusa [3293] proved that dragons, crowns, kites, coronas of certain graphs,$C_n \odot \overline{K_2}$ and $C_n \odot \overline{K_3}$, and graphs obtained by attaching each vertex of P_n to the central vertex of $K_{1,2}$ and to $K_{1.3}$ are root square graphs.

Abdul Saleem, Vanitha Jothy, and Haajira [8] showed that the following graphs admit root square mean labelings: paths, cycles, combs, ladders, triangular and quadrilateral snakes, stars, complete graphs, and arrow graphs. Periaswamy [2827] proved that the following graphs admit root square mean labelings $C_h \cup P'_h, (C_h \odot K_1) \cup P'_h, (C_h \odot K_2) \cup P'_h, (C_h \odot K_3) \cup P'_h, (C_h \odot K_1) \cup (P'_h \odot K_1), (C_h \odot K_2) \cup (P'_h \odot K_1)$, and $(C_h \odot K_3) \cup (P'_h \odot K_1)$.

[8] new

[2827] new

Meena and Mani [2475] investigated the existence of root square mean labelings for

certain cycle related graphs. They showed that the following graphs admit root square mean labelings: the path union of two or three copies of C_n, the P_k path union of two copies of C_n, the path union of two or three copies of $C_n \odot K_1$, and the P_k path union of two copies of $C_n \odot K_1$. In [2476] they proved that various theta related graphs are root square mean graphs.

Arockiaraj, Durai Baskar, and Rajesh Kannan [289] calls a graph G with p vertices and q edges a *F-root square mean graph* if there is an injective function f from the vertices of $V(G)$ to $\{1, 2, \ldots, q+1\}$ such that for each edge uv the induced function $f^*(uv) = \lfloor \sqrt{(f(u)^2 + f(v)^2)/2} \rfloor$ is bijective. They proved that the following are F-root square mean graphs: paths, the graph $P_n \circ S_m$ obtained from the path P_n by attaching m pendent vertices to each vertex of P_n, twigs $TW(P_n)$, the graph $[P_n; S_m]$ obtained from n copies of S_m and the path P_n by joining u_i with the central vertex $v_1^{(i)}$ of the i^{th} copy of S_m with of an edge for $i \leq n$, the mirror graph of P_n, the total graph of P_n, P_n^2, ladders, and slanting ladders. In [290], Arockiaraj, Durai Baskar, and Rajesh Kannan analyzed that the line graph operation preserves the F-root square meanness of line graph of the path, cycle, star, $P_n \circ S_1, P_n \circ S_2, [P_n; S_1], S(P_n \circ S_1)$, ladder, slanting ladder, the crown graph $C_n \circ S_1$ and the arbitrary subdivision of S_3. Gopi [1310] proved that triangular snakes T_n $(n \geq 2)$, $A(T_n)$ $(n \geq 3), D(T_n)$ $(n \geq 2)$, quadrilateral snakes, $A(Q_n), D(Q_n)$ $(n \geq 3)$ are F-root square mean graphs.

Rajesh Kannan, Vikrama Prasad, and Gopi [1887] call a graph G with p vertices and q edges a *super root mean graph* if there is an injective function f from the vertices of G to $\{1, 2, \ldots, p+q\}$ such that for each edge uv the induced function $f^*(uv) = \lfloor \sqrt{(f(u)^2 + f(v)^2)/2} \rfloor$ or $f^*(uv) = \lceil \sqrt{(f(u)^2 + f(v)^2)/2} \rceil$ yields the set of vertex labels and edge labels $\{1, 2, \ldots, p+q\}$. They proved the following are super root square mean graphs: $P_m \cup P_m$ $(m, n \geq 3)$; $P_m \cup (P_n \odot K_1)$ $(m, n \geq 3)$; $(P_m \odot K_1) \cup (P_n \odot K_1)$ $(m, n \geq 3)$; the union of a path and a triangular snake; and the union of $P_n \odot K_1$ and a triangular snake. Gopi and Kalaiyarasi [1312] prove that the following graphs have a super root square mean labeling: P_n^2 $(n \geq 4)$, slanting ladders $SL_n (n \geq 3)$, triangular snakes with a pendent edge attached to each vertex, and quadrilateral snakes with a pendent edge attached to each vertex.

Chitra devi and Saravana Kumar [809] proved that $nP_m, nK_3, P_n \odot \overline{K_2}$, the middle graph of path P_n (graphs obtained by starting with a path and joining every consecutive pair of vertices excluding the two end vertices of the path to a new isolated vertex), and $C_m \odot P_n$ admits super root square mean labelings. Venkatesan and Thirugnanasambandam [4080] proved that the following graphs admit super root square mean labelings: P_n^2 $(n \geq 4)$, slanting ladders, $T_n \odot K_1$ $(n \geq 3)$ (T_n is a triangular snake), and $Q_n \odot K_1$ (Q_n is a quadrilateral snake). Orias and Pedrano [2741] determined the super root square mean labelings of $C_n \odot K_1$, middle cycles (graphs obtained by starting with a cycle and joining every consecutive pair of vertices of the cycle to a new isolated vertex), polygonal chains, alternate polygonal chains, and kayak paddles.

In [160] Akilandeswari calls a graph is a *k-super root square mean* graph if it is possible to label the vertices with distinct elements from $\{k, k+1, k+2, \ldots, p+q+k-1\}$ in such a way that when each edge uv is labeled with $\lfloor \sqrt{(f(u)^2 + f(v)^2)/2} \rfloor$ or

$\lceil\sqrt{(f(u)^2+f(v)^2)/2}\rceil$, the union of the vertex and edge labels is the set $\{k, k+1, k+2, \ldots, p+q+k-1\}$. He proved that $P_n \odot K_{1,2}, P_n \odot K_{1,3}$, the corona product of the quadrilateral snake and K_1, double triangular snakes, and the corona product of the double triangular snake and K_1 admit k-super root square mean labelings.

A *radio mean square* labeling of a connected graph is an injective map h from the set of vertices of the graph G to the set of positive integers $\mathbb{N}$, such that for any two distinct vertices x, y, the inequality $d(x,y) + \lceil (h(x))^2 + (h(y))^2/2\rceil \geq \dim(G) + 1$ holds. For a particular radio mean square labeling h, the maximum number of $h(v)$ taken over all vertices of G is called its spam, denoted by $\mathrm{rmsn}(h)$, and the minimum value of $\mathrm{rmsn}(h)$ taking over all radio mean square labeling h of G is called the *radio mean square number* of G, denoted by $\mathrm{rmsn}(G)$. In [434] Badr, Nada, Al-Shamiri, Abdel-Hay, and Elrokh investigated the radio mean square numbers for paths and cycles. For a graph G, they present an approximate algorithm to determine $\mathrm{rmsn}(G)$. Finally, they introduced a new mathematical model to find the upper bound of $\mathrm{rmsn}(G)$ for graph G.

In [2053] Kulandhai Therese and Romila introduced the notion of cube root cube mean labeling as follows. For a graph $G(V,E)$ with $|(V(G)| = p$ and $|E(G)| = q$, let $f : V(G) \to \{1, 2, \ldots, q+1\}$ be an injective function. The induced edge labeling f^* for a vertex labeling f is defined by $f^*(e) = \lfloor \sqrt[3]{(f(u)^3 + f(v)^3)/2}\rfloor$ or $\lceil \sqrt[3]{(f(u)^3 + f(v)^3)/2}\rceil$ for all $e = uv \in E(G)$. If f is a bijection it is called a *cube root cube mean labeling*. If such labeling exists, G is said to be a *k-cube root cube mean* graph. They proved that paths, combs, ladders, and quadrilateral snakes admit cube root cube mean labelings.

In [3044] Princy Kala introduced the notion of k-super cube root cube mean labeling as follows. For a graph $G(V,E)$ with $|(V(G)| = p$ and $|E(G)| = q$ let $f : V(G) \to \{k, k+1, k+2, \ldots, p+q+k-1\}$ be an injective function. The induced edge labeling f^* for a vertex labeling f is defined by $f^*(e) = \lfloor \sqrt[3]{(f(u)^3 + f(v)^3)/2}\rfloor$ or $\lceil \sqrt[3]{(f(u)^3 + f(v)^3)/2}\rceil$ for all $e = uv \in E(G)$. If f is a bijection and $f(V(G)) \cup f^*(E(G)) = \{k, k+1, k+2, \ldots, p+q+k-1\}$, f is called a *k-super cube root cube mean labeling*. If such labeling exists, then G is said to be a *k-super cube root cube mean* graph. He proved the existence k-super cube root mean labelings for triangular snakes, double triangular snakes, quadrilateral snakes, double quadrilateral snakes, alternate triangular snakes, alternate double triangular snakes, alternate quadrilateral snakes, and alternate double quadrilateral snakes. In [3045] Princy Kala proved that the corona products of a triangular snake and K_1, an alternate triangular snake and K_1, an alternate triangular snake and $2K_1$, an alternate quadrilateral snake and K_1, $P_n \odot K_{1,2}$, and $P_n \odot K_{1,3}$ are k-super cube root cube mean graphs.

Let G be a graph and let $f : V(G) \to \{1, 2, \ldots, n\}$ be a function such that the label of the edge uv is $(f(u)+f(v))/2$ or $(f(u)+f(v)+1)/2$ according as $f(u)+f(v)$ is even or odd and $f(V(G)) \cup \{f^*(e) : e \in E(G)\} \subseteq \{1, 2, \ldots, n\}$. If n is the smallest positive integer satisfying these conditions together with the condition that all the vertex and edge labels are distinct and there is no common vertex and edge labels, then n is called the *super mean number* of a graph G and it is denoted by $S_m(G)$. Nagarajan, Vasuki, and Arockiaraj [2642] proved that for any graph of order p, $S_m(G) \leq 2^p - 2$ and provided an upper bound of the super mean number of the graphs: $K_{1,n}$ $n \geq 7$; $tK_{1,n}$, $n \geq 5$, $t > 1$;

the bistar $B(p,n),\ p>n$; the graphs obtained by identifying a vertex of C_m and the center of $K_{1,n},\ n\geq 5$; and the graphs obtained by identifying a vertex of C_m and the vertex of degree 1 of $K_{1,n}$. They also gave the super mean number for the graphs $C_n, tK_{1,4}$, and $B(p,n)$ for $p=n$ and $p=n+1$.

Manickam and Marudai [2411] defined a graph G with q edges to be an *odd mean graph* if there is an injective function f from the vertices of G to $\{1,3,5,\ldots,2q-1\}$ such that when each edge uv is labeled with $(f(u)+f(v))/2$ if $f(u)+f(v)$ is even, and $(f(u)+f(v)+1)/2$ if $f(u)+f(v)$ is odd, then the resulting edge labels are distinct. Such a function is called a *odd mean* labeling. For integers a and b at least 2, Vasuki and Nagarajan [4066] use P_a^b to denote the graph obtained by starting with vertices $y_1,y_2,\ldots,y_a$ and connecting y_i to y_{i+1} with b internally disjoint paths of length $i+1$ for $i=1,2,\ldots,a-1$ and $j=1,2,\ldots,b$. For integers $a\geq 1$ and $b\geq 2$ they use $P^b_{\langle 2a\rangle}$ to denote the graph obtained by starting with vertices $y_1,y_2,\ldots,y_{a+1}$ and connecting y_i to y_{i+1} with b internally disjoint paths of length $2i$ for $i=1,2,\ldots,a$ and $j=1,2,\ldots,b$. They proved that the graphs $P_{2r,m}, P_{2r+1,2m+1}$, and $P^m_{\langle 2r\rangle}$ are odd mean graphs for all values of r and m. Jeyanthi and Gomathi [1667] proved the edge linked cyclic snake $EL(kC_n)$ $(n\geq 6)$ is an odd mean graph. Sasikala and Harikumaran [3314] proved that double triangular snakes, jelly fish, and alternate triangular snakes are vertex odd mean graphs. [3314] new

For a T_p-tree T with m vertices $T@P_n$ is the graph obtained from T and m copies of P_n by identifying one pendent vertex of ith copy of P_n with ith vertex of T. For a T_p-tree T with m vertices $T@2P_n$ is the graph obtained from T by identifying the pendent vertices of two vertex disjoint paths of equal lengths $n-1$ at each vertex of T. Ramya, Selvi and Jeyanthi [3126] prove that $P_m\odot\overline{K_n}$ $(m\geq 2, n\geq 1)$ is an odd mean graph, T_p trees are odd mean graphs, and, for any T_p tree T, the graphs $T@P_n$, $T@2P_n$, $\langle T\tilde{o}K_{1,n}\rangle$ are odd mean graphs.

For a T_p-tree T with m vertices let $T\hat{o}C_n$ denote the graph obtained from T and m copies of C_n by identifying a vertex of i^{th} copy of C_n with i^{th} vertex of T and $T\tilde{o}C_n$ denote the graph obtained from T and m copies of C_n by joining a vertex of i^{th} copy of C_n with i^{th} vertex of T by an edge. In [1761] Selvi, Ramya, and Jeyanthi prove that for a T_p tree T the graphs $T\hat{o}C_n$ $(n>3, n\neq 6)$ and $T\tilde{o}C_n,\ (n>3,\ n\neq 6)$ are odd mean graphs.

Ramya, Selvi, and Jeyanthi [3125] prove that for a T_p-tree T the following graphs are odd mean graphs: $T@P_n,\ T@2P_n,\ P_m\odot\overline{K_n}$, and the graph obtained from T and m copies of $K_{1,n}$ by joining the central vertex of ith copy of $K_{1,n}$ with ith vertex of T by an edge.

For a graph G and some fixed vertex v of G, Pooranam, Vaski, and Suganthi [2993] proved the following graphs have odd mean labelings: graphs obtained from a path $P_m: u_1u_2\cdots u_m$ and G by joining u_i to v to the ith copy of G. Their results include the cases where $G=C_{4n}$, a star, or the cube Q_3. For a graph G and some fixed vertex v of G they also proved the existence of odd mean labelings for graphs obtained from a path $P_m: u_1u_2\cdots u_m$ and G by identifying u_i with v in the ith copy of G, where G is Q_3 or $C^{(2)}_{4n}$ and v is the vertex of $C^{(2)}_{4n}$ of degree 4. Vijaya kumar, Ulaganathan, and Thirusangu [4096] proved that nontrivial cases of duplicate graphs of ladders, extended duplicate graphs of stars, extended duplicate graph of bistars, and the extended duplicate graph of double stars admit mean labelings, odd mean labelings, and even mean labelings. [4096] new

A graph G is said to be *vertex odd mean* graph if there exist an injective function $f : V(G)$ to $\{1, 3, 5, \ldots, 2|E(G)| - 1\}$ such that the induced mapping $f^* : E(G)$ to the set of positive integers defined by $f^*(uv) = (f(u) + f(v))/2$ is injective. Such a function is called a *vertex odd mean* labeling. A graph G is called a *vertex even mean* graph if there exist an injective function $f : V(G)$ to $\{2, 4, 6, \ldots, 2|E(G)|\}$ such that the induced mapping $f^* : E(G)$ to the set of positive integers defined by $f^*(uv) = (f(u) + f(v))/2$ is injective. Such a function is called a *vertex even mean* labeling. Revathi [3165] gave vertex odd and even mean labelings for nontrivial umbrella graphs, mongolian tents, and $K_1 + C_n$. Anitha, Selvam, and Thirusangu [260] proved that the extended duplicate graph of kite graph admits mean, even mean, and odd mean labelings, Prajapati and Raval [3021] and [3022] proved that cyclic snakes for $n = 3, 4, 5$ are vertex even mean and vertex odd mean graphs. They have proved that double cyclic snake and alternating cyclic snake for $n = 3, 4, 5$ are vertex odd mean and vertex even mean graph. Prajapati and Raval [3023] proved that every vertex even mean graph is vertex odd mean graph and vice versa. They also proved that triangular snakes TS_n, $TS_n \odot P_2$, quadrilateral snakes QS_n, $QS_n \odot P_2$, double quadrilateral snakes, alternating double quadrilateral snakes, jelly fish, crowns, shells, and fans are vertex even mean and vertex odd mean graphs. Raval and Prajapati [3137] proved that cyclic snakes (obtained by replacing each vertex of a path by a cycle), quadrilateral snakes, pentagonal snakes, and alternating quadrilateral snakes admit even vertex and odd mean labelings. They also proved that even vertex odd mean graphs are even mean graphs. An *extended vertex odd mean* labeling of $G(V, E)$ is an injection $f : V \to \{2t - 1 : 1 \leq t \leq q + 1\}$ so that the edge labels are assigned by f^* defined by $f^*(uv) = (f(u) + f(v)/2$ are distinct. Komalamba and Vijayakumar [1993] [1993] new a proved the existence of extended vertex odd mean labeling for the duplicate graphs of the paths, combs, twigs, stars, and bistars.

A graph $G(V, E)$ with q edges is called an *even vertex odd mean* graph if there exists an injective labeling $f : V(G)$ to $\{0, 2, 4, 6, \ldots, 2q\}$ such that for the induced edge labeling of E given by $f^*(uv) = (f(u) + f(v))/2$, we have $f^*(E) = \{1, 3, 5, \ldots, 2q - 1\}$. Basher [552] proved that the graphs obtained by super subdivisions of paths, cycles, combs, crowns, and planar grids are even vertex odd mean graphs. In [554] Basher gave sufficient conditions for certain uniform theta graphs to be even vertex odd mean graphs.

Maheswari, Azhagarasi, and Samuvel [2389] prove that the path union P_m $(m \geq 2)$ of even cycles C_{2n} $(n \geq 3)$ with parallel P_3 chords and the path union P_m $(m \geq 2)$ of odd cycles C_{2n+1} $(n \geq 3)$ with parallel P_3 chords are vertex even mean and vertex odd mean graph. They also prove that crowns $C_{2n} \odot K_1$ $(n \geq 3)$ with parallel P_3 chords and crowns $C_{2n+1} \odot K_1$ $(n \geq 3)$ with parallel P_3 chords possesses vertex even mean labeling and vertex odd mean labeling.

In [224] Amuthavalli and S. Dineshkumar say a (p, q) graph G has a *k-odd edge mean* labeling if there exists an injection f from the edges of G to $\{0, 1, 2, 3, \ldots, 2k+2p-3\}$ such that the induced map f^* defined on $V(G)$ by $f^*(v) = \lceil \sum(f(vu)/\deg(v) \rceil$ is a bijection from V to $\{2k-1, 2k+1, 2k+3, \ldots, 2k+2p-3\}$. A graph that admits a k-odd edge mean labeling is called a *k-odd edge mean* graph. They proved that P_n $(n \neq 4)$, $K_{1,2n}$ $(n \geq 2)$, and C_n $(\neq 6, 7)$ are k-odd edge mean graphs for all k.

Thirugnanasambandam, Chitra, and Vishnupriya [3891] introduced a notion of prime odd mean graphs as follows. A graph G with p vertices and q edges for which there exists an injective function $f : V(G) \to \{1, 3, 5, \ldots, 2q + 1\}$ such that gcd $(f(u), f(v)) = 1$ and the induced edge labeling $f^* : E(G) \to \{2, 4, 6, \ldots, 2p - 2\}$ defined by $f^*(uv) = \lceil (f(u) + f(v))/2 \rceil$ is injective is called a *prime odd mean* graph. They proved that caterpillars, cycles, kites, ladders, spiders, and stars are prime odd mean graphs.

In [808] Chitra, Priya, and Vishnupriya [808] introduced the notion of *odd Fibonacci mean* labeling of graph G as an injective f from the vertices of G to the odd Fibonacci numbers such that the induced edge labeling f^* defined by $f^*(uv) = (f(u) + f(v))/2$ is injective. A graph that admits an odd Fibonacci mean labeling is called an *odd Fibonacci mean* graph. They proved the following graphs admit odd Fibonacci mean labelings: wheels, helms, gears, cycles, banana trees, tadpoles, fire crackers, fans, stars, ladders, complete tripartite graphs, and Dutch windmills.

Gayathri and Amuthavalli [1203] (see also [223]) say a (p, q)-graph G has a (k, d)-*odd mean* labeling if there exists an injection f from the vertices of G to $\{0, 1, 2, \ldots, 2k - 1 + 2(q - 1)d\}$ such that the induced map f^* defined on the edges of G by $f^*(uv) = \lceil (f(u)+f(v))/2 \rceil$ is a bijection from edges of G to $\{2k-1, 2k-1+2d, 2k-1+4d, \ldots, 2k-1+2(q-1)d\}$. When $d = 1$ a (k, d)-odd mean labeling is called k-*odd mean*. For $n \geq 2$ they prove the following graphs are k-odd mean for all k: P_n; combs $P_n \odot K_1$; crowns $C_n \odot K_1$ $(n \geq 4)$; bistars $B_{n,n}$; $P_m \odot \overline{K_n}$ $(m \geq 2)$; $C_m \odot \overline{K_n}$; $K_{2,n}$; C_n except for $n = 3$ or 6; the one-point union of C_n $(n \geq 4)$ and an endpoint of any path; grids $P_m \times P_n$ $(m \geq 2)$; $(P_n \times P_2) \odot K_1$; arbitrary unions of paths; arbitrary unions of stars; arbitrary unions of cycles; the graphs obtained by joining two copies of C_n $(n \geq 4)$ by any path; and the graph obtained from $P_m \times P_n$ by replacing each edge by a path of length 2. They prove the following graphs are not k-odd mean for any k: K_n; K_n with an edge deleted; $K_{3,n}$ $(n \geq 3)$; wheels; fans; friendship graphs; triangular snakes; Möbius ladders; books $K_{1,m} \times P_2$ $(m \geq 4)$; and webs. For $n \geq 3$ they prove $K_{1,n}$ is k-odd mean if and only if $k \geq n - 1$. Gayathri and Amuthavalli [1204] prove that the graph obtained by joining the centers of stars $K_{1,m}$ and $K_{1,n}$ are k-odd mean for $m = n, n + 1, n + 2$ and not k-odd mean for $m > n + 2$. For $n \geq 2$ the following graphs have a (k, d)-mean labeling [1232]: $C_m \cup P_n$ $(m \geq 4)$ for all k; arbitrary unions of cycles for all k; P_{2m}; P_{2m+1} for $k \geq d$; (P_{2m+1} is not (k, d)-mean when $k < d$); combs $P_n \odot K_1$ for all k; $K_{1,n}$ for $k \geq d$; $K_{2,n}$ for $k \geq d$; bistars for all k; nC_4 for all k; and quadrilateral snakes for $k \geq d$.

In [3399] Seoud and Salim [3396] proved that a graph has a k-odd mean labeling if and only if it has a mean labeling. In [3399] Seoud and Salim give upper bounds of the number of edges of graphs with a (k, d)-odd mean labeling

Pricilla [3040] defines an *even mean labeling* of a graph G as an injective function f from the vertices of G to $\{2, 4, \ldots, 2|E(G)|\}$ such that the edge labels given by $(f(u) + f(v))/2$ are distinct. Vaidya and Vyas [4032] proved that $D_2(P_n)$, $M(P_n)$, $T(P_n)$, $S'(P_n)$, P_n^2, P_n^3, switching of pendent vertex in P_n, $S'(B_{n,n})$, double fans, and duplicating each vertex by an edge in paths are even mean graphs.

Gayathri and Gopi [1212] defined a graph G with q edges to be an k-*even mean graph* if there is an injective function f from the vertices of G to $\{0, 1, 2, \ldots, 2k + 2(q - 1)\}$

such that when each edge uv is labeled with $(f(u)+f(v))/2$ if $f(u)+f(v)$ is even, and $(f(u)+f(v)+1)/2$ if $f(u)+f(v)$ is odd, then the resulting edge labels are $\{2k, 2k+2, 2k+4, \ldots, 2k+2(q-1)\}$. Such a function is called a *k-even mean* labeling. In [1212] they proved that the graphs obtained by joining two copies of C_n with a path P_m are k-even mean for all k and all $m, n \geq 3$ when $n \equiv 0, 1 \pmod 4$ and for all $k \geq 1,\ m \geq 7$, and $n \geq 3$. In [1214] Gayathri and Gopi proved that various graphs obtained by joining two copies of stars $K_{1,m}$ and $K_{1,n}$ with a path by identifying the one endpoint of the path with the center of one star and the other endpoint of the path with the center of the other star are k-even mean. In [1213] they proved that various graphs obtained by appending a path to a vertex of a cycle are k-even mean. In [1215] they proved that $C_n \cup P_m$, $n \geq 4,\ m \geq 2$, is k-even mean for all k. Gayathri and Gopi [1218] proved the following are k-even mean graphs: shadow graphs of stars with at least 3 vertices; edge duplication graphs of cycles with at least 4 vertices; and vertex duplication graphs of paths and cycles with at least 4 vertices.

Gayathri and Gopi [1216] say graph G with q edges has a (k,d)-*even mean* labeling if there exists an injection f from the vertices of G to $\{0, 1, 2, \ldots, 2k+2(q-1)d\}$ such that the induced map f^* defined on the edges of G by $f^*(uv) = (f(u)+f(v))/2$ if $f(u)+f(v)$ is even and $f^*(uv) = (f(u)+f(v)+1)2$ if $f(u)+f(v)$ is odd is a bijection from edges of G to $\{2k, 2k+2d, 2k+4d, \ldots, 2k+2(q-1)d\}$. A graph that has a (k,d)-even mean labeling is called a (k,d)-*even mean graph.* They proved that $P_m \oplus nK_1 (m \geq 3, n \geq 2)$ has a (k,d)-even mean labeling in the following cases: all (k,d) when m is even; all (k,d) when m is odd and n is odd; and m is odd, n is even and $k \geq d$.

Kalaimathy [1811] investigated conditions under which a mean labeling for a graph G will yield a (k,d)-even mean labeling for G and vice versa. He also gave conditions under which two graphs that have $(1,1)$-mean labelings can be joined by an single edge to obtain a new graph that has a $(1,1)$-even mean labeling. Gopi's Ph. D. thesis [1307] has a large number of results about mean, k-mean, k-odd mean, k-even mean, (k,d)-odd mean , and (k,d)-mean labelings.

In [4070] Vasuki, Nagarajan, and Arockiaraj introduced the notion of even vertex odd mean graphs as follows. A (p,q)-graph is said to have an *even vertex odd mean* labeling if there exists an injective function f from $V(G)$ to $\{0, 2, 4, \ldots, 2q-2, 2q\}$ such that the induced map $f^* : E(G)$ to $\{1, 3, 5, \ldots, 2q-1\}$ defined by $f^*(uv) = (f(u)+f(v))/2$ is a bijection. A graph that admits an even vertex odd mean labeling is called an *even vertex odd mean* graph. They proved that paths, C_{4n}, $K_{2,n}$, bistars $B_{m,n}$ for $n = m, m+1$, quadrilateral snakes Q_n, combs, $P_m \times P_n$, $C_{4m} \times P_n$, ladders, and dragons are even vertex odd mean graphs. Rajesh Kannan, Vikrama Prasad, Gopi [1888] proved the following graphs have an even vertex odd mean labeling: slanting ladders $SL_n (n \geq 3)$; double triangular snakes; alternative double triangular snakes; graphs obtained by starting with a tree G with at least 3 vertices and a mean labeling and a copy G' of G by joining each vertex of G to its corresponding vertex in G' with an edge; graphs obtained by starting with a path $v_1 v_2 \cdots v_n$ $(n \geq 4)$ and joining v_1 and v_3 to an isolated vertex; graphs obtained by starting with a path $v_1 v_2 \cdots v_n$ $(n \geq 4)$ and appending two edges to each of $v_2, v_3, \ldots, v_{n-1}$; and graphs obtained from a quadrilateral snake and appending an edge

at each vertex. The *H-graph of a path* P_n is the graph obtained from two copies of P_n with vertices $v_1, v_2, \ldots, v_n$ and $u_1, u_2, \ldots, u_n$ by joining the vertices $v_{n/2+1}$ and $u_{n/2+1}$ by an edge if n is odd and the vertices $v_{(n+1)/2}$ and $u_{(n/2}$ by an edge if n is even. Rajesh Kannan, Vikrama Prasad, and Gopi [1889] prove that the H-graph of P_n $(n \geq 3)$ and the graph $H \odot K_1$ have even vertex odd mean labelings where H is the H graph of P_n $(n \geq 3)$. In [1890] and [1891] Rajesh Kannan, Vikrama Prasad, and Gopi proved the following are even vertex odd mean graphs: graphs obtained by joining the centers of two stars $K_{1,m}$ and $K_{1,n}$ by a path P_t $(m, n, t \geq 2)$, graphs obtained by duplicating an edge of C_n $(n \geq 4)$, graphs obtained by joining each endpoint of P_3 to n isolated vertices, shadow graphs of stars, shadow graphs of bistars $B(n, n)$, mirror graphs of paths, and the graphs obtained taking two copies of $P_n \times P_2$ and joining each vertex of one with the matching vertex in the other with an edge. Kannan, Vikrama Prasad, and Gopi [1892] proved that the following graphs admit even vertex odd mean labelings: slanting ladders SL_n $(n \geq 3)$, twigs $\mathrm{TW}(n)$ $(n \geq 4$, double triangular snakes, alternative double quadrilateral snakes, and $Q_n \cdot K_1$. Prasad, Rejesh Kannan, and Gopi [3034] proved that $C_{4m} \odot K_{1,4n}$, $P_m \odot P_n$, and $\overline{K_2} + \overline{K_n}$ have even vertex odd mean labelings. In [3021] Prajapati and Raval proved that quadrilateral snakes, pentagonal snakes, and alternating quadrilateral snakes are vertex even and odd mean graphs. They also proved that even vertex odd mean graphs are even mean graphs. Jeyanthi, Ramya, and Selvi [1735] prove that T_P-trees (transformed trees), $T@P_n$, $T@2P_n$, and $\langle T \hat{o} K_{1,n} \rangle$ (where T is a T_P tree) are even vertex odd mean graphs.

Basher [556] investigated an even vertex labeling for the calendula graphs and introduced an arbitrary *calendula graph* as one in which if every edge from C_m is attached by an edge from arbitrary C_{n_i}, where n_i may vary for each $1 \leq i \leq m$. He proved that these graphs are also even vertex odd mean graphs. In [558] Basher proved several cycle related graphs are even vertex odd mean graphs. Basher and Kamran Siddiqui [560] proved that paths, cycles, combs, crowns, and planar grid super subdivisions are even vertex odd mean graphs.

For a graph $G(V, E)$ a bijection f from $V(G) \cup E(G)$ onto $\{1, 2, \ldots, |V(G)| + |V(E)|\}$ is said to be a *total mean labeling* if the values of $f^*(uv) = \lceil (f(u) + f(v) + f(uv))/3 \rceil$ taken over all edges are distinct. A graph G is said to be a *total mean labeling* graph if it admits a total mean labeling. Karuppasamy and Kaleeswari [1914] proved that $P_n, P_n^+, K_{1,n}, K_{2,n}, C_n, B_{m,n}$, triangular snakes, and alternate triangular snakes are total mean labeling graphs.

Murugan and Subramanian [2591] say a (p, q)-graph G has a *Skolem difference mean* labeling if there exists an injection f from the vertices of G to $\{1, 2, \ldots, p+q\}$ such that the induced map f^* defined on the edges of G by $f^*(uv) = (|f(u) - f(v)|)/2$ if $|f(u) - f(v)|$ is even and $f^*(uv) = (|f(u) - f(v)| + 1)/2$ if $|f(u) + f(v)|$ is odd is a bijection from edges of G to $\{1, 2, \ldots, q\}$. A graph that has a Skolem difference mean labeling is called a *Skolem difference mean* graph. They show that the graphs obtained by starting with two copies of P_n with vertices $v_1, v_2, \ldots, v_n$ and $u_1, u_2, \ldots, u_n$ and joining the vertices $v_{(n+1)/2}$ and $u_{(n+1)/2}$ if n is odd and the vertices $v_{n/2+1}$ and $u_{n/2}$ if n is even are Skolem difference mean. Parmar and Vaghela [2778] proved the following graphs have Skolem difference

mean labelings: brooms $B_{n,d}$ $(n \geq 4, d \geq 2)$ (the graph with n vertices obtained from P_d by appending $n-d$ edges at an endpoint), combs $P_n \odot K_1$ $(n \geq 2)$, $K_{1,m} \cup K_{1,n}$ $(m, n \geq 2)$, and $K_{1,3} * K_{1,n}$ $(n \geq 2)$ obtained from $K_{1,3}$ by attaching root of a star $K_{1,n}$ at each pendent vertex of $K_{1,3}$. Jeyanthi [1658] proved $\cup P_{n_i}$ $(n_i \geq 2)$; $\cup K_{1,n_i}$ $(n_i \geq 2)$; $C_n \cup P_m$ $(n \geq 3, m \geq 2)$; $K_{m,n} \cup (m-1)(n-1)K_2$; and $K_{1,1,n} \cup (n-1)K_2$ are Skolem difference mean graphs. She also proved that if G is a Skolem difference mean graph, then $G \cup nK_2$ is a Skolem difference mean graph.

Let $L_0, L_1, \ldots$ denote the sequence of Lucas numbers. In [2846] Ponmoni, Navaneetha Krishnan, and Nagarajan introduce the following graph labeling method. A graph G with p vertices and q edges is said to have a *Skolem difference Lucas mean* labeling if there is an injective function f from the vertices to $\{1, 2, \ldots, L_{p+q}\}$ such that when the edge uv is labeled with $|f(u) - f(v)|/2$ if $|f(u) - f(v)|$ is even, and $(|f(u) - f(v)| + 1)/2$ if $|f(u) - f(v)|$ is odd, then the resulting edge labels are distinct and belong to $\{L_1, L_2, \ldots L_q\}$. A graph that admits a Skolem difference Lucas mean labeling is called a *Skolem difference Lucas mean* graph. They proved the graphs obtained from $K_{1,m}$ by identifying the center of $K_{1,n}$ with the endpoint of each non-center vertex of $K_{1,m}$, bistars, $K_{1,m} \odot 2P_n$ and $\langle K_{1,n}^{(1)}, K_{1,n}^{(2)}, \ldots, K_{1,n}^{(m)} \rangle$ are Skolem difference Lucas mean graphs.

In [3256] Salisa and Peldrano introduced the notion of Lucas cordial labeling as follows. An injective function $f : V(G) \to \{L_1, L_2, \ldots, L_n\}$, where L_j is the jth Lucas number, is said to be a *Lucas cordial labeling* of G if the induced function $f^* : E(G)$ to $\{0, 1\}$ satifies the condition that the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1 In [3256] Salisa and Peldrano demonstrated the existence of Lucas cordial labelings for non-trivial cases of alternate triangular snakes, quadrilateral snakes, double alternate quadrilateral snakes, and the cycle quadrilateral snakes. Pedrano, and Salise [2818] demonstrated the existence of Lucas cordial labelings for alternate triangular snakes $A(T_n)$, $n \geq 2$, quadrilateral snakes Q_n, $n \geq 2$, double alternate quadrilateral snakes $DA(QS_n)$, $n \geq 2$, and cycle quadrilateral snakes CQ_n, $n \geq 3$. [3256] new [3256] new [2818] new

Selvi, Ramya, and Jeyanthi [3353] prove that $C_n @ P_n$ $(n \geq 3,\ m \geq 1)$, $K_n (n \leq 3)$, the shrub $St(n_1, n_2, \ldots, n_m)$, and the banana tree $Bt(n, n, \ldots, n)$ are Skolem difference mean graphs. They show that if G is a (p, q) graph with $q > p$ then G is not a Skolem difference mean graph and prove that K_n $(n \geq 4)$ is not a Skolem difference mean graph. A skolem difference mean labeling for which all the labels are odd is called an *extra Skolem difference mean* labeling. They also prove that the graph $T\langle K_{1,n_1} : K_{1,n_2} : \cdots : K_{1,n_m} \rangle$, obtained from the stars K_{1,n_1}, K_{1,n_2}, ..., K_{1,n_m} by joining the central vertex of K_{1,n_j} and $K_{1,n_{j+1}}$ to a new vertex w_j for $1 \leq j \leq m-1$ and the graph $T\langle K_{1,n_1} \circ K_{1,n_2} \circ \cdots \circ K_{1,n_m} \rangle$, obtained from K_{1,n_1}, K_{1,n_2}, ..., K_{1,n_m} by joining a leaf of $K_{1,n_{j+1}}$ to a new vertex w_j for $1 \leq j \leq m-1$ by an edge are extra Skolem difference mean graphs. Jeyanthi, Selvi, and Ramya [1759] proved that the union of any number of paths, any number of stars, $G \cup nK_2$ where G is an extra Skolem difference mean tree, $C_n \cup P_m$ $(n \geq 3, m \geq 2)$, $K_{m,n} \cup (m-1)(n-1)K_2$, and $K_{1,1,n} \cup (n-1)K_2$ have Skolem difference mean labelings. Vaghela and Parmar [3943] provided an extra Skolem difference mean labeling for combs, twigs of P_n, H graphs of P_n, graphs obtained from $K_{1,2}$ by attaching the root of $K_{1,n}$

at each pendent vertex of $K_{1,2}$, graphs obtained from $K_{1,3}$ by attaching the root of $K_{1,n}$ at each pendent vertex of $K_{1,3}$, and related graphs. Gross and Yellen [1341] proved that T_p-trees and caterpillars are extra Skolem difference mean graphs.

Let $G(V,E)$ be a graph with p vertices and q edges. Ramya, Kalaiyarasi, and Jeyanthi [3123] say G is a *Skolem odd difference mean* if there exists an injective function $f : V(G) \to \{0,1,2,3,\ldots,p+3q-3\}$ such that the induced map $f^* : E(G) \to \{1,3,5,\ldots,2q-1\}$ denoted by $f^*(uv) = \lceil |f(u)-f(v)|/2 \rceil$ is a bijection. A graph that admits a Skolem odd difference mean labeling is called a *odd difference mean* graph. They prove that P_n, C_n $(n \geq 4)$, $K_{1,n}$, $P_n \odot K_{1,n}$, coconut trees $T(n,m)$ obtained by identifying the central vertex of the star $K_{1,m}$ with a pendent vertex of P_n, $B_{m,n}$, caterpillars $S(n_1,n_2,\ldots,n_m)$, $P_m@P_n$ and $P_m@2P_n$ are Skolem odd difference mean graphs. ($P_m@P_n$ is obtained from P_m and m copies of P_n by identifying one pendent vertex of the i-th copy of P_n with the i-th vertex of P_m; $P_m@2P_n$ is defined analogously.) They establish that K_n, $n > 3$ and $K_{2,n}$ $(n \geq 3)$ are not Skolem odd difference mean graphs. They also prove that $K_{2,n}$ is a Skolem odd difference mean graph if $n \leq 2$. In [1685] Jeyanthi, Kalaiyarasi, Ramya, and Saratha Devi prove that bistars $B(m,n)$, mP_n, $mP_n \cup tP_s$, $mK_{1,n} \cup tK_{1,s}$ and the graph $\langle P_m \tilde{o} S_n \rangle$ obtained from P_m and m copies of $K_{1,n}$ by joining the central vertex of i^{th} copy of $K_{1,n}$ with i^{th} vertex of P_m by an edge admit Skolem odd difference mean labelings. They also prove that if $G(p,q)$ is a Skolem odd differences mean graph then $p \geq q$ and that wheels, umbrellas, books, and ladders are not Skolem odd difference mean graphs. They call a Skolem odd difference mean labeling a *Skolem even vertex odd difference mean* labeling if all the vertex labels are even. They prove that P_n, $K_{1,n}$, $P_n \odot K_1$, the coconut tree $T(n,m)$ obtained by identifying the central vertex of $K_{1,m}$ with a pendent vertex of a path P_n, $B(m,n)$, caterpillars $S(n_1,n_2,\ldots,n_m)$, $P_m@P_n$ are $P_m@2P_n$ are even vertex odd difference mean and C_n is not a Skolem even vertex odd difference mean graph. In [1813] Kalaiyarasi, Ramya, and Jeyanthi prove the following graphs have Skolem odd difference mean labelings: graphs obtained from a T_p tree with m vertices and m copies of $K_{1,n}$ by identifying the central vertex of ith copy of $K_{1,n}$, with ith vertex of T; graphs obtained by connecting an isolated vertex to central vertex of each of a number of stars; the banana trees obtained by connecting an isolated vertex to one leaf of each of any number of $K_{1,n}$; graphs obtained from $K_{1,n_1}, K_{1,n_2}, \ldots, K_{1,n_m}$ by joining the central vertices of K_{1,n_j} and $K_{1,n_{j+1}}$ to a new vertex w_j for $1 \leq j \leq m-1$; graphs obtained from $K_{1,n_1}, K_{1,n_2}, \ldots, K_{1,n_m}$ by joining a leaf of K_{1,n_j} and a leaf of $K_{1,n_{j+1}}$ to a new vertex w_j for $1 \leq j \leq m-1$.

In [1765] Jeyanthi, Selvi, and Ramya proved $C_m \cup C_n$ $(m,n \geq 3$ and $m \leq n)$ admit Skolem difference mean labelings. Kalaiyarasi and Ramya [1812] showed that $T \odot \overline{K_n}$, banana trees $B_t(\underbrace{n,n,\ldots,n}_{m \text{ times}})$, and regular bamboo trees $(S_{n,m})$ are Skolem odd difference mean graphs. [1765] new [1812] new

Murugan and Farzana [2590] introduced the concept of a 2-Skolem difference mean labeling of a graph as follows. A graph $G = (V,E)$ with p vertices and $2q$ edges is said to have a 2-Skolem difference mean labeling if it is possible to label the vertices $x \in V$ with distinct elements $f(x)$ from the set $\{1,2,\ldots,p\}$ in such a way that the edge $e = uv$ is labeled with $\frac{|f(u)-f(v)|}{2}$ if $|f(u)-f(v)|$ is even, and $\frac{|f(u)-f(v)|+1}{2}$ if $|f(u)-f(v)|$ is odd, [2590] new

and the resulting labels of the edges are distinct and are from $\{1, 1, 2, 2, 3, 3, \ldots, q, q\}$. A graph that admits 2-Skolem difference mean labeling is called a 2-*Skolem difference mean* graph. They proved the following graphs admit 2-Skolem difference mean labelings: $K_{1,2n}$ P_{2n+1} $B_{m,n}$ where m and n are of opposite parity, coconut trees graph $C_{m,n}$ where m and n are of opposite parity, caterpillars $S(X_1, X_2, \ldots, X_n)$, twigs obtained from the path by attaching exactly two pendent edges to each internal vertex of the path, and $P_n \odot K_1 - e_1$, $n \geq 2$, where e_1 is an edge incident with the vertex v_1 in G.

Lau, Jeyanthi, Ramya, and Kalaiyarasi [2099] say a (p, q)-graph $G(V, E)$ is a *Skolem even difference mean* if there exists an injective function $f : V(G) \to \{0, 1, 2, 3, \ldots, p + 3q - 1\}$ such that the induced map $f^* : E(G) \to \{2, 4, \ldots, 2q\}$ defined by $f^*(uv) = \lceil |f(u) - f(v)|/2 \rceil$ is a bijection. A graph that admits a Skolem even difference mean labeling is called a *even difference mean*. They prove: the disjoint union of paths of length at least 2 and $K_{2,n} \cup (n-1)K_2$ $(n \geq 2)$ are Skolem even vertex odd difference mean graphs; if G is a Skolem even vertex odd difference mean $(q + 1, q)$-graph, then $G \cup nK_2$, $G \cup P_n$, and $G \cup K_{1,n}$ are Skolem odd difference mean graphs; $C_m \cup P_n$ $(n \geq 2)$ is a Skolem odd difference mean graph for $m = 4$ and 6; the caterpillar $S(n_1, n_2, \ldots, n_m)$ is a Skolem even vertex even difference mean graph; $P_m@P_n$, mP_n, $K_{m,n} \cup (m-1)(n-1)K_2$ $(m, n \geq 2)$, $K_{1,n} \cup nK_2$, and $K_{1,1,n} \cup nK_2$ are Skolem even difference mean graphs; and if G is a Skolem even vertex even difference mean $(q+1, q)$-graph, then $G \cup nK_2$ is a Skolem even difference mean graph. They conclude with the following open problem: Establish that $G \cup nK_2$ where G is a (complete) multipartite graph is a Skolem even difference mean graph.

Kalaiyarasi, Ramya, and Jeyanthi [1684] say a graph $G(V, E)$ with p vertices and q edges has a *centered triangular mean* labeling if it is possible to label the vertices with distinct elements $f(x)$ from S, where S is a set of non-negative integers in such a way that for each edge $e = uv$, $f^*(e) = \lceil (f(u) + f(v))/2 \rceil$ and the resulting edge labels are the first q centered triangular numbers. A graph that admits a centered triangular mean labeling is called a *centered triangular mean* graph. They prove that P_n, $K_{1,n}$, bistars $B_{m,n}$, coconut trees, caterpillars $S(n_1, n_2, n_3, \ldots, n_m)$, $St(n_1, n_2, n_3, \ldots, n_m)$, banana trees $Bt(n, n, \ldots, n)$ and $P_m@P_n$ are centered triangular mean graphs.

Selvi, Ramya, and Jeyanthi [3354] define a *triangular difference mean* labeling of a graph $G(p, q)$ as an injection $f : V \longrightarrow Z^+$, such that when the edge labels are defined as $f^*(uv) = \lceil |f(u) - f(v)|/2 \rceil$ the values of the edges are the first q triangular numbers. A graph that admits a triangular difference mean labeling is called a *triangular difference mean* graph. They prove that the following are triangular difference mean graphs: P_n, $K_{1,n}$, $P_n \odot K_1$, bistars $B_{m,n}$, graphs obtained by joining the roots of different stars to the new vertex, trees $T(n, m)$ obtained by identifying a central vertex of a star with a pendent vertex of a path, the caterpillar $S(n_1, n_2, \ldots, n_m)$ and the graph $C_n@P_m$.

A graph $G(V, E)$ with p vertices and q edges is said to have *centered triangular difference mean* labeling if there is an injective mapping f from V to Z^+ such that the edge labels induced by $f^*(uv) = \lceil |f(u) - f(v)|/2 \rceil$ are the first q centered triangular numbers. A graph that admits a centered triangular difference mean labeling is called a *centered triangular difference mean* graph. Ramya, Selvi, and Jeyanthi [1762] prove that P_n, $K_{1,n}$, $C_n \odot K_1$, bistars $B_{m,n}$, C_n $(n > 4)$, coconut trees, caterpillars

$S(n_1, n_2,\ n_3, \ldots, n_m)$, $C_n@P_m$ $(n > 4)$ and $S_{m,n}$ are centered triangular difference mean graphs.

Gayathri and Tamilselvi [1232] say a (p, q)-graph G has a (k, d)-*super mean* labeling if there exists an injection f from the vertices of G to $\{k, k+d, \ldots, k+(p+q)d\}$ such that the induced map f^* defined on the edges of G by $f^*(uv) = \lceil (f(u)+f(v))/2 \rceil$ has the property that the vertex labels and the edge labels together are the integers from k to $k+(p+q)d$. When $d = 1$ a (k, d)-super mean labeling is called k-*super mean*. For $n \geq 2$ they prove the following graphs are k-super mean for all k: odd cycles; P_n; $C_m \cup P_n$; the one-point union of a cycle and the endpoint of P_n; the union of any two cycles excluding C_4; and triangular snakes. For $n \geq 2$ they prove the following graphs are (k, d)-super mean for all k and d: P_n; odd cycles; combs $P_n \odot K_1$; and bistars. In [1738] Jeyanthi, Ramya, and Thangavelu proved the following graphs have k-super mean labelings: C_{2n}, $C_{2n+1} \times P_m$, grids $P_m \times P_n$ with one arbitrary crossing edge in every square, and antiprisms on $2n$ vertices $(n > 4)$. (Recall an antiprism on $2n$ vertices has vertex set $\{x_{1,1}, \ldots, x_{1,n}, x_{2,1}, \ldots, x_{2,n}\}$ and edge set $\{x_{j,i}, x_{j,i+1}\} \cup \{x_{1,i}, x_{2,i}\} \cup \{x_{1,i}, x_{2,i-1}\}$ where subscripts are taken modulo n). Jeyanthi, Ramya, Thangavelu [1737] give k-super mean labelings for a variety of graphs. Jeyanthi, Ramya, Thangavelu, and Aditanar [1736] show how to construct k-super mean graphs from existing ones. For $n \geq 3$ Gayathri and Tamilselvi [1232] proved the following graphs are k-super edge mean for all k: paths; cycles; combs $P_n \odot K_1$; triangular snakes; crowns $C_n \odot K_1$; the one-point union of C_3 and an endpoint of P_n; and $P_n \odot K_2$.

In [3296] Sandhya, Somasundaram, and Ponraj call a graph with q edges a *harmonic mean* graph if there is an injective function f from the vertices of the graph to the integers from 1 to $q+1$ such that when each edge uv is labeled with $\lceil 2f(u)f(v)/(f(u)+f(v)) \rceil$ or $\lfloor 2f(u)f(v)/(f(u)+f(v)) \rfloor$ the edge labels are distinct. They prove the following graphs have such a labeling: paths, ladders, triangular snakes, quadrilateral snakes, $C_m \cup P_n$ $(n > 1)$; $C_m \cup C_n$; nK_3; $mK_3 \cup P_n$ $(n > 1)$; mC_4; $mC_4 \cup P_n$; $mK_3 \cup nC_4$; and $C_n \odot K_1$ (crowns). They also prove that wheels, prisms, and K_n $(n > 4)$ with an edge deleted are not harmonic mean graphs. In [3294] Sandhya, Somasundaram, and Ponraj investigated the harmonic mean labeling for a polygonal chain, square of the path and dragon and enumerate all harmonic mean graph of order at most 5. In [1592] Jayasekaran and David Raj prove that some disconnected graphs are harmonic mean graphs. In [3081] Raj, Jayasekaran, and Sandhya investigate some new families of harmonic mean graphs. Seoud and Salim [3401] provided upper bounds of the number of edges of graphs of given orders with harmonic mean labelings and showed that all graphs of order at most 9 have harmonic mean labelings using the floor function portion of the definition. Meena and Sivasakth [2477] prove that subdivision graphs of $P_n \odot K_1, P_n \odot \overline{K_2}$, H-graphs, $C_n \odot K_1, C_n \odot \overline{K_2}$, quadrilateral snakes, and triangular snakes are harmonic mean.

Sandhya, Somasundaram, and Ponraj [3295] proved that the following graphs have harmonic mean labelings: graphs obtained by duplicating an arbitrary vertex or an arbitrary edge of a cycle; graphs obtained by joining two copies of a fixed cycle by an edge; the one-point union of two copies of a fixed cycle; and the graphs obtained by starting with a path and replacing every other edge by a triangle or replacing every other edge by a quadrilateral.

In [3291] Sandhya, Somasundaram, and Ponraj investigated the harmonic mean labeling for polygonal chains and squares of the paths and dragons. They also enumerate all harmonic mean graph of order at most 5. Sandhya, Jayasekaran, and Raj [3285] showed that the following graphs are harmonic mean: graphs obtained by attaching a pendent vertex alternatively to the vertices of P_n, step ladders, and graphs obtained by attaching pendent edges to both sides of each vertex of P_n. Somasundaram, Ponraj, and Sandhya [3661] proved that paths, cycles, combs, and ladders are harmonic mean graphs. In [3723] Suganya, Maheswari, and Mary investigated the existence of harmonic mean labeling of degree splitting graph of some standard graphs.

Vaidya and Barasara [3949] proved that the following graphs have harmonic mean labelings: graphs obtained by the duplication of an arbitrary vertex or arbitrary edge of a path or a cycle; the graphs obtained by the duplication of an arbitrary vertex of a path or cycle by a new edge; and the graphs obtained by the duplication of an arbitrary edge of a path or cycle by a new vertex.

In [2654] Narasimhan and Sampathkumar called a graph with p vertices a *contra harmonic mean* graph if there is an injective function f from the vertices of the graph to the integers from 1 to p such that when each edge uv is labeled with $f(uv) = \lceil (f(u))^2 + (f(v))^2/(f(u) + f(v)) \rceil$ or $f(uv) = \lfloor (f(u))^2 + (f(v))^2/(f(u) + f(v)) \rfloor$ the edge labels are distinct. They prove the following graphs have such a labeling: paths, cycles, $C_m \cup P_n, C_m \cup C_n, nK_3, nK_3 \cup P_m$, and $nK_3 \cup C_m$. Gopi [1311] called a graph with q edges a *k-contra harmonic mean* graph if there is an bijective function f from the edges of the graph to the integers from $k-1$ to $k+q+1$ such that each edge uv is labeled with $f(uv) = \lceil (f(u))^2+(f(v))^2/(f(u)+f(v)) \rceil$ or $f(uv) = \lfloor (f(u))^2+(f(v))^2/(f(u)+f(v)) \rfloor$. He proves that triangular snakes, double triangular snakes, quadrilateral snakes, and double quadrilateral snakes have k-contra harmonic mean labelings.

Gopi and Suba [1318] say a graph G with p vertices and q edges is a *super Lehmer-3 mean graph* if there is an injective function f from the vertices of G to $\{1, 2, \ldots, q+1\}$ such that for each edge uv the induced function $f^*(uv) = \lfloor (f(u)^3 + f(v)^3)/(f(u)^2 + f(v)^2) \rfloor$ or $f^*(uv) = \lceil (f(u)^3 + f(v)^3)/(f(u)^2 + f(v^2)) \rceil$ yields that the set of vertex labels and edge labels is $\{1, 2, \ldots, p\}$. They prove that $P_m \odot K_{1,n}$ and the graph obtained by identifying each endpoint of a path with an endpoint of the star $K_{1,n}$ have a super Lehmer-3 labeling. In [1315] Gopi and Nirmala provide Lehmer-3 mean labelings for $P_m \odot C_n$ $(m, n \geq 3)$ and $P_m \odot K_1 \odot C_n$ $(m, n \geq 3)$. In [3662] Somasundaram, Sandhya, and Pavithra proved that paths, combs, ladders, and crowns are super Lehmer-3 mean graphs. In [3663] Somasundaram, Sandhya, and Pavithra investigated the existence of Lehmer-3 mean labelings of graphs of the form $C_m \odot K_1, C_m \odot K_{1,2}, P_m \odot K_n$, and $P_m \odot K_{1,n}$. In [3664] they investigated the existence of Lehmer-3 mean labelings of graphs of the form $mC_n, mC_n \cup P_k, mC_n \cup C_k, mC_n \cup (P_k \odot K_1), mC_m \cup (P_k \odot K_{1,2}), mC_m \cup (P_k \odot K_{1,3})$ and $mC_m \cup (P_k \odot K_3)$. Somasundaram, Sandhya, and Pavithra [3666] proved that paths, cycles, combs, crowns, triangular snakes, and quadrilateral snakes admit Lehmer-3 mean labelings. Paramaguru, and Manirathnam [2764] investigated the existence of Lehmer-3 mean labeling of bicyclic-related graphs. A *dragon* is obtained by joining an end point of the path to a vertex of a cycle. In [1567] Jaisankar, Manivel, and Keerthana investigated

the existence of Lehmer-3 mean labelings of various dragon-related graphs. In [3037] Praveen Prabhu and Shanmuga Sundaram proved that the standard theta graphs and their structural variations obtained by subdivisions and attachments of pendent vertices admit super Lehmer-3 mean labelings.

In [1319] Gopi and Prakash investigated the k-super Lehmer-3 meanness of triangular snakes, double triangular snakes, alternative triangular snakes, quadrilateral snakes, double quadrilateral snakes, and alternative quadrilateral snakes. In [3666] Somasundaram, Sandhya, and Pavithra proved that the following graphs admit k-super Lehmer-3 mean labelings: paths, combs, nP_m, $P_n \cup (P_m \odot K_1)$, graphs obtained by attaching an end vertex of P_n to a vertex of C_3, and the union of graphs obtained by attaching an end vertex of P_n to a vertex of C_3 with a comb.

In [3479] Shalini and Meena introduced the notion of Lehmer-4 mean labelings as follows. A graph $G(V, E)$ with p vertices and q edges is called a *Lehmer-4 mean graph*, if there is an injection from V to $\{2, 4, 6, \ldots, 2p\}$ and the induced function $f^*(uv) = \lfloor (f(u)^4 + f(v)^4)/(f(u)^2 + f(v)^2) \rfloor$ or $f^*(uv) = \lceil (f(u)^4 + f(v)^4)/(f(u)^2 + f(v^2)) \rceil$ in an injection. In this case, f is called a *Lehmer-4 mean labeling* mean of G. They proved paths, combs, $P_n \odot K_{1,2}$ and $P_n \odot K_{1,3}$ are Lehmer-4 mean graphs.

An *F-geometric mean* labeling of a graph G with q edges, is an injective function from the vertex set of G to $\{1, 2, \ldots, q+1\}$ such that the edge labels obtained from the floor function of geometric mean of the vertex labels of the end vertices of each edge, are all distinct and the set of edge labels is $\{1, 2, \ldots, q\}$. Durai Baskar, Arockiaraj, and Rajendran [983] proved that the following graphs are F-geometric mean: graphs obtained by identifying a vertex of consecutive cycles (not necessarily of the same length) in a particular way; graphs obtained by identifying an edge of consecutive cycles (not necessarily of the same length) in a particular way; graphs obtained by joining consecutive cycles (not necessarily of the same length) by paths (not necessarily of the same length) in a particular way; $C_n \odot K_1$; $P_n \odot K_1$; $L_n \odot K_1$; $G \odot K_1$ where G is the graph obtained by joining two copies of P_n by an edge in a particular way; graphs obtained by appending two edges at each vertex of graphs obtained by joining two copies of P_n by an edge in a particular way; graphs obtained from C_n by appending two edges at each vertex of C_n; graphs obtained from ladders by appending two edges at each vertex of the ladders; graphs obtained from P_n by appending an end point of the star S_2 to each vertex of P_n; and graphs obtained from P_n by appending an end point of the star S_3 to each vertex of P_n. Sandhya and Somasundaram [3290] proved that double quadrilateral snakes and alternate double quadrilateral snake graphs are geometric mean graphs.

Durai Basker and Arockiaraj [982] study the F-geometric meanness of cycles, stars, complete graphs, combs, ladders, triangular ladders, middle graphs of paths, graphs obtained from duplicating arbitrary vertex by a vertex as well as arbitrary edge by an edge in cycles, and subdivisions of combs and stars.

A *C-geometric mean* labeling of a graph G with q edges, is an injective function from the vertex set of G to $\{1, 2, 3, \ldots, q+1\}$ such that the edge labels obtained from the ceiling function of the geometric mean of the vertex labels of the end vertices of each edge are all distinct and the set of edge labels is $\{2, 3, 4, \ldots, q+1\}$. A graph is said to

be a *C-geometric mean graph* if it admits a C-geometric mean labeling. In [985] Durai Baskar and Arockiaraj study the C-geometric meanness of some cycle related graphs such as cycle, union of a path and a cycle, unions of two cycles, the graphs $C_3 \times P_n$, corona of cycle, the graphs $P_{a,b}, P_b^a$ and some chain graphs.

A geometric mean labeling f of $G(V, E)$ is called a *super geometric mean* labeling if $f(V) \cup f(E) = \{1, 2, \ldots, |V| + |E|\}$. Sandhya, Merly, and Shiny [3287] [3288] prove that the subdivision graphs of the following graphs have super geometric mean labelings: alternate quadrilateral snakes, double quadrilateral snakes, alternate double quadrilateral snakes, triple quadrilateral snakes, and subdivisions of alternate triple quadrilateral snakes. In [3289] they prove that the following graphs have super geometric mean labelings: triangular ladders, triangular snakes, alternate triangular snakes, quadrilateral snakes, and alternate quadrilateral snakes. Hemalatha and Selvi [1432] prove that following graphs have super geometric mean labelings: flags, kayak paddles, dumbbells, polygonal snakes, and graphs obtained by connecting any number of copies of C_n where each joined to the next with an edge.

In 2026 Nirmala Irudayam and Sherin [2682] introduced the notion of *generalized geometric mean labeling* (GGML) as follows. For a simple graph $G = (V, E)$ and a real number $k \in (0, 1)$ a k-GML consists of an injective vertex labelling $f : V \to \mathbb{N}$ with the induced edge labeling $g_k(uv) = \left\lfloor (f(u)f(v))^k \right\rfloor$, $0 < k < 1$, such that all edge labels are distinct. This notion is a continuous generalization of classical geometric mean labeling, which is the special case that $k = 1/2$. They proved that paths, cycles, stars, and m–shadow graphs admit k–GML for all $0 < k < 1$. [2682] new

In 2017 Hemalatha, Mohanaselvi, and Amuthavalli [1434] defined a *radio geometric mean* labeling of a graph G as a mapping from the vertex set to the set of natural numbers such that for two distinct vertices u and v of G, $d(u, v) + \lceil f(u)f(v) \rceil \geq 1 + diam(G)$. The *radio geometric mean number* of f is the maximum number assigned to any vertex of G. The *radio geometric mean number* of G is the minimum value taken over all radio geometric mean labeling f of G. A *radio antipodal geometric mean* labeling of a graph G is a mapping f from the vertex set of G to the set of natural numbers such that for two distinct vertices u and v of G, $d(u, v) + \lceil \sqrt{f(u)f(v)} \rceil \geq diam(G)$. If $d(u, v) = diam(G)$, the vertices u and v can be given the same label and if $d(u, v) \neq diam(G)$, the vertices u and v are assigned different labels. The *radio antipodal geometric mean number* of f is the maximum number assigned to any vertex of G. The *radio antipodal geometric mean number* of G is the minimum value taken over all radio geometric mean labeling f of G. Hemalatha, Mohanaselvi, and Amuthavalli [1434] provided the radio geometric mean numbers for stars, bistars, complete k-ary trees with 3 levels, and the graph lotus inside a circle. In [1433] Hemalatha and Amuthavalli determined the radio geometric mean number of splitting graphs of stars and bistars. Hemalatha and Mohanaselvi [1435] found the radio geometric mean numbers of the subdivision graphs of complete graphs, complete bipartite graphs, books, and windmills. In [1274] Giridaran, Jose, and Jeony provided upper bounds for the radio antipodal geometric mean number of ladders, triangular ladders, circular ladders ($C_n \times K_2$), and pagoda graphs (graphs obtained by joining two adjacent endpoints of a ladder to new vertex.

Arockiaraj and Meena Kumari introduced the F-face magic mean labeling of graphs in [983]. This motivated Meena Kumari and Arockiaraj [2471] to introduce the (1,0,0)-F-face magic mean labeling of graphs as follows. A bijection ϕ from $V(G)$ to $\{1, 2, \ldots, |V(G)|\}$ is called a (1,0,0)-F-*face magic mean* labeling of G if the induced face labeling $\phi^*(f_i) = \lfloor$(sum of the labels of the vertices in the boundary of f_i)/deg$(f_i)\rfloor$ is a constant for each face f_i, including the exterior face of G, where deg f_i is the number of edges that bound the face. A graph that admits an (1,0,0)-F-face magic mean labeling is called (1,0,0)-F-*face magic mean*. In [2471] Arockiaraj and Meena Kumar showed that $P_n + K_1$ $(n \geq 2)$, cycles with certain cords, $C_m \times P_n$ where m and n are even, and graphs obtained by duplicating every edge of a cycle by a vertex admit (1,0,0)-F-face magic mean labelings.

In [3792] Sundaram, Ponraj, and Somasundaram introduced a new labeling parameter called the *mean number* of a graph. Let f be a function from the vertices of a graph to the set $\{0, 1, 2, \ldots, n\}$ such that the label of any edge uv is $(f(u) + f(v))/2$ if $f(u) + f(v)$ is even and $(f(u) + f(v) + 1)/2$ if $f(u) + f(v)$ is odd. The smallest integer n for which the edge labels are distinct is called the *mean number* of a graph G and is denoted by $m(G)$. They proved that for a graph G with p vertices $m(tK_{1,n}) \leq t(n + 1) + n - 4$; $m(G) \leq 2^{p-1} - 1$; $m(K_{1,n}) = 2n - 3$ if $n > 3$; $m(B(p, n)) = 2p - 1$ if $p > n + 2$ where $B(p, n)$ is a bistar; $m(kT) = kp - 1$ for a mean tree T, $m(W_n) \leq 3n - 1$, and $m(C_3^{(t)}) \leq 4t - 1$.

Let f be a function from $V(G)$ to $\{0, 1, 2\}$. For each edge uv of G, assign the label $\lceil (f(u) + f(v))/2 \rceil$. Ponraj, Sivakumar, and Sundaram [2984] say that f is a *mean cordial labeling* of G if $|v_{f(i)} - v_{f(j)}| \leq 1$ for i and j in $\{0, 1, 2\}$ where $v_{f(x)}$ and $e_{f(x)}$ denote the number of vertices and edges labeled with x, respectively. A graph with a mean cordial labeling is called a *mean cordial graph*. Observe that if the range set of f is restricted to $\{0, 1\}$, a mean cordial labeling coincides with that of a product cordial labeling. Ponraj, Sivakumar, and Sundaram [2984] prove the following: every graph is a subgraph of a connected mean cordial graph; $K_{1,n}$ is mean cordial if and only $n \leq 2$; C_n is mean cordial if and only $n \equiv 1, 2 \pmod 3$; K_n is mean cordial if and only $n \leq 2$; W_n is not mean cordial for all $n \geq 3$; the subdivision graph of $K_{1,n}$ is mean cordial; the comb $P_n \odot K_1$ is mean cordial; $P_n \odot 2K_1$ is mean cordial; and $K_{2,n}$ is a mean cordial if and only $n \leq 2$. Seoud and Salim [3401] provided upper bounds of the number of edges of graphs of given orders with mean cordial labelings and proved that $P_{2t} \times P_2$ is mean cordial if and only if $t \equiv 2 \bmod 3$ and $C_n \odot K_1$ is mean cordial if and only if $n \equiv 1$ or $2 \bmod 3$.

In [2972] Ponraj and Sivakumar proved the following graphs are mean cordial: mG where $m \equiv 0 \pmod 3$; $C_m \cup P_n$; $P_m \cup P_n$; $K_{1,n} \cup P_m$; $S(P_n \odot K_1)$; $S(P_n \odot 2K_1)$; P_n^2 if and only if $n \equiv 1 \pmod 3$ and $n \geq 7$; and the triangular snake T_n $(n > 1)$ if and only if $n \equiv 0 \pmod 3$. They also proved that if G is mean cordial then mG, $m \equiv 1 \pmod 3$ is mean cordial. Deshmukh and Shaikh [931] prove the graph $\langle K_{1,n} : 2\rangle$ and the path union of n copies of $K_{1,m}$ are mean cordial graphs.

In [2943] Ponraj and Sathish Narayanan proved double triangular snakes $D(T_n)$ are mean cordial if and only if $n > 3$ and obtained partial results on mean cordial labelings of alternate triangular snakes, double alternate triangular snakes. Shendra shainy, Vinothkumar, and Balaji [3503] proved that for $\beta_1 < \beta_2 < \beta_3$, the three star graph $K_{1,\beta_1} \wedge K_{1,\beta_2} \wedge K_{1,\beta_3}$ is a mean cordial graph if and only if $|\beta_2 - \beta_3| \leq 3\beta_1 + 7$. Further,

in [2400] Maheshwari, Vinothkumar and Balaji proved that for $\beta_1 = \beta_2 < \beta_3$ the three star graph $K_{1,\beta_1} \wedge K_{1,\beta_2} \wedge K_{1,\beta_3}$ is a mean cordial graph if and only if $|\beta_2 - \beta_3| \leq 3\beta_1 + 6$.

In [2960] Ponraj, Sathish Narayanan, and Ramasamy introduced the notion of total mean cordial labeling. A *total mean cordial* labeling of a graph $G(V, E)$ is a function $f : V(G) \to \{0, 1, 2\}$ such that when each edge xy is assigned the label $\lceil (f(x) + f(y))/2 \rceil$ we have $|ev_f(i) - ev_f(j)| \leq 1$, $i, j \in \{0, 1, 2\}$, where $ev_f(x)$ denotes the total number of vertices and edges labeled with x. A graph with a total mean cordial labeling is called *total mean cordial*. In [2960], [2961], and [2962], Ponraj, Sathish Narayanan, and Ramasamy determined the total mean cordiality of the following graphs: P_n; C_n; $K_{1,n}$; W_n; $K_2 + mK_1$; combs $P_n \odot K_1$; double combs $P_n \odot 2K_1$; crowns; flowers; lotuses inside a circle; bistars; quadrilateral snakes; $K_{2,n}$; olive trees; $S(P_n \odot K_1)$; $S(K_{1,n})$ ($S(G)$ denotes the subdivision of G); triangular snakes; P_n^2; fans F_n; umbrellas; butterflies; and dumbbells. In [2942], [2944], and [2945], Ponraj and Sathish Narayanan determined the total mean cordiality of $K_n^c + 2K_2$; prisms; gears; helms; $P_1 \cup P_2 \cup \cdots \cup P_n$; $L_n \odot K_1$; $S(W_n)$; $S(P_n \odot 2K_1)$; and graphs obtained by subdividing each step of a ladder exactly once.

Let G be a (p, q)-graph. Ponraj and Sathish Narayanan [2947] and [2948] proved the following. If G satisfies any one of the following three conditions then $G \odot 2K$ is total mean cordial: (1) G is a tree, (2) G is a unicycle, (3) $q = p + 1$. If G satisfies any one of the following three conditions then the shadow graph of G is total mean cordial: (1) G is a tree, (2) G is a unicycle, (3) $q = p + 1$. They also proved that the following are total mean cordial graphs: $C_n \odot K_2$, $C_n^{(2)}$, dragons, splitting graphs of stars, splitting graphs of combs, books, ladders, $P_n \odot K_2$ if and only if $n \neq 1$, and $G \cup P_n$ ($n \neq 3$).

In [2323] Lourdusamy and Joy Beaula define the notion of S_3 (all permutations of $\{1, 2, 3\}$) mean cordial graphs as follows. Let $G(V, E)$ be a graph and $g : V(G) \to S_3$ be a function. For each edge xy assign the label 1 if $\lceil (|g(x)| + |g(y)|)/2 \rceil$ is odd, and 0 otherwise. Such a g is said to be a *group S_3 mean cordial* labeling if the number of vertices labeled with i and j differ by at most 1 where i, j are the elements in S_3 and the number of edges labeled with 0 and 1 differ by at most 1. A graph with a group S_3 mean cordial labeling is called a *group S_3 mean cordial* graph. They prove that $K_{2,n}$, subdivisions of fans, double fans, wheels, and helms are S_3 mean cordial graphs.

Ponraj, Sathish Narayanan, and Kala introduced the concept of radio mean labeling in [2958]. A *radio mean* labeling of a connected graph G is a one-to-one map f from $V(G)$ to the set of natural numbers such that for each pair of distinct vertices u and v of G, $d(u, v) + \left\lceil \frac{f(u)+f(v)}{2} \right\rceil \geq 1 + \text{diam } (G)$. The *radio mean number* of f, $rmn(f)$, is the maximum number assigned to any vertex of G. The radio mean number of G, $rmn(G)$, is the minimum value of $rmn(f)$ taken over all radio mean labelings f of G. They proved $rmn(G) \geq |V(G)|$; if G is a (p, q)-graph with diameter $d \geq 2$, then $rmn(G) \leq p + d - 2$; and if G is a (p, q)-connected graph with diameter 2 or 3, then $rmn(G) = p$. They also determine the radio mean number of K_n, $K_{m,n}$, sunflowers, helms, gears, lotuses inside a circle, and graphs obtained by identifying any two vertices of two wheels of the same size, Aasi, Asif, Iqbal, and Ibrahim [1] determine the radio number and the radio mean number for the lexicographic product of a path with a path, a path with a cycle, and a cycle with

a cycle. Moreover, they present computer programs for finding such radio labelings of these families of graphs. Sunitha, Raj, and Subramanian [3797] investigated radio mean labeling of path and cycle related graphs. Lavanya, Dhanyashree, and Meera [2130] determined the radio mean number of subdivision graphs of complete graphs, Mongolian tents, subdivision of friendship graphs, and diamond graphs. If the minimum value of the radio mean number taken over all radio mean labelings of G is $|V(G)|$, the graph is said to be *radio mean graceful*. They proved that the graphs they studied are radio mean graceful.

Augustine and Roy [94] provided upper bounds of radio numbers and radio mean numbers for honeycomb and honeycomb torus networks. [94] new

In [2957] and [2959] Ponraj, Sathish Narayanan, and Kala determine the radio mean numbers of $S(K_{m,n})$ $(m > 1, n > 1)$; $K_{m,n} \odot P_t$; $C_6^{(t)}$; $W_n \odot P_m$; graphs obtained by joining the rim vertices of the two wheels with an edge; and graphs obtained from a wheel by subdividing each spoke by a vertex. In [2963] Ponraj, Sathish Narayanan, and Kala give the radio mean number of graphs with diameter three, lotuses inside a circle, helms, and sunflower graphs.

In [2969] and [2949] Ponraj and Sathish Narayanan give the radio mean number of the following graphs: subdivisions of stars, subdivisions of wheels, subdivisions of $K_2 + mK_1$, subdivisions of bistars, jelly fish, subdivisions of jelly fish, books with pentagonal pages, graphs obtained by taking m disjoint copies of $K_{1,n}$ and joining a new vertex to the centers of the m copies of $K_{1,n}$.

A *radio mean D-distance* labeling of a connected graph G is an injective map f from $V(G)$ to the natural numbers such that for two distinct vertices u and v of G, $d^D(u,v) + \lceil (f(u) + f(v))/2 \rceil \geq 1 + diam^D(G)$, where $d^D(u,v)$ denotes the distance D between u and v and $diam^D(G)$ denotes the D-diameter of G. The *radio mean D-distance number* of f, $rmn^D(f)$, is the maximum label assigned to any vertex of G. The *radio mean D-distance number* of G, $rmn^D(G)$, is the minimum value of $rmn^D(f)$ taken over all radio mean D-distance labeling f of G. Nicholas and Bosco [2670] determined the radio mean D-distance number of cycles, wheels, gears, helms, fans, and friendship graphs.

In [2946] Ponraj and Sathish Narayanan proved that the following graphs are not mean cordial: $K_2 + \overline{K}_m$; $\overline{K}_n + 2K_2$; $P_n \times K_2$; flower graphs; sunflower graphs; $C_n \odot K_2$. Also they proved the following: the Mongolian tent $MT_{m,n}$ is mean cordial if and only if $m \equiv 0 \pmod 3$ or $n \equiv 0 \pmod 3$ ($MT_{m,n}$ is the graph obtained from $P_m \times P_n$, n odd, by adding one extra vertex above the grid and joining every other vertex of the top row of $P_m \times P_n$ to the new vertex); the book B_m is mean cordial if and only if $m = 1$; books with n pentagonal pages are mean cordial if and only if $n \equiv 1 \pmod 3$; $P_n \odot K_2$ is mean cordial if and only if $n \equiv 0 \pmod 3$; quadrilateral snakes are mean cordial; alternate quadrilateral snakes $A(Q_n)$ are mean cordial if and only if the square starts from second vertex of the path P_n, ends with $(n-1)^{th}$ vertex and $n \equiv 0, 2 \pmod 3$, or the square starts from first vertex, ends with n^{th} vertex and $n \equiv 0, 2 \pmod 3$, or the square starts from second vertex, ends with n^{th} vertex and $n \equiv 0, 1 \pmod 3$.

Let G be a graph and let $f : V(G) \to \{0, 1, \ldots, k-1\}$ $k > 1$. For each edge uv, assign the label $f(uv) = \lceil \frac{f(u)+f(v)}{2} \rceil$. Ponraj, Subbulakshmi, and Somasundaram

[2987] say that f is a *k-total mean cordial* labeling of G if $|t_{mf}(i) - t_{mf}(j)| \leq 1$, for all $i, j \in \{0, 1, \ldots, k-1\}$, where $t_{mf}(x)$ denotes the total number of vertices and edges labeled with $x \in \{0, 1, \ldots, k-1\}$. A graph with admit a k-total mean cordial labeling is called *k-total mean cordial graph*. Ponraj, Subbulakshmi, and Somasundaram [2987] investigated the 4-total mean cordial labeling behavior of fans, wheels, jelly fish, jewel graphs, ladders, and triangular snakes.

A *balanced mean cordial* labeling is a mean cordial labeling f with the property that the number of vertices labeled with i is the same as the number labeled with j for $i, j \in \{0, 1, 2\}$ and likewise for the edges. Kaneria, Khoda, and Karavadiya [1839] prove: the path union of n copies of a graph G is a mean cordial when $n \equiv 0 \pmod 3$; if G is balanced mean cordial, then $P_n \times G$ and $C_n \times G$ are balanced mean cordial; and if $f : V(G) \longrightarrow \{0, 1, 2\}$ is a balanced mean cordial labeling for G, then G^* is also a balanced mean cordial graph.

In [1692] Jeyanthi and Maheswari define a *one modulo three mean* labeling of a graph G with q edges as an injective function ϕ from the vertices of G to $\{a \mid 0 \leq a \leq 3q-2$ where $a \equiv 0 \pmod 3$ or $a \equiv 1 \pmod 3\}$ and ϕ induces a bijection ϕ^* from the edges of G to $\{a \mid 1 \leq a \leq 3q-2$ where $a \equiv 1 \pmod 3\}$ given by $\phi^*(uv) = \lceil(\phi(u) + \phi(v))/2\rceil$. They proved that P_{2n}, combs, bistars $B_{n,n}$, T_p-trees with an even number of vertices, C_{4n+1}, ladders, $K_{1,2n} \times P_2$ are one modulo three mean graphs. They also proved that bistars $B_{m,n}$ $(m \neq n), K_{1,n}$ $(n > 3)$, and K_n, $(n > 3)$ are not one modulo three mean graphs. In [1699] Jeyanthi, Maheswari, and Pandiaraj [1699] proved that $DA(Q_n), DA(Q_2) \odot nK_1, DA(Q_m) \odot nK_1, DA(T_2) \odot nK_1, DA(T_m) \odot nK_1, \overline{S}(DA(T_n)), \overline{S}(DA(Q_n))$, and mP_n are one modulo three mean graphs.

Jeyanthi, Maheswari, and Pandiaraj [1698] prove that following graphs have one modulo three mean labelings: books $K_{1,2n} \times P_2$; splitting graphs $S'(P_{2n})$; vertex duplication graphs $D(G, v')$; edge duplication graphs $D(G, e')$; nth alternate quadrilateral snake graphs $NA(Q_m)$; graphs obtained by joining the endpoints of paths P_{4m} to n isolated vertices; and extended jewel graphs EJ_n with vertex set $\{u, v, x, y, w, z, u_i : 1 \leq i \leq n\}$ and edge set $\{uv, ux, xy, yz, vw, wz, vu_i, zu_i : 1 \leq i \leq n\}$. For graphs G_1 and G_2, $G_1 \widehat{\circ} G_2$ is the graph obtained from G_1 and $|V(G_1)|$ copies of G_2 by joining a vertex of ith copy of G_2 with the ith vertex of G_1 by an edge. Jeyanthi, Maheswari, and Pandiaraj [1701] prove that the graphs $T \odot \overline{K_n}$, $T \widehat{\circ} K_{1,n}$, $T \widehat{\circ} P_n$, and $T \widehat{\circ} 2P_n$ are one modulo three mean graphs. Sudarvizhi and Balasangu [3712] provide one modulo three mean labelings for triangular books, duplication subdivisions of the central edge of bistars, and slanting ladders (graphs obtained from two paths $u_1, u_2, u_3, \ldots, u_n$ and $v_1, v_2, v_3, \ldots, v_n$ by joining each u_i with v_{i+1} for $1 \leq i \leq n_1$).

In [1224] and [1230] Gayathri and Prakash proved the following graphs admit one modulo three mean graphs: the mirror graph of P_n, $\mathrm{Spl}(P_n)$, $P_m \times P_n, (P_m \times P_2) \odot K_{1,n}$, bistars, $C_n @ P_m, C_n \odot P_m$, and $H \odot \overline{K_m}$, where H is the graph is obtained from two copies of P_n with vertices $u_1, u_2, \ldots, u_n$ and $v_1, v_2, \ldots, v_n$ by joining $u_{(n+1)/2}$ to $v_{(n+1)/2}$ with an edge when $n > 1$ is odd, and $u_{n/2}$ to $v_{n/2+1}$ with an edge when n is even. In [1227] Gayathri and Prakash investigated the existence of one modulo three mean labelings of disconnected graphs. In [1225] they provided a necessary condition for a graph to admit a one modulo

three mean labeling. In [1226] Gayathri and Prakash provided one modulo three mean labelings for various trees. In [1228] and [1229] they proved the following graphs admit are k-one modulo three mean labelings: $P_n, K_{1,n}$, combs, $P_m \odot K_{1,n}, B_{m,n}, L_n = P_n \times P_2, K_{1,n} \times P_2$, $A(Q_n), D(Q_n), A(D(Q_n)), A(D(Q_n)) \odot \overline{K_m}, A(Q_n) \odot \overline{K_m}$, and the graphs obtained by joining the end points of P_m to n isolated points.

A graph G is said to be *one modulo three root square mean* graph if there is an injective function ϕ from the vertex set of G to the set $\{0, 1, 3, \ldots, 3q-2, 3q\}$ where q is the number of edges of G and ϕ induces a bijection ϕ^* from the edge set of G to $\{1, 4, \ldots, 3q-2\}$ given by $\phi^*(uv) = \left\lceil \sqrt{\frac{[\phi(u)]^2+[\phi(v)]^2}{2}} \right\rceil$ or $\left\lfloor \sqrt{\frac{[\phi(u)]^2+[\phi(v)]^2}{2}} \right\rfloor$. The function ϕ is called a *one modulo three root square mean* labeling of G. In [1596] Jayasekaran and Jaslin Melbha investigated some path related graphs that have one modulo three root square mean labelings.

Somasundaram, Vidhyarani, and Ponraj [3668] introduced the concept of a *geometric mean* labeling of a graph G with p vertices and q edges as an injective function $f : V(G) \to \{1, 2, \ldots, q+1\}$ such that the induced edge labeling $f^* : E(G) \to \{1, 2, \ldots, q\}$ defined as $f^*(uv) = \left\lceil \sqrt{f(u)f(v)} \right\rceil$ or $\left\lfloor \sqrt{f(u)f(v)} \right\rfloor$ is bijective. Among their results are: paths, cycles, combs, ladders are geometric mean graphs and K_n $(n > 4)$ and $K_{1,n}$ $(n > 5)$ are not geometric mean graphs. Somasundaram, Vidhyarani, and Sandhya [3669] proved $C_m \cup P_n$, $C_m \cup C_n$, nK_3, $nK_3 \cup P_n$, $nK_3 \cup C_m$, P_n^2, and crowns are geometric mean graphs. Vaidya and Barasara [3952] investigated geometric mean labelings in context of duplication of graph elements in cycle C_n and path P_n. Durai Baskar, Arockiyaraj, and Rajendran investigate the geometric meanness of some graphs obtained from paths.

In Jeyanthi, Maheswari, and Pandiaraj [1700] define a graph G to be a one modulo three geometric mean graph if there is an injective function ϕ from the vertex set of G to the set $\{a | 1 \leq a \leq 3q-2$ and either $a \equiv 0 \pmod 3$ or $a \equiv 1 \pmod 3\}$ where q is the number of edges of G and ϕ induces a bijection ϕ^* from the edge set of G to $\{a | 1 \leq a \leq 3q-2$ and $a \equiv 1 \pmod 3\}$ given by $\phi^*(uv) = \left\lceil \sqrt{\phi(u)\phi(v)} \right\rceil$ or $\left\lfloor \sqrt{\phi(u)\phi(v)} \right\rfloor$ the function ϕ is called one modulo three geometric mean labeling of G. They proved paths, cycles with length at least 5, ladders, $P_n \odot K_1, P_n \odot P_2, P_n \odot \overline{P_2}$, subdivision graphs $S(P_n \odot K_1)$, and subdivision graphs $S(P_n \odot K_2)$ are one modulo three geometric graphs. They also prove that $K_{1,n}$ $(n \geq 3)$ and graphs in which every edge lies on a triangle are not one modulo three geometric mean graph.

In [1766] Jeyanthi, Selvi and Ramya introduced a new graph labeling as follows. A graph $G(p, q)$ is said to be a *one modulo* N-*difference mean* graph if there is an injection f from the vertex set of G to the set $\{0 \leq a \leq 2(q-1)N + 1\}$ and either $a = 0 \pmod N$ or $a = 1 \pmod N$, where N is a positive integer and f induces a bijection f^* from the edge set of G to $\{1 \leq a \leq (q-1)N + 1\}$ and $a = 1 \pmod N$ given by $f^*(uv) = \lceil |f(u) - f(v)|/2 \rceil$. Such a function f is called a *one modulo* N-*difference mean labeling* of G. They establish that $B_{m,n}$, $S_{m,n}$, $P_n@P_m$, $B(l, m, n)$, $T(n, m)$, shrubs, caterpillars, and $K_{1,n}$ are one modulo N-difference mean graph. In addition, they show that C_3 is not a one modulo N-difference mean graph.

Jeyanthi, Selvi, and Ramya [1760] define a *restricted triangular difference mean* label-

ing of a graph G with p vertices and q edges as an injection $f : V \to \{1, 2, 3, \ldots, pq\}$ such that for each edge uv, the edge labels defined by $f^*(uv) = \lceil |f(u) - f(v)|/2 \rceil$ are the first q triangular numbers. A graph that admits a restricted triangular difference mean labeling is called a *restricted triangular difference mean graph*. Jeyanthi, Selvi, and Ramya [1760] investigate the restricted triangular difference mean behaviors of the paths, combs, K_n, bistars $B_{m,n}$, caterpillars $S(n_1, n_2, \ldots, n_m)$, $K_{m,n}$, wheels, and graphs obtained by joining the centers of different stars to the new vertex. They also give a necessary condition for a graph to be a restricted triangular difference mean graph.

A (p, q) graph $G(V, E)$ is said to be an *analytic mean* graph if it is possible to injectively label the vertices with $\{0, 1, 2, \ldots, p-1\}$ in such a way that when each edge uv is labeled with $|(f(v)^2 - (f(u))^2|/2$ when $|(f(v)^2 - (f(u))^2|$ is even and $|(f(v)^2 - (f(u))^2 + 1|/2$ when $|(f(v)^2 - (f(u))^2| + 1$ is odd and the edge labels are distinct. In this case, f is called an *analytic mean* labeling of G. Raj and Vivek [3087] proved that $P_m \cup C_n \cup K_{1,s}$ $(m, s \geq 2,\ n \geq 3)$, $(P_m + K_1) \cup K_{1,n}$ $(m, n \geq 2)$, graphs obtained by identifying the apex vertices of $K_{1,m}$ and $K_{1,n}$ and one vertex of two copies of C_s where $m, n \geq 2,\ c \geq 3$ are analytic mean graphs.

Let $G = (V, E)$ be a graph with p vertices and q edges. A graph G is *analytic odd mean* if there exist an injective function $f : V \to \{0, 1, 3, 5 \ldots, 2q - 1\}$ with an induce edge labeling $f^* : E \to Z$ such that for each edge uv with $f(u) < f(v)$, $f^*(uv) = \left\lceil \frac{f(v)^2 - (f(u)+1)^2}{2} \right\rceil$ if $f(u) \neq 0$, and $f^*(uv) = \left\lceil \frac{f(v)^2}{2} \right\rceil$ if $f(u) = 0$ is injective. In this case we say that f is an *analytic odd mean labeling* of G. Jeyanthi, Gomathi, and Lau [1682] proved that fans, double fans, double wheels, closed helms, total graphs of cycles, total graphs of paths, armed crowns $C_n \Theta P_m$, generalized Petersen graphs $GP(n, 2)$ are analytic odd mean graphs. In [1670] they prove that wheels, flower graphs, some splitting graphs, and multiples of graphs are analytic odd mean graphs. In [1671] they prove that quadrilateral snakes, double quadrilateral snakes, coconut trees, fire cracker graphs, some star graphs, splitting graphs, complete bipartite graphs, unicyclic graphs, and the graphs obtained from a path of vertices $v_1, v_2, v_3, \ldots, v_n$ by joining i pendent vertices at each of ith vertex $1 \leq i \leq n$ (denoted $P_n(1, 2, \ldots, n)$) are analytic odd mean graphs. Jeyanthi, Gomathi, and Lau [1683] proved the square graphs of $P_n, C_n, B_{n,n}$ H-graphs, and $H \odot mK_1$ admit analytic odd mean labelings.

In [1767] Jeyanthi, Selvi, and Rama proved that the following graphs admit analytic odd mean labeling: arbitrary union of paths, $M_2(P_n)(n \geq 2)$, ladders, slanting slanting ladders, diamond snakes, quadrilateral snakes, alternately quadrilateral snakes, $Jl_n(P_3)(n \geq 1)$, $C_4 \odot K_{1,n}(n \geq 1)$, $DUP_2(K_{1,n})$, $DUP_2(B_{n,n})$, friendship graphs, and $nC_4(n \geq 1)$. Jeyanthi, Gomathi, Lau, and Shiu [1673] proved that the following graphs admit analytic odd mean labeling: $P_n + C_m$, tadpoles $T(m, n)$, $C_{m_1} + C_{m_2}$, $JC(m_1, m_2)$, edge linked cycle snakes $EL(kC_n)$, triangular snakes T_n, double triangular snakes $D(T_n)$, $P_n(C_m)$, nC_m, $P_n(T(m, 1))$, shell graphs Sh_n. They propose the conjecture that a (p, q)-graph is analytic odd mean when $p \leq q + 1$.

In [1668] Jeyanthi and Gomathi proved that the subdivision and super subdivision of the following graphs are analytic odd mean: cycles, stars, combs, and graphs obtained from $P_n \odot K_1$ by subdividing the each edge of P_n. Jeyanthi, Gomathi, and Lau [1671]

proved that quadrilateral snakes, double quadrilateral snakes, coconut trees, fire cracker graphs, and $P_n(1,2,\ldots,n)$ are analytic odd mean graphs. They also proved that $C_n \odot K_1$, prisms, helms, banana trees, perfect binary trees, unicyclic graphs, certain caterpillars, and spiders are analytic odd mean graphs. Jeyanthi and Gomathi [1669] proved that the graphs TL_n, (the triangular ladder obtained from L_n), $TL_n \odot K_1$, $T_n \odot K_1$, and $Q_n \odot K_1$ admit an analytic odd mean labelings. Simaringa and Thirunavukkarasu [3591] proved that triangular books, double triangular books, triangular snakes, double triangular snakes, butterflies, drums, $C_c \bigodot P_c$, $F_c \bigodot K_{1,d}$, $1 \le d \le 2c-1$, $W_c \odot K_{1,d}$, $1 \le d \le 2c$, $P_c^2 \odot K_{1,d}$, $c \ge 3$ and $1 \le d \le 2c-3$ are analytic odd mean graphs.

Let G be a (p,q) graph and f a injective function from $V(G)$ to $\{k, k+1, \ldots, p+q+k-1\}$ For each edge uv, let $f^* = \lceil (2f(u)f(v)/(f(u)+f(v)) \rceil$ or $\lfloor (2f(u)f(v)/(f(u)+f(v)) \rfloor$. We say f is a *k-super harmonic mean* if $f(V) \cup \{f^*(uv) \mid uv \in E(G)\} = \{k, k+1, \ldots, p+q+k-1\}$. A graph that admits a k-super harmonic mean labeling is called a *k-super harmonic mean* graph. In the case that $k=1$ the labeling is called a *super harmonic mean* labeling. For all $n > 1$ Tamilselvi and Revathi [3863] prove that the following graphs have k-super harmonic mean labelings: P_n, nP_m $(m>1), P_n \odot K_1, P_n \odot \overline{K_2}, P_n \odot \overline{K_3}, P_n^2$ $(n \ge 4)$, the subdivision graph of $P_n \odot K_1$, and the middle graph of P_n.

In [3286] Sandya, Merly, and Deepa introduced a heronian mean labeling of graphs as follows. A graph $G(V,E)$ with p vertices and q edges is said to be a *heronian mean graph* if there exists an injective function f from V to $1,2,3,\ldots,q+1$ such that when each edge uv is labeled with $f^*(uv) = \lfloor (f(u)+f(v)+\sqrt{f(u)f(v)})/3 \rfloor$ or $\lceil (f(u)+f(v)+\sqrt{f(u)f(v)})/3 \rceil$, the resulting edge labels are distinct. In this case f is called a *heronian mean labeling* of G. A heronian mean labeling f is called a *super heronian mean* labeling if $\{f(V(G)) \cup f(e) : e \in E(G)\} = \{1,2,3,\ldots,p+q\}$ In [4043] Valencerina and Pedrano proved that the subdivision of the triangular snake T_n, the quadrilateral snake Q_n, the double triangular snake $D(T_n)$, and the double quadrilateral snake $D(Q_n)$ are super heronian mean graphs for all $n \ge 3$.

Sampath, Narasimhan, and Nagaraja [3266] proved cycles, $K_{1,n}$ if and only if $n \le 4$, $C_m \cup P_n$, $C_m \cup C_n$, nK_3, $nK_3 \cup P_m$, $nK_3 \cup C_m$, mC_4, crowns $C_n \odot K_1$, dragons $C_n @ P_m$, and P_n^2 admit heronian mean labelings. Anitha, Selvam, and Thirusangu [261] provide heronian mean labelings for the extended duplicate graph of the kite graph.

A graph $G=(V,E)$ with p vertices and q edges is said to be a (k,d)-*heronian mean graph* if it is possible to label the vertices $x \in V$ with distinct labels $f(x)$ from $k, k+d, k+2d, \ldots, k+qd$ in such a way that when each edge uv is labeled with $f^*(uv) = \lfloor (f(u)+f(v)+\sqrt{f(u)f(v)})/3 \rfloor$ or $\lceil (f(u)+f(v)+\sqrt{f(u)f(v)})/3 \rceil$, then the resulting edge labels are distinct. In this case f is called a (k,d)-*heronian mean labeling* of G. Note that the case $k=1$ and $d=1$ is a heronian mean labeling. Akilandeswari and Tamilselvi [161] proved that paths, ladders, and $P_n \odot mK_1$ for $n \ge 2$, $1 \le m \le 4$, are k-heronian mean graphs. In [161] Akilandeswari and Tamilselvi proved that the following graphs have (k,d)-heronian mean labelings: paths, $(P_n \times P_2) \odot K_1$, $T_n \odot K_1$ (T_n is the triangular snake obtained from P_n), $Q_n \odot K_1$, $TL_n \odot K_1$ (TL_n is the triangular ladder obtained from L_n), Peterson graphs, and the graphs obtained from two copies of P_n with vertices $v_1, v_2, \ldots, v_n$ and $u_1, u_2, \ldots, u_n$ by joining the vertices $u_{(n+1)/2}$ and $v_{(n+1)/2}$ if n is

odd and $u_{n/2+1}$ and $v_{n/2+1}$ if n is even.

A *pronic number* is one of the form $n(n+1)$, where n is a positive integer. In [2996] Porchelvi and Devi defined a *pronic Heron mean* labeling of a graph G with n vertices as a bijection $f : V(G) \to \{0, 2, 6, 12, \ldots, n(n+1)\}$ such that the resulting edge labels obtained by $f^*(uv) = \lceil (f(u)+f(v)+\sqrt{f(u)f(v)}/3 \rceil$ for every edge uv are distinct. In [942] Devi and Porchevi gave pronic Heron mean labelings for the generalized Peterson graphs $P(6,2)$, $P(8,3)$,$P(10,2)$, $P(10,3)$, and $P(12,5)$. They also investigated the existence of pronic Heron labelings for mK_3, the unions of cycles and paths, and the union of combs and paths. They described an algorithm to label the vertices for the pronic Heron mean labeling for certain disconnected graphs. They further investigated the existence of pronic labelings for mK_3, the unions of cycles and paths, and the union of combs and paths.

Arockiaraj and Meena [296] say a planar graph has an F-*face magic mean* labeling it there exists an assignment of labels to the edges that induces an assignment of labels to the faces of the graph such that the mean weight of each face is constant. They proved that the following graphs have F-face magic mean labelings: $P_{2n}+K_1$, the one-point union of m copies of C_n, mC_n-snakes, and graphs obtained by identifying the endpoints of any number of copies P_n. In [297] and [2472] Arockiaraj and Meena Kumari investigated the existence of $(1,0,0)$ F-face magic mean labelings for paths, butterflies, and middle graphs of cycles, P_n+K_1, $C_m \times P_n$, latitude graphs, cyclic ladders, graphs obtained by duplicating every edge of P_n ($n \geq 2$), even cycles, the middle graphs of cycles, and the middle graphs of butterflies. In [2473] they investigated $(0,1,0), (1,1,0)$-F-face magic mean labelings of slanting ladders, $P_m \times P_n$, $K_{2,n}$, and certain graphs obtained by applying graph operations. Vani Shree and Dhanalakshmi [4045] investigated the existence of (1, 0, 0)-F-face magic mean labelings of ladders, tortoises, and the middle graph of paths. They also provided (1,0,0)- and (1,1,0)-F-face magic mean labelings for ortho chain square cactus graphs, para chain square cactus graphs, triangular snakes, and quadrilateral snakes.

Amara Jothi, Baskar Babujee, and David [211] investigated face magic labeling of planar graphs of types $(1,0,1), (1,1,0)$, $(0,1,1)$ and (1, 1, 1) on duplication graphs. For a graph $G(V,E)$, Rajesh Kannan, Manivannan, and Durai Basker [1894] defined an *exponential* labeling as injective function χ from V to $\{1, 2, 3, \ldots, |E|+1\}$ such that the induced function χ^* from E to $\{1, 2, \ldots, |E|\}$ given by $\chi(uv)^* = \lfloor (1/e)(\chi(v)^{\chi(v)}/\chi(u)^{\chi(u)})^{(1/\chi(v)-\chi(u))} \rfloor$ for all $uv \in E$ is a bijection. A graph that allows an exponential mean labeling is called an *exponential.* They investigated the exponential mean labeling of graphs obtained by duplication of an edge or vertex of paths and path related graphs.

In 2022 Baskaran and Ganapathy [590] introduced a new variant of mean labeling as follows. (In prior editions of this survey the definition was stated incorrectly.) A function h is said to be a C-*exponential mean* labeling of a graph $G(V,E)$ if $h : V \to \{1, 2, \ldots, |E|+1\}$ is injective and the induced function $h^* : E \to \{2, 3, \ldots, |E|+1\}$ defined by $h^*(uv) = \lceil \frac{1}{e}((h(v)^{h(v)}/(h(u)^{h(u)}))^{1/(h(v)-h(u))} \rceil$ is bijective. They investigated the exponential meanness of paths, triangular trees (see Section 3.1), $C_m \odot P_n$, $P_m \times P_n$, one-sided step graphs (the graph obtained by starting with the path P_n with consecutive edges $e_1, e_2, \ldots, e_{n-1}$ and erecting a ladder that has $n-i-1$ steps with e_i at the bottom), double-sided step

graphs (the graph obtained by starting with the path P_{2n} with edges $e_1, e_2, \ldots, e_{2n-1}$, where on every edge e_i and erecting a ladder that has $i+1$ steps for $i = 1, 2, \ldots, n$ and on every edge and for $i = n+1, n+2, \ldots, 2n-1$ a ladder on edge e_i that has $2n+1-i$ steps), one-sided arrow graphs (the graphs obtained from joining the vertices of the bottom row of $P_m \times P_n$ to a new vertex), double-sided arrow graphs (the graphs obtained from joining the vertices of the bottom row of $P_m \times P_n$ to a new vertex and by joining the vertices of the right column of $P_m \times P_n$ to a new vertex), and subdivisions of ladders. Unaware of the Baskaran and Ganapathy's 2022 paper, in 2025 [1314] Gopi and Kannan introduced a different labeling using the term "C-exponential" labeling. To avoid confusion, I will use adifferent name as follows. A function h is said to be a *near* C-*exponential mean* labeling of a graph $G(V, E)$ if $h : V \to \{1, 2, \ldots, |E|\}$ is injective and the induced function $h^* : E \to \{1, 2, 3, \ldots, |E| + 1\}$ defined by $h^*(uv) = \lceil \frac{1}{e}((h(v)^{h(v)}/(h(u)^{h(u)}))^{1/(h(v)-h(u))} \rceil$ or $h^*(uv) = \lfloor \frac{1}{e}((h(v)^{h(v)}/(h(u)^{h(u)}))^{1/(h(v)-h(u))} \rfloor$ is bijective. (Note the addition of 1 as an edge label.) Gopi and Kannan proved the following quadrilateral-related graphs admit C-exponential mean labelings: Q_n $(n \geq 2), T(Q_n)$ $(n \geq 2), A(Q_n)$ $(n \geq 4), A(T(Q_n))$ $(n \geq 4), Q_n \odot \overline{K_1}$ $(n \geq 2)$, and $Q_n \odot \overline{K_2}$ $(n \geq 2)$. Madhumitha, Rao, Prabhu, Nayak, and D'Souza [2406] proposed a new encryption technique to encode messages in a way that leverages the mathematical properties of the C-exponential mean labeling. [1314] new [2406] new

In [1885] Kannan uses the term C_{exp} *average assignment* labeling rather than C-exponential labeling. He proved the following graphs admit C-exponential labelings: $C_m \cup P_n$ $(n \geq 2), T_m \cup C_n$ (T_m is obtained by attaching a pendent vertex to a neighbor of the pendent vertex of P_{n-1}), chains of C_n, graphs obtained from $C_{n_1}, C_{n_2}, \ldots, C_{n_k}$ by connecting each C_{n_i} to $C_{n_{i+1}}$ with some path, and tadpoles. [1885] new

A function f is said to be an *absolute mean graceful* of a graph G, if $f : V(G) \to \{0, \pm 1, \pm 2, \ldots, \pm|E|\}$ is injective and the edge labeling function $f^* : E(G) \to \{1, 2, \ldots, |E|\}$ defined as $f^*(uv) = \lceil (f(u) - f(v))/2 \rceil$ is bijective. A graph that admits such labeling is called *absolute mean graceful* graph. In [1831] and [1832] Kaneria, Chudasama, and Andharia proved that the path unions of a finite number of copies of trees, paths, cycles, complete bipartite graphs, grid graphs, step grid graphs, and double step grid graphs are absolute mean graceful graphs. In [158] Akbari1, Kaneria, and Parmar prove that jewel graphs, jelly fish graphs, and the extended jewel graphs admit absolute mean graceful labelings. In [1868] Kaneria and Shah investigated the absolute mean graceful labeling of m-splitting and degree splitting graphs of paths, bistarsm and stars. Jani and Kaneria [1573] gave absolute mean graceful labelings for m-splitting and m-shadow graphs of even cycles. In [2779] Parmar, Akbari, Rupani gave explicit labeling constructions to show that the eight-sprockets, octopus graphs, flower pots, triangular books, kayak paddle graphs $KP(3, 4, n)$, armed crowns, and the corona product of $P_n \odot \overline{K_n}$ admit absolute mean graceful labelings. [1868] new [1573] new [2779] new

In [2747] Palani and Shunmugapriya defined the concept of near mean labeling in digraphs as follows. Let $D(V, A)$ be a digraph where V is the vertex is set and A is the arc set. Let $f : V \to \{0,\ 1,\ 2,\ \ldots, q\}$ be a 1-1 map. Define $f^* : A \to \{1,\ 2,\ \ldots, q\}$ by $f^*(e = u\vec{v}) = \left\lceil \frac{f(u)+f(v)}{2} \right\rceil$. Let $f^*(v) = \left| \sum_{w \in V} f^*(v\vec{w}) - \sum_{w \in V} f^*(w\vec{v}) \right|$. If $f^*(v) \leq 2$ for all $v \in A(D)$, f is said to be a *near mean* labeling of D and D is said to be a *near mean*

digraph. They investigated the existence of near mean labeling in dragon digraphs (the one-point union of cycle and a path where the edges of a cycle are directed clockwise or anti-clockwise and the edges of the path edges are directed towards the cycle). In [2748] they investigated the existence of near mean labelings for various dicyclic snakes.

7.15 Pair Sum and Pair Mean Graphs

For a (p, q) graph G Ponraj and Parthipan [2901] define an injective map f from $V(G)$ to $\{\pm 1, \pm 2, \ldots, \pm p\}$ to be a *pair sum* labeling if the induced edge function f_{em} from $E(G)$ to the nonzero integers defined by $f_e(uv) = f(u) + f(v)$ is one-one and $f_e(E(G))$ is either of the form $\{\pm k_1, \pm k_2, \ldots, \pm k_{\frac{q}{2}}\}$ or $\{\pm k_1, \pm k_2, \ldots, \pm k_{\frac{q-1}{2}}\} \cup \{k_{\frac{q+1}{2}}\}$, according as q is even or odd. A graph with a pair sum labeling is called *pair sum graph*. In [2901] and [2903] they proved the following are pair sum graphs: P_n, C_n, K_n if and only if $n \leq 4$, $K_{1,n}$, $K_{2,n}$, bistars $B_{m,n}$, combs $P_n \odot K_1$, $P_n \odot 2K_1$, and all trees of order up to 9. Also they proved that $K_{m,n}$ is not pair sum graph if $m, n \geq 8$ and enumerated all pair sum graphs of order at most 5.

In [2905], [2906], [2907], and [2908] Ponraj, Parthipan, and Kala proved the following are pair sum graphs: $K_{1,n} \cup K_{1,m}$, $C_n \cup C_n$, mK_n if $n \leq 4$, $(P_n \times K_1) \odot K_1$, $C_n \odot K_2$, dragons $D_{m,n}$ for n even, $\overline{K_n} + 2K_2$ for n even, $P_n \times P_n$ for n even, $C_n \times P_2$ for n even, $(P_n \times P_2) \odot K_1$, $C_n \odot K_2$ and the subdivision graphs of $P_n \times P_2, C_n \odot K_1, P_n \odot K_1$, triangular snakes, and quadrilateral snakes. Ponraj, Parthipan, and Kala [2902] prove the following graphs are pair sum graphs: the union of two stars, the union of a path and a star, ladders, $C_n \odot K_1$, $C_m \cup C_m, mK_n$ if and only if $n \leq 4$, quadrilateral snake $Q_n(n \text{ odd})$, and triangular snakes.

In [2925] Ponraj and Prabhu investigated the existence of pair mean cordial labelings for copies of paths, copies of cycles, spider graphs, generalized theta graphs, tunjung graphs (obtained from a helm graph $H_n (n \geq 3)$ by adjoining a new pendent vertex w_i at the pendent vertex v_i of H_n, adding an edge $v_i v_{i+1}$ where $v_{n+1} = v_1$, and connecting v_i and w_i to the central vertex v_0 for $i = 1, 2, \ldots, n$), and volcano graphs (a triangle with pendant vertices attached to one vertex of the triangle). [2925] new

A (p, q)-graph G is said to be a *super pair sum* if there exists a bijection f from $V(G) \cup E(G)$ to $\{0, \pm 1, \pm 2, \ldots, \pm(\frac{p+q-1}{2})\}$ when $p + q$ is odd and from $V(G) \cup E(G)$ to $\{\pm 1, \pm 2, \ldots, \pm(\frac{p+q}{2})\}$ when $p + q$ is even such that $f(uv) = f(u) + f(v)$. A graph that admits a super pair sum labeling is called a *super pair sum graph*. Vasuki, Velmurugan, and Sugirtha [4074] prove that the graphs $H_n \odot mK_1$, (H_n is obtained from two copies of P_n $(n \geq 3)$ with vertices $v_1, v_2, \ldots, v_n$ and $u_1, u_2, \ldots, u_n$ by joining $v_{(n+1)/2}$ and $u_{(n+1)/2}$ if n is odd and $v_{n/2}$ and $u_{(n+2)/2}$ if n is even); $(P_{2n}; S_m)$, $S'(P_{2n})$, $< B_{m,n} : P_k >$ for $m \geq 1$, $n \geq 1$, $k \equiv 2 \pmod 4$, $< B(m) : P_k >$ for $m \geq 1$ $k \equiv 0, 2 \pmod 4$ and $2B_{m,n}$ $(m \geq 1, n \geq 1)$ are super pair sum graphs.

Jeyanthi and Sarada Devi [1739] define an injective map f from $E(G)$ to $\{\pm 1, \pm 2, \ldots, \pm q\}$ as an *edge pair sum labeling* of a graph $G(p, q)$ if the induced function of f^* from $V(G)$ to $Z - \{0\}$ defined by $f^*(v) = \sum f(e)$ taken over all edges e incident to v is one-one and $f^*(V(G))$ is either of the form $\{\pm k_1, \pm k_2, \ldots, \pm k_{p/2}\}$ or

$\{\pm k_1, \pm k_2, \ldots, \pm k_{(p-1)/2}\} \cup \{k_{p/2}\}$ according as p is even or odd. A graph with an edge pair sum labeling is called an *edge pair sum graph*. They proved that P_n, C_n, triangular snakes, $P_m \cup K_{1,n}$, and $C_n \odot \overline{K_m}$ are edge pair sum graphs.

Jeyanthi, Sarada Devi, and Lau [1749] proved that the following graphs have edge pair sum labelings: triangular snakes T_n, $C_n \cup C_n$, $K_{1,n} \cup K_{1,m}$, and bistars $B_{m,n}$. They also proved that every graph is a subgraph of a connected edge pair sum graph. Jeyanthi and Sarada Devi [1740] showed that $P_{2n} \times P_2$ and the graphs $P_n(+)N_m$ obtained from a path P_n by joining its endpoints to m isolated vertices are edge pair sum graphs. Jeyanthi and Sarada Devi [1742] proved that the following graphs have edge pair sum labeling: shadow graphs $S_2(P_n)$, $S_2(K_{1,n})$, total graphs $T(C_{2n})$ and $T(P_n)$, the one-point union of any number of copies of C_n, the one-point union of C_m and C_n, P^2_{2n-1}, and full binary trees in which all leaves are at the same level and every parent has two children. Jeyanthi and Sarada Devi [1741] proved the spiders $SP(1^m, 2^t)$, $SP(1^m, 2^t, 3)$, $SP(1^m, 2^t, 4)$, and for t even $SP(1^m, 3^t, 3)$ are edge pair sum graphs. In [1740] Jeyanthi and Sarada Devi prove some cycle related graphs are edge pair sum graphs. In [1742] they prove that the one point union of cycles, perfect binary trees, shadow graphs, total graphs, and P^2_n admit edge pair sum graph. In [1748] Jeyanthi and Sarada provide edge pair sum labelings for jewel graphs, gears, triangular ladders, balanced lobsters, and double wheels $2C_n + K_1$.

The tree $WT(n)$ is obtained from $K_{1,n+2}$ with central vertex c_1 and end vertices $x_i : 1 \leq i \leq n+2$ and another $K_{1,n+2}$ with central vertex c_2 and end vertices $y_j : 1 \leq j \leq n+2$ by identifying vertex x_{n+2} and y_{n+2} and denoting the identified vertices by w. A w-tree $WT(n : k)$ is obtained from k copies of $WT(n)$ by joining a new vertex a to vertex w of each copy of $WT(n)$. Jeyanthi, Sarada Devi, and Lau [1750] proved that the graphs $WT(n : k)$ trees have edge pair sum labelings (see also [1751]).

In [1744], [1750], [1743], [1747] Jeyanthi and Sarada Devi prove the following graphs are edge pair sum graphs: shell graphs; some butterfly graphs; jelly fish; Y-trees; theta graphs; wheels with subdivided spokes, $P_m + 2K_1$; $C_4 \times P_m$; $P_n \odot \overline{K}_m$; $(P_2 \times P_m) \odot \overline{K_n}$; $P_m \times C_3$; books; graphs obtained from the path P_n having an even fixed even number quadrilaterals on each edge of the path; $K_2 + mK_1$; graphs obtained by identifying one end point from each of m copies of P_n; closed helms; graphs that are two copies of generalized Petersen graphs joined by a path P_n, $n \geq 5$; and graphs that two copies of fan $P_n \odot K_1$ joined by a path P_n, $n \geq 5$.

In [1745] Jeyanthi and Sarada Devi prove the following graphs admit edge pair sum labelings: $K_{2,n}$, double triangular snakes, wheels, flowers, $\langle C_m, K_{1,n} \rangle$ ($m \geq 4$, n odd) obtained from C_m and $K_{1,n}$ by identifying any vertex of C_m with the central vertex of $K_{1,n}$, and $\langle C_m * K_1 \rangle$ ($m \geq 4$) the graphs obtained from C_m and $K_{1,n}$ by identifying any vertex of C_m with an endpoint vertex of $K_{1,n}$. In [1746] they prove that the subdivision of graph of bistars $B_{m,n}$, $P_n \odot K_1$, triangular snakes when the path has an odd number of vertices, double triangular snakes, double quadrilateral snakes, double alternative triangular snakes, and double alternative quadrilateral snakes are edge pair sum graph.

In [215] Amudha and Jayapriya introduced the notion of labeling pair sum modulo labelings as follows. An undirected simple graph G with p vertices and q edges is said to be a *pair sum modulo* graph if there is an injective function f from $V(G)$ to $\{\pm 1, \pm 2, \ldots, \pm p\}$

such that the induced edge labeling g from $E(G)$ to $\{0, 1, 2, \ldots, q-1\}$ defined by $g(uv) = (f(u) + f(v))(\mathrm{mod}\, q)$ is injective. They proved the Petersen graphs, a certain family of coconut trees, $K_{2,n}$, and bistars have pair sum modulo labelings. Jayapriya [1590] proved spider graphs with at most five legs admit pair sum modulo labelings. Ponraj, Parthipan, and Kala [2902] prove the following graphs are pair sum graphs: the union of two stars, the union of a path and a star, ladders, $C_n \odot K_1$, $C_m \cup C_m, mK_n$ if and only if $n \leq 4$, quadrilateral snake $Q_n(n$ odd), and triangular snakes.

For a (p, q) graph G Ponraj and Parthipan [2904] define an injective map f from $V(G)$ to $\{\pm 1, \pm 2, \ldots, \pm p\}$ to be a *pair mean* labeling if the induced edge function f_{em} from $E(G)$ to the nonzero integers defined by $f_{em}(uv) = (f(u) + f(v))/2$ if $f(u) + f(v)$ is even and $f_{em}(uv) = (f(u) + f(v) + 1)/2$ if $f(u) + f(v)$ is odd is one-one and $f_{em}(E(G)) = \{\pm k_1, \pm k_2, \ldots, \pm k_{q/2}\}$ or
$f_{em}(E(G)) = \{\pm k_1, \pm k_2, \ldots, \pm k_{(q-1)/2}\} \cup \{k_{(q+1)/2}\}$, according as q is even or odd. A graph with a pair mean labeling is called a *pair mean* graph. They proved the following graphs have pair mean labelings: P_n, C_n if and only if $n \leq 3$, K_n if and only if $n \leq 2$, $K_{2,n}$, bistars $B_{m,n}$, $P_n \odot K_1$, $P_n \odot 2K_1$, and the subdivision graph of $K_{1,n}$. Also they found a relation between pair sum labelings and pair mean labelings.

In [2578] Muthuselvi and Saratha Devi introduced the notion of distance pair mean labeling as follows. For a graph $G(V, E)$ with p vertices and q edges they define an injective map f from V to $\{\pm 1, \pm 2, \ldots, \pm p\}$ to be a *distance pair mean* labeling if the induced edge function $w(v)$ from the nonzero integers defined by $w(v) = \lceil |S|/|N(v)| \rceil \neq 0$, where S is the sum of all $f(u)$ over all vertices u adjacent to v is one-one and $w(V) = \{\pm k_1, \pm k_2, \ldots, \pm k_{q/2}\}$ or $w(v)) = \{\pm k_1, \pm k_2, \ldots, \pm k_{(q-1)/2}\} \cup \{k_{(q+1)/2}\}$, according to whether q is even or odd. A graph with a distance pair mean labeling is called a *distance pair mean* graph. They showed that P_n $(n \geq 3)$, C_n $(n \geq 5)$, $P_n \times P_2$ $(n \geq 3)$, and $P_n \odot K_1$ $(n \geq 3)$ admit distance pair mean labelings. They also observed that the star graph does not admit a distance pair mean labeling.

The graph $G@P_n$ is obtained by identifying an end vertex of a path P_n with any vertex of G. A graph $G(V, E)$ with q edges is called a $(k+1)$-*equitable mean* graph if there is a function f from V to $\{0, 1, 2, \ldots, k\}$ $(1 \leq k \leq q)$ such that the induced edge that labeling f^* from E to $\{0, 1, 2, \ldots, k\}$ given by $f^*(uv) - \lceil (f(u)+f(v))/2 \rceil$ has the properties $|v_f(i) - v_f(j)| \leq 1$ and $|e_{f^*}(i) - e_{f^*}(j)| \leq 1$ for $i, j = 0, 1, 2, \ldots, k$ where $v_f(x)$ and $e_{f^*}(x)$ are the number of vertices and edges of G respectively with the label x. In [1657] Jeyanthi proved the following: a connected graph with q edges is a $(q+1)$-equitable mean graph if and only if it is a mean graph; a graph is 2-equitable mean graph if and only if it is a product cordial graph; for every graph G, the graph $3mG$ is a 3-equitable mean graph; for every 3-equitable mean graph G, the graph $(3m+1)G$ is a 3-equitable mean graph; C_n is a 3-equitable mean graph if and only if $n \not\equiv 0 \pmod 3$; P_n is a 3-equitable mean graph for all $n \geq 2$; if G is a 3-equitable mean graph then $G@P_n$ is a 3-equitable mean graph for $n \equiv 1 \pmod 3$; the bistar $B(m, n)$ with $m \geq n$ is a 3-equitable mean graph if and only if $n \geq \lfloor q/3 \rfloor$; $K_{1,n}$ is a 3-equitable mean graph if and only if $n \leq 2$; and for any graph H and $3m$ copies $H_1, H_2, \ldots, H_{3m}$ of H, the graph obtained by identifying a vertex of H_i with a vertex of H_{i+1} for $1 \leq i \leq 3m-1$ is a 3-equitable mean graph.

In [2085] Lakshmi and Nagarajan introduced the notion of geometric mean cordial labeling of graphs as follows. Let $G = (V, E)$ be a graph and f be a mapping from $V(G)$ to $\{0, 1, 2\}$. The graph G is called *geometric mean cordial* if each edge uv can be assigned the label $\lceil \sqrt{f(u)f(v)} \rceil$ in such a way that and $|v_f(i) - v_f(j)| \leq 1$ and $|e_f(i) - ef_(j)| \leq 1$, where $v_f(x)$ and $e_f(x)$ denote the number of vertices and edges labeled with x and $x \in \{0, 1, 2\}$ They proved that P_n, C_n $(n \equiv 1, 2 \pmod 3)$ and $K_{1,n}$ are geometric mean cordial graphs and $K_n (n > 2)$, $K_{2,n}$ $(n > 2)$, $K_{n,n}$ $(n > 2)$ and wheels are not geometric mean cordial graphs. In [1859] Kaneria, Meera, and Maulik call these graphs *geometric mean 3-equitable*. They proved: K_{mn} $(m, n \geq 4)$ is not a geometric mean 3-equitable graph, caterpillars $S(x_1, x_2, \ldots, x_t)$ and $C_n \odot K_1$ $(t \geq 2)$ are geometric mean 3-equitable graphs, and $C_n \odot K_1$ is a geometric mean 3-equitable graph if and only if $n \equiv 1, 2 \pmod 3$.

7.16 Irregular Total Labelings

In 1988 Chartrand, Jacobson, Lehel, Oellermann, Ruiz, and Saba [783] defined an *irregular labeling* of a graph G with no isolated vertices as an assignment of positive integer weights to the edges of G in such a way that the sums of the weights of the edges at each vertex are distinct. The minimum of the largest weight of an edge over all irregular labelings is called the *irregularity strength* $s(G)$ of G. If no such weight exists, $s(G) = \infty$. Chartrand et al. gave a lower bound for $s(mK_n)$. Faudree and Lehel [1071] obtained $s(G) \leq \lceil n/2 \rceil + 9$ for regular graphs. For every graph G of order $n \geq 4$ and with finite irregularity strength, Nierhoff [2676] proved $s(G) \leq n-1$. Using deterministic and mostly probabilistic techniques, Frieze, Gould, Karonski, and Pfender [1122] obtained the following bounds: $s(G) \leq c_1 \cdot n/\delta)$ if $\Delta \leq n^{1/2}$; and $s(G) \leq c_2 \cdot (\log n)n/\delta$ if $\Delta > n^{1/2}$, where c_1 and c_2 are explicit constants. Bohman and Kravitz found an infinite sequence of trees with strength converging to $(11 - \sqrt{5})/8$. Faudree, Jacobson, and Lehel [1073] gave an upper bound for $s(mK_n)$ when $n \geq 5$ and proved that for graphs G with $\delta(G) \geq n - 2 \geq 1$, $s(G) \leq 3$. They also proved that if G has order n and $\delta(G) = n - t$ and $1 \leq t \leq \sqrt{n/18}$, $s(G) \leq 3$. They asked for the optimal lower bound for the minimum degree of a graph G of order n that implies $s(G) \leq K$ for any given constant K. More generally, they also posed a similar question regarding the upper bound $S(G) \leq K$ for any given constant K. In 2024 Przybylo [3052] provided asymptotically tight solutions of these problems by proving that such optimal lower bound is of order $n/(K-1)$ for every fixed integer $K \geq 3$. Aigner and Triesch proved $s(G) \leq n + 1$ for any graph G with $n \geq 4$ vertices for which $s(G)$ is finite. In [3049] Przybylo proved that $s(G) < 112n/\delta + 28$, where δ is the minimum degree of G and G has n vertices. The best bound of this form is currently due to Kalkowski, Karoński, and Pfender, who showed in [1814] that $s(G) \leq 6\lceil n/\delta \rceil < 6n/\delta + 6$. In Przybylo [3047] showed that for d-regular graphs $s(G) < 16n/d + 6$. In 1991 Cammack, Schelp, and Schrag [753] proved that the irregularity strength of a full d-ary tree $(d = 2, 3)$ is its number of pendent vertices and conjectures that the irregularity strength of a tree with no vertices of degree two is its number of pendent vertices. This conjecture was proved by Amar and Togni [210] in 1998. In [3051] Przybylo proved that the conjecture of Faudree and Lehel holds asymptotically for the cases in which d is neither very small nor very

close to n.

The *edge irregularity strength* of a graph G, denoted by $es(G)$, is the smallest positive integer k for which G possesses an edge irregular k-labeling. Muthu Guru Packiam, Manimaran, and Thuraiswamy [2614] prove the following: $s(C_n \odot mK_1) = mn, s(P_n \odot K_2) = n+1, s(C_n \odot K_2) = n+1, s(P_n \odot K_3) = n+1$, and $sC_n \odot K_3) = n+1$. In [1785] Jinnah and Kumar determined the irregularity strength of triangular snakes and double triangular snakes. Hasni, Tarawneh, Siddiqui, Raheem, and Asim [1384] determined the exact value of edge irregularity strength of disjoint union of cycles and generalized prisms. These results provide an upper bound Hashi, Tarawneh, and Husin [1381] determined the edge irregularity strength of $P_n \odot P_m$ for $n \geq 2$ and $m = 3, 4$, and 5. Imran, Aslam, Zafar, and Nazeer [1521] determined the exact value of the edge irregularity strength of caterpillars, n-star graphs, (n, t)-kite graphs, cycle chains, and friendship graphs. Tarawneh, Hasni, Ahmad, and Asim [3868] determined the exact value of the edge irregularity strength for some classes of plane graphs. In [130] Ahmad, Khan, Ahmad, Nadeem, and Siddiqui determined the exact value of the edge irregularity strength for categorical product of two paths. In [1526] Imran, Cancan, and Ali determined the edge irregularity strength of certain splitting graphs, shadow graphs, jewel graphs, jellyfish graphs, and m copies of 4-pan graphs. Uma Devi, Kamaraj, and Arockiaraj [940] determined the odd Fibonacci edge irregularity strength for P_n $(n \geq 2), K_{1,n}, P_n \odot K_1$ $(n \geq 2)$, bistars $B_{(m,n)}$ and proved the nonexistence of an odd Fibonacci edge irregular labeling for K_n $(n \leq 3)$ and $K_{m,n}$ $(m \geq 2, n \geq 4)$. In [941] Devi, Kamaraj, and Arockiaraj determined the odd Fibonacci edge irregularity strength for P_n $(n \geq 2)$, stars, subdivision of stars, subdivision of fans, $P_n \odot mK_1$ $(m, n \geq 2)$, and the graphs obtained by joining i pendent vertices to the ith vertex of P_n $(n \geq 2)$. Muthukkumar [2610] proved that the edge irregularity strength of the corona product of a tree T with K_1 is $2es(T)$. Further, he proved that the edge irregularity strength of binomial tree B_k is 2^{k-1}. Nadeem, Cancan, Imran, and Ali [2631] determined the exact value of edge irregularity strength of triangular snakes with and without pendent edges and alternate triangular snakes with and without pendent edges. In [1524] Imran, Cancan, Ali, and Aslam determined the exact values of edge irregularity strength of linear phenylene graphs and various families of snake graphs. Ahmad [100] determined the edge irregularity strength of the uniform subdivision of stars, fire crackers, and banana trees. In [4318] Zhang, Cancan, Nadeem, and Imran gave the exact value of the edge irregularity strength of various families of comb related graphs.

Salma, Nagesh, Vijaya Chandra Kumar, and Narahari [3241] determined the exact values of the edge irregularity strength for the series composition of the isomorphic uniform theta graphs, $S(\Theta_{m,n})$, for the cases where $m = 2$ and $m = 3$ with $n \geq 2$. Additionally, they gave the edge irregularity strength for the shadow graph of paths with odd length, and the square of paths and theta graphs each extended by adding a single pendent edge. They also discussed an application of edge irregularity strength in optimizing path selection within communication networks. Nagesh and Girish [2644] determined bounds of edge irregularity strength of the line graphs of combs. They also determined bounnds for the edge irregularity strength of the line cut-vertex graphs of combs. Salma, Nagesh, and Prahlad [3240] gave the exact value of the edge irregularity strength of firefly graphs [3241] new [2644] new [3240] new

of order n that consist of s triangles, t pendent paths of length 2, and $n-2s-2t-1 \geq 1$ pendent edges sharing a vertex in common. Salma and Nagesh [3238] determined the edge irregularity strength of ladders, triangular ladders, and diagonal ladders. Salma, Nagesh, Azghar Pasha, and Narahari [3239] determined the edge irregularity strength of the Mycielskian graphs of paths and cycles. Tarawneh, Hasni, and Ahmad [3869] determined the exact value of the edge irregularity strength for triangular grids, the zigzag graphs, and $P_n \times P_m \times P_2$. Putra and Susanti [3058] determined the total edge irregularity strength of centralized uniform theta graphs.

[3238] new

[3239] new

[3869] new

[3058] new

Motivated by the notion of the irregularity strength of a graph and various kinds of other total labelings, Bača, Jendroľ, Miller, and Ryan [379] introduced the *total edge irregularity strength* of a graph as follows. For a graph $G(V,E)$ a labeling $\partial : V \cup E \to \{1,2,\ldots,k\}$ is called an *edge irregular total k-labeling* if for every pair of distinct edges uv and xy, $\partial(u)+\partial(uv)+\partial(v) \neq \partial(x)+\partial(xy)+\partial(y)$. Similarly, ∂ is called an *vertex irregular total k-labeling* if for every pair of distinct vertices u and v, $\partial(u)+\sum \partial(e)$ over all edges e incident to $u \neq \partial(v)+\sum \partial(e)$ over all edges e incident to v. The minimum k for which G has an edge (vertex) irregular total k-labeling is called the *total edge (vertex) irregularity strength* of G. The total edge (vertex) irregular strength of G is denoted by $\mathrm{tes}(G)$ ($\mathrm{tvs}(G)$). They prove: for $G(V,E)$, E not empty, $\lceil (|E|+2)/3 \rceil \leq \mathrm{tes}(G) \leq |E|$; $\mathrm{tes}(G) \geq \lceil (\Delta(G)+1)/2 \rceil$ and $\mathrm{tes}(G) \leq |E| - \Delta(G)$, if $\Delta(G) \leq (|E|-1)/2$; $\mathrm{tes}(P_n) = \mathrm{tes}(C_n) = \lceil (n+2)/3 \rceil$; $\mathrm{tes}(W_n) = \lceil (2n+2)/3 \rceil$; $\mathrm{tes}(C_3^n)$ (friendship graph) $= \lceil (3n+2)/3 \rceil$; $\mathrm{tvs}(C_n) = \lceil (n+2)/3 \rceil$; for $n \geq 2$, $\mathrm{tvs}(K_n) = 2$; $\mathrm{tvs}(K_{1,n}) = \lceil (n+1)/2 \rceil$; and $\mathrm{tvs}(C_n \times P_2) = \lceil (2n+3)/4 \rceil$. In [3257] Salma1, Nagesh1, and Narahari determined the exact values of the edge irregular k-labeling for the one-point union of C_k and $K_{1,n-k}$ $(n > k)$ for $3 \leq k \leq 7$. Ahmad, Nurdin, and Baskoro [134] determined the exact value of the total edge (vertex) irregularity strength of generalized Halin graphs. Al-Mushayt, Ahmad, and Siddiqui [197] determined the exact values of the total edge irregular strength of hexagonal grid graphs. The (m,n)-lollipop graph denoted by $L_{m,n}$ is a graph obtained by joining a complete graph K_m to a path graph P_n with a bridge. In [320] Attiya, Ahmed, and Salama define a *triple star snake graph*, $PS_{3,n}$, as the graph obtained from the path P_n by replacing every edge in P_n with $K_{1,3}$. The graph obtained from the path P_n by replacing every edge in P_n with $K_{1,m}$ is called an *m-star snake graph* and is denoted by $PS_{m,n}$. They proved $tes(PS_{3,n}) = n-1$ and $tes(PS_{m,n}) = \lceil (mn+2)/3 \rceil$.

For positive integers $l \geq 2, k \geq 2$, and $d \geq 1$, the *arithmetic ladder graph* $T_{l,k,d}$ with l levels, k columns, and difference d, is the graph

$$\begin{aligned}
V(T_{l,k,d}) &= \{t_{i,0}^0 \mid 1 \leq i \leq l\} \\
&\quad \cup \{t_{i,j}^r \mid 1 \leq i \leq l,\ 1 \leq j \leq 1+(r-1)d,\ 1 \leq r \leq k\}, \\
E(T_{l,k,d}) &= \{t_{i,0}^0 t_{i,1}^1 \mid 1 \leq i \leq l\} \\
&\quad \cup \{t_{i,j}^r t_{i,1}^{r+1} \mid 1 \leq i \leq l,\ j = 1+(r-1)d,\ 1 \leq r \leq k-1\} \\
&\quad \cup \{t_{i,j}^r t_{i,j+1}^r \mid 1 \leq i \leq l,\ 1 \leq j \leq (r-1)d,\ 2 \leq r \leq k\} \\
&\quad \cup \{t_{i,0}^0 t_{i+1,0}^0 \mid 1 \leq i \leq l-1\} \\
&\quad \cup \{t_{i,j}^r t_{i+1,j}^r \mid 1 \leq i \leq l-1,\ j = 1+(r-1)d,\ 1 \leq r \leq k\}.
\end{aligned}$$

Wulandari and Susanti [4219] determined the exact value of the total edge irregularity strength of arithmetic ladder graphs. [4219] new

Ni'mah and Indriati [2677] determined $\text{tvs}(L_{m,n})$ for $m \geq 3$ and $n \geq 1$. Aftiana and Indriati [79] proved that for $n \geq 3$ the total edge irregularity strength of the graph obtained by joining two copies of K_n (barbell graph) with an edge is $\lceil (n^2 - n + 3)/3 \rceil$. In [2721] Nurdin and Hye consider the splitting graph of stars as a land transportation system and give the exact value of their total vertex irregularity strength. For $m, n \geq 3$ Indriati, Widodo, and Sugeng [1539] determined the exact value of the total vertex irregularity strength for generalized helm graphs H_n^m (obtained from W_n by attaching P_m vertices at each vertex of the n-cycle) and for prisms with outer pendent edges. Nurdin [2722] determined the total vertex irregularity strength of banana trees and quad trees. Susanti, Puspitasari and Khotimah [3815] determined the exact value of total edge irregularity strength for staircase graphs, double staircase graphs, and mirror-staircase graphs. In [3816] Susanti, Wahyuni, Sutjijana, Sutopo, and Ernanto provide the total edge irregularity strength of generalized arithmetic staircase graphs and generalized double-staircase graphs. Susanto, Simanjuntak, and Baskoro [3822] determined $\text{tvs}(T)$ for trees T where the maximum degree is at least three and $\text{tvs}(T)$ for trees T with maximum degree at least three and large number of exterior vertices.

In [3823] Ramalakshmi and Kathiresan the introduced following two notions. A total labeling λ is said to be an *edge irregular total absolute difference k-labeling* of the graph G if for every two different edges e and f of G we have $\text{wt}(e) \neq \text{wt}(f)$, where the weight of an edge $e = xy$ is defined as $\text{wt}(e) = |\lambda(e) - \lambda(x) - \lambda(y)|$. The minimum k for which the graph G has an edge irregular total absolute difference labeling is called the *total absolute difference edge irregularity strength* of the graph G, $\text{tades}(G)$. Ramalakshmi and Kathiresan obtained estimations on the parameter tades and determined the precise values of tades for paths, cycles, stars, wheels, friendship graphs, and T_p-tree graphs. They also showed that for any graph $G, \text{tades}(G) \geq \max\left(\frac{|E(G)|}{3} + 2, \Delta(G) + 1\right)$. Susanto and Wijaya [3824] determined the total absolute difference edge irregularity strength of triangular ladders, alternating triangulated ladders, and double triangulated ladders. [3823] new [3824] new

In [1441] Hinding, Sunusi, Syahrani, and Buhari defined the *modified helm* $\mathcal{H}_n$ as the graph constructed from helm graph H_n by removing its central vertex and non-central vertices, $x_1, x_2, \ldots, x_n, y_1, y_2, \ldots, y_n$ and adding $2n$ vertices $w_1, v_1, w_2, v_2, \ldots, w_n, v_n$ such that the edges of $\mathcal{H}_n$ are: $\{x_1 w_, x_n v_i\}(i = 1, 2, \ldots, n) \cup \{x_{i+1} w_i\}(i = 1, 2, \ldots, n-1) \cup \{x_{i-1} v_1\}(i = 2, \ldots, n) \cup \{x_i y_i\}(i = 1, 2, \ldots, n) \cup \{x_1 x_2, x_2 x_3, \ldots, x_{n-1} x_n, x_n x_1\}$. They determined the total vertex irregularity strength of the modified helm graph and provided an application of total vertex irregularity strength of the modified helm graph to the study of the topology of the transportation routes.

For R be a commutative ring R with the set of all zero divisor set $Z(R)$ of R the zero divisor graph of $R, \Gamma(R)$, has vertex set $Z(R)$ and (x, y) is an edge if and only if $xy = 0$. Ahmad [101] determined the total vertex irregularity strength of zero divisor graphs associated with the commutative rings $Z_{p^2} \times Z_q$ where p and q are distinct primes.

In [126] Ahmad, Ibrahim, and Siddiqui determined the total irregularity strength of generalized Petersen graphs. In [199] Al-Mushayt, Ahmad, and Siddiqui [199] deter-

mined the total edge (vertex) irregularity strength for convex polytope graphs having the same diameter. In [3577] Siddiqui determined the irregularity strength of six classes of convex polytope graphs with pendent edges. Ramdani, Salman, Assiyatum, and Semaničová-Feňovčíková [3118] establish upper bounds for the total vertex (edge) irregularity strength and total irregularity strength for disjoint union of arbitrary graphs. Naeem and Siddiqui [2634] determined the total irregularity strength of disjoint union of isomorphic copies of the generalized Petersen graph. In [4246] Yang, Siddiqui, Ibrahim, Ahmad, and Ahmad determined the exact value of the total irregularity strength of three planar graphs. In [3237] Salama introduced three kinds of snake graphs and investigated their total irregularity strength. Ibrahim, Khan, Asim, and Waseem [1480] determined the exact value of the total irregularity strength of cubic graphs. Tilukay, Tomasouw, Rumlawang, and Leleuray in [3904] proved that K_n and $K_{n,n}$ are both totally irregular total graphs with their ts equal to their tes. Tilukay, Taihuttu, Salman, Rumlawang, and Leleury [3901] proved that $K_{m,n}$ is a totally irregular total graph for any positive integer m and n.

Jendroľ, Miškul, and Soták [1645] (see also [1646]) proved: $\text{tes}(K_5) = 5$; for $n \geq 6$, $\text{tes}(K_n) = \lceil (n^2 - n + 4)/6 \rceil$; and that $\text{tes}(K_{m,n}) = \lceil (mn + 2)/3 \rceil$. They conjecture that for any graph G other than K_5, $\text{tes}(G) = \max\{\lceil (\Delta(G) + 1)/2 \rceil, \lceil (|E| + 2)/3 \rceil\}$. Ivančo and Jendroľ [1556] proved that this conjecture is true for all trees. Jendroľ, Miškuf, and Soták [1645] prove the conjecture for complete graphs and complete bipartite graphs. The conjecture has been proven for the categorical product of two paths [111], the categorical product of a cycle and a path [3575], the categorical product of two cycles [119], the Cartesian product of a cycle and a path [427], the subdivision of a star [3576], and the toroidal polyhexes [388]. In [137] Ahmad, Siddiqui, and Afzal proved the conjecture is true for graphs obtained by starting with m vertex disjoint copies of P_n $(m, n \geq 2)$ arranged in m horizontal rows with the jth vertex of row $i + 1$ directly below the jth vertex row i for $1 = 1, 2, \ldots, m-1$ and joining the jth vertex of row i to the $j+1$th vertex of row $i+1$ for $1 = 1, 2, \ldots, m-1$ and $j = 1, 2, \ldots, n-1$ (the *zigzag* graph). Siddiqui, Ahmad, Nadeem, and Bashir [3579] proved the conjecture for the disjoint union of p isomorphic sun graphs (i. e., $C_n \odot K_1$) and the disjoint union of p sun graphs in which the orders of the n-cycles are consecutive integers. They pose as an open problem the determination of the total edge irregularity strength of disjoint union of any number of sun graphs. Brandt, Misšku f, and Rautenbach [702] proved the conjecture for large graphs whose maximum degree is not too large relative to its order and size. In particular, using the probabilistic method they prove that if $G(V, E)$ is a multigraph without loops and with nonzero maximum degree less than $|E|/10^3\sqrt{8|V|}$, then $\text{tes}(G) = (\lceil |E| + 2)/3 \rceil$. As corollaries they have: if $G(V, E)$ satisfies $|E| \geq 3 \cdot 10^3 |V|^{3/2}$, then $\text{tes}(G) = \lceil (|E| + 2)/3 \rceil$; if $G(V, E)$ has minimum degree $\delta > 0$ and maximum degree Δ such that $\Delta < \delta\sqrt{|V|}/10^3 \cdot 4\sqrt{2}$ then $\text{tes}(G) = \lceil (|E| + 2)/3 \rceil$; and for every positive integer Δ there is some $n(\Delta)$ such that every graph $G(V, E)$ without isolated vertices with $|V| \geq n(\Delta)$ and maximum degree at most Δ satisfies $\text{tes}(G) = \lceil (|E| + 2)/3 \rceil$. Notice that this last result includes d-regular graphs of large order. They also prove that if $G(V, E)$ has maximum degree $\Delta \geq 2|E|/3$, then G has an edge irregular total k-labeling with $k = \lceil (\Delta + 1)/2 \rceil$. Pfender [2831] proved the

conjecture for graphs with at least 7×10^{10} edges and proved for graphs $G(V, E)$ with $\Delta(G) \leq E(G)/4350$ we have $\text{tes}(G) = (\lceil |E| + 2)/3 \rceil$. Susanti and Haq [3812] determined the minimum k such that an odd staircase graph can be labeled by an edge irregular total k-labeling. Murhu Guru Packiam, Manimaran, and Thuraiswamy [2571] investigate how the addition of a new edge affects the total edge irregularity strength of a graph. Laurence and Kathiresan [2127] determined the total edge irregular strength of path union of cycles. Sivakumar, Vidyanandini, Sreedevi, Nayak, and Bhoi [3623] determined the total edge irregularity strength of complete tripartite graphs.

In [1764] Jeyanthi and Sudha investigated the total edge irregularity strength of the disjoint union of wheels. They proved the following: $\text{tes}(2W_n) = \lceil (4n+2)/3 \rceil, n \geq 3$; for $n \geq 3$ and $p \geq 3$ the total edge irregularity strength of the disjoint union of p isomorphic wheels is $\lceil (2(pn+1)/3 \rceil$; for $n_1 \geq 3$ and $n_2 = n_1 + 1$, $\text{tes}(W_{n_1} \cup W_{n_2}) = \lceil (2(n_1 + n_2 + 1)/3) \rceil$; for n_1, n_2, n_3 where $n_1 \geq 3$ and $n_{i+1} = n_1 + i$ for $i = 1, 2$, $\text{tes}(W_{n_1} \cup W_{n_2} \cup W_{n_3}) = \lceil (2(n_1 + n_2 + n_3 + 1)/3) \rceil$; the total edge irregularity strength of the disjoint union of $p \geq 4$ wheels $W_{n_1} \cup W_{n_2} \cup \cdots \cup W_{n_p}$ with $n_{i+1} = n_1 + i$ and $N = \sum_{j=1}^{p} n_j + 1$ is $\lceil 2N/3) \rceil$; and the total edge irregularity strength of $p \geq 3$ disjoint union of wheels $W_{n_1} \cup W_{n_2} \cup \cdots \cup W_{n_p}$ and $N = \sum_{j=1}^{p} n_j + 1$ is $\lceil (2N/3 \rceil$ if $\max\{n_i \mid 1 \leq i \leq p\} \leq \frac{1}{2} \lceil (2N/3 \rceil$.

In [2562] Mughal and Jamil determined the tight lower bounds for the total face irregular strength of type $(1, 1, 0)$ of grids and wheels. Mughal, Jamil, and Virk [2563] determined the exact tight lower bounds for the face irregular strength of generalized plane graphs under a ρ-labeling of class $(\alpha_1, \beta_1, \gamma_1)$ for the vertex ρ-(1, 0, 0), the edge ρ-(0, 1, 0), the face ρ-(0, 0, 1), the vertex-face ρ-(1, 0, 1), the edge-face ρ-(0, 1, 1) and the ρ-(1, 1, 1) for grid graphs.

A k-labeling ϕ of a planar graph G is defined to be a face irregular k-labeling of type (α, β, γ) if for every two different faces f and g of G we have $wt_{\phi(\alpha,\beta,\gamma)}(f) \neq wt_{\phi(\alpha,\beta,\gamma)}(g)$. The face irregularity strength of type (α, β, γ) of a planar graph G, denoted $\text{fs}(\alpha, \beta, \gamma)(G)$, is the smallest integer k such that G admits a face irregular k-labeling of type (α, β, γ). In [420] Bača, Ovais, Semaničová-Feňovčíková, and Nengah Suparta, estimated the lower bounds and the upper bounds of the face irregularity strength of type (α, β, γ) for 2-connected planar graphs, where $\alpha, \beta, \gamma \in \{0, 1\}$, and determined the precise values of these parameters for ladders and fan graphs and proved the sharpness of the lower bounds.

The *complete star* of a graph G is the graph obtained from $p+1$ copies of the graph G by joining each vertex of $G^{(0)}$ with all corresponding vertices of all the copies $G^{1)}, \ldots, G^{(p)}$. Susanti, Khotimah, Hidayati, and Wahyujati [3813] determined the total edge irregularity strength of snowflake graphs, water bears graphs, the complete star of C_n, and two other families of ladder related graphs.

In [1768], [1770], [1771], and [1769] Jeyanthi and Sudha determine the total edge irregularity strength of fans, helms, closed helms, webs, flowers, gears, sun flowers, tadpoles, armed crowns, split graphs of cycles, split graph of paths, disjoint unions of isomorphic double wheels, and disjoint unions of consecutive non-isomorphic double wheels. Bokhary, Ali, and Maqbool [682] determined the exact values for the total vertex and edge irregularity strength of three wheel related families of graphs. Bokhary, Imran, and Ali [684] determined the exact value of convex polytopes generated by prisms and antiprisms and

determined their total vertex irregularity strength and total edge irregularity strength. Ibrahim, Asif, Ahmad, and Siddiqui [1478] investigated the total irregularity strength of fans, helms, closed helms, webs, flower graphs, gears, and sunflowers. In [3148] Ratnasari1, Wahyuni1, Susanti1, and Palupi1 determined the total edge irregularity strength of book graphs, double and triple book graphs, and gave the exact value of the total edge irregularity strength of quadruplet book graphs and quintuplet book graphs. Nurdini, Rosyida, and Mulyono [2726] determined the total edge irregularity strength of the chain graph that consists of tadpole graph $T(6, n)$ on each block and constructed an algorithm to find it.

A generalized helm H_n^m is a graph obtained by inserting m vertices in every pendent edge of a helm H_n. Indriati, Widodo, and Sugeng [1539] proved that for $n \geq 3$, $\text{tes}(H_n^1) = \lceil (4n+2)/3 \rceil$, $\text{tes}(H_n^2) = \lceil (5n+2)/3 \rceil$, and $\text{tes}(H_n^m) = \lceil ((m+3)n+2))/3 \rceil$ for $m \equiv 0 \bmod 3$. They conjecture that $\text{tes}(H_n^m) = \lceil ((m+3)n+2))/3 \rceil$, for all $n \geq 3$ and $m \geq 10$.

A *cactus* graph G is a connected graph where no edge lies in more than one cycle. A cactus graph consisting of some blocks where each block is C_n with same n is called an *n-uniform cactus graph*. If each cycle of the cactus graph has no more than two cut-vertices and each cut-vertex is shared by exactly two cycles, then G is called *n-uniform cactus chain* graph. Rosyida and Indriati [3212] determined the tes of n-uniform cactus chain graphs of length r for some $n \equiv 0 \bmod 3$. They also investigated the tes of tadpole chain graphs. Rosyida and Indriati [3211] determined the total edge irregularity strength of the triangular cactus chain with length r and $r+1$ pendent vertices (TC_r^{r+1}) is $\lceil (4r+3)/3 \rceil$. A *para-squares cactus* is a graph each of whose blocks is a 4-cycle and two or more squares share a cut vertex. The *para-squares cactus chain graph* is a cactus graph such that each of the two squares has one common cut vertex. Rosyida and Indriati determined that total edge irregularity strength of para square cactus chain graph with length r and r pendent vertices (Q_r^r) is $\lceil (5r+2)/3 \rceil$. Rosyida, Ningrum, Mulyono, and Indriati [3213] determined the total edge irregularity strength of general uniform cactus chain graphs having $(n-2)r$ pendent vertices and length r.

Nurdin, Baskoro, Salman, and Gaos [2724] determine the total vertex irregularity strength of trees with no vertices of degree 2 or 3; improve some of the bounds given in [379]; and show that $\text{tvs}(P_n) = \lceil (n+1)/3) \rceil$. They also made the following two conjectures. (A) For every tree T, $\text{tvs}(T) = \max \lceil (n_1+1)/2 \rceil, \lceil (n_1+n_2+1)/3 \rceil, \lceil (n_1+n_2+n_3+1)/4 \rceil$. (B) For every graph G with minimum degree δ and maximum degree Δ, $\text{tvs}(G) = \max \lceil (\delta + \sum_{j=1}^{i} n_j)/(i+1) \rceil : i \in [\delta, \Delta]$, where n_j denotes the number of vertices of degree j. In a 2023 Susanto, Simanjuntak, and Baskoro [3820] disproved these conjectures by giving infinite families of counterexamples.

In [2727] Nurdin, Salman, Gaos, and Baskoro prove that for $t \geq 2$, $\text{tvs}(tP_1) = t$; $\text{tvs}(tP_2) = t+1$; $\text{tvs}(tP_3) = t+1$; and for $n \geq 4$, $\text{tvs}(tP_n) = \lceil (nt+1)/3) \rceil$. Ahmad, Bača, and Bashir [112] proved that for $n \geq 3$ and $t \geq 1$, $\text{tvs}((n,t)\text{-kite}) = \lceil (n+t)/3 \rceil$, where the (n,t)-kite is a cycle of length n with a t-edge path (the tail) attached to one vertex. In [1344] Guo, Chen, Wang, and Yao give the total vertex irregularity strength

of certain complete m-partite graphs. In [3829] Susilawati, Baskaro, and Simanjuntak determined the total vertex irregularity strength of a particular subdivision of any tree. In [3831], [3832], and [3830] Susilawati, Baskoro and Simanjuntak determined the total vertex irregularity strength for trees with maximum degree four or five, for subdivision of several classes of caterpillars, for the subdivision of fire crackers, and for the subdivision of an amalgamation of stars. In [3833] they studied the vertex irregularity strength for trees having many vertices of degree 2. Nurdin, Baskoro, Salman and Gaos [2725] determined the total vertex irregularity strength for firecrackers and banana trees.

Anholcer, Kalkowski, and Przybylo [257] prove that for every graph with $\delta(G) > 0$, $\text{tvs}(G) \leq \lceil 3n/\delta \rceil + 1$. Majerski and Przybylo [2407] prove that the total vertex irregularity strength of graphs with n vertices and minimum degree $\delta \geq n^{0.5}\ln n$ is bounded from above by $(2 + o(1))n/\delta + 4$. Their proof employs a random ordering of the vertices generated by order statistics. Anholcer, Karonński, and Pfender [256] prove that for every forest F with no vertices of degree 2 and no isolated vertices $\text{tvs}(F) = \lceil (n_1 + 1)/2 \rceil$, where n_1 is the number of vertices in F of degree 1. They also prove that for every forest with no isolated vertices and at most one vertex of degree 2, $\text{tvs}(F) = \lceil (n_1 + 1)/2 \rceil$. Anholcer and Palmer [258] determined the total vertex irregularity strength C_n^k, which is a generalization of the circulant graphs $C_n(1, 2, \ldots, k)$. They prove that for $k \geq 2$ and $n \geq 2k + 1$, $\text{tvs}(C_n^k = \lceil (n + 2k)/(2k + 1) \rceil$. Przybylo [3049] obtained a variety of upper bounds for the total irregularity strength of graphs as a function of the order and minimum degree of the graph.

In [3909] Tong, Lin, Yang, and Wang give the exact values of the total edge irregularity strength and total vertex irregularity strength of the toroidal grid $C_m \times C_n$. In [3580] Siddiqui, Miller, and Ryan determine the exact values of the total edge irregularity strength of octagonal grid graph. In [120] Ahmad, Bača, and Siddiqui gave the exact value of the total edge and total vertex irregularity strength for disjoint union of prisms and for disjoint union of cycles. In [117] Ahmad, Bača, and Numan showed that $\text{tes}(\bigcup_{j=1}^{m} F_{n_j}) = 1 + \sum_{j=1}^{m} n_j$ and $\text{tvs}(\bigcup_{j=1}^{m} F_{n_j}) = \lceil (2 + 2\sum_{j=1}^{m} n_j)/3 \rceil$, where $\bigcup_{j=1}^{m} F_{n_j}$ denotes the disjoint union of friendship graphs. Chunling, Xiaohui, Yuansheng, and Liping, [831] showed $\text{tvs}(K_p) = 2$ $(p \geq 2)$ and for the generalized Petersen graph $P(n, k)$ they proved $\text{tvs}(P(n, k)) = \lceil n/2 \rceil + 1$ if $k \leq n/2$ and $\text{tvs}(P(n, n/2)) = n/2 + 1$. They also obtained the exact values for the total vertex strengths for ladders, Möbius ladders, and Knödel graphs. For graphs with no isolated vertices, Przybylo [3047] gave bounds for $\text{tvs}(G)$ in terms of the order and minimum and maximum degrees of G. For d-regular $(d > 0)$ graphs, Przybylo [3049] gave bounds for $\text{tvs}(G)$ in terms d and the order of G.

Ahmad, Ahtsham, Imran, and Gaig [102] determined the exact values of the total vertex irregularity strength for five families of cubic plane graphs. In [109] Ahmad and Bača determine that the total edge-irregular strength of the categorical product of C_n and P_m where $m \geq 2$, $n \geq 4$ and n and m are even is $\lceil (2n(m - 1) + 2)/3 \rceil$. They leave the case where at least one of n and m is odd as an open problem. In [119] and [120] Ahmad, Bača, and Siddiqui determine the exact values of the total edge irregularity strength of the categorical product of two cycles, the total edge (vertex) irregularity strength for the disjoint union of prisms, and the total edge (vertex) irregularity strength for the

disjoint union of cycles. In [108] Ahmad, Awan, Javaid, and Slamin study the total vertex irregularity strength of flowers, helms, generalized friendship graphs, and web graphs. Indriati, Widodo, Wijayanti, Sugeng, and Bača [1541] determine the exact value of the total edge irregularity strength of the generalized web graph $W(n,m)$ and two families of related graphs. Ahmad, Bača, and Numan [117] determined the exact values of the total vertex irregularity strength and the total edge irregularity strength of a disjoint union of friendship graphs. Bokhary, Ahmad, and Imran [681] determined the exact value of the total vertex irregularity strength of Cartesian and categorical product of two paths. Koam and Ahmad [1972] determined the total vertex irregularity strength for all theta graphs and certain values of the total vertex irregularity strength of the centralized uniform theta graphs. They provide a conjecture for the lower bound of total vertex irregularity strength of the centralized uniform theta graphs. In [683] Bokhary and Faheem proved the conjecture of Bokhary, Ahmam, and Imran [681] that the $tvs(P_m \Box P_n) = \lfloor \frac{mn+2}{5} \rfloor$ for $m, n \geq 2$ for $5 \leq m \leq 10$ and $n \geq 1$. the state graph for Tower of Hanoi problems with three towers. Farida and Indriati [1068] determined the total edge irregularity strength of the state graph for Tower of Hanoi problems with three towers.

In [2723] Nurdin, Salman, and Baskoro determine the total edge-irregular strength of the following graphs: for any integers $m \geq 2,\ n \geq 2$, $\text{tes}(P_m \odot P_n) = \lceil (2mn+1)/3 \rceil$; for any integers $m \geq 2,\ n \geq 3$, $\text{tes}(P_m \odot C_n) = \lceil ((2n+1)m+1)/3 \rceil$; for any integers $m \geq 2,\ n \geq 2$, $\text{tes}(P_m \odot K_{1,n}) = \lceil (2m(n+1)+1)/3 \rceil$; for any integers $m \geq 2$ and $n \geq 3$, $\text{tes}(P_m \odot G_n) = \lceil (m(5n+2)+1)/3 \rceil$ where G_n is the gear graph obtained from the wheel W_n by subdividing every edge on the n-cycle of the wheel; for any integers $m \geq 2,\ n \geq 2$, $\text{tes}(P_m \odot F_n) = \lceil m(5n+2)+1 \rceil$, where F_n is the friendship graph obtained from W_{2m} by subdividing every other rim edge; for any integers $m \geq 2$ and $n \geq 3$; and $\text{tes}(P_m \odot W_n) = \lceil ((3n+2)m+1)/3 \rceil$.

In [3094], [3096], and [3093] Rajasingh, Rajan, Teresa Arockiamary, and Quadras provide the total edge irregularity strengths of honeycomb mesh networks, hexagonal networks, butterfly networks, benes networks, and series compositions of uniform theta graphs. In [3095] Rajasingh and Teresa Arockiamary determined the total edge irregularity strength of honeycomb torus network, circulant networks, and hexagonal networks.

Ratnasari and Susanti [3136] determined the exact value of the total edge irregularity strength of triangular ladders, diagonal ladders, and circular triangular ladders. In [2725] Nurdin, Baskoro, Salman, and Gaos proved: the total vertex-irregular strength of the complete k-ary tree ($k \geq 2$) with depth $d \geq 1$ is $\lceil (k^d+1)/2 \rceil$ and the total vertex-irregular strength of the subdivision of $K_{1,n}$ for $n \geq 3$ is $\lceil (n+1)/3 \rceil$. They also determined that if G is isomorphic to the caterpillar obtained by starting with P_m and m copies of P_n denoted by $P_{n,1}, P_{n,2}, \ldots, P_{n,m}$, where $m \geq 2,\ n \geq 2$, then joining the i-th vertex of P_m to an end vertex of the path $P_{n,i}$, $\text{tvs}(G) = \lceil (mn+3)/3 \rceil$. They conjectured that the total vertex irregularity strength of any tree T is determined only by the number of vertices of degrees 1, 2 and 3 in T. This conjecture was confirmed by Susilawati, Baskoro, and Simanjuntak [3828] by considering all trees with maximum degree five. They also characterized all such trees having the total vertex irregularity strength either t_1, t_2, or t_3, where $t_i = \lceil (1 + \sum_{j=1}^{i} n_j)/(i+1) \rceil$ and n_i is the number of vertices of degree i.

Ahmad and Bača [110] proved $\text{tvs}(J_{n,2}) = \lceil (n+1)/2 \rceil$ $(n \geq 4)$ and conjectured that for $n \geq 3$ and $m \geq 3$, $\text{tvs}(J_{n,m}) = \max\{\lceil (n(m-1)+2)/3 \rceil, \lceil (nm+2)/4 \rceil\}$. They also proved that for the circulant graph (see §5.1 for the definition) $C_n(1,2)$, $n \geq 5$, $\text{tvs}(C_n(1,2)) = \lceil (n+4)/5 \rceil$. They conjecture that for the circulant graph $C_n(a_1, a_2, \ldots, a_m)$ with degree r at least 5 and $n \geq 5$, $1 \leq a_i \leq \lfloor n/2 \rfloor$, $\text{tvs}(C_n(a_1, a_2, \ldots, a_m) = \lceil (n+r)/(1+r) \rceil$. Ahmad, Arshadb, and Ižaríková [107] determine $\text{tes}(G)$ where G is the generalized helm and $\text{tvs}(G)$ where G is the generalized sun graph.

Slamin, Dafik, and Winnona [3630] consider the total vertex irregularity strengths of the disjoint union of isomorphic sun graphs, the disjoint union of consecutive nonisomorphic sun graphs, $\text{tvs}(\cup_{i=1}^{t} S_{i+2})$, and disjoint union of any two nonisomorphic sun graphs. (Recall $S_n = C_n \odot K_1$.) Rajasingh and Annamma [3092] determine the total vertex irregularity strength of 1-fault tolerant Hamiltonian graphs $CH(n), H(n)$, and $W(m)$. Indriati, Widodo, Wijayanti, Sugeng, Bača, and Semaničová-Feňovčíková [1542] determine the exact value of the total vertex irregularity strength for generalized helm graphs and for prisms with outer pendent edges. In [317] Asim and Hasni provided an upper bound for $\text{es}(K_n)$ that is far better than the previously known upper bound.

In [96] Ahmad shows that the total vertex irregularity strength of the antiprism graph A_n $(n \geq 3)$ is $\lceil (2n+4)/5 \rceil$ (see §5.7 for the definition) and gives the vertex irregularity strength of three other families convex polytope graphs. Al-Mushayt, Arshad, and Siddiqui [198] determined an exact value of the total vertex irregularity strength of some convex polytope graphs. Ahmad, Baskoro, and Imran [123] determined the exact value of the total vertex irregularity strength of disjoint union of helm graphs.

For $n \geq 3, m \geq 2$ Jeyanthi and Sudha [1772] determine the total vertex irregularity strength of $P_n \odot K_1, P_n \odot K_2$, $C_n \odot K_2$, $L_n \odot K_1$, $P_2 \odot C_n$, $P_n \odot \overline{K_m}$, $(C_n \times P_2) \odot K_1$, and $C_n \odot \overline{K_m}$. In [1773] they determine the total vertex irregularity strength for the graph obtained from a cycle by identifying the endpoint of a path and the vertex of a cycle, $C_n \odot P_m$, the split graph of a cycle, and split graph of a path. In [1773] they determine the total vertex irregularity strength for quadrilateral snakes, sunflowers, double wheels, triangular books, quadrilateral books, and graphs obtained from the wheel W_n and attaching n pendent edges to the center. In [1775] Jeyanthi and Sudha determined the total irregularity strength of the n-crossed prism, m copies of crossed prism, necklace and m copies of necklace graph and that these graphs admit totally irregular total k-labeling.

A total edge Fibonacci irregular labeling $f : V(G) \cup E(G) \to \{1, 2, \ldots, k\}$ of a graph G is a labeling of vertices and edges of G in such a way that for distinct edges xy and $x'y'$ their weights $f(x) + f(xy) + f(y)$ and $f(x') + f(x'y') + f(y')$ and are distinct Fibonacci numbers. A graph that has a total edge Fibonacci irregular labeling is called a *total edge Fibonacci irregular* graph. Karthikeyan, Navanaeethakrishnan, and Sridevi [1912] proved that stars, bistars $B_{n,n}$ $(n \geq 2)$, and two particular families of star related and bistar related graphs are total edge Fibonacci irregular graphs. The total edge Fibonacci irregularity strength, $\text{tefs}(G)$ is the minimum k for which G has total edge Fibonacci irregular labeling. Amutha and Uma Devi [220] determined the exact values of the total edge Fibonacci irregularity strength of fans, double fans, umbrella, and wheels.

In [88] Agustin, Dafik, Marsidi, and Albirri introduced a natural extension of the

notation of the total H-irregularity strength of graphs by considering the evaluation of the weight that is not only for each edge but also the weight on each subgraph H of G. They say a total α-labeling is a *total H-irregular α-labeling* of the graph G if for a subgroup H of G, the total H-weights $W(H) = \sum_{v\in V(H)} f(v) + \sum_{e\in E(H)} f(e)$ are distinct. The minimum α for which the graph G has a total H-irregular α-labeling is called the *total H-irregularity strength* of G, denoted by $tHs(G)$. They study the tHs of shackles and amalgamations of any graphs and their bounds. In [313] Ashraf, Bača, Semaničová-Feňovčíková, and Siddiqui determined the exact value of the total (vertex, edge) H-irregularity strengths for the ladders and fan graphs. Mursalina and Susanti [2573] gave the exact total H-irregularity strength values for balloons, double balloons, and double balloon ladder graphs. [2573] new

The sqaure chain graph has vertex set $\{a_i : 1 \le i \le n+1\} \cup \{b_i, c_i : 1 \le i \le n\}$ and edge set $\{a_ib_i, a_ic_i, a_{i+1}b_i, a_{i+1}c_i : 1 \le i \le n\}$. Susanto [3818] gave a complete characterization of H-irregular labeling behavior in square chain graphs. [3818] new Bača, Lascsákov'a, and Semaničová-Feňovčíková [387] determined vertex H-irregularity strength of join product graphs. [387] new Labane, Bouchemakh, and Semaničová-Feňovčíková [2084] gave the exact values for the vertex H-irregularity strength of diamond graphs. [2084] new Naeem, Siddiqui, Bača, Semaničová-Feňovčíková, and Ashraf [2632] determined the exact values of H-irregularity strength for prisms, antiprisms, triangular ladders, diagonal ladders, wheels and gear graphs. [2632] new Wahyujati and Susanti [4118] investigated comb product graphs and established exact values for their total H-irregularity strength. [4118] new Shulhany, Rukmayadi, Maharani, and Syaifara [3574] studied the total H-irregularity strength for generalized butterfly graphs and eclipse graphs. [3574] new

The notion of an irregular labeling of an Abelian group Γ was introduced Anholcer, Cichacz and Milanič in [250]. They defined a Γ-*irregular labeling* of a graph G with no isolated vertices as an assignment of elements of an Abelian group Γ to the edges of G in such a way that the sums of the weights of the edges at each vertex are distinct. The *group irregularity strength* of G, denoted $s_g(G)$, is the smallest integer s such that for every Abelian group Γ of order s there exists Γ-irregular labeling of G. They proved that if G is connected, then $s_g(G) = n+2$ when $\cong K_{1,3^{2q+1}-2}$ for some integer $q \ge 1$; $s_g(G) = n+1$ when $n \equiv 2 \pmod 4$ and $G \ncong K_{1,3^{2q+1}-2}$ for any integer $q \ge 1$; and $s_g(G) = n$ otherwise. Moreover, Anholcer and Cichacz [246] showed that if G is a graph of order n with no component of order less than 3 and with all the bipartite components having both color classes of even order. Then $s_g(G) = n$ if $n \equiv 1 \pmod 2$; $s_g(G) = n+1$ if $n \equiv 2 \pmod 4$; and $s_g(G) \le n+1$ if $n \equiv 0 \pmod 4$. In [249] Anholcer, Cichacz, Jura, Marczyk presented some upper bound on group the irregularity strength for all graphs. Moreover, they gave the exact values and bounds on $s_g(G)$ for disconnected graphs without a star as a component. Anholcer, Cichacz, and Przybyło [254] proved that $s_g(G) \le 2n$. They also considered locally irregular labelings where only sums of adjacent vertices are required to be distinct. For the corresponding graph invariant they proved the general upper bound: $\Delta(G)+\mathrm{col}(G)-1$ (where $\mathrm{col}(G)$ is the coloring number of G) in the case when the identity element is not used as an edge label, and a slightly worse one if additionally the identity is forbidden as the sum of labels around a vertex. In both cases they provided a sharp

upper bound for trees and a constant upper bound for planar graphs. In [851] Cichacz and Krupińska gave a bound and exact values of the group irregular strength for graphs without small stars as components.

Marzuki, Salman, and Miller [2457] introduced a new irregular total k-labeling of a graph G called *total irregular total k-labeling*, denoted by $ts(G)$, which is required to be at the same time both vertex and edge irregular. They gave an upper bound and a lower bound of $ts(G)$; determined the total irregularity strength of cycles and paths; and proved $ts(G) \geq \max\{tes(G), tvs(G)\}$. For $n \geq 3$, Ramdani and Salman [3115] proved $ts(S_n \times P_2) = n+1$; $ts((P_n + P_1) \times P_2) = \lceil (5n+1)/3 \rceil$, $ts(P_n \times P_2) = n$; and $ts(C_n \times P_2) = n$. In [3116] Ramdani, Salman, and Assiyatun prove that for a regular graph G $ts(mG) \leq m(ts(G)) - \lfloor (m-1)/2 \rfloor$, $ts(mC_n) = \lceil (mn+2)/3 \rfloor$ for $n \equiv 3 \bmod 3$, and $ts(m(C_n \times P_2) = mn+1$.

In [3905] Titawanno, Tilukay, Leleury, Taihuttu, and Loves examined the total irregular total labeling of double star and triple star graphs, which is a class of caterpillar graphs, and obtained the exact values of the total disjoint irregularity strength of those graphs corresponding to the lower bound for the total irregularity strength of graph G, $ts(G)$. Tilukay and Salman [3903] introduced a new labeling method as follows. A *total disjoint irregular total labeling* of a graph $G = (V, E)$, with the edge weight set $W(E)$ and the vertex weight set $W(V)$ on G, is a totally irregular total labeling for which $W(E) \cap W(V) = \emptyset$. Its parameter is called the *total disjoint irregularity strength* of G, denoted by $ds(G)$. They gave a lower bound for the values of $ds(G)$ and determined the exact value of $ds(G)$ for path graphs, cycle graphs, star graphs, and complete graphs. [3905] new [3903] new

In [3117] Ramdani, Salman, Assiyatun, Semaničová-Feňovčíková, and Bača estimate the upper bound of the total irregularity strength of graphs and determine the exact value of the total irregularity strength for three families of graphs. In [1776] Jeyanthi and Sudha determined the total irregularity strength of double fans DF_n $(n \geq 3)$, double triangular snakes DT_p $(p \geq 3)$, joint-wheel graphs WH_n $(n \geq 3)$, and $P_m + \overline{K_m}$ $(m \geq 3)$. In addition, they show that these graphs admit totally irregular total k-labeling and they determined the exact ts value for each. In [1540] Indriati, Widodo, Wijayanti, and Sugeng provided the total irregularity strength of some caterpillars.

In [3900] Tilukay, Salman, and Persulessy proved that fans, wheels, triangular books, friendship graphs, double fans DF_n, $(n \geq 3)$, double triangular snakes DT_p $(p \geq 3)$, joint-wheel graphs, $P_m + K_m$ $(m \geq 3)$, stars, double-stars, and caterpillars (see also [1543]) are totally irregular total graphs. Ramdani, Salman, Assiyatun, Semaničová-Feňovčíková, and Bača [3117] proved that for any positive integer $n \geq 2$, $\mathrm{ts}(K_{n,n}) = \lceil (n^2+2)/3 \rceil$. Tilukay, Taihuttu, Salman, Rumlawang, and Leleury [3901] proved that $K_{m,n}$ is a totally irregular total graph for any positive integer m and n. In [3900] Tilukay, Salman, and Persulessy proved that fans, wheels, triangular books, friendship graphs are totally irregular total graphs, double fans DF_n, $(n \geq 3)$, double triangular snakes DT_p, $(p \geq 3)$, joint-wheel graphs, $P_m + K_m$ $(m \geq 3)$, stars, double-stars, and caterpillars (see also [1543]).

In [2613] Muthgu Guru Packiam defines a *face irregular total k-labeling* f from $V \cup E \cup F$ to $\{1, 2, \ldots, k\}$ of a 2-connected plane graph $G(V, E, F)$ as a labeling of vertices and edges

such that different faces have different weights. The minimum k for which a plane graph G has a face irregular total k-labeling is called *total face irregularity strength* of G and is denoted by $tfs(G)$. He provides a bound on this parameter and the exact values for shell graphs and a family of planar graphs consisting of an even number of 5-sided faces and one external infinite face. In [385] Bača, Lascsková, Naseem, and Semaničová-Feňovčíková estimate the lower and upper bounds of the entire face irregularity strength for the disjoint union of multiple copies of a plane graph and prove the sharpness of the lower bound in two cases. Tilukay, Salman, Ilwaru, and Rumlawang [3902] estimated the bounds of $tfs(G)$ and prove that the lower bound is sharp for cycles, books with m polygonal pages, and wheels.

For a graph G, Tanna, Ryan, and Semaničová-Feňovčíková [3864] define a k-labeling ρ as a labeling such that the edges of G are labeled with $\{1, 2, \ldots, k_e\}$ and the vertices of G are labeled with $\{0, 2, \ldots, 2k_v\}$, where $k = \max\{k_e, 2k_v\}$. The labeling ρ is called an *edge irregular reflexive k-labeling* if distinct edges have distinct weights, where the edge weight is defined as the sum of the label of that edge and the labels of its endpoints. The smallest k for which such a labeling exists is called the *reflexive edge strength* of G. They gave exact values for the reflexive edge strength for prisms, wheels, baskets (graphs obtained by removing a spoke from a wheel), and fans. Guirao, Ahmad, Siddiqui, and Ibrahim [1343] investigated the exact value of the reflexive edge strength for disjoint union of s isomorphic copies of generalized Peterson graphs. Zhang, Ibrahim, ul Haq Bokhary, and Siddiqui [4319] provided exact value of the reflexive edge strength for disjoint union of gears and prisms. Bača, Irfan, Ryan, Semaničová-Feňovčíková, and Tanna [377] determined the exact value of the reflexive edge strength for cycles, the Cartesian product of two cycles, and for join graphs of the path and cycle with $2K_2$. Wang, Khan, Ibrahim, Bonyah, Siddiqui, and Khalid [4125] determined the reflexive edge irregularity strength for the Cartesian product of a path and a cycle. In [4270] Yoong, Hasni, Lau, Asim, and Ahmad obtained the exact reflexive edge strength for antiprisms, two kinds of convex polytopes, and $C_n \odot P_m$ $(n \geq 3, m \geq 2)$. Yoong, Hasni, Lau, and Irfan [4271] provided the exact value of the reflexive edge strength of three classes of plane graphs. In [553] Basher examined two types of eight-sided and four-faced or six-sided and four-faced planar maps that have an edge irregular reflexive k-labeling. He gave the precise value of the reflexive edge strength for these two classes. In [376] Bača, Irfan, Ryan, Semaničová-Feňovčíková, and Tanna provided the precise values of the reflexive edge strength for the generalized friendship graphs $f_{n,m}$ for $n = 3, 4, 5, m \geq 1$ and made a conjecture of the value for the remaining cases. In [377] they determined the exact value of the reflexive edge strength for cycles, the Cartesian product of two cycles, and for join graphs of the path and cycle with $2K_2$. Bača, Kovář, and Semaničová-Feňovčíková [1546] provided the precise value of the reflexive edge strength of $C_n \times P_m$ where $n \geq 3$ and $m \geq 2$. Basher [559] determined the exact value of the reflexive edge strength of toroidal polyhexes. Mato and Wichianpaisarn [2458] provided the exact value of the reflexive edge strength of cycles plus one edge that contains a triangle. In [1943] Khan, Akram, Ishtiaq, Garayev, and Popa investigated tight lower bound for non-planar prisms, cross prisms, cross-particle modified graphs, and cross-particle glowing graphs under edge-irregular reflexive k-labelings.

The reflexive edge strength of the following graphs have been determined by various authors: banana tree graphs $B_{2,n}$ and $B_{3,n}$ [2717]; the categorical product $P_a \times P_b$ for $a \geq 3$ and $b \geq 3$ [1941]; palm trees $C_p - B_{q,r}$ with $p = 3, q = 2, 3$ and $r \geq 3$ [1805]; $P_n \odot K_1$ and $P_n \odot P_2$ [1538]; sun graphs, and the corona graphs of cycles and null graphs that have one or two vertices [3456]; $P_n \odot K_1$ and $P_n \odot P_2$ [1538]; and the triangular chain graphs mC_3-path for $m \geq 2$ [725]

In [1537] Indriati and Azzahra determined the edge irregular reflexive k-labelings of Mongolian tents and double quadrilateral snakes. Zalzabila, Indriati, and Martini [4304] investigated edge irregular reflexive k-labelings of alternate triangular snakes and the double alternate quadrilateral snakes. In [4303] Zalsa, Indriati, and Martini determined the edge irregular reflexive k-labeling for dumbbell graphs and the corona of open ladders and the null graph.

A *vertex irregular reflexive k-labeling* of a graph G is total k-labeling such that for every two distinct vertices the sums of labels of edges that are incident to each vertex and the vertex label itself are distinct. The *reflexive vertex strength* of a graph G is a minimum k such that G has a vertex irregular reflexive k-labeling. Agustin, Iman Utoyo, Dafik, and Venkatachalam [92] determined the exact value of reflexive vertex strength of ladders and $K_{2,n}$.

The *reflexive vertex strength* of a graph G is the minimum k for which G has a vertex irregular reflexive k-labeling. In [93] Agustin, Susilowati, Dafik, Cangul, and Mohanapriya determine the exact value of the reflexive vertex strength of regular graphs and regular-like graphs.

A total k-labeling of a graph is a function f_e from the edge set to $\{1, 2, \ldots, k_e\}$ and a function f_v from the vertex set to $\{0, 2, 4, \ldots, 2k_v\}$, where $k = \max k_e, 2k_v$. A *distance irregular reflexive k-labeling* of graph G is a total k-labeling if for every two distinct vertices u and u' of G, $wt(u) \neq wt(u')$, where $wt(u) = f_v(u) +$ the sum of $f_e(uv)$ over all edges uv of G. The minimum k for graph G that has a distance irregular reflexive k-labeling is called the *distance reflexive strength* of the graph G. In [89] Agustin, Dafik, Mohanapriya, Marsidi, and Cangul determine the lower bound of distance reflexive strength of any graph and the exact value of distance reflexive strength of paths, stars, and friendship graphs.

Recall that an *edge-covering* of G is a family of subgraphs $H_1, H_2, \ldots, H_t$ such that each edge of $E(G)$ belongs to at least one of the subgraphs $H_i, i = 1, 2, \ldots, t$. In this case we say that G admits an $(H_1, H_2, \ldots, H_t)$-(edge) covering. If every subgraph H_i is isomorphic to a given graph H, we say that G admits an *H-covering*. Motivated by the irregularity strength and the edge irregularity strength of a graph G, Ashraf, Bača, Kimáková, and Semaničová-Feňovčíková [309] introduced two new parameters, edge (vertex) H-irregularity strengths, as the natural extensions of the parameters s(G) and es(G) as follows. Let G be a graph admitting an H-covering. For the subgraph H of G under the edge k-labeling β from $E(G)$ to $\{1, 2, \ldots, k\}$, the associated H-weight is defined as $\text{wt}_\beta(H) = \sum \beta(e)$ over all edges e. An edge k-labeling β is called an *H-irregular edge k-labeling* of the graph G if for every two different subgraphs H' and H'' isomorphic to H we have $wt_\beta(H') \neq wt_\beta(H'')$. The edge *H-irregularity strength* of a graph G, denoted by ehs(G, H), is the smallest integer k such that G has an H-irregular edge k-labeling.

Ibrahim, Gulzar, Fazil, and Azhar [1479] compute the exact value of edge H-irregularity strength of hexagonal and octagonal grid graphs. Ashraf et al. define the *vertex H-irregularity strength* of a graph G, $\mathrm{vhs}(G,H)$, analogously. They estimate the bounds of the parameters $\mathrm{ehs}(G,H)$ and $\mathrm{vhs}(G,H)$ and determine the exact values of the edge (vertex) H-irregularity strength for paths, ladders, and fans in order to prove the sharpness of lower bounds of these parameters. Nisviasari, Dafik, and Agustin [2690] determined the total H-irregularity strength of triangular ladders when H is a windmill or triangular ladder. For a graph G the minimum k for which G has an H-irregular reflexive k-labeling is called the *reflexive H strength* of graph G and is denoted by $rHs(G)$. In [2436] Marsidil, Agustin, Dafik, Rahman, and Sullystiawati initiated the study the lower bound of the reflexive H strength of graphs and the reflexive H strength of flowers, $Shack(C_t, v, n)$, and books, where H is isomorphic to C_3, C_t, and C_4, respectively. In [3770] Sullystiawati, Marsidi, Putra, and Agustin determined lower bounds of the reflexive H-strength of paths, wheels, double fans, triangular ladders, and ladders.

In [311] Ashraf, Bača, Semaničová-Feňovčíková, and Saputro determined the exact value of the cycle-irregularity strength of ladders and fan graphs. Ashraf, Bača, Lascsáková, and Semaničová-Feňovčíková [310] estimated the bounds for the total H-irregularity strength of a graph and determined the exact values of the total H-irregularity strength for paths ladders and fans. Ashrafa, Bača, Semaničová-Feňovčíková, and Shabbirc [312] investigated the total (respectively, edge and vertex) G-irregularity strengths of the graphs that contains exactly n subgraphs isomorphic to G. Ahmad, Bača, and Semaničová-Feňovčíková [118] determined the exact values of $\mathrm{ehs}(G, C_4)$ for grids and generalized prisms.

In [105] Ahmad, Al-Mushayt, and Bača define a vertex k-labeling ϕ of a graph G from $V(G)$ to $\{1, 2, \ldots, k\}$ to be *edge irregular k-labeling* if for every two distinct edges e and f, there is $w_\phi(e) \neq w_\phi(f)$, where the weight of an edge $e = xy$ is $w_\phi(xy) = \phi(x) + \phi(y)$. The minimum k for which the graph G has an edge irregular k-labeling is called the *edge irregularity strength* of G, denoted by $es(G)$. They estimated the bounds of the edge irregularity and determined its exact values for paths, cycles, stars, double stars and $P_m \times P_n$. Tarawneh, Hasni, and Ahmad [3866] determined the exact value of the edge irregularity strength of the corona product of graphs with paths. Tarawneh, Hasni, and Ahmad [3867] determine the exact value of edge irregularity strength of corona graphs $C_n \odot mK_1$ ($m \geq 2$). Ahmad [98] determined the exact value of $\mathrm{es}(C_n \odot K_1)$. In [115] Ahmad, Bača, and Nadeen determined the exact value of the edge irregularity strength for several classes of Toeplitz graphs. In [3457] Seyam and Nadeem generalized results of Ahmad, Bača, and Nadeem and provided the exact values of the edge irregularity strength for some generalized classes of Toeplitz graphs.

Tarawneh, Hasni, Siddiqui, and Asim [3871] determined the exact value of edge irregularity strength of disjoint union of zigzag graphs, grids, and generalized sun graphs. Ahmed, Omar, and Bača [135] obtained estimations of the edge irregularity strength of graphs and determined the precise values for paths, stars, double stars, and the Cartesian product of two paths. Suparta and Suharta [3810] determined the edge irregularity strength of the chain graph mK_3-path for $m \geq 3 \pmod 4$ and the chain graph $C[C_n^{(m)}]$

for $n = 0 \pmod 4$ and provided bounds for the edge irregularity strength of $P_m + \overline{K_n}$ for $m, n \geq 3$. In [4323] Zhang, Mehmood, Rehman, Hussain, and Zhang provided the exact values of the edge irregularity strengths of C_4-snakes, complete m-partite graphs, and the middle graphs of paths and cycles.

A *chain graph* $C[B_1, B_2, \ldots, B_n]$ is a graph with blocks $B_1, B_2, \ldots, B_n$ such that B_i and B_{i+1} have a common vertex in such a way that the block-cut vertex graph is a path. Ahmad, Gupta, and Simanjuntak [124] prove the following: $es(C[C_4^{(n)}]) = 2n + 1$; if H_m is an mK_3-path, then $es(H_m)$ has lower bound $\lceil \frac{3m+3}{2} \rceil$ and upper bound $2m + 1$; $es(mK_4\text{-path}) = 3m + 2$; and $es(K_{1,n} + \overline{K_1}) = n + 2$ for $n \geq 3$. They obtained bounds for $es(P_m + \overline{K_n})$ and determined that the edge irregularity strength of a graph obtained by joining the vertex of degree m in $K_{1,m}$ to each vertex in $K_{1,n}$, and the vertex of degree n in $K_{1,n}$ to each vertex in $K_{1,m}$ is $m + n + 2$. They posed the open problems of determining $es(mK_3\text{-path})$, $es(mK_n\text{-path})$ $(m \geq 2)$ and $n \geq 5$, and $es(P_m + \overline{K_n})$ for $n \geq 1$ and $m \geq 7$. Tarawneh, Hasni, and Asim [3870] determined the exact value of edge irregularity strength for disjoint union of a star graph and the subdivision of a star graph.

The *strong product* of graphs G_1 and G_2 has as vertices the pairs (x, y) where $x \in V(G_1)$ and $y \in V(G_2)$. The vertices (x_1, y_1) and (x_2, y_2) are adjacent if either x_1x_2 is an edge of G_1 and $y_1 = y_2$ or if $x_1 = x_2$ and y_1y_2 is an edge of G_2. For $m, n \geq 2$ Ahmad, Bača, Bashir, Siddiqui [113] proved that the total edge irregular strength of the strong product of P_m and P_n is $\lceil 4(mn + 1)/3 \rceil - (m + n)$. Al-Mushayt [196] determined the edge irregularity strength of Cartesian product of a star and P_2 and a cycle and P_2, and the strong product of path P_n with P_2. Conjectures for the exact value of $K_{1,n} \times P_m$ and $C_n \times P_m$ are stated. Bača and Siddiqui [428] determine the exact value of the total edge irregularity strength of the strong product of any two cycles.

An edge $e \in \overline{G}$ is called a *total positive edge* or *total negative edge* or *total stable edge* of G if $\text{tvs}(G + e) > \text{tvs}(G)$ or $\text{tvs}(G + e) < \text{tvs}(G)$ or $\text{tvs}(G + e) = \text{tvs}(G)$, respectively. If all edges of $\overline{G}$ are total stable (total negative) edges of G, then G is called a *total stable* (*total negative*) graph. Otherwise G is called a *total mixed* graph.

Muthu Guru Packiam and Kathiresan [2744] showed that $K_{1,n}$ $n \geq 4$, and the disjoint union of $t \geq 2$ copies of K_3 are total negative graphs and that the disjoint union of $t \geq 2$ copies of P_3 is a total mixed graph.

For a simple graph G with no isolated edges and at most one isolated vertex Anholcer [244] calls a labeling $w : E(G) \to \{1, 2, \ldots, m\}$ *product-irregular*, if all product degrees $pd_G(v) = \prod_{e \ni v} w(e)$ are distinct. Analogous to the notion of irregularity strength the goal is to find a product-irregular labeling that minimizes the maximum label. This minimum value is called the *product irregularity strength* of G and is denoted by $ps(G)$. He provides bounds for the product irregularity strength of paths, cycles, Cartesian products of paths, and Cartesian products of cycles. In [245] Anholcer gives the exact values of $ps(G)$ for $K_{m,n}$ where $2 \leq m \leq n \leq (m + 2)(m + 1)/2$, some families of forests including complete d-ary trees, and other graphs with $d(G) = 1$. Darda and Hujdurović [897] proved that $ps(X) \leq |V(X)| - 1$ for any graph X with more than 3 vertices and gave a connection between the product irregularity strength and the multidimensional multiplication table problem. Skowronek-Kaziów [3627] proves that for the complete graphs $ps(K_n) = 3$. In

2023 Bensmail, Hocquard, Lajou, and Sopena [619] proved that the product version of the 1-2-3 conjecture, raised by Skowronek-Kazió [3627] in 2012, that for every connected graph with order at least 3, there is an assignment of the labels 1, 2, 3 to the edges in such a way that no two adjacent vertices are incident to the same product of labels.

In [9] Abdo and Dimitrov introduced the total irregularity of a graph. For a graph G, they define $\mathrm{irr}_t(G) = (1/2)\sum_{u,v\in V} |d_G(u)-d_G(v)|$, where $d_G(w)$ denotes the vertex degree of the vertex w. For G with n vertices they proved $\mathrm{irr}_t(G) \leq (1/12)(2n^3 - 3n^2 - 2n + 3)$. For a tree G with n vertices they prove $\mathrm{irr}_t(G) \leq (n-1)(n-2)$ and equality holds if and only if $G \approx S_n$. You, Yang, and You [4272] determined the graph with the maximal total irregularity among all unicyclic graphs.

Inspired by the concept of distant chromatic numbers Przybylo [3050] calls a labeling f from the edges of a graph G to $\{1, 2, 3, \ldots, k\}$ *r-distant irregular*, if for every vertex v, the weights of the set of all vertices that are at distance less than or equal to r from v are pairwise distinct, where the weight of the vertex is the sum of the labels of the edges that are incident with that vertex. The minimum k for which there exists an r-distant irregular labeling of G is called *r-distant irregularity strength* of G and is denoted by $s_r(G)$. Muthu Guru Packiam, Manimaran, and Thuraiswamy [2615] proved the following: $s_1(P_n) = 2$ for $n = 3, 4, 5$; $s_1(P_n) = 3$ if $n > 5$; $s_1(C_n) = 3$; $s_1(K_{m,n}) = s(K_{m,n})$; $s_1(F_n) = s(F_n) = \lceil (n+1)/3 \rceil$ for $n > 2$; $s_1(K_{m,n}) = 3$ when $1 < n/2 \leq m < n$; $s_1(P_n \times K_2) = 3$; $s_1(C_n \times K_2) = 3$; $s_1(K_{m,n}) = 3$ when $1 < n/2 \leq m < n$; and provide the exact value for $s_1(P_m \odot \overline{K_n})$ for all m and n. They also prove that if G is d-regular with n vertices, then $s_1(G) = s(G) \leq \lceil n/2 \rceil + 1$ for $d \geq n/2$. Susanto, Wijaya, Purnama, and Slamin [3825] derived a new lower bound of distance irregularity strength for graphs with pendent vertices. They also determined the distance irregularity strength of some families of disjoint unions of paths, suns, helms and friendships graphs. In [618] Bensmail proved that determining the distant irregularity strengths are NP-hard problems.

In a preprint [4187] Wijayanti, Noor, Indriati, Alghofari, and Slamin defined a *distance vertex irregular total k-labeling* of a simple undirected graph $G(V, E)$, as a function $f : V(G) \cup E(G) \to \{1, 2, \ldots, k\}$ such that for every pair of distinct vertices $u, v \in V(G)$ the weights of u and v are distinct. The *weight* of vertex $v \in V(G)$ is the sum of the labels of vertices in neighborhood of v and the labels of all incident edges to v. The *total distance vertex irregularity strength* of G (denoted by $tdis(G)$) is the minimum k for which G has a distance vertex irregular total k-labeling. In [4188] Wijayanti et al. generalized to distance D where D is any subset of $\{1, 2, \ldots, \mathrm{diam}(G)\}$. Wijayanti, Noor, Indriati, Alghofari [4188] determined the exact total distance vertex irregularity strength of fan and wheel graphs for $D = \{1\}$. In [4186] Wijayanti, Noor, Indriati, Alghofari, and Slamin determined the exact value of the total distance vertex irregularity strength of $\overline{K_m} \odot \overline{K_n}, C_n \odot \overline{K_1}$, and $P_n \odot \overline{K_1}$. Wijayanti, Wijayanti, and Surono [4190] provided a lower bound and determined the value of total distance vertex irregularity strength of the hairy cycle C_m^n and proved that for $m = 2, 3, 4$ and $n \geq 5$ odd, the hairy cycle C_m^n graphs admits an D-distance vertex irregular total k-labelings with total distance vertex irregularity strength $tdis(C_m^n) \geq \lceil (mn+1)/2 \rceil$.

An *inclusive distance vertex irregular labeling* of a graph G is an assignment of elements

of the set $\{1, 2, \ldots, k\}$ to the vertices of G such that the sums of numbers assigned to the closed neighborhoods of all vertices are distinct. The minimum number k for which there exists an inclusive distance vertex irregular labeling of G is denoted by $\widehat{dis}(G)$. Bača, Semaničová-Feňovčíková, Slamin, and Sugeng [423] establish a lower bound for the inclusive distance vertex irregularity strength for any graph and determine the exact value of this parameter for several families of graphs. Cichacz, Gömrlich, and Semaničová-Feňovčíková [841] prove that for a simple graph G on n vertices in which no two vertices have the same closed neighborhood is $\widehat{dis}(G) \leq n^2$.

A mapping $\phi : V(G) \to \{1, 2, \ldots, k\}$ of a simple graph G is said to be an *non-inclusive distance vertex irregular k-labeling* of G if the sums of labels of vertices in the open neighborhood of every vertex are distinct. The minimum k for which G has a non-inclusive distance vertex irregular k-labeling is called a *non-inclusive*distance irregularity strength and is denoted by dis(G)). Susanto, Wijaya, Slamin, and Semaničová-Feňovčíková [3826] developed new lower bounds for non-inclusive distance irregularity strength of graphs and proved that such bounds are sharp for some classes of graphs with pendent vertices. They also discussed some properties for non-inclusive distance irregularity strength of trees.

Susanto, Wijaya, Sudarsana, and Slamin [3827] studied non-inclusive and inclusive distance irregularity strength for the join product of two given graphs. They gave bounds on non-inclusive and inclusive distance irregularity strength for $G+H$ and showed that in some conditions, the bounds are achievable. They determined the exact value of dis($G + K_1$) for any given graph G and developed an algorithm for calculating the upper bound on inclusive distance irregularity strength of complete multipartite graphs.

Susanto, Betistiyan, Halikin, and Wijaya [3819] derived new lower bounds on inclusive distance irregularity strength for arbitrary graph and showed that the bounds are sharp for two stars $2K_{1,n}$ and three stars $3K_{1,n}$. For a integer $m \geq 2$ and non-trivial connected graph G of order p and size q, the *m-shadow of a graph* G, denoted by $D_m(G)$, is the graph obtained by taking m copies of G, say G_1, G_2,..., G_m, and joining each vertex u in G_i, $i = 1, 2, \ldots, m-1$, to the neighbors of the corresponding vertex u' in G_{i+1}. In [4183] Wijaya, Aulia, Halikin, and Kusbudiono showed that dis$(T_{n,2}) = (n^2 + 2)/2$, where $T_{n,2}$ is a complete n-ary tree to level two.

For a graph G and an integer $m \geq 3$, the *closed m-shadow of a graph* G, denoted by $CD_m(G)$, is a graph obtained from the m-shadow of G by joining an edge from every vertex u in G_1 to the neighbors of the corresponding vertex u' in G_m.

Inayah, Susanto and Semaničová-Feňovčíková [1535] proved the following: for every non-trivial connected graph G of order p and size q, and integers m, t with $2 \leq t \leq m-1$, $D_m(G)$ is super (a, d)-$D_t(G)$-antimagic for $d \in \{0, 2, \ldots, p+3q\}$ when $p+q$ is even and for $d \in \{1, 3, \ldots, p+3q\}$ when $p+q$ is odd; for every non-trivial connected graph G of order p and size q, and integers m, t with $2 \leq t \leq m-1$ and $\gcd(m, t) = 1$, $CD_m(G)$ is super (a, d)-$D_t(G)$-antimagic for $d \in \{0, 2, \ldots, p+q\}$ when $p+q$ is even and for $d \in \{1, 3, \ldots, p+q\}$ when $p+q$ is odd; for every non-trivial connected graph G of order p and size q, and for any integer $m \geq 3$, $CD_m(G)$ is super (a, d)-$D_m(G)$-antimagic for $d \in \{0, 2, \ldots, 2q\}$.

A natural modification of an irregular labeling is a modular irregular labeling introduced by Muthugurupackiam and Ramya [2576] in 2018. They call an edge k-labeling

$\phi : E(G) \to \{1, 2, \ldots, k\}$ of positive integers to the edges of a graph G of order n a *modular irregular* labeling of G if the weight function $\theta : V(G) \to Z_n$ defined by $\theta(v) = wt\phi(v) = \sum \phi(uv)$ over all $u \in N(v)$ is bijective. The *modular irregularity strength* is defined as the minimum k for which G has a modular irregular labeling. If there is no such labeling for a graph then its modular irregularity strength is defined as ∞. They proved the modular irregularity of the tadpole graph and double-cycle graph. Bača, Muthugurupackiam, Kathiresan, and Ramya in [414] gave a lower bound of the modular irregularity strength, and the exact values for paths, cycles, stars, triangular graphs, and gear graphs are determined. In [381] Bača, Kimáková, Lascsáková, and Semaničová-Feňovčikově determine the exact value of the irregularity strength and the modular irregularity strength of fan graphs. Bačca, Imran, and Semaničově-Feňovčcíkově showed the existence of an irregular labeling scheme that proves the exact value of the irregularity strength of wheels. By modifying this irregular mapping in six cases, they obtained labelings that determine the exact value of the modular irregularity strength of wheels as a natural modification of the irregularity strength. Ali, Bača, Lascsaková, Semaničová-Feňovčıková, ALoqaily, and Mlaiki [183] obtained estimations on the modular total vertex irregularity strength, and gave the precise values of this invariant for certain graphs. In [3725] Sugeng, Barack, Hinding, and Simanjuntak constructed a modular irregular labeling and determined its modular irregularity strength of regular double-star graphs and friendship graphs. Haryeni, Awanis, Bača, and Semaničová-Feňovčikově [1380] estimated the bounds of the (modular) edge irregularity strength for $P_n + \overline{K_m}$ and $C_n + \overline{K_m}$ and determined the corresponding exact value of the (modular) edge irregularity strength for some fans and wheels in order to prove the sharpness of the presented bounds. In [3729] Sugeng, John, Lawrence, Anwar, Bača, and Semaničová-Feňovčíková determined the exact values of the modular irregularity strength of rose graphs, daisy graphs and sunflower graphs.

In [271] Ariyanti, Barack, John, and Bustamam investigated the modular irregularity strength of the corona product of a circulant graph with a null graph of order p. They also showed the modular irregularity strength of $G \odot C_3$ is $n + 1$ and that the modular irregularity strength of $G \odot (P_2 \cup P_1)$ is $3n/2$. They further proved that the modular irregularity strength of $G \odot P_5$ is $\lceil (5n+1)/3 \rceil$. Sugeng, Sofyan, Sy, Hinding, and Simanjuntak [3739] determined the modular irregularity strength of Dutch windmills and obtained the exact value of the modular irregularity strength of the vertex amalgamation and the comb product of paths with cycle-related graphs. [271] new [3739] new

In [1242] Gayathri and Saravanakumar define a *k-edge-weighting* of a graph $G(V, E)$ as a map $\phi : E(G)$ to $\{1, 2, 3, \ldots, k\}$. They say that a k-edge weighting of G is an *equitable irregular* if for every pair of adjacent vertices u and v in G the sum of edge-weights appearing on the edges incident at the vertex v under ϕ and the sum of edge-weights appearing on the edges incident at the vertex u under ϕ differ by at most 1. They proved that polar grid and Mongolian tent are equitable irregular.

In [3821] Susanto, Simanjuntak, and Baskoro introduced the notion of D-irregular labelings that generalizes both non-inclusive and inclusive d-distance irregular labelings of graphs as follows. Let $G(V, E)$ be a graph, D a set of distances, and k a positive integer.

A mapping ϕ from V to $\{1, 2, \ldots, k\}$ is called a *D-irregular k-labeling* of G if every two distinct vertices have distinct weights, where the weight of a vertex x is defined as the sum of labels of vertices whose distance from x belongs to D. The smallest integer k for which G admits a D-irregular labeling is the *D-irregularity strength* of G and denoted it by $s_D(G)$. They established several fundamental properties on D-irregularity strength for arbitrary graphs and determined the exact D-irregularity strength for families of graphs with small diameter or small maximum degree.

7.17 Geometric Labelings

If a and r are positive integers at least 2, we say a (p, q)-graph G is (a, r)-*geometric* if its vertices can be assigned distinct positive integers such that the value of the edges obtained as the product of the endpoints of each edge is $\{a, ar, ar^2, \ldots, ar^{q-1}\}$. Hegde [1405] has shown the following: no connected bipartite graph, except the star, is (a, a)-geometric where a is a prime number or square of a prime number; any connected (a, a)-geometric graph where a is a prime number or square of a prime number, is either a star or has a triangle; $K_{a,b},\ 2 \le a \le b$ is (k, k)-geometric if and only if k is neither a prime number nor the square of a prime number; a caterpillar is (k, k)-geometric if and only if k is neither a prime number nor the square of a prime number; $K_{a,b,1}$ is (k, k)-geometric for all integers $k \ge 2$; C_{4t} is (a, a)-geometric if and only if a is neither a prime number nor the square of a prime number; for any positive integers t and $r \ge 2$, C_{4t+1} is (r^{2t}, r)-geometric; for any positive integer t, C_{4t+2} is not geometric for any values of a and r; and for any positive integers t and $r \ge 2$, C_{4t+3} is (r^{2t+1}, r)-geometric. Hegde [1407] has also shown that every T_p-tree and the subdivision graph of every T_p-tree are (a, r)-geometric for some values of a and r (see Section 3.2 for the definition of a T_p-tree). He conjectures that all trees are (a, r)-geometric for some values of a and r.

Hegde and Shankaran [1417] prove: a graph with an α-labeling (see §3.1 for the definition) where m is the fixed integer that is between the endpoints of each edge has an (a^{m+1}, a)-geometric for any $a > 1$; for any integers m and n both greater than 1 and m odd, mP_n is (a^r, a)-geometric where $r = (mn + 3)/2$ if n is odd and (a^r, a)-geometric where $r = (m(n + 1) + 3)/2$ if n is even; for positive integers $k > 1, d \ge 1$, and odd n, the generalized closed helm (see §5.3 for the definition) $CH(t, n)$ is (k^r, k^d)-geometric where $r = (n - 1)d/2$; for positive integers $k > 1, d \ge 1$, and odd n, the generalized web graph (see §5.3 for the definition) $W(t, n)$ is (k^r, a)-geometric where $a = k^d$ and $r = (n - 1)d/2$; for positive integers $k > 1, d \ge 1$, the generalized n-crown $(P_m \times K_3) \odot K_{1,n}$ is (a, a)-geometric where $a = k^d$; and $n = 2r + 1$, $C_n \odot P_3$ is (k^r, k)-geometric.

If a and r are positive integers and r is at least 2 Arumugan, Germina, and Anadavally [278] say a (p, q)-graph G is *additively* (a, r)-*geometric* if its vertices can be assigned distinct integers such that the value of the edges obtained as the sum of the endpoints of each edge is $\{a, ar, ar^2, \ldots, ar^{q-1}\}$. In the case that the vertex labels are nonnegative integers the labeling is called *additively* $(a, r)*$-*geometric*. They prove: for all a and r every tree is additively $(a, r)*$-geometric; a connected additively (a, r)-geometric graph is either a tree or unicyclic graph with the cycle having odd size; if G is a connected unicyclic

graph and not a cycle, then G is additively (a,r)-geometric if and only if either a is even or a is odd and r is even; connected unicyclic graphs are not additively $(a,r)*$-geometric; if a disconnected graph is additively (a,r)-geometric, then each component is a tree or a unicyclic graph with an odd cycle; and for all even a at least 4, every disconnected graph for which every component is a tree or unicyclic with an odd cycle has an additively (a,r)-geometric labeling.

Vijayakumar [4092] calls a graph G (not necessarily finite) *arithmetic* if its vertices can be assigned distinct natural numbers such that the value of the edges obtained as the sum of the endpoints of each edge is an arithmetic progression. He proves [4091] and [4092] that a graph is arithmetic if and only if it is (a,r)-geometric for some a and r.

7.18 Strongly Multiplicative Graphs

Beineke and Hegde [613] call a graph with p vertices *strongly multiplicative* if the vertices of G can be labeled with distinct integers $1, 2, \ldots, p$ such that the labels induced on the edges by the product of the end vertices are distinct. They prove the following graphs are strongly multiplicative: trees; cycles; wheels; K_n if and only if $n \leq 5$; $K_{r,r}$ if and only if $r \leq 4$; and $P_m \times P_n$. They then consider the maximum number of edges a strongly multiplicative graph on n vertices can have. Denoting this number by $\lambda(n)$, they show: $\lambda(4r) \leq 6r^2; \lambda(4r+1) \leq 6r^2+4r; \lambda(4r+2) \leq 6r^2+6r+1$; and $\lambda(4r+3) \leq 6r^2+10r+3$. Adiga, Ramaswamy, and Somashekara [75] give the bound $\lambda(n) \leq n(n+1)/2 + n - 2 - \lfloor (n+2)/4 \rfloor - \sum_{i=2}^{n} i/p(i)$ where $p(i)$ is the smallest prime dividing i. For large values of n this is a better upper bound for $\lambda(n)$ than the one given by Beineke and Hegde. It remains an open problem to find a nontrivial lower bound for $\lambda(n)$.

Seoud and Zid [3412] prove the following graphs are strongly multiplicative: wheels; rK_n for all r and n at most 5; rK_n for $r \geq 2$ and $n = 6$ or 7; rK_n for $r \geq 3$ and $n = 8$ or 9; $K_{4,r}$ for all r; and the corona of P_n and K_m for all n and $2 \leq m \leq 8$. In [3390] Seoud and Mahran [3390] give some necessary conditions for a graph to be strongly multiplicative.

In Kanani and Chhaya [1823] and [1824] prove the following graphs are strongly multiplicative: the total graph, splitting graph, and shadow graph of paths; triangular snakes; splitting graphs of stars and bistars, the degree splitting graph of the bistars $B_{n,n}$, and restricted square graph $B^2_{m,n}$. In [1827] and [1828] Kanani and Chhaya prove the following graphs are strongly multiplicative: helms, flowers, fans, friendship graphs, bistars, gears, double triangular snakes, double fans, double wheels, snakes, double alternate quadrilateral snakes, double quadrilateral snakes, braid graphs, and triangular ladders.

Germina and Ajitha [1243] (see also [49]) prove that $K_2 + \overline{K_t}$, quadrilateral snakes, Petersen graphs, ladders, and unicyclic graphs are strongly multiplicative. Acharya, Germina, and Ajitha [49] have shown that $C_k^{(n)}$ (see §2.2 for the definition) is strongly multiplicative and that every graph can be embedded as an induced subgraph of a strongly multiplicative graph. Germina and Ajitha [1243] define a graph with q edges and a strongly multiplicative labeling to be *hyper strongly multiplicative* if the induced edge labels are $\{2, 3, \ldots, q+1\}$. They show that every hyper strongly multiplicative graph has exactly one nontrivial component that is either a star or has a triangle and every graph can be

embedded as an induced subgraph of a hyper strongly multiplicative graph.

Vaidya, Dani, Vihol, and Kanani [3973] prove that the arbitrary supersubdivisions of tree, K_{mn}, $P_n \times P_m$, $C_n \odot P_m$, and C_n^m are strongly multiplicative. Vaidya and Kanani [3979] prove that the following graphs are strongly multiplicative: a cycle with one chord; a cycle with twin chords (that is, two chords that share an endpoint and with opposite endpoints that join two consecutive vertices of the cycle; the cycle C_n with three chords that form a triangle and whose edges are the edges of two 3-cycles and a $n-3$-cycle. duplication of an vertex in cycle (see §2.7 for the definition); and the graphs obtained from C_n by identifying of two vertices v_i and v_j where $d(v_i, v_j) \geq 3$. In [3982] the same authors prove that the graph obtained by an arbitrary supersubdivision of path, a star, a cycle, and a tadpole (that is, a cycle with a path appended to a vertex of the cycle.

Krawec [2004] calls a graph G on n edges *modular multiplicative* if the vertices of G can be labeled with distinct integers $0, 1, \ldots, n-1$ (with one exception if G is a tree) such that the labels induced on the edges by the product of the end vertices modulo n are distinct. He proves that every graph can be embedded as an induced subgraph of a modular multiplicative graph on prime number of edges. He also shows that if G is a modular multiplicative graph on prime number of edges p then for every integer $k \geq 2$ there exist modular multiplicative graphs on p^k and kp edges that contain G as a subgraph. In the same paper, Krawec also calls a graph G on n edges *k-modular multiplicative* if the vertices of G can be labeled with distinct integers $0, 1, \ldots, n+k-1$ such that the labels induced on the edges by the product of the end vertices modulo $n+k$ are distinct. He proves that every graph is k-modular multiplicative for some k and also shows that if $p = 2n+1$ is prime then the path on n edges is $(n+1)$-modular multiplicative. He also shows that if $p = 2n+1$ is prime then the cycle on n edges is $(n+1)$-modular multiplicative if there does not exist $\alpha \in \{2, 3, \ldots, n\}$ such that $\alpha^2 + \alpha - 1 \equiv 0 \bmod p$. He concludes with four open problems. In [2005] Krawec shows that every graph is a subgraph of a modular multiplicative graph. He also defines k-modular multiplicative graphs and proves that certain families of paths and cycles admit such a labeling.

In [2657] Nasir, Idrees, Sadiq, Farooq, Kanwal, and Imran show that the join of K_1 and a triangular ladder TL_n ($n \geq 3$), umbrella graphs, and generalized Petersen graphs $GP(n, k)$ for ($n \leq 3$) and $1 \leq k \leq n/2$, double combs, and sunflower planar graphs (obtained by appending one edge to each vertex of the rim of a wheel) admit strongly multiplicative labelings.

7.19 Line-graceful Labelings

Gnanajothi [1276] has defined a concept similar to edge-graceful. She calls a graph with n vertices *line-graceful* if it is possible to label its edges with $0, 1, 2, \ldots, n$ such that when each vertex is assigned the sum modulo n of all the edge labels incident with that vertex the resulting vertex labels are $0, 1, \ldots, n-1$. A necessary condition for the line-gracefulness of a graph is that its order is not congruent to 2 (mod 4). Among line-graceful graphs are (see [pp. 132–181] [1276]) P_n if and only if $n \not\equiv 2 \pmod 4$; C_n if and only if $n \not\equiv 2 \pmod 4$; $K_{1,n}$ if and only if $n \not\equiv 1 \pmod 4$; $P_n \odot K_1$ (combs) if and only if n is even; $(P_n \odot K_1) \odot K_1$

if and only if $n \not\equiv 2 \pmod 4$; (in general, if G has order n, $G \odot H$ is the graph obtained by taking one copy of G and n copies of H and joining the ith vertex of G with an edge to every vertex in the ith copy of H); mC_n when mn is odd; $C_n \odot K_1$ (crowns) if and only if n is even; mC_4 for all m; complete n-ary trees when n is even; $K_{1,n} \cup K_{1,n}$ if and only if n is odd; odd cycles with a chord; even cycles with a tail; even cycles with a tail of length 1 and a chord; graphs consisting of two triangles having a common vertex and tails of equal length attached to a vertex other than the common one; the complete n-ary tree when n is even; trees for which exactly one vertex has even degree. She conjectures that all trees with $p \not\equiv 2 \pmod 4$ vertices are line-graceful and proved this conjecture for $p \leq 9$.

Gnanajothi [1276] has investigated the line-gracefulness of several graphs obtained from stars. In particular, the graph obtained from $K_{1,4}$ by subdividing one spoke to form a path of even order (counting the center of the star) is line-graceful; the graph obtained from a star by inserting one vertex in a single spoke is line-graceful if and only if the star has $p \not\equiv 2 \pmod 4$ vertices; the graph obtained from $K_{1,n}$ by replacing each spoke with a path of length m (counting the center vertex) is line-graceful in the following cases: $n = 2$; $n = 3$ and $m \not\equiv 3 \pmod 4$; and m is even and $mn + 1 \equiv 0 \pmod 4$.

Gnanajothi studied graphs obtained by joining disjoint graphs G and H with an edge. She proved such graphs are line-graceful in the following circumstances: $G = H$; $G = P_n, H = P_m$ and $m+n \not\equiv 0 \pmod 4$; and $G = P_n \odot K_1$, $H = P_m \odot K_1$ and $m+n \not\equiv 0 \pmod 4$.

In [3986] and [3987] Vaidya and Kothari proved the following graphs are line graceful: fans F_n for $n \not\equiv 1 \pmod 4$; W_n for $n \not\equiv 1 \pmod 4$; bistars $B_{n,n}$ if and only if for $n \equiv 1, 3 \pmod 4$; helms H_n for all n; $S'(P_n)$ for $n \equiv 0, 2 \pmod 4$; $D_2(P_n)$ for $n \equiv 0, 2 \pmod 4$; $T(P_n)$, $M(P_n)$, alternate triangular snakes, and graphs obtained by duplication of each edge of P_n by a vertex are line graceful graphs.

In 2012, [1793] Jones, Kolasinski, and Zhang independently introduced the notion of *modular edge-graceful*, which was first given in 1991 by Gnanajothi [1276] with the name line-graceful graph. Gnanajothi conjectured that every tree of order $n \neq 2 \pmod 4$ is line-graceful (that is, modular edge-graceful). Jones, Kolasinski, and Zhang showed that this conjecture is true and furthermore that a nontrivial connected graph of order n is modular edge-graceful if and only if $n \neq 2 \pmod 4$. They defined the *modular edge-gracefulness* $\mathrm{meg}(G)$ of a connected graph G order $n \geq 3$ as the smallest integer $k \geq n$ for which there exists an edge labeling $f : E(G) \to Z_k$ such that the induced vertex labeling $f' : V(G) \to Z_k$ is one to-one. Jones, Kolasinski, and Zhang is showed that $\mathrm{meg}(G) = n + 1$ for every connected graph G that is not modular edge-graceful. [1793] new

In [1794] Jones and Zhang called a modular edge-graceful labeling f of G *nowhere-zero* if $f(e) \neq 0$ for all e in $E(G)$ and, called such a graph *nowhere zero modular edge-graceful* . They showed showed that a connected graph G of order $n \geq 3$ is nowhere-zero modular edge-graceful if and only if $n \neq 2 \pmod 4$, G is not K_3, and G is not a star of even order. For a connected graph G of order $n \geq 3$, the smallest integer $k \geq n$ for which there exists an edge labeling $f : E(G) \to Z_k \setminus \{0\}$ such that the induced vertex labeling f' is one-to-one is referred to as the *nowhere-zero modular edge-gracefulness* of G and [1794] new

determined this number for every connected graph of order at least 3.

7.20 Graceful k-colorings

A *graceful k-coloring* of a graph G is a proper vertex coloring f of G that assigns colors from a set to the vertices of G that induces a proper edge coloring of G, where the color for each edge uv is $|f(u) - f(v)|$. The minimum positive integer k for which G has a graceful k-coloring is called the *graceful chromatic number* of G and is denoted by $\chi_g(G)$. It follows from the definitions of gracefulness and graceful chromatic number of a graph G with n vertices that $\chi(G) \leq 2^{n-1}$. In [2762] Pantoja, Dantas, and Luiz proved that $\chi_g(G) \leq 2.5\Delta^2 - 1.5\Delta + 1$. Bi, Byers, and Zhang [657] obtained upper bounds for the graceful chrommatic numbers of various families of trees. In [656] Bi, Byers, English, Laforge, and Zhang determined the graceful chromatic numbers of cycles, wheels, and caterpillars and gave an upper bound for the graceful chromatic number of trees in terms of their maximum degrees. For $\Delta > 2$ and h at most 4, English and Znang [1036] determined the graceful chromatic numbers of a class of rooted trees T, h with maximum degree Δ and height h. They further showed that for $\Delta > 2$, $\lim_{n\to\infty} \chi_g(T_{\Delta,h} = \lceil 5\Delta/3 \rceil$.

In [2763] Pantoja, Dantas, and Luiz showed that under certain restrictions, a tree with maximum degree Δ has graceful chromatic number $\Delta + 1$ or $\Delta + 2$. Moreover, they identify a family of lobsters with maximum degree 3 that have graceful chromatic number of 4 and a family of lobsters that have graceful chromatic number of 5, thereby showing that both lower and upper bounds for this parameter are attained within this class of graphs. Furthermore, they gave a polynomial-time algorithm to determine whether a given tree with maximum degree three has a graceful chromatic number of 4. Finally, they proved that the Graceful 4-Coloring Problem is NP-complete. [2763] new

7.21 k-sequential Labelings

In 1981 Bange, Barkauskas, and Slater [471] defined a *k-sequential* labeling f of a graph $G(V, E)$ as one for which f is a bijection from $V \cup E$ to $\{k, k+1, \ldots, |V \cup E| + k - 1\}$ such that for each edge xy in E, $f(xy) = |f(x) - f(y)|$. This generalized the notion of *simply sequential* where $k = 1$ introduced by Slater. Bange, Barkauskas, and Slater showed that cycles are 1-sequential and if G is 1-sequential, then $G + K_1$ is graceful. Hegde and Shetty [1416] have shown that every T_p-tree (see §4.4 for the definition) is 1-sequential. In [3633], Slater proved: K_n is 1-sequential if and only if $n \leq 3$; for $n \geq 2$, K_n is not k-sequential for any $k \geq 2$; and $K_{1,n}$ is k-sequential if and only if k divides n. Acharya and Hegde [54] proved: if G is k-sequential, then k is at most the independence number of G; P_{2n} is n-sequential for all n and P_{2n+1} is both n-sequential and $(n+1)$-sequential for all n; $K_{m,n}$ is k-sequential for $k = 1, m$, and n; $K_{m,n,1}$ is 1-sequential; and the join of any caterpillar and $\overline{K_t}$ is 1-sequential. Acharya [40] showed that if $G(E, V)$ is an odd graph with $|E| + |V| \equiv 1$ or 2 (mod 4) when k is odd or $|E| + |V| \equiv 2$ or 3 (mod 4) when k is even, then G is not k-sequential. Acharya also observed that as a consequence of results of Bermond, Kotzig, and Turgeon [632] we have: mK_4 is not k-sequential for any

k when m is odd and mK_2 is not k-sequential for any odd k when $m \equiv 2$ or 3 (mod 4) or for any even k when $m \equiv 1$ or 2 (mod 4). He further noted that $K_{m,n}$ is not k-sequential when k is even and m and n are odd, whereas $K_{m,k}$ is k-sequential for all k. Acharya points out that the following result of Slater's [3634] for $k = 1$ linking k-graceful graphs and k-sequential graphs holds in general: A graph is k-sequential if and only if $G + v$ has a k-graceful labeling f with $f(v) = 0$. Slater [3633] also proved that a k-sequential graph with p vertices and $q > 0$ edges must satisfy $k \leq p - 1$. Hegde [1402] proved that every graph can be embedded as an induced subgraph of a simply sequential graph. In [40] Acharya conjectured that if G is a connected k-sequential graph of order p with $k > \lfloor p/2 \rfloor$, then $k = p-1$ and $G = K_{1,p-1}$ and that, except for $K_{1,p-1}$, every tree in which all vertices are odd is k-sequential for all odd positive integers $k \leq p/2$. In [1402] Hegde gave counterexamples for both of these conjectures.

In [1414] Hegde and Miller prove the following: for $n > 1$, K_n is k-sequentially additive if and only if $(n, k) = (2, 1), (3, 1)$ or (3,2); $K_{1,n}$ is k-sequentially additive if and only if k divides n; caterpillars with bipartition sets of sizes m and n are k-sequentially additive for $k = m$ and $k = n$; and if an odd-degree (p, q)-graph is k-sequentially additive, then $(p+q)(2k+p+q-1) \equiv 0$ (mod 4). As corollaries of the last result they observe that when m and n are odd and k is even $K_{m,n}$ is not k-sequentially additive and if an odd-degree tree is k-sequentially additive then k is odd.

In [3387] Seoud and Jaber proved the following graphs are 1-sequentially additive: graphs obtained by joining the centers of two identical stars with an edge; $S_n \cup S_m$ if and only if nm is even; $C_n \odot \overline{K_m}$; $P_n \odot \overline{K_m}$; kP_3; graphs obtained by joining the centers of k copies of P_3 to each vertex in $\overline{K_m}$; and trees obtained from K by replacing each edge by a path of length 2 when $n \equiv 0, 1$ (mod 4). They also determined all 1-sequentially additive graphs of order 6.

7.22 IC-colorings

For a subgraph H of a graph G with vertex set V and a coloring f from V to the natural numbers define $f_s(H) = \Sigma f(v)$ over all $v \in H$. The coloring f is called an IC-*coloring* if for any integer k between 1 and $f_s(G)$ there is a connected subgraph H of G such that $f_s(H) = k$. The IC-*index* of a graph G, $M(G)$, is $\max\{f_s |\ f_s$ is an IC-coloring of $G\}$. Salehi, Lee, and Khatirinejad [3252] obtained the following: $M(K_n) = 2^n - 1$; for $n \geq 2$, $M(K_{1,n}) = 2^n + 2$; if Δ is the maximum degree of a connected graph G, then $M(G) \geq 2^\Delta + 2$; if $ST(n; 3^n)$ is the graph obtained by identifying the end points of n paths of length 3, then $ST(n; 3^n)$ is at least $3^n + 3$ (they conjecture that equality holds for $n \geq 4$); for $n \geq 2$, $M(K_{2,n}) = 3 \cdot 2^n + 1$; $M(P_n) \geq (2 + \lfloor n/2 \rfloor)(n - \lfloor n/2 \rfloor) + \lfloor n/2 \rfloor - 1$; for $m, n \geq 2$, the IC-index of the double star $DS(m, n)$ is at least $(2^{m-1} + 1)(2^{n-1} + 1)$ (they conjecture that equality holds); for $n \geq 3$, $n(n + 1)/2 \leq M(C_n) \leq n(n - 1) + 1$; and for $n \geq 3$, $2^n + 2 \leq M(W_n) \leq 2^n + n(n-1) + 1$. They pose the following open problems: find the IC-index of the graph obtained by identifying the endpoints of n paths of length b; find the IC-index of the graph obtained by identifying the endpoints of n paths; and find the IC-index of $K_{m,n}$. Shiue and Fu [3561] completed the partial results by Penrice [2820]

Salehi, Lee, and Khatirinejad [3252] by proving $M(K_{m,n}) = 3 \cdot 2^{m+n-2} - 2^{m-2} + 2$ for any $2 \leq m \leq n$.

7.23 Minimal k-rankings

A *k-ranking* of a graph is a labeling of the vertices with the integers 1 to k inclusively such that any path between vertices of the same label contains a vertex of greater label. The *rank number of a graph* G, $\chi_r(G)$, is the smallest possible number of labels in a ranking. A k-ranking is *minimal* if no label can be replaced by a smaller label and still be a k-ranking. The concept of the rank number arose in the study of the design of very large scale integration (VLSI) layouts and parallel processing (see [906], [2232] and [3361]). Ghoshal, Laskar, and Pillone [1271] were the first to investigate minimal k-rankings from a mathematical perspective. Laskar and Pillone [2093] proved that the decision problem corresponding to minimal k-rankings is NP-complete. It is HP-hard even for bipartite graphs [926]. Bodlaender, Deogun, Jansen, Kloks, Kratsch, Müller, Tuza [672] proved that the rank number of P_n is $\chi_r(P_n) = \lfloor \log_2(n) \rfloor + 1$ and satisfies the recursion $\chi_r(P_n) = 1 + \chi_r(P_{\lceil (n-1)/2 \rceil})$ for $n > 1$. The following results are given in [926]: $\chi_r(S_n) = 2$; $\chi_r(C_n) = \lfloor \log_2(n-1) \rfloor + 2$; $\chi_r(W_n) = \lfloor \log_2(n-3) \rfloor + 3 (n > 3)$; $\chi_r(K_n) = n$; the complete t-partite graph with n vertices has ranking number $n+1$ - the cardinality of the largest partite set; and a split graph with n vertices has ranking number $n+1$ - the cardinality of the largest independent set (a *split graph* is a graph in which the vertices can be partitioned into a clique and an independent set.) Wang proved that for any graphs G and H $\chi_r(G + H) = \min\{|V(G)| + \chi_r(H), |V(H) + \chi_r(G)\}$.

In 2009 Novotny, Ortiz, and Narayan [2719] determined the rank number of P_n^2 from the recursion $\chi_r(P_n^2) = 2 + \chi_r(P(_{\lceil (n-2)/2 \rceil})$ for $n > 2$. They posed the problem of determining $\chi_r(P_m \times P_n)$ and $\chi(P_n^k)$. In 2009 [204] and [203] Alpert determined the rank numbers of P_n^k, C_n^k, $P_2 \times C_n$, $K_m \times P_n$, $P_3 \times P_n$, Möbius ladders and found bounds for rank numbers of general grid graphs $P_m \times P_n$. About the same time as Alpert and independently, Chang, Kuo, and Lin [767] determined the rank numbers of P_n^k, C_n^k, $P_2 \times P_n, P_2 \times C_n$. Chang et al. also determined the rank numbers of caterpillars and proved that for any graphs G and H $\chi_r(G[H]) = \chi_r(H) + |V(H)|(\chi_r(G) - 1)$.

In 2010 Jacob, Narayan, Sergel, Richter, and Tran [1564] investigated k-rankings of paths and cycles with pendent paths of length 1 or 2. Among their results are: for any caterpillar G $\chi_r(P_n) \leq \chi_r(G) \leq \chi_r(P_n) + 1$ and both cases occur; if $2^m \leq n \leq 2^{m+1}$ then for any graph G obtained by appending edges to an n-cycle we have $m+2 \leq \chi_r(G) \leq m+3$ and both cases occur; if G is a lobster with spine P_n then $\chi_r(P_n) \leq \chi_r(G) \leq \chi_r(P_n) + 2$ and all three cases occur; if G a graph obtained from the cycle C_n by appending paths of length 1 or 2 to any number of the vertices of the cycle then $\chi_r(P_n) \leq \chi(G) \leq \chi(P_n) + 2$ and all three cases occur; and if G the graph obtained from the comb obtained from P_n by appending one path of length m to each vertex of P_n then $\chi_r(G) = \chi_r(P_n) + \chi_r(P_{m+1}) - 1$.

Sergel, Richter, Tran, Curran, Jacob, and Narayan [3414] investigated the rank number of a cycle C_n with pendent edges, which they denote by CC_n, and call a *caterpillar cycle*. They proved that $\chi(CC_n) = \chi_r(C_n))$ or $\chi(CC_n) = \chi_r(C_n)) + 1$ and showed that both cases

occur. A *comb tree*, denoted by $C(n,m)$, is a tree that has a path P_n such that every vertex of P_n is adjacent to an end vertex of a path P_m. In the comb tree $C(n,m)$ $(n \geq 3)$ there are 2 pendent paths P_{m+2} and $n-2$ paths P_{m+1}. They proved $\chi_r(C(n,m)) = \chi_r(P_{m+1}) - 1$. They define a *circular lobster* as a graph where each vertex is either on a cycle C_n or at most distance two from a vertex on C_n. They proved that if G is a lobster with longest path P_n, then $\chi_r(P_n) \leq \chi_r(G) \leq \chi_r(P_n) + 2$ and determined the conditions under which each true case occurs. If G is circular lobster with cycle C_n, they showed that $\chi_r(C_n) \leq \chi_r(G) \leq \chi_r(C_n) + 2$ and determined the conditions under which each true case occurs. An *icicle graph* I_n $(n \geq 3)$ has three pendent paths P_2 and is comprised of a path P_n with vertices $v_1, v_2, \ldots, v_n$ where a path P_{i-1} is appended to vertex v_i. They determine the rank number for icicle graphs.

Richter, Leven, Tran, Ek, Jacob, and Narayan [3172] define a *reduction* of a graph G as a graph G_S^* such that $V(G_S^*) = V(G) \setminus S$ and, for vertices u and v, uv is an edge of G_S^* if and only if there exists a $u-v$ path in G with all internal vertices belonging to S. A vertex *separating set* of a connected graph G is a set of vertices whose removal disconnects G. They define a *bent ladder* $BL_n(a,b)$ as the union of ladders L_a and L_b (where $L_n = P_n \times P_2$) that are joined at a right angle with a single L_2 so that $n = a+b+2$. A staircase ladder SL_n is a graph with $n-1$ subgraphs $G_1, G_2, \ldots, G_{n-1}$ each of which is isomorphic to C_4. (They are ladders with a maximum number of bends.) Richter et al. [3172] prove: $\chi_r(BL_n(a,b)) = \chi_r(L_n) - 1$ if $n = 2^k - 1$ and $a \equiv 2$ or $3 \pmod 4$ and is equal to $\chi_r(L_n)$ otherwise; $\chi_r(SL_n) = \chi_r(L_{n+1})$ if $n = 2^k + 2^{k-1} - 2$ for some $k \geq 3$ and is equal to $\chi_r(L_n)$ otherwise; and for any ladder L_n with multiple bends, the rank number is either $\chi_r(L_n)$ or $\chi_r(L_n) + 1$).

The *arank number* of a graph G is the maximum value of k such that G has a minimal k-ranking. Eyabi, Jacob, Laskar, Narayan, and Pillone [1059] determine the arank number of $K_n \times K_n$, and investigated the arank number of $K_m \times K_n$.

7.24 Set Graceful and Set Sequential Graphs

The notions of set graceful and set sequential graphs were introduced by Acharaya in 1983 [41]. A graph is called *set graceful* if there is an assignment of nonempty subsets of a finite set to the vertices and edges of the graph such that the value given to each edge is the symmetric difference of the sets assigned to the endpoints of the edge, the assignment of sets to the vertices is injective, and the assignment to the edges is bijective. A graph is called *set sequential* if there is an assignment of nonempty subsets of a finite set to the vertices and edges of the graph such that the value given to each edge is the symmetric difference of the sets assigned to the endpoints of the edge and the assignment of sets to the vertices and the edges is bijective. The following has been shown: P_n $(n > 3)$ is not set graceful [1406]; C_n is not set sequential [55]; C_n is set graceful if and only if $n = 2^m - 1$ [1408] and [41]; K_n is set graceful if and only if $n = 2, 3$ or 6 [2542]; K_n $(n \geq 2)$ is set sequential if and only if $n = 2$ or 5 [1408]; $K_{a,b}$ is set sequential if and only if $(a+1)(b+1)$ is a positive power of 2 [1408]; a necessary condition for $K_{a,b,c}$ to be set sequential is that a, b, and c cannot have the same parity [1406]; $K_{1,b,c}$ is not set sequential

when b and c even [1408]; $K_{2,b,c}$ is not set sequential when b and c are odd [1406]; no theta graph is set graceful [1406]; the complete nontrivial n-ary tree is set sequential if and only if $n+1$ is a power of 2 and the number of levels is 1 [1406]; a tree is set sequential if and only if it is set graceful [1406]; the nontrivial plane triangular grid graph G_n is set graceful if and only if $n = 2$ [1408]; every graph can be embedded as an induced subgraph of a connected set sequential graph [1406]; every graph can be embedded as an induced subgraph of a connected set graceful graph [1406], every planar graph can be embedded as an induced subgraph of a set sequential planar graph [1408]; every tree can be embedded as an induced subgraph of a set sequential tree [1408]; and every tree can be embedded as an induced subgraph of a set graceful tree [1408]. Hegde conjectures [1408] that no path is set sequential. Hegde's conjecture [1409] that every complete bipartite graph that has a set graceful labeling is a star was proved by Vijayakumar [4093]. Shahida and Sunitha [3463] prove that the concept of set-gracefulness is equivalent to topologically set-gracefulness in trees and almost all finite trees are not set-graceful. Using this they characterize topologically set-graceful stars and topologically set-graceful paths. In [46] Acharya and Germina survey results on set-valuations of graphs and give open problems and conjectures.

Germina, Kumar, and Princy [1241] prove: if a (p, q)-graph is set-sequential with respect to a set with n elements, then the maximum degree of any vertex is $2^{n-1} - 1$; if G is set-sequential with respect to a set with n elements other than K_5, then for every edge uv with $d(u) = d(v)$ one has $d(u) + d(v) < 2^{n-1} - 1$; $K_{1,p}$ is set-sequential if and only if p has the form $2^{n-1} - 1$ for some $n \geq 2$; binary trees are not set-sequential; hypercubes Q_n are not set-sequential for $n > 1$; wheels are not set-sequential; and uniform binary trees with an extra edge appended at the root are set-graceful and set graceful.

Vijayakumar [4093] and Gyri, Balister, and Schelp [366] proved that if a complete bipartite graph G has a set-graceful labeling, then it is a star. Abhishek [11] described a method for constructing a set-graceful bipartite graph of arbitrarily large order and size beginning with a set-graceful bipartite graph. Acharya, Germina, Princy, and Rao [51] proved that $K_{1,m,n}$ is set-graceful if and only if $m = 2^s - 1$ and $n = 2^t - 1$ and almost all graphs are not set-graceful. In [12] Abhishek surveys results on set-valued graphs. Many open problems and conjectures are included.

Acharya [41] has shown: a connected set graceful graph with q edges and $q+1$ vertices is a tree of order $p = 2^m$ and for every positive integer m such a tree exists; if G is a connected set sequential graph, then $G + K_1$ is set graceful; and if a graph with p vertices and q edges is set sequential, then $p+q = 2^m - 1$. Acharya, Germina, Princy, and Rao [51] proved: if G is set graceful, then $G \cup \overline{K_t}$ is set sequential for some t; if G is a set graceful graph with n edges and $n+1$ vertices, then $G + \overline{K_t}$ is set graceful if and only if m has the form $2^t - 1$; $P_n + \overline{K_m}$ is set graceful if $n = 1$ or 2 and m has the form $2^t - 1$; $K_{1,m,n}$ is set graceful if and only if m has the form $2^t - 1$ and n has the form $2^s - 1$; $P_4 + \overline{K_m}$ is not set graceful when m has the form $2^t - 1$ $(t \geq 1)$; $K_{3,5}$ is not set graceful; if G is set graceful, then graph obtained from G by adding for each vertex v in G a new vertex v' that is adjacent to every vertex adjacent to v is not set graceful; and $K_{3,5}$ is not set graceful.

Acharya, Germina, Abhishek, and Slater [48] prove C_m is set-graceful if and only if $m = (4^n - 1)/3$; mK_2 is set-sequential if and only if $m = (4^n - 1)/3$; and, for $r + s = 2^{n-1}$ the bistar $B(r, s)$ is set-sequential if and only if r and s are odd. They also prove that connected planar graphs with an even number of faces, regular polyhedrons, and cacti containing an odd number of cycles are not set-sequential.

Abhishek [11] proved that if G is a set-sequential bipartite graph and H is $2k$-set-sequential, then $4^k G \cup H$ is set-sequential. As a corollary, he gets mP_3 is set-sequential if and only if $m = (16^n - 1)/5$. Abhishek and Agustine [14] characterized the set-sequential caterpillars of diameter four and give a necessary condition for a graph to be set-sequential. Abhishek [13] characterized the set-sequential caterpillars of diameter five. Golowich and Kim [1289] showed that all caterpillars T of diameter k such that $k \leq 18$ or $|V(T)| \geq 2^{k-1}$ are set-sequential, where T has only odd-degree vertices and $|V(T)| = 2^{n-1}$ for some n. They also provided a new method of recursively constructing set-sequential trees.

In [2482] Mehra and Puneet introduce a topological integer additive set-labeling of signed graphs as follows. Let $S = (V, E, s)$ be a signed graph with corresponding graph $G = (V, E)$ and the signature function s. Here, G is an integer additive set-labeled graph having an injective function $f : V(G) \to P(X) - \{\emptyset\}$ that produces another injective function $g_f : E(G) \to P(X) - \{\emptyset, \{0\}\}$ defined by $g_f(uv) = f(u) + f(v)$ for every edge uv, where X is the subset of non-negative integers, $P(X)$ is its power set, and the signature function defined as $s : E(G) \to \{+, -\}$ is such that $s(uv) = -1^{|f(u)+f(v)|}$ for all edges uv. If $f(V(G)) \cup \{\emptyset\}$ forms a topology on X then the signed graph S is called a *topological integer additive set-labeled signed graph* (T-IASL). They proved the following graphs have T-IASL labelings: paths, stars, double stars, tadpoles, and graphs obtained by identifying an end of a path with the center of a star.

7.25 Vertex Equitable Graphs

Given a graph G with q edges and a labeling f from the vertices of G to the set $\{0, 1, 2, \ldots, \lceil q/2 \rceil\}$ define a labeling f^* on the edges by $f^*(uv) = f(u) + f(v)$. If for all i and j and each vertex the number of vertices labeled with i and the number of vertices labeled with j differ by at most one and the edge labels induced by f^* are $1, 2, \ldots, q$, Lourdusamy and Seenivasan [2342] call a f a *vertex equitable* labeling of G. They proved the following graphs are vertex equitable: paths, bistars, combs, n-cycles for $n \equiv 0$ or 3 (mod 4), $K_{2,n}$, ${C_3}^t$ for $t \geq 2$, quadrilateral snakes, $K_2 + mK_1$, $K_{1,n} \cup K_{1,n+k}$ if and only if $1 \leq k \leq 3$, ladders, arbitrary super divisions of paths, and n-cycles with $n \equiv 0$ or 3 (mod 4). They further proved that $K_{1,n}$ for $n \geq 4$, Eulerian graphs with n edges where $n \equiv 1$ or 2 (mod 4), wheels, K_n for $n > 3$, triangular cacti with $q \equiv 0$ or 6 or 9 (mod 12), and graphs with p vertices and q edges, where q is even and $p < \lceil q/2 \rceil + 2$ are not vertex equitable. Lourdusamy and Patrick [2335] prove that triangular ladders TL_n, $L_n \odot mK_1$, $Q_n \odot K_1$, $TL_n \odot K_1$, and alternate triangular snakes $A(T_n)$ are vertex equitable graphs. In [64] Acharya, Jain, and Kansal introduced vertex equitable labelings of signed graphs and studied vertex equitable behavior of signed paths, signed stars, and signed complete bipartite graphs $K_{2,n}$.

Jeyanthi and Maheswari [1696] proved that the following graphs have vertex equitable labelings: the square of the bistar $B_{n,n}$; the splitting graph of the bistar $B_{n,n}$; C_4-snakes; connected graphs for in which each block is a cycle of order divisible by 4 (they need not be the same order) and whose block-cut point graph is a path; $C_m \odot P_n$; tadpoles; the one-point union of two cycles; and the graph obtained by starting friendship graphs, $C_{n_1}^{(2)}, C_{n_2}^{(2)}, \ldots, C_{n_k}^{(2)}$ where each $n_i \equiv 0 \pmod 4$ and joining the center of $C_{n_i}^{(2)}$ to the center of $C_{i+1}^{(2)}$ with an edge for $i = 1, 2, \ldots, k-1$. In [1687] Jeyanthi and Maheswari prove that T_p trees, bistars $B(n, n+1)$, $C_n \odot K_m$, ${P_n}^2$, tadpoles, certain classes of caterpillars, and $T \odot \overline{K_n}$ where T is a T_p tree with an even number of vertices are vertex equitable. Jeyanthi and Maheswari [1690] gave vertex equitable labelings for graphs constructed from T_p trees by appending paths or cycles.

In [1686] Jeyanthi and Maheswari show a number of families of graphs have vertex equitable labelings. Their results include: armed crowns $C_m \odot P_n$, shadow graphs $D_2(K_{1,n})$; the graph $C_m * C_n$ obtained by identifying a single vertex of a cycle graph C_m with a single vertex of a cycle graph C_n if and only if $m + n \equiv 0, 3 \pmod 4$; for $n \equiv 0 \pmod 4$ the graph obtained from m copies of $C_n * C_n$ and P_m by joining each vertex of P_m with the cut vertex in one copy of $C_n * C_n$; and graphs obtained by duplicating an arbitrary vertex and an arbitrary edge of a cycle; the total graph of P_n; the splitting graph of P_n; and the fusion of two edges of C_n.

For a graph H with vertices $v_1, v_2, \ldots, v_n$ and n copies of a graph G, $H \mathbin{\hat{o}} G$ is a graph obtained by identifying a vertex u_i of the ith copy of G with a vertex v_i of H for $1 \le i \le n$. The graph $H \mathbin{\tilde{o}} G$ is a graph obtained by joining a vertex u_i of the ith copy of G with a vertex v_i of H by an edge for $1 \le i \le n$. Jeyanthi, Maheswari, and Laksmi prove [1711] that the graphs $L_m \mathbin{\hat{o}} nC_4$, $L_m \mathbin{\tilde{o}} nC_4$, $C_m \mathbin{\tilde{o}} nC_4$, and $P_m \mathbin{\tilde{o}} nC_4$ are vertex equitable graphs. The graph $S^*(G)$ is obtained from a graph G by replacing every edge e of G with $K_{2,m}$ $(m \ge 2)$ with the endpoints of e merged with the two vertices of the 2-vertices part of $K_{2,m}$ after removing the edge e from G. Jeyanthi, Maheswari, and Vijaya Laksmi [1707] prove the graphs $S^*(P_n \cdot K_1)$, $S^*(B(n,n))$, $S^*(P_n \times P_2)$, and $S^*(Q_n)$ of the quadrilateral snake are vertex equitable.

In [1694] Jeyanthi and Maheswari proved the double alternate triangular snake $DA(T_n)$ obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to two new vertices v_i and w_i is vertex equitable; the double alternate quadrilateral snake $DA(Q_n)$ obtained from a path $u_1, u_2, \ldots, u_n$ by joining u_i and u_{i+1} (alternatively) to two new vertices v_i, x_i and w_i, y_i respectively and then joining v_i, w_i and x_i, y_i is vertex equitable; and $NQ(m)$ the n^{th} quadrilateral snake obtained from the path $u_1, u_2, \ldots, u_m$ by joining u_i, u_{i+1} with $2n$ new vertices v_j^i and $w_j^i, 1 \le i \le m-1, 1 \le j \le n$ is vertex equitable. Jeyanthi and Maheswari [1705] prove $DA(T_n) \odot K_1$, $DA(T_n) \odot 2K_1$, $DA(T_n)$, $DA(Q_n) \odot K_1$, $DA(Q_n) \odot 2K_1$, and $DA(Q_n)$ are vertex equitable.

Jeyanthi, Maheswari, and Vijayalaksmi [1706] proved the following graphs are vertex equitable: jewel graphs J_n with vertex set $\{u, v, x, y, u_i : 1 \le i \le n\}$ and edge set $\{ux, uy, xy, xv, yv, uu_i, vu_i : 1 \le i \le n\}$; jelly fish graphs $(JF)_n$ with vertex set $\{u, v, u_i, v_j : 1 \le i \le n, 1 \le j \le n-2\}$ and edge set $\{uu_i : 1 \le i \le n\} \cup \{vv_j : 1 \le j \le n-2\} \cup \{u_{n-1}u_n, vu_n, vu_{n-1}\}$; lobsters constructed from the path $a_1, a_2, \ldots, a_n$ with

vertices a_{i1} and a_{i2} adjacent to a_i for $1 \leq i \leq n$ and pendent vertices $a_{ij}^1, a_{ij}^2, \ldots, a_{ij}^k$ joining a_{ij} for $1 \leq i \leq n$ and $j = 1, 2$; $L_n \odot \overline{K_m}$; and the graph obtained from ladder a L_n and $2n$ copies of $K_{1,m}$ by identifying a non-central vertex of ith copy of $K_{1,m}$ with ith vertex of L_n.

Jeyanthi, Mahewari, and Vijaya Laksmi [1703] prove the following graphs are vertex equitable: graphs obtained by joining a vertex of a cycle to a degree 2 vertex of a comb $(P_n \odot K_1)$ with an edge; path unions of quadrilateral snakes; cycle unions of n copies of mC_4-snakes where $n \equiv 0, 3 \bmod 4$; the graphs obtained from a path $u_1, u_2, \ldots, u_m$ by joining the end points of each edge $u_i u_{i+1}$ to $2n$ isolated vertices v_j^i, w_j^i for $1 \leq m-1,\ 1 \leq j \leq n$, where n is even (the nth quadrilateral snake).

Jeyanthi, Maheswari, and Vijaya Laksmi [1703] prove that subdivisions of double triangular snakes $S(D(Tn))$, subdivisions of double quadrilateral snakes $S(D(Q_n))$, subdivisions of double alternate triangular snakes $S(DA(T_n))$, subdivisions of double alternate quadrilateral snakes $S(DA(Q_n))$, $DA(Q_m) \odot nK_1$, and $DA(T_m) \odot nK_1$ admit vertex equitable labelings.

The *super subdivision* graph $S^*(G)$ of a graph G is the graph obtained from G by replacing every edge uv of G by $K_{2,m}$ (m may vary for each edge) and identifying u and v with the two vertices in $K_{2,m}$ that form the partite set with exactly two members. Jeyanthi, Maheswari, and Vijayalaksmi [1707] prove that super subdivision graphs of $P_n \odot K_1$, bistars $B(n, n)$, $P_n \times P_2$, and quadrilateral snakes are vertex equitable.

For a graph H with vertices $v_1, v_2, \ldots, v_n$ and n copies of a graph G, $H \ \widehat{o}\ G$ is a graph obtained by identifying a vertex u_i of the ith copy of G with a vertex v_i of H for $1 \leq i \leq n$. The graph $H \ \widetilde{o}\ G$ is a graph obtained by joining a vertex u_i of the ith copy of G with a vertex v_i of H by an edge for $1 \leq i \leq n$. Jeyanthi, Maheswari, and Laksmi [1711] prove that the graphs $L_m \ \hat{o}\ nC_4$, $L_m \ \tilde{o}\ nC_4$, $C_m \ \tilde{o}\ nC_4$ and $P_m \ \tilde{o}\ nC_4$ are vertex equitable graphs.

For a graph G with p vertices and q edges and $A = \{1, 3, \ldots, q\}$ if q is odd or $A = \{1, 3, \ldots, q+1\}$ if q is even, Jeyanthi, Maheswari and Vijaya Laksmi [1702] say a vertex labeling f from $V(G)$ to A is an *odd vertex equitable even labeling* if the induced edge labeling f^* defined by $f^*(uv) = f(u) + f(v)$ for all edges uv has the property that for all u and v in A the number of vertices labeled with u and the number of vertices labeled with v differ by at most 1 and the induced edge labels are $2, 4, \ldots, 2q$. A graph that admits odd vertex equitable even labeling is called an *odd vertex equitable even* graph. They show that the following graphs have odd vertex equitable even labelings: paths, graphs obtained by identifying an endpoint of P_m with each vertex of P_n, $K_{1,n}$ if and only if $n = 1$ or 2, $K_{1,n} \cup K_{1,n-2}$ $(n \geq 3)$, $K_{2,n}$, T_p-trees, C_n when $n \equiv 0$ or $1 \bmod 4$, quadrilateral snakes, ladders L_n, $L_n \odot K_1$, and arbitrary super subdivision of paths. They prove that if every edge of a graph G is an edge of a triangle, then G is not an odd vertex equitable even graph. As a corollary of this they get that the following are not odd vertex equitable even graphs: K_n $(n \geq 3)$, wheels, triangular snakes, double triangular snakes, triangular ladders, flower graphs, fans $P_n \odot K_1$ $(n \geq 2)$, double fans $P_n \odot K_2$, $(n \geq 2)$, friendship graphs C_n^3, windmills K_m^n $(m > 3)$, $K_2 + mK_1$, $B_{n,n}^2$, total graphs $T(P_n)$, and composition graphs $P_n[P_2]$. They also show that if G is a (p, q) graph with $p \leq \lceil q/2 \rceil + 1$, then G is

not an odd vertex equitable even graph. Jeyanthi and Maheswari [1697] proved that the subdivision of double triangular snakes and the subdivision of double quadrilateral snakes are odd vertex equitable even graphs.

Lourdusamy and Patrick [2333] proved that $P_n \odot mK_1$, the quadrilateral snake attached to each vertex of path P_n, the super splitting graph $S^*(P_n \odot K_1)$, the super splitting graphs of ladders and the bistars $B_{n,n}$, $B^2_{n,n}$, and the splitting $S'(B_{n,n})$ admit even vertex equitable even labelings. Maheswari and Jeyanthi [2391] have shown the following graphs admit odd vertex equitable even labelings: duplicate graphs of ladders; duplicate graphs of the subdivision of ladders; duplicate graphs of a quadrilaterals; and $P_3 \times P_n$ for odd $n > 2$. Lourdusamy, Wency, and Patrick [2352] prove that $S(D(Q_n)), S(D(T_n)), DA(Q_m) \odot nK_1, DA(T_m) \odot nK_1, S(DA(Q_n))$ and $S(DA(T_n))$ are an even vertex equitable even graphs.

For graphs G_1 and G_2 that graph $G_1\hat{O}G_2$ is obtained from G_1 and $|VG_1)|$ copies of G_2 by identifying one vertex of ith copy of G_2 with ith vertex of G_1. Jeyanthi, Maheswari, and Vijayalakshmi [1713] proved the following graphs have odd vertex equitable even labelings: subdivision graphs of ladders, $L_m\hat{O}P_n$, $L_n \odot \overline{K_m}$ $(m > 1)$, C_n if and only if $n \equiv 0$ or 1 (mod 4), $K_{1,n+k} \cup K_{1,n}$ i if and only if $k = 1, 2$, and $\langle L_n\hat{O}K_{1,m}\rangle$.

Motivated by the concept of vertex equitable labeling first defined by Lourdusamy and Seenivasan in [2342], Lourdusamy, Mary, and Patrick [2331] introduced the concept of even vertex equitable even labeling as follows. Let G be a graph with p vertices and q edges and $A = \{0, 2, 4, \ldots, q+1\}$ if q is odd or $A = \{0, 2, 4, \ldots, q\}$ if q is even. A graph G is said to be an *even vertex equitable even* labeling if there exists a vertex labeling f from $V(G)$ to A that induces an edge labeling f defined by $f^*(uv) = f(u)+f(v)$ for all edges uv such that for all a and b in A, $|v_f(a) - v_f(b)| \leq 1$ and the induced edge labels are $2, 4, \ldots, 2q$, where $v_f(a)$ is the number of vertices v with $f(v) = a$ for $a \in A$. A graph that admits even vertex equitable even labeling is called an *even vertex equitable even* graph. They proved that paths, combs, complete bipartite graphs, cycles, $K_2 + mK_1$, bistars, ladders, $(P_n \times P_2) \odot K_1$, and the subdivision graphs of ladders and bistars $B_{n,n}$ admit an even vertex equitable even labeling. In [2338] Lourdusamy and Patrick proved that $C_m \odot P_n$, C_{4n} and C_{4n+3} with a quadrilateral snake attached to each vertex of the cycle, the graphs obtained by indentifying an edge of C_m and C_n, and the graphs obtained by duplicating an arbitrary vertex and edge of a cycle admit an even vertex equitable even labeling. Lourdusamy, Shobana Mary, and Patrick [2341] proved P^2_n, $S(P_n \odot K_1)$, $S'(P_n)$, $T(P_n)$, graphs obtained by duplication of an edge of a path, quadrilateral snakes, $D(Q_n)$, $A(T_n)$, and $DA(T_n)$ have even vertex equitable even labelings. Lourdusamy and Patrick [2333] proved that $P_n \odot mK_1$, the quadrilateral snake attached to each vertex of path P_n, the super splitting graph $S^*(P_n \odot K_1)$, the super splitting graphs of ladders and the bistars $B_{n,n}$, $B^2_{n,n}$, and the splitting $S'(B_{n,n})$ admit even vertex equitable even labelings.

7.26 Representations of Graphs modulo n

In 1989 Erdős and Evans [1040] defined a *representation* modulo n of a graph G with vertices $v_1, v_2, \ldots, v_r$ as a set $\{a_1, \ldots, a_r\}$ of distinct, nonnegative integers each less than

n satisfying $\gcd(a_i - a_j, n) = 1$ if and only if v_i is adjacent to v_j. They proved that every finite graph can be represented modulo some positive integer. The *representation number*, $\mathrm{Rep}(G)$, is smallest such integer. Obviously the representation number of a graph is prime if and only if a graph is complete. Evans, Fricke, Maneri, McKee, and Perkel [1056] have shown that a graph is representable modulo a product of a pair of distinct primes if and only if the graph does not contain an induced subgraph isomorphic to $K_2 \cup 2K_1$, $K_3 \cup K_1$, or the complement of a chordless cycle of length at least five. Nešetřil and Pultr [2666] showed that every graph can be represented modulo a product of some set of distinct primes. Evans et al. [1056] proved that if G is representable modulo n and p is a prime divisor of n, then $p \geq \chi(G)$. Evans, Isaak, and Narayan [1057] determined representation numbers for specific families as follows (here we use q_i to denote the ith prime and for any prime p_i we use $p_{i+1}, p_{i+2}, \ldots, p_{i+k}$ to denote the next k primes larger than p_i): $\mathrm{Rep}(P_n) = 2 \cdot 3 \cdot \cdots \cdot q_{\lceil \log_2(n-1) \rceil}$; $\mathrm{Rep}(C_4) = 4$ and for $n \geq 3$, $\mathrm{Rep}(C_{2n}) = 2 \cdot 3 \cdot \cdots \cdot q_{\lceil \log_2(n-1) \rceil + 1}$; $\mathrm{Rep}(C_5) = 3 \cdot 5 \cdot 7 = 105$ and for $n \geq 4$ and not a power of 2, $\mathrm{Rep}(C_{2n+1}) = 3 \cdot 5 \cdot \cdots \cdot q_{\lceil \log_2 n \rceil + 1}$; if $m \geq n \geq 3$, then $\mathrm{Rep}(K_m - P_n) = p_i p_{i+1}$ where p_i is the smallest prime greater than or equal to $m - n + \lceil n/2 \rceil$; if $m \geq n \geq 4$, and p_i is the smallest prime greater than or equal to $m - n + \lceil n/2 \rceil$, then $\mathrm{Rep}(K_m - C_n) = q_i q_{i+1}$ if n is even and $\mathrm{Rep}(K_m - C_n) = q_i q_{i+1} q_{i+2}$ if n is odd; if $n \leq m - 1$, then $\mathrm{Rep}(K_m - K_{1,n}) = p_s p_{s+1} \cdots p_{s+n-1}$ where p_s is the smallest prime greater than or equal to $m - 1$; $\mathrm{Rep}(K_m)$ is the smallest prime greater than or equal to m; $\mathrm{Rep}(nK_2) = 2 \cdot 3 \cdot \cdots \cdot q_{\lceil \log_2 n \rceil + 1}$; if $n, m \geq 2$, then $\mathrm{Rep}(nK_m) = p_i p_{i+1} \cdots p_{i+m-1}$, where p_i is the smallest prime satisfying $p_i \geq m$, if and only if there exists a set of $n - 1$ mutually orthogonal Latin squares of order m; $\mathrm{Rep}(mK_1) = 2m$; and if $t \leq (m-1)!$, then $\mathrm{Rep}(K_m + tK_1) = p_s p_{s+1} \cdots p_{s+m-1}$ where p_s is the smallest prime greater than or equal to m. Narayan [2653] proved that for $r \geq 3$ the maximum value for $\mathrm{Rep}(G)$ over all graphs of order r is $p_s p_{s+1} \cdots p_{s+r-2}$, where p_s is the smallest prime that is greater than or equal to $r - 1$. Agarwal and Lopez [84] determined the representation numbers for complete graphs minus a set of stars.

Evans [1055] used matrices over the additive group of a finite field to obtain various bounds for the representation number of graphs of the form nK_m. Among them are $\mathrm{Rep}(4K_3) = 3 \cdot 5 \cdot 7 \cdot 11; \mathrm{Rep}(7K_5) = 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23$; and $\mathrm{Rep}((3q-1)/2)K_q) \leq p_q p_{q+1} \cdots p_{(3q-1)/2)}$ where q is a prime power with $q \equiv 3 \pmod 4$, p_q is the smallest prime greater than or equal to q, and the remaining terms are the next consecutive $(3q-3)/2$ primes; $\mathrm{Rep}(2q-2)K_q) \leq p_q p_{q+1} \cdots p_{(3q-3)/2)}$ where q is a prime power with $q \equiv 3 \bmod 4$, and p_q is the smallest prime greater than or equal to q; $\mathrm{Rep}((2q-2)K_q) \leq p_q p_{q+1} \cdots p_{2q-3}$.

In [2652] Narayan asked for the values of $\mathrm{Rep}(C_{2^k+1})$ when $k \geq 3$ and $\mathrm{Rep}(G)$ when G is a complete multipartite graph or a disjoint union of complete graphs. He also asked about the behavior of the representation number for random graphs. Yahyaei and Katre [4236] gave upper and lower bounds for the representation number of a caterpillar and exact values in some cases.

Akhtar, Evans, and Pritikin [164] characterized the representation number of $K_{1,n}$ using Euler's phi function, and conjectured that this representation number is always of the form 2^a or $2^a p$, where $a \geq 1$ and p is a prime. They proved this conjecture for "small"

n and proved that for sufficiently large n, the representation number of $K_{1,n}$ is of the form $2^a,\ 2^a p$, or $2^a pq$, where $a \geq 1$ and p and q are primes. In [165] they showed that for sufficiently large $n \geq m$, $\text{rep}(K_{m,n}) = 2^a, 3^a, 2^a p^b$, or $2^a pq$, where $a, b \geq 1$ and p and q are primes; and for sufficiently large order, $\text{rep}(K_{n_1,n_2,\ldots,n_t} = p^a, p^a q^b$, or $p^a q^b u$, where p, q, u are primes with $p, q < u$. Akhtar [166] determined the representation number of graphs of the form $K_2 \cup nK_1$ (he uses the notation $K_2 + nK_1$) and studies their prime decompositions. Using relations between representation modulo r and product representations, he determined representation number of binary trees and gave an improved lower bound for hypercubes.

7.27 Sequentially Additive Graphs

Bange, Barkauskas, and Slater [472] defined a *k-sequentially additive labeling* f of a graph $G(V, E)$ to be a bijection from $V \cup E$ to $\{k, \ldots, k + |V \cup E| - 1\}$ such that for each edge $xy,\ f(xy) = f(x) + f(y)$. They proved: K_n is 1-sequentially additive if and only if $n \leq 3$; C_{3n+1} is not k-sequentially additive for $k \equiv 0$ or 2 (mod 3); C_{3n+2} is not k-sequentially additive for $k \equiv 1$ or 2 (mod 3); C_n is 1-sequentially additive if and only if $n \equiv 0$ or 1 (mod 3); and P_n is 1-sequentially additive. They conjecture that all trees are 1-sequentially additive. Hegde [1404] proved that $K_{1,n}$ is k-sequentially additive if and only if k divides n.

Hajnal and Nagy [1355] investigated 1-sequentially additive labelings of 2-regular graphs. They prove: kC_3 is 1-sequentially additive for all k; kC_4 is 1-sequentially additive if and only if $k \equiv 0$ or 1 (mod 3); $C_{6n} \cup C_{6n}$ and $C_{6n} \cup C_{6n} \cup C_3$ are 1-sequentially additive for all n; C_{12n} and $C_{12n} \cup C_3$ are 1-sequentially additive for all n. They conjecture that every 2-regular simple graph on n vertices is 1-sequentially additive where $n \equiv 0$ or 1 (mod 3).

Acharya and Hegde [56] have generalized k-sequentially additive labelings by allowing the image of the bijection to be $\{k, k+d, \ldots, (k+|V \cup E|-1)d\}$. They call such a labeling *additively (k, d)-sequential*.

7.28 Difference Graphs

Analogous to a sum graph, Harary [1368] calls a graph a *difference graph* if there is a bijection f from V to a set of positive integers S such that $xy \in E$ if and only if $|f(x) - f(y)| \in S$. Bloom, Hell, and Taylor [666] have shown that the following graphs are difference graphs: trees, $C_n,\ K_n,\ K_{n,n},\ K_{n,n-1}$, pyramids, and n-prisms. Gervacio [1247] proved that wheels W_n are difference graphs if and only if $n = 3, 4$, or 6. Sonntag [3677] proved that cacti (that is, graphs in which every edge is contained in at most one cycle) with girth at least 6 are difference graphs and he conjectures that all cacti are difference graphs. Sugeng and Ryan [3736] provided difference labelings for cycles; fans; cycles with chords; graphs obtained by the one-point union of K_n and P_m; and graphs made from any number of copies of a given graph G that has a difference labeling by identifying one vertex the first with a vertex of the second, a different vertex of the second with the third

and so on. Vaithilingam [4036] proved that gears, ladders, fans, friendship graphs, helms, and wheels admit difference labelings.

Hegde and Vasudeva [1429] call a simple digraph a *mod difference digraph* if there is a positive integer m and a labeling L from the vertices to $\{1, 2, \ldots, m\}$ such that for any vertices u and v, (u, v) is an edge if and only if there is a vertex w such that $L(v) - L(u) \equiv L(w) \pmod{m}$. They prove that the complete symmetric digraph and unidirectional cycles and paths are mod difference digraphs.

In [3386] Seoud and Helmi provided a survey of all graphs of order at most 5 and showed the following graphs are difference graphs: K_n, $(n \geq 4)$ with two deleted edges having no vertex in common; K_n, $(n \geq 6)$ with three deleted edges having no vertex in common; gear graphs G_n for $n \geq 3$; $P_m \times P_n$ $(m, n \geq 2)$; triangular snakes; C_4-snakes; dragons (that is, graphs formed by identifying the end vertex of a path and any vertex in a cycle); graphs consisting of two cycles of the same order joined by an edge; and graphs obtained by identifying the center of a star with a vertex of a cycle.

7.29 Square Sum Labelings and Square Difference Labelings

A bijective mapping $f : V(G)$ to $\{0, 1, 2, \ldots, |V(G)| - 1\}$ is said to be a *square sum labeling* if the induced function f^* from $E(G)$ to the positive integers defined by $f^*(xy) = (f(x)^2 + (f(y))^2$ is injective. A graph that has a square sum labeling is called a *square sum* graph. A *square difference* graph is defined the same way with $f^*(xy) = (f(x)^2 + (f(y))^2$ replaced with $f^*(xy) = |(f(x)^2 - (f(y))^2|$. Maheswari and Srividya [2396] proved that every cycle C_n $(n \geq 6)$ with parallel P_3 chords admit a vertex odd mean labeling, a vertex even mean labeling, and a square sum labeling. Maheswari, Azhagarasi, and Samuvel [2388] proved cycles with at least 6 vertices with parallel P_4 chords are vertex odd mean graphs and vertex even mean graphs and they admit square sum labelings, and square difference labelings.

In [2395] they proved that the following graphs are square sum graphs: cycles with parallel chords, graphs obtained by attaching an arbitrary number of pendent edges at a vertex of degree 2 of a cycle with parallel chords, duplication of a vertex of degree 2 of a cycle with parallel chords, crowns with parallel chords, chains of even cycles with parallel chords, and graphs obtained from copies of C_n by joining a vertex from each copy of C_n to a common vertex. Patel and Ghodasara [2793] proved that the graph obtained by joining two copies of a specific graph by a path of arbitrary length admits a square sum labeling. They gave some results about the square sum graphs of arbitrary super subdivisions. Zhang, Naeem, Tariq, and Zhao [4324] investigated the square sum labeling of generalized Petersen graphs and double generalized Petersen graphs. Shiama [3510] proved that the total graph of paths and cycles, and the middle graphs of paths and cycles, admit square sum labelings. Parameswari and Margret [2768] proved that the octopus graph and Vanessa graph admit square sum labelings. Krithika and Keerthana [2039] proved that shell bow graphs admit square difference labelings and square difference prime labelings.

Kulli [2056] defined a *semitotal-block* graph $T_b(G)$ of a graph G as the graph whose

set of vertices is the union of the set of vertices and blocks of G and in which two points are adjacent if and only the corresponding vertices of G are adjacent or the corresponding blocks are incident. Mirajkar and Sthavarmath [2509] proved the following graphs admit square sum and square difference labelings: $T_b(P_n \odot K_1)$ $(n \geq 3)$; $T_b(C_n \odot K_1)$ $(n \geq 3)$ except for $n = 0 \pmod 8$; and $T_b(F_n \odot K_1)$ $(n \geq 4)$. In [876] Cynthia and Poorani proved circulant networks and a torus are square difference graphs. In [880] Cynthia, Poorani, and Mercy provide an algorithm to find the square difference labelings of bloom graphs.

Ajitha, Arumugam, and Germina [154] prove the following graphs have square sum labelings: trees; cycles; $K_2 + mK_1$; K_n if and only if $n \leq 5$; $C_n^{(t)}$ (the one-point union of t copies of C_n); grids $P_m \times P_n$; and $K_{m,n}$ if $m \leq 4$. They also prove that every strongly square sum graph except K_1, K_2, and K_3 contains a triangle.

In [3509] Shiama proved that the total graphs of paths and cycles and the middle graphs of paths and cycles are square sum graphs. A *lilly* graph is obtained by identifying one endpoint of each of two copies of P_n with the center of $K_{1,2n}$ $(n > 1)$. Samuvel and Kalaivani proved [3274] the following graphs are square sum graphs: the graph obtained by the duplication of the center vertex and another vertex of a lilly graph; the graph obtained by identifying any two vertices of a lilly graph, and the graph obtained by the switching of the center vertex and another vertex of a lilly graph.

Subhashini, Ramanathan, and Manimekalai [3706] proved that pyramids with at least 3 rows, hanging pyramids, graphs obtained from starting with r copies of a hanging pyramid and joining each copy with the next one with an edge, and the one point union of r copies of a pyramid and hanging pyramids admit square sum labelings. Maheswari, Azhagarasi, and Samuvel [2390] showed that the following graphs are square sum graphs: the corona product of P_m $(m \geq 2)$ and C_{2n} $(n \geq 3)$ with P_3 parallel chords; the corona product P_m $(m \geq 2)$ and C_{2n+1} $(n \geq 3)$ with P_3 parallel chords; $C_{2n} \odot K_1$ $(n \geq 3)$ with P_3 parallel chords; $C_{2n+1} \odot K_1$ with P_3 parallel chords; the chain of cycle $C_{2n,m}$ $(n \geq 3)$ with P_3 parallel chords (a chain of m copies of C_{2n} with P_3 parallel chords where each copy shares exactly one vertex of the next copy); and edge connected cycle C_{2n} $(n \geq 3)$ with P_3 chords.

In [85] Agasthi and Narvathi provided ways to construct square sum, square difference, root mean square, strongly multiplicative, even mean and odd mean labelings for triangular snakes, and the central graph of triangular snake graphs.

In [1256] Ghodasara and Patel gave a counterexample to the conjecture by Germina and Sebastian [1246] that if G_1 and G_2 are square sum graphs then $G_1 \cup G_2$ is a square sum graph. They proved that the duplication graphs of any vertex of the following graphs are square sum graphs: K_n if and only if $n \leq 7$, the Petersen graph $P(5, 2)$, $K_{1,n}$, and C_n They also proved that cycle C_n with $[\frac{n}{2}]$ concurrent chords is a square sum graph.

In [1253] Ghodasara and Patel proved that the following constructions based on the bistar $B_{n,n}$ are square sum graphs: the restricted square, the splitting graph, the shadow graph, the degree splitting graph, the arbitrary super subdivision graph, and the duplication of any vertex of $B_{n,n}$. They defined restricted total graph of $B_{n,n}$ as a graph with vertex set $= V(B_{n,n}) \cup E(B_{n,n}) = \{u, v, w, u_i, v_i, u_i', v_i'/1 \leq i \leq n\}$, where u and v are apex vertices, u_i and v_i are pendent vertices, w, u_i' and v_i' are vertices corresponding

to the edges of $B_{n,n}$ and edge set $= E(B_{n,n}) \cup \{uw, vw, wu_i', wv_i', uu_i', vv_i', u_iu_i', v_iv_i', /1 \leq i \leq n\}$. They also defined restricted middle graph of $B_{n,n}$ as a graph with vertex set $= V(B_{n,n}) \cup E(B_{n,n}) = \{u, v, w, u_i, v_i, u_i', v_i'/1 \leq i \leq n\}$, where u and v are apex vertices, u_i and v_i are pendent vertices, w, u_i' and v_i' are vertices corresponding to the edges of $B_{n,n}$ and edge set $= \{uw, vw, wu_i', wv_i', uu_i', vv_i', u_iu_i', v_iv_i', /1 \leq i \leq n\}$. They proved that restricted total graph and restricted middle graph of $B_{n,n}$ are square sum graphs.

Germina and Sebastian [1245] proved that the following graphs are square sum graphs: trees; unicyclic graphs; mC_n; cycles with a chord; the graphs obtained by joining two copies of cycle C_n by a path P_k; and graphs that are a path union of k copies of C_n and the path is P_2. In [3372] Seoud and Al-Harere give several necessary conditions for a graph to be a square sum graph and show that $2C_n, P_{2n}$, and C_{2n} are square sum graphs. Huilgol and Sriram [1470] prove that if G_1 and G_2 are square sum, then $G_1 \cup G_2 \cup G_3$ is also square sum, where G_3 is a set of isolated vertices.

In [3655] Somashekara and Veena used the term "square sum labeling" to mean "strongly square sum labeling." They proved that the following graphs have strongly square sum labelings: paths, $K_{1,n_1} \cup K_{1,n_2} \cup \cdots \cup K_{1,n_k}$, complete n-ary trees, and lobsters obtained by joining centers of any number of copies of a star to a new vertex. They observed that that if every edge of a graph is an edge of a triangle then the graph does not have strongly square sum labeling. As a consequence, the following graphs do not have a strongly square sum labelings: $K_n, n \geq 3$; wheels; fans $P_n + K_1$ $(n \geq 2)$; double fans $P_n + K_2$ $(n \geq 2)$; friendship graphs $C_3^{(n)}$; windmills $K_m^{(n)}$ $(m > 3)$; triangular ladders; triangular snakes; double triangular snakes; and flowers. They also proved that helms are not strongly square sum graphs and the graphs obtained by joining the centers of two wheels to a new vertex are not strongly square sum graphs.

Govindan, Pinelas, and Dhivya [1322] introduced the notion a *cube sum* graph as a graph $G(V, E)$ with p vertices and q edges such that if there exists a bijection $f : V \to \{0, 1, 2, \ldots, p-1\}$, then the induced function $f^* : E \to N$, defined by $f(^*uv) = (f(u))^3 + (f(v))^3$, is injective. They proved that paths, cycle, stars, wheels, and fans are cube sum graphs. Patel and Ghodasara [2791] proved that the following graphs are cube sum graphs: trees, gears, shell graphs, helms, K_n if and only if n is at most 11, and $K_{2,n}$ for all n. Krithika and Keerthana [2039] proved that shell bow graphs, the path union of two subdivided shell graphs, and the disjoint union of two copies of shell bow graphs admit cube difference labelings.

Let $G_1, G_2, \ldots, G_n$ be subdivided shell graphs of any order. The graph $SSG(n)$ is obtained by adding an edge to apexes of G_i and G_{i+1} for $i = 1, 2, \ldots, n-1$. The graph $SSG(n)$ is called a *path union of n subdivided shell graphs*. In [2040] Krithika and Keerthana proved that $SSG(2)$ admits square difference labeling and square difference prime labeling.

In [3671] Sonchhatra and Ghodasara call a (p, q)-graph $G = (V, E)$ *sum perfect square* if there exists a bijection f from V to $\{0, 1, 2, \ldots, p-1\}$ such that the function f^* from E defined by $f^*(uv) = (f(u)) + (f(v))^2$ for all edges uv is an injection. Such an f is called a *sum perfect square* labeling of G. In a series of four papers the following graphs are proved to be sum perfect square graphs: cycles, cycles with one chord, cycles with twin

chords, trees [3671]; several snake related graphs [3672]; $K_{1,n}+K_1,\ K_2+mK_1,\ C_n\odot K_1$, graphs obtained from $K_{1,n}$ with endpoint vertices $v_1, v_2, \ldots, v_n$ by joining v_i and v_{i+1} with an edge for $i = 1, 2, \ldots, v_{\lfloor n/2 \rfloor}$ ("half wheel"), the middle graphs of paths, the total graphs of paths [3674]; P^2 $(n > 1)$, $mK_{1,n}$, mC_n,, and the splitting graph and the shadow graph of a star [3670]. In [3674] they prove that the union of two stars and that for any sum perfect square graph G, $G \cup P_n$ is sum perfect square. They conjecture that the union of any two sum perfect square graphs is sum perfect square.

In [3671] Sonchhatra, Pandya, and Chhaya call a (p, q)-graph $G(V, E)$ *sum perfect cube* if there exists a bijection f from V to $\{0, 1, 2, \ldots, p-1\}$ such that the function f^* defined by $f^*(uv) = f(u)^3+3f(u)f(v)^2+3f(u)^2f(v)+f(v)^3$ for all edges uv is an injection. Such an f is called a *sum perfect cube* labeling of G. They proved that trees, C_n $(n \geq 4)$ with one cord or twin cords, and tadpoles admit sum perfect cube labelings

Ajitha, Princy, Lokesha, and Ranjini [152] defined a graph $G(p, q)$ to be a *square difference graph* if there exist a bijection f from $V(G)$ to $\{0, 1, 2, \ldots, p-1\}$ such that the induced function f^* from $E(G)$ to the natural numbers given by $f^*(uv) = |(f(u))^2 - (f(v))^2|$ for every edge uv of G is a bijection. Such a the function is called a *square difference labeling* of the graph G. They proved that following graphs have square difference labelings: paths, stars, cycles, K_n if and only if $n \leq 5$, $K_{m,n}$ if $m \leq 4$, friendship graphs $C_3^{(n)}$, triangular snakes, and $K_2 + mK_1$. They also prove that every graph can be embedded as a subgraph of a connected square difference graph and conjecture that trees, complete bipartite graphs and $C_k^{(n)}$ are square difference graphs.

Kulli [2056] defined a *semitotal-block* graph $T_b(G)$ of a graph G as the graph whose set of vertices is the union of the set of vertices and blocks of G and in which two points are adjacent if and only the corresponding vertices of G are adjacent or the corresponding blocks are incident. Mirajkar and Sthavarmath [2509] proved the following graphs admit square sum and square difference labelings: $T_b(P_n \odot K_1)$ $(n \geq 3)$; $T_b(C_n \odot K_1)$ $(n \geq 3)$ except for $n = 0 \pmod 8$; and $T_b(F_n \odot K_1)$ $(n \geq 4)$. In [880] Cynthia, Poorani, and Mercy provide an algorithm to find the square difference labelings of bloom graphs.

Tharmaraj and Sarasija [3884] proved that following graphs have square difference labelings: fans F_n $(n \geq 2)$; $P_n + \overline{K_2}$; the middle graphs of paths and cycles; the total graph of a path; the graphs obtained from m copies of an odd cycle and the path P_m with consecutive vertices $v_1, v_2, \ldots, v_m$ by joining the vertex v_i to a vertex of the i^{th} copy of the odd cycle; and the graphs obtained from m copies of the star S_n and the path P_m by joining the vertex v_i of P_m to the center of the i^{th} copy of S_n. Sebastian and Germina [3332] proved that certain planar graphs and higher order level joined planar grid admit square sum labeling. They also study square sum properties of several classes of graphs with many odd cycles.

In [1236] Geetha and Kalamani showed that the following graphs have square difference labelings: two copies of the same star whose centers are joined by a path; and two copies of the same cycles whose centers are joined by a path, the restricted square of bistar $B_{n,n}$; the restricted total graph of bistar $B_{n,n}$; the restricted middle graph of $B_{n,n}$ Shiama [3512] showed that cycles, complete graphs, cycle cacti, ladders, lattice grids, quadrilateral snakes, $K_2 + mK_1$ admit square difference labelings.

In [3804] Sunoj and Mathew Varkey defined the *square difference prime labeling* of a graph $G(V, E)$ as a bijection f from V to $\{0, 1, 2, \ldots, |V| - 1\}$ such that the induced function f^*_{spd} from E to the natural numbers given by $f^*_{spd}(uv) = |(f(u)^2 - (f(v))^2|$ has the property that the greatest common divisor of the labels of the edges incident with a vertex of degree greater than one is 1. In [3804], [3805], [3806], and [3807] they proved the following graphs admit square difference prime labelings: certain triangular snakes, double triangular snakes and alternate triangular snakes, centipedes, twigs, coconut trees, combs, stars, subdivision of stars, bistars, cycles, middle graphs of cycles, total graphs of cycles, lilies graph, and some star related graphs.

Vaghela and Parmar [3940] say a graph G admits a *difference perfect square cordial* labeling if there is a bijection $f : V(G) \to \{1, 2, \ldots, |V(G|\}$ such that for each edge uv the induced map $f^* : E(G) \to \{0, 1\}$ defined by $f^*(uv) = 1$ if $u^2 - 2uv + v^2 = 1$, and 0 otherwise, has the property that the number of edges labeled with 0 and the number of edges labeled with 1 difference by at most 1. A graph that admits a difference perfect square cordial labeling is said to be a *difference perfect square cordial* graph. They obtained difference perfect square cordial labelings for paths, cycles, wheels, fans, combs, crowns, $(C_m \odot K_1) \cup (P_n \odot K_1)$, $D_2(P_n)$, P_n^2, $K_2 \odot C_n$, graphs consisting of two copies of C_n that share a common edge, the vertex switching of C_n, and the graph obtained by starting with P_n ($n \geq 6$) and two new vertices u and v of P_n and joining v to first two vertices and last two vertices of P_n and joining u to the remaining vertices of P_n (called the *shipping graph*) In [3941] Vaghela and Parmar provided difference perfect square cordial labelings of the H-graphs of paths, some corona graphs, total graphs of paths, and graphs obtained from $P_n \times P_2$ where P_n has consecutive vertices $v_1, v_2, \ldots, v_n$ by joining v_i in left P_n to v_{i+1} in the right copy of P_n with an edge for $i = 1, 2, \ldots, n - 1$. In [3942] Vaghela and Parmar obtain difference perfect square cordial labeling of triangular snake graphs, quadrilateral snake graphs, alternate triangular snake graphs, alternate quadrilateral snake graphs, irregular triangular snake graphs, irregular quadrilateral snake graphs, double triangular snake graphs, double quadrilateral snake graphs, double alternate triangular snake graphs, and double alternate quadrilateral snake graphs.

A graph G is said to admit a *square difference labeling* if there exists an injection $f : V(G) \to \{0, 1, 2, \ldots, n - 1\}$ for some n such that when each edge of G is assigned the absolute square difference of its end-vertices, the resulting labels are distinct. *Cube difference labeling* is defined analogously.

Shiana [3511] proved that paths, cycles, stars, fans, wheels, crowns, helms, dragons, coconut trees, and shell graphs admit cube difference labelings. Sharon Philomena and Thirusangu [3493] proved the cycle cactus graph $C_n^{(3)}$, the tree of diameter 4 obtained from the bistar $B_{n,n}$ by subdividing the middle edge with a new vertex, and the graph obtained by joining one vertex of a cycle and one vertex of degree 2 of a comb by an edge have square and cube difference labelings (that is, the absolute cube difference of end-vertices of the edges are distinct). Sherman [3507] proved the path union of nC_3 and the disjoint union of m stars $K_{1,n_1}, K_{1,n_2}, \ldots, K_{1,n_m}$ are square difference graphs

Subashini, Bhuvaneswari, and Manimekalai [3702] proved the following graphs have square difference labelings: theta graphs, the duplication of any vertex of degree 3 in the

cycle of a theta graph, the one point union of any number of theta graphs, the path union of any number of copies of a theta graph, the fusion of any two vertices in the cycle of a theta graph, and the switching of a central vertex of a theta graph.

7.30 Permutation and Combination Graphs

Hegde and Shetty [1423] define a graph G with p vertices to be a *permutation graph* if there exists a injection f from the vertices of G to $\{1, 2, 3, \ldots, p\}$ such that the induced edge function g_f defined by $g_f(uv) = f(u)!/|f(u) - f(v)|!$ is injective. They say a graph G with p vertices is a *combination graph* if there exists a injection f from the vertices of G to $\{1, 2, 3, \ldots, p\}$ such that the induced edge function g_f defined as $g_f(uv) = f(u)!/|f(u) - f(v)|!f(v)!$ is injective. They prove: K_n is a permutation graph if and only if $n \leq 5$; K_n is a combination graph if and only if $n \leq 5$; C_n is a combination graph for $n > 3$; $K_{n,n}$ is a combination graph if and only if $n \leq 2$; W_n is a not a combination graph for $n \leq 6$; and a necessary condition for a (p, q)-graph to be a combination graph is that $4q \leq p^2$ if p is even and $4q \leq p^2 - 1$ if p is odd. They strongly believe that W_n is a combination graph for $n \geq 7$ and all trees are combinations graphs. Baskar Babujee and Vishnupriya [588] prove the following graphs are permutation graphs: P_n; C_n; stars; graphs obtained adding a pendent edge to each edge of a star; graphs obtained by joining the centers of two identical stars with an edge or a path of length 2); and complete binary trees with at least three vertices. Seoud and Salim [3397] determine all permutation graphs of order at most 9 and prove that every bipartite graph of order at most 50 is a permutation graph. Seoud and Mahran [3389] give an upper bound on the number of edges of a permutation graph and introduce some necessary conditions for a graph to be a permutation graph. They show that these conditions are not sufficient for a graph to be a permutation graph.

Ghodasara and Patel [1255] proved that the following graphs are permutation graphs: the Petersen graph $P(5, 2)$, trees, $K_{3,n}$ $(n \geq 1)$ for $n+3$ prime, W_n $(n \geq 3)$ for $n+1$ prime, shell graph S_n $(n \geq 3)$ for prime n, dumbbell graph $D_{n,k,2}$ $(n, k \geq 3)$, $C_n \odot K_1$ $(n \geq 3)$, and the one point union $C_n^{(k)}$ $(k \geq 2, n \geq 3)$ of k copies of cycle C_n. A t-ply $P_t(u, v)$ is a graph with t paths, each of length at least two and such that no two paths have a vertex in common except for the end vertices u and v. Ghodasara and Patel defined t^*-ply $P_{t^*}(u, v)$ as a special case of t-ply $P_t(u, v)$ graph with every t path have same length and proved that t^*-ply $P_{t^*}(u, v)$ is a permutation graph.

The graph obtained from two copies of an (m, n) kite graph by connecting the degree 1 vertex of one copy to the vertex of degree 3 and the second copy is called the 1-*join (m,n) kite*. The graph obtained by repeating this construction with t copies of an (m, n) kite is called the *t-join (m,n) kite*. Sriramr and Govindarajan [3691] proved t-join (m, n) kites are permutation graphs.

Ghodasara and Patel [1254] proved that the following graphs are combination graphs: $C_n \times P_2$ for $n \geq 6$, umbrella graph $U(m, n)$ for $m, n > 2$, armed crown $C_n \oplus P_m$ for $n \geq 4$ and $m \geq 1$, the graphs obtained by joining C_{2m} $(m \geq 2)$ to each pendent vertex of $K_{1,n}(n \geq 2)$, the duplication of any rim vertex of W_n for $n \geq 7$, C_n with $[\frac{n-4}{2}]$ concurrent

chords for $n \geq 6$, and the duplication of vertex in C_n for $n \geq 5$.

Hegde and Shetty [1423] say a graph G with p vertices and q edges is a *strong k-combination graph* if there exists a bijection f from the vertices of G to $\{1, 2, 3, \ldots, p\}$ such that the induced edge function g_f from the edges to $\{k, k+1, \ldots, k+q-1\}$ defined by $g_f(uv) = f(u)!/|f(u) - f(v)|!f(v)!$ is a bijection. They say a graph G with p vertices and q edges is a *strong k-permutation graph* if there exists a bijection f from the vertices of G to $\{1, 2, 3, \ldots, p\}$ such that the induced edge function g_f from the edges to $\{k, k+1, \ldots, k+q-1\}$ defined by $g_f(uv) = f(u)!/|f(u) - f(v)|!$ is a bijection. Seoud and Anwar [3375] provided necessary conditions for combination graphs, permutation graphs, strong k-combination graphs, and strong k-permutation graphs.

Seoud and Al-Harere [3374] showed that the following families are combination graphs: graphs that are two copies of C_n sharing a common edge; graphs consisting of two cycles of the same order joined by a path; graphs that are the union of three cycles of the same order; wheels W_n $(n \geq 7)$; coronas $T_n \odot K_1$, where T_n is the triangular snake; and the graphs obtained from the gear G_m by attaching n pendent vertices to each vertex which is not joined to the center of the gear. They proved that a graph $G(n, q)$ having at least 6 vertices such that 3 vertices are of degree $1, n-1, n-2$ is not a combination graph, and a graph $G(n, q)$ having at least 6 vertices such that there exist 2 vertices of degree $n-3$, two vertices of degree 1 and one vertex of degree $n-1$ is not a combination graph.

Seoud and Al-Harere [3373] proved that the following families are combination graphs: unions of four cycles of the same order; double triangular snakes; fans F_n if and only if $n \geq 6$; caterpillars; complete binary trees; ternary trees with at least 4 vertices; and graphs obtained by identifying the pendent vertices of stars S_m with the paths P_{n_i}, for $1 \leq n_i \leq m$. They include a survey of trees of order at most 10 that are combination graphs and proved the following graphs are not combination graphs: bipartite graphs with two partite sets with $n \geq 6$ elements such that $n/2$ elements of each set have degree n; the splitting graph of $K_{n,n}$ $(n \geq 3)$; and certain chains of two and three complete graphs. Seoud and Anwar [3375] proved the following graphs are combination graphs: dragon graphs (the graphs obtained from by joining the endpoint of a path to a vertex of a cycle); triangular snakes T_n $(n \geq 3)$; wheels; and the graphs obtained by adding k pendent edges to every vertex of C_n for certain values of k.

In [3371] and [3373] Seoud and Al-Harere proved the following graphs are non-combination graphs: $G_1 + G_2$ if $|V(G_1)|, |V(G_2)| \geq 2$ and at least one of $|V(G_1)|$ and $|V(G_2)|$ is greater than 2; the double fan $\overline{K_2} + P_n$; $K_{l,m,n}$; $K_{k,l,m,n}$; $P_2[G]$; $P_3[G]$; $C_3[G]$; $C_4[G]$; $K_m[G]$; $W_m[G]$; the splitting graph of K_n $(n \geq 3)$; K_n $(n \geq 4)$ with an edge deleted; K_n $(n \geq 5)$ with three edges deleted; and $K_{n,n}$ $(n \geq 3)$ with an edge deleted. They also proved that a graph $G(n, q)$ $(n \geq 3)$ is not a combination graph if it has more than one vertex of degree $n-1$.

In [3886] and [3885] Tharmaraj and Sarasija defined a graph $G(V, E)$ with p vertices to be a *beta combination* graph if there exist a bijection f from $V(G)$ to $\{1, 2, \ldots, p\}$ such that the induced function B_f from $E(G)$ to the natural numbers given by $B_f(uv) = (f(u) + f(v))!/f(u)!f(v)!$ for every edge uv of G is injective. Such a function is called a *beta combination* labeling. They prove the following graphs have beta combination label-

ings: K_n if and only if $n \leq 8$; ladders L_n $(n \geq 2)$; fans F_n $(n \geq 2)$; wheels; paths; cycles; friendship graphs; $K_{n,n}$ $(n \geq 2)$; trees; bistars; $K_{1,n}$ $(n > 1)$; triangular snakes; quadrilateral snakes; double triangular snakes; alternate triangular snakes (graphs obtained from a path $v_1, v_2, \ldots, v_n$, where for each odd $i \leq n-1$, v_i and v_{i+1} are joined to a new vertex $u_{i,i+1}$; alternate quadrilateral snakes (graphs obtained from a path $v_1, v_2, \ldots, v_n$, where for each odd $i \leq n-1$, v_i and v_{i+1} are joined to two new vertices $u_{i,i+1,1}$ and $u_{i,i+1,2}$); helms; gears; combs $P_n \odot K_1$; and coronas $C_n \odot K_1$.

7.31 Strongly *-graphs

A variation of strong multiplicity of graphs is a strongly *-graph. A graph of order n is said to be a *strongly *-graph* if its vertices can be assigned the values $1, 2, \ldots, n$ in such a way that, when an edge whose vertices are labeled i and j is labeled with the value $i + j + ij$, all edges have different labels. Adiga and Somashekara [76] have shown that all trees, cycles, and grids are strongly *-graphs. They further consider the problem of determining the maximum number of edges in any strongly *-graph of given order and relate it to the corresponding problem for strongly multiplicative graphs. In [3391] and [3392] Seoud and Mahan give some technical necessary conditions for a graph to be strongly *-graph,

Baskar Babujee and Vishnupriya [588] have proved the following are strongly *-graphs: $C_n \times P_2$, $(P_2 \cup \overline{K}_m) + \overline{K}_2$, windmills $K_3^{(n)}$, and *jelly fish* graphs $J(m, n)$ obtained from a 4-cycle v_1, v_2, v_3, v_4 by joining v_1 and v_3 with an edge and appending m pendent edges to v_2 and n pendent edges to v_4.

Baskar Babujee and Beaula [568] prove that cycles and complete bipartite graphs are vertex strongly *-graphs. Baskar Babujee, Rajesh Kannan, and Vishnupriya [580] prove that wheels, paths, fans, crowns, $(P_2 \cup mK_1) + \overline{K_2}$, and umbrellas (graphs obtained by appending a path to the central vertex of a fan) are vertex strongly *-graphs.

In [3393] Seoud, Roshdy, and AboShady gave an upper bound for the number of edges of any graph in terms of the number of vertices to be a strongly *-graph and some new families to be strongly*- graphs. They also provided an algorithm for checking if a graph is a strongly *-graph or not.

7.32 Triangular Sum Graphs

Hegde and Shankaran [1418] call a labeling of graph with q edges a *triangular sum labeling* if the vertices can be assigned distinct non-negative integers in such a way that, when an edge whose vertices are labeled i and j is labeled with the value $i + j$, the edges labels are $\{k(k+1)/2 |\ k = 1, 2, \ldots, q\}$. They prove the following graphs have triangular sum labelings: paths, stars, complete n-ary trees, and trees obtained from a star by replacing each edge of the star by a path. They also prove that K_n has a triangular sum labeling if and only if n is 1 or 2 and the friendship graphs $C_3^{(t)}$ do not have a triangular sum labeling. They conjecture that K_n $(n \geq 5)$ are forbidden subgraphs of graph with triangular sum labelings. They conjectured that every tree admits a triangular sum labeling. They

show that some families of graphs can be embedded as induced subgraphs of triangular sum graphs. They conclude saying "as every graph cannot be embedded as an induced subgraph of a triangular sum graph, it is interesting to embed families of graphs as an induced subgraph of a triangular sum graph". In response, Seoud and Salim [3395] showed the following graphs can be embedded as an induced subgraph of a triangular sum graph: trees, cycles, nC_4, and the one-point union of any number of copies of C_4 (friendship graphs).

Vaidya, Prajapati, and Vihol [4000] showed that cycles, cycles with exactly one chord, and cycles with exactly two chords that form a triangle with an edge of the cycle can be embedded as an induced subgraph of a graph with a triangular sum labeling. They proved that several classes of graphs do not have triangular sum labelings. Among them are: helms, graphs obtained by joining the centers of two wheels to a new vertex, and graphs in which every edge is an edge of a triangle. As a corollary of the latter result they have that $P_m + \overline{K_n}$, $W_m + \overline{K_n}$, wheels, friendship graphs, flowers, triangular ladders, triangular snakes, double triangular snakes, and flowers. do not have triangular sum labelings.

Seoud and Salim [3395] proved the following are triangular sum graphs: $P_m \cup P_n,\ m \geq 4$; the union of any number of copies of $P_n,\ n \geq 5$; $P_n \odot \overline{K_m}$; symmetrical trees; the graph obtained from a path by attaching an arbitrary number of edges to each vertex of the path; the graph obtained by identifying the centers of any number of stars; and all trees of order at most 9.

For a positive integer i the ith *pentagonal number* is $i(3i-1)/2$. Somashekara and Veena [3656] define a *pentagonal sum labeling* of a graph $G(V, E)$ as one for which there is a one-to-one function f from $V(G)$ to the set of nonnegative integers that induces a bijection f^+ from $E(G)$ to the set of the first $|E|$ pentagonal numbers. A graph that admits such a labeling is called a *pentagonal sum graph*. Somashekara and Veena [3656] proved that the following graphs have pentagonal sum labelings: paths, $K_{1,n_1} \cup K_{1,n_2} \cup \cdots \cup K_{1,n_k}$, complete n-ary trees, and lobsters obtained by joining centers of any number of copies of a star to a new vertex. They conjecture that every tree has a pentagonal sum labeling and as an open problem they ask for a proof or disprove that cycles have pentagonal labelings. They observed that if every edge of a graph is an edge of a triangle then the graph does not have pentagonal sum labeling. As was the case for triangular sum labelings the following graphs do not have a pentagonal sum labeling: $P_m + \overline{K_n}$, and $W_m + \overline{K_n}$ wheels, friendship graphs, flowers, triangular ladders, triangular snakes, double triangular snakes, and flowers. Somashekara and Veena [3656] also proved that helms and the graphs obtained by joining the centers of two wheels to a new vertex are not pentagonal sum graphs.

7.33 Divisor Graphs

Santhosh and Singh [3309] call a graph $G(V, E)$ a *divisor graph* if V is a set of integers and $uv \in E$ if and only if u divides v or vice versa. They prove the following are divisor graphs: trees; mK_n; induced subgraphs of divisor graphs; cocktail party graphs $H_{m,n}$ (see

Section 7.1) for the definition); the one-point union of complete graphs of different orders; complete bipartite graphs; W_n for n even and $n > 2$; and $P_n + \overline{K_t}$. They also prove that C_n $(n \geq 4)$ is a divisor graph if and only if n is even and if G is a divisor graph then for all n so is $G + K_n$.

Chartrand, Muntean, Saenpholphat, and Zhang [786] proved complete graphs, bipartite graphs, complete multipartite graphs, and joins of divisor graphs are divisor graphs. They also proved if G is a divisor graph, then $G \times K_2$ is a divisor graph if and only if G is a bipartite graph; a triangle-free graph is a divisor graph if and only if it is bipartite; no divisor graph contains an induced odd cycle of length 5 or more; and that a graph G is divisor graph if and only if there is an orientation D of G such that if (x, y) and (y, z) are edges of D then so is (x, z).

In [169] and [171] Al-Addasi, AbuGhneim, and Al-Ezeh determined precisely the values of n for which P_n^k $(k \geq 2)$ are divisor graphs and proved that for any integer $k \geq 2$, C_n^k is a divisor graph if and only if $n \leq 2k + 2$. In [34] they gave a characterization of the graphs G and H for which $G \times H$ is a divisor graph and a characterization of which block graphs are divisor graphs. (Recall a graph is a *block graph* if every one of its blocks is complete.) They showed that divisor graphs form a proper subclass of perfect graphs and showed that cycle permutation graphs of order at least 8 are divisor graphs if and only if they are perfect. (Recall a graph is *perfect* if every subgraph has chromatic number equal to the order of its maximal clique.) In [170] Al-Addasi, AbuGhneim, and Al-Ezeh proved that the contraction of a divisor graph along a bridge is a divisor graph; if e is an edge of a divisor graph that lies on an induced even cycle of length at least 6, then the contraction along e is not a divisor graph; and they introduced a special type of vertex splitting that yields a divisor graph when applied to a cut vertex of a given divisor graph.

AbuHijleh, AbuGhneim, and Al-Ezeh [36] prove that for any tree T, T^2 is a divisor graph if and only if T is a caterpillar and the diameter of T is less than six. For any caterpillar T and a positive integer k with $\text{diam}(T) < 2k$, they show that T^k is a divisor graph. Moreover, for a caterpillar T and $k \geq 3$ with $\text{diam}(T) = 2k$ or $\text{diam}(T) = 2k + 1$, they show that T^k is a divisor graph if and only if the centers of T have degree two. In [35] AbuHijleh, AbuGhneim, and Al-Ezeh prove that the k-th power Q_n^k of Q_n is a divisor graph if and only if $n = 2, 3$ or $n \geq 4$ and $k \geq n - 1$ hold. In the case of the n-dimensional folded-hypercube FQ_n (that is, the graph obtained from Q_n by adding to it a perfect matching that connects opposite pairs of the vertices of Q_n) they show that FQ_n is a divisor graph for odd n, but not for even $n \geq 4$. They also prove $(FQ_n)^k$ is not a divisor graph if and only if $2 \leq k \leq \lceil n/2 \rceil$, where $n \geq 5$.

Ganesan and Uthayakumar [1180] proved that $G \odot H$ is a divisor graph if and only if G is a bipartite graph and H is a divisor graph. Frayer [1110] proved $K_n \times G$ is a divisor graph for each n if and only if G contains no edges and $\overline{K_n \times K_2}$ $(n \geq 3)$ is a divisor graph. Vinh [4114] proved that for any $n > 1$ and $0 \leq m \leq n(n-1)/2$ there exists a divisor graph of order n and size m. She also gave a simple characterization of divisor graphs due to Chartrand, Muntean, Saenpholphat, and Zhang [786]. Gera, Saenpholphat, and Zhang [1239] established forbidden subgraph characterizations for all divisor graphs that contain at most three triangles. Tsao [3919] investigated the vertex-chromatic number,

the clique number, the clique cover number, and the independence number of divisor graphs and their complements. In [3381] Seoud, El Sonbaty, and Mahran discuss here some necessary and sufficient conditions for a graph to be a divisor graph.

In [1915] Kasthuri, Karuppasamy, and Nagarajan introduced the notions of SD-divisor labelings and SD-divisor cordial labelings as follows. Let G be finite, simple graph G with n vertices and f be a bijection from $V(G)$ to $\{1, 2, \ldots, n\}$. Define a function f' induced by f by assigning $f'(uv) = 1$ if $S = f(u) + f(v)$ divides $D = |f(u) - f(v)|$ and $f'(uv) = 0$, otherwise. They say that f' is a SD-*divisor* labeling if $f'(uv) = 1$ for all edges uv. In [3168] Revathi and Mary Jeya Jothi [3168] provided SD-divisor labelings for paths and cycles and certain path and cycle related graphs. They further find SD-divisor labelings of some subdivision graphs, coronas, and splitting graphs, and prove that stars, complete graphs, and wheels do not admit SD-divisor labelings.

A graph G with n vertices has a *modular multiplicative divisor* (MMD) labeling if there exist a bijection f from vertices of G to the set of all natural numbers from 1 to n such that when the edge uv is labeled $f(u)f(v) (\mathrm{mod}\, n)$, then n divides the sum of all edge labels of G. They prove that r-dimensional butterfly networks admit MMD labelings. Revathi, Dharmakkan, and Jeya Jothi [3166] proved that $K_{l,m,n}$ admits a modular multiplicative divisor labeling. Revathi and Rajeswari [3169] proved P_n, $P_{a,b}$ (the graph that connects two vertices by means of b internally disjoint paths of length a each), the shadow graph of a path, and $P_n \times P_1$, (where n is not a multiple of 6) admit modular multiplicative divisor labelings. They also discuss the upper bound for the number of edges in a modular multiplicative divisor graphs. In [3167] Revathi and Rajeswari proved that the split graphs of cycles, helms, and flower graphs admit modular multiplicative divisor labelings.

A *divisor 3-equitable* labeling of a graph G is a bijection $d : V(G) \to \{1, 2, \ldots, |V(G)|\}$ such that induced map d^* defined on the edges of G by, for any edge uv with $d(u) \leq d(v)$, $d^*(uv)) = 1$ if $d(v)/d(u) = 1$; $d^*(uv) = 2$ if $d(v)/d(u) = 2$; and $d^*(uv) = 0$, otherwise, has the property that the number of edges labeled with i and the number of edges labeled with j differ by at most 1 for all i and j. A graph that admits a divisor 3-equitable labeling is called a *divisor 3-equitable* graph. The existence and non-existence of divisor 3-equitable labelings have been determined for wheels [1597], complete graphs and stars [1598], ladders, triangular snakes, and lollipop graphs [3298], and the degree splitting graph of ladders [3130].

7.34 Other Kinds of Labelings

Zumkeller labelings

A positive integer n is said to be a *Zumkeller* number if all the positive factors of n can be partitioned into two disjoint parts so that the sum of the two parts are equal. (For example, 6 partitions as $\{1, 2, 3\}$ and $\{6\}$ and 20 partitions as $\{2, 12\}$ and $\{1, 3, 4, 6\}$. An injective function f from the vertices of a graph G to the natural numbers N is said to be a *Zumkeller* labeling of G, if the induced function $f^* : E(G) \to N$ defined by $f^*(xy) = f(x)f(y)$ is a Zumkeller number for all edges xy. A graph that admits a Zumkeller labeling is called a *Zumkeller* graph. Balamurugan, Thirusangu, Thomas, and

Murali [461] investigated the existence of Zumkeller labelings of paths, cycles, and ladders. Wilson and Bebincy [4203] proved that the splitting graph of paths, the total graphs of paths, the shadow graphs of paths, and the middle graphs of paths admit Zumkeller labelings. Wilson and Bebincy investigated the existence of Zumkeller labelings for closed helms, double wheels, sunflowers, flower graphs, and the prisms of wheels. Patodia and Saikia [2807] provided algorithms to label complete bipartite graphs, wheels, cycles, and paths with m-Zumkeller numbers.

A simple graph $G(V, E)$ is said to be a *k-Zumkeller* graph if there is an injection f from the vertices of G to the natural numbers such that when each edge xy is assigned the label $f(x)f(y)$, the resulting edge labels are k distinct Zumkeller numbers. Balamurugan, Thirusangu, and Thomas [460] proved that twig graphs admit a 4-Zumkeller labelings. Basher [557] showed that the super subdivision of paths, cycles, combs, ladders, crowns, circular ladders, grids, and prism are k-Zumkeller graphs. An injection is called a *t-m-Zumkeller* labeling of a graph G if, for every edge uv, the induced function defined by $f^*(uv) = f(u)f(v)$ is an m-Zumkeller number and $|f^*(E)| = t$, where t denotes the number of distinct m-Zumkeller numbers on the edges of G. A graph that admits a $t-m$-Zumkeller labeling is called a *t-m-Zumkeller graph.*. In [2808] Patodia and Saikia gave t-m-Zumkeller labelings for paths, cycles, combs graphs, ladders graphs, and twigs.

A positive integer is said to be a *half-Zumkeller* number if its proper positive divisors can be partitioned into two disjoint non-empty subsets of equal sum. A *half-Zumkeller* labeling of a graph is an injective mapping f from the vertex set into the set of natural numbers N such that the induced mapping f^* from the edges to N given by $f^*(uv) = f(u)f(v)$ is a half-Zumkeller number. A graph that admits a half-Zumkeller labeling is called a *half-Zumkeller* graph. Zeen El Deen, Elmahdy, Elkholy, and El Sherbiny [4310] gave half-Zumkeller labelings for stacked books $(K_{m,1} \times P_n)$, $P_m \times P_n$, prisms of ladders, grids, gears, and flowers. They also showed that if G is a half-Zumkeller graph and H is a non-totally disconnected subgraph of G, then H is a half-Zumkeller graph.

In 2015 Murali, Thirusangu, and Meenakshi [2570] introduced *Zumkeller cordial* labeling of graphs as an injective function f from the vertices to the natural numbers such that the induced function f^* from the edges to $\{0, 1\}$ defined by $f^*(xy) = f(x)f(y)$ is 1 if $f(x)f(y)$ is a Zumkeller number and 0 otherwise with the condition that the number of edges labeled with 1 and the number labeled with 0 differ by at most 1. They proved the existence of Zumkeller cordial labeling for paths, cycles, and stars. Murali, Thirusangu, and Balamurugan [2568] showed the existence of Zumkeller cordial labelings for helms, wheels, flower graphs, and crowns.

Lucky labelings

A labeling of the vertices of a graph with natural numbers is said to be *lucky* if the sum of labels over all neighbors of vertex v and the sum of labels over all neighbors of vertex u are distinct whenever u and v are adjacent. (An isolated vertex is defined to have sum 0). The smallest nonzero label of a lucky labeling of a graph G is called the *lucky number* of the graph and is denoted by $\eta(G)$. A lucky labeling of a graph that is injective is called a *proper lucky* graph. The proper lucky number of graph G is denoted

by $\eta_p(G)$. Sateesh Kumar and Meenakshi [3316] determined the proper lucky numbers for $K_{m,n}$, friendship graphs, certain triangular books, and certain rectangular books. Chiranjilal Kujur, Xavier, and Raja [2052] gave proper lucky labelings for K-identified triangular meshes and K-identified Sierpinski gasket graphs. Ahadi, Dehghan, Kazemi, and Mollaahmadi [95] proved that for a given planar 3-colorable graph G, determining whether $\eta(G) = 2$ is NP-complete, and that for every $k \geq 2$, it is NP-complete to decide whether $\eta(G) = k$ for a given graph G. Using algebraic methods Czerwiński, Grytczuk, and Zelazny [882] proved that $\eta(T) \leq 2$ for every tree T, and $\eta(G) \leq 3$ for every bipartite planar graph G. They obtained a bound for the lucky number in terms of the acyclic chromatic number and conjecture that for every graph G, $\eta(G) \leq \chi(G)$. In [2051] Kujur gave the lucky number and proper lucky number for the bloom graph (a particular planar, tripartite, 4-regular graph). Elrokh, Sateesh Kumar, and Meenaksh [1015] computed the lucky numbers for jelly fish graphs, cocktail party graphs, and crowns. Sateesh Kumar and Meenakshi [3315] determined the lucky numbers and proper lucky numbers for quadrilateral snakes, double quadrilateral snakes, alternate quadrilateral snakes, and double alternate quadrilateral snakes. Akila Devi, Vembu, Jasmine, and Deva Suba [159] gave an algorithm for verifying lucky and proper lucky labelings of complete multipartite graphs with up to three partitions. In [1476] Hussaina, Shabbira, and Ahmada determined the lucky numbers of pentagonal, alternate pentagonal, and double pentagonal snakes; pentagonal books; hairy cycles; Jahangir graphs; and quadrilateral snakes with pendant edges and their alternate forms.

[159] new

[1476] new

In 2015 [2501] Miller, Rajasingh, Emilet, and Jemilet introduced the notion of d-lucky labeling of graphs as follows. For a graph $G(V, E)$, let $f : V(G) \to N$ and $c(u) = \sum_{v \in N(u)} f(v) + d(u)$, where $d(u)$ denotes the degree of u and $N(u)$ denotes the open neighborhood of u. They say that f is *d-lucky* if $c(u) \neq c(v)$ for every pair of adjacent vertices u and v in G. The *d-lucky number* of G, denoted by $\eta_d(G)$, is the least positive k such that G has a d-lucky labeling with $\{1, 2, \ldots, k\}$. They proved that $\eta_d((G) = 2$ for hypercube networks, butterfly networks, benes networks, mesh networks, hypertrees, and X-trees.

Xavier and Rathi [4221] determined the proper lucky numbers of hexagonal meshes and honeycomb networks. Indira, Selvam, and Thirusangu [1536] gave proper lucky labelings for extended duplicate graphs of quadrilateral snakes. In [1528] Imran, Farahani, Cancan, and Alaeiyan showed that various families of combs and snakes admit lucky labelings and determined the exact value of lucky number for the graphs. Kumar and Meenakshi [3318] determined the lucky number of triangular graphs. In [3317] Kumar and Meenakshi proved that jewels, combs, and fans are lucky and proper lucky. Ashwini, Selvam, and Ghanajothi [315] determined the lucky number for complete graphs and complete bipartite graphs and studied the lucky numbers for K_n with an edge removed. In [1549] Irudhaya, Chitra, and Murugan determined the lucky numbers of some star related graphs.

A labeling of the vertices of a graph with natural numbers is said to be a *lucky edge* labeling if for any two adjacent edges the sums of the vertex labels incident to the two edges are distinct. The smallest positive integer k for which a graph G has a lucky edge labeling from the set $\{1, 2, \ldots, k\}$ is called the *lucky number* of G and denoted by $\eta(G)$.

A graph that admits a lucky edge labeling is called a *lucky edge graph*. Esakkiammal, Thirusangu, and Seethalakshmi [1043] showed that the super subdivisions of stars and wheels are lucky edge labeled graphs and determined their lucky numbers. Esakkiammal, Thirusangu and Seethalakshmi [1042] we showed that the H-super subdivision of paths, cycles, and the corona of C_n are lucky edge graphs and they provided their lucky numbers. In [2667] Nellaimurugan and Chitra proved that the planar grid are lucky edge graphs. [1042] new [2667] new

In [2579] Murugan and Chitra provided lucky edge labelings for P_n, C_n and $P_n \odot C_n$. In [2581] they proved that triangular snakes, book with triangular pages, and $P_n \times C_3$ (triangular prisms) are lucky edge graphs. Indira, Selvam, and Thirusangu [1536] gave lucky edge labelings for extended duplicate graphs of quadrilateral snakes. Chitra and Murugan proved that graphs obtained from two copies P_n in which the ith vertex of one path is joined with the $(i+1)$th vertex of the second copy, fish graphs subindexgraphsfish (the one-point union of a vertex of C_n and a vertex of K_3), butterfly graphs, double triangular snakes, flower graphs, and P_n^2 are lucky edge graphs. Nagarajan and Priyadharsini [2643] provided lucky edge labelings and lucky numbers for cars, lotus graphs, prisms, and $C_{2m}@P_t$. Mohana priya and Santhiya [2540] gave lucky edge labelings for graphs obtained by joining C_m and C_n with an edge, bistars, wheels, fans, and butterfly graphs. Murugan and Chitra [2580] gave lucky edge labeling of fans, spider, and twig graphs. Aishwarya [153] provided lucky edge labelings and lucky numbers for ladders, shell graphs, and books with triangular pages.

Murugan, Irudhaya, and Chitra [2587] proved that triangular snakes, books with triangular pages, and triangular prisms are lucky edge graphs. In [2583–2586] they proved that bistars, wheels, parachutes, complete graphs, complete bipartite graphs, fans, spiders, twigs, and H_n graphs, and planar grids are lucky edge graphs. Aishwarya [147,148] proved that helms, flowers, and double stars are lucky edge graphs. Ramya and Shalini [3127] proved that fish tail graphs are lucky edge graphs. [2587] new [2583–2586] new [147,148] new [3127] new

Shalini Rajendra Babu and Ramya [1548] proved that fish graphs, butterflies, double triangular snakes, flowers, and P_n^2 are lucky edge graphs. Ramya and Shalini [3127] proved that fish tail graphs are lucky edge labeled graphs. Murugan, Irudhaya, Chitra [2587] proved that triangular snakes, books with triangular pages, triangular prisms ($P_n \times C_3$) are lucky edge graphs. Sridevi and Ragavi [3688] proved that K_n, tadpoles and rectangular books are lucky edge graphs. Nagarajan and Priyadharshini [2643] proved that lotus graphs, prisms, $C_n@K_3$ are lucky edge graphs. Murugan, Irudhaya, and Chitra [2582] showed that paths, combs, cycles, and crowns are lucky edge graphs. Shalini Rajendra Babu, Ramya, and Rangarajan [3476] proved that splitting graphs of stars, alternative triangular snakes, alternative quadrilateral snakes, and double alternative quadrilateral snakes graphs are lucky edge graphs. Shalini Rajendra Babu and Ramya [3475, 3476] proved that H–graphs, n copies of H-graphs, theta graphs, the path union of theta graphs, and the duplication of theta graphs are lucky edge graphs. [1548] new [3127] new [2587] new [3688] new [2643] new [2582] new [3476] new [3475, 3476] new

Sugapriya and Aishwariya [3724] proposed the concept of lucky edge mean labeling as follows. For a graph $G(V,E)$, a function $f : V \to Z^+$ is said to be a *lucky edge mean* labeling of G if the induced function $f^* : E \to N$ defined by $f^*(uv) = \lceil (f(u) + f(v))/2 \rceil$ has the property that $f(u \neq f(v)$, whenever edges uv and vw are adjacent. They proved [3724] new

that the corona product of paths with paths, the corona product of paths with stars, the corona product of paths with wheels, and the corona product of paths graphs with complete graphs are lucky edge graphs.

Sugapriya, and Vaanmathi [217] introduced the notion of lucky edge geometric mean labeling as follows. For a graph $G(V, E)$, a function $f : V \to Z^+$ is said to be a *lucky edge geometitc mean* labeling of G if the induced function $f^* : E \to Z^+$ defined by $f^*(uv) = \lceil \sqrt{f(u)f(v)} \rceil$ or $\lfloor \sqrt{f(u)f(v)} \rfloor$ has the property that $f(u) \neq f(v)$ whenever edges uv and vw are adjacent. They proved that paths ladders, open ladders, slanting laddesr, stars, degree splitting graphs of stars, bistars, and barycentric edge subdivisions of stars are lucky edge geometric mean graphs. [217] new

In [1944] Khalid, Zhang, Atkinswestley, Sugapriya, Farahani, Alaeiyan, Cancan, Lone, and Akgül introduced the concept of lucky edge odd mean labeling as follows. For a graph $G(V, E)$ with q edges, a *lucky edge odd mean* labeling of G is an injective function $f : V(G) \to \{1, 3, 5, \ldots, 2q + 1\}$ such that each edge uv is assigned the label $f^*(uv) = \frac{f(u)+f(v)}{2}$, and the induced edge labels are exactly $\{2, 4, 6, \ldots, 2q\}$. If such a labeling exists, the graph G is called a *lucky edge odd mean* graph. The least positive odd integer k for which a graph G has a lucky edge odd mean labeling from the set $\{1, 3, 5, \ldots, k\}$ is called the *lucky edge odd mean number* of G. They showed that the middle graph of stars, the total graph of stars, the central graph of stars, the middle graph of bistars, the total graph of bistars, and the central graph of bistars g admit lucky edge odd mean labelings and determined the lucky edge odd mean number of these graphs. [1944] new

S_3 cordial remainder labelings

In [2353] Lourdusamy, Jenifer Wency, and Patrick introduced the concept of the group S_3 cordial remainder labeling as follows. Let $G(V, E)$ be a graph and let $g : V \to S_3$ be a function. For each edge xy assign the label r where r is the remainder when $o(g(x))$ is divided by $o(g(y))$ or $o(g(y))$ is divided by $o(g(x))$ according as $|g(x)| \geq |g(y)|$ or $|g(y)| \geq |g(x)|$. The function g is called a *group S_3 cordial remainder* labeling of G if $|v_g(x) - v_g(y)| \leq 1$ and $|e_g(1) - e_g(0)| \leq 1$, where $v_g(x)$ denotes the number of vertices labeled with x and $e_g(i)$ denotes the number of edges labeled with i $(i = 0, 1)$. A graph G that admits a group S_3 cordial remainder labeling is called a *group S_3 cordial remainder* graph . They prove the following graphs admit a group S_3 cordial remainder labeling: paths, cycles, stars, bistars, complete bipartite graphs, wheels, fans, combs, crowns, the lotus inside a circle, double fans, ladders, slanting ladders, and triangular ladders. In [1647] they proved that shadow graph of cycle and path, splitting graph of cycle, armed crown, umbrella graphs, and dumbbell graphs admit a group S_3 cordial remainder labeling. Also they proved that certain snake related graphs are a group S_3 cordial remainder graphs. In [2354–2356], they investigated the behavior of group S_3 cordial remainder labelings of subdivision of stars, subdivision of bistars, subdivision of wheels, subdivision of combs, subdivision of crowns, subdivision of fans, subdivision of ladders, helms, flower graphs, closed helms, gears, sunflowers, triangular snakes, quadrilateral snakes, square of paths, the duplication of a vertex by a new edge in path and cycle graphs, the duplication of an edge by a new vertex in paths and cycles, and total graphs of cycles and paths.

Pair difference cordial labelings

For a (p,q)-graph $G(V,E)$, let $\rho = \frac{p}{2}$ if p is even and $\frac{p-1}{2}$, if p is odd, and $L = \{\pm 1, \pm 2, \pm 3, \ldots, \pm\rho\}$. Ponraj, Gayathri, and Somasundaram [2875] say G is a *pair difference cordial* if the mapping $f : V \longrightarrow L$ obtained by assigning different labels in L to the different elements of V when p is even, and different labels in L to $p-1$ elements of V and repeating a label for the remaining one vertex when p is odd, has the property that for each edge uv of G labeled with $|f(u) - f(v)|$ it holds that $\left|\Delta_{f_1} - \Delta_{f_1^c}\right| \leq 1$, where Δ_{f_1} and $\Delta_{f_1^c}$ respectively denote the number of edges labeled with 1 and number of edges not labeled with 1. They investigated the pair difference cordial labeling behavior of paths, cycles, stars, bistars, snakes, butterfly graphs, some wheels and path related graphs, splitting graphs, shadow graphs, ladder related graphs, subdivision of graphs the corona of some graphs, mongolian tents, planar grids, star related graphs, $m-$copies of some graphs, Franklin graphs, Heawood graphs, Tietze graphs, and Durer graphs. They further investigated the pair difference cordial labeling behavior of certain trees, certain graphs derived from cube graphs, Petersen graphs $p(n,k)$, and broken wheels in [2872–2881], [2882–2884, 2886]. In [2870, 2871] Ponraj and Gayathri investigated the pair difference snake graphs in cordial labeling behaviors of double alternate snake graphs. Ponraj, Gayathri, and Sivakumar investigated the pair difference cordial labelings of m-copies of paths, cycles, stars, ladders, and the subdivision of wheels and combs. Ponraj, Gayathri, and Somasundaram [2866, 2867] investigated the pair difference cordiality of subdivision of ladders, triangular snakes, alternate triangular snakes, product related to prisms, and torus grids. In [2868, 2869] Ponraj and Gayathri determined the pair difference cordial number of degree splitting graph of bistar, complete bipartite graphs, ladders, wheels, diamond ladder graphs, latitude to ladders, octopus graphs, pagoda graphs, volcano graphs, planter graphs, and semi Jahangir graphs.

In [2909] Ponraj, Prabhu, and Sivakumar introduced the notion of a *pair mean cordial labeling* (PMC-labeling) as follows. Let $G(V,E)$ be a graph with p vertices and q edges. Let $\rho = p/2$, if p is even, $\rho = (p-1)/2$, if p is odd, and let $M = \{\pm 1, \pm 2, \ldots, \pm\rho\}$. Let λ be an injection from V to M, when p is even, and an injection from $p-1$ elements of V to $p-1$ elements of M, when p is odd, and allow a repeat of a label in M for the remaining one vertex. Such a labeling is said to be a *pair mean cordial labeling* if, for each edge uv of G, there exists a label $(\lambda(u) + \lambda(v))/2$, if $\lambda(u) + \lambda(v)$ is even and a label $(\lambda(u) + \lambda(v) + 1)/2$, if $\lambda(u) + \lambda(v)$ is odd such that the number of edges labeled with 1 and the number of edges labeled otherwise differ by at most 1. A graph for which there is a pair mean cordial labeling (PMC-labeling) is called a *pair mean cordial* graph (PMC-graph). In [2909–2913] Ponraj and Prabu investigated the pair mean cordial labeling behavior of paths, stars, cycles, bistars, triangular snakes, alternate triangular snakes, quadrilateral snakes, alternate quadrilateral snakes, middle graphs, splitting graphs, friendship graphs, crowns, double fans, and some corona graphs. They investigated the pair mean cordial labelings of ice cream graphs, closed webs, circulant graphs, zig-zag chord graphs, pentagonal circular ladders, djembe graphs, quadrilateral

friendship graphs, and origami graphs. In [2927] Ponraj, Prabhu, and Sivakumar investigated the pair mean cordial labeling behavior of some graphs that contain cycles. Ponraj and Prabhu [2914–2917] studied the pair mean cordial labeling behavior of Cartesian products of two graphs with components consisting of paths, cycles, wheels, and stars. They also studied the pair mean cordial labeling behavior of hexagonal snakes, irregular quadrilateral snakes, triple triangular snakes, and pair mean cordial graphs paired with ladders, hurdles, key, lotuses and necklace graphs. In [2918–2921, 2936] Ponraj, Prabhu, and Sivakumar investigated pair mean cordial labeling of the total graphs of prism related graphs, some classes of graphs containing cycles, pair mean cordial labeling of pizza graphs, snail graphs, sun graphs, kusudama flower graphs, tree related graphs, the prism of wheel graph, udukkai, octopuses, drums, and fire cracker graphs. In [2919, 2920, 2922–2924] Ponraj, and Prabhu studied pair mean cordiality of certain ladder graphs and some results on pair mean cordial labeling and pair mean cordial labelings of diamond snake graphs, banana graphs, and tortoise graphs. In [2926] Ponraj, Prabhu, and Ramasamy studied pair mean cordial graphs derived from cycles. In [2900] Ponraj, Prabhu, and Ramasamy investigated the PMC-labeling behavior of double fans, triple fans, m–enriched fans, C_n–snakes, stripe blade graphs, armed helms, alternate armed helms, and spectrum graphs. [2900] new

Even sum labelings

Andharia and Kaneria [235] introduced the concept of even sum labeling as follows. A graph $G = (V, E)$ is said to admit *even sum* labeling if there exist an injective function $f : V(G) \to \{0, \pm 2, \pm 4, \ldots, \pm 2|V(G)|\}$ such that the induced mapping $f^* : E(G) \to \{2, 4, \ldots, 2|E(G)|\}$ defined by $f^*(uv) = f(u) + f(v), \forall uv \in E(G)$ is bijective. The function f is called an *even sum* labeling of G. The graph that admits even sum labeling is called an *even sum* graph. In [235], [1829], [1830], and [236] Andharia and Kaneria proved the following graphs are even sum graphs: slanting ladders SL_n $(1 < n < 9)$, P_n, C_{4n}, the complete bipartite graphs, $P_m \times P_n$, mirror graphs $M(P_n)$ $(2 \leq n \leq 6)$, jelly fish graphs, the splitting graph of stars, the degree splitting graph of stars, the splitting graph of $K_{2,n}$, the splitting graph of $K_{1,n,n}$, jewel graphs, triangular books, the triangular book graph with a bookmark, P_m $(+)$ $\overline{K_n}$, and $(\overline{K_n} \cup P_3) + 2K_1$. In [3083] Raj, Krishna, and Mary showed that combs, crowns, $K_m \cup (m-2)K_1$ $(m > 1)$, $K_{m,n} \cup (m-1)K_1$ $(m > 1$ and $W_n \cup 2K_1$ are even sum graphs. In [3082] they proved that paths, stars, and $C_n \cup K_1$ are even sum graphs, whereas cycles are not and in [2030] they obtained some further results on even sum graphs. [3083] new [3082] new

Permuted cordial labelings

In [1009] Elrokh, Al-Shamiri, Almazah, and El-hay used permutation groups together with the concept of cordiality to introduce a new method of labeling. For any set of three distinct even permutations they show how the vertices of a graph can be labeled in such a way that there is an induced edge labeling with the property that for any two distinct vertex labels and any two distinct edge labels, the difference of their labels is at most 1. They call these *permuted cordial* labelings. They proved that paths, cycles, fans,

and wheels, the union of any two paths, and the union of any two cycles admit permuted cordial labelings. They further investigated the permuted cordiality for the union of any path with cycle. Alsatami, Algrawani, and Abd El-hay [205] proved the following graphs admit permuted cordial labelings: $P_n \odot P_m$ $(n, m \geq 2)$; $C_n \odot C_{3m}$ $(n, m \geq 3)$ and $n \neq 2 \pmod 3), m \neq 2 \pmod 3))$; $P_n \odot C_m$; $C_n \odot P_m$; $P_n \odot P_m^2$ $(n, m \geq 2)$; and $P_n \odot C_m^2$ $(n \geq 2),\ m > 4)$.

Odd-even sum labelings

Monika and Murugan [2544] proved the graphs obtained by the following duplications admit odd-even sum labelings: the apex vertex of stars, the pendent vertices of stars, the apex vertex of stars by an edge, a vertex of P_n $(n \geq 2)$, an edge by a vertex of P_n $(n \geq 3)$, a pendent vertex by an edge of P_n $(n \geq 2)$, and a pendent edge of P_n $(n \geq 2)$ by edge, They also proved that C_3 is not an odd-even sum graph.

Prime distance labelings

Eggleton, Erdős, and Skilton, [998] call a graph a *prime distance* graph if its vertices can be labeled with distinct integers in such a way that for any two adjacent vertices, the absolute difference of their labels is a prime number. Laison, Starr, and Walker [2090] prove that trees, cycles, and bipartite graphs are prime distance graphs, and that Dutch windmill graphs and paper mill graphs are prime distance graphs if and only if the Twin Prime Conjecture and the Polignac's Conjecture that for every positive even integer n there are infinitely many cases of two consecutive prime numbers with difference n are true, respectively. Parthiban and David [2783] proved the following graphs admit prime distance labelings: the graph obtained by the duplication of every vertex by an edge in any prime distance graph H if the Twin prime conjecture is true; the graph obtained by fusing any two vertices v_i and v_j, where $d(v_i, v_j) \geq 3$, of C_n; the graph obtained by duplicating any vertex of C_n $(n \geq 6)$; and the graph obtained from m copies of C_n by joining a vertex in the ith copy of C_n to a vertex in the $(i+1)$th copy of C_n where $1 \leq i \leq m-1$.

Liouville difference cordial labelings

In [18] Abiramasundari and Senthil Vadivu introcdued the notion of Liouville difference cordial labelings as follows. Let $G = (V, E)$ be a simple graph with vertex set V and edge set E. A 1-1 function $f : V \to \mathbb{N}$ is said to be a *Liouville difference cordial labeling* of the graph G if the induced edge function $f^* : E \to \{-1, 1\}$ defined by $f^*(uv) = \lambda_{|f(u)-f(v)|}$, satisfies the condition $|e_{f^*}(-1) - e_{f^*}(1)| \leq 1$ for all $uv \in E$, where $e_{f^*}(k)$ is the number of edges labeled with k. They established the existence of such labelings for stars, bistars, twigs, sparklers, and lobsters. In [20] Abiramasundari, Bhuvaneswari, Ilayaraja and Imran proved that $C_n, C_n \odot K_1$ and Dutch windmills admit Liouville difference cordial labelings. Abiramasundari and Senthil Vadivu [19] established that the splitting graphs of paths, combs, ladders, stars, and bistars admit Liouville difference cordial labelings. [18] new [20] new [19] new

2-odd labelings Laison, Starr, and Walker [2090] say a graph G is a 2-*odd graph*

if its vertices can be labeled with distinct integers such that for any two adjacent vertices, the absolute difference of their labels is either an odd integer or 2. They give a characterization of 2-odd graphs in terms of edge colorings and use this characterization to determine which circulant graphs of a particular form are 2-odd and to prove results on circulant prime distance graphs.

Pir and Parathiban [2844] proved that diamonds, double fans, double alternate triangular snakes, *king* graphs, which consist of the $m \times n$ chessboard with edges being the moves that a king chess piece can make, and antiprisms admit 2-odd labelings.

In [22] Abirami, Parthiban, and Srinivasan prove the following graphs admit 2-odd labelings: W_n $(n \geq 4)$, fans, graphs with $2n$ vertices obtained from an n-cycle and joining every two consecutive edges to a new vertex, and shell graphs $C(n, n-3)$ $(n \geq 4)$. Abirami, Parthiban, and Srinivasan [21] proved the following graphs admit 2-odd labelings: helms H_n $(n \geq 4)$, double wheels, umbrellas, graphs obtained by performing the duplication of a vertex by an edge at all the vertices of 2-odd graph if the Twin Prime conjecture is true, $K_a + mK_1$, butterfly graphs $B_{n,m}$ $(n \geq 3)$ if Goldbach's conjecture is true, and graphs obtained from a cycle by performing duplication of a vertex by a vertex at all the vertices of the cycle if Goldbach's conjecture is true.

Absolute difference labelings

In [3477] Shalini and Dhayaran introduced the notion of *absolute differences of cubic and square difference* labelings of a graph as labelings for which every edge label is the absolute difference of the cubes of the vertices and the difference of the squares of the vertices. They proved that paths, stars, cycles, fans, windmills, and wheels admit differences of cubic and squared labelings. They also observed that the labels of the edges must be even. Shalini, Gowri, and Dhayabaran [3478] proved that barbells, cycle cactuses, coconut trees, shells, and dragons admit absolute differences of cubic and squared labelings. In [4051], [4052], [4053], [4054], [4055], [4056], [4057], [4058], [3802], [4059], and [3803] Mathew Varkey and Sunoj proved that planar grids, web graphs, kayak paddle graphs, snake graphs, armed crowns, fans, friendship graphs, windmill graphs, cycles, wheels, gears, helms, 2-tuple graphs, middle graphs, total graphs, shadow graphs barbells, $K_{2,n}^{(m)}$, and $K_{1,n} \odot 2P_m$ admit absolute difference of cubic and square sum labelings.

Quotient labelings

In [3774] Sumathi and Rathi introduced the notion of quotient labeling as follows. Let $G(V, E)$ be a finite, non-trivial, simple,undirected graph. For a one-one assignment $f : V \to \{1, 2, \ldots, |V|\}$ let $f^*(uv) = \lfloor f(u)/f(v) \rfloor$. Then f^* is a said to be a *quotient labeling* of G if $f^* : E \to \{1, 2, \ldots, |V|\}$ and $f(u) > f(v)$. (The edge labels need not be distinct.) The maximum value of $f^*(E(G))$, denoted $ql(f^*)$, is called the *q-labeling number* of G. The *quotient labeling number* of G, $QL(G)$, is the minimum value among all $ql(f^*)$. They found the quotient labeling number of the following graphs: the graph obtained by duplicating an arbitrary vertex of a cycle by an edge, the graph obtained by duplicating an arbitrary edge of cycle by a vertex, two copies C_n sharing a common edge, the twig graphs, the splitting graph of a path, the composition graph $P_m[P_2]$ and joint

sum of two copies of C_n. In [3775] Sumathi and Rathi determine the quotient number of various families of ladder-related graphs. In [3776] Sumathi and Rathi provided the quotient labeling number of quadrilateral snakes, double quadrilateral snakes, alternate triangular snakes, alternate double triangular snakes, and the subdivision of triangular and quadrilateral snakes along the main path.

Edge-sum distinguishing game

The edge-sum distinguishing game (ESD game) is a graph labeling game proposed by Tuza [3922] in 2017. In such a game, the players, traditionally called Alice and Bob, alternately assign an unused label $f(v) \in \{1, 2, \ldots, s\}$ to an unlabeled vertex v of a graph G, and the induced edge label $\phi(uv)$ of an edge $uv \in E(G)$ is given by $\phi(uv) = f(u) + f(v)$. Alice's goal is to end up with an injective vertex labeling of all vertices of G that induces distinct edge labels, and Bob's goal is to prevent this. Tuza also posed the following questions about the ESD game: Given a simple graph G, for which values of s can Alice win the ESD game? And if Alice wins the ESD game with the set of labels $\{1, 2, \ldots, s\}$, can she also win with $\{1, 2, \ldots, s+1\}$? Oliveira, Artigas, Dantas, and Luiz [911] partially answer these questions by presenting bounds on the number of consecutive non-negative integer labels necessary for Alice to win the ESD game on general and classical families of graphs.

Oblong labelings

Oblong numbers are the numbers of dots that can be placed in rows and columns in a rectangular array, each row containing one more dot than each column. (The first five oblong numbers are 2, 6, 12, 20, and 30.) In [3038] Prema and Murugan proved the following graphs have oblong labelings: H-graphs, jelly fish, shrubs, $G \odot K_1$ where G is an H-graph that has an oblong labeling, banana trees, and graphs obtained by identifying an endpoint of $K_{1,3}$ with an end vertex of P_n. Muthumanickavel and Murugan [2616] investigated the existence of oblong sum labelings of the union of graphs involving stars and subdivisions of stars and bistars. They also prove that helms do not admit oblong labelings.

Signed cordial labelings

In 2011 two versions of cordial labelings utilizing -1 and 1 as vertex and edge labels were introduced. Devaraj and Delphy [938] defined the notion of signed cordial graphs as follows. A graph $G(V, E)$ is called *signed cordial* if the edges can be assigned -1 and 1 in such a way that when each vertex v is assigned the product of the labels of the edges incident with v the resulting graph has the property that the number of vertices labeled with i and the number of edges labeled with i differ by at most 1 for $i = -1$ and 1. A graph is called a *signed cordial* graph if it admits a signed cordial labeling. Similarly, Babujee and Loganathan [581] say graph $G(V, E)$ is *signed product cordial* if the vertices can be assigned -1 and 1 in such a way that when each edge assigned the product of its endpoints, the number of vertices labeled with i and the number of edges labeled with i differ by at most 1 for $i = -1$ and 1. A graph is called *signed product cordial* if it admits a signed product cordial labeling. In [2628] Nada, Elrokh, Elmshtaye,

and El-hay gave necessary and sufficient conditions for which cones $(\overline{K_n} + C_m)$ and their second powers are signed product cordial. In [581] Babujee and Loganathan proved that paths, cycles, stars, and bistars are signed product cordial. Devaraj and Delphy [938] investigated signed-cordiality of complete graphs, books, Jahangir graphs, flower graphs, and Petersen graphs. Cynthia and Padmavathy [878] proved that $C_n \times P_n$ and certain banana trees admit signed cordial and signed product cordial labelings. In [5] Abdel-Aal and Bashammakh proved that a graph G is cordial if and only if it is signed product cordial.

In [879] Cynthia and Padmavathy proved that Bloom graphs and torus networks admit signed cordial labelings and signed product cordial labelings. Santhi and Albert [3305] gave signed product cordial labeling for some cycle related graph operations. Rilwan and Hussain [3173] investigated the signed product cordial, square difference, and analytic mean labelings for the identity graph. In [**?**] Ghosh and Pal proved that combs, webs, complete bipartite graphs, jewels, gears, quadrilateral snakes, triangular snakes, tadpoles, and shells are signed product cordial. Raj and Manoharan [3085] proved that the graph obtained by arbitrary supersubdivisions of $(P_n \times P_m) \odot P_s$ are signed product cordial. They also gave conditions for which tadpoles and $C_n \odot P_m$ are signed product cordial. Ghosh and Pal [1270] provided signed product cordial labelings and total signed product cordial labelings for the Cartesian product of balanced bipartite graphs and paths. They showed that the time complexity of their algorithms is superlinear. Nada, Elrayes, Elrokh, and Rabie [2626] showed that the union and the join of any two fourth power of paths are always signed product cordial. They also showed the join and union of fourth power of cycles are signed cordial in some cases. In [3924] Ulaganathan, Selvam and Kumar proved that the duplicate graph of triangular ladders, the extended duplicate graph of bistars, and the extended duplicate graph of the double stars admit signed product cordial labelings.

k-vertex weighting

In [3523] Shiu, Laub, and Ng [3523] introduced the following notion. For a simple, finite and undirected graph $G(V, E)$ of order n, a k-vertex weighting of a graph G is a mapping $w : V(G) \to \{1, \ldots, k\}$. A k-vertex weighting induces an edge labeling $f_w : E(G) \to N$ such that $f_w(uv) = w(u) + w(v)$. Such a labeling is called an *edge-coloring k-vertex weighting* if $f_w(e) \neq f_w(e')$ for any two adjacent edges e and e'. They determined the minimum k that admit an edge-coloring k-vertex weighting for the following graphs: wheels, gears, grids, $P_2 \times C_n$, lollipops, theta graphs, and two cycles joined by a path (*long dumbbells*).

Harmonic mean cordial

Let f be a function from $V(G)$ to $\{1, 2\}$. For each edge uv of G, assign the label $\lfloor 2(f(uf(v))/(f(u) + f(v))\rfloor$. Such an f is a *harmonic mean cordial labeling* of G if $|v_{f(i)} - v_{f(j)}| \leq 1$ for i and j in $\{1, 2\}$, where $v_{f(x)}$ and $e_{f(x)}$ denote the number of vertices and edges labeled with x, respectively. A graph with a harmonic mean cordial labeling

is called a *harmonic mean cordial graph*. In [2772] Parejiya, Jani, and Hathi, proved that $CH_n \odot K_1$ (CH_n is the closed helm obtained from W_n) and the tensor product of two paths admit harmonic mean cordial labelings. They also proved that $K_{m,n}, K_n \vee C_m$ (join of $K_{m,n}$ and C_n) and $C_n \vee C_m$ are not do not admit harmonic mean cordial labelings. Parejika, Lalchandani, Jani, and Mundadiya [2773] proved the splitting graphs of the following graphs are harmonic mean cordial: P_n (n even), stars, bistars, combs, and $P_n \odot \overline{K_m}$. In [1574] Jani, Lalchandani, and Parejiya investigated the existence of [1574] new harmonic mean cordial labelings of $C_n \vee W_m$ and $K_n \vee W_m$.

Tri sum perfect square cordial labeling

A graph $G(V, E)$ is said to be *tri sum perfect square cordial* if there exists a function f from V to $\{-1, 0, 1\}$ such that the function f^* from E to $\{0, 1\}$ defined by $f^*(uv) = f(u)^2 + 2f(u)f(v) + f(v)^2$ has the property that the number of edges labeled with 0 and the number of edges labeled with 1 differ by at most 1. Such an f is called a *tri sum perfect square cordial* labeling of G. In [3945] Vaghela and Parmar gave tri sum perfect square cordial labelings for paths, combs, stars, fans, the H-graphE of P_n, the $H \odot K_1$ graph of P_n, and the union of two stars.

Radio contra harmonic mean labeling

In [308] Ashika and Asha defined the *radio contra harmonic mean labeling* of a connected graph $G(V, E)$ as an injective function $f : V(G) \to Z$ such that for each pair of distinct vertices u and v of G, $d(u, v) + \left\lceil \frac{f(u)+f(v)}{2} \right\rceil \geq 1 + \text{diam } (G)$ or $d(u, v) + \left\lfloor \frac{f(u)+f(v)}{2} \right\rfloor \geq 1 + \text{diam } (G)$. The *radio contra harmonic mean number* of f is the maximum number assigned to any vertex of G. The *radio contra harmonic mean number* of G is the minimum value of the radio contra harmonic mean number taken over all radio contra harmonic mean labelings f of G. Ashika and Asha determined the radio contra harmonic mean numbers and labelings of $G_n \odot K_1$ (G_n is the gear graph), $(W_n \odot mK_1) \odot K_1$, the subdivision of jellyfish, and the path union for two copies of the bistar graph $B_{m,n)}$).

SD-harmonious labelings

For a graph G with q edges, an injection $f : V(G) \to \{0, 1, 2, \ldots, q\}$ is said to be SD-*harmonious labeling* if the induced function $f^* : E(G) \to \{0, 2, \ldots, 2q - 2\}$ defined by $f(uv) = S + D \pmod{2q}$ is bijective, where $S = f(u) + f(v)$ and $D = |f(u) - f(v)|$, for every edge uv in $E(G)$. In [2351] Lourdusamy, Wency, and Patrick investigated SD-harmonious labelings of paths, trees, star related graphs, and the disjoint union of graphs.

Dividing graceful labelings

Zahraa, Nabeel, and Fawzi [4300] say a graph $G(V, E)$ has a *dividing graceful labeling* if there is a one-to-one mapping ϕ from V to $\{1, 2, 3, \ldots, |E|\}$ such that for every edge uv the induced function ϕ^* defined by $\phi^*(uv) = \lceil (\phi(u) + \phi(v))/|E| \rceil$ has the property that $\phi^*(E) = \{1, 2, 3, \ldots, |E|\}$. They showed the following graph have dividing graceful

labelings: stars, graphs obtained by joining the centers of two copies of $K_{1,t}$, and spiders with diameter 4.

Square product labelings

In [2511] Mirajkar and Sthavarmath introduced the notion of a *square product* graph as one for which there exists a bijection f from $V(G)$ to $\{1, 2, 3, \ldots, p\}$ that induces an injective labeling f^* from $E(G)$ to N defined by $f^*(uv) = f(u)^2 f(v)^2$. In [2510] they proved that $C_n \odot K_1$ and certain kinds of cactus graphs are square product graphs.

Numbering of graphs

In [1517] Ichishima, Oshima, and Takahashi define a *numbering* f of a graph G of order n as a labeling that assigns distinct elements of the set $\{1, 2, \ldots, n\}$ to the vertices of G. The strength $\mathrm{str}_f(G)$ of a numbering $f : V(G) \to \{1, 2, \ldots, n\}$ of G is defined by $\mathrm{str}_f(G) = \max\{f(u) + f(v) \mid uv \in E(G)\}$, that is, $\mathrm{str}_f(G)$ is the maximum edge label of G and the strength $\mathrm{str}(G)$ of a graph G itself is $\mathrm{str}(G) = \min\{\mathrm{str}_f(G) \mid f \text{ is a numbering of } (G)\}$. They proved necessary and sufficient conditions for the strength of a graph G of order n to meet $\mathrm{str}(G) = 2n - 2\beta(G) + 1$ and $\mathrm{str}(G) = n + \delta(G) = 2n - 2\beta(G) + 1$, where $\beta(G)$ and $\delta(G)$ denote the independence number and the minimum degree of G, respectively. They answer open problems posed by Gao, Lau and Shiu, and the earlier result leads us to determine a formula for the strength of graphs containing a particular class of graphs as a subgraph. They also extend what is known in the literature about k-stable properties. In [1484] Ichishima, López,

In [1502] Ichishima, Muntaner-Batle, Oshima, and Takahashi gave a sharp lower bound for the strength of a graph in terms of its domination number as well as its (edge) covering and (edge) independence number. They also provided a necessary and sufficient condition for the strength of a graph to attain a bound in terms of their subgraph structure and established a sharp lower bound for the domination number of a graph under certain conditions. In [1507] Ichishima, Muntaner-Batle, and Takahashi used the concept of independence number of a graph to determine formulas for the strength of powers of paths and cycles. In [1508] Ichishima, Muntaner-Batle, and Takahashi study general conditions for graphs that allow one to determine which graphs have the property that lower and upper bounds for the strength coincide. In [1510] Ichishima1, Muntaner-Batle, and Takahashi use the concept of independence number of a graph to determine formulas for the strength of powers of paths and cycles.

In [1518] Ichishima, Oshima, and Takahashi define an *edge numbering* f of a graph G of size m as a labeling that assigns distinct elements of the set $\{1, 2, \ldots, m\}$ to the edges of G. The *edge-strength*, $\mathrm{estr}(G)$, of G is defined as the minimum $\mathrm{estr}_f(G)$ where $\mathrm{estr}_f(G) = \max\{f(e_1) + f(e_2) \mid e_1, e_2 \text{ are adjacent edges of } G\}$ over all edge numberings of G. They present formulas for $\mathrm{estr}(G)$ when G is the forest whose components are stars of order at least three and the complete bipartite graph whose partite sets consist of at least two vertices. The edge-strength of a graph G is the strength of the line graph of G, and thus this work extends what was known about the edge-strength and strength.

Fibonacci range labelings

A simple finite graph G with p vertices and q edges is said to be a *Fibonacci range* graph if there is a bijective $f : V(G) \to \{F_2, F_3, F_4, \ldots, F_{p+1}\}$, where F_i is the ith Fibonacci number, such that the induced edge labels given by $f^*(uv) = \lceil (f(u)^2 + f(v)^2)/(f(a) + f(b)) \rceil$ or $f^*(uv) = \lfloor (f(u)^2 + f(v)^2)/(f(a) + f(b)) \rfloor$ can be assigned subscripts $1, 2, \ldots, n$ in such a way that the ratios $e - 1/e_2, e_2/e_3, \ldots, e_{n-1}/e_n$ approach the golden ratio $\phi = 1.618\ldots$. In [2735] Odyuo and Mercy classified k-copies of shell graph $C[n, (n-3)]k$ with a union of K_2 for $n = 4$, $2K_2$ for $n = 6$, and $3K_2$ for $n = 8$ having a common end vertex joined to the apex of the shell are Fibonacci range graphs.

Super arithmetic graceful labelings

A connected graph is said to be *highly irregular* if each of its vertices is adjacent only to vertices with distinct degrees. Let $H_i(m, n)$ denote the bipartite graph of order $n = 2m$ $(m \geq 2)$ having partite sets, $V_1 = \{u_1, u_2, \ldots, u_m\}$ and $V_2 = \{v_1, v_2, \ldots, v_m\}$ and edge set $E(H) = \{u_i v_j : 1 \leq i \leq m, 1 \leq j \leq m + 1 - i\}$ with $\deg H(u_i) = \deg H(v_i) = m + 1 - i$ for $i = 1, 2, \ldots, m$. The highly irregular graph $H_i^{(1)}(m, m)$ of order $2m + 1 \geq 9$ is obtained by subdividing the edge $u_2 v_{m-1}$ of $H_i(m, m)$ for $m \geq 4$. An *E-super arithmetic graceful* labeling of a graph G is a bijection f from the union of the vertex set and edge set to $\{1, 2, 3, \ldots, |V(G) \cup E(G)|\}$ such that the edges have the labels from the set $\{1, 2, 3, \ldots, |E(G)|\}$ and the induced mapping given by $f(uv) = f(u) + f(v) - f(uv)$ for $uv \in E(G)$ has the range $\{|V(G) \cup E(G)| + 1, |V(G) \cup E(G)| + 2, \ldots, |V(G)| + 2|E(G)|\}$. In [268] Anubala and Ramachandran proved that $H_i(m, m), H_i^{(1)}(m, m)$, and chain of even cycles $C_{4,n}, C_{6,n}$ are E-super arithmetic graceful.

E-super arithmetic graceful labelings

In [3339] Sekar and Varatharajaprumal define a (p, q) graph G to be *E-super arithmetic graceful* if there exists a bijection f from $V(G) \cup E(G)$ to $\{1, 2, \ldots, p + q\}$ such that $f(E(G)) = \{1, 2, \ldots, q\}$, $f(V(G) = \{q + 1, q + 2, \ldots, q + p\}$, and the induced mapping f^* given by $f^*(uv) = f(u) + f(v) - f(uv)$ for $uv \in E(G)$ has the range $\{p + q + 1, p + q + 2, \ldots, p + 2q\}$. They provided E-super arithmetic graceful labelings for four families of cycle related graphs.

Gaussian tribonacci r-graceful labelings

For natural number r, Sunitha and Sheriba [3799] say an injective function ϕ from $V(G)$ to $\{0, ki, 1 + ki, 2 + ki, \ldots, GT_{q+r-1}\}$ for all k, where GT_{q+r-1} is the $(q + r - 1)^{th}$ Gaussian tribonacci number in the Gaussian tribonacci sequence is a *Gaussian tribonacci r-graceful labelings* if the induced edge labeling $\phi^* : E(G) \to \{GT_1, GT_2, \ldots, GT_{q+r-1}\}$ is bijective. A graph that admits Gaussian tribonacci r-graceful labeling is called *Gaussian tribonacci r-graceful* A graph is said to be *Gaussian tribonacci arbitrarily graceful* if it is Gaussian tribonacci r-graceful for all r. They prove that paths, combs, coconut trees, regular caterpillars, bistars, and the subdivision of bistars are Gaussian tribonacci arbitrarily

graceful for all nontrivial cases.

Radio heronian mean k-graceful labelings

For a graph G and a positive integer k a mapping g from $V(G)$ to $\{k, k+1, \ldots, k+N-1\}$ is a *radio heronian mean k-labeling* if for any two distinct vertices s and t of G, $d(s,t) + \lceil (g(s) + g(t) + \sqrt{g(s)g(t)}\ )/3 \rceil \geq 1 + D$, where D is the diameter of G. The radio heronian mean k-number of g, $rrhmn_k(g)$, is the maximum number assigned to any vertex of G. The *radio heronian mean number* of $G, rhmn_k(g)$, is the minimum value of $rhmn_k(g)$ taken overall radio heronian mean labelings g of G. If $rhmn_k(g) = |V(G)| + k - 1$, G is said to be a *radio heronian mean k-graceful.*. In [3800] Sunitha and Rani investigated the radio heronian mean number of $T_n \odot K_1$ (obtained by joining a single pendent edge to each vertex of a triangular snake), $IT_n \odot K_1$ (IT_n is the irregular triangular snake), and $Q_n \odot K_1$ (Q_n is the quadrilateral snake). In [3801] Sunitha and Rani proved that the degree splitting of graphs of combs, rooted trees, graphs obtained from a path by attaching a pendent edge to every internal vertex of the path (*hurdle graphs*) and twigs admit radio heronian mean k-graceful labelings.

Heronian Dd-distance mean labelings

In [692] Bosco and Dinesh introduced the concept of radio heronian Dd-distance mean labeling graphs as follows. A *radio heronian Mean Dd-distance labeling* of a connected graph G is an injective map f from the vertex set to the positive integers such that for two distinct vertices u and v of G we have $D^{Dd}(u,v) + \lceil (f(u) + \sqrt{f(u)f(v)} + f(v))/3 \rceil \geq 1 + diam^{Dd}(G)$, where $D^{Dd}(u,v)$ denotes the Dd-distance between u and v and $diam^{Dd}(G)$ denotes the Dd-diameter of G. The *radio heronian Dd-distance number* of f ($rhmnDd(f)$) is the maximum label assigned to any vertex of G. The *radio heronian Dd-distance number* of G ($rhmnDd(G)$ is the minimum value of $rhmnDd(f)$ taken over all radio heronian Dd distance labeling of G. They determined radio heronian mean Dd-distance number of complete graphs, paths, stars, subdivisions of stars, and fans. In [691] Bosco and Dinesh determined the radio heronian mean Dd-distance number of degree splitting stars, flower graphs, shadow graphs, switching wheels, and switching helms.

Fibonacci graceful antimagic graph

In [953] Dharman and Murugesan call *Fibonacci graceful antimagic graph* G with q edges as one that has an injective function $\phi : V(G) \to \{0, 1, 2, \ldots, F_{q+1}\}$ (F_i is the ith Fibonacci number) such that the induced edge labeling $\phi^*(uv) = |\phi(u) - \phi(v)|$ is a bijection onto the set $\{F_2, F_3, \ldots, F_{q+1}\}$, the vertex sums are pairwise distinct, and all the edges labels are distinct. They showed that the following graphs admit Fibonacci graceful antimagic labelings: $C_n \odot K_1$, double stars, bistars, and graphs obtained by joining the centers of two copies of $K_{1,n}$ with an edge. some star related graphs have such labelings. In [2526] Mitra and Bhoumik proved that $C_n \odot K_t$ is Fibonacci graceful for $t = 1, 2$, and 3

In [3480] Shanmuga and Amuthavalli introduced the concept of a *super Fi-*

bonacci graceful antimagic graph G with q edges as one that has an injective function $\phi : V(G) \to \{0, F_2, F_3, \ldots, F_{q+1}\}$ (F_i is the ith Fibonacci number) such that the induced edge labeling $\phi^*(uv) = |\phi(u) - \phi(v)|$ is a bijection onto the set $\{F_2, F_3, \ldots, F_{q+1}\}$, the vertex sums are pairwise distinct, and all the edges labels are distinct. They proved that following graphs have such labelings: graphs obtained by joining n copies of $C_4 + e$ and m copies K_2 with a common vertex (*clematis flower*); graphs obtained by joining n copies of C_3 and m copies K_2 with a common vertex (*cherry blossom*), and graphs obtained by joining n copies of C_6 with a common vertex (*rose*).

Radio even mean graceful graphs

A *radio even mean graceful labeling* of a connected graph G is a bijection π from the vertex set $V(G)$ to $\{2, 4, 6, \ldots, 2|V|\}$ satisfying the condition $(d(s,t) + \lceil \phi(s) + \phi(t) \rceil)/2 \geq 1 + diam(G)$ for every s and t in $V(G)$. A graph that admits radio even mean graceful labeling is called a *radio even mean graceful* graph. radio even mean. In [710] Brindha Mary, Raj, and Jayasekaran investigate the radio even mean graceful labeling on degree splitting of some special graphs.

Sudoku number of graphs

In [2100] Jeyaseeli, Lau, Shiu, and Arumugam introduced a new concept of Sudoku number as follows. Let G be a graph with chromatic number k. The minimum number of vertices in G that have been pre-colored such that this pre-coloring can be extended uniquely to a k-coloring of G is called the *Sudoku number* of the graph G. They gave the Sudoku numbers of the amalgamations of complete graphs, tadpole graphs, lollipop graphs, and graphs obtained by identifying every alternate edge of C_{2n} with an edge of a distinct complete graph K_m and its related graph.

Prime graceful

In 2018 Selvarajan and Subramoniam [3345] introduced the notion of prime graceful labelings. For a graph $G(V, E)$ with p vertices and q edges they say an injection from V to $\{1, 2, \ldots, k\}$, where $k = \min(2p, 2q)$, that has the property that, for each edge uv, $\gcd(f(u), f(v)) = 1$ and the induced function f^* from E to $\{1, 2, \ldots, k-1\}$ defined by $f^*(uv) = |f(u)) - f(v))|$ is an injection is a *prime graceful* labeling A graph that admits a prime graceful labeling is said to be a *prime graceful* graph. They proved that paths, cycles, stars, friendship graphs, bistars, $C_4 \cup P_n$, $K_{m,2}$ and $K_{m,2} \cup P_n$ admit prime graceful labelings. Nandhini and Lakshmi [2650] proved that cycles with one pendent edge, helms, and certain cases of triangular snakes are prime graceful.

In 2024, the notion of total prime graceful labeling technique was introduced by Keerthika and Kowsalya [1947] as follows. A *total prime graceful labeling* of a graph G is prime graceful labeling such that for each edge uv, the labels assigned to u and v and uv are relatively prime. Total prime graceful labeling facilitates the identification of security threats within the network infrastructure. In [2002] Keerthika and Kowsala discussed total prime graceful labelings of a star graph in the context of a VPN clients and access control systems.

In 2021 Panma and Rochanakul [2759] introduced a different version for prime graceful graphs. A graph G with n vertices and m edges is said to be *prime-graceful* if there is an injection $\psi : V(G) \to \{1, 2, \ldots, m+1\}$, where $\gcd(\psi(u), \psi(v)) = 1$ for all $e = \{u, v\} \in E(G)$ and the induced function $\psi^* : E(G) \to \{1, 2, \ldots, m\}$ defined as $\psi^*(e) = |\psi(u) - \psi(v)|$ is injective. They showed stars, bistars $B_{n,n}$, bistars $B_{n,p-2}$, where p is an odd prime, $K_{2,n}$, tristars $SL(3, n)$, triangular books $B_n^{(3)}$, and some spiders are prime graceful. They also study k-prime-graceful labelings where the range of ψ is extended to $\min\{kn, km\}$ for $k > 1$.

LH graphs

In [1069] Farisa and Parvathy introduced the notion of LH graphs as follows. A graph G with n vertices is said to have an *LH* labeling if there exists a bijective function $f : V(G)$ to $\{1, 2, 3, \ldots, n\}$ such that the induced map f^* from $E(G)$ to set of natural numbers defined by $f^*(uv) = LCM(f(u), f(v))/HCF(f(u), f(v))$ is injective (LCM and HCF denote the least common multiple and highest common factor, respectively). A graph that admits an *LH* labeling is called an *LH graph*. They proved that theta graphs, sparklers, H graphs, and the splitting graphs of paths and combs are LH graphs. In [1070] they proved that combs, helms, triangular snakes, quadrilateral snakes, twigs, and bistars are LH graphs. Obaed and Arif [2734] proved that cycles, cactus graphs, coconut trees, combs, triangular snakes, and $K_2 \odot C_n$ admit LH-labelings.

k-distant edge total labelings; k-distant vertex total labelings; k-distant edge chromatic number

In [3129] Rana introduced the following graph labelings. Given an undirected connected graph $G(V, E)$, a labeling f from $V \cup E$ to $\{1, 2, \ldots, r\}$ is called a k-distant vertex total labeling if $|w(x) - w(y)| \geq k$ for every edge xy, where $w(x) = f(x) + \sum(x, y)$ over all edges xy is the weight of the vertex x. The minimum r for which a graph G has a k-distant vertex total labeling is called the *total k-distant chromatic number* of G and is denoted by $\gamma_{kd}(G)$. A labeling f from $V \cup E$ to $\{1, 2, \ldots, r\}$ is called a *k-distant edge total labeling* if for any two adjacent edges e_1 and e_2 we have $|w(e_1) - w(e_2)| \geq k$, where $w(e) = f(e) + f(x) + f(y)$ and $x, y \in V$ are the two end vertices of the edge e. The minimum r for which a graph G has a k-distant edge total labeling is called the *total k-distant edge chromatic number* of G and is denoted by $\gamma'_{kd}(G)$. Rana determines the k-distant vertex total labeling for paths, cycles, complete graphs, stars, bistars, friendship graphs, and the value of the parameter γ_{kd} for these graph classes. Moreover, k-distant edge total labelings for paths, cycles and stars are investigated and an upper bound of γ_{kd} and a lower bound of γ'_{kd} are presented.

Super totient labelings

A positive integer n is called *super totient* if the residues of n that are prime to n can be partitioned into two disjoint subsets of equal sums. An injective function f defined on the vertices of a graph $G(V, E)$ to a subset of integers is called a *super totient labeling* if the function f^* from E to the natural numbers defined by $f^*(xy) = f(x)f(y)$ assigns

a super totient number for all edges xy of G. A graph that admits such a labeling is called a *super totient graph*. In [2403] Mahmood and Ali proved that every graph with at least one edge admits a super totient labeling. If the range of a super totient labeling f is $\{1, 2, \ldots, |V|\}$, then f is said to be a *restricted super totient labeling*. They provided super totient labelings for friendship graphs, wheels, complete graphs, and complete bipartite graphs. In [1375] Harrington and Wong classified all restricted super totient labelings of complete bipartite graphs, trees, wheel graphs, and friendship graphs. Furthermore, they introduced the notion of the *sum index* of a graph G, which is the minimum positive integer k such that there exists an injective vertex labeling f of G with k being the cardinality of the range of f. They showed that the sum index is related to super totient labelings, and determined the sum index of complete graphs, complete bipartite graphs, and certain families of trees including caterpillar graphs.

Harrington, Henninger-Voss, Karhadkar, Robinson, and Wong [1377] defined the *difference index* as the minimum value of $|f(u) - f(v)|$ taken over all injective labelings f of the vertices of the graph. They provided upper and lower bounds on the sum index and difference index, and determine the sum index and difference index of various families of graphs. In [1383] Haslegrave investigated two conjectures made by Harrington, Henninger-Voss, Karhadkar, Robinson and Wong in [1376] concerning relationships between the sum index, the difference index, and the exclusive sum number of graphs. One conjecture posited an exact relationship between the first two invariants. Haslegrave showed that the predicted value may be arbitrarily far from the truth in either direction. He also established some new bounds on both the sum index and the difference index. The other conjecture, that the exclusive sum number can exceed the sum index by an arbitrarily large amount, follows from known non-constructive results, whereas Haslegrave gave an explicit construction demonstrating it. Harrington et al. [1377] updated their preprint [1376] with two counterexamples to the first conjecture giving a discrepancy of 1, and only in one direction and therefore modified the conjecture from an equality to an inequality. Haslegrave results showed that the modified is still false in general. Harrington, Henninger-Voss, Karhadkar, Robinson, and Wong [1377] defined the *difference index* as the minimum value of $|f(u) - f(v)|$ taken over all injective labelings f of the vertices of the graph. They provided upper and lower bounds on the sum index and difference index, and determine the sum index and difference index of various families of graphs.

[1377] new
[1383] new
[1376] new
[1377] new
[1376] new
[1377] new

Zhang and Wang [4322] improved some bounds in given by Harrington et al.[1377] on the sum and difference indices concerning the relationships betweeen these two indices between a graph and its induced subgraph. For regular graphs, they made slight generalizations of Haslegrave's bounds on the sum and difference indices in [1383]. They gave the sum and difference indices of necklace graphs, the complements of matchings, cycles, paths and compared the sum indices with (integral) sum numbers and analyzed the sum and difference indices of regular graphs. They conclued with some conjectures and open questions.

[4322] new

In [929] Desai and Wang determined the sum indices of the complete multipartite graphs, hypercubes, and some cluster graphs. They also studied the maximum number of edges in a graph with a fixed sum index.

[929] new

Fuzzy vertex-antimagic labelings

A *fuzzy vertex anti-magic* (FVAM) labeling of a graph G is a 1-1 correspondence $f : E(G)$ to $\{1, 2, \ldots, |E(G)\}$ such that for any two different vertices v and w, the total of the labels on edges incident to v are distinct from the total of labels on edges incident to w. In [3352] Selvarasu and Murugan proved that a graph that is formed by joining two terminal edges to every internal vertex of the path (*twing*) and combs admit FVAM labeling by providing an algorithm.

Relatively prime edge labelings

For a graph $G(V, E)$ Janani, and Ramachandran [1569] say a bijection from E to $\{1, 2, \ldots, |E|\}$ is a *relatively prime edge* labeling of G if, for each vertex v, the labels of the edges incident to v are pairwise relatively prime. A graph that admits a relatively prime edge labeling is called a *relatively prime edge labeled* graph. For a relatively prime edge labeled graph G the *edge prime index* of G is the minimum number of edges needed to be removed from G to result in a relatively prime edge labeled graph. They showed that certain trees admit relatively prime edge labelings. In [1571] Janani and Ramachandran provided the edge prime indexes for $K_n \odot K_1, K_{2,n}$ $(n > 2), K_{3,n}$ $(n > 3)$, ladders L_n $(n > 3)$, and generalized Petersen graphs $P(n, k)$ where n and k are even. In [**?**] Janani and Ramachandran gave the following definition. A graph $G(V, E)$ is called a *coprime edge* labeling if there is a bijection $f : E$ to $\{1, 2, \ldots, k\}$ such that for $k \geq q$ and each vertex v, the labels of the edges incident to v are pairwise relatively. The *prime index* of a coprime graph G is the smallest number of edges needed to be removed from G to result in a prime graph. They determine the edge prime indexes of K_n, wheels $C_n \odot K_1$ (n odd), $K_n \odot K_1$, and $P_{2n} + P_2$. In [1974] Komarullah, Wijaya, Krisnawati, Hidayat investigated prime edge labeling and coprime edge labeling for tadpoles, combs, disjoint unions of combs, disjoint unions of two cycles, disjoint unions of paths, volcano graphs, and regular caterpillars. Furthermore, they provided precise values for the minimum coprime edge number of $C_m \cup C_n$ with odd m and n, volcano graphs, and regular caterpillars. [1974] new

Mean harmonic energy

In [178] Al-Roainee and Mahde introduced the concept of mean harmonic energy of a graph as follows. For a simple connected graph $G(V, E)$ with vertices $v_1, v_2, \ldots, v_n$, let d_i be the degree of v_i. The *mean harmonic matrix* of G is the $n \times n$ matrix whose (i, j)-entry is $a_{ij} = 2/(1/d_i + 1/d_j)$ if $v_i v_j$ is an edge of G and 0 otherwise. The *mean harmonic energy* of G is the sum of the absolute values of the set of eigenvalues of the mean harmonic matrix of G. They obtained bounds for mean harmonic energy of graphs and gave there exact value for complete graphs, complete bipartite graphs, stars, crowns, and friendship graphs.

Diophantine prime graphs

In [2658] Nasr, Elsonbaty, Seoud, and Anwar gave a generalization for the concept of prime graphs as follows. A graph $G(V, E)$ with n vertices is said to be a *Diophantine*

prime graph if there exists a bijective map f from V to $\{1, 2, \ldots, n\}$ such that for every edge uv, $\gcd(f(u), f(v))$ divides n. Clearly, any prime graph is a diophantine graph and every diophantine graph with a prime number of vertices is a prime graph. For a maximal Diophantine graph with n vertices, they gave explicit formulas of the number of edges and the degree of a vertex and determined the full degree vertices. They also provided necessary and sufficient and conditions that determine the equality of degrees among non-full degree vertices in the maximal ophantine graphs.

δ-labelings

In 2021 [3434] Sethuraman and Murugan introduced the notion of a δ-labeling of a simple graph in their investigation of decomposing complete graphs into arbitrary trees as follows. Let G be a graph with m edges. An injection $f : V(G)$ to $\{0, 1, 2, \ldots, 4m\}$ is said to be a *δ-labeling* of G if the induced function $f' : E(G)$ to $\{1, 2, \ldots, 2m\}$ has the property that for each edge uv, $f'(uv) = |f(u) - f(v)|$, if $|f(u) - f(v)| \leq 2m$ and $4m + 1 - |f(u) - f(v)|$, if $|f(u) - f(v)| > 2m$ is injective. They used δ-labelings to study how to decompose complete graphs into arbitrary trees. They proved that all trees admit δ-labeling and asked what graphs admit δ-labelings. In [3415] Sethukkarasi, Vidyanandini, and El-Mesady proved the following graphs admit δ-labeling: double stars; double fans; graphs obtained by joining the apexes of two stars with an edge; $P_{2n-1} \odot 2nK_1$; complete bipartite graphs; cycles; cycles with one appended edge; graphs obtained by joining two copies of C_n by an edge; $C_n \times P_2$; and $C_n \odot K_1$ admit δ-labelings. They discussed how δ-labelings can be used for DNA sequence alignment.

Super vertex sum labeling

For a graph G, let f be a bijective function from $V(G)$ onto $\{1, 2, \ldots, |V(G)|\}$. Joseph and Kureethara [1798] say f is a *super vertex sum labeling* of G if, for every pair of adjacent vertices $u, v \in V(G)$, there is some $w \in V(G)$ such that $f(w) = f(u) + f(v)$ and the maximum value of $f(u)$ over all vertices $u \in V(G)$ is $|V(G)|$. The smallest number of isolated vertices that is needed for a graph G to have a super vertex sum labeling is called *super vertex sum number* of G and is denoted by $\mu_{sv}(G)$. A super vertex sum labeling that uses $\mu_{sv}(G)$ isolated vertices is said to be *optimal* . They determined the lower bound of $\mu_{sv}(G)$ and gave optimal super vertex sum labelings for the super subdivision of paths, cycles, stars, and spiders. In [1799], they proved that the lower bound of $\mu sv(G)$ is $\delta(G)$ and showed that super subdivision of paths, cycles, stars, and spiders are super vertex sum graphs with $\mu_{sv}V(G) = 2$.

AUM block labelings

In [3930] Uma Maheswari and Azhagarasi introduced the notion of AUM block labeling as follows. Let G be a graph with p vertices, q edges, and b blocks, where $p, q, b \geq 1$. Let $V(G) = \{v_1, v_2, \ldots, v_p\}$, $E(G) = \{e_1, e_2, \ldots, e_q\}$, and $B(G) = \{B_1, B_2, \ldots, B_b\}$ denote the vertex set, edge set, and block set of G, respectively. A graph G is said to have an *AUM block labeling* if there exists a bijection $f : V(G) \to \{1, 2, \ldots, p\}$ and an induced function $f^* : E(G) \to \mathbb{Z}^+$ defined by $f^*(uv) = f(u) + f(v)$, together with [3930] new

a function $f^{**} : B(G) \to \mathbb{Z}^+$ defined as follows. Let B_j be incident with the vertices $v_{j_1}, v_{j_2}, \ldots, v_{j_k}$, where $1 \leq j_\ell \leq p$, and the edges $e_{j_1}, e_{j_2}, \ldots, e_{j_m}$, where $1 \leq j_r \leq q$. Then $f^{**}(B_j) = \sum_{i=1}^{k} f(v_{j_i}) + \sum_{i=1}^{m} f^*(e_{j_i})$, and $f^{**}(B_j) \neq f^{**}(B_i)$ whenever $1 \leq i, j \leq b$ and $i \neq j$. They prove the following graphs admit AUM block sum labelings: paths, triangular snakes, alternate triangular snakes, double triangular snakes, double alternate triangular snakes, quadrilateral snakes, and alternate quadrilateral snakes. Uma Maheswari and Azhagasari [3931] gave AUM block labelings for cycle cacti and friendship graphs. Uma Maheswari and Purnalakshimi [3933] showed that paths, tadpoles, friendship graphs, and cycle cactus graphs admit AUM block labelings. In [3934] they gave AUM block labelings for kites, combs, and diamond snake graphs. In [3932] Uma Maheswari and Azhagarasi gave AUM block labelings for stars, bistars, combs, sunlet graphs, diamond snakes, pentagonal snakes, and alternate pentagonal snakes. [3931] new [3933] new [3934] new [3932] new

In [3935] Uma Maheswari and Sumathi introduced a labeling called AUM *block geometric mean* labeling in which labels are assigned to the blocks of a graph based on the geometric mean of its vertices and edges. They obtained AUM block geometric mean labelings for triangular snakes, alternate triangular snakes, double triangular snakes, quadrilateral snakes, alternate quadrilateral snakes, pentagonal snakes, and alternate pentagonal snakes. Uma Maheswari and Sumathi [3937] gave AUM block geometric mean labelings for olive trees, coconut trees, Y-trees, and provided a characterization of AUM block geometric mean labelings. In [3936] they provided AUM block geometric mean labeling for tadpoles, kayak paddles, and bull graphs. [3935] new [3937] new [3936] new

Random walk labeling

Fang, Pang, Tardos, and Zuo [1064] and Fried and Mansour [1119], independently and almost at the same time, defined a new kind of graph labeling, called a successive vertex ordering in [1064] and random walk labeling in [1119]. A *random walk labeling* of a graph G with vertex set V is a labeling that is obtains by performing the following process: Set $i = 1$ and let $v \in V$. Label v with i. Then, as long as there are vertices that are not labeled, pick $w \in V$ that is adjacent to v and replace v with w. If v is not labeled, increase i by 1 and label v with i. Clearly, in a random walk labeling, for each i, the vertices labeled by $1, 2, \ldots, i$ constitute a subgraph of G. This is how a successive vertex ordering is defined. Fang et al. [1064] concentrated on fully regular graphs, while Fried and Mansour [1119], [1120], [1121] studied random walk labelings of specific graph families such as grid graphs, torus graphs and king graphs, all of size $n \times 2$, perfect trees, barbells, lollipops, and tadpole graphs.

,

References

[1] M. Aasi, M. Asif, T. Iqbal, and M. Ibrahim, Radio labelings of lexicographic product of some graphs, *J. Math.,* (2021) Art. ID 9177818, 6 pp. https://www.hindawi.com/journals/jmath/2021/9177818/

[2] M. E. Abdel-Aal, Odd harmonious labelings of cyclic snakes, *Internat. J. Appl. Graph Theory Wireless Adhoc Networks and Sensor Networks*, **5** (3) (2013) 1-13. doi:10.5121/jgraphoc.2013.5301

[3] M. E. Abdel-Aal, New classes of odd graceful graphs, *Internat. J. Appl. Graph Theory in Wireless ad hoc Networks and Sensor Networks,* **5**(2) (2013) 1-12. doi:10.5121/jgraphoc.2013.5201

[4] M. E. Abdel-Aal, New families of odd harmonious graphs, *Internat. J. Soft Comput., Math. Control*, **3**(1) (2014) 1-13.

[5] M. E. Abdel-Aal and S. A. Bashammakh, A study on the varieties of equivalent cordial labeling graphs *AIMS Math.,* **9**(12) (2024) 34720–34733. doi:DOI:10.3934/math.20241653

[6] M. E. Abdel-Aal, N. M. NourEldeen, E. Badr, and H. M. Shabana, Symmetric cryptography approach based on odd graceful labeling, *Sci. African*, (2026) 24 pp. doi:10.1016/j.sciaf.2026.e03592

[7] M. E. Abdel-Aal1 and M. A. Seoud, Further results on odd harmonious graphs, *Internat. J. Appl. Graph Theory in Wireless Adhoc Networks and Sensor Networks*, **8**(3/4) (2016) 1-14. doi:10.5121/jgraphoc.2016.8401

[8] R. Abdul Saleem, N. Vanitha Jothy, and M. Haajira, Root square mean labeling of uncoordinated graphs, *Fund. Appl. Math. aSoft Computing*, (2022) 152–160. https://www.researchgate.net/publication/410790937_Root_Square_Mean_Labeling_of_Uncoordinated_Graphs

[9] H. Abdo, S. Brandt, D. Dimitrov, The total irregularity of a graph, *Disc. Math. Theor. Comput. Sci.,* **16**(1) (2014) 201-206. https://www.researchgate.net/publication/229438159_The_total_irregularity_of_a_graph

[10] R. J. R. Abel and M. Buratti, Some progress on $(v, 4, 1)$ difference families and optical orthogonal codes, *J. Combin. Theory Ser. A,* **106** (2004) 59-75. https://www.researchgate.net/publication/220078766_Some_progress_on_v_4_1_difference_families_and_optical_orthogonal_codes

[11] K. Abhishek, A note on set-indexed graphs, *J. Disc. Math. Sci. Crypt.* **18** (2015) 31-40.

[12] K. Abhishek, Set-valued graphs: a survey, *J. Disc. Math. Sci. Crypt.* **18** (2015) 55-80.

[13] K. Abhishek, Set-valued graphs II, *J. Fuzzy Set Valued Analysis*, **2013** (2013) 1-16.

[14] K. Abhishek and G. K. Agustine, Set-valued graphs, *J. Fuzzy Set Valued Analysis*, **2012** (2012) 17 pages. https://d-nb.info/1027620361/34

[15] V. J. Abhyankar, Direct Methods of Gracefully Labeling Graphs, Ph. D. Thesis, University of Mumbai, 2002.

[16] V. J. Abhyankar and V. N. Bhat-Nayak, Easiest graceful labeling of olive trees, *Bull., Bombay Math. Coll.*, **14** (2000) 16-25. NOT ON WEB 12/22/24

[17] V. J. Abhyankar and V. N. Bhat-Nayak, personal communication (2001).

[18] C. Abiramasundari and P. Senthil Vadivu, Liouville difference cordial labeling of some acyclic graphs, *Internat. J. Stat. Appl. Math.*, **11**(1) Part A (2026) 35–40. https://www.mathsjournal.com/pdf/2026/vol11issue1/PartA/11-1-4-309.pdf

[19] C. Abiramasundari and P. Senthil Vadivu, On Liouville difference cordial labeling of some splitting graphs, *Int. J. Stat. Appl. Math.*, **11**(7) (2026) 36-39. https://www.mathsjournal.com/pdf/2026/vol11issue7/PartA/11-6-271-568.pdf

[20] C. Abiramasundari, R. Bhuvaneswari, M. Ilayaraja, and R. Imran, A study of Liouville difference cordial labeling in cyclic graph classes, *JAAFR - J. ADV, Future Res.* **4**(5) (2026) e955-e958. https://rjwave.org/jaafr/papers/JAAFR2605964.pdf

[21] P. Abirami, A. Parthiban, and N. Srinivasan, On 2-odd labeling of graphs, *European J. Molecular & Clinical Medicine,* **07**(07) (2020) 3914-3918. https://api.semanticscholar.org/CorpusID:235238605

[22] P. Abirami, A. Parthiban, and N. Srinivasan, Some results on 2-odd labeling of graphs, *Internat. Recent Tech. Eng.*, (IJRTE), **8**(5) (2020) 5644-5646. doi:DOI:10.35940/ijrte.E7041.018520

[23] T. Abraham and S. Jose, Fibonacci product cordial graphs, *J. Emerging Tech. Innovative Res.,* **6**(1) (2019) 58-63. https://www.jetir.org/papers/JETIR1901808.pdf

[24] J. Abrham, Perfect systems of difference sets-A survey, *Ars Combin.,* **17A** (1984) 5-36.

[25] J. Abrham, Graceful 2-regular graphs and Skolem sequences, *Discrete Math.*, **93** (1991) 115-121.

[26] J. Abrham, M. Carter, and K. Eshghi, personal communication (circa 1995).

[27] J. Abrham and A. Kotzig, Extensions of graceful valuations of 2-regular graphs consisting of 4-gons, *Ars Combin.,* **32** (1991) 257-262.

[28] J. Abrham and A. Kotzig, Two sequences of 2-regular graceful graphs consisting of 4-gons. Fourth Czechoslovakian Symposium on Combinatorics, Graphs and Complexity (Prachatice, 1990), 1–4, *Ann. Discrete Math.*, **51**, North-Holland, Amsterdam, 1992.

[29] J. Abrham and A. Kotzig, All 2-regular graphs consisting of 4-cycles are graceful, *Discrete Math.*, **135** (1994) 1-14. https://doi.org/10.1016/0012-365X(93)E0095-L

[30] J. Abrham and A. Kotzig, Graceful valuations of 2-regular graphs with two components, *Discrete Math.*, **150** (1996) 3-15. https://doi.org/10.1016/0012-365X(95)00171-R

[31] A. Abueida and K. Roblee, Harmonious labelings of disconnected graphs involving cycles and multiple components consisting of starlike trees, In: F. Hoffman, (eds) Combinatorics, Graph Theory and Computing. SEICCGTC 2020. Springer Proceedings in Mathematics & Statistics, **388**. Springer, Cham. 131-140. https://link.springer.com/chapter/10.1007/978-3-031-05375-7_10

[32] A. Abueida and K. Roblee, More harmonious labelings of families of disjoint unions of an odd cycle and certain trees, *J. Combin. Math. Combin. Comput.*, **115** (2020) 61-68. https://combinatorialpress.com/jcmcc-articles/volume-119/harmonious-labelings-of-families-of-disjoint-unions-of-an-odd-\cycle-and-certain-trees/

[33] B. Abughazaleh and O. Abughneim, Prime labeling of graphs constructed from wheel graph, *Heliyon* **10**(2) (2024) e23979 1-9. https://www.sciencedirect.com/science/article/pii/S2405844024000100

[34] E. A. AbuHijleh, O. A. AbuGhneim, and H. Al-Ezeh, Characterizing when powers of a caterpillar are divisor graphs, *Ars Combin.,* **113A** (2014) 85-89. https://www.researchgate.net/publication/270956712_Characterizing_When_Powers_of_a_Caterpillar_are_Divisor_Graphs

[35] E. A. AbuHijleh, O. A. AbuGhneim, and H. Al-Ezeh, Characterizing which powers of hypercubes and folded hypercubes are divisor graphs, *Discuss. Math. Graph Theory,* **35**(2) (2015) 301-311. doi:10.7151/dmgt.1801

[36] E. A. AbuHijleh, O. A. AbuGhneim and H. Al-Ezeh, Characterizing when powers of a caterpillar are divisor graphs, *Ars Combin.*, **113** (2014) 85-95. https://combinatorialpress.com/article/ars/Volume%20113-A/volume-113-A-paper-7.pdf

[37] R. Aceska, N. Arcila-Maya, J. Carlson, A. Marr, M. Parnes, K. Ryan, H. Schuerger, and J. F. Vasquez, New results on difference distance magic labelings, https://doi.org/10.48550/arXiv.2601.20492

[38] R. Aeska, L. Mckailyn, and A. Ripperger, Ornaments and difference distance magic oriented graphs, https://doi.org/10.48550/arXiv.2606.16001

[39] B. D. Acharya, Construction of certain infinite families of graceful graphs from a given graceful graph, *Def. Sci. J.,* **32** (1982) 231-236. NOT ON WEB 12/22/24

[40] B. D. Acharya, On d-sequential graphs, *J. Math. Phy. Sci.*, **17** (1983) 21-35. NOT ON WEB 12/22/24

[41] B. D. Acharya, Set valuations of a graph and their applications, MRI Lecture Notes in Applied Mathematics, No. 2, Mehta Research Institute, Allahabad, 1983.

[42] B. D. Acharya, Are all polyominoes arbitrarily graceful?, *Proc. First Southeast Asian Graph Theory Colloquium*, Ed. K. M. Koh and H. P. Yap, Springer-Verlag, N. Y. 1984, 205-211.

[43] B. D Acharya and S. Arumugarn, Labeling of discrete structures *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 1-14.

[44] B. D. Acharya, S. Arumugarn, and A. Rosa, Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008.

[45] B. D. Acharya and K. A. Germina, Strongly k-indexable unicyclic graphs, *Graph Theory Notes N. Y.,* **55** (2008) 45-49.

[46] B. D. Acharya and K. A. Germina, Set-valuations of graphs and their applications: A survey, *Annals Pure Appl. Math.*, **4**(1) (2013) 8-42. https://pdfs.semanticscholar.org/e438/81bc8a4f04508693916f527945ac0b7b1b6a.pdf

[47] B. D. Acharya and K. A. Germina, Maximal strongly k-indexable graphs, *J. Combin. Math. Combin. Comput.,* **76** (2011) 75-99. https://combinatorialpress.com/article/jcmcc/Volume%20076/vol-076-paper%206.pdf

[48] B. D. Acharya, K. A. Germina, K. Abhishek, and P. J. Slater, Some new results on set-graceful and set-sequential graphs, *Combin. System Sci.*, **37**(2-4) (2012) 229-239.

[49] B. D. Acharya, K. A. Germina, and V. Ajitha, Multiplicatively indexable graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 29-40.

[50] B. D. Acharya, K. A. Germina, and T. M. K. Anandavally, Some new perspectives on arithmetic graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 41-46.

[51] B. D. Acharya, K. A. Germina, K. L. Princy, and S. B. Rao, On set-valuations of graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 47-56.

[52] B. D. Acharya, K. A. Germina, K. L. Princy, and S. B. Rao, Graph labellings, embedding and *NP*-completeness theorems, *J. Combin. Math. Combin. Comput.,* **67** (2008) 163-180.

[53] B. D. Acharya and M. K. Gill, On the index of gracefulness of a graph and the gracefulness of two-dimensional square lattice graphs, *Indian J. Math.,* **W23** (1981) 81-94.

[54] B. D. Acharya and S. M. Hegde, Further results on k-sequential graphs, *Nat. Acad. Sci. Lett.,* **8** (1985) 119-122.

[55] B. D. Acharya and S. M. Hegde, Set sequential graphs, *Nat. Acad. Sci. Lett.,* **8** (1985) 387-390.

[56] B. D. Acharya and S. M. Hegde, Arithmetic graphs, *J. Graph Theory*, **14** (1990) 275-299.

[57] B. D. Acharya and S. M. Hegde, Strongly indexable graphs, *Discrete Math.*, **93** (1991) 123-129.

[58] B. D. Acharya and S. M. Hegde, On certain vertex valuations of a graph I, *Indian J. Pure Appl. Math.,* **22** (1991) 553-560.

[59] B. D. Acharya, S. B. Rao, and S. Arumugan, Embeddings and NP-complete problems for graceful graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 57-62.

[60] B. D. Acharya, S. B. Rao, T. Singh, and V. Parameswaran, Some sigma labelled graphs, Presented at 19th Annual Conference of Ramanujan Mathematical Society, held at Dr. B. R. Ambedkar University, Agra, July 21-24, 2004.

[61] B. D. Acharya, S. B. Rao, T. Singh, and V. Parameswaran, Neighborhood magic graphs, 2004, unpublished.

[62] M. Acharya and V. Bhat-Nayak, Minimal 4-equitability of $C_{2n} \odot K_1$, *Ars Combin.,* **65** (2002) 209-236.

[63] M. Acharya and V. Bhat-Nayak, k-equitability of $C_{2n} \odot K_1$, $k = 2, 2n$ and associated graphs. *Util. Math.,* **68** (2005) 109-130.

[64] M. Acharya, R. Jain, and S. Kansal, Vertex equitable labeling of signed graphs, *Elect. Notes Discr. Math.,* **63** (2017) 461-468.

[65] M. Acharya and T. Singh, Graceful signed graphs, *Czechoslovak Math. J.,* **54**(129) (2004) 291-302.

[66] D. G. Adalja, Sum divisor cordial labeling for vertex switching of special graphs, *Internat. J. Applied Graph Th.*, **5**(2) (2021) 34–43. https://www.ijagt.com/upload/Sum_Divisor.pdf

[67] D. G. Adalja, Sum divisor cordial labeling in the context of graphs operations on bistar, *TWMS J. Appl. Engin. Math.*,**12**(2) (2022) 588–599. https://jaem.isikun.edu.tr/web/images/articles/vol.12.no.2/19.pdf

[68] D. G. Adalja and G. V. Ghodasara, Sum divisor cordial labeling of snakes related graphs, *J. Comput. Math. Sci.*, **9**(7) (2018) 754-768. https://www.researchgate.net/publication/376514102_Sum_Divisor_Cordial_Labeling_on_Some_Cycle_and_Snake_Related_Graphs

[69] D. G. Adalja and G. V. Ghodasara, Sum divisor cordial labeling in the context of duplication of graph elements, *Mathematics Today,* **35** (2019) 61-83.

[70] D. G. Adalja and G. V. Ghodasara, Some new sum divisor cordial graphs, *Internat. J. Appl. Graph Th.*, textbf2 (2018) 19–33. https://www.ijagt.com/upload/Some_New_Sum_Divisor_Cordial_Graphs.pdf

[71] D. G. Adalja and P. Patel, Sum divisor cordial labeling for path union of two copies of cycle related graphs, *Palestine J. Math.*, **12**(4) (2023) 125–132. https://pjm.ppu.edu/sites/default/files/papers/PJM_12%284%29_2023_125_to_132.pdf

[72] M. Adamaszek, Efficient enumeration of graceful permutations, *J. Combin. Math. Combin. Comput.,* **87** (2013) 191-197. https://arxiv.org/pdf/math/0608513.pdf

[73] R. Adawiyah, I. I. Makhfudloh, and R. M. Prihandini, Local edge (a, d)-antimagic coloring on sunflower, umbrella graph and its application, *Alifmatika J. Pendidikan dan Pembelajaran Matematika,* https://www.researchgate.net/publication/371884100_Local_edge_a_d_-antimagic_coloring_on_sunflower_umbrella_graph_and_its_application

[74] T. C. Adefokun and D. O. Ajayim On the bounds of the radio numbers of stacked-book graph, *Internat. J. Math. Combin.,* **4** (2022) 66-73. https://www.researchgate.net/publication/370937821_Vol42022_of_International_Journal_of_Mathematical_Combinatorics/

[75] C. Adiga, H. Ramashekar, and D. Somashekara, A note on strongly multiplicative graphs, *Discuss. Math.,* **24** (2004) 81-83.

[76] C. Adiga and D. Somashekara, Strongly $\star$-graphs, *Math. Forum,* **13** (1999/00) 31-36.

[77] C. Adiga, C. K. Subbaraya, A. S. Shrikanth and M. A. Sriraj, On vertex balance index set of some graphs, *Bull. Iran. Math. Soc., 39* (4) (2013) 627-634.

[78] J. Afeltra, M. Dias, H.-H. Su, and H. Todt, On the edge-balance index sets of broken wheels, *Congr. Numer.,* **223** (2015) 205-218.

[79] M. Aftiana and D. Indriati, On edge irregular total k-labeling and total edge irregularity etrength of barbell graphs, *J. Physics Conference Ser.,* (2019) 1306:012031 1-7.

[80] H. U. Afzal On super edge-magicness of two special families of graphs, *Util. Math.* **97** (2015) 97-108.

[81] H. U. Afzal, M. Ather, A. Q. Baig, and A. Maheshwari, Further results on super edge-magic graphs, *Util. Math.,* **110** (2019) 283-291.

[82] H. U. Afzal and M. Aslam, On the deficiency of disjoint union of further planar classes of graphs, *Util. Math.,* **105** (2017) 207-218.

[83] H. U. Afzal, M. Javaid, A. M. Alanazi, and M. G. Alshehri, Computing edge weights of symmetric classes of networks, Mathematical Problems in Engineering, **2021** 2021 Article ID 5562544, 22 pages. https://doi.org/10.1155/2021/5562544

[84] A. Agarwal and M. Lopez, Representation number for complete graphs minus stars, *J. Combin. Math. Combin. Comput.,* **87** (2013) 33-41.

[85] P. Agasthi and N. Narvathi, On some labeling of belttriangular snake and central graph of triangular snake graph, In: *Nat. Conf. Math. Tech. Appl.,* (NCMTA) (2018) 1-19. https://iopscience.iop.org/article/10.1088/1742-6596/1000/1/012170

[86] A. Aguado and S. El-Zanati, On σ-labeling the union of three cycles, *J. Combin. Math. Combin. Comput.,* **64** (2008) 33-48.

[87] A. Aguado, S. El-Zanati, H. Hake, J. Stob, and H. Yayla, On ρ-labeling the union of three cycles, *Australas. J. Combin.,* **37** (2007) 155-170.

[88] I. H. Agustin, Dafik, Marsidi, E. R. Albirri, On the total H-irregularity strength of graphs: A new notion, *J. Physics: Conf. Series,* **855** (2017) 012004 doi:10.1088/1742-6596/855/1/012004

[89] I. H. Agustin, D. Dafik, N. Mohanapriya, M. Marsidi, and I. N. Cangul, The distance irregular reflexive k-labeling of graphs, *CAUCHY –J. Mat. Murni dan Aplikasi*, **7**(4) (2023) 622-630. https://www.researchgate.net/publication/371005283_The_Distance_Irregular_Reflexive_k-Labeling_of_Graphs

[90] I. H. Agustin, M. Hasan, Dafik, R. Alfarisi and R. M. Prihandini, Local edge antimagic colorings of graphs, *Far East J. Math. Sci.,* (FJMS) **102**(9) (2017) 1925-1941. doi:10.17654/MS102091925

[91] I. H. Agustin, F. Susanto, Dafik, R. M. Prihandini, R. Alfarisi, and I. Sudarsana, On H-supermagic labelings of m-shadow of paths and cycles, *Int. J. Math. Math. Sci.,* (2019) Art. ID 8780329, 7 pp. https://doi.org/10.1155/2019/8780329

[92] I. H. Agustin, M. Iman Utoyo, Dafik, and M. Venkatachalam, The reflexive edge strength on some almost regular graphs, *Heliyon* **7** (2021) https://doi.org/10.1016/j.heliyon.2021.e06991

[93] I. H. Agustin, L. Susilowati, Dafik, I. N. Cangul, and N. Mohanapriya, On the vertex irregular reflexive labeling of several regular and regular-like graphs, *J. Discrete Math. Sci. Crytography* **25**(5) (2022), 1457-1473. https://scholar.unair.ac.id/en/publications/on-the-vertex-irregular-reflexive-labeling-of-several-regular-and

[94] T. Augustine and S. Roy, Radio labeling of certain networks, *Palestine J. Math.*, **12**(1) (2023) 75–80. https://www.researchgate.net/publication/373895536_Radio_Labeling_of_Certain_Networks

[95] A. Ahadi, A. Dehghan, M. Kazemi, and E. Mollaahmadi, Computation of lucky number of planar graphs is NP-hard. *Inform. Process. Let.,* **112**(4) (2012) 109-112.

[96] A. Ahmad, On vertex irregular total labeling of convex polytope graphs, *Util. Math.,* **89** (2012) 69-78.

[97] S. Ahmad, On the evaluation of a subdivision of the ladder graph, *Punjab Univ. J. Math. (Lahore),* **47**(1) (2015) 15-19. http://pu.edu.pk/images/journal/maths/PDF/1_4-02-09.pdf

[98] A. Ahmad, Computing the edge irregularity strength of certain unicyclic graphs, unpublished.

[99] A. Ahmad, Computing 3-total edge product cordial labeling of generalized Petersen graphs $P(n, m)$, *Ars Combin.,* **137** (2018) 263-271.

[100] A. Ahmad, Computing an edge irregular k-labeling of star related trees, *Ars Combin.* **155** (2021) 169-179.

[101] A. Ahmad, Computing the total vertex irregularity strength associated with zero divisor graph of commutative ring, *Kragujevac J. Math.,* **46**(5) (2022) 711-719. https://imi.pmf.kg.ac.rs/kjm/pub/kjom465/kjm_46_5-4.pdf

[102] A. Ahmad, S. Ahtsham, H. B. M. Imran, and A. Q. Baig, Vertex irregular labelings of cubic graphs, *Util. Math.,* **91** (2013) 287-299.

[103] Y. Ahmad, U. Ali, M. Bilal, S. Zafar, Z. Zahid, Some new standard graphs labeled by 3-total edge product cordial labeling, *Appl. Math. Nonlinear Sci.*,**2**(1) (2017) 61-72.

[104] A. Ahmad, K. Ali, M. Bača, P. Kovář, and A. Semaničová-Feňovčíková, Vertex-antimagicness of regular graphs, *Acta Math. Sinica,* English series, **28**(9) (2012) 1865-1874. https://www.researchgate.net/publication/257481499_Vertex-antimagic_labelings_of_regular_graphs

[105] A. Ahmad, O. Al-Mushayt, M. Bača, On edge irregular strength of graphs, *Appl. Math. Comput.,* **243** (2014) 607-610.

[106] A. Ahmad, K. Ali, and E. Baskoro, On super edge-magic total labelings of a forest of banana trees, *Util. Math.,* **83** (2010) 323-332

[107] A. Ahmad, M. Arshadb, and G. Ižaríková, Irregular labelings of helm and sun graphs, *AKCE Internat. J. Graphs Combin.,* **12** (2015) 161-168. https://doi.org/10.1016/j.akcej.2015.11.010

[108] A. Ahmad, K. M. Awan, I. Javaid, and Slamin, Total vertex irregularity strength of wheel related graphs, *Australas. J. Combin.,* **51** (2011) 147-156. https://ajc.maths.uq.edu.au/pdf/51/ajc_v51_p147.pdf

[109] A. Ahmad and M. Bača, On edge irregular total labeling of certain family of graphs, *AKCE J. Graphs Combin.,* **6** (2009) 21-29. https://www.researchgate.net/publication/261951606_Edge_irregular_total_labeling_of_certain_family_of_graphs#fullTextFileContent

[110] A. Ahmad and M. Bača, On vertex irregular total labeling, *Ars Combin.*, **112** (2013) 129-139.

[111] A. Ahmad and M. Bača, Total edge irregularity strength of the categorical product of two paths, *Ars Combin.*, **114** (2014) 203-212. https://www.researchgate.net/publication/262685108_Total_edge_irregularity_strength_of_a_categorical_product_of_two_paths

[112] A. Ahmad, M. Bača, and Y. Bashir, Total vertex irregularity strength of certain classes of unicyclic graphs, *Bull. Math. Soc. Sci. Math. Roumanie*, **57**(2) (2014) 147-152. https://www.researchgate.net/publication/262933043_Total_vertex_irregularity_strength_of_certain_classes_of_unicyclic_graphs

[113] A. Ahmad, M. Bača, Y. Bashir, and M. K. Siddiqui, Total edge irregularity strength of strong product of two paths, *Ars Combin.,* **106** (2012) 449-459.

[114] A. Ahmad, M. Bača, M. Lascsáková, and A. Semaničová-Feňovčíková, Super magic and antimagic labelings of disjoint union of plane graphs, *Science Internat.*, **24**(1) (2012) 21-25. https://www.researchgate.net/publication/262692776_Super_magic_and_antimagic_labelings_of_disjoint_union_of_plane_graphs

[115] A. Ahmad, M. Bača, and M. F. Nadeen, On edge irregularity strength of Toeplitz graphs, *U.P.B. Sci. Bull., Series A,* **78**(4) (2016) 155-162. https://www.scientificbulletin.upb.ro/rev_docs_arhiva/full15e_507690.pdf

[116] A. Ahmada, M. Bača, M. Naseem, and A. Semaničová-Feňovčíková, On 3-total edge product cordial labeling of honeycomb, *AKCE Internat. J. Graphs Combin.,* **14** (2017) 149-157. https://doi.org/10.1016/j.akcej.2017.02.002

[117] A. Ahmad, M. Bača and M. Numan, On irregularity strength of disjoint union of friendship graphs, *Elect. J. Graph Th. App.,* **1**(2) (2013) 100-108. https://www.ejgta.org/index.php/ejgta/article/viewFile/2/9

[118] A. Ahmad, M. Bača, and A. Semaničová-Feňovčíková, Computation of edge C_4-irregularity strength of Cartesian product of graphs, *J. Discrete Math. Sci. and Crypt.*, (2022)

[119] A. Ahmad, M. Bača, and M. K. Siddiqui, On edge irregular total labeling of categorical product of two cycles, *Th. Comp. Systems*, **54** (2014) 1-12. https://link.springer.com/article/10.1007/s00224-013-9470-3

[120] A. Ahmad, M. Bača, and M. K. Siddiqui, Irregular total labeling of disjoint union of prisms and cycles, *Australas. J. Combin.*, **59**(1) (2014) 98-106. https://www.researchgate.net/publication/261913345_Irregular_total_labeling_of_disjoint_union_of_prisms_and_cycles

[121] A. Ahmad, A. Q. Baig, and M. Imran, On super edge-magicness of graphs, *Util. Math.,* **89** (2012) 373-380. https://www.researchgate.net/publication/267145329_On_Super_Edge-magicness_of_Graphs

[122] M. A. Ahmed and J. B. Babujee, On face bimagic labeling of graphs, *IOSR J. Math.,* **12**(6) VI (2016) 1-7. http://www.iosrjournals.org/iosr-jm/papers/Vol12-issue6/Version-6/A1206060107.pdf

[123] A. Ahmad, E. T. Baskoro, and M. Imran, Total vertex irregularity strength of disjoint union of Helm graphs, *Discuss. Math. Graph Theory,* **32**(3) (2012) 427-434. https://www.researchgate.net/publication/257922360_Total_vertex_irregularity_strength_of_disjoint_union_of_Helm_graphs

[124] A. Ahmad, A. Gupta, and R. Simanjuntak, Computing the edge irregularity strengths of chain graphs and the join of two graphs, *Electron. J. Graph Theory Appl.,* (EJGTA) **6**(1) (2018) 201-207.

[125] A. Ahmad, R. Hasni, M. Irfan, M. Naseem, and M. K. Siddiqui, On 3-total edge product cordial labeling of grid, *Asian-Eur. J. Math.,* **4**(6) (2021) Paper No. 2150096, 10 pp. https://doi.org/10.1016/j.akcej.2017.02.002

[126] A. Ahmad, M. Ibrahim, and M. K. Siddiqui, On the total irregularity strength of generalized Petersen graph, *Math. Rep. (Bucur.),* **18**(68) no. 2 (2016) 197-204. http://imar.ro/journals/Mathematical_Reports/Pdfs/2016/2/5.pdf

[127] A. Ahmad, M. Imran, and A. Semaničová-Feňovčíková, Relation between mean labeling and $(a;d)$-edge antimagic vertex labeling, *Sci. Internat.,* **24**(1)(2012) 7-9.

[128] A. Ahmad, I. Javaid, and M. F. Nadeem, Further results on super edge magic deficiency of unicyclic graphs, *Ars Combin.,* **99** (2011) 129-138.

[129] A. Ahmad, I. Javaid, M. F. Nadeem, and R. Hasni, On super edge magic deficiency of some families related to ladder graphs, *Australas. J. Combin.,* **51** (2011) 201-208. http://ajc.maths.uq.edu.au/pdf/51/ajc_v51_p201.pdf

[130] A. Ahmad, S. S. Khan, S. Ahmad, M. F. Nadeem, and M. K. Siddiqui, Edge irregularity strength of categorical product of two paths, *Ital. J. Pure Appl. Math.*, **45** (2021) 801-813. https://ijpam.uniud.it/online_issue/202145/63%20Siddiqui_et_all_(edge).pdf

[131] A. Ahmad, S.C. López, F.A. Muntaner-Batle, and M. Rius-Font, Enumerating super edge-magic labeling for the union of non-isomorphic graphs, *Bull. Aust. Math. Soc.,* **84** (2011)(2) 310-321.

[132] A. Ahmad and R. Marinescu-Ghemeci, Radio labeling of some ladder-related graphs, *Math. Rep. (Bucur.)* **19**(69) (2017), no. 1, 107-119. http://imar.ro/journals/Mathematical_Reports/Pdfs/2017/1/11.pdf

[133] A. Ahmad, M. Nadeem, and A. Gupta, On super edge-magic deficiency of certain Toeplitz graphs, *Hacet. J. Math. Stat.,* **47**(3) (2018) 513-519. http://www.hjms.hacettepe.edu.tr/uploads/45dfc139-090e-4fde-b21f-6a880aef6761.pdf

[134] A. Ahmad, Nurdin, and E. T. Baskoro, On total irregularity strength of generalized Halin graph, *Ars Comb.,* **122** (2015) 319-332. https://www.researchgate.net/publication/296415305_On_total_irregularity_strength_of_generalized_Halin_graph

[135] A. Ahmed, Al-M. Omar, and M. Bač, On edge irregularity strength of graphs, On edge irregularity strength of graphs, *Appl. Math. Comput.* **1** (2014) 16-28.

[136] A. Ahmad, A. Semaničová-Feňovčíková, and M. K. Siddiqui, Construction of α-tree from smaller graceful trees, *Util. Math.,* **99** (2016) 175-186.

[137] A. Ahmad, M. K. Siddiqui, and D. Afzal, On the total edge irregularity strength of zigzag graphs, *Australas. J. Combin.,* **54** (2012) 141-149.

[138] A. Ahmad, M. K. Siddiqui, M. F. Nadeem, and M. Imran, On super edge magic deficiency of kite graphs, *Ars Combin.,* **107** (2012) 201-208. https://www.researchgate.net/publication/265504052_On_super_edge_magic_deficiency_of_kite_graphs

[139] A. Ahmad and I. Tomescu, On vertex-magic total labelings of some families of rotationally-symmetric graphs, *Util. Math.*, **86** (2011) 347-357. https://www.researchgate.net/publication/268442776_On_vertex-magic_total_labeling_of_some_families_of_rotationally-symmetric_graphs

[140] M. A. Ahmed, J. Babujee, M. Bača, and A. Semaničová-Feňovčíková, On totally antimagic, edge-magic and vertex-antimagic total graphs, *Util. Math.,* **111** (2019) 161-173. http://www.rjm-cs.ro/Ahmed_Babujee_2017_3.pdf

[141] M. A. Ahmed and J. Baskar Babujee, Totally antimagic total labeling of complete bipartite graphs, *Rom. J. Math. Comput. Sci.,* **7**(1) (2017) 21-28.

[142] M. A. Ahmed, and J. Baskar Babujee, Encryption through labeled graphs using strong face bimagic labeling, *Internat. Math. Forum*, (2017) 151-158.

[143] M. A. Ahmed and J. B. Babujee, On face antimagic labeling of strong face plane graphs, **Appl. Math. Sci.,** (2017) **11** 77-91. doi:10.12988/AMS.2017.610262

[144] M. A. Ahmed, A. Semaničová-Feňovčíková, M. Bača, J. Baskar Babujee, and L. Shobana, On graceful antimagic graphs, *Aequat. Math.,* Springer Nature Switzerland AG 2022. https://doi.org/10.1007/s00010-022-00930-1

[145] T. Ahmed and H. Snevily, The α-labeling number of comets is 2, *Bull. Inst. Combin. Appl.,* 72 (2014) 25-40.

[146] M. Aigner and E. Triesch, Irregular assignments of trees and forests, *SIAM J. Discrete Math.*, **3**(4) (1990) 439-449.

[147] A. Aishwarya, A study on lucky-edge labeling, *Internat. J. Innov. Res. Sci. Engin. Tech.*, **4**(3) (2015) 1266-1269. https://www.ijirset.com/upload/2015/march/40_aishwarya_paper_3(5_PAGES)_3.pdf

[148] A. Aishwarya, Lucky-edge labeling of different graphs, *Internat. J. Innov. Res. Sci. Engin. Tech.*, **4**(2) (2015) 469-471. https://periodicos.uem.br/ojs/index.php/BSocParanMat/article/view/76930/751375160670

[149] D. Ajayi, Totally antimagic total labeling of ladders, prisms and generalised Petersen graphs, https://doi.org/10.48550/arXiv.1609.04299

[150] D. O. Ajayi and T. C. Adefokun, On bounds of radio number of certain product graphs, *J. Nigerian Math. Soc.*, **48**(2) (2018) 71-78. https://ojs.ictp.it/jnms/index.php/jnms/article/view/330/57

[151] D. O. A. Ajayi and A. D. Akwu, Totally antimagic total labelings of ladders, prisms and generalized Petersen graphs, https://arxiv.org/pdf/1609.04299

[152] V. Ajitha, K. L. Princy, V. Lokesha, and P. S. Ranjini, On square difference graphs, *Math. Combin. Internat. Book Ser.*, **1** (2012) 31-40.

[153] A. Aishwarya, Lucky edge labeling of certain graphs, *IJIRSET,* **4**(1) (2015) 18708-18711..
https://docslib.org/doc/4217723/lucky-edge-labeling-of-certain-graphs

[154] V. Ajtha, S. Arumugam, and K. A. Germina, On square sum graphs, *AKCE J. Graphs Combin.,* **6** (2009) 1-10. https://www.researchgate.net/publication/266713407_On_square_sum_graphs

[155] S. Akbari, N. Ghareghani, G. Khosrovshahi, and S. Zare, A note on zero-sum 5-flows in regular graphs, *Electron. J. Combin.,* **19**(2) (2012), Paper 7, 5 pp.

[156] S. Akbari, M. Kano, S. Zare, 0-Sum and 1-sum flows in regular graphs, *Electronic J. Combin.,* **23**(2) (2016) #P2.37 10pp. https://www.combinatorics.org/ojs/index.php/eljc/article/view/v23i2p37/pdf

[157] S. Akbari, F. Rahmati, and S. Zare, Zero-sum magic labelings and null sets of regular graphs. *Electronic J. Combin.,* **21**(2) (2014) Paper 2.17, 9 pp.

[158] P. Z. Akbari1, V. J. Kaneria, and N. A. Parmar, Absolute mean graceful labeling of jewel graph and jelly fish graph, *Internat. J. Math. Trends Tech.,* **68**(1) (2022) 86-93. 10.14445/22315373/IJMTT-V68I1P510

[159] S. Akila Devi, S. Vembu, G. Jasmine, and S. C. Deva Suba, Graph-theoretic verification of lucky and proper lucky labeling invariants, *Interna. J. Math. Trends Tech.* (IJMTT), **72**(7) (2026) 10-13. https://ijmttjournal.org/public/assets/volume-72/issue-7/IJMTT-V72I7P102.pdf

[160] K. Akilandeswari, k-Super root square mean labeling of some graphs, *Internat. J. Sci. Res.,* (IJSR) **6**(2) (2017). https://bit.ly/2YxOkZB

[161] K. Akilandeswari and M. Tamilselvi, On k-Heronian mean labeling, *Internat. J. Innovative Res. Sci., Eng. Tech.* **6**(9) (2017) 18019-18023.

[162] K. Akilandeswari and M. Tamilselvi, k-Heronian mean labeling of some graphs, *Internat. J. Pure Appl. Math.,* **117**(12) (2017) 173-180.

[163] D. G. Akka and N. S. Warad, Super magic strength of a graph, *Indian J. Pure Appl. Math.,* **41**(4) (2010) 557-568.

[164] R. Akhtar, A. B. Evans, and D. Pritikin, Representation numbers of stars, *Integers,* **10** (2010), A54, 733-745.

[165] R. Akhtar, A. B. Evans, and D. Pritikin, Representation numbers of complete multipartite graphs, *Discrete Math.,* **312**(6) (2012) 1158-1165.

[166] R. Akhtar, The representation number of some sparse graphs, *Discrete Math.,* **312**(22) (2012) 3417-3423.

[167] A. D. Akwu and D. O. A. Ajayi, On totally antimagic total labelings of complete bipartite graphs, *Roman. J. Math. Comput. Sci.,* **7**(1) 21-28. arXiv:1601.02112v1 [math. CO] 9 Jan 2016. https://www.researchgate.net/publication/290263695_On_totally_antimagic_total_labeling_of_complete_bipartite_graphs

[168] A. D. Akwu and D. O. A. Ajayi, Totally antimagic total labeling of ladders, prisms and generalised Petersen graphs, *J. Disc. Math. Sci. Cryp.*, **27**(1) (2024) 31-44. DOI:10.47974/JDMSC-1229

[169] S. Al-Addasi, O. A. AbuGhneim and H. Al-Ezeh, Divisor orientations of powers of paths and powers of cycles. *Ars Combin.* **94** (2010) 371-380. https://www.researchgate.net/publication/264964064_Divisor_orientations_of_powers_of_paths_and_powers_of_cycles

[170] S. Al-Addasi, O. A. AbuGhneim and H. Al-Ezeh, Merger and vertex splitting in divisor graphs, *Internat. Math. Forum*, **5**(38) (2010) 1861-1869. https://www.researchgate.net/publication/260351030_Merger_and_Vertex_Splitting_in_Divisor_Graphs

[171] S. Al-Addasi, O. A. AbuGhneim and H. Al-Ezeh, Characterizing powers of cycles that are divisor graphs, *Ars Combin.,* **97A** (2010) 447-451.

[172] S. Al-Addasi, O. A. AbuGhneim and H. Al-Ezeh, Further new properties of divisor graphs, *J. Combin. Math. Combin. Comput.,* **81** (2012) 261-272.

[173] L. Alam, A. Semaničová-Feňovčíková, and I.-L. Popa, Graceful local antimagic labeling of graphs: A pattern analysis using Python, *Symmetry* **17**(108) (2025) 1-12. https://doi.org/10.3390/sym17010108

[174] M. Aljohanis and S. N. Daoud, Edge odd graceful labeling in some wheel-related graphs, *Math.,* **12**(8) (2024) 1203 https://www.mdpi.com/2227-7390/12/8/1203

[175] M. Aljohani and S. N. Daoud, The edge odd graceful labeling of water wheel graphs, *Axioms* **14**(5) (2025) https://www.mdpi.com/2075-1680/14/1/5

[176] A. Aljouiee, On prime cordial labeling of graphs, *Kyungpook Math. J.*, **56** (2016) 41-46. http://dx.doi.org/10.5666/KMJ.2016.56.1.41

[177] M. G. Almeida and T. Singh, Some results on local distance antimagic chromatic number, arXiv:2410.9674v1

[178] G. A. N. Al-Roainee and S. S. Mahde, Mean harmonic energy of a graph, *Internat. J. Comp. Appl.*, (0975 - 8887) **186**(20) (2024) https://www.researchgate.net/publication/381003035_Mean_Harmonic_Energy_of_a_Graph

[179] R. E. L. Aldred and B. D. McKay, Graceful and harmonious labellings of trees. http://users.cecs.anu.edu.au/~bdm/papers/graceful.pdf

[180] R. E. L. Aldred, J. Širáň and M. Širáň, A note on the number of graceful labelings of paths, *Discrete Math.,* **261** (2003) 27-30.

[181] M. Alfalayleh, L. Brankovic, H. Giggins, M. Z. Islam, Towards the Graceful Tree Conjecture: A survey, In: *Proceed. AWOCA 2004*, 7-9 July, Ballina, Austrtalia (2004).

[182] G. Ali, M. Bača, F. Bashir, and A. Semaničová, On super vertex-antimagic labeling of disjoint union of paths, *AKCE J. Graphs Combin.,* **6** (2009) 11-20.

[183] G. Ali, M. Bača, M. Lascsaková, A Semaničová-Feňovčıková, A. ALoqaily, and N. Mlaiki, Modular total vertex irregularity strength of graphs, *AIMS Mathematics,* **8**(4) (2023) 7662-7671 doi:10.3934/math.2023384

[184] G. Ali, M. Bača, F. Bashir, and A. Semaničová, On face antimagic labeling of disjoint union of prisms, *Util. Math.,* **85** (2011) 97-112.

[185] G. Ali, M. Bača, Y. Lin, and A. Semaničová-Feňovčiková, Super-vertex-antimagic total labelings of disconnnected graphs, *Discrete Math.,* **309** (2009) 6048-6054.

[186] K. Ali, M. Hussain, A. Ahmad and M. Miller, Magic Labelings of type (a, b, c) of families of wheels, *Math. Computer Sci.,* **7** (2013) 315-319.

[187] K. Ali, M. Hussain, and Razzaq, Super edge-magic total labeings of a tree, *Util. Math.,* **91** (2013) 355-364.

[188] K. Ali, M. Hussain, H. Shaker, and M. Javaid, Super edge-magic total labeling of subdivided stars, *Ars Combin.*, **120** (2015) 161-167.

[189] K. Ali, S. T. R. Rizvi, and A. Semaničová-Feňovčiková, C_4-supermagic labelings of disjoint union of prisms, *Math. Rep. (Bucur.),* **18**(68) (2016) no. 3, 315-320.

[190] M. Ali, M.T Rahim, G. Ali, and M. Farooq, An upper bound for the radio number of generalized gear graph, *Ars Combin.,* **107** (2012) 161-168.

[191] S. Alikhani, W. Kocay, G.-C. Lau, and S.-M. Lee, On the k-edge magic graphs, *Elect. Notes Disc. Math.,* **45** (2014) 35-41.

[192] M. M. Ali Al-Shamiri, A. Elrokh, Y. El–Mashtawye, and S. Euat Tallah, The cordial labeling for the Cartesian product between paths and cycles, *Internat. J. Res.,*- Granthaalayah, **8**(3) (2020) 331-341. https://doi.org/10.5281/zenodo.3742934.

[193] A. H. Alkasasbeh, · E. Badr, · H. Attiya, and · H. M. Shabana, Radio number for friendship communication networks, *Mathematics*, **11** (2023) 4232. https://doi.org/10.3390/math11204232

[194] A. H. Alkasasbeh and D. Dyer, Graceful labellings of various cyclic snakes, https://doi.org/10.48550/arXiv.2012.10341 19 pages.

[195] A. H. Alkasasbeh, D. Dyer, and J. Howell, Graceful labellings of variable windmills using skolem-type sequences, *Ars Combin.,* (2024) **159** 109-131. https://combinatorialpress.com/ars-articles/volume-159/graceful-labellings-of-variable-windmills-using-skolem-type-sequences/

[196] O. Al-Mushayt, On the edge irregularity strength of products of certain families with P_2, *Ars Combin.,* **135** (2017) 323-334.

[197] O. Al-Mushayt, A. Ahmad, and M. K. Siddiqui, On the total edge irregularity strength of hexagonal grid graphs, *Australas. J. Combin.,* **53** (2012) 263-271.

[198] O. Al-Mushayt, A. Arshad, and M. K. Siddiqui, Total vertex irregularity strength of convex polytope graphs, *Acta Math. Univ. Comenian. (N.S.),* **82**(1) (2013) 29-37.

[199] O. Al-Mushayt, A. Ahmad, and M. K. Siddiqui, Irregular total labelings of convex polytope graphs having the same diameter, *Util. Math.,* **101** (2016) 295-307.

[200] N. Alon, Combinatorial nullstellensatz, *Combinatorics, Probabability and Computing*, **8** (1999) 7-29.

[201] N. Alon, G. Kaplan, A. Lev, Y. Roditty, and R. Yuster, Dense graphs are antimagic, *J. Graph Theory,* **47** (2004) 297-309.

[202] N. Alon and E. Scheinerman, Generalized sum graphs, *Graphs and Combin.,* **8** (1992) 23-29.

[203] H. Alpert, Rank numbers of grid graphs, *Discrete Math.,* **310** (2010) 3324-3333.

[204] H. Alpert, Rank numbers of path powers and grid graphs, personal communication (2013).

[205] K. A. Alsatami, Y. Algrawani, and A. Abd El-hay, A novel problem and algorithm for solving permuted cordial labeling of corona product between two graphs, *Math. Models Engin.,* (2025) 1-12. doi:DOI: 10.21595/mme.2024.24609

[206] F. Alyani, F. Firmansah, W. Giyarti, and K. A. Sugeng, The odd harmonious labeling of kC_n snake graphs for specific values, that is, for $n = 4$ and $n = 8$. *Proceed. IICMA 2013 Graph Combin.*, (2013) 225-230. https://www.researchgate.net/publication/282355295_THE_ODD_HARMONIOUS_LABELING_OF_kCn-_SNAKE_GRAPHS_FOR_SPESIFIC_VALUES_OF_n_THAT_IS_FOR_n_4_AND_n_8

[207] K. Al Azizu, L. Yulianti, S. Sy, and N. Narwen, On super edge magic total labeling of the branched-prism graph, *Internat. J. Progressive Sci. Tech.*, **16**(2) (2019) 203-206. https://ijpsat.org/index.php/ijpsat/article/view/1288/659

[208] K. Al Azizu, L. Yulianti, N. Narwen, and S. Sy, On super (a, d)-edge antimagic total labeling of branched-prism graph, *Indonesian J. Combin.*, **5**(1) (2021) 11-16. doi:10.19184/ijc.2021.5.1.2

[209] R. Amalia and Masruroh, Local antimagic vertex total coloring on fan graph and graph resulting from comb product operation, *J. Physics:Conf. Ser.*, **1836** (2021) 012012 https://www.researchgate.net/publication/350332971_Local_antimagic_vertex_total_coloring_on_fan_graph_and_graph_resulting_from_comb_product_operation

[210] D. Amar and O. Togni, Irregularity strength of trees, *Discrete Math.,* **190**(1-3) (1998) 15-38.

[211] A. Amara Jothi, J. Baskar Babujee, and N. G. David, On face magic labeling of duplication graphs. http://oaji.net/articles/2017/296-1504963989.pdf

[212] A. Amara Jothi, N. G. David, J. Baskar Babujee, Super edge bimagic labeling graphs, *Int. J. Math. Sci. Comput.,* **5**(1) (2015) 39-43.

[213] B. Ambili and G. Suresh Singh, Graceful labeling of arbitrary strong supersubdivisions of graphs, *Graph Theory Notes N. Y.,* **61** (2011) 31-36.

[214] M. Amini and K. Eshghi, Constraint programming models and population-based simulated annealing algorithm for finding graceful and α-labeling of quadratic graphs, *Iran J. Comput. Sci.,* **1**(3) (2018) 155-164. https://link.springer.com/article/10.1007/s42044-018-0012-7

[215] P. Amudha and J. Jayapriya, Pair sum modulo labeling of graph, *Adv. Math.: Sci. J.*, **10**(2) (2021) 723-727. DOI:https://doi.org/10.37418/amsj.10.2

[216] P. Amudha, J. Jayapriya, and J. Gowri, An algorithmic approach for encryption using graph labeling,
textitJ. Phys.: Conf. Ser., (2021) 1770 012072, 1-10. https://downloads.hindawi.com/journals/mpe/2021/5562544.pdf

[217] R. S. Amutha, P. Sugapriya, and S. Vaanmathi, Lucky-edge geometric mean labeling of ladder and star graph, *GIS Sci. J.*, **8**(2) (2021) 1752–1762.

[218] S. Amutha and K. M. Kathiresan, Pendant edge extensions of two families of graphs, *Proceed. Nat. Seminar on Algebra and Discrete Math.,* Univ. Kerala, Thiruvananthapuram, India, 146-150.

[219] S. Amutha and M. Uma Devi, Super graceful labeling for some families of fan graphs, *J. Comput. Math. Sci.*, **10**(8) (2019) 1551-1562. http://www.compmath-journal.org/dnload/S-Amutha-and-M-Uma-Devi-/CMJV10I08P1551.pdf

[220] S. Amutha and M. Uma Devi, Total edge Fibonacci irregular labeling for fan, wheel and umbrella graph, *J. Comp. Math. Sci*, **10**(12) (2019) 1654-1664. http://www.compmath-journal.org/dnload/S-Amutha-and-M-Uma-Devi-/CMJV10I12P1654.pdf

[221] D. Amuthavalli and M. Premkumar, Super Fibonacci graceful labeling of some cycle related graphs, *Internat. J. Math. Trends Tech.,***38**(3) (2016) https://www.researchgate.net/publication/348002618_Super_Fibonacci_Graceful_Labeling_Of_Some_Cycle_Related_Graphs

[222] D. Amuthavalli and O. V. S. Sundaram, Fibonacci graceful anti-magic labeling for Zodiac signs in astronomy, *J. Astatic Soc. Mumbai*, **XCVI**(26) (2023) 7-9. https://www.researchgate.net/publication/374062123_FIBONACCI_GRACEFUL_ANTI-MAGIC_LABELING_FOR_ZODIAC_SIGNS_IN_ASTRONOMY

[223] K. Amuthavalli, Graph Labeling and its Applications-Some Generalization of Odd Mean Labelings, Ph. D. Thesis, (2010) Mother Teresa Women's University, Kodaikanal Tamilnadu, India.

[224] K. Amuthavalli and S. Dineshkumar, k-Odd edge mean labeling of some basic graphs, *J. Informatics Math. Sci.*, **9**(3) (2017) 915-921. https://www.rgnpublications.com/journals/index.php/jims/article/view/1012/757

[225] K. Amuthavalli and P. Sugapriya, Reverse super edge bimagic labeling of star related graphs, *Internat. J. Res. Engin. Appl. Management (IJREAM)*, **05** (2019) 480-484.. https://ijream.org/papers/IJREAMV05I0250124.pdf

[226] K. Amuthavalli and P. Sugapriya, Reverse super edge-trimagic labeling labeling for star related graphs, *Malaya J. Mat.,* **7**(3) (2019) 519-525. https://www.malayajournal.org/articles/MJM07030025.pdf

[227] S. Anusa, S. S. Sandhya, and S. Somasundaram, New classes of mean graphs, *J. Disc. Math. Sci. Crypt.*,, **22**(3) (2019) 351-360. DOI:10.1080/09720529.2018.1534369

[228] M. Andar, S. Boxwala, and N. Limaye, Cordial labelings of some wheel related graphs, *J. Combin. Math. Combin. Comput.,* **41** (2002) 203-208.

[229] M. Andar, S. Boxwala, and N. Limaye, A note on cordial labelings of multiple shells, *Trends Math.,* (2002) 77-80.

[230] M. Andar, S. Boxwala, and N. Limaye, On the cordiality of the t-uniform homeomorphs–I, *Ars Combin.*, **66** (2003) 313-318.

[231] M. Andar, S. Boxwala, and N. Limaye, On the cordiality of the t-uniform homeomorphs–II (Complete graphs), *Ars Combin.*, **67** (2003) 213-220.

[232] M. Andar, S. Boxwala, and N. Limaye, New families of cordial graphs, *J. Combin. Math. Combin. Comput.,* **53** (2005) 117-154.

[233] M. Andar, S. Boxwala, and N. Limaye, On the cordiality of the t-ply $P_t(u, v)$, *Ars Combin.,* **77** (2005) 245-259.

[234] M. Andar, S. Boxwala, and N. Limaye, On the cordiality of corona graphs, *Ars Combin.,* **78** (2006) 179-199.

[235] P. P. Andharia and V. J. Kaneria, Even sum labeling of some graphs, *Int. J. Comp. Math. Sci.,* **7**(5) (2018) 199-202.

[236] P. P. Andharia and V. J. Kaneria, Even sum property of J_n, $B(3, n)$, TB_n, P_m $(+)$ $\overline{K_n}$ and $(\overline{K_n} \cup P_3) + 2K_1$, *IOSR J. Math.,* **18**(1) Ser. II (2022) 24-30.

[237] J. D. Andoyo, On Legendre cordial labeling of complete graphs, https://doi.org/10.48550/arXiv.2509.09528

[238] J. D. Andoyo, On arithmetic cordial labeling of product graphs, https://doi.org/10.48550/arXiv.2608.11837

[239] J. D. Andoyo, J. Clarisse, C. Prudencio, and R. F. Rulete, On arithmetic cordial labeling of some graphs, *Open J. Math. Sci.*, **10** (2026) 438-454. doi:DOI:10.30538/oms2026.0297

[240] J. D. Andoyo and R. Rulete On Legendre cordial labeling of some graphs, *Eur. J. Math. Stat.,* **6**(5) (2025) 1-7. https://ej-math.org/index.php/ejmath/article/view/409/215

[241] A. D. C. Angel, J. R. M. Antalan, J. Loureynz, F. Gamurot, and R. P. Tagle, On the edge-gracefulness of water wheel-type graphs and their splitting graphs, *European J. Prue App. Math.*, **18**(4) (2025) Article Number 6425 33pp. https://www.ejpam.com/index.php/ejpam/article/view/6425

[242] D. A. Angel Sherin and V. Maheswari, Encoding and decryption process using edge magic labeling, *J. Physics Conf. Series* **1362**:012024 (2019). doi:10.1088/1742-6596/1362/1/012024

[243] D. A. Angel Sherin, V. Maheswari, and V. Balaji, Encryption of dual numbers using edge injective labeling, *Internat. J. Multidisciplnary Res. Arts, Sci. Commerce,* (IJMRASC) **1**(40 (2021) 54-60. https://www.sdnbvc.edu.in/wp-content/uploads/2022/01/PAPERID7.pdf

[244] M. Anholcer, Product irregularity strength of graphs, *Discrete Math.,* **309** (2009) 6434-6439.

[245] M. Anholcer, Product irregularity strength of certain graphs, *Ars Math. Contemp.,* **7**(1) (2014) 23-29.

[246] M. Anholcer and S. Cichacz, Group irregular labelings of disconnected graphs, *Contributions Discrete Math.,* **12**(2) (2017) 158-166.

[247] M. Anholcer, S. Cichacz, Note on distance magic products $G \circ C_4$, *Graphs and Combin.,* **31**(5) (2015) 1117-1124. doi:10.1007/s00373-014-1453-x.

[248] M. Anholcer, S. Cichacz, D. Froncek, R. Simanjuntak, and J. Qiu, Group distance magic and antimagic hypercubes,*Discrete Math.* (2021) http://kbo.ue.poznan.pl/anholcer/publ/20210829_hypercubes.pdf

[249] M. Anholcer, S. Cichacz, R. Jura, A. Marczyk, Note on group irregularity strength of disconnected graphs, *Open Math.,* **16** (2018) 154-160, doi:10.1515/math-2018-0017.

[250] M. Anholcer, S. Cichacz, and M. Milanič, Group irregularity strength of connected graphs, *J. Combin. Optim.,* (2013) **30**(1) (2015) 1-17. doi:10.1007/s10878-013-9628-6

[251] M. Anholcer, S. Cichacz, I. Peterin, and A. Tepeh, Distance magic labeling and two products of graphs, *Graphs and Combin.,* **31**(5) (2015) 1117-1124. doi:10.1007/s00373-014-1455-8

[252] M. Anholcer, S. Cichacz, I. Peterin, and A. Tepeh, Distance magic labeling and two products of graphs, *Graphs Combin.,* **31**(5) (2015) 1125-1136.

[253] M. Anholcer, S. Cichacz, I. Peterin, and A. Tepeh, Group distance magic labeling of direct product of graphs, *Ars Math. Contemp.* **9** (2015) 93-108.

[254] M. Anholcer, S. Cichacz, and J. Przybyło, Linear bounds on nowhere-zero group irregularity strength and nowhere-zero group sum chromatic number of graphs, *Appl. Math. Comput.*, *Appl. Math. Comput.*, **343** (2019) 149-155. arXiv:1805.05204v1

[255] M. Anholcer, S. Cichacz, I. Peterin, and A. Tepeh, Group distance magic labeling of direct product of graphs,]Ars Math. Contemp., **9**(1) (2015) 93-107.

[256] M. Anholcer, M. Karoński, and F. Pfender, Total vertex irregularity strength of forests, *Discrete Math.,* **312**(2) (2012) 229-237. arXiv:1103.2087v1 [math.CO] 10 Mar. 2011.

[257] M. Anholcer, M. Karoński, and J. Przbylo, A new bound for the total vertex irregularity strength of graphs, *Discrete Math.,* **309** (2009) 6316-6317.

[258] M. Anholcer and C. Palmer, Irregular labelings of circulant graphs, *Discrete Math.,* **312**(23) (2012) 3461-3466.

[259] D. J. Anick, Counting graceful labelings of trees: a theoretical and empirical study, *Discrete Appl. Math.,* **198** (2016) 65-81.

[260] K. Anitha, B. Selvam, and K. Thirusangu, Mean, odd mean and even mean labeling for the extended duplicate graph of kite graph, *Inter. J. Math. Trends Tech.*, **42**(1) (2017) 39-44. http://www.ijmttjournal.org/2017/Volume-42/number-1/IJMTT-V42P506.pdf

[261] K. Anitha, B. Selvam, and K. Thirusangu, k-Graceful, odd-even graceful, Heronian mean and analytic mean labeling for the extended duplicate graph of kite graph, *Internat. J. Engin. Tech.* **7**(4.10) (2018) 355-359.

[262] V. Annamma and P. Vijitha, Sum divisor cordial labeling of certain graphs, *J. Alg. Stat.*, **13**(2) (2022) 3238-3242. https://publishoa.com/index.php/journal/article/view/1054/911

[263] K. Annathurai, R. Ponraj, and R. Kala, Some 3-remainder cordial graphs,*Global J. Engin. Sci. Res.*, **5**(6) (2018) 169-175.

[264] K. Annathurai, R. Ponraj, and R. Kala, Further results on 3-remainder cordial labeling of graphs, *Internat. J. Management, IT Engin.,* **8**(8) (2018) 120-129.

[265] K. Annathurai and M. Gomathi, Absolutely k-harmonious labeling of product graphs, *Amer. J. Appl. Math.*, (2026) **14**(5) 303-306. doi:10.11648/j.ajam.20261405.14

[266] J. R. M. Antalan, A. Angel, J. L. Gamurot, and R. Tagle, Edge-graceful usual fan graphs, arXiv:2412.08338

[267] A. Anto Kinsley and P. Siva Ananthi, DdDistance in graphs, *Imperial J. Interdisc. Res.*, **3**(2) (2017) 2454-1362.

[268] S. Anubala and · V. Ramachandran, E-super arithmetic graceful labelling of $H_i(m,m)$, $H_i^{(1)}(m,m)$ and chain of even cycles, *Acta Universitatis Sapientiae* **15**(1) (2023) 81-90. https://api.semanticscholar.org/CorpusID:260705405

[269] S. Anusa, S. S. Sandhya, and S. Somasundaram, New classes of mean graphs, *J. Discrete Math. Sci. Cryptogr.,* **22**(3) (2019) 351-360.

[270] R. Aravamudhan and M. Murugan, Numbering of the vertices of $K_{a,1,b}$, unpublished.

[271] K. Ariyanti, Z. Z. Barack, P. John, and A. Bustamam, Modular irregularity strength on some corona products of graphs, *J. Indonesian Math. Soc.*, **32**(1) (2026) 1-15. https://jims-a.org/index.php/jimsa/article/view/2248/473

[272] I. C. Arkut, R. C. Arkut, and A. Basak, Topology constrained label switching for multicast routing, *Proceed. Eighth IEEE Internat. Sympos. Comput. Comm. (ISCC'03),* 1530-1346/03 1-8. http://www.academia.edu/4370947/

[273] I. Arkut, R. Arkut, and N. Ghani, Graceful label numbering in optical MPLS networks, In: *Proc. SPIE,* **4233** (2000) 1-8 OptiComm 2000: Optical Networking and Communications. Imrich Chlamtac: Ed (2000).

[274] A. Armstrong and D. McQuillan, Vertex-magic total labelings of even complete graphs, *Discrete Math.,* **311** (2011) 676-683. https://www.sciencedirect.com/science/article/pii/S0012365X11000148

[275] S. Arumugam and J. Bagga, Graceful labeling algorithms and complexity—a survey, *Indones. Math. Soc.,* Special Edition (2011) 1-9.

[276] S. Arumugam, D. Fronček, and N. Kamatchi, Distance magic graphs—a survey, *J. Indonesian Math. Soc.*, Special Edition, 11–26 (2011).

[277] S. Arumugam and K. Germina, On indexable graphs, *Discrete Math.*, **161** (1996) 285-289.

[278] S. Arumugam, K. Germina, and T. Anandavally, On additively geometric graphs, *J. Combin. Math. Combin. Comput.,* **70** (2009) 207-216.

[279] S. Arumugam and N. Kamatchi, On (a, d)-distance antimagic graphs, *Australas. J. Combin.,* **54** (2012) 279-287.

[280] S. Arumugam, N. Kamatchi, and P. Kovář, Distance magic graphs, *Util. Math.,* **99** (2016) 131-142. https://www.researchgate.net/publication/305426259_Distance_Magic_Graphs/link/58df56cf4585153bfe947e3e/download

[281] S. Arumugam, Y.-C. Lee, K. Premalatha, T.-M. Wang, On local antimagic vertex coloring for corona products of graphs, arXiv:1808.04956.pdf.

[282] S. Arumugam, M. Miller, O. Phanalasy, and J. Ryan, Antimagic labeling of generalized pyramid graphs, *English Series Acta Mathematica Sinica*, **30** (2) (2014) 283-290. https://actamath.cjoe.ac.cn/Jwk_sxxb_en/EN/10.1007/s10114-014-2381-7

[283] S. Arumugam and M. Nalliah, Super (a, d)-edge antimagic total labelings of friendship and generalized friendship graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), 103-110,

[284] S. Arumugam and M. Nalliah, Super (a, d)-edge antimagic total labelings of friendship graphs, *Australas. J. Combin.,* **53** (2012) 237-243. https://www.researchgate.net/publication/269010612_Super_ad-edge_antimagic_total_labelings_of_friendship_graphs/link/5779f9d708ae4645d611f51d/download

[285] S. Arumugam and M. Nalliah, Super $(a, 3)$-edge-antimagic total labelings for union of two stars, *Util. Math.,* **101** (2016) 337-349.

[286] S. Arumugam, K. Premalatha, M. Bača, and A. Semaničová-Feňovčíková, Local antimagic vertex coloring of a graph, *Graphs Combin.*, **33**(2) (2017) 275-285. https://link.springer.com/article/10.1007/s00373-017-1758-7

[287] M. Arnold and F. Vitor, A flexible edge search algorithm for graceful tree labeling, *Internat. J. Comput. Sci. Math. Theory*, (IJCSMT) **12**(3) (2026) 295-333. https://www.iiardjournals.org/get/IJCSMT/VOL.%2012%20NO.%203%202026/A%20FLEXIBLE%20EDGE%20SEARCH%20ALGORITHM%20295-333.pdf

[288] S. T. Arockiamary, J. Baskar Babujee, and G. Vijayalakshmi, k-prime total labeling of graphs, *Advances Appl. Disc. Math.,* **35** (2022) 45-59. http://dx.doi.org/10.17654/0974165822051

[289] S. Arockiaraj, A. Durai Baskar and A. Rajesh Kannan, F-root square mean labeling of graphs obtained from paths, *Internat. J. Math. Combin.,* **2**(2017), 92-104. https://fs.unm.edu/IJMC/FRootSquareMeanLabelingOfGraphs.pdf

[290] S. Arockiaraj, A. Durai Baskar and A. Rajesh Kannan, F-root square mean labeling of line graph of some graphs, *Utilitas Math.,* **112**(2019), 11-32. https://fs.unm.edu/IJMC/FRootSquareMeanLabelingOfGraphs.pdf

[291] S. Arockiaraj and P. Mahalakshmi, On odd sum graphs, *Internat. J. Math. Combin.,* **4** (2013) 59-77. https://fs.unm.edu/IJMC/OnOddSumGraphs.pdf

[292] S. Arockiaraj, P. Mahalakshmi, and P. Namasivayam, Odd sum labeling of some subdivision graphs, *Kragujevac J. Math.,* **38**(1) (2014) 203-222. http://elib.mi.sanu.ac.rs/files/journals/kjm/42/kjom3801-16.pdf

[293] S. Arockiaraj, P. Mahalakshmi, and P. Namasivayam, Odd sum labelings of some splitting graphs, *Util. Math.,* **105** (2017) 53-73.

[294] S. Arockiaraj, P. Mahalakshmi, and P. Namasivayam, Odd sum labeling of graphs obtained by duplicating any edge of some graphs. *Electron. J. Graph Theory Appl. (EJGTA),* **3**(2) (2015) 197-215. https://www.ejgta.org/index.php/ejgta/article/view/133/pdf_7

[295] S. Arockiaraj, P. Mahalakshmi, and P. Namasivayam, Odd sum labeling of graphs obtained from some graph operations, *J. Graph Labeling,* **2**(2) (2015) 217-233.

[296] S. Arockiaraj and A. Meena, On F-face magic mean labeling of some planar graphs, *Internat. J. Pure Appl. Math.,* **117**(6) (2017) 1-8. https://acadpubl.eu/jsi/2017-117-5-6/articles/6/1.pdf

[297] S. Arockiaraj and A. Meena Kumari, On F-face magic mean labeling of some duplicated graphs, *Internat. J. Appl. Math.*, **31**(2) (2018) 231-240. https://www.diogenes.bg/ijam/contents/2018-31-2/5/5.pdf

[298] S. Arockiaraj, A. Rajesh Kannan, and A. Durai Baskar, Super F-centroidal mean graphs, *Internat. J. Math. Combin.,* **3** (2019) 113-127. https://fs.unm.edu/IJMC/SuperF-CentroidalMeanGraphs.pdf

[299] S. Arockiaraj, A. Rajesh Kannan, and A. Durai Baskar, F-centroidal mean labeling of graphs obtained from paths, *Internat. J. Math. Combin.,* **4** (2019) 122-135.

[300] S. Arockiaraj, A. R. Kannan, P. Manivannan, and A. Durai Baskar, FCM labeling of some graphs and its line graph, *J. Physics: Confer. Ser.*, **1597** (2020) 012033 1-8. doi:DOI:10.26637/MJM0703/0025

[301] S. Arockiamary and G. Vijayalakshmi, k-prime total labeling of shell related graphs, *Adv. Appl. Discr. Math.,* **39**(1) (2023) 1-27. http://dx.doi.org/10.17654/0974165823033

[302] S. T. Arockiamary and G. Vijayalakshmi, Vertex k-prime labeling of cyclic snakes, *Comm. Math. Appl.,* **14**(1) (2023) 9-20. https://www.researchgate.net/publication/370927885_Vertex_k-Prime_Labeling_of_Cyclic_Snakes

[303] S. T. Arockiamary and G. Vijayalakshmi, Vertex k-prime labeling on graphs, *European Chem. Bull.,* **12**(4) (2023) 9627-9633. https://www.researchgate.net/publication/370927953_Vertex_k-Prime_Labeling_on_graphs

[304] S. T. Arockiamary and G. Vijayalakshmi, Vertex k-prime labeling of theta graphs, *Indian J. Sci. Tech.*, **16**(26) (2023) 2008-2015. https://www.researchgate.net/publication/372405676_Vertex_k-Prime_Labeling_of_Theta_Graphs

[305] S. T. Arockiamary and P. Vijayalakshmi, A comparative study on prime vertex k-prime labelings of one point union of path graphs, *TWMS J. App. Eng. Math.*,**15**(9) (2025) 2398-2405. https://www.researchgate.net/publication/395334933_A_COMPARATIVE_STUDY_ON_PRIME_AND_VERTEX_k-PRIME_LABELING_OF_ONE_POINT_UNION_OF_PATH_GRAPHS

[306] J. Arthy, Cycle and path related graphs on L-cordial labeling, *Internat. J. Innovative Sci. Res. Tech.*, **4**(6) (2019). https://ijisrt.com/assets/upload/files/IJISRT19JU676_(2).pdf.pdf

[307] Y. F. Ashari and A. N. M. Salman, (H_1, H_2)-supermagic labelings for some shackles of connected graphs H_1 and H_2, (2019) *J. Phys.: Conf. Ser.,* 1127 012059. https://iopscience.iop.org/article/10.1088/1742-6596/1127/1/012059/pdf

[308] T. S. Ashika and S. Asha, Radio contra harmonic mean number of graphs, *Indian J. Sci. Tech.,* **17**(16) (2024) 1647-1653. https://www.researchgate.net/publication/379968316_Radio_Contra_Harmonic_Mean_Number_of_Graphs

[309] F. Ashraf, M. Bača, Z. Kimáková, and A. Semaničová-Feňovčíková, On vertex and edge H-irregularity strengths of graphs, *Discrete Math. Algorithms Appl.,* **8**(4) (2016) 1650070, 13 pp.

[310] F. Ashraf, M. Bača, M. Lascsáková, and A. Semaničová-Feňovčíková, On H-irregularity strengths of graphs, *Disc. Math. Graph Th.,* **37** (2017) 1067-1078.

[311] F. Ashraf, M. Bača, A. Semaničová-Feňovčíková, and S. W. Saputro, On cycle-irregularity strength of ladders and fan graphs, *Elect. J. Graph Th. Appl.,* **8**(1) (2020) 181-194. https://www.ejgta.org/index.php/ejgta/article/view/449/pdf_133

[312] F. Ashraf, M. Bača, A. Semaničová-Feňovčíková, A. Shabbirc, On H-irregularity strengths of G-amalgamationm of graphs, *Elect. J. Graph Th. Appl.,* **5** (2) (2017) 325-334. https://www.ejgta.org/index.php/ejgta/article/view/391/pdf_58

[313] F. Ashraf, M. Bača, A. Semaničová-Feňovčíková, and M. K. Siddiqui, On H-irregularity strength of ladders and fan graphs, *AKCE Int. J. Graphs Comb.,* **17**(1) (2020) 213-219. https://doi.org/10.1016/j.akcej.2019.04.002

[314] Y. F. Ashari, A. N. M. Salman, and R. Simanjuntak, On forbidden subgraphs of (K_2, H)-sim-(super) magic graphs, *Symmetry* **13** (2021) 1346. https://doi.org/10.3390/sym13081346

[315] J. Ashwini, S. P. Selvam, and R. B. Gnanajothi, Some new results on lucky labeling, *Baghdad Sci. J.*, **20**(1) (2023) 365-370. https://bsj.uobaghdad.edu.iq/index.php/BSJ/article/view/8569/4331

[316] M. Asif, G. Ali, M. Numan, and A. Semaničová-Feňovčíková, Cycle-supermagic labeling for some families of graphs, *Util. Math.,* **103** (2017) 51-59. https://www.researchgate.net/publication/343336247_Cycle-supermagic_labeling_for_some_families_of_graphs

[317] M. A. Asim and R. Hasni, Iteractive algorithm for computing irregularity strength of complete graph, *Ars Combin.,* **138** (2018) 17-24.

[318] J. Asplund and N. B. Fox, Minimum coprime labelings for operations on graphs, *Integers.* **19** (2019) Paper No. A24, 26 pp. https://www.semanticscholar.org/reader/77ac086965b32f63ec2684163a0434c45a64b58d

[319] E. Asumpta, Purwanto, and T. D. Chandra, Odd harmonious labeling of some family of snake graphs, AIP Conference Proceedings 2639(1):040002 (2022) 7 pp. Conference: Proceedings of the II International Scientific Conference on Advances in Science, Engineering,and Digital Education: (ASEDU-II 2021 doi:10.1063/5.0111278

[320] H. Attiya, N. Ahmed, and F. Salama, Total edge irregularity strength of star snake graphs, *European J. Pure Appl. Math.,* **18**(2) (2025) Article Number 5673. https://www.researchgate.net/publication/375464789_Total_Edge_Irregularity_Strength_of_Star_Snake_Graphs

[321] S. Avadayappan, P. Jeyanthi, and R. Vasuki, Super magic strength of a graph, *Indian J. Pure Appl. Math.,* **32** (2001) 1621-1630. https://www.researchgate.net/publication/257305617_2_SAvadayappanPJeyanthi_and_RVasuki_Super_Magic_Strength_of_a_Graph_Indian_Journal_of_Pure_and_Applied_Mathematics3211_20011621-_1630

[322] S. Avadayappan, R. Vasuki, and P. Jeyanthi, Magic strength of a graph, *Indian J. Pure Appl. Math.,* **31** (2000) 873-883. https://www.researchgate.net/publication/257305454_1_SAvadayappanPJeyanthi_and_RVasuki_Magic_Strength_of_a_Graph_Indian_Journal_of_Pure_and_Applied_Mathematics_307_2000_873-883

[323] S. Avadayappan and R. Vasuki, New families of mean graphs, *Internat. J. Math. Combin.,* **2** (2010) 68-80. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=a78f64610c786c013e6f0a4c9ab8caf3dc8f9e84

[324] J. Ayel and O. Favaron, Helms are graceful, in Progress in Graph Theory (Waterloo, Ont., 1982), Academic Press, Toronto, Ont. (1984) 89-92.

[325] A. Azaizeh, R. Hasni, G. C. Lau, and A. Ahmad, 3-total edge product cordial labeling of complete, bipartite and generalized friendship graphs, *Util. Math.,* **106** (2018) 259-270. https://www.researchgate.net/publication/324827903_3-total_edge_product_cordial_labeling_of_complete_bipartite_and_generalised_friendship_graphs

[326] M. Azeem, Cycle-super magic labeling of polyomino linear and zig-zag chains, *J. Operations Intell.*, **1**(1) (2023) 67-81. https://jopi-journal.org/index.php/jopi/article/view/5/5

[327] S. Babitha and J. Baskar Babujee, Prime cordial labeling on graphs, *Internat. Scholarly Sci. Res. Innovation* **7**(5) (2013) 848-853. https://www.semanticscholar.org/paper/Prime-Cordial-Labeling-on-Graphs-Babitha-Bab422aff9b3081b135bc04190ec7a117ed37f67197

[328] M. Baća, P. Jeyanthi, P. Selvagopal, N. T. Muthu Raja, and A. Semaničová-Feňovčíková, Ladders and fan graphs are cycle-antimagic, *Hacettepe J. Math. & Stat.*, **49**(3) (2020) 1093-1106. https://www.researchgate.net/publication/338275842_Ladders_and_fan_graphs_are_cycle-antimagic

[329] M. Bača, On magic and consecutive labelings for the special classes of plane graphs, *Util. Math.,* **32** (1987) 59-65. https://www.researchgate.net/publication/262723400_On_magic_and_consecutive_labelings_for_the_special_classes_of_plane_graphs

[330] M. Bača, Labelings of two classes of convex polytopes, *Util. Math.,* **34** (1988) 24-31. https://www.researchgate.net/profile/Martin-Baca/publication/262723554_Labelings_of_two_classes_of_convex_polytopes/links/0f317538edd86d4a82000000/Labelings-of-two-classes-of-convex-polytopes.pdf

[331] M. Bača, Labelings of m-antiprisms, *Ars Combin.,* **28** (1989) 242-245. https://www.researchgate.net/profile/Martin-Baca/publication/262723393_Labelings_of_m-antiprisms/links/00b7d538edd56d09f8000000/Labelings-of-m-antiprisms.pdf

[332] M. Bača, On magic labelings of Möbius ladders, *J. Franklin Inst.,* **326** (1989) 885-888. https://www.sciencedirect.com/science/article/pii/0016003289900100

[333] M. Bača, On magic labelings of type $(1, 1, 1)$ for three classes of plane graphs, *Math. Slovaca,* **39** (1989) 233-239. https://www.sciencedirect.com/science/article/pii/001600329290053J

[334] M. Bača, On certain properties of magic graphs, *Util. Math.,* **37** (1990) 259-264. https://www.researchgate.net/publication/262723457_On_certain_properties_of_magic_graphs

[335] M. Bača, On magic labelings of m-prisms, *Math. Slovaca,* **40** (1990) 11-14. https://www.researchgate.net/publication/262723194_On_magic_labellings_of_m-prisms

[336] M. Bača, On consecutive labelings of plane graphs, *J. Franklin Inst.,* **328** (1991) 249-253. https://www.sciencedirect.com/science/article/pii/001600329190033Y

[337] M. Bača, On magic labelings of honeycomb, *Discrete Math.,* **105** (1992) 305-311. https://doi.org/10.1016/0012-365X(92)90153-7

[338] M. Bača, On magic labelings of grid graphs, *Ars Combin.,* **33** (1992) 295-299. https://www.researchgate.net/publication/262723621_On_magic_labelings_of_grid_graphs

[339] M. Bača, On magic labelings of type $(1,1,1)$ for the special class of plane graphs, *J. Franklin Inst.,* **329** (1992) 549-553. https://www.sciencedirect.com/science/article/pii/001600329290053

[340] M. Bača, On magic labellings of convex polytopes, *Ann. Disc. Math.,* **51** (1992) 13-16.

[341] M. Bača, Labelings of two classes of plane graphs, *Acta Math. Appl. Sinica,* **9** (1993) 82-87. https://www.researchgate.net/publication/225996946_Labelings_of_two_classes_of_plane_graphs

[342] M. Bača, Face antimagic labelings of convex polytopes, *Util. Math.,* **55** (1999) 221-226. https://www.researchgate.net/publication/266240246_Face_antimagic_labelings_of_convex_polytopes

[343] M. Bača, Antimagic labelings of antiprisms, *J. Combin. Math. Combin. Computing,* **35** (2000) 217-224. https://combinatorialpress.com/article/jcmcc/Volume%20035/vol-035-paper%2015.pdf

[344] M. Bača, Special face numbering of plane quartic graphs, *Ars Combin.,* **57** (2000) 285-292. https://www.researchgate.net/publication/220620268_Special_Face_Numbering_of_Plane_Quartic_Graphs

[345] M. Bača, Consecutive-magic labeling of generalized Petersen graphs, *Util. Math.,* **58** (2000) 237-241. https://www.researchgate.net/publication/262723294_Consecutive-magic_labeling_of_generalized_Petersen_graphs

[346] M. Bača, On face antimagic labelings of plane graphs, *J. Combin. Math. Combin. Comput.,* **53** (2005) 33-38.

[347] M. Bača and C. Barrientos, Graceful and edge-antimagic labelings, *Ars Combin.,* **96** (2010) 505-513. https://www.researchgate.net/publication/262696982_Graceful_and_edge-antimagic_labeling

[348] M. Bača and C. Barrientos, On super edge-antimagic total labelings of mK_n, *Discrete Math.,* **308** (2008) 5032-5037. https://www.sciencedirect.com/science/article/pii/S0012365X07007212

[349] M. Bača and F. Bashir, On super d-antimagic labelings of disjoint union of prisms, *AKCE J. Graphs Combin.,* **6** (2009) 31-39. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088869

[350] M. Bača, Y. Bashir, M. F. Nadeem, and A. Shabbir, On super edge-antimagic total labeling of Toeplitz graphs. Mathematics in the 21st century, 1–10, Springer Proc. Math. Stat., 98, Springer, Basel, 2015. https://link.springer.com/content/pdf/10.1007/978-3-0348-0859-0_1?pdf=chapter+toc

[351] M. Bača, F. Bashir, and A. Semaničová, Face antimagic labeling of antiprisms, *Util. Math.,* **84** (2011) 209-224. https://www.researchgate.net/publication/262693003_Face_antimagic_labelings_of_antiprisms

[352] M. Bača E. T. Baskoro, L. Brankovic, S. Jendrol', Y. Lin, O. Phanalasy, J. Ryan, A. Semaničová-Feňovčíková, Slamin, K. A. Sugeng, A survey of face-antimagic evaluations of graphs, *Australa. J. Combin.,* **69**(3) (2017) 382-393. https://www.researchgate.net/publication/320285837_A_survey_of_face-antimagic_evaluations_of_graphs

[353] M. Bača, B. Baskoro, and Y. Cholily, Face antimagic labelings for a special class of plane graphs C_a^b, *J. Combin. Math. Combin. Comput.,* **55** (2005) 5-15. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%201.pdf

[354] M. Bača, B. Baskoro, and Y. Cholily, On d-antimagic labelings for a special class of plane graphs, *J. Combin. Math. Combin. Comput.,* **61** (2007) 21-32. https://combinatorialpress.com/article/jcmcc/Volume%20061/vol-061-paper%202.pdf

[355] M. Bača, B. Baskoro, Y. Cholily, S. Jendroľ, Y. Lin, M. Miller, J. Ryan, R. Simanjuntak, Slamin, and K. Sugeng, Conjectures and open problems in face antimagic evaluations of graphs, *J. Indonesian Math. Soc. (MIHMI),* **11** (2005) 175-192.

[356] M. Bača, B. Baskoro, S. Jendroľ and M. Miller, Antimagic labelings of hexagonal plane maps, *Util. Math.,* **66** (2004) 231-238. https://www.researchgate.net/publication/268664044_Antimagic_labelings_of_hexagonal_plane_maps

[357] M. Bača, B. Baskoro, and M. Miller, On d-antimagic labelings of honeycomb, *Proceedings 12th Australian Workshop on Combinatorial Algorithms,* Bandung, Indonezia (2001) 45-50.

[358] M. Bača, E. T. Baskoro, and M. Miller, Antimagic valuations for the special class of plane graphs, *Lecture Note Computer Science-LNCS,* **3330** (2005) 58-64. https://link.springer.com/content/pdf/10.1007/978-3-540-30540-8_6.pdf

[359] M. Bača, E. T. Baskoro, M. Miller, J. Ryan, R. Simanjuntak, and K. A. Sugeng, Survey of edge antimagic labelings of graphs, *J. Indonesian Math. Soc.,* (MIHMI) **12** (2006) 113-130. https://www.researchgate.net/publication/228738097_Survey_of_edge_antimagic_labelings_of_graphs

[360] M. Bača, E. T. Baskoro, R. Simanjuntak, and K. A. Sugeng, Super edge-antimagic labelings of the generalized Petersen graph $P(n,(n-1)/2)$, *Util. Math.,* **70** (2006) 119-127.

[361] M. Bača, F. Bertault, J. MacDougall, M. Miller, R. Simanjuntak, and Slamin, Vertex-antimagic total labelings of graphs, *Discuss. Math. Graph Theory,* **23** (2003) 67-83. https://repository.unej.ac.id/bitstream/handle/123456789/813/Pages%20from%20DMGT_23_VAM_Graph_2003.pdf?sequence=1&isAllowed=y

[362] M. Bača, F. Bertault, J. MacDougall, M. Miller, R. Simanjuntak, and Slamin, Vertex-antimagic total labelings of (a,d)-antimagic and (a,d)-face antimagic graphs, preprint.NOT on Web 12/27/24

[363] M. Bača and L. Brankovic, Edge antimagicness for a class of disconnected graphs, *Ars Combin.*, **97A** (2010) 145-152. https://www.researchgate.net/publication/262693043_Edge-antimagicness_for_a_class_of_disconnected_graphs

[364] M. Bača, L. Brankovic, M. Lascsáková, O. Phanalasy, and A. Semaničová-Feňovčíková, On d-antimagic labelings of plane graphs, *Electronic J. Graph Theory Appl.*, **1**(1) (2013) 28-39. https://www.ejgta.org/index.php/ejgta/article/view/16/3

[365] M. Bača, L. Brankovic, and A. Semaničová-Feňovčikově, Labelings of plane graphs with determined face weights, *Acta Mechanica Slovaca*, **13**(2) (2009) 64-71.

[366] M. Bača, L. Brankovic, and A. Semaničová-Feňovčikově, Labelings of plane graphs containing Hamilton path, *Acta Math. Sinica,* **27**(4) (2011) 701-714. https://link.springer.com/article/10.1007/s10114-011-9451-x

[367] M. Bača, Dafik, M. Miller, and J. Ryan, Edge-antimagic total labeling of disjoint unions of caterpillars, *J. Combin. Math. Combin. Computing*, **65** (2008) 61-70. https://combinatorialpress.com/article/jcmcc/Volume%20065/vol-065-paper%206.pdf

[368] M. Bača, Dafik, M. Miller, and J. Ryan, Antimagic labeling of disjoint unions of s-crowns, *Util. Math.,* **79** (2009) 193-205.

[369] M. Bača and I. Holländer, Prime-magic labelings of $K_{n,n}$, *J. Franklin Inst.,* **327** (1990) 923-926. https://www.researchgate.net/profile/Martin-Baca/publication/245184672_Prime-magic_labelings_of_Knn/links/5a0b09830f7e9b0cc0251c13/Prime-magic-labelings-of-Kn-n.pdf

[370] M. Bača and I. Holländer, Labelings of a certain class of convex polytopes, *J. Franklin Inst.,* **329** (1992) 539-547. https://colab.ws/articles/10.1016%2F0016-0032%2892%2990052-i

[371] M. Bača and I. Holländer, On (a, d)-antimagic prisms, *Ars Combin.,* **48** (1998) 297-306. https://www.researchgate.net/publication/268251381_On_ad-antimagic_prisms

[372] M. Bača and I. Holländer, On (a, b)-consecutive Petersen graphs, *Acta Math. Appl. Sinica (English Ser.),* **14** (1998) 265-270. https://www.academia.edu/18622776/On_a_b_consecutive_petersen_graphs

[373] M. Bača, I. Holländer, and K. W. Lih, Two classes of super-magic quartic graphs, *J. Combin. Math. Combin. Comput.,* **23** (1997) 113-120. https://www.researchgate.net/publication/262723614_Two_classes_of_super-magic_graphs.

[374] M. Bača, M. Imran, and A. Semaničová-Feňovčcíková, Irregularity and modular irregularity strength of wheels, *Mathematics,* **9**(2710) (2021) 1-14. https://doi.org/10.1016/j.akcej.2018.01.013

[375] M. Bača, M. Irfan, A. Javad, and A. Semaničová-Feňočová, On total edge product cordial labeling of fullerenes, *Electron. J. Graph Theory Appl.,* (EJGTA) **6**(2) (2018) 238-249. http://www.ejgta.org/index.php/ejgta/article/view/395/pdf_79

[376] M. Bača, M. Irfan, J. Ryan, A. Semaničová-Feňovčíková, and D. Tanna, On edge irregular ´reflexive labellings for the generalized friendship graphs, *Mathematics*, **5**(4) (2017) 67. doi:10.3390/math5040067

[377] M. Bača, M. Irfan, J. Ryan, A. Semaničová-Feňovčíková, and D. Tanna, Note on edge irregular reflexive labelings of graphs, *AKCE Internat. J. Graphs Combin.*, **16**(2) (2019) 145-157. https://doi.org/10.1016/j.akcej.2018.01.013

[378] M. Bača, J. Jendrol, M. Miller, and J. Ryan, Antimagic labelings of generalized Petersen graphs that are plane, *Ars Combin.,* **73** (2004) 115-128.

[379] M. Bača, J. Jendrol, M. Miller, and J. Ryan, On irregular total labellings, *Discrete Math.,* **307** (2007) 1378-1388.

[380] M. Bača, P. Jeyanthi, P. Selvagopal, N. T. Muthu Raja, A. Semaničová-Feňovcıková, Ladders and fans are cycle-antimagic, *Hacettepe J. Math. Stat.*, **49**(3) (2020) 1093-1106.

[381] M. Bača, Z. Kimáková, M. Lascsáková, and A. Semaničová-Feňovčiková, The irregularity and modular irregularity strength of fan graphs, *Symmetry* **13**(605) (2021) 1-13. https://doi.org/10.3390/sym13040605

[382] M. Bača, Z. Kimáková, A. Semaničová-Feňovčikovă, and M. A. Umar, Tree-antimagicness of disconnected graphs, *Math. Probl. Eng.*, 2015, Art. ID 504251, 4 pp. https://onlinelibrary.wiley.com/doi/10.1155/2015/504251

[383] M. Bača, P. Kovář, T. Kovář, and A. Semaničová-Feňovčíková,On vertex in-out-antimagic total digraphs, *Disc. Math.,* (2023) 2023, 113758. https://doi.org/10.1016/j.disc.2023.113758

[384] M. Bača, P. Kovář, A. Semaničová, and M. K. Shafiq, On super $(a,1)$- edge-antimagic total labelings of regular graphs, *Discrete Math.,* **310** (2010) 1408-1412. https://core.ac.uk/reader/82056410

[385] M. Bača, M. Lascsḱová, M. Naseem, A. Semaničová-Feňovčíková, On entire face irregularity strength of disjoint union of plane graphs, *Appl. Math. Comput.,* **307** (2017) 232-238. https://www.researchgate.net/publication/315600510_On_entire_face_irregularity_strength_of_disjoint_union_of_plane_graphs

[386] M. Bača, M. Lascsáková, and A. Semaničová, On connection between α-labelings and edge-antimagic labeling of disconnected graphs, *Ars Combin.*, **106** (2012) 321-336. https://www.researchgate.net/publication/262692851_On_connection_between_a-labelings_and_edge-antimagic_labelings_of_disconnected_graphs

[387] M. Bača, M. Lascsáková, and A. Semaničová-Feňovčíková, Modular H-irregularity strength of graphs, *Mathematics*, **13**(16) (2025) 2599. https://www.researchgate.net/publication/395014870_Modular_H-Irregularity_Strength_of_Graphs

[388] M. Bača, M. Lascsáková and M. K. Siddiqui, Total edge irregularity strength of toroidal fullerene, *Math. Comput. Sci.*, **7**, (2013) 487-492. https://link.springer.com/article/10.1007/s11786-014-0172-2

[389] M. Bača, Y. Lin, and M. Miller, Valuations of plane quartic graphs, *J. Combin. Math. Combin. Computing*, **41** (2002) 209-221. https://www.researchgate.net/publication/262723100_Valuations_of_plane_quartic_graphs

[390] M. Bača, Y. Lin, and M. Miller, Antimagic labelings of grids, *Util. Math.,* **72** (2007) 65-75. https://www.researchgate.net/publication/262722982_Antimagic_labelings_of_grids

[391] M. Bača, Y. Lin, M. Miller, and J. Ryan, Antimagic labelings of Möbius grids, *Ars Combin.,* **78** (2006) 3-13. https://www.researchgate.net/profile/Martin-Baca/publication/220620291_Antimagic_labelings_of_Msbius_grids/links/0c960538da988ba8c2000000/Antimagic-labelings-of-Msbius-grids.pdf

[392] M. Bača, Y. Lin, M. Miller, and R. Simanjuntak, New constructions of magic and antimagic graph labelings, *Util. Math.,* **60** (2001) 229-239. https://www.researchgate.net/publication/262723106_New_constructions_of_magic_and_antimagic_graph_labelings

[393] M. Bača, Y. Lin, M. Miller, and M. Z. Youssef, Edge-antimagic graphs, *Discrete Math.,* **307** (2007) 1232-1244. https://core.ac.uk/download/pdf/82480238.pdf

[394] M. Bača, Y. Lin, and F. A. Muntaner-Batle, Super edge-antimagic of path-like trees, *Util. Math.,* **73** (2007) 117-128. https://www.researchgate.net/publication/262723240_Super_edge-antimagic_labelings_of_the_path-like_trees

[395] M. Bača, Y. Lin, and F. A. Muntaner-Batle, A note on even disjoint union of paths, *AKCE Int. J. Graphs Comb.*, **6**(1) (2009) 41-46. https://www.researchgate.net/publication/262698010_A_note_on_even_disjoint_union_of_paths

[396] M. Bača, Y. Lin, F. Muntaner-Batle, and M. Rius-Font, Strong labeling of linear forests, *Acta Math. Sinica, English Series*, **25** (2009) 1951-1964. https://www.researchgate.net/publication/225323894_Strong_labelings_of_linear_forests

[397] M. Bača, Y. Lin, and F. Muntaner-Batle, Edge-antimagic labeling of forests, *Util. Math.*, **81** (2010) 31-40. https://www.researchgate.net/publication/262697772_Edge-antimagic_labelings_of_forest

[398] M. Bača, J. MacDougall, M. Miller, Slamin, and W. Wallis, Survey of certain valuations of graphs, *Discussiones Mathematicae Graph Theory,* **20** (2000) 219-229. https://www.researchgate.net/publication/264312004_A_Survey_of_Certain_Valuations_of_Graphs

[399] M. Bača, Y.Lin, and A. Semaničová-Feňovčíková, Note on super antimagicness of disconnected graphs, *AKCE J. Graphs Combin.,* **6** (2009) 47-55. https://www.tandfonline.com/doi/epdf/10.1080/09728600.2009.12088871?needAccess=true

[400] M. Bača and M. Miller, Antimagic face labeling of convex polytopes based on biprisms, *J. Combin. Math. Combin. Comput.,* **36** (2001) 229-236. https://www.researchgate.net/publication/262723422_Antimagic_face_labeling_of_convex_polytopes_based_on_biprisms

[401] M. Bača and M. Miller, Valuations of a certain class of convex polytopes, *J. Combin. Math. Combin. Comput.,* **43** (2002) 207-218. https://www.researchgate.net/publication/262723286_Valuations_of_a_certain_class_of_convex_polytopes

[402] M. Bača and M. Miller, On d-antimagic labelings of type $(1, 1, 1)$ for prisms, *J. Combin. Math. Combin. Comput.,* **44** (2003) 199-207. https://combinatorialpress.com/article/jcmcc/Volume%20044/vol-044-paper%2018.pdf

[403] M. Bača and M. Miller, Super Edge-Antimagic Graphs: A Wealth of Problems and Some Solutions, BrownWalker Press, 2007, Boca Raton, FL, USA

[404] M. Bača, M. Miller, O. Phanalasy, J. Ryan, A. Semaničová-Feňovčíková, and A. Abildgaard Sillasen, Totally antimagic total graphs, *Australas. J. Combin.,* **61** (2015) 42-56. https://www.academia.edu/es/66812029/Totally_antimagic_total_graphs

[405] M. Bača, M. Miller, O. Phanalasy, and A. Semaničová-Feňovčíková, Super d-antimagic labelings of disconnected plane graphs, *Acta Math. Sin. (Engl. Ser.),* **26** (2010) 2283-2294. https://link.springer.com/article/10.1007/s10114-010-9502-8

[406] M. Bača, M. Miller, O. Phanalasy, and A. Semaničová-Feňovčíková, Constructions of antimagic labelings for some families of regular graphs, *J. Algor. Computation*, **44**, (2013) 1-7. https://www.researchgate.net/publication/262692172_Constructions_of_antimagic_labelings_for_some_families_of_regular_graphs

[407] M. Bača, M. Miller, O. Phanalasy and A. Semaničová-Feňočová, Antimagic labeling of join graphs,*Math. Comput. Sci.,* **9**(2) (2015) 139-143. https://www.researchgate.net/publication/276424030_Antimagic_Labelings_of_Join_Graphs

[408] M. Bača, M. Miller, and J. Ryan, On d-antimagic labelings of prisms and antiprisms, *Proceedings 12th Australian Workshop on Combinatorial Algorithms,* Bandung, Indonezia (2001) 51-58. https://www.researchgate.net/publication/220620227_On_d-antimagic_labelings_of_prism

[409] M. Bača, M. Miller, J. Ryan, and A. Semaničová-Feňovčíková, On H-antimagicness of disconnected graphs, *Bull. Aust. Math. Soc.,* **94**(92) (2016) 201-207. https://www.cambridge.org/core/services/aop-cambridge-core/content/view/D93AA8E8B8A9D1F6C6570E46C93DB9E0/S0004972716000204a.pdf/on-dollarhdollar-antimagicness-of-disconnected-graphs.pdf

[410] M. Bača, M. Miller, J. Ryan, and A. Semaničová-Feňovčíková, Edge-magic total labelings. In: Magic and Antimagic Graphs. Developments in Mathematics, **60** (2-19). Springer, Cham.

[411] M. Bača, M. Miller, J. Ryan, and A. Semaničová-Feňovčíková, Magic and Antimagic Graphs Attributes, Observations, and Challenges in Graph Labelings, *60* Springer Nature Switzerland AG 2019

[412] M. Bača, M. Miller, and Slamin, Vertex-magic total labeling of generalized Petersen graphs, *Int. J. Comput. Math.,* **79** (2002) 1259-1263. https://www.tandfonline.com/doi/epdf/10.1080/00207160214650?needAccess=true

[413] M. Bača, F. Muntaner-Batle, A. Semaničová-Feňovčíková, and M. K. Shafiq, On super $(a, 2)$-antimagic total labeling of disconnected graphs, *Ars Combin.*, **113**, (2014) 129-137. https://www.researchgate.net/publication/262685288_On_super_a2-edge-antimagic_total_labeling_of_disconnected_graphs

[414] M. Bača, K. Muthugurupackiam, K. M. Kathiresan, and S. Ramya, Modular irregularity strength of graphs, *Electron. J. Graph Theory Appl.* **8** (2020) 435-443. https://www.researchgate.net/publication/346505290_Modular_irregularit_strength_of_graphs

[415] M. Bača, M. Numan, and A. Semaničová-Feňovčíková, Super d-antimagic labelings of generalized prism, *Util. Math.,* **99** (2016) 101-119. https://www.researchgate.net/profile/Martin-Baca/publication/301776062_Super_d-antimagic_labelings_of_generalized_prism/links/5732c69608ae9f741b2361db/Super-d-antimagic-labelings-of-generalized-prism.pdf

[416] M. Bača, M. Numan, and A. Semaničová-Feňovčíková, Super d-antimagic labelings of disjoint union of generalized prisms, *Util. Math.,* **103** (2017) 299-310. https://www.researchgate.net/publication/318124883_Super_d-antimagic_labelings_of_disjoint_union_of_generalized_prisms

[417] M. Bača, M. Numan, and A. Shabbir, Labelings of type $(1, 1, 1)$ for toroidal fullerenes, *Turkish J. Math.*, **37** (2013) 899-907. https://journals.tubitak.gov.tr/cgi/viewcontent.cgi?article=2275&context=math

[418] M. Bača, M. Numan, and M. K. Siddiqui, Super face antimagic labelings of union of antiprisms, *Math. Comput. Sci.*, **7**(2) (2013) 245-253. https://www.researchgate.net/publication/257759497_Super_Face_Antimagic_Labelings_of_Union_of_Antiprisms

[419] M. Bača and M. Murugan, Super edge-antimagic labelings of cycle with a chord, *Australas. J. Combin.,* **35** (2006) 253-261.

[420] M. Bača, A. Ovais, A. Semaničová-Feňovčíková, and I. Nengah Suparta, On face irregular evaluations of plane graphs, *Discuss. Math. Graph Th.*, **42**(2) (2022) 549-568. https://doi.org/10.7151/dmgt.2294

[421] M. Bača, A. Semaničová-Feňovčíková, R.-T. Lai, and T.-M. Wang, On local antimagic vertex coloring for complete full t-ary trees, *Fundamenta Informaticae*, **XXI** (2020) 1001-1015. https://www.researchgate.net/publication/359891015_On_local_antimagic_vertex_coloring_for_complete_full_t-ary_trees

[422] M. Bača, A. Semaničová-Feňovčíková, and M. K. Shafiq, A method to generate large classes of edge-antimagic trees, *Util. Math.,* **86** (2011) 33-43.

https://www.researchgate.net/publication/262692858_A_method_to_generate_large_classes_of_edge-antimagic_trees

[423] M. Bača, A. Semaničová-Feňovčíková, S. Slamin, and K. A. Sugeng, On inclusive distance vertex irregular labelings, *Electron. J. Graph Theory Appl.,* **6**(1) (2018) 61-83. https://www.ejgta.org/index.php/ejgta/article/view/463/pdf_65

[424] M. Bača, A. Semaničová-Feňovčikově, M. A. Umar, and D. Welyyanti, On H-antimagicness of Cartesian product of graphs, *Turkish J. Math.,* **42**(1) (2018) 339-348.

[425] M. Bača, A. Semaničová-Feňovčíková, T. M. Wang, and G. H. Zhang, On $(a, 1)$-vertex-antimagic edge labeling of regular graphs, *J. Appl. Math.,* (2015), Art. ID 320616, 7 pp. https://www.researchgate.net/publication/277963549_On_1-Vertex-Antimagic_Edge_Labeling_of_Regular_Graphs

[426] M. Bača, and A. Shabbir, Total labelings of toroidal polyhexes, *Science Internat.*, **24**(3,) (2012) 239-241. https://www.researchgate.net/publication/262692538_Total_labelings_of_toroidal_polyhexes

[427] M. Bača, and M. K. Siddiqui, Total edge irregularity strength of generalized prism, *Applied Math. Comput.*, **235** (2014) 168-173. https://www.researchgate.net/publication/261102860_Total_edge_irregularity_strength_of_generalized_prism

[428] M. Bača and M. K. Siddiqui, On total edge irregularity strength of strong product of two cycles, *Util. Math.,* **104** (2017) 255-275. https://www.scientificbulletin.upb.ro/rev_docs_arhiva/full8d8_693905.pdf

[429] M. Bača and M. Z. Youssef, Further results on antimagic graph labeling, *Australas. J. Combin.,* **38** (2007) 163-172. https://ajc.maths.uq.edu.au/pdf/38/ajc_v38_p163.pdf

[430] M. Bača and M. Z. Youssef, On harmonious labelings of corona graphs, *J. Appl. Math.*, **2014** Art. ID 627248, 4 pp. https://onlinelibrary.wiley.com/doi/pdf/10.1155/2014/627248

[431] E. M. Badr, On graceful labeling of the generalization of cyclic snakes, *J. Discrete Math. Sci. Cryptogr.,* **18**(6) (2015) 773-783. https://www.researchgate.net/publication/273418550_On_Graceful_Labeling_of_the_Generalization_of_Cyclic_Snakes

[432] E. M. Badr and M. I. Moussa, An upper bound of radio k-coloring problem and its integer linear programming model, *Wireless Networks*, **26**(7) (2020) 4955-4964. https://www.researchgate.net/publication/331842382_An_upper_bound_of_radio_k-coloring_problem_and_its_integer_linear_programming_model

[433] E. M. Badr, M. I. Moussa, and K. Kathiresan, Crown graphs and subdivision of ladders are odd graceful, *Internat. J. Computer Math.,* **88**(17) (2011) 3570-3576. DOI:10.1080/00207160.2011.610893

[434] E. M. Badr, S. Nada, M. Al-Shamiri, A. Abdel-Hay, and A. Elrokh, A novel mathematical model for radio mean square labeling problem, *J. Math.*, 2022, Art. ID 3303433, 9 pp. https://doi.org/10.1155/2022/3303433

[435] E.-S. Badr, S. Alaybani, A. Elrokh, A. A. El-hay, and B. Almutairi, An integer linear programming model for solving radio mean labeling problem, *IEEEAcess* **8** 2020 162343-16234. https://www.researchgate.net/publication/344443208_An_Integer_Linear_Programming_Model_for_Solving_Radio_Mean_Labeling_Problem

[436] J. Bagga, L. P. Fotso, P. B. Max, and S. Arumugam, New classes of graceful unicyclic graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 27-32. https://colab.ws/articles/10.1016%2Fj.endm.2015.05.005

[437] J. Bagga and A. Heinz, Graceful labelings of unicyclic graphs. Advances in discrete mathematics and applications: Mysore, 2008, 69–79, *Ramanujan Math. Soc. Lect. Notes Ser.*, **13**, Ramanujan Math. Soc., Mysore, 2010.

[438] P. Bahl, S. Lake, and A. Wertheim, Gracefulness of families of spiders, *Involve*, **3** (2010) 241-247. https://msp.org/involve/2010/3-3/involve-v3-n3-p01-s.pdf

[439] D. R. Bai and J. Senbagamalar, Cubic roots cordial labeling of sunlet and friendship graphs, In book: Proceed. 4th Internat. Conf. Math. Modeling Comput. Sci., (2025) 107-115.

[440] A. Q. Baig, A. Ahmad, E. Baskoro, and R. Simanjuntak, On the super edge-magic deficiency of forests, *Util. Math.,* **86** (2011) 147-159. https://www.researchgate.net/publication/265534281_On_the_super_edge-magic_deficiency_of_forests

[441] A. Q. Baig, E. Baskoro, and A. Semaničová-Feňovčiková, On the super edge-magic deficiency of a star forest, *Ars Combin.,* **115** (2014) 3-12. https://www.researchgate.net/publication/287059730_On_the_super_edge-magic_deficiency_of_a_star_forest

[442] A. Q. Baig, M. Imran, I. Javaid, and A. Semaničová-Feňovčiková, Super edge-magic deficiency of graphs, *Util.Math.,* **87** (2012) 355-364. https://www.researchgate.net/publication/264554566_Super_edge-magic_deficiency_of_graphs

[443] A. Q. Baig and H. U. Afzal, Some super edge-magic (p, q)-graphs with magic constant $3p$, *Util. Math.,* **98** (2015) 53-63. https://www.researchgate.net/publication/265466409_Super_Edge_Magic_p_q_Graphs_with_maximum_magic_constant_3p

[444] A. Q. Baig, H. U. Afzal, M. Imran, and I. Javaid, Super edge-magic labeling of volvox and pancyclic graphs, *Util. Math.,* **93** (2014) 49-56. https://www.researchgate.net/publication/264554381_Super_edge-magic_labeling_of_volvox_and_pancyclic_graph

[445] A. Q. Baig, E. T. Baskoro, and A. Semaničová-Feňovčíková, On the super edge-magic deficiency of a star forest, *Ars Combin.,* **115** (2014) 3-12. https://www.researchgate.net/profile/A-Baig/publication/287059730_On_the_super_edge-magic_deficiency_of_a_star_forest/links/57c6f5d708ae9d64047e92ea/On-the-super-edge-magic-deficiency-of-a-star-forest.pdf

[446] S. Bailey and C. Barrientos, Operations with mean graphs, *Congr. Numer.*, **217** (2013) 5-19.

[447] V. Balaji, D. S. T. Ramesh, and A. Subramanian, Some results on relaxed Skolem mean graphs, *Bull. Kerala Math. Assoc.,* **5**(2) (2009) 33-44.

[448] V. Balaji, D. S. T. Ramesh and A. Subramanian, Relaxed Skolem mean labeling, *Adv. Appl. Disc. Math.,* **5**(1) (2010) 11-22.

[449] S. Balamoorthy, S. V. Bharanedhar, and N. Kamatchi, On the products of group vertex magic graphs, *AKCE Int. J. Graphs Comb.*, **19**(3) (2022) 268-275. https://www.tandfonline.com/doi/full/10.1080/09728600.2022.2136021

[450] S. Bala, S. Sundarraj, and K. Thirusangu, Cordial labelings for comb related graph, *Bull. Pure Appl. Sci. Sect. E Math. Stat.,* 38E (Special Issue) (2S) (2019) 92-98. https://bpasjournals.com/admin/upload/dynamic2/92-98-bpas-e-dec.2019-special%20issue.pdf

[451] S. Bala, V. Krithika Devi, and K. Thirusangu, Some cordial labelings on square graph of a comb, *Bull. Pure Appl. Sci. Sect. E Math. Stat.,* **38E** (Special Issue)(2S) (2019) 112-118. https://bpasjournals.com/admin/upload/dynamic2/112-118-bpas-e-dec.2019-special%20issue.pdf

[452] S. Bala, S. Sundarraj, and K. Thirusangu, Cordial labelings for comb related graph, *Bull. Pure Appl. Sci. Sect. E Math. Stat.,* 38E (Special Issue) (2S) (2019) 92-98 (2019) https://bpasjournals.com/admin/upload/dynamic2/92-98-bpas-e-dec.2019-special%20issue.pdf

[453] V. Balaji and V. Maheswari, Some results on relaxed mean labeling, *Math. Combin.*, **3** (2015) 73-80. http://fs.unm.edu/IJMC/SomeResultsOnRelaxedMeanLabeling.pdf

[454] V. Balaji, D. S. T. Ramesh, and A. Subramanian, Some results on relaxed mean graphs, *Internat. J. Math. Combin.,* **3** (2015) 73-80. https://fs.unm.edu/IJMC/SomeResultsOnRelaxedMeanLabeling.pdf

[455] V. Balajia, D. S. T. Ramesh, and S. Sudhakar, Further results on relaxed mean labeling, *Int. J. Adv. Appl. Math. and Mech.,* **3**(3) (2016) 92-99. http://www.ijaamm.com/uploads/2/1/4/8/21481830/v3n3p8_92-99.pdf

[456] V. Balaji, Solution of a conjecture on Skolem mean graph of stars $K_{1,l} \cup K_{1,m} \cup K_{1,n}$, *Internat. J. Math. Combin.,* **4** (2011) 115-117. https://fs.unm.edu/IJMC/SolutionOfAConjecture.pdf

[457] V. Balaji, D. S. T. Ramesh and A. Subramanian, Skolem mean labeling, *Bull. Pure and Applied Sci.,* **26E** (2) (2007) 245-248.

[458] V. Balaji, D. S. T. Ramesh and A. Subramanian, Some results on Skolem mean graphs, *Bull. Pure and Applied Sci.*, **27E** (1)(2008) 67-74.

[459] R. Balakrishnan, Graph labelings, unpublished.

[460] B. J. Balamurugan, K. Thirusangu, and D. G. Thomas, Zumkeller labeling for twig graphs, *Electron. Notes Disc. Math.*, **48** (2015) 119-126. https://www.sciencedirect.com/science/article/pii/S1571065315000189

[461] B. J. Balamurugan, K. Thirusangu, and D. G. Thomas, B. J. Murali, k-Zumkeller labeling of graphs, *Int. J. Eng. Technol.*, **7**(4) (2018) 460-463. https://www.sciencepubco.com/index.php/ijet/article/view/21040

[462] C. Balbuena, E. Barker, K. C. Das, Y. Lin, M. Miller, J. Ryan, Slamin, K. Sugeng, and M. Tkac, On the degrees of a strongly vertex-magic graph, *Discrete Math.,* **306** (2006) 539-551. https://www.academia.edu/87364855/On_the_degrees_of_a_strongly_vertex_magic_graph

[463] R. Balakrishnan and R. Kumar, Decomposition of complete graphs into isomorphic bipartite subgraphs, *Graphs and Combin.,* **10** (1994) 19-25. https://link.springer.com/article/10.1007/BF01202466

[464] R. Balakrishnan and R. Kumar, Existence and nonexistence of certain labellings for the graph $K_n^c \bigvee 2K_2$, *Util. Math.*, **46** (1994) 97-102.

[465] R. Balakrishnan and R. Sampathkumar, Decompositions of regular graphs into $K_n^c \vee 2K_2$, *Discrete Math.*, **156** (1996) 19-28. https://www.sciencedirect.com/science/article/pii/0012365X9400026F

[466] R. Balakrishnan, A. Selvam, and V. Yegnanarayanan, On felicitous labelings of graphs, *Proceed. National Workshop on Graph Theory and its Appl. Manonmaniam Sundaranar Univ., Tiruneli* (1996) 47-61.

[467] R. Balakrishnan, A. Selvam, and V. Yegnanarayanan, Some results on elegant graphs, *Indian J. Pure Appl. Math.*, **28** (1997) 905-916.

[468] C. Balbuena, E. Barker, Y. Lin, M. Miller, and K. Sugeng, Consecutive magic graphs, *Discrete Math.,* **306** (2006) 1817-1829.

[469] C. Balbuena, P. García-Vázquez, X. Marcote, and J. C. Valenzuela, Trees having an even or quasi even degree sequence are graceful, *Applied Math. Letters,* **20** (2007) 370-375. https://www.researchgate.net/publication/220317839_Trees_having_an_even_or_quasi_even_degree_sequence_are_graceful

[470] D. Bange and A. E. Barkauskas, Fibonacci graceful graphs, *Fibonacci Quarterly*, **71**(3) (1983) 174-188. https://www.fq.math.ca/Scanned/21-3/bange.pdf

[471] D. Bange, A. Barkauskas, and P. Slater, Simply sequential and graceful graphs, *Proc. of the 10th S.E. Conf. on Combinat., Graph Theory, and Computing*, (Util. Math. Winnipeg, 1979) 155-162.

[472] D. Bange, A. Barkauskas, and P. Slater, Sequentially additive graphs, *Discrete Math.*, **44** (1983) 235-241. https://www.sciencedirect.com/science/article/pii/0012365X83901875?via%3Dihub

[473] J. R. Bansari, A. G. Jekil, and A. S. Mahadityasinh, Cordial labeling of Goldberg snark and its related graphs, https://doi.org/10.48550/arXiv.2511.08664

[474] D. Bantva, Further results on the radio number of trees, *Elect. Notes Discr. Math.,* **63** (2017) 85-91. https://www.sciencedirect.com/science/article/pii/S1571065317303049

[475] D. Bantva, Radio number for middle graph of paths, *Elect. Notes Discr. Math.,* **63** (2017) 93-100. https://www.semanticscholar.org/reader/dd30df1c0937bca9030a0be0bce8153dd72c98d9

[476] D. Bantva, A lower bound for the radio number of graphs, *Algorithms and Discrete Applied Mathematics*. CALDAM 2019. Lecture Notes in Computer Science, vol 11394, 161-173. Springer, Cham https://www.researchgate.net/publication/330490931_A_Lower_Bound_for_the_Radio_Number_of_Graphs_5th_International_Conference_CALDAM_2019_Kharagpur_India_February_14-16_2019_Proceedings

[477] D. Bantva, Optimal radio labelings of graphs, arXiv:2212.13578v1

[478] D. Bantva and D. Der-Fen Liu, Radio number for the cartesian product of two trees, arXiv:2202.13983

[479] D. Bantva, S. Vaidya, and S. Zhou, Radio number of trees, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 135-141. https://www.semanticscholar.org/reader/4531ff12ec3aee0bfc3497620fae94971f91a712

[480] J. Bao, L. Zhao, Y. Yang, W. Feng, and Jirimutu, The generalized Petersen graph $P(n, 7)$ is $(\frac{3n+6}{2}, 3)$-antimagic, *Util. Math.,* **100** (2016) 33-41.

[481] M. V. Bapat, Some vertex prime graphs and a new type of graph labeling, *Internat. J. Math. Trends and Tech.,* **47**(1) (2017) 49-55. https://ijmttjournal.org/public/assets/volume-47/number-1/IJMTT-V47P507.pdf

[482] M. V. Bapat, Some complete-graph related families of product cordial (pc) graphs, *Aryabhatta J. Math. Informatics*, **09**(02) (2017) 133-140. https://www.academia.edu/38688975/Some_Complete_Graph_Related_Families_Of_Product_Cordial_pc_Graphs

[483] M. V. Bapat, Product cordial labeling of some fusion graphs, *Internat. J. Math. Trends and Tech.,* **50**(2) (2017) 125-129. https://ijmttjournal.org/public/assets/volume-50/number-2/IJMTT-V50P519.pdf

[484] M. V. Bapat, Extended vertex edge additive cordial (EVEAC) labeling-a new labeling method for graphs, *Internat. J. Stat. Appl. Math.,* **2**(6) (2017) 92-94. https://www.mathsjournal.com/pdf/2018/vol3issue1/PartD/2-6-27-690.pdf

[485] M. V. Bapat, Extended edge vertex cordial labelings of graph, *Internat. J. Math. Archive,* **8**(9) (2017) 17-19. https://www.researchgate.net/publication/342815445_EXTENDED_EDGE_VERTEX_CORDIAL_LABELING_OF_GRAPH

[486] M. V. Bapat, A note on L-Cordial labeling Of graphs, *Internat. J. Math. Appl.*, **5**(4-D) (2017) 457-460.

[487] M. V. Bapat, New results on L-cordial labeling of graphs, *Internat. J. Stat. Appl. Math.*, **3**(2) (2018) 641-644. https://www.mathsjournal.com/pdf/2018/vol3issue2/PartH/3-2-44-901.pdf

[488] M. V. Bapat and N. B. Limaye, Some families of 3-equitable graphs, *J. Combin. Math. Combin. Comput.,* **48** (2004) 179-196. https://combinatorialpress.com/jcmcc-articles/volume-048/some-families-of-3-equitable-graphs/

[489] M. V. Bapat and N. B. Limaye, A note on 3-equitable labelings of multiple shells, *J. Combin. Math. Combin.,* Comput. **51** (2004) 191-202. https://combinatorialpress.com/article/jcmcc/Volume%20051/vol-051-paper%2013.pdf

[490] M. V. Bapat and N. B. Limaye, Edge-three cordial graphs arising from complete graphs, *J. Combin. Math. Combin. Comput.*, **56** (2006) 147-169. https://combinatorialpress.com/article/jcmcc/Volume%20056/vol-056-paper%2010.pdf

[491] M. V. Bapat and N. B. Limaye, E_3-cordiality of some helm-related graphs, *Ars Combin.,* **119** (2015) 429-443.

[492] C. M. Barasara, Edge and total edge product cordial labeling of some new graphs, *Internat. J. Engineering, Sci. Math.*, **7**(2) (2018) 263-273. https://ijesm.co.in/uploads/68/4847_pdf.pdf

[493] C. M. Barasara, Edge product cordial and total edge product cordial labeling of some wheel related graphs. In: N. P. Shrimali N. H. Shah (eds.), *Recent Adv. Graph Th.*, (1st ed.), CRC Press, 2020.

[494] C. M. Barasara, Edge and total edge product cordial labeling of some new graphs, *Internat. Engin., Sci. Math.*, **7**(2) (2018) 263-273. https://www.ijesm.co.in/uploads/68/4847_pdf.pdf

[495] C. M. Barasara and Y. B. Thakkar, Divisor cordial labeling of graphs in the context of some graph operations, *Adv. Appl. Discrete Math.*, **28**(1) (2021) 31-48. https://research.ebsco.com/c/x2uggr/viewer/pdf/ixooagtt5f?route=details

[496] C. M. Barasara and P. J. Prajapati, Prime cordial labeling for some path, cycle and wheel graphs, *Adv. Appl. Discr. Math.*, **30** (2022) 35-58. https://www.mililink.com/upload/article/25925498aams_vol_217_may_2022_a1_p3577-3594_c._m._barasara_and_y._b._thakkar.pdf

[497] C. M. Barasara and P. J. Prajapati, Antimagic labeling for some snake graphs, *Proyecciones J. Math.,* **43**(2) (2024) 521-537. https://www.researchgate.net/publication/379668848_Antimagic_labeling_for_some_snake_graphs

[498] C. M. Barasara and Y. B. Thakkar, Divisor cordial labeling for ladders and total graph of some graphs, *Adv. Appl. Math. Sci.*, **21**(7) (2022) 3577-3594. https://www.mililink.com/upload/article/25925498aams_vol_217_may_2022_a1_p3577-3594_c._m._barasara_and_y._b._thakkar.pdf

[499] C. M. Barasara and Y. B. Thakkar, Divisor cordial labeling for some cycle and wheel related graphs, *Malaya J. Mat.*, **8**(3) (2020) 966-972. http://malayajournal.org/articles/MJM08030041.pdf

[500] C. M. Barasara and Y. B. Thakkar, Divisor cordial labeling for some snakes and degree splitting related graphs, *South East Asian J. Math. Math. Sci.*, **19(1)** (2023) 211-224. https://rsmams.org/journals/articleinfo.php?articleid=826&tag=seajmams

[501] C. M. Barasara and Y. B. Thakkar, Further results on divisor cordial graph, *South East Asian J. Math. Math. Sci.*, **20** (Proceedings) (2022) 115-138. https://rsmams.org/download/proceedings/2_20_0_1122293033_FURTHER%20RESULTS%20ON.pdf

[502] C. M. Barasara and Y. B. Thakkar, Some new results on power cordial labeling, *J. Appl. Math. Informatics*, **41**(3) (2023) 615-631. https://www.researchgate.net/publication/374947066_FURTHER_RESULTS_ON_DIVISOR_CORDIAL_GRAPH

[503] C. M. Barasara and Y. B. Thakkar, Some characterizations and NP-complete problems for power cordial graphs, *J. Math.*, **2023**(Article ID 2257492) (2023) 5 pages. https://doi.org/10.1155/2023/2257492

[504] C. M. Barasara and Y. B. Thakkar, Characterizations for some diviso cordial type labelings, Conference: National Conference on Advancement in Mathematical Research and Development - 2024 At: Pilvai, Gujarat, INDIA. https://www.researchgate.net/publication/379511650_CHARACTERIZATIONS_FOR_SOME_DIVISOR_CORDIAL_TYPE_LABELINGS

[505] C. M. Barasara and P. J. Prajapati, Antimagic labeling of some degree splitting graphs, *Ratio Math.*, **48** (2023) 12 pages. http://eiris.it/ojs/index.php/ratiomathematica/article/view/1253

[506] C. M. Barasara, S. K. Vaidya, and P. L. Vihol, 3-equitable labeling in the context of some graph operations, *J. Applied Computer Science & Math.,* **11**(5) (2011) 69-75.

[507] C. Barrientos, New families of equitable graphs, *Util. Math.,* **60** (2001) 123-137. https://www.researchgate.net/publication/266424539_New_families_of_equitable_graphs

[508] C. Barrientos, Graceful labelings of cyclic snakes, *Ars Combin.,* **60** (2001) 85-96. https://www.arxiv.org/pdf/2012.10341v2

[509] C. Barrientos, Equitable labelings of corona graphs, *J. Combin. Math. Combin. Comput.,* **41** (2002) 139-149. https://combinatorialpress.com/article/jcmcc/Volume%20041/vol-041-paper%2010.pdf

[510] C. Barrientos, Graceful labelings of chain and corona graphs, *Bull. Inst. Combin. Appl.,* **34** (2002) 17-26.

[511] C. Barrientos, Difference Vertex Labelings, Ph.D. Thesis, Universitat Politécnica de Catalunya, Barcelona, 2004. Notb on web 1/12/25

[512] C. Barrientos, The gracefulness of unions of cycles and complete bipartite graphs, *J. Combin. Math. Combin. Comput.* **52** (2005) 69-78. https://combinatorialpress.com/article/jcmcc/Volume%20052/vol-052-paper%206.pdf

[513] C. Barrientos, Graceful graphs with pendant edges, *Australas. J. Combin.* **33** (2005) 99-107. https://ajc.maths.uq.edu.au/pdf/33/ajc_v33_p099.pdf

[514] C. Barrientos, Graceful arbitrary supersubdivisions of graphs, *Indian J. Pure Appl. Math.,* **38** (2007) 445-450. https://paperzz.com/doc/7700166/graceful-arbitrary-supersubdivisions-of-graphs?utm_source=chatgpt.com

[515] C. Barrientos, Odd-graceful labelings of trees of diameter 5, *AKCE Internat. J. Graphs and Combin.*, **6** (2009) 307–313. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088894?utm_source=chatgpt.com

[516] C. Barrientos, Unicylic graceful graphs, unpublished.

[517] C. Barrientos, On graceful chain graphs, *Util. Math.,* **78** (2009) 55-64.

[518] C. Barrientos, Odd-graceful labelings of trees of diameter 5, *AKCE Int. J. Graphs Comb.,* **6** (2009) 307-313. arXiv:0807.3738

[519] C. Barrientos, Alpha graphs with different pendent paths, *Elect. J. Graph Th. Appl.,* (EJGTA) **8**(2) (2020) 301-317. https://www.ejgta.org/index.php/ejgta/article/view/1036/pdf_143

[520] C. Barrientos, On additive vertex labelings, *Indonesian J. Combin.,* **4**(1) (2020) 34-52. https://www.ijc.or.id/index.php/ijc/article/view/121/83

[521] C. Barrientos, On the generation of alpha graphs, *J. Algebra Combin. Discrete Struct. Appl.*, **9**(2) (2022) 101-114. https://doi.org/10.13069/jacodesmath.1111733

[522] C. Barrientos, Mean trees, *Bull. Inst. Combin. Appl.,* **75** (2015) 8-18.

[523] C. Barrientos, M. E. Abdel-Aal, S. Minion, and D. Williams, The mean labeling of some crowns, *J. Algorithms and Comput.*, **45** (2014) 43-54. https://jac.ut.ac.ir/article_7920_54ecbd311317abd1ff2dc3518ec27a83.pdf

[524] C. Barrientos and S. Barrientos, On graceful supersubdivisions of graphs, *Bull. Inst. Combin. Appl.,* **70** (2014) 77-85.

[525] C. Barrientos, I. Dejter, and H. Hevia, Equitable labelings of forests, *Combin. and Graph Theory*, **1** (1995) 1-26.

[526] C. Barrientos and H. Hevia, On 2-equitable labelings of graphs, *Notas de la Sociedad de Matemática de Chile,* **XV** (1996) 97-110.

[527] C. Barrientos and E. Krop, Mean graphs, *AKCE Int. J. Graphs Comb.,* **11**(1) (2014) 13-26. https://www.researchgate.net/publication/268640068_Mean_graphs

[528] C. Barrientos and E. Krop, Improved bounds for relaxed graceful trees, *Graphs Combin.,* **33** (2017) 287-305. arXiv:1402.0196

[529] C. Barrientos and S. Minion, Alpha labelings of snake polyominoes and hexagonal chains, *Bull. Inst. Combin. Appl.,* **74** (2015) 73-83.

[530] C. Barrientos and S. Minion, Three graceful operations, *J. Algorithms and Comput.,* **45** (2014) 13-24.

[531] C. Barrientos and S. Minion, Enumerating families of labeled graphs, *J. Integer Sequences,* **18** (2015) Article 15.1.7, 14 pages. https://cs.uwaterloo.ca/journals/JIS/VOL18/Minion/minion3.pdf

[532] C. Barrientos and S. Minion, New attack on Kotzig's Conjecture, *Electron. J. Graph Theory Appl.*, **4**(2) (2016) 119-131.

[533] C. Barrientos and S. Minion, On the graceful Cartesian product of alpha-trees, *Theory Appl. Graphs,* **4**(1) (2017) Art. 3, 12 pp. https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1030&context=tag

[534] C. Barrientos and S. Minion, On the number of α-labeled graphs, *Discuss. Math. Graph Th.,* **38**(1) (2018) 177-188. https://www.researchgate.net/publication/317253638_On_the_Number_of_a-Labeled_Graphs

[535] C. Barrientos and S. Minion, Constructing graceful graphs with caterpillars, *J. Algor. Computation,* **48** (2016) 117-125.

[536] C. Barrientos and S. Minion, New α-trees and graceful unions of α-graphs and linear forests, *J. Combin. Math. Combin. Comput.,* **108** (2019) 205-220.

[537] C. Barrientos and S. Minion, Snakes: from graceful to harmonious, *Bull. Institute Combin. Appl.,* CA **79** (2017) 95-107. https://bica.the-ica.org/Volumes/79//Reprints/LBICA1Final-Reprint.pdf

[538] C. Barrientos and S. Minion, Snakes and caterpillars in graceful graphs, *J. Algorithms and Comput.*, **50** (2018) 37-47.

[539] C. Barrientos and S. Minion, Folding trees gracefully, *AKCE Int. J. Graphs Comb.*, **17**(3) (2020) 796-800. https://www.mobt3ath.com/uplode/book/book-110037.pdf?ver=accessable

[540] C. Barrientos, Special graceful labelings of irregular fences and lobsters, *Universal J. Math. Appl.*, **2**(1) (2019) 1-10. https://www.researchgate.net/publication/331906482_Special_Graceful_Labelings_of_Irregular_Fences_and_Lobsters

[541] C. Barrientos and S. Minion, New advances in Kotzig's conjecture, *Fundamental J. Math. Appl.,* **2**(2) (2019) 186-194 https://www.researchgate.net/publication/338081246_New_Advances_in_Kotzig's_Conjecture

[542] C. Barrientos and S. Minion, The graceful coalescence of alpha cycles, *Comm. Adv. Math. Sc.,*, **2**(2) (2019) 114-120. https://dergipark.org.tr/en/download/article-file/745297

[543] C. Barrientos and S. Minion, Series-parallel operations with alpha-graphs, *Theory Appl. Graphs*, **6**(1), Article 4 (2019). https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1082&context=tag

[544] C. Barrientos and S. Minion, Counting and labeling grid related graphs, *Electron. J. Graph Th. Appl.*, **7**(2) (2019) 349-363. https://ejgta.org/index.php/ejgta/article/view/753/pdf_117

[545] C. Barrientos and S. Minion, Some families of α-labeled subgraphs of the integral grid, *Commun. Comb. Optim.*, preprint.

[546] C. Barrientos and S. Minion, Broader families of cordial graphs, *Indonesian J. Combin.*, **4**(1) (2020) 34-52. https://www.ijc.or.id/index.php/ijc/article/view/145/111

[547] C. Barrientos and S. Minion, Harmonious graphs from α-trees, *J. Graph Th. Appl.*, **9**(2) (2021) 357-375. https://ejgta.org/index.php/ejgta/article/viewFile/1200/pdf_184

[548] C. Barrientos, M. Z. Youssef, Relaxing the injectivity constraint on arithmetic and harmonious labelings, *Electronic Graph Th. Appl.,* **10**(2) (2022) 523-539. doi:http://dx.doi.org/10.5614/ejgta.2022.10.2.13

[549] M. Barrus, Antimagic labeling and canonical decomposition of graphs, *Inform. Process. Lett.,* **110** (2010) 261-263. https://math.uri.edu/~barrus/papers/AntimagicPreprint.pdf

[550] A. Basak, MPLS multicasting using caterpillars and a graceful labelling scheme, IEEE Conference Publications, Information Visualisation, 2004. IV 2004. Proceedings. Eighth International Conference. 382-387. doi:10.1109/IV.2004.1320172

[551] S. S. Basha, Reverse super edge-magic strength of banana trees, *Int. J. Comput. Sci. Math.,* **9**(5) (2018) 515-523.

[552] M. Basher, On even vertex odd mean labeling of super subdivision of some graphs, *Ars Combin.,* **155** (2021) 95-105. https://www.researchgate.net/publication/363865534_On_even-odd_meanness_of_super_subdivision_of_some_graphs

[553] M. Basher, On the edge irregular reflexive labeling for some classes of plane graphs, *Soft Comput.*, (2023). https://link.springer.com/article/10.1007/s00500-023-07973-9

[554] M. Basher, Even vertex odd mean labeling of uniform the theta graphs, *Proyecciones J. Math.,* **43**(1) (2024) 153-162. https://www.revistaproyecciones.cl/index.php/proyecciones/libraryFiles/downloadPublic/91

[555] S. A. Bashammakh, W. C. Shiu, R. S. Sabibha, P. Jeyanthi, and M. E. Abdel-Aal, Further results on bijective product k-cordial labeling, *Mathematics*, **13**(15) (2025) 1-12.

[556] M. Basher, On even vertex odd mean labeling of the calendula graphs, *Proyecciones,* **39**(6) (2020) 1515-1535. https://www.researchgate.net/publication/346871598_On_even_vertex_odd_mean_labeling_of_the_calendula_graphs#fullTextFileContent

[557] M. Basher, k-Zumkeller labeling of super subdivision of some graphs, *J. Egypt Math. Soc.*, **29**(12) (2021). https://doi.org/10.1186/s42787-021-00121-y

[558] M. Basher, Even vertex odd mean labeling of some cycle related graphs, *J. Discrete Math. Sci. Crypt.,* (2021) 2331-2340.

[559] M. Basher, The reflexive edge strength of toroidal fullerene, *AKCE Internat. J. Graphs Combin.,* DOI:10.1080/09728600.2022.2150587

[560] M. Basher and M. Kamran Siddiqui, On even-odd meanness of super subdivision of some graphs, *Proyecciones* (Antofagasta) **41**(5) (2022) 1213-1228. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/5302/4150

[561] J. Baskar Babujee, Bimagic labelings in path graphs, *Math. Education,* **38** (2004) 12-16.

[562] J. Baskar Babujee, On edge bimagic labeling, *J. Combin. Inf. Syst. Sci.,* **28** (2004) 239-244.

[563] J. Baskar Babujee, Euler's phi function and graph labeling, *Int. J. Contemp. Math. Sci.,* **5** (2010) 977-984. https://www.m-hikari.com/ijcms-2010/17-20-2010/babujeeIJCMS17-20-2010.pdf z

[564] J. Baskar Babujee, Prime labelings on graphs, *Proc. Jangjeon Math. Soc.,* **10** (2007) 121-129.

[565] J. Baskar Babujee and S. Babitha, Encrypting and decrypting number using labeled graphs, *European J. Sci. Res.*, (2012) 14-24.

[566] J. Baskar Babujee and S. Babitha, On 1-vertex bimagic vertex labeling, *Tamkang J. Math., 45* (2014)(3) 259-272. https://journals.math.tku.edu.tw/index.php/TKJM/article/view/1004/894

[567] J. Baskar Babujee, S. Babitha, and V. Vishnupriya, New constructions of super edge bimagic labeling, *JCMCC,* **79** (2011) 23-31. https://combinatorialpress.com/article/jcmcc/Volume%20079/vol-079-paper%202.pdf

[568] J. Baskar Babujee and C. Beaula, On vertex strongly*-graph, *Proceed. Internat. Conf. Math. and Comput. Sci.,* 25-26 July 2008, Loyola College, Chennai.

[569] J. Baskar Babujee and R. Jagadesh, Super vertex total edge bimagic labeling for graphs with cycles, *Pacific-Asian J. Math.,* **2** (2008) 113-122.

[570] J. Baskar Babujee and R. Jagadesh, Super edge bimagic labeling for disconnected graphs, *Inter. Journal Appl. Math. Eng. Sci.*, **2** (2008) 171-175.

[571] J. Baskar Babujee and R. Jagadesh, Super edge bimagic labeling for trees, *Inter. J. Analyzing Methods Components Combin. Biol. Math.*, **1** (2008) 107-116.

[572] J. Baskar Babujee and R. Jagadesh, Super edge bimagic labeling for some classes of connected graphs derived from fundamental graphs, *Inter. J. Combin. Graph Theory Appl.*, **1** (2008) 85-92.

[573] J. Baskar Babujee and R. Jagadesh, Vertex consecutive edge bimagic labeling for star like graphs *Global J. Appl. Math. and Math. Sci.,* **1** (2008) 197-202.

[574] J. Baskar Babujee and R. Jagadesh, Prime labeling for some class of acyclic graphs, *Internat. J. Analyzing of Components and Combin. Biology in Math*, **1** (2009) 27-35.

[575] J. Baskar Babujee and R. Jagadesh, On a-vertex consecutive edge bimagic total labeling for graphs with cycles, *Indian J. Math. and Math. Sci.,* **5** (2009) 149-158.

[576] J. Baskar Babujee and R. Jagadesh, On a-vertex consecutive edge bimagic total labeling for certain classes of graphs, *Internat. J. Math. Sci.,* **9** (2010) 49-61.

[577] J. Baskar Babujee and R. Jagadesh, On edge vertex prime labeling, *Int. J. Pure Appl. Math.. 114*(6) (2017) 209-218. https://acadpubl.eu/jsi/2017-114-5/articles/2/24.pdf

[578] J. Baskar Babujee, R. Jagadesh, and V. Vishnupriya, On a-vertex consecutive edge bimagic total labeling for some class of graphs derives from fundamental graphs, *Pacific-Asian J. Math.*, **3** (2009) 283-291.

[579] J. Baskar Babujee and R. Jagadesh, Super edge bimagic labeling for some classes of connected graphs derived from fundamental graphs, *Internat. J. Combin. Graph Th. Appl.,* **3**(2) (2019) 119-126. https://www.ashwinanokha.com/resources/IJCGTA4-2-4-n.pdf

[580] J. Baskar Babujee, K. Kannan, and V. Vishnupriya, Vertex strongly*-graphs, *Internat. J. Analyzing Components and Combin. Biology in Math.,* **2** 19-25.

[581] J. Baskar Babujee and S. Loganathan, On signed product cordial labeling, **Applied Math.,** (2011) **2**(12) 1525-1530. https://www.scirp.org/pdf/AM20111200015_85335006.pdf

[582] J. Baskar Babujee and N. Rao, Edge-magic trees, *Indian J. Pure Appl. Math.,* **33** (2002) 1837-1840.

[583] J. Baskar Babujee and L. Shobana, Prime cordial labelings, *Int. Review on Pure and Appl. Math.,* **5** (2009) 277-282.

[584] J. Baskar Babujee and L. Shobana, Prime and prime cordial labeling for some special graphs, *Int. J. Contemp. Math. Sciences,* **5** (2010) 2347-2356.

[585] J. Baskar Babujee and L. Shobana, On Z_3-magic labeling and Cayley digraphs, *Int. J. Contemp. Math. Sciences*, **5** (2010) 2357-2368.

[586] J. Baskar Babujee and B. Suresh, New constructions of edge bimagic graphs from magic graphs, *Appl. Math.*, **(**2) (2011) 1393-1396. https://www.scirp.org/pdf/AM20111100014_81402625.pdf

[587] J. Baskar Babujee and V. Vishnupriya, Prime labelings on trees, *Internat. Review Pure Appl. Math.,* **2** (2006) 159-162.

[588] J. Baskar Babujee and V. Vishnupriya, Permutation labelings for some trees, *Internat. J. Math. Comput. Sci.,* **3** (2008) 31-38.

[589] J. Baskar Babujee, V. Vishnupriya, and R. Jagadesh, On a-vertex consecutive edge bimagic labeling for trees, *Internat. J. Comput. Math. and Numerical Simulation,* **2** 67-78.

[590] T. Baskaran and R. Ganapathy, A study on C-exponential mean labeling of graphs, *J. Math.,* **(7-8** (2022) 1-7. https://www.researchgate.net/publication/364248905_A_Study_on_C_-Exponential_Mean_Labeling_of_Graphs

[591] E. Baskoro and Y. Cholily, Expanding super edge-magic graphs, *J. Math. Fund. Sci.,* S.l., **36**(2) (2013) 117-125. https://typeset.io/pdf/expanding-super-edge-magic-graphs-113ma56nrt.pdf

[592] E. Baskoro and A. A. G. Ngurah, On super edge-magic total labelings, *Bull. Inst. Combin. Appl.,* **37** (2003) 82-87.

[593] E. T. Baskoro, I. W. Sudarsana, and Y. M. Cholily, How to construct new super edge-magic graphs from some old ones, *J. Indones. Math. Soc.,* **11**(2) (2005) 155-162. https://www.researchgate.net/publication/228940967_How_to_construct_new_super_edge-magic_graphs_from_some_old_ones

[594] J. Bass, personal communication (2013).

[595] A. R. Basunia, S. Das, L. Saha, and K. Tiwary, Antipodal number of full m-ary trees, *Theoret. Comput. Sci.*, **88**(5) (2021) 131–145. doi:(10.1016/j.tcs.2021.06.03

[596] A. R. Basunia, L. Saha, and K. Tiwary, Radio number of some path related graph, *Internat. J. Math. Trends Tech.*, **67**(2) (2021) 15-19. https://www.researchgate.net/publication/350695685_Radio_Number_of_Some_Path_Related_Graph

[597] D. Bantva, S. Vaidya, and S. Zhou, Some techniques to find large lower bound trees for the radio number, *Discrete Applied Math.,* **217** (2017) 110-122. https://www.researchgate.net/publication/369199805_Some_techniques_to_find_large_lower_bound_trees_for_the_radio_number

[598] R. Beals, J. Gallian, P. Headley, and D. Jungreis, Harmonious groups, *J. Combin. Th., Series A*, **56** (1991) 223-238. https://deepblue.lib.umich.edu/bitstream/handle/2027.42/29431/0000512.pdf;sequence=1

[599] A. F. Beardon, The maximum degree in a vertex-magic graph, *Austral. J. Combin.*, **30** (2004) 113-116.

[600] A. F. Beardon, Magic labellings of infinite graphs, *Austral. J. Combin.*, **30** (2004) 117-132. https://ajc.maths.uq.edu.au/pdf/30/ajc_v30_p117.pdf

[601] A. F. Beardon, The average degree in vertex-magic graph, *Austral. J. Combin.,* **35** (2006) 51-56. https://ajc.maths.uq.edu.au/pdf/35/ajc_v35_p051.pdf

[602] N. A. Beatress and P. B Sarasija, Even-odd harmonious graphs, *Internat. J. Math. Soft Comput.,* **5**(1) (2015) 23-29. https://www.researchgate.net/publication/349843792_Even-odd_Harmonious_Labeling_of_Some_Graphs

[603] N. A. Beatress, P. B. Sarasija, Square harmonious graphs, *Internat. J. Adv. Res. Sci., Engin. and Tech.,* **3**(2) (2016) 1428-1430. http://www.ijarset.com/upload/2016/february/10_IJARSET_sarasija.pdf

[604] G. Beaudoire, C. Bentz, and C. Picouleau, Antimagic labelling of graphs with maximum degree $\Delta(G) = n - 4$, https://arxiv.org/pdf/2603.02956 26pp.

[605] G. Beaudoire, C. Bentz and C. Picouleau, Antimagicness of join graphs, https://doi.org/10.48550/arXiv.2609.35245

[606] B. Beavers, Golomb rulers and graceful graphs, https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=a0bb24cd9b8ec1bbc17b7511321045b859e7636d z

[607] M. Beck and M. Farahmand, Partially magic labelings and the antimagic graph conjecture, *Sém. Lothar. Combin.,* **78B** (2017) Art. 86, 11 pp. https://arxiv.org/abs/1511.04154

[608] M. Beck and M. Jackanich, Bipartite graphs are weakly antimagic, arXiv:1306.1763v2 [math.CO] 18 Jun 2013.

[609] A. Bedi, M. Ollis, and S. Ramesh, On prime labelings of uniform cycle snake graphs, *PUMP J. Undergrad. Res.*, **6** (2023) 151-171. https://doi.org/10.46787/pump.v6i0.3633

[610] S. Beena, On Σ and Σ' labelled graphs, *Discrete Math.,* **309** (2009) 1783-1787. https://www.sciencedirect.com/science/article/pii/S0012365X08001489

[611] R. Raziya Begam, M. Palanivelrajan, K. Gunasekaran, and A. R. Shahul Hameed, Graceful labeling of some theta related graphs, *Internat. J. Fuzzy Mathematical Archives*, **2** (2013) 78-84. http://www.researchmathsci.org/IJFMAart/ijfma-v2-10.pdf

[612] A. K. Behera, D. Mishra, and P. C. Nayak, Some new classes of even-even and odd-even graceful graphs, *Internat. J. Appl. Engin. Res.,* **10**(19) (2015) 40171-40176. https://www.researchgate.net/publication/284896532_Some_new_classes_of_even_even_and_odd_even_graceful_graphs

[613] L. W. Beineke and S. M. Hegde, Strongly multiplicative graphs, *Discuss. Math. Graph Theory,* **21** (2001) 63-75. https://www.researchgate.net/publication/265360915_Strongly_multiplicative_graphs

[614] J. R. Bell, Products of graceful bipartite graphs, *Aequat. Math.,* **93** (2019) 1245-1260. https://link.springer.com/article/10.1007/s00010-019-00649-6

[615] J. R. Bell and D. F. Cummins, Graceful pairings, *A*ustralas. J. Combin., **75** (2019) 343-356. https://ajc.maths.uq.edu.au/pdf/75/ajc_v75_p343.pdf

[616] A. Benini and A. Pasotti, Decompositions of complete multipartite graphs, via generalized graceful labelings, *Australasian J. Combin.,* **59**(1) (2014) 120-143. https://ajc.maths.uq.edu.au/pdf/59/ajc_v59_p120.pdf

[617] A. Benini and A. Pasotti, α-labelings of a class of generalized Petersen graphs, *Discuss. Math. Graph Th.,* **35** (2015) 43-53. https://www.researchgate.net/publication/273311578_a-Labelings_of_a_Class_of_Generalized_Petersen_Graphs

[618] J. Bensmail, On the hardness of determining the irregularity strength of graphs, *Theoretical Computer Sci.*, **937** (2022) 96-107. https://www.sciencedirect.com/science/article/abs/pii/S0304397522005746

[619] J. Bensmail, H. Hocquard, D. Lajou, and E. Sopena, A proof of the multiplicative 1-2-3 conjecture, *Combinatorica,* (2023) https://doi.org/10.1007/s00493-023-00003-0

[620] K. Benson, M. Porter, and M. Tomova, The radio numbers of all graphs of order n and diameter $n-2$, *Matematiche (Catania),* **68**(2) (2013) 167-190. https://core.ac.uk/download/pdf/236004047.pdf

[621] M. Benson and S. M. Lee, On cordialness of regular windmill graphs, *Congr. Numer.*, **68** (1989) 45-58.

[622] K. Bérczi, A. Bernäth, and M. Vizer, Regular graphs are antimagic, *Electron. J. Combin.,* **22**(3) (2015) Paper 3.34, 1-6. arXiv:1504.08146

[623] C. Berge, Regularisable graphs II, *Discrete Math.,* **23** (1978) 91-95. https://www.sciencedirect.com/science/article/pii/0012365X78901085

[624] D. Bergstrand, F. Harary, K. Hodges, G. Jennings, L. Kuklinski, and J. Wiener, The sum numbering of a complete graph, *Bull. Malaysian Math. Soc.,* **12** (1989) 25-28.

[625] D. Bergstrand, K. Hodges, G. Jennings, L. Kuklinski, J. Wiener, and F. Harary, Product graphs are sum graphs, *Math. Magazine,* **65** (1992) 262-264. https://www.jstor.org/stable/pdf/2691455.pdf

[626] Z. Berikkyzy, A. Brandt, S. Jahanbekam, V. Larsen, and D. Rorabaugh, List-antimagic labeling of vertex-weighted graphs, *Discrete Math. Theor. Comput. Sci.,* **23**(3) (2021), Paper No. 5, 12 pp. https://dmtcs.episciences.org/8792/pdf

[627] O. Berkman, M. Parnas, and Y. Roditty, All cycles are edge-magic, *Ars Combin.,* **59** (2001) 145-151. https://www.researchgate.net/publication/220620173_All_Cycles_are_Edge-Magic

[628] A. H. Berliner, N. Dean, J. Hook, A. Marr, A. Mbirika, and C. D. McBee, Coprime and prime labelings of graphs, *J. Integer Seq.,* **19**(5) (2016) article 16.5.8, 14 pp. https://arxiv.org/abs/1604.07698

[629] J. C. Bermond, Graceful graphs, radio antennae and French windmills, *Graph Theory and Combinatorics*, Pitman, London (1979) 18-37. https://hal.science/hal-02340680/

[630] J. C. Bermond, A. E. Brouwer, and A. Germa, Systemes de triplets et differences associèes, *Problems Combinatories et Thèorie des Graphs*, Colloq. Intern. du Centre National de la Rech. Scient., 260, Editions du Centre Nationale de la Recherche Scientifique, Paris (1978) 35-38. https://hal.science/hal-02322623

[631] J. C. Bermond and G. Farhi, Sur un probleme combinatoire d'antennes en radioastronomie II, *Annals of Discrete Math.,* **12** (1982) 49-53.

[632] J. C. Bermond, A. Kotzig, and J. Turgeon, On a combinatorial problem of antennas in radioastronomy, in Combinatorics, A. Hajnal and V. T. Sós, eds., *Colloq. Math. Soc. János Bolyai*, **18**, 2 vols. North-Holland, Amsterdam (1978) 135-149. https://hal.science/hal-02322841/file/23-BKT76-antennas-hal.pdf

[633] J. C. Bermond and D. Sotteau, Graph decompositions and G-design, *Proc. 5th British Combin. Conf., 1975*, *Congr. Numer.,* **XV** (1976) 53-72. https://inria.hal.science/hal-02361407/document

[634] F. Bertault, M. Miller, H. Pé-Rosés, R. Feria-Puron, and E. Vaezpour, A heuristic for magic and antimagic graph labellings, *Proceed. VII Spanish Congress on Metaheuristics, and Evolutive and Bioinspired Algorithms,* (MAEB 2010) V. Campos, A. Duarte, M. Gallego, F. Cortazar, R. Marti (eds). Ibergarceta Publicaciones, S.L., Madrid. 677-684. arXiv:1305.1880

[635] M.-L. Beste, J.-H. de Wiljes, and M. Kreh, Sum and mod sum graphs of arithmetic sets, *Integers* **22** (2022) #51 23pp. http://math.colgate.edu/~integers/w51/w51.pdf

[636] D. Beutner and H. Harborth, Graceful labelings of nearly complete graphs, *Result. Math.,* **41** (2002) 34-39. https://link.springer.com/article/10.1007/BF03322754

[637] L. Bezegová and J. Ivančo, An extension of regular supermagic graphs, *Discrete Math.,* **310**(24) (2010) 3571-3578. https://www.sciencedirect.com/science/article/pii/S0012365X10003481

[638] L. Bezegová and J. Ivančo, On conservative and supermagic graphs, *Discrete Math.,* **311**(21) (2011) 2428-2436. https://www.sciencedirect.com/science/article/pii/S0012365X11003232

[639] L. Bezegová and J. Ivančo, A characterization of complete tripartite degree-magic graphs, *Discuss. Math. Graph Theory* **32**(2) (2012) 243-253.

[640] L. Bezegová and J. Ivančo, Number of edges in degree-magic graphs, *Discrete Math.,* **313** (2013) 1349-1357. https://www.sciencedirect.com/science/article/pii/S0012365X13001039

[641] P. Bhals, S. Lake, and A. Werthem, Gracefulness of families of spiders, *Involve,* **3**(3) (2010) 241-247. https://msp.org/involve/2010/3-3/involve-v3-n3-p01-s.pdf

[642] V. Bhat-Nayak and U. Deshmukh, New families of graceful banana trees, *Proc. Indian Acad. Sci. Math. Sci.*, **106** (1996) 201-216. https://www.ias.ac.in/article/fulltext/pmsc/106/02/0201-0216

[643] V. Bhat-Nayak and U. Deshmukh, Gracefulness of $C_{4t} \cup K_{1,4t-1}$ and $C_{4t+3} \cup K_{1,4t+2}$, *J. Ramanujan Math. Soc.,* **11** (1996) 187-190.

[644] V. Bhat-Nayak and U. Deshmukh, Skolem-graceful labelings of unions of paths, personal communication (1998).

[645] V. Bhat-Nayak and U. Deshmukh, Gracefulness of $C_3 \cup P_n$, unpublished.

[646] V. Bhat-Nayak and U. Deshmukh, Gracefulness of $C_{2x+1} \cup P_{x-2\theta}$, *Proc. International Conf. on Graph Theory and Number Theory, Trichy 1996.*

[647] V. N. Bhat-Nayak and S. K. Gokhale, Validity of Hebbare's conjecture, *Util. Math.*, **29** (1986) 49-59.

[648] V. N. Bhat-Nayak and A. Selvam, Gracefulness of n-cone $C_m \vee K_n^c$, *Ars Combin.,* **66** (2003) 283-298.

[649] V. N. Bhat-Nayak and S. Telang, Cahit-k-equitability of $C_n \circ K_1$, $k = n$ to $2n-1$, $n \geq 3$, *Congr. Numer.,* **155** (2002) 131-213.

[650] V. N. Bhat-Nayak and S. Telang, Cahit-equitability of coronas, *Ars Combin.,* **71** (2004) 3-32.

[651] A. A. Bhatti, M. Javaid, and M. Hussain, On super (a, d)-edge-antimagic total labeling of subdivided caterpillar, *Util. Math.,* **98** (2015) 227-241.

[652] A. A. Bhatti, M. Tahir and M. Javaid, Super (a, d)-EAT labeling of some wheel like graphs, *J. Graph Labeling,* **1**(2) (2015) 137-149.

[653] A. A. Bhatti, Q. Zahra, and M. Javaid, Further results on super (a, d)-EAT labeling of subdivided stars, *Util. Math.,* **98** (2015) 113-126.

[654] A. H. Bhavale, Antimagic labelings of a complete graph, *Math. Student*, **94** (3-4) (2025) 113-120. https://www.researchgate.net/publication/392839051_Antimagic_labelings_of_a_complete_graph

[655] A. H. Bhavale, Equivalence of labeled graphs and lattices, *Congressus Numerantium*, (2025) 13 pages.. https://arxiv.org/abs/2501.05064

[656] Z. Bi, A. Byers, S. English, E. Laforge, and P. Zhang, Graceful colorings of graphs, *J. Combin. Math. Combin. Comput.,* **101** (2017) 101-119.

[657] Z. Bi, A. Byers and P. Zhang, Revisiting graceful labelings of graphs, *J. Combin. Math. Combin. Comput.*, (2017) 141-158. https://combinatorialpress.com/article/jcmcc/Volume%20102/vol-102-paper%2010.pdf

[658] M. P. Biatch, J. S. Baggab, and S. Arumugam, A survey and a new class of graceful unicylic graphs, *AKCE Inter. J. Graphs and Comb. 17*(2) 2020, 673-678. https://doi.org/10.1080/09728600.2020.1832853

[659] A. Bigham, E.A. Donovan, J. Pack, J. Turley, and L. W. Wiglesworth, Prime labelings of snake graphs, *Pump J. Undergraduate research*, **2** (2019) 131-149. https://journals.calstate.edu/pump/article/view/1274

[660] L. Bijukumar, Some path and star related discrete graphs, *Math. Stat.* **13**(5) (2025) 269-278. doi:DOI:10.13189/ms.2025.130502

[661] R. Binthiya and P. B. Sarasija, Some new even harmonious graphs, *Internat. Math. Soft Comput.*, **4**(2) (2014) 105-111. https://oaji.net/articles/2017/296-1504713618.pdf

[662] A. Blinco, S. El-Zanati, and C. Vanden Eynden, On the cyclic decomposition of complete graphs into almost-bipartite graphs, *Discrete Math.,* **284** (2004) 71-81. https://www.sciencedirect.com/science/article/pii/S0012365X04001244

[663] G. S. Bloom, A chronology of the Ringel-Kotzig conjecture and the continuing quest to call all trees graceful, *Ann. N.Y. Acad. Sci.*, **326** (1979) 32-51. https://nyaspubs.onlinelibrary.wiley.com/doi/abs/10.1111/j.1749-6632.1979.tb17766.x

[664] G. S. Bloom and S. W. Golomb, Applications of numbered undirected graphs, *Proc. IEEE,* **65** (1977) 562-570. https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=1454786

[665] G. S. Bloom and S. W. Golomb, Numbered complete graphs, unusual rulers, and assorted applications, in *Theory and Applications of Graphs, Lecture Notes in Math.*, **642**, Springer-Verlag,New York (1978) 53-65. https://link.springer.com/chapter/10.1007/BFb0070364

[666] G. S. Bloom, P. Hell, and H. Taylor, Collecting autographs: n-node graphs that have n-integer signatures, *Annals N.Y. Acad. Sci.,* **319** (1979) 93-102.

[667] G. S. Bloom and D. F. Hsu, On graceful digraphs and a problem in network addressing, *Congr. Numer.,* **35** (1982) 91-103.

[668] G. S. Bloom and D. F. Hsu, On graceful directed graphs that are computational models of some algebraic systems, *Graph Theory with Applications to Algorithms and Computer Science*, Ed. Y. Alavi, Wiley, New York (1985) 89-102.

[669] G. S. Bloom and D. F. Hsu, On graceful directed graphs, *SIAM J. Alg. Discrete Meth.,* **6** (1985) 519-536.

[670] G. S. Bloom, A. Marr, W. D. Wallis, Magic digraphs, *J. Combin. Math. Combin. Comput.,* **65** (2008) 205-212.

[671] C. Bloomfield, D. D.-F. Liu, and J. Ramirez, Radio-k-labeling of cycles for large k, *Discrete Appl. Math.,* **316** (2022) 60-70. https://www.sciencedirect.com/science/article/pii/S0166218X22000981

[672] H. Bodlaender, J. Deogun, K. Jansen, T. Kloks, D. Kratsch, H. Müller, and Z. Tuza, Rankings of graphs, *SIAM J. Discrete Math.,* **11** (1998) 168-181.

[673] R. Bodendiek, H. Schumacher, and H. Wegner, Űber graziőse Graphen, *Math.-Phys. Semesterberichte,* **24** (1977) 103-106.

[674] R. Bodendiek and G. Walther, Arithmetisch antimagische Graphen, *Graphentheorie III*, K. Wagner and R. Bodendiek (eds.), Mannhein, 1993.

[675] R. Bodendiek and G. Walther, On number theoretical methods in graph labelings, *Res. Exp. Math.,* **21** (1995) 3-25. https://www.academia.edu/75834676/On_number_theoretical_methods_in_graph_labelings

[676] R. Bodendiek and G. Walther, (A, D)-antimagic parachutes, *Ars Combin.*, **42** (1996) 129-149.

[677] R. Bodendiek and G. Walther, (a, d)-antimagic parachutes II, *Ars Combin.*, **46** (1997) 33-63. https://combinatorialpress.com/article/ars/Volume%20042/volume-42-paper-10.pdf

[678] R. Bodendiek and G. Walther, On arithmetic antimagic edge labelings of graphs, *Mitt. Math. Ges. Hamburg,* **17** (1998) 85-99. https://www.researchgate.net/publication/266363684_On_arithmetic_antimagic_edge_labelings_of_graphs

[679] R. Bodendiek, H. Schumacher, and H. Wegner, Űber eine spezielle Klasse groziőser Eulerscher Graphen, *Mitt. Math. Gesellsch. Hamburg*, **10** (1975) 241-248.

[680] T. Bohman and D. Kravitz, On the irregularity strength of trees, *J. Graph Theory,* **45** (2004) 241-254. https://onlinelibrary.wiley.com/doi/epdf/10.1002/jgt.10158

[681] S. A. H. Bokhary, A. Ahmad, and M. Imran, On vertex irregular total labelings of cartesian products of two paths, *Util. Math.,* **90** (2013) 239-249.

[682] S. A. H. Bokhary, U. Ali, and S. Maqbool, Irregular total labeling of wheel related graphs, *Util. Math.,* **107** (2018) 231-242.

[683] S. A. Bokhary and H. Faheem, Vertex irregular total labeling of grid graph, *Palest. J. Math.,* **8**(1) (2019) 52-62. https://pjm.ppu.edu/sites/default/files/papers/PJM_Oct2018_52to62.pdf

[684] S. A. Bokhary, M. Imran, and U. Ali, On the total irregularity strength of convex polytope graphs, *Proyecciones,* **40**(5) (2021) 1267-1277. https://www.scielo.cl/pdf/proy/v40n5/0716-0917-proy-40-05-1267.pdf

[685] J. Boland, R. Laskar, C. Turner, and G. Domke, On mod sum graphs, *Congr. Numer.,* **70** (1990) 131-135.

[686] J. Bondy and U. Murty, *Graph Theory with Applications*, North-Holland, New York (1976).

[687] C. P. Bonnington and J. Širáň, Bipartite labelings of trees with maximum degree three, *J. Graph Theory,* **31** (1999) 79-91. https://onlinelibrary.wiley.com/doi/epdf/10.1002/%28SICI%291097-0118%28199905%2931%3A1%3C7%3A%3AAID-JGT2%3E3.0.CO%3B2-Y

[688] R. Boonklurb, A. Narissayaporn, and S. Singhun, Super edge-magic labeling of m-node k-uniform hyperpaths and m-node k-uniform hypercycles, *AKCE Internat. J. Graphs and Combin.,* **13** (2016) 218-229. https://www.sciencedirect.com/science/article/pii/S097286001530058X

[689] R. Boonklurb, N. Ruamkaew, and S. Singhun, Directed edge-graceful labeling of digraph consisting of c cycles of the same size, *J. Discrete Math. Sci.Cryptography*, **25**(1) (2022) 53-72.

[690] I. Borosh, D. Hensley, and A. Hobbs, Vertex prime graphs and the Jacobsthal function, *Congr. Numer.,* **127** (1997) 193-222.

[691] K. J. Bosco and M. Dinesh, On radio Heronian Dd-distance mean of some standard graphs, *Internat. J. Health Sci.*, 6(S3) (2020) 3038-3043. https://media.neliti.com/media/publications/429722-on-radio-heronian-dd-distance-mean-of-so-82441092.pdf

[692] K. J. Bosco and M. Dinesh, On radio Heronian DD-distance mean labeling of some basic graphs, *J. Emerging Tech. Innovative Res. (JETIR)*, **8**(1) (2021) 441-445. https://www.jetir.org/papers/JETIR2101264.pdf

[693] M. I. Bosmia and K. K. Kanani, Divisor cordial labeling in the context of graph operations on bistar, *Global J. Pure and Appl. Math.,* **12**(3) (2016) 2605-2618. https://www.researchgate.net/publication/306214795_Divisor_cordial_labeling_in_the_context_of_graph_operations_on_bistar

[694] M. I. Bosmia and K. K. Kanani, Various graph labeling techniques for the line graph of bistar, *Internat. J. Tech. Innovat. Mod. Eng. Sci.* **4**(09) (2018) 851–858. https://ijtimes.com/paper-view.php?id=IJT-2021-026BBA&utm_source=chatgpt.com

[695] M. I. Bosmia and K. K. Kanani, Divisor cordial labeling in the context of corona product, 9th National Level Science Symposium. February 14, 2016, Organized by Christ College, Rajkot, Sponsored by GUJCOST, Gandhinagar, Mathematics & Statistics, **3** (2016) 178-182. https://www.researchgate.net/publication/316830482_Divisor_Cordial_Labeling_in_the_Context_of_Corona_Product

[696] M. I. Bosmia, M. A. Patel, Y. M. Parmar, and P. L. Vihol, Cordial labeling of the line graph of bistar *Commun. Mathe. Appl.*, **16**(3) (2025) 895–907. doi:DOI:10.26713/cma.v16i3.3317

[697] D. Bouchard, P. Clark, and H.-H. Su, On edge-balance index sets of L-product of cycles, *J. Indones. Math. Soc.,* Special Edition (2011) 27-38. https://www.researchgate.net/publication/266582070_On_the_edge-balance_index_sets_of_L-product_of_cycles

[698] D. Bouchard, P. Clark, S. M. Lee, S.-P. Lo, and H.-H. Su, On balance index sets of generalized book and ear expansion graphs, 24th MCCCC. *J. Combin. Math. Combin. Comput.,* **82** (2012) 3-15. https://combinatorialpress.com/article/jcmcc/Volume%20082/vol-082-paper%201.pdf

[699] D. Bouchard, P. Clark, and H.-H. Su, On edge-balance index sets of L-product of cycles with stars, Part II, 24th MCCCC. *J. Combin. Math. Combin. Comput.,* **82** (2012) 199-209. https://combinatorialpress.com/article/jcmcc/Volume%20082/vol-082-paper%2013.pdf

[700] S. Boxwala, A. S. Phadke, P. N. Shinde, and N. Mundlik, On group A-cordial labelling of double uniform $(t_1^{\ell-1}, t_2^{\ell_2})$-ply, *Indian J. Adv. Math. (IJAM)*, **6**(1) 44-48. https://www.ijam.latticescipub.com/wp-content/uploads/papers/v6i1/A124206010426.pdf

[701] S. A. Boxwala and P. Vashishta, Some new families of graceful graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014) 127-133, *Electron. Notes Discrete Math.*, **48**, Elsevier Sci. B. V., Amsterdam, 2015. https://www.sciencedirect.com/science/article/pii/S1571065315000190

[702] S. Brandt, J. Miškuf, and D. Rautenbach, On a conjecture about edge irregular total labelings, *J. Graph Theory*, **57** (2008) 333-343. https://onlinelibrary.wiley.com/doi/epdf/10.1002/jgt.20287

[703] L. Brankovic, C. Murch, J. Pond, and A. Rosa, Alpha-size of trees with maximum degree three and perfect matching. In: *Proceedings of AWOCA2005*, 18-21 September, Ballarat, Australia, (2005) 47-56.

[704] L. Brankovic and M. J. Reynolds, Computer search for graceful labeling: a survey, *Elect. J. Graph Th. Appl.,* **10**(1) (2022) 319-336. https://www.ejgta.org/index.php/ejgta/article/view/1057/pdf_225

[705] L. Brankovic, A. Rosa, and J. Širáň, Labellings of trees with maximum degree three-an improved bound, *J. Combin. Math. Combin. Comput.,* **55** (2005) 159-169. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2015.pdf

[706] L. Brankovic and I. M. Wanless, Graceful labelling: State of the art, applications and future directions, *Math Comput. Sci.*, **5** (2011) 11-20. https://www.researchgate.net/publication/220437473_Graceful_Labelling_State_of_the_Art_Applications_and_Future_Directions

[707] R. L. Bras, C. P. Gomes, and B. Selman, Double-wheel graphs are graceful, *IJCAI'13 Proceedings of the Twenty-Third International Joint Conference on Artificial Intelligence*, AAAI Press (2013) 587-593. https://www.ijcai.org/Proceedings/13/Papers/094.pdf

[708] L. Branson, A. Sutton, and X. Yan, Finding antimagic labelings of trees by evolutionary search, *Conference: FOGA '23: Foundations Genetic Algor., XVII*, (2023) 27-37. https://dl.acm.org/doi/pdf/10.1145/3594805.3607133

[709] B. Brešar and S. Klavžar, Θ-graceful labelings of partial cubes, *Discrete Math.* **306** (2006) 1264-1271. https://www.sciencedirect.com/science/article/pii/S0012365X06001609

[710] V. T. Brindha Mary, C. David Raj, and C. Jayasekaran, Radio even mean graceful labeling on some special graphs, *Malaya J. Mat.*, **8**(04) (2020) 2323-2328. https://www.malayajournal.org/index.php/mjm/article/view/1912/1293

[711] G. Brinkmann, S. Crevals, H. Mélot, L.J. Rylands, and E. Steffan, α-labelings and the structure of trees with nonzero α-deficit. *Discrete Math. Theor. Comput. Sci.*, **14**(1) (2012) 159-174. https://www.researchgate.net/publication/265628588_alpha-Labelings_and_the_Structure_of_Trees_with_Nonzero_alpha-Deficit

[712] H. Broersma and C. Hoede, Another equivalent of the graceful tree conjecture, *Ars Combin.,* **51** (1999) 183-192. https://combinatorialpress.com/article/ars/Volume%2051/volume-51-paper-17.pdf

[713] C. Bu, Gracefulness of graph $K_n + \overline{K_m}$, *J. Harbin Shipbuilding Eng. Inst.*, **15** (1994) 91-93.

[714] C. Bu, On edge-gracefulness of graphs, *Heilongjiang Daxue Ziran Kexue Xuebao,* **12** (1995) 18-20.

[715] C. Bu, Sequential labeling of the graph $C_n \odot \overline{K}_m$, unpublished.

[716] C. Bu and C. Cao, The gracefulness for a class of disconnected graphs, *J. Natural Sci. Heilongjiang Univ.,* **12** (1995) 6-8.

[717] C. Bu and L. Chen, Some conclusions about graceful graphs, *J. Harbin Shipbuilding Eng. Inst.*, **14** (1993) 103-108.

[718] C. Bu and W. Feng, Some composite theory about graceful graphs, *J. Harbin Eng. Univ.*, **16** (1995) 94-97.

[719] C. Bu, Z. Gao, and D. Zhang, On k-gracefulness of $r - p_n \times p_2$, *J. Harbin Shipbuilding Eng. Inst.*, **15** (1994) 85-89.

[720] C. Bu and B. He, The k-gracefulness of some graphs, *J. Harbin Eng. Univ.*, **14** (1993) 92-95.

[721] C. Bu and J. Shi, A class of (k, d)-arithmetic graphs, *J. Harbin Eng. Univ.*, **16** (1995) 79-81.

[722] C. Bu and J. Shi, Some conclusions about indexable graphs, *J. Harbin Eng. Univ.*, **16** (1995) 92-94.

[723] C. Bu, D. Zhang, and B. He, k-gracefulness of C_n^m, *J. Harbin Shipbuilding Eng. Inst.*, **15** (1994) 95-99.

[724] C. Bu and J. Zhang, The properties of (k, d)-graceful graphs, unpublished.

[725] N. I. S. Budi, Kekuatan Sisi Refleksif pada Graf Tadpole, Graf Diamond Snake, Graf Quadrilateral Snake dan Graf K1n P3, Skripsi, Prodi Matematika FMIPA UNS, 2021.

[726] R. Bunge, S. El-Zanati, W. O'Hanlon, and C. Vanden Eynden, On γ-labeling almost-bipartite graphs, *Ars Combin.,* **107** (2012) 65-80. https://combinatorialpress.com/article/ars/Volume%20107/volume-107-paper-5.pdf

[727] R. Bunge, A. Chantasartrassmee, S. El-Zanati, and C. Vanden Eynden, On cyclic decompositions of complete graphs into tripartite graphs, *J. Graph Theory,* **72** (2013) 90-111. https://onlinelibrary.wiley.com/doi/10.1002/jgt.21632

[728] M. Buratti and A. Del Fra, Existence of cyclic k-cycle systems of the complete graph, *Discret. Math.*, **261** (2003) 113-125. https://www.sciencedirect.com/science/article/pii/S0012365X02004636

[729] M. Buratti, G. Rinaldi, and T. Traetta, Some results on 1-rotational Hamiltonian cycle systems, *J. Combin. Designs,* **22**(6) (2014) 231-251.

[730] A. C. Burgess, P. Danziger, and T. Traetta, On the Oberwolfach problem for single-flip 2-factors via graceful labelings, *J. Combin. Theory*, Series A, **189** (2022), 105611, 39 pages. arXiv:2010.07231

[731] M. Burzio and G. Ferrarese, The subdivision graph of a graceful tree is a graceful tree, *Discrete Math.*, **181** (1998) 275-281. https://www.sciencedirect.com/science/article/pii/S0012365X97000691?via%3Dihub

[732] S. I. Butt, M. Numan, I. A. Shah, and S. Ali, Face labelings of type (1,1,1) for generalized prism, *Ars Combin.,* **137** (2018) 41-52.

[733] A. Byers and A. O'Mellan, An amalgamation of graceful and harmonious labelings, *Bull. Inst. Combin. Appl.*, **96** (2022) 53-67. http://bica.the-ica.org/Volumes/96//Reprints/BICA2021-31-Reprint.pdf

[734] S. Cabaniss, R. Low, and J. Mitchem, On edge-graceful regular graphs and trees, *Ars Combin.*, **34** (1992) 129-142. https://combinatorialpress.com/article/ars/Volume%20034/volume-34-paper-14.pdf?utm_source=chatgpt.com

[735] L. Caccetta and R. Jia, Positive binary labelings of graphs, *Austral. J. Combin.*, **14** (1996) 133-148. https://ajc.maths.uq.edu.au/pdf/14/ocr-ajc-v14-p133.pdf

[736] R. Čada, J. Ekstein, P. Holub, and O. Togni, Radio labelings of distance graphs, *Discrete Appl. Math.*, **161**(18) (2013) 2876-2884. https://www.sciencedirect.com/science/article/pii/S0166218X13003028

[737] I. Cahit, Elegant valuation of the paths, *Ars Combin.*, **16** (1983) 223-227. https://www.researchgate.net/publication/266984918_Elegant_valuation_of_the_paths

[738] I. Cahit, Cordial graphs: a weaker version of graceful and harmonious graphs, *Ars Combin.,* **23** (1987) 201-207. https://www.researchgate.net/publication/233792958_Cordial_Graphs_A_Weaker_Version_of_Graceful_and_Harmonious_Graphs

[739] I. Cahit, On cordial and 3-equitable labellings of graphs, *Util. Math.*, **37** (1990) 189-198.

[740] I. Cahit, Status of graceful tree conjecture in 1989, in *Topics in Combinatorics and Graph Theory*, R. Bodendiek and R. Henn (eds), *Physica-Verlag*, Heidelberg (1990) 189-198. https://www.researchgate.net/publication/233792994_On_cordial_and_3-equitable_labelings_of_graphs

[741] I. Cahit, Recent results and open problems on cordial graphs, *Contemporary Methods in Graph Theory*, R. Bodendiek (ed.), Wissenschaftsverlag, Mannheim, 1990, 209-230.

[742] I. Cahit, Equitable tree labellings, *Ars Combin.,* **40** (1995) 279-286. https://combinatorialpress.com/article/ars/Volume%20040/volume-40-paper-22.pdf

[743] I. Cahit, On harmonious tree labellings, *Ars Combin.,* **41** (1995) 311-317. https://combinatorialpress.com/article/ars/Volume%20041/volume-41-paper-29.pdf

[744] I. Cahit, H-cordial graphs, *Bull. Inst. Combin. Appl.*, **18** (1996) 87-101.

[745] I. Cahit, Some totally modular cordial graphs, *Discuss. Math. Graph Theory,* **22** (2002) 247-258. doi:10.7151/dmgt.1173

[746] I. Cahit, Graceful labelings of rooted complete trees, personal communication (circa 2002). https://www.academia.edu/4376717/Graceful_Labeling_of_the_Rooted_Complete_Trees

[747] I. Cahit, A note on graceful directed trees, (2003) Science Direct Working Paper No S1574-0358(04)70939-9, https://ssrn.com/abstract=3177648

[748] I. Cahit and R. Yilmaz, E_3-cordial graphs, *Ars Combin.,* **54** (2000) 119-127. https://www.researchgate.net/publication/265367290_E_3_-cordial_graph

[749] H. Cai, L. X. Wei, X. R. Lu, Gracefulness of unconnected graphs $(P_1 \vee P_n) \cup G_r$, $(P_1 \vee P_n) \cup (P_3 \vee \overline{K}_r)$ and $W_n \cup St(m)$, *J. Jilin Univ. Sci.,* **45** (2007) 539-543.

[750] B. Calhoun, K. Ferland, L. Lister, and J. Polhill, Totally magic labelings of graphs, *Australas. J. Combin.,* **32** (2005) 47-59. https://ajc.maths.uq.edu.au/pdf/32/ajc_v32_p047.pdf

[751] N. Cairnie and K. Edwards, The computational complexity of cordial and equitable labelling, *Discrete Math.,* **216** (2000) 29-34. https://www.sciencedirect.com/science/article/pii/S0012365X99002952

[752] J. M. Campbell, On a conjecture concerning antipodal labelings for cycles, doi:10.48550/arXiv.2609.29477

[753] L. A. Cammack, Schelp and G. Schrag, Irregularity strength of full d-ary trees *Congr. Numer.,* **81** (1991) 113-119.

[754] D. Canales, M. Tomova, and C. Wyels, A gap in the achievable radio number line, *AKCE Int. J. Graphs Combin.,* **10**(4) (2013) 349-357. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088751

[755] M. R. Cerioli, C. G. Fernandes, O. Lee, C. N. Lintzmayer, On edge-magic labelings of forests, *Elect. Notes Theor. Comput. Sci.,* **346**(30) (2019) 299-307. https://www.sciencedirect.com/science/article/pii/S1571066119300775

[756] K. Carlson, Generalized books and C_m-snakes are prime graphs, *Ars Combin.* **80** (2006) 215-221. https://combinatorialpress.com/article/ars/Volume%20080/volume-80-paper-20.pdf

[757] Y. Caro, Y. Roditty, and J. Schőnheim, Starters for symmetric $(n, G, 1)$-designs. ρ-labelings revisited, unpublished.

[758] H. Carter and N. B. Fox, Odd prime graph labelings, arXiv:2208.08488 22 pages.

[759] A. Catral, New constructs toward the Graham–Sloane harmonious labelling conjecture, preprint. https://www.researchgate.net/publication/395346735

[760] R. Cattell, Vertex magic total labelings of complete multipartite graphs, *J. Combin. Math. Combin. Computing,* **55** (2005) 187-197. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2017.pdf

[761] R. Cattell, Graceful labellings of paths, *Discrete Math.,* **307** (2007) 3161-3176. https://www.sciencedirect.com/science/article/pii/S0012365X07001215

[762] N. Cavenagh, D. Combe, and A. M. Nelson, Edge-magic group labellings of countable graphs, *Electronic J. Combin.,* **13** (2006) #R92 (19 pages). https://www.combinatorics.org/ojs/index.php/eljc/article/view/v13i1r92/pdf

[763] D. Chakraborty, S. Nandi, S. Sen, and D. K. Supraja, A linear algorithm for radio k-coloring powers of paths having small diameter. In: Hsieh, SY., Hung, LJ., Lee, CW. (eds) *Combin. Alg.,*. IWOCA 2023. Lecture Notes in Computer Science, **13889** (2023) Springer, Cham. https://doi.org/10.1007/978-3-031-34347-6_13

[764] P. Chalise and R. M. Low, Application of the combinatorial nullstellensatz to magic-type graph labelings, https://arxiv.org/pdf/2607.20724

[765] W. H. Chan, R. M. Low, and W. C. Shiu, Group-antimagic labelings of graphs, *Congr. Numberrantim,* **217** (2013) 21-31. https://www.researchgate.net/publication/264947717_Group-antimagic_labelings_of_graphs

[766] B. Chandra and R. Kala, Group S_3 cordial prime labeling of graphs, *Malaya J. Matematik,* **S**(1) (2019) 403-407. https://www.researchgate.net/publication/334700018_Group_S_3_cordial_prime_labeling_of_graphs

[767] C.-W. Chang, D. Kuo, and H.-C. Lin, Ranking numbers of graphs, *Inform. Process. Letters* **110** (2010) 711-716. https://www.sciencedirect.com/science/article/pii/S0020019010001481?via%3Dihub

[768] F.-H. Chang, H.-B. Chen, W.-T. Li, and Z. Pan, Shifted-antimagic labelings for graphs, *Graphs Combin.,* **37** (2021) 1065–1182. arXiv:1806.06019.

[769] F.-H. Chang, P. Chin, W.-T. Li, and Z. Pan, The strongly antimagic labelings of double spiders, arXiv:1712.09477

[770] F.-H. Chang, W.-T. Li, D.-F. Daphne Liu, and Z. Pan, On k-shifted antimagic spider forests, *Disc. Appl. Math.,* **358**(15) (2024) 468-476.

[771] F. Chang, Y.-C. Liang, Z. Pan, and X. Zhu, Antimagic labeling of regular graphs, *J. Graph Theory,* **82** (2016) 339-349. arXiv:1505.07688

[772] G. J. Chang, Strong sum graphs, *Bull. Inst. Combin. Appl.,* **7** (1993) 47-52.

[773] G. J. Chang, D. F. Hsu, and D. G. Rogers, Additive variations on a graceful theme: some results on harmonious and other related graphs, *Congr. Numer.,* **32** (1981) 181-197.

[774] Y.-W. Chang and S.-P. Liu, Cartesian products of some regular graphs admitting antimagic labeling for arbitrary sets of real numbers, *J. Math.,* 2021, Art. ID 4627151, 8 pp. https://www.researchgate.net/publication/357475062_Cartesian_Products_of_Some_Regular_Graphs_Admitting_Antimagic_Labeling_for_Arbitrary_Sets_of_Real_Numbers

[775] R. Charishma and P. Nageswari, Cordial labeling in the context of some graphs, *J. Informa. Sys. Engin. Management*, **10**(35a) (2025) 1-31. https://jisem-journal.com/index.php/journal/article/view/6813/3151

[776] R. Charishma and P. Nageswari, L-Fibonacci cordial labeling to create new L-Fibonacci cordial families, *Math. Stat.,* **13**(4) (2025) 194-200. https://www.hrpub.org/download/20250830/MS2-13441462.pdf

[777] G. Chartrand, D. Erwin, D. VanderJagt, and P. Zhang, γ-labelings of graphs, *Bull. Inst. Combin Appl.,* **44** (2005) 51-68. https://www.researchgate.net/publication/292377888_ON_gamma-LABELINGS_OF_GRAPHS

[778] G. Chartrand, D. Erwin, D. VanderJagt, and P. Zhang, On γ-labeling of trees, *Discuss. Math. Graph Theory,* **25** (2005) 363-383. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1289

[779] G. Chartrand, D. Erwin, and P. Zhang, Radio antipodal colorings of graphs, *Math. Bohem.,* **127** (2002) 57-69. https://dml.cz/bitstream/handle/10338.dmlcz/133978/MathBohem_127-2002-1_7.pdf

[780] G. Chartrand, D. Erwin, and P. Zhang, A graph labeling problem suggested by FM channel restrictions, *Bull. Inst. Combin. Appl.,* **43** (2005) 43-57.

[781] G. Chartrand, D. Erwin, P. Zhang, and F. Harary, Radio labelings of graphs, *Bull. Inst. Combin. Appl.,* **33** (2001) 77-85.

[782] G. Chartrand, H. Hevia, and O.R. Oellermann, The chromatic number of a factorization of a graph, *Bull. Inst. Combin. Appl.,* **20** (1997) 33-56.

[783] G. Chartrand, M. Jacobson, J. Lehel, O. Oellermann, S. Ruiz, and F. Saba, Irregular networks, *Congr. Numer.,* **64** (1988) 187-192. https://www.researchgate.net/publication/265701559_Irregular_networks

[784] G. Chartrand, S. M. Lee, and P. Zhang, Uniformly cordial graphs, *Discrete Math.,* **306** (2006) 726-737. https://www.sciencedirect.com/science/article/pii/S0012365X0600094X

[785] G. Chartrand and L. Lesniak, *Graphs & Digraphs* 4th ed. CRC Press (2005).

[786] G. Chartrand, R. Muntean, V. Seanpholphat and P. Zhang, Which graphs are divisor graphs, *Congr. Numer.,* **151** (2001) 180-200. https://www.researchgate.net/publication/268319541_Which_graphs_are_divisor_graphs

[787] G. Chartrand, L. Nebesky, and P. Zhang, Radio k-colorings of paths, *Discuss. Math. Graph Theory,* **24** (2004) 5-21. https://www.academia.edu/108888081/Radio_k_colorings_of_paths

[788] M. Chaurasiya, A. Limda, N. Parmar, and Rupani, Note on k-prime labeling of some graphs, *IRE Journals,* **4**(12) (2021) https://www.irejournals.com/formatedpaper/1702749.pdf

[789] A. Chavez, P. Le, D. Der-Fen Liu, and M. Shurman, Antimagic labeling for unions of graphs with many three-paths, arXiv:2203.14842v1 16 pages.

[790] A. Chavez, D. D.-F. Liu, and M. Shurman, Optimal radio-k-labelings of trees, *European J. Combin.*, **91** (2001) 103203. https://www.sciencedirect.com/science/article/pii/S0195669820301244

[791] P. D. Chawathe and V. Krishna, Odd graceful labelings of countably infinite locally finite bipartite graphs, *Conference on Graph Theory and its Applications*, March 2001, School of Mathematics, Anna University, Chennai.

[792] P. D. Chawathe and V. Krishna, Antimagic labeling of complete m-ary trees, Number theory and discrete mathematics (Chandigarh, 2000), 77-80, Trends Math., Birkhäuser, Basel, 2002.

[793] T. T. Chelvam, N. M. Rilwan, and G. Kalaimurugan, Antimagic and magic labelings in Cayley digraphs, *Australas. J. Combin.,* **55** (2013) 65-71. https://ajc.maths.uq.edu.au/pdf/55/ajc_v55_p065.pdf

[794] K.-J. Chen, S.-M. Lee, and Y.-C. Wang, On the edge-graceful indices of the $L-$product of $(p, p+1)$ graphs and K_2, unpublished.

[795] W. C. Chen, H. I. Lü, and Y. N. Yeh, Operations of interlaced trees and graceful trees, *Southeast Asian Bull. Math.*, **21** (1997) 337-348.

[796] L.-C. Chen, On harmonious labelings of the amalgamation of wheels, Master's Thesis, Chung Yuan Christian University, Taiwan.

[797] Z. Chen, Harary's conjectures on integral sum graphs *Discrete Math.*, **160** (1996) 241-244. https://www.sciencedirect.com/science/article/pii/0012365X9500163Q?via|%3Dihub

[798] Z. Chen, Integral sum graphs from identification, *Discrete Math.*, **181** (1998) 77-90. https://www.sciencedirect.com/science/article/pii/S0012365X97000460?via%3Dihub

[799] Z. Chen, On super edge-magic graphs. *J. Combin. Math. Combin. Comput.,* **38** (2001) 55-64. https://combinatorialpress.com/article/jcmcc/Volume%20038/vol-038-paper%205.pdf

[800] Z. Chen, On integral sum graphs, *Discrete Math.*, **306** (2006) 19-25. https://www.sciencedirect.com/science/article/pii/S0012365X05005935?via%3Dihub

[801] Z.-Z. Chen, A generalization of the Bodendiek conjecture about graceful graphs, *Topics in Combinatorics and Graph Theory*, R. Bodendiek and R. Henn, eds., Physica-Verlag, Heidelberg, 1990, 737-746.

[802] Z. Chen, On integral sum graphs with a saturated vertex, *Czechoslovak Math. J.,* **60**(135) (2010) 669-674. https://dml.cz/bitstream/handle/10338.dmlcz/140597/CzechMathJ_60-2010-3_6.pdf

[803] W-T Chen and J-M Kuo, Caterpillars with diameter at least 5 are k-shifted antimagic for any integers k, *AIMS Math.*, **11**(4) (2026) 9529–9540. doi:DOI:10.3934/math.2026394

[804] Y. Cheng, Lattice grids and prisms are antimagic, *Theoret. Comput. Sci.,* **374** (2007) 66-73. arXiv:0603106 [math.CO].

[805] Y. Cheng, A new class of antimagic Cartesian product graphs, *Discrete Math.,* **308** (2008) 6441-6448. https://www.sciencedirect.com/science/article/pii/S0012365X07010175?via%3Dihub

[806] H. Cheng, B. Yao, X. Chen, and Z. Zhang, On graceful generalized spiders and caterpillars, *Ars Combin.* **87** (2008) 181-191. https://combinatorialpress.com/article/ars/Volume%20087/volume-87-paper-15.pdf

[807] C. Chithra, G. Marimuthu, and G. Kumar, Some cycle-supermagic labeling of graphs, *Internat. J. Comput. Eng. Res.,* (IJCER) **8**(9) (2018) 79-83. http://www.ijceronline.com/papers/Vol8_issue9/Version-2/M0809027983.pdf

[808] G. Chitra, J. Priya, and Y. Vishnupriya, Odd Fibonacci mean labeling of some special graphs, *Int. J. Math. Trends Technol.*, **66**(1) (2020) 115-126. http://www.ijmttjournal.org/Volume-66/Issue-1/IJMTT-V66I1P515.pdf

[809] R. Chitra Devi and S. Saravana Kumar, Super root square mean labeling of some graphs, *Internat. J. Math. Trends Tech.*, (IJMTT) **30**(2) (2016) 92-94. https://www.researchgate.net/publication/317304532_Super_Root_Square_Mean_Labeling_of_Some_Graphs

[810] A. M. Chitre and N. B. Limaye, On 5-equitabilty of one point union of shells, *AKCE J. Graphs Combin.*, **6** (2009) 57-68. https://www.tandfonline.com/doi/epdf/10.1080/09728600.2009.12088872?needAccess=true&role=button

[811] A. M. Chitre and N. B. Limaye, On edge-3-equitability of $\overline{K_n}$-union of gears, *J. Combin. Math. Combin. Comput.,* **83** (2012) 129-150. https://combinatorialpress.com/article/jcmcc/Volume%20083/vol-083-paper%2012.pdf?utm_source=chatgpt.com

[812] A. M. Chitre and N. B. Limaye, On edge-3-equitability of $\overline{K_n}$-union of helms, *J. Combin. Math. Combin. Comput.*, **94** (2015) 283–299. https://combinatorialpress.com/article/jcmcc/Volume%20094/vol-094-paper%2022.pdf

[813] J. O. Choi, J. P. Georges, and D. Mauro, David, Relating edge-coverings to the classification of Z_2^k-magic graphs, *Discrete Math.,* **312**(19) (2012) 2938-2945. https://www.sciencedirect.com/science/article/pii/S0012365X12002543?via%3Dihub

[814] D. Chopra, R. Dios, and S. M. Lee, On the edge-magicness of Zykov sum of graphs, personal communication (2013).

[815] D. Chopra and S. M. Lee, On super edge-magic graphs which are weak magic, *J. Combin. Math. Combin. Comput.,* **62** (2007) 177-187. https://combinatorialpress.com/article/jcmcc/Volume%20062/vol-062-paper%2017.pdf

[816] D. Chopra and S. M. Lee, On $Q(a)P(b)$-super edge-graceful graphs, *J. Combin. Math. Combin. Comput.*, **58** (2006) 135-152. https://www.researchgate.net/publication/229076137_On_the_k-edge_magic_graphs

[817] D. Chopra and S. M. Lee, On the integer magic spectra of the generalized theta graphs, unpublished.

[818] D. Chopra, S. M. Lee, and H. H. Su, On edge-balance index sets of fans and wheels, unpublished

[819] C.-C. Chou and S. M. Lee, On Z_3-magic graphs, unpublished.

[820] C.-C. Chou, M. Galiardi, M. Kong, S. M. Lee, and D. Perry, On edge-balance index sets of L-product of cycles with stars, Part I. *J. Combin. Math. Combin. Comput.,* **78** (2011) 195-211. https://combinatorialpress.com/article/jcmcc/Volume%20078/vol-078-paper%2016.pdf?utm_source=chatgpt.com

[821] D. Chopra, S. M. Lee, and H. H. Su, On edge-balance index sets of fans and broken fans, Proceedings of the Fortieth Southeastern International Conference on Combinatorics, Graph Theory and Computing, *Congr. Numer.*, **196** (2009) 183-201.

[822] S. A. Choudum and S. P. M. Kishore, All 5-stars are Skolem graceful, *Indian J. Pure and Appl. Math.*, **27** (1996) 1101-1105. https://www.researchgate.net/publication/255910562_All_5-stars_are_Skolem_Graceful

[823] S. A. Choudum and S. P. M. Kishore, Graceful labellings of the union of paths and cycles, *Discrete Math.,* **206** (1999) 105-117. https://www.sciencedirect.com/science/article/pii/S0012365X98003951

[824] S. A. Choudum and S. P. M. Kishore, On Skolem gracefulness of k-stars, *Ramanujan Mathematical Society Inter. Conf. Discrete Math. and Number Theory*, 1996.

[825] S. A. Choudum and S. P. M. Kishore, Graceful labelling of the union of cycles and stars, unpublished.

[826] F. R. K. Chung and F. K. Hwang, Rotatable graceful graphs, *Ars Combin.,* **11** (1981) 239-250. https://mathweb.ucsd.edu/~fan/mypaps/fanpap/fc22rotatable.pdf

[827] P.-T. Chung and S. M. Lee, On the super edge-graceful spiders of even order, *J. Combin. Math. Combin. Comput.,* **64** (2008) 3-17. https://combinatorialpress.com/article/jcmcc/Volume%20064/vol-064-paper%201.pdf

[828] P.-T. Chung and S. M. Lee, A note on the edge-balance index sets for graphs, *Cong. Numer.,* **203** (2010) 105-129.

[829] P.-T. Chung, S. M. Lee, W.-Y. Gao, More trees of even orders which are super edge-graceful, *Congr. Numer.,* **206** (2010) 41-58.

[830] P.-T. Chung, S. M. Lee, W.-Y. Gao, and K. Schaffer, On the super edge-graceful trees of even orders, *Congr. Numer.,* **181** (2006) 5-17.

[831] T. Chunling, L. Xiaohui, Y. Yuansheng, and W. Liping, Irregular total labellings of some families of graphs, *Indian J. Pure Appl. Math.*, **40**(3)(2009) 155-181.

[832] P. Chuaephon, Sum divisor cordial labeling of bipartite, fan, and related graphs, *Internat. J. Applied Graph Th.*, **10**(1) (2026) 1-17. https://www.ijagt.com/upload/173088f277240ec61233e1b33c47c72b.pdf

[833] S. Cichacz, Group distance magic labeling of some cycle-related graphs, *Australas. J. Combin.,* **57** (2013) 235-243. https://ajc.maths.uq.edu.au/pdf/57/ajc_v57_p235.pdf

[834] S. Cichacz, Note on group distance magic complete bipartite graphs, *Central European J. Math.,* **12** (3) (2014) 529-533. https://link.springer.com/article/10.2478/s11533-013-0356-z

[835] S. Cichacz, Note on group distance magic graphs $G[C_4]$, *Graphs Combin.,* **30** (3) (2014) 565-571. doi:10.1007/s00373-013-1294-z https://link.springer.com/article/10.1007/s00373-013-1294-z

[836] S. Cichacz, Group distance magic graphs $G \times C_n$, *Discrete Appl. Math.* **177** (20) (2014) 80-87. doi:10.1016/j.dam.2014.05.044 https://arxiv.org/abs/1302.6561

[837] S. Cichacz, Distance magic (r,t)-hypercycles, *Util. Math.*, **101** (2016) 283-294. https://home.agh.edu.pl/~cichacz/preprints/MD_57.pdf

[838] S. Cichacz, A Γ-magic rectangle set and group distance magic labeling, *Combinatorial algorithms*, 122-127, Lecture Notes in Comput. Sci., 8986, Springer, Cham, 2015. doi:10.1007/978-3-319-19315-1_11

[839] S. Cichacz, On zero sum-partition of Abelian groups into three sets and group distance magic labeling, *Ars Math. Contemporanea,* **13**(2) (2017) 417-425. https://amc-journal.eu/index.php/amc/article/view/1054 zek

[840] S. Cichacz, Zero sum partition into sets of the same order and its applications, *Electronic J. Combin.,* **25**(1) (2018) #P1.20.

[841] S. Cichacz, A. Gömrlich, and A. Semaničová-Feňovčíková, Upper bounds on inclusive distance vertex irregularity strength, *Graphs Combin.,* **37**(6) (2021) 2713-2721.

[842] S. Cichacz, On some graph-cordial Abelian groups, *Disc. Math.*, **345**(5) (2022) 112815
https://www.sciencedirect.com/science/article/abs/pii/S0012365X22000218?via%3Dihub

[843] S. Cichacz, B. Freyberg, D. Fronček, Orientable ζ_n-distance magic graphs, *Discuss. Math. - Graph Theory,* (2019) 533-546. https://www.dmgt.uz.zgora.pl/publish/article.php?doi=2094

[844] S. Cichacz and D. Fronček, Distance magic circulant graphs, *Discrete Math.,* **339**(1) (2016) 84-94. http://www.ii.uj.edu.pl/documents/12980385/26042491/MD_71.pdf

[845] S. Cichacz and D. Fronček, E. Krop, and C. Raridan, Distance magic Cartesian products of graphs, *Discuss Math. Graph Theory,* **36** (2016) 299-308. doi:10.7151/dmgt.1852

[846] S. Cichacz, D. Fronček, I. Singgih, Vertex magic total labelings of 2-regular graphs, *Discrete math.,* **340** (1) (2017) 3117-3124. doi:10.1016/j.disc.2016.06.022

[847] S. Cichacz, D. Fronček, K. Sugeng, and S. Zhou, Group distance magic and antimagic graphs, *Acta Math. Sinica*, English Series **32**(10) (2016) 1159-1176. https://actamath.cjoe.ac.cn/Jwk_sxxb_en/EN/10.1007/s10114-016-4646-9

[848] S. Cichacz, D. Fronček, A. Khodkar, and W. Xu, Super edge-graceful paths and cycles, *Bull. ICA.* **57** (2009) 79-90. https://www.academia.edu/18955302/Super_edge_graceful_paths_and_cycles

[849] S. Cichacz, A. Görlich, Constant sum partition of set of integers and distance magic graphs, *Discuss. Math. Graph Theory,* **38**(1) (2018) 97-106. https://www.researchgate.net/publication/255749063_Constant_sum_partition_of_sets_integers_and_distance_magic_graphs

[850] S. Cichacz, A. Görich, and Z. Tuza, Cordial labeling of hypertrees, *Disc. Math.,* **313** (22) (2013) 2518-2524. doi:10.1016/j.disc.2013.07.025

[851] S. Cichacz and B. Krupińska, Group irregularity strength of disconnected graphs, arXiv:2306.11914

[852] S. Cichacz and M. Nikodem, Union of distance magic graphs, *Discuss. Math. Graph The.,* **37**(1) (2017) 239-249. https://www.dmgt.uz.zgora.pl/publish/article.php?doi=1932

[853] J. Clarisse, C. Prudencio, and R. F. Rulete, On Legendre product cordial labeling of some special graphs, *European J. Math. Stat.*, **6**(6) (2025) 12pp. https://www.eu-opensci.org/index.php/ejmath/article/view/14418/3351

[854] N. E. Clarke and A. Sanaei, Skolem labelling of generalised Dutch windmills, *Australasian J. Combin.,* **66**(3) (2016) 407-419. https://ajc.maths.uq.edu.au/pdf/66/ajc_v66_p407.pdf

[855] B. Clemens, R. Coulibaly, J. Garvens, J. Gonnering, J. Lucas, and S. Winters, An introduction to the gracefulness of one-point and two-point union of graphs, *Congr. Numer.*, **171** (2004) 167-171.

[856] A. Clifton and A. Khodkar Super edge-graceful labelings of kites, *J. Combin. Math. Combin. Comput.,* **102** (2017) 277-295. https://combinatorialpress.com/article/jcmcc/Volume%20102/vol-102-paper%2018.pdf

[857] Z. Coles, A. Huszar, J. Miller, and Z. Szaniszlo, 4-equitable tree labelings, *Congressus Numerantium* **228** (2017) 51-63. https://scholar.valpo.edu/cgi/viewcontent.cgi?article=1052&context=math_stat_fac_pubs

[858] N. Cohen and Kovše, Not all partial cubes are Θ-graceful, *Discrete Math.*, **343**(10) (2020) 112031 arXiv:1809.05320

[859] D. Combe, Magic labelings of graphs over finite abelian groups, *Austral. J. Comb.* **29** (2004) 259-271. https://ajc.maths.uq.edu.au/download/29/ajc_v29_p259.pdf

[860] A. Collins, C. Magnant, H. Wang, Tight super-edge-graceful labelings of trees and their applications, *AKCE Internat. J. Graphs Combin.,* **12** (2015) 113-118. https://ajc.maths.uq.edu.au/pdf/29/ajc_v29_p259.pdf

[861] C. Constantine and E. Suwastika, Graceful labeling of zero-divisor graph $\Gamma)p^2q)$ and $\Gamma(p^3q)$, Electronic Journal of Graph Theory and Applications 13 (2) (2025), 309–322 https://ejgta.org/index.php/ejgta/article/view/2143/pdf_334

[862] D. Craft and E. H. Tesar, On a question by Erdős about edge-magic graphs, *Discrete Math.*, **207** (1999) 271-276. https://www.sciencedirect.com/science/article/pii/S0012365X99001107

[863] D. W. Cranston, Regular bipartite graphs are antimagic, *J. Graph Theory,* **60** (2009) 173-182. https://www.researchgate.net/publication/227763356_Regular_Bipartite_Graphs_Are_Antimagic http://front.math.ucdavis.edu/1303.4580. arXiv:1303.4850.

[864] D. W. Cranston, Y. Liang and X. Zhu, Regular graphs of odd degree are antimagic, *J. Graph Theory*, **80**(1) (2015) 28-33. https://www.researchgate.net/publication/267457132_Regular_Graphs_of_Odd_Degree_Are_Antimagic

[865] S. J. Curran, Prime labelings on planar grid graphs, *Theory Appl. Graphs*, **9**(1), Article 6, (2022), 14 pages. https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1224&context=tag

[866] S. J. Curran, C_4-face-magic labelings on odd order projective grid graphs, *Australas. J. Combin.*, **83**(3) 361-396. https://ajc.maths.uq.edu.au/pdf/83/ajc_v83_p361.pdf

[867] S. J. Curran, Odd order C_4-face-magic projective grid graphs, arXiv:2206.03586.

[868] S. J. Curran and S. C. Locke, C_4-face-magic labelings on even order projective grid graphs, In: Hoffman, F., Holliday, S., Rosen, Z., Shahrokhi, F., Wierman, J. (eds) *Combin., Graph Th. Comput.*, (2024) SEICCGTC 2021. Springer Proceedings in Mathematics &Statis https://link.springer.com/chapter/10.1007/978-3-031-52969-6_17#Abs1

[869] S. J. Curran and S. C. Locke, C_4-face-magic labelings on even order projective grid graphs, In: Hoffman, F., Holliday, S., Rosen, Z., Shahrokhi, F., Wierman, J. (eds) Combinatorics, Graph Theory and Computing. SEICCGTC 2021. Springer Proceed. Math. Stat., **448** Springer, Cham.

[870] S. J. Curran and R. M. Low, C_4-face-magic torus labelings on $C_4 \times C_4$, *Cong. Numer.,* **233** (2019) 79-94. https://combinatorialpress.com/article/cn/Volume%20233/vol-233-paper%207.pdf

[871] S. J. Curran, R. M. Low, and S. C. Locke, C_4-face-magic labelings on Klein bottle grid graphs, arXiv:2206.02028 (2022).

[872] S. J. Curran, R. Low, and S. C. Locke, C_4-face-magic toroidal labelings on $C_m \times C_n$, *Art Discr. Appl. Math.,* **4** (2021) #P1.04, 33 page https://adam-journal.eu/index.php/ADAM/article/view/1368/1191

[873] S. J. Curran, R. M. Low and S. C. Locke, Equatorially balanced C_4-face-magic labelings on Klein bottle grid graphs, https://doi.org/10.48550/arXiv.2206.02028

[874] S. J. Curran and M. A. Ollis, A comment on Dean's construction of prime labelings on ladders, arXiv:2306.04088v1

[875] N. J. Cutinho, S. Sudha, and S. Arumugam, Distance antimagic labelings of Cartesian product of graphs, *AKCE Int. J. Graphs Comb.,* **17**(3) (2020) 940-942. https://www.tandfonline.com/doi/epub/10.1016/j.akcej.2019.08.005?needAccess=true

[876] J. A. Cynthia and P. Poorani, Square difference labeling of circulant network $G(n; \pm\{1, 2\}$, *Int. J. Pure Appl. Math.*, **109**(7) (2016) 109-113. https://www.acadpubl.eu/jsi/2016-109-si/7/14/14.pdf

[877] J. A. Cynthia and P. Poorani, Square difference labeling of torus network, *Int. J. Pure Appl. Math.*, **120**(8) (2018) 9-17. https://acadpubl.eu/hub/2018-120-8/1/2.pdf

[878] J. A. Cynthia and E. Padmavathy, Signed and signed product cordial labeling of cylinder graphs and banana tree, *Internat. J. Math. Trends Tech.,* **65**(3) (2019) 36-43. https://ijmttjournal.org/public/assets/volume-65/issue-3/IJMTT-V65I3P505.pdf

[879] V. J. A. Cynthia and E. Padmavathy, Signed cordial labelings and signed product labeling of some interconnected netwohks, *Adv. Appl. Discr. Math.,* **26**(1) (2021) 35-51. https://www.pphmj.com/abstract/13630.htm

[880] J. A. Cynthia, P. Poorani, and M. H. Mercy, Square difference labeling of bloom graph, *J. Adv. Comput. Intell. Intell. Informatics,* **27**(6) (2023) :1108-1112. https://www.jstage.jst.go.jp/article/jaciii/27/6/27_1108/_pdf/-char/en

[881] J. Cyrile, G. Valdehueza, and A. C. Pedrano, On the total edge product cordial labeling of some corona graphs, *Internat. J. Math. Trends Tech.,* **67**(10) (2021) 1-6. arXiv:2111.08838 https://www.researchgate.net/publication/356293406_On_the_Total_Edge_Product_Cordial_Labeling_of_Some_Corona_Graphs

[882] S. Czerwiński, J. Grytczuk, and W. Zelazny, Lucky labelings of graphs, *Infor. Process. Let.,*, **109**(18) (2009) 1078-1081. https://www.sciencedirect.com/science/article/pii/S0020019009002117?via%3Dihub

[883] Dafik, A. I. Kristiana, S. Setiawani, and K. M. F. Azizah, Generalized shackle of fans is a super (a, d)-edge-antimagic total graph, *J. Graph Label.,* **2** (1) (2016) 59-68.

[884] Dafik, M. Miller, J. Ryan, and M. Bača, On antimagic labelings of disjoint union of complete s-partite graphs, *J. Combin. Math. Combin. Comput.,* **65** (2008) 41-49. https://www.researchgate.net/publication/262698625_On_antimagic_labelings_of_disjoint_union_of_complete_s-partite_graphs

[885] Dafik, M. Miller, J. Ryan, and M. Bača, On super (a, d)-edge-antimagic total labelings of disconnected graphs, *Discrete Math.,* **309** (2009) 4909-4915. https://core.ac.uk/reader/82595010

[886] Dafik, M. Miller, J. Ryan, and M. Bača, Super edge-antimagic labeling of $mK_{n,n,n}$, *Ars Combin.,* **101** (2011) 97-107. https://www.researchgate.net/publication/235903025_Super_edge-antimagic_total_labelings_of_mKn

[887] Dafik, M. Miller, J. Ryan, and M. Bača, Antimagic labelings of the union of two stars, *Australas. J. Combin.,* **42** (2008) 35-44. https://ajc.maths.uq.edu.au/pdf/42/ajc_v42_p035.pdf

[888] Dafik, Slamin, F. Eka, and L. Sya'diyah, Super antimagicness of triangular book and diamond ladder graphs, *Proceed. of IICMA 2013 Research Topic*, 8 pp. https://www.researchgate.net/publication/290998038_Super_Antimagicness_of_Triangular_Book_and_Diamond_Ladder_Graphs

[889] Dafik, S. Slamin, R. W. Romdhani, and I. Y. Arianti, On super antimagicness of generalized flower and disk brake graphs, *Util. Math.,* **100** (2016) 127-135.

[890] Dafik, Slamin, D. Tana, A. Semaničová-Feňovčíková,and M. Bača, Constructions of H-antimagic graphs using smaller edge-antimagic graphs, *ArsComb.,* **133** (2017) 233-245.

[891] H. Dao, H. Hua, M. Ngo, and C. Raridan, On the edge-balanced index sets of complete even bipartite graphs, *J. Graph Labeling,* **2**(1) (2016) 79-87. arXiv:1509.01841

[892] H. Dao, H. Hua, and C. Raridan, The conclusion to the edge-balance index set problem for complete bipartite graphs, In book: Combinatorics, Graph Theory and Computing, (2025) 95-106.

[893] S. N. Daoud, edge-odd-graceful labeling of some path and cycle related graphs, *AKCE Int. J. Graphs Comb.*, **14** (2017) 178–203. https://www.tandfonline.com/doi/pdf/10.1016/j.akcej.2017.03.001

[894] S. N. Daoud, Edge odd graceful labeling of cylinder and torus grid graphs, *IEEE*, **7** (2019) 10568-10592. https://ieeexplore.ieee.org/document/8587242

[895] S. N. Daoud, Edge even graceful labeling of polar grid graphs, *Symmetry,* **11**(38):22 (2019) 1-28. https://www.mdpi.com/2073-8994/11/1/38

[896] S. N. Daoud, Vertex odd graceful labeling, *Ars Combin.,* **142** (2019) 65-87.

[897] R. Darda and A. Hujdurović, On bounds for the product irregularity strength of graphs, *Graphs and Combin.*, **31**(5) (2015) 1347-1357. https://link.springer.com/article/10.1007/s00373-014-1458-5

[898] D. Darmaji, S. Wahyudi, R. Rinurwatia, and S. W. Saputro, On the construction of super edge-magic total graphs, *Elect. J. Graph Th. Appl.,* **10**(1) (2022) 301-309. http://dx.doi.org/10.5614/ejgta.2022.10.1.21

[899] W. S. Darmaji and S. W. Saputro, On the construction of super edge-magic total graphs, *Electron. J. Graph Theory Appl.*, (EJGTA) **10**(1) (2022) 301-309. http://dx.doi.org/10.5614/ejgta.2022.10.1.21

[900] S. Das, S. Ghosh, S. Nandi, S. Sen, A lower bound technique for radio k-coloring, *Discrete Math.*, **340** (2017) 855-861. https://www.sciencedirect.com/science/article/pii/S0012365X16304320

[901] T. Das, T. Lehtilä, S. Nandi, S. Sen, and D. K. Supraja, On radio k-labeling of the power of the infinite path, *Inform. Process. Lett.*, **182** (2023) 106386. https://homepages.iitdh.ac.in/~sen/Supraja_ICGT.pdf

[902] S. S. Dash, A. K. Behera, and D. P. Bhatta, Some explorations on odd-even graceful labeling of graphs, *J. Disc. Math. Sci. Cryp.*, **29**(8) (2026) 2989–2999. https://tarupublications.com/journals/jdmsc/volume/29/issue/8/some-explorations-on-odd-even-graceful-labeling-of-graphs/pdf

[903] G. K. Dayanand and S. Ahmed, The super magic properties of connected and disconnected graphs, *J. Inf. Optim. Sci.,* **36**(3) (2015) 231-246.

[904] G. K. Dayanand, K. Ahmed, and D. G. Akka, Cahit-8-equitablity of coronas $C_n \cdot K_1$, *IJRET: Internat. J. Res. Engin. Tech.,* **03** Special Issue: 03 (2014) 956-960. https://ijret.org/volumes/2014v03/i15/IJRET20140315181.pdf

[905] N. Dean, Proof of the prime ladder conjecture, *Integers,* **A40** (2017) 9 pages. https://math.colgate.edu/~integers/r40/r40.pdf

[906] P. de la Torre, R. Greenlaw, and T. M. Przytycka, Optimal tree ranking is in NC, *Parallel Process. Lett.*, **2** (1992) 31-41. https://www.researchgate.net/publication/220439635_OPTIMAL_TREE_RANKING_IS_IN_mathcalNC

[907] G. K. Dayanand and S. Ahmed, The super magic properties of connected and disconnected graphs, *J. Inf. Optim. Sci.,* **36**(3) (2015) 231-246.

[908] Daili, Z-H Wang, and Z. Xie, Gracefulness of $P_{2r,2m}$, *Ars Combin.,* **104** (2012) 185-191.

[909] J. W. Daykin, · C. S. Iliopoulos, ·M. Miller, and · O. Phanalasy, Antimagicness of generalized corona and snowflake graphs, *Math. Comput. Sci.,* **9**(1) (2015) 105-111. https://link.springer.com/content/pdf/10.1007/s11786-014-0213-x.pdf?pdf=button

[910] D. L. de Oliveira, D. Artigas, S. Dantas, L. Frickes, and A. G. Luiz, Graceful game on some graph classes, *RAIRO - Operat. Res.*, (2023). https://www.rairo-ro.org/articles/ro/pdf/2024/01/ro220747.pdf

[911] D. L. de Oliveira, D. Artigas, S. Dantas, and A. Luiz, Bounds on the edge-sum distinguishing game, *Mate. Contempornea,* **55**(8) (2023) 67-76. doi:DOI: 10.21711/231766362023/rmc558
https://mc.sbm.org.br/wp-content/uploads/sites/9/sites/9/2023/12/Article-7.pdf

[912] D. L. de Oliveira, S. Dantas, and A. G. Luiz, Results on the graceful game and range-relaxed graceful game, In: J. Nešetřil, G. Perarnau, J. Rue, and O. Serra, O. Extended Abstracts EuroComb 2021. *Trends Math.,* (2021) **14** Birkhauser, Cham. https://link.springer.com/chapter/10.1007/978-3-030-83823-2_34

[913] P. Deb and N. B. Limaye, On elegant labelings of triangular snakes, *J. Combin. Inform. System Sci.,* **25** (2000) 163-172.

[914] P. Deb and N. B. Limaye, Some families of elegant and harmonious graphs, *Ars Combin.*, **61** (2001) 271-286. https://combinatorialpress.com/article/ars/Volume%20061/volume-61-paper-24.pdf

[915] P. Deb and N. B. Limaye, On harmonious labelings of some cycle related graphs, *Ars Combin.,* **65** (2002) 177-197. https://combinatorialpress.com/article/ars/Volume%20065/volume-65-paper-15.pdf

[916] M. R. Z. El Deen, I. M. Hanafy, S. G. Shehda, and N. A. Omar, Super edge bimagic and trimagic total labeling for some graphs, *Alfarama J. Basic App. Sci.*, **6**(IV) (2025) 431-432. https://ajbas.journals.ekb.eg/article_449211_7230705e8199f04ea1335a053ee67320.pdf

[917] B. Deepa and M. Veerasamy, An enhanced DNA structure for one-time pad together with graph labeling techniques, *AIP Confer. Proceed.,* 2385, 130045 (2022) 1-9. https://aip.scitation.org/doi/pdf/10.1063/5.0071283

[918] N. S. Deepthi, Super edge-magic total strength of some unicyclic graphs, *Ars Combin.*, **159** (2024) 73-85. https://combinatorialpress.com/article/ars/super-edge-magic-total-strength-of-some-unicyclic-graphs.pdf

[919] A. Delman, S. Koilraj, and P. L. Raj, A neighbourhood prime labeling of some special graphs, *Inter. J. Pure Appl. Math.,* **120**(4) (2018) 601-607. https://ijpam.eu/contents/2018-120-4/10/10.pdf

[920] C. Delorme, Two sets of graceful graphs, *J. Graph Theory*, **4** (1980) 247-250.

[921] C. Delorme, M. Maheo, H. Thuillier, K. M. Koh, and H. K. Teo, Cycles with a chord are graceful, *J. Graph Theory*, **4** (1980) 409-415.

[922] K. Deng, Bipartite graphs with minimum degree at least 15 are antimagic, *J. Graph Th.* (2026). arXiv:2507.17302v2

[923] K. Deng and Y. Li, Caterpillars with maximum degree 3 are antimagic, *Discrete Math.,* **342**(6) (2019) 1799-1801. https://www.sciencedirect.com/science/article/pii/S0012365X19300792

[924] Y. X. Deng, Harmoniousness of the graphs $C_n + \overline{K}_t$, *J. Math. Res. Exposition*, **15** (1995), suppl., 79-81.

[925] G. Denham, M. G. Leu, and A. Liu, All 4-stars are Skolem graceful, *Ars Comb.,* **36** (1993) 183-191. https://combinatorialpress.com/article/ars/Volume%20036/volume-36-paper-17.pdf

[926] D. Dereniowski, Rank Coloring of Graphs, in: M. Kubale (Ed.), Graph Colorings, *Contemporary Mathematics,* **352**, Amer. Math. Soc., (2004) 79-93.

[927] T. Deretsky, S. M. Lee, and J. Mitchem, On vertex prime labelings of graphs, in *Graph Theory, Combinatorics and Applications Vol. 1*, J. Alavi, G. Chartrand, O. Oellerman, and A. Schwenk, eds., Proceedings 6th International Conference Theory and Applications of Graphs (Wiley, New York, 1991) 359-369.

[928] U. Derings and H. Hünten, Magic graphs - A new characterization, Report No. 83265 - OR, Universität Bonn April 1983, ISSN 0724-3138.

[929] D. N. Desai and R. Wang, New developments on graph sum index, *Graphs and Combin.*, **42** (2026) 37 https://arxiv.org/abs/2410.16494?utm_source=chatgpt.com

[930] U. Deshmukh, Skolem graceful graphs - A survey, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 165-170.

[931] U. Deshmukh and V. Y. Shaikh, Mean cordial labelling of some star-related graphs, *Internat. J. Math. Combin.*. **3** (2016) 146-157. https://fs.unm.edu/IJMC/MeanCordialLabellingOfSomeStarRelatedGraphs.pdf

[932] D. Devakirubanithi and J. Jeba Jesintha, Odd graceful labeling of few graphs, *Internat. J. Engin. Res. Develop.,* **20**(3) (2024) 49-55. https://www.ijerd.com/paper/vol20-issue3/G20034955.pdf

[933] D. Devakirubanithi, J. Jeba Jesintha, and M. V. Abigail, Felicitous labeling of varied bistar, braid and globe graph, *Indust. Engin. J.*, (2023) **52**(4), No. 2. 155-160. https://www.researchgate.net/publication/374233992_FELICITOUS_LABELING_OF_VARIED_BISTAR_BRAID_AND_GLOBE_GRAPH

[934] D. Devakirubanithi, J. Jeba Jesintha, and B. D. Benita, Odd harmonious labeling of arrow and $T_p \odot K_{1,n}$ graphs, *Indust. Engin. J.* (2023) **52**(4) No. 2. 142-148. https://www.researchgate.net/publication/374234143_ODD_HARMONIOUS_LABELING_OF_ARROW_AND_GRAPHS

[935] D. Devakirubanithi, J. Jeba Jesintha, and R. Charishma, Cordial labeling on families of blade graph, *Sci. Math. Japonicae,* **85**(3) (2023) 161-170. https://www.researchgate.net/publication/374234424_CORDIAL_LABELING_ON_FAMILIES_OF_BLADE_GRAPH

[936] J. Devaraj, On edge-cordial graphs, *Graph Theory Notes of New York,* **XLVII** (2004) 14-18.

[937] J. Devaraj, On consecutive labelings of ladder graphs, *Bull. Pure Appl. Sci. Sect. E Math. Stat.,* **26** (2007) 1-10.

[938] J. Devaraj and P. Delphy, On signed cordial graph, *Int. J. Math. Sci. Appl.*, (2011) **1**(3) (2011) 1156-1167. https://ijmsa.yolasite.com/resources/14--sep.pdf

[939] B. Devi, E. Bernie, S. Faridha, and N. Rajeshwaran, Total PD mean edge-balanced labeling of graphs, *Internat. J. Appl. Math.*, **38**(12s) (2025). https://ijamjournal.org/ijam/publication/index.php/ijam/article/view/1648/1499

[940] M. U. Devi, M. Kamaraj, and S. Arockiaraj, Odd Fibonacci edge irregular labeling for some simple graphs, *J. Alg. Stat.*, **13**(3) (2022) 13(3) 1230-1238. https://www.publishoa.com/index.php/journal/article/view/742

[941] M. U. Devi, M. Kamaraj, and S. Arockiaraj, Odd Fibonacci edge irregular labeling for some trees obtained from subdivision and vertex identification operations, *Baghdad Sci. J.*, **20**(1)(SI)) (2023) 332-338.

[942] S. A. Devi and R. S. Porchelvi, Pronic graceful labeling on special cases of generalized Petersen graphs,*Wesleyan J. Res.,* **13**(58) (2021) 86-89. https://dx.doi.org/10.21123/bsj.2023.8420

[943] S. K. Devi and B. Gayathri, k-even edge-graceful labeling of the graph $P_n \otimes K_{1,m}$, *Bull. Pure Appl. Sci.,* **29 E**(2) 387-407.

[944] S. A. Devi, R. S. Porchelvi, and T. Arjun, Pronic Heron mean labeling on special cases of generalized Petersen graph $P(n,k)$ and disconnected graphs, *Malaya J. Matematik*, *9*(1) (2021) 262-266. https://doi.org/10.26637/MJM0901/0044

[945] J. DeVito, A. Niedzialomski, and J. Warren, Radio number of Hamming graphs of diameter 3, *Theory Appl. Graphs*, **9**(2) (2022) Art. 10, 13 pp. https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1166&context=tag

[946] E. Dhananjaya and W.-T. Li, Antimagic labeling of forests with sets of consecutive integers, *Discrete Appl. Math.,* **309** (2022) 75-84. https://www.sciencedirect.com/science/article/pii/S0166218X21004546?via%3Dihub

[947] E. Dhananjaya and W.-T. Li, The antimagic orientations for graphs obtained by some graph operations, *Taiwanese J. Math.,* **29**(4) (2025) 687-703. https://arxiv.org/pdf/2012.10087

[948] S. Dhanalakshmi and N. Parvathi, Mean square cordial labeling related to some cyclic graphs, *Indian J. Engin. Tech.*, **7**, no. 4.10, (2018), 797–800. https://www.researchgate.net/publication/332681831_Mean_Square_Cordial_Labeling_of_Some_Cycle_Related_Graphs

[949] S. Dhanalakshmi and N. Parvathi, Mean square cordial labeling related to some acyclic graphics and its rough approximations, *J. Physics: Conference Series*, 1000, (2018), 012040 doi:10.1088/1742-6596/1000/1/01204.

[950] S. Dhanalakshmi and N. Parvathi, Mean square cordial labeling on star related graphs, *J. Physics: Conference Series,* **1377** (2019) 012027 14 pages, doi:10.1088/1742-6596/1377/1/012027.

[951] S. Dhanalakshmi, Mean square cordial labeling of a shell-butterfly graph, *Internat. J. Math. Comp Sci.,* **15**(4), (2020) 1235–1242, http://ijmcs.future-in-tech.net/15.4/R-Dhanalakshmi.pdf

[952] S. Dhanalakshmi1 and S. Thirunavukkarasu, Mean square cordial labeling of some pentagonal snake graphs, *Proceed. First Internat. Confer. Comput. Communication and Control Sys.*, I3CAC 2021, 7-8 June 2021, Bharath University, Chennai, India.
https://eudl.eu/doi/10.4108/eai.7-6-2021.2308859

[953] A. Dharman and S. Murugesan, Python programming on Fibonacci graceful anti-magic labeling for star graphs, *Internat. J. Res. Appl. Sci. Engin. Tech.*, **13**(1) (2025) 1923-1928. https://www.ijraset.com/best-journal/python-programming-on-fibonacci-graceful-anti-magic-labeling-for-star-graphs-

[954] A. Dharman and O. V. Shanmuga Sundaram, Fibonacci graceful labelings for some special graphs, *Advan. Engin. Sci.*, (2022). https://www.researchgate.net/publication/374053745_FIBONACCI_GRACEFUL_LABELING_FOR_SOME_SPECIAL_GRAPHS

[955] A. T. Diab, Study of some problems of cordial graphs, *Ars Combin.,* **92** (2009) 255-261. https://combinatorialpress.com/article/ars/Volume%20092/volume-92-paper-18.pdf

[956] A. T. Diab, On cordial labelings of the second power of paths with other graphs, *Ars Combin.,* **97A** (2010) 327-343. https://combinatorialpress.com/article/ars/Volume%20097-A/volume-97-A-paper-26.pdf

[957] A. T. Diab, On cordial labelings of wheels with other graphs, *Ars Combin.,* **100** (2011) 265-279. https://combinatorialpress.com/article/ars/Volume%20100/volume-100-paper-22.pdfm

[958] A. T. Diab, Generalization of some results on cordial graphs, *Ars Combin.,* **99** (2011) 161-173. https://combinatorialpress.com/article/ars/Volume%2099/volume-99-paper-16.pdf

[959] A. T. Diab, On cordial labelings of the second power of cycles with other graphs, *Util. Math.,* **97** (2015) 65-84. https://combinatorialpress.com/article/ars/Volume%20097-A/volume-97-A-paper-26.pdf

[960] A. T. Diab and E. Elsakhawi, Some results on cordial graphs, *Proc. Math. Phys. Soc. Egypt,* No. 77 (2002) 67-87.

[961] A. T. Diab and S. A. E. Mohammed, On cordial labelings of fans with other graphs, *Ars Combin.,* **106** (2012) 263-275. https://combinatorialpress.com/article/ars/Volume%20106/volume-106-paper-23.pdf

[962] A. T. Diab and S. Nada, Some results on specific graphs, *Ars Combin.* **132** (2017) 93-103. https://combinatorialpress.com/article/ars/Volume%20132/volume-132-paper-8.pdf

[963] N. Diefenderfer, M. Hastings, L. N. Heath, H. Prawzinsky, B. Preston, J. Rushall, E. White, and A. Whittemore, Prime vertex labelings of families of unicyclic graphs, *Rose-Hulman Undergrad. Math J.,* **16**(1) (2015) 253-269. https://scholar.rose-hulman.edu/cgi/viewcontent.cgi?article=1027&context=rhumj

[964] N. Diefenderfer, D. C. Ernst, M. G. Hastings, L. N. Heath, H. Prawzinsky, B. Preston, E. White, and A. Whittemore, Prime vertex labelings of several families of graphs, *Involve,* **9**(4) (2016) 667-688. https://www.researchgate.net/publication/290810930_Generalized_Shackle_of_Fans_is_a_Super_ad-Edge-Antimagic_Total_Graph

[965] J. Dinitz and P. Rodney, Disjoint difference families with block size 3, *Util. Math.,* **52** (1997) 153-160. https://site.uvm.edu/jdinitz/files/2021/04/disjoint.pdf

[966] D.M.T.B. Dissanayake, R.A.S.T. Abeysekara, K.D.E. Dhananjaya, A.A.I. Perera, and P.G.R.S. Ranasinghe, Prime labeling of complete tripartite graphs of the form $K_{1,m,n}$, *Elixir Appl. Math.*, **130** (2019) 53092-53094. https://www.elixirpublishers.com/articles/1673250759_201905001.pdf

[967] J. Doma, Unicyclic Graceful Graphs, M. S. Thesis, McMaster Univ., 1991.

[968] D. Donovan, S. El-Zanati, C. Vanden Eynden, and S. Sutinuntopas, Labelings of unions of up to four uniform cycles, *Australas. J. Combin.,* **29** (2004) 323-336. https://www.researchgate.net/publication/268073557_Labelings_of_unions_of_up_to_four_uniform_cycles

[969] M. Doob, On the construction of magic graphs, *Congr. Numer.,* **10** (1974) 361-374.

[970] M. Doob, Generalizations of magic graphs, *J. Combin. Theory, Ser. B,* **17** (1974) 205-217. https://www.sciencedirect.com/science/article/pii/0095895674900276

[971] M. Doob, Characterizations of regular magic graphs, *J. Combin. Theory, Ser. B*, **25** (1978) 94-104. https://www.sciencedirect.com/science/article/pii/S0095895678800136

[972] W. Dou, On the mod sum number of $H_{m,n}$, *J. Comb. Optim.,* **26**(3) (2013) 465-471. https://link.springer.com/article/10.1007/s10878-011-9432-0

[973] W. Dou and J. Gao, The (mod, integral) sum numbers of fans and $K_{n,n} - E(nK_2)$, *Discrete Math.,* **306** (2006) 2655-2669. https://www.sciencedirect.com/science/article/pii/S0012365X06003906

[974] W. Dou and J. Gao, Some results on mod (integral) sum graphs, *Ars Combin.,* **82** (2007) 3-31. https://combinatorialpress.com/article/ars/Volume%20082/volume-82-paper-1.pdf

[975] C. Dorathy and N. Nagamani, Graceful labeling on cycle related graphs, In: *Futuristic Trends in Contemporary Mathematics & Applications*, V3B4, IIP Series, Iterative Internat. Publ. (IIP), **3** (2024) 210-220. doi:10.58532/V3BBCM4P2CH10

[976] S. Drajnová, J. Ivančo, A. Semaničová, Numbers of edges in supermagic graphs, *J. Graph Theory,* **52** (2006) 15-26. https://onlinelibrary.wiley.com/doi/epdf/10.1002/jgt.20144

[977] G. M. Du, Cordiality of complete k-partite graphs and some special graphs, *Neimenggu Shida Xuebao Ziran Kexue Hanwen Ban*, (1997) 9-12.

[978] G. M. Du, On the cordiality of the union of wheels, *J. Inn. Mong. Norm. Univ. Nat. Sci.,* **37** (2008) 180-181, 184.

[979] Z.-t. Du and H.-q. Sun, The gracefulness of the digraph $n \cdot \vec{C}_{2p}$, *J. Beijing Univ. Posts and Telecommunications,* **17** (1994) 85-88.

[980] G. Duan and Y. Qi, k-gracefulness of two classes of graphs, (Chinese) *J. Zhengzhou Univ. Nat. Sci. Ed.,* **34** (2002) 36-38, 49.

[981] M. Dufour, Sur la Decomposition d'un Graphe Complet en Arbres Isomorphes, Ph.D. Thesis, Universite de Montreal, 1995.c

[982] A. Durai Baskar and S. Arockiaraj, F-geometric mean graphs, *Appl. Appl. Math.,* **10**(2) (2015) 937-952 https://www.pvamu.edu/mathematics/wp-content/uploads/sites/49/20_r766_arockiaraj_aam_r766_sa_110114_ready_amh.pdf

[983] A. Durai Baskar, S. Arockiaraj, and B. Rajendran, F-geometric mean labeling of some chain graphs and thorn graphs, *Kragujevac J. Math.* **37**(1) (2013) 163-186. https://imi.pmf.kg.ac.rs/kjm/pub/13681225790724_geometric_meanness_of_some_chain_graphs_and_thorn_graphs-kjm.pdf

[984] A. Durai Baskar, S. Arockiyaraj, and B. Rajendran, Geometric meanness of graphs obtained from paths, *Util. Math.,* **101** (2016) 45-68.

[985] A. Durai Baskar and S. Arockiaraj, C-Geometric mean labeling of some cycle related graphs, *Internat. J. Math. Combin.* **2** (2019) 69-85. https://search.proquest.com/openview/16137f5f82c5f19a9ad9700490d080e5/1?cbl=51283&pq-origsite=gscholar

[986] K. Driscoll, E. Krop, and M. Nguyen, All trees are six-cordial, *Elect. J. Graph Theory Appl.*, **5**(1) (2016) 21-35. https://www.researchgate.net/publication/301875801_All_trees_are_six-cordial

[987] K. Driscol, All trees are seven-cordial, arXiv:1909.12351.

[988] A. Dsouza, Saumya Y M, Kumudakshi, Antimagic labeling of digraphs, *Math. Stat.* **13**(2) (2025) 105-109. doi:DOI: 10.13189/ms.2025.130205

[989] A. Durai Baskar, S. Arockiyaraj, and B. Rajendran, Geometric meanness of graphs obtained from paths, *Util. Math.,* **101** (2016) 45-68.

[990] B. D'Souza and J. Pereira, Characterizations of additively graceful signed paths and cycles, *Commum. Combin. Optim.*, (2025), to appear.

[991] B. D'Souza and J. Pereira, Some results on additively graceful signed stars and double stars, https://doi.org/10.48550/arXiv.2604.21135 15 pp.

[992] B. D'Souza and J. Pereira, Additively graceful signed graphs on complete graphs and some open problems resolved, *Ars Combin.*, **167** (2026) 29-43. https://doi.org/10.61091/ars167-02

[993] D. Dyer, I. Payne, N. Shalaby, and B. Wicks, On the graceful conjecture for triangular cacti, *Australas. J. Combin.,* **53** (2012) 151-170. https://ajc.maths.uq.edu.au/pdf/53/ajc_v53_p151.pdf

[994] P. Dyrlaga, K. Szopa, Orientable Z_n-distance magic regular graphs, arXiv:1712.02676 [math.CO] (2017).

[995] E. Ebrahem, S. Hoory, and D. Kotlar, An infinite family of counterexamples to a conjecture on distance magic labeling, arXiv:2401.00807v2 [math.CO] 14 pp. arXiv:2401.00807v2

[996] T. Eccles, Graphs of large linear size are antimagic, *J. Graph Theory,* **81**(3) (2016) 236-261. arXiv:1409.3659

[997] M. Edwards and L. Howard, A survey of graceful trees, *Atlantic Elect. J. Math.,* **1** (2006) 5-30. http://euclid.trentu.ca/aejm/V1N1/Edwards.v1n1.pdf

[998] R. B. Eggleton, P. Erdős, and D. K. Skilton, Colouring the real line, *J. Combin. Theory Ser. B*, **39**(1) (1985) 86-100. https://reader.elsevier.com/reader/sd/pii/0095895685900395?token=7F112C8B1EEB8BA4D4551FDFBF02251A3D021F203348A41135D32AF968D5B36A75C81BBEDE7C065963originRegion=us-east-1&originCreation=20220727122943

[999] S. Ehard, S. Glock, and F. Joos, A rainbow blow-up lemma for almost optimally bounded edge-colourings, arXiv:907.09950 [math.CO], 28 pages. https://onlinelibrary.wiley.com/doi/epdf/10.1002/rsa.20907

[1000] P. Eldergill, Decomposition of the Complete Graph with an Even Number of Vertices, M. Sc. Thesis, McMaster University, 1997.

[1001] A. A. El-hay, K. A. Alsatami, A. ELrokh, and A. Rabie, Cordial labeling of corona product of paths and fourth order of lemniscate graphs, *European J. Prue Appl. Math.* **18**(1) Article Number 5470 (2025) 1-14. https://www.ejpam.com/index.php/ejpam/article/view/5470

[1002] A. A. El-hay, K. A. Alsatami, and A. ELrokh, A novel problem and algorithm for solving cordial cabeling of some fifth powers of graphs *European J. Pure Appl. Math.*, **18**(1) Article Number 5812, (2025). https://www.ejpam.com/index.php/ejpam/article/view/5812 1-14.

[1003] A. A. El-hay, Y. Elmshtaye, and A. Elrokh, Solving singed product cordial labeling of corona products of paths and the third power of lemniscate graphs, *Turkish J. Comput. Math. Edu.,* (TURCOMAT) **14**(2) (2023)) 806-823. https://www.ejpam.com/index.php/ejpam/article/view/5812

[1004] E. F. Elizeche and A. Tripathi, A characterization of the sum and integral sum labellings of some classes of graphs, *Integers* **20** (2020) Paper No. A33, 24 pp. http://math.colgate.edu/~integers/u33/u33.pdf

[1005] M. N. Ellingham, Sum graphs from trees, *Ars Combin.*, **35** (1993) 335-349. https://combinatorialpress.com/article/ars/Volume%20035/volume-35-paper-33.pdf

[1006] H. Elmorsy, Study on labeling matrix for super vertex-magic total labeling of cycle and complete graphs, *Adv. Dynamical Sys. Appl.*, **18**(1) (2023) 41-50, https://www.ripublication.com/adsa23/v18n1p03.pdf

[1007] Y. Elmshtaye, A. Elrokh, and A. A. El-hay, Total cordial for the corona product of paths and the third power of double fans-generalized fans, *Advances and Appl. Disc. Math.*, **42**(4) (2025) 303-334. https://doi.org/10.17654/0974165825020

[1008] A. Elrokh, K. A. Alsatami, and R. Hamza, On 1-Near mean cordial labeling of the strong product of two paths: characterization and a verification algorithm, *J. Math.*, (2026) 3959342. https://doi.org/10.1155/jom/3959342

[1009] A. Elrokh, M. M. A. Al-Shamiri, M. M. A. Almazah, and A. El-hay, A novel problem for solving permuted cordial labeling of graphs, *Symmetry* **15**(4) (2023) 825. https://www.researchgate.net/publication/369629154_A_Novel_Problem_for_Solving_Permuted_Cordial_Labeling_of_Graphs

[1010] A. Elrokh, M. M. Ali Al-Shamiri, A. A. El-hay, A novel problem for solving cordial labeling of corona product between path and third order of cone graphs, *Turkish J. Comput. Math. Edu.,* **13**(3) (2022) 970-986. https://www.researchgate.net/publication/366138234_A_NOVEL_PROBLEM_FOR_SOLVING_CORDIAL_LABELING_OF_CORONA_PRODUCT_BETWEEN_PATH_AND_THIRD_ORDER_OF_CONE_GRAPHS

[1011] A. Elrokh, M. M. Ali Al-Shamiri, S. Nada, and A. A. El-hay, Cordial and total cordial labeling of corona product of paths and second order of lemniscate graphs, *J. Math.*, (2022), Article ID 8521810, 9 pages. https://www.hindawi.com/journals/jmath/2022/8521810/

[1012] A. Elrokh, E. Badr, M. Al-Shamiri, and S. Ramadhan, Upper bounds of radio number for triangular snake and double triangular snake graphs, *J. Math.,* 2022, Art. ID 3635499, 8 pp. https://www.hindawi.com/journals/jmath/2022/3635499/

[1013] A. Elrokh and S. Elkom, The cordial labeling for the four-leaved rose graph, *Appl. Comput. Math.,* **7**(4) (2018) 203-211. doi:10.11648/j.acm.20180704.14

[1014] A. I. Elrokh, Y. Elmshtaye, and Abd El-hay, The cordiality of cone and lemniscate graphs, *Appl. Math. Inform. Sci.* **16**(6) (2022)) 1027-1034. https://www.naturalspublishing.com/files/published/3rb6lq6p83nq33.pdf

[1015] A. Elrokh, T. V. Sateesh Kumar, and S. Meenaksh, Lucky labeling of jelly fish graph $J(m, n)$, cocktail party graphs, and crowns. *AIP Confer. Proceed. 2516*, 210038 (2022). https://doi.org/10.1063/5.0108660

[1016] A. Elrokh, S. Nada, and E. El-Shafey, Cordial labeling of corona product of path graph and second power of fan graph, *Open J. Discr. Math.*, **11** (2021) 31-42. https://www.scirp.org/journal/paperinformation.aspx?paperid=108431

[1017] A. Elrokh and A. Rabie, The cordiality of the sum and union of two fourth power of paths and cycles, *J. Egypt Math. Soc.,* (2021) **29**(3), 1-13 https://doi.org/10.1186/s42787-020-00111-6

[1018] A. N. Elsawy and R. N. Almohammadi, Paley, cubic Paley, quadruple Paley, and generalized Paley graphs with an edge graceful labeling, *European J. Pure Appl. Math.*, **18**(4) (2025) Article Number 6600. https://www.ejpam.com/index.php/ejpam/article/view/6600

[1019] A. Elsonbaty and S. N. Daoud, Edge even graceful labeling of some path and cycle related graphs, *Ars Combin.*, **30** (2017) 79-96. https://www.researchgate.net/publication/315116297_Edge_Even_Graceful_Labeling_of_Some_Path_and_Cycle_Related_Graphs

[1020] A. Elsonbaty and K. Mohamed, On some variants of gracefulness of cycle graphs, *Ars Combin.,* **135** (2017) 39-50. https://combinatorialpress.com/article/ars/Volume%20135/volume-135-paper-4.pdf?utm_source=chatgpt.com

[1021] A. Elsonbaty and S. Nagy Daoud, Edge even graceful labeling of cylinder grid graph, *Symmetry* (2019), 11, 584. 30 pages. https://www.mdpi.com/2073-8994/11/4/584

[1022] A. Elumalai, On Graceful, Cordial and Elegant Labelings of Cycles Related and Other Graphs, Ph.,D. dissertation, Anna University, 2004, Chennai, India.

[1023] A. Elumalai, Graceful and felicitous labeling of stem-lotus graph, *Malaya J. Mat.,* **11**(04) (2023) 465–469. http://doi.org/10.26637/mjm1104/010

[1024] A. Elumalai and G. Sethuraman, Cordialness of cycles with parallel P_k-chords and multiple subdivision graphs, *Ars Combin.,* **85** (2007) 85-98. https://combinatorialpress.com/article/ars/Volume%20085/volume-85-paper-8.pdf

[1025] A. Elumalai and G. Sethuraman, Gracefulness of union of cycle with parallel chords and complete bipartite graphs or paths, *J. Discrete Math. Sci. Cryptogr.,* **12** (2009) 245-252. https://connectjournals.com/file_full_text/240902H_JDMSC290_245-252.pdf?utm_source=chatgpt.com

[1026] A. Elumalai and G. Sethuraman, Elegant labeled graphs, *J. Inform. Math. Sci.,* **2**(1) (2010) 45-49. https://www.rgnpublications.com/journals/index.php/jims/article/view/26/25

[1027] A. Elumali and G. Sethuraman, Gracefulness of a cycle with parallel chords and parallel P_k-chords of different lengths, *Ars Combin.,* **104** (2012) 143-148. https://combinatorialpress.com/article/ars/Volume%20104/volume-104-paper-11.pdf

[1028] S. El-Zanati, H.-L. Fu, and C.-L. Shiue, A note on the α-labeling number of bipartite graphs, *Discrete Math.,* **214** (2000) 241-243. https://www.sciencedirect.com/science/article/pii/S0012365X99003131?via%3Dihub

[1029] S. El-Zanati and D. I. Gannon, On ρ-labeling 2-regular graphs consisting of 5-cycles, *Int. J. Math. Comput. Sci.,* **6**(1) (2011) 13-20. https://bpb-us-w2.wpmucdn.com/about.illinoisstate.edu/dist/b/222/files/2019/11/Art057.RhoLabelingC5s-IJMCS.pdf

[1030] S. El-Zanati, M. Kenig, and C. Vanden Eynden, Near α-labelings of bipartite graphs, *Australas. J. Combin.,* **21** (2000) 275-285. https://ajc.maths.uq.edu.au/pdf/21/ocr-ajc-v21-p275.pdf?utm_source=chatgpt.com

[1031] S. El-Zanati and C. Vanden Eynden, Decompositions of $K_{m,n}$ into cubes, *J. Combin. Designs,* **4** (1996) 51-57.

[1032] S. El-Zanati and C. Vanden Eynden, On graphs with strong α-valuations, *Ars Combin.*, **56** (2000) 175-188. https://combinatorialpress.com/article/ars/Volume%20056/volume-56-paper-17.pdf

[1033] S. El-Zanati and C. Vanden Eynden, On α-valuations of disconnected graphs, *Ars Combin.*, **61** (2001) 129-136. https://combinatorialpress.com/article/ars/Volume%20061/volume-61-paper-10.pdf

[1034] S. El-Zanati and C. Vanden Eynden, On Rosa-type labelings and cyclic graph decompositions, *Math. Slovaca,* **59** (2009) 1-18. https://link.springer.com/content/pdf/10.2478/s12175-008-0108-x.pdf

[1035] S. El-Zanati, C. Vanden Eynden, and N. Punnim, On the cyclic decomposition of complete graphs into bipartite graphs, *Australas. J. Combin.,* **24** (2001) 209-219. https://ajc.maths.uq.edu.au/pdf/24/ocr-ajc-v24-p209.pdf

[1036] S. English and P. Znang, On graceful colorings of trees, *Math. Bohem.*, **142**(1) (2017) 57-73. https://www.researchgate.net/publication/314251162_On_graceful_colorings_of_trees

[1037] H. Enomoto, A. S. Llado, T. Nakamigawa, and G. Ringel, Super edge-magic graphs, *SUT J. Math.,* **34** (1998) 105-109. https://projecteuclid.org/journals/sut-journal-of-mathematics/volume-34/issue-2/SUPER-EDGE-MAGIC-GRAPHS/10.55937/sut/991985322.full

[1038] H. Enomoto, K. Masuda, and T. Nakamigawa, Induced graph theorem on magic valuations, *Ars Combin.,* **56** (2000) 25-32. https://combinatorialpress.com/article/ars/Volume%20056/volume-56-paper-3.pdf

[1039] A. A. Ephremnath and A. Elumalai, Every cycle with chord Hamiltonian path is harmonious and elegant, *Int. J. Adv. Engg. Tech.,***VII** (II) (2016) 1-4. https://www.researchgate.net/publication/311970017_Every_cycle_with_chord_Hamiltonian_path_is_Harmonious_and_Elegant

[1040] P. Erdős and A. B. Evans, Representations of graphs and orthogonal Latin squares, *J. Graph Theory,* **13** (1989) 593-595. https://old.renyi.hu/~p_erdos/1989-31.pdf

[1041] W. Q. Erickson, D. Herden, J. Meddaugh, M. R. Sepanski, I. Echols, C. Hammon, J. Marchena-Menendez, J. Mohn, B. Radillo-Murguia, and I. Ruiz-Bolanos, Klein cordial trees and odd cyclic cordial friendship graphs, (2022) arXiv:2211.10044 29 pages.

[1042] E. Esakkiammal, K. Thirusangu, and S. Seethalakshmi, Lucky-edge labeling of super subdivision of path graphs, *Recent Trends Graph Th. Combin.*, (2016) 42–49. https://books.google.com/books?hl=en&lr=&id=3Q5pEAAAQBAJ&oi=fnd&pg=PA42&ots=zjzPpP1lUg&sig=Pf1aBOSNiQalKcw3cvh3kjJBPJs#v=onepage&q&f=false

[1043] E. Esakkiammal, K. Thirusangu, and S. Seethalakshmi, Lucky edge labeling of super subdivision of star and wheel graphs, *Internat. J. Pure Appl. Math.*, **113**(7) (2017) 84-92. https://acadpubl.eu/jsi/2017-113-pp/articles/7/11.pdf

[1044] E. Esakkiammal, K. Thirusangu, and S. Seethalakshmi, Lucky-edge labeling of H-super subdivision of graphs, *Annals Pure App. Math.* **14**(3) (2017) 601–610. http://www.researchmathsci.org/apamart/apam-v14n3-27.pdf

[1045] E. Esakkiammal, K. Thirusangu, and S. Seethalakshmi, Lucky-edge labeling of super subdivision of planar grid graph, *Global J. Pure Appl. Math.*, **13**(1) (2017) 226–228.

[1046] K. Eshghi, The Existence and Construction of α-valuations of 2-Regular Graphs with 3 Components, Ph. D. Thesis, Industrial Engineering Dept., University of Toronto, 1997.

[1047] K. Eshghi, α-valuations of special classes of quadratic graphs, *Bull. Iranian Math. Soc.,* **28** (2002) 29-42. http://bims.iranjournals.ir/article_158_b47971249d3793df17f61e78fcd65885.pdf

[1048] K. Eshghi, Holey α-labelings of graphs, *Util. Math.,* **64** (2003) 139-147.

[1049] K. Eshghi, Extension of α-labelings of quadratic graphs, *IJMMS* **11** (2004) 571-578. https://www.researchgate.net/publication/27359173_Extension_of_a-labelings_of_quadratic_graphs

[1050] K. Eshghi, An introduction to the graceful graphs, Sharif University of Technology 2002) 65 pp. https://sharif.edu/~eshghi/Graceful%20Graphs-Final%20Edition-89-12-15.pdf

[1051] K. Eshghi and P. Azimi, Applications of mathematical programming in graceful labeling of graphs, *J. Appl. Math.,* **1**(1) (2004) 1-8. https://onlinelibrary.wiley.com/doi/epdf/10.1155/S1110757X04310065

[1052] K. Eshghi and P. Azimi, An algorithm for finding a feasible solution of graph labeling problems, *Util. Math.,* **72** (2007) 163-174.

[1053] K. Eshghi and M. Carter, Construction of α-valuations of special classes of 2-regular graphs, Topics in Applied and Theoretical Mathematics and Computer Science, *Math. Comput. Sci. Eng.*, WSEAS, Athens (2001) 139-154. https://www.wseas.us/e-library/conferences/cairns2001/papers/624.pdf?utm_source=chatgpt.com

[1054] K. Eshghi and M. Salarrezaie, Existence and construction of α-labeling for quadratic graph $Q(7, 4k)$ and its extensions, *Int. J. Graph. Theory Appl.,* (IJGTA), unpublished.

[1055] A. B. Evans, Representations of disjoint unions of complete graphs, *Disc. Math.*, 307 (2007) 1191–1198. doi:DOI: 10.1016/j.disc.2006.07.029

[1056] A. B. Evans, G. H. Fricke, C. C. Maneri, T. A. McKee, and M. Perkel, Representations of graphs modulo n, *J. Graph Theory* **18** (1994) 801-815. https://www.lehigh.edu/~gi02/modd.pdf

[1057] A. B. Evans, G. Isaak, and D. A. Narayan, Representations of graphs modulo n, *Discrete Math.,* **223** (2000) 109-123. https://www.sciencedirect.com/science/article/pii/S0012365X00000625

[1058] G. Exoo, A. Ling, J. McSorley, N. Phillips, and W. Wallis, Totally magic graphs, *Discrete Math.,* **254** (2002) 103-113. https://www.sciencedirect.com/science/article/pii/S0012365X01003673?via%3Dihu

[1059] G. Eyabi, J. Jacob, R. Laskar, D. Narayan, and D. Pillone, Minimal rankings of the Cartesian product $K_n \square K_m$, *Discuss. Math. Graph Th.,* **32**(4) (2012) 725-735. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1634

[1060] A. Ezhil, Prime labelling for some bipartite related graphs, *Internat. J. Math. Comput. Sci.,* **11**(7) (2023) 3571-3573. https://www.researchgate.net/publication/372666196_PRIME_LABELLING_FOR_SOME_BIPARTIATE_RELATED_GRAPHS

[1061] J. A. J. Ezhil, M. Regees, and T. S. I. Mary, Edge trimagic graceful labeling of some star graphs, *Webology* **17**(1) (2020) 628-631. https://webology.org/data-cms/articles/20230209053243pmWEBOLOGY%2017%20(1)%20-%201.pdf

[1062] E. M. Fachriza and K. A. Sugeng, The local antimagic total chromatic number on barabell wheel graphs, *Jurnal Matematika UNAND* **14**(3) (2025) 267-274. https://jmua.fmipa.unand.ac.id/index.php/jmua/article/view/1440/1022

[1063] L. Fan and Z. Liang, On Lee's conjecture and some results, *Discuss. Math. Graph Theory,* **29** (2009) 481-498. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1459

[1064] L. Fang, H. Huang, J. Pach, G. Tardos, and J. Zuo, Successive vertex orderings of fully regular graphs *J. Combin. Theory Ser. A* **199** (2023). https://doi.org/10.48550/arXiv.2206.13592

[1065] W. Fang, A computational approach to the graceful tree conjecture, arXiv:1003.3045 [cs.DM].

[1066] W. Fang, New computational result on harmonious trees, arXiv:1106.3490v1 [cs.DM] 17 June 2011.

[1067] A. Farida, R. P. Indah, and N. A. Sudibyo, Magic covering and edge magic labelling and its application magic covering and edge magic labelling and its application, *J. Physics: Confer. Ser.,* 1657 (2020) 012051 1-5. https://iopscience.iop.org/article/10.1088/1742-6596/1657/1/012051

[1068] H. Farida and D. Indriati, On the total edge irregularity strength of three towers hanoi graph, *J. Physics Confer. Ser.,* (2019) 1211:012010 https://iopscience.iop.org/article/10.1088/1742-6596/1211/1/012010/pdf

[1069] M. Farisa and K. S. Parvathy, Lh labeling of some graphs, *Internat. J. Recent Tech. Engin.,* (IJRTE) **7** (2019) 874-878. https://www.ijrte.org/wp-content/uploads/papers/v7i6s2/F11020476S219.pdf

[1070] M. Farisa and K. S. Parvathy, LH labeling of graphs, *Adv. Math. Sci. J.*, **10**(4) (2021) 2167-2179. https://research-publication.com/amsj/uploads/papers/vol-10/iss-04/AMSJ-2021-N04-31.pdf

[1071] R. J. Faudree and J. Lehel, Bound on the irregularity strength of regular graphs, *Combinatorics (Eger, 1987), Colloq. Math. Soc. Jănos Bolyai*, 52 (1988) 247–256.

[1072] R. J. Faudree, M. S. Jacobson, J. Lehel and R. H. Schelp, Irregular networks, regular graphs and integer matrices with distinct row and column sums, *Discrete Math.*, **76** (1989) 223-240. https://www.sciencedirect.com/science/article/pii/0012365X8990321X

[1073] R. J. Faudree, M. S. Jacobson, L. Kinch, and J. Lehel, Irregularity strength of dense graphs. *Discrete Math.,* **91**(1) (1991) 45-59. https://www.sciencedirect.com/science/article/pii/0012365X9190161T

[1074] F. Febriana and K. A. Sugeng, Odd harmonious labeling on squid graph and double squid graph, *J. Physics: Conference Ser.,* **1538**(1) (2020) 1–5. https://iopscience.iop.org/article/10.1088/1742-6596/1538/1/012015/pdf

[1075] W. Feng, On the gracefulness of the digraphs $n-\overrightarrow{C_m}$, *Internat. J. Pure Appl. Math.,* **78**(3) (2012) 429-434. https://www.ijpam.eu/contents/2012-78-3/13/13.pdf

[1076] W. Feng, C. Xu, and Jirimutu, A survey of the gracefulness of digraphs, *Int. J. Pure Appl. Math.,* **69** (2011) 245-253. https://www.ijpam.eu/contents/2011-69-3/2/2.pdf

[1077] W. Feng, X. Hong, Y. Yang, and Jirimutu, On (a, d)-antimagic labelings of generalized Petersen graphs $P(n, 5)$, *Util. Math.* **95** (2014) 349-356.

[1078] H. Fernau and K. Gajjar, Sum labelling graphs of maximum degree two, *Discrete Math.* **347**(8) (2024), 113893. doi:10.1016/j.disc.2024.113893

[1079] H. Fernau and K. Gajjar, The space complexity of sum labelling, *Theory of Computing Systems,* (2023) https://doi.org/10.1007/s00224-023-10130-2

[1080] B. Fern/'andez, R. H. Maleki, S. Miklavič, and A. S. Razafimahatratra, On closed distance magic circulants of valency up to 5. *Discrete Math.*, **346** (2023), 113581. doi:10.1016/j.disc.2023.113581

[1081] C. Fernandez, A. Flores, M. Tomova, and C. Wyels, The radio number of gear graphs, arXiv:0809.2623

[1082] H. Fernau, J. F. Ryan, and K. A. Sugeng, A sum labelling for the generalised friendship graph, *Discrete Math.,* **308** (2008) 734-740. https://www.sciencedirect.com/science/article/pii/S0012365X07005122

[1083] J. Ferraro, G. Newkirk, and S. Shan, Antimagic orientation of subdivided caterpillars, *Discrete Appl. Math.,* 313 (2022) 45-52. arXiv:2106.08430.

[1084] R. Figueroa-Centeno and R. Ichishima, The n-dimensional cube is felicitous, *Bull. Inst. Combin. Appl.,* **41** (2004) 47-50.

[1085] R. M. Figueroa-Centeno and R. Ichishima, On the sequential number and super edge-magic deficiency of graphs, Ars Combin., **129** (2016) 157-163. https://combinatorialpress.com/article/ars/Volume%20129/volume-129-paper-12.pdf

[1086] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, Bertrand's postulate and magic product labelings, *Bull. Instit. Combin. Appl.,* **30** (2000) 53-65. https://www.researchgate.net/publication/266214893_Bertrand's_postulate_and_magical_product_labelings

[1087] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, The place of super edgemagic labelings among other classes of labelings, *Discrete Math.,* **231** (2001) 153-168. https://www.sciencedirect.com/science/article/pii/S0012365X00003149

[1088] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, On super edge-magic graphs, *Ars Combin.,* **64** (2002) 81-95. https://www.researchgate.net/publication/220620148_On_Super_Edge-Magic_Graphs

[1089] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, Magical coronations of graphs, *Australas. J. Combin.,* **26** (2002) 199-208. https://ajc.maths.uq.edu.au/pdf/26/ajc_v26_p199.pdf

[1090] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, Labeling the vertex amalgamation of graphs, *Discuss. Math. Graph Theory*, **23** (2003) 129-139. https://www.researchgate.net/publication/268613774_Labeling_the_vertex_amalgamation_of_graphs

[1091] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, On edge-magic labelings of certain disjoint unions of graphs, *Australas. J. Combin.* **32** (2005) 225-242. https://www.researchgate.net/publication/268891852_On_edge-magic_labelings_of_certain_disjoint_unions_of_graphs z

[1092] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, Some new results on the super edge-magic deficiency of graphs, *J. Combin. Math. Combin. Comput.,* **55** (2005) 17-31. https://www.researchgate.net/publication/266952486_Some_new_results_on_the_super_edge-magic_deficiency_of_graphs

[1093] R. Figueroa-Centeno, R. Ichishima, and F. Muntaner-Batle, On the super edge-magic deficiency of graphs, *Ars Combin.,* **78** (2006) 33-45. https://combinatorialpress.com/article/ars/Volume%20078/volume-78-paper-4.pdf

[1094] R. Figueroa-Centeno, R. Ichishima, F. Muntaner-Batle and A. Oshima, A magical approach to some labeling conjectures, *Discussiones Math. Graph Theory,* **31** (2011) 79-113. https://www.researchgate.net/publication/220469087_A_magical_approach_to_some_labeling_conjectures

[1095] R. Figueroa-Centeno, R. Ichishima, F. Muntaner-Batle, and A. Oshima, Gracefully cultivating trees on a cycle, *Electron. Notes Discrete Math.*, **48** (2015) 143-150. https://www.sciencedirect.com/science/article/pii/S1571065315000219

[1096] R. Figueroa-Centeno, R. Ichishima, F. Muntaner-Batle, and M. Rius-Font, Labeling generating matrices, Labeling generating matrices, *J. Combin. Math. Combin. Comput.,* **67** (2008) 189-216. https://www.researchgate.net/publication/33421013_Labeling_Generating_Matrices

[1097] F. Firmansah, The odd harmonious labeling on variation of the double quadrilateral windmill graphs, *Jurnal Ilmu Dasar*, **18**(2) (2017) 109-118. https://doi.org/10.19184/jid.v18i2.5648

[1098] F. Firmansah, Pelabelan armonis ganjil pada graf ular jaring berlipat, *Sainmatika: J. Ilmiah Mate. Dan Ilmu Pengetahuan Alam*, **17**(1) (202) 1–8. https://doi.org/10.31851/sainmatika.v17i1.3182

[1099] F. Firmansah, Odd harmonious labeling on some string graph classes, *BAREKENG: J. Ilmu Mat. Dan Terapan*, **16**(1) (2022) 315-322. https://doi.org/10.30598/barekengvol16iss1pp313-320

[1100] F. Firmansah, The odd harmonious labeling of layered graphs, *JTAM (Jurnal Teori Dan Aplikasi Matematika)*, **7**(2) (2023) 339-348. https://journal.ummat.ac.id/index.php/jtam/article/view/12506

[1101] F. Firmansah and W. Giyarti, Odd harmonious labeling on the amalgamation of the generalized double quadrilateral windmill graph, *Desimal: J. Mat.*, **4**(3) (2021) 373-378. http://ejournal.radenintan.ac.id/index.php/desimal/article/view/10823

[1102] F. Firmansah, Tasari, and M. R. Yuwono, Odd harmonious labeling of lotus flower graphs, *CAUCHY – J. Matematika Murni Aplikasi,* **10**(2) (2025) 1133-1146. https://ejournal.uin-malang.ac.id/index.php/Math/article/view/35272/pdf

[1103] F. Firmansah, Tasari, and M. R. Yuwono, Odd harmonious labeling of the zinnia flower graphs, *Jurnal Ilmiah Sains*, **23**(1) (2023) 40-46. https://doi.org/10.35799/jis.v23i1.46771

[1104] F. Firmansah and T. Tasari, Odd harmonious labeling on edge amalgamation from double quadrilateral graphs, *Desimal: J. Mat.,* **3**(1) (2020) 65-72. DOI:10.24042/djm.v3i1.5712

[1105] F. Firmansah and M. R. Yuwono, Odd harmonious labeling of pleated of the Dutch windmill graphs, *Cauchy J. Mate. Murni dan Aplikasi*, **4**(4) (2017) 161-166. doi:10.18860/ca.v4i4.4043

[1106] N. B. Fox, B. Bonney, and S. Patten, Odd graceful labelings of prism graphs, *Math Exchange*, (2025) **19**(1) 16-30. https://openjournals.bsu.edu/mathexchange/article/view/6083/3320

[1107] N. B. Fox and C. P. Mooney, On prime labelings of zero-divisor graphs, *Congress. Numerant.*, **236** (2025) 73-100. https://combinatorialpress.com/cn-articles/volume-236/on-prime-labelings-of-zero-divisor-graphs/

[1108] N. B. Fox and J. Spaeth, Total prime labelings of various graphs. arXiv:2601.14535

[1109] S. Francis and V. Balaji, On relaxed mean labeling for wheel graphs, *Internat. J. Pure Appl. Math,*, **114** (2017) 173-182. https://acadpubl.eu/jsi/2017-114-5/articles/21.pdf

[1110] C. Frayer, Properties of divisor graphs, *Rose-Hulman Undergraduate Math. J.*, **2** (2003) Article 4. https://scholar.rose-hulman.edu/rhumj/vol4/iss2/4/?utm_source=chatgpt.com

[1111] S. Freeda and R. S. Chellathurai, H- and H_2-cordial labeling of some graphs, *Open J. Discrete Math.*, **2** (2012) 149-155. doi:10.4236/ojdm.2012.24030

[1112] B. J. Freyberg, Face-magic labelings of type (a, b, c) from edge-magic labelings of type (α, β). *Bull. Instit. Combin. Appl.*, **93** (2021) 81-103. http://bica.the-ica.org/Volumes/93/Reprints/BICA2021-05-Reprint.pdf

[1113] B. J. Freyberg, Corrigendum: Face-magic labelings of type (a, b, c) from edge-magic labelings of type (α, β), **Bull. Inst. Combin. Appl.,** 94 (2022) 71-72. https://doi.org/10.26493/2590-9770.1658.c96

[1114] Bryan Freyberg, On face-magic labelings of regular tesselations, *The Art Disc. Appl. Math.,* to appear, 9 pp. https://doi.org/10.26493/2590-9770.1658.c96

[1115] B. Freyberg, G. Jensen, and K. Scott, Type (a, b, c) face-magic labelings of calendula graphs, *Combin. Combinatorial Comput.*, : MCCCC34, Springer, (2026) 129–143.

[1116] B. Freyberg and M. Keranen, Orientable Z_n-distance magic labeling of the Cartesian product of many cycles, *Electron. J. Graph Theory Appl.,* **5**(2) (2017) 304-311. https://media.neliti.com/media/publications/196196-EN-orientable-z_n-distance-magic-labeling-o.pdf

[1117] B. Freyberg and M. Keranen, Orientable Z_n-distance magic labeling of the Cartesian product of two cycles, *Australas. J. Combin.,* **69** (2017) 222-235. https://ajc.maths.uq.edu.au/pdf/69/ajc_v69_p222.pdf

[1118] B. Freyberg and M. Keranen, Orientable Z_n-distance magic magic graphs via products, *Australasian J. Combin.,* **70**(3) (2018) 319-328. https://ajc.maths.uq.edu.au/pdf/70/ajc_v70_p319.pdf

[1119] S. Fried and T. Mansour, Graph labelings obtainable by random walks, *Discrete Appl. Math.* (2023). https://www.semanticscholar.org/reader/687105698f5da6d63ecf2ce382fa8044d00e683a

[1120] S. Fried and T. Mansour, Further results on random walk labelings, *Discrete Appl. Math.* **353** (2024) 211-221. https://doi.org/10.48550/arXiv.2305.09971

[1121] S. Fried and T. Mansour, Random walk labelings of perfect trees and other graphs, to appear in *Quaest. Math.* (2025). https://doi.org/10.2989/16073606.2025.2458800

[1122] A. Frieze, R. J. Gould, M. Karonski and F. Pfender, On graph irregularity strength, *J. Graph Theory* **41** (2002) 120-137. https://onlinelibrary.wiley.com/doi/epdf/10.1002/jgt.10056

[1123] D. Fronček, Bi-cyclic decompositions of complete graphs into spanning trees, *Discrete Math.,* **307** (2007) 1317-1322. https://www.sciencedirect.com/science/article/pii/S0012365X06007266

[1124] D. Fronček, Recent advances in Rosa-type labelings of graphs, *AKCE J. Graphs Combin.,* **6** (2009) 69-78. https://www.tandfonline.com/doi/epdf/10.1080/09728600.2009.12088873?needAccess=true

[1125] D. Fronček, α_2-labeling of graphs, *Opuscula Math.,* **29** (2009) 393-397. https://www.researchgate.net/publication/266909857_a_2_-labeling_of_graphs

[1126] D. Fronček, Group distance magic labeling of Cartesian product of cycles, *Australas. J. Combin.,* **55** (2013) 167-174. https://ajc.maths.uq.edu.au/pdf/55/ajc_v55_p167.pdf

[1127] D. Fronček, Handicap distance antimagic graphs and incomplete tournaments, *AKCE Int. J. Graphs Comb.,* **16**(2) (2013) 119-127. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088729

[1128] D. Fronček, Alpha labelings of full hexagonal caterpillars, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 85-89. https://www.tandfonline.com/doi/epdf/10.1016/j.akcej.2016.03.003?needAccess=true

[1129] D. Fronček, Handicap incomplete tournaments and ordered distance antimagic graphs, https://www.researchgate.net/publication/267092003_Handicap_incomplete_tournaments_and_ordered_distance_antimagic_graphs

[1130] D. Fronček, Regular incomplete tournaments with handicap two, *Congr. Numer.,* **227** (2016) 277-285.

[1131] D. Fronček, Regular handicap graphs of odd order, *J. Combin. Math. Combin. Comput.,* **102** (2017) 253-266. https://combinatorialpress.com/article/jcmcc/Volume%20102/vol-102-paper%2016.pdf

[1132] D. Fronček, Full spectrum of regular incomplete 2-handicap tournaments of order $n \equiv 0 \pmod{16}$, *J. Combin. Math. Combin. Comput.,* **106** (2018) 175-184. https://www.d.umn.edu/~dfroncek/preprints/2-handicap%200mod16-II-02a.pdf

[1133] D. Fronček, Supermagic labeling of $C_n \square C_m$, *Utilitas Math.*, **126** (2026) 171-185. https://combinatorialpress.com/um-articles/vol-126/supermagic-labeling-of-c_nbox-c_m/

[1134] D. Fronček, $(K_4 - e)$ supermagic labeling of triangular ladders, *J. Combin. Math. Combin. Comput.*, **128** (2025) 215-219. https://combinatorialpress.com/article/jcmcc/Volume%20128/k_4-e-supermagic-labeling-of-triangular-ladders.pdf

[1135] D. Fronček, C_n-supermagic labeling of rich calendula graphs, *Congressus Numer.*, **237** (2026) 9-35. https://doi.org/10.61091/cn237-02

[1136] D. Fronček, O. Kingston, and K. Vezina, Alpha labelings of straight simple polynomial caterpillars, *Congr. Numer.,* **222** (2014) 57-64. https://www.d.umn.edu/~dfroncek/preprints_soubory/polyominal%20caterpillars%2001e.pdf

[1137] D. Fronček, P. Kovář, and T. Kovářová, Vertex magic total labeling of products of cycles, *Australas. J. Combin.*, **33** (2005) 169-181. https://ajc.maths.uq.edu.au/pdf/33/ajc_v33_p169.pdf

[1138] D. Fronček, P. Kovář, and T. Kovářová, Constructing distance magic graphs from regular graphs, *J. Combin. Math. Combin. Comput.,* **78** (2011) 349-354. https://combinatorialpress.com/article/jcmcc/Volume%20078/vol-078-paper%2027.pdf

[1139] D. Fronček and M. Kubesa, Factorizations of complete graphs into spanning trees, *Congr. Numer.,* **154** (2002) 125-134. https://www.tandfonline.com/doi/epub/10.1016/j.akcej.2019.08.009?needAccess=true

[1140] D. Fronček, J. McKeown, M. McKeown, Z_{2nm}-supermagic labeling of $C_n \square C_m$, Indones. J. Combin., 2(2) (2018), 57-71. https://www.ijc.or.id/index.php/ijc/article/view/62

[1141] D. Fronček and M. McKeown, Note on diagonal construction of Z_{2nm}-supermagic labeling of $C_n \square C_m$, *AKCE Int. J. Graphs Comb.*, **17**(3) (2020) 952-954. https://www.tandfonline.com/doi/epdf/10.1016/j.akcej.2019.08.009?needAccess=true

[1142] D. Fronček, P. Paananen, L. Sorensen, Group-supermagic labeling of Cartesian product of two odd cycles, *Bull. Inst. Combin. Appl.*, **101** (2024) 39-53. https://bica.the-ica.org/Volumes/101/Reprints/BICA2022-17-Reprint.pdf

[1143] D. Fronček, P. Paananen, and L. Sorensen, Group-supermagic labeling of Cartesian products of two even cycles, *Discrete Math.,* (2023) 113741. https://www.sciencedirect.com/science/article/pii/S0012365X23004272

[1144] D. Fronček and J. Qiu, Supermagic graphs with many odd degrees, *Combin. Algor.*, 229–236, Lecture Notes in Comput. Sci., 11638, Springer, Cham, 2019.ivan https://link.springer.com/chapter/10.1007/978-3-030-25005-8_19

[1145] D. Fronček and A. Shepanik, Regular handicap graphs of order $n \equiv 0 \pmod 8$, *Electronic J. Graph Th. Appl.* **6** (2) (2018) 208-218. https://www.ejgta.org/index.php/ejgta/article/view/506/pdf_67

[1146] D. Fronček and A. Shepanik, Regular handicap graphs of order $n \bmod 4 \pmod 8$, *Elec. J. Graph Theory Appl.,* **10**(1), (2022) 259-273. https://www.ejgta.org/index.php/ejgta/article/view/1362

[1147] D. Fronček and L. Tollefson, Decompositions of complete graphs into kayak paddles, *J. Indones. Math. Soc.,* Special Edition (2011) 30-44. https://jims-a.org/index.php/jimsa/article/view/17/pdf_1

[1148] D. Fronček and L. Tollefson, Decompositions of complete graphs into kayak paddles with two odd cycles, Proceedings of the Forty-Second Southeastern International Conference on Combinatorics, Graph Theory and Computing. *Congr. Numer.,* **207** (2011) 161-169. https://www.researchgate.net/publication/266832130_

Decompositions_of_complete_graphs_into_kayak_paddles_with_two_odd_cycles

[1149] D. Fronček and S. Winters, Another family of gracious bipartite graphs, *Congr. Numer.,* **170** (2004) 65-70.

[1150] R. Frucht, Graceful numbering of wheels and related graphs, *Ann. N.Y. Acad. Sci.*, **319** (1979) 219-229.

[1151] R. Frucht, On mutually graceful and pseudograceful labelings of trees, *Scientia Series A,* **4** (1990/1991) 31-43.

[1152] R. Frucht, Nearly graceful labelings of graphs, *Scientia*, **5** (1992-1993) 47-59.

[1153] R. Frucht and J. A. Gallian, Labeling prisms, *Ars Combin.*, **26** (1988) 69-82.

[1154] R. Frucht and L. C. Salinas, Graceful numbering of snakes with constraints on the first label, *Ars Combin.*, **20** (1985), B, 143-157.

[1155] C.-M. Fu, N.-H. Jhuang and Y.-L. Lin, Integer-magic spectra of sun graphs, *Ars Combin.,* **103** (2012) 55-64. https://www.researchgate.net/publication/265494198_Integer-Magic_Spectra_of_Sun_Graphs

[1156] H. L. Fu and K. C. Huang, On prime labelling, *Discrete Math.,* **127** (1994) 181-186. https://www.researchgate.net/publication/220192761_On_prime_labellings

[1157] H. L. Fu and S. L. Wu, New results on graceful graphs, *J. Combin. Info. Sys. Sci.*, **15** (1990) 170-177. https://www.researchgate.net/publication/269008380_New_results_on_graceful_graphs

[1158] Y. Fukuchi, Graph labelings in elementary abelian groups, *Discrete Math.,* **189** (1998) 117-122. https://www.sciencedirect.com/science/article/pii/S0012365X98000065

[1159] Y. Fukuchi, A recursive theorem for super edge-magic labelings of trees, *SUT J. Math.,* **36** (2000) 279-285.

[1160] Y. Fukuchi, Edge-magic labelings of generalized Petersen graphs $P(n, 2)$, *Ars Combin.,* **59** (2001) 253-257. https://combinatorialpress.com/ars-articles/volume-059-ars-articles/edge-magic-labelings-of-generalized-petersen-graphs-pn-2/

[1161] Y. Fukuchi, Edge-magic labelings of wheel graphs, *Tokyo J. Math.,* **24** (2001) 153-167. https://projecteuclid.org/journals/tokyo-journal-of-mathematics/volume-24/issue-1/Edge-Magic-Labelings-of-Wheel-Graphs/10.3836/tjm/1255958319.full

[1162] Y. Fukuchi and A. Oshima, Super-edge-magic labelings of a family of trees with diameter 4, *Adv. Appl. Discrete Math.,* **1** (2008) 149-157. https://projecteuclid.org/journals/tokyo-journal-of-mathematics/volume-24/issue-1/Edge-Magic-Labelings-of-Wheel-Graphs/10.3836/tjm/1255958319.full

[1163] Y. Fukuchi, and A. Oshima, Super-edge-magic labeling of trees with large diameter, *Far East J. Math. Sci.,* **28** (2008) 497-571. https://www.rs.tus.ac.jp/egawa_60th_birthday/abstract/contributed_talk/Akito_Oshima.pdf

[1164] A. Graf, Graceful labelings of pendant graphs, *Rose-Hulman Undergrad. Math. J.*, **15**(1) (2014), Article 10. https://scholar.rose-hulman.edu/cgi/viewcontent.cgi?article=1060&context=rhumj

[1165] R. M. Gajjar and K. K. Raval, Some labeling of north star and lotus star graphs, *METSZET J.* **7**(12) (2023) 379-387. https://www.researchgate.net/publication/371538693_Some_Labeling_of_North_Star_and_Lotus_Star_Graphs

[1166] S. J. Gajjar, Prime cordial labeling of generalized web graph $W(t, n)$, *Internat J. Res. Analytical Reviews,* (IJRAR) **6**(2) (2019) 697-710. https://www.researchgate.net/publication/366659959_Prime_cordial_labeling_of_generalized_web_Graph

[1167] S. J. Gajjar and A. K. Des, Cordial labeling of generalized Jahangir graph, *Internat. J. Math. Appl.*, **4**(1–D) (2016) 21-33. https://www.researchgate.net/publication/329626405_Cordial_Labeling_of_Generalized_Jahangir_Graph

[1168] J. A. Gallian, Labeling prisms and prism related graphs, *Congr. Numer.*, **59** (1989) 89-100. https://www.researchgate.net/publication/265929297_Labeling_prisms_and_prism_related_graphs#fullTextFileContent

[1169] J. A. Gallian, A survey: recent results, conjectures and open problems on labeling graphs, *J. Graph Theory*, **13** (1989) 491-504. 10.1002/jgt.3190130410

[1170] J. A. Gallian, Open problems in grid labeling, *Amer. Math. Monthly*, **97** (1990) 133-135. https://www.researchgate.net/publication/265701091_Open_Problems_in_Grid_Labeling

[1171] J. A. Gallian, A guide to the graph labeling zoo, *Discrete Appl. Math.,* **49** (1994) 213-229. https://www.researchgate.net/publication/222843108_A_guide_to_the_graph_labeling_zoo

[1172] J. Gallian, unpublished.

[1173] J. A. Gallian and D. S. Jungreis, Labeling books, *Scientia*, **1** (1988) 53-57. https://www.researchgate.net/publication/265314475_Labeling_books

[1174] J. A. Gallian, J. Prout, and S. Winters, Graceful and harmonious labelings of prisms and related graphs, *Ars Combin.*, **34** (1992) 213-222. https://combinatorialpress.com/article/ars/Volume%20034/volume-34-paper-21.pdf

[1175] J. A. Gallian and L. A. Schoenhard, Even harmonious graphs, *AKCE J. Graphs Combin.,* **11**(1) (2014) 27-49. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2014.12088761

[1176] J. A. Gallian and D. Stewart, Properly even harmonious labelings of disconnected graphs, *AKCE J. Graphs Combin.,* **12**(2-3) (2015) 193-203. https://www.tandfonline.com/doi/pdf/10.1016/j.akcej.2015.11.015

[1177] J. A. Gallian and D. Stewart, Even harmonious labelings of disjoint graphs with a small component, *AKCE J. Graphs Combin.,* **12**(2-3) (2015) 204-215. https://www.tandfonline.com/doi/pdf/10.1016/j.akcej.2015.11.016

[1178] J. A. Gallian and D. Stewart, Properly even harmonious labelings of disjoint unions with even sequential graphs, *J. Graph Labeling,* **1**(1) (2015) 1-10. https://www-users.cse.umn.edu/~dkstewar/publications/GalSteP3.pdf

[1179] R. Ganesan, A. Bhaalamurugan, and Christy, Prime labeling for some new classes of graphs, *Int. J. Res. Anal.*, Rev. 6 (2019) 232–235. https://ijrar.org/papers/IJRAR1ARP040.pdf?utm_source=chatgpt.com

[1180] S. Ganesan and D. Uthayakumar, Corona of bipartite graphs with divisor graphs produce new divisor graphs, *Bull. Kerala Math. Assoc.,* **9**(1) (2012) 219-226.

[1181] G. S. Ganeshwari and G. Sudhana, Even-odd average harmonious graph, *Proceed. Internat. Virtual Confer.*, (2021) 165-176. https://www.researchgate.net/profile/Ather-Raina/publication/355475969_Proceedings_of_International_Virtual_Conference/links/6172866ca767a03c14919b5d/Proceedings-of-International-Virtual-Conference.pdf#page=178

[1182] T. Gangopadhyay and S. P. Rao Hebbare, Bigraceful graphs-I, *Util. Math.*, **17** (1980) 271-275.

[1183] D. I. Gannon and S. I. El-Zanati, All 2-regular graphs with uniform odd components admit ρ-labelings, *Australas. J. Combin.,* **53** (2012) 207-219. https://ajc.maths.uq.edu.au/pdf/53/ajc_v53_p207.pdf

[1184] A-B. Gao, X-D. Zhang, and L-J. Xu, On (super) vertex-graceful labeling of graphs, *Ars Combin.,* **126** (2016) 121-131. https://combinatorialpress.com/article/ars/Volume%20126/volume-126-paper-11.pdf?utm_source=chatgpt.com

[1185] M. Gao, The edge different sets of the direct product of two paths, MS thesis, Fuzhou University, 2010.

[1186] Y. Z. Gao, On gracefulness of $C_m \cup P_2$, *J. Hebei Teachers College*, No. 3 (1993) 13-21.

[1187] Y. Z. Gao and Z. H. Liang, private communication.

[1188] Y. Gao, Yuping, and S. Shan, Antimagic orientation of lobsters, *Discrete Appl. Math.,* **287** (2020) 21-26. https://www.sciencedirect.com/science/article/abs/pii/S0166218X20303644

[1189] Z.-B. Gao, The labelings of a variation of banana trees, *Ars Combin.,* **94** (2010) 175-181. https://combinatorialpress.com/article/ars/Volume%20094/volume-94-paper-17.pdf?utm_source=chatgpt.com

[1190] Z.-B. Gao, Odd graceful labelings of some union graphs, *J. Nat. Sci. Heilongjiang Univ.,* **24** (2007) 35-39.

[1191] Z.-B. Gao, The super vertex-graceful labeling of graphs, *Ars Combin.,* **124** (2016) 389-400.
https://combinatorialpress.com/ars-articles/volume-124-ars-articles/the-super-vertex-graceful-labeling-of-graphs/
z

[1192] Z.-B. Gao, G. C. Lau, and S. M. Lee, On friendly index and product-cordial index sets of Möbius-liked graph, *J. Discrete Math. Sci. Crypt.,* **21**(3) (2018) 647-659. https://www.academia.edu/99215890/On_friendly_index_and_product_cordial_index_sets_of_M%C3%B6bius_liked_graph

[1193] Z.-B. Gao, W. C. Shiu, S-M. Lee and G. C. Lau, Full edge-friendly index sets of one point union of cycles, *Ars Combin.*, **159** (2024) 21-30. https://doi.org/10.15672/hujms.1266085

[1194] Z.-B. Gao and C. Fan, Some discussion on super edge-magic labelings of $St(a_1, \ldots, a_n)$, *Ars Combin.,* **108** (2013) 187-192.

[1195] Z.-B. H. Gao, Ruo-Yuan, S.-M. Lee, H.-N. Ren, and G.-Y. Sun, Discussions on full friendly index sets of a class of cubic graphs,*Ars Combin.,* **146** (2019) 375-389.

[1196] Z.-B. Gao, W. C. Shiu, S.-M. Lee, and G-C. Lau, Full edge-friendly index sets of one point union of cycles, *Ars Comb.*, **159** (2024) 21-30. https://combinatorialpress.com/article/ars/full-edge-friendly-index-sets-of-one-point-union-of-cycles.pdf

[1197] Z-B. Gao, G-Y. Sun, S-M. Lee, On full friendly index sets of 1-level and 2-levels N-grids, *Discrete Appl. Math.,* **211** (2016) 68-78. https://www.academia.edu/119847087/On_full_friendly_index_sets_of_1_level_and_2_levels_N_grids

[1198] Z.-B. Gao, G.-Y. Sun, Y.-N. Zhang, Y. Meng, and G. C. Lau, Product cordial and total product cordial labelings of P_{n+1}^n, *J. Discrete Math.,* (2015) Art. ID 512696, 6 pp. heDOIis10.1155/2015/512696

[1199] Z.-B. Gao and X. Zhang, A note on super edge-graceful labelings of caterpilars, *J. Shanghai Jiaotong Univ.,* **42** (2008) 493-495.

[1200] Z.- B. Gao, X.-D. Zhang, and L.-J. Xu, Odd graceful labelings of graphs, *Discrete Math. Algor. Appl.*, **1**(3) (2009) 377-388. https://doi.org/10.1142/S1793830909000300?utm_source=chatgpt.com

[1201] M. Gardner, Mathematical games: the graceful graphs of Solomon Golomb, or how to number a graph parsimoniously, *Scientific Amer.,* **226**3 (1972) 108-112; **226**4 (1972) 104; **226**(6) (1972) 118.

[1202] M. Gardner, Wheels, Life and Other Mathemaical Amusements, W. H. Freeman & Company, 1983.

[1203] B. Gayathri and K. Amuthavalli, personal communication (2008).

[1204] B. Gayathri and K. Amuthavalli, k-odd mean labeling of the graph $\langle K_{1,n} : K_{1,m}\rangle$, *Acta Ciencia Indica Math.,* **XXIV M** (2008) 827-834.

[1205] B. Gayathri and K. Amuthavalli, k-odd mean labeling of prism, *Trans. Comb.,* **4**(1) (2015) 49-56. https://toc.ui.ac.ir/article_5388_c3d1298da0efeb3c0c28f5a89756531d.pdf

[1206] B. Gayathri and S. K. Devi, k-edge-graceful labeling and k-global edge-graceful labeling of some graphs, *Internat. J. Engin. Sci., Adv. Comput. and Bio-Tech.,* **2**(1) (2010) 25-37. https://ijesacbt.com/archive/Volume2issue1003.pdf

[1207] B. Gayathri and S. K. Devi, k-even edge-graceful labeling of the bistar graph $B_{n,n}$, *Proceed. Internat. Conf. Math. and Comput. Sci. ICMCS 2011,* 525-532.

[1208] B. Gayathri and S. K. Devi, k-even edge-graceful labeling of some graphs, *Internat. J. Math. and Comput. Sci.,* **6**(1) (2011) 27-34.

[1209] B. Gayathri and S. K. Devi, k-even edge-graceful labeling of some cycle related graphs, *Internat. J. Engin. Sci. Invention,* **2**(10) (2013) 1-17. https://ijesi.org/papers/Vol%202(10)/A0210101017.pdf

[1210] B. Gayathri and M. Duraisamy, personal communication (2008)

[1211] B. Gayathri, M. Duraisamy, and M. Tamilselvi, Even edge graceful labeling of some cycle related graphs, *Int. J. Math. Comput. Sci.,* **2** (2007) 179-187.

[1212] B. Gayathri and R. Gopi, k-even mean labeling of $D_{m,n}@C_n$, *Internat. J. Enger. Sci., Adv. Comput. Bio-Tech.,* **1**(3) (2010) 137-145. https://ijesacbt.com/archive/Volume1issue3004.pdf

[1213] B. Gayathri and R. Gopi, k-even mean labeling of $D_{m,n}$, *Acta Ciencia Indica,* **XXXVII M**(2) (2011) 291-300.

[1214] B. Gayathri and R. Gopi, k-even mean labeling of $T_{n,m,t}$, *Internat. J. Enger. Sci., Adv. Comput. Bio-Tech.,* **2**(2) (2011) 77-87. https://ijesacbt.com/archive/Volume2issue2003.pdf

[1215] B. Gayathri and R. Gopi, k-even mean labeling of $C_n \cup P_m$, *Elixir Appl. Math.,* **36** (2011) 3102-3105. https://www.elixirpublishers.com/articles/1686029139_201107017.pdf?utm_source=chatgpt.com

[1216] B. Gayathri and R. Gopi, (k,d)-even mean labeling of $P_m \oplus nK_1$, *Internat. J. Math. Soft Comput.*, **1**(1) (2011) 17-23. https://www.academia.edu/101314061/_K_D_Even_Mean_Labeling_of_P_M_Odot_NK_1_

[1217] B. Gayathri and R. Gopi, Necessary condition for mean labeling, *Internat. J. Engin. Sci., Adv. Comput. Bio-Tech.,* **4**(3) July-Sept. (2013) 43-52. https://www.researchgate.net/publication/293062475_Necessary_condition_for_mean_labeling

[1218] B. Gayathri and R. Gopi, k-even mean labeling of some graph operations, *Internat. J. Engin. Sci., Adv. Comput. and Bio-Tech.,* **4**(3) Oct-Dec. (2013) 59-70. https://www.researchgate.net/publication/293062405_k-even_mean_labeling_of_some_graph_operations

[1219] B. Gayathri and R. Gopi, Cycle related mean graphs, *Elixir Internat. J. Applied Sci.,* No. 71, (2014) 25116-25124. https://www.researchgate.net/publication/363090190_Cycle_Related_Mean_Graphs

[1220] B. Gayathri and R. Gopi, Some properties of k-mean labeling, *Internat. J. Multidisciplinary Res. Publication (IJMRAP),* **1**(3) (2018) 1-2. https://ijmrap.com/wp-content/uploads/2018/09/IJMRAP-PP2257Y18.pdf

[1221] B. Gayathri and R. Gopi, Some general results on restricted k-mean graph, *J. Discrete Math. Sci. Cryptography*, (2019) 1-11. https://www.researchgate.net/publication/334242208_Some_general_results_on_restricted_k_-mean_graph

[1222] B. Gayathri and V. Hemalatha, Even sequential harmonious graphs, personal communication (2008).

[1223] B. Gayathri and D. Muthurramakrishnan, Some results on k-even sequential harmonious labeling of graphs, *Elixir Internat. J. Applied Sci.*, **47** (2012) 9054-9057. https://www.elixirpublishers.com/articles/1681285358_201206075.pdf

[1224] B. Gayathri and P. Vadivel, One modulo three mean labeling of some special graphs, *Math. Sci. Internat. Res. J.,* **4**(2) (2015) 434-440.

[1225] B. Gayathri and V. Vadivel Necessary condition for one modulo three mean labeling of graphs, *Internat. J. Engineering Sci., Adv. Comput. Bio-Tech.,* **6**(2) (2015) 41-61. https://ijesacbt.com/archive/Volume6issue2002.pdf?utm_source=chatgpt.com

[1226] B. Gayathri and P. Vadivel, One modulo three mean labeling of some family of trees, *Internat. J. Engin. Sci., Adv. Comput. Bio-Tech.,* **6**(4) (2015) 141-161. https://www.researchgate.net/publication/368257734_61480_95_61481_95_61501_95_61501_95_61501_95_61480_95_61481_95_61480_95_61481_95_61485_95_61538

[1227] B. Gayathri and P. Vadivel, One modulo three mean labeling of disconnected graphs, *Jamal Acad. Res. J.: Interdecipilinary,* **Special Issue** (2016) 21-26. https://www.researchgate.net/publication/363069935_ONE_MODULO_THREE_MEAN_LABELING_OF_DISCONNECTED_GRAPHS

[1228] B. Gayathri and P. Vadivel, k-one modulo three mean labeling of some path related graphs, *Inter. J. Res. Advent Tech.,* **6**(10) (2018) 2790-2795. https://www.researchgate.net/publication/368255121_ONE_MODULO_THREE_MEAN_LABELING_OF_SOME_PATH_RELATED_GRAPHS

[1229] B. Gayathri and P. Vadivel, k-one modulo three mean labeling of snake related graphs, *Internat. J. Sci. Res. Review* **7**(9) (2018) 169-177. http://www.dynamicpublisher.org/gallery/23-ijsrr-d817.pdf

[1230] B. Gayathri and P. Vadivel, One modulo three mean labeling of some path related graphs, *Adalya J.,* **8**(9) (2019) 176-181. https://www.researchgate.net/publication/368255121_ONE_MODULO_THREE_MEAN_LABELING_OF_SOME_PATH_RELATED_GRAPHS

[1231] B. Gayathri and M. Subbiah, Strong edge graceful labelings of some graphs, *Bull. Prue Appl. Sci.,* **27E** (2008) 1-10. https://www.thefreelibrary.com/Strong+edge+graceful+labeling+of+some+graphs.-a0199684928

[1232] B. Gayathri and M. Tamilselvi, personal communication (2008).

[1233] C. Gayathri and S. Saravanakumar, Equitable irregular edge-weighting of polar grid graph and Mongolian tent graph, *Proyecciones (Antofagasta)* **43**(3) (2024) 483-494. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/4677/4499

[1234] K. Gayathri, V. Thanjavur, P. Maniammai, and C. Sakar, Prime cordial labeling of some graphs, *J. Physics: Confer. Ser.,* 1947 (2021) 012016, 1-7. https://iopscience.iop.org/article/10.1088/1742-6596/1947/1/012016/pdf

[1235] G. Ge, Y. Miao and X. Sun, Perfect difference families, perfect difference matrices and related combinatorial structures, *J. Combin. Des.*, **18** (2010) 415-449. https://onlinelibrary.wiley.com/doi/10.1002/jcd.20259?msockid=11ad4ff45bd568ea3e475d365afe6977

[1236] T. Geetha and D. Kalamani, Square difference labeling for some graphs, *J. Computer Math. Sci.*, textbf10(4) 695-704. http://www.compmath-journal.org/dnload/T-Geetha-and-D-Kalamani/CMJV10I04P0695.pdf

[1237] S. Geethalakshmi, D. Saraswathi, and B. S. Divya, Super magic labelling of complete bipartite graphs, *J. Comput. Math.*, **8**(2) (2024) 69-73. https://www.shcpub.in/index.php/cm/article/view/387/208

[1238] J. P. Georges, D. W. Mauro, and Y. Wang, On the structures of V_4-magic and Z_4-magic graphs, *J. Combin. Math. Combin. Comput.,* **75** (2010) 137-152.

[1239] R. Gera, V. Saenpholphat, and P. Zhang, Divisor graphs with triangles, *Congr. Numer.,* **165** (2003) 51-64.

[1240] K. A. Germina, More on classes of strongly indexable graphs, *European J. Pure and Applied Math.*, **3-2** (2010) 269-281. https://www.ejpam.com/index.php/ejpam/article/view/577

[1241] K. A. Germina, A. Kumar, and K. L. Princy, Further results on set-valued graphs, *J. Discrete Math. Sci. Cryptogr.,* **11** (2008) 559-566. https://www.connectjournals.com/file_full_text/112305H_jdmsc259_559-566.pdf?utm_source=chatgpt.com

[1242] C. Gayathri and S. Saravanakumar, Equitable irregular edge-weighting of polar grid graph and Mongolian tent graph, *Proyecciones (Antofagasta)* **43**(3) (2024) 483-494. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/4677/4499

[1243] K. A. Germina and V. Ajitha, Strongly multiplicative graphs, unpublished.

[1244] K. A. Germina and T. M. K. Anandavally, On arithmetic embeddings of graphs, unpublished.

[1245] K. A. Germina and R. Sebastian, On square sum graphs, *Proyecciones* **32**(2) (2013) 107-117. https://www.scielo.cl/pdf/proy/v32n2/art02.pdf

[1246] K. A. Germina and R. Sebastian, Futher results on square sum graph, *Internat. Math. Forum*, **8**(1) (2013) 47-57. http://www.m-hikari.com/imf/imf-2013/1-4-2013/sebastianIMF1-4-2013.pdf

[1247] S. V. Gervacio, Which wheels are proper autographs?, *Sea Bull. Math.,* **7** (1983) 41-50.

[1248] S. V. Gervacio, personal communication (2006).

[1249] H. V. Getzimah and K. Palani, Antimagic total labeling on Z_{p+q}, *J. Comp. Math. Sci.*, **9**(11) (2018) 1653-1660. https://www.researchgate.net/publication/334655120_Antimagic_Total_Labeling_on

[1250] M. Ghebleh and R. Khoeilar, A note on: "H-cordial graphs," *Bull. Inst. Combin. Appl.,* **31** (2001) 60-68. https://www.researchgate.net/publication/2141958_A_note_on_H-cordial_graphs

[1251] G. V. Ghodasara and I. I. Jadav, New grid related cordial graphs, *Internat. J. Appl. Math.*, **28**(2) (2013) 1244-1248. https://www.researchgate.net/publication/280008859_New_Grid_Related_Cordial_Graphs

[1252] G. V. Ghodasara and J. P. Jena, Prime cordial labeling of the graphs related to cycle with one chord, twin chords and triangle, *Internat. J. Pure and Appl. Math.*, **89**(1) (2013) 79-87. https://www.ijpam.eu/contents/2013-89-1/9/9.pdf

[1253] G. V. Ghodasara and M. Patel, Some bistar related square sum graphs, *Internat. J. Math. Trend and Tech.* **47**(3) 172-177. https://ijmttjournal.org/public/assets/volume-47/number-3/IJMTT-V47P521.pdf

[1254] G. V. Ghodasara and M. Patel, Some new combination graphs, *Internat. J. Math. Appl.*, **5** (2A) (2017) 153-161.

[1255] G. V. Ghodasara and M. Patel, More on permutation labeling of graphs, *Internat. J. Appl. Graph Th.*, **1**(2) (2017) 30-42. https://ijmttjournal.org/2017/Volume-47/number-3/IJMTT-V47P521.pdf

[1256] G. V. Ghodasara and M. Patel, Innovative results on square sum labeling of graphs, *J. Graph Labeling*, **4**(1) (2018) 15-24. https://www.researchgate.net/publication/363403125_INNOVATIVE_RESULTS_ON_SQUARE_SUM_LABELING_OF_GRAPHS

[1257] G. V. Ghodasara, A. H. Rokad and I. I. Jadav, Cordial labeling of grid related graphs, *Internat. J. Comb. Graph Th. and App.*, **6**(2)(2013) 55-62. https://www.researchgate.net/publication/279950647_CORDIAL_LABELING_OF_GRID_RELATED_GRAPHS

[1258] G. V. Ghodasara and A. H. Rokad, Cordial labeling of $K_{n,n}$ related graphs, *Internat. J. Sci. Res.*, **2**(5) (2013) 74-77. https://www.researchgate.net/publication/302025526_Cordial_Labeling_of_K_nn_related_graphs

[1259] G. V. Ghodasara and A. H. Rokad, Cordial labeling in context of vertex switching of special graphs, *Internat. J. Math. Sci.,* **33**(2) (2013) 1389-1396. https://www.researchgate.net/publication/302025614_Cordial_Labeling_in_Context_of_Vertex_Switching_of_Special_Graphs

[1260] G. V. Ghodasara and A. H. Rokad, Cordial Labeling in context of barycentric subdivision of special graphs, *Internat. J. Math. Res.*, **5**(5) (2013) 475-483. http://www.irphouse.com/ijmr/ijmrv5n5_05.pdf

[1261] G. V. Ghodasara and S. G. Sonchhatra, Cordial labeling of fan related graphs, *Internat. J. Sci. Eng. Res.*, **4**(8) (2013) 470-476. https://www.ijser.org/researchpaper/Cordial-Labeling-of-Fan-Related-Graphs.pdf

[1262] G. V. Ghodasara and S. G. Sonchhatra, Further results on 3-equitable labeling, *IAENG Internat. J. Appl. Math.* IJAM, **45**(1) (2015) IJAM_45_1_01. https://www.iaeng.org/IJAM/issues_v45/issue_1/IJAM_45_1_01.pdf

[1263] G. V. Ghodasara and S. M. Vaghasiya, Product cordial labeling of graphs related to cycle with one chord, twin chords and triangle, *Internat. J. Math. Comput. Modelling*, **18**(2) (2013) 1109-1113.

[1264] G. V. Ghodasara and S. M. Vaghasiya, Product cordial labeling of graphs related to helm, closed helm and gear graph, *Internat. J. Pure and Appl. Math.*, **91**(4) (2014) 495-504. http://dx.doi.org/10.12732/ijpam.v91i4.6

[1265] E. Ghorbani and S. Kamali, Prime labeling of ladders, arXiv:1610.08849v1 [math.CO].

[1266] S. Ghosh, On certain classes of graceful lobsters, *Ars Combin.,* **136** (2018) 67-96. arXiv:1306.2932 23 pages

[1267] S. Ghosh, S. Mohanty, and A. Pal, Cordial labeling of Cartesian product between two balanced bipartite graphs, *J. Informatics Math. Sci.*, **11**(2) (2019) 179-193. https://www.rgnpublications.com/journals/index.php/jims/article/view/1132/914

[1268] P. Ghosh and A. Pal, Some new Fibonacci divisor cordial graphs, *AMO - Adv. Modeling Optim.,* **17**(2) (2015) 221-232. https://camo.ici.ro/journal/vol17/v17b5.pdf

[1269] P. Ghosh and A. Pal, Special classes of signed product cordial graphs, *Interna. J. Comput. Sci. Inform. Security* (IJCSIS) **15** (2017) 14-23. https://www.academia.edu/33409783/Special_Classes_of_Signed_Product_Cordial_Graphs

[1270] S. Ghosh and A. Pal, Signed product and total signed product cordial labeling of Cartesian product between balanced bipartite graph and path, *Adv. Comput. Comm. Paradigms,* **706** (2018) 515-522.

[1271] J. Ghoshal, R. Laskar, and D. Pillone, Minimal rankings, *Networks,* **28** (1996) 45-53. https://onlinelibrary.wiley.com/doi/epdf/10.1002/%28SICI%291097-0037%28199608%2928%3A1%3C45%3A%3AAID-NET6%3E3.0.CO%3B2-D

[1272] J. Ghoshal, R. Laskar, D. Pillone, and G. Fricke, Further results on mod sum graphs, *Congr. Numer.,* **101** (1994) 201-207. https://combinatorialpress.com/article/ars/Volume%20052/volume-52-paper-16.pdfhttps://combinatorialpress.com/article/ars/Volume%20052/volume-52-paper-16.pdf

[1273] M. Giridaran, Application of super magic labeling in cryptography, *Internat. J. Innovative Research in Sci., Engin. Tech.,* (IJIRSET)**9**(6) 2020 7 48-16-4822. http://www.ijirset.com/upload/2020/june/134_Application_DOI.PDF

[1274] M. Giridaran, T. A. Jose, and E. A. Jeony, On the radio antipodal geometric mean number of ladder related graphs, *Malaya J. Mat.*, **10**(01) (2022) 98-109. http://doi.org/10.26637/mjm1001/009

[1275] L. Girija and M. Karthikeyan, Magic and antimagic labeling Of copies of jelly fish graph, *J. Huazhong U. Sci. Tech.*, **50**(1) (2021) 1-6. https://www.researchgate.net/publication/349126147_Magic_and_Antimagic_Labeling_Of_Copies_Of_Jelly_Fish_Graph/references

[1276] R. B. Gnanajothi, Topics in Graph Theory, Ph. D. Thesis, Madurai Kamaraj University, 1991.

[1277] E. K. Gnang, On graceful labelings of trees, arXiv:1811.07614 [math.CO]. 17 pages.

[1278] E. K. Ganag, A proof of the Kotzig-Ringel-Rosa Conjecture, arXiv:2202.03178. 17 pages

[1279] E. K. Gnang and M. P. Williams, A proof of the Graham Sloane conjecture, arXiv:2201.02240. 10 pages.

[1280] R. Godbold and P. J. Slater, All cycles are edge-magic, *Bull. Inst. Combin. Appl.,* **22** (1998) 93-97. https://www.researchgate.net/publication/220620173_All_Cycles_are_Edge-Magic

[1281] L. Goddyn, R. B. Richter, and J. Širáň, Triangular embeddings of complete graphs from graceful labellings of paths, *J. Combin. Theory Ser. B,* **97** (2007) 964-970. https://www.sciencedirect.com/science/article/pii/S0095895607000226?via%3Dihub

[1282] A. Godinho and T. Singh, Group distance magic labeling of C_n^r, *Algorithms and Discrete Applied Math.,* 187-192, Lecture Notes in Comput. Sci., 10156, Springer, Cham, 2017. https://link.springer.com/chapter/10.1007/978-3-319-53007-9_17

[1283] A. Godinho and T. Singh, Some distance magic graphs, *Elect. Notes Discr. Math.,* **63** (2017) 311-315. https://www.sciencedirect.com/science/article/pii/S1571065317303293

[1284] A. Godinho, T. Singh, and S. Arumugam, On S-magic graphs, *Elect. Notes Discr. Math.*, **48** (2015) 267-273. http://dx.doi.org/10.1016/j.endm.2015.05.040

[1285] A. Godinho, T. Singh, and S. Arumugam, On nearly distance magic graphs, *Theoretical Comput. Sci. Disc. Math.*, 76-82, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017.

[1286] C. G. Goh and C. K. Lim, Graceful numberings of cycles with consecutive chords, *Electronic J. Combin.*, 17 (2010), #R12.

[1287] S. W. Golomb, How to number a graph, in *Graph Theory and Computing*, R. C. Read, ed., Academic Press, New York (1972) 23-37.

[1288] S. W. Golomb, Polyominoes: Puzzles, Patterns, Problems, and Packings, second ed. Princeton University Press, Princeton, NJ, 1994.

[1289] L. Golowich and C. Kim, New classes of set-sequential trees, *Discrete Math.,* **343**(3) (2020), 111741, 13 pp. doi:10.1016/j.disc.2019.111741

[1290] S. Gomathi, P. Venugopal, and T. Arputha Jose, Ladder related graphs using radio labeling, *Adv. Appl. Math. Sci.*, **21**(1) (2021) 79-92. https://www.mililink.com/upload/article/921916354aams_vol_211_november_2021_a6_p79-92_s._gomathi_and_t._jose_arputha.pdf?utm_source=chatgpt.com

[1291] S. Gomathi and P. Venugopal, Channel assignment of triangular grid and ladder related graphs using radio antipodal labeling, *Solid State Tech.*, **64**(2) (2021). https://www.researchgate.net/publication/357310601_Channel_assignment_of_triangular_grid_and_ladder_related_graphs_using_radio_labeling

[1292] S. Gomathi and P. Venugopal, Channel assignment of iriangular and rhombic honeycomb networks using radio labeling techniques, *Comm. Math. Appl.*, **12**(3) (2021) 665-676. https://www.rgnpublications.com/journals/index.php/cma/article/view/1621

[1293] S. Gomathi and P. Venugopal, Radio antipodal number of honeycomb derived networks, *Scientific Reports,* (2022) **12**(1) 10pp. doi:10.1038/s41598-022-23618-7

[1294] V. L. Gomathi, Felicitous labelings of some graphs, *Academic J. Applied Math. Sci.,* **2**(8) (2016) 93-97. https://web.archive.org/web/20180411055931/https://arpgweb.com/pdf-files/ajams2(8)93-97.pdf

[1295] V. L. Gomathi, A. Nagarajan, A. NellaiMurugan, On felicitous labelings of $P_{r,2m+1}$, P_r^{2m+1} and $C_n \times P_m$, *Internat. J. Math. Soft Computing,* **3**(1) (2012) 97-105.

[1296] V. L. Gomathi and A. Nagarajan, Felicitous labelings of some graphs, *OUTREACH* **IX** (2016) 132-139. https://web.archive.org/web/20180411055931/https://arpgweb.com/pdf-files/ajams2(8)93-97.pdf

[1297] J. Gómez, Solution of the conjecture: If $n \equiv 0 \pmod 4$, $n > 4$, then K_n has a super vertex-magic total labeling, *Discrete Math.*, **307** (2007) 2525-2534. https://www.sciencedirect.com/science/article/pii/S0012365X07000374

[1298] J. Gómez, Two new methods to obtain super vertex-magic total labelings of graphs, *Discrete Math.*, **308** (2008) 3361-3372. https://www.sciencedirect.com/science/article/pii/S0012365X07004864

[1299] J. Gómez and P. Kovář On super vertex-magic total labeling of the disjoint union of k copies of K_n, *Ars Combin.,* **113** (2013) 175-192.

[1300] J. T. Gondalia, Divisor cordial labelings of Herschel graph in context of various graph operations, *Harbin Gongye Daxue Xuebao/J. Harbin Inst. Tech.*, **54**(6) (2022) 185–190.

[1301] J. T. Gondalia, Mutiply divisor cordial labeling of special graphs, *Advances Appl. Math. Sci.*, **22**(11) (2023) 2213-2226. https://www.researchgate.net/profile/Jatin-Gondalia/publication/375916243_MULTIPLY_DIVISOR_CORDIAL_LABELING_OF_SPECIAL_GRAPHS/links/65632e023fa26f66f42924f4/MULTIPLY-DIVISOR-CORDIAL-LABELING-OF-SPECIAL-GRAPHS.pdf

[1302] J. T. Gondalia, On Fibonacci product cordial labeling in context of vertex switching of graphs, *Adv. Appl. Disc. Math.*, **35** (2022) 25-35.https://www.pphmj.com/abstract/14874.htm?utm_source=chatgpt.com

[1303] J. T. Gondalia and A. H. Rokad, Multiply divisor cordial labeling, *Internat. J. Engin. Adv. Tech.*, **9**(2) (2019) 1901-1904. https://www.researchgate.net/publication/344312265_Multiply_Divisor_Cordial_Labeling

[1304] J. T. Gondalia and A. H. Rokad, Multiply divisor cordial labeling in context of ring sum of graphs, *Adv. Math.: Sci. J.*, **9**(11) (2020) 9037-9044. https://www.research-publication.com/amsj/uploads/papers/vol-09/iss-11/AMSJ-2020-N11-8.pdf

[1305] J. T. Gondalia and A. H. Rokad, Integer cordial labeling of some star and bistar related graphs, *South East Asian J. Math. Math. Sci.,* **17**)2) (2021) 277-285. https://www.researchgate.net/publication/354126647_INTEGER_CORDIAL_LABELING_OF_SOME_STAR_AND_BISTAR_RELATED_GRAPHS

[1306] J. Goodell, A. Beveridge, M. Gallagher, D. Goodwin, J. Gyori, and A. Joseph, Sum graphs, (1990) unpublished.

[1307] R. Gopi, A Study on Different Kinds of Mean Graphs, Ph. D. Thesis, Bharathiasan University, Trichy, 2013.

[1308] R. Gopi, Odd sum labeling of alternative quadrilateral snake, *Internat. J. Engin. Sci., Advan. Comput. and Bio-Tech.*, **7**(3) (2016) 73-77. https://www.researchgate.net/publication/314222286_Odd_Sum_Labeling_of_Alternative_Quadrilateral_Snake

[1309] R. Gopi, Odd sum labeling of tree related graphs, *Internat. J. Math. and its Applications.*, **4**(4) (2016) 11-16. https://www.researchgate.net/publication/312070408_Odd_Sum_Labeling_of_Tree_Related_Graphs

[1310] R. Gopi, F-root square mean labeling of some graphs, *Internat. J. Math. Combin., 1* (2019) 146-149. https://www.researchgate.net/publication/332173176_F-Root_Square_Mean_Labeling_of_Some_Graphs

[1311] R. Gopi, k-Contra harmonic mean labeling of snake related graphs, *Math. J. Interdiscip. Sci.,* **7**(2) (2019) 77-80. https://mjis.chitkara.edu.in/index.php/mjis/article/view/175/114

[1312] R. Gopi, Super root square mean labeling of some graphs, *Internat. J. Engin. Sci., Advanced Comput. and Bio-Tech.*, **8**(2) (2017) 75-81. DOI:10.26674/ijesacbt/2017/49175

[1313] R. Gopi and A. Irudaya Mary, Odd sum labelings of some more graphs, *Internat. J. Engin. Sci., Adv. Comput. and Bio-Tech.,* **7**(4) (2016) 95-103. https://www.researchgate.net/publication/314222391_Odd_Sum_Labeling_of_Some_More_Graphs

[1314] R. Gopi and M. Kannan, C-Exponential mean labeling of quadrilateral reelated graphs, *Indian J. Nat. Sci.*, **16**(93) (2025) 105659. https://www.researchgate.net/publication/400315569_C_-Exponential_Mean_Labeling_of_Quadrilateral_Related_Graphs

[1315] R. Gopi and C. Nirmala, Lehmer-3 mean labeling of cycle related graphs, *Int. J. Res. Instinct.,* **5** (2018) 30-35. https://www.researchgate.net/publication/326367621_LEHMER-3_MEAN_LABELING_OF_CYCLE_RELATED_GRAPHS

[1316] R. Gopi, B. Sathish Kumar, F-Centroidal mean labeling of some graphs, *AEGAEUM J.*, **9**(1) (2021) 143-153. https://www.researchgate.net/publication/348648621_F-Centroidal_Mean_Labeling_of_Some_Graphs

[1317] R. Gopi, B. Sathish Kumar, F-Centroidal mean labelling of trees and cycle related graphs, *Internat. J. Aquatic Sci.*, **12**(2 (2021) 3623-3632. https://www.researchgate.net/publication/354236539_F-Centroidal_Mean_Labelling_of_Trees_and_Cycle_Related_Graphs

[1318] R. Gopi and C. Suba, Super Lehmer-3 mean labeling of tree related graphs, *Internat. J. Math. Appl.,* **5** 3-A (2017) 25-27. https://www.researchgate.net/

publication/319187098_Super_Lehmer-3_Mean_Labeling_of_Tree_Related_Graphs

[1319] R. Gopi and P. Vadivel, k-super contra harmonic mean labeling of graphs, *Asian J. Math. Sci.*, **7**(3)(2023) 112-116. https://www.researchgate.net/publication/375660813_K-SUPER_CONTRA_HARMONIC_MEAN_LABELING_OF_GRAPHS

[1320] R. Gopi and P. Vadivel, k-Super Lehmer-3 mean labeling of some graphs, *Indian J. Natural Sci.,* **14**(81) (2023) 66785-66789. https://www.researchgate.net/publication/377967140_k-Super_Lehmer-3_Mean_Labeling_of_Some_Graphs

[1321] R. J. Gould and V. Rödl, Bounds on the number of isolated vertices in sum graphs, *Graph Theory, Combin. and Appl.*, **1** (1991) 553-562. https://www.math.emory.edu/~rg/P38Y91.pdf

[1322] V. Govindan, S. Pinelas, and S. Dhivya, Cube sum labeling of graphs, *Annals Commun. Math.,* **3**(2) (2020) 171-176. https://www.technoskypub.com/wp-content/uploads/2023/09/6-V3-2-GPD-Cube-sum-labeling-of-graphs.pdf

[1323] T. Grace, Graceful, Harmonious, and Sequential Graphs, Ph. D. Thesis, University Illinois at Chicago Circle, 1982.

[1324] T. Grace, On sequential labelings of graphs, *J. Graph Theory*, **7** (1983) 195-201.

[1325] T. Grace, K_4 snakes are sequential, Technical Report, Department of Computer Science, Illinois Institute of Technology (1986).

[1326] A. J. Graham, D. A. Pike, and N. Shalaby, Skolem labelled trees and $P_s \square P_t$ Cartesian products, *Australas. J. Combin.,* **38** (2007) 101-115. https://ajc.maths.uq.edu.au/pdf/38/ajc_v38_p101.pdf

[1327] R. L. Graham and N. J. A. Sloane, On additive bases and harmonious graphs, *SIAM J. Alg. Discrete Methods,* **1** (1980) 382-404. https://www.researchgate.net/publication/237778431_On_Additive_Bases_and_Harmonious_Graphs

[1328] M. Grannell, T. Griggs, and F. Holroyd, Modular gracious labelings of trees, *Discrete Math.,* **231** (2001) 199-219. https://www.sciencedirect.com/science/article/pii/S0012365X00003186?via%3Dihub

[1329] I. Gray, New Construction Methods for Vertex-magic Total Labelings of Graphs, Ph.D. thesis, University of Newcastle, 2006.

[1330] I. Gray, Vertex-magic total labelings of regular graphs, *SIAM J. Discrete Math.* **21** (2007) 170-177. doi:10.1137/050639594

[1331] I. Gray and J. MacDougall, Sparse semi-magic squares and vertex-magic labelings, *Ars Combin.,* **80** (2006) 225-242. https://www.researchgate.net/publication/220620238_Sparse_Semi-Magic_Squares_and_Vertex-Magic_Labeling

[1332] I. Gray and J. MacDougall, Sparse anti-magic squares and vertex-magic labelings of bipartite graphs, *Discrete Math.,* **306** (2006) 2878-2892. https://www.researchgate.net/publication/220620238_Sparse_Semi-Magic_Squares_and_Vertex-Magic_Labelings

[1333] I. Gray and J. A. MacDougall, Vertex-magic labelings of regular graphs. II, *Discrete Math.,* **309** (2009) 5986-5999. https://www.sciencedirect.com/science/article/pii/S0012365X09002647

[1334] I. Gray and J. MacDougall, Vertex-magic labeling of non-regular graphs, *Australas. J. Combin.,* **46** (2010) 173-183. https://ajc.maths.uq.edu.au/pdf/46/ajc_v46_p173.pdf

[1335] I. Gray and J. A. MacDougall, Vertex-magic labelings: Mutations, *Austral. J. Combin.,* **45** (2009) 189-206. https://ajc.maths.uq.edu.au/pdf/45/ajc_v45_p189.pdf

[1336] I. Gray and J. MacDougall, Vertex-magic labeling of regular graphs: disjoint unions and assemblages. *Discrete Appl. Math.,* **160** (2012) 1114-1125. https://www.sciencedirect.com/science/article/pii/S0166218X11004781

[1337] I. Gray, J. MacDougall, R. Simpson, and W. Wallis, Vertex-magic total labeling of complete bipartite graphs, *Ars Combin.,* **69** (2003) 117-127. https://www.researchgate.net/publication/220620332_Vertex-Magic_Total_Labelings_of_Complete_Bipartite_Graphs

[1338] I. Gray, J. MacDougall, J. McSorley, and W. Wallis, Vertex-magic labeling of trees and forests, *Discrete Math.,* **261** (2003) 285-298. https://opensiuc.lib.siu.edu/cgi/viewcontent.cgi?article=1090&context=math_articles

[1339] I. Gray, J. MacDougall, and W. Wallis, On vertex-magic labeling of complete graphs, *Bull. Inst. Combin. Appl.,* **38** (2003) 42-44. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=efe828a43efd9c017c3558c6e2e67de790c612d7

[1340] R. P. Grimaldi, Graphs from rings, *Congr. Numer.*, **71** (1990) 95-104.

[1341] J. L. Gross and J. Yellen, Graph Theory and its Applications, CRC press, 2005.

[1342] M. V. S. Gudin and A. Pedrano, Tribonacci cordial labeling of some snake graphs, *Annals Communications Math.*, **8**(1) (2025) 393-405. doi:10.62072/acm.2025.080306

[1343] J. L. G. Guirao, S. Ahmad, M. K. Siddiqui, and M. Ibrahim, Edge irregular reflexive labeling for the disjoint union of generalized Petersen graph, *Mathematics*, **6** 304. https://www.mdpi.com/2227-7390/6/12/304

[1344] J. Guo, X. Chen, Z. Wang, and B. Yao, Total vertex irregularity strength of certain equitable complete m-partite graphs, *Ars Comb.*, **123** (2015) 407-418. https://combinatorialpress.com/article/ars/Volume%20123/volume-123-paper-32.pdf

[1345] W. F. Guo, Gracefulness of the graph $B(m,n)$, *J. Inner Mongolia Normal Univ.*, (1994) 24-29.

[1346] W. F. Guo, Gracefulness of the graph $B(m,n,p)$, *J. Math. (PRC)*, **15** (1995) 345-351.

[1347] S. Guruswamy and M. Varadhan, Generating graceful unicyclic graph from a given graceful tree, *Ars Combin.*, **151** (2020) 25–36.

[1348] A. Gutiérrez and A. Lladó, Magic coverings, *J. Combin. Math. Combin. Comput.*, **55** (2005) 43-56. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%204.pdf

[1349] P. Gvozjak, On the Oberwolfach Problem for Cycles with Multiple Lengths, Ph.D. thesis, Burnaby: Simon Fraser University, 2004. https://summit.sfu.ca/item/2738

[1350] A. Gyárfás and J. Lehel, A method to generate graceful trees, in *Colloque C.N.R.S. Problèmes Combinatories et Théorie des Graphes, Orsay, 1976* (1978) 207-209. https://www.renyi.hu/~gyarfas/Cikkek/08_GyarfasLehel_AMethodToGenerateGracefulTrees.pdf

[1351] E. Gyri, P. N. Balister, and R. H. Schelp, Coloring vertices and edges of a graph by nonempty subsets of a set, *Eur. J. Comb.*, **4** (2011) 533-537. https://core.ac.uk/download/pdf/82481312.pdf

[1352] H. M. Hafez, R. El-Shanawany, and A. A. El Atik, Odd harmonious labeling of the converse skew product of graphs, *Bull. Inst. Comb. Appl.*, **98** (2023) 90–110. http://bica.the-ica.org/Volumes/98/Reprints/BICA2022-39-Reprint.pdf

[1353] M. Hailat, On the cordial labeling of certain trigraphs, *J. Math. Comput. Sci.*, **11**(6) (2021) 6936-6948. https://scik.org/index.php/jmcs/article/view/6068

[1354] M. Hailat, On the cordial of weak labeling of an ℓ-fold cycle craph, $T_\ell(C_n)$, *J. Math. Res.* **17**(5) (2025) 1-9. https://www.ccsenet.org/journal/index.php/jmr/article/view/0/52644

[1355] P. Hajnal and G. Nagy, Simply sequentially additive labelings of 2-regular graphs, *Discrete Math.,* **310** (2010) 922-928. https://core.ac.uk/download/pdf/81968314.pdf

[1356] V. Halasza and Z. Tuza, Distance-constrained labeling of complete trees, *Discrete Math.,* **338** (2015) 1398-1406. https://www.sciencedirect.com/science/article/pii/S0012365X15000874?via%3Dihub

[1357] S. Hall, K. Hillesheim, E. Kocina, and M. Schmit, personal communication (2013).

[1358] A. C. Hamm, J. A. Hamm, and A. Way, A note on prime labeling k-partite k-graphs, *Involve* **15**(2) (2022) 233–239. doi:10.2140/involve.2022.15.233

[1359] N. Han and Z. Liang, On the graceful permutation graphs conjecture, *J. Discrete Math. Sci. Cryptogr.,* **11** (2008) 501-526. https://connectjournals.com/file_full_text/111805H_jdmsc238_501-526.pdf

[1360] A. K. Handa, A. Godinho, and T. Singh, Distance antimagic labeling of the ladder graph, *Elect. Notes Discr. Math.,* **63** (2017) 317-322. https://www.researchgate.net/publication/321447534_Distance_Antimagic_Labeling_of_the_Ladder_Graph

[1361] A. K. Handa, A. Godinho, and T. Singh, Some distance antimagic labeled graphs. In: S. Govindarajan and A. Maheshwari A. (eds), Algorithms and Discrete Applied Mathematics. Lecture Notes in Computer Science, **9602** 190-200. Springer, Cham 2016. https://link.springer.com/chapter/10.1007/978-3-319-29221-2_16

[1362] T. Hao, On sum graphs, *J. Combin. Math. Combin. Computing,* **6** (1989) 207-212. https://combinatorialpress.com/article/jcmcc/Volume%2006/vol-006-paper%2021.pdf

[1363] K. M. M. Haque, X. Lin, Y. Yang, and J. Zhang, Prime cordial labeling of flower snark and related graphs, *Ars Combin.,* **105** (2012) 45-52. https://combinatorialpress.com/article/ars/Volume%20105/volume-105-paper-5.pdf

[1364] K. M. M. Haque, X. Lin, Y. Yang, and P. Zhao, On the prime labeling of generalized Petersen graph $P(n, 1)$, *Util. Math.,* **83** (2010) 95-106.

[1365] K. M. M. Haque, X. Lin, Y. Yang, and P. Zhao, On the prime cordial labeling of generalized Petersen graph, *Util. Math.,* **82** (2010) 71-79. https://www.researchgate.net/publication/266005155_On_the_prime_cordial_labeling_of_generalized_Petersen_graph

[1366] K. M. M. Haque, X. Lin, Y. Yang, and P. Zhao, On the prime labeling of generalized Petersen graph $P(n, 3)$, *Int. J. Contemp. Math. Sci.,* **6** (2011) 1783-1800. https://m-hikari.com/ijcms-2011/33-36-2011/haqueIJCMS33-36-2011.pdf

[1367] K. M. M. Haque, X. Lin, Y. Yang, and P. Zhao, Prime labeling on Knödel graphs $W_{3,n}$, *Ars Combin.,* **109** (2013) 113-128. https://combinatorialpress.com/article/ars/Volume%20109/volume-109-paper-8.pdf

[1368] F. Harary, Sum graphs and difference graphs, *Congr. Numer.*, **72** (1990) 101-108.

[1369] F. Harary, Sum graphs over all the integers, *Discrete Math.,* **124** (1994) 99-105. https://core.ac.uk/download/pdf/82165654.pdf

[1370] F. Harary, I. Hentzel, and D. Jacobs, Digitizing sum graphs over the reals, *Caribb. J. Math. Comput. Sci.,* **1** (1991) 1-4. https://www.researchgate.net/publication/259715778_Digitizing_Sum_Graphs_Over_the_Real

[1371] F. Harary and D. Hsu, Node-graceful graphs, *Comput. Math. Appl.*, **15** (1988) 291-298. https://www.sciencedirect.com/science/article/pii/0898122188902143

[1372] P. Hariprabakaran, A. Manshath, V. Maheswari, and V. Balaji, Coding strategies through three star graphs and mean labeling, *AIP Conference Proceedings* 2519, 020004 (2022); https://aip.scitation.org/doi/pdf/10.1063/5.0109709

[1373] M. Harminc, On a characterization of graphs by average labelings, *Discuss. Math. Graph Theory,* **17** (1997) 133-136. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1046

[1374] M. Harminc and R. Soták, Graphs with average labellings, graph theory (Prague, 1998) *Discrete Math.,* **233** (2001) 127-132. doi:10.1016/S0012-365X(00)00232-6

[1375] J. Harrington and T. W. H. Wong, On super totient numbers and super totient labelings of graphs, *Disc. Math.,* **343**(2):111670 (2020), DOI:10.1016/j.disc.2019.111670 https://www.sciencedirect.com/science/article/pii/S0012365X19303486?via%3Dihub

[1376] J. Harrington, E. Henninger-Voss, K. Karhadkar, E. Robinson, and T. W. H. Wong, Sum index and difference index of graphs, (2020). https://arxiv.org/abs/2008.09265v2

[1377] J. Harrington, E. Henninger-Voss, K. Karhadkar, E. Robinson, and T. W. H. Wong, Sum index and difference index of graphs, Sum index and difference index of graphs, *Discrete Appl. Math.*, **325** (2023) 262–283. https://arxiv.org/abs/2104.06959

[1378] N. Hartsfield and G. Ringel, *Pearls in Graph Theory*, Academic Press, San Diego, 1990.

[1379] N. Hartsfield and W. F. Smyth, The sum number of complete bipartite graphs, in *Graphs and Matrices* (ed. R. Rees), Marcel Dekker (1992) 205-211.

[1380] D. O. Haryeni, Z. Y. Awanis, M. Bača, and A. Semaničová-Feňovčiková, Modular version of edge irregularity strength for fan and wheel graphs, *Symmetry*, **14**(12):2671 (2022) doi:10.3390/sym14122671

[1381] R. Hashi, I. T. Tarawneh, and M. N. Husin, A remark on the edge irregularity strength of corona proguct of two paths, *J. Math. Sci. and Information,* **2**(1) (2022) 51-58. https://journal.umt.edu.my/index.php/jmsi/article/view/288/214

[1382] J. Haslegrave, Proof of a local antimagic conjecture, *Discrete Math. Theor. Comput. Sci.,* **20**(1) (2018), Paper No. 18, 14 pp. https://dmtcs.episciences.org/4550/pdf

[1383] J. Haslegrave, Sum index, difference index and exclusive sum number of graphs, *Graphs Combin.,* **39**(2), Paper No. 32, 12 (2023) 12pp. doi:10.1007/s00373-023-02624-0

[1384] R. Hasni, I. Tarawneh, M. K. Siddiqui, A. Raheem, and M. A. Asim, Edge irregular k-labeling for disjoint union of cycles and generalized prisms, *Malaysian J. Math. Sci.,* **15**(1) (2021) 79-90. https://mjms.upm.edu.my/fullpaper/2021-January-15-1/Hasni%2C%20R.-79-90.pdf?utm_source=chatgpt.com

[1385] H. Hatima, N. Hinding, and M. Nur, Harmonious labeling on helm graphs, *Daya Matematis: J. Inovasi Pendidikan Matematika*, **10**(3) (2022) 240-245. doi:DOI:10.26858/jdm.v10i3.40683

[1386] M. Haviar and S. Kurtulík, A new class of graceful graphs: k-enriched fan graphs and their characterisations, *CUBO, A Math. J.,* **23**(02)(2021) 313-331. https://www.scielo.cl/pdf/cubo/v23n2/0719-0646-cubo-23-02-313.pdf

[1387] P. Haxell, O. Pikhurko, and A. Taraz, Primality of trees *J. Combinatorics,* **2** (2011) 481-500. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=3e5162fe9a9740541e5b225c0181ade0c6772c1c

[1388] M. He, The gracefulness of the graph $2C_n$, *Neimenggu Daxue Xuebao Ziran Kexue*, **26** (1995) 247-251.

[1389] W. He, L. Wang, H. Mi, Y. Shen, and X. Yu, Integral sum graphs from a class of trees, *Ars Combin.,* **70** (2004) 197-205. https://combinatorialpress.com/article/ars/Volume%20070/vol-70-paper%2018A.pdf

[1390] W. He, X. Yu, H. Mi, Y. Sheng, and L. Wang, The (integral) sum number of the graph $K_n \backslash E(K_r)$ for $K_r \subset K_n$, *Discrete Math.,* **243** (2002) 241-252. doi:10.1016/S0012-365X(01)00218-7

[1391] Y. He, L. Shen, Y. Wang, Y. Chang, Q. Kang, and X. Yu, The integral sum number of complete bipartite graphs $K_{r,s}$, *Discrete Math.,* **239** (2001) 137-146. https://www.sciencedirect.com/science/article/pii/S0012365X00001722?via%3Dihub

[1392] R. Hebbar and B. Sooryanarayana, Radio multiplicative number of certain classes of transformation graphs, *Int. J. Math. Appl.*, **7**(4) (2019) 127-148. https://www.researchgate.net/publication/338477914_Radio_Multiplicative_Number_of_Certain_Classes_of_Transformation_Graphs

[1393] F. F. Hadiputra and T. Maryati, A note on local edge antimagic chromatic number of graphs, *Proyecciones J. Math.,* **43**(2) (2024) 447-458. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/6014/4479

[1394] F. F. Hadiputra, K. A. Sugeng, D. Silaban, T. K. Maryati, and D. Froncek, Chromatic number of super vertex local antimagic total labelings of graphs, *Electron. J. Graph Theory Appl.,* (EJGTA) **9**(2) (2021) 485-498. https://www.ejgta.org/index.php/ejgta/article/view/1236/pdf_194

[1395] P. T. Headley, *Elect. J. Graph Th. Appl.*, The distance magic property and two families of Cartesian product graphs, **13**(1) (2025). https://www.ejgta.org/index.php/ejgta/article/view/858

[1396] D. Hefetz, Anti-magic graphs via the combinatorial nullstellensatz, *J. Graph Theory,* **50** (2005) 263-272. https://www.researchgate.net/publication/220549101_Anti-magic_graphs_via_the_Combinatorial_NullStellenSatz

[1397] D. Hefetz, T. Mütze, and J. Schwartz, On antimagic directed graphs, *J. Graph Theory,* **64** (2010) 219-232. https://www.researchgate.net/publication/220548802_On_Antimagic_Directed_Graphs

[1398] D. Hefetz, A. Saluz, and H. Tran, An application of the combinatorial nullstellensatz to a graph labeling problem, *J. Graph Theory,* **65** (2010) 70-82. https://onlinelibrary.wiley.com/doi/epdf/10.1002/jgt.20466

[1399] A. I. Hefnawy, E. A. Elsid, and S. Euat Tallah, Cordial labeling for the sum and union of the second power of paths and stars, *Util. Math.,* **108** (2018) 283-291.

[1400] S. M. Hegde, Additively graceful graphs, *Nat. Acad. Sci. Lett.*, **12** (1989) 387-390. https://www.researchgate.net/publication/266697941_Additively_graceful_graphs

[1401] S. M. Hegde, On indexable graphs, *J. Combin. Inf. Sci. Sys.,* **17** (1992) 316-331. https://www.researchgate.net/publication/265359577_On_indexable_graphs

[1402] S. M. Hegde, On K-sequential graphs, *Nat. Acad. Sci. Lett.*, **16** (1993) 299-301.

[1403] S. M. Hegde, On harmonious labelings of graphs, *KREC Research Bull.,* **5** (1996) 15-18.

[1404] S. M. Hegde, On (k, d)-graceful graphs, *J. Combin., Inform. Sys. Sci.*, **25** (2000) 255-265. https://www.researchgate.net/publication/305329191_On_k_d_graceful_graphs

[1405] S. M. Hegde, Geometric graphs, *KREC Research Bulletin*, **9** (2000) 27-34.

[1406] S. M. Hegde, Further results on set sequential and set graceful graphs, *Electronic Notes Discrete Math.*, **15** (2003) 100-104. https://www.sciencedirect.com/science/article/pii/S157106530400544X

[1407] S. M. Hegde, On multiplicative labelings of graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 83-96.

[1408] S. M. Hegde, On set labelings of graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 97-108

[1409] S. M. Hegde, Set colorings of graphs, *Europ. J. Combin.,* **30** (2009) 986-995. https://www.sciencedirect.com/science/article/pii/S0195669808001492?utm_source=chatgpt.com

[1410] S. M. Hegde, On sequentially additive graphs, unpublished.

[1411] S. M. Hegde and Kumudakshi, Construction of graceful digraphs using algebraic structures, *J. Discrete Math. Sci. Cryptogr.* **99** (2016) 103-116.

[1412] S. M. Hegde and Kumudakshi, Graceful digraphs and complete mappings, *Elect. Notes Discr. Math.*, **48** (2015) 151-156. https://doi.org/10.1016/j.endm.2015.05.021

[1413] S. M. Hegde and Kumudakshi, Further results on graceful directed graphs, *Electronic Notes Discrete Math.,* **53** (2016) 149-154. https://www.sciencedirect.com/science/article/pii/S1571065316300671

[1414] S. M. Hegde and M. Miller, Further results on sequentially additive graphs, *Discuss. Math. Graph Theory,* **27** (2007) 251-268. https://www.researchgate.net/publication/220468862_Futher_results_on_sequetially_additive_graphs

[1415] S. M. Hegde and T. S. Murthy, A partial solution to cordial tree conjecture, *J. Discrete Math. Sci. Cryptogr.,* **17**(3) (2014) 257-263. https://www.researchgate.net/publication/265390236_A_Partial_Solution_to_Cordial_Tree_Conjecture

[1416] S. M. Hegde and S. Shetty, Sequential and magic labeling of a class of trees, *Nat. Acad. Sci. Letters,* **24** (2001) 137-141. https://www.researchgate.net/publication/266240373_Sequential_and_magic_labeling_of_a_class_of_trees

[1417] S. M. Hegde and P. Shankaran, Geometric labeled graphs, *AKCE J. Graphs Combin.,* **5** (2008) 83-97.

[1418] S. M. Hegde and P. Shankaran, On triangular sum labelings of graphs, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 109-115. https://www.researchgate.net/publication/343058448_On_Triangular_Sum_Labelings_of_Graphs

[1419] S. M. Hegde and S. Shetty, On graceful trees, *Appl. Math. E-Notes,* **2** (2002) 192-197. https://www.researchgate.net/publication/242444232_On_Graceful_Trees

[1420] S. M. Hegde and S. Shetty, On arithmetic graphs, *Indian J. Pure Appl. Math.,* **33** (2002) 1275-1283. https://www.researchgate.net/profile/Suresh-Hegde/publication/265359361_On_arithmetic_graphs/links/541314600cf2788c4b35900a/On-arithmetic-graphs.pdf

[1421] S. M. Hegde and S. Shetty, On magic strengths of graphs, *KREC Research Bull.,* **11** (2002) 15-21.

[1422] S. M. Hegde and S. Shetty, On magic graphs, *Australas. J. Combin.,* **27** (2003) 277-284. https://ajc.maths.uq.edu.au/pdf/27/ajc_v27_p277.pdf

[1423] S. M. Hegde and S. Shetty, Combinatorial labelings of graphs, *Applied Math. E-Notes,* **6** (2006) 251-258. https://www.researchgate.net/publication/239464525_Combinatorial_Labelings_Of_Graphs

[1424] S. M. Hegde and S. Shetty, Strongly indexable graphs and applications, *Discrete Math.,* **309** (2209) 6160-6168. https://www.sciencedirect.com/science/article/pii/S0012365X09002854

[1425] S. M. Hegde and S. Shetty, Strongly k-indexable and super edge magic labelings are equivalent, unpublished

[1426] S. M. Hegde, S. Shetty, and P. Shankran, Further results on super edge-magic deficiency of graphs, *Ars Combin.,* **99** (2011) 487-502. https://combinatorialpress.com/article/ars/Volume%2099/volume-99-paper-42.pdf

[1427] S. M. Hegde and Shivarajkumar, On graceful unicyclc wheels, *Ars Combin.,* **117** (2014) 47-64. https://combinatorialpress.com/article/ars/Volume%20117/volume-117-paper-4.pdf

[1428] S. M. Hegde and Shivarajkumar, On k-graceful digraphs, *Util. Math.* **95** (2014) 161-173. https://www.researchgate.net/publication/283252990_On_k-graceful_digraphs

[1429] S. M. Hegde and Vasudeva, On mod difference labelings of digraphs, *AKCE J. Graphs Combin.,* **6** (2009) 79-84. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088874

[1430] S. M. Hegde and Vasudeva, An algorithm for sum labeling of graphs, (2009) unpublished. https://www.academia.edu/62997890/An_algorithm_for_sum_labeling_of_graphs

[1431] K. Heinrich and P. Hell, On the problems of bandsize, *Graphs and Combin.*, **3** (1987) 279-284. https://link.springer.com/article/10.1007/BF01788550

[1432] V. Hemalatha and V. M. Selvi, Super geometric mean labeling of some cycle related graphs, *Internat. J. Sci. Engin. Res.*, **6**(11) (2015) 522-529. https://www.ijser.org/researchpaper/SUPER-GEOMETRIC-MEAN-LABELING-OF-SOME-CYCLE-RE pdf

[1433] V. Hemalatha and K. Amuthavalli, Radio geometric mean number of splitting of star and bistar, *Internat. J. Res. Advent Tech.*, **6**(7) (2018) 1696-1700. https://ijrat.org/downloads/Vol-6/july-2018/paper%20ID-67201899.pdf

[1434] V. Hemalatha, V. Mohanaselvi, and K. Amuthavalli, Radio geometric mean labeling of some star like graphs, *J. Informatics Math. Sci.*, **9**(3) (2017) 969-977. https://www.rgnpublications.com/journals/index.php/jims/article/view/1017/761

[1435] V. Hemalatha and V. Mohanaselvi, Radio geometric mean number of some subdivision graphs, *Internat. J. Pure Appl. Math.,* **113**(6) (2017) 362-374. https://acadpubl.eu/jsi/2017-113-pp/articles/6/40.pdf

[1436] H. Hendy, K. A. Sugeng, A. N. M. Salman, and N. Ayunda, Another H-super magic decompositions of the lexicographic product of graphs, *Indonesian J. Combin.,* **2**(2) (2018) 72-81. http://dx.doi.org/10.19184/ijc.2018.2.2.2

[1437] A. Hertz and C. Picouleau, On graceful difference labelings of disjoint unions of circuits, https://pisrt.org/psrpress/j/odam/2019/3/5/on-graceful-difference-labelings-of-disjoint-unions-of-circuits.pdf

[1438] K. E. Hilda and J. Jeba Jesintha, Shell flower graphs: felicitous, *Internat. J. Pure Appl. Math.,* **120**(8) (2018) 179-188. https://acadpubl.eu/hub/2018-120-8/2/19.pdf https://www.acadpubl.eu/hub/2018-120-8/2/19.pdf

[1439] K. E. Hilda and S. Krithika, New results on cycle related sum divisor cordial graphs, *Proceed. Internat. Confer. Indian Contribution Math.*, – Ancient & Modern, First Edition, (2022) 64-67. https://www.researchgate.net/publication/374738914_NEW_RESULTS_ON_CYCLE_RELATED_SUM_DIVISOR_CORDIAL_GRAPHS

[1440] Z. R. Himami, D. R. Silaban, On local antimagic vertex coloring of corona products related to friendship and fan graph, *Indon. J. Combin.*, **5**(2) (2021) 110-121. https://www.ijc.or.id/index.php/ijc/article/view/161/129

[1441] N. Hinding, N. Sunusi, Y. Syahrani, and H. M. Buhari, Development of navigation systems in optimal utilization of transportation routes in topology helm graphs using graph labeling, *Edelweiss Appl. Sci. Tech.,* **9**(1) (2025) 481-492. https://doi.org/10.55214/25768484.v9i1.4161/1623

[1442] M.-H. Ho and W. C. Shiu, Full Friendly index sets and full product-cordial index sets of some permutation Petersen graphs, *J. Combin. and Number Theory,* **5**(3) 227-244. https://www.researchgate.net/publication/324727649_Full_friendly_index_sets_and_full_product-cordial_index_sets_of_some_permutation_Petersen_graphs

[1443] Y. S. Ho and S. M. Lee, Some initial results on the supermagicness of regular complete k-partite graphs, *J. Combin. Math. Combin. Computing,* **39** (2001) 3-17. https://combinatorialpress.com/article/jcmcc/Volume%20039/vol-039-paper%201.pdf

[1444] Y. S. Ho, S. M. Lee, and H. K. Ng, On friendly index sets of root-unions of stars by cycles, *J. Combin. Math. Combin. Comput.,* **62** (2007) 97-120. https://combinatorialpress.com/article/jcmcc/Volume%20062/vol-062-paper%209.pdf

[1445] Y. S. Ho, S. M. Lee, and H. K. Ng, On the friendly index sets of regular windmills, unpublished.

[1446] Y. S. Ho, S. M. Lee, and E. Seah, on the edge-graceful (n, kn)-multigraphs conjecture, *J. Combin. Math. Combin. Computing,* (1991) 141-147. https://combinatorialpress.com/article/jcmcc/Volume%2009/vol-009-paper%2013.pdf?utm_source=chatgpt.com

[1447] Y. S. Ho, S. M. Lee, and S. C. Shee, Cordial labellings of unicyclic graphs and generalized Petersen graphs, *Congr. Numer.*, **68** (1989) 109-122.

[1448] Y. S. Ho, S. M. Lee, and S. C. Shee, Cordial labellings of the cartesian product and composition of graphs, *Ars Combin.*, **29** (1990) 169-180.

[1449] C. Hoede and H. Kuiper, All wheels are graceful, *Util. Math.*, **14** (1978) 311.

[1450] J. Holden, D. McQuillan, and J. M. McQuillan, A conjecture on strong magic labelings of 2-regular graphs, *Discrete Math.,* **309** (2009) 4130-4136. https://www.sciencedirect.com/science/article/pii/S0012365X08007103?utm_source=chatgpt.com

[1451] M. Horton, Graceful Trees: Statistics and Algorithms, 2003, Univ. Tasmania, http://eprints.utas.edu.au/19/1/GracefulTreesStatisticsAndAlgorithms.pdf.

[1452] M. Hovey, *A*-cordial graphs, *Discrete Math.*, **93** (1991) 183-194. https://www.sciencedirect.com/science/article/pii/0012365X9190254Y?via%3Dihub

[1453] P. Hrnčiar and A. Haviar, All trees of diameter five are graceful, *Discrete Math.,* **233** (2001) 133-150. https://www.sciencedirect.com/science/article/pii/S0012365X00002338?via%3Dihub

[1454] P. Hrnčiar and G. Monoszova, A new family of graceful trees, preprint.

[1455] C-C. Hsiao, On Graph Labeling Problems of Antimagic Type, M. S. Thesis, TungHai University, 2006.

[1456] C. H. Hsieh, K.-Y. Chen, M.-Y. Jiang, J.-J. Liaw, and J. Shin, Estimation of $PM_{2.5}$ concentration based on support vector regression with improved dark channel prior and high frequency information in images, *IEEE Access,* **10** (2022) 48486-48498. https://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=9767851

[1457] D. F. Hsu, Harmonious labelings of windmill graphs and related graphs, *J. Graph Theory*, **6** (1982) 85-87.

[1458] D. F. Hsu and A. D. Keedwell, Generalized complete mappings, neofields, sequenceable groups and block designs, I, *Pacific J. Math.*, **111** (1984) 317-332. https://www.academia.edu/103942656/Generalized_complete_mappings_neofields_sequenceable_groups_and_block_designs_I

[1459] D. F. Hsu and A. D. Keedwell, Generalized complete mappings, neofields, sequenceable groups and block designs, II, *Pacific J. Math.,* **117** (1985) 291-312. https://www.academia.edu/119815759/Generalized_complete_mappings_neofields_sequenceable_groups_and_block_designs_II

[1460] H. L. Hu, New graceful labelings of the n-cycle C_n and its r-crowns, *Pure Appl. Math.,* (Xi'an) **26** (2010) 454-457.

[1461] H. Hua and C. Raridan, On the edge-balanced index sets of odd/even complete bipartite graphs, *Congr. Numer.,* **219** (2014) 227-232. https://arxiv.org/abs/1405.1673

[1462] N. B. Huamaní, M. Atoche Bravo, and F. Poma, Families of gracefuls spiders with $\ell(2k+1)-k, \ell(2k+1)-k+1$ and $\ell(2k+1)+k+1$ legs, *Cont. Math.,* **6**(1) (2025) 730-742. https://ojs.wiserpub.com/index.php/CM/article/view/5497/2924

[1463] C. Huang, A. Kotzig, and A. Rosa, Further results on tree labellings, *Util. Math.*, **21c** (1982) 31-48.

[1464] C. Huang, J. Li, and L. Sun, Edge-magic total labeling of two joined graphs, *Conference: 2024 IEEE 6th Adv. Inform. Management, Communicates, Elect. Automation Control Confer.* (IMCEC) (2024) https://ieeexplore.ieee.org/document/10575492

[1465] Q. Huang, Harmonious labeling of crowns $C_n \odot K_1$, unpublished.

[1466] T.-Y. Huang, Antimatic Labeling on Spiders, Master's Thesis, Department of Mathematics, National Taiwan University, 2015.

[1467] J. Huang and S. Skiena, Gracefully labeling prisms, *Ars Combin.,* **38** (1994) 225-242. https://combinatorialpress.com/article/ars/Volume%20038/volume-38-paper-23.pdf

[1468] P.-Y. Huang, T.-L. Wong, and X. Zhu, Weighted-1-antimagic graphs of prime power order, *Discrete Math.,* **312**(14) (2012) 2162-2169. https://core.ac.uk/download/pdf/82488747.pdf

[1469] R. Huang, M. F. Hanif, M. K. Siddiqui, and M. F. Nadeem, On super magic algorithm for union of comb and star graph, *Konuralp J. Math.*, **11** (1) (2023) 82-89. https://dergipark.org.tr/en/pub/konuralpjournalmath/issue/76134/929700

[1470] M. I. Huilgol and V. Sriram, Square sum labeling of disjoint union of graphs, *J. Graph Labeling,* **2**(2) (2016) 103-106. https://www.researchgate.net/publication/291691469_Square_sum_labeling_of_disjoint_union_of_graphs

[1471] N. S. Hungund and D. G. Akka, Reverse super edge-magic strength of some new classes of graphs, *J. Discrete Math. Sci. Cryptography*, **16**(1) (2013) 19-29, DOI: 10.1080/09720529.2013.778462 http://dx.doi.org/10.1080/09720529.2013.778462

[1472] F. H. Husna and D. Rahmadani, Elegant labeling of the lotus graph and its variations, *Jambura J. Math.*, **8**(2)(2026) 257–264. https://doi.org/10.37905/jjom.v8i2.39844

[1473] M. Hussain, K. Ali, M.T. Rahim, and E. T. Baskoro, On (a, d)-vertex-antimagic labelings of Harary graphs, *Util. Math.,* **83** (2010) 73-80. https://www.researchgate.net/publication/266701597_On_ad-vertex-antimagic_total_labeling_of_Harary_graphs

[1474] M. Hussain, E.T. Baskoro, and K. Ali, On super antimagic total labeling of Harary graph, *Ars Combin.,* **104** (2012) 225-233. https://combinatorialpress.com/article/ars/Volume%20104/volume-104-paper-19.pdf

[1475] M. Hussain, E. Baskoro, and Slamin, On super edge-magic total labeling of banana trees, *Util. Math.,* **79** (2009) 243-251. https://www.researchgate.net/publication/235798012_On_super_edge-magic_total_labeling_of_banana_trees

[1476] Z. Hussaina, K. Shabbira, and K. Ahmada, Computation for lucky number of some families of planar graphs, *Mathematica Universalis*, **1**(1) (2026) 26–45. https://mathematicauniversalis.com/index.php/MU/article/view/3/2

[1477] M. Hussainn and A. Tabraiz, Super d-anti-magic labeling of subdivided kC_5, *Turkish J. Math.,* **39**(5) (2015) 773-783. https://journals.tubitak.gov.tr/cgi/viewcontent.cgi?article=2133&context=math

[1478] M. Ibrahim, M. Asif, A. Ahmad, and M. K. Siddiqui, Computing the total irregularity strength of wheel related graphs, *Util. Math.,* **108** (2018) 321-338. https://www.researchgate.net/publication/286456461_Total_Edge_Irregularity_Strength_of_Wheel_Related_Graphs

[1479] M. Ibrahim, A. Gulzar, M. Fazil, and M. N. Azhar, On edge H-irregularity strength of hexagonal and octagonal grid graphs, *Hindawin J. Math.*, **2022** Article ID 6047926 (2022) 10 pages.
https://doi.org/10.1155/2022/6047926

[1480] M. Ibrahim, S. Khan, M. A. Asim, and M. Waseem, Total irregularity strength of some cubic graphs, *Proyecciones,* **40**(4) (2021) 905-918. https://www.researchgate.net/publication/350981082_Total_irregularity_strength_of_some_cubic_graphs

[1481] R. Ichishima, S. C. López, and F. A. Muntaner-Batle, On the beta-number of forests with isomorphic components, *Discuss. Math. Graph Theory,* **38** (2018) 683-701. https://upcommons.upc.edu/bitstream/handle/2117/121742/Beta_upccommons.pdf?sequence=1&isAllowed=y

[1482] R. Ichishima, S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, The power of digraph products applied to labelings, *Discrete Math.,* **312** (2012) 221-228. https://core.ac.uk/download/pdf/81955487.pdf

[1483] R. Ichishima, S. C. López, F. A. Muntaner-Batle, and Y. Takahashi, Recent studies on the super edge-magic deficiency of graphs, *Theory Appl. Graphs, 11*(1), Article 1.https://digitalcommons.georgiasouthern.edu/tag/vol11/iss1/1

[1484] R. Ichishima, S. C. López, F. A. Muntaner-Batle, and Y. Takahashi, On the strength and domination number of graphs, *Contrib. Math.,* **8** (2023) 11–15. https://shahindp.com/Contrib_Math/v8/CM23_v8_pp11-15.pdf

[1485] R. Ichishima and F. A. Muntaner-Batle, On the strong sequential number of graphs, *Util. Math.*, 116 (2020) 247-259.

[1486] R. Ichishima and F. A. Muntaner-Batle, On the strong beta-number of galaxies with five components, preprint.

[1487] R. Ichishima and F. A. Muntaner-Batle, On the super edge-magicness of graphs with a specific degree sequence, *Open J. Discret. Appl. Math.,* (2023) **6**(3), 22—25. arXiv:2308.10661v1 https://pisrt.org/psrpress/j/odam/2023/3/on-the-super-edge-magicness-of-graphs-with-a-specific-degree-sequence.pdf

[1488] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, Enlarging the classes of super edge-magic 2-regular graphs, *AKCE Int. J. Graphs Comb.*, **10**(2) (2013) 129-146. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088730?utm_source=chatgpt.com

[1489] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, The measurements of closeness to graceful graphs, *Australas. J. Combin.*, **62** (2015) 197-210. https://ajc.maths.uq.edu.au/pdf/62/ajc_v62_p197.pdf

[1490] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, The consecutively super edge-magic deficiency of graphs and related concepts, *Electron. J. Graph Theory Appl.,* (EJGTA) **8**(1) (2020) 71-92. https://www.ejgta.org/index.php/ejgta/article/view/785

[1491] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, On the beta-number of joins of graphs, *Australas. J. Combin.*, **69** (3) (2017) 402-409. https://ajc.maths.uq.edu.au/pdf/69/ajc_v69_p402.pdf

[1492] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, The harmonious number of graphs and related graph related parameters, *Bull. Inst. Combin. Appl.*, **82** (2018) 80-95. https://bica.the-ica.org/Volumes/82//Reprints/BICA2017-19-Main-Reprint.pdf

[1493] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, New parameters for studying graceful properties of graphs, *Elect. Notes Discrete Math.,* **60** (2017) 3-10. https://www.sciencedirect.com/science/article/pii/S1571065317300872

[1494] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, Bounds for the strength of graphs, *Australasian J. Combin.,* **72**(3) (2018) 492-508. https://ajc.maths.uq.edu.au/pdf/72/ajc_v72_p492.pdf

[1495] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, On the strength of some trees, *AKCE Internat. J. Graphs Combin.,* **17**(1) 486-494. https://www.tandfonline.com/doi/epub/10.1016/j.akcej.2019.06.002?needAccess=true

[1496] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, On the strong beta-number of galaxies with three and four components, *AKCE Internat. J. Graphs Combin.,* **17**(1) (2019) 314-323. https://www.tandfonline.com/doi/epub/10.1016/j.akcej.2019.03.004?needAccess=true

[1497] R. Ichishima, F. A. Muntaner-Batle, A. Oshima, A result on the strength of graphs by factorizations of complete graphs, *Discrete Math. Lett.,* 8 (2022) 78-82. https://www.semanticscholar.org/reader/1ad7349ac30e24b07bef0fe930f781f25056ba42

[1498] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, A result on the strength of graphs by factorizations of complete graphs, *Discrete Math. Lett.,* **8** (2022) 78-82. https://www.dmlett.com/archive/v8/DML22_v8_pp78-82.pdf

[1499] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, On the beta-number of the coronas of graphs, *Memoirs Kokushikan Univer. Inform. Sci.*, **43** 1-7. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088730?utm_source=chatgpt.com

[1500] R. Ichishima, F. A. Muntaner-Batle, and A. Oshima, On the beta-number and gamma-number of galaxies, *Discrete Math. Lett.,* **1** (2019) 49-53. https://www.dmlett.com/archive/DML19_v1_p.4953.pdf

[1501] R. Ichishima, F. A. Muntaner-Batle, and O. Oshima, Minimum degree conditions for the strength and bandwidth of graphs, *Discrete Appl. Math.,* **320** (2022) 191-198. https://www.sciencedirect.com/science/article/abs/pii/S0166218X22001597

[1502] R. Ichishima, F. A. Muntaner-Batle, A. Oshima, and Y. Takahashi, The strength of graphs and related invariants, *Memoirs Kokushikan Univ. Inf. Sci.,* **41** (2020), 1–8. https://comb-opt.azaruniv.ac.ir/article_14720_9c26bd36c60a54859c209dab3a0cd6ee.pdf

[1503] R. Ichishima, F.-A. Muntaner-Batle, A. Oshima, and Y. Takahashi, On certain labelings of disjoint unions of cycles and stars, *Bull. Math. Soc. Sci. Math. Roumanie,* **69**(2) (117) (2026) 199–217.

[1504] R. Ichishima, F. A. Muntaner-Batle, and M. Rius-Font, Embedding trees into super edge-magic trees, *Bull. Inst. Combin. Appl.,* **52** (2008) 51-60. https://www.researchgate.net/publication/268711436_Embedding_trees_into_super_edge-magic_trees

[1505] R. Ichishima, F. A. Muntaner-Batle, and M. Rius-Font, Bounds on the size of super edge-magic graphs depending on the girth, *Ars Combin.,* **119** (2016) 129-133. https://www.researchgate.net/publication/43244477_Bounds_on_the_size_of_super_edge-magic_graphs_depending_on_the_girth

[1506] R. Ichishima, A. Oshima, and Y. Takamashi, The strength sum of graphs and related invariants, (2020), https://www.researchgate.net/publication/357754018_The_strength_of_graphs_and_related_invariants

[1507] R. Ichishima, F. A. Muntaner-Batle, and Y. Takahashi, On the strength and independence number of powers of paths and cycles, *Comm. in Comb. Optimi,*, **10**(4) (2025) 717-728. https://comb-opt.azaruniv.ac.ir/article_14720_9c26bd36c60a54859c209dab3a0cd6ee.pdf

[1508] R. Ichishima, F. A. Muntaner-Batle and Y. Takahashi, A method to compute the strength using bounds, arXiv:2311.14923v1

[1509] R. Ichishima, F. Muntaner-Batle, and Y. Takahashi, Some new results concerning the valences of (super) edge-magic graphs, *Bull. Math. Soc. Sci. Math. Roumanie*, **68** (116)(2) (2025) 185–200.

[1510] R. Ichishima, F. A. Muntaner-Batle, and Y. Takahashi, On the strength and independence number of powers of paths and cycles, *Communications Combin. Optimization,* to appear. https://comb-opt.azaruniv.ac.ir/article_14720.html

[1511] R. Ichishima and A. Oshima, On the beta-number of linear forests with an even number of components, *AKCE Int. J. Graphs Comb.,* **15**(3) (2018) 238-241. https://www.tandfonline.com/doi/pdf/10.1016/j.akcej.2017.11.003

[1512] R. Ichishima and A. Oshima, On partitional labelings of graphs, *Math. Comput. Sci.,* **3** (2010) 39-45. https://link.springer.com/content/pdf/10.1007/s11786-009-0008-7.pdf?pdf=previewhttps://link.springer.com/content/pdf/10.1007/s11786-009-0008-7.pdf?pdf=preview

[1513] R. Ichishima and A. Oshima, On partitional and other related graphs, *Math. Comput. Sci.,* **5** (2011) 41-50. https://www.researchgate.net/publication/43244477_Bounds_on_the_size_of_super_edge-magic_graphs_depending_on_the_girth

[1514] R. Ichishima and A. Oshima, On the super edge-magic deficiency of 2-regular graphs with two components, *Ars Combin.,* **129** (2016) 437-447. https://combinatorialpress.com/article/ars/Volume%20129/volume-129-paper-34.pdf

[1515] R. Ichishima and A. Oshima, On the super edges-magic deficiency and α-valuations of graphs, *J. Indones. Math. Soc.,* Special Edition (2011) 59-69. https://www.researchgate.net/publication/265779875_On_the_super_edge-magic_deficiency_and_a-valuations_of_graphs

[1516] R. Ichishima and A. Oshima, Bounds for the gamma-number of graphs, *Util. Math.,* **109** (2018) 313-325.

[1517] R. Ichishima, A. Oshima, and Y. Takahashi, On the strength and independence number of graphs, *Contrib. Math.,* **6** (2022) 25-29. https://pdfs.semanticscholar.org/e61b/db557758d50073b6cd17b78143d89baf0861.pdf

[1518] R. Ichishima, A. Oshima, and Y. Takahashi, Some new results on the edge-strength and strength of graphs, *Discrete Math. Lett*, **12** (2023) 22-25. https://www.dmlett.com/archive/v12/DML23_v12_pp22-25.pdf

[1519] R. Ichishima, A. Oshima, and Y. Takahashi, On the beta-number of galaxies with symmetric tructure, *Southeast Asian Bull. Math.*, **47** (2023) 205-218. https://www.researchgate.net/publication/369269254_Southeast_Asian_Bulletin_of_Mathematics_On_the_Beta-Number_of_Galaxies_with_Symmetric_Structure

[1520] M. Imran, H. U. Afzal, and A. Q. Baig, On super edge-magic deficiency of volvox and dumbbell graphs, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 112-119. https://www.researchgate.net/publication/304366919_On_super_edge-magic_deficiency_of_volvox_and_dumbbell_graphs

[1521] M. Imran, A. Aslam, S. Zafar, and W. Nazeer, Further results on the edge irregularity strength of graphs, *Indonesian J. Combin.,* **1**(2) (2017) 36-45. doi:10.19184/ijc.2017.1.2.5

[1522] M. Imran, A. Q. Baig, and A. S. Feňovčiková, Some results on super edge-magic deficiency of graphs, *Kragujevac J. Math.,* **44**(2) (2020) 237-249. http://elib.mi.sanu.ac.rs/files/journals/kjm/60/kjmn60p237-249.pdf

[1523] M. Imran, M. Cancan, and Y. Ali, Some comb related mean graphs, *Palestine J. Math.,* **13** 4)(2024) 556-560. https://pjm.ppu.edu/sites/default/files/papers/PJM_13%284%29_2024_556_to_560.pdf

[1524] M. Imran, M. Cancan, Y. Ali, and J. Aslam, On edge irregularity strength of different families of graphs, *Proyecciones J. Math.,* **42**(6) (2023) 1549-1566. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/5873/4401

[1525] M. Imran, M. Cancan, Y. Ali, M. Nadeem, S. Mushtaq, A. Aslam, and R. Riaz, Some comb related cordial graphs, *Inter. J. Res. Publ. Reviews*, **3**(10) (2022) 2334-2339. https://www.researchgate.net/publication/364958790_Some_Comb_Related_Cordial_Graphs

[1526] M. Imran, M. Cancan, M. Nadeem, and Y. Ali, Further results on edge irregularity strength of some graphs, *Proyeccione J. Math.,* **43**(1) (2024) 133-151. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/6175/4458

[1527] M. Imran, M. Cancan, M. Nadeem, and M. F. Nadeem, Results on super edge magic deficiency of some well known classes of finite graphs, *Proyecciones J. Math.,* **43**(5) (2024) 1075-1096. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/6329/4572

[1528] M. Imran, M. R. Farahani, M. Cancan, and M. Alaeiyan, Computation of lucky number of comb graphs Cf_w, Cg_w, Ch_w and triangular snake, alternate triangular snake graphs, *Power System Tech.,* **48**(1) (2024) 1381-1394. https://powertechjournal.com/index.php/journal/article/view/399

[1529] M. Imran, M. K. Siddiqui, and M. Numan, Super d-antimagic labeling of uniform subdivision of wheel, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.,* **77** (2015)(2) 227-240. https://www.scientificbulletin.upb.ro/rev_docs_arhiva/fulle56_708694.pdf

[1530] N. Inayah, Another antimagic decomposition of generalized Petersen graph, *Indonesian J. Pure and Appl. Math.,* **3**(2) (2021) 92-100. http://journal.uinjkt.ac.id/index.php/inprime/article/view/19954

[1531] N. Inayah, A. Lladó, J. Moragas, Magic and antimagic H-decompositions, *Discrete Math.,* **312**(7) (2012) 1367-1371. https://www.sciencedirect.com/science/article/pii/S0012365X11005486?via%3Dihub

[1532] N. Inayah, A. N. M. Salman, and R. Simanjuntak, On (a,d)-H-antimagic coverings of graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 273-281. https://combinatorialpress.com/article/jcmcc/Volume%20071/vol-071-paper%2021.pdf

[1533] N. Inayah, R. Simanjuntak and A. N. M. Salman, Super $(a,d)-H$-antimagic total labelings for shackles of a connected graph H, *Australasian J. Combin.,* **57** (2013) 127-138. https://ajc.maths.uq.edu.au/pdf/57/ajc_v57_p127.pdf

[1534] N. Inayah, I. W. Sudarsana, S. Musdalifah, and N. D. Mangesa, On super mean labeling for total graph of path and cycle, *Internat. J. Math. Math. Sci.,* **7-8** (2018) 1-5. https://www.semanticscholar.org/reader/3268bd724eeac0ed61db4cc09ffc709bfee4dffa

[1535] N. Inayah, F. Susanto, A. Semaničová-Feňovčíková On H-antimagic coverings for m-shadow and closed m-shadow of connected graphs, *Heliyon* **7** (10) (2021), e08203. https://www.researchgate.net/publication/355414881_On_H-antimagic_coverings_for_m-shadow_and_closed_m-shadow_of_connected_graphs

[1536] P. Indira, B. Selvam, and K. Thirusangu, Proper lucky labeling and lucky edge labeling for the extended duplicate graph of quadrilateral snake, *Math. Internat. J. Math. Trends Tech.,* **67** (2021) 60-65. doi:10.14445/22315373/IJMTT-V67I1P50 Corpus ID: 234050223

[1537] D. Indriati and T. Azzahra, Edge irregular reflexive labelings on Mongolian tent graph $(M_{m,3})$ and double quadrilateral snake graph, *Bareking J. Ilmu Matematika dan Terapan*, **17**(4) (2023) 1933-1940. https://www.researchgate.

net/publication/376836681_EDGE_IRREGULAR_REFLEXIVE_LABELING_ON_MONGOLIAN_TENT_GRAPH_M_m3_AND_DOUBLE_QUADRILATERAL_SNAKE_GRAPH

[1538] D. Indriati, I. E. W. Widodo, and I. Rosyida, Edge irregular reflexive labeling on corona of path and other graphs, *J. Phys. Conf. Ser.*, **1489**(1) (2020). https://iopscience.iop.org/article/10.1088/1742-6596/1489/1/012004/pdf

[1539] D. Indriati, I. E. W. Widodo, and K. A. Sugeng, On the total edge irregularity strength of generalized helm, *AKCE Int. J. Graphs Comb.*, **10**(2) (2013) 147-155. https://www.researchgate.net/publication/260021098_On_the_total_edge_irregularity_strength_of_generalized_helm

[1540] D. Indriati, I. E. W. Widodo, I. Wijayanti, and K. A. Sugeng, Totally irregular total labeling of some caterpillar graphs, *Electron. J. Graph Theory Appl.,* (EJGTA) **8**(2) (2020) 247-254. https://www.ejgta.org/index.php/ejgta/article/view/1062/pdf_140

[1541] D. Indriati, I. E. W. Widodo, I. E. Wijayanti, K. A. Sugeng, and M. Bača, On total edge irregularity strength of generalized web graphs and related graphs, *Math. Comput. Sci.,* **9**(2) (2015) 161-167. https://www.ejgta.org/index.php/ejgta/article/view/1062/pdf_140

[1542] D. Indriati, I. E. W. Widodo, I. E. Wijayanti, K. A. Sugeng, M. Bača, and A. Semaničová-Feňovčíková, The total vertex irregularity strength of generalized helm graphs and prisms with outer pendant edges, *Austral. J. Combin.,* **65**(1) (2016) 14-26. https://ajc.maths.uq.edu.au/pdf/65/ajc_v65_p014.pdf

[1543] D. Indriati, I. E. W. Widodo, I. E. Wijayanti, and K. A. Sugeng, On total irregularity strength of star graphs, double-stars, and caterpillar, *AIP Confer. Proceed.*, **1707**(1) (2016) 020008(1)-020008(6).

[1544] P. Inpoonjai, Degree-magic labelings on the join and composition of complete tripartite graphs, *Comm. Math. Appl.,* textbf10(3) (2019) 391-402. https://www.rgnpublications.com/journals/index.php/cma/article/view/1157/0

[1545] P. Inpoonjai, Supermagic triple graphs, *Int. J. Anal. App.*, (2026) **24**(35). 10pp. https://etamaths.com/index.php/ijaa/article/view/4764

[1546] M. Irfan, M. Bača, P. Kovář, and A. Semaničová-Feňovčíková, On reflexive edge strength of generalized prism graphs, *Electronic J. Graph Theory Appl.*, **10**(2) 415-424. doi:10.5614/ejgta.2022.10.2.6

[1547] M. Irfan and A. Semaničová-Feňovčová, On total labelings of graphs with prescribed weights, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 191-199. https://www.researchgate.net/publication/304713348_On_total_labelings_of_graphs_with_prescribed_weights

[1548] R. M. Irudhaya, A. Chitra, and A. N. Murugan, Lucky edge labeling of some special graphs, *Internat. J. Recent Res. Aspects, Special Issue: Conscientious Comput. Tech.,* (2018) 274-277. www.ijrra.net/April2018/ConsComp2018_66.pdf

[1549] R. M. Irudhaya, A. Chitra, A. N. Murugan, Lucky edge labeling of star related graphs, *J. Comput. Math. Sci.*, **9**(9) (2018) 1124-1131.

[1550] G. Isaak, R. Jamison, and D. Narayan, Greedy rankings and arank numbers, *Inform. Process. Lett.,* **109** (2009) 825-827. https://www.sciencedirect.com/science/article/pii/S0020019009001306

[1551] J. Ivančo, On supermagic regular graphs, *Math. Bohemica,* **125** (2000) 99-114. https://dml.cz/bitstream/handle/10338.dmlcz/126259/MathBohem_125-2000-1_6.pdf

[1552] J. Ivančo, Magic and supermagic dense bipartite graphs, *Discuss. Math. Graph Theory,* **27** (2007) 583-591. https://www.researchgate.net/publication/220468778_Magic_and_supermagic_dense_bipartite_graph

[1553] J. Ivančo, A construction of supermagic graphs, *AKCE J. Graphs Combin.,* **6** (2009) 91-102. https://combinatorialpress.com/article/jcmcc/Volume%20042/vol-042-paper%2010.pdf

[1554] J. Ivančo, Supermagic generalized double graphs, *Discuss. Math. Graph Th.,* **36** (2016) 211-225. https://www.researchgate.net/publication/284204510_Supermagic_generalized_double_graphs

[1555] J. Ivančo, On k-total edge product cordial graphs,*Australasian J. Comb.,* **67**(3) (2017) 476-485. https://ajc.maths.uq.edu.au/pdf/67/ajc_v67_p476.pdf

[1556] J. Ivančo, and S. Jendroľ, Total edge irregularity strength of trees, *Discuss. Math. Graph Theory,* **26** (2006) 449-456. https://www.researchgate.net/publication/220468817_Total_edge_irregularity_strength_of_trees

[1557] J. Ivančo, P. Kovář, and A. Semaničová-Feňovčová, On the existence of regular supermagic graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 49-64. https://combinatorialpress.com/jcmcc-articles/volume-071/on-the-existence-of-regular-supermagic-graphs/

[1558] J. Ivančo, Z. Lastivkova, and A. Semaničova, On magic and supermagic line graphs, *Math. Bohemica,* **129** (2004) 33-42. https://www.researchgate.net/publication/267441957_On_magic_and_supermagic_line_graphs

[1559] J. Ivančo and I. Lučkaničová, On edge-magic disconnected graphs, *SUT J. Math.,* **38** (2002) 175-184. https://www.researchgate.net/publication/228940969_On_edge-magic_disconnected_graphs

[1560] Ivančo and Polláková, Supermagic graphs having a saturated vertex, *Discuss. Math. Graph Th.,* **34**(1) (2014) 75-84. https://www.researchgate.net/publication/362594315_Supermagic_graphs_having_a_saturated_vertex

[1561] J. Ivančo and A. Semaničova, Some constructions of supermagic non-regular graphs, *Australas. J. Combin.,* **38** (2007) 127-139. https://ajc.maths.uq.edu.au/pdf/38/ajc_v38_p127.pdf

[1562] J. Ivančo and A. Semaničova, Some constructions of supermagic graphs using antimagic graphs, *SUT J. Math.,* **42** (2007) 177-186. https://projecteuclid.org/journals/sut-journal-of-mathematics/volume-42/issue-2/Some-constructions-of-supermagic-graphs-using-antimagic-graphs/10.55937/sut/1262445107.full

[1563] K. Iyappan, A. Loganathan, D. Kumar, $H - E$-Supermagic labelings of some families of graphs, *Nonlinear Studies*, **31**(3) (2024) 957-963. https://www.researchgate.net/publication/383738774_H-E-Supermagic_labelings_of_some_families_of_graphs

[1564] J. Jacob, D. Narayan, E. Sergel, P. Richter, and A. Tran, personal communication (2010).

[1565] R. Jagadesh and J. Baskar Babujee, Super edge bimagic labeling, *Proceed. 3rd Inter. Conf. Math. Comp. Sci.*, 5-6th Jan. 2009, Loyola College, Chennai, India, (2009) 79-82.

[1566] R. Jagadesh and J. Baskar Babujee, Edge vertex prime labeling for some class of graphs, National Conference on Recent Trends in Mathematics and its Applications, 24-25 February 2017, SRM University, Vadapalani, Chennai, India.

[1567] C. Jaisankar, M. Manivel, and E. Keerthana, Lehmer-3 mean labeling of dragon graphs, *Malaya J. Mat.*, **5**(1) (2021) 290-293. https://www.malayajournal.org/articles/MJMS210062.pdf

[1568] P. Jampachon, K. Nakprasit, and T. Poomsa-ard, Graceful labeling of some classes of spider graphs with three legs greater than one, *Thai J. Math.,* **12**(3) (2014) 621-630. https://thaijmath2.in.cmu.ac.th/index.php/thaijmath/article/view/477/473

[1569] R. Janani and T. Ramachandran, On relatively prime edge labeling of graphs, *Engin. Letters,* **30**(2) (2022) 659-665. https://www.engineeringletters.com/issues_v30/issue_2/EL_30_2_30.pdf

[1570] R. Janani and T. Ramachandran, On prime index of a graph, *Ratio Math.*, **48** (2023) 516-527. https://eiris.it/ojs/index.php/ratiomathematica/article/view/1315/pdf

[1571] R. Janani and T. Ramachandran, On edge prime index of a graph, *Commun. Appl. Nonlinear Anal.,* **31**(2) (2024) 42-54. doi:DOI: 10.52783/cana.v31.512

[1572] O. Jane and A. Myom, Totally antimagic total labelling of a complete bipartite graph and its application in end-to-end encryption, *American Internat. J. Res. Sci., Tech., Engin. Math. AIJRSTEM* 17-404. https://www.researchgate.net/publication/377399091_Totally_Antimagic_Total_Labelling_of_a_Complete_Bipartite_Graph_and_its_Application_in_End-to-End_Encryption

[1573] J. H. Jani and V. J. Kaneria, Absolute mean graceful labeling of various m-shadow and m-splitting related graphs, *Utilitas Math.* **127** 93-104. https://combinatorialpress.com/um-articles/vol-127/absolute-mean-graceful-labeling-of-various-m-shadow-and-m-splitting-related-graph

[1574] D. Jani, P. T. Lalchandani, and J. Parejiya, Harmonic mean cordial labeling of join of some graphs, *TANGENCE*, #138 (2026) 207-218.

[1575] M. M. Jariya, Index of labeling, *Internat. J. Math. Trends Tech.*, **657** (2019) 87-96. https://www.ijmttjournal.org/Volume-65/Issue-8/IJMTT-V65I8P509.pdf

[1576] I. Javaid, A. Ismail, M. Salman, A. Ahmad, and Slamin, Labeling of chordal rings, *Util. Math.,* **90** (2013) 61-75. https://www.researchgate.net/publication/258004777_Labeling_of_Chordal_Rings

[1577] I. Javaid, F. Khalid, A. Ahmad, and M. Imran, On a weaker version of sum labeling of graphs, *Math. Rep. (Bucur.)* **16**(66) (2014), no. 3, 413-420. https://www.researchgate.net/publication/264554480_On_a_weaker_version_of_sum_labeling_of_graphs

[1578] M. Javaid, On super edge-antimagic total labeling of subdivided stars, *Discuss. Math. Graph Theory,* **34**(4) (2014) 691-706. https://www.dmgt.uz.zgora.pl/publish/article.php?doi=1764

[1579] M. Javaid, Super (a, d)-EAT labeling of subdivided stars, *AKCE Int. J. Graphs Comb.,* **12**(1) (2015) 14-18. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2015.06.002?scroll=top&needAccess=true#d1e278

[1580] M. Javaid and A. A. Bhatti, On anti-magic vertex labeling of disjoint union of star hyper-graphs, *Canadian J. Sci. and Eng. Math.,* **3**(5) (2012) 202-209.

[1581] M. Javaid, A. A. Bhatti, and M. K. Aslam, Super (a, d)-edge antimagic total labeling of a subclass of trees, *AKCE Internat. J. Graphs Combin.,* **14** (2017) 158-164. https://www.sciencedirect.com/science/article/pii/S0972860016300937

[1582] M. Javaid, A. A. Bhatti, and M. Hussain, On (a, d)-edge-antimagic total labelings of extended w-trees, *Util. Math.,* **87** (2012) 293-303. https://www.researchgate.net/publication/266860718_On_ad-edge-antimagic_total_labelings_of_extended_w-trees

[1583] M. Javaid and A. A. Bhatti, On super (a, d)-edge-antimagic total labelings of generalized extended w-trees, *AKCE Int. J. Graphs Comb.,* **11** (2014) 115-126. https://www.researchgate.net/publication/321078581_On_Super_Edge-Antimagic_Total_Labeling_of_Generalized_Extended_W-Trees

[1584] M. Javaid, A. A. Bhatti, M. Hussain, and K. Ali, Super edge-magic total labeling on a forest of extended w-trees, *Util. Math.,* **91** (2013) 155-162. https://www.researchgate.net/publication/265875516_Super_edge-magic_total_labeling_on_forest_of_extended_w-trees

[1585] M. Javaid, M. Hussain, K. Ali, and K. H. Dar, Super edge-magic total labeling on w-trees, *Util. Math.,* **86** (2011) 183-191. https://www.researchgate.net/publication/266192975_Super_edge-magic_total_labeling_on_w-trees

[1586] M. Javaid, M. Hussain, K. Ali, and M. Shaker, On super edge-magic total labeling on subdivision of trees, *Util. Math.* **89** (2012) 169-177. https://www.researchgate.net/profile/Kashif-Ali-27/publication/267445971_Super_edge-magic_total_labeling_of_subdivision_of_stars/links/54aa6ad70cf274dc0b5c2716/Super-edge-magic-total-labeling-of-subdivision-of-stars.pdf

[1587] A. Javed and M. K. Jamil, 3-total edge product cordial labeling of rhombic grid, *AKCE Int. J. Graphs Comb.,* **16**(2) (2019) 213–221. https://www.researchgate.net/publication/322380597_3-total_edge_product_cordial_labeling_of_rhombic_grid

[1588] S. Javed, A. Riasat, and S. Kanwal, On super edge-magicness and deficiencies of forests, *Util. Math.,* **98** (2015) 149-169. https://www.researchgate.net/publication/292377948_On_super_edge-magicness_and_deficiencies_of_forests

[1589] J. Jayapriya, 0-edge magic labeling of slitting graph, *Global J. Pure and Appl. Math.*, **12**(1) (2016) 421-426. https://www.ripublication.com/gjpam16/gjpamv12n1_39.pdf

[1590] J. Jayapriya, Pair sum modulo labelling of spider graph, *Internat. J. Special Edu.,* **37**(3) (2022) 4205-4211. https://www.researchgate.net/publication/360168225_Pair_Sum_Modulo_Labelling_of_Spider_Graph

[1591] J. Jayapriya and K. Thirusangu, 0-edge magic labeling for some class of graphs, *Indian J. Comput. Sci. Engin.(IJCSE)*, **3**(3) (2012) 425-427. https://www.researchgate.net/publication/267940060_0-EDGE_MAGIC_LABELING_FOR_SOME_CLASS_OF_GRAPHS

[1592] C. Jayasekaran and C. David Raj, Harmonic mean labeling of disconnected graphs, *J. Discrete Math. Sci. Cryptogr.,* **19**(1) (2016) 1-12.

[1593] C. Jayasekaran and J. L. Flower, On edge trimagic labeling of umbrella, dumb bell and circular ladder graphs, *Annals Pure Appl. Math.*, **13**(1) (2017) 73-87. https://www.researchgate.net/publication/320798252_On_Edge_Trimagic_Labeling_of_Umbrella_Dumb_Bell_and_Circular_Ladder_Graphs

[1594] C. Jayasekaran and J. L. Flower, Edge trimagic total labeling of Möbius ladder, book and dragon graphs, *Annals Pure Appl. Math.*, **13**(2) (2017) 151-163. http://www.researchmathsci.org/apamart/apam-v13n2-1.pdf

[1595] C. Jayasekaran and J. L. Flower, Edge trimagic total labelings of some shadow and splitting graphs, *Internat. J. Pure Appl. Math.*, **120**(3) (2018) 303-313. https://www.ijpam.eu/contents/2018-120-3/2/2.pdf

[1596] C. Jayasekaran and M. Jaslin Melbha, Some more results on one modulo three root square mean graphs, *Internat. Elect. J. Pure Appl. Math.,* **13**(1) (2019) 47-55. https://www.e.ijpam.eu/contents/articles/201901301004.pdf

[1597] K. T. JebiNivathitha, N. Srinivasan, A. Parthiban, and Sangeeta, On D3EL of complete and star graphs, *J. Xidian Univer.*, **14**(5) (2020) 4448-4454.

[1598] K. T. JebiNivathitha, N. Srinivasan, A. Parthiban, and Sangeeta, On D3EL of wheel graphs, *J. Gujarat Res. Soc.,* **8**(5) (2020) 5480-5483.

[1599] J. Jeba Jesintha, New Classes of Graceful Trees, Ph. D. Thesis, Anna University, Chennai, India, 2005.

[1600] J. Jeba Jesintha and D. W. Devakirubanithi, Cordial labeling on different types of nested triangular graphs, *Bull. Pure Appl. Sci., Sect. E Math. Stat.*, **40E**(2) (2021) 135-139. https://sdbindex.com/documents/00000419/00001-32780.pdf

[1601] J. Jeba Jesinthan, D. Devakirubanithi, and B. Aathry, Odd graceful labelings of SSG with star related graphs, *Adv. Appl. Math. Sci.,* **212** (2021) 519-529. https://www.mililink.com/upload/article/10125aams_vol_212_december_2021_a1_p519-529_j._jeba_jesintha_and_b._arthy.pdf

[1602] J. Jeba Jesintha, D. Devakirubanithi, and M. Angel, Odd graceful labeling of stalked subdivided shell graph, In book: Multidisciplinary Research in Academics: The Way Forward, Publisher: Emerald Publishers, 2023 https://www.researchgate.net/publication/398106138_Odd_Graceful_Labeling_of_Stalked_Subdivided_Shell_Graph

[1603] J. Jeba Jesintha, D. Devakirubanithi, and R. Charishma, New types of graphs on cordial labeling, *Bull. Pure Appl. Sci.,* Sect. E Math. Stat., **42E**(1) (2023) 36-42. https://www.bpasjournals.com/math-and-stat/index.php/journal/article/view/28?utm_source=chatgpt.com

[1604] J. Jeba Jesintha, D. Devakirubanithi, and E. A. Jasma Pershi, New classes of difference cordial graphs, *J. Information and Computational Science*, *10*(7) (2020) 636-643. https://www.researchgate.net/publication/374234073_New_Classes_of_Difference_Cordial_Graphs

[1605] J. Jeba Jesintha, D. Devakirubanithi, and L. Monisha, Edge product cordial labeling on duplication of prism graph, *Bull. Pure Appl. Sci.,* Sect. E Math. Stat., **39E**(2) (2020) 265–270. https://api.semanticscholar.org/CorpusID:239722931

[1606] J. Jeba Jesintha, D. Devasagayam, and J. M. Sherlin, Odd graceful labeling of super subdivision of few graphs, *Bull. Pure Appl. Sci.- Math. Stat.,* **41e**(2) (2022) 167-171. https://www.researchgate.net/publication/374234245_ODD_GRACEFUL_LABELING_OF_SUBDIVISION_AND_SUPER_SUBDIVISIOIN_OF_FEW_TYPES_OF_GRAPHS

[1607] J. Jeba Jesintha and K. Ezhilarasi Hilda, Butterfly graphs with shell orders m and $2m + 1$ are graceful, *Bonfring Internat. J. Research Communication Engin.,* **2**(2) (2012) 1-5. https://www.researchgate.net/publication/271305096_Butterfly_Graphs_with_Shell_Orders_m_and_2m1_are_Graceful

[1608] J. Jeba Jesintha and K. E. Hilda, Graceful labeling of bow graphs and shell-flower graphs, *Internat. J. Comput. Alg.,* **2** (2013) 214-218. DOI:10.20894/IJCOA.101.002.001.017

[1609] J. Jeba Jesintha and K. E. Hilda, Shell-butterfly graphs are harmonious, *Adv. Appl. Disc. Math.*, **14**(1) (2014) 39-51. https://www.researchgate.net/publication/346528833_Advances_and_Applications_in_Discrete_Mathematics_SHELL-BUTTERFLY_GRAPHS_ARE_HARMONIOUS

[1610] J. Jeba Jesintha and K. E. Hilda, Variation of graceful labeling on disjoint union of two subdivided shell graphs, *Annals Pure Applied Math.,* **8**(2) (2014) 19-25. http://www.researchmathsci.org/apamart/apam-v8n2-3.pdf

[1611] J. Jeba Jesintha and K. E. Hilda, Subdivided uniform shell bow graphs are one modulo three graceful, *Math. Sci. Internat. Res. J.,* *3*(2) (2014) 645-646. https://www.researchgate.net/publication/346528719_subdivided_uniform_shell_bow_graphs_are_odd_graceful

[1612] J. Jeba Jesintha and K. E. Hilda, Subdivided uniform shell bow graphs and subdivided shell graphs are odd graceful, *Eduventure Res. J. Queen Mary's College,* **7**(2) (2014) 103-105.

[1613] J. Jeba Jesintha and K. Ezhilarasi Hilda, All uniform bow graphs are graceful, *Math. Comput. Sci.,* **9** (2015)(2) 185-191. https://www.researchgate.net/publication/346528719_subdivided_uniform_shell_bow_graphs_are_odd_graceful

[1614] J. Jaba Jesintha and K. E. Hilda, Double shells with two pendant edges at the apex are k-graceful, *J. Combin. Math. Combin. Comput.,* **92** (2015) 47-57. https://combinatorialpress.com/article/jcmcc/Volume%20092/vol-092-paper%205.pdf

[1615] J. Jeba Jesintha and K. E. Hilda, Shell butterfly graphs are graceful, *Internat. J. Pure Appl. Math.,* **101**(6) (2015) 949-956. https://www.researchgate.net/publication/346528963_P_A_SHELL-BUTTERFLY_GRAPHS_ARE_GRACEFUL

[1616] J. Jeba Jesintha and K. E. Hilda, $\rho^{\star}$ labeling of paths and shell butterfly graphs, *Internat. J. Pure and Appl. Math.,* **101**(5) (2015) 645-653. [J. Jeba Jesintha and K. E. Hilda, $\rho^{\star}$ labeling of paths and shell butterfly graphs, *Internat. J. Pure and Appl. Math.,* **101**(5) (2015) 645-653.

[1617] J. Jeba Jesintha and K. E. Hilda, Shell butterfly graphs are edge odd graceful, *Internat. J. Pure and Appl. Math.,* **109**(7) (2016) 159-166. https://www.inderscienceonline.com/doi/abs/10.1504/IJCAET.2021.111639

[1618] J. Jeba Jesintha and K. E. Hilda, The graph SSG(2) is odd graceful and odd harmonious, *Internat. J. Comp. Aided Engin. and Tech.*, **14**(1) (2019) 93-101. https://www.academia.edu/88330489/The_graph_SSG_2_is_odd_graceful_and_odd_harmonious

[1619] J. Jeba . Jesintha and K. E. Hilda, Subdivided shell flower graphs are odd graceful, *Internat. J. Innovation Sci. Math.*, **2**(4)(2014) 402-404. https://www.researchgate.net/publication/346647817_Subdivided_Shell_Flower_Graphs_are_Odd_Graceful

[1620] J. Jeba. Jesintha and K. E. Hilda, Disjoint union of two SSG (2) is odd graceful, *Internat. J. Math. Comput. Sci,*, **15**(4) (2020) 1143-1148. https://future-in-tech.net/15.4/R-Hilda.pdf

[1621] J. Jeba Jesintha, R. Jaya Glory, and A. Elakiya Solai, Odd graceful labeling of path union of caterpillars, *Advan. Appl. Disc. Math.,* **28**(2) (2021) 287-294. NOW Chat 9/30/26

[1622] J. Jeba Jesintha and G. Sethuraman, All arbitrary fixed generalized banana trees are graceful, *Math. Comput. Sci.*, **5**(1) (2011) 51-62.

[1623] J. Jeba Jesintha and K. Subashini, On gracefulness of vertex switching of even cycles, *International J. Pure Appl. Math.,*(IJPAM), **109**(10) (2016) 11-18. http://acadpubl.eu/jsi/2016-109-si/10/2.pdf

[1624] J. Jeba Jesintha and K. Subashini, On cordial labeling of vertex switching of cycles, *Middle East J. Sci. Res.,*(MEJSR), **25**(04) (2017) 791-794. http://www.idosi.org/mejsr/mejsr25(4)17/13.pdf

[1625] J. Jeba Jesintha and K. Subashini, Gracefulness in the path union of vertex switching of odd cycle graphs, *Math. Sci. Internat. Res. J*(IMRFJ), **07**(Special Issue 2) (2018) 71-75.

[1626] J. Jeba Jesintha and K. Subashini, Cordial labeling for new class of graphs, *IAETSD J. for Adv. Res. in Appl. Sci.,*(IAETSDJARAS), **05**(04) (2018), 98-101. https://app.box.com/s/9wi68sa59stvz7yxby77g4ywe9olbkg1

[1627] J. Jeba Jesintha and K. Subashini, Gracefulness in the path union of vertex switching of even cycles in increasing order, *South East Asian J. Math. and Mathl. Sci.,* (SEAJMAMS), **14**(3) (2018)15-20. http://rsmams.org/journals/articleinfo.php?articleid=317&tag=seajmams

[1628] J. Jeba Jesintha and K. Subashini, Cordiality in the path union of vertex switching of cycles in increasing order, *South East Asian J. Math.Mathl. Sci.,* (SEAJMAMS), **14**(3) (2018) 111-118. http://rsmams.org/journals/articleinfo.php?articleid=327&tag=seajmams

[1629] J. Jeba Jesintha and K. Subashini, Gracefulness of the join of cycle with different graphs, *Internat. J of Pure and Appl. Math.,* (IJPAM), **120** (Special Issue)(08) (2018) 49-57. https://acadpubl.eu/hub/2018-120-8/2/6.pdf

[1630] J. Jeba Jesintha and K. Subashini, Gracefulness in the path union of vertex switching of cycles, *Internat. J Res. Eng., IT Social Sci.,* (IJREISS), **09**(Spl. issue) (2019) 06-10. http://indusedu.org/pdfs/IJREISS/IJREISS_3388_99004.pdf

[1631] J. Jeba Jesintha and K. Subashini, On gracefulness of quadrilateral snakes *Internat. J. Res. Eng. IT Social Sci.,* (IJREISS), **09**(Spl. issue) (2019) 15-23. http://indusedu.org/pdfs/IJREISS/IJREISS_3390_59454.pdf

[1632] J. Jeba Jesintha and K. Subashini, On gracefulness in the path union and cycle of $P_m\theta S_n$, *Bull. Pure Appl. Sci. Sec. E. Math. Stat.,*(BPAS) **38E**(Special Issue)(2S) (2019) 85-91. https://www.bpasjournals.com/admin/upload/dynamic2/85-91-bpas-e-dec.2019-special%20issue.pdf

[1633] J. Jeba Jesintha and K. Subashini, Cordiality in the path union of vertex switching of wheel graphs in increasing order,*Bull. Pure Appl. Sci. Sec. E. Math. Stat.,* (BPAS) **38E**(Special Issue)(2S) (2019) 79-84. https://www.bpasjournals.com/admin/upload/dynamic2/79-84-bpas-e-dec.2019-special%20issue.pdf

[1634] J. Jeba Jesintha and K. Subashini, Cordiality in the path union and cycle of jelly fish graphs *J. Inf. Comp. Sci.,*(JOICS) **10**(7) (2020) 740-749. http://www.joics.org/gallery/ics-3920.pdf

[1635] J. Jeba Jesintha, K. Subashini, and Allu Merin Sabu, Graceful labeling on a new family of graphs, *Bull. Pure Appl. Sci. Sec. E. Math. Stat.,*(BPAS) **38E**(Special Issue)(2S) (2019) 1-6. https://bpasjournals.com/admin/upload/dynamic2/1-6-bpas-e-dec2019-special-issue.pdf

[1636] J. Jeba Jesintha, K. Subashini, and Allu Merin Sabu, Graceful labeling on twig diamond graph with pendant edges, *Bull. Pure Appl. Sci. Sec. E. Math. Stat.,* (BPAS) **39E**(2) (2020) 188-192. https://bpasjournals.com/admin/upload/dynamic2/188-192-bpas-e-dec.2020.pdf

[1637] J. Jeba Jesintha, K. Subashini, and H. Aysha Siddiqa, Gracefulness in the path union of vertex switching of odd and even cycle graphs *J. Maharaja Sayajirao Univ. Baroda,* **54**(2)(XI) (2020) 140-144.

[1638] J. Jeba Jesintha, K. Subashini, and J. R. Rashmi Beula, Graceful labeling of some new graphs, *Bull. Pure Appl. Sci. Sec. E. Math. Stat.,*(BPAS) **38E**(Special Issue)(2S) (2019) 60-64. https://bpasjournals.com/admin/upload/dynamic2/60-64-bpas-e-dec.2019-special%20issue.pdf

[1639] J. Jeba Jesintha and G. Sethuraman, A new class of graceful rooted trees, A new class of graceful rooted trees, *J. Discrete Math. Sci. Cryptogr.,* **11** (2008) 421-435. https://connectjournals.com/file_full_text/84904H_jdmsc235_421-435.pdf

[1640] J. Jeba Jesintha and G. Sethuraman, Generation of graceful trees, *Internat. J. Algor. Comput. Math.,* **2** (2009) 33-38.

[1641] J. Jeba Jesintha and G. Sethuraman, All arbitrarily fixed generalized banana trees are graceful, *Math. Comput. Sci.,* **5**(1) (2011) 51-62.

[1642] J. Jeba Jesintha and K. Subashini, Gracefulness in path union of vertex switching of even cycles in increasing order, *South East Asian J. Math. Math. Sci.,* **14**(3) (2018) 15-20. https://indusedu.org/pdfs/IJREISS/IJREISS_3388_99004.pdf

[1643] J. Jeba Jesintha, N. K. Vinodhini, and S. Divya Lakshmi, Antimagic labeling of triangular book graph and double fan graph, *Bull. Pure Appl. Sci., Sect. E Math. Stat.*, **42E**(1) (2023) 27-30. https://www.bpasjournals.com/math-and-stat/index.php/journal/article/view/26/10

[1644] J. Jeba Jesintha, and C. Yogalakshmi, Edge-odd graceful labeling of Jahangir graph, *South East Asian J. Math. Math. Sci.,* **17**(3) (2021) 347-354. https://rsmams.org/journals/articleinfo.php?articleid=644&tag=seajmams

[1645] S. Jendroľ, J. Misšuf, and R. Soták, Total edge irregularity strength of complete graphs and complete bipartite graphs, *Elec. Notes Discrete Math.,* **28** (2007) 281-285.

[1646] S. Jendroľ, J. Misšuf, and R. Soták, Total edge irregularity strength of complete graphs and complete bipartite graphs, *Discrete Math.,* **310** (2010) 400-407. https://www.sciencedirect.com/science/article/pii/S157106530700042X

[1647] S. Jenifer Wency, A. Lourdusamy, and F. Patrick, Several result on group S_3 cordial remainder labeling, *AIP Conference Proceed.*, 2261, 030035 (2020).

[1648] R. H. Jeurissen, Magic graphs, a characterization, *Europ. J. Combin.,* **9** (1988) 363-368. https://www.sciencedirect.com/science/article/pii/S0195669888800660?via%3Dihub

[1649] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi, and M. Z. Youssef, k-Product cordial behaviour of union of graphs, *J. Indones. Math. Soc.,* **28**(1) (2022) 1-7. https://www.researchgate.net/publication/359210439_k-Product_Cordial_Behaviour_of_Union_of_Graphs

[1650] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi and M. Z. Youssef, k-Product cordial labeling of Napier bridge graphs, *Nepal J. Math. Sci.*, **3**(2) (2022) 59-70. doi:10.3126/njmathsci.v3i2.49201

[1651] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi, and M. Z. Youssef, k-Product cordial labeling of cone graphs, *Internat. J. Math. Combin.*, **2** (2022) 72-80. http://fs.unm.edu/IJMC/k-Product_Cordial_Labeling_of_Cone_Graphs.pdf

[1652] K. Jeya Daisy, R. S. Sabibha, P. Jeyanthi, and M. Z. Youssef, k-product cordial labeling of fan graphs, *TWMS J. App. Eng. Math.*, **13**(1) (2023) 11-20. https://www.researchgate.net/publication/366822667_K-PRODUCT_CORDIAL_LABELING_OF_FAN_GRAPHS

[1653] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi, and M. Z. Youssef, k-Product cordial labeling of cone graphs, *Internat. J. Math. Combin.,* **2** (2022) 72-80 https://fs.unm.edu/IJMC/k-Product_Cordial_Labeling_of_Cone_Graphs.pdf

[1654] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi, and M. Z. Youssef, k-Product cordial labeling of product of graphs, *Discrete Math., Algorithms and Appl.*, **16**(1) (2024) 2250187 (8 pages).

[1655] K. Jeya Daisy, R. S. Sabibhab, P. Jeyanthi, and M. Z. Youssef, k-Product cordial labeling of splitting graph of star graphs, *Commun. Combin., Cryptogr. & Computer Sci.*, **1** (2024) 68-80. https://www.researchgate.net/publication/376358019_k-Product_Cordial_Labeling_of_Splitting_Graph_of_Star_Graphs

[1656] K. Jeya Daisy, R. Santrin Sabibha, P. Jeyanthi and M. Z. Youssef, Further results on k-product cordial labeling, *TWMS J. Appl. Engin. Math.*, **14**(3) (2024) 981-990. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.3/08.pdf

[1657] P. Jeyanthi, k-equitable mean labeling, *J. Algor. Computation*, **44**(1) (2013) 21-30. https://doi.org/10.22059/jac.2013.7913

[1658] P. Jeyanthi, M. Selvi and D. Ramya, Skolem difference mean labeling of union of graphs, *Util. Math.,* **111** (2019) 139-150.

[1659] P. Jeyanthi, N. Angel Benseera and M. Immaculate Mary, On totally magic cordial labeling, *SUT J. Math.*, **49**(1) (2013) 13-18. https://www.researchgate.net/publication/266540964_On_totally_magic_cordial_labeling

[1660] P. Jeyanthi and N. Angel Benseera, Totally vertex-magic cordial labeling, *SUT J. Math.*, **49**(1) (2013) 1-12. https://www.researchgate.net/publication/257422541_PJeyanthi_and_NAngel_Benseera_Totally_vertex-magic_cordial_labeling_SUT_Journal_of_Mathematics_Vol_49_No1_2013_1-12

[1661] P. Jeyanthi and N. Angel Benseera, Totally magic cordial labeling of one-point union of n copies of a graph, *Opuscula Math.*, **34**(1) (2014) 115-133. https://pdfs.semanticscholar.org/7b6f/e45aa7f4dffcccbd6276a15d67cef9a7387c.pdf

[1662] P. Jeyanthi and N. Angel Benseera, Totally magic cordial labeling of some graphs, *J. Algorithms Comput.*, **46** (2015) 1-8. https://jac.ut.ac.ir/article_7921_5e2b6a274667fa1b3976387dd2ecb005.pdf

[1663] P. Jeyanthi, N. Angel Benseera, and I. Cahit, Totally magic cordial labeling of mP_n and mK_n, *Proyecciones J. Math.*, **35**(4) (2016) 371-381. https://www.scielo.cl/pdf/proy/v35n4/art01.pdf

[1664] P. Jeyanthi, N. Angel Benseera and G.-C, Lau, On k-totally magic cordial labeling of graphs, *Disc. Math., Algor. Appl.*, **7**(3)(2015) 1550024, 7 pp. https://www.researchgate.net/publication/277938227_On_k-totally_magic_cordial_labeling_of_graphs

[1665] P. Jeyanthi and N. Angel Benseera, Totally magic cordial deficiency of some graphs, *Utilitas Math.,* **105** (2017) 87-94.

[1666] P. Jeyanthi, N. Angel Benseera, and G.-C. Lau, Hypo-k-totally magic cordial labeling of graphs, *Proyecciones J. Math.*, **34**(4) (2015) 347-355. https://www.scielo.cl/pdf/proy/v34n4/art04.pdf

[1667] P. Jeyanthi and R. Gomathi, Odd mean labeling of chain of graphs, *Electron. Notes Disc. Math.*, **48** (2015) 157-160. http://dx.doi.org/10.1016/j.endm.2015.05.022

[1668] P. Jeyanthi and R. Gomathi, Analytic odd mean labeling of super subdivision and H-super subdivision of graphs, *J. Emerging Tech. Innov. Res. (JETIR)*, **6**(6) (2019) 541-551. http://www.jetir.org/papers/JETIR1908A25.pdf

[1669] P. Jeyanthi and R. Gomathi, Analytic odd mean labeling of corona graphs, *Malaya J. Mat.,* **8**(3)(2020) 898-902. https://www.researchgate.net/publication/342493786_Analytic_odd_mean_labeling_of_Corona_graphs

[1670] P. Jeyanthi, R. Gomathi, and G.-C. Lau, Analytic odd mean labelings of some graphs, *Palestine J. Math.,* **8**(2) (2019) 392-399. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_392to399.pdf

[1671] P. Jeyanthi, R. Gomathi, and G.-C. Lau, Some results on analytic odd mean labeling of graph, *Bull. Internat. Math. Virtual Inst.*, **9** (2019) 487-499. http://www.imvibl.org/buletin/bulletin_imvi_9_3_2019/bulletin_imvi_9_3_2019_493_500.pdf

[1672] P. Jeyanthi, R. Gomathi, and G.-C. Lau, Further results on analytic odd mean labeling of graphs, *Internat. J. Math. Combin.,* **4** (2019) 136-148. https://www.researchgate.net/publication/338448124_Further_Results_on_Analytic_Odd_Mean_Labeling_of_Graphs

[1673] P. Jeyanthi, R. Gomathi, G. C. Lau, and W. C. Shiu, Analytic odd mean labeling of union and identification of some graphs, *Proyecciones J. Mathematics*, **42**(5) (2023) 1141-1155. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_392to399.pdf

[1674] P. Jeyanthi and K. Jeya Daisy, Z_k-magic labeling of subdivision graphs, *Discrete Math., Algor. Appl.,* **8**(3) (2016) 1650046 (19 pages).

[1675] P. Jeyanthi and K. Jeya Daisy, Certain classes of magic graphs, *J. Graph Labeling*, **4**(1) (2018) 38-47. https://www.researchgate.net/publication/341313267_Certainclassesofmagicgraphs

[1676] P. Jeyanthi and K. Jeya Daisy, Z_k-magic labeling of some families of graphs, *J. Algor. Computation,* **50**(2) (2018) 1-12. https://www.researchgate.net/publication/341313196_Z_k_-Magic_Labeling_of_Some_Families_of_Graphs

[1677] P. Jeyanthi and K. Jeya Daisy, Z_k-Magic labeling of star of graphs, *Proyecciones J. Math.*, **39**(1) (2020) 31-51. https://www.researchgate.net/publication/341313315_Zk_-_Magic_Labeling_of_Star_of_Graphs

[1678] P. Jeyanthi and K. Jeya Daisy, Z_k-Magic labeling of open star of graphs, *Bull. Int. Math. Virtual Inst.,* **7**(2) (2017) 243-255. https://www.researchgate.net/publication/341313315_Zk_-_Magic_Labeling_of_Star_of_Graphs

[1679] P. Jeyanthi, K. Jeya Daisy, and A. Semaničová-Feňovčíková, Z_k-Magic labeling of path union of graphs, *CUBO J. Math.*, **21**(2) (2019) 15-35. https://www.researchgate.net/publication/341312908_Z_k_-Magic_Labeling_of_Path_Union_of_Graphs

[1680] P. Jeyanthi and K. Jeya Daisy, Some results on Z_k-magic labeling, *Palestine J. Math.,* **8**(2) (2019) 400-412. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_400to412.pdf

[1681] P. Jeyanthi and K. Jeya Daisy, Z_k-magic labeling of cycle of graphs, *Internat. J. Math. Combin.,* **1** (2019) 88-102. https://fs.unm.edu/IJMC/ZkMagicLabeling.pdf

[1682] P. Jeyanthi, R. Gomathi and G.-C. Lau, Analytic odd mean labeling of some standard graphs, *Palestine J. Math.,* **8**(1) (2019) 444-450. https://pjm.ppu.edu/sites/default/files/papers/PJM_Dec_2019_444to450.pdf

[1683] P. Jeyanthi, R. Gomathi, and G.-C. Lau, Analytic odd mean labeling of square and H graphs, *Internat. J. Math. Combin.,* Special Issue 1 (2018) 61-67. https://www.researchgate.net/profile/P-Jeyanthi/publication/330239773_Analytical_odd_mean_labeling_of_Square_and_H_graphs/links/5c35539c92851c22a3660bd8/Analytical-odd-mean-labeling-of-Square-and-H-graphs.pdf

[1684] P. Jeyanthi, R. Kalaiyarasi and D. Ramya, Centered triangular mean graphs, *Inter. J. Math. Comb.,* **1** (2019) 126-133. https://www.researchgate.net/publication/332230070_Centered_Triangular_Mean_Graphs

[1685] P. Jeyanthi, R. Kalaiyarasi, D. Ramya, and T. Saratha Devi, Some results on Skolem odd difference mean labeling, *Proyecciones J. Math.,* **35**(4) (2016) 405-415. https://www.scielo.cl/pdf/proy/v35n4/art04.pdf

[1686] P. Jeyanthi and A. Maheswari, Vertex equitable labeling of cycle and path related graphs, *Util. Math.,* **98** (2015) 215-226.

[1687] P. Jeyanthi and A. Maheswari, Some results on vertex equitable labeling, *Open J. Discrete Math.,* **2** (2012) 51-57. http://dx.doi.org/10.4236/ojdm.2012.22009 NEW Nov. 2016

[1688] P. Jeyanthi and A. Maheswari, 3-product cordial labeling, *SUT J. Math.,* **48** (2) (2012) 231-240. https://www.researchgate.net/publication/267086781_3-product_cordial_labeling

[1689] P. Jeyanthi and A. Maheswari, 3-product cordial labeling of some graphs, *Internat. J. Math. Combin.,* **1** (2012) 96-105. https://www.researchgate.net/publication/257422341_PJeyanthi_and_AMaheswari_3-product_cordial_labeling_of_some_graphs_International_Journal_of_Mathematical_Combinatorics_Vol12012_96-105

[1690] P. Jeyanthi and A. Maheswari, Vertex equitable labeling of transformed trees, *J. Algor. Computation*, **44**(1) (2013) 9-20. https://jac.ut.ac.ir/article_7912_b7a385d110f5ec4c2d805c82bcf3079e.pdf

[1691] P. Jeyanthi and A. Maheswari, Some results on 3-product cordial labeling, *Util. Math.,* **99** (2016) 215-229. https://www.researchgate.net/publication/301543696_Some_results_on_3-product_cordial_labeling

[1692] P. Jeyanthi and A. Maheswari, One modulo three mean labeling of graphs, *Amer. J. Appl. Math. Stat.*, **2**(5) (2014) 302-306. https://pubs.sciepub.com/ajams/2/5/2/ajams-2-5-2.pdf

[1693] P. Jeyanthi and A. Maheswari, Vertex equitable labeling of graphs, *Graph Theory Notes New York*, **LXVII** (2014) 34-42. https://www.researchgate.net/publication/265547093_Vertex_equitable_labeling_of_graphs

[1694] P. Jeyanthi and A. Maheswari, Further results on vertex equitable graphs, *Electron. Notes Disc. Math.*, **48** (2015) 161-164. http://dx.doi.org/10.1016/j.endm.2015.05.023

[1695] P. Jeyanthi and A. Maheswari, 3-product cordial labeling of star graphs, *Southeast Asian Bull. Math.*, **39**(3) (2015) 429-437. https://www.researchgate.net/publication/281505995_3-Product_Cordial_Labeling_of_Star_Graphs

[1696] P. Jeyanthi and A. Maheswari, Vertex equitable labeling of cyclic snakes and bistar graphs, *J. Scientific Res.,* **6**(1) (2014) 79-85. https://www.researchgate.net/publication/259562636_Vertex_Equitable_Labeling_of_Cyclic_Snakes_and_Bistar_Graphs

[1697] P. Jeyanthi and A. Maheswari, Odd vertex equitable even labeling of cycle related graphs, *CUBO A Math. J.,* **20** (02) (2018) 13-21. doi:10.4067/S0719-06462018000200013

[1698] P. Jeyanthi, A. Maheswari, and P. Pandiaraj, One modulo three mean labeling of cycle related graphs, *Internat. J. Pure and Appl. Math.,* **103**(4)(2015) 625-633. https://www.researchgate.net/publication/281853316_One_Modulo_Three_Mean_Labeling_of_Cycle_Related_Graphs

[1699] P. Jeyanthi, A. Maheswari, and P. Pandiaraj, On one modulo three mean labeling of graphs, *J. Discrete Math. Science & Cryptography*, **19:2** (2016) 375-384. https://www.researchgate.net/publication/303970865_P_Jeyanthi_A_Maheswari_P_Pandiaraj_2016_On_one_modulo_three_mean_labeling_of_graphs_Journal_of_Discrete_Mathematical_Sciences_and_Cryptography_192_375-384_DOI_10108009720529201511028 38

[1700] P. Jeyanthi, A. Maheswari, and P. Pandiaraj, One modulo three geometric mean graphs, *J. Algor. Comput.,* **50**(1) (2018) 101-108. https://jac.ut.ac.ir/article_68342_e53cdfd892a4cf9b99d51c533e8902a6.pdf

[1701] P. Jeyanthi, A. Maheswari, and P. Pandiaraj, One modulo three mean labeling of transformed trees, *Proyecciones J. Math.*, **35**(3) (2016) 277-289. https://www.scielo.cl/pdf/proy/v35n3/art05.pdf

[1702] P. Jeyanthi, A. Maheswari and M. VijayaLaksmi, Odd vertex equitable even labeling of graphs, *Proyecciones J. Math.,* **36**(1) (2017) 1-11. https://scielo.conicyt.cl/pdf/proy/v36n1/art01.pdf

[1703] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, New results on vertex equitable labeling, *J. Algebra Combin. Discrete Structures Appl.,* **3**(2)(2016) 97-104. https://dergipark.org.tr/en/download/article-file/267792

[1704] P. Jeyanthi, A. Maheswari and M. VijayaLaksmi, Vertex equitable labeling of union of cyclic snake related graphs, *Proyecciones J. Mathematics*, **35**(1) (2016) 177-186. https://scielo.conicyt.cl/pdf/proy/v35n2/art03.pdf

[1705] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, Vertex equitable labeling of double alternate snake graph, *J. Algor. Comput.*, **46** (2015) 27-34. https://jac.ut.ac.ir/article_7923_76d1a6298f68c081c967627653edc287.pdf

[1706] P. Jeyanthi, A. Maheswari and M. VijayaLaksmi, Vertex equitable labeling of cycle and star related graphs, *J. Scientific Res.*, **7**(3)(2015) 33-42. https://www.banglajol.info/index.php/JSR/article/view/22810

[1707] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, Vertex equitable labelings of super subdivision graphs, *Scientific Internat.,* **27**(5) (2015) 3881-3883. https://www.researchgate.net/profile/P_Jeyanthi2/publication/284182322_VERTEX_EQUITABLE_LABELING_OF_SUPER_SUBDIVISION_GRAPHS/links/564ee37a08aefe619b103a26.pdf

[1708] P. Jeyanthi, A. Maheswari, and M. VijayaLakshmi, 3-Product cordial labeling of some snake graphs, *Proyecciones J. Math.* **38**(1) (2019) 13-30. https://www.scielo.cl/pdf/proy/v38n1/0716-0917-proy-38-01-00013.pdf

[1709] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, Further results on 3-product cordial labeling, *Proyecciones J. Math.,* **38**(2) (2019) 191-202. https://www.scielo.cl/pdf/proy/v38n2/0717-6279-proy-38-02-00191.pdf

[1710] P. Jeyanthi, A. Maheswari, and M. VijayaLakshmi, Vertex switching in 3-product cordial graphs, *J. Algor. Computation*, **50**(1) (2018) 185-193. https://jac.ut.ac.ir/article_68965_547e5985e086535b9ff0e6861eab88e3.pdf

[1711] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, Vertex equitable labeling of cyclic snake related graphs, *Proyecciones J. Math.*, **37**(4) (2018) 613-625.

[1712] P. Jeyanthi, A. Maheswari, and M. VijayaLaksmi, Odd vertex equitable even labeling of ladder graphs, *Jordon J. Math. Stat.*, **12**(1) (2019) 75-87. https://www.researchgate.net/publication/332062041_Odd_vertex_equitable_even_labeling_of_ladder_graphs

[1713] P. Jeyanthi, A. Maheswari, and M. Vijayalakshmi, Odd vertex equitable even labeling of ladder graphs, *Jordan J. Math. Stat.,* **12**(1) (2019) 75-87. https://www.researchgate.net/publication/332062041_Odd_vertex_equitable_even_labeling_of_ladder_graphs

[1714] P. Jeyanthi and N. T. Muthuraja, Some cycle-super magic graphs, *Internat. J. Math. Soft Comput.,* **4**(2) (2014) 137-144.

[1715] P. Jeyanthi, N. T. Muthuraja, A. Semaničová-Feňovčíková and S. J. Dharshikha, More classes of cycle super antimagic graphs, *Australasian J. Combin.*, **67**(1) (2017) 46-64. https://ajc.maths.uq.edu.au/pdf/67/ajc_v67_p046.pdf

[1716] P. Jeyanthi, S. Philo, and Kiki A.Sugeng, Odd harmonious labeling of some new families of graphs, *SUT J. Math.,* **51**(2) (2015) 181-193. https://www.rs.tus.ac.jp/sutjmath/_userdata/51-2/03-jeyanthi-philo-sugeng.pdf

[1717] P. Jeyanthi and S. Philo, Odd harmonious labeling of some cycle related graphs, *Proyecciones J. Math.*, **35**(1) (2016) 85-98. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/1236/949

[1718] P. Jeyanthi and S. Philo, Odd harmonious labeling of plus graphs, *Bull. Int. Math. Virtual Inst.*, **7** (2017)(3) 515-526. https://www.researchgate.net/publication/316472719_ODD_HARMONIOUS_LABELING_OF_PLUS_GRAPHS

[1719] P. Jeyanthi and S. Philo, Odd harmonious labeling of some new graphs, *Southeast Asian Bull. Math.,* **43** (2019) 509-523. https://www.researchgate.net/publication/329626405_Cordial_Labeling_of_Generalized_Jahangir_Graph

[1720] P. Jeyanthi and S. Philo, Odd harmonious labeling of some super subdivision graphs, *Proyecciones J. Math.,* **38**(1) (2019) 1-11. https://scielo.conicyt.cl/pdf/proy/v38n1/0716-0917-proy-38-01-00001.pdf

[1721] P. Jeyanthi and S. Philo, Odd harmonious labeling of complete bipartite graphs, *Southeast Asian Bull. Math.,* **46** (2022) 23-34. https://www.researchgate.net/publication/359462364_Southeast_Asian_Bulletin_of_Mathematics_Odd_Harmonious_Labeling_of_Complete_Bipartite_Graphs

[1722] P. Jeyanthi and S. Philo, New results on odd harmonious labeling of graphs, *TWMS J. App. Eng. Math.,* **12**(4) (2022) 1301-1310. https://jaem.isikun.edu.tr/web/images/articles/vol.12.no.4/14.pdf

[1723] P. Jeyanthi and S. Philo, Odd harmonious labeling of line and disjoint union of graphs, *Chinese J. Math. Sci.,* (1)(1) (2021) 61-68 https://romanpub.com/resources/cjms%20v1-1-paper6tex.pdf

[1724] P. Jeyanthi and S. Philo, Odd harmonious labeling of subdivided shell graphs, *Internat. J. Comput. Sci. Eng.,* **7**(5) (2019) 77-80. https://www.researchgate.net/publication/332564163_Odd_Harmonious_Labeling_of_Subdivided_Shell_Graphs

[1725] P. Jeyanthi and S. Philo, Odd harmonious labeling of step ladder graphs, *Util. Math.,* **115** (2020) 89-103.

[1726] P. Jeyanthi and S. Philo, Odd harmonious labelings of certain graphs, *JASC: J. Appl. Sci. Comput.,* **VI**(IV) (2019) 1224-1232. https://www.researchgate.net/profile/P_Jeyanthi2/publication/332320709_ODD_HARMONIOUS_LABELING_OF_CERTAIN_GRAPHS/links/5cadc56492851ccd4ac087b5/ODD-HARMONIOUS-LABELING-OF-CERTAIN-GRAPHS.pdf

[1727] P. Jeyanthi and S. Philo, Some results on odd harmonious labeling of graphs, *Bull. Internat. Math. Virtual Inst.,* **9** (2019) 567-576. https://www.researchgate.net/publication/332902831_SOME_RESULTS_ON_ODD_HARMONIOUS_LABELING_OF_GRAPHS

[1728] P. Jeyanthi and S. Philo, Odd harmonious labeling of some classes of graph, *CUBO, A Math. J.,* **22**(3) (2020) 299-314. https://www.researchgate.net/publication/347489135_Odd_Harmonious_Labeling_of_Some_Classes_of_Graphs

[1729] P. Jeyanthi, S. Philo and M. Z. Youssef, Odd harmonious labeling of grid graphs, *Proyec. J. Math.,* **38**(3) (2019) 411-428. https://scielo.conicyt.cl/pdf/proy/v38n3/0716-0917-proy-38-03-00411.pdf

[1730] P. Jeyanthi and D. Ramya, Some results on super mean graphs, *Util. Math.,* **92** (2013) 149-160.

[1731] P. Jeyanthi and D. Ramya, On construction of mean graphs, *J. Scientific Research* **5**(2) (2013) 265-273. https://www.researchgate.net/publication/257364403_PJeyanthiand_DRamya_On_construction_of_mean_graphs_Journal_of_Scientific_Research_Vol5_2_2013_265-273

[1732] P. Jeyanthi and D. Ramya, Super mean graphs, *Util. Math.,* **96** (2015) 101-109.

[1733] P. Jeyanthi, D. Ramya, and A. Maheswari, On mean labeling of some trees, *Util. Math.*, **93** (2014) 161-191.

[1734] P. Jeyanthi and D. Ramya Super mean labeling of some classes of graphs, *International J. Math. Combin.,* **1** (2012) 83-91. https://fs.unm.edu/IJMC/SuperMeanLabelingOfSomeClasses.pdf

[1735] P. Jeyanthi, D. Ramya, and M. Selvi, Even vertex odd mean labeling of transformed trees, *TWMS J. App. Eng. Math.*, **10**(2) 2020 338-345. https://dergipark.org.tr/en/download/article-file/1181243?utm_source=chatgpt.com

[1736] P. Jeyanthi, D. Ramya, and P. Thangavelu, On super mean graphs, *AKCE J. Graphs Combin.,* **6**(1) (2009) 103-112. https://www.researchgate.net/publication/257365261_PJeyanthi_DRamya_and_PThangaveluOn_Super_mean_graphs_AKCE_Graphs_Combin6_No1_2009_103-112

[1737] P. Jeyanthi, D. Ramya, and P. Thangavelu, Some constructions of k-super mean graphs, *Inter. J. Pure Applied Math.,* **56** (2009) 77-86. https://www.researchgate.net/publication/257364411_PJeyanthi_DRamya_and_PThangavelu_Some_Construction_of_k-super_mean_graphs_International_Journal_of_Pure_and_Applied_MathematicsVol56_No1_200977-86

[1738] P. Jeyanthi, D. Ramya, and P. Thangavelu, On super mean labeling of some graphs, *SUT J. Math.,* **46** (2010) 53-66. 83-91 https://vixra.org/pdf/1403.0437v1.pdf

[1739] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling, *J. Scientific Research*, **5** (3) (2013) 457-467. https://www.banglajol.info/index.php/JSR/article/view/15001

[1740] P. Jeyanthi and T. Sarada Devi, On edge pair sum labeling of graphs, *Internat. J. Math. Trends Tech.,* **7**(2) (2014) 106-113. https://ijmttjournal.org/public/assets/volume-7/number-2/IJMTT-V7P514.pdf

[1741] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of spider graph, *J. Algor. Comput.*, **45** (2014) 25-34. https://journals.ut.ac.ir/article_7918_d0a0b362799482703ea6296c4b91f013.pdf

[1742] P. Jeyanthi and T. Sarada Devi, Some edge pair sum graphs, *J. Discr. Math. Science & Crypt.,* **18**(5) (2015) 481-493.

[1743] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of some Cartesian product graphs, *Discrete Math., Algor. Appl.,* **8** (23) (2016) 1650024 (10 pages). https://www.worldscientific.com/doi/abs/10.1142/S1793830916500245

[1744] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of butterfly graph with shell order, *Malaya J. Matematik*, **4**(2) (2016) 205-210. https://www.malayajournal.org/articles/MJM_April_3.pdf

[1745] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of cycle related graphs, *J. Algor. Comput.*, **48** (2016) 57-68. https://journals.ut.ac.ir/article_7940_56568b68b988fc59429af15e748b7a64.pdf

[1746] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of some subdivision of graphs, *Bull. Internat. Math. Virtual Inst.,* **8** (2018) 67-80. http://www.imvibl.org/buletin/bulletin_imvi_8_1_2018/bulletin_imvi_8_1_2017_67_80.pdf

[1747] P. Jeyanthi and T. Saraha Devi, Some new families of edge pair sum graphs, *Palestine J. Math.,* **8**(2) (2019) 61-69. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_61to69.pdf

[1748] P. Jeyanthi and T. Sarada Devi, Edge pair sum labeling of some classes of graphs, *J. Discr. Math. Sci. Crypt.,***24**(4) (2021) 1075-1084.. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_61to69.pdf

[1749] P. Jeyanthi, T. Sarada Devi, and G.-C Lau, Some results on edge pair sum labeling, *Electronic Notes Disc. Math.*, **48** (2015) 169-173. http://dx.doi.org/10.1016/j.endm.2015.05.025

[1750] P. Jeyanthi, T. Sarada Devi, and G.-C. Lau, New results on edge pair sum graphs, *Internat. J. Math. Appl.,* **4**(1-B) (2016) 57-65. https://www.researchgate.net/publication/300165834_New_Results_on_Edge_Pair_Sum_Graphs

[1751] P. Jeyanthi, T. Saratha Devi, and G.-C. Lau, Edge pair sum labeling of $WT(n : k)$ tree, *Global J. Pure and Appl. Math.*, **11**(3) (2015) 1523-1539. https://www.ripublication.com/gjpam%202015/gjpamv11n3_40.pdf

[1752] P. Jeyanthi and P. Selvagopal, Magic covering of chain of an arbitrary 2-connected simple graph, *SUT J. Math.,* **44** (2008) 23-31. https://www.researchgate.net/publication/257365101_PJeyanthi_and_PSelvagopal_Magic_covering_of_chain_of_an_arbitrary_2-connected_Simple_graph_SUT_Journal_of_Mathematics_Vol_44_No12008_23-31

[1753] P. Jeyanthi and P. Selvagopal, More classes of H-supermagic graphs, *Internat. J. Algor., Comput. and Math.,* **3** (2010) 93-108. https://www.researchgate.net/publication/257306209_9_PJeyanthi_and_PSelvagopal_More_Classes_of_H-Supermagic_Graphs_International_Journal_of_Algorithms_Computing_and_Mathematics_Vol3_12010_93-108

[1754] P. Jeyanthi and P. Selvagopal, H-supermagic strength of some graphs, *Tokyo J. Math.,* **33** (2010) 499-507. https://www.researchgate.net/publication/48419529_H-Supermagic_Strength_of_Some_Graphs

[1755] P. Jeyanthi and P. Selvagopal, Some C_4-supermagic graphs, *Ars Combin.,* **111** (2013) 129-136. https://combinatorialpress.com/article/ars/Volume%20111/volume-111-paper-11.pdf

[1756] P. Jeyanthi and P. Selvagopal, Supermagic coverings of some simple graphs, *Internat. J. Math. Combin.,* **1** (2011) 33-48. https://fs.unm.edu/IJMC/SupermagicCoveringsOfSomeSimpleGraphs.pdf

[1757] P. Jeyanthi, P. Selvagopal, and S. Soma Sundaram, Some C_3-supermagic graphs, *Util. Math.,* **89** (2012) 357-366.

[1758] P. Jeyanthi, M. Selvi, and D. Ramya, Skolem difference mean labeling of disconnected graphs, *Proyecciones J. Math.,* **36**(2) (2017) 347-361. https://www.scielo.cl/scielo.php?script=sci_arttext&pid=S0716-09172017000200347

[1759] P. Jeyanthi, M. Selvi, and D. Ramya, Skolem difference mean labeling of union of graphs, *Util. Math.*, **111** (2019) 139-150.

[1760] P. Jeyanthi, M. Selvi, and D. Ramya, Restricted triangular difference mean graphs, *Proyecciones J. Math.,* **39**(2) (2020) 275-286. https://www.researchgate.net/publication/341045505_Restricted_triangular_difference_mean_graphs

[1761] P. Jeyanthi, M. Selvi, and D. Ramya, Odd mean labeling of $T\hat{o}C_n$ and $T\tilde{o}C_n$, *Util. Math.*, **107** (2018) 115-130.

[1762] P. Jeyanthi, M. Selvi, and D. Ramya, Centered triangular difference mean graphs, *Util. Math.,* **112** (2019) 125-138.

[1763] P. Jeyanthi, M. Selvi and D. Ramya, Triangular difference mean graphs, *Internat. J. Math. Combin.,* **2** (2020) 109-117. https://www.researchgate.net/publication/342803089_Triangular_Difference_Mean_Graphs

[1764] P. Jeyanthi and A. Sudha, Total edge irregularity strength of disjoint union of wheel graphs, *Elect. Notes Disc. Math.*, **48** (2015) 175-182. http://dx.doi.org/10.1016/j.endm.2015.05.026

[1765] P. Jeyanthi, M. Selvi and D. Ramya, Skolem difference mean labeling of disconnected graphs, *Proyecciones* **36**(2) (2017) 347-361. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/1650/1331

[1766] P. Jeyanthi, M. Selvi and D. Ramya, One modulo N-difference mean graphs, *Palestine J. Math.*, **12**(3) (2023) 163-169. https://pjm.ppu.edu/sites/default/files/papers/PJM_12%283%29_2023_163_to_169.pdf

[1767] P. Jeyanthi, M. Selvi, and D. Ramya, New results on one modulo N-difference mean graphs, *Open J. Discrete Math.*, **13** (2023) 100-112. https://www.scirp.org/pdf/ojdm_2023102315360487.pdf

[1768] P. Jeyanthi and A. Sudha, Total edge irregularity strength of wheel related graphs, *J. Graph Labeling,* **2**(1) (2015) 45-57. https://www.researchgate.net/publication/286456461_Total_Edge_Irregularity_Strength_of_Wheel_Related_Graphs

[1769] P. Jeyanthi and A. Sudha, On the total irregularity strength of wheel related graphs, *Utilitas Math.*, **110** (2019) 131-134. https://www.researchgate.net/publication/286456461_Total_Edge_Irregularity_Strength_of_Wheel_Related_Graphs

[1770] P. Jeyanthi and A. Sudha, Some results on edge irregular total labeling, *J. Internat. Math. Virtual Institute,* **9** (2019) 73-91. http://www.imvibl.org/journal/9_19/journal_imvi_89_2019_73_91.pdf

[1771] P. Jeyanthi and A. Sudha, Total edge irregularity strength of disjoint union of double wheel graphs, *Proyecciones J. Math.* **35**(3) (2016) 251-262. https://www.scielo.cl/pdf/proy/v35n3/art03.pdf

[1772] P. Jeyanthi and A. Sudha, Total vertex irregularity strength of corona product of some graphs, *J. Algorithms Comput.*, **48**(1) (2016) 127-140. https://journals.ut.ac.ir/article_7948_75d3525b3126762d9d4dfccb440cd4e7.pdf

[1773] P. Jeyanthi and A. Sudha, Total edge irregularity strength of some families of graphs, *Util. Math.*, **109** (2018) 139-153.

[1774] P. Jeyanthi and A. Sudha, Total vertex irregularity strength of some graphs, *Palestine J. Math.*, **7**(2) (2018) 725-733. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2018_725to733.pdf

[1775] P. Jeyanthi and A. Sudha, Total irregularity strength of disjoint union of crossed prism and necklace graphs, *Util. Math.*, **114** (2020) 13-31.

[1776] P. Jeyanthi and A. Sudha, On the total irregularity strength of some graphs, *Bull. Inter. Math. Virtual Inst.,* **9**(2) (2019) 393-401. http://www.imvibl.org/buletin/bulletin_imvi_9_2_2019/bulletin_imvi_9_2_2019_393_401.pdf

[1777] S. Jezný and M. Trenklér, Characterization of magic graphs, *Czechoslovak Math. J.*, **33** (1983) 435-438. https://dml.cz/bitstream/handle/10338.dmlcz/101893/CzechMathJ_33-1983-3_11.pdf

[1778] Y. Ji, J. Liu, Y. Bai, and S. Wu, Full friendly index sets of mC_n, *Frontiers Comput. Sci.,* (electronic) **16**(3) (2021). https://academic.hep.com.cn/fcs/CN/article/downloadArticleFile.do?attachType=PDF&id=30039

[1779] T.-S. A. Jiang, The radio number of grid graphs, arXiv:1401.6583v1 [math.CO] 25 Jan 2014. Chceked web Apr 2025 NOW Chat 10/1/26

[1780] D. J. Jin, S. Z. Liu, S. H. Lee, H. L. Liu, X. G. Lu, and D. Zhang, The joint sum of graceful trees, *Comput. Math. Appl.*, **26** (1993) 83-87. https://core.ac.uk/download/pdf/82770208.pdf

[1781] D. J. Jin, F. H. Meng, and J. G. Wang, The gracefulness of trees with diameter 4, *Acta Sci. Natur. Univ. Jilin.*, (1993) 17-22.

[1782] M. I. Jinnah and S. Beena, On E_4-cordial graphs, *Ars Combin.,* **119** (2015) 413-422.

[1783] M. I. Jinnah and A. V. Nair, On cordial labelling of graphs, *Adv. Appl. Disc. Math.*, **14**(2) 75-94.

[1784] M. I. Jinnah and G. Suresh Singh, A note on arthmetic numberings of graphs, *Proc. Symposium on Graphs and Combinatorics, Kochi, Kerala, India* (1991) 83-87.

[1785] M. I. Jinnah and K. R. Santhosh Kumar, Irregularity strength of triangular snake and double triangular snake, *Adv. Appl. Discrete Math.* **9** (2012)(2) 83-92.

[1786] Jirimutu, On k-gracefulness of r-crown $I_r(K_{1,n})$ $(n \geq 2, r \geq 2)$ for complete bipartite graph, *J. Inner Mongolia Univ. Nationalities,* **2** (2003) 108-110.

[1787] Jirimutu, Study on Graph Labelings and Hypergraph Decomposition, Dalian, PhD dissertation, Dalian University of Techonology (2006).

[1788] Jirimuta, Y-L Bao, and F-L Kong, On k-gracefulness of r-crown for complete bipartite graphs, *International J. Pure and Applied Math.,* **15** (2004) 75-80. https://www.ijpam.eu/contents/2004-15-1/10/10.pdf?utm_source=chatgpt.com

[1789] Jirimuta, J. Wang, and X. Xirong, On the gracefulness of the digraphs $n \cdot \overrightarrow{C_m}$, *Internat.J. Pure and Appl. Math.,* **23** (2005) 393-400.

[1790] Jirimuta and J. Wang, On (a, d)-antimagic labelings of generalized Petersen graphs $P(n, 2)$, *Ars Combin.,* **90** (2009) 161-174.

[1791] Jirimutu, X. Xu, W. Feng, and Y. Bao, Proof of a conjecture on the gracefulness of a digraph, *Util. Math.,* **81** (2010) 255-264.

[1792] P. John, B. Lubis, and L. A. Sugeng, Distance antimagic labeling for copies of graphs, *J. Matematika UNAND* **15**(3) (2026) 319–324. https://jmua.fmipa.unand.ac.id/index.php/jmua/article/view/1655/1062

[1793] R. Jones, K. Kolasinski, and P. Zhang, A proof of the modular edge-graceful trees conjecture, *J. Combin. Math. Combin. Comput.*, **080** (2012) 445-455. https://combinatorialpress.com/article/jcmcc/Volume%20080/vol-080-paper%2030.pdf

[1794] R. Jones and P. Zhang, Nowhere-zero modular edge-graceful graphs, *Discuss. Math. Graph Th.*, **32** (2012) 487–505. https://doi.org/10.7151/dmgt.1621

[1795] R. P. Jones, Gracelessness, Proc. 10th S-E Conf. Combinatorics, Graph Theory, and Computing, Congr. Numer., **XXIII-XXIV**, *Util. Math.,* (1979) 547-552.

[1796] T. A. Jose and M. Giridaran, Radial radio mean labeling of Mongolian tent and diamond graphs, *Internat. J. Res. Appl. Sci. Engin. Tech.*, **8**(7) (2020) 2078-2083. https://www.ijraset.com/fileserve.php?FID=30759

[1797] T. A. Jose, R. Prabakaran, and R. Gomathy, On the radio antipodal mean number of some grid related graphs, *Comm. Math. Appl.*, **14**(2) (2023) 745-758. https://www.rgnpublications.com/journals/index.php/cma/article/view/2043/1565

[1798] G. R. Joseph and J. V. Kureethara, Super vertex sum labeling of graphs, *Int. J. Pure Appl. Math.*, **118**(10) (2018) 293-300. https://www.acadpubl.eu/jsi/2018-118-10-11/articles/10/70.pdf

[1799] G. R. Joseph and J. V. Kureethara, Some graphs with super vertex sum number 2, *Int. J. Appl. Math.*, **31**(4) 641–648. https://www.researchgate.net/publication/327362687_Some_graphs_with_super_vertex_sum_number_2

[1800] J. R. Joshi and D. Parmar, H_k — Cordial labeling of m -polygonal snake graphs, *Alochang Chakra J.*, **IX**(IV) (2020) 1924-1938. https://www.researchgate.net/publication/340923339_HK-_Cordial_labeling_of_m_-11990-1-12234-129054Polygonal_Snake_Graphs/references#fullTextFileContent

[1801] J. R. Joshi and D. Parmar, H_k-cordial labeling of some snake graph, *J. Xidian University*, **14**(3) (2020) 844-885. https://www.researchgate.net/publication/340115606_HK-_Cordial_labeling_of_Some_Snake_Graphs

[1802] A. Jovanna, Relaxed Skolem mean labeling of 4-star graph with partition (3,1), *Internat. J. Math. Comp. Res.,* **13** (2025) 5084-5086. https://zenodo.org/records/15233852

[1803] G. Jothi, Highly vertex prime graphs, personal communication (2011).

[1804] J. S-T. Juan and D-F. Liu, Antipodal labelings of cycles, *Ars Combin.*, **103** (2012) 81-96. https://combinatorialpress.com/ars-articles/volume-103-ars-articles/antipodal-labelings-for-cycles/

[1805] R. Junetty, D. Indriati, and B. Winarno, Edge irregular reflexive labeling of palm tree graph $C_3 - B_{2,r}$ and $C_3 - B_{3,r}$, *AIP Conf. Proc.*, **2566**(1) (2022) https://aip.scitation.org/doi/pdf/10.1063/5.0116566

[1806] D. Jungreis and M. Reid, Labeling grids, *Ars Combin.,* **34** (1992) 167-182. https://www.cflmath.com/~reid/Research/Grids/grids.pdf

[1807] M. K. A. Kaabar and K. Yenoke, Radio and radial radio numbers of certain sunflower extended graphs, *Hindawi Internat. J. Math. Math. Sci.,* (2022) Article ID 9229409, 1-9. https://onlinelibrary.wiley.com/doi/epdf/10.1155/2022/9229409

[1808] M. Kalaimathi and B. J. Balamurugan, Computation of even-odd harmonious labeling of certain family of acyclic graphs, *Internat. J. Engin. Adv. Tech.,* **9**(1S3) (2019) 414-419. https://www.ijeat.org/wp-content/uploads/papers/v9i1s3/A10751291S319.pdf

[1809] M. Kalaimathi and B. J. Balamurugan, Biconditional cordial labeling of cycle related graphs, *AIP Confer. Proc.*, (2020) 2282(1):020017 doi:10.1063/5.0028659

[1810] M. Kalaimathi, B.J. Balamurugan, and J. V. Rao, Computation of biconditional cordial labeling of super subdivision of graphs, In: R.R. Raje, F. Hussain, and R.J. Kannan (eds) Artificial Intelligence and Technologies. Lecture Notes in Electrical Engineering, **806** Springer, Singapore, (2020) https://doi.org/10.1007/978-981-16-6448-9_60.

[1811] S. Kalaimathy, Some results on (k, d)-even mean labeling, *Inter. J. Math. Soft Comput.,* **2**(2) (2012) 103-112. https://oaji.net/articles/2017/296-1502656938.pdf

[1812] R. Kalaiyarasi and D. Ramya, Skolem odd difference mean labeling of trees, *Global J. Pure Appl. Math.*, **11**(2) N (2015) 887-898. https://ripublication.com/gjpam%202015/gjpamv11n2_29.pdf

[1813] R. Kalaiyarasi, D. Ramya, and P. Jeyanthi, Skolem odd difference mean labeling of trees, *Global J. Pure and Appl. Math.*, **11**(2) (2015) 887-898. https://www.ripublication.com/gjpam%202015/gjpamv11n2_29.pdf?utm_source=chatgpt.com

[1814] M. Kalkowski, M. Karoński, and F. Pfender, A new upper bound for the irregularity strength of graphs, *SIAM J. Discrete Math.,* **25** (2011) 1319-1321. https://www.math.uni-rostock.de/~pfender/papers/24.pdf

[1815] S. Kalyan and R. Kempepatil, Note on advanced labeling and Fibonacci graceful graphs, *Pramana Res. J.*, **9**(5) (2019) 295-317. https://www.pramanaresearch.org/gallery/prj-p810.pdf

[1816] N. Kamachi and S. Arumugam, Distance antimagic graphs, *J. Combin. Math. Combin. Comput.*, **84** (2013) 61–67. https://combinatorialpress.com/article/jcmcc/Volume%20084/vol-084-paper%206.pdf

[1817] N. Kamatchi, K. Paramasivam, A. V. Prajeesh, K. M. Sabeel, and S. Arumugam, On group vertex magic graphs, *AKCE Int. J. Graphs Comb.*, **17**(1) (2020) 461-465. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2019.04.001

[1818] N. Kamatchi, A. Ramalakshmi, and S. Nilavarasi, A note on distance magic and distance antimagic graphs, *Elect. Notes Discr. Math.*, **48** (2015) 183-187. http://dx.doi.org/10.1016/j.endm.2015.05.027

[1819] T. Kamaraj and J. Thangakani, Edge even and edge odd graceful labelings of Paley graphs, *J. Phys.: Conf. Ser.,* (2021 1770 012068 https://iopscience.iop.org/article/10.1088/1742-6596/1770/1/012068/pdf

[1820] N. Kamatchi, G. R. Vijayakumar, A. Ramalakshmi, S. Nilavarasi, and S. Arumugam, Distance antimagic labelings of graphs, *Theoretical Comput. Sci. Discr. Math.*, 113-118, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017.

[1821] J. C. Kanani and V. J. Kaneria, Graceful labeling for some snake related graphs, *Ganita,* **71** (1) (2021) 243-256. https://bharataganitaparisad.com/wp-content/uploads/2021/10/711-ch025.pdf

[1822] K. K. Kanani and M. I. Bosmia, On cube divisor cordial graphs, *Internat. J. Math. Comput. Appl. Res. (IJMCAR)*, **5**(4) (2015) 117-128. https://www.researchgate.net/publication/281078200_ON_CUBE_DIVISOR_CORDIAL_GRAPHS

[1823] K. K. Kanani and T. M. Chhaya, Strongly multiplicative labeling of some path related graphs, *Internat. J. Math. Comput. Appl. Res. (IJMCAR)*, *5*(5) (2015) 1-6. https://www.researchgate.net/publication/288642835_STRONGLY_MULTIPLICATIVE_LABELING_OF_SOME_PATH_RELATED_GRAPHS

[1824] K. K. Kanani and T. M. Chhaya, Some new families of strongly multiplicative graphs, 9th National Level Science Symposium. February 14, 2016, Organized by Christ College, Rajkot, Sponsored by GUJCOST, Gandhinagar, Mathematics & Statistics, **3** (2016) 197-200.

[1825] K. K. Kanani and M. V. Modha, 7-cordial labeling of standard graphs, *Internat. J. Appl. Math. Res.,* **3**(4) (2014) 547-560. https://www.sciencepubco.com/index.php/ijamr/article/view/3371

[1826] K. K. Kanani and N. B. Rathod, Some new 4-cordial graphs, *J. Math. Comput. Sci.,* **4**(5) (2014) 834-848. https://www.researchgate.net/publication/319351983_SOME_NEW_4-CORDIAL_GRAPHS

[1827] K. K. Kanani and T. M. Chhaya, Strongly multiplicative labeling of some standard graphs, *Internat. J. Math. Soft Comput,* **7**(1) (2017) 13-21. https://www.researchgate.net/publication/317168353_Strongly_Multiplicative_Labeling_of_some_Standard_Graphs

[1828] K. K. Kanani and T. M. Chhaya, K. K. Kanani and T. M. Chhaya, Strongly multiplicative labeling of some snake related graphs, *Internat. J. Math. Trends and Tech. (IJMTT),* **45**(1) (2017) 53-56. https://www.iosrjournals.org/iosr-jm/papers/Vol18-issue1/Ser-2/B1801022430.pdf

[1829] V. J. Kaneria and P. P. Andharia, Some results on even sum labeling of graphs, *J. Calcutta Math. Soc.,* **15**(2) (2019) 129-138.

[1830] V. J. Kaneria and P. P. Andharia, Further results on even sum labeling of graphs, *Global J. Pure and Appl. Math.,* **16**(2) (2020) 155-166. https://www.ripublication.com/gjpam20/gjpamv16n2_02.pdf

[1831] V. J. Kaneria and H. P. Chudasama, Absolute mean graceful labeling in various graphs, *Int. J. Math. Appl.,* **5**(4) (2017) 723-726. https://mjis.chitkara.edu.in/index.php/mjis/article/view/15/12

[1832] V. J. Kaneria, H. P. Chudasama, and P. P. Andharia, Absolute mean graceful labeling in path union of various graphs, *Math. J. Interdiscip. Sci.,* **7**(1) (2018), 51-56. https://www.researchgate.net/publication/345371202_Absolute_Mean_Graceful_Labeling_in_Path_Union_of_Various_Graphs

[1833] V. J. Kaneria, A. M. Gohil, and H. M. Makadia, Graceful related labeling and its applications, *Int. J. Math. Res.*, **7**(1) (2015) 47-54. http://www.irphouse.com/ijmr/ijmrv7n1_09.pdf

[1834] V. J. Kaneria, M. M. Jariya, and H. Karavadiya, Cordially of the complete graphs, *J. Graph Labeling,* **2**(2) (2016) 89-101.

[1835] V. J. Kaneria and M. M. Jariya, Semi smooth graceful graph and construction of new graceful trees, *Elixir Appl. Math.,* **76** (2014) 28536-28538. https://www.elixirpublishers.com/articles/1678942777_201411087.pdf

[1836] V. J. Kaneria and M. M. Jariya, Smooth graceful graphs and its applications to construct graceful graphs, *Int. J. Sci. and Res.,* (IJSR) **3**(8) (2014) 909-912. https://www.ijsr.net/archive/v3i8/MDIwMTU0ODg=.pdf

[1837] V. J. Kaneria, M. M. Jariya, and H. M. Makadia, Graceful labeling of arrow graphs and double arrow graphs, *Malaya J. Matematik*, **3**(4)(2015) 382-386. https://www.malayajournal.org/articles/paper2_2015.pdf

[1838] V. J. Kaneria, M. M. Jariya, and M. Meghpara, Graceful labeling for some star related graphs, *Int. Math. Forum,* **9**(26) (2014) 1289-1293. https://www.researchgate.net/publication/318636729_Graceful_Labeling_for_Some_Star_Related_Graphs

[1839] V. J. Kaneria, M J Khoda, and H M Karavadiya, Balanced mean cordial labeling and graph operations, *Int. J. Math. Appl.,* **4**(3–A) (2016) 181-184. https://ijmaa.in/index.php/ijmaa/article/view/983/967

[1840] V. J. Kaneria, M. Meghpara and M. Khoda, Semi smooth graceful labeling and its application to produce α-labeling, *J. Graph Labeling*, **2**(2) (2016) 153-160.

[1841] V. J. Kaneria and H. M. Makadia, Some graceful graphs, *J. of Math. Res.,* **4**(1) (2012) 54-57. https://www.ccsenet.org/journal/index.php/jmr/article/view/12190

[1842] V. J. Kaneria and H. M. Makadia, Graceful labeling for tensor product of two path of odd lengths and some grid related graphs, *Int. J. Innovation Sci. Math.,* **2**(5) (2014) 470-472. https://www.ijism.org/administrator/components/com_jresearch/files/publications/IJISM_260_Final.pd

[1843] V. J.c Kaneria and H. M. Makadia, Graceful labeling for step grid graph, *J. Advances Math.*, **9**(5) (2014) 2647-2654. https://rajpub.com/index.php/jam/article/view/2335

[1844] V. J. Kaneria and H. M. Makadia, Some results on graceful labeling for step grid related graphs, *AKCE Int. J. Math. Trends Tech. (IJMTT)* **65** (11) (2019) 29-38. doi:DOI: 10.14445/22315373/IJMTT-V65I11P503

[1845] V. J. Kaneria and H. M. Makadia, Graceful labeling for double step grid graph, *Int. J. Math. Appl.,* **3**(1) (2015) 33-38. https://ijmaa.in/index.php/ijmaa/article/view/400

[1846] V. J. Kaneria and H. M. Makadia, Graceful labeling for plus graph, *Int. J. Current Rec. Sci. Tech.*, **1**(3) (2015) 15-20. https://crst.gfer.org/index.php/crst/article/view/9

[1847] V. J. Kaneria and H. M. Makadia, Some results on graceful labeling for families of plus graph, *Int. J. Current Rec. Sci. Tech.*, **1**(4) (2015) 17-23. https://crst.gfer.org/index.php/crst/article/view/18/18

[1848] V. J. Kaneria and H. M. Makadia, Graceful labeling for swastik graph, *Int. J. Math. Appl.*, **3**(3-D) (2015) 25-29 http://ijmaa.in/v3n3-d/25-29.pdf

[1849] V. J. Kaneria and H. M. Makadia, Some results on graceful labeling for families of swastik graphs, *Advances and Appl. Discrete Math.*, **16**(2) (2015) 161-172. https://doi.org/10.17654/AADMOct2015_161_172

[1850] V. J. Kaneria, H. M. Makadia, and M. M. Jariya, Graceful labeling for cycle of graphs, *Int. J. Math. Res.,* **6** (2) (2014) 173-178. http://www.irphouse.com/ijmr/ijmrv6n2_08.pdf

[1851] V. J. Kaneria, H. M. Makadia, M. M. Jariya, and M. Meghpara, Graceful labeling for complete bipartite graphs, *Appl. Math. Sci.,* **8** (103) (2014) 5099-5104. https://www.m-hikari.com/ams/ams-2014/ams-101-104-2014/makadiaAMS101-104-2014.pdf

[1852] V. J. Kaneria, H. M. Makadia, and M. Meghpara, Some graceful graphs, *Int. J. Math. Soft Comp.,* **4**(2) (2014) 165-172. https://www.ccsenet.org/journal/index.php/jmr/article/view/12190

[1853] V. J. Kaneria, H. M. Makadia, and M. Meghpara, Gracefulness of cycle of cycles and cycle of complete bipartite graphs, *Int. J. Math. Trend Tech.,* **12**(1) (2014) 19-26. https://ijmttjournal.org/archive/ijmtt-v12p503

[1854] V. J. Kaneria, H. M. Makadia, and M. Meghpara, Cordiality of a star of the complete graph and a cycle graph $C(n \cdot K_n)$, *J. Math. Res.,* **6**(4) (2014) 18-28. https://ccsenet.org/journal/index.php/jmr/article/view/39732

[1855] V. J. Kaneria, H. M. Makadia, and M. Meghpara, Graceful labeling for grid related graphs, *Int. J. Math. Soft Comput.,* **5**(1) (2015), 111-117. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=ecd8a1787bf69016ddf21eedbf5ea1276fce7fd9

[1856] V. J. Kaneria, H. M. Makadia and R. V. Viradia, Graceful labeling for disconnected grid related graphs, *Bull. Math. Sci. Appl.*, **4**(1) (2015) 6-11. https://www.semanticscholar.org/reader/012cf0654bf4ee03acf635c55f0d52c3cdfa595d

[1857] V. J. Kaneria, H. M. Makadia and R. V. Viradia, Some results on graceful labeling for double step grid related graphs, *Int. J. Math. Sci. Appl.*, **9**(1) (2015) 117-127.

[1858] V. J. Kaneria, H. M. Makadia and R. V. Viradia, Various graph operation on semi smooth graceful graphs, *Int. J. Math. and Soft Computing*, **6**(1) (2016 57-79. https://www.researchgate.net/publication/321164406_Various_Graph_Operations_on_Semi_Smooth_Graceful_Graphs

[1859] V. J. Kaneria, M. Meera, and K. Maulik, Geometric mean 3-equitable labeling of some graphs, *Internat. J. Sci. Res. Reviews*, IJSRR (2018) **7**(1) Suppl. 245-250. https://www.researchgate.net/publication/327718616_Geometric_mean_3-equitable_labeling_of_some_graphs

[1860] V. J. Kaneria and M. Meghpara, Mean labeling for some cycle of graphs, *Int. J. Math. Sci. Eng. Appl.*, **9**(2) (2015) 267-274. https://www.researchgate.net/publication/318653709_mean_labeling_for_some_cycle_of_graphs

[1861] V. J. Kaneria and M. Meghpara, Semi smooth graceful labeling on some graphs, *Int. J. Math. Appl.*, **3** (3C) (2015) 1-5. https://ijmaa.in/index.php/ijmaa/article/view/460/458

[1862] V. J. Kaneria and M. Meghpara, Graceful labeling for one point union for path of graphs, *Int. J. Math. Appl.*, **3**(1) (2015) 49-54. https://www.researchgate.net/profile/Meera-Meghpara/publication/318636593_Graceful_labeling/links/59731bbaa6fdcc8348826e5d/Graceful-labeling.pdf

[1863] V. J. Kaneria, M. Meghpara, and H. M. Makadia, Mean labeling for step grid graph, *Advan. Appl. Math. Sci.*, **14**(1) (2015) 111-114. https://www.researchgate.net/publication/318636734_MEAN_LABELING_FOR_STEP_GRID_GRAPH

[1864] V. J. Kaneria, M. Meghpara, and H. M. Makadia, Graceful labeling for one point union of path and barycentric subdivision of a grid graph, *Int. J.*

Math. Comp. Res., **2**(9) (2014) 624-629. https://www.researchgate.net/publication/318636575_GRACEFUL_LABELING_FOR_ONE_POINT_UNION_OF_PATH_AND_BARYCENTRIC_SUBDIVISION_OF_A_GRID_GRAPH

[1865] V. J. Kaneria, M. Meghpara, and H. M. Makadia, Graceful labeling for open star of graphs, *Int. J. Math. Stat. Invention,* (IJMSI), **2**(9) (2014) 19-23. https://ijmsi.org/Papers/Volume.2.Issue.9/E029019023.pdf

[1866] V. J. Kaneria, M. Meghpara, and H. M. Makadia, Cordial labeling for cycle of complete bipartite graphs and cycle of wheels, *Int. J. Pure Appl. Math.,* (IJPAM), **101**(1) (2015) 1-8. https://www.researchgate.net/publication/275240630_Cordial_labeling_for_cycle_of_complete_bipartite_graphs_and_cycle_of_wheels

[1867] V. J. Kaneria, K. M. Patadiya, and J. R. Teraiya, Balanced cordial labeling and its application to produce new cordial families, *Int. J. Math. Appl.*, **4** (1-C) (2016) 65-68. https://ijmaa.in/index.php/ijmaa/article/view/595

[1868] V. J. Kaneria and J. M. Shah, Absolute mean graceful labelings in the context of m[splitting and degre spitting graphs, *South East Asian J. Math. Math. Sci.*, **19**(3) (2023) 359-370. https://doi.org/10.56827/SEAJMMS.2023.1903.28

[1869] V. J. Kaneria, O. Teraiya and M. Meghpara, Double path union of α-graceful graph and its α-labeling, *J. of Graph Labeling*, **2**(2) (2016) 107-114.

[1870] V. J. Kaneria and J. R. Teraiya, Complete star of a graph and its balanced cordial labeling, *Internat. J. Math. Soft Comput.,* **7**(1) (2017) 89-94.

[1871] V. J. Kaneria, J. R. Teraiya, and K. M. Patadiya, Some result on balanced cordial graphs, *Int. J. Math. Appl.*, **4** (2-A), (2016) 85-87. https://ijmaa.in/index.php/ijmaa/article/view/1009/994

[1872] V. J. Kaneria and S. K. Vaidya, Index of cordiality for complete graphs and cycle, *Inter. J. Applied Math. and Computation,* **2(4)** (2010) 38-46. http://www.darbose.in/ojs/index.php/ijamc/article/view/2.4.5

[1873] V. J. Kaneria, S. K. Vaidya, G. V. Ghodasara, and S. Srivastav, Some classes of disconnected graceful graphs, *Proc. First Internat. Conf. Emerging Technologies and Appl. Engin. Tech. Sci.,* (2008) 1050-1056.

[1874] V. J. Kaneria, R. V. Viradia, M. M. Jariya, and H. M. Makadia, Various labeling for the graph $C(t \cdot P_n)$, *Int. J. Math. Comp. Res., 2*(11) (2014) 745-751.

[1875] V. J. Kaneria, R. V. Viradia and H. M. Makadia, Mean labeling for path union and cycle of graphs, unpublished.

[1876] A. Kanetkar, Prime labeling of grids, *AKCE J. Graphs Combin.,* **6** (2009) 135-142. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088880

[1877] A. Kanetkar, S. S. Sane, Graceful labeling of a family of quasistars with paths in arithmetic progression, *Ars Combin.,* **83** (2007) 307-320. https://combinatorialpress.com/article/ars/Volume%20083/volume-83-paper-23.pdf

[1878] N. Kang, S. Chen, Z. Li, and L. Hou, D-magic labelings of the halved n-cube, *Discrete Math.*, **345**(11) (2022) 113044. 8 pages. https://www.sciencedirect.com/science/article/pii/S0012365X22002503?via%3Dihub

[1879] S. M. Kang, S. Nazeer, W. Nazeer, I. Kousar and C. Y. Jung, Radio labeling and radio number of caterpillar related graphs, *Mitteilungen Klosterneuburg,* **65**(5) (2015) 149-159. https://www.researchgate.net/publication/276937980_Radio_Labeling_and_Radio_Number_of_Caterpillar_Related_Graphs

[1880] S. M. Kang, S. Nazeer, W. Nazeer, and B.-Y. Lee, Radio number of caterpillar graphs, *Wulfenia,* **22**(5) (2015) 48-58. https://www.researchgate.net/publication/276937980_Radio_Labeling_and_Radio_Number_of_Caterpillar_Related_Graphs

[1881] Q. D. Kang, The k-gracefulness of the product graphs $P_m \times C_{4n}$, *J. Math. Res. Exposition*, **9** (1989) 623-627.

[1882] Q. Kang, S. M. Lee, and L. Wang, On the edge-graceful spectra of the wheel graphs, unpublished.

[1883] Q. D. Kang, Z.-H. Liang, Y.-Z. Gao, and G.-H. Yang, On the labeling of some graphs, *J. Combin. Math. Combin. Comput.*, **22** (1996) 193-210. https://combinatorialpress.com/article/jcmcc/Volume%20022/vol-22-paper%2019.pdf

[1884] Q. D. Kang and X. Zhao, Strongly harmonious labelings of windmill graphs, *J. Hebei Normal College*, **2** (1992) 1-7.

[1885] A. R. Kannan, A novel approach to C_{exp} average assignments on chain graphs, *Baghdad Sci. J.* (2025) **22**(4) 1317–1327. https://www.researchgate.net/publication/394363897_A_novel_approach_to_Cexp_average_assignments_on_chain_graphs

[1886] A. R. Kannan, P. Manivannan, K. Loganathan, and S. Gyeltshen, A new notion of classical mean graphs based on duplicating operations, *Hindawi J. Math.*, Vol. 2022, Article ID 9253512, 13 pages. https://www.hindawi.com/journals/jmath/2022/9253512/

[1887] M. Kannan, R. Vikrama Prasad, and R. Gopi, Super root square mean labeling of disconnected graphs, *Internat. J. Math. Appl.,* **4**(1–C) (2016) 93-98. https://www.researchgate.net/publication/303814124_Super_Root_Square_Mean_Labeling_Of_Disconnected_Graphs

[1888] M. Kannan, R. Vikrama Prasad, and R. Gopi, Even vertex odd mean labeling of some graphs, *Global J. Pure and Applied Math.,* **13**(3) (2017) 1019-1034. https://www.researchgate.net/publication/315809319_Even_Vertex_Odd_Mean_Labeling_of_Some_Graphs

[1889] M. Kannan, R. Vikrama Prasad, and R. Gopi, Even vertex odd mean labelings of H-graph, *Internat. J. Math. Archive,* **8**(8) (2017) 162-167. https://www.researchgate.net/publication/320195593_EVEN_VERTEX_ODD_MEAN_LABELING_OF_H-GRAPH

[1890] M. Kannan, R. Vikrama Prasad, and R. Gopi, A notion of even vertex odd mean labeling graphs, *Internat. J. Math. Trends Tech.*, (IJMTT) **50**(3) (2017) 153. https://ijmttjournal.org/2017/Volume-50/number-3/IJMTT-V50P524.pdf

[1891] M. Kannan, R. Vikrama Prasad, and R. Gopi, Some graph operations of even vertex odd mean labeling graphs, *Internat. J. Appl. Eng. Res.,* **12**(18) (2017) 7749-7753. https://www.ripublication.com/ijaer17/ijaerv12n18_86.pdf

[1892] M. Kannan, R. Vikrama Prasad, and R. Gopi, Even vertex odd mean labeling of some graph, *Global Pure Appl. Math.*, **13**(2) (2017) 1019-1034. https://www.ripublication.com/gjpam17/gjpamv13n3_13.pdf

[1893] M. Kannan, R. Vikrama Prasad, and R. Gopi, A notion of even vertex odd mean labeling graphs, *Internat. J. Math. Trends Tech.* (IJMTT), **50**(3) (2017) 153-156. https://ijmttjournal.org/public/assets/volume-50/number-3/IJMTT-V50P524.pdf

[1894] A. Rajesh Kannan, P. Manivannan and A. Durai Baskar, Exponential mean labeling of some graphs obtained from duplicating operations, *J. Physics Confer. Ser.,* **1597**(1):012028 (2020) 1-10. https://iopscience.iop.org/article/10.1088/1742-6596/1597/1/012028/pdf

[1895] A. N. Kansagara and S. K. Patel, Prime labeling in the context of web graphs without center, *AKCE Int. J. Graphs Comb.,* **18**(3) (2021) 132-142. https://www.tandfonline.com/doi/epub/10.1080/09728600.2021.1974286?needAccess=true

[1896] S. Kanwal, Z. Iftikhar, and A. Azam, SEMT labelings and deficiencies of forests with two components (I), *Punjab Univ. J. Math. (Lahore)* **51**(5) (2019) 137-149. https://pu.edu.pk/images/journal/maths/PDF/Paper-10_51_5_2019.pdf

[1897] S. Kanwal, A. Azam, and Z. Iftikhar, SEMT labelings and deficiencies of forests with two components (II), *Punjab U. of Math.,* **51**(4)(2019) 1-12. https://pu.edu.pk/images/journal/maths/PDF/Paper-1_51_4_2019.pdf

[1898] S. Kanwal, M. Imtiaz, Z. Iftikhar, R. Ashraf, M. Arshad, R. Irfan, and I. Sumbal, Embedding of supplementary results in strong EMT valuations and strength, *Open Math., 17*(1) (2019) 527-543. https://www.degruyterbrill.com/document/doi/10.1515/math-2019-0044/html?utm_source=chatgpt.com

[1899] S. Kanwal, S. Javed, and A. Riasat, On the super edge-magicness and the deficiency of some families of acyclic graphs, *Util. Math.,* **100** (2016) 323-356. https://www.researchgate.net/publication/314950411_On_the_Super_Edge-Magicness_and_the_Deficiency_of_Some_Families_of_Acyclic_Graphs

[1900] S. Kanwal and I. Kanwal, SEMT valuations of disjoint union of combs, stars and banana trees, *Punjab Univ. J. Math. (Lahore)*, **50**(3) (2018) 131-144. https://pu.edu.pk/images/journal/maths/PDF/Paper-11_50_3_2018.pdf

[1901] S. Kanwal, A. Riasat, M. Imtiaz, Z. Iftikhar, S. Javed, and S. Ashraf, Bounds of strong EMT strength for certain subdivision of star and bistar, *Open Math.,* **16** (2018) 1313-1325. https://www.researchgate.net/publication/329165156_Bounds_of_Strong_EMT_Strength_for_certain_Subdivision_of_Star_and_Bistar

[1902] R. R. Kanzariya, M. A. Chaurasia, and M. P. Rupani, Prime labeling of subdivided helm graphs, *Adv. Appl. Disc. Math.*, **43**(4) (2026) 401-418. https://doi.org/10.17654/0974165826026

[1903] R. R. Kanzariya and M. P. Rupani, New families of prime graphs via vertex expansions, *Internat. J. Math. Trends Tech.*, **72**(6) (2026), 59-70. https://ijmttjournal.org/archive/IJMTT-V72I6P107

[1904] G. Kaplan, A. Lev, and Y. Roditty, Bertrand's postulate, the prime number theorem and product anti-magic graphs, *Discrete Math.,* **308** (2008) 787-794. https://doi.org/10.1016/j.disc.2007.07.049

[1905] G. Kaplan, A. Lev, and Y. Roditty, On zero-sum partitions and anti-magic trees, *Discrete Math.,* **309** (2009) 2010-2014. https://core.ac.uk/download/pdf/82696115.pdf

[1906] N. Karst, J. Langowitz, J. Oehrlein, and D. S. Troxell, Radio k-chromatic number of cycles for large k, *Discrete Math. Algorithms Appl.,* **9**(3) (2017), 1750031, 20pp. https://doi.org/10.1142/S1793830917500318

[1907] M. K. Karthik Chidambaram, On prime labeling of some special graphs, *Internat. J. Food and Nutritional Sci.*, **11**(11) (2022) 2068-2073. https://www.slideshare.net/slideshow/prime-labelling-of-some-special-graphs/55406128

[1908] M. K. Karthik Chidambaram, S. Athisayanathan, and R. Ponraj, Group $\{1, -1, i, -i\}$ cordial labeling of some splitting graphs, *Global J. Engin. Sci. Researches*, **4**(11) (2017) 83-88. https://www.academia.edu/en/36038631/Group_1_1_i_i_Cordial_Labeling_of_Some_Splitting-_Graphs

[1909] M. K. Karthik Chidambaram, S. Athisayanathan, and R. Ponraj, Group $\{1, -1, i, -i\}$ cordial labeling of product related graphs, *Internat. J. Adv. Sci. Res. Devel.,* **04** (06/I), (2017) 27-33. https://www.academia.edu/33810064/Group_1_1_i_i_Cordial_Labeling_of_Product_Related_Graphs

[1910] S. Karthik, K. M. Sabeel, and K. Paramasivama, Group vertex magicness of product graphs and trees, https://www.researchgate.net/publication/368688402_Group_vertex_magicness_of_product_graphs_and_trees

[1911] C. Karthikeyan, S. Arthi, M. Abinaya, R. Swathi, A. Madhumathi, Super Fibonacci graceful labelling of some cycle related graphs, *JSRD -Internat. J. Sci. Res. Develop.,* **5**(12) (2018) 196-198. https://www.researchgate.net/publication/348002618_Super_Fibonacci_Graceful_Labeling_Of_Some_Cycle_Related_Graphs

[1912] S. Karthikeyan, S. Navanaeethakrishnan, and R. Sridevi, Total edge Fibonacci irregular labeling of some star graphs, *Int. J. Math. Soft Comput.*, **5**(1) (2015) 73-78. https://oaji.net/articles/2017/296-1504962802.pdf

[1913] G. P. Karthiyayini1 and D. A. Jovanna, Study on relaxed Skolem mean labeling of 4-star graph with partition (3,1), *Internat. J. Math. Comput. Res.*, **14**(03) (2026) 6219-6221. https://ijmcr.in/index.php/ijmcr/article/view/1257/965

[1914] K. Karuppasamy and S. Kaleeswari, Total mean labeling graphs, *Internat. J. Recent Tech. Engin.* (IJRTE), **8**(4S4) (2019) 90-92. https://www.ijrte.org/wp-content/uploads/papers/v8i4s4/D10361284S419.pdf

[1915] K. Kasthuri, K. Karuppasamy, and K. Nagarajan, SD-divisor labeling of path and cycle related graphs, *AIP Conference Proceed.,* 2463, 030001 (2022) 030001-1-030001-10.

[1916] F. Kastrati1, W. Myrvold, L. D. Panjer, and A. Williams, Cordial forests, In: Fernau, H., Jansen, K. (eds) Fundamentals of Computation Theory. FCT 2023. Lecture Notes in Computer Science, **14292** Springer, Cham., 291-303. https://doi.org/10.1007/978-3-031-43587-4_21

[1917] F. Kastrati, W. Myrvold, L. D. Panjer, and A. Williams, Cordial forests, Conference: 24th International Symposium on Fundamentals of Computation Theory (FCT 2023) At: Trier, Germany, September 2023. https://www.researchgate.net/publication/373140098_Cordial_Forests

[1918] K. Kathiresan, Subdivisions of ladders are graceful, *Indian J. Pure Appl. Math.,* **23** (1992) 21-23.

[1919] K. Kathiresan, Two classes of graceful graphs, *Ars Combin.*, **55** (2000) 129-132.

[1920] K. Kathiresan, Graceful labeling of ladders with pendant edges, *Internat. J. Appl. Graph Th. in Wireless Ad hoc Networks and Sensor Networks* **8**(1) (2016) 1-8. https://www.researchgate.net/publication/300084258_Ladder_and_Subdivision_of_Ladder_Graphs_with_Pendant_Edges_are_Odd_Graceful

[1921] K. Kathiresan, Odd graceful graphs, unpublished.

[1922] K. Kathiresan and S. Amutha, Arbitrary supersubdivisions of stars are graceful, *Indian J. Pure Appl. Math.,* **35** (2004) 81-84.

[1923] K. M. Kathiresan and S. Amutha, Fibonacci Graceful Graphs, Ph. D. Thesis, Madurai Kamaraj University, October 2006.

[1924] K. M. Kathiresan and S. Amutha, Fibonacci graceful graphs, *Ars Combin.*, **97** (2010) 41-50. https://combinatorialpress.com/ars-articles/volume-097-ars-articles/fibonacci-graceful-graphs/

[1925] K. Kathiresan and R. Ganesan, A labeling problem on the plane graphs $P_{a,b}$, *Ars Combin.,* **73** (2004) 143-151. https://combinatorialpress.com/ars-articles/volume-073-ars-articles/a-labeling-problem-on-the-plane-graphs-p_ab/

[1926] K. Kathiresan and R. Ganesan, d-antimagic labelings of plane graphs P_a^b, *J. Combin. Math. Combin. Comput.,* **52** (2005) 89-96. https://combinatorialpress.com/jcmcc-articles/volume-052/b-antimagic-labelings-of-plane-graphs-p_ba/

[1927] K. Kathiresan and S. Gokulakrishnan, On magic labelings of type $(1, 1, 1)$ for the special classes of plane graphs, *Util. Math.,* **63** (2003) 25-32.

[1928] K. M. Kathiresan and S. D. Laurence, On super (a, d)-H-antimagic total covering of star related graphs, *Discuss. Math. Graph Theory,* **35**(4) (2015) 755-764. https://sciendo.com/pdf/10.7151/dmgt.1832

[1929] K. M. Kathiresan, S. Muthumari, and R. Ramalakshmi, Edge odd graceful labelings of certain new classes of graphs, *Bull. Inst. Combin. Appl.,* **95** (2022) 57-69. https://bica.the-ica.org/Volumes/95/Reprints/BICA2022-03-Reprint.pdf

[1930] K. Kathiresan, S. Muthuvel, and V. Nagasubbu, Consecutive labelings for two classes of plane graphs, *Util. Math.,* **55** (1999) 237-241. NOW Chat 10/1/26

[1931] K. Kathiresan and S. Sabarimalai Madha, Star super edge-magic deficiency of graphs, *Contrib. Discrete Math.,* **12**(1) (2017) 143-156. https://cdm.ucalgary.ca/article/view/62318/46780

[1932] K. Kathiresan and R. Sumathi, Solution to an open problem in gracefulness of arbitrary supersubdivisions of graphs, *Util. Math.,* **84** (2011) 333-338.

[1933] S. Kavitha and G. Jayalalitha, Finite and neighborhood Gaussian prime distance graph, *J. Data Acquisition Processing*, **38**(1) (2023) 4062-4073. https://www.researchgate.net/publication/370717959_FINITE_AND_NEIGHBORHOOD_GAUSSIAN_PRIME_DISTANCE_GRAPHS_1

[1934] S. Kavitha and V. L. Stella Arputha Mary, Results on hefty V_4-vertex magic graph labeling, *J. Pharmaceutical Negative Results* **14** Special Issue 1 (2023) 864-871. https://www.pnrjournal.com/index.php/home/article/view/7413/9724

[1935] K. Kayathri and R. Amutha, Edge-graceful labelings of connected graphs, *Electronic Notes Discrete Math.,* **53** (2016) 287-296. https://www.sciencedirect.com/science/article/pii/S1571065316300786

[1936] M. Kchikech, R. Khennoufa, and O. Togni, Linear and cyclic radio k-labelings of trees, *Discuss. Math. Graph Theory,* **27** (2007) 105-123. https://bibliotekanauki.pl/articles/743693

[1937] M. Kchikech, R. Khennoufa, and O. Togni, Radio k-labelings for cartesian products of graphs, *Discuss. Math. Graph Theory,* **28** (2008) 165-178. https://www.researchgate.net/publication/220081789_Radio_k-Labelings_for_Cartesian_Products_of_Graphs

[1938] E. Khairunnisa, D. E. Nurvazly, and N. P. Indah, Pelabelan graceful graf hasil korona bunga mawar, *FARABI J. Matematika dan Pendidikan Matematika*, **8**(1) (2025) 169-176. https://www.researchgate.net/publication/408156668_Pelabelan_Graceful_Graf_Hasil_Korona_Bunga_Mawar

[1939] E. Khairunnisa and K. A. Sugeng, Graceful labelling of corona product of aster flower graph, *Adv. Intel. Systems Res.*, (AISR), **157** (2018) 68-71. Math., Informa., Sci., Edu. Internat. Conf. (MISEIC 2018). https://download.atlantis-press.com/article/25905010.pdf

[1940] M. Khalid, S. T. R. Rizvi, and K. Ali, Note on cycle-(super)magic labelings of disconnected graphs, *Util. Math.,* **104** (2017) 315-320. arXiv:1506.06087

[1941] M. J. A. Khan, M. Ibrahim, and A. Ahmad, On edge irregular reflexive labeling of categorical product of two paths, *Comput. Syst. Sci. Eng.*,**36**(3)(2021) 485-492. https://www.researchgate.net/publication/348654478_On_Edge_Irregular_Reflexive_Labeling_of_Categorical_Product_of_Two_Paths

[1942] N. Khan, Cordial labelling of cycles, *Annals Pure Appl. Math.,* **1**(2) (2012) 117-130. doi:doi: 10.32604/csse.2021.014810

[1943] S. Khan, M. W. Akram, U. Ishtiaq, M. Garayev, and I.-L. Popa, Edge-irregular reflexive strength of non-planar graphs, *Symmetry* **17** (386) (2025) 1-16. https://www.mdpi.com/2073-8994/17/3/386

[1944] M. Khalid, X. Zhang, A. Atkinswestley, P. Sugapriya, M. R. Farahani, M. Alaeiyan, M. Cancan, S. Y. Lone, and A. Akgül, Innovative lucky-edge odd mean labeling of star-related graphs: bridging theory and circuit Design, *Bol. Soc. Paran. Mat. (3s)* **43** (2025) 1–19. https://periodicos.uem.br/ojs/index.php/BSocParanMat/article/view/76930/751375160670

[1945] S. Khatun and Sk. Md. Abu Nayeem, Graceful labeling of some zero divisor graphs, *Elect. Notes Discr. Math.,* **63** (2017) 189-196. https://www.sciencedirect.com/science/article/pii/S1571065317303153

[1946] J. Keene and A. Simoson, Balanced strands for asymmetric, edge-graceful spiders, *Ars Combin.*, **42** (1996) 49-64. https://combinatorialpress.com/article/ars/Volume%20042/volume-42-paper-3.pdf

[1947] M. Keerthika and V. Kowsalya, Total prime graceful labeling, *Internat. Confer. on Emerging Trends Math. Stat.*, Sri Ramakrishna College of Arts & Science, Tamil Nadu, India, (2024) 714-718. https://www.researchgate.net/publication/381324921_TOTAL_PRIME_GRACEFUL_LABELING

[1948] P. Keevash and K. Staden, Ringel's tree packing conjecture in quasirandom graphs, *J. Eur. Math. Soc.,* **27** (2025) 1769-1826. https://ems.press/content/serial-article-files/50361

[1949] A. Kézdy, ρ-valuations for some stunted trees, *Discrete Math.,* **306** (2006) 2786-2789. https://www.sciencedirect.com/science/article/pii/S0012365X06004183?via%3Dihub

[1950] R. Khennoufa and O. Togni, The radio antipodal and radio number of the hypercube, *Ars Combin.,* **102** (2011) 447-461. https://www.researchgate.net/publication/253321032_The_Radio_Antipodal_and_Radio_Numbers_of_the_Hypercube

[1951] R. Khennoufa and O. Togni, A note on radio antipodal colourings of paths, *Math. Bohem.,* **130** (2005) 277-282. http://o.togni.u-bourgogne.fr/KhennoufaAntipodal.pdf

[1952] A. Khodkar, S. Nolen, and J. Perconti, Super edge-graceful labelings of complete bipartite graphs, *Australas. J. Combin.,* **46** (2010) 241-261. https://ajc.maths.uq.edu.au/pdf/46/ajc_v46_p241.pdf

[1953] A. Khodkar, Super edge-graceful labelings of complete tripartite graphs, *J. Combin. Math. Combin. Comput.,* **76** (2011) 137-158. https://combinatorialpress.com/article/jcmcc/Volume%20076/vol-076-paper%209.pdf

[1954] A. Khodkar, The union of vertex disjoint 3-cycles is super edge-graceful, *Bull. Inst. Combin. Appl.,* **69** (2013) 43-46.

[1955]

[1956] A. Khodkar, R. Rasi, and S. M. Sheikholeslami, Super edge-gracefulness of complete graphs, *Util. Math.,* **87** (2012) 207-233.

[1957] A. Khodkar and K. Vinhage, Super edge-graceful labelings of total stars and total cycles, *J. Combin. Math. Combin. Comput.,* **76** (2011) 233-247. https://combinatorialpress.com/article/jcmcc/Volume%20076/vol-076-paper%2016.pdf

[1958] M. H. Khuluq, V. H. Krisnawati, and N. Hidayat, On group-vertex-magic labeling of simple graphs, *CAUCHY –J. Matematika Murni dan Aplikasi*, **8**(2) (2023) 167-174. https://www.semanticscholar.org/reader/a9a706d90b9a547a9c2393f0d807d41c24344aab

[1959] M. H. Khuluq, V. H. Krisnawati, and N. Hidayat, On Z_k-vertex-magic labeling of prime graph $PG(Z_n)$, *J. Alg. Related Topics*, **13**(2) (2025) 27-37. https://doi.org/10.22124/jart.2024.26022.1601

[1960] B. M. Kim, W. Hwang, and B. C. Song, Radio number for the product of a path and a complete graph, *J. Comb. Optim.,* **30**(1) (2015) 139-149.

[1961] R. Y. Kim, S-M. Lee, and H. K. Ng, On balancedness of some graph constructions, *J. Combin. Math. Combin. Comp.,* **66** (2008) 3-16. https://www.researchgate.net/publication/238617521_On_Balancedness_of_Some_Graph_Constructions

[1962] R. Y. Kim, S-M. Lee, and H. K. Ng, On balancedness of some families of graphs, unpublished.

[1963] S.-R. Kim and J. Y. Park, On super edge-magic graphs, *Ars Combin.,* **81** (2006) 113-127. https://www.researchgate.net/publication/220620148_On_Super_Edge-Magic_Graphs

[1964] Y.-H. Kim, On constructions of new super edge-magic graphs from some old ones by attaching some pendants, *Commun. Korean Math. Soc.,* **32**(1) (2017 225-231. https://koreascience.kr/article/JAKO201711656577502.pdf

[1965] J. S. Kimberley and J. A. MacDougall, Mutation of vertex-magic regular graphs, 24th MCCCC. *J. Combin. Math. Combin. Comput.,* **82** (2012) 157-177. https://combinatorialpress.com/article/jcmcc/Volume%20082/vol-082-paper%2011.pdf

[1966] W. W. Kirchherr, On the cordiality of some specific graphs, *Ars Combin.,* **31** (1991) 127-138. https://combinatorialpress.com/ars-articles/volume-031-ars-articles/on-the-cordiality-of-some-specific-graphs/

[1967] W. W. Kirchherr, NEPS operations on cordial graphs, *Discrete Math.,* **115** (1993) 201-209. https://www.sciencedirect.com/science/article/pii/0012365X9390489G

[1968] S. P. Kishore, Graceful Labellings of Certain Disconnected Graphs, Ph. D. Thesis, Indian Institute of Technology, Madras, 1996.

[1969] S. Klee, H. Lehmann, and A. Park, Prime labeling of families of trees with Gaussian integers, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 165-176. https://www.sciencedirect.com/science/article/pii/S0972860015300402

[1970] D. Knuth, personal communication (2021).

[1971] D. E. Knuth, The Art of Computer Programming Volume, 4B 2021.

[1972] A. N. Koam and A. Ahmad, Computation of total vertex irregularity strength of theta graphs, *IEESAccess,* **7** (2019) 113826-113831. https://ieeexplore.ieee.org/stamp/stamp.jsp?arnumber=8786105

[1973] H. Komarullah, K. Wijaya, N. Hidayat, and V. H. Krisnawati, On the minimum coprime number of some graphs,*Sci. Tech. Indonesia*, **11**(4) (2026) 1300-1310. https://sciencetechindonesia.com/index.php/jsti/article/view/2407

[1974] H. Komarullah, K. Wijaya, V. H. Krisnawati, and N. Hidayat, On prime and coprime edge labeling of some graphs, *J. Matematika UNAND¿ 15*(3) (2026) Hal. 291–301. https://jmua.fmipa.unand.ac.id/index.php/jmua/article/view/1658

[1975] K. M. Koh, M. Miller, W. F. Smyth, and Y. Wang, On optimum summable graphs, *AKCE Int. J. Graphs Comb.,* **3**(1) (2006) 45-57 https://core.ac.uk/download/pdf/77136167.pdf

[1976] K. M. Koh, L. Y. Phoon, and K. W. Soh, The gracefulness of the join of graphs, *Electron. Notes Discrete Math.,* **48** (2015) 57-64. https://www.sciencedirect.com/science/article/pii/S1571065315000104

[1977] K. M. Koh, L. Y. Phoon, and K. W. Soh, The gracefulness of the join of graphs (II), *AKCE Internat. J. Graphs Combin.,* **12** (2015) 180-185. https://www.sciencedirect.com/science/article/pii/S0972860015000407

[1978] K. M. Koh and N. Punnim, On graceful graphs: cycles with 3-consecutive chords, *Bull. Malaysian Math. Soc.,* **5** (1982) 49-63.

[1979] K. M. Koh, D. G. Rogers, P. Y. Lee, and C. W. Toh, On graceful graphs V: unions of graphs with one vertex in common, *Nanta Math.,* **12** (1979) 133-136.

[1980] K. M. Koh, D. G. Rogers, and C. K. Lim, On graceful graphs: sum of graphs, Research Report 78, College of Graduate Studies, Nanyang University (1979).

[1981] K. M. Koh, D. G. Rogers, and T. Tan, On graceful trees, *Nanta Math.*, **10** (1977) 27-31.

[1982] K. M. Koh, P. Y. Lee, and T. Tan, Fibonacci trees, *S*outheast Asian Bull. Math., **2**(1) (1978) 45-47.

[1983] K. M. Koh, D. G. Rogers, and T. Tan, A graceful arboretum: a survey of graceful trees, in *Proceed. Franco-Southeast Asian Confer.,* Singapore, May 1979, **2** 278-287.

[1984] K. M. Koh, D. G. Rogers, and T. Tan, Products of graceful trees, *Discrete Math.*, **31** (1980) 279-292. https://www.sciencedirect.com/science/article/pii/0012365X80901399

[1985] K. M. Koh, D. G. Rogers, and T. Tan Another class of graceful trees, *J. Austral. Math. Soc. Ser. A,* **31** (1981) 226-235. doi:10.1017/S1446788700033516

[1986] K. M. Koh, D. G. Rogers, H. K. Teo, and K. Y. Yap, Graceful graphs: some further results and problems, *Congr. Numer.*, **29** (1980) 559-571.

[1987] K. M. Koh, T. Tan, and D. R. Rogers, Interlaced trees: a class of graceful trees, *Combinatorial Mathematics, VI (Proc. Sixth Austral. Conf., Univ. New England, Armidale, 1978)*, 65-78, Lecture Notes in Math., **748** Springer, Berlin, 1979. https://link.springer.com/chapter/10.1007/BFb0102685

[1988] K. M. Koh, T. Tan, and D. G. Rogers, Two theorems on graceful trees, *Discrete Math.,* **25** (1979) 141-148. https://www.sciencedirect.com/science/article/pii/0012365X79900165

[1989] K. M. Koh and K. Y. Yap, Graceful numberings of cycles with a P_3-chord, *Bull. Inst. Math. Acad. Sinica,* **12** (1985) 41-48.

[1990] T. Kojima On C_4-supermagic labelings of the Cartesian product of paths and graphs, *Discrete Math.,* **313** (2013) 164-173. https://www.sciencedirect.com/science/article/pii/S0012365X12004001?via%3Dihub

[1991] S. R. Kola and P. Panigrahi, Radio numbers of some caterpillars, The Eighth International Workshop on Graph Labelings (IWOGL 2014) 289-296, *Electron. Notes Discrete Math.*, **48**, Elsevier Sci. B. V., Amsterdam, 2015. http://dx.doi.org/10.1016/j.endm.2015.05.043

[1992] S. R. Kola and P. Panigrahi, Nearly antipodal chromatic number $ac'(P_n)$ of the path P_n, *Math. Bohem.*, **134** (2009) 77-86. https://dml.cz/bitstream/handle/10338.dmlcz/140642/MathBohem_134-2009-1_8.pdf

[1993] K. Komalamba and P. Vijayakumar, Extended vertex odd mean labeling in certain graphs, *Bol. Soc. Paran. Mat.*, (3s) **43**(4) (2025) 1–11. https://www.jetir.org/papers/JETIRC006365.pdf

[1994] H. Komarullah, N. Hidayat, V. H. Krisnawati, and K. Wijaya, Prime and odd prime labelings of broom graphs and some related graphs, *Pan-Amer. J. Math.*, **5** (2026) 9 pp. https://www.researchgate.net/publication/404617052_Prime_and_Odd_Prime_Labelings_of_Broom_Graphs_and_Some_Related_Graphs

[1995] H. Komarullah, N. Hidayat, V. H. Krisnawati, and K. Wijaya, Prime and odd prime labelings on cycle-related graphs, *Sci. Tech. Indonesia*, **112**) (2026) 553-558. https://sciencetechindonesia.com/index.php/jsti/article/view/2232/645

[1996] M. Konečný, S. Kučera, J. Novotná, J. Pekárek, Š. Šimsa, and M. Töpfer, Minimal sum labeling of graphs, Combinatorial algorithms, 252-263, *Lecture Notes in Comput. Sci.*, 10765, Springer, Cham, 2018.

[1997] M. Kong, A. C. Lee, and S. M. Lee, On the balance index sets of the homeomorph of regular graphs, *Congr. Numer.,* **204** (2010) 193-203.

[1998] M. C. Kong, Y.-C. Wang, and S.-M. Lee. On edge-balanced index sets of some complete k-partite graphs. *Proceed. Fourteeth Southeastern Internat. Confer. Combin., Graph Th. and Comput.*, **196** (2009) 71-94.

[1999] M. Koppendrayer, personal communication (2012).

[2000] V. Kostyuk, D. Narayan, and V. Williams, Minimal rankings and the arank number of a path, *Discrete Math.,* **306** (2006) 1991-1996. https://repository.rit.edu/article/1438/

[2001] P. Kovář, K. Rozman, and P. Šparl, Self-reverse labelings of distance magic graphs, https://doi.org/10.1007/s40840-026-02042-4

[2002] V. Kowsalya and M. Keerthika, Network security utilizing total prime graceful lablings, *J. Dynamics Control*, **9**(10) (2025) 21-28. https://jodac.org/wp-content/uploads/2025/10/JoDaC-9-10-21-28.pdf

[2003] V. Kowsalya and M. Keerthika, Odd-even prime graceful labeling and k-odd-even prime graceful labeling approaches for diverse graph families, *Fuzzy Sys. Soft Comput.*, 100 **20**(02II) (2025)

[2004] W. O. Krawec, Modular multiplicative graphs, *Graph Theory Notes N. Y.,* **64** (2013) 45-48.

[2005] W. O. Krawec, Modular multiplicative graphs, *Ars Combin.,* **124** (2016) 33-40. https://combinatorialpress.com/article/ars/Volume%20124/volume-124-paper-3.pdf

[2006] H. K. Krishnappa, K. Kopthapalli, and V. C. Venkaiah, Vertex magic total of complete graphs, *AKCE J. Graphs Combin.,* **6** (2009) 143-154. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088881

[2007] H. K. Krishnappa, N. K. Srinath, and S. Manjunath, Vertex magic total labeling of complete graphs and their application for public-key cryptosystem, *Internat. J.Innovative Res. Comput. Commun. Engin.*, **1**(2) (2013). https://www.researchgate.net/publication/292971834_Vertex_Magic_Total_Labeling_of_Complete_Graphs_and_their_application_for_Public-Key_Cryptosystem

[2008] M. C. Kong, S. M. Lee, E. Seah, and A. Tang, A complete characterization of balanced graphs, *J. Combin. Math. and Combin. Comput.,* **66** (2008) 225-236.

[2009] M. C. Kong, S. M. Lee, and H. S. H. Sun, On magic strength of graph, *Ars Combin.,* **45** (1997) 193-200.
https://www.sciencedirect.com/science/article/pii/S0166218X13003028

[2010] D. Korze, Z. Shao, and A. Vesel, New results on radio k-labelings of distance graphs, *Discrete Appl. Math.*, **319** (2022) 472-479. https://www.sciencedirect.com/science/article/pii/S0166218X13003028

[2011] D. Kotlar, Distance magic labeling in complete 4-partite graphs, *Graphs Combin.,* **32** (2016) 1027-1038. https://doi.org/10.48550/arXiv.1410.6916

[2012] A. Kotzig, Decompositions of a complete graph into $4k$-gons (in Russian), *Matematický Casopis*, **15** (1965) 229-233.

[2013] A. Kotzig, On certain vertex valuations of finite graphs, *Util. Math.,* **4** (1973) 67-73.

[2014] A. Kotzig, β-valuations of quadratic graphs with isomorphic components, *Util. Math.*, **7** (1975) 263-279.

[2015] A. Kotzig, Decomposition of complete graphs into isomorphic cubes, *J. Combin. Theory, Series B,* **31** (1981) 292-296. https://www.sciencedirect.com/science/article/pii/0095895681900319?via%3Dihub

[2016] A. Kotzig, Recent results and open problems in graceful graphs, *Congr. Numer.,* **44** (1984) 197-219.

[2017] A. Kotzig and A. Rosa, Magic valuations of finite graphs, *Canad. Math. Bull.*, **13** (1970) 451-461. https://doi.org/10.4153/CMB-1970-084-1

[2018] A. Kotzig and A. Rosa, Magic valuations of complete graphs, Centre de Recherches Mathematiques, Universite de Montreal, (1972) CRM-175.

[2019] A. Kotzig and J. Turgeon, β-valuations of regular graphs with complete components, *Colloq. Math. Soc. János Bolyai 18, Combinatorics*, Keszthély, Hungary, 1976.

[2020] P. Kovář, Vertex magic total labeling of products of regular VMT graphs and regular supermagic graphs, *J. Combin. Math. Combin. Comput.,* **54** (2005) 21-31. https://combinatorialpress.com/article/jcmcc/Volume%20054/vol-054-paper%202.pdf

[2021] P. Kovář, Magic labelings of regular graphs, *AKCE Inter. J. Graphs and Combin.,* **4** (2007) 261-275. https://www.tandfonline.com/doi/epdf/10.1080/09728600.2007.12088841?needAccess=true

[2022] P. Kovář, Unified approach to magic labeling of copies of regular graphs, *Proceed. Thirty-Fifth Southeastern International Conference on Combinatorics, Graph Theory and Computing. Congr. Numer.,* **168** (2004) 197-205.

[2023] P. Kovář, D. Fronček, and T. Kovářová, A note on 4-regular distance magic graphs, *Australas. J. Combin.*, **54** (2012) 127-132. https://ajc.maths.uq.edu.au/pdf/54/ajc_v54_p127.pdf

[2024] P. Kovář, M. Kravčenko, M. Krbeček, and A. Silber, Supermagic graphs with many different degrees, *Discuss. Math. Graph Theory,* **41** (2021) 1041-1050. doi:10.7151/dmgt.2227

[2025] P. Kovár and A. Silber, Distance magic graphs of high regularity, *AKCE Int. J. Graphs Comb.,* **9**(2) (2012) 213-219. https://www.researchgate.net/publication/267676999_Distance_magic_graphs_of_high_regularity

[2026] P. Kováča, A. Silber, P. Kabelíkova—Hrušsková, and M. Kravčenko, On regular distance magic graphs of odd order, *J. Combin. Math. Combin. Comput.,* **117** (2023) 55-64. https://combinatorialpress.com/article/jcmcc/On-Regular-Distance-Magic-Graphs-of-Odd-Order.pdf

[2027] V. Kowsalya and M. Keerthika, Prime labeling of corona product of graphs, *Indian J. Natural Sci.,* **14**(82) (2024). https://www.researchgate.net/publication/378156933_Prime_Labeling_of_Corona_Product_of__Graphs

[2028] J. Kratochvil, M. Miller, and N. H. Minh, Sum graph labels-An upper bound and related problems. In: *Proceed. Twelfth Australasian Workshop Combin. Algor.,* Bandun, Indonesia (2001).

[2029] J. Kratochvil, M. Miller, and H. Nguyen, Sum graph labels–An upper bound and related problems, *Proc. of AWOCA 2001*, Institut Teknologi Bandung, Indonesia, (2001) 126-131.

[2030] K. C. Krishna, C. D. Raj, and K. R. Mary, Some standard results on even sum graphs, *Malaya J. Mat.,* **8**(04) (2020) 2282-2284. https://www.malayajournal.org/index.php/mjm/article/view/1899/1282

[2031] A. K. Krishnaa, A study of the major graph labelings of trees, *Informatica (Vilnius)* **15** (2004) 515-524. https://www.researchgate.net/publication/220073731_A_Study_of_the_Major_Graph_Labelings_of_Trees

[2032] A. K. Krishnaa, On antimagic labelings of some cycle related graphs, *J. Discrete Math. Sci. Cryptogr.,* **15** (2012) (4-5) 225-235. https://www.researchgate.net/publication/263110445_On_antimagic_labellings_of_some_cycle_related_graphs

[2033] A. K. Krishnaa, Formulas and algorithms of antimagic labelings of some helm related graphs, *J. Discrete Math. Sci. Cryptogr.,* **19**(2) (2016) 425-434. https://www.researchgate.net/publication/303982265_Formulas_and_algorithms_of_antimagic_labelings_of_some_helm_related_graphs

[2034] A. K. Krishnaa and M. S. Dulawat, Labelling paths, *J. Rajasthan Acad. Phy. Sci.,* **5** (2006) 99-104. https://www.researchgate.net/publication/368478154_Labelling_of_paths

[2035] A. K. Krishnaa, M. S. Dulawat, and G. S. Rathore, Computational complexity in decision problems, presented in Conf. of Raj. Parishad, Dec. 14-15, 2001, Udaipur, India.

[2036] V. H. Krisnawati, A. A. G. Ngurah, N. Hidayat, and A. R. Alghofari, On the super edge-magic deficiency of forests, December 2019AIP Conference Proceedings 2192(1):040007 https://aip.scitation.org/doi/10.1063/1.5139133

[2037] V. H. Krisnawati, A. A. G Ngurah, N. Hidayat, and A. R. Alghofari, On the (consecutively) super edge-magic deficiency of subdivision of double stars. *J. Math.,* (2020) Art. ID 4285238, 1-16. https://www.researchgate.net/publication/347543608_On_the_Consecutively_Super_Edge-Magic_Deficiency_of_Subdivision_of_Double_Stars

[2038] S. Krithika and K. E. Hilda, Sum divisor cordial labeling on some cycle and snake related graphs, *Confer. : Internat. Confere. Math. Models Comput. Intell.,* - ICMMCI 2023 At: Ethiraj College for women (Autonomous), Chennai, Tamil Nadu, India. https://www.researchgate.net/publication/376514102_Sum_Divisor_Cordial_Labeling_on_Some_Cycle_and_Snake_Related_Graphs

[2039] S. Krithika and R. Keerthana, Cube difference labeling on shell related graphs, *Proceed. Internat. Conf. Indian Contributions Math. Ancient & Modern* (ICICM 2022), **01** (2022) 68-73. https://www.researchgate.net/publication/377470609_Cube_Difference_Labeling_on_Shell_Related_Graphs

[2040] S. Krithika and R. Keerthana, Square difference and square difference prime labeling of $SSG(2)$, *Iternat. Conf. Applied Math. Comput.,*

(ICAMC2023) – Proceedings, Doss Vaishnav College, Chennai, Tamilnadu https://www.researchgate.net/publication/377471224_SQUARE_DIFFERENCE_AND_SQUARE_DIFFERENCE_PRIME_LABELING_OF

[2041] S. Krithika, E. Kowsalya, and N. Lavanya, Sum divisor cordial labeling of ice cream graph and circular fans with chords, *Internat. Confer. Math. Models Comput. Intell.*, - ICMMCI. https://www.researchgate.net/publication/376514302_Sum_Divisor_Cordial_Labeling_of_Ice_Cream_Graph_and_Circular_Fans_with_Chords

[2042] S. Krithika and E. H. Stanley, Sum divisor cordial labeling on shell related graphs, *Adv. Appl. Math. Sci.,* (2021) **21**(1) 205-212. https://www.researchgate.net/publication/374328416_Sum_Divisor_Cordial_Labeling_on_Shell_Related_Graphs

[2043] E. Krop, Lobsters with an almost perfect matching are graceful, *Bull. Inst. Combin. Appl.,* **74** (2015) 21-24. arXiv:1402.3994

[2044] E. Krop, S. M. Lee, and C. Raridan, On the number of unlabeled vertices in edge-friendly labelings of graphs, *Discrete Math.*, **312** (2012) 574-577. arXiv:1103.5127

[2045] E. Krop, S. M. Lee, and C. Raridan, On the edge-balanced index sets of product graphs, *J. Indones. Math. Soc.,* (2011) Special edition, 71-78. arXiv:1103.4994

[2046] E. Krop, S. Minion, P. Patel, and C. Raridan, A solution to the edge-balanced index set problem for complete odd bipartite graphs, *Bull. Inst. Combin. Appl.,* **70** (2014) 119-125. arXiv:1307.8071

[2047] E. Krop, F. Mutiso, and C. Raridan, On super edge-graceful trees of diameter four, *Congr. Numer.,* **210** (2011) 151-167 arXiv:1106.3063

[2048] E. Krop and K. Sikes, On the edge-balanced index sets of complete bipartite graphs, *Congr. Numer.,* **207** (2011) 23-32. arXiv:1106.1085

[2049] J. W. Krussel, Equitable labelling of complete bipartite graphs, unpublished.

[2050] Q. Kuang, S. M. Lee, J. Mitchem, and A-G. Wang, On edge-graceful unicyclic graphs, *Congr. Numer.,* **61** (1988) 65-74.

[2051] C. Kujur, Lucky labeling and proper lucky labeling for bloom graph, *IOSR J. Math.,* **13** (2017) 52-59. https://doi.org/10.9790/5728-1302025259 https://www.academia.edu/33809434/Lucky_Labeling_and_Proper_Lucky_Labeling_for_Bloom_Graph

[2052] C. Kujur, S. A. Xavier, and A. Raja, Proper lucky labeling of K-identified triangular mesh and K-identified Sierpinski Gasket graphs, *Internat. J. Sci. Engin. Res.,* (IJSER), **8**(4) (2017) 249-261. https://www.ijser.org/researchpaper/

Proper-Lucky-Labeling-of-k-Identified-Triangular-Mesh-and-k-Identified-Sierpinski-pdf

[2053] S. Kulandhai Therese and K. Romila, Cube root cube mean labeling of graphs, *Internat. J. Math. Trends Tech.,* (IJMTT) **65**(2) (2019) 130-135. http://ijmttjournal.org/Volume-65/Issue-2/IJMTT-V65I2P520.pdf

[2054] A. Kulkarni and J. Bagga, Graceful line graphs of nongraceful graphs: a windmill counterexample, preprint October 2026.

[2055] V. R. Kulli and M. H. Muddebihal, Lict and litact graph of graph, *J. Appl. Anal. Comput.*, **2**(1) (2006) 33-43. https://www.researchgate.net/publication/289117576_Lict_graph_and_litact_graph_of_a_graph

[2056] V. R. Kulli, The semitotal-block graph and the total-block graph of a graph, *Indian J. Pure Appl. Math.*, **7**(6) (1976) 625-630. https://www.researchgate.net/publication/268998873_The_semitotal-block_graph_and_the_total-block_graph_of_a_graph

[2057] D. S. Kumar and T. Gayathri, Edge odd graceful labeling of cartesian product of two paths, *Internat. J. Pure Appl. Math.,* **119**(14) (2018) 51-57. https://www.researchgate.net/publication/380375817_EDGE_ODD_GRACEFUL_LABELING_OF_CARTESIAN_PRODUCT_OF_TWO_PATHS

[2058] D. Kumar, V. Nirmala, and J. S. Lawrence, *E*-super vertex magic graceful labeling of graphs, *Palestine J. Math.,* **14** (Special Issue II)(2025) 174–179. https://pjm.ppu.edu/sites/default/files/papers/PJM_14%28Special%20Issue%20II%29_2025_174_to_179.pdf

[2059] D. Kumar, S. V. Kumar, and J. S. Lawrence, Even vertex in-magic total labeling of digraphs, *Proyecciones J. Math.,* **44**(1) (2025) 1-8. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/5866

[2060] A. Kumar, D. Mishra, A. Kumar, and V. Kumar, Alpha labeling of cyclic graphs, *Int. J. Appl. Comput. Math.,* **7**(4) (2021) Paper No. 151, 7 pp. https://doi.org/10.1007/s40819-021-01084-5

[2061] A. Kumar, D. Mishra, A. Verma, and V. K. Srivastava, Alpha labeling of cyclic graphs - I, *Ars Combin.,* **154** (2021) 257-263. https://link.springer.com/article/10.1007/s40819-021-01084-5

[2062] D. Kumar, V. Nirmala, and S. Mutharasu, *V*-Super vertex in-antimagic total graceful labelings of digraphs, *Math. Engin. Sci., Aerosoace, MESA*, **15**(1) (2024) 267-272. https://www.researchgate.net/publication/378547864_V_-Super_vertex_in-antimagic_total_graceful_labelings_of_digraphs

[2063] D. Kumar, E. Mohankumar, J. S. Lawrance, and M. Bhuvaneshwari, $H-E$-super magic graceful labelings of graphs,*Commu. Math. Appl.*, **16**(2) (2025) 641-648. https://www.rgnpublications.com/journals/index.php/cma/article/view/2854/1888

[2064] J. S. Kumar, Hilbert graceful labeling on complete multipartite graph, *Asian Res. Math.,* **20**(9) (2024) 132-139. https://journalarjom.com/index.php/ARJOM/article/view/836

[2065] S. S. Kumar and G. Marimuthu, H-E-super magic decomposition of complete bipartite graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 297-300. http://dx.doi.org/10.1016/j.endm.2015.05.044

[2066] H. N. Kumar, G. Sathiamoorthy, and S. Balachandran, Vertex antimagic total labeling of modified fan graphs, *Stat. Optim. Inform. Comput.*, **16** (2026) 3747–3752. https://www.researchgate.net/publication/414419646_Vertex_Antimagic_Total_Labeling_of_Modified_Fan_Graphs

[2067] R. Kuppan, L. Shobana and I. N. Cangul, Encrypting and decrypting algorithms using strong face graph of a tree, *Inter. J. Comp. Math.: Comp. Sys. Th.,* **5**(4) (2020) 225-233. https://www.researchgate.net/publication/343516675_Encrypting_and_Decrypting_Algorithms_Using_Strong_Face_Graph_of_a_Tree

[2068] R. Kuppan and L. Shobana, Double duplication of special classes of cycle related graphs, *Math. Stat.*, **10**(4)(2022) 773-781. https://www.hrpub.org/download/20220630/MS8-13427481.pdf

[2069] T. J. Rajesh Kumar, Total neighborhood prime labeling of some trees, *Proyecciones J. Math.,* **41**(1) (2022) 101-110. https://www.scielo.cl/pdf/proy/v41n1/0716-0917-proy-41-01-101.pdf

[2070] N. Kumari and S. Mehra, On vertex product cordial labeling, *Internat. J. Pure Appl. Math.*,**114**(3) (2017) 553-570. https://ijpam.eu/contents/2017-114-3/11/index.html

[2071] D. Kuo, G. Chang, and Y.-H. Kwong, Cordial labeling of mK_n, *Discrete Math.*, **169** (1997) 121-131. https://www.academia.edu/22024108/Cordial_labeling_of_mKn

[2072] S. Kuppusamy and S. Guruswamy, Graceful and cordial labeling of subdivision of graphs, *Electronic Notes Disc. Math.,* **53** (2016) 123-131. https://www.sciencedirect.com/science/article/pii/S1571065316300658

[2073] H. Kwong, On balance index sets of rooted trees, *Ars Combin.,* **91** (2009) 373-382. https://combinatorialpress.com/ars-articles/volume-091-ars-articles/on-balance-index-sets-of-rooted-trees/

[2074] H. Kwong, and S. M. Lee, On the integer-magic spectra of the corona of two graphs, 36th Southeastern International Conference on Combinatorics, Graph Theory, and Computing, *Congr. Numer.* **174** (2005) 207-222. https://www.researchgate.net/publication/262864029_On_the_integer-magic_spectra_of_the_corona_of_two_graphs

[2075] H. Kwong, and S. M. Lee, On friendly index sets of generalized books, *J. Combin. Math. Combin. Comput.,* **66** (2008) 43-58. https://www.researchgate.net/publication/262864452_On_friendly_index_sets_of_generalized_books

[2076] H. Kwong, and S. M. Lee, On edge-balance index sets of generalized theta graphs, *Cong. Numer.,* **198** (2009) 157-168. https://www.researchgate.net/publication/262840831_On_edge-balance_index_sets_of_generalized_theta_graphs

[2077] H. Kwong, and S. M. Lee, On edge-balance index sets of flower graphs, *Bull. Instit. Combin. Appl.,* **62** (2011) 107-116. https://www.academia.edu/22024114/On_edge_balance_index_sets_of_flower_graphs

[2078] H. Kwong, and S. M. Lee, S.-P. Lo, and Y.C. Wang, On uniformly balanced graphs, *Australasian J. Combin.,* **53** (2012) 83-95. https://ajc.maths.uq.edu.au/pdf/53/ajc_v53_p083.pdf

[2079] H. Kwong, S. M. Lee, and K. K. Ng, On friendly index set of 2-regular graphs, *Discrete Math.* **308** (2008) 5522-5532. https://www.academia.edu/99215874/On_friendly_index_sets_of_2_regular_graphs

[2080] H. Kwong, S. M. Lee, and H. K. Ng, On product-cordial index sets of cylinders, *Congr. Numer.,* **206** (2010) 139-150. https://www.researchgate.net/publication/262866447_On_product-cordial_index_sets_of_cylinders

[2081] H. Kwong, S. M. Lee, H. K. Ng, On product-cordial index sets and friendly index sets of 2-regular graphs and generalized wheels, *Acta Math. Sinica,*(Engl. Ser.) **28** (4) (2012) 661-674. https://link.springer.com/article/10.1007/s10114-011-0515-8

[2082] H. Kwong, S.-M. Lee, and H. K. Ng, On product-cordial index sets of Möbius ladders and their bridge join with K_4, *J. Graph Labeling*, **2**(1) (2016) 25-43.

[2083] H. Kwong, S-M. Lee, and D. G. Sarvate, On balance index sets of one-point unions of graphs, *J. Combin. Math. Combin. Comput.,* **66** (2008) 113-127. https://www.researchgate.net/publication/262864720_On_balance_index_sets_of_one-point_unions_of_graphs

[2084] H. Labane, I. Bouchemakh, and A. Semaničová-Feňovčíková, On H-irregularity strength of cycles and diamond graphs, *Disc. Math., Alg. Appl.*, **14**(5) (2022) 2150157. https://www.worldscientific.com/doi/10.1142/S1793830921501573

[2085] K. C. Lakshmi and K. Nagarajan, Geometric mean cordial labeling of graphs, *Intnat. J. Math.Soft Comput,*, **7**(1) (2017)75-87. https://oaji.net/articles/2017/296-1504971638.pdf

[2086] D. R. Lakshmi and S. Vangipuram, A note on the graceful numbering of a class of trees, *Def. Sci. J.,* **35** (1985) 65-70. https://core.ac.uk/download/pdf/333721467.pdf

[2087] M. Lakshmi and D. Bharathi, SDC labeling on path union and cycle of zero divisor graphs, *Internat. J. Math. Trends Tech.* (IJMTT), **71**(4) (2025) 8-20. https://ijmttjournal.org/public/assets/volume-71/issue-4/IJMTT-V71I4P102.pdf

[2088] N. Lakshmi, P. Sudhakar, and N. Sudhakar, An application of graph labeling cryptographic technique with magic labeling, In: J. R. Szymanski, C. K. Chanda, P. K. Mondal, and K. A. Khan, (eds,) Energy Systems, Drives and Automations. ESDA 2021. *Lecture Notes in Electrical Engineering*,**1057**. Springer, Singapore, August 2023.

[2089] P. Lalitha and L. Tamilselvi, Hexagonal snakes with alpha labeling along a path, *J. Physics Confer.,* Series 1504:012009 2020 https://www.researchgate.net/publication/341760424_Hexagonal_Snakes_With_Alpha_Labeling_Along_A_Path

[2090] J. D. Laison, C. Starr, and A. Walker, Finite prime distance graphs and 2-odd graphs, *Disc. Math.* **313**(20) (2013) 2281-2291. https://reader.elsevier.com/reader/sd/pii/S0012365X13002665?token=3B3684E8787CAE012B93F0D27F86B1474F6F0FBA7642A9F975200A47662F41CB576BDBC71704C71B7C&originRegion=us-east-1&originCreation=20220726202356

[2091] D. R. Lashmi and S. Vangipuram, An α-valuation of quadratic graph $Q(4, 4k)$, *Proc. Nat. Acad. Sci. India* Sec. A, **57** (1987) 576-580.

[2092] A. Lasim, I. Halikin and K. Wijaya, The harmonious, odd harmonious, and even harmonious labeling, *Math. App.*, **16**(4) (2022) 1131-1138. doi:10.30598/barekengvol16iss4pp1131-1138

[2093] R. Laskar and D. Pillone, Theoretical and complexity results for minimal rankings, Recent advances in interdisciplinary mathematics (Portland, ME, 1997). *J. Combin. Inform. System Sci.,* **25** (2000) 17-33.

[2094] V. Latchoumanane and M. Varadhan, Antimagicness of tensor product for some wheel related graphs with star, *Asian-European J. Math.,* **17**(10) (2024) 2450049. https://doi.org/10.48550/arXiv.2310.07114

[2095] G.-C. Lau, Every graph is local antimagic total and its applications, *Opuscula Math.*, **43**(6) (2023) 841–864. doi:10.7494/OpMath.2023.43.6.841

[2096] G.-C .Lau, S. Alikhani, S. M. Lee, and W. Kocay, On k-edge-magic labelings of maximal outerplanar graphs, *AKCE Int. J. Graphs Comb.,* **12**(1) (2015) 40-46. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2015.06.006?scroll=top&needAccess=true#d1e451

[2097] G.-C. Lau, H. H. Chu, N. Suhadak, F. Y. Foo, and H. K. Ng, On SD-prime cordial graphs, *Internat. J. Pure Appl. Math.,* (2016) 1017-1028. https://ijpam.eu/contents/2016-106-4/4/4.pdf

[2098] G.-C. Lau, Z.-B. Gao, S. M. Lee, and G.-Y. Sun, On friendly index sets of the edge-gluing of complete graph and cycles, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 107-111. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2016.06.004#d1e264

[2099] G.-C. Lau, P. Jeyanthi D. Ramya and R. Kalaiyarasi, On Skolem odd and even difference mean graphs, *J. King Saud Univer.– Science*, **30** (2018) 286-291. https://jksus.org/content/185/2018/30/2/pdf/10.1016_j.jksus.2017.09.018.pdf?utm_source=chatgpt.com

[2100] G.-C. Lau, J. M. Jeyaseeli, W. C. Shiu, and S. Arumugam, Sudoku number of graphs, *AKCE Int. J. Graphs Comb.*, **20**(2) (2023) 209-216. https://www.tandfonline.com/doi/full/10.1080/09728600.2023.2218917

[2101] G.-C. Lau, S. M. Lee, and W. C. Shiu, On edge-prime cubic graphs with small components, *Open J. Discret. Appl. Math.,* **2**(2) (2019) 48-58. https://www.researchgate.net/publication/335502722_On_edge-prime_cubic_graphs_with_small_components

[2102] G. C. Lau, J. Li, and W. C. Shiu, Approaches which output infinitely many graphs with small local antimagic chromatic number, *Discrete Math. Algorithms Appl.*, **15**(2), 2250079 (2023). https://www.researchgate.net/publication/344150155_Approaches_Which_Output_Infinitely_Many_Graphs_With_Small_Local_Antimagic_Chromatic_Number

[2103] G.-C. Lau and M. Nalliah, On local antimagic chromatic number of a corona product graph, *Bull. ICA*, **98** (2023) 111–121. https://bica.the-ica.org/Volumes/98/Reprints/BICA2022-35-Reprint.pdf?utm_source=chatgpt.com

[2104] G.-C. Lau, H.-K. Ng, and W.-C. Shiu, Affirmative solutions on local antimagic chromatic number, *Graphs and Combin.,* **36**(5) (2020) 1337-1354.

[2105] G. C. Lau, K. Premalatha, S. Arumugam, and W. C. Shiu, On local antimagic chromatic number of cycle-related join graphs II, *Discrete Math. Algorithms Appl.*, **16(3)** (2024) 2350022. https://doi.org/10.1142/S1793830923500222

[2106] G. C. Lau, K. Premalatha, W. C. Shiu, and M. Nalliah, On local antimagic chromatic numbers of circulant graphs join with null graphs or cycles, *Proyecciones (Antofagasta)*, **42(**5) (2023) 1307–1332, https://www.scielo.cl/pdf/proy/v42n5/0716-0917-proy-42-05-1307.pdf

[2107] G. C. Lau, W. C. Shiu and H. K. Ng, On local antimagic chromatic number of graphs with cut-vertices, *Iran. J. Math. Sci. Inform.*, **19(1)** (2024) 1-19. doi:DOI:10.61186/ijmsi.19.1.1

[2108] G. C. Lau, K. Schaffer, and W. C. Shiu, Every graph is local antimagic total and its applications, *Opuscula Math.*, **43**(6) (2023) 841-864. https://www.opuscula.agh.edu.pl/vol43/6/art/opuscula_math_4341.pdf

[2109] G.-C. Lau and W. C. Shiu, On local antimagic total labeling of amalgamation graphs, (2022) 16 pages. https://www.researchgate.net/publication/359228369_On_local_antimagic_total_labeling_of_amalgamation_graphs

[2110] G. C. Lau and W. C. Shiu, On local antimagic total labeling of complete graphs amalgamation, *Opuscula Mathematica*, **43(3)** (2023), 429-453. https://doi.org/10.7494/OpMath.2023.43.3.429

[2111] G.-C. Lau, W.-C. Shiu, K. Premalatha, and M. Nalliah, Constructions of local antimagic 3-colorable graphs of fixed odd size | matrix approach. https://doi.org/10.48550/arXiv.2403.16484

[2112] G.-C. Lau, W. C. Shiu, and H.-K. Ng, Further results on cube divisor cordial labeling, *AKCE Internat. J. Graphs and Combin.* **13** (2016) 200-209.

[2113] G.-C. Lau, W.-C. Shiu, and H.-K. Ng, On local antimagic chromatic number of cycle-related join graphs, *Discuss. Math. Graph Th.,* **41** (2021) 133-152. arXiv:1805.04888

[2114] G.-C. Lau, W. C. Shiu, H.-K.-Ng, and P. Jeyanthi, Further results on SD-prime labeling, *JCMMCC* **98** (2016) 151-170.

[2115] G.-C Lau, W.-C. Shiu, and H.-K. Ng, On number of pendants in local antimagic chromatic numbers, *J. Discrete Math. Sci. Cryptogr.*, (2021) arXiv:2001.05138.

[2116] G.-C Lau, W.-C. Shiu, and H.-K. Ng, On local antimagic chromatic number of graphs with cut-vertices, *Iran. J. Maths. Sci. Inform.*, **19**(1) (2024) 1-17. https://ijmsi.ir/article-1-1726-en.html

[2117] G.-C. Lau, W.-C. Shiu, and H.-K. Ng, On k-super graceful labeling of graphs, *Thai J. Math.*, **20**3 (2022) 1375-1387. http://thaijmath.in.cmu.ac.th/index.php/thaijmath/article/view/2624/354355122

[2118] G.-C. Lau, W.-C. Shiu, H.-K. Ng, Z. B. Gao and K. Schaffer, On k-super graceful graphs with extremal maximum vertex degree, *J. Discrete Math. Sci. Cryptogr.*, **27**(6) (2024) 1765-1783. https://tarupublications.com/journals/jdmsc/volume/27/issue/6/on-k-super-graceful-graphs-with-extremal-maximum-vertex-degree/pdf

[2119] G.-Ch. Laua, W. Ch. Shiub, M. Ch. Nalliahc, R. Zhangd, and K. Premalathae, Complete characterization of bridge graphs with local antimagic chromatic number 2, *Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Komp'yuternye Nauki*, **34**(3) (2024) 375–396. https://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=vuu&paperid=896&option_lang=eng

[2120] G. C. Lau, K. Premalatha, W. C. Shiu, and M. Nalliah, On local antimagic chromatic numbers of circulant graphs join with null graphs or cycles, *Proyecciones J. Math.*, **42**(5) (2023) 1307-1332. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/5834

[2121] G. C. Lau, W. C. Shiu and H. K. Ng, On number of pendants in local antimagic chromatic number, *J. Discrete Math. Sci. Cryptogr.*, **25**(8) (2022) 2673-2683. https://doi.org/10.1080/09720529.2021.1920190

[2122] G. C. Lau, W. C. Shiu, K. Premalatha, R. Zhang and M. Nalliah, On the local antimagic chromatic number of the lexicographic product of graphs, *Discrete Math. Lett.*, **11** (2023) 76-83. z doi:DOI:10.47443/dml.2022.149

[2123] G. C. Lau and W. C. Shiu, On join product and local antimagic chromatic number of regular graphs, *Acta Math. Hungar.*, **169**(1) (2023) 108-133. https://doi.org/10.1007/s10474-023-01298-7

[2124] G. C. Lau and W.C. Shiu, On local antimagic chromatic number of lexicographic product graphs, *Acta Math. Hungar.*, **169**(1) (2023) 158-170. https://link.springer.com/article/10.1007/s10474-023-01305-x

[2125] G. C. Lau, W. C. Shiu and C. X. Soo, On local antimagic chromatic number of spider graphs, *J. Discrete Math. Sci. Cryptogr.*, **26**(2) (2023) 303-339. https://doi.org/10.1080/09720529.2021.1892270

[2126] S. D. Laurence and K. M. Kathiresan, On super (a, d)-P_h-antimagic total labeling of stars, *AKCE Int. J. Graphs Comb.*, **12**(1) (2015) 54-58. https://www.researchgate.net/publication/283802053_On_super_a_d-Ph-antimagic_total_labeling_of_Star

[2127] S. D. Laurence and K. M. Kathiresan, The total edge irregular strength of path union of cycles, *Util. Math.,* **105** (2017) 125-131. https://www.researchgate.net/publication/328048050_The_Total_Edge_Irregular_strength_of_the_path_union_of_cycles

[2128] Q. Laurejas and A. Pedrano, On 0-edge magic labeling of some graphs, *Internat. J. Math. Appl.,* **5** Issue 2C (2017) 329-337. https://ijmaa.in/index.php/ijmaa/article/view/825/814

[2129] S. Lavanya and V. Ganesa, Different operations of sunlet graph S_n, *Malaya J. Mat.*, **9**(1) (2021) 272-282. http://malayajournal.org/articles/MJM09010046.pdf

[2130] Y. Lavanya, Dhanyashree, and K. N. Meera, Radio mean graceful graphs, *Internat. Confer. Applied Physics, J. Physics: Conf. series*, 1172 (2019) 012071. https://iopscience.iop.org/article/10.1088/1742-6596/1172/1/012071/pdf

[2131] H.-F. Law, Full friendly index sets of spiders, *Ars Combin.,* **119** (2015) 23-31. https://combinatorialpress.com/article/ars/Volume%20119/volume-119-paper-3.pdf

[2132] V. M. Lawas and Y. F. Lim, Stars are square harmonious graphs, *DLSU Research Congress 2017*, De La Salle University, Manila, Philippines, (2017) 1-6. https://www.researchgate.net/publication/359659568_Stars_are_Square_Harmonious_Graphs

[2133] A. N-T. Lee and S. M. Lee, On $(k, 1)$-strongly indexable graphs associated with sequences of positive integers, *Congr. Numer.,* **199** (2009) 205-215. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=f75f5f29397d93b7dd4074c1acce72f5f44a9bdb

[2134] A. N-T. Lee and S. M. Lee, On a construction of (k, d)-strongly indexable graphs, unpublished

[2135] A. N-T. Lee, S. M. Lee, and H. K. Ng, On balance index sets of graphs, *J. Combin. Math. Combin. Comput.,* **66** (2008) 135-150. https://www.academia.edu/4978178/On_The_Balance_Index_Sets_of_Graphs

[2136] A. C. Lee, S. M. Lee, and H.-H. Su, On the balance index sets of generalized friendship graphs, envelope graphs of cycles and cubic trees, *Proceed. Fortieth Southeastern Internat. Conf. Combin., Graph Th. Comput. Congr. Numer.,* **196** (2009) 3-22.

[2137] C. Lee, Minimum coprime graph labelings, *J. Integer Sequences*, **23** (2020) Article 20.11.4. https://cs.uwaterloo.ca/journals/JIS/VOL23/Lee/lee7.pdf?utm_source=chatgpt.com

[2138] H. Y. Lee, H. M. Lee, and G. J. Chang, Cordial labelings of graphs, *Chinese J. Math.,* **20** (1992) 263-273.

[2139] L. M. Lee, S. M. Lee, and G. Murthy, On edge-graceful labelings of complete graphs–solution of Lo's conjecture, *Congr. Numer.*, **62** (1988) 225-233.

[2140] M.-J. Lee, On super $(a, 1)$-edge-antimagic total labeling of grids and crowns, *Ars Combin.,* **104** (2012) 97-105. https://combinatorialpress.com/article/ars/Volume%20104/volume-104-paper-8.pdf

[2141] M.-J. Lee, W,-H. Tsai, and C. Lin, Super $(a, 1)$-cyclic-antimagic labelings of the grid, *Ars Combin.,* **112** (2013) 3-12. NOW Chat 10/2/26

[2142] M.-J. Lee, C. Lin, and W.-H. Tsai, On antimagic labeling for power of cycles, *Ars Combin.,* **98** (2011) 161-165.

[2143] M.-J. Lee, W.-H. Tsai, and C. Lin, On super $(a, 1)$-edge-antimagic total labelings of subdivision of stars, *Util. Math.,* **88** (2012) 355-365.

[2144] P-S Lee, On α-labelings of prism graphs and gear graphs, unpublished.

[2145] S. M. Lee, k-graceful labelling of Mongolian tents and related graphs, *Congr. Numer.*, **50** (1985) 85-96.

[2146] S. M. Lee, A conjecture on edge-graceful trees, *Scientia*, **3** (1989) 45-57.

[2147] S. M. Lee, New directions in the theory of edge-graceful graphs, *Proc. 6th Caribbean Conf. Combin. & Computing* (1991) 216-231.

[2148] S. M. Lee, On constructions which yield fully magic graphs, *Ars Combin.* **107** (2012) 385-409. https://combinatorialpress.com/article/ars/Volume%20107/volume-107-paper-31.pdf

[2149] S. M. Lee, E. Chen, E. Yera, and L. Wang, On super edge-graceful $(p, p+1)$-graphs, *Congr. Numer.,* **171** (2004) 51-65.

[2150] S. M. Lee, K-J. Chen, and Y-C. Wang, On the edge-graceful spectra of cycles with one chord and dumbbell graphs, *Congr. Numer.,* **170** (2004) 171-183.

[2151] S. M. Lee, K. Ho, and H.-H. Su, On balance index sets of k-level wheel graphs, *Congr. Numer.,* **229** (2017) 313-324.

[2152] S. M. Lee, Y. S. Ho, S. K. Tan, and H. H. Su, Edge-magic indices of stars, Proceedings of the Forty-Third Southeastern International Conference on Combinatorics, Graph Theory and Computing. *Congr. Numer.,* **213** (2012) 15-26.

[2153] S. M. Lee, M. Kitagaki, J. Young, and W. Kocay, On edge-graceful and edge-magic maximal outerplanar graphs, *J. Combin. Math. Combin. Comput.,* **59** (2006) 119-129. https://www.researchgate.net/publication/265584874_On_edge-graceful_and_edge-magic_maximal_outerplanar_graphs

[2154] S. M. Lee and M. C. Kong, On super edge-magic n-stars, *J. Combin. Math. Combin. Comput.* **42** (2002) 87-96. https://combinatorialpress.com/article/jcmcc/Volume%20042/vol-042-paper%205.pdf

[2155] S. M. Lee, M. Kong, and Y. C. Wang, On edge-balance index sets of some complete k-partite graphs, *Congr. Numer.*, **196** (2009) 71-94.

[2156] S. M. Lee, K. Y. Lai, Y. S. Wang, and M. K. Kiang, On the graceful permutation graphs conjecture, *Congr. Numer.*, **103** (1994) 193-201.

[2157] S. M. Lee and A. N-T Lee, On super edge-magic unicyclic graphs, unpublished.

[2158] S. M. Lee and A. N-T Lee, On super edge-magic graphs with many odd cycles, Proceed. Thirty-Fourth Southeastern Internat. Confer. Comb., Graph Th. Comput. *Congr. Numer.,* **163** (2003) 65-80.

[2159] S. M. Lee, A. Lee, H. Sun, and I. Wen, On the integer-magic spectra of graphs, *J. Combin. Math. Combin. Comput.,* **42** (2002) 77-86. https://combinatorialpress.com/article/jcmcc/Volume%20042/vol-042-paper%204.pdf

[2160] S. M. Lee and E. Leung, and H. K. Ng, On super vertex-graceful trees, Proceedings of the Thirty-Fifth Southeastern International Conference on Combinatorics, Graph Theory and Computing, *Congr. Numer.,* **167** (2004) 3-26.

[2161] S. M. Lee, E. Leung, and H. K. Ng, On super vertex-graceful unicylic graphs, *Czechoslovak Math. J.*, **59**(134) (2009) 1-22. https://www.academia.edu/8380644/On_super_vertex_graceful_unicyclic_graphs#loswp-work-container

[2162] S. M. Lee, C. Levesque, S-P. B. Lo, and K. Schaffer, On the edge-graceful spectra of the cylinder graphs (I), *J. Combin. Math. Combin. Comput.,* **66** (2008) 195-214. https://combinatorialpress.com/article/jcmcc/Volume%20066/vol-066-paper%2017.pdf

[2163] S. M. Lee and A. Liu, A construction of k-graceful graphs from complete bipartite graphs, *SEA Bull. Math.,* **12** (1988) 23-30.

[2164] S. M. Lee and A. Liu, A construction of cordial graphs from smaller cordial graphs, *Ars Combin.*, **32** (1991) 209-214. https://combinatorialpress.com/article/ars/Volume%20032/volume-32-paper-21.pdf

[2165] S. M. Lee, A. Liu, and S. K. Tan, On balanced graphs, Proceedings of the Twenty-first Manitoba Conference on Numerical Mathematics and Computing (Winnipeg, MB, 1991), *Congr. Numer.,* **87** (1992) 59-64.

[2166] S. M. Lee and S-P. Lo, On (1,2)-strongly indexable spiders, *J. Combin. Math. Combin. Comput.,* **72** (2010) 101-113. https://combinatorialpress.com/article/jcmcc/Volume%20072/vol-072-paper%207.pdf?utm_source=chatgpt.com

[2167] Sin-Min Lee, R. M. Low, H. K. Ng, and Y-C. Wang, On friendly index sets of k-galaxies, *Elect. J. Graph Theory and Appl.,* **7**(1) (2019) 1–10. DOI:10.5614/ejgta.2019.7.1.1

[2168] S. M. Lee, P. N. Ma, L. Valdés, and S.-M Tong, On the edge-graceful grids, *Congr. Numer.,* **154** (2002) 61-77.

[2169] S. M. Lee, T. Min-Fang, and S. P. B. Lo, On the edge-balance index set of some trees, unpublished.

[2170] S. M. Lee and K. C. Ng, Every Young tableau graph is d-graceful, *Combinatorial Math. Annal., New York Acad. Sci.,* **555** (1989) 296-302.

[2171] S. M. Lee and H. K. Ng, On friendly index sets of total graphs of trees, *Util. Math.,* **73** (2007) 81-95.

[2172] S. M. Lee and H. K. Ng, On friendly index sets of bipartite graphs, *Ars Combin.,* **86** (2008) 257-271. https://combinatorialpress.com/article/ars/Volume%20086/volume-86-paper-23.pdf

[2173] S. M. Lee and H. K. Ng, A class of k-graceful bipartite planar graphs, unpublished.

[2174] S. M. Lee and H. K. Ng, On friendly index sets of cycles with parallel chords, *Ars Combin.,* **97A** (2010) 253-267. https://combinatorialpress.com/article/ars/Volume%20097-A/volume-97-A-paper-19.pdf

[2175] S. M. Lee and H. K. Ng, On friendly index sets of graphs, unpublished.

[2176] S. M. Lee and H. K. Ng, On friendly index sets of prisms and Möbius ladders, *J. Combin. Math. Combin. Comput.,* **90** (2014) 59-74. https://combinatorialpress.com/article/jcmcc/Volume%20090/vol090-paper%205.pdf

[2177] S. M. Lee, H. K. Ng, Y.-S. Ho, and F. Saba, On edge-graceful edge-splitting extensions of paths and spiders, unpublished.

[2178] S. M. Lee, H. K. Ng, and G.-C. Lau, On friendly index sets of spiders, *Malays. J. Math. Sci.,* **8**(1) (2014) 47-68. https://scispace.com/pdf/on-friendly-index-sets-of-spiders-5a8zj3cuhf.pdf

[2179] S. M. Lee, H. K. Ng, and H. Sun, On super vertex-graceful caterpillars, Proceed. Forty-First Southeastern Inter. Conf. Combin., Graph Th. and Compu., *Congr. Numer.,* **204** (2010) 33-44.

[2180] S. M. Lee, H. K. Ng, and S. M. Tong, On the balance index of the chain-sum graphs of cycles, *Util. Math.,* **77** (2008) 113-123.

[2181] S. M. Lee, H. K. Ng, S. M. Tong, On friendly index sets of broken wheels with three spokes, *J. Combin. Math. Combin. Comput.,* **74** (2010) 13-31. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=bfdec084e4ac60437ab76cb0a2f510504462ed0e

[2182] S. M. Lee, H. K. Ng, and Y. Wen, On the edge-magic indices of $(v, v+1)$-graphs, *Util. Math.,* **72** (2007) 97-110.

[2183] S. M. Lee, Y. C. Pan, and M. C. Tsai, On vertex-graceful $(p, p+1)$-graphs, *Congr. Numer.,* **172** (2005) 65-78.

[2184] S. M. Lee, W. M. Pigg, and T. J. Cox, On edge-magic cubic graphs conjecture, *Congr. Numer.*, **105** (1994) 214-222.

[2185] S. M. Lee, L. Quach, and S. Wang, On Skolem-gracefulness of graphs which are disjoint union of paths and stars, *Congr. Numer.*, **61** (1988) 59-64.

[2186] S. M. Lee, F. Saba, E. Salehi, and H. Sun, On the V_4-magic graphs, *Congr. Numer.,* **156** (2002) 59-67.

[2187] S. M. Lee, F. Saba, and G. C. Sun, Magic strength of the kth power of paths, *Congr. Numer.*, **92** (1993) 177-184.

[2188] S. M. Lee and E. Salehi, Integer-magic spectra of amalgamations of stars and cycles, *Ars Combin.,* **67** (2003) 199-212. https://salehi.faculty.unlv.edu/pioneer/IMS-ASC.pdf

[2189] S. M. Lee, E. Salehi and H. Sun, Integer-magic spectra of trees with diameter at most four, *J. Combin. Math. Combin. Comput.,* **50** (2004) 3-15. https://www.salehi.faculty.unlv.edu/IMS-TDF.pdf

[2190] S. M. Lee, E. Schmeichel, and S.C. Shee, On felicitous graphs, *Discrete Math.,* **93** (1991) 201-209. https://www.sciencedirect.com/science/article/pii/0012365X91902562?via%3Dihub

[2191] S. M. Lee and E. Seah, On edge-gracefulness of kth power cycles, *Congr. Numer.*, **71** (1990) 237-242.

[2192] S. M. Lee and E. Seah, Edge-gracefulness labelings of regular complete K-partite graphs, *Congr. Numer.,* **75** (1990) 41-50.

[2193] S. M. Lee and E. Seah, On edge-gracefulness of composition of step graphs and null graphs, *Graph Theory, Combinatorics, Algorithms, and Applications (San Francisco, 1989)*, SIAM (1991) 325-330.

[2194] S. M. Lee and E. Seah, On edge-graceful triangular snakes and sunflower graphs, unpublished.

[2195] S. M. Lee, E. Seah, and S.-P. Lo, On edge-graceful 2-regular graphs, *J. Combin. Math. Combin. Comput.,* **12** (1992) 109-117.

[2196] S. M. Lee, E. Seah, and S. K. Tan, On edge-magic graphs, *Congr. Numer.,* **132** (1992) 179-191.

[2197] S. M. Lee, E. Seah, and P.-C. Wang, On edge-gracefulness of kth power graphs, *Bull. Inst. Math. Acad. Sinica,* **18** (1990) 1-11.

[2198] S. M. Lee, and Q. X. Shan, All trees with at most 17 vertices are super edge-magic, *16th MCCCC Conference*, Carbondale, University Southern Illinois, Nov. 2002.

[2199] S. M. Lee and S.C. Shee, On Skolem-graceful graphs, *Discrete Math.,* **93** (1991) 195-200. https://www.sciencedirect.com/science/article/pii/0012365X9190255Z

[2200] S. M. Lee, S. L. Song, and L. Valdeés, On $Q(a)P(b)$-super edge-graceful wheels, unpublished.

[2201] S. M. Lee and H.-H. Su, On the balance index sets of permutation graphs, unpublished.

[2202] S. M. Lee, H.-H. Su, and H. Todt, On the edge-balance index sets of broken wheels, *Congr. Numer.,* **221** (2014) 31-42.

[2203] S. M. Lee, H.-H. Su, and Y.-C. Wang, On the integer-magic spectra of honeycomb graphs, *Cong. Numer.,* **193** (2008) 49-65.

[2204] S. M. Lee, H.-H. Su, and Y.-C. Wang, On balance index sets of disjoint graphs, unpublished.

[2205] S. M. Lee, H.-H. Su, and Y.-C. Wang, On k-edge-magic Halin graphs, Proceed. Forty-First Southeastern Inter. Conf. Combin., Graph Th. Comput., *Congr. Numer.*, **204** (2010) 129-145.

[2206] S. M. Lee, H.-H. Su, and Y.-C. Wang, On the edge-balance index sets of $(p, p+1)$-graphs, *Int. J. Contemp. Math. Sci.,* **7** (2012) **29** 1429-1447. https://www.m-hikari.com/ijcms/ijcms-2012/29-32-2012/suhsinhaoIJCMS29-32-2012.pdf

[2207] S. M. Lee, H.-H. Su, and Y.-C. Wang, On k-edge-magic cubic graphs, 24th MCCCC. *J. Combin. Math. Combin. Comput.,* **82** (2012) 87-103. https://combinatorialpress.com/article/jcmcc/Volume%20082/vol-082-paper%206.pdf

[2208] S. M. Lee, H.-H. Su, and W. Wei, On a family of the super edge-graceful trees, *J. Combin. Math. Combin. Comput.,* **106** (2018) 23-36.

[2209] S. M. Lee, H. Sun, W. Wei, Y. Wen and P. Yiu, The super edge-gracefulness of two infinite families of trees, *Congr. Numer.,* **190** (2008) 109-128.

[2210] S. M. Lee, H. Sun, and I Wen, On group-magic graphs, *J. Combin. Math. Combin. Computing,* **38** (2001) 197-207. https://combinatorialpress.com/jcmcc-articles/volume-038/on-group-magic-graphs/

[2211] S. M. Lee and S. K. Tan, A class of arbitrarily graceful planar bipartite graphs, *J. Combin. Math. Combin. Comput.,* **9** (1991) 119-127. https://combinatorialpress.com/jcmcc-articles/volume-009/a-class-of-arbitrarily-graceful-planar-bipartite-graphs/

[2212] S. M. Lee, M.-F. Tao, and B. Lo, On the edge-balance index sets of some trees, unpublished.

[2213] S. M. Lee, S. M. Tong, and E. Seah, On the edge-magic and edge-graceful total graphs conjecture, *Congr. Numer.,* **141** (1999) 37-48.

[2214] S. M. Lee, L. Valdés, and Y. S. Ho, On group-magic trees, double trees and abbreviated double trees, *J. Combin. Math. Combin. Computing,* **46** (2003) 85-95. https://combinatorialpress.com/jcmcc-articles/volume-046/on-group-magic-trees-double-trees-and-abbreviated-double-trees/

[2215] S. M. Lee and G. Wang, All pyramids, lotuses and diamonds are k-graceful, *Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.),* **32** (1988) 145-150.

[2216] S. M. Lee and L. Wang, On k-edge-graceful trees, unpublished.

[2217] S. M. Lee and P. Wang, On the k-gracefulness of the sequential join of null graphs, *Congr. Numer.*, **71** (1990) 243-254.

[2218] S. M. Lee, J. Y-C Wang, On super edge-magicness of chain graphs whose blocks are complete, *Congr. Numer.* **162** (2003). https://www.academia.edu/3533157/On_Super_Edge_magicness_of_Chain_Graphs_whose_Blocks_are_Complete_Graphs

[2219] S. M. Lee, L. Wang, H. Ng, Y-C Wang, On the edge-graceful spectra of the corona of $(p, p+1)$-graphs with K_1, unpublished.

[2220] S. M. Lee, L. Wang, K. Nowak, and W. Wei, On the edge-graceful trees conjecture, *J. Combin. Math. Combin. Comput.,* **54** (2005) 83-98. https://combinatorialpress.com/article/jcmcc/Volume%20054/vol-054-paper%206.pdf

[2221] S. M. Lee, L. Wang, and Y. Wen, On the edge-magic cubic graphs and multigraphs, *Congr. Numer.,* **165** (2003) 145-160.

[2222] S. M. Lee, S. Wang, and I. Wui, On Skolem-gracefulness of 4-stars, *Congr. Numer.,* **62** (1988) 235-239.

[2223] S. M. Lee and W. Wei, On the super vertex-gracefulness of cartesian products of graphs, *Congr. Numer.,* **180** (2006) 175-192.

[2224] S. M. Lee, L. Wang, and E. R. Yera, On super edge-graceful Eulerian graphs, *Congr. Numer.,* **174** (2005) 83-96.

[2225] S. M. Lee, I. Wen, and H. Hugo, On group-magic graphs, *J. Combin. Math. Combin. Comput.* **38** (2001) 197-207. https://combinatorialpress.com/jcmcc-articles/volume-038/on-group-magic-graphs/

[2226] S. M. Lee and H. Wong, On the integer spectra of the power of paths, *J. Combin. Math. Combin. Comput.,* **42** (2002) 187-194. https://combinatorialpress.com/jcmcc-articles/volume-042/on-the-integer-magic-spectra-of-the-power-of-paths/

[2227] S. M. Lee and R. Wong, On $P(a)Q(1)$-super vertex-graceful unicyclic graphs, *Congr. Numer.,* **173** (2005) 79-96.

[2228] S. M. Lee, Y. S. Wong, and M. K. Kiang, On graceful permutations graphs conjecture, *Congr. Numer.,* **103** (1994) 193-201.

[2229] S. M. Lee and I. Wui, On Skolem-gracefulness of 2-stars and 3-stars, *Bull. Malaysian Math. Soc.*, **10** (1987) 15-20.

[2230] S. M. Lee, I. Wui and J. Yeh, On the amalgamation of prime graphs, *Bull. Malaysian Math. Soc. (Second Series),* **11** (1988) 59-67.

[2231] S. Leena, V. Sudhakar, P. Anuradha, B. Senthil, and V. Balaj, *Internat. J. Appl. Math.*, **38**(6s) (2025) 1019-1040. https://ijamjournal.org/ijam/publication/index.php/ijam/article/view/450/414

[2232] C. E. Leiserson, Area efficient graph layouts for VLSI, in: *Proc. 21st Ann. IEEE Symposium, FOCS*, (1980) 270-281. https://people.csail.mit.edu/cel/resume/papers/AreaEffGraphVLSI.pdf

[2233] S. Letzter, A. Pokrovskiy, and E. Williams, On the gracesize of trees, https://doi.org/10.48550/arXiv.2511.11331

[2234] C. H. Li, The Antimagic Labeling of Balanced Double Star Union cP_3, Master Thesis, National Chung Hsing University, Taiwan, 2019.

[2235] J. Li, G. C. Lau and W. C. Shiu, On modulo local antimagic chromatic number of graphs, *J. Discrete Math. Sci. Cryptogr.*, **25**(8) (2022) 2519-2533. https://doi.org/10.1080/09720529.2021.1880691

[2236] P. C. Li, Antimagic labelings of cycle powers, *Ars Combin.*, **124** (2016) 341-351. https://combinatorialpress.com/article/ars/Volume%20124/volume-124-paper-29.pdf

[2237] R. Li, The spum and sum-diameter of graphs: labelings of sum graphs, *Discrete Math.*, **345**(5) (2022), 112806. https://www.sciencedirect.com/science/article/pii/S0012365X22000127?utm_source=chatgpt.com

[2238] T. Li, Z-X. Song, G. Wang, D. Yang, and C-Q. Zhang, Antimagic orientations of even regular graphs, *J. Graph Theory,* **90** (2019) 46-53. https://www.researchgate.net/publication/318392393_Antimagic_orientations_of_even_regular_graphs

[2239] W.-T. Li, Antimagic labeling for subdivisions of graphs, *Discrete Appl. Math.,* **363**(4) 215-223. https://www.sciencedirect.com/science/article/abs/pii/S0166218X24005432?via%3Dihub

[2240] W.-T. Li and Y.-S. Wang, Labeling trees of small diameters with consecutive integers, *Taiwanese J. Math. Adv. Pub.,* (2022) 1-23. https://doi.org/10.11650/tjm/221103

[2241] W. T. Li and P. W. Yang, Constructing the antimagic labelings for double stars union paths on three vertices, *J. Comb. Optim.*, **51**(3) (2026). https://doi.org/10.1007/s10878-025-01377-1

[2242] W. Z. Li, G. H. Li, and Q. T. Yan, Study on some labelings on complete bipartite graphs, *Adv. Comput. Sci., Envir., Ecoinforma., and Ed., Comm. Comput. Inf. Sci.,* **214** (2011) 297-301.

[2243] X. Li, V. Mak, and S. Zhou, Optimal radio labellings of complete m-ary trees. *D*iscrete Appl. Math., **158**(5) (2010) 507-515. https://www.academia.edu/20201853/Optimal_radio_labellings_of_complete_ary_trees?uc-g-sw=69609859

[2244] H. X. Liang, and C. F. Liu, On k-gracefulness of graphs, *Dongbei Shida Xuebao*, **33** (1991) 41-44.

[2245] Y.-C. Liang, T.-L. Wong, and X. Zhu, Anti-magic labeling of trees, *Discrete Math.,* **331** (2014) 9-14. https://www.academia.edu/75115503/Anti_magic_labeling_of_trees

[2246] Y.-C. Liang and X. Zhu, Anti-magic labelling of Cartesian product of graphs, *Theoret. Comput. Sci.* **477** (2013) 1-5. https://www.sciencedirect.com/science/article/pii/S0304397512011383?via%3Dihub

[2247] Y.-C. Liang and X. Zhu, Antimagic labeling of cubic graphs, *J. Graph Th.,* **75** (2014) 31-36.

[2248] Z. Liang, The harmoniousness of book graph $St(4k+1) \times P_2$, *Southeast Asian Bull. Math.*, **21** (1997) 181-184.

[2249] Z. Liang, On the gracefulness of the graph $C_m \cup P_n$, *Ars Combin.,* **62** (2002) 273-280. https://combinatorialpress.com/article/ars/Volume%20062/volume-62-paper-26.pdf

[2250] Z. Liang, On the graceful conjecture of permutation graphs of paths, *Ars Combin.,* **91** (2009) 65-82. https://combinatorialpress.com/article/ars/Volume%20091/volume-91-paper-6.pdf

[2251] Z. Liang, Cycle-supermagic decompositions of complete multipartite graphs, *Discrete Math.,* **312**(22) (2012) 3342-3348. https://www.sciencedirect.com/science/article/pii/S0012365X12003470

[2252] Z. Liang, On the strongly c-harmoniousness of cycle with some chords, *Ars Combin.,* **109** (2013) 143-160. https://combinatorialpress.com/ars-articles/volume-109-ars-articles/on-the-strongly-c-harmoniousness-of-cycles-with-some-chords/

[2253] Z. -H. Liang, G-supermagic coverings of graphs, *Acta Math. Appl. Sin.,* **37**(5) (2014) 857-864.

[2254] Z. Liang and Z.-L. Bai, On the odd harmonious graphs with applications, *J. Appl. Math. Comput.,* (2009) **29** 105-116. https://link.springer.com/article/10.1007/s12190-008-0101-0

[2255] Z. Liang and Y. Miao, On Lee's conjecture, *Util. Math.,* **87** (2012) 305-329. https://www.researchgate.net/publication/260833501_On_a_conjecture_of_JM_Lee

[2256] Z. Liang, D. Q. Sun, and R. J. Xu, k-graceful labelings of the wheel graph W_{2k}, *J. Hebei Normal College*, **1** (1993) 33-44.

[2257] Z. Liang, H. Zhang, N. Xu, S. Ye, Y. Fan, and H. Ge, Gracefulness of five permutation graphs of paths. *Util. Math.,* **72** (2007) 241-249.

[2258] Z. Liang and H. Zuo, On the gracefulness of the graph $P_{2m,2n}$,*Appl. Anal. Discrete Math.,* **4** (2010) 175-180. https://scispace.com/pdf/on-the-gracefulness-of-the-graph-p-2m-2n-4xvpscvsu7.pdf

[2259] Q. Liao and W. Liu, The group vertex magicness of unicyclic graphs and bicyclic graphs, arXiv:2303.04588.pdf

[2260] S.-C. Liaw, D. Kuo, and G. Chang, Integral sum numbers of graphs, *Ars Combin.,* **54** (2000) 259-268. https://combinatorialpress.com/article/ars/Volume%20054/volume-54-paper-19.pdf

[2261] K. B., Libeeshkumar, Edge induced V_4-magic labeling of subdivision graphs, *Asian Res. J. Math.,* **21**(8) (2025) 119-128. https://www.journalarjom.com/index.php/ARJOM/article/view/976/2091

[2262] K. B. Libeeshkumar and V. Anil Kumar, Induced V_4-magic labeling of cycle related graphs, *Malaya J. Mat.*, **8**(2) (2020) 473-477. https://www.malayajournal.org/articles/MJM08020025.pdf

[2263] K. B. Libeeshkumar and V. Anil Kumar, Induced V_4-magic labeling of some subdivision graphs, *Nonlinear Stud.,* **28**(3) (2021) 761-768.

[2264] M. Licona and J. Tey, Gracefulness of two nested cycles: a first approach, *Disc. Appl. Math.,* **376** (2025) 72-87. https://doi.org/10.48550/arXiv.2411.12998

[2265] M. Licona and J. Tey, Gracefulness of two nested cycles: A first approach, *Discrete Appl. Math.*, **376** (2025) 72-87. https://arxiv.org/pdf/2411.12998

[2266] K.-W Lih, On magic and consecutive labelings of plane graphs, *Util. Math.,* **24** (1983) 165-197.

[2267] N. B. Limaye, k-equitable graphs, $k = 2, 3$, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 117-133.

[2268] C.-M. Lin and T-M. Wang, On zero magic sums of integer magic graphs, *Ars Combin.,* **118** (2015) 119-134. https://www.researchgate.net/publication/295558785_On_Zero_Magic_Sums_of_Integer_Magic_Graphs

[2269] Y. Lin, A. Ahmad, M. Miller, K. Sugeng, and M. Bača, Further results on d-antimagic labelings of antiprisms, *Proceed. 15-th Australasian Workshop Combin. Alg.,* (2004) 103-108, Ballina, Australia.

[2270] Y. Lin and M. Miller, Vertex-magic total labelings of complete graphs, *Bull. Inst. Combin. Appl.,* **33** (2001) 68-76.

[2271] Y. Lin, M. Miller, R. Simanjuntak, and Slamin, Magic and antimagic labelings of wheels, unpublished.

[2272] Y. Lin, Slamin, M. Bača, and M. Miller, On d-amtimagic labelings of prisms, *Ars Combin.,* **72** (2004) 65-76. https://combinatorialpress.com/article/ars/Volume%20072/volume-72-paper-6.pdf

[2273] Y. Lin and K. Sugeng, Face antimagic labelings of plane graphs p_a^b, *Ars Combin.,* **80** (2006) 259-273. https://combinatorialpress.com/article/ars/Volume%20080/volume-80-paper-24.pdf

[2274] Z. Lingqi, Z. Linna, R. Yuan, The generalized Petersen graph $P(n,3)$ is $(5n + 5/2, 2)$-antimagic, *Util. Math.,* **96** (2015) 17-25.

[2275] W. Lestari1, Dafik, Susanto, and A. A. Wahab, On the strong rainbow antimagic coloring of some special graph, *Proceed. 6th Internat. Conf. Comb., Graph Th., and Network Topology*, (ICCGANT 2022) https://www.atlantis-press.com/proceedings/iccgant-22/125986152

[2276] Y. Listiana, Darmaji, and Slamin, Vertex magic total lablings on sun di-graphs, *Proceed. Internat. Sem. Math. Edu. Graph Th.*, Unisma (2014) 583-589. https://www.researchgate.net/publication/370938854_Proceedings_of_International_Seminar_on_Mathematics_Education_and_Graph_Theory_-_Unisma_2014

[2277] A. Litersky, Graceful kayak paddles, M.S. Thesis, University of Minnesota Duluth, 2011.

[2278] B. Liu, Some new results on graceful graphs, *Math. Appl.,* **3** (1990) 108-110.

[2279] B. Liu, Sums of squares and labels of graphs, *Math. Practice Theory*, (1994) 25-29.

[2280] Z. Liu, J. Liu, and Y. Ji, The full m index sets of $P_2 \times P_n$, *J. Math.,* **1** (2023) 1-8. https://www.researchgate.net/publication/376690054_The_Full_m_Index_Sets_of_P_2_P_n

[2281] X. Liu, Y. Liu, B. Yao, Y. Ma and H. Lian, On odd-graceful labelings of irregular dragon graphs, 2014 *IEEE Internat. Confer. Progress Informa. Comput.*,, Shanghai, China, 2014, 415-418, https://ieeexplore.ieee.org/document/6972368

[2282] B. Liu and X. Zhang, On a conjecture of harmonious graphs, *Systems Science and Math. Sciences*, **4** (1989) 325-328.

[2283] B. Liu and X. Zhang, On harmonious labelings of graphs, *Ars Combin.*, **36** (1993) 315-326. https://combinatorialpress.com/article/ars/Volume%20036/volume-36-paper-34.pdf

[2284] D. Liu, Radio number for trees, *Discrete Math.,* **308** (2008) 1153-1164. https://www.calstatela.edu/sites/default/files/treenewdm.pdf

[2285] D. Liu, Multi-level distance labelings for trees, unpublished.

[2286] D. Liu, L. Saha, and S. Das, Improved lower bounds for the radio number of trees, *Theoretical Comp. Sci.,* **851**, (2020) doi:DOI: 10.1016/j.tcs.2020.05.023

[2287] D. Liu and M. Xie, Radio number for square of cycles, *Congr. Numer.,* **169** (2004) 105-125. https://www.researchgate.net/publication/255628456_Radio_Number_for_Square_of_Cycles

[2288] D. Liu and M. Xie, Radio number for square of paths, *Ars Combin.,* **90** (2009) 307-319. https://www.researchgate.net/publication/267666823_Radio_number_for_square_paths

[2289] D. Liu and X. Zhu, Multi-level distance labelings for paths and cycles, *SIAM J. Discrete Math.,* **19** (2005) 610-621. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=3ad49efc463458fc514a25fbdb091baa6fa09e64

[2290] J.-B. Liu, M. K. Aslam, M. Javaid, and A. Raheen, Computing edge-weight bounds of antimagic labeling on a class of trees, *IEEE Access*, **4** (2016) 1-11. https://scispace.com/pdf/computing-edge-weight-bounds-of-antimagic-labeling-on-a-5ew4ispckd.pdf

[2291] J. B. Liu, T. Zou, Y. Lu, Gracefulness of Cartesian product graphs, *Pure Appl. Math. (Xi'an),* **28** (2012)(3) 329-332, 362. NOW 7/17/25 or Chat 10/3/26

[2292] J. B. Liu, L. Wang, and Y. Lu, Odd graceful labeling and algorithm of bicyclic graphs with a common edge, *J. Hefei Univ. Technol. Nat. Sci.,* **35**(6) (2012) 857-859.

[2293] R. Y. Liu, On Bodendiek's conjecture for graceful graphs, *Chinese Quart. J. Math.,* **4** (1989) 67-73.

[2294] Y. Liu, The gracefulness of the star graph with top sides, *J. Sichuan Normal Univ.*, **18** (1995) 52-60.

[2295] Y. Liu, Proof of a conjecture concerning gracefulness of graphs, *Huaihua Shizhuan Xuebao*, **15** (1996) 13-25. NOW 7/17/25 or Chat 10/3/26

[2296] Y. Liu, Crowns graphs Q_{2n} are harmonious graphs, *Hunan Annals Math.*, **16** (1996) 125-128.

[2297] Y. Liu, All crowns and helms are harmonious, unpublished.

[2298] Z. S. Liu, A class of graceful graphs and two classes of graceful graphs derived from it, *Neimenggu Daxue Xuebao,* **16** (1985) 183-186.

[2299] Q. Liu and Z. Liu, A necessary and sufficient condition for a graph to be E_2-cordial, *Ars Combin.,* **123** (2015) 159-167. https://combinatorialpress.com/ars-articles/volume-123-ars-articles/a-necessary-and-sufficient-condition-for-a-graph-to-be-e_2-cordial/

[2300] Q. Liu, Z. Liu, and H. Wu, On the E_k-cordiality of some graphs, *Ars Combin.,* **120** (2015) 193-198. https://combinatorialpress.com/article/ars/Volume%20120/volume-120-paper-17.pdf

[2301] Z. Liu and B. Zhu, A necessary and sufficient condition for a 3-regular graph to be cordial, *Ars Combin.,* **84** (2007) 225-230. https://combinatorialpress.com/ars-articles/volume-084-ars-articles/a-necessary-and-sufficient-condition-for-a-3-regular-graph-to-be-cordial/

[2302] M. C. Liunokas, G. L. Putra, R. M. Pangaribuan, and I. G. Pasangka, Pelabelan anti elabelan, anti ajaib pada graf hasil kali sisir, *J. Matematika Unand*, **14**(3) (2025) 240-252. https://jmua.fmipa.unand.ac.id/index.php/jmua/article/view/1408/1020

[2303] A. Lladó and J. Moragas, Cycle-magic graphs, *Discrete Math.,* **307** (2007) 2925-2933. https://www.sciencedirect.com/science/article/pii/S0012365X07000957?via%3Dihub

[2304] M.-L. Lo and L. V. Alegria, Radio number for fourth power paths, *Involve* **9**(2) (2016) 317-332. https://projecteuclid.org/journals/involve-a-journal-of-mathematics/volume-9/issue-2/Radio-number-for-fourth-power-paths/10.2140/involve.2016.9.317.full

[2305] S. Lo, On edge-graceful labelings of graphs, *Congr. Numer.*, **50** (1985) 231-241.

[2306] S. Locke and A. Niedzialomski, $K_n \square P$ is radio graceful, *Matematiche (Catania)* **73** (2018) (1) 127-137. https://lematematiche.dmi.unict.it/index.php/lematematiche/article/view/1518

[2307] S. C. López and F. A. Muntaner-Batle, A new application of the $\otimes_h$-product to α-labelings, *Discrete Math.,* **338** (2015) 839-843. https://www.researchgate.net/publication/264424913_A_new_application_of_the_otimes_h-product_to_alpha-labelings

[2308] S. C. López and F. A. Muntaner-Batle, Langford sequences and a product of digraphs, *European J. Combin.,* **53** (2016) 86-95. https://www.sciencedirect.com/science/article/pii/S0195669815002504

[2309] S. C. López, F. A. Muntaner-Batle, and M. Prabu, A new labeling construction from the $\otimes_h$-product. *Discrete Math.,* **340**(8) (2017) 1903-1909. https://arxiv.org/pdf/1607.07313

[2310] S. C. López, F. A. Muntaner-Batle, and M. Prabu, On the super edge-magicness of graphs of equal order and size, *Ars Combin.,* **154** (2021) 31-44. https://arxiv.org/pdf/1706.00211.pdf

[2311] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Super edge-magic models, *Math. in Comput. Sci.,* **5** (2011) 63-68. https://repositori.udl.cat/server/api/core/bitstreams/b5e7567d-5711-4ab7-8197-8096d3809902/content

[2312] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Bi-magic and other generalizations of super edge-magic labeling, *Bull. Aust. Math. Soc.,* **84** (2011) 137-152. https://www.researchgate.net/publication/231767725_BI-magic_and_other_generalizations_of_super_edge-magic_labelings

[2313] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Open problems involving super edge-magic labelings and related topics, *Bull. Inst. Combin. Appl.,* **65** (2012) 43-56. https://mat.upc.edu/en/people/susana.clara.lopez/publications/openprob.pdf

[2314] S. C. López, F. A. Muntaner-Batle, M. Rius-Font, Perfect super edge-magic graphs, *Bull. Math. Soc. Sci. Math. Roumanie,* **55** (103)(2) (2012) 199-208. https://www.researchgate.net/publication/235922779_Perfect_super_edge-magic_graphs

[2315] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, On super edge-magic decomposable graphs, *Indian J. Pure Appl. Math.* **43**(5) (2012) 455-473. https://repositori.udl.cat/server/api/core/bitstreams/fc9ba01f-82e0-47b9-b355-f05ae0f83583/content

[2316] S. C. López, F. A. Muntaner-Batle and M. Rius-Font, New problems related to the valences of (super) edge-magic labelings, *AKCE Int. J. Graphs Combin.,* **10**(2) (2013) 169-181. https://repositori.udl.cat/server/api/core/bitstreams/bb1b16fe-e3e3-4ec9-b6c0-0a7f449921b1/content

[2317] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, A problem on edge-magic labelings of cycles, *Canadian Math. Bull.,* **57** (2014) 375-380. doi:10.4153/CMB-2013-036-1

[2318] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Labeling constructions using digraph products, *Discrete Appl. Math.,* **161** (2013) 3005-3016. https://www.sciencedirect.com/science/article/pii/S0166218X13002849?via%3Dihub

[2319] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Perfect edge-magic graphs, *Bull. Math. Soc. Sci. Math. Roumanie,* **57** (105)(1) (2014) 81-91. https://ssmr.ro/bulletin/pdf/57-1/articol_8.pdf?utm_source=chatgpt.com

[2320] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, The jumping knight and other (super) edge-magic constructions, *Mediterr. J. Math.,* **11**(2) (2014) 217-235. https://scispace.com/pdf/the-jumping-knight-and-other-super-edge-magic-constructions-3j1lvscn44.pdf?utm_source=chatgpt.com

[2321] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Magic coverings and the Kronecker product, *Util. Math.,* **95** (2014) 73-84. https://www.researchgate.net/publication/41799168_Magic_coverings_and_the_Kronecker_product

[2322] S. C. López, F. A. Muntaner-Batle, and M. Rius-Font, Enumerating super edge-magic labelings for some types of pathlike trees, *Utilitas Math.* **96** (2015) 1 -15. https://upcommons.upc.edu/server/api/core/bitstreams/4d3f8986-19d2-4a20-8de3-eb5a8bfdef9c/content

[2323] A. Lourdusamy and F. J. Beaula, Further results on group S_3 mean cordial labeling, *Sciencia Acta Xaveriana*, **13**(1) (2022) 83-94 https://www.researchgate.net/publication/362325517_Further_results_onGroup_S3_Mean_Cordial

[2324] A. Lourdusamy and F. J. Beaula, Total absolute difference edge irregularity strength of some families of graphs, *TWMS J. Appl. Engin. Math.*, **13**(3) (2023) 1005–1012. https://jaem.isikun.edu.tr/web/images/articles/vol.13.no.3/17.pdf

[2325] A. Lourdusamy and F. J. Beaula, Total absolute difference edge irregularity strength of some families of graphs, *TWMS J. Appl. Engin. Math.*, **13**(3) (2023) 1005–1012. https://jaem.isikun.edu.tr/web/images/articles/vol.13.no.3/17.pdf

[2326] A. Lourdusamy, F. J. Beaula, and F. Patrick, Total absolute difference edge irregularity strength of T_p-tree graphs,*Proyecciones* **42**(6) (2023) 1597–1614. https://ejurnal.ung.ac.id/index.php/Euler/article/view/39334/14020

[2327] A. Lourdusamy and S. George, Super vertex mean labeling of cycles through different ways, *Proyecciones,* **37**(2) (2018) 181-198. https://www.scielo.cl/pdf/proy/v37n2/0716-0917-proy-37-02-00181.pdf

[2328] A. Lourdusamy and S. George, Super vertex mean graphs of order ≤ 7, *J. Appl. Math. Inform.,* **35**(5-6) (2017) 565-586. https://koreascience.or.kr/article/JAKO201730048780377.page

[2329] A. Lourdusamy and S. George, Linear cyclic snakes as super vertex mean graphs, *Internat. J. Math. Sci. Comput.,* **1** (2018) 109-126. https://fs.unm.edu/IJMC/LinearCyclicSnakesAsSuperVertex.pdf

[2330] A. Lourdusamy, S. George and M. Seenivasan, An extension of mean labeling, preprint.

[2331] A. Lourdusamy, J. S. Mary and F. Patrick, Even vertex equitable even labeling, *Sciencia Acta Xaveriana,* **7** (2016) 37-46. https://www.researchgate.net/profile/Lourdusamy_Arockiam2/publication/303564979_Even_Vertex_Equitable_Even_Labeling/links/574876d908ae2e0dd30162ed/Even-Vertex-Equitable-Even-Labeling.pdf

[2332] A. Lourdusamy and F. Patrick, Sum divisor cordial graphs, *Proyecciones,* **35**(1) (2016) 119-136. https://www.scielo.cl/pdf/proy/v35n1/art08.pdf

[2333] A. Lourdusamy and F. Patrick, Even vertex equitable even labeling of path and bistar related graphs, *Informa. Sci. Comput.,* **2016** (2016) 1-9. https://www.researchgate.net/publication/313160627_Even_Vertex_Equitable_Even_Labeling_of_Path_and_Bistar_Related_Graphs

[2334] A. Lourdusamy, and F. Patrick, Sum divisor cordial labeling for star and ladder related graphs, *Proyecciones J. Math.*, textbf35(4) (2016) 437-455. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/1320/1032

[2335] A. Lourdusamy and F. Patrick, Vertex equitable labeling for ladder and snake related graphs, *J. Prime Research Math.,* **13** (2017) 1-7. https://www.researchgate.net/publication/321875188_Vertex_equitable_labeling_for_ladder_and_snake_related_graphs

[2336] A. Lourdusamy and F. Patrick, New construction on SD-prime cordial labeling, *Theoretical Computer Science and Discrete Mathematics*, 134-143, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017. https://www.researchgate.net/publication/323239523_On_SD-Prime_Cordial_Labeling

[2337] A. Lourdusamy and F. Patrick, Some results on SD-prime cordial labeling, *Proyecciones,* **36**(4) (2017) 601-614. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/2538/2140

[2338] A. Lourdusamy and F. Patrick, Even vertex equitable even labeling for cycle related graphs, Kragujevac J. Math. **43**(3) (2019) 427-441. https://imi.pmf.kg.ac.rs/kjm/pdf/accepted-finished/db52db9c7ff1618dd35125495d743075_2536_07152018_043122/kjm_43_3-7.pdf

[2339] A. Lourdusamy and F. Patrick, Sum divisor cordial labeling for path and cycle related graphs, *J. Prime Res. Math.*, **15** (2019) 101-114. https://jprm.sms.edu.pk/index.php/jprm/article/view/145/148

[2340] A. Lourdusamy and F. Patrick, Several results on sum divisor cordial graph, *Jordan J. Math. Stat.*, **15**(3b) (2022) 643-660. https://journals.yu.edu.jo/jjms/issues/vol15no32022pdf/16.PDF

[2341] A. Lourdusamy, J. Shobana Mary, and F. Patrick, Even vertex equitable even labeling for path related graphs, *Sciencia Acta Xaveriana*, **7**(1) (2018) 47-56. https://www.researchgate.net/publication/321884242_EVEN_VERTEX_EQUITABLE_EVEN_LABELING_FOR_PATH_RELATED_GRAPH

[2342] A. Lourdusamy and M. Seenivasan, Vertex equitable labeling of graphs, *J. Discrete Math. Sci, Crypt.,* **11** (2008) 727-735. https://connectjournals.com/file_full_text/122706H_jdmsc274_727-735.pdf

[2343] A. Lourdusamy and M. Seenivasan, Mean labeling of cyclic snakes, *AKCE Int. J. Graphs Combin.,* **8** (2011) 105-113. https://www.researchgate.net/publication/265831306_Mean_labelings_of_cyclic_snakes

[2344] A. Lourdusamy and M. Seenivasan, Vertex-mean graphs, *Internat. J. Math. Combin.,* **3** (2011) 114-120. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=bc9bf79eb801c1df7bef13c67fbd88683f35bc18

[2345] A. Lourdusamy and M. Seenivasan, Mean labeling of edge linked cyclic snakes, *AKCE Int. J. Graphs Combin.,* **10**(4) (2013) 391-403. https://www.researchgate.net/publication/265782446_Mean_labeling_of_edge_linked_cyclic_snakes

[2346] A. Lourdusamy, M. Seenivasan, S. George and R. Revathy, Super vertex-mean graphs, *Sciencia Acta Xaver.,* **5**(2) (2014) 39-46. https://www.researchgate.net/publication/303565010_Super_Vertex_Mean_Graph

[2347] A. Lourdusamy, E. Veronisha, and F. J. Beaula, Totally magic cordial labeling on zero divisor graphs, *Adv. Appl. Math. Sci.,* **21**(12) (2022) 6827-6842. https://www.mililink.com/upload/article/1890981335aams_vol_2112_october_2022_a18_p6827-6842_a._lourdusamy,_e._vernonisha_and_f._joy_beaula.pdf

[2348] A. Lourdusamy, S. J. Wency, and F. J. Beaula, SDC labeling on zero divisor graphs, *Sciencia Acta Xaveriana*, **12**(2) (2021) 11-19. https://www.researchgate.net/publication/362325187_SDC_Labeling_on_Zero_Divisor_Graphs

[2349] A. Lourdusamy, S. J. Wency, and F. Patrick, On SD-prime cordial labeling, *Internat. J. Pure Appl. Math.,* (2017) 221-228. https://www.researchgate.net/profile/Lourdusamy-Arockiam/publication/323239523_On_SD-Prime_Cordial_Labeling/links/5a87c4f8458515b8af8d6c69/On-SD-Prime-Cordial-Labeling.pdf

[2350] A. Lourdusamy, S. J. Wency, and F. Patrick, Sum divisor cordial labeling of T_p-tree related graphs, *Ars Combin.,* **157** (2023) 3-22. doi:DOI:10.61091/ars157-01 https://combinatorialpress.com/ars-articles/volume-157/sum-divisor-cordial-labeling-of-t_p-tree-related-graphs/

[2351] A. Lourdusamy, S. J. Wency, and F. Patrick, On SD-harmonious labeling, *Mapana J. Sci.*, **17**(3) (2018) 1–15. https://doi.org/10.12723/mjs.46.1

[2352] S. J. Lourdusamy, S. J. Wency, and F. Patrick, Even vertex equitable even labeling for snake related graphs, *Proyecciones,* **38**(1) (2019) 177-189. https://scielo.conicyt.cl/scielo.php?script=sci_arttext&pid=S0716-09172019000100177

[2353] A. Lourdusamy, S. J. Wency, and F. Patrick, Group S_3 cordial remainder labeling, *Internat. J. Recent Tech. Engin.*, **8**(4) (2019) 8276–8281.

[2354] A. Lourdusamy, S. J. Wency, and F. Patrick, Group S_3 cordial remainder labeling for subdivision of graphs, *J. Appl. Math. Informatics*, **38**(3-4) (2020) 221–238. https://koreascience.kr/article/JAKO202016151586225.pdf

[2355] A. Lourdusamy, S. J. Wency, and F. Patrick, Group S_3 cordial remainder labeling for wheel and snake related graphs, *Jordan J. Math. Stat.*, **14**(2) (2021) 267–286. https://journals.yu.edu.jo/jjms/Issues/Vol14No22021PDF/6.pdf

[2356] A. Lourdusamy, S. J. Wency, and F. Patrick, Group S_3 cordial remainder labeling for path and cycle related graphs, *J. Appl. Math. Informatics*, **39**(1-2) (2021) 223–237. https://koreascience.kr/article/JAKO202109950461292.pdf

[2357] A. Lourdusamy, S. J. Wency, and F. Patrick, Some results on group S_3 cordial remainder labeling, *Palest. J. Math.* **11**(2) (2022) 511–520. https://pjm.ppu.edu/sites/default/files/papers/PJM_May_2022_511_to_520.pdf

[2358] R. M. Low and S. M. Lee, On group-magic Eulerian graphs, *J. Combin. Math. Compin. Comput.,* **50** (2004) 141-148.

[2359] R. M. Low and S. M. Lee, On integer-magic Eulerian graphs, personal communication (2006).

[2360] R. M. Low and D. Roberts, Application of the combinatorial nullstellensatz to integer magic graph labelings, *Theory Appl. Graphs,* **9**(1) (2022) Article 3, 14 pages. digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1211&context=tag

[2361] R. M. Low, D. Roberts, and J. Zheng, The integer-antimagic spectra of graphs with a chord, *Theory Appl. Graphs*, **8**(1) (2021) Article 1. https://doi.org/10.20429/tag.2021.080101

[2362] R. M. Low and W. C. Shiu, On the integer-magic spectra of graphs, *Congr. Numer.,* **191** (2008) 193-203.

[2363] R. M. Low and L. Sue, Some new results on the integer-magic spectra of tessellation graphs, *Australas. J. Combin.,* **38** (2007) 255-266. https://ajc.maths.uq.edu.au/pdf/38/ajc_v38_p255.pdf

[2364] A. Lozano, M. Mora, C. Seara, and J. Tey, Caterpillars are antimagic, *Mediterr. J. Math.,* **18**(39) (2021). doi:10.1007/s00009-020-01688-z

[2365] A. Lozano, M. Mora, and C. Seara, Antimagic labeling of caterpillas, *Appl. Math. Comput.,* **347** (2017) 734-740. https://upcommons.upc.edu/bitstream/handle/2117/125781/AMC-D-18-01012R1-Antimagic+Labelings+of+Caterpillars-Nov-2018.pdf

[2366] A. Lozano, M. Mora, C. Seara, and J. Tey, Caterpillars are antimagic, *Mediterr. J. Math.,* (2021) 18:39 arXiv:1812.06715

[2367] A. Lozano, M. Mora, C. Seara, and J. Tey, Trees whose even-degree vertices induce a path are antimagic, arXiv:1905.06595 [math.CO] 7pp.

[2368] A. Lozano, M. Mora, C. Seara, and J. Tey, Trees whose even-degree vertices induce a path are antimagic, *Discuss. Math. Graph Th.* **42** (2022) 959–966. https://doi.org/10.48550/arXiv.1905.06595

[2369] H.-C. Lu, On the constructions of sequential graphs, *Taiwanese J. Math.,* **10** (2006) 1095-1107. https://www.researchgate.net/publication/239852722_On_the_constructions_of_sequential_graphs

[2370] H.-C. Lu, On large harmonious graph, *Ars Combin.,* **91** (2009) 447-458. https://combinatorialpress.com/article/ars/Volume%20091/volume-91-paper-41.pdf

[2371] X. Lu and X. F. Li, $P_1 \bigvee T_m$ graphs and a certification of its gracefulness, *Gongcheng Shuxue Xuebao*, **13** (1996) 109-113.

[2372] X. Lu, W. Pan, and X. Li, k-gracefulness and arithmetic of graph $St(m) \cup K_{p,q}$, *J. Jilin Univ.,* **42** (2004) 333-336.

[2373] K. J. Ma and C. J. Feng, About the Bodendiek's conjecture of graceful graph, *J. Math. Research and Exposition*, **4** (1984) 15-18.

[2374] K. J. Ma and C. J. Feng, On the gracefulness of gear graphs, *Math. Practice Theory*, (1984) 72-73.

[2375] X. D. Ma, Some classes of graceful graphs, *J. Xinjiang Univ. Nat. Sci.,* **3** (1986) 106-107.

[2376] X. Ma, A graceful numbering of a class of graphs, *J. Math. Res. and Exposition*, (1988) 215-216.

[2377] X. Ma, Y. Liu, and W. Liu, Graceful graphs: cycles with $(t-1)$ chords, *Math. Appl.*, **9** (1990), supp l., 6-8

[2378] J. A. MacDougall, M. Miller, Slamin, and W. D. Wallis, Vertex-magic total labelings of graphs, *Util. Math.,* **61** (2002) 3-21. https://www.researchgate.net/publication/235798060_Vertex-magic_total_labelings_of_graphs

[2379] J. A. MacDougall, M. Miller, and K. Sugeng, Super vertex-magic total labelings of graphs, *Proceedings Australasian Workshop Combin. Algorithm 2004*, Balina, NSW (2004) 222-229. https://www.researchgate.net/publication/277188358_Super_Vertex-magic_Total_Labelings_of_Graphs

[2380] J. A. MacDougall, M. Miller, and W. D. Wallis, Vertex-magic total labelings of wheels and related graphs, *Util. Math.,* **62** (2002) 175-183. https://www.researchgate.net/publication/263929935_Vertex-Magic_Total_Labelings_of_Wheels_and_Related_Graphs

[2381] J. A. MacDougall and W. D. Wallis, Strong edge-magic labeling of a cycle with a chord, *Australas. J. Combin.,* **28** (2003) 245-255. https://ajc.maths.uq.edu.au/pdf/28/ajc_v28_p245.pdf

[2382] J. A. MacDougall and W. D. Wallis, Strong edge-magic graphs of maximum size, *Discrete Math.,* **308** (2008) 2756-2763. https://www.sciencedirect.com/science/article/pii/S0012365X07004050?utm_source=chatgpt.com

[2383] L. Mader and S. Mitra, Exploring fibonacci cordiality in corona graphs, *Util. Math.*, **126** 277-296. https://doi.org/10.61091/um126-14

[2384] K. Mahalingam and H. P. Rajendran, m-Bonacci graceful labeling, *AKCE Int. J. Graphs Comb,* **18**(1) (2021) 7-15. https://www.tandfonline.com/doi/epub/10.1080/09728600.2021.1876505?needAccess=true

[2385] A. E. S. H. Maharani, M. U. Romdhini, Abdurahim, L. G. Dwikasari. and N. D. Lestari, Graceful labeling of the upstairs graph, Journal of Hunan University Natural Sciences, Vol 53, No 8 (2026) 165-175. doi:10.55463/issn.1674-2974.53.8.11

[2386] M. Maheo, Strongly graceful graphs, *Discrete Math.,***29** (1980) 39-46. https://core.ac.uk/download/pdf/82643761.pdf

[2387] M. Maheo and H. Thuillier, On d-graceful graphs, *Ars Combin.,* **13** (1982) 181-192.

[2388] A. U. Maheswari, S. Azhagarasi, and B. Samuel, Some labelings on cycle with parallel P_4 chord, *Turkish J. Comput. Math. Edu.*, **12**(10) (2021) 5500-5508. https://turcomat.org/index.php/turkbilmat/article/view/5355

[2389] A. U. Maheswari, S. Azhagarasi, and J. B. Samuvel, Vertex even mean and vertex odd mean labeling for path union and crown on cycle with parallel P_3 chords, *Design Engin.*, **6** (2021) 5775-5792. https://www.researchgate.net/publication/361103423_VERTEX_EVEN_MEAN_AND_VERTEX_ODD_MEAN_LABELING_FOR_PATH_UNION_AND_CROWN_ON_CYCLE_WITH_PARALLEL_P_3_CHORDS

[2390] A. U. Maheswari, S. Azhagarasi, and J. B. Samuvel, Square sum labelings on new families of cycle (C_n) with parallel P_3 chords, *Stochastic Modeling Appl.,* **26**(3) Special Issue 2022 Part - 4) 432-440, Special Issue on Recent Research on Management, Applied Sciences and Technology. https://www.researchgate.net/publication/361101964_SQUARE_SUM_LABELING_ON_NEW_FAMILIES_OF_CYCLE_Cn_WITH_PARALLEL_CHORDS

[2391] A. Maheswari and P. Jeyanth, Odd vertex equitable even labeling of duplication and product graphs, *Internat. J. Math. Combin.,* **1** (2022) 56-64. https://fs.unm.edu/IJMC/Odd_Vertex_Equitable_Even_Labeling_of_Duplication_and_Product_Graphs.pdf

[2392] G. U. Maheswari, M. J. Jebarani and V. Balaji, Coding techniques through Fibonacci webs, difference cordial labeling and GMJ code method, *J. Physics, Conf Series* 1139 (2018) 012077 https://www.researchgate.net/publication/359864620_Odd_Vertex_Equitable_Even_Labeling_of_Duplication_and_Product_Graphs

[2393] G. U. Maheswari, G. Margaret, J. Jebarani, and V. Balaji, Coding techniques through Fibonacci webs, difference cordial labeling and GMJ code method, *J. Physics: Confer. Ser.*, **1139** (2018), 012077.

[2394] G. M. Maheswari, S. J. Obaiys, N. A. B. Jamaludin, and N. A. Alwan, Coding and descriptions for the subset of twin star mean labeling, *Asia Pac. J. Math.,* (2024) 11:59. https://apjm.apacific.org/PDFs/11-59.pdf

[2395] A. U. Maheswari and V. Srividya, New families of square sum graphs, *J. Comput. Math. Sci.,* **10**(4) (2019) 907-914.

[2396] A. U. Maheswari and V. Srividya, Some labelings on cycles with parallel P_3 chords, *JASC: J. Appl. Sci. Comput.,* **VI** (I) (2019) 469-475. https://pdfs.semanticscholar.org/f1f2/bc5bc33c50a47e0c3bea04d6957c6536cf96.pdf?_ga=2.208011094.537058225.1571009368-2077171013.1571009368

[2397] G. Uma Maheswari, G. M. J. Jebarani and V. Balaji, On coding through a two star and super mean labeling, Springer - Applied Mathematics and Scientific Computing, **2** 2017). Trends in Mathematics, 469-478. doi:10.1007/978-3-030-01123-9_46

[2398] V. Maheswari, D. S. T. Ramesh, and V. Balaji, Relaxed mean labeling, *Internat. J. of Math. Res.*, **4**(3) (2012) 217-224.

[2399] M. Maheo, Strongly graceful graphs, *Discrete Math.,* **29** (1980) 39-46. https://www.sciencedirect.com/science/article/pii/0012365X9090285

[2400] V. Maheshwari, L. Vinothkumar and V. Balaji, Further results on mean cordial labeling for three star graph, *Proc. Jangjeon Math. Soc.*, **23** 2 (2020) 241-251. https://jangjeonopen.or.kr/public/upload/1589423526-pjms23-2-11-fin.pdf

[2401] G. U. Maheswari, P. Hariprabakaran, and V. Balaji, Coding techique with two and three stars, $SSML(V_4, S_2)$ and GMJ code, *J. Xi'an Shiyou Univee., Natural Science Ed.*, **16**(11) 31-37. https://www.xisdxjxsu.asia/V16I11-8.pdf

[2402] A. Mahmoody, A note on graceful graphs with large chromatic numbers, *Ars Combin.,* **90** (2009) 423-424. https://combinatorialpress.com/article/ars/Volume%20090/volume-90-paper-36.pdf a

[2403] M. K. Mahmood and S. Ali, A novel labeling algorithm on several classes of graphs, *Punjab Univ. J. Math.,* **49**(2) (2017) 23-35. https://www.researchgate.net/publication/330674081_A_Novel_Labeling_Algorithm_on_Several_Classes_of_Graphs

[2404] H. Mahmoudzadeh and K. Eshghi, Metaheuristic approach to the graceful labeling problem, *Int. J. Appl. Metaheuristic Comput.,* IJAMC **1** (2010) 42-57. http://vigir.missouri.edu/~gdesouza/Research/Conference_CDs/IEEE_SSCI_2007/Swarm%20Intelligence%20Symposium%20-%20SIS%202007/data/papers/SIS/S003P004.pdf

[2405] H. Mahmoudzadeh and K. Eshghi, A metaheuristic approach to the graceful labeling problem, Modeling, Analysis, and Applications in Metaheuristic Computing: Advancements and Trends. IGI Global, 2012, (2014) 217-232. doi:10.4018/978-1-4666-0270-0.ch013 https://pu.edu.pk/images/journal/maths/PDF/Paper-3_49_2_17.pdf

[2406] K. V. Madhumitha, A. Rao, T. P. Prabhu, S. Nayak, and S. D'Souza, Secure communication via C-exponential mean-labeled graphs: A novel message encryption scheme, *J. Discrete Math. Sci. Cryptography*, **29**(6) (2026) 2393–2408. https://www.tarupublications.com/doi/10.47974/JDMSC-2653

[2407] P. Majerski and J. Przybylo, Total vertex irregularity strength of dense graphs, *J. Graph Th.,* **76**(1) (2014) 34-41.

[2408] P. Majerski and J. Przybyo, On the irregularity strength of dense graphs, *SIAM J. Discrete Math.,* **28**(1) (2014) 1, 197-205. https://arxiv.org/pdf/2406.09584

[2409] H. M. Makadia, H. M. Karavadiya, and V. J. Kaneria, Some results on α-graceful graphs, *Internat. J. of Math. Combin.*, **2** (2017) 116-122. https://fs.unm.edu/IJMC/SomeResultsOnAlfaGracefulGraphs.pdf

[2410] H. M. Makadia, H. M. Karavadiya, and V. J. Kaneria, Graceful centers of graceful graphs and universal graceful graphs, *Proyecciones J. Math.,* **38** (2019)(2) 305-314. https://www.scielo.cl/pdf/proy/v38n2/0717-6279-proy-38-02-00305.pdf

[2411] K. Manickam and M. Marudai, Odd mean labelings of graphs, *Bull. Pure Appl. Sci.,* 25E **1** (2006) 149-153.

[2412] K. Manickam and M. Marudai, Edge magic labeling of graphs *Util. Math.,* **79** (2009) 181-187.

[2413] K. Manickam, M. Marudai, and R. Kala, Some results on felicitous labeling of graphs, *J. Combin. Math. Combin. Comput.,* **81** (2012) 273-279. https://combinatorialpress.com/article/jcmcc/Volume%20081/vol-081-paper%2023.pdf

[2414] F. A. Manico and A. C. Pedrano, On vertex balance index set of some snake graphs, *Adv. Appl. Discrete Math.,* **28**(2) (2021) 253-265. http://dx.doi.org/10.17654/DM028020253

[2415] A. Manonmani and R. Savithiri, Double quadrilateral snakes on K-odd sequential harmonious labeling of graphs, *Malaya J. Math.*, **3** (2015) 607–611. https://www.malayajournal.org/articles/paper19_2015.pdf

[2416] A. Manshath, P. Hariprabakaran, M. Veerasamy, and V. Balaji, GMJ coding using even felicitous labeling and bistar graphs, July 2021 *Journal of Physics Conference Series* 1964(2):022004. doi:10.1088/1742-6596/1964/2/022004

[2417] J. M. Manulang and K. A. Sugeng, Graceful labeling on torch graph, *Indon. J. Combin.,* **2**(1) (2018) 14-19. https://www.researchgate.net/publication/325876565_Graceful_labeling_on_torch_graph

[2418] H. T. Marbun and A. N. M. Salman, Wheel-supermagic labelings for a wheel k-multilevel corona with a cycle, *AKCE Int. J. Graphs Comb.*, **10**(2)(2013) 183-191. https://www.tandfonline.com/doi/abs/10.1080/09728600.2013.12088734

[2419] B. Mari and R. S. Jeyaraj, Radio labeling of supersub–division of path graphs, *IEEE* **11** PP(99):1-1 (2023) https://www.researchgate.net/publication/375033736_Radio_labeling_of_supersub-division_of_path_graphs

[2420] B. Mari and R. S. Jeyaraj, Further results on the radio number for some construction of the path, complete, and complete bipartite graphs, Heliyon 2024, doi:DOI:10.1016/j.heliyon.2024.e34434

[2421] G. T. Marimuthu and M. Balakrishnan, E-super vertex magic labelings of graphs, *Discrete Appl. Math.,* **160**(12) (2012) 1766-1774.

[2422] G. T. Marimuthu and M. Balakrishnan, Super edge magic graceful graphs, *Information Sciences*, **287**(10) (2014) 140-151.

[2423] G. T. Marimuthu, S. Kavitha, and M. Balakrishnan, Super edge magic graceful labeling of generalized Petersen graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 235-241. http://dx.doi.org/10.1016/j.endm.2015.05.035

[2424] G. T. Marimuthu and P. Krishnaveni, Super edge-antimagic graceful labeling of graphs, Malaya J. Mat. 3(3)(2015) 312–317. https://www.researchgate.net/publication/280729329_Super_Edge-antimagic_Graceful_labeling_of_Graphs

[2425] G. Marimuthu and P. Krishnaveni, Super edge-antimagic gracefulness of disconnected graphs, *Theoretical Comput. Sci. Disc. Math.*, 152-155, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017.

[2426] G. T. Marimuthu and M. Balakrishnan, E-super vertex magic labelings of graphs, *Discrete Appl. Math.,* **160**(12) (2012) 1766-1774.

[2427] G. T. Marimuthu and G. Kumar, On consecutive magic graphs, *Elect. Notes Discr. Math.*, **48** (2015) 213-215. http://dx.doi.org/10.1016/j.endm.2015.05.031

[2428] G. T. Marimuthu and G. Kumar, Solution to some open problems on E-super vertex magic labeling of disconnected graphs, *Appl. Math. Comput.,* **268** (2015) 657-663. https://ideas.repec.org/a/eee/apmaco/v268y2015icp657-663.html

[2429] G. T/ Marimuthu, B. Suganya, S. Kalaivani, and M. Balakrishnan, E-super vertex magic labelling of graphs and some open problems, *Appl. Appl. Math.,***10**(1) (2015) 536-543. https://www.pvamu.edu/sites/mathematics/journal/aam/2015/vol-10-issue-1/31_r722-marimuthu-vol.-10-issue-1-posted-06-20-15.pdf

[2430] G. T. Marimuthu and G. Kumar, V-super and E-super vertex-magic total labelings of graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 223-230. http://dx.doi.org/10.1016/j.endm.2015.05.033

[2431] G. T. Marimuthu and G. Kumar, Vertex N-magic total labeling of graphs, *J. Graph Label.*, **2**(2) (2016) 123-133. https://www.researchgate.net/publication/308158754_Vertex_N-_magic_total_labeling_of_graphs

[2432] G. T. Marimuthu and S. S. Kumar, H-V-super magic decomposition of complete bipartite graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 231-233. http://dx.doi.org/10.1016/j.endm.2015.05.034

[2433] G. T. Marimuthu, M. S. Raja Durga, and G. Durga Devi, Solution to some open problems on E-super vertex magic total labeling of graphs, *Appl.*

Appl. Math., **10**(2) (2015) 1104-1112. https://digitalcommons.pvamu.edu/cgi/viewcontent.cgi?article=1474&context=aam

[2434] G. T. Marimuthu, M. S. Raja Durga, and G. Durga Devi, V-Super vertex in-antimagic total labelings of digraphs, *J. Graph Labeling,* **1**(1) (2015) 21-30. https://www.researchgate.net/publication/378547864_V_-Super_vertex_in-antimagic_total_graceful_labelings_of_digraphs

[2435] R. Marinescu-Ghemeci, Radio number for some thron graphs, *Discuss. Math. Graph Theory,* **30** (2010) 210-222. https://eudml.org/doc/271026

[2436] Marsidi1, I. H. Agustin, Dafik, Md. S. Rahman, and L. H. Sullystiawati, On irregular reflexive k-labeling of graphs, *CAUCHY—-J. Mate Murni dan Aplikasi*, **8**(2) https://ejournal.uin-malang.ac.id/index.php/Math/article/viewFile/23753/pdf

[2437] T. S. Martini, M. Roswitha, and D. A. Lestari, Cycle-supermagic covering on grid graph and $K_1 + \overline{K_2}$. Proceed. 3rd Internat. Sympos. Current Prog. Math. Sci. 2017 (ISCPMS2017) *AIP Conf. Proc.,* (2023) 020197-1–020197-4; https://doi.org/10.1063/1.5064194

[2438] T. S. Martini, On reverse super edge bimagic labeling of broom graph and generalized butterfly graph, *Presented on Internat. Confer. Math. : Edu., Th. Appl.* 2022 (in *Progress Publishing in AIP Confer. Proceed.*), 2022. doi:DOI:10.1063/5.0262383

[2439] T. S. Martini, D. Indriati, and T. A. Kusmayadi, On reverse super edge bimagic labeling of gear graph, hibiscus graph and dove tail graph, *Internat. J. Comput. Sci. Appl. Math.,* **10**(2) (2024). https://iptek.its.ac.id/index.php/ijcsam/article/view/21944/8713

[2440] T. S. Martini and Y. Susanti, On reverse super edge trimagic labeling of dove tail related graphs, *AIP Conf. Proc.¿* 3201, 030009 (2024) https://pubs.aip.org/aip/acp/article/3201/1/030009/3320733/On-reverse-super-edge-trimagic-labeling-of-dove

[2441] A. M. Marr, Labelings of Directed Graphs, Ph.D. thesis, Southern Illinois University at Carbondale, 2007, 63 pages.

[2442] A. M. Marr, Graceful labelings of directed graphs, *J. Combin. Math. Combin. Comput.,* **66** (2008) 97-103. https://combinatorialpress.com/article/jcmcc/Volume%20066/vol-066-paper%208.pdf

[2443] A. M. Marr, N. C. K. Phillips, and W. D. Wallis, Bimagic labelings, *AKCE J. Graphs Combin.*, **6** (2009) 155-160. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088882?utm_source=chatgpt.com

[2444] A. M. Marr, S. Ochel, and B. Perez, In-magic total labelings of digraphs, *J. Graph Labeling,* **1**(2) (2015) 81-93. https://eudml.org/doc/270953

[2445] P. Martinez, J. Ortiz, M. Tomova, and C. Wyels, Radio numbers for generalized prism graphs, *Discuss. Math. Graph Theory* **31** (2011) 45-62. arXiv:1007.5346v1 [math.CO].

[2446] A. Marr and R. Simanjuntak, D-Magic oriented graphs, *Symmetry* **13** (2021) 2261 1-9. www.mdpi.com/2073-8994/13/12/2261

[2447] V. L. Stella Arputha Mary, S. Navaneetha Krishnan, and A. Nagarajan, Z_{4p}-magic labeling for some special graphs, *Internat. Math. Soft Comput.*, **3**(3)(2013) 61-70. https://oaji.net/articles/2017/296-1504259768.pdf

[2448] T. K. Maryati, E. T. Baskoro, and A. N. M. Salman, P_h-(super) magic labelings of some trees, *J. Combin. Math. Combin. Comput.,* **65** (2008) 197-204. https://www.researchgate.net/publication/260032961_Ph-supermagic_labelings_of_some_trees

[2449] T. K. Maryati, E. T. Baskoro, A. N. M. Salman, and Irawati, On the path-(super)magicness of a cycle with some pendants, *Util. Math.,* **96** (2015) 319-330.

[2450] T. K. Maryati and F. F. Hadiputra, Local edge antimagic chromatic number of join product of graphs, *Indonesian J. Combin.*, **9**(2) (2025) 69–77. https://www.ijc.or.id/index.php/ijc/article/view/248/177

[2451] T. K. Maryati, A. N. M. Salman, and E. T. Baskoro, and Irawati, The supermagicness of a disjoint union of isomorphic connected graphs, *Proceed. 4th IMT-GT Internat. Conf. Math. Stat. Appl.,* **3** (2008) 1-5. https://www.researchgate.net/publication/260032786_The_Supermagicness_of_A_Disjoint_Union_of_Isomorphic_Connected_Graphs

[2452] T. K. Maryati, A. N. M. Salman, and E. T. Baskoro, Supermagic coverings of the disjoint union of graphs and amalgamations, *Discrete Math.,* **313**(4) (2013) 397-405. https://www.sciencedirect.com/science/article/pii/S0012365X12004876

[2453] T. Maryati, A. N. M. Salman, E. T. Baskoro, J. Ryan, M. Miller, On H-supermagic labelings for certain shackles and amalgamations of a connected graph, *Util. Math.,* **83** (2010) 333-342. https://www.researchgate.net/publication/259974741_On_H-supermagic_labelings_for_certain_shackles_and_amalgamations_of_a_connected_graph

[2454] M. Matamala and J. Zamora, Graphs admitting antimagic labeling for arbitrary sets of positive integers, LAGOS'17—IX Latin and American Algorithms, Graphs and Optimization, *Electron. Notes Discrete Math.*, **62** (2017) 159-164.

[2455] M. Matamala and J. Zamora, Weighted antimagic labeling, *Discrete Appl. Math.,* **245** (2018) 194-201. https://repositorio.uchile.cl/bitstream/handle/2250/152934/Weighted-antimagic-labeling.pdf?sequence=1

[2456] M. Matamala and J. Zamora, Graphs admitting antimagic labeling for arbitrary sets of positive numbers, *Discrete Appl. Math.,* **281** (2020) 246-251. https://repositorio.uchile.cl/bitstream/handle/2250/175692/Graphs-admitting-antimagic-labeling.pdf?sequence=1&isAllowed=y

[2457] C. C. Marzuki, A. N. M. Salman, and M. Miller, On the total irregularity strength of cycles and paths, *Far East J. Math. Sci.,* **82**(1) (2013) 1-21.

[2458] U. Mato and T. Wichianpaisarn, The reflexive edge strength of cycles plus one edge, *WSEAS Trans. Math.,* (2024) 1-5. https://www.researchgate.net/publication/377310944_The_Reflexive_Edge_Strength_of_Cycles_Plus_One_Edge

[2459] M. Mavronicolas and L. Michael, A substitution theorem for graceful trees and its applications *Discrete Math.,* **309** (2009) 3757-3766. https://www.sciencedirect.com/science/article/pii/S0012365X08005992

[2460] O. Mbianda, Properly Even Harmonious Graphs, M. S. Thesis, University of Minnesota Duluth, 2016.

[2461] D. McQuillan, A technique for constructing magic labelings of 2-regular graphs, *J. Combin. Math. Combin. Comput.,* **75** (2010) 129-135. https://combinatorialpress.com/article/jcmcc/Volume%20075/vol-075-paper%2010.pdf

[2462] D. McQuillan, Edge-magic and vertex-magic total labelings of certain cycles, *Ars Combin.,* **90** (2009) 257-266. https://www.researchgate.net/publication/262260997_Edge-magic_and_Vertex-magic_Total_Labelings_of_Certain_Cycles

[2463] D. McQuillan and J. McQuillan, Magic labelings of triangles, *Discrete Math.,* **309** (2009) 2755-2762. https://www.sciencedirect.com/science/article/pii/S0012365X08004287?via%3Dihub

[2464] D. McQuillan and J. McQuillan, Strong vertex-magic and edge-magic labelings of 2-regular graphs of odd order using Kotzig completion, *Discrete Math.,* **341**(1) (2018) 194-202. https://www.sciencedirect.com/science/article/pii/S0012365X1730287X

[2465] D. McQuillan and K. Smith, Vertex-magic total labeling of odd complete graphs, *Discrete Math.,* **305** (2005) 240-249. https://www.sciencedirect.com/science/article/pii/S0012365X05005108

[2466] D. McQuillan and K. Smith, Vertex-magic total labeling of multiple complete graphs, *Congr. Numer.,* **180** (2006) 201-205. https://www.researchgate.net/publication/262014457_Vertex-magic_total_labeling_of_multiple_complete_graphs

[2467] J. McSorley and W. Wallis, On the spectra of totally magic labelings, *Bull. Inst. Combin. Appl.,* **37** (2003) 58-62. http://lagrange.math.siu.edu/mcsorley/papers/nspectrafin.pdf

[2468] J. McSorley and J. Trono, On k-minimum and m-minimum edge-magic injections of graphs, *Discrete Math.,* **310** (2010) 56-69. https://www.sciencedirect.com/science/article/pii/S0012365X09003884

[2469] K. McTavish, personal communication (circa 1990).

[2470] H R Medini, S. D'souza, D. Nayak , and P. G. Bhat, Encoding and decoding of messages using graph labeling and the complement of a graph, *Global Stochastic Analysis*, **1**(1) (2024) 1-13. https://www.mukpublications.com/resources/1.%20Dr.%20Sabitha%20D'Souza_pagenumber.pdf

[2471] A. Meena Kumari and S. Arockiaraj, On (1,1,0)-F-face magic mean labeling of some graphs, *Utilitas Math.,* **103** (2017) 139-159. https://iopscience.iop.org/article/10.1088/1742-6596/1139/1/012049/pdf

[2472] A. Meena Kumari and S. Arockiaraj, On (1,0,0)-F-face magic mean labeling of some graphs, *J. Physics: Conference Series*, **1139** (2018) 7 pp. https://iopscience.iop.org/article/10.1088/1742-6596/1139/1/012049/pdf

[2473] A. Meena Kumari and S. Arockiaraj, Further results on F-face magic mean graphs, *Songklanakarin J. Sci. Tech.,* **42**(6) (2020) 1397-1406.
bibitemMeNa14 S. Meena and J. Naveen, Some results on prime labeling of graphs, *Internat. J. Math. Comput. Res.*, **2**(9) (2014) 606-618. http://ijmcr.in/index.php/ijmcr/article/view/167/166

[2474] S. Meena and G. Gajalakshmi, Odd prime labeling of graphs related to circular ladder, *Comm. Math. Appl.,* **13**(4) (2022) 1307-1315. https://www.rgnpublications.com/journals/index.php/cma/article/view/2173/1484

[2475] S. Meena, and R. Mani, Root square mean labeling of some cycle related graphs, *Internat. J. Science Adv. Res. Tech.*, **5**(7) (2019) 786-789. https://ijsart.com/public/storage/paper/pdf/IJSARTV5I733090.pdf

[2476] 8. S. Meena, and R. Mani, Root square mean labeling of theta related graphs, *Math. Stat. Engin. Appl.*, **71**(4) (2022) 13049-13070. https://www.philstat.org/index.php/MSEA/article/view/2678/2127

[2477] S. Meena and M. Sivasakthi, Harmonic mean labeling Of subdivision graphs, *IJRAR- Internat. J. Res. Analytical Reviews,* **6**(1) (2019) 196-204. http://ijrar.com/upload_issue/ijrar_issue_20542941.pdf

[2478] S. Meena and S. Usharani, Product cordial labelling for some bicyclic graphs, *Commun. Math. Appl.,* 13(4) (2022) 1329-1336. doi:10.26713/cma.v13i4.2171

[2479] S. Meena and K. Vaithilingam, Prime labeling of friendship graphs, *Internat. J. Engin. Res. Tech.*, (IJERT) **1**(10) (2012). https://www.ijert.org/research/prime-labeling-of-friendship-graphs-IJERTV1IS10257.pdf

[2480] S. Meena and K. Vaithilingam, Prime labeling for some helm related graphs, *Internat. J. Innovative Res. in Sci., Engin. Tech.*, **2**(4) (2013) 10-75-1085. https://www.ijirset.com/upload/april/31_PRIME%20LABELING.pdf

[2481] D. E. W. Meganingtyas, Dafik, and Slamin, Super (a, d)-vertex antimagic total labelings on directed cycles, *Proceed. Internat. Sem. Math. Edu. Graph Th.*, Unisma (2014) 509-515. https://www.researchgate.net/publication/370938854_Proceedings_of_International_Seminar_on_Mathematics_Education_and_Graph_Theory_-_Unisma_2014

[2482] S. Mehra and Puneet, 2017, Topological integer additive set-labeling of signed graphs, *Int. J. Recent Sci. Res.* **8**(12) (2017) 22423-22427. https://recentscientific.com/sites/default/files/9355.pdf

[2483] A. Meissner and K. Zwierzyński, Vertex-magic total labeling of a graph by distributed constraint solving in the Mozart System, *in* Parallel Processing and Applied Mathematics, Lecture Notes in Computer Science, **3911** Springer Berlin/Heidelberg, 2006.

[2484] L. S. Melnikov and A. V. Pyatkin, Regular integral sum graphs, *Discrete Math.*, **252** (2002) 237-245. https://www.sciencedirect.com/science/article/pii/S0012365X01004265

[2485] E. Mendelsohn and N. Shalaby, Skolem labelled graphs, *Discrete Math.,* **97** (1991) 301-317. https://www.sciencedirect.com/science/article/pii/0012365X91904469

[2486] E. Mendelsohn and N. Shalaby, On Skolem labelling of windmills, *Ars Combin.*, **53** (1999) 161-172. https://combinatorialpress.com/article/ars/Volume%20053/volume-53-paper-12.pdf

[2487] L. Meng, Trees of odd order with at most two vertices of degree two are edge-graceful, https://arxiv.org/abs/2608.23881

[2488] Y. Miao and Z. Liang, On the strongly c-harmoniousness cycle with P_2 or P_3 chord, *Ars Combin.,* **102** (2011) 101-128. https://combinatorialpress.com/article/ars/Volume%20102/volume-102-paper-9.pdf

[2489] G. Michael and M. Z. Youssef, On prime self-complementary graphs, *J. Disc. Math. Sci. and Cryptography*, **17**(3) (2014) 239-256.

[2490] S. Miklavic and P. Ŝparl, On distance magic labelings of Hamming graphs and folded hypercubes, *Discuss. Math. Graph Th.*, (2021) 1-17. https://sciendo.com/de/article/10.7151/dmgt.2430

[2491] Ŝ. Miklavi and P. Ŝparl, On distance magic circulants of valency 6, *Discrete Appl. Math.,* **329** (2023), 35-48. https://www.sciencedirect.com/science/article/abs/pii/S0166218X22004875?via%3Dihub

[2492] M. Miller, Open problems in graph theory: labelings and extremal graphs, personal communication (2008).

[2493] M. Miller and M. Bača: Antimagic valuations of generalized Petersen graphs, *Australas. J. Combin.,* **22** (2000) 135-139. https://ajc.maths.uq.edu.au/pdf/22/ocr-ajc-v22-p135.pdf

[2494] M. Miller, M. Bača, and Y. Lin, On two conjectures concerning (a, d)-antimagic labelings of antiprisms, *J. Combin. Math. Combin. Comput.,* **37** (2001) 251-254. https://combinatorialpress.com/article/jcmcc/Volume%20037/vol-037-paper%2021.pdf

[2495] M. Miller, M. Bača, and J. A. MacDougall, Vertex-magic total labeling of generalized Petersen graphs and convex polytopes, *J. Combin. Math. Combin. Comput.,* **59** (2006) 89-99. https://combinatorialpress.com/article/jcmcc/Volume%20059/vol-059-paper%204.pdf

[2496] M. Miller, D. Patel, J. Ryan, K. Sugeng, Slamin, and M. Tuga, Exclusive sum labeling of graphs, *J. Combin. Math. Combin. Comput.,* **55** (2005) 137-148. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2013.pdf

[2497] M. Miller, O. Phanalasy, J. Ryan, All graphs have antimagic total labelings, *Electronic Notes Disc. Math.*, **38**(1) (2011) 645-650.

[2498] M. Miller, O. Phanalasy, J. Ryan, and L. Rylands, Antimagicness of some families of generalized graphs, *Australas. J. Combin.,* **53** (2012) 179-190. https://ajc.maths.uq.edu.au/pdf/53/ajc_v53_p179.pdf

[2499] M. Miller, O Phanalasy, J. Ryan, and L. Rylands, A note on antimagic labelings of trees, *Bull. Inst. Combin. Appl.,* **72** (2014) 94-100.

[2500] M. Miller, O. Phanalasy, J. Ryan, and L. Rylands, Sparse graphs with vertex antimagic edge labelings, *AKCE Int. J. Graphs Comb.*, **10**(2 (2013) 193-198. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088735

[2501] M. Miller, I. Rajasingh, D. A. Emilet, and D. A. Jemilet, d-Lucky labeling of graphs, *Procedia Comput. Sci.,* **57** (2015) 766-771. https://api.semanticscholar.org/CorpusID:61910700

[2502] M. Miller, J. Ryan, K. Sugeng, Slamin, and M. Tuga, Exclusive sum labeling of graphs, *J. Combin. Math. Combin. Comput.,* **55** (2005) 137-148. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2013.pdf

[2503] M. Miller, C. Rodger, and R. Simanjuntak, Distance magic labelings of graphs, *Australas. J. Combin.,* **28** (2003) 305-315. https://ajc.maths.uq.edu.au/pdf/28/ajc_v28_p305.pdf

[2504] M. Miller, J. Ryan, and Slamin, Integral sum numbers of cocktail party graphs and symmetric complete bipartite graphs, *Bull. Inst. Combin. Appl.,* **25** (1999) 23-28.

[2505] M. Miller, J. Ryan, Slamin, and W. Smyth, Labelling wheels for the minimum sum number, *J. Combin. Math. Combin. Comput.,* **28** (1998) 289-297. https://combinatorialpress.com/article/jcmcc/Volume%20028/vol-28-paper%2020.pdf

[2506] M. Miller, J. Ryan, Slamin, K. Sugeug, and M. Tuga, Open problems in exclusive sum graph labeling, unpublished.

[2507] M. Miller, J. Ryan, and W. Smyth, The sum number of the cocktail party graph, *Bull. Inst. Combin. Appl.*, **22** (1998) 79-90. https://researchportal.murdoch.edu.au/esploro/outputs/journalArticle/The-sum-number-of-the-cocktail/991005545338407891

[2508] K. G. Mirajkar and P. G. Sthavarmath, Edge-odd gracefulness of lict and litact graphs for few types of graphs, *AIP Confer. Proceed.,* 2385, 130039 (2022) https://www.researchgate.net/publication/357648727_Edge-odd_gracefulness_of_lict_and_litact_graphs_for_few_types_of_graphs_Edge-Odd_Gracefulness_of_Lict_and_Litact_Graphs_for_Few_Types_of_graphs

[2509] K. G. Mirajkar and P. G. Sthavarmath, Square sum and square difference labelings of semitotal-block graph for some class of graphs, *Ratio Mathematica*, **47** (2023) 101-125. https://www.researchgate.net/publication/367016888_Square_sum_and_square_difference_labelings_of_semitotal-block_graph_for_some_class_of_graphs

[2510] K. G. Mirajkar and P. G. Sthavarmath, Square product labeling for few types of cactus graphs, *J. Pharmaceutical Negative Results*, **13**(10) Special Issue (2022) 2735-2743. https://www.pnrjournal.com/index.php/home/article/view/9193/12812

[2511] K. G. Mirajkar and P. G. Sthavarmath, On square product labeling, *South East Asian J. Math. Math. Sci.*, unpublished.

[2512] D. Mishra and S. Bhattacharjee, Some new classes of lobsters exhibiting graceful labeling, *Global J. Pure Appl. Math.*, **11**(2) (2015) 909-930. https://www.ripublication.com/gjpam%202015/gjpamv11n2_31.pdf

[2513] D. Mishra and A. C. Panda, Some new graceful generalized classes of lobsters, *Internat. J. Pure Appl. Math.,* (IJPAM, **91**(1) (2014) 57-76. https://ijpam.eu/contents/2014-91-1/7/7.pdf

[2514] D. Mishra, A. C. Panda, and S. Bhattacharjee, Some new classes of graceful lobsters, *Internat. J. Scientific Res. Math. Stat. Sci.,* **8**(6) (2021) 30-39. https://www.isroset.org/pdf_paper_view.php?paper_id=2649&5-ISROSET-IJSRMSS-07036.pdf

[2515] D. Mishra, A. C. Panda, and R. B. Dash, A class of graceful lobsters with even number of branches incident on the central path, *Internat. J. Math. Sci. Appl.*, **1**(2) (2011) 463-472. http://ijmsa.yolasite.com/resources/5-may.pdf

[2516] D. Mishra, S. K. Rout, and P. C. Nayak, Some new graceful lobsters with pendant vertices with central paths, *Indian J. Sci. Tech.*, **10**(6) (2017) 1-9.. doi:10.17485/ijst/2017/v10i6/86164

[2517] D. Mishra and A. C. Panda, Some new family of graceful lobsters, *Adv. Appl. Discrete Math.,* **14** (2014)(1) 1-24.

[2518] D. Mishra and A. C. Panda, A class of diameter six graceful trees, *J. Advances Math.,* **9**(5) (2014) 2677- 2686. https://rajpub.com/index.php/jam/article/view/2346

[2519] D. Mishra and A. C. Panda, A class of diameter six trees with graceful labelings, *Inter. J. Math. Trends Tech.,* **9**(1) (2014) 1-11. doi:10.14445/22315373/IJMTT-V9P508

[2520] D. Mishra and A. C. Panda, A family of graceful diameter six trees generated by component moving techniques, *British J. Math. Comput. Sci.,* **21**(1) Article no.BJMCS.3107, (2017) 1-15. https://journaljamcs.com/index.php/JAMCS/article/view/268/536

[2521] D. Mishra and P. Panigrahi, Graceful lobsters obtained by component moving of diameter four trees, *Comput. Math. Appl.,* **50** (2005) 367-380. https://www.sciencedirect.com/science/article/pii/S0898122105002816?utm_source=chatgpt.com

[2522] D. Mishra and P. Panigrahi, Some graceful lobsters with all three types of branches incident on the vertices of the central path, *Comput. Math. Appl.* **56** (2008) 1382-1394. https://www.sciencedirect.com/science/article/pii/S0898122108001624?via%3Dihub

[2523] D. Mishra and P. Panigrahi, Some new classes of graceful lobsters obtained from diameter four trees, *Math. Bohem.,* **135** (2010) 257-278. https://dml.cz/bitstream/handle/10338.dmlcz/140703/MathBohem_135-2010-3_3.pdf

[2524] J. Mitchem and A. Simoson, On edge-graceful and super-edge-graceful graphs, *Ars Combin.*, **37** (1994) 97-111. https://combinatorialpress.com/article/ars/Volume%20037/volume-37-paper-8.pdf

[2525] S. Mitra and S. Bhoumik, Graceful labeling of triangular extension of complete bipartite graph, *Electron. J. Graph Theory Appl. (EJGTA),* **7**(1) (2019) 11-30. https://www.ejgta.org/index.php/ejgta/article/view/241/pdf_92

[2526] S. Mitra and S. Bhoumik, Fibonacci cordial labeling of some special families of graphs, *Annals Pure Appl. Math.,* **21**(2) (2020) 135-140. http://www.researchmathsci.org/apamart/APAM-v21n2-9.pdf

[2527] S. Mitra and S. Bhoumik, Tribonacci cordial labeling of graphs, *J. Appl. Math. Physics*, **10**(4) (2022) https://www.scirp.org/journal/paperinformation?paperid=117007

[2528] S. Mitra and S. Bhoumik, A study of fibonacci cordial labeling in structured graph families, *J. Combin. Math. Combin. Comput.*, **127** (2025) 245–261. https://doi.org/10.61091/jcmcc127-18

[2529] S. Mitra, A. Pritchard, and S. Bhoumik, On fibonacci cordial labeling of some planar graphs, *TWMS J. App. Eng. Math.,* **15**(5) (2025) 1153-1163. https://jaem.isikun.edu.tr/web/images/articles/vol.15.no.5/09.pdf

[2530] M. V. Modha and K. K. Kanani, Some new families of 5-cordial graphs, *Internat. J. Math. and Soft Computing,* **5**(1) (2015) 129-141. https://www.researchgate.net/publication/270898974_Some_New_Families_of_5-Cordial_Graphs

[2531] M. V. Modha and K. K. Kanani, k-cordial labeling of fan and double fan, *Internat. J. Applied Math. Res.*, **4**(2) (2015) 362-369. https://www.sciencepubco.com/index.php/ijamr/article/view/4530

[2532] M. V. Modha and K. K. Kanani, Some wheel related 7-cordial graphs, *Internat. J. Emerging Tech. Appl. Eng., Tech., Sci.,* **9** (2016) 71-78.

[2533] M. V. Modha and K. K. Kanani, k-cordiality of wheel, path related and cycle related graphs, *Internat. J. Math. Sci. Comput.,* **5**(2) (2015) 79-82. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/Paper-04-15.pdf

[2534] M. V. Modha and K. K. Kanani, k-cordial labeling of some prisms, *Internat. J. Math. Trends and Tech. (IJMTT),* **35**(3) (2016) 174-176. https://www.researchgate.net/publication/308576347_k-cordial_labeling_of_some_prisms

[2535] M. V. Modha and K. K. Kanani, On k-cordial labeling of some graphs, *British J. Math. Comput Sci.,* **13**(3) (2016) 1-7. https://journaljamcs.com/index.php/JAMCS/article/view/564/1130

[2536] M. V. Modha and K. K. Kanani, 5-cordiality of some path, cycle and wheel related graphs, 9th National Level Science Symposium. February 14, 2016, Organized by Christ College, Rajkot, Sponsored by GUJCOST, Gandhinagar, Mathematics & Statistics, **3** (2016) 183-187.

[2537] V. Mohan and A. Sugumaran, Some new divisor cordial graphs, *J. Emerging Tech. and Innovative Res.,* **5**(8)(2018) 229-238. https://ijmttjournal.org/Volume-12/number-2/IJMTT-V12P513.pdf

[2538] V. Mohan and A. Sugumaran, Difference cordial labeling of some graphs related to square graph, *J. Emerging Tech. and Innovative Res.*, **5**(11)(2018) 589-594. https://www.jetir.org/papers/JETIR1811077.pdf

[2539] V. Mohan and A. Sugumaran, Difference cordial labeling for plus and hanging pyramid graphs, *Internat. J. Res. Analytical Reviews*, **6**(1)(2019) 1106-1114.

[2540] M. Mohana priya and S. M. Santhiya, Lucky edge labeling for some graphs, *The Internat. J Analytical Experimental Modal Analysis,* **XII**(IX) (2020) 1224-1231. http://www.ijaema.com/gallery/146-ijaema-september-4595.pdf

[2541] M. Mollard and C. Payan, Elegant labelings and edge-colorings: A proof of two conjectures of Hartman, and Chang, Hsu, Rogers, *Ars Combin.,* **36** (1993) 97-106. https://combinatorialpress.com/article/ars/Volume%20036/volume-36-paper-9.pdf

[2542] M. Mollard, C. Payan, and S. Shixin, Graceful problems, *Seventh Hungarian Colloquium on Finite and Infinite Combinatorics*, Budapest July 1987.

[2543] K. M. Mominul Haque, X. Lin, Y. Yang, and J. Zhang, Prime cordial labeling of flower snark and related graphs, *Ars Combin.,* **105** (2012) 45-52. https://combinatorialpress.com/article/ars/Volume%20105/volume-105-paper-5.pdf

[2544] K. Monika and K. Murugan, Odd-even sum labeling in the context of duplication of graph elements, *Mapana J. of Sci.*, **17**(3) (2018) 17-28. https://doi.org/10.12723/mjs.46.2

[2545] M. Mora and J. Tey, Antimagic and product antimagic graphs with pendant edges, *Mediterr. J. Math.* (2026) **23** article 27. 20pp. https://doi.org/10.1007/s00009-025-03020-z

[2546] R. Montgomery, A. Pokrovskiy, and B. Sudakov, Embedding rainbow trees with applications to graph labelling and decomposition, *J. European Math. Soc.*, **22**(10) (2020) 3101–3132.

[2547] R. Montgomery, A. Pokrovskiy and B. Sudakov, A proof of Ringel's conjecture, *Geom. Funct. Anal.,* **31** (2021) 663-720. arXiv:2001.02665 37 pages. https://doi.org/10.1007/s00039-021-00576-2

[2548] C. P. Mooney, On Gracefully and Harmoniously Labeling Zero-Divisor Graphs, Rings, Monoids, and Module (Springer) **382** (2021) 239–260 (editor by A. Badawi and J. Coykendall).

[2549] D. Morgan, All lobsters with perfect matchings are graceful, *Electron. Notes Discrete Math.,* **11** (2002) 6 pp. https://www.researchgate.net/publication/275041986_All_lobsters_with_perfect_matchings_are_graceful

[2550] D. Morgan, Gracefully labeled trees from Skolem sequences, *Congr. Numer.,* **142** (2000) 41-48.

[2551] D. Morgan and R. Rees, Using Skolem and Hooked-Skolem sequences to generate graceful trees, *J. Combin. Math. Combin. Comput.,* **44** (2003) 47-63. https://combinatorialpress.com/article/jcmcc/Volume%20044/vol-044-paper%205.pdf

[2552] M. Morris-Rivera, M. Tomova, C. Wyels, A. Yeager, The radio number of $C_n \square C_n$, *Ars Combin.,* **120** (2015) 7-21.

[2553] M. H. Bani Mostafa A. and E. Ghorbani, Hamiltonicity of a coprime graph, *Graphs Combin.,* **37** (2021) 2387-2395. arXiv:2005.06308.pdf

[2554] M. I. Moussa and E. M. Badr, Odd graceful labelings of crown graphs, *1st Internat. Conf. Comp. Sci. Algor. Appl.,* **2009** 1-5. Cairo, Egypt. https://www.researchgate.net/publication/215648009_Odd_Graceful_Labelings_of_Crown_Graph

[2555] M. I. Moussa, Some simple algorithms for some odd graceful labeling graphs, *Proceed. 9th WSEAS Internat. Conf. Applied Informatics and Communications (AIC '09)* August, 2009, Moscow, Russia. 399-404. https://www.wseas.us/e-library/conferences/2009/moscow/AIC/AIC63.pdf

[2556] M. I. Moussa, An algorithm for odd graceful labeling of the union of paths and cycles,*Internat. J. Appl. Graph Theory in Wireless ad hoc Networks (Graph Hoc)*, **2**(1) (2010) 112-119. https://www.researchgate.net/publication/

215648040_An_Algorithm_for_Odd_Graceful_Labeling_of_the_Union_of_Paths_and_Cycles

[2557] M. I. Moussa, An algorithm for odd gracefulness of the tensor product of two line graphs, *Inter. J. Applications of Graph Theory in Wireless ad hoc Networks and Sensor Networks (GRAPH-HOC),* **3** (2011). https://www.researchgate.net/publication/215648091_An_Algorithm_for_Odd_Gracefulness_of_the_Tensor_Product_of_Two_Line_Graphs

[2558] M. I. Moussa and E. M. Badr, Ladder and subdivision of ladder graphs with pendant edges are odd graceful, *Internat. J. Appl. Graph Theory Wireless Ad hoc Networks and Sensor Networks,* (GRAPH-HOC) textbf8(1) (2016) 1-8. arXiv:1604.02347.pdf

[2559] D. Moulton, Graceful labelings of triangular snakes, *Ars Combin.*, **28** (1989) 3-13.

[2560] H. Mukherjee and R. Pawar, Congruence distance magic graphs, https://arxiv.org/pdf/2302.05652.pdf

[2561] Z. Mujtahid, M. Salman, A. T. Hidayat, H. Qausar, and Y. Zahara, How to construct super edge-magic total labeling of theta graph $\theta(2, b, c)$, *Indonesian J. Math. Edu.,* **7**(1) (2024) 13-19. https://journal.untidar.ac.id/index.php/ijome/article/view/990/706

[2562] A. Mughal and N. Jamil, Total face irregularity strength of grid and wheel graph under k-labeling of type $(1, 1, 0)$, *J. Math.,* (2021) Art. ID 1311269, 16 pp. https://www.hindawi.com/journals/jmath/2021/1311269/#conflicts-of-interest

[2563] A. A. Mughal, R. N. Jamil, A. R. Virk, Characterizing irregularity in planar graph structures, *J. Prime Res. Math.,* **19**(2) (2023) 72-95. https://jprm.sms.edu.pk/media/pdf/jprm/volume_19/issue-2/Characterizing-Irregularity-in-Planar-Graph-Structures.pdf

[2564] J. Mülbacher, Magische Quadrate und ihre Verallgemeinerung: ein graphentheoretisches Problem in: Graphs, Data Structures, Algorithms, Hensen Verlag, München, 1979.

[2565] K. Mumtaz, P. John, and D. R. Silaban, The odd harmonious labeling of matting graph, *J. Phys.: Conf. Ser.,* 1722 012050 (2021) 1-5. https://iopscience.iop.org/article/10.1088/1742-6596/1722/1/012050/pdf

[2566] K. Mumtaz and D. R. Silaban, The odd harmonious labelling of n hair-kC_4-snake graph, 2021 *J. Phys.: Conf. Ser.,* 1725 012089. https://ojs3.unpatti.ac.id/index.php/barekeng/article/view/5091/5129

[2567] A. Munia, J. Maowa, S. Tania, M. Kaykobad, A new class of graceful tree, *Internat. J. Eng. Sci. and Res.,* **5** (2014) 1112-1115. https://www.researchgate.net/publication/359058415_A_new_class_of_Graceful_Tree

[2568] B. J. Murali, K. Thirusangu, and B. J. Balamurugan, Zumkeller cordial labeling of cyclie related graphs, *Internat. J. Pure Appl. Math.,* **116**(3) (2017) 617-627. https://www.ijpam.eu/contents/2017-116-3/7/7.pdf

[2569] B. J. Murali, K. Thirusangu, R. Madura Meenakshi, Biconditional cordial labeling of cycles, *Internat. J. Appl. Eng. Res.* (IJAER), **10**(80) (2015) 188-191.

[2570] B. J. Murali, K. Thirusangu, R. M. Meenakshi, Zumkeller cordial labeling of graphs, *Adv. Intelligent Systems Comput.*, Springer **412** (2015) 533-541.

[2571] K. Murhu Guru Packiam, T. Manimaran, and A. Thuraiswamy, On total edge irregularity strength of graph, *Ars Combin.,* **129** (2016) 173-183. https://www.researchgate.net/publication/318529446_On_Total_Edge_Irregularity_Strength_of_Graphs

[2572] F. A. Muntaner-Batle, Special super edge-magic labelings of bipartite graphs, *J. Combin. Math. Combin. Comput.*, **39** (2001) 107-120.

[2573] M. Mursalina and Y. Susanti, On constructing H-irregularity labeling with minimum label for certain balloon graphs, *Jurnal Matematika UNAND*, **15**(3) Hal (2026) 346–357. https://jmua.fmipa.unand.ac.id/index.php/jmua/article/view/1422/1065

[2574] A. N. Murugan and A. Mathubala, Cycle-related homo-cordial graphs, *Internat. J. Multidisciplinary Res. Develop.*, **2**(10) (2015) 9201%) 84–88. http://www.allsubjectjournal.com/archives/2015/vol2/issue10/2-8-145

[2575] K. Muruganantham and V. Kowsalya, Odd-prime graceful labelling and odd-even prime graceful labelling, *Indian J. Nat. Sci.*, **5**(86) (2024) 82326–82331. https://www.researchgate.net/publication/387798480_Odd-Prime_Graceful_Labelling_and_Odd-Even_Prime_Graceful_Labelling

[2576] K. Muthugurupackiam and S. Ramya, Modular irregularity strength of two classes of graphs, *J. Comp. Math. Sci.* **9**(9) (2018) 1132-1141. https://onlinelibrary.wiley.com/doi/epdf/10.1155/2021/4746609

[2577] I. Muthuselvam, K. Muthugurupackiam, S. Anbalagan, P. Mala, and Y. Shang, Further results on sum index and difference index of graphs, *Res. Math.*, **13**(1) 2026. https://doi.org/10.1080/27684830.2026.2692183

[2578] N. Muthuselvi and T. Saratha Devi, On distance pair mean labeling of graphs, *Indian J. Nat. Sci.,* **15**(86) (2024) https://www.researchgate.net/publication/378745698_On_Distance_Pair_Mean_Labeling_of_Graphs_NMUTHUSELVI

[2579] A. N. Murugan and R. M. I. A. Chitra, Lucky edge labeling of P_n, C_n and corona of P_n, C_n, *IJSIMR*, **2**(8) (2014) 710-718.

[2580] A. N. Murugan and R. M. I. A. Chitra, Lucky edge labeling of fans, spider, and twig graphs, *Proceed. UGC Sponsored 2 day Nat. Seminar Recent Devel. Functional Analysis, Topology and Graph Theory* (2015) 243-250.

[2581] A. N. Murugan and R. M. I. A. Chitra, Lucky edge labeling of triangular graphs, *Internat. J. Math. Trends Tech.,* (IJMTT) **36** (2016) 116-118. https://doi.org/10.14445/22315373/IJMTT-V36P516

[2582] A. N. Murugan, R. M. Irudhaya, and A. Chitra, Lucky-edge labeling of P_n, C_n and corona of P_n, C_n, *Internat. J. Sci. Innov. Math. Res.*, **2**(8) (2014) 710–718. https://www.arcjournals.org/pdfs/ijsimr/v2-i8/6.pdf

[2583] A. Nellai Murugan, R. Maria Irudhaya Aspin Chitra, Lucky edge labeling of bi-star, wheel, parachute, complete and complete bipartite graph, *Proc. UGC Sponsored National Conference on Advances in Fuzzy Algebra, Fuzzy Topology and Fuzzy Graphs, A.P.C. Mahalaxmi College for Women*, Thoothukudi, (2015) 112–121.

[2584] A. N. Murugan, R. M. Irudhaya, and A. Chitra, Lucky-edge labeling of fans, spider and twig graphs, *Proc. UGC Sponsored 2-Day Nat. Sem. Recent Develop. Funct. Anal. Top. Graph Th.* (2015) 243–250.

[2585] A. N. Murugan, R. M. Irudhaya, and A. Chitra, Lucky-edge labeling of H-graph, *OUTREACH* **VIII** (2015) 75–85. http://www.researchmathsci.org/apamart/apam-v14n3-27.pdf

[2586] A. N. Murugan, R. M. Irudhaya, and A. Chitra, Lucky-edge labeling of planar grid graphs, *Internat. J. Mod. Sci. Engin. Tech.*, **2**(9) (2015) 1–8.

[2587] A. N. Murugan, R. M. Irudhaya, and A. Chitra, Lucky-edge labeling of triangular graphs, *Internat. J. Math. Trends Tech.*, **36**(2) (2016) 116–118. https://ijmttjournal.org/archive/ijmtt-v36p516

[2588] A. N. Murugan and P. I. Selvaraj, Path related cup cordial graphs, *Indian J. Applied Res.,* **4**(8) (2011) 433-436. https://www.researchgate.net/publication/276195461_Path_Related_Cup_Cordial_Graphs

[2589] A. N. Murugan and V. Selva Vidhya, Path related hetro-cordial graphs, *Internat. J. Emerging Tech. Engineering Res.*, (2015) **2**(3) (2015) 9-14. https://www.ijeter.everscience.org/Manuscripts/Volume-2/Issue-3/Vol-2-issue-3-M-03.pdf

[2590] K. Murugan and S. Farzana, 2-Skolem difference mean labeling of some graphs, *World Scientific News*, **215** (2026) 11-15. https://worldscientificnews.com/wp-content/uploads/2026/05/WSN-215-2026-1-15.pdf

[2591] K. Murugan and A. Subramanian, Skolem difference mean labeling of H-graphs, *Internat. J. Math. Soft Comput.*, **1**(1) (2011) 115-129. http://oaji.net/articles/2015/296-1433770540.pdf

[2592] M. Murugan, Almost-magic, relaxed-magic and magic strength of a graph, *Util. Math.,* **65** (2004) 53-64.

[2593] M. Murugan and G. Arumugan, Bi-graceful graphs, mumber theory and discrete mathematics (Chandigarh, 2000), Trends in Mathematics, Birkhäuser, Basel (2002) 243–249.

[2594] M. Murugan and G. Arumugan, Are banana trees graceful?, *Math. Ed.* (Siwan), **35** (2001) 18-20.

[2595] M. Murugan and G. Arumugan, On graceful numberings of nC_5 with a common edge, unpublished.

[2596] M. Murugan and G. Arumugan, An algorithm to find graceful numberings of a Spl. class of banana trees, unpublished.

[2597] S. Murugan and S. Chandra Kumar, $H-V$-super magic labeling of H-factorable graphs, *Malaya J. Mat.,* **7**(2) (2019) 138-141. https://www.malayajournal.org/index.php/mjm/article/view/691/435

[2598] S. Murugan and S. Chandra Kumar, H-V-super magic labeling of H-factorable graphs, *Malaya J. Mat.*, **7**(2) (2019) 138-141. https://www.malayajournal.org/archive/selected_article.php?id=571

[2599] S. Murugan and S. C. Kumar, $C_m - E$-super magic graceful labeling of graphs, Malaya Journal of Matematik 7(4) (2019), 716-719. https://www.malayajournal.org/articles/MJM07040015.pdf

[2600] N. Murugesan and R. Senthil Amutha, Bimagic total labeling for bistar $B_{n,n}$, *Internat. J. Sci. Innov. Math. Res. (IJSIMR)*, **2**(9) (2014) 764-769. https://arcjournals.org/article/7208?utm_source=chatgpt.com

[2601] S. Murugesan, D. Jayaraman, and J. Shiama 3-Equitable prime cordial labeling of graphs, *Internat. J. Appl. Inform. Systems* (IJAIS), **5**(9) (2013) 1-4. https://research.ijais.org/volume5/number9/ijais13-450974.pdf

[2602] S. Murugesan, D. Jayaraman, and J. Shiama, Square divisor cordial graphs, *Internat. J. Comput. Appl.*, **64** (22) (2013) 1-4. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=fbb1c787655c3c14afc298bc1af7fca110c65ef0

[2603] A. Muthaiyan and P. Pugalenthi, Vertex odd divisor cordial labeling of graphs, *IJISET - Internat. J. Innovative Sci., Engin. Tech.,* **2**(10) (2015) 395-400. http://ijiset.com/vol2/v2s10/IJISET_V2_I10_53.pdf

[2604] S. Mutharasu, N. M. Bernard, and D. Kumar, V_k-Super vertex magic graceful labeling of graphs, *Malaya J. Mat.,* **8**(3) (2020) 950-954. doi:10.26637/MJM0803/0037

[2605] S. Mutharasu, N. Mary Bernard, D. Kumar, V_k- Super vertex in-magic labelings of directed graphs, *J. Physics: Confer. Ser.*, **1850** (2021) 012067 pages 1-7. https://iopscience.iop.org/article/10.1088/1742-6596/1850/1/012067/pdf

[2606] S. Mutharasu and D. Kumar, V_k-Super vertex magic labeling of graphs, *Malaya J. Mat.,* **6**(4) (2018) 795-799. https://www.malayajournal.org/articles/MJM06040014.pdf

[2607] S. Mutharasu and D. Kumar, E_k-super vertex magic labeling of graphs, *Int. J. Math. Appl.*,**6** (1)(E) (2018) 1005-1012. https://www.semanticscholar.org/paper/E-k-Super-Vertex-Magic-Labeling-of-Graphs-Mutharasu-Kumar/6a89fd6fab91c2b9d799825d8e09f64a0b717788

[2608] S. Mutharasu and D. Kumar, E_k-super vertex in-magic labeling of digraphs, *Math. Eng. Sci. Aerospace,* **11**(1) (2020) 193-199. http://nonlinearstudies.com/index.php/mesa/article/view/2159

[2609] S. Mutharasu, D. Kumar, and N. M. Berrard, V_k-super vertex out-magic labeling of digraphs, *Punjab U. J. Math.,* **57**(7) (2019) 35-41. http://pu.edu.pk/images/journal/maths/PDF/Paper-4_51_7_2019.pdf

[2610] S. Muthukkumar, Edge irregularity strength of binomial trees, May 2024*Commun. Appl. Nonlinear Analy.,* **31**(2) (2024) 214-217. https://internationalpubls.com/index.php/cana/article/view/536/418

[2611] S. Muthukkumar and K. Rajendran, Every tree is an integral sum graph, *Commun. Appl. Nonlinear Analy.,* **31**(2) (2024) 404-408. https://www.researchgate.net/publication/391930529_Every_Tree_is_an_Integral_Sum_Graph

[2612] S. Muthukkumar and K. Rajendran, Generation of anti-magic graphs, *Int. J. Anal. Appl.,* **23**(30) (2025) 1-5. https://etamaths.com/index.php/ijaa/article/view/3202

[2613] K. Muthgu Guru Packiam, Total face irregularity strength of plane graphs, *J. Graph Label.,* **2**(1) (2016) 69-77. https://www.researchgate.net/publication/308992297_Total_face_irregularity_strength_of_plane_graphs

[2614] K. Muthu Guru Packiam, T. Manimaran, and A. Thuraiswamy, Irregularity strength of corona of two graphs, *Theoretial Comput. Sci. Disc. Math.*, 175-181, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017. https://www.researchgate.net/publication/319139269_Irregularity_Strength_of_Corona_of_Two_Graphs

[2615] K. Muthu Guru Packiam, T. Manimaran, and A. Thuraiswamy, 1-distant irregularity strength of graphs, *Theoretical Comput. Sci. Disc. Math.*, 182-190, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017.

[2616] G. Muthumanickavel and K. Murugan, Oblong sum labeling of union of some graphs, *World Scientific News,* **145** (2020) 85-94. http://www.worldscientificnews.com/wp-content/uploads/2020/04/WSN-145-2020-85-94.pdf

[2617] G. Muthumanickavel and M. Nalliah, Optimizing factory workers work shifts scheduling using local antimagic vertex coloring, *Contemp.Math.*, **5**(4) (2024) 5621-5640. https://ojs.wiserpub.com/index.php/CM/article/view/5635

[2618] D. Muthuramakrishnan, k-even Sequential Harmonious Labeling of Graphs, PhD thesis, Bharathidasan University, 2013.

[2619] D. Muthuramakrishnan and S. Sutha, Pell graceful labeling of graphs, *Malaya J. Matematik*, **7**(3) (2019) 508-512. http://www.malayajournal.org/articles/MJM07030023.pdf

[2620] I. Muthuselvam, K. Muthugurupackiam, S. Anbalagan, P. Mala, and Y. Shang, Further results on sum index and difference index of graphs, *Res. Math.*, **13**(1) 2026. https://doi.org/10.1080/27684830.2026.2692183

[2621] N. Muthuselvi and T. Saratha Devi, Local super (a, d) edge antimagic total labeling of graphs, *IAENG Internat. J. Appl. Math.*, **55**(1) (2025) 254-259. https://www.iaeng.org/IJAM/issues_v55/issue_1/IJAM_55_1_28.pdf

[2622] T. Myers and S. Curran, C_4-face-magic labeling on a 4×4 Klein bottle grid graph, https://doi.org/10.48550/arXiv.2606.06817

[2623] H. Nagamochi, M. Miller, and Slamin, Bounds on the number of isolates in sum graph labeling, *Discrete Math.,* **240** (2001) 175-185. https://www.sciencedirect.com/science/article/pii/S0012365X00003903

[2624] S. Nada, A. T. Diab, A. Elrokh, and D. E. Sabra, The corona between cycles and paths, *J. Egyptian Math. Soc.*, **25**(2) (2017) 111-118. https://www.sciencedirect.com/science/article/pii/S1110256X16300542

[2625] S. Nada, A. T. Diab, A. Elrokh, and D. E. Sabra, The corona between paths and cycles, *Ars Combin.*, **139** (2018) 269-281.

[2626] S. Nada, A. Elrayes, A. Elrokh, and A. Rabie, Signed product cordial of the sum and union of two fourth power of paths and cycles, *Machine Learn. Res.,* **4** (2019) 45-50. https://www.academia.edu/100037644/Signed_Product_Cordial_of_the_Sum_and_Union_of_Two_Fourth_Power_of_Paths_and_Cycles

[2627] S. Nada, A. Elrokh, A. Elrayes, and A. Rabie, Cordial labeling of corona product between paths and fourth power of paths, *Iranian J. Math. Sci. Information,*, **18**(2) (2023) 107-115. http://ijmsi.ir/files/site1/user_files_529145/ayaebrahim828_yahoo-A-10-4681-1-d34f166.pdf

[2628] S. Nada, A. I. Elrokh, Y. Elmshtaye, and Abd El-hay, On singed product cordial of cone graph and its second power, *Turkish J. Computer Math. Edu.,* (TURCOMAT) **13**(03) (2022) 597-606. https://www.turcomat.org/index.php/turkbilmat/article/view/13070/9364

[2629] S. Nada, A. I. Elrokh, and E. Elshafey, The cordiality of the join and union of the second power of fans, *Appl. Comput. Math.,* **7**(6) (2018) 219-224. doi:10.11648/j.acm.20180706.11

[2630] S. Naduvtath, A study on set-cordial graphs, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, **71**(2) (2022) 339-348. https://dergipark.org.tr/en/download/article-file/684619

[2631] M. F. Nadeem, M. Cancan, M. Imran, and Y. Ali, On edge irregularity strength of certain families of snake graph, *J. Prime Res. Math.,* **19**(1) (2023) 92-101. https://www.researchgate.net/publication/374081679_On_edge_irregularity_strength_of_certain_families_of_snake_graph

[2632] M. F. Naeem, M. K. Siddiqui, M. Bača, A. Semaničová-Feňovčíková, and F. Ashraf, On edge H-irregularity strengths of some graphs, *Discuss. Math. Graph Th.*, **41**(4) (2021) 949–961. https://www.researchgate.net/publication/333749610_On_edge_H-irregularity_strengths_of_some_graphs

[2633] U. Nadzima and T. S. Martini, Super $(a, d) - H$-antimagic total labeling on wheel edge corona product with a path and a cycle, ICMETA 2018 IOP Conf. Series: *J. Physics: Conf. Series* 1306 (2019) 012007 2414-2423. doi:10.1088/1742-6596/1306/1/012007

[2634] M. Naeem and M. K. Siddiqui, Total irregularity strength of disjoint union of isomorphic copies of generalized Petersen graph, *Discrete Math. Algor. Appl.,* **9**(6) (2017) 1750071, 9 pp. https://www.researchgate.net/publication/320507355_Total_irregularity_strength_of_disjoint_union_of_isomorphic_copies_of_generalized_Petersen_graph

[2635] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaran, Even vertex magic total labeling, *Internat. J. Pure Appl. Math.*, **115** (9)(2017) 363-375. https://www.acadpubl.eu/jsi/2017-115-9/articles/9/32.pdf

[2636] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaran, *Internat. J. Math Trends Tech.,* Odd vertex magic total labeling of trees, **52**(6) (2017) 374-379. https://ijmttjournal.org/public/assets/volume-52/number-6/IJMTT-V52P553.pdf

[2637] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaran, Even vertex magic total labeling of isomorphic and non isomorphic suns, *Internat. J. Math Trends Tech.,* **52** (7) (2017) 458-467. https://ijmttjournal.org/archive/ijmtt-v52p566?utm_source=chatgpt.com

[2638] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaan, Even vertex magic total labeling of some 2-regular graphs, *Internat. J. Math Trends Tech.,* **54**(1) (2018) 52-59. https://ijmttjournal.org/public/assets/volume-54/number-1/IJMTT-V54P506.pdf

[2639] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaan, Odd vertex magic total labeling of some graphs, *Internat. J. Pure Appl. Math.*, **115** (10)(2018) 97-109. https://www.acadpubl.eu/jsi/2018-118-10-11/articles/10/53.pdf

[2640] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakan, Odd vertex magic total labeling of isomorphic and non isomorphic suns, *Internat. J. Math Trends Tech.,* **52**(1) (2018) 34-41. https://www.researchgate.net/publication/322577783_EVEN_VERTEX_MAGIC_TOTAL_LABELING_OF_ISOMORPHIC_AND_NON_ISOMORPHIC_SUNS

[2641] CT. Nagaraj, C. Y. Ponnappan, and G. Prabakaran, Multiple magic graphs, *Int. J. Math. Appl.*, **6**(1) (2018) 109-115. (Special Issue) https://ijmaa.in/index.php/ijmaa/article/view/1403/1361

[2642] A. Nagarajan, R. Vasuki, and S. Arockiaraj, Super mean number of a graph, *Kragujevac J. Math.* **36**(1) (2012) 93-107. https://imi.pmf.kg.ac.rs/kjm/pub/13476270796413_kjom3601-11.pdf

[2643] S. Nagarajan and G. Priyadharsini, Lucky edge labeling of new graphs, *Internat. J. Math. Trends Tech.,* (IJMTT) **65**(8) (2019) 26-30. http://www.ijmttjournal.org/Volume-65/Issue-8/IJMTT-V65I8P505.pdf

[2644] H. M. Nagesh and V. R. Girish, On edge irregularity strength of line graph and line cut-vertex graph of comb graph, *Not. Numb. Theory Discrete Math.*, **28**(3) (2022) 517–524. https://doi.org/10.7546/nntdm.2022.28.3.517-524

[2645] M. Nalliah, Antimagic labeling of digraphs, *J. Indones. Math. Soc.,* **22**(1) (2016) 61-69. https://jims-a.org/index.php/jimsa/article/view/216/pdf

[2646] M. Nalliah, A survey on recent open problems on super (a, d)-edge antimagic total labelings *J. Graph Labeling,* **1**(2) (2015) 129-136.

[2647] M. Nalliah and S. Arumugam, Super $(a, 3)$-edge antimagic total labeling for union of two stars, *Theoretical Computer Science and Discrete Mathematics*, 203-211, Lecture Notes in Comput. Sci., 10398, Springer, Cham, 2017.

[2648] M. Nalliah, R. Shankar, and T.-M. Wang, Local antimagic vertex coloring for generalized friendship graphs, *J. Disc. Math. Sci. Crypt.*, **26**(4) (2023) 1063–1078. https://tarupublications.com/journals/jdmsc/volume/26/issue/4/local-antimagic-vertex-coloring-for-generalized-friendship-graphs/pdf

[2649] A. K. Handa, A. Godinho, T. Singh, and S. Arumugam, Distance antimagic labeling of join and corona of two graphs, *AKCE Internat. J. Graphs Combin.*, **14**(2) (2017) 172–177. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2017.04.003?scroll=top&needAccess=true#d1e1353

[2650] S. P. Nandhini and B. Pooja Lakshmi, Study on prime graceful labeling for some special graphs, *Nat. Volatiles & Essent. Oils* (2021) 13161-13171.

[2651] S. A. A. Naqvi, F. Ashraf, A. Ovais, M. W. Rasheed, A. Asif, and A. Alameri, Optimizing hybrid network topologies in communication networks through irregularity strength, *Scientific Reports,* **15**(1) (2025) doi:DOI:10.1038/s41598-025-05631-8

[2652] D. Narayan, Problem 380. Representations of graphs modulo n, *Discrete Math.,* **257** (2002) 614.

[2653] D. Narayan, An upper bound for the representaion number of graphs of fixed order, *Integers,* **3** (2003) A12 4 pages. https://repository.rit.edu/cgi/viewcontent.cgi?article=1291&context=article

[2654] G. Narasimhan and R. Sampathkumar, Contra harmonic mean labeling of graphs, *Mapana-J. Sciences*,**12**(8) (2013) 23–29. https://doi.org/10.12723/mjs.26.7

[2655] S. Naresh kumar and D. Bharathi, Even-odd cordial labeling on join of zero divisor graphs, *Internat. J. Math. Trends Tech.*, **72**(7) (2026) 53-68. https://ijmttjournal.org/public/assets/volume-72/issue-7/IJMTT-V72I7P107.pdf

[2656] A. Naseem, K. Shabbir, and H. Shaker, The radio number of edge-joint graphs, *Ars Combin.,* **139** (2018) 337-351. https://www.scribd.com/document/775453703/2018-The-Radio-Number-of-Edge-Joint-Graphs

[2657] S. Nasir, N. Idrees, A. Sadiq, F. B. Farooq, S. Kanwal, and M. Imran, Strongly multiplicative labeling of diamond graph, generalized Petersen graph, and some other graphs, *Hindawi J. Math.*, **2022**, Article ID 3203108, 5 pages. https://www.hindawi.com/journals/jmath/2022/3203108/

[2658] A. Nasr, A. Elsonbaty, M. A. Seoud, and M. Anwar, Diophantine graphs, (2025) 13pp. https://www.researchgate.net/publication/387670539_Diophantine_Graphs

[2659] K. R. A. Navas, V. Ajitha, and T. K. M. Varkey, On vertex integer-magic spectra of caterpillar graphs, *Malaya J.Mat.*, **8**(4) (2020) 1543-1546. https://www.malayajournal.org/articles/MJM08040035.pdf

[2660] S. Nazeer, M. S. Khan, I. Kousar, and W. Nazeer, Radio labeling and radio number for generalized caterpillar graphs, *J. Appl. Math. Informatics,* **34** (5-6) (2016) 451-465. https://koreascience.kr/article/JAKO201628741000716.page

[2661] S. Nazeer, I. Kousar, and W. Nazeer, Radio and radio antipodal labelings for circulant graphs $G(4k+2;\{1,2\})$, *J. Appl. Math. Inform.,* **33**(1-2) (2015) 173-183. https://www.koreascience.or.kr/article/JAKO201505557791732.pdf

[2662] W. Nazeer, S. Nazeer, I. Kousar, and M. Munir, The radio and radio antipodal numbers of non-bipartite cubic graphs of order 2^k, *Ars Combin.,* **145** (2019) 301-312. https://www.researchgate.net/publication/315498239_The_radio_and_radio_antipodal_numbers_of_non-bipartite_cubic_graphs_of_order_2_k

[2663] P. Nedumaran, K. Thirusangu, and V. Celin Mary, 1-Near mean cordial and biconditional cordial labeling of certain classes of graphs, *Internat. J, Pure and Appl. Math.*, **117**(12) (2017), 297-307. https://acadpubl.eu/jsi/2017-117-11-14/articles/12/33.pdf

[2664] N. Neela and C. Selvaraj, Total magic cordial labeling of complete multipartite graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), 251–254, *Electron. Notes Discrete Math.*, **48**, Elsevier Sci. B. V., Amsterdam, 2015. https://www.researchgate.net/publication/279635915_Total_Magic_Cordial_Labeling_of_Complete_Multipartite_Graphs

[2665] N. Neela and C. Selvaraj, Conjecture on odd graceful graphs, *J. Combin. Math. Combin. Comput.,* **97** (2016) 65-82. https://www.researchgate.net/publication/305266174_Conjecture_on_odd_graceful_graphs

[2666] J. Nešetřil and A. Pultr, A Dushnik-Miller type dimension of graphs and its complexity, in: M. Karpinski, Ed., Fundamentals of Computation Theory, Lecture Notes in Computer Science, **56**, Springer, Berlin, 1977, 482-493.

[2667] A. NellaiMurugan and R. M. I. A. Chitra, Lucky edge labeling of planar grid graph, *Int. J. Mod. Sci. Eng. Technol.*, **2** (2015), 1–8.

[2668] S. Nemani and G. Joshi, E-super vertex magic labeling of cartoon flowers and wounded flowers, *AKCE Int. J. Graphs Comb.,* **17**(3) (2020) 1039-1044. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2019.09.001

[2669] C.-m Ni Z.-s Liu, and F.-l Lu, On the E_3-cordiality of some graphs, *J. Combin. Math. Combin. Comput.,* **97** (2016) 3-10.

[2670] T. Nicholas and K. J. Bosco, Radio mean D-distance labeling of cycle-relted graphs, *Internat. J. Math. Trends Tech.,* (IJMTT) **40**(3) (2016) 247-251. https://ijmttjournal.org/public/assets/volume-40/number-3/IJMTT-V40P530.pdf

[2671] T. Nicholas and P. Maya, Some results on integer cordial graph, *J. Progressive Res. Math.*, (JPRM) **8**(1) (2016) 1183-1194. https://scitecresearch.com/journals/index.php/jprm/article/view/786

[2672] T. Nicholas and P. and Maya, Some results on integer edge cordial graph, *J. Eng. Appl. Math.,* **3**(1) 2017) 1-12. doi:10.18831/djmaths.org/2017011001

[2673] T. Nicholas, S. Somasundaram, and V. Vilfred, Some results on sum graphs, *J. Comb. Inf. System Sci.*, **26** (2001) 135-142. https://www.researchgate.net/profile/V-Vilfred-Kamalappan/publication/342804207_P3-NSV-sum-graph/links/5f06f5ff92851c52d6249871/P3-NSV-sum-graph.pdf

[2674] A. J. Niedzialomski, Consecutive radio labelings and the Cartesian product of graphs, Ph. D. Thesis, University of Iowa, August 2013.

[2675] A. J. Niedzialomski, Radio graceful Hamming graphs, *Discuss. Math. Graph Theory,* **36**(4) (2016) 1007-1020. https://www.researchgate.net/publication/305270422_Radio_Graceful_Hamming_Graphs

[2676] T. Nierhoff, A tight bound on the irregularity strength of graphs, *SIAM J. Discrete Math.,* **13** (2000) 313-323.

[2677] S. A. N. Ni'mah and D. Indriati, On vertex irregular total k-labeling and total vertex irregularity strength of lollipop graphs, *J. Physics Confer. Series* 1306:012025 July 2019. https://iopscience.iop.org/article/10.1088/1742-6596/1306/1/012025/pdf

[2678] S. Nila Steffi and J. Subhashini, Product binary L-cordial labeling, *Adv. Appl. Math. Sci.* **21**(3 (2022) 1455-1462. https://www.mililink.com/upload/article/1547354525aams_vol_213_january_2022_a33_p1455-1462_s._nila_steffi__and_j._subhashini.pdf

[2679] S. Nila Steffi and J. Subhashini, Sum binary L-cordial labeling, *AIP Conf. Proceed,*, 2385, 130006 (2022).

[2680] S. Nila Steffi and J. Subhashini, Mean binary L-cordial labeling, *Internat. J. Artificial Intelligence and Machine Learning,* **6**(9s) (2026) 12-17 https://www.svedbergopen.com/index.php/ijaiml/article/view/1484/923

[2681] P. K. Niranjan and S. Rao Kola, On the radio number for corona of paths and cycles, *AKCE Internat. J. Graphs Combin.*, **17**(1) 2020) 269-275. https://www.sciencedirect.com/science/article/pii/S0972860019300684

[2682] F. Nirmala Irudayam and F. Sherin, Generalized geometric mean-labelling of m-shadow graphs and its operators, *Bol. Soc. Paran. Mat.*, (3s.) **44**(12) (2026) 1–14. https://periodicos.uem.br/ojs/index.php/BSocParanMat/article/view/80436/751375163125

[2683] R. Nirmalasari Wijaya, J. Ryan, and T. Kalinowski, Cube-magic labelings of grids, *Ars Combin.*, textit151 (2020) 247–256. arXiv:1702.02639.

[2684] R. W. Nirmalasari, Wijaya, J. Ryan, and T. Kalinowski, Total edge irregularity strength of the Cartesian product of bipartite graphs and paths, *J. Indones. Math. Soc.,* **29**(02) (2023) 156-165. https://dergipark.org.tr/en/download/article-file/5376785

[2685] R. Nishanthini, Vertex N-magic total labeling of Jahangir, octopus, and cycle-star graphs with applications to chemical reaction networks and water pollution, *Nonlinear Studies*, **33**(1) (2026) 441-446.

[2686] R. Nishanthini and R. Jeyabalan, Sharp bounds on vertex N-magic total labeling of graphs, *Math. Stat.*, **12**(3) (2024) 234-239. https://www.hrpub.org/journals/article_info.php?aid=14234

[2687] R. Nishanthini and R. Jeyabalan, Strong VNMT labelings on graph families and their applications in disease transmission networks, *Math. Eng., Sci. Aerospace*, **16**(3) (2025) 743-752.

[2688] R. Nishanthini1 and R. Jeyabalan, Extension of vertex N-magic total graphs with duality approach, *Palestine J. Math.*, **15**(1) (2026) 584–592. https://www.researchgate.net/publication/403401945_EXTENSION_OF_VERTEX_N_-MAGIC_TOTAL_GRAPHS_WITH_DUALITY_APPROACH

[2689] R. Nishanthini and R. Jeyabalan, Vertex N-magic total labelings in graph families: Existence, constructions, and constraints, *Math. Engin. Sci. Aerospace (MESA) 17*(2) (2026) 781–793.

[2690] R. Nisviasari, Dafik, and I. H. Agustin, The total H-irregularity strength of triangular ladder graphs, *J. Phys.: Conf. Ser.,* 1465 012026 (2020) https://iopscience.iop.org/article/10.1088/1742-6596/1465/1/012026

[2691] R. Nithya and K. Anitha, Even vertex ζ-graceful labeling on rough graph, *Math. Stat.*, **11**(1) (2023) 71–77. doi:10.13189/ms.2023.110108

[2692] D. Nivedha and S. Yamini, On the antimagicness of generalized edge corona graphs, *Heliyon* **10** (2024) e24002. doi:10.1016/j.heliyon.2024.e24002

[2693] H. K. Ng, α-valuations and k-gracefulness, *Notices AMS,* **7** (1986) 247.

[2694] H. K. Ng, Gracefulness of a class of lobsters, *Notices AMS,* **7** (1986) 825-05-294.

[2695] S. Ng, F. Alwie, T. P. Marjadi, and K. A. Sugeng, Harmonious labeling of the union of m-cycle and n-path graph, *J. Phys.: Conf. Ser.,* **1725** (2021) 012088 1-8. https://iopscience.iop.org/article/10.1088/1742-6596/1725/1/012088

[2696] A. A. G. Ngurah, On (a, b)-edge-antimagic total labeling of odd cycle. *J. Indones. Math. Soc.,* **9** (2003) 9-12.

[2697] A. A. G. Ngurah, On magic and antimagic total labelings of graphs, *J. Phys.: Conf. Ser.,* 1503 012006 2020. 7pp. https://iopscience.iop.org/article/10.1088/1742-6596/1503/1/012006/pdf

[2698] A. A. G. Ngurah and Adiwijaya, New results on the (super) edge-magic deficiency of chain graphs, *Int. J. Math. Math. Sci.,* (2017), Art. ID 5156974, 6 pp. https://onlinelibrary.wiley.com/doi/epdf/10.1155/2017/5156974

[2699] A. A. G. Ngurah, On the (super) edge-magic deficiency of chain graphs, *J. Combin. Math. Combin. Comput.,* **103** (2017) 225-236. https://combinatorialpress.com/article/jcmcc/Volume%20103/vol-103-paper-17.pdf

[2700] A. A. G. Ngurah and E. Baskoro, On magic and antimagic total labelings of generalized Petersen graph, *Util. Math.,* **63** (2003) 97-107.

[2701] A. A. G. Ngurah, E. Baskoro, and R. Simamjuntak, On antimagic total labelings of generalized Petersen graph, *J. Combin. Math. Combin. Comput.,* **55** (2005) 57-70. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%205.pdf

[2702] A. A. G. Ngurah, E. Baskoro, and R. Simamjuntak, On (a, d)-edge-antimagic total labelings of mC_n, *Bull. Inst. Combin. Appl.,* **48** (2006) 35-44.

[2703] A. A. G. Ngurah, E. Baskoro, and R. Simamjuntak, On the super edge-magic deficiencies of graphs, *Australas. J. Combin.,* **40** (2008) 3-14. http://ajc.maths.uq.edu.au/pdf/40/ajc_v40_p003.pdf

[2704] A. A. G. Ngurah, E. Baskoro, and R. Simamjuntak, On new families of (super) edge-magic graphs, *Util. Math.,* **74** (2007) 111-120.

[2705] A. A. G. Ngurah, E. T. Baskoro, and I. Tomescu, Magic graphs with pendant edges, *Ars Combin.,* **99** (2011) 149-160. https://combinatorialpress.com/article/ars/Volume%2099/volume-99-paper-15.pdf

[2706] A. A. G. Ngurah, A. N. M. Salman, and I. W. Sudarsana, On supermagic coverings of fans and ladders, *SUT J. Math.,* **46** (2010) 67-78. https://projecteuclid.org/journals/sut-journal-of-mathematics/volume-46/issue-1/On-supermagic-coverings-of-fans-and-ladders/10.55937/sut/1280430657.full

[2707] A. A. G. Ngurah, A. N. M. Salman, and L. Susilowati, H-supermagic labelings of graphs, *Discrete Math.,* **310** (2010) 1293-1300. https://www.sciencedirect.com/science/article/pii/S0012365X09006219

[2708] A. A. G. Ngurah and R. Simanjuntak, Super edge-magic deficiency of join-product graphs, *Util. Math.,* **105** (2017) 279-289. https://www.researchgate.net/publication/259803975_Super_edge-magic_deficiency_of_join-product_graphs

[2709] A. A. G. Ngurah and R. Simanjuntak, On the super edge-magic deficiency of join product and chain graphs, *Electron. J. Graph Theory Appl. (EJGTA),* **7**(1) (2019) 157-167. https://www.ejgta.org/index.php/ejgta/article/view/487

[2710] A. A. G. Ngurah, R. Simamjuntak, and E. Baskoro, On (super) edge-magic total labeling of subdivision of $K_{1,3}$, *SUT J. Math.,* **43** (2007) 127-136.

[2711] T. Nicholas and S. Somasundaram, More results on integral sum graphs, Graph Theory and its Applications, Editors: R. Balakrishnan et al., Narosa Publishing House, New Delhi, India (2004) 75-83.

[2712] T. Nicholas and S. Somasundaram, More results on sum graphs, *Proceedings of the Conference on Graph Theory and its Applications held at Anna University, Chennai,* (2001) 73-83.

[2713] T. Nicholas, S. Somasundaram, and V. Vilfred, On (a, d)-antimagic special trees, unicyclic graphs and complete bipartite graphs, *Ars Combin.,* **70** (2004) 207-220. https://combinatorialpress.com/article/ars/Volume%20070/volume-70-paper-19.pdf

[2714] T. Nicholas and V. Vilfred, Sum graph and edge reduced sum number, unpublished.

[2715] R. W. Nirmalasari Wijaya, A. Semaničová-Feňovčíková, J. Ryan, and T. Kalinowski, H-supermagic labelings for firecrackers, banana trees and flowers, *Australasian J. Combin.,* **69**(3) (2017) 442-451. https://ajc.maths.uq.edu.au/pdf/69/ajc_v69_p442.pdf?utm_source=chatgpt.com

[2716] R. Nistyawati and T. S. Martini, On super $(a, d) - H$-antimagic total labeling of edge corona product and corona product of gear graph with path and cycle, IOP Conf. Series: *J. Physics: Conf.,* Series 1306 (2019) 012010 1-7. https://iopscience.iop.org/article/10.1088/1742-6596/1306/1/012010/pdf

[2717] J. A. Novelia and D. Indriati, Edge irregular reflexive labeling on banana tree graphs $B_{2,n}$ and $B_{3,n}$, *AIP Conf. Proc.*, **2326** (2021) doi:doi:10.1063/5.0039316

[2718] T. Noviati, T. S. Martini, and D. Indriati, On H-supermagic labeling of friendship edge corona product with path and sun edge corona product with path, *J. Physics Conference Series,* 1306:012008 (2019) 1-6. https://iopscience.iop.org/article/10.1088/1742-6596/1306/1/012008/pdf

[2719] S. Novotny, J. Ortiz, and D. Narayan, Minimal k-rankings and the rank number of P_n^2, *Inform. Process. Lett.,* **109** (2009) 193-198. https://repository.rit.edu/cgi/viewcontent.cgi?article=2439&context=article

[2720] R. Nowakowski and C. Whitehead, Ordered graceful labellings of the 2-star, Graph theory (Prague, 1998), *Discrete Math.,* **233** (2001) 183-191. https://www.sciencedirect.com/science/article/pii/S0012365X00002375?via%3Dihub

[2721] K. Nurdin and K. Hye, Irregular labeling on transportation network of splitting graphs of stars, *Proc. Jangjeon Math. Soc.,* **22**(1) (2019) 103-108. http://jangjeonopen.or.kr/public/upload/1549515921-pjms22-1-11.pdf

[2722] N. Nurdin, On the total edge irregularity strength of quadtrees and all banana trees, *J. Indones. Math. Soc.,* **18**(1) (2012) 31-36. https://media.neliti.com/media/publications/93485-EN-on-the-total-vertex-irregularity-strengt.pdf

[2723] S. Nurdin, A. N. M. Salman, and E. T. Baskoro, The total edge-irregular strengths of the corona product of paths with some graphs, *J. Combin. Math. Combin. Comput.,* **65** (2008) 163-175. https://www.researchgate.net/publication/261216173_The_total_edge-irregular_strengths_of_the_corona_product_of_paths_with_some_graphs

[2724] S. Nurdin, E. T. Baskoro, A. N. M. Salman,and N. N. Gaos, On the total vertex irregularity strength of trees. *Discrete Math.,* **310** (2010) 3043-3048. https://www.researchgate.net/publication/220193224_On_the_total_vertex_irregularity_strength_of_trees

[2725] S. Nurdin, E. T. Baskoro, A. N. M. Salman, and N. N. Gaos, On the total vertex irregular labelings for several types of trees, *Util. Math.*, **83** (2010) 277-290. https://www.researchgate.net/publication/264943737_On_total_vertex-irregular_labellings_for_several_types_of_trees

[2726] E. Nurdini, I. Rosyida, and Mulyono, On total edge irregularity strength of tadpole chain graph $Tr(6, n)$, *J. Phys.: Conf.,* Ser. 1538 (2020) 012001 1-9. https://iopscience.iop.org/article/10.1088/1742-6596/1538/1/012001

[2727] S. Nurdin, A. N. M. Salman, N. N. Gaos, and E. T. Baskoro, On the total vertex-irregular strength of a disjoint union of t copies of a path, *J. Combin. Math. Combin. Comput.,* **71** (2009) 227-233. https://combinatorialpress.com/article/jcmcc/Volume%20071/vol-071-paper%2016.pdf

[2728] D. E. Nurvazly and K. A. Sugeng, Graceful labelling of edge amalgamation of cycle graph, *J. Physics Confer. Ser.,* **1108** 012047 (2018) http://iopscience.iop.org/article/10.1088/1742-6596/1108/1/012047/pdf

[2729] D. Nurvazly, S. L. Chasanah, and A. R. Wiranto, Variations of graceful labelling of subgraph of millipede graph, *AIP Conference Proceedings 2563*, 050018 (2022). https://doi.org/10.1063/5.0105447

[2730] U. Odabasi, · D. Roberts, and · R. M. Low, The integer-antimagic spectra of Hamiltonian graphs, *Elect. J. Graph Th. and Appl.,* **9**(2) (2021) 301-308. https://www.ejgta.org/index.php/ejgta/article/view/1097/pdf_180

[2731] U. Odabasi, D. Roberts, and R. M. Low, The integer-antimagic spectra of a disjoint union of Hamiltonian graphs, *Turkish J. Math.*, **46** (2022) 1310–1317. https://doi.org/10.55730/1300-0098.3161

[2732] U. Odabasi, D. Roberts, and R. M. Low, The integer-antimagic spectra of a weak join of Hamiltonian graphs, *Theory Appl. Graphs*, **12**(1) (2025) Article 5. https://doi.org/10.20429/tag.2025.120105

[2733] U. Odabas¸ D. Roberts, and R. M. Low, Preservation of the $\mathbb{Z}_k$ property of a graph under edge addition, *Discuss. Math. Graph Th.* (2026) 13pp. doi:DOI:10.7151/dmgt.2628

[2734] S. H. Obaed and N. E. Arif, LH labelling of some certain graphs, AIP Conf. Proc. 2834, 080057 (2023) 080057-1-9. https://doi.org/10.1063/5.0161943

[2735] A. S. Odyuo and P. Mercy, On Fibonacci range labeling for standard shell graphs, *J. Adv. Math. Comput. Sci,* **38**(7) (2023) 67-75. https://www.researchgate.net/publication/370248581_On_Fibonacci_Range_Labeling_for_Standard_Shell_Graphs

[2736] M. A. Ollis, New complete Latin squares of odd order, *European J. Combin.,* **41** (2014) 35-46. https://www.sciencedirect.com/science/article/pii/S019566981400050X?via%3Dihub

[2737] T. Öner and E. Ate, Investigation of C_m-supermagic properties in the union of C_n and C_m graph structures, *J. New Theory*, **50** (2025) 9-29. https://dergipark.org.tr/en/download/article-file/4548449

[2738] T. Öner and E. Erol, On C_m-supermagicness of book-snake graphs, *Punjab Univ. J. Math.*, (Lahore) **53**(4) (2021) 221-230. http://acikerisim.mu.edu.tr:8080/xmlui/bitstream/handle/20.500.12809/9305/oner.pdf?sequence=1&isAllowed=y

[2739] T. Öner, M. Hussain, and S. Banaras, C_m-supermagic labeling of polygonal snake graphs, *J. Math. Comput. Sci.*, **20**(3) (2020) 20(3) 189-195. https://www.isr-publications.com/jmcs/articles-8292-c-m-supermagic-labeling-of-polygonal-snake-graphs

[2740] P. H. Ong, H. V. Chen, and W. S. Ng, Antimagic labeling of graph unions of trees and 4-cycles, *J. Indones. Math. Soc.*, **32**(03) (2026) 1–16. https://jims-a.org/index.php/jimsa/article/view/2238/497

[2741] M. J. M. Orias and A. C. Pedrano, On super root square mean labeling of some cycle related graphs, *Acta Sci. Comput. Sci.,* **5**(1) (2023) 40-48. https://www.gjesr.com/Issues%20PDF/Archive-2018/November-2018/23.pdf

[2742] A. Oshima, Personal communication, 2024.

[2743] A. Ovais, M. Umar, M. Bača, and A. Semaničová-Feňovčíková, Fans are cycle-antimagic, *Australas. J. Combin.,* **68** (2017) 94-105. https://ajc.maths.uq.edu.au/pdf/68/ajc_v68_p094.pdf

[2744] L. Packiam and K. Kathiresan, On total vertex irregularity strength of graphs, *Discuss. Math. Graph Theory,* **32**(1) (2012) 39-45. https://bibliotekanauki.pl/articles/743655

[2745] J. L. Painter, H. C. Landers, and W. M. Mattox, Harmonious labelings via cosets and subcosets, *Theory Appl. Graphs,* **9**(2) Article 4 (2022) 24 pages. https://digitalcommons.georgiasouthern.edu/tag/vol9/iss2/4

[2746] R. N. Pakpahan, I. Mursidah, I. Novitasari, and K. A. Sugeng, Graceful labeling for some supercaterpillar graphs, *AIP Confer.Proceed.*, **1862**(1) 030121 (2107). https://pubs.aip.org/aip/acp/article/1862/1/030121/649927/Graceful-labeling-for-some-supercaterpillar-graphs

[2747] K. Palani and A. Shunmugapriya, Near mean labeling in dragon digraphs, *J. Xidian Univer.*, **14**(3) (2020) 1298-1307. https://www.researchgate.net/publication/340979735_NEAR_MEAN_LABELING_IN_DRAGON_DIGRAPHS

[2748] K. Palani and A. Shunmugapriya, Near mean labeling in dicyclic snakes, *Ratio Math.*, **45** (2023) 238-247. http://eiris.it/ojs/index.php/ratiomathematica/article/view/238-247/0

[2749] K. Palani and J. Surya, 1-Near mean cordial labeling of wheel, globe, binary trees, odd cycle with a chord, *Indian J. App. Res.*, **5** (2015). https://www.worldwidejournals.com/indian-journal-of-applied-research-(IJAR)/recent_issues_pdf/2015/October/October_2015_1492512934__50.pdf

[2750] K. Palani and J. Surya, 1-Near mean cordial labeling of graphs, *Internat. J. Math. Archive* **6**, no. 7 (2015) 15–20. https://www.researchgate.net/publication/334604938_1-NEAR_MEAN_CORDIAL_LABELING_OF_GRAPHS

[2751] W. Pan and X. Lu, The gracefulness of two kinds of unconnected graphs $(P_2 \vee \overline{K_n}) \cup St(m)$ and $(P_2 \vee \overline{K_n}) \cup T_n$, *J. Jilin Univ.*, **41** (2003) 152-154.

[2752] A. C. Panda and D. Mishra, Some new classes of graceful diameter six trees, *Turkic World Math. Soc. J. Appl. Engin. Math.,* **5**(2) (2015) 269-275. https://dergipark.org.tr/tr/download/article-file/1179779

[2753] A. C. Panda, D. Mishra, R. B. Dash, A class of diameter six trees exhibiting graceful labeling. *J. Discrete Math. Sci. Cryptogr.,* **19**(5-6) (2016) 947-963.

[2754] J. Pandimadevi and S. P. Subbiah, Vertex antimagic total labeling of digraphs, *Kyungpook Math. J.,* **55**(2) (2015) 267-277. https://pdfs.semanticscholar.org/4ae7/c43db787fc9282ef5730d3efea848dfde0e0.pdf

[2755] P. Pandya and N. P. Shrimali, Vertex-edge neighborhood prime labeling of some graphs, *Internat. J. Scien. Res. Review,* **7**(10) 2018 735-743. http://www.dynamicpublisher.org/gallery/82-ijsrr-d1002.pdf

[2756] P. Panigrahi, A survey on radio k-colorings of graphs, *AKCE J. Graphs Combin.,* **6** (2009) 161-169. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2009.12088883?utm_source=chatgpt.com

[2757] P. Panigrahi and D. Mishra, Graceful lobsters obtained from diameter four trees using partitioning technique, *Ars Combin.,* **87** (2008) 291-320. https://combinatorialpress.com/article/ars/Volume%20087/volume-87-paper-24.pdf

[2758] P. Panigrahi and J. Saha, On harmoniousness of hypercubes, *AKCE J. Graphs Combin.,* **5** (2008) 189-198. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2008.12088864?utm_source=chatgpt.com

[2759] S. Panma and P. Rochanakul, Prime-graceful graphs, *Thai J. Math.*, **19**(4)(2021) 1685-1697. https://thaijmath2.in.cmu.ac.th/index.php/thaijmath/article/view/1264

[2760] A. Panpa, S. Imnang, and T. Wasuanankul, Graceful labeling of spider graphs with at most five legs, *J. Appl. Math.,* **2025** Article ID 5826777, 5 pages. https://onlinelibrary.wiley.com/doi/epdf/10.1155/jama/5826777

[2761] A. Panpa and T. Poomsa-ard, On graceful spider graphs with at most four legs of lengths greater than one, *J. Appl. Math.,* **3** (2016) 1-5. https://onlinelibrary.wiley.com/doi/epdf/10.1155/2016/5327026

[2762] P. T. Pantoja, S. Dantas, and A. G. Luiz, Graceful colorings of graphs with maximum degree three, *Discrete Appl. Math.*, **374** (2025) 75-96.

[2763] P. T. Pantoja, S. Dantas, and A. G. Luiz, On graceful colorings in subcubic trees, *Discrete Appl. Math.*, **381** (2026) 98-111. https://doi.org/10.1016/j.dam.2025.11.027

[2764] N. Paramaguru, and D. Manirathnam, Lehmer-3 mean labeling of bicyclic graphs, *Adv. Mechanics*, **12**(1) (2024) 274-286. https://advancesinmechanics.com/pdf/2024-274.pdf

[2765] K. M. Paramasivam, K. Yenoke, B. S. K. Muralidharan, Radial radio number of chess board graph and king's graph, *TELKOMNIKA Telecommunication, Comput., Electronics Control*, **20**(1) (2022) 9-18. https://www.semanticscholar.org/reader/5ae484e43f627baf7e79ca061dee9d58151e81c0

[2766] R. Parameswari, Total magic cordial labeling of square and shadow graphs, *Appl. Math. Sci.*, **9**(25) (2015 1229-1234. http://dx.doi.org/10.12988/ams.2015.4121058

[2767] R. Parameswari and C. Jayalakshmi, On cordial totally magic labeling of flower graphs, *AIP Confer. Proceed.*, 2639, 040002 (2022) 151-160. doi:10.1007/978-981-19-5224-1_17

[2768] R. Parameswari and R. A. Margret, Square sum labeling of Octopus graph and Vanessa graph, *November 2022AIP Conference Proceed.,* **2516**(1):210042 doi:10.1063/5.0119754

[2769] R. Parameswari and R. Rajeswari, Total bimagic labeling and total magic cordial labeling of Paley digraphs, 2013 International Conference on Pattern Recognition, Informatics and Mobile Engineering, 2013, 506-509, doi:10.1109/ICPRIME.2013.6496723

[2770] R. Parameswari and R. Rajeswari, Total magic cordial labeling of (P_n, K_1), gear and shadow graphs, *Int. J. Pure Appl. Math.,* **101**(6) (2015) 993-1001. https://acadpubl.eu/jsi/2015-101-5-6-7-8/2015-101-6/17/17.pdf

[2771] R. Parameswari, K. Saradha Pritha, and R. Rajeswari, Integer cordial and face integer cordial labeling of some flower graphs, *Internat. Conf. Math. Sci.*, (ICMS 2020), *J. Physics: Confer. Ser.,* 1770 (2021) 012079. https://iopscience.iop.org/article/10.1088/1742-6596/1770/1/012079

[2772] J. Parejiya, D. Jani, and Y. Hathi, Harmonic mean cordial labeling of some graphs, *TWMS J. App. and Eng. Math.,* **14**(1) (2023) 284-297. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.1/24.pdf

[2773] J. Parejika, P. T. Lalchandani, D. B. Jani, and S. Mundadiya, Harmonic mean cordial labelings of some well known graphs, *TWMS J. App. Eng. Math.,* **15**(3) (2025) 651-660. https://jaem.isikun.edu.tr/web/images/articles/vol.15.no.3/13.pdf

[2774] A. Parestu, D. R. Silaban, and K. A. Sugeng, Pelabelan simpul-ajaib total dari gabungan graf matahari, *prosising Seminar Nasional Matematika, Universitas Parahyangan, Bandung,* **3** (2008) 407-414.

[2775] A. Parestu, D. R. Silaban, and K. A. Sugeng, Vertex antimagic total labelings of the union of suns, *J. Combin. Math. Combin. Comput.,* **71** (2009) 179-188. https://combinatorialpress.com/article/jcmcc/Volume%20071/vol-071-paper%2011.pdf

[2776] J. Y. Park, J. H. Choi, and J. H. Bae, On super edge-magic labeling of some graphs, *Bull. Korean Math. Soc.,* **45** (2008) 11-21. https://www.researchgate.net/publication/238883645_On_super_edge-magic_labeling_of_some_graphs

[2777] D. Parmar and J. Joshi, H_k -Cordial labeling of triangular snake graph, *J. Appl. Sci. Computat.*, **6**(3) (2019) 2118-2123. https://www.researchgate.net/publication/331952232_HK-_Cordial_Labeling_of_Triangular_Snake_Graph

[2778] D. Parmar and U. Vaghela, Skolem difference mean labeling of some path related graphs, *Pramana Res. J.*, **9**(5) (2019) 651-657. https://www.researchgate.net/publication/333261117_Skolem_Difference_Mean_Labeling_of_Some_Path_Related_Graphs

[2779] N. A. Parmar, P. Z. Akbari, M. P. Rupani, An exploration of absolute mean graceful labeling across distinct graphs, *Internat. J. Math. Trends Tech. (IJMTT)*, **72**(4) (2026) 40-46, 2026. https://ijmttjournal.org/public/assets/volume-72/issue-4/IJMTT-V72I4P106.pdf

[2780] Y. Parmar, Edge vertex prime labeling for wheel, fan and friendship graph, *Internat. J. Math. Stat. Invention (IJMSI)*, **5**(8) (2017) 23-29. https://www.ijmsi.org/Papers/Volume.5.Issue.8/E05082329.pdf

[2781] Y. Parmar, Edge vertex prime labeling for $K_{2,n}$ and $K_{3,n}$ graphs, *Math. J. Interdisciplinary Sci.,* **6**(2) (2018) 167-180. https://core.ac.uk/download/484084709.pdf

[2782] Y. M. Parmar, M. A. Patel, M. I. Bosmia, P. L. Vihol, and D. J. Prajapati, Constructions of total neighborhood magic labeling for disjoint union of complete graphs, *J. Appl. Math. Informatics* **44**(5) (2026) 2063-2075.

[2783] A. Parthiban and N. G. David, On prime distance labeling of graphs, in: S. Arumugam, J. Bagga, L. Beineke, B. Panda, (eds.), *Th. Comput. Sci. Discr. Math.*, ICTCSDM 2016 Lecture Notes in Computer Science 10398 **31** (2017) 238-241. NOW 8/24/25

[2784] A. Parthiban and V. Sharma, A comprehensive survey on prime cordial and divisor cordial labeling of graphs, April 2020 *J. Physics Conf.,* Series 1531:012074 https://www.researchgate.net/publication/341874840_A_comprehensive_survey_on_prime_cordial_and_divisor_cordial_labeling_of_graphs

[2785] A. Parthiban and Vishally Sharma, Double divisor cordial labeling of graphs, Journal of Physics: Conference Series 2267 (2022) 012026 1-12. https://iopscience.iop.org/article/10.1088/1742-6596/2267/1/012026/pdf

[2786] A. Pasotti, Constructions for cyclic Möbius ladder systems, *Discrete Math.,* **310** (2010) 3080-3087. https://www.sciencedirect.com/science/article/pii/S0012365X09001770

[2787] A. Pasotti, On d-divisible α-labelings of $C_{4k} \times P_m$, *Util. Math.,* **90** (2013) 135-148. https://www.researchgate.net/publication/51915575_On_d-divisible_graceful_alpha-labelings_of_C_4ktimes_P_m

[2788] A. Pasotti, On d-graceful labelings, *Ars Combin.,* **111** (2013) 207-223. https://combinatorialpress.com/article/ars/Volume%20111/volume-111-paper-17.pdf

[2789] A. M. Pastel, and H. Raynaud, Numerotation gracieuse des oliviers, in *Colloq. Grenoble*, Publications Université de Grenoble, (1978) 218-223.

[2790] M. A. Patel, Y. M. Parmar, M. I. Bosmia, P. L. Vihol, H. S. Shukla, and D. M. Diwan, Total neighborhood magic labeling for complete graphs, *IAENG Internat. J. Applied Math.*, **56**(2) (2026) 769-776. https://www.iaeng.org/IJAM/issues_v56/issue_2/IJAM_56_2_28.pdf

[2791] M. J. Patel and V. Ghodasara, Cube sum labeling (taxi-cab labeling) of graphs, *Internat. J. Math. Combin.,* **2** (2023) 78-86. https://www.researchgate.net/publication/372344264_Cube_Sum_Labeling_Taxi-Cab_Labeling_of_Graphs

[2792] M. J. Patel and G. V. Ghodasara, Neighborhood-prime labeling of some cycle related graphs, *Math. Today*, **34** (2018) 159-173.

[2793] M. J. Patel and G. V. Ghodasara, Square sum labeling in context of some graph operations, *Internat. J. Math. Trends and Tech.* **66**(12) (2020), 134-147. http://ijmttjournal.org/archive/ijmtt-v66i12p519

[2794] M. J. Pate1 and G. V. Ghodasara, On permutation labeling of graphs, *Internat. J. Math. Appl.*, **9**(1) (2021) 27-33. https://ijmaa.in/index.php/ijmaa/article/view/98/98

[2795] S. K. Patel, Neighborhood-prime labeling of some generalized Petersen graphs, *Internat. J. Math. Soft Comput.,* **7**(2) (2017) 111-118.

[2796] S. K. Patel and A. N. Kansagara, Prime labeling of union graphs, *Internat. J. Sci. Research Math. Stat. Sci.*, **6**(1) (2019) 86-93. https://www.isroset.org/pub_paper/IJSRMSS/10-IJSRMSS-01696.pdf?utm_source=chatgpt.com

[2797] S. K. Patel and A. N. Kansagara, Neighborhood-prime labeling of snake graphs, *Acta Math. Universitatis Comenianae*, **90**(4) (2021) 353-376. http://www.iam.fmph.uniba.sk/amuc/ojs/index.php/amuc/article/view/1509/904

[2798] S. K. Patel, A. N. Kansagara and J. B. Vasava, Prime labeling in the context of subdivision of some cycle related graphs, *TWMS J. App. Eng. Math.*, **11**(2) (202) pp. 524-540. http://jaem.isikun.edu.tr/web/images/articles/vol.11.no.2/21.pdf

[2799] S. K. Patel, U. M. Prajapati, and A. N. Kansagara, Product cordial labeling of extensions of barbell graph, *Math. Combin. Book Ser.,* **4** (2018) 146-159. https://fs.unm.edu/IJMC/IJMC-4-2018.pdf?utm_source=chatgpt.com

[2800] S. K. Patel and N. P. Shrimali, Neighborhood-prime labeling, *Internat. J. Math. and Soft Comput.*, **5**(2) (2015) 135-143. https://oaji.net/articles/2017/296-1504965852.pdf

[2801] S. K. Patel and N. P. Shrimali, Neighborhood-prime labeling of some union graphs, *Internat. J. Math. Soft Comput.*, **6**(1) (2016) 39-47. https://oaji.net/articles/2017/296-1504967246.pdf

[2802] S. K. Patel and N. P. Shrimali, Neighborhood-prime labeling of some product graphs, *Algebra Discrete Math.*, **25**(1) (2018) 118-129. https://m.mathnet.ru/links/6e6958617c2e2a8c903252e92f896b6e/adm647.pdf

[2803] S. K. Patel and J. B. Vasava, On prime labeling of some union graphs, *Kragujevac J. Math.,* **42**(3) (2018) 441-452. https://scindeks-clanci.ceon.rs/data/pdf/1450-9628/2018/1450-96281803441P.pdf

[2804] S. K. Patel and J. B. Vasava, On prime labeling of some union graphs and circulant graphs, *Inter. J. Sci. Res. Math. Stat. Sci.*, **5**(6) (2018) 248-254. https://www.isroset.org/pdf_paper_view.php?paper_id=1000&34-IJSRMSS-01559.pdf

[2805] S. K. Patel and J. B. Vasava, Some results on (a, d)-distance antimagic labeling, *Proy. J. Math.*, **39**(2) (2020) 361-381. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/3475/3366

[2806] M. B. Paterson and D. R. Stinson, Circular external difference families, graceful labellings and cyclotomy, *Discrete Math.,* **347**(10) (2024) 114103. https://www.sciencedirect.com/science/article/pii/S0012365X24002346

[2807] H. Patodia and H. K. Saikia, m-Zumkeller graphs, *Adv. Math. Sci. J.* **9**(7) (2020) 4687-4694. https://www.research-publication.com/amsj/uploads/papers/vol-09/iss-07/AMSJ-2020-N7-35.pdf

[2808] H. Patodia and H. K. Saikia, t-m-Zumkeller labeling of graphs, *Adv. Appl. Disc. Math.*, **27**(1) (2021) 1-14. doi:10.17654/DM027010001

[2809] R. Patrias and O. Pechenik, Path-cordial abelian groups, *Australas. J. Combin.,* **80** (2021) 157-166. https://ajc.maths.uq.edu.au/pdf/80/ajc_v80_p157.pdf

[2810] L. Pattabiraman, T. Loganathan, and C. V. Rao, Graceful labeling of hexagonal snakes, Graceful Labeling of Hexagonal Snakes *AIP Confer. Proceed.* 2112, 020056 (2019) 1-3. https://doi.org/10.1063/1.5112241

[2811] B. Pattanaik, A. Behera, C. Bohidar, D. P. Bhatta, and B. K. Sethi, Graceful labeling of paths, cycles and caterpillars via inverse transformation, March 2026 Journal of Discrete Mathematical Sciences and Cryptography, **29**(3) (2026) 1441-1459. https://www.researchgate.net/publication/402078388_Graceful_labeling_of_paths_cycles_and_caterpillars_via_inverse_transformation/references

[2812] B. Pattanaik, A. K. Behera, C. Bohidar, and B. Biswal, Different classes of graceful coronas and their second kind, *J. Disc. Math. Sci. Crypt.,* **29**(8) (2026) 3037–3049. https://tarupublications.com/journals/jdmsc/volume/29/issue/8/different-classes-of-graceful-coronas-and-their-second-kind

[2813] R. Pawar and T. Singh, Distance magic labeling of generalised Mycielskian graphs, arXiv:2306.07578v1. 9pp.

[2814] O. Pechenik and J. Wise, Generalized graph cordialty, *Discuss. Math. Graph Th.,* **32**(3) (2012) 557-567. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1626

[2815] A. C. Pedrano and R. Rulete, On the total product cordial labeling on the cartesian product of $P_m \times C_n$, $C_m \times C_n$ and the generalized Petersen graph $P(m,n)$, *Malaya J. Mat.,* **5**(2) (2017) 194-201. https://www.malayajournal.org/articles/Paper_7_2017.pdf

[2816] A. Pedrano and R. Rulete, On total product cordial labeling of some crown graphs, *Internat. J. Math. Appl.*, **5**(2B) (2017) 273-284. https://www.researchgate.net/publication/317596528_On_Total_Product_Cordial_Labeling_of_Some_Crown_Graphs

[2817] A. C. Pedrano and R. F. Rulete, On the total product cordial labeling on the cartesian product of $P_m \times C_n, C_m \times C_n$ and the generalized Petersen graph $P(m,n)$, *Malaya J. Mat.*, **5**(3) (2017) 531-539. https://www.researchgate.net/publication/317913334

[2818] A. C. Pedrano, and E. R. Salise Jr., On Lucas cordial labeling of some snake graphs, https://www.technoskypub.com/wp-content/uploads/2025/12/3-P3-V8N4-On-Lucas-Cordial-Labeling.pdf

[2819] E. Pegg Jr., personnal communication, 2021

[2820] S. G. Penrice, Some new graph labeling problems: a preliminary report, DIMACS Tech. Rep. 95-29 (1995). https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=b59595bf0d2f2b7173b80ef13b227f462757c2a9

[2821] J. Pereira, T. Singh, and S. Arumugam, On (k, d)-Skolem graceful graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), 81-88, *Electron. Notes Discrete Math.*, **48**, Elsevier Sci. B. V., Amsterdam, 2015. https://arxiv.org/pdf/1705.06736v1

[2822] J. Pereira, T. Singh, and S. Arumugam, m-gracefulness of graphs, *Algor. Discrete Appl. Math., Lecture Notes in Comput. Sci.*, 9602, Springer, [Cham], (2016) 289-298.

[2823] J. Pereira, T. Singh, and S. Arumugam, A new measure for gracefulness of graphs, *Elect. Notes Discr. Math.*, **48** (2015) 275-280.

[2824] J. Pereira, T. Singh, and S. Arumugam, On (k, d)-hooked Skolem graceful graphs, *Elect. Notes Discr. Math.*, **48** (2015) 81-88. https://doi.org/10.1016/j.endm.2015.05.012 arXiv:1705.06736v1

[2825] J. Pereira, T. Singh, and S. Arumugam, Additively graceful signed graphs, *AKCE Internat. J. Graphs Combin.* **20** (3) (2023) 300-307. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2023.2243619

[2826] K. Periasamy, K. Venugopal, and P. L. Rozario Raj, kth *Fibonacci prime labeling of graphs Internat. J. Math. Trends Tech.,* **68**(5) (2022) 61-67. https://www.researchgate.net/publication/361207921_K_th_Fibonacci_Prime_Labeling_of_Graphs/references#fullTextFileContent

[2827] J. Periaswamy, Root square mean labeling of uncoordinated graphs, *Neuroquantology*, 2022 **20**(11) (2022) 6290-6296. https://www.neuroquantology.com/media/article_pdfs/6290-6296.pdf

[2828] M. A. Perumal, S. Navaneethakrishnan, S. Arockiaraj, A. Nagarajan, Super graceful labeling for some special graphs, *Int. J. Res. Rev. Appl. Sci.* **9**(3) (2011) 382-404.

[2829] M. A. Perumal, S. Navaneethakrishnan, A. Nagarajan, S. Arockiaraj, Super graceful labeling for some simple graphs, *Internat. J. Math. Soft Computing,* **2** (2012) 35-49. https://oaji.net/articles/2017/296-1502655319.pdf

[2830] K. Petrie and B. Smith, Symmetry breaking in graceful graphs, in *Proc.CP'03, LNCS,* Springer (2003) 930-934. https://www.researchgate.net/publication/225127936_Symmetry_Breaking_in_Graceful_Graphs

[2831] F. Pfender, Total edge irregularity strength of large graphs, *Discrete Math.,* **312**(2) 229-237. arXiv:1006.4501

[2832] O. Phanalasy, Antimagic labeling of generalized sausage graphs, *J. Indones. Math. Soc.,* **20**(2) (2014) 95-110. https://media.neliti.com/media/publications/94307-EN-antimagic-labeling-of-generalized-sausag.pdf

[2833] O. Phanalasy, M. Miller, C. S. Iliopoulos, S. P. Pissis, and E. Vaezpour, Construction of antimagic labeling for the Cartesian product of regular graphs, *Math. in Comput. Sci.*, **5**(1) (2011) 81-87. https://www.academia.edu/62629909/Construction_of_antimagic_labeling_for_the_Cartesian_product_of_regular_graphs_NOVA_The_University_of_Newcastles_Digital_Repository?uc-g-sw=52813124

[2834] O. Phanalasy, M. Miller, J. Ryan and S. Arumugam, Antimagic labeling of generalized pyramid graphs, *Acta. Math. Sin.-English Ser.,* **30** (2014) 283-290. https://actamath.cjoe.ac.cn/Jwk_sxxb_en/EN/abstract/abstract22175.shtml

[2835] O. Phanalasy, M. Miller, L. J. Rylands, and P. Lieby, On a relationship between completely separating systems and antimagic labeling of regular graphs, In C. S. Iliopoulos and W. F. Smyth, editors, *Proceedings of IWOCA10, Lecture Notes in Computer Science*, **6460** (2011) 238-241.

[2836] N. C. K. Phillips, R. S. Rees, and W. D. Wallis, Edge-magic total labelings of wheels, *Bull. Inst. Combin. Appl.*, **31** (2001) 21-30.

[2837] N. C. K. Phillips, R. S. Rees, and W. D. Wallis, personal communication (circa 2002).

[2838] D. A. Pike, A. Sanaei and N. Shalaby, Pseudo-Skolem sequences and graph Skolem labelling, *Math. Scandinavica*, **120** (2017) 17-38. https://www.mscand.dk/article/view/25502/22451

[2839] O. Pikhurko, Every tree with at most 34 vertices is prime, *Util. Math.*, **62** (2002) 185-190. https://pikhurko.github.io/E/Pikhurko02um.pdf

[2840] O. Pikhurko, Dense edge-magic graphs and thin additive bases, *Discrete Math.,* **306** (2006) 2097-2107. https://pikhurko.github.io/E/Pikhurko06dm.pdf

[2841] O. Pikhurko, Trees are almost prime, *Discrete Math.,* **307** (2007) 1455-1462. https://pikhurko.github.io/E/Pikhurko07dm.pdf

[2842] O. Pikhurko, Characterization of product anti-magic graphs of large order, *Graphs and Combin.,* **23** (2007) 681-689. https://pikhurko.github.io/E/Pikhurko07gc.pdf?utm_source=chatgpt.com

[2843] D. Pipaliya1 and M. Chaurasia1, Mean labeling of graphs associated with paths and cycles, *Int. J. Math. Appl.*, **14**(2)(2026) 215–223. https://ijmaa.in/index.php/ijmaa/article/view/1712/1567

[2844] A. A. Pir and A. Parathiban, Further results on 2-odd labelings of graphs, *Adv. Appl. Math. Sci.*, **21**(8) (2022) 4265-4171. https://www.mililink.com/upload/article/2137325821aams_vol_218_june_2022_a5_p4265-4271_ajaz_ahmad_pirand_and_a._parthiban.pdf

[2845] S. Poljak and M. Sûra, An algorithm for graceful labeling of a class of symmetrical trees, *Ars Combin.*, **14** (1982) 57-66.

[2846] A. Ponmoni, S. Navaneetha Krishnan and A. Nagarajan, Skolem difference Lucas mean labeling for star related graphs, *Internat. J. Math. Trends Tech.,* **62**(2) (2018) 104-109. http://www.ijmttjournal.org/archive/ijmtt-v62p515

[2847] C. Y. Ponnappan, CT. Nagaraj, and G. Prabakaran, Odd vertex magic labeling, *Internat. J. Comput. Organization Trends,* **9**(1) (2014) 86-89. https://pdfs.semanticscholar.org/88c4/f87f384255dddab2fca95e4fc0644424f8cc.pdf

[2848] R. Ponraj, $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial labeling of graphs, *Varahmihir J. Math. Sci.,* **8** (2008) 137-145.

[2849] R. Ponraj, Further results on $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial labeling of graphs, *J. Indian Acad. Math.,* **31** (2009) 157-163.

[2850] R. Ponraj and M. Maria Adaickalam, Quotient cordial labeling of star related graphs, *J. Indian Acad. Math.,* **37**(2)(2015) 313-324.

[2851] R. Ponraj and M. Maria Adaickalam, 3-difference cordiality of some graphs, *Palestine. J. Mathematics*, **6**(II) (2017) 141-148. https://pjm.ppu.edu/sites/default/files/papers/PJM_JUNE_2017_6.pdf

[2852] R. Ponraj and M. Maria Adaickalam, 3-difference cordiality of some union of graphs, *Palestine J. Math.*, **2** (2017) 202-218. https://www.researchgate.net/publication/313859755_3-Difference_cordial_labeling_of_some_union_of_graphs

[2853] R. Ponraj and M. Maria Adaickalam, 3-difference cordial labeling of some cycle related graphs, *J. Algor. Comput.*, **47** (2016) 1-10. journals.ut.ac.ir/article_7927_2058a8520b6be55a66b6d88cf1ff676f.pdf

[2854] R. Ponraj, M. Maria Adaickalam, and R. Kala, 3-difference cordiality of some special graphs, *Jordan J. Math. Stat.,* **11**(2) (2018) 143-166. https://jjms.yu.edu.jo/index.php/jjms/article/view/975/388

[2855] R. Ponraj, M. Maria Adaickalam and R. Kala, Quotient cordial labeling of graphs, *Inter. J. Math. Combin.,* **1** (2016) 101-108. https://vixra.org/pdf/1604.0183v1.pdf

[2856] R. Ponraj, M. Maria Adaickalam and R. Kala, k-difference cordial labeling of graphs, *Internat. J. Math. Comb.*, **2** (2016) 121-131. https://vixra.org/pdf/1607.0542v1.pdf

[2857] R. Ponraj, M. Adaickalam, and R. Kala, 3-difference cordiality of corona of double alternate snake graph, *Bull. Int. Math. Virtual Inst.,* **8**(2) (2018) 245-258. http://www.imvibl.org/buletin/bulletin_imvi_8_2_2018/bulletin_imvi_8_2_2018_245_258.pdf

[2858] R. Ponraj, K. Annathurai, and R. Kala, Remainder cordial labeling of graphs, *J. Algor. Comput.*, **49** (2017) 17-30. https://jac.ut.ac.ir/article_7965_9bf96b3c6575b2e98518384f18e0de66.pdf

[2859] R. Ponraj, K. Annathurai, and R. Kala, k-remainder cordial graphs, *J. Algor. Comput.,* **49**(2) (2017) 41-52. https://jac.ut.ac.ir/article_7976_cc187cb94e0d46cf3e46d78a35564977.pdf

[2860] R. Ponraj, K. Annathurai, and R. Kala, 4-remainder cordial labeling of some special graphs, *Internat. J. Pure Appl. Math.*, **118**(6) (2018) 399-405. https://acadpubl.eu/jsi/2018-118-6/articles/6/44.pdf

[2861] R. Ponraj, K. Annathurai, and R. Kala, 4-remainder cordial labeling of some graphs, *Internat. J. Math. Combin.*, **1** (2018) 138-145. https://fs.unm.edu/IJMC/4RemainderCordialLabelingOfSomeGraphs.pdf

[2862] R. Ponraj, K. Annathurai, and R. Kala, 4-reminder cordial labeling graphs obtained from ladder, *Internat.l J. Mathematical Combin.,* personal communication.

[2863] R. Ponraj, K. Annathurai, and R. Kala, Remainder cordiality of some graphs, *Palestine J. Math.*, **8**(1) (2019) 367-372. https://pjm.ppu.edu/sites/default/files/papers/PJM_Oct2018_367to372.pdf

[2864] R. Ponraj, K. Annathurai, and R. Kala, 4-remainder cordial labeling of some new families graphs, *Palestine J. Math.*, **8**(2) (2019) 78-86. https://pjm.ppu.edu/sites/default/files/papers/PJM_April2019_78to86.pdf

[2865] R. Ponraj, K. Annathurai, and R. Kala, Some results on remainder cordial labeling of graph, *Ars Combin.*, **162** (2025) 39-49. https://combinatorialpress.com/article/ars/Volume%20162/some-results-on-4-remainder-cordial-labeling-of-graphs.pdf

[2866] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of certain subdivision graphs, *J. Appl. Math. & Informatics Eng. Math.*, **42**(1) (2024) 1-14. https://www.koreascience.kr/article/JAKO202410743214826.pdf

[2867] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of some product related graphs, *TWMS J. App. Eng. Math.*, **14**(4) (2024) 1676-1688. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.4/28.pdf

[2868] R. Ponra and A. Gayathri, Pair difference cordial number of some degree splitting graph, *International J. Mathematical Combinatorics*, **3** (2024) 99-107. https://fs.unm.edu/IJMC/Pair_Difference_Cordial_Number_of_Some_Degree_Splitting_Graph.pdf

[2869] R. Ponraj and A. Gayathri, New families of pair difference cordial graphs, *J. Algor. Comput.,* **56**(2) (2024) 113-122. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.4/28.pdf

[2870] R. Ponraj, A. Gayathri, and M. Sivakumar, Pair difference cordial labeling of m-copies of path, cycle, star and ladder graphs,*J. Algor. Comput.*, **54**(2) (2022) 37-47. https://jac.ut.ac.ir/article_90409_bb72c11885a76aab1d04bf9e4bae4ad1.pdf

[2871] R. Ponraj, A. Gayathri, and M. Sivakumar, Pair difference cordial labeling of subdivision of wheel and comb graph, *Internat. J. Math. Combin.*, **1** (2023) 116-125. https://fs.unm.edu/IJMC/Pair_Difference_Cordial_Labeling_of_Subdivision_of_Wheel_and_Comb_Graphs.pdf

[2872] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of some snake and butterfly graphs, *J. Algor. Comput.*, **53**(1) (2021) 149-163. https://jac.ut.ac.ir/article_81649_9670058e7708586f959c57de1e3434f5.pdf

[2873] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial graphs obtained from the wheels and the paths, *J. Appl. Pure Math.*, **3**(3-4) (2021) 97-114.

[2874] R. Ponraj, A. Gayathri, and S. Somasundaram, Some pair difference cordial graphs, *Ikonion J. Math..*, **3**(2) (2021) 17-26. https://dergipark.org.tr/en/download/article-file/1729055

[2875] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of graphs, *J. Math. Comp. Sci.***11**(3) (2021) 2551-2567. https://scik.org/index.php/jmcs/article/view/5601

[2876] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of some union of graphs, *TWMS J. App. Eng. Math.*, **13**(3) (2023) 1083-1095. https://jaem.isikun.edu.tr/web/images/articles/vol.13.no.3/23.pdf

[2877] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of some graphs derived from ladder graph, *J. Math. Comp. Sci.*, **11**(5) (2021) 6105-6124. https://scik.org/index.php/jmcs/article/view/6132

[2878] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of planar grid and mongolian tent, *J. Algor. Comput.*, **53**(2) (2021) 47-56. https://jac.ut.ac.ir/article_85484_e3023a85223c575225e1ae0d841c5933.pdf

[2879] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of mirror graph, shadow graph and splitting graph of Certain graphs, *Maltepe J. Math.*, **4**(1) (2022) 24-32.

[2880] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of some star related graphs,*J. Mehani Math. Res.*, **12**(2), (2023), 255-266. https://jmmrc.uk.ac.ir/article_3470_0c80908562505a9c0fe2abd9f3338bac.pdf

[2881] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of $m-$ copies of some graphs, *J. Appl. Pure Math.*, **4** (5-6) (2022) 317-329. https://koreascience.kr/article/JAKO202206757623002.pdf

[2882] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of Franklin graph, Heawood graph, Tietze graph and Durer graph, *Internat. J. Math. Combin.*, **4** (2022) 74-81. https://fs.unm.edu/IJMC/Pair_Difference_Cordial_Labeling_of_Franklin_Graph,%20Heawood_Graph,%20Tietze_Graph_and_Durer_Graph.pdf

[2883] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of some trees and some graphs derived from cube graph, *Internat. J. Math. Combin.*, **2** (2023) 65-73. https://fs.unm.edu/IJMC/Pair_Difference_Cordial_Labeling_of_Some_Trees_and_Some_Graphs_Derived_From_Cube_Graph.pdf

[2884] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of certain broken wheel graphs, *J. Universal Math.*, **6**(1) (2023) 39-48. https://dergipark.org.tr/en/download/article-file/2404828

[2885] R. Ponraj, A. Gayathri, and S. Somasundaram, 4-remainder cordial labeling of parachute, twig, kayak paddle graph, *Palest. J. Math.* **11**(2) (2022) 26-37. https://pjm.ppu.edu/sites/default/files/papers/PJM_May_2022_26_to_37.pdf

[2886] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordial labeling of Petersen graphs $p(n,k)$, *J. Appl. Pure Math.*, **5**(1-2) (2023) 41-53. https://koreascience.kr/article/JAKO202315154844634.pdf

[2887] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of certain subdivision graphs, *J. Appl. Math. & Informatics Eng. Math.*, **42**(1) (2024) 1-14. https://www.koreascience.kr/article/JAKO202410743214826.pdf

[2888] R. Ponraj, A. Gayathri, and S. Somasundaram, Pair difference cordiality of some product related graphs, *TWMS J. App. Eng. Math.*, **14**(4) (2024) 1676-1688. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.4/28.pdf

[2889] R. Ponraj and R. Jeya, Vector basis S-cordial labeling of graphs, *J. Math. Comput. Sci.*, **15:5**(1 (2025) 1-13. https://scik.org/index.php/jmcs/article/view/9116

[2890] R. Ponraj and R. Jeya, Certain VB $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$-cordial thorn graphs, *Global J. Pure and App. Math.*, **21**(1) (2025) 1-14. https://www.ripublication.com/gjpam23/gjpamv21n1_01.pdf

[2891] R. Ponraj and R. Jeya, Vector basis $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$-cordial labeling of generalized friendship graph, tadpole graph and gear graph, *Global J. Pure and App. Math.*, **21**(1) (2025) 81-94. https://www.ripublication.com/gjpam23/gjpamv21n1_06.pdf

[2892] R. Ponraj, and R. Jeya, Vector basis $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$-cordial labeling of $D(T_n)\odot mK_1$ and $D(Q_n)\odot mK_1$, *Internat. J. Math. Appl.*, **3**(2) (2025) 57-69. https://ijmaa.in/index.php/ijmaa/article/view/1565/1479

[2893] R. Ponraj, and R. Jeya, Vector basis S-cordial labeling of friendship graph, fan graph and lilly graph, *J. Indian Acad. Math.*, **47**(1) (2025) 169-180. https://www.researchgate.net/publication/395357993_11_MSS_2300

[2894] R.Ponraj and R. Jeya, Vector basis S-cordial graph, *Internat. J. Environmental Sci.*, **11**(18s) (2025) 1539-1551. https://theaspd.com/index.php/ijes/article/view/4369/3189

[2895] R. Ponraj and R. Jeya, Vector basis $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$-cordial labeling of certain trees and flower graphs, *Algebra Letters*, **2025**(2) (2025) 1-16. https://www.scik.org/index.php/abl/article/viewFile/9243/4304

[2896] R. Ponraj and R. Jeya, Vector basis $\{(1,1,1,1),(1,1,1,0),(1,1,0,0),(1,0,0,0)\}$-cordial labeling of $L_n \odot mK_1$ and $T(P_n) \odot mK_1$, *J. Algor. Comput.*, **57**(1) (2025) 30-40. https://journals.ut.ac.ir/article_103143_3332a4ed8ea14929da0255ba6adf17a0.pdf

[2897] R. Ponraj and R. Jeya, Vector basis S-cordial labeling of graphs, *J. Math. Comput. Sci.*, **15** (2025) ID5. https://doi.org/10.28919/jmcs/9116

[2898] R. Ponraj and R. Jeya, Vector basis S-cordial labeling of balloon of triangular snake graph, *J. Appl. Pure Math.*, **8**(1-2) (2026) 89-99.

[2899] R. Ponraj and R. Jeya, Vector basis S-cordial labeling of graphs with applications in biological models, *Commun. Math. Biology Neurosci.* (2026) https://doi.org/10.28919/cmbn/9614

[2900] R. Ponraj, S. Prabhu, and A. M. S. Ramasamy, PMC-labeling of certain classes of graphs, *J. Fund. Math. Appl.* (JFMA) **8**(2) (2025) 175-189. https://ejournal2.undip.ac.id/index.php/jfma/article/view/26263/13513

[2901] R. Ponraj and J. V. X. Parthipan, Pair sum labeling of graphs, *J. Indian Acad. Math.,* **32**(2) (2010) 587-595. https://www.researchgate.net/publication/268494646_Pair_Sum_Labeling_of_Graphs

[2902] R. Ponraj, J. V. X. Parthipan, and R. Kala, Some results on pair sum labeling of graphs, *Internat. J. Math. Combin.*. **4** (2010) 55-61. https://www.researchgate.net/publication/267129811_Some_results_on_pair_sum_labeling_of_graphs

[2903] R. Ponraj and J. V. X. Parthipan, Further results on pair sum labeling of trees, *Appl. Math.*, **2**(10) (2011) 1270-1278. https://www.scirp.org/pdf/AM20111000011_22898282.pdf

[2904] R. Ponraj and J. V. X. Parthipan, Pair mean labeling of graphs, *J. Indian Acad. Math.,* **33**(2) (2011) 375-383. https://www.researchgate.net/publication/267655602_Pair_mean_labeling_of_graphs

[2905] R. Ponraj, J. V. X. Parthipan, and R. Kala, Some results on pair sum labeling of graphs, *Inter. J. Mathematical Combin.,* **4** (2010) 53-61. https://fs.unm.edu/IJMC/SomeResultsOnPairSumLabeling.pdf

[2906] R. Ponraj, J. V. X. Parthipan, and R. Kala, A note on pair sum graphs, *J. Scientific Research*, **3**(2) (2011) 321-329. https://www.researchgate.net/publication/270111862_A_Note_on_Pair_Sum_Graphs

[2907] R. Ponraj, J. V. X. Parthipan and R. Kala, Further results on pair sum graphs, *Appl. Math.,* **3**(3) (2012) 267-275. https://www.scirp.org/journal/PaperInformation?PaperID=18106

[2908] R. Ponraj, J. V. X. Parthipan and R. Kala, Some new families of pair sum graphs, *J. Math. Comput. Sci.,* **2**(3) (2012) 747-758. https://scik.org/index.php/jmcs/article/view/175

[2909] R. Ponraj and S. Prabhu, Pair mean cordiality of some snake graphs, *Global J. Pure Apl. Math.*, **18**(1) (2022) 283-295.

[2910] R. Ponraj and S. Prabhu, Pair mean cordial labeling of graphs, *J. Algor. Comput.*, **54**(1) (2022) 1-10. https://jac.ut.ac.ir/article_87392_1826c7cf2b60089d62aa5b172956101a.pdf

[2911] R. Ponraj and S. Prabhu, Pair mean cordial labeling of graphs obtained from path and cycle, *J. Appl. Pure Math.* **4**(3-4) (2022) 85-97. https://koreascience.or.kr/article/JAKO202222357137062.page

[2912] R. Ponraj and S. Prabhu, Pair mean cordial labeling of some corona graphs, *J. Indian Acad. Math.*,**44**(1) (2022) 45-54.

[2913] R. Ponraj and S. Prabhu, On pair mean cordial graphs, *J. Appl. Pure Math.*, **5**(3-4) (2023) 237-253. https://koreascience.or.kr/article/JAKO202323459641777.pdf

[2914] R. Ponraj and S. Prabhu, Pair mean cordial labeling of some union of graphs, *J. Appl. and Pure Math.*, **6**(1-2) (2024) 55-69.

[2915] R. Ponraj and S. Prabhu, Pair mean cordial labeling of hexagonal snake, irregular quadrilateral snake and triple triangular snake, *Indian J. Nat. Sci.*, **15)(83** (2024) 73075-73081.

[2916] R. Ponraj and S. Prabhu, Pair mean cordial graphs paired with ladder, *Internat. J. Math. Combin.*, **3** (2024) 49-64. https://fs.unm.edu/IJMC/Pair_Mean_Cordial_Graphs_Paired_with_Ladder.pdf

[2917] R. Ponraj and S. Prabhu, Pair mean cordial labeling of hurdle, key, lotus and necklace graphs, *BAREKENG J. Math. Appl.*, **18**(4) (2024) 2795-2804. https://doaj.org/article/d51831d94d484401b5ae4910c22ee894?utm_source=chatgpt.com

[2918] R. Ponraj, S. Prabhu, and M. Sivakumar, Pair mean cordial labeling of total graph of some graphs and prism related graphs, *Comm. Combin. Cryptography & Comput. Sci.*, **02** (2024)181-192.

[2919] R. Ponraj and S. Prabhu, Pair mean cordiality of certain ladder graphs, *J. Appl. & Pure Math.*, **6**(5-6) (2024) 327-336.

[2920] R. Ponraj and S. Prabhu, Further results on pair mean cordial graphs, *TWMS J. App. Eng. Math.*, **15**(2) (2025) 431-442 https://jaem.isikun.edu.tr/web/images/articles/vol.15.no.2/16.pdf?utm_source=chatgpt.com

[2921] R. Ponraj, S. Prabhu, and M. Sivakumar, PMC-labeling of some classes of graphs containing cycles, *BAREKENG J. Math. Appl.*, **19**(2) (2025) 1445-1456. https://ojs3.unpatti.ac.id/index.php/barekeng/article/view/16053/9986

[2922] R. Ponraj, and S. Prabhu, Pair mean cordial labeling of diamond snake graph, banana tree and tortoise graph, *TWMS J. App. Eng. Math.*, **15**(5) (2025) 1190-1199. https://jaem.isikun.edu.tr/web/images/articles/vol.15.no.5/12.pdf?utm_source=chatgpt.com

[2923] R. Ponraj, S. Prabhu, and M. Sivakumar, PMC-labeling of certain tree related graphs and prism of wheel graph, *Internat. J. Math. Combin.*, (2025) 1–2 (2025) 93–102. https://fs.unm.edu/IJMC/PMC-Labeling_of_Certain_Tree_Related_Graphs_and_Prism_of_Wheel_Graph.pdf?utm_source=chatgpt.com

[2924] R. Ponraj, and S. Prabhu, Pair mean cordial labeling of udukkai, octopus, drum and fire cracker graphs, *Disc. Math., Algor. Appl.* **17**(6) (2025) 2450096 12 pp. https://www.worldscientific.com/doi/10.1142/S1793830924500964

[2925] R. Ponraj and S. Prabhu, New results on pair mean cordial labeling of certain graph classes, *Open Disc. Appl. Math.*, (ODAM) **9**(1) (2026) 46-70. https://doi.org/10.30538/psrp-odam2026.0133

[2926] R. Ponraj, S. Prabhu and A. M. S. Ramasamy, PMC-graphs derived from cycles, *Bol. Soc. Paran. Mat.*, **43**(3s) (2025) 1-11. https://periodicos.uem.br/ojs/index.php/BSocParanMat/article/view/76332/751375160334

[2927] R. Ponraj, S. Prabhu, and M. Sivakumar, PMC-labeling of some classes of graphs containing cycles, *BAREKENG: J. Math. App.*, **19**(2) (2025) 1445-1456. https://ojs3.unpatti.ac.id/index.php/barekeng/article/view/16053/9986

[2928] R. Ponraj, Rajpal Singh and S. Sathish Narayanan, 4-Prime cordiality of some classes of graphs, *J. Algor. Comput.,* (2016) **48** (1) 69-79. https://jac.ut.ac.ir/article_7941_01050790c89557f0425ee26385c3b6b9.pdf

[2929] R. Ponraj, Rajpal Singh and S. Sathish Narayanan, 4-Prime cordiality of some cycle related graphs, *Appl. Appled Math.,: An Internat. J.,* (AAM) **12**(1) (2017) 230-240. https://www.pvamu.edu/aam/wp-content/uploads/sites/182/2017/06/15_R916_AAM_Ponraj_RP_061816_Posted_061217_pp_230_240.pdf

[2930] R. Ponraj, Rajpal Singh and R. Kala, Some more 4-prime cordial graphs, *Internat. J. Math. Combin.,* **2** (2017) 105-115. https://fs.unm.edu/IJMC/SomeMore4PrimeCordialGraphs.pdf

[2931] R. Ponraj, Rajpal Singh and R. Kala, Some new families of 4-prime cordial graphs, *Internat. J. Math. Combin.,* **2** (2018) 125-135. https://www.researchgate.net/publication/267433635_Some_New_Families_of_Prime_Cordial_Graphs

[2932] R. Ponraj and Rajpal Singh, 4-Prime cordial labeling Of some triangular snake graphs and diamond snake graphs, *Math. Sci. Internat. Res. J.,* **7** (2018) 305-308.

[2933] R. Ponraj, Rajpal Singh, and R. Kala, 4-Prime cordiality of some cycle and star related graphs, *Internat. J. Pure Appl. Math.,* **118**(6)(2018) 235-241. https://www.acadpubl.eu/jsi/2018-118-6/articles/6/26.pdf

[2934] R. Ponraj, and Rajpal Singh, and R. Kala 3-Prime cordiality of some cycle related special graphs, *Global J. Engin. Sci. Res.,* **5**(7) (2018) 98-104. https://www.gjesr.com/Issues%20PDF/Archive-2018/June-2018/15.pdf

[2935] R. Ponraj, Rajpal Singh and R. Kala, 4-Prime oordial labeling of some degree splitting graphs, *Internat. J. Management, IT Engin.,* **6** (6)(2018) 227-236. https://www.ijmra.us/project%20doc/2018/IJMIE_JULY2018/IJMRA-14100.pdf

[2936] R. Ponraj, S. Prabhu, and M. Sivakumar, Pair mean cordial labeling of pizza graph, snail graph, sun graph and kusudama flower graph, *Indian J. Nat. Sci.*, **16**(89) (2025) 91844-91851.

[2937] R. Ponraj and S. Sathish Narayanan, Further results on difference cordial labeling of corona graphs, *J. Indian Academy Math.,* **35** (2013) 217-235.

[2938] R. Ponraj and S. Sathish Narayanan, Difference cordiality of some graphs obtained from double alternate snake graphs, *Global J. Math. Sciences: Theory and Practical,* **5** (2013) 167-175. http://www.irphouse.com/gjms/gjmsv5n3_04.pdf

[2939] R. Ponraj and S. Sathish Narayanan, Difference cordiality of some snake graphs, *J. Appl. Math. and Informatics*, **32** (2014) 377-387. https://koreascience.kr/article/JAKO201416747605232.pdf

[2940] R. Ponraj and S. Sathish Narayanan, Difference cordiality of some derived graphs, *Internat. J. Math. Combin.*, **4** (2013) 37-48. https://fs.unm.edu/IJMC/DifferenceCordialityOfSomeDerivedGraphs.pdf

[2941] R. Ponraj and S. Sathish Narayanan, Difference cordial labeling of graphs obtained from triangular snakes, *Applications Applied Math.*, **9**(2) (2014) 811-825. https://digitalcommons.pvamu.edu/cgi/viewcontent.cgi?article=1403&context=aam

[2942] R. Ponraj and S. Sathish Narayanan, Total mean cordiality of $K_n^c + 2K_2$, *Palestine J. Math.,* **4**(2) (2015) 431-438. https://pjm.ppu.edu/sites/default/files/papers/26%20%20Total%20Mean%20Cordiality%20of.pdf

[2943] R. Ponraj and S. Sathish Narayanan, Mean cordiality of some snake graphs, *Palestine J. Math.,* **4**(2) (2015) 439-445. https://pjm.ppu.edu/sites/default/files/papers/27%20%20Mean%20cordiality%20of%20some%20snake%20graphs.pdf

[2944] R. Ponraj and S. Sathish Narayanan, Total mean cordial labeling of some cycle related graphs, *J. Appl. Math. & Informatics*, **33** (1-2) (2015) 101-110. https://www.researchgate.net/publication/276395672_TOTAL_MEAN_CORDIAL_LABELING_OF_SOME_CYCLE_RELATED_GRAPHS

[2945] R. Ponraj and S. Sathish Narayanan, Some results on total mean cordial labeling of graphs, *International J. Math. Combin.*, **2** (2015) 122-132. https://fs.unm.edu/IJMC/SomeResultsOnTotalMeanCordialLabeling.pdf

[2946] R. Ponraj and S. Sathish Narayanan, Existence and non existence of mean cordial labeling of certain graphs, *J. Prime Research Math.*, **11** (2015) 123-136. https://www.researchgate.net/publication/302383520_Existence_and_Nonexistence_of_mean_cordial_labeling_of_graphs

[2947] R. Ponraj and S. Sathish Narayanan, Total mean cordiality of some derived graphs, *Bull. inter. Math. Virtual Institute*, **5** (2015) 181-190. http://www.imvibl.org/buletin/bulletin_imvi_5_2_2015/bulletin_imvi_5_1_2015_181_190.pdf

[2948] R. Ponraj and S. Sathish Narayanan, Further results on total mean cordial labeling of graphs, *J. Algor. Comput.*, **46** (2015) 73-83. https://jac.ut.ac.ir/article_7926_9d2173db725a3759d46d6f1e33486b61.pdf

[2949] R. Ponraj and S. Sathish Narayanan, Radio mean number of certain graphs, *Inter. J. Math. Combin.,* **2** (2016) 51-64. https://fs.unm.edu/IJMC/RadioMeanNumberOfCertainGraphs.pdf

[2950] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordial labeling of graphs, *Global J. Math. Sciences: Theory and Practical,* **5**(3) (2013) 185-196. https://www.researchgate.net/publication/302385037_Difference_Cordial_Labeling_of_Graphs

[2951] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordial labeling of graphs obtained from double snakes, *Internat. J. of Math. Research,* **5** (2013) 317-322. http://www.irphouse.com/ijmr/ijmrv5n3_03.pdf

[2952] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordial labeling of corona graphs, *J. Math. Comput. Sci.*, **3** (2013) 1237-1251. https://www.researchgate.net/publication/302384278_DIFFERENCE_CORDIAL_LABELING_OF_CORONA_GRAPHS

[2953] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordial labeling of subdivided graphs, *Scientia Magna,* **9**(3) (2013) 57-66. https://www.thefreelibrary.com/Difference+cordial+labeling+of+subdivided+graphs.-a0370395380

[2954] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordial labeling of subdivision of snake graphs, *Universal J. Appl. Math.*, **2**(1) (2014) 40-45. https://www.academia.edu/105681635/Difference_Cordial_Labeling_of_Subdivision_of_Snake_Graphs

[2955] R. Ponraj, S. Sathish Narayanan, and R. Kala, Difference cordiality of product related graphs, *Tbilisi Math. J.*, **8**(2) (2015) 41-47. https://projecteuclid.org/journals/advanced-studies-euro-tbilisi-mathematical-journal/volume-8/issue-2/Difference-cordiality-of-product-related-graphs/10.1515/tmj-2015-0009.full

[2956] R. Ponraj, S. Sathish Narayanan, and R. Kala, A note on difference cordial graphs, *Palestine J. Math.,* **4**(1) (2015) 189-197. https://pjm.ppu.edu/sites/default/files/papers/23_0.pdf?utm_source=chatgpt.com

[2957] R. Ponraj, S. Sathish Narayanan, and R. Kala, On radio mean number of some graphs, *Internat. J. Math. Combin.*, **3** (2014) 41-48. https://fs.unm.edu/IJMC/OnRadioMeanNumberOfSomeGraphs.pdf

[2958] R. Ponraj, S. Sathish Narayanan, and R. Kala, Radio mean labeling of a graph, *AKCE Internat. J. Graphs Combin.*, **12** (2015) 224-228. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2015.11.019#d1e381

[2959] R. Ponraj, S. Sathish Narayanan, and R. Kala, Radio mean number of some wheel related graphs, *Jordan J. Math. and Stat. (JJMS),* **7**(4) (2014) 273-286. https://jjms.yu.edu.jo/index.php/jjms/article/view/1074/465

[2960] R. Ponraj, S. Sathish Narayanan and A. M. S. Ramasamy, Total mean cordial labeling of graphs, *International J.Math. Combin.*, **4** (2014) 56-68. https://fs.unm.edu/IJMC/TotalMeanCordialLabelingOfGraphs.pdf

[2961] R. Ponraj, S. Sathish Narayanan and A. M. S. Ramasamy, Total mean cordial labeling of some graphs, *Util. Math.*, **106** (2018) 209–221.

[2962] R. Ponraj, S. Sathish Narayanan and A. M. S. Ramasamy, Total mean cordiality of umbrella, butterfly and dumbbell graphs, *Jordan J. Math. and Stat. (JJMS)*, **8**(1)(2015) 59-77. https://jjms.yu.edu.jo/index.php/jjms/article/view/1071/462

[2963] R. Ponraj, S. Sathish Narayanan, and R. Kala, Radio mean labeling of a graph *AKCE Internat. J. Graphs Combin.,* **12** (2015) 224-228. https://www.researchgate.net/publication/287374769_Radio_mean_labeling_of_a_graph

[2964] R. Ponraj and Rajpal Singh, 4-Prime cordial graphs obtained from 4-prime cordial graphs, *Bull. Int. Math. Virtual Inst.* **8**(1) (2018) 1-9. http://www.imvibl.org/buletin/bulletin_imvi_8_1_2018/bulletin_imvi_8_1_2017_1_9.pdf

[2965] R. Ponraj, Rajpal Singh, R. Kala and S. Sathish Narayanan, k-Prime cordial graphs, *J. Appl. Math. & Informatics*, **34**(3-4) (2016) 227-237. https://www.researchgate.net/publication/305763538_k-PRIME_CORDIAL_GRAPHS

[2966] R. Ponraj, Rajpal Singh, and S. Sathish Narayanan, A note on 3-prime cordial graphs, *J. Algor. Comput.,* **48** (2016) 45-55. https://journals.ut.ac.ir/article_7939_495c61efe2289c038edc6c7099386507.pdf

[2967] R. Ponraj, Rajpal Singh, and R. Kala, 4-Prime cordial labeling of some special graphs, *Bull. Int. Math. Virtual Inst.* **8**(1) (2018) 89-97. https://www.researchgate.net/publication/323689277_4-PRIME_CORDIAL_LABELING_OF_SOME_SPECIAL_GRAPHS

[2968] R. Ponraj, Rajpal Singh,and R. Kala, 4-Prime cordiality of m copies of some graphs, *Palestine J. Math.*, **8**(1) (2018) 307-311. https://pjm.ppu.edu/sites/default/files/papers/PJM_Oct2018_307to311.pdf

[2969] R. Ponraj, S. Sathish Narayanan and R. Kala, Radio mean number of some subdivision graphs, *Jordan J. Math. and Stat. (JJMS),* **9**(1) (2016) 45-64. https://jjms.yu.edu.jo/index.php/jjms/article/view/1046/438

[2970] R. Ponraj, Rajpal Singh, and S. Sathish Narayanan, On parity combination cordial graphs, *Palestine J. Math.*, *6*(1) (2017) 211-218. https://pjm.ppu.edu/sites/default/files/papers/PJM_MAR_2017_25.pdf

[2971] R. Ponraj, S. Sathish Narayanan and A. M. S. Ramasamy, Parity combination cordial labeling of graphs, *Jordan J. Math. and Stat. (JJMS)*, **8**(4)(2015) 293-308. https://journals.yu.edu.jo/jjms/Issues/Vol8No42015PDF/3.pdf

[2972] R. Ponraj and M. Sivakumar, On mean cordial graphs, *Internat. J. Math. Combin.,* **3** (2013) 78-84. https://fs.unm.edu/IJMC/OnMeanCordialGraphs.pdf

[2973] R. Ponraj, M. Sivakumar, and M. Sundaram, k-Product cordial labeling of graphs, *Int. J. Contemp. Math. Sci.,* **7** (2012) 733-742. https://www.m-hikari.com/ijcms/ijcms-2012/13-16-2012/ponrajIJCMS13-16-2012.pdf

[2974] R. Ponraj, M. Sivakumar, and M. Sundaram, On 4-product cordial graphs, *Inter. J. Math. Archive,* **7** (2012) 2809-2814.

[2975] R. Ponraj, M. Sivakumar, and M. Sundaram, k-total product cordial labeling of graphs, *Appl. Appl. Math.: An Inter. J.*, **7** (2012) 708-716. https://digitalcommons.pvamu.edu/cgi/viewcontent.cgi?article=1304&context=aam

[2976] R. Ponraj, M. Sivakumar, and M. Sundaram, On 3-total product cordial graphs, *Int. Math. Forum,* **7**(29-32) (2012) 1537-1546. https://www.m-hikari.com/imf/imf-2012/29-32-2012/ponrajIMF29-32-2012.pdf

[2977] R. Ponraj, M. Sivakumar, and M. Sundaram, New families of 3-product cordial graphs, *Inter. J. Math. Archive,* **3**(5) (2012) 1985-1990. NOW 8/28/25 or Chat 10/8/26

[2978] R. Ponraj, M. Sivakumar, and M. Sundaram, 3-Total product cordial labeling of some subdivided graphs, *Inter. J. Math. Research,* **5** (2012) 517-526.

[2979] R. Ponraj and M. Sivakumar, A note on k-Total product cordial graphs, *Global J. Math. and Math. Sciences,* **1** (2012) 37-44.

[2980] R. Ponraj, M. Sivakumar, and M. Sundaram, On 4-product cordial graphs, *Inter. J. Math. Archive,* **7** (2012) 2809-2814.

[2981] R. Ponraj, M. Sivakumar, and M. Sundaram, k-Total product cordial labeling of graphs, *Appl. Appl. Math.: An Inter. J.*, **7** (2012) 708-716. https://digitalcommons.pvamu.edu/cgi/viewcontent.cgi?article=1304&context=aam

[2982] R. Ponraj, M. Sivakumar, and M. Sundaram, On 3-total product cordial graphs, *Int. Math. Forum,* **7**(29-32) (2012) 1537-1546. https://www.m-hikari.com/imf/imf-2012/29-32-2012/ponrajIMF29-32-2012.pdf

[2983] R. Ponraj, M. Sivakumar, and M. Sundaram, New families of 3-product cordial graphs, *Inter. J. Math. Archive,* **3**(5) (2012) 1985-1990.

[2984] R. Ponraj, M. Sivakumar, and M. Sundaram, Mean cordial labeling of graphs, *Open J. Discrete Math.,* **2** (2012) 145-148. doi:10.4236/ojdm.2012.24029

[2985] R. Ponraj and S. Somasundaram, Further results on mean graphs, *Proc. SACOEFERENCE, National Level Conference, Dr. Sivanthi Aditanar College of Engineering*, (2005) 443-448.

[2986] R. Ponraj and S. Somasundaram, Mean labeling of graphs obtained by identifying two graphs, *J. Discrete Math. Sci. Cryptogr.,* **11**(2) (2008) 239-252.

[2987] R. Ponraj, S. Subbulakshmi, and S. Somasundaram, On 4-total mean cordial graphs, *J. Appl. Math. Inform.,* **39**(3-4) (2021) 497-506. https://www.koreascience.or.kr/article/JAKO202116954748838.pdf

[2988] R. Ponraj, S. Subbulakshmi, and S. Somasundaram, Further results on 4-difference cordial graphs, *Palest. J. Math.,* **11**(1) (2022) 142-148. https://pjm.ppu.edu/sites/default/files/papers/PJM_November_2021_142_to_148.pdf

[2989] R. Ponraj and T. Sutharson, Fractional product cordial labeling of graph, *J. Appl. Pure Math.*, **7**(1-2) (2025) 49-58. https://koreascience.kr/article/JAKO202512036003896.pdf

[2990] R. Ponraj and T. Sutharson, Fractional product cordial labeling of web graph, dumbbell graph, jellyfish graph, jewel graph and tadpole graph,*Indian Acad. Math.*, **46**(2) (2024) 303-312.

[2991] R. Ponraj and T. Sutharson, Some fractional product cordial graphs, *Internat. J. Math. Combin.*, **1-2** (2025) 67-77. https://fs.unm.edu/IJMC/Some_Fractional_Product_Cordial_Graphs.pdf

[2992] R. Ponraj and T. Sutharson, FP-cordial labeling of kayak paddle, parachute, and certain corona graphs, *J. Appl. Pure Math.*, **7**(5-6) (2025) 365-382 . https://koreascience.or.kr/article/JAKO202502339627576.pdf

[2993] G. Pooranam, R. Vasuki, and S. Suganthi, New families of odd mean graphs, *Internat. J. Math. Combin.,* **1** (2019) 134-145. https://www.researchgate.net/publication/338178781_International_Journal_of_Mathematical_Combinatorics_vol_12019

[2994] V. Poovila and C. Vimala, Application of vertex odd power mean labeling in cryptography, *AIP Conf. Proc.,* 3252, 020216 (2025) 1-5. https://pubs.aip.org/aip/acp/article/3252/1/020216/3338071/Application-of-vertex-odd-power-mean-labeling-in

[2995] R. S. Porchelvi and S. A. Devi, Pronic graceful graphs, *J. Xidian University*, **14**(3) (2020) 895-898.

[2996] R. S. Porchelvi and S. A. Devi, Pronic Heron mean graphs, *The Internat. J. Analytical and Experimental Modal Analysis,* **XII**(III) (2020) 2207-2212. https://docslib.org/doc/2974969/pronic-heron-mean-graphs-1r#google_vignette

[2997] P. Pradhan and K. Kamesh, On k-graceful labeling of some graphs, *Math. Inform.,* **34**(1-2) (2016) 9-17. https://koreascience.kr/article/JAKO201614734990648.pdf

[2998] P. Pradhan and A. Kumar, Graceful hairy cycles with pendent edges and some properties of cycles and cycle related graphs, *Bull. Calcutta Math. Soc.,* **103**(3) (2011) 233-246.

[2999] T. R. Pradipta and A. N. M. Salman, Some cycle-supermagic labelings of the calendula graphs, *J. Physics Conference Series,* **948**(1):012071 (2018) doi:10.1088/1742-6596/948/1/012071

[3000] U. M. Prajapati, R. Asad, and R. M. Gajjar, Cordiality in the context of duplication in quadrilateral snake graph, *Internat. J. Sci. Res. Math. Stat. Sci.*, **10**(5) (2023) 22-25. https://www.isroset.org/pdf_paper_view.php?paper_id=3295&4-ISROSET-IJSRMSS-09035.pdf

[3001] U. M. Prajapati and S. J. Gajjar, Some results on prime labeling *Open J. Discrete Math.,* **4** (2014) 60-66. http://dx.doi.org/10.4236/ojdm.2014.43009

[3002] U. M. Prajapati and R. M. Gajjar, personal commuication (2011).

[3003] U. M. Prajapati and R. M. Gajjar, Cordial labeling for complement of some graphs, *Math. Today*, **30** (2015) 99-118.

[3004] U. M. Prajapati and R. M. Gajjar, Cordial labeling in the context of duplication of some graph elements, *Internat. J. Math. Soft Comput.*, **6**(2) (2016) 65-73. https://oaji.net/articles/2017/296-1504970018.pdf

[3005] U. M. Prajapati and R. M. Gajjar, Cordiality in the context of duplication in crown related graphs, *J. Math. Comput. Sci.,* **6**(6) (2016) 1058-1073. http://scik.org/index.php/jmcs/article/view/2896

[3006] U. M. Prajapati and R. M. Gajjar, Cordiality in the context of duplication in flower related graphs, *Internat. J. Math. Soft Comput.*, **7**(2) (2017) 9-101. www.ijmsc.com/index.php/ijmsc/article/view/ijmsc-7-2-8/pdf

[3007] U. M. Prajapati and R. M. Gajjar, Cordiality in the context of duplication in web and armed helm, *Internat. J. Math. Combin.*, **3** (2017) 90-105. https://vixra.org/pdf/1809.0414v1.pdf

[3008] U. M. Prajapati and R. M. Gajjar, Some labeling techniques of holiday star graph, *Internat. J. Math. Trends Tech.*, **49**(6) (2017) 339-344. www.ijmttjournal.org/2017/Volume-49/number-6/IJMTT-V49P554.pdf

[3009] U. M. Prajapati and R. M. Gajjar, Some labeling techniques of Kusudama flower graph, *Math. Today*, **34**(A) (2018) 211-217.

[3010] . U. M. Prajapati and S. J. Gaijjar, Some labeling techniques of braided star graph, *Internat. J. Math. Appl.,* **5**(4-C) (2017) 361-369 https://ijmaa.in/index.php/ijmaa/article/view/1281/1245

[3011] U. M. Prajapati and R. M. Gajjar, Cordiality in the context of duplication in helm and closed helm, *Internat. J. Math. Appl.*, **6**(1-A) (2018) 139-145. https://ijmaa.in/index.php/ijmaa/article/view/907/892

[3012] U. M. Prajapati and R. M. Gajjar, Some labeling techniques of boreale star graph, *Internat. J. Res. Analytical Reviews*, **6**(1) (2019) 668-671. https://ijrar.org/papers/IJRAR19J2514.pdf

[3013] U. M. Prajapati and S. J. Gajjar, Prime labeling of generalized Petersen graph, *Internat. J. Math. Soft Comput.,* **5**(1) (2015) 65-71. https://www.researchgate.net/publication/321078817_Prime_Labeling_of_Generalized_Petersen_Graph

[3014] U. M. Prajapati and S. J. Gajjar, Prime cordial labeling of generalized prism graph $Y_{m,n}$, *Ultra Scientist,* **27**(3)A (2015) 189-204. https://www.researchgate.net/publication/329626528_Prime_Cordial_Labeling_of_Generalized_Prism_Graph

[3015] U. M. Prajapati and N. B. Patel, Edge product cordial labeling of some cycle related graphs, *Open J. Discrete Math.*, **6**(6) (2016) 268-278. http://dx.doi.org/10.4236/ojdm.2016.64023

[3016] U. M. Prajapati and N. B. Patel, Edge product cordial labeling of some graphs, *J. Appl. Math. Comput. Mech.,* **18**(1) (2019) 69-76. http://amcm.pcz.pl/?id=view&volume=18&issue=1&article=6

[3017] U. M. Prajapati and N. B. Patel, Edge product cordial labelings of switching operation on some graphs, *TWMS J. App. Eng. Math.*, **12**(1) (2022) 191-199. https://jaem.isikun.edu.tr/web/images/articles/vol.12.no.1/17.pdf

[3018] U. M. Prajapati and K. K. Raval, Product cordial labeling in context of some one point union graphs, *IJRAR- Internat. Res. Analytical Reviews,* **5I**(4) (2018) 490-495. https://www.ijrar.com/upload_issue/ijrar_issue_20542226.pdf

[3019] U. M. Prajapati and K. K. Raval, Product cordial graph in the context of some graph operations on gear graph, *Open J. Discrete Math.*, **6** (2016) 259-267. https://www.scirp.org/pdf/ojdm_2016090916212031.pdf?utm_source=chatgpt.com

[3020] U. M. Prajapati and K. K. Raval, Some graphs in the context of product cordial labeling, *Math. Today*, **33** (2017) 58-66. http://mathematicstoday.org/archives/V33_2017_9.pdf

[3021] U. M. Prajapati and K. K. Raval, Vertex even and odd mean labeling in the context of some cyclic snake graphs, *Internat. J. Emerging Tech. Innovat, Res.,* **5**(7) (2018) 319-324. http://www.jetir.org/papers/JETIRC006365.pdf

[3022] U. Prajapati and K. Raval, Vertex odd and even mean labeling of triangular snake, *J. Appl. Sci. Computations,* **5**(12) (2018) 1273-1280. http://j-asc.com/gallery/182-december-1554.pd

[3023] U. Prajapati and K. Raval, Several results of vertex even mean and vertex odd mean graphs, *Adv. Appl. Math. Sci.*, **19**(4) (2020) 259-272. https://www.researchgate.net/publication/357867081_Several_Results_of_Vertex_Even_Mean_and_Vertex_Odd_Mean_Graphs

[3024] U. M. Prajapati and P. D. Shah, Some edge product cordial graphs in the context of duplication of some graph elements, *Open J. Discrete Math.*, **6** (2016) 248-258. http://dx.doi.org/10.4236/ojdm.2016.64021

[3025] U. M. Prajapati and K. P. Shah, On odd prime labeling, *IJRAR- Internat. J. Res. Analy. Rev.,* **5**(4) (2018) 2348-1269. https://ijrar.com/upload_issue/ijrar_issue_20542373.pdf

[3026] U. M. Prajapati and K. P. Shah, Odd prime labeling of graphs with duplication of graph elements, *JASC: J. Appl. Sci.Comp.*, **V**(XII) (2018) 1281-1288 http://j-asc.com/gallery/183-december-1553.pdf.

[3027] U. M. Prajapati and K. P. Shah, Odd rime labeling of graphs with duplication of graph elements, *JASC: J. Appl. Sci. Comput.,* **5**(12) (2018) 1281-1288.

[3028] U. M. Prajapati and B. M. Suthar, Prime labeling in the context of duplication of graph elements in $K_{2,n}$, *Internat. J. Math. Soft Comput.*, **7**(1) (2017) 117-125. https://oaji.net/articles/2017/296-1504971936.pdf

[3029] U. M. Prajapati and A. V. Vantiya, SD-prime cordial labeling of some snake graphs, *JASC: J. Appl. Sci. Comput.,* **VI**(IV) (2019) 1857-1868. https://www.researchgate.net/publication/335685524_SD-Prime_Cordial_Labeling_of_Some_Snake_Graph

[3030] U. M. Prajapati and A. V. Vantiya, SD-prime cordial labeling of K_4-snake and related graphs, *J. Xidian U.,* **14**(4) (2020) 543-551. https://www.researchgate.net/publication/340600650_SD-Prime_Cordial_Labeling_of_K4-Snake_and_Related_Graphs

[3031] U. Prajapati and A. V. Vantiya, SD-Prime cordial labeling of alternate k-polygonal snake of various types, *Proyecciones,* **40**(3) (2021) 619-634. https://www.scielo.cl/pdf/proy/v40n3/0716-0917-proy-40-03-619.pdf

[3032] V. Prakash, R. Gopi, and P. Shalini, Anti skolem mean labeling of quardrilateral snake related graphs, *Tuijin Jishu/J. Propulsion Tech.*, **44**(5) (2023) 495-500. https://comb-opt.azaruniv.ac.ir/article_14428_29f3bfe45780bf0345829025de1c755a.pdf

[3033] G. Prasad and G. U. Maheswari, Matrix coding technique on sunflower graphs with edge product cordial labeling, *IOP Conf. Series: Materials Sci. Engin.,* **872** (2020) 012004 1-11. https://www.researchgate.net/publication/342502155_Matrix_Coding_Technique_on_Sunflower_Graphs_with_Edge_Product_Cordial_Labeling

[3034] R. V. Prasad, M. Kannan, and R. Gopi, Some results on even vertex odd mean labeling graphs, *Internat. J. Mechanical Enginer. Tech.*, (IJMET), **9**(2) (2018) 615-621. https://iaeme.com/MasterAdmin/Journal_uploads/IJMET/VOLUME_9_ISSUE_2/IJMET_09_02_062.pdf?utm_source=chatgpt.com

[3035] D. A. B. Prasetyo, Strong edge antimagic total labeling on multistar, *J. Phys.: Conf. Ser.,* (2021) 1778 012009 1-7. https://iopscience.iop.org/article/10.1088/1742-6596/1778/1/012009/meta

[3036] D. Prathap and J. Baskar Babujee, Magic and bimagic labeling for star graphs, *Internat. Review Pure Appl. Math.,* **5**(1) (2009) 67-76.

[3037] D. Praveen Prabhu and O. V. Shanmuga Sundaram, Investigation of super Lehmer–3 mean labeling in theta related graphs, *Stat. Optim. Informat. Comput.*, **14**(4) (2025) 2070-2090. http://www.iapress.org/index.php/soic/article/view/2633/1440

[3038] M. Prema and K. Murugan, Oblong sum labeling of some special graphs. World Scientific News, 98 (2018) 12-22. http://www.worldscientificnews.com/wp-content/uploads/2018/02/WSN-98-2018-12-22-1.pdf

[3039] K. Premalatha, G. C. Lau, S. Arumugam and W. C. Shiu, On local antimagic chromatic number of various join graphs, *Commun. Comb. Optim.*, **8**(4) (2023) 693-714. https://comb-opt.azaruniv.ac.ir/article_14428_29f3bfe45780bf0345829025de1c755a.pdf

[3040] B. N. G. Pricilla, A Study On New Classes Of Graphs In Variations Of Graceful Graph, Ph.D. Thesis, Bharath University, Chennai, 2008. http://shodhganga.inflibnet.ac.in/handle/1w0603/33

[3041] R. M. Prihandini and R. Adawiyah1, On super $3(n+5,2)$-edge antimagic total labeling and its application to construct Hill cipher algorithm, *BAREKENG: J. Math. Its Applications*,**17**(1) (2023) 0029–0036. https://ojs3.unpatti.ac.id/index.php/barekeng/article/view/6427/5506

[3042] R. M. Prihandini, Dafik, I. H. Agustin, R. Alfarisi, R. Adawiyah, and K. A. Santoso, On super $(a,d) - P_2 \triangleright H$ antimagic total labeling of disjoint union of comb product graphs, *J. Physics Conf. Ser.* 1211:012012 (2019) https://www.researchgate.net/publication/332916076_On_super_ad-P_2unicodex22B3H_antimagic_total_labeling_of_disjoint_union_of_comb_product_graphs

[3043] G. Siva Prijith, M. Subbulakshmi, and S. Chandrakala, On topological cordial labelling of some graphs, *Mapana J. Sciences,* **22**(1) (2023) 139-150. Special Issue. https://journals.christuniversity.in/index.php/mapana/article/view/3884/2409

[3044] V. Princy Kala, k-super cube root cube mean labeling of graphs, *Proyecciones*, (on line), **40**(5) (2021) 1097-1116. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/4258/3885

[3045] V. Princy Kala, k-super cube root cube mean labeling of some corona graphs, *Proyecciones* **41**(4) (2022) 903-921. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/5200/4093

[3046] V. Priyadharshini and M. Nalliah, Local distance antimagic chromatic number for the union of complete bipartite graphs, *Tamkang J. Math.*, **54**(4) (2023) 281-291. https://www.researchgate.net/publication/375612296_Local_distance_antimagic_chromatic_number_for_the_union_of_complete_bipartite_graphs

[3047] J. Przybylo, Irregularity strength of regular graphs, *Electron. J. Combin.,* **15**(1) (2008), P82, 10 pp. https://www.combinatorics.org/ojs/index.php/eljc/article/view/v15i1r82

[3048] J. Przybylo, Linear bound on the irregularity strength and the total vertex irregularity strength of graphs, *SIAM J. Discrete Math.*, **23** (2009) 511-516.

[3049] J. Przybylo, Linear bound on the irregularity strength and the total vertex irregularity strength of graphs, *SIAM J. Discrete Math.,* **23**(1) (2008/09) 511-516.

[3050] J. Przybylo, Distant irregularity strength of graphs, *Discrete Math.,* **313**(24) (2013) 2875-2880. https://www.sciencedirect.com/science/article/pii/S0012365X13003865?via%3Dihub

[3051] J. Przybyło, Asymptotic confirmation of the Faudree–Lehel conjecture on irregularity strength for all but extreme degrees, *J. Graph. Theory,* **100** 2022 189-204. https://onlinelibrary.wiley.com/doi/10.1002/jgt.22772?msockid=039e6c86024a607b2a3c793c033e61b1

[3052] J. Przybyło, The irregularity strength of dense graphs — on asymptotically optimal solutions of problems of Faudree, Jacobson, Kinch and Lehel, *European J. Combin.,* (2024). https://www.sciencedirect.com/science/article/pii/S0195669824000982

[3053] Q. Pu, M. Numan, S. I. Butt, M. Asif, F. Rafique, and Z. Shao, Cyclic super magic labelings for toroidal and Klein-bottle fullerene, *IEEE Access*, **7** (2019) 145928-145933. https://www.researchgate.net/publication/336029087_Cyclic_Super_Magic_Labelings_for_Toroidal_and_Klein-Bottle_Fullerenes

[3054] J.-F. Puget, Breaking symmetries in all different problems, *in* Proceedings of SymCon04, the 4th International Workshop on Symmetry in Constraints, 2004, 6 pp. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=de5f6b75b7620c9b4aff3ec464a6b3767c485e65

[3055] D. A. Pujiwati, I. Halikin, and K. Wijaya, Odd harmonious labeling of two graphs containing star, *AIP Conference Proceedings 2326, 020019* (2021) 1-5. https://aip.scitation.org/doi/pdf/10.1063/5.0039644

[3056] N. Punnim and N. Pabhapote, On graceful graphs: cycles with a P_k-chord, $k \geq 4$, *Ars Combin.,* **23A** (1987) 225-228. NOW 8/30/25 or Chat 10/9/28

[3057] C. Purcell, J. Ryan, Z. Ryjáček, and M. Skyvová, On exclusive sum labellings of hypergraphs, *Graphs and Combin.,* **38**(2) (2022) 16 pp. DOI:10.1007/s00373-021-02432-4

[3058] R. W. Putra and Y. Susanti, On total edge irregularity strength of centralized uniform theta graphs, *AKCE Int. J. Graphs Comb.*, **15**(1) (2018) 7–13. https://www.sciencedirect.com/science/article/pii/S0972860017300555

[3059] P. R. L. Pushpam and A. Saibulla, On super (a, d)-edge antimagic total labeling of certain families of graphs, *Discuss. Math. Graph Theory* **32**(3) (2012) 535-543. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1623

[3060] A. Puthussery and G. Muneeswari (2021) Faulty node detection using vertex magic total labelling, in distributed system. In: P. Karuppusamy, I. Perikos, F. Shi, and T. N. Nguyen (eds) Sustainable Communication Networks and Application.

Lecture Notes on Data Engineering and Communications Technologies, Springer, Singapore. **55** (23021 619-630. https://doi.org/10.1007/978-981-15-8677-4_50

[3061] A. Pyatkin, New formula for the sum number for the complete bipartite graphs, *Discrete Math.,* **239** (2001) 155-160. https://www.sciencedirect.com/science/article/pii/S0012365X01001881

[3062] J. Quadras and S. Teresa Arockiamary, Total edge irregularity strength of hexagonal networks, *J. Combin. Math. Combin. Comput.,* **92** (2015) 131-138. https://combinatorialpress.com/article/jcmcc/Volume%20092/vol-092-paper%2013.pdf

[3063] J. Qian, On some conjectures and problems in graceful labelings graphs, unpublished.

[3064] J. Radha, Y. Venkatesan, K. Vitaldas, and V. Perumal, Group $\{1, -1, i, -1\}$ cordial labeling in some classes of graphs, *5th International Conference on Intelligent Computing: IConIC* 2K22 25–26 March 2022, **2790**(1) (2023) https://doi.org/10.1063/5.0152417

[3065] J. Rafael, M. Antalan, A. DC. Angel, J. Loureynz, F. Gamurot, and R. P. Tagle, Edge-graceful usual fan graphs and the edge-gracefulness of degree-splitted usual fan graphs, *IAENG Internat. J. Appl. Math.* **56**(2) (2026) 559-572. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4998369

[3066] J. Rafael, M. Antalan, R. P. Tagle, A. DC Angel, J. Loureynz, and F. Gamurot, An investigation on the edge-gracefulness of bubble sort, bubble sort star, and Kulli-cycle windmill graphs, *Sci. Tech. Indonesia*, **11**(4) (2026) 1735-1743. https://sciencetechindonesia.com/index.php/jsti/article/view/2466

[3067] P. Ragukumara and G. Sethuraman, Binomial trees are graceful, *AKCE Internat. J. Graphs Combin.,* **17**11 632-636. https://www.tandfonline.com/doi/pdf/10.1016/j.akcej.2018.06.005

[3068] A. Raheem, On super (a, d)-edge-antimagic total labeling of a subdivided stars, *Ars Combin.,* **136** (2018) 169-179. https://www.researchgate.net/publication/266859985_On_super_ad-edge-antimagic_total_labeling_of_subdivided_stars

[3069] A. Raheem and A. Q. Baig, Super edge-antimagicness of subdivided star trees, *Discuss. Math. Graph Th.,* **1** (2016) 27-33. https://fs.unm.edu/IJMC/SuperEdgeAntimagicLabelingOfSubdivided.pdf

[3070] A. Raheem, M. Javaid, and A. Q. Baig, On super edge-antimagicness of subdivided stars, *Discuss. Math. Graph Th.,* **4** (2015) 663-673.

https://www.researchgate.net/publication/282062714_On_Super_Edge-Antimagicness_of_Subdivided_Stars

[3071] A. Raheem, M. Javaid, M. Numan, and R. Hasni, Sum divisor cordial labeling of disjoint union of paths and subdivided star, *J. Disc. Math. Sci. Crypt.,* (2021) 2467-2478. url-https://www.researchgate.net/publication/353662299_Sum_divisor_cordial_labeling_of_disjoint_unio

[3072] M. T. Rahim, K. Ali, Kashif and I. Javaid, On antimagic total labeling of some families of graphs, *Ars Combin.,* **95** (2010) 225-234. https://combinatorialpress.com/article/ars/Volume%20095/volume-95-paper-20.pdf

[3073] M. T. Rahim, M. Farooq, M. Ali, and S. Jan, Multi-level distance labelings for generalized gear graphs *Internat. J. Math. and Soft Comput.,* **2**(1) (2012) 57-63. https://www.researchgate.net/publication/234059451_Multi-Level_Distance_Labelings_for_Generalized_Gear_Graph

[3074] M. T. Rahim and Slamin, Most wheel related graphs are not vertex magic, *Util. Math.,* **77** (2008) 193-199. https://repository.unej.ac.id/bitstream/handle/123456789/775/Pages%20from%20UM_77_VM_Wheel_2008.pdf?sequence=1&isAllowed=y

[3075] M. T. Rahim and Slamin, Vertex-magic total labelings of the union of suns, *Ars Combin.,* **103** (2012) 305-310. https://combinatorialpress.com/article/ars/Volume%20103/volume-103-paper-25.pdf

[3076] M. T. Rahim and Slamin, Vertex-magic total labeling of the union of suns, *Ars Combin.,* 103 (2012) 305-310. https://combinatorialpress.com/article/ars/Volume%20103/volume-103-paper-25.pdf

[3077] M. T. Rahim, I. Tomescu, and Slamin, On vertex-magic total labeling of some wheel related graphs, *Util. Math.,* **73** (2007) 97-104. https://www.researchgate.net/publication/235798020_On_vertex-magic_total_labeling_of_some_wheel_related_graphs

[3078] D. Rahmadani, A. Aldiansyah, D. Pratiwi, M. Yunus, and V. Kusumasari, Prime labeling of amalgamation flower graphs, *BAREKENG: J. Math. & App.* **19**(4) (2025) 2431-2442. https://www.researchgate.net/publication/395261706_PRIME_LABELING_OF_AMALGAMATION_OF_FLOWER_GRAPHS

[3079] D. S. Rahmadani, D. E. Putri, Wasono, and H. Sandariria, Local antimagic edge coloring of gear graphs and semi parachute graphs, *J. Math. Edu. Sci.*, **8**(2) (2025) 123-132. https://journal.unugiri.ac.id/index.php/JaMES/article/view/4775/1842

[3080] S. Rahmawati, K. A. Sugeng, D. R. Silaban, M. Miller, and M. Bača, Construction of new larger (a, d)-edge antimagic vertex graphs by using adjacency matrices, *Austral. J. Combin.*, **56**, (2013) 257-272. https://ajc.maths.uq.edu.au/pdf/56/ajc_v56_p257.pdf

[3081] C. D. Raj, C. Jayasekaran, and S. S. Sandhya, Few families of harmonic mean graphs, *J. Combin. Math. Combin. Comput.,* **96** (2016) 235-243. https://combinatorialpress.com/article/jcmcc/Volume%20096/vol-096-paper%2016.pdf?utm_source=chatgpt.com

[3082] C. D. Raj, K. R. Mary, and K. C. Krishna, Sum graphs & its related concepts, *Internat. J.Aquatic Sci.*, **12**(02) (2021) 126-129. https://www.researchgate.net/publication/397869637_Sum_Graphs_Its_Related_Concepts

[3083] C. D. Raj, K. C. Krishna, and K. R. Mary, A study of even sum graphs, *South East Asian J. Math. Math. Sci. Proceeding*, **18** (2022) 119-124. https://www.researchgate.net/publication/397869735_A_STUDY_ON_EVEN_SUM_GRAPHS

[3084] P. L. R. Raj and S. Koilraj, Cordial labeling for the splitting graph of some standard graphs, *Internat. J. Math. Soft Comput.*, **1**(1)(2011) 105-114. http://oaji.net/articles/2015/296-1433770428.pdf

[3085] P. L. R. Raj and R. L. J. Manoharan, Signed product cordial graphs in the context of arbitrary supersubdivision, *Scientia Magna.* **8** (2012) 77-87. https://www.academia.edu/26344047/Signed_product_cordial_graphs_in_the_context_of_arbitrary_supersubdivision

[3086] P. L. R. Raj and R. Valli, Some new families of divisor cordial graphs, *Internat. J. Math. Trends Tech.,* **7** (2) (2014) 94-102. http://ijmttjournal.org/Volume-7/number-2/IJMTT-V7P512.pdf

[3087] P. L. R. Raj and K. Vivek, Some new families of analytic mean graphs, *Internat. J. Anal. Exper. Modal Anal.,* **VI**(X) (2019) 358-361. https://www.researchgate.net/publication/336847676_Some_New_Families_of_Analytic_Mean_Graphs

[3088] S. Soundar Rajan and J. Baskar Babujee, Further results on signed product cordial labeling, *Revista Argentina de Clinica Psicologica*, **XXXII** (2023) N*1,01-04. https://www.revistaclinicapsicologica.com/data-cms/articles/20230206034702pm7001.pdf

[3089] I. Rajasingh and P. R. L. Pushpam, Strongly harmonious labeling of helms, personal communication (circa 1996).

[3090] I. Rajasingh and P. R. L. Pushpam, On graceful and harmonious labelings of t copies of $K_{m,n}$ and other special graphs, personal communication (circa 1996).

[3091] I. Rajasingh, B. Rajan, and V. Annamma, On total vertex irregularity strength of triangle related graphs, *Annals Pure Appl. Math.,* **1**(2) (2012) 108-116.

[3092] I. Rajasingh and V. Annamma, Total vertex irregularity strength of 1-fault tolerant hamiltonian graphs, *Math. Comput. Sci.,* **9**(2) (2015) 151-160.

[3093] I. Rajasingh, B. Rajan, and S. Teresa Arockiamary, Irregular total labeling of butterfly and benes networks, In: Abd Manaf A., Sahibuddin S., Ahmad R., Mohd Daud S., El-Qawasmeh E. (eds), Informatics Eng. Infor. Sci., ICIEIS 2011, *Comm. Comput. Information Sci.,* **253** (2011) Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-25462-8_25

[3094] I. Rajasingh and S. Teresa Arockiamary, Total edge irregularity strength of honeycomb networks, *Internat. J. Comput. and Applied Math.*, **8**(3) (2013) 213-220.

[3095] I. Rajasingh and S. Teresa Arockiamary, Total edge irregularity strength of honeycomb torus networks, *Global J. Pure Appl. Math.*, **13**(4) (2017) 1135-1142. https://www.ripublication.com/gjpam17/gjpamv13n4_01.pdf

[3096] I. Rajasingh and S. Teresa Arockiamary, Total edge irregularity strength of series parallel graphs, *Internat. J. Pure Appl. Math.*, **99**(1) (2015) 11-21. https://www.researchgate.net/publication/275242200_Total_edge_irregularity_strength_of_series_parallel_graphs

[3097] T. J. Rajesh Kumar, A. S. Jerome,, and K. R. Santhosh Kumar, A note on Gaussian total neighborhood prime labeling, *Adv. Math.: Sci. J.,* **10**(3) (2021) 1821-1832. https://www.researchgate.net/publication/350536922_A_NOTE_ON_GAUSSIAN_TOTAL_NEIGHBOURHOOD_PRIME_LABELING

[3098] T. J. Rajesh Kumar T.J. and T. K. Mathew Varkey, A note on total neighborhood labelings, *Internat. J. Pure and Appl. Math.*, **118**(4) (2018) 1007-1013. https://www.ijpam.eu/contents/2018-118-4/14/14.pdf

[3099] T. J. Rajesh Kumar and T. K. Mathew Varkey, Gaussian neighborhood-prime labeling of some trees and other graphs, *Annals Pure and Appl. Math.*, **16**(1) (2018) 133-140. http://www.researchmathsci.org/APAMart/apam-v16n1-15.pdf

[3100] T. J. Rajesh Kumar and T. K. Mathew Varkey, A note on total neighborhood labelings, *Internat. J. Pure and Appl. Math.*, **118**(4) (2018) 1007-1013. https://ijpam.eu/contents/2018-118-4/14/14.pdf

[3101] S. Rajkumar and M. Nalliah, On local edge antimagic chromatic number of graphs, *Proyecciones (Antofagasta)* **41**(6) (2022) 1397-1412. doi:10.22199/issn.0717-6279-5339

[3102] S. Rajkumar, M. Nalliah, G. Uma Maheswari, Super $(a, d) - G + e$-antimagic total labeling of $G_u[S_n]$, *Proyecc. J. Math.,* **41**(1) (2022) 249-262. https://www.revistaproyecciones.cl/article/view/5014

[3103] S. Rajkumar, M. Nalliah, and M. Venkataraman, Note on super $(a, 1)$-P_3-antimagic total labeling of star S_n, *Ural Math. J.* **7**(2) (2022) 86-93. https://umj.uran.ru/index.php/umj/article/view/357/pdf

[3104] V. Ramachandran and C. Sekar, One modulo N gracefulness of acyclic graphs, *Ultra Scientist,* **25**(3)A (2013) 417-424. NOW 8/31/25 orvChat 10/9/26

[3105] V. Ramachandran and C. Sekar, One modulo N gracefulness of splitting graphs and subdivision of double triangle graphs, *Scientia* **125** (2014) 1-9.

[3106] V. Ramachandran and C. Sekar, One modulo N gracefulness of arbitrary supersubdivisions of graphs, *Internat. J. Math. Combin.,* **2** (2014) 36-46. https://vixra.org/pdf/1411.0280v1.pdf

[3107] V. Ramachandran and C. Sekar, Gracefulness and one modulo N gracefulness of $L_n \otimes S_m$, *Scientia Magna,* **10**(3) (2014) 66-76.

[3108] V. Ramachandran and C. Sekar, One modulo N gracefulness of regular bamboo tree and coconut tree, *Internat. J. Appl. Graph Theory Wireless ad hoc Networks Sensor Networks*, (GRAPH-HOC) **6**(2) (2014) 1-10. doi:10.5121/jgraphoc.2014.6201

[3109] V. Ramachandran and C. Sekar, One modulo N gracefulness of n-polygonal snakes, $C_n^{(t)}$ and $P_{a,b}$, *Internat. J. Engineering Res. & Tech.*, (IJERT) **2**(10) (2013) 3514-3529. https://www.ijert.org/research/one-modulongracefulness-of-n-polygonal-snakes-c-t-n-andpab-IJERTV2IS100899.pdf

[3110] V. Ramachandran and C. Sekar, Graceful labelling of supersubdivision of ladder, *Internat. J. Math. Appl.,* **2**(2) (2014) 29-36. https://oaji.net/articles/2014/1167-1408377254.pdf

[3111] V. Ramachandran and C. Sekar, One modulo N gracefulness of crowns, armed crowns and chain of even cycles, *Ars Combin.,* **138** (2018) 143-159.

[3112] K. Ramanjaneyulu, V. C. Venkaiah, and K. Kothapalli, Cordial labelings of a class of planar graphs *AKCE J. Graphs Combin.,* **6** (2009) 171-181. https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=e4e18fed7037a7a1b35f57671540744523cf7d21

[3113] M. Ramalakshmi and K. M. Kathiresan, Total absolute difference edge irregularity strength of the graphs, *Kragujevac J. Math.*, **46**(6) (2022) 895–903. https://www.researchgate.net/publication/363366676_TOTAL_ABSOLUTE_DIFFERENCE_EDGE_IRREGULARITY_STRENGTH_OF_GRAPHS

[3114] J. L. Ramírez-Alfonsín, Gracefulness of replicated paths and cycles, *Ars Combin.,* **53** (1999) 257-269.

[3115] R. Ramdani and A. N. M. Salman, On the total irregularity strength of some Cartesian product graphs, *AKCE Int. J. Graphs Comb.,* **10**(2) (2013) 199-209. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088736

[3116] R. Ramdani, A. N. M. Salman, and H. Assiyatun, On the total irregularity strength of regular graphs, *J. Math. Fundam. Sci.,* **47**(3) (2015) 281-295. https://journal-itb.org/jmfs/volumes/2015/47/issues/3/on-the-total-irregularity-strength-of-regular-graphs.html

[3117] R. Ramdani, A. N. M. Salman, H. Assiyatun, A. Semaničová-Feňovčíková, and Bača, Total irregularity strength of three families of graphs, *Math. Comput. Sci.,* **9**(2) (2015) 229-237. https://www.researchgate.net/publication/276424159_Total_Irregularity_Strength_of_Three_Families_of_Graph

[3118] R. Ramdani, A, N. M. Salman, A. Assiyatum, and A. Semaničová-Feňovčíková, On the total irregularity strength of disjoint union of graphs, *Math. Reports*, **18**(68)(4) (2016) 469-482. https://imar.ro/journals/Mathematical_Reports/Pdfs/2016/4/4.pdf

[3119] D. S. T. Ramesh, D. Angel Jovanna, Relaxed Skolem mean labeling of 5–star graphs with partition 3, 2, *Internat. J. Engin. Adv. Tech.,* **11**(2) (2021) 9-11. https://www.ijeat.org/wp-content/uploads/papers/v11i2/B32501211221.pdf

[3120] D. S. T. Ramesh, S. O. Sopna, I. Gnanaselvi, Skolem mean labeling of four star graphs, *IOSR J. Engin.,* **07**(09) (2017) 6-10. https://www.iosrjen.org/Papers/vol7_issue9/Version-1/B0709010610.pdf

[3121] D. Ramya and P. Jeyanthi, Mean labeling of some graphs, *SUT J. Math.,* **47**(2) (2011) 129-141. https://www.researchgate.net/publication/257365108_DRamya_and_PJeyanthi_Mean_Labeling_of_Some_Graphs_SUT_Journal_of_Mathematics_Volume_47_No2_2011_129_-141

[3122] D. Ramya and P. Jeyanthi, New super mean graphs, *J. Disc. Math. Sci. Cryptogr.*, **17**(5-6) (2014) 395-418.

[3123] D. Ramya, R. Kalaiyarasi and P. Jeyanthi, Skolem odd difference mean graphs, *J. Algor. and Comput.,* **45** (2014) 1-12. https://jac.ut.ac.ir/article_7916_727d297ed723f77921f5f0cb0b62cd65.pdf

[3124] D. Ramya, R. Ponraj, and P. Jeyanthi, Super mean labeling of graphs, *Ars Combin.,* **112** (2013) 65-72. https://combinatorialpress.com/article/ars/Volume%20112/volume-112-paper-5.pdf

[3125] D. Ramya, M. Selvi, and P. Jeyanthi, Odd mean labeling of some trees, *Util. Math.,* **106** (2018) 65-78.

[3126] D. Ramya, M. Selvi and P. Jeyanthi, Odd mean labeling of some graphs, unpublished.

[3127] N. Ramya and R. Shalini, On Lucky-edge Labeling of some graphs, *Internat. J. Innovative Res. Sci. Engin. Tech.*, **4**(9) (2015) 8146–8148. https://www.ijirset.com/upload/2015/september/30A_Ramya-4(5_pages)_9.pdf

[3128] M. Ramya and S. Meenakshi, Labeling techniques of some special graphs, *Internat. J. Pure Appl. Math.*, **116**(24) (2017) 93 –102. https://acadpubl.eu/jsi/2017-116-23-24/articles/24/10.pdf

[3129] A. Rana, On the k-distant total labeling of graphs, *Malaya J. Mat.*, **8**(2) (2020) 556-560. https://www.malayajournal.org/index.php/mjm/article/view/1455

[3130] V. Jeba Rani, S. Sunder Raj, and T. Shyla Isaac Mary, Vertex polynomial of ladder graphs, *Infokara Research*, **8**(11) 2019) 169-176.

[3131] S. B. Rao, Sigma graphs: A survey, *in* Labeling of Discrete Structures and Applications, Narosa Publishing House, New Delhi, 2008, 135-140.

[3132] S. B. Rao and U. K. Sahoo, Embeddings in Eulerian graceful graphs, *Australasian J. Comb.,* **62**(1) (2015) 128-139. https://ajc.maths.uq.edu.au/pdf/62/ajc_v62_p128.pdf

[3133] S. B. Rao, T. Singh, and V. Parameswaran, Some sigma labelled graphs: I, Graphs, Combinatorics, Algorithms and Applications, S. Armugam, B. D. Acharya and S. B. Rao, eds., (2004), 125-133, Narosa Publishing House, New Delhi. https://www.researchgate.net/publication/290485423_Some_sigma_labelled_graphs_I

[3134] S. N. Rao, Prime labelling, In: R. C. Bose Centenary Symposium on Discrete Math. and Applications, Kolkata (2002).

[3135] S. P. Rao Hebbare, Graceful cycles, *Util. Math.*, **10** (1976) 307-317.

[3136] L. Ratnasari and Y. Susanti, Total edge irregularity strength of ladder-related graphs, *Asian-Eur. J. Math.,* **13**(4) (2020), no. 2050072, 16 pp.

[3137] K. K. Raval and U. M. Prajapati, Vertex even and odd mean labeling in the context of some cyclic snake graphs, *J. Emerging Tech. Innovative Res. (JETIR)*, **4**(6) (2017) 319-324. https://www.jetir.org/view?paper=JETIRC006365

[3138] K. K. Raval and M. Prajapati, Signed product cordial labelings of some fractal graphs, *Gradiva Review J.* **8**(11) 2022 537-546. https://www.researchgate.net/publication/368273533_SIGNED_PRODUCT_CORDIAL_LABELING_OF_SOME_FRACTAL_GRAPHS

[3139] M. Ravi and R. Kala, Total prime graph, *Internat. J. Comput. Eng. Res.*, **2**(5) (2012) 1588-1593. https://www.ijceronline.com/papers/Vol2_issue5/BR02515881593.pdf

[3140] N. B. Rathod and K. K. Kanani, Some path related 4-cordial graphs, *Internat. J. Math. and Soft Computing,* **5**(2) (2015) 21-27. https://www.researchgate.net/publication/280530536_Some_path_related_4-cordial_graphs

[3141] N. B. Rathod and K. K. Kanani, V_4-cordial labeling of quadrilateral snakes,*Internat. J. Emerging Tech. Appl. Eng., Tech., Sci.,***9** (2016) 45-51. https://www.researchgate.net/publication/319352168_V_4-CORDIAL_LABELING_OF_QUADRILATERAL_SNAKES

[3142] N. B. Rathod and K. K. Kanani, 4-cordial labeling of star, book and fan related graphs, *Proceedings of 8th National Level Science Symposium*, Rajkot, India, **2** (2015) 38-42. https://www.researchgate.net/publication/290440448_4-CORDIAL_LABELING_OF_STAR_BOOK_AND_FAN_RELATED_GRAPHS

[3143] N. B. Rathod and K. K. Kanani, On V_4-cordial labeling of graphs, *British J. Math. Comput Sci.,* **13**(4) (2016) 1-15. https://www.researchgate.net/publication/288072970_On_V4-cordial_Labeling_of_Graphs

[3144] N. B. Rathod and K. K. Kanani, 5-cordial labeling of some standard graphs, *Proceedings of 8th National Level Science Symposium*, Rajkot, India, **2** (2015) 43-48. https://www.researchgate.net/publication/290440616_5-CORDIAL_LABELING_OF_SOME_STANDARD_GRAPHS

[3145] N. B. Rathod and K. K. Kanani, 4-cordiality of some new path related graphs, *Internat. J. Math. Trends Techno. (IJMTT)* **34**(1) (2016) 5-8. https://ijmttjournal.org/public/assets/volume-34/number-1/IJMTT-V34P502.pdf

[3146] N. B. Rathod and K. K. Kanani, V_4-cordial labeling of some alternate triangular snake related graphs, 9th National Level Science Symposium. February 14, 2016, Organized by Christ College, Rajkot, Sponsored by GUJCOST, Gandhinagar, Mathematics & Statistics, **3** (2016) 188-196. https://www.researchgate.net/publication/319352316_V_4-cordial_Labeling_of_Some_Alternate_Triangular_Snake_Related_Graphs

[3147] J. Ratilal and P. Parmar, H_k-Cordial labeling of some snake graphs, *J. Xidian Univ.,* **14**(3) (2020) 844-858. https://www.researchgate.net/publication/340115606_HK_Cordial_labeling_of_Some_Snake_Graphs/references

[3148] L. Ratnasari1, S. Wahyuni1, Y. Susanti1, and D. J. E. Palupi1, Total edge irregularity strength of quadruplet and quintuplet book graphs, *ITM Web of Conferences* 36, 03004 ICMSA 2020 (2021) 1-10. https://www.itm-conferences.org/articles/itmconf/pdf/2021/01/itmconf_icmsa2021_03004.pdf

[3149] B. Rayjada, M. Sarvaiya and J. Gadhiya, A computational perspective on cryptographic models using prime cordial labeling, *Artificial Intell. Smart Tech.* Sustainability Conf. (AISTS), Rajkot, India, 2025, 1-6. https://ieeexplore.ieee.org/document/11233014

[3150] M. A. Razzaq, S. T. Rizvi, and K. Ali, H-group magic total labelings of families of fan graphs, *Util. Math.,* **108** (2018) 213-219.

[3151] K. A. Reddy and S. Sharief Basha, New classes of reverse super edge magic graphs, *AIMS Math.,* **7**(3) (2022) 3590-3602. http://www.aimspress.com/article/doi/10.3934/math.2022198

[3152] T. A. Redl, Graceful graphs and graceful labelings: two mathematical programming formulations and some other new results, *Congressus Numer.,* **164** (2003) 17-32. https://repository.rice.edu/items/90406cd2-e2b5-4a02-8f29-d2d7dfab910e?utm_source=chatgpt.com

[3153] S. Redmond, An ideal-based zero-divisor graph of a commutative ring, *Comm. Algebra.* **9** (2003)n 4425-4443. https://doi.org/10.1081/AGB-120022801

[3154] M. Reid, personal communication (1992).

[3155] M. Regees, Super edge trimagic total labeling of some star type graphs, *J. Discrete Math. Sci. Cryptogr.,* **20**(3) (2017) 747-754.

[3156] M. Regees, L. M. Anisha, and T. Nicholas, Edge magic and bimagic harmonious labeling of ladder graphs, *Malaya J. Matematik*, **8**(1) (2020) 206-215. https://www.malayajournal.org/index.php/mjm/article/view/1375

[3157] M. Regees and C. Jayasekaran, More results on edge trimagic labeling of graphs, *Internat. J. Math. Archive*, **4**(11) (2013) 247-255.

[3158] M. Regees and C. Jayasekaran, Super edge trimagic total labeling of square of a cycle and gear graphs, *Math. Sci. Internat. Res. J.*, **4**(2) (2015) 113-115.

[3159] M. Regees and C. Jayasekaran, Super edge trimagic total labeling of generalized prism and web graphs, *J. Discr. Math. Sci. Cryp.*, **19**(1) (2016) 81-92.

[3160] A. ur Rehman Virk and A. Riasat, Odd graceful labeling of W-Tree $WT_(n,k)$ and its disjoint union, *Utilitas Math.*, **118** (2024) 51-62. doi:DOI: 10.61091/um118-05

[3161] P. Reka, and S. P. Soundariya, Palindromic antimagic labeling of products of paw and banner graph, *Contemp. Math.,* **5**(2) (2024) 2705-1056 (Online) https://ojs.wiserpub.com/index.php/CM/article/view/4180/2326

[3162] J. Renuka, P. Balaganesan, P. Selvaraju, On harmonious labeling, *Internat. J. Advance in Math. Math. Sci.,* **1**(2) (2012) 65-70.

[3163] J. Renuka and B. Palanivelu, Odd harmonious and strongly odd harmonious labeling of some classes of graphs, *Indian J. Public Health Res. Devel.,* **9**(9) (2018) 397-402. https://www.ijour.net/ijor.aspx?target=ijor:ijphrd&volume=9&issue=9&article=075&type=pdf

[3164] N. Revathi, Mean labeling of some graphs, *Internat. J. Sci. Res. (IJSR),* **4**(10) (2015) 1921-1923. https://www.ijsr.net/archive/v4i10/23101501.pdf

[3165] N. Revathi, Vertex odd mean and even mean labeling of some graphs, *IOSR J. Math.,* **11**(2) (2015) 70-74. https://www.iosrjournals.org/iosr-jm/papers/Vol11-issue2/Version-4/J011247074.pdf

[3166] R. Revathi, A. Dharmakkan, and R. M. Jeya Jothi, MMD labeling of complete tripartite graphs, *J. Physics Confer. Ser.*, (2010) **1770**(1):012083. https://iopscience.iop.org/article/10.1088/1742-6596/1770/1/012083/pdf

[3167] R. Revathi and R. Rajeswari, On modular multiplicative divisor graphs, in *IEEE Proceed. Internat. Conf. Pattern Recognition, Informatics Mobile Engin.,* (PRIME), 2013.

[3168] R. Revathi and R. Mary Jeya Jothi, MMD labeling of butterfly networks, *Proceed. 2018 IEEE Internat. Conf. Current Trends toward Converging Tech.,* Coimbatore, India

[3169] R. Revathi and R. Rajeswari, Modular multiplicative divisor labeling of some path related graphs, *Indian J. Sci. Technol.*, **7**(12) (2014) 1967-1972. https://i-scholar.in/index.php/indjst/article/view/59430/50407

[3170] K. Rhys, Every spider with at most six legs is graceful, preprint, https://zenodo.org/records/22837484

[3171] A. Riasat, S. Kanwal, and S. Javed, On odd-graceful labeling of disjoint union of graphs, *Util. Math.,* **101** (2016) 189-241. https://www.researchgate.net/publication/314950487_On_odd-graceful_labeling_of_disjoint_union_of_graph

[3172] P. Richter, E. Leven, A. Tran, B. Ek, J. Jacob, and D. A. Narayan, Rank numbers for bent ladders, *Discuss. Math. Graph Th.* **34**(2) (2014) 309-329. https://www.researchgate.net/publication/265770583_Rank_numbers_for_bent_ladders

[3173] N. M. Rilwan and A. S. Hussain, Prime cordial and signed product cordial labeling on identity graph, *Adv. Math.: Sci. J.,* **8** (2019) 749-753. https://www.researchgate.net/publication/351098253_PRIME_CORDIAL_AND_SIGNED_PRODUCT_CORDIAL_LABELING_ON_IDENTITY_GRAPH

[3174] N. M. Rilwan and R. Radha, Edge vertex prime labeling of Cayley (di)graphs, *Malaya J. Mat.,* **8**(2) (2020) 531-535. https://www.researchgate.net/publication/341936051_Edge_vertex_prime_labeling_of_Cayley_digraphs

[3175] G. Ringel, Problem 25, in Theory of Graphs and its Applications, *Proc. Symposium Smolenice 1963*, Prague (1964) 162.

[3176] G. Ringel and A. Llado, Another tree conjecture, *Bull. Inst. Combin. Appl.,* **18** (1996) 83-85.

[3177] G. Ringel, A. Llado, and O. Serra, Decomposition of complete bipartite graphs into trees, DMAT Research Report 11/96, Univ. Politecnica de Catalunya.

[3178] A. Riskin, Cordial deficiency, *Bull. Malays. Math. Sci. Soc.,* **30** (2007) 201-204. arXiv:math/0610760v1 [math.CO] 25 Oct 2006.

[3179] A. Riskin, On the cordial deficiency of complete multipartite graphs, arXiv:0706.2431v1 [math. CO] 16 June 2007.

[3180] A. Riskin, Z_2^2-cordiality of K_n and $K_{m,n}$, arXiv:0709.0290v1 [math.CO] 3 Sep. 2007.

[3181] A. Riskin and G. Weidman, On edge graceful labelings of disjoint unions of $2r$-regular edge graceful graphs, *Bull. Inst. Combin. Appl.,* **52** (2008) 45-50. arXiv:math/0605234v1.

[3182] R. Ristiawan, Graf pendulum dengan pelabelan graceful super Fibonacci, *Faktor Exacta,* **12**(1) (2019) 51-57. https://journal.lppmunindra.ac.id/index.php/Faktor_Exacta/article/view/3365/2492

[3183] S. T. R. Rizvi, K. Ali, and M. Hussain, Cycle-supermagic labelings of the disjoint union of graphs, *Util. Math.,* **104** (2017) 215-226. arXiv:1506.06087.

[3184] S. T. R. Rizvi, K. Ali, N. Iqbal, and A. Gulraze, Super edge-magicness of stars like graphs, *Ars Combin.,* **136** (2018) 405-417.

[3185] S. T. R. Rizvi, M. Khalid, K. Ali, M. Miller, and J. Ryan, On cycle-supermagicness of subdivided graphs, *Bull. Aust. Math. Soc.,* **92**(1) (2015) 11-18. https://www.researchgate.net/publication/276509799_ON_CYCLE-SUPERMAGICNESS_of_SUBDIVIDED_GRAPHS

[3186] D. Roberts and R. M. Low, Group-antimagic labelings of multi-cyclic graphs, *Theory Appl. Graphs*, **3**(1) (2016) Article 6. https://digitalcommons.georgiasouthern.edu/tag/vol3/iss1/6

[3187] L. Robertson and B. Small, On Newman's conjecture and prime trees, *Integers: The Elect. J. Combin. Number Theory,* **9** (2009) A10 112-128. https://math.colgate.edu/~integers/j10/j10.pdf

[3188] E. Robeva, An Extensive Survey of Graceful Trees, Undergraduate Honors Thesis, Stanford University, 2011.

[3189] K. Roblee and A. A. Abueida, More harmonious labelings of families of disjoint unions of an odd cycle and certain trees, *J. Combin. Math. Combin. Comput.,* **115**(1) (2020) 61-68. https://doi.org/10.61091/jcmcc119-30

[3190] Y. Roditty and T. Bachar, A note on edge-magic cycles, *Bull. Inst. Combin. Appl.,* **29** (2000) 94-96.

[3191] R. I. Rofa, A functional definition of graceful labelling of rooted symmetric trees, arXiv:2109.09511.

[3192] R. Rofa, Graceful and strongly graceful permutations. arXiv:2107.14007.

[3193] R. I. Rofa, On number representation, arXiv:1310.8277.

[3194] D. G. Rogers, A graceful algorithm, *Southeast Asian Bull. Math.*, **2** (1978) 42-44.

[3195] A. H. Rokad, 3-Equitable labeling in context of ring sum of graphs *Research & Reviews: Discrete Math. Structures: Open Journal Systems*, **2**(3) (2015) 6272-6275. https://www.researchgate.net/publication/302025630_3-Equitable_Labeling_in_Context_of_Ring_Sum_of_Graphs

[3196] A. H. Rokad, Fibonacci cordial labeling of some special graphs, *Oriental J. Comput. Sci. Tech,*. (2017) **10**(4) (2017) 824-828. https://www.computerscijournal.org/pdf/vol10no4/OJCST_Vol10_N4_p_824-828.pdf

[3197] A. H. Rokad, Product cordial labeling of double wheel and double fan related graphs, *Kragujevac J. Math.,* **43**(1) (2019) 7-13. http://elib.mi.sanu.ac.rs/files/journals/kjm/55/kjmn55p7-13.pdf

[3198] A. H. Rokad and G. V. Ghodasara, Fibonacci cordial labeling of some special graphs, *Annals Pure Appl. Math.*, **11**(1) (2016) 133-144. http://www.researchmathsci.org/apamart/apam-v11n1-16.pdf

[3199] A. H. Rokad and K. M. Patadiya, Cordial labeing of some graphs, *Aryabhatta J. Math. Informatics,* **09** (01) (2017) 589-597. https://www.researchgate.net/publication/319553115_Cordial_Labeling_of_Some_Graphs

[3200] Q. Rong and D. Xiong, The gracefulness of graph $P_{2r,b}$, *J. Systems Sci. Math. Sci.,* **30**(5) (2010) 703-709.

[3201] B. Roopa and L. Shobana, Face bimagic labeling on some graphs, *Turkish J. Comput. Math. Edu.*, (TURCOMAT) **12**(6) (2021) 101-107. https://turcomat.org/index.php/turkbilmat/article/view/1273/1051

[3202] D. Ropp, Graceful labelings of cycles and prisms with pendant points, *Congress. Numer.,* **75** (1990) 218-234.

[3203] A. Rosa, On certain valuations of the vertices of a graph, *Theory of Graphs (Internat. Symposium, Rome, July 1966)*, Gordon and Breach, N. Y. and Dunod Paris (1967) 349-355. https://www.researchgate.net/publication/244474213_On_certain_valuations_of_the_vertices_of_a_graph

[3204] A. Rosa, Labelling snakes, *Ars Combin.*, **3** (1977) 67-74.

[3205] A. Rosa, Cyclic Steiner triple systems and labelings of triangular cacti, *Scientia*, **1** (1988) 87-95.

[3206] A. Rosa and J. Širáň, Bipartite labelings of trees and the gracesize, *J. Graph Theory*, **19** (1995) 201-215.

[3207] T. Rose and H.-H. Su, An algorithm to calculate the balance index set of a graph, Proceedings of the Forty-Third Southeastern International Conference on Combinatorics, Graph Theory and Computing. *Congr. Numer.,* **213** (2012) 107-121.

[3208] M. Roswitha, E.T. Baskoro, T. K. Maryati, N. A. Kurdhi, and I. Susanti, Further results on cycle-supermagic labeling, *AKCE Int. J. Graphs Comb.*, **10**(2) (2013) 211-220. https://www.researchgate.net/publication/264909824_Further_results_on_cycle-supermagic_labeling

[3209] M. Roswitha and E.T. Baskoro, H-magic covering of some classes of graphs, *AIP Conf. Proc. on ICREM5,* ITB Bandung, **1450** (2012) 135-138. NOW 9/1/25v

[3210] M. Roswitha, T. S. Martini, and S. A. Nugroho, $(a, d) - H$-Anti magic total labeling on double cones graph, ICMETA 2018 IOP Conf. Series: *J. Physics: Conf. Series* 1306 (2019) 012012 1-5. https://www.researchgate.net/publication/335697411_a_d_-_H_-Anti_Magic_Total_Labeling_on_Double_Cones_Graph

[3211] I. Rosyida and D. Indriati, On total edge irregularity strength of some cactus chain graphs with pendant vertices, *IOP Conf. Series: J. Physics: Conf. Series* 1211 (2019) 012016. doi:10.1088/1742-6596/1211/1/012016

[3212] I. Rosyida and D. Indriati, Computing total edge irregularity strength of some n-uniform cactus chain graphs and related chain graphs, *Indon. J. Combin.,* **4**(1) (2020) 53-75. https://www.researchgate.net/publication/342523731_Computing_total_edge_irregularity_strength_of_some_n-uniform_cactus_chain_graphs_and_related_chain_graphs/comments#fullTextFileContent

[3213] I. Rosyida, E. Ningrum, Mulyono, and D. Indriati, On the total edge irregularity strength of general uniform cactus chain graphs with pendant vertices, *J. Discrete Math. Sci. Crypt.*, (2020) **23**(6) 1335-1358. DOI:10.1080/09720529.2020.1820704

[3214] P. L. Rozario Raj and M. M. Sheriff, Some more results on neighbourhood-prime labeling of graphs, *Internat. J. Anal. Exper. Modal Anal.,* **XI**(X) Iss (2019) 38-43. https://www.researchgate.net/publication/336847767_Some_More_Results_on_Neighbourhood-Prime_Labeling_of_Graphs

[3215] K. Rozmana and Šparl, Distance magic labelings of Cartesian products of cycles, https://arxiv.org/abs/2403.03279

[3216] K. Rozman and P. Sparl, Tetravalent distance magic graphs of small order and an infinite family of examples, *Ars Math. Contemp.,* (2025) 1-13. https://doi.org/10.26493/1855-3974.3424.d5e

[3217] J. Ryan, Exclusive sum labelings of graphs: A survey, *AKCE J. Graphs Combin.,* **6** (2009) 113-126.

[3218] R. Ryan, A. Marquez, and A. C. Pedrano, On total product cordial labeling of some snake graphs, *Annals Commun. Math.*, **9** (2026) (8) 23 pp. https://www.technoskypub.com/wp-content/uploads/2026/06/acm-09024.pdf

[3219] J. Ryan, O. Phanalasy, M. Miller and L. Rylands, On antimagic labeling for generalized web and flower graphs, In C. S. Iliopoulos and W. F. Smyth, editors, *Proceed. IWOCA10, Lecture Notes in Computer Science*, **6460** (2011) 303-313.

[3220] J. Ryan, O. Phanalasy, L. Rylands and M. Miller, On antimagic labeling for generalized web and flower graphs, Combinatorial algorithms, 303-313, Lecture Notes in Comput. Sci., 6460, Springer, Heidelberg, 2011.

[3221] L. Rylands, O. Phanalasy, J. Ryan and M. Miller, Construction for antimagic generalized web graphs, *Int. J. Graphs and Combinatorics*, **8** (2011) 141-149. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2011.12088938

[3222] K. M. Sabeel, A. V. Prajeesh, and K. Paramasivam, A characterization for V_4-vertex magicness of trees with diameter 5, In *Computat. Sci. - Modelling, Comput. Soft Comput.*, Springer, (2021) 243-249.

[3223] K. M. Sabeel, K. Paramasivam, A. V. Prajeesh, N. Kamatchi, and S. Arumugam, A characterization of group vertex magic trees of diameter up to 5, *Australas. J. Combin.*, **85** (2023) 49-60. https://ajc.maths.uq.edu.au/pdf/85/ajc_v85_p049.pdf

[3224] R. S. Sabibha, K. Jeya Daisy, P. Jeyanthi, M. Z. Youssef, k-product cordial labeling of path graphs, *Open J. Disc. Math.*, **15**(1) 2025 https://www.scirp.org/journal/paperinformation?paperid=138493

[3225] K. Saengsura and T. Poomsa-ard, Graceful labeling of some spider graphs, *European J. Pure Appl. Math.*, **18**(2) , (2025) Article Number 5305. https://doi.org/10.29020/nybg.ejpam.v18i2.5305

[3226] A. Saibulla, B. Raj, S. Suganya, and M. G. F. Kareem, Total edge-cordial labeling of Koch curve and Koch snowflake graphs, *Eur. Chem. Bull.,* **12**(6) (2023) 2733-2740. https://www.researchgate.net/publication/383282861_TOTAL_EDGE-CORDIAL_LABELING_OF_KOCH_CURVE_AND_KOCH_SNOWFLAKE_GRAPHS

[3227] L. Saha and A. R. Basunia, Radio graceful labelling of graphs, *Theory Appl. Graphs,* **7**(1) (2020), Art. 7, 8 pp. https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1106&context=tag

[3228] L. Saha, A. R. Basunia, S. Das, and K. Tiwary, Radio k-chromatic number of full m-ary trees, *Theory Comput. Syst.,* **66**(1) (2022) 114-142. https://link.springer.com/article/10.1007/s00224-021-10056-7

[3229] L. Saha and P. Panigrahi, Antipodal number of some powers of cycles, *Discrete Math.,* **312**(9) (2012) 1550-1557. https://www.sciencedirect.com/science/article/pii/S0012365X11004985

[3230] L. Saha and P. Panigrahi, A graph radio k-coloring algorithm, *Combin. Algorithms*, (IWOCA 2012): Lecture Notes in Comput. Sci., Springer, Berlin, Germany, **7643**, 2012. https://link.springer.com/content/pdf/10.1007/978-3-642-35926-2_15.pdf

[3231] L. Saha and P. Panigrahi, On the radio number of toroidal grids, *Australasian J. Combin.,* **55** (2013) 273-288. https://ajc.maths.uq.edu.au/pdf/55/ajc_v55_p273.pdf

[3232] L. Saha and P. Panigrahi, On the radio number of square of graphs, The Eighth International Workshop on Graph Labelings (IWOGL 2014), 205-212, *Electron. Notes Discrete Math.,* **48**, Elsevier Sci. B. V., Amsterdam, 2015. http://dx.doi.org/10.1016/j.endm.2015.05.030

[3233] L. Saha and P. Panigrahi, A new graph radio k-coloring algorithm, *Disc. Math., Algor. and Appl.*, **11**(1) Article ID 1950005, p. 10, 2019. https://www.researchgate.net/publication/328807166_A_New_Graph_Radio_k_-coloring_Algorithm

[3234] A. R. Sahaya, P. Maya, and T. Nicholas, Product integer cordial labeling of some well known graphs, *J. Emerging Tech. Innovative Res.*, **05**(11) (2018) 132-138. https://www.jetir.org/papers/JETIR1811022.pdf

[3235] S. S. Sarasvati, I. Halikin, and K. Wijaya, Odd harmonious labeling of $P_n \trianglerighteq C_4$ and $P_n \trianglerighteq D_2(C_4)$, *Indonesian J. Combin.*, **5**(2) (2021) 94-101. https://doi.org/10.19184/ijc.2021.5.2.5

[3236] U. Sarkar, A note on obtaining bipartite radio graceful graphs of arbitrarily large radio numbers with radio graceful complements, *Discrete Appl. Math.*, **377** (2025) 350-355. https://colab.ws/articles/10.1016%2Fj.dam.2025.07.012

[3237] F. Salama, Computing the total edge irregularity strength for quintet snake graph and related graphs, *J. Discrete Math. Sci. Crypt.*, (2021)

[3238] U. Salma and H. M. Nagesh, On edge irregularity strength of ladder-related graphs, *TWMS J. App. Eng. Math.*, **14**(2) (2024) 696–702. https://jaem.isikun.edu.tr/web/images/articles/vol.14.no.2/21.pdf

[3239] U. Salma, H. M. Nagesh, B. Azghar Pasha, and N. Narahari, On edge irregularity strength of Mycielskian of paths and cycles, *J. Mines Metals Fuels*, **71**(12A) (2022) 208–213. https://mail.informaticsjournals.co.in/index.php/jmmf/article/view/43610/29692

[3240] U. Salma, H. M. Nagesh, and D. Prahlad, A note on edge irregularity strength of firefly graph, *Not. Numb. Theory Discrete Math.* **29**(1) (2023) 147–153. https://nntdm.net/papers/nntdm-29/NNTDM-29-1-147-153.pdf

[3241] U. Salma, H. M. Nagesh, U. Vijaya Chandra Kumar, and N. Narahari, Edge irregularity strength of Graphs: Theoretical results and applications to communication networks, *Acta Univ. Sapientiae Inform.*, **18**(20) (2026). https://link.springer.com/article/10.1007/s44427-026-00036-5

[3242] M. Salarrezaei and K. Eshghi, An integer programming model and a Tabu search algorithm to generate α-labeling of special classes of quadratic qraphs, *Iranian J. Operations Res.,* **7**(1) (2016) 1-17. https://www.researchgate.net/publication/324559673_An_Integer_Programming_Model_and_a_Tabu_Search_Algorithm_to_Generate_a-labeling_of_Special_Classes_of_Quadratic_Graphs

[3243] M. Salarrezaei, Mathematical programming models for graceful labeling problem of 2-regular graphs. Master of Science thesis, Sharif University of Technology (2012).

[3244] E. Salehi, Zero-sum magic graphs and their null sets, *Ars Combin.,* **82** (2007) 41-53. https://combinatorialpress.com/article/ars/Volume%20082/volume-82-paper-4.pdf

[3245] E. Salehi, PC-labelings of a graphs and its PC-sets, *Bull. Inst. Combin. Appl.,* **58** (2010) 112-121. https://www.salehi.faculty.unlv.edu/PCL1.pdf

[3246] E. Salehi and D. Bayot, The friendly index set of $P_m \times P_n$, *Util. Math.,* **81** (2010) 121-130. https://salehi.faculty.unlv.edu/FIS-G.pdf

[3247] E. Salehi and P. Bennett, On integer-magic spectra of caterpillars, *J. Combin. Math. Combin. Comput.,* **61** (2007) 65-71. https://salehi.faculty.unlv.edu/pioneer/IMS-Caterpillars.pdf

[3248] E. Salehi, S. Churchman, T. Hill, and J. Jordan, Product cordial sets of trees, *Congr. Numer.,* **220** (2014) 183-193. https://salehi.faculty.unlv.edu/PCLT.pdf

[3249] E. Salehi and S. De, On a conjecture concerning the friendly index sets of trees, *Ars Combin.,* **90** (2009) 371-381. https://salehi.faculty.unlv.edu/pioneer/FIS-C.pdf

[3250] E. Salehi and S.-M. Lee, On friendly index sets of trees, *Cong. Numer.,* **178** (2006) 173-183. https://salehi.faculty.unlv.edu/pioneer/FIS-Trees.pdf

[3251] E. Salehi and S. M. Lee, Integer-magic spectra of functional extensions of graphs, *J. Combin. Math. Combin. Comput.,* **64** (2008) 127-139. https://salehi.faculty.unlv.edu/IMS-FEG.pdf

[3252] E. Salehi, S. M. Lee, and M. Khatirinejad, IC-colorings and IC-indices of graphs, *Discrete Math.,* **299** (2005) 297-310. https://www.researchgate.net/publication/220187396_IC-Colorings_and_IC-Indices_of_graphs

[3253] E. Salehi and D. McGinn, On friendly index set of hypercube Q_n, *J. Combin. Math. Combin. Comput.,* **106** (2018) 109-124. https://www.researchgate.net/publication/329400621_On_Friendly_Index_Sets_of_Hypercube_Qn

[3254] E. Salehi and Y. Mukhin, Product cordial sets of long grids, *Ars Combin.,* **107** (2012) 339-351. https://salehi.faculty.unlv.edu/PCLG.pdf

[3255] E. Salehi, Y. Mukhin, and S. W. Saputro, Cordial sets of hypercubes, *Bull. Instit. Combin. Appl.,* **75** (2015) 95-106. https://salehi.faculty.unlv.edu/CS-Cubes.pdf

[3256] E. R. Salisa JR. and A. Peldrano, On Lucas cordial labelings of some snake graphs, *Annals Commun. Math.*, **8**(4) (2025) 451-458. https://www.technoskypub.com/journals/acm-2025-080403/

[3257] U. Salma, H. M. Nagesh, and Narahari N, On edge irregularity strength of cycle-star graphs, *Proyecciones J. Math.,* **43**(3) (2024) 725-741. https://scispace.com/pdf/on-edge-irregularity-strength-of-cycle-star-graphs-4yef6atnt8.pdf

[3258] A. N. M. Salman and T. K. Maryati, On graph-(super)magic labeling of a path-amalgamation of isomorphic graphs, in: The Proceedings of the 6th IMT-GT International Conference on Mathematics, Statistics and its Applications, 2010, 228-233. https://www.ijour.net/ijor.aspx?target=ijor:ijphrd&volume=9&issue=9&article=075&type=pdf

[3259] A. N. M. Salman, A. A. G. Ngurah, and N. Izzati, On (super) edge-magic total labelings of a subdivision of a star S_n, *Util. Math.,* **81** (2010) 275-284.

[3260] H. Salmasian, A result on prime labelings of trees, *Bull. Inst. Combin. Appl.,* **28** (2000) 36-38.

[3261] S. K. O. D. Samarathunge, A. M. U. M. Athapattu, and A. A. I. Perera, Characterization of E-super vertex magic labelling of odd order binary trees, *Ceylon J. Sci.*, **51**(4) (2022) 389-393. http://doi.org/10.4038/cjs.v51i4.8056

[3262] A. Selvaganapathy and E. J. Lalithkumar, Gaussian anti magic labeling on joining of cricket graph, dart graph, diamond graph and domino graph, *Phyton-Annales Rei Botnicae,* **66**(1) (2026) 258-262. https://phyton-annales.org/index.php/p/article/view/47/52

[3263] A. Selvaganapathy and P. Ramesh, Gaussian anti magic labeling on tree related graphs, *Com. Res. Dev.*, **25**(2) (2025) 140-148. https://journalcrd.org/wp-content/uploads/14-CRD2639.pdf

[3264] A. Selvaganapathy and K. Thirusangu, Gaussian anti magic labeling on path union of graphs, *J. Asiatic Soc. Mumbai*, **XCV**(39) (2022) 45-51. https://www.researchgate.net/publication/408633647_GAUSSIAN_ANTI_MAGIC_LABELING_ON_PATH_UNION_OF_GRAPHS

[3265] A. Selvaganapathy and K. Thirusangu, Gaussian anti magic labeling on N-join graphs, *J. Asiatic Soc. Mumbi*, **XCVI**(32) (2023) 145-153. https://www.researchgate.net/profile/Selvaganapathy-Aniyappan/publication/408634132_Gaussian_Anti_Magic_Labeling_on_N_-_Join_Graphs/links/6a4f6501f6cec25ad2d7e1da/Gaussian-Anti-Magic-Labeling-on-N-Join-Graphs.pdf

[3266] R. K. Sampath, G. Narasimhan, and K. M. Nagaraja, Heron mean labelling of graphs, *Internat. J. Recent Sci. Res.,* **8**(9) (2017) 19808-19811. https://recentscientific.com/sites/default/files/8458-A-2017.pdf

[3267] A. E. Samuel and S. Kalaivani, Prime labeling for some planter related graphs, *Internat. J. Math. Res.* (IJMR), **8**(3) 221-231, 2016.

[3268] A. E. Samuel and S. Kalaivani, prime labeling for some vanessa related graphs, *Indian J. Appl. Res.*, (IJAR), **7**(4) (2017) 136-145.

[3269] A. E. Samuel and S. Kalaivani, Prime labeling for some octopus related graphs, *Annals Pure and Appl. Math.*, **16**(2) (2018) 317-312. https://iosrjournals.org/iosr-jm/papers/Vol12-issue6/Version-3/J1206035764.pdf

[3270] A. E. Samuel and S. Kalaivani, Square sum labeling for some lilly related graphs, *Internat. J. Adv. Tech. Engin. Exploration*, **4**(29) (2017) 68–72.

[3271] A. E. Samuel and S. Kalaivani, Prime labeling to drums graphs, *Annals Pure Appl. Math.* (APAM),**16**(2) (2018) 307-312. http://researchmathsci.org/APAMart/apam-v16n2-7.pdf

[3272] A. E. Samuel and S. Kalaivani, Prime labeling to brush graphs, *Internat. J. Math. Trends Tech.*, **5**(4) (2018) 259–262. https://ijmttjournal.org/public/assets/volume-55/number-4/IJMTT-V55P533.pdf

[3273] A. K. Samuel, J. Vinolin, and D. S. T. Ramesh, Relaxed Skolem mean labeling of four star graphs $K_{1,\ell_1} \cup K_{1,\ell_2} \cup K_{1,\ell_3} \cup K_{1,l}$ *where* $\ell_1 \leq \ell_2 \leq \ell_3$ *with* $|\ell - \ell_1\ell_2 - \ell_3| = 2$, *Internat. J. Engin. Adv. Tech.,* (IJEAT) **9**(3) (2020) 3577-3588. https://www.ijeat.org/wp-content/uploads/papers/v9i3/C6394029320.pdf

[3274] E. Samuvel and S. Kalaivani, Square sum labeling for some lilly related graph, *Internat. J. Adv. Tech. Exploration*, **4**(29) online, (2017) 68-72. https://www.accentsjournals.org/PaperDirectory/Journal/IJATEE/2017/4/3.pdf

[3275] Ľ. Šándorová and M. Trenklér, On a generalization of magic graphs, *Colloquia Math. Societatis J. Bolyai*, **52** Combinatorics, North-Holland, Amsterdam 1988, 447-452. http://math.ku.sk/~trenkler/1988-On%20a%20generalization%20of%20magic%20graphs.pdf

[3276] Y. Sanaka, On γ-labelings of complete bipartite graphs, *Ars Combin.,* **111** (2013) 251-256. https://combinatorialpress.com/article/ars/Volume%20111/volume-111-paper-20.pdf

[3277] S. S Sandhya, S. Somasundaram, and S. Anusa, Root square mean labeling of some new disconnected graphs, *Internat. J. Math. Trends Tech.*, **15** (2014) 85-92. https://ijmttjournal.org/public/assets/volume-15/number-2/IJMTT-V15P511.pdf

[3278] S. S Sandhya, S. Somasundaram, and S. Anusa, Root square mean labeling of subdivision of some more graphs, *Internat. J. Math. Res.*, **6** (2014) 253-266. http://www.irphouse.com/ijmr/ijmrv6n3_04.pdf

[3279] S. S Sandhya, S. Somasundaram, and S. Anusa, Some new results on root square mean labeling, *Internat. J. Math. Arch.*, **5** (2014) 130-135.

[3280] S. S Sandhya, S. Somasundaram, and S. Anusa, Root square mean labeling of some more disconnected graphs, *Internat. Math. Forum*, **10** (2015) 25-34. https://www.m-hikari.com/imf/imf-2015/1-4-2015/anusaIMF1-4-2015.pdf

[3281] S. S. Sandhya, S. Somasundaram, and S. Anusa, Some results on root square mean graphs, *Internat. J. Math. Res*, **7**(2) (2015) 125-134 http://www.irphouse.com/ijmr/ijmrv7n2_04.pdf

[3282] S. Saduakdee and K. Khemmani, The unique γ-min labelings of graphs, *Thai J. Math.,* 2018, Special issue, 187-203. https://thaijmath2.in.cmu.ac.th/index.php/thaijmath/article/view/768/765

[3283] S. Saduakdee and V. Khemmani, The even vertex magic total labelings of t-fold wheels, *AIMS Math.*, **8**(11) (2023) 27513-27527. doi:10.3934/math.20231407

[3284] V. Sandhiya and M. C. Nalliah, Local total antimagic chromatic number for the disjoint union of star graphs, *European J. Prue Appl. Math.*, **17**(4) (2024) 2828-2842. https://www.ejpam.com/ejpam/article/view/5383?utm_source=chatgpt.com

[3285] S. S. Sandhya, C. Jayasekaran, and C. S. David Raj, Some new families of harmonic mean graphs, *Internat. J. Math. Res.*,**5**(1) (2013) 223-232. http://www.irphouse.com/ijmr/ijmrv5n2_03.pdf

[3286] S. S. Sandya, E. R. Merly and S. D. Deepa, Heronian mean labeling of graphs, textitInternat. Math. Forum, **12**(15) (2017) 705-713. https://www.m-hikari.com/imf/imf-2017/13-16-2017/p/deepaIMF13-16-2017.pdf

[3287] S. S. Sandhya, E. R. Merly, and B. Shiny, Subdivision of super geometric mean labeling for quadrilateral snake graphs, *Internat. J. Math. Trends Tech.*, **24**(1) (2015) 1-16. https://ijmttjournal.org/public/assets/volume-24/number-1/IJMTT-V24P501.pdf

[3288] S. S. Sandhya, E. R. Merly, and B. Shiny, Super geometric mean labeling on double quadrilateral snake graphs, *Asian Pacific J. Res.,* **1** (XXI) (2015) 128-135.

[3289] S. S. Sandhya, E. R. Merly, and B. Shiny, New classes of super geometric mean graphs, *J. Discrete Math. Sci. Cryptogr.,* **19**(1) (2016) 117-140. https://www.researchgate.net/publication/303358146_New_classes_of_super_geometric_mean_graph

[3290] S. S. Sandhya and S. Somasundaram, Geometric mean labeling on double quadrilateral snake graphs, *Internat. J. Math. Res.*, **6**(2) (2014) 179-182 http://irphouse.com/ijmr/ijmrv6n2_09.pdf

[3291] S. S. Sandhya, S. Somasundaram, and R. Ponraj, Some more results on harmonic mean graphs, *J. Math. Res.*, **4**(1) (2012) 21-29. https://pdfs.semanticscholar.org/6b4b/befb514517c02b7e6f6d92d63fc12762872a.pdf

[3292] S. S. Sandhya, S. Somasundaram, and S. Anusa, Root square mean labeling of graphs, *Internat. J. Contemp. Math. Sci.*, **9**(14) (2014) 667-676. https://www.m-hikari.com/ijcms/ijcms-2014/13-16-2014/anusaIJCMS13-16-2014.pdf

[3293] S. S. Sandhya, S. Somasundaram, and S. Anusa, Some more results on root square mean graphs, *J. Math. Res.*, **7**(1) (2015) 72-81, 2015. https://www.ccsenet.org/journal/index.php/jmr/article/view/41459

[3294] S. S. Sandhya, S. Somasundaram, and R. Ponraj, Some more results on harmonic mean graphs, *J. Math. Res.,* **4**(1) (2012) 21-29. https://ccsenet.org/journal/index.php/jmr/article/view/13019

[3295] S. S. Sandhya, S. Somasundaram, and R. Ponraj, Harmonic mean labeling of some cycle related graphs. *Int. J. Math. Anal.,* (Ruse) **6**(37-40) (2012) 1997-2005. http://www.m-hikari.com/ijma/ijma-2012/ijma-37-40-2012/sandhyaIJMA37-40-2012.pdf

[3296] S. S. Sandhya, S. Somasundaram, R. Ponraj, Some results on harmonic mean graphs, *Int. J. Contemp. Math. Sci.,* **7** (2012) 197-208. https://m-hikari.com/ijcms/ijcms-2012/1-4-2012/sandhyaIJCMS1-4-2012.pdf

[3297] I. P. Sandy, A. Rizal, E. N. Manurung, and K. A. Sugeng, Alternative construction of graceful symmetric trees, *J. Phys.: Conf. Ser.,* **1008** (2018) 012031 https://iopscience.iop.org/article/10.1088/1742-6596/1008/1/012031/pdf

[3298] A. P. Sangeeta, Analyzing the applications of graph theory in communication networks through the divisor 3-equitable labeling of graphs. In: D. Singh, R. K. Chaudhary, and K. Dev Kumar,(eds) Computer Aided Constellation Management and Communication Satellites. Lecture Notes in Electrical Engineering, Springer, Singapore, **987**, (2023) 105-112. https://doi.org/10.1007/978-981-19-8555-3_12

[3299] A. Sangeeta, R. Parthiban, and P. Selvaraju, 3-equitable and prime labeling of some classes of graphs, *Math. Stat.,* **11**(3) (2023) 534-540. https://www.researchgate.net/publication/370598062_3-Equitable_and_Prime_Labeling_of_Some_Classes_of_Graphs

[3300] S. Sangeetha and U. Usharani, Relaxed mean labeling of some corona graphs, *AIP Confer. Proceed.,* (2019) 2112, 020067 (2019) https://aip.scitation.org/doi/pdf/10.1063/1.5112252

[3301] K. Sankar and G. Sethuramam, Graceful and cordial labeling of subdivision of graphs, *Electronic Notes Discrete Math.,* **53** (2016) 123-131.

[3302] G. Santhosh, Sequential coronations of graphs, *Natl. Acad. Sci. Lett.,* **28** (2005) 269-270.

[3303] G. Santhosh and G. Suresh Singh, A note on the super edge-magic deficiency of graphs, *Far East J. Appl. Math.* **24** (2006) 233-242.

[3304] M. Santhi and J. Albert, Signed product cordial labeling and signed total product cordial labeling for some new graphs, *IMPACT: International J. Res. Appl., Nat. Social Sci.,* **3**(1) (2015) 133-138.

[3305] M. Santhi and J. Albert, Signed product cordial in cycle related graphs, *Internat. J. Math. Comput. Appl. Res.*, **5** (2016) 29-36. https://www.academia.edu/11778889/SIGNED_PRODUCT_CORDIAL_IN_CYCLE_RELATED_GRAPHS

[3306] G. Santhosh and G. Suresh Singh, On super magic strength of graphs, *Far East J. Appl. Math.,* **18** (2005) 199-207.

[3307] G. Santhosh, On weak magic graphs, *Ars Combin.,* **92** (2009) 245-254. https://combinatorialpress.com/article/ars/Volume%20092/volume-92-paper-17.pdf

[3308] G. Santhosh and G. Suresh Singh, A note on subdivision of integral sum graphs, unpublished.

[3309] G. Santhosh and G. Suresh Singh, On divisor graphs, unpublished.

[3310] R. Santrin Sabibha, K. Jeya Daisy, P. Jeyanthi, and M. Z. Youssef, On bijective product square k-cordial labeling, *Discrete Math., Alg. Appl.,* **17**(01) 2450013 (2025) 13pp. https://www.worldscientific.com/doi/epdf/10.1142/S1793830924500137

[3311] G. A. Saputri, K. A. Sugeng, and D. Fronček, The odd harmonious labeling of dumbbell and generalized prism graphs, *AKCE Int. J. Graphs Comb.*, **10**(2) (2013) 221-228. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2013.12088738?needAccess=true

[3312] P. B. Sarasija and R. Binthiya, Even harmonious graphs with applications, *Internat. J. Comput. Sci. Infor. Security,* **9** (2011) 161-163.

[3313] R. Sarita, A. Sehgal, and A. Malik, Graceful labeling of prime index graph of group $Z_p \times Z_p^n$, *Contemp. Math.,* [Internet], **4**(3) (2024) 612-619. https://ojs.wiserpub.com/index.php/CM/article/view/2727/1630

[3314] A. Sasikala and M. Harikumaran, Vertex odd mean labeling of double triangular snake $(D(kC_n) + 2nP_2)$, jelly fish $J(M, N)$, $P_n \circ H_1$, and an alternate triangular snake $A(T_n)$. *Archeo Sci. J.* **17**(1) (2024) 106-116. https://archeosciences.org/wp-content/uploads/14-1.pdf

[3315] T. V. Sateesh Kumar and S. Meenakshi, Lucky and proper lucky labeling of quadrilateral snake graphs, *IOP Conf. Ser.: Material Sci. Engin.*, **1085** (2021) 012039 doi:10.1088/1757-899X/1085/1/012039

[3316] T. V. Sateesh Kumar and S. Meenakshi, Proper lucky labeling of graph, S.-L. Peng et al. (eds.), *Proceed. First Internat. Conf. Math. Modeling Comput. Sci., Adv. Intell. Sys. Comput.,* **1292** (2021) 341-351. https://doi.org/10.1007/978-981-33-4389-4_31

[3317] T. V. Sateesh Kumar and S. Meenakshi, Jewel, comb and fan graph are lucky and proper lucky, *AIP Conf. Proceed.,* **2451**(1) (2022) 20-44. https://doi.org/10.1063/5.0095423

[3318] T. V. Sateesh Kumar and S. Meenakshi, Triangular graphs are proper lucky and lucky, *AIP Conf. Proceed.,* **2451**(1) (2022) 104-122. https://www.researchgate.net/publication/364255497_Triangular_graphs_are_proper_lucky_and_lucky

[3319] A. A. Sathakathullaa, M. Akrama, M. and P. G. Rajeswari, Cordial, total cordial, edge cordial, total edge cordial labeling of some box type fractal graphs, *Internat. J. Algebra Stat.,* **1**(2) (2012) 99-106. https://www.researchgate.net/publication/258257334_On_cordial_Total_cordial_Edge_cordial_Total_Edge_cordial_labeling_of_some_box_type_fractal_graphs

[3320] A. A. Sathakathulla, A. A. and P. G. Rajeswari, Edge cordial, total edge cordial labeling of some ladder type fractal graphs, *Internat. J. Applied Sci. Eng. Res.*, *1*(6) (2012) 793-801. https://www.researchgate.net/publication/272796839_Edge_Cordial_Total_Edge_Cordial_Labeling_Of_Some_Ladder_Type_Fractal_Graphs

[3321] G. Sathiamoorthy, C. Natarajan, S. K. Ayyaswamy, and T. N. Janakiraman, Graceful labeling of splitting graph of a caterpillar, Internat. J. Mech. Engin. Tech., (IJMET) **8**(11) (2017) 663-667. https://iaeme.com/MasterAdmin/Journal_uploads/IJMET/VOLUME_8_ISSUE_11/IJMET_08_11_067.pdf

[3322] G. Sathiamoorthy, Graceful labeling of nova cloud connected in a path, *Internat. J. Mech. Engin. Tech.,* (IJMET) **8**(11) (2017) 1-9. https://iaeme.com/MasterAdmin/Journal_uploads/IJMET/VOLUME_8_ISSUE_11/IJMET_08_11_001.pdf

[3323] S. Sathish Narayanan and M. Vijayaragavan, Some 3-divisor cordial graphs derived from path, *Jordan J. Math. and Stat.,* (JJMS) **14**(2) (2021) 335-350. https://journals.yu.edu.jo/jjms/Issues/Vol14No22021PDF/9.pdf

[3324] K. Savitha and A. Sasikala, On families of some graphs admitting felicitous labeling, *Bol. Soc. Paran. Mat.*, (3s.) **44** (2026) 1–11. doi:10.5269/bspm.80717

[3325] K. Schaffer and S. M. Lee, Edge-graceful and edge-magic labelings of Cartesian products of graphs, *Congr. Numer.,* **141** (1999) 119-134.

[3326] S. A. Schluchter, J. Z. Schroeder, K. Cokus, R. Ellingson, H. Harris, E. Rarity, and T. Wilson, Prime labelings of generalized Petersen graphs, *Involve,* **10**(1) (2017) 109-124. https://msp.org/involve/2017/10-1/involve-v10-n1-p08-p.pdf

[3327] S. A. Schluchter, and T. Wilson, Prime labelings of bipartite generalized Petersen graphs and other prime cubic bipartite graphs, *Congr. Numer.,* **226** (2016) 227–241.

[3328] J. Z. Schroeder, Every cubic bipartite graph has a prime labeling except $K_{3,3}$, *Graphs Combin.,* **35**(1) (2019) 119-140. https://www.researchgate.net/publication/329195937_Every_Cubic_Bipartite_Graph_has_a_Prime_Labeling_Except_K_33K33

[3329] J. Z. Schroeder, Correction to: Every cubic bipartite graph has a prime labeling except $K_{3,3}$, *Graphs and Combinat.,* (2022) 38:148 https://doi.org/10.1007/s00373-022-02555-2

[3330] J. Z. Schroeder, *J. Combin.,* A 2-regular graph has a prime labeling if and only if it has at most one odd component. **12**(3) (2021) 379-388.

[3331] J. Z. Schroeder, Correction to: Every cubic bipartite graph has a prime labeling except $K_{3,3}$, *Graphs and Combinat.,* (2022) 38:148 https://link.springer.com/article/10.1007/s00373-022-02555-2

[3332] R. Sebastian and K. A. Germina, Square sum labeling of class of planar graphs, *Proyecciones,* **34**(1) (2015) 55-68. https://www.researchgate.net/publication/275413038_Square_Sum_Labeling_of_Class_of_Planar_Graphs

[3333] J. Sedláček, Problem 27, *in* Theory of Graphs and its Applications, *Proc. Symposium Smolenice,* June, (1963) 163-167.

[3334] J. Sedláček, On magic graphs, *Math. Slov.,* **26** (1976) 329-335. https://dml.cz/bitstream/handle/10338.dmlcz/133034/MathSlov_26-1976-4_8.pdf

[3335] M. Seenivasan and A. Lourdusamy, Absolutely harmonious labeling of graphs, *Internat. J. Math. Combin.*, **2** (2011) 40-51. https://dml.cz/bitstream/handle/10338.dmlcz/133034/MathSlov_26-1976-4_8.pdf

[3336] A. Sehgal, N. Takshak, P. Maan, and A. Malik, Graceful labeling of power graph of group $Z_{2^{k-1}} \times Z_4$, *Asian-European J. Math.* (2021)

[3337] C. Sekar, Studies in Graph Theory, Ph. D. Thesis, Madurai Kamaraj University, 2002.

[3338] C. Sekar and V. Ramachandren, Graceful labelling of arbitrary supersubdivision of disconnected graph, *Ultra Scientist,* **25**(2)A (2013) 315-318. /fs.unm.edu/IJMC/OneModuloNGracefullness.pdf?utm_source=chatgpt.com

[3339] A. Sekar and R. Varatharajaperumal, *E*-super arithmetic graceful labelling of some standard classes of cubic graphs related to cycles, *Bull. Int. Math. Virtual Inst.*,**12**(2) (2022) 353-367. http://fs.unm.edu/IJMC/ESuperArithmeticGraceful.pdf

[3340] L. S. Selistin and S. Asha, On topological cordial graphs, *J. Sci. Tech.* **5**(1) 25–28.

[3341] L. S. Selistin ans S. Asha, Topological cordial labelling of some graphs, *Malaya J. Mat.,* **9**(1) (2021) 861-863. https://doi.org/10.12723/mjs.sp1.11

[3342] S. P. Selva and G. Lathamaheswari, Total 3-sum cordial labeling, *IJMA*, **5**(5) (2014) 248-253. https://www.ijma.info/index.php/ijma/article/view/2899/0

[3343] P. Selvagopal and P. Jeyanthi, On C_k-supermagic graphs, *Inter. J. Math. Comput. Sci.,* **3** (2008) 25-30. https://www.researchgate.net/publication/257365099_PSelvagopal_and_PJeyanthi_On_Ck_-_super_magic_graphs_International_Journal_of_Mathematics_and_Computer_Science_Vol3_No12008_25-30

[3344] P. Selvagopal, P. Jeyanthi, N. P. T. Muthuraja, and A. Semaničová-Feňovčiková Super (a, d)-star-antimagic graphs, *Hacet. J. Math. Stat.,* **48** (2019)(3) 700-710. https://dergipark.org.tr/tr/download/article-file/735107

[3345] T. M. Selvarajan and R. Subramoniam, Prime graceful labeling, *Internat. J. Engin. Tech.*, **7**(4.36) (2018) 750-752. https://www.researchgate.net/publication/332672920_Prime_Graceful_Labeling

[3346] P. Selvaraju, New classes of graphs with α-valuation, harmonious and cordial labelings, Ph. D. Thesis, Anna University, 2001. Madurai Kamaraj University, 2002.

[3347] P. Selvaraju, P. Balaganesan, and J. Renuka, Odd harmonious labeling of some path related graphs, *Internat. J. Math. Sci. Engg. Appls.*, **7**(3) (2013) 163-170. doi:org/10.4067/S0716-09172016000100006

[3348] P. Selvaraju, P. Balaganesan, J. Renuka, and M. L. Suresh, Harmonious and vertex graceful labeling on path and star related graphs, *Internat. J. Pure Appl. Math.*, **93**(4) (2014) 501-509. doi:http://dx.doi.org/10.12732/ijpam.v93i4.2

[3349] P. Selvaraju, P. Balaganesan, J. Renuka, and M. L. Suresh, Vertex graceful labeling of $C_j \cup C_k \cup C_l$, *Appl. Math. Sci.*, **8**(82 (2014) 4047-4052. https://www.m-hikari.com/ams/ams-2014/ams-81-84-2014/balaganesanAMS81-84-2014.pdf

[3350] P. Selvaraju and C. Moha, Prime valuation of union of cycles and cycle related graphs, *Internat. J. Advan. Res. Comput. Sci. Software Engin.,* **7**(6) (2017) 839-842. https://www.researchgate.net/publication/319124529_Prime_Valuation_of_Union_of_Cycles_and_Cycle_Related_Graphs

[3351] P. Selvaraju and G. Sethurman, Decomposition of complete graphs and complete bipartitie graphs into copies of P_n^3 or $S_2(P_n^3)$ and harmonious labeling of $K_2 + P_n$, *J. Indones. Math. Soc.,* Special Edition (2011) 109-122.

[3352] S. Selvarasu and S. S. Murugan, Fuzzy anti-magic labeling on comb and twing graphs, *Modelling Engin. Problems*, **11**(2) (2024) 500-506. https://doi.org/10.18280/mmep.110222

[3353] M. Selvi, D. Ramya and P. Jeyanthi, Skolem difference mean graphs, *Proyecciones J. Math.*, **34**(3) (2015) 243-254. https://www.scielo.cl/pdf/proy/v34n3/art04.pdf

[3354] M. Selvi, D. Ramya and P. Jeyanthi, Triangular difference mean graphs, *Internat. J. Math. Combin.,* **2** (2020) 109-117. https://www.researchgate.net/publication/342803089_Triangular_Difference_Mean_Graphs

[3355] A. Semaničová, On magic and supermagic circulant graphs, *Discrete Math.,* **306**(18) (2006) 2263-2269. https://www.sciencedirect.com/science/article/pii/S0012365X06003165?via%3Dihub

[3356] A. Semaničová-Feňovčíková, M. Bača, M. Lascsáková, M. Miller, and J. Ryan, Wheels are cycle-antimagic, *Electronic Notes Disc. Math.*, **48** (2015) 11-18. The Eighth International Workshop on Graph Labelings (IWOGL 2014

[3357] A. Semaničová-Feňovčíková, M. Bača, and M. Lascsáková, Two constructions of H-antimagic graphs *AKCE Internat. J. Graphs Combin.,* **14**(1) (2017) 42-47. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2016.11.004?scroll=top&needAccess=true#d1e1402

[3358] M. F. Semenyuta, (a, d)-Distance antimagic labeling of some types of graphs, *Cybernetics Systems Anal.,* **52**(6) (2016) 950-955. https://www.researchgate.net/publication/310742802_a_d-Distance_Antimagic_Labeling_of_Some_Types_of_Graphs

[3359] M. F. Semenyuta, On the (a, d)-distance antimagic and 1-vertex bimagic vertex labelling of some types of graphs, *Cybernet. Systems Anal.,* **54(2)** (2018) (2) 295-301.

[3360] M. F. Semenyuta, S. M. Nedilko, and V. M. Nedilko, Bimagic vertex labelings, *Cybernet. Systems Anal.,* **54**(5) (2018) 771-778. https://www.researchgate.net/publication/327870019_Bimagic_Vertex_Labelings

[3361] A. Sen, H. Deng, and S. Guha, On a graph partition problem with application to VLSI Layout, *Inform. Process. Lett.,* **43** (1992) 87-94. https://www.academia.edu/7994513/On_a_Graph_Partition_Problem_with_Application_to_VLSI_Layout

[3362] R. Senthil Amutha and N. Murugesan, Characterization of connected vertex magic total labeling graphs in topological ideals, *J. Inform. Eng. Appl.*, *1*(1) (2011) 22-25. https://www.researchgate.net/publication/277986795_Characterization_of_connected_Vertex_Magic_Total_labeling_graphs_in_Topological_idealsm

[3363] M. Seoud and M. Aboshady, Further results on parity combination cordial labeling, *J. Egyptian Math. Soc.,* **28**(1) (2020) Paper No. 25, 10 pp. https://joems.springeropen.com/articles/10.1186/s42787-020-00082-8

[3364] M. A. Seoud, G. M. Abd El Hamid, and M. S. Abo Shady, Indexable and strongly indexable graphs, *Proceed. Pakistan Acad. Sci.,* **49**(2) (2012) 139-144. https://paspk.org/wp-content/uploads/proceedings/49,%20No.%202/28a0a32aIndexable%20and%20Strongly%20Indexable%20Graphs.pdf

[3365] M. A. Seoud and M. E. Abdel-Aal, On odd graceful graphs, *Ars Comb.,* **108** (2013) 161-185. https://combinatorialpress.com/article/ars/Volume%20108/volume-108-paper-14.pdf

[3366] M. A. Seoud and A. E. I. Abdel Maqsoud, On 3-equitable and magic labelings, *Proc. Math. Phys. Soc. Egypt,* **7** (2000) 67-76.

[3367] M. A. Seoud and A. E. I. Abdel Maqsoud, On cordial and balanced labelings of graphs, *J. Egyptian Math. Soc.,* **7** (1999) 127-135.

[3368] M. A. Seoud, A. E. I. Abdel Maqsoud and Y. I. Aldiban, New classes of graphs with and without 1-vertex magic vertex labeling, *Proc. Pakistan Acad. Sci.,* **46** (2009) 159-174.

[3369] M. A.Seoud, A. E. I. Abdel Maqsoud and J. Sheehan, Harmonious graphs, *Util. Math.*, **47** (1995) 225-233.

[3370] M. A. Seoud, A. E. I. Abdel Maqsoud, and J. Sheehan, Gracefulness of the union of cycles and paths, *Ars Combin.,* **54** (2000) 283-292. https://combinatorialpress.com/article/ars/Volume%20054/volume-54-paper-21.pdf

[3371] M. A. Seoud and M. N. Al-Harere, Some notes on non combination graphs, *Applied Math. Sciences*, **6**(131) (2012) 6515-6520. https://www.m-hikari.com/ams/ams-2012/ams-129-132-2012/seoudAMS129-132-2012-1.pdf

[3372] M. A. Seoud and M. N. Al-Harere, Further results on square sum graphs, *Nat. Acad. Sci. Lett.,* **37**(5) (2014) 473-475.

[3373] M. A. Seoud and M. N. Al-Harere, Some notes on combination graphs, *Ars Combin.*, **129** (2016) 95-106. https://combinatorialpress.com/article/ars/Volume%20129/volume-129-paper-8.pdf

[3374] M. A. Seoud and M. A. Anwar, On combination and permutation graphs, *Util. Math., 98* 243-255. https://www.researchgate.net/publication/295579013_On_Combination_and_Permutation_Graphs

[3375] M. A. Seoud and M. A. Anwar, Some families of combination and permutation graphs, *Ars Combin.,* **127** (2016) 435-443. https://www.researchgate.net/publication/307928064_Some_Families_of_Combination_and_Permutation_Graphs

[3376] M. A. Seoud, A. T. Diab, and E. A. Elsahawi, On strongly-C harmonious, relatively prime, odd graceful and cordial graphs, *Proc. Math. Phys. Soc. Egypt,* no. 73, (1998) 33-55.

[3377] M¿ A. Seoud, A. El Sonbaty, and S. S. Abd El Rehim, Some methods of labeling graphs, *J. Egyptian Math. Soc.,* **18**(2) (2010) 199-221.

[3378] M. A. Seoud and E. A. Elsahawi, On almost graceful, felicitous and elegant graphs, *J. Egyptian Math. Soc.,* **7** (1999) 137-149.

[3379] M. A. Seoud and E. A. El Sahawi, On variations of graceful labelings, *Ars Combin.,* **87** (2008) 127-138. https://combinatorialpress.com/article/ars/Volume%20087/volume-87-paper-10.pdf

[3380] M. A. Seoud and E. A. El Sahawi, On strongly c-elegant graphs, personal comunication.

[3381] M. A. Seoud, A. El Sonbaty, and A. E. A. Mahran, On divisor graphs, *J. Egyptian Math. Soc.,* **18**(2) (2010) 187-198.

[3382] M. A. Seoud A. El Sonbaty, and A. E. A. Mahran, Primality of some graphs, *Ars Combin.,* **112** (2013) 459-469. https://combinatorialpress.com/article/ars/Volume%20112/volume-112-paper-38.pdf

[3383] M. A. Seoud and H. M. Hafez, Odd harmonious and strongly odd harmonious graphs, *KYUNGPOOK Math. J.*, **58**(4) (2018) 747-759. https://kmj.knu.ac.kr/journal/view.html?volume=58&number=4&spage=747

[3384] M. A. Seoud and E. F. Helmi, On product cordial graphs, *Ars Combin.,* **101** (2011) 519-529. https://combinatorialpress.com/article/ars/Volume%20101/volume-101-paper-40.pdf

[3385] M. A. Seoud and E. F. Helmi, Some α-graphs and odd graceful graphs, *Ars Comb.,* **101** (2011) 385-404. https://combinatorialpress.com/article/ars/Volume%20101/volume-101-paper-29.pdf

[3386] M. A. Seoud and E. F. Helmi, On difference graphs, *J. Combin. Math. and Combin. Comput.* **76** (2011) 189-199. https://combinatorialpress.com/article/jcmcc/Volume%20076/vol-076-paper%2012.pdf

[3387] M. A. Seoud and H. Jaber, On 1-sequentially additive graphs, personal comunication.

[3388] M. A. Seoud and H. Jaber, Prime cordial and 3-equitable prime cordial graphs, *Util. Math.,* **111** (2019) 95–125.

[3389] M. A. Seoud and A. E. A. Mahran, On permutation graphs, *J. Egyptian Math. Soc.* **20** (2012) 57-63. https://joems.journals.ekb.eg/article_386388_0401513efbe8972a9f1d7a9db91fb6e7.pdf

[3390] M. A. Seoud and A. E. A. Mahran, On strongly multiplicative graphs, *Ars Combin.,*, **118** (2015) 155-165. https://combinatorialpress.com/article/ars/Volume%20118/volume-118-paper-12.pdf

[3391] M. A. Seoud and A. E. A. Mahran, Necessary conditions for strongly *-graphs, *AKCE Internal. J. Graphs Comb.*, **9**(2) (2012) 115-122. https://www.tandfonline.com/doi/pdf/10.1080/09728600.2012.12088953

[3392] M. A. Seoud and A. E. A. Mahran, Some notes on strongly *-graphs, *Proceed. Pakistan Acad. Sci.,: Pakistan Acad. Sci.,A. Physical Comput. Sci.,* **54**(2) (2017) 179-195. https://www.paspk.org/wp-content/uploads/2017/07/A-06-PS-410-On-Strongly-Graphs.pdf

[3393] M. A. Seoud, E. M. Roshdy, and M. S. AboShady, On strongly *-graphs, *Proc. Pak. Acad. Sci.,* A **54**(2) (2017) 179-195. https://www.researchgate.net/publication/337534591_On_Strongly_-Graph

[3394] M. A. Seoud and M. A. Salim, Two upper bounds of prime cordial graphs, *JCMCC,* **75** (2010) 95-103. https://combinatorialpress.com/article/jcmcc/Volume%20075/vol-075-paper%207.pdf

[3395] M. A. Seoud and M. A. Salim, Further results on triangular sum graphs, *Internat. Math. Forum,* **7**(48) (2012) 2393-2405. http://www.m-hikari.com/imf/imf-2012/45-48-2012/seoudIMF45-48-2012.pdf

[3396] M. A. Seoud and M. A. Salim, On odd mean graphs, *Util. Math.,*, **84** (2011) 119-130.

[3397] M. A. Seoud and M. A. Salim, On permutation labeling, *J. Egyptian Math. Soc.,* **19**(3) (2011) 134-136. https://www.researchgate.net/publication/257498209_On_permutation_labeling

[3398] M. A. Seoud and M. A Salim, On mean graphs, *Ars Combin.,* **115** (2014) 13-34. https://combinatorialpress.com/article/ars/Volume%20115/volume-115-paper-2.pdf

[3399] M. A. Seoud and M. A. Salim, Upper bounds of four types of graph labelings, *127* (2016) 271-278. https://combinatorialpress.com/article/ars/Volume%20127/volume-127-paper-24.pdf

[3400] M. A. Seoud and M. A. Salim, Further results on edge-odd graceful graphs, *Turkish J. Math.,* **40**(3) (2016) 647-656. https://journals.tubitak.gov.tr/cgi/viewcontent.cgi?article=2056&context=math

[3401] M. A. Seoud and M. A. Salim, Some results and remarks on harmonic mean and mean cordial labelings, *Ars Combin.,* **133** (2017) 329-340. https://combinatorialpress.com/article/ars/Volume%20133/volume-133-paper-23.pdf

[3402] M. A. Seoud, A. El Sonbaty, and A. E. A. Mahran, On prime graphs, *Ars Combin.,* **104** (2012) 241-260. https://combinatorialpress.com/article/ars/Volume%20104/volume-104-paper-21.pdf

[3403] M. A. Seoud and R. J. Wilson, Some disgraceful graphs, *Int. J. Math. Ed. Sci. Tech.,* **24** (1993) 435-441.

[3404] M. A. Seoud and M. Z. Youssef, On labelling complete tripartite graphs, *Int. J. Math. Ed. Sci. Tech.*, **28** (1997) 367-371.

[3405] M. A. Seoud and M. Z. Youssef, On prime labelings of graphs, *Congr. Numer.,* **141** (1999) 203-215.

[3406] M. A. Seoud and M. Z. Youssef, Families of harmonious and non-harmonious graphs, *J. Egyptian Math. Soc.,* **7** (1999) 117-125.

[3407] M. A. Seoud and M. Youssef, New families of graceful disconnected graphs, *Ars Combin.*, **57** (2000) 233-245. https://combinatorialpress.com/article/ars/Volume%20057/volume-57-paper-22.pdf

[3408] M. A. Seoud and M. Z. Youssef, On harmonious graphs of order 6, *Ars Combin.,* **65** (2002) 155-176. https://combinatorialpress.com/article/ars/Volume%20065/volume-65-paper-14.pdf

[3409] M. A. Seoud and M. Z. Youssef, The effect of some operations on labelling of graphs, *Proc. Math. Phys. Soc. Egypt*, **73** (2000) 35-49. https://www.researchgate.net/publication/291125488_The_effect_of_some_operations_on_labelling_of_graphs

[3410] M. A. Seoud and M. Z. Youssef, Harmonious labellings of helms and related graphs, https://www.researchgate.net/publication/312218305_Harmonious_labelling_of_helms_and_related_graphs.

[3411] M. A. Seoud and M. Z. Youssef, On gracefulness of disconnected graphs, https://www.researchgate.net/publication/312218547_On_gracefulness_of_disconnected_graphs.

[3412] M. A. Seoud and A. Zid, Strong multiplicativity of unions and corona of paths and complete graphs, *Proc. Math. Phys. Soc. Egypt,* **74** (1999) 59-71.

[3413] B. J. Septory, M. I. Utoyo, Dafik, B. Sulistiyono, and I. H. Agustin, On rainbow antimagic coloring of special graphs, *J. Phys.: Conf. Ser.,* 1836 012016 (2021). https://iopscience.iop.org/article/10.1088/1742-6596/1836/1/012016/pdf

[3414] E. Sergel, P. Richter, A. Tran, P. Curran, J. Jacob, and D. A. Narayan, Rank numbers for some trees and unicycle graphs, *Aequat. Math.,* **82**(1-2) (2011) 65-79.

[3415] A. Sethukkarasi, S. Vidyanandini, and A. El-Mesady, Effective delta-labeling exploration algorithms for graph representation and DNA sequence alignment, *Int. J. Inf. Tecnol.,* (2025) https://www.researchgate.net/publication/387861211_Effective_delta-labeling_exploration_algorithms_for_graph_representation_and_DNA_sequence_alignment

[3416] E. Setiawan, K. A. Sugeng, Silaban, and D. Riama, On b-edge consecutive edge magic total labeling on trees, *Electron. J. Graph Theory Appl.*, (EJGTA) **10**(2) (2022) 553-563. https://www.ejgta.org/index.php/ejgta/article/view/1585/pdf_248

[3417] G. Sethuraman and R. Dhavamani, Graceful numbering of an edge-gluing of shell graphs, *Discrete Math.,* **218** (2000) 283-287. https://www.sciencedirect.com/science/article/pii/S0012365X9900360X

[3418] G. Sethuraman and R. Dhavamani, Graceful numbering of union of shell graphs, unpublished.

[3419] G. Sethuraman and A. Elumalai, On graceful graphs: Pendant edge extensions of a family of complete bipartite and complete tripartite graphs, *Indian J. Pure Appl. Math.,* **32** (2001) 1283-1296.

[3420] G. Sethuraman and A. Elumalai, Gracefulness of a cycle with parallel P_k-chords, *Australas. J. Combin.,* (2005) **32**) 205-211. https://ajc.maths.uq.edu.au/pdf/32/ajc_v32_p205.pdf

[3421] G. Sethuraman and A. Elumalai, Packing of any set of graphs into a graceful/harmonious/elegant graph, *Ars Combin.,* **76** (2005) 297-301. https://combinatorialpress.com/article/ars/Volume%20076/volume-76-paper-26.pdf

[3422] G. Sethuraman and A. Elumalai, Graceful, harmonious and elegant labellings on star extension graphs, unpublished.

[3423] G. Sethuraman and A. Elumalai, Every graph is a vertex induced subgraph of a graceful graph and elegant graph, unpublished.

[3424] G. Sethuraman and J. Jesintha, A new class of graceful lobsters, *J. Combin. Math. Combin. Computing,* **67** (2008) 99-109. https://combinatorialpress.com/article/jcmcc/Volume%20067/vol-067-paper%2010.pdf

[3425] G. Sethuraman and J. Jesintha, Gracefulness of a family of rooted trees, *Far East J. Appl. Math.,* **30** (2008) 143-159.

[3426] G. Sethuraman and J. Jesintha, A new family of graceful rooted trees, *Proc. National Conf. Appl. Math.,* (2008) 74-80.

[3427] G. Sethuraman and J. Jesintha, A new class of graceful rooted trees, *J. Disc. Math. Sci. Crypt.,* **11** (2008) 421-435. https://connectjournals.com/file_full_text/84904H_jdmsc235_421-435.pdf

[3428] G. Sethuraman and J. Jesintha, Generating new graceful trees, *Proc. Inter. Conf. Math. Comput. Sci.,* July (2008) 67-73.

[3429] G. Sethuraman and J. Jesintha, Generation of graceful trees, *Proc. Inter. Conf. Math. Comput Sci.,* **1** (2009) 1-3.

[3430] G. Sethuraman and J. Jesintha, All extended banana trees are graceful, *Proc. Internat. Conf. Math. Comput. Sci.,* **1** (2009) 4-8.

[3431] G. Sethuraman and J. Jesintha, All banana trees are graceful, *Advances Appl. Disc. Math.,* **4** (2009) 53-64. https://www.pphmj.com/article.php?act=art_abstract_show&art_id=4224&flag=prev

[3432] G. Sethuraman and S. P. M. Kishore, On graceful graphs: Union of n copies of edge deleted subgraphs of K_4, *Indian J. Pure Appl. Math.,* **30** (1999) 801-808.

[3433] G. Sethuraman and V. Murugan, Generating graceful trees from caterpillars by recursive attachment, *Electronic Notes Discrete Math.,* **53** (2016) 133-147. https://www.academia.edu/42249036/Generating_Graceful_Trees_from_Caterpillars_by_Recursive_Attachment

[3434] G. Sethuraman and · V. Murugan, Decomposition of complete graphs into arbitrary trees, *Graphs Combin.,* **37** (2021) 1191-1203. https://www.researchgate.net/publication/266709410_Decomposition_of_complete_graphs_into_tree

[3435] G. Sethuraman and V. Murugan, Generating graceful unicyclic graph from a given graceful tree, *Ars. Combin.,* February (2019)

[3436] G. Sethuraman and V. Murugan, Generating graceful unicyclic graphs from a given forest, *AKCE Internat. J. Graphs Combin.,* **17**(10 (2020) 592-608. https://www.researchgate.net/publication/332025576_Generating_graceful_unicyclic_graphs_from_a_given_forest

[3437] G. Sethuraman and V. Murugan, Generating graceful trees from caterpillars by recursive attachment, *Elect. Notes Disc. Math.*, **5**3 (2016) 133–147.

[3438] G. Sethuraman and P. Ragukumar, Every tree is a subtree of graceful tree, graceful graph and alpha-labeled graph, *Ars Combin.,* **132** (2017) 27-47. https://combinatorialpress.com/article/ars/Volume%20132/volume-132-paper-3.pdf

[3439] G. Sethuraman and P. Ragukumar, Towards optimal embedding of an arbitrary tree in a graceful tree, The Eighth International Workshop on Graph Labelings (IWOGL 2014), 73-80, *Electron. Notes Discrete Math.,* **48**, Elsevier Sci. B. V., Amsterdam, 2015.

[3440] G. Sethuraman, P. Ragukumar, and P. J. Slater, Embedding an arbitrary tree in a graceful tree, *Bull. Malays. Math. Sci. Soc.,* **39**(1) (2016) suppl., S341-S360.

[3441] G. Sethuraman, P. Ragukumar, and P. J. Slater, Any tree with m edges can be embedded in a graceful tree with less than $4m$ edges and in a graceful planar graph, *Discrete Math.,* **340**(2) (2017) 96-106. https://www.sciencedirect.com/science/article/pii/S0012365X16302308

[3442] G. Sethuraman and K. Sankar, On graceful and cordial labeling of shell graphs, *Ars Combin.,* **99** (2011) 225-242. https://combinatorialpress.com/article/ars/Volume%2099/volume-99-paper-21.pdf

[3443] G. Sethuraman and K. Sankar, Subdivision of wheel is graceful and cordial, *Internat. J. Math. Anal.*, textbf9(17) (2015) 817-822. http://www.m-hikari.com/ijma/ijma-2015/ijma-17-20-2015/01IsankarIJMA17-20-2015-08.pdf

[3444] G. Sethuraman and P. Selvaraju, Gracefulness of arbitrary supersubdivisions of graphs, *Indian J. Pure Appl. Math.,* **32** (2001) 1059-1064.

[3445] G. Sethuraman and P. Selvaraju, On graceful graphs: one vertex unions of non-isomorphic complete bipartite graphs, *Indian J. Pure Appl. Math.,* **32** (2001) 975-980. https://www.researchgate.net/publication/265424682_On_graceful_graphs_One_vertex_unions_of_non-isomorphic_complete_bipartite_graphs

[3446] G. Sethuraman and P. Selvaraju, One edge union of shell graphs and one vertex union of complete bipartite graphs are cordial, *Discrete Math.,* **259** (2002) 343-350. https://www.sciencedirect.com/science/article/pii/S0012365X02003059

[3447] G. Sethuraman and P. Selvaraju, Decompositions of complete graphs and complete bipartite graphs into isomorphic supersubdivision graphs, *Discrete Math.,* **260** (2003) 137-149. https://www.sciencedirect.com/science/article/pii/S0012365X02004557?via%3Dihub

[3448] G. Sethuraman and P. Selvaraju, Super-subdivisions of connected graphs are graceful, unpublished.

[3449] G. Sethuraman and P. Selvaraju, On graceful graphs I: Union of non-isomorphic complete bipartite graphs with one vertex in common, *J. Combin. Inform. System Sci.,* **26** (2001) 23-32.

[3450] G. Sethuraman and P. Selvaraju, New classes of graphs on graph labeling, unpublished.

[3451] G. Sethuraman and P. Selvaraju, On harmonious and felicitous graphs: Union of n-copies of edge deleted subgraphs of K_4, unpublished.

[3452] G. Sethuraman, P. Selvaraju, and A. Elumalai, On harmonious, felicitous, elegant and cordial graphs: Union of n copies of edge deleted subgraphs of K_4. unpublished.

[3453] G. Sethuraman and S. Venkatesh, All trees are integral sum graphs, unpublished.

[3454] G. Sethuraman and K. M. Shermily, Antimagic labeling of new classes of trees, *AKCE Int. J. Graphs Comb.,* **18**(2) (2021) 110-116. https://www.tandfonline.com/doi/full/10.1080/09728600.2021.1964334

[3455] G. Sethuraman and S. Venkatesh, Decomposition of complete graphs and complete bipartite graphs into α-labelled trees, *Ars Combin.,* **93** (2009) 371-385. https://combinatorialpress.com/article/ars/Volume%20093/volume-93-paper-36.pdf

[3456] I. Setiawan and D. Indriati, Edge irregular reflexive labeling on sun graph and corona of cycle and null graph with two vertices, *Indones. J. Comb.*, *5*(1) (2021). https://www.ijc.or.id/index.php/ijc/article/view/143/110

[3457] N. M. Seyam and M. F. Nadeem, On edge irregularity strength of some classes of Toeplitz graphs, *Open J. Discret. Appl. Math.,* textbf2 (2024) 23-34. https://pisrt.org/psrpress/j/odam/2024/2/On-edge-irregularity-strength-of-some-classes-of-Toeplitz-graphs.pdf

[3458] H. Shaabani and I. N. Kamalabadi, An efficient population-based simulated annealing algorithm for the multi-product multi-retailer perishable inventory routing problem, *Comput. Ind. Eng.,* **99** (2016)

[3459] M. K. Shafiq, G. Ali, and R. Simanjuntak, Distance magic labeling of a union of graphs, *AKCE J. Graph Combin.,* **6** (2009) 191-200. https://www.researchgate.net/publication/265582771_Distance_magic_labelings_of_a_union_of_graph

[3460] P. V. Shah, D. B. Parmar, Integer cordial labeling of triangular snake graph, *Inter. J. Sci. Res. Reviews*, **08**(1) (2019) 3118-3126. https://www.researchgate.net/publication/332319553_Integer_Cordial_Labeling_of_Triangular_Snake_Graph

[3461] P. V. Shah, and D. B. Parmar, Integer cordial labeling of some different snake graph, *J. Xidian Univer.,* **14**(4) (2020) 1361-1375. doi:10.37896/jxu14.4/159

[3462] D. Shah and D. Parmar, Integer cordial labeling of alternate snake graph and irregular snake graph, *Applications Appl. Math.* (AAM), *17*(3) Art. 5 (2022) 59-71. https://digitalcommons.pvamu.edu/cgi/viewcontent.cgi?article=2038&context=aam

[3463] A. T. Shahida and M. S. Sunitha, Characterization for topologically set-graceful stars and paths, *Adv. Appl. Discrete Math.*, **15**(1) (2015) 1-5 https://www.researchgate.net/publication/276912798_CHARACTERIZATION_FOR_TOPOLOGICALLY_SET-GRACEFUL_STARS_AND_PATHS

[3464] R. Shankar and M. C. Nalliah, Local antimagic chromatic number for the corona product of wheel and null graphs, *Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Komp'yuternye Nauki,* **32**(3) (2022) 463–485. https://www.mathnet.ru/links/d3cb9333330b463cfddc6d828032bb9f/vuu821.pdf

[3465] S. Shan and Y. Zhong, Graceful labeling of two families of spiders, *Graphs and Combin.*, **42**(5) (2026). arXiv:2605.14295v2

[3466] R. Shankar and M. C. Nalliah, Local vertex antimagic chromatic number of some wheel related graphs, *Proyecciones (Antofagasta)*, **41**(1) (2022) 319–334. https://www.scielo.cl/pdf/proy/v41n1/0716-0917-proy-41-01-319.pdf

[3467] V. Sandhiya and M. C. Nalliah, Local total antimagic chromatic number for the disjoint union of star graphs, *European J. Prue Appl. Math.*, **17**(4) (2024) 2828-2842. https://www.ejpam.com/ejpam/article/view/5383?utm_source=chatgpt.com

[3468] V. Shendra Shainy and V. Balaji, On limits of Skolem mean labeling for star graphs $K_{1,a_1} \cup K_{1,b_1} \cup K_{1,c,1}$. *JASC: J. Appl. Sci. Comput.*, **V**(XII) (2018) 1-6.

[3469] A. R. Shahul Hameed, M. Palanivelrajan, K. Gunasekaran, and R. Raziya Begam, On graceful labeling of some bicyclic graphs, *Internat. J. Fuzzy Mathematical Archives*, **3** (2013) 1-8. http://www.researchmathsci.org/IJFMAart/ijfma-v3-1.pdf

[3470] A. R. Shahul Hameed, M. Palanivelrajan, K. Gunasekaran and R. Raziya Begam, Graceful labeling of extended komodo dragon graphs, *Bull. Math. Stat. Res.,* **2**(4) (2014) 454-459. http://www.bomsr.com/Vol.2.Issue.4.2014/R.%20RAZIYA%20BEGAM%20454-459.pdf

[3471] A. R. Shahul Hameed, M. Palanivelrajan, K. Gunasekaran and R. Raziya Begam, Graceful labeling of extended squid graphs, *Bull. Math. Stat. Res.,* **2**(4) (2014) 465-473. http://www.bomsr.com/Vol.2.Issue.4.2014/R.%20RAZIYA%20BEGAM%20465-473.pdf

[3472] A. R. Shahul Hameed, M. Palanivelrajan, K. Gunasekaran and R. Raziya Begam, Graceful labeling of komodo dragon graphs, *J. Sci. Res. Physical & Math. Sci.*, **2**(2) (2015) 17-27.

[3473] A. R. Shahul Hameed, M. Palanivelrajan, K. Gunasekaran and R. Raziya Begam, Graceful labeling of squid graphs, *J. Sci. Res. Physical & Math. Sci.*, **2**(2) (2015) 28-38.

[3474] H. Shaker, A. Rana, M. H. Zobair, and M. Hussain, Super edge-magic total labeling of subdivision of stars, *Ars Combin.,* **116** (2014) 177-183. https://combinatorialpress.com/article/ars/Volume%20116/volume-116-paper-14.pdf

[3475] A. Shalini Rajendra Babu and N. Ramya, Lucky-edge labeling of H graph, N copies of H-graph, theta graph, duplication of theta graphs and path union of theta graphs, *Annals Romanian Soc. Cell Biology* **25**(2)(2021) 4480–4497. http://annalsofrscb.ro/index.php/journal/article/view/1471/1225

[3476] A. Shalini Rajendra Babu, N. Ramya, and K. Rangarajan, On lucky-edge labeling of splitting graphs and snake graphs, *Internat. J. Innov. Techn. Exploring Engin.*, **8**(5) (2019) 638–641. https://www.ijitee.org/wp-content/uploads/papers/v8i5/E3299038519.pdf

[3477] P. Shalini and D. P. Dhayabaran, An absolute differences of cubic and square difference labelings, *Internat. J. Adv. Sci. Tech. Res.*, **3**(5) (2015) 1-8. https://www.researchgate.net/publication/362174849_An_Absolute_Differences_of_Cubic_and_Square_Difference_Labelings

[3478] P. Shalini, R. Gowri, and D. P. Dhayabaran, An absolute differences of cubic and square difference labeling for some families of graphs, *Internat. J. Analytical Experimental Modal Analysis*, **XI**(X) (2019) 538-544. http://www.ijaema.com/gallery/59-october-2570.pdf

[3479] P. Shalini and S. A. Meena, Lehmer-4 mean labeling of graphs, *Internat. J. Res. Appl. Sci. Engin. Tech.,* (IJRASET) **10**(XII) (2022) doi:10.22214/ijraset.2022.48176

[3480] O. V. Shanmuga and S. D. Amuthavalli, Super fibonacci graceful anti–magic labeling for flower graphs and python coding, *Tuijin Jishu J. Propulsion Tech.*, **44**(3) (2023) 3407-3412. https://www.researchgate.net/publication/376197575_Super_fibonacci_graceful_anti_-_magic_labeling_for_flower_graphs_and_python_coding

[3481] S. Shan and X. Yu, Antimagic orientation of biregular bipartite graphs, *Electron. J. Combin.,* **24** (2017) Paper 4.31. https://www.combinatorics.org/ojs/index.php/eljc/article/view/v24i4p31/pdf

[3482] S. Shan, Antimagic orientation of graphs with minimum degree at least 33, *J. Graph Th.,* **98**(4) (2021) 676-690. arXiv:2005.01829

[3483] J.-L. Shang, Spiders are antimagic, *Ars Combin.,* **118** (2015) 367-372. https://combinatorialpress.com/article/ars/Volume%20118/volume-118-paper-29.pdf

[3484] J.-L. Shang, P_2, P_3, P_4-free linear forests are antimagic, *Util. Math.,* **101** (2016) 13-22. https://www.researchgate.net/publication/372888907_P2P3P4-free_linear_forests_are_antimagic

[3485] J.-L. Shang, Antimagic labeling of linear forests, *Util. Math.,* **106** (2018) 23-37 https://www.researchgate.net/publication/378880250_Antimagic_Labeling_of_Forests

[3486] J.-L Shang, C. Lin, and S-C Liaw, On the antimagic labeling of star forests, *Util. Math.,* **97** (2015) 373-385. https://www.researchgate.net/publication/286778546_On_the_antimagic_labeling_of_star_forests

[3487] Z. Shao, F. Deng, Z. Li, and A. Vese, Graceful labelings of the generalized Petersen graphs, *Commun. Combin. Opt.*, **2**(2) (2017) 149-159. https://comb-opt.azaruniv.ac.ir/article_13646_07d33d001066dc9b0e695120e6125c8a.pdf

[3488] A. Sharary, Integal sum graphs from complete graphs, cycles and wheels, *Arab Gulf Sci. Res.,* **14-1** (1996) 1-14. https://agjsr.agu.edu.bh/uploads/images/papers/pdfs/d1be586421b5108967097d58047f537d_611e14dbe4a38.pdf

[3489] V. Sharma and A. Parthiban, On divisor cordial labeling of certain classes of planar graphs, *J. Alg. Stat.*, **13**(2) (2022) 1421-1425.

[3490] V. Sharma and A. Parthiban, On double divisor cordial labeling of star related graphs, *Adv. Appl. Math. Sci.*, **21**(8) (2022) 4389-4398. https://iopscience.iop.org/article/10.1088/1742-6596/2267/1/012026/pdf

[3491] S. Philo, P. Jeyanthi, and B. Davazz, Odd harmonious labeling of complete bipartite graphs, *Southeast Asian Bull. Math.,* **46**(1) (2022) 23-34. https://www.researchgate.net/publication/359462364_Southeast_Asian_Bulletin_of_Mathematics_Odd_Harmonious_Labeling_of_Complete_Bipartite_Graphs

[3492] V. Sharon Philomena and K. Thirusangu, Some new classes of 3-total product cordial graphs, *Internat. J. Comput. Algor.,* **3** (2014) 128-130

[3493] V. Sharon Philomena and K. Thirusangu, Square and cube difference labeling of cycle cactus, special tree and a new key graph, *Annals Pure Appl. Math.,* **8**(2) (2014) 115-122. http://www.researchmathsci.org/apamart/apam-v8n2-14.pdf

[3494] N. K. Shawkat and M. A. Ahmed, On antimagic labeling for some families of graphs, *Ibn AL-Haitham J. Pure Appl. Sci.,* **36**(1) (2023) 284-291. doi:10.30526/36.1.3014

[3495] S. C. Shee, On harmonious and related graphs, *Ars Combin.,* **23** (1987) A, 237-247.

[3496] S. C. Shee, Some results on λ-valuation of graphs involving complete bipartite graphs, *Discrete Math.,* **87**(1) (1991) 73-80. https://www.sciencedirect.com/science/article/pii/0012365X9190072A?via%3Dihub

[3497] S. C. Shee and Y. S. Ho, The cordiality of one-point union of n-copies of a graph, *Discrete Math.,* **117** (1993) 225-243.

[3498] S. C. Shee and Y. S. Ho, The cordiality of the path-union of n copies of a graph, *Discrete Math.,* **151** (1996) 221-229. https://www.sciencedirect.com/science/article/pii/0012365X94000995?via%3Dihub

[3499] S. Shee and S. Lee, On harmonious and felicitious labeling of graphs, *Congr. Numer.,* **68** (1989) 155-170.

[3500] Y. Shen, J. Dong, G. Zheng, and L. Guo, The radio number of standard caterpillars, *Ars Combin.*, **125** (2016) 257-269. https://combinatorialpress.com/ars-articles/volume-125-ars-articles/the-radio-number-of-standard-caterpillars/?utm_source=chatgpt.com

[3501] V. Shendra Shainy and V. Balaji, Even felicitous labeling, *Malaya J. Mat.*, **S**(1) (2020) 45-47. https://doi.org/10.26637/MJM0S20/0009

[3502] V. Shendra shainy, P. Hariprabakaran, V. Swathy, and V. Balaji, MGL coding techniques with Skolem mean like labeling and two star graphs, *J. Physics Conference Series,* **1964**(2):022009 (2021) 1-6. https://iopscience.iop.org/article/10.1088/1742-6596/1964/2/022009/pdf

[3503] V. Shendra shainy, L. Vinothkumar and V. Balaji, Mean cordial labeling for three star graph, *J. Phys.: Conf. Ser.,* **1362** (2019) 012036. https://iopscience.iop.org/article/10.1088/1742-6596/1362/1/012036

[3504] D. A. Sheppard, The factorial representation of major balanced labelled graphs, *Discrete Math.,* **15** (1976) 379-388. https://www.sciencedirect.com/science/article/pii/0012365X76900510?via%3Dihub

[3505] M. M. Sheriff, A. F. Abbas and P. L. R. Raj, Face integer cordial labeling of graphs, *Internat. J. Math. Trends Tech.,* (IJMTT) **41**(2) (2017) 177-185. https://www.researchgate.net/publication/315917668_Face_Integer_Cordial_Labeling_of_Graphs

[3506] M. M. Sheriff and A. F. Abbas, A study on face integer edge cordial graphs, *Internat. J. Fuzzy Math. Archive* **15**(2) (2018) 209-216. http://www.researchmathsci.org/IJFMAart/IJFMA-v15n2-12.pdf

[3507] Z. Sherman, Square difference labeling of some union and disjoint union graphs, *Algebra Discrete Math.,* **25**(2) (2018) 294-302. https://dspace.luguniv.edu.ua/xmlui/bitstream/handle/123456789/4503/269-3445-1-PB.pdf?isAllowed=y&sequence=1&utm_source=chatgpt.com

[3508] V. S. Shigehalli and Usmanbasha, Radio labeling of Mycielskian graph of certian graphs, *Internat. J. Appl. Math.*, **38**(7a) (2025) 1774-1790. https://ijamjournal.org/ijam/publication/index.php/ijam/article/view/1636/1488

[3509] J. Shiama, Square sum labeling for middle and total graphs, *Internat. J. Comp. Appl.*, **37**(4) (2005) 205-211.

[3510] J. Shiama, Square sum labeling for some middle and total graphs, *Internat. J. Comput. Appl.*, **37**(4) (2012) 6-8. https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.736.5400&rep=rep1&type=pdf

[3511] J. Shiama, Cube difference labeling of some graphs, *Internat. J. Engin. Sci. Innovative Tech.* (IJESIT), **2**(6) (2013) 200-205. http://www.ijesit.com/Volume%202/Issue%206/IJESIT201306_25.pdf

[3512] J. Shiama, Square difference labeling for some graphs, *Internat. J. Comput. Appl.*, **44**(4) (2012) 30-33. https://research.ijcaonline.org/volume44/number4/pxc3878399.pdf

[3513] H. Shimazu, Graceful labelling of the union of cycles and paths, *Internat. J. Physics Appl.,* **3**(2) (2011) 205-208.

[3514] Shisheng Li, personal communication. Oct. 2026 https://github.com/daizisheng/notes/blob/main/taocp-7.2.2.3-ex81-ungraceful-Kn-x-Pr-and-Kn-x-Cr.pdf

[3515] W. C. Shiu, Super-edge-graceful labelings of some cubic graphs, *Acta Math. Sin.,* **22** (2006) 1621-1628. https://www.researchgate.net/publication/226808487_Super-edge-graceful_Labelings_of_Some_Cubic_Graphs

[3516] W. C. Shiu, Edge-magic labeling matrices of the composition of paths and null graphs, *Congr. Numer.,* **187** (2007) 55-68.

[3517] W. C. Shiu, The non-edge-magic simple connected cubic graph of order 10, *JCMCC* **83** (2012) 225-241. https://combinatorialpress.com/article/jcmcc/Volume%20083/vol-083-paper%2021.pdf

[3518] W. C. Shiu, Product-cordial index set for Cartesian product of a graph with a path, *J. Combin. Number Th.,* **4**(3) (2012) 177-194. https://shiu1958.github.io/PDF/P1230pub.pdf?utm_source=chatgpt.com

[3519] W. C. Shiu, Extreme edge-friendly indices of complete bipartite graphs, *Trans. Combin.,* **5**(3) (2016) 11-21. https://toc.ui.ac.ir/article_12473_5cd474dbde30ab0cf85638c76d56808a.pdf

[3520] W. C. Shiu, Full edge-friendly index sets of complete bipartite graphs, *Trans. Combin.,* **6**(2) (2017) 7-17. https://toc.ui.ac.ir/article_20739_045d1f054d5254d803c8df777bac0d71.pdf

[3521] W. C. Shiu, Integer-antimagic spectra of disjoint unions of cycles, *Theory Appl. Graphs,* **5**(2) (2018) Art. 3, 12 pp. https://digitalcommons.georgiasouthern.edu/cgi/viewcontent.cgi?article=1067&context=tag

[3522] W. C. Shiu and M.-H. Ho, Full friendly index sets of slender and flat cylinder graphs, *Transactions Combin.,* **2** (4) (2013) 63-80. https://scholars.hkbu.edu.hk/en/publications/full-friendly-index-sets-of-slender-and-flat-cylinder-graphs-7

[3523] W.-C. Shiu, G.-C. Laub, H.-K. Ng, Edge-coloring vertex-weighting of graphs, *Iranian J. Math. Sci. Informatics,* **16**(1) (2021) 1-13. https://www.researchgate.net/publication/226610853_On_vertex-coloring_edge-weighting_of_graphs

[3524] W. C. Shiu and H. Kwong, Full friendly index sets of $P_2 \times P_n$, *Discrete Math.,* **308** (2008) 3688-3693. https://www.sciencedirect.com/science/article/pii/S0012365X0700489X?via%3Dihub

[3525] W. C. Shiu and H. Kwong, An algebraic approach for finding balance index sets, *Australasian J. Combin.,* **45** (2009) 139-155. https://ajc.maths.uq.edu.au/download/45/ajc_v45_p139.pdf

[3526] W. C. Shiu and H. Kwong, Product-cordial index and friendly index of regular graphs, *Transactions Combin.*, **1** (2012) 15-20. https://toc.ui.ac.ir/article_482_0f0636556a2c2f345a1e9f8e54e0934d.pdf

[3527] W. C. Shiu, P. C. B. Lam, Super-edge-graceful labelings of multi-level wheel graphs, fan graphs and actinia graphs, *Congr. Numer.,* **174** (2005) 49-63. https://shiu1958.github.io/PDF/P0540pub.pdf

[3528] W. C. Shiu, P. C. B. Lam, and H. L. Cheng, Supermagic labeling of an s-duplicate of $K_{n,n}$, *Congr. Numer.,* **146** (2000) 119-124.

[3529] W. C. Shiu, P. C. B. Lam, and H. L. Cheng, Edge-gracefulness of the composition of paths with the null graphs, *Discrete Math.,* **253** (2002) 63-76. https://shiu1958.github.io/PDF/p0015fin.pdf

[3530] W. C. Shiu, P. C. B. Lam, and S. M. Lee, Edge-magic index sets of (p, p)-graphs, *Congr. Numer.,* **137** (1999) 97-107.

[3531] W. C. Shiu, P. C. B. Lam, and S. M. Lee, On a construction of supermagic graphs, *J. Combin. Math. Combin. Comput.,* **42** (2002) 147-160. https://combinatorialpress.com/article/jcmcc/Volume%20042/vol-042-paper%2010.pdf

[3532] W. C. Shiu, P. C. B. Lam, and S. M. Lee, Edge-magic indices of $(n, n-1)$-graphs, *Electron. Notes Discrete Math.,* **11** (2002) 443-458. https://www.academia.edu/8380632/Edge_magic_Indices_of_n_n_1_graphs

[3533] W. C. Shiu, P. C. B. Lam, and S. M. Lee, Edge-magic index sets of square of paths, *Util. Math.,* **97** (2015) 271-286.

[3534] W. C. Shiu, P. C. B. Lam, and P. K. Sun, Construction of group-magic graphs and some A-magic graphs with A of even order, *Congr. Numer.,* **167** (2004) 97-107.

[3535] W. C. Shiu and G.-C. Lau, Some results on k-edge-magic broken wheel graphs, *J. Graph Labeling,* **1**(2) (2015) 65-79. https://shiu1958.github.io/PDF/P1570pub.pdf

[3536] W.-C. Shiu, G.-C. Lau, and S.-M. Lee, On (semi-)edge-primality of graphs, *Iranian J. Math. Sci. and Informatics,* **12**(2) (2017) 1-14. http://ijmsi.ir/article-1-924-en.pdf

[3537] W. C. Shiu, G. C. Lau, and M. Nalliah, Local distance antimagic chromatic number of join product of graphs with cycles or paths, *Hacet. J. Math. Stat.*, **53(3)** (2024) 788-802. https://www.researchgate.net/publication/379828052_Local_Distance_Antimagic_Chromatic_Number_of_Join_Product_of_Graphs_with_Cycles_or_Paths

[3538] W.C. Shiu and G-C. Lau, On SD-prime labelings of some one point unions of gears, *Symmetry* **13**(849) (2021) 18 pp. https://doi.org/10.3390/sym13050849

[3539] W. C. Shiu and S. M. Lee, Some edge-magic cubic graphs, *J. Combin. Math. Combin. Comput.,* **40** (2002) 115-127. https://combinatorialpress.com/article/jcmcc/Volume%20040/vol-040-paper%208.pdf

[3540] W. C. Shiu and S. M. Lee, Full friendly index sets and full product-cordial index sets of twisted cylinders, *J. Combin. Number Thoery,* **3** (3) (2012) 209-216. https://shiu1958.github.io/PDF/P1150pub.pdf

[3541] W. C. Shiu, S. M. Lee, and K. Schaffer, Some k-fold edge-graceful labelings of $(p, p-1)$-graphs, *J. Combin. Math. Combin. Comput.,* **38** (2001) 81-95. https://combinatorialpress.com/article/jcmcc/Volume%20038/vol-038-paper%208.pdf

[3542] W. C. Shiu and M. H. Ling, Full friendly index sets of Cartesian products of two cycles, *Acta Mathematica Sinica, English Series,* **26** (2010) 1233-1244. https://www.researchgate.net/publication/238848457_Full_friendly_index_sets_of_Cartesian_products_of_two_cycles

[3543] W. C. Shiu, M. H. Ling, and R. M. Low, The entire edge-graceful spectra of cycles with one chord, Proc. Thirty-Seventh Southeastern International Conf. on Combin., Graph Theory and Comput., *Congr. Numer.,* **183** (2006) 213-219.

[3544] W. C. Shiu, M. H. Ling, and R. M. Low, The edge-graceful spectra of connected bicyclic graphs without pendant, *JCMCC,* **66** (2008) 171-185. https://combinatorialpress.com/article/jcmcc/Volume%20066/vol-066-paper%2015.pdf

[3545] W. C. Shiu and R. M. Low, Group magicness of complete N-partite graphs, *J. Combin. Math. Combin. Comput.,* **58** (2006) 129-134. https://combinatorialpress.com/article/jcmcc/Volume%20058/vol-058-paper%2013.pdf

[3546] W. C. Shiu and R. M. Low, Integer-magic spectra of sun graphs, *J. Comb. Optim.,* **14** (2007) 309-321. https://www.researchgate.net/publication/265494198_Integer-Magic_Spectra_of_Sun_Graphs

[3547] W. C. Shiu and R. M. Low, Ring-magic labelings of graphs, *Australasian J. Combin.,* **41** (2008) 147-158. https://ajc.maths.uq.edu.au/pdf/41/ajc_v41_p147.pdf

[3548] W. C. Shiu and R. M. Low, Z_k-magic labeling of fans and wheels with magic-value zero, *Australasian J. Combin.,* **45** (2009) 309-316. https://ajc.maths.uq.edu.au/pdf/45/ajc_v45_p309.pdf

[3549] W. C. Shiu and R. Low, The integer-magic spectra of bicyclic graphs without pendant, Proceedings of the Forty-Third Southeastern International Conference on Combinatorics, Graph Theory and Computing. *Congr. Numer.,* **214** (2012) 65-73

[3550] W. C. Shiu and R. M. Low, Group-magic labelings of graphs with deleted edges, *Australasian J. Combin.,* **57** (2013) 3-19. https://ajc.maths.uq.edu.au/pdf/57/ajc_v57_p003.pdf

[3551] W. C. Shiu and R. M. Low, Integer-antimagic spectra of complete bipartite graphs and complete bipartite graphs with a deleted edge, *Bull. Inst. Combin. Appl.,* **76** (2016) 54-68.

[3552] W. C. Shiu and R. M. Low, The integer-antimagic spectra of dumbbell graphs, *Bull. ICA,* **77** (2016) 89-110.

[3553] W. C. Shiu and R. M. Low, The integer-magic spectra and null sets of the Cartesian product of trees, *Austral. J. Combin.,* **70**(1) (2018) 157-167. https://ajc.maths.uq.edu.au/pdf/70/ajc_v70_p157.pdf

[3554] W. C. Shiu and R. M. Low, On the integer-antimagic spectra of non-Hamiltonian graphs, *Theory Appl. Graphs,* **9**(2) (202a2) Article 8 1-19. https://digitalcommons.georgiasouthern.edu/tag/vol9/iss2/8

[3555] W. C. Shiu, R. Santrin Sabibha, P. Jeyanthi, K. Jeya Daisy, and G. C. Lau, Bijective product and product square k-cordial labeling of graphs, *Jordan J. Math. Stat.,* **17**(3) (2024) 391-400. https://jjms.yu.edu.jo/index.php/jjms/article/view/442/40

[3556] W. C. Shiu, P. K. Sun, and R. M. Low, Integer-antimagic spectra of tadpole and lollipop graphs, *Congr. Numer.*, (2015) **225** (2015) 5-22. https://www.researchgate.net/publication/283085888_Integer-antimagic_spectra_of_tadpole_and_lollipop_graphs

[3557] W. C. Shiu and F. S. Wong, Extreme friendly indices of $C_m \times P_n$, *Cong. Numer.,* **197** (2009) 65-75.

[3558] W. C. Shiu and F. S. Wong, Strong vertex-graceful labeling for some double cycles, *Congr. Numer.,* **20** (2010) 17-24. https://shiu1958.github.io/PDF/P1050pub.pdf

[3559] W. C. Shiu and F. S. Wong, Full friendly index sets of cylinder graphs, *Australasian J. Combin.,* **52** (2012) 141-162. https://ajc.maths.uq.edu.au/pdf/52/ajc_v52_p141.pdf

[3560] C.-L. Shiue and H.-L. Fu, α-labeling number of trees, *Discrete Math.,* **306** (2006) 3290-3296. https://www.sciencedirect.com/science/article/pii/S0012365X06004766?via%3Dihub

[3561] C.-L. Shiue and H.-L. Fu, The IC-indices of complete bipartite graphs, *Electron. J. Combin.,* **15**(1) (2008) Research paper 43, 13 pages. https://www.combinatorics.org/ojs/index.php/eljc/article/view/v15i1r43/pdf

[3562] C.-L. Shiue and H.-C. Lu, Trees which admit no α-labelings, *Ars Combin.,* **103** (2012) 453-463. https://combinatorialpress.com/article/ars/Volume%20103/volume-103-paper-38.pdf

[3563] M. A. Shivarajkumar, Sriraj, and S. M. Hegde, Graceful labeling of digraphs—a survey, *AKCE Internat. J. Graphs Combin.*, (2021) 1-5. https://doi.org/10.1080/09728600.2021.1978014

[3564] S. Shixin and S. Yu, On Hamming-graceful graphs, unpublished.

[3565] N. P. Shrimali and P. B. Pandya, Total neighborhood prime labeling of some graphs, *Internat. J. Sci. Res. Math. Stat. Sci.*, **5**(6) (2018) 157-163. https://www.researchgate.net/publication/330338335_Total_Neighborhood_Prime_Labeling_of_Some_Graphs

[3566] N. P. Shrimali and Y. M. Parmar, Edge vertex prime labeling for some trees, *J*ASC: J. Appl. Sci. Comput., **VI**(I) (2019) 2136-1249. https://www.researchgate.net/publication/330959323_Edge_vertex_prime_labeling_for_some_trees

[3567] N. P. Shrimali and Y. M. Parmar, Edge vertex prime labeling for some graphs, *South East Asian J. Math. Math. Sci.,* **16**(1) (2020), 149-156. https://rsmams.org/download/articles//2_16_1_1231044381_Paper%2013%20EDGE%20VERTEX%20PRIME%20LABELING%20FOR%20SOME%20GRAPHS.pdf

[3568] N. P. Shrimali and Y. M. Parmar, Distance magic and distance antimagic labeling of some product graphs, *Recent Advancements in Graph Theory*, CRC Press, (2020) 181-191. https://www.researchgate.net/publication/346580733_Distance_Magic_and_Distance_Antimagic_Labeling_of_Some_Product_Graphs

[3569] N. P. Shrimali, Y. M. Parmar, and M. A. Patel, Total neighborhood magic labeling for complete graph, path and $C_n \odot K_1$, *Internat. J. Sci. Res. Math. Stat. Sci.,* **11**(3) (2024) 01-05. https://www.isroset.org/pdf_paper_view.php?paper_id=3531&1-ISROSET-IJSRMSS-09614.pdf

[3570] N. P. Shrimali and A. K. Rathod, Vertex-edge neighborhood prime labeling of some trees, *South East Asian J. Math. Math. Sci.,* **16**(3) (2020) 207-218. http://rsmams.org/download/articles/2_16_3_838091879_Paper%2017%20%20VERTEX-EDGE%20NEIGHBORHOOD%20PRIME%20LABELING%20OF%20SOME%20TREES.pdf

[3571] N. P. Shrimali, A. K. Rathod, and P. L. Vihol, Neighborhood-prime labeling for some graphs, *Malaya J. Mat.,***7**(1) (2019) 108-112. http://www.malayajournal.org/selected_article.php?id=538

[3572] N. P. Shrimali and S. K. Singh, Gaussian vertex prime labeling of some graphs, *J. Chungcheong Math. Soc.*. **34**(3) (2021) 285–294. https://www.koreascience.or.kr/article/JAKO202123758839918.pdf

[3573] N. P. Shrimali and S. T. Trivedi, Gaussian neighborhood-prime labeling of graphs containing Hamiltonian cycle, *Math. J. Interdiscip. Sci.,* **7**(2) (2019) 161-165. https://mjis.chitkara.edu.in/index.php/mjis/article/view/190/125

[3574] A. Shulhany, Y. Rukmayadi, A. D. Maharani, and Z. Syaifara, The total H-irregularity strength of some graph classes, AIP Conference Proceedings 2468 (2022), 070005

[3575] M. K. Siddiqui, On the total edge irregularity strength of a categorical product of a cycle and a path, *AKCE J. Graphs, Combin.,* **9**(1) (2012) 43-52.

[3576] M. K. Siddiqui, On edge irregularity strength of subdivision of star S_n, *Internat. J. Math. Soft Comput.,* **2**(1) (2012) 75-82. https://www.researchgate.net/publication/280628183_On_Edge_Irregularity_Strength_of_Subdivision_of_Star_SN

[3577] M. K. Siddiqui, On irregularity strength of convex polytope graphs with certain pendent edges added, *Ars Combin.,* **129** (2016) 199-210. https://combinatorialpress.com/article/ars/Volume%20129/volume-129-paper-16.pdf

[3578] M. K. Siddiqui, On Z_{3k}-magic labeling of certain families of graphs, unpublished.

[3579] M. K. Siddiqui, A. Ahmad, M. F. Nadeem, and Y. Bashir, Total edge irregularity strength of the disjoint union of sun graphs, *Internat. J. Math. Soft Comput.,* **3**(1) (2013) 21-27. https://www.academia.edu/29127460/Total_edge_irregularity_strength_of_the_disjoint_union_of_sun_graphs

[3580] M. K. Siddiqui, M. Miller, and J. Ryan, Total edge irregularity strength of octagonal grid graph, *Util. Math.,* **103** (2017) 277-287. https://www.researchgate.net/publication/320466762_Muhammad_Kamran_Siddiqui_M_Miller_J_Ryan_Total_edge_irregularity_strength_of_octagonal_grid_graph_Utilitas_Mathematica_Volume_103_Pages_277-287_Published_JUN_2017

[3581] M. K. Siddiqui, M. Numan, and M. A. Umar, Face antimagic labeling of Jahangir graph, *Math. Comput. Sci.,* **7**(2) (2013) 237-243. https://www.researchgate.net/publication/257759496_Face_Antimagic_Labeling_of_Jahangir_Graph

[3582] D. R. Silaban, A. Parestu, B. N. Herawati, K. A. Sugeng, and Slamin, Vertex-magic total labelings of unions of generalized Petersen graphs and union of special circulant graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 201-207.

[3583] D. R. Silaban and K. A. Sugeng, Edge antimagic total labelings on paths and uncycles, *J. Combin. Math. Combin. Comput.,* **65** (2007) 1-8. https://www.researchgate.net/publication/265587900_Edge_antimagic_total_labeling_on_paths_and_unicycles#fullTextFileContent

[3584] M. Simanihuruk, T. A. Kusmayadi, B. Swita, M. Romala, and F. Damanik, A conjecture on super edge-magic total labeling of 4-cycle books, *Int. J. Math. Math. Sci.,* (2021) Art. ID 8483926, 8 pp. https://www.hindawi.com/journals/ijmms/2021/8483926/

[3585] R. Simanjuntak, F. Bertault, and M. Miller, Two new (a, d)-antimagic graph labelings, *Proc. Eleventh Australia Workshop Combin. Algor.*, Hunrer Valley, Australia (2000) 179-189.

[3586] R. Simanjuntak, T. Nadeak, F. Yasin, K. Wijaya, N. Hinding, and K. A. Sugeng, Another antimagic conjecture, *Symmetry* (2021) **13** 2071 https://www.mdpi.com/2073-8994/13/11/2071

[3587] R. Simanjuntak and K. Wijaya, On distance antimagic graphs, arXiv:1312.7405.

[3588] M. Simaringa and S. Muthukumaran, Edge vertex prime labeling of some graphs, *Malaya J. Matematik,* **7**(2) (2019) 264-268. http://www.malayajournal.org/articles/MJM07020018.pdf

[3589] M. Simaringa and S. Muthukumaran, Edge vertex prime labeling of graphs, *Malaya J. Mat.,* **7**(3) (2019) 572-578. https://www.malayajournal.org/articles/MJM07030034.pdf

[3590] M. Simaringa and S. Muthukumaran, Edge vertex prime labeling of union of graphs, *Internat. J. Engin. Adv. Tech,;* **9**(1s6) (2019) 89-97. DOI:10.35940/ijeat.A1017.1291S619

[3591] M. Simaringa and S. Thirunavukkarasu, An analytic odd mean labeling of some new results of graphs, *Nonlinear Stud.,* **28**(3) (2021) 795-799.

[3592] N. Simarmata, I. P. Sandy, and K. A. Sugeng, Graceful labeling construction for some special tree graph using adjacency matrix, Elect. J. Graph Th. Appl., **11**(2) (2023) 343-356. https://www.ejgta.org/index.php/ejgta/article/view/1677/pdf_289

[3593] G. J. Simmons, Synch-sets: a variant of difference sets, *Proc. 5th Southeastern Conference on Combinatorics, Graph Theory and Computing*, Util. Math. Pub. Co., Winnipeg (1974) 625-645.

[3594] M. Sindhu and S. C. Kumar, Even vertex in-magic total labeling of some 2-regular digraphs, *J. Elect., Comput. Networking Appl. Math.*, **01**(02) (2021) 1-11. https://journal.hmjournals.com/index.php/JECNAM/article/view/161/933

[3595] S. Singla, A. Tiwari, and A. Tripathi, Some results on the spum and the integral spum of graphs, *Discrete Math.,* **344**(5) (2021) 112311. https://doi.org/10.1016/j.disc.2021.112311

[3596] D. Sinha and J. Kaur, Edge-friendly labelings of graph–II, *Util. Math.,* **99** (2016) 251-271. https://www.researchgate.net/publication/316284683_Edge_friendly_labelings_of_the_graph-II

[3597] D. Sinha and J. Kaur, Region indices of friendly labeling of a planar graph, *Util. Math.,* **101** (2016) 149-178. https://www.researchgate.net/publication/312138142_Region_indices_of_friendly_labelling_of_planar_graph

[3598] I. Singgih, New Methods for Magic Total Labelings of Graphs, Masters Thesis, Department of Mathematics and Statistics, U. Minnesota Duluth, 2015.

[3599] S. Singla, A. Tiwari, and A. Tripathi, Some results on the spum and the integral spum of graphs,*Discrete Mathematics* **344** (2021) 11231 https://iopscience.iop.org/article/10.1088/1742-6596/1778/1/012009/pdf

[3600] G. Suresh Singh, A note on graceful prisms, *Nat. Acad. Sci. Lett.,* **15** (1992) 193-194.

[3601] G. Suresh Singh, Subdivisions of ladders are arithmetic, *Multidiscplinary Research Review,* **2** (1992) 23-25.

[3602] G. Suresh Singh, A note on sequential crowns, *Nat. Acad. Sci. Lett.*, **16** (1993) 243-245.

[3603] G. Suresh Singh, A note on labeling of graphs, *Graphs and Combin.*, **14** (1998) 201-207.

[3604] G. Suresh Singh, A note on labeling of graphs, *Graphs and Combin.*, **14** (1998) 201-207.

[3605] G. Suresh Singh, Some generalities on arithmetic graphs, *Graph Theory Notes of New York,* **XXXVIII** (2000) 12-16.

[3606] G. Suresh Singh, personal communication (2015)

[3607] G. Suresh Singh, Divisor graphs-I, unpublished.

[3608] G. Suresh Singh, and J. Devaraj, On triangular graceful graphs, unpublished.

[3609] G. Suresh Singh, and G. Santhosh, A note on integral sum crowns, *Ars Combin.,* **66** (2003) 65-77. https://combinatorialpress.com/article/ars/Volume%20066/volume-66-paper-5.pdf

[3610] G. Singh, N. Takshak, A. Sehgal, and A. Malik, On graceful labeling of some power graphs, *J. Indones. Math. Soc.*, **32**(03) (2026) 1–13. https://jims-a.org/index.php/jimsa/article/view/2000/500

[3611] G. Suresh Singh, and V. Vilfred, Some results on arithmetic trees, unpublished.

[3612] N. M. Singhi, G. R. Vijayakumar and N. Usha Devi, Set-magic labelings of infinite graphs. *Ars Combin.,* **63** (2002) 305-310. https://combinatorialpress.com/article/ars/Volume%20063/volume-63-paper-25.pdf

[3613] S. Singhun, Graphs with edge-odd graceful labelings, *Int. Math. Forum,* **8**(9-12) (2013) 577-582. http://www.m-hikari.com/imf/imf-2013/9-12-2013/singhunIMF9-12-2013.pdf

[3614] S. Singhun, R. Boonklurb, and C. Charnsamorn, A supermagic labeling of finite copies of Cartesian product of cycles, *East-West J. Math.,* **17**(1) (2015) 61-69.

[3615] D. Sinha and J. Kaur, Full friendly index set-I, *Discrete Appl. Math.,* **161**(9) (2013) 1262-1274. https://www.researchgate.net/publication/257001055_Full_friendly_index_set_-_I

[3616] D. Sinha and J. Kaur, Full friendly index set-II, *J. Combin. Math. Combin. Comput.,* **79** (2011) 65-75. https://www.researchgate.net/publication/274378729_Full_Friendly_index_set-_II

[3617] D. Sinha and J. Kaur, Edge friendly labelings of graph-I, *Ars Combin.,* **129** (2016) 139-156. https://www.researchgate.net/publication/312138495_Edge_friendly_labelings_of_the_graph-I

[3618] D. Sinha and J. Kaur, Region indices of friendly labeling of a planar graph, *Util. Math.,* **101** (2016) 149-178. https://www.researchgate.net/publication/312138142_Region_indices_of_friendly_labelling_of_planar_graph

[3619] D. Sinha and J. Kaur, Region indices of edge-friendly labeling of a plane graph, *Util. Math.,* **99** (2016) 187-213. https://www.researchgate.net/publication/301205528_Region_Indices_of_Edge-friendly_labeling_of_a_Plane_Graph

[3620] Siqinbate and W. Feng, Wei, Gracefulness of the digraph $3\vec{C}_{2p}$, *Pure Appl. Math.* (Xi'an) **29**(2) (2013) 111-117.

[3621] Siqinqimuge, Jirimutu, and L. Zhao, On the gracefulness of the digraphs n-$\vec{C}_{19}$ for even n, *Util. Math.,* **85** (2011) 193-199.

[3622] M. Sivakumar, On 4-total product cordiality of some corona graphs, *Internat. J. Math. Combin.,* **3** (2016) 99-106. https://fs.unm.edu/IJMC/On4TotalProductCordialityOfSomeCorona.pdf

[3623] S. Sivakumar, S. Vidyanandini, E. Sreedevi, S. R. Nayak, and A. K. Bhoi, Edge irregularity strength exists in the anti-theft network, In: P. K. Mallick, A. K. Bhoi, P. Barsocchi, and V. H. C. de Albuquerque, (eds), *Cognitive Informatics Soft Comput.*, Lecture Notes in Networks and Systems, **375** (2022) Springer, Singapore.

[3624] V. Sivakumaran, K. Sankar, and S. Prabhu, Distance antimagic labeling of zero-divisor graphs, arXiv:2407.08211v1 https://www.researchgate.net/publication/382178711_Distance_Antimagic_Labeling_of_Zero-Divisor_Graph

[3625] R. Sivaraman, Graceful graphs and its applications, *Internat. J. Current Research,* **8**(11) (2016) 41062-41067. https://www.journalcra.com/sites/default/files/issue-pdf/18683.pdf

[3626] T. Skolem, On certain distribution of integers into pairs with given differences, *Math. Scand.*, **5** (1957) 57-68. https://gdz.sub.uni-goettingen.de/id/PPN35397434X_0005?tify=%7B%22view%22:%22info%22,%22pages%22:%5B60%5D%7D

[3627] J. Skowronek-Kaziów, Multiplicative vertex-colouring weightings of graphs, *Inform. Process. Lett.,* **112**(5) (2012) 191-194.

[3628] S. Slamet, K. Sugeng, and M. Miller, Sum graph based access structure in a secret sharing scheme, *J. Prime Research Math.,* **2** (2006) 113-119. https://www.researchgate.net/publication/228869655_Sum_graph_based_access_structure_in_a_secret_sharing_scheme

[3629] Slamin, M. Bača, Y. Lin, M. Miller, and R. Simanjuntak, Edge-magic total labelings of wheels, fans and friendship graphs, *Bull. ICA,* **35** (2002) 89-98. https://www.researchgate.net/publication/234058521_Edge-magic_total_labelings_of_wheels_fans_and_friendship_graphs

[3630] Slamin, Dafik, and W. Winnona, Total vertex irregularity strength of the disjoint union of sun graphs, *Int. J. Comb.,* **2012** Art. ID 284383, 9 pp. https://www.emis.de/journals/HOA/IJCT/Volume2012/284383.pdf

[3631] Slamin and M. Miller, On two conjectures concerning vertex-magic total labelings of generalized Petersen graphs, *Bull. Inst. Combin. Appl.,* **32** (2001) 9-16. https://www.researchgate.net/publication/234057840_On_two_conjectures_concerning_vertex_magic_total_labelings_of_generalized_Petersen_graphs

[3632] Slamin, A. C. Prihandoko, T. B.Setiawan, V. Rosita, and B. Shaleh, Vertex-magic total labelings of disconnected graphs, *J. Prime Research in Math.,* **2** (2006) 147-156. https://www.researchgate.net/publication/234058013_Vertex-magic_Total_Labelings_of_Disconnected_Graphs

[3633] P. J. Slater, On k-sequential and other numbered graphs, *Discrete Math.,* **34** (1981) 185-193. https://www.sciencedirect.com/science/article/pii/0012365X81900662?via%3Dihub

[3634] P. J. Slater, On k-graceful graphs, *Proc. of the 13th S. E. Conf. on Combinatorics, Graph Theory and Computing,* (1982) 53-57.

[3635] P. J. Slater, On k-graceful, locally finite graphs, *J. Combin. Theory, Series B,* **35** (1983) 319-322. https://www.sciencedirect.com/science/article/pii/0095895683900588

[3636] P. J. Slater, Problems in graph theory: graceful and sequential numbering of infinite graphs, *Southeast Asian Bull. Math.,* **9** (1985) 15-22.

[3637] P. J. Slater, On k-graceful, countably infinite graphs, *Discrete Math.,* **61** (1986) 293-303. https://www.sciencedirect.com/science/article/pii/0012365X86901007

[3638] R. Slíva, Antimagic labeling graphs with a regular dominating subgraph, *Inform. Process. Lett.,* **112**(21) (2012) 844-847.

[3639] D. Small, Regular (even) spider graphs are edge-graceful, *Congr. Numer.,* **74** (1990) 247-254.

[3640] B. M. Smith, Set of symmetry breaking constraints, (2005) https://pdfs.semanticscholar.org/c2cf/ccdb6f07044b87bb4e5a1a152d24939c7540.pdf

[3641] B. M. Smith and J.-F Puget, Constraint models for graceful graphs, *Constraints,* **15** (1) (2010) 64-92. https://www.researchgate.net/publication/220202433_Constraint_models_for_graceful_graphs

[3642] W. Smyth, Sum graphs of small sum number, *Colloquia Mathematica Societatis János Bolyai*, **60** (1991) 669-678. https://korandi.org/docs/misc/setsgraphsnumbers/setsgraphsnumbers46.pdf

[3643] W. Smyth, Sum graphs: New results, new problems, *Bull. Inst. Combin. Appl.,* **2** (1991) 79-81.

[3644] H. Snevily, Combinatorics of Finite Sets, Ph. D. Thesis, U. Illinois, 1991.

[3645] H. Snevily, New families of graphs that have α-labelings, *Discrete Math.*, **170** (1997) 185-194. https://www.sciencedirect.com/science/article/pii/0012365X9500159T

[3646] B. Sokolowsky, Achieving all radio numbers, *AKCE Int. J. Graphs Comb.,* **10**(1) (2013) 77-95. https://www.tandfonline.com/doi/epdf/10.1080/09728600.2013.12088726?needAccess=true

[3647] A. Solairaju and M. A. Arockiasamy, Gracefulness of k-step staircase graphs, *J. Analysis Computation,* **6** (2010) 109-114.

[3648] A. Solairaju and M. A. Arockiasamy, Graceful mirror-staircase graphs, *Inter. J. Contemporary Math. Sci.,* **5** (2010) 2433-2441. https://www.m-hikari.com/ijcms-2010/49-52-2010/arockiasamyIJCMS49-52-2010.pdf

[3649] A. Solairaju and M. A. Arockiasamy, A study of labeling techiques in relations to a new family of staircase graphs, unpublished.

[3650] A. Solairaju and K. Chithra, New classes of graceful graphs by merging a finite number of C_4, *Acta Cienc. Indica Math.,* **34** (2008) 959-965.

[3651] A. Solairaju and K. Chithra, Edge-odd graceful labeling of some graphs, *Proceedings of the ICMCS,* **1** (2008) 101-107.

[3652] A. Solairaju and K. Chithra, Edge odd graceful graphs, *Electronic Notes in Discrete Math.,* **33** (2009) 15-20. https://www.sciencedirect.com/science/article/pii/S1571065309000225

[3653] M. Soleha, Purwanto and D. Rahmadani, Edge odd graceful of alternate snake graphs, 2022 J. Phys.: Conf. Ser. 2157 012002 https://iopscience.iop.org/article/10.1088/1742-6596/2157/1/012002/pdf

[3654] D. D. Somashekara and C. R. Veena, On strong vertex graceful graphs, *Internat. Math. Forum,* **5**(56) (2010) 2751-2757. https://scispace.com/pdf/on-strong-vertex-graceful-graphs-39sthdlffi.pdf

[3655] D. D. Somashekara and C. R. Veena, On square sum labelings of graphs, *Proc. Jangjeon Math. Soc,* **15**(1) (2012) 69-78. https://www.kci.go.kr/kciportal/landing/article.kci?arti_id=ART001629880 cleck on PDF button.

[3656] D. D. Somashekara and C. R. Veena, On pentagonal sum labelings of graphs, *Adv. Studies Contemp. Math.*, **22**(3) (2012) 421-432.

[3657] S. Somasundaram and R. Ponraj, Mean labelings of graphs, *Natl. Acad, Sci. Let.*, **26** (2003) 210-213.

[3658] S. Somasundaram and R. Ponraj, Non-existence of mean labeling for a wheel, *Bull. Pure and Appl. Sciences (Mathematics & Statistics)*, **22E** (2003) 103-111.

[3659] S. Somasundaram and R. Ponraj, Some results on mean graphs, *Pure and Applied Math. Sci.*, **58** (2003) 29-35.

[3660] S. Somasundaram and R. Ponraj, On mean graphs of order < 5, *J. Decision Math. Sci.*, **9** (2004) 47-58.

[3661] S. Somasundaram, R. Ponraj, and S. S. Sandhya, Harmonic mean labeling of graphs, unpublished.

[3662] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, Super Lehmer-3 mean labeling, *J. Math. Res.,* **8**(5) (2016) 29-36. https://pdfs.semanticscholar.org/88ca/6b1c41a575da4d72524c9084262d21d3c976.pdf

[3663] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, Lehmer-3 mean labeling of some new disconnected graphs, *Internat. J. Math. Trends Tech.* (IJMTT), **35**(1) (2016) 1-14. https://ijmttjournal.org/public/assets/volume-35/number-1/IJMTT-V35P501.pdf

[3664] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, Lehmer-3 mean labeling of some disconnected graphs, *Internat. J .Math. Res.*, **8**(2) 2016) 133-142. http://www.irphouse.com/ijmr16/ijmrv8n2_08.pdf

[3665] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, Lehmer-3 mean labeling of graphs, *Internat. J. Math. Forum*, **12**(17) (2017) 819-825. https://www.m-hikari.com/imf/imf-2017/17-20-2017/p/pavithraIMF13-16-2017.pdf

[3666] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, k-Super Lehmer-3 mean graph, *Internat. J. Sci. Innov. Math. Res.* (IJSIMR), **5**(5) (2017) 1-7. https://www.arcjournals.org/pdfs/ijsimr/v5-i5/1.pdf

[3667] S. Somasundaram, S. S. Sandhya, and T. S. Pavithra, Super Lehmer-3 mean number of graphs, *Global J. Pure Appl. Math.*, **13**(10) (2017) 7377-7385. https://www.ripublication.com/gjpam17/gjpamv13n10_32.pdf

[3668] S. Somasundaram, P. Vidhyarani and R. Ponraj, Geometric mean labelings of graphs, *Bull. Pure Appl. Sci,*, **30E**(2) (2011) 153-160.

[3669] S. Somasundaram, P. Vidhyarani and S. S. Sandhya, Some results on geometric mean graphs, *Internat. Math. Forum*, **7**(28) (2012) (28) 1381-1391. https://www.m-hikari.com/imf/imf-2012/25-28-2012/vidhyaraniIMF25-28-2012.pdf

[3670] S. Sonchhatra and G. V. Ghodasara, Sum perfect square graphs in context of some graph operations, *Internat. J. Math. Trends Tech. (IJMTT),* **46**(2) (2017) 62-65. https://ijmttjournal.org/public/assets/volume-46/number-2/IJMTT-V46P512.pdf

[3671] S. G. Sonchhatra and G. V. Ghodasara, Sum perfect square labeling of graphs, *Internat. J. Scient. Innovative Math. Res.,* **4** (2016) 64-70. https://www.arcjournals.org/pdfs/ijsimr/v4-i1/9.pdf

[3672] S. G. Sonchhatra and G. V. Ghodasara, Snakes related sum perfect square graphs, *Internat. J. Mathematics Appl.*, **4** (2016) 157-168. https://ijmaa.in/index.php/ijmaa/article/view/980/964

[3673] S. G. Sonchhatra and G. V. Ghodasara, Some new permutation graphs, *Internat. J. Math. Archive*, **7** (2016) 209-215. https://www.researchgate.net/publication/318815286_Some_New_Permutation_Graphs

[3674] S. G. Sonchhatra and G. V. Ghodasara, Some new sum perfect square graphs, *Internat. J. Pure Appl. Math.*, **113**(3) (2017) 489-499. https://www.researchgate.net/publication/317340424_SOME_NEW_SUM_PERFECT_SQUARE_GRAPHS

[3675] S. G. Sonchhatra, D. D. Pandya, and T. M. Chhaya, Sum perfect cube labeling of graphs, *The Sci. Temper,* **15**(2) (2024) 2397-2400, doi:Doi:10.58414/SCIENTIFICTEMPER.2024.15.2.57

[3676] M. Sonntag, Antimagic vertex-labelling of hypergraphs, *Discrete Math.,* **247** (2002) 187-199. https://www.sciencedirect.com/science/article/pii/S0012365X01001753

[3677] M. Sonntag, Difference labelling of cacti, *Discuss. Math. Graph Theory,* **23** (2003) 55-65. http://dml.mathdoc.fr/item/bwmeta1.element.bwnjournal-article-doi-10_7151_dmgt_1185/

[3678] M. Sonntag and H.-M. Teichert, Sum numbers of hypertrees, *Discrete Math.,* **214** (2000) 285-290. https://www.sciencedirect.com/science/article/pii/S0012365X99003076?via%3Dihub

[3679] M. Sonntag and H.-M. Teichert, On the sum number and integral sum number of hypertrees and complete hypergraphs, *Discrete Math.,* **236** (2001) 339-349. https://www.sciencedirect.com/science/article/pii/S0012365X00004520?via%3Dihub

[3680] B. Sooryanarayana and P. Raghunath, Radio labeling of cube of a cycle, *Far East J. Appl. Math.,* **29** (2007) 113-147.

[3681] B. Sooryanarayana, M. Vishu Kumar, K. Manjula, Radio number of cube of a path, *Int. J. Math. Comb.*, **1** (2010) 5-29. https://fs.unm.edu/IJMC/IJMC-1-2010.pdf?utm_source=chatgpt.com

[3682] D. Speyer and Z. Szaniszló, Every tree is 3-equitable, *Discrete Math.,* **220** (2000) 283-289. https://www.sciencedirect.com/science/article/pii/S0012365X00000339?via%3Dihub

[3683] R. Sridevi, S. Navaneethakrishnan, and K. Nagarajan, Almost super Fibonacci graceful labeling, *Scientia Magna*, **6**(3) (2010) 99-125. Not on web https://www.sciencedirect.com/science/article/pii/S0012365X00000339

[3684] R. Sridevi, S. Navaneethakrishnan, and K. Nagarajan, Super Fibonacci graceful labeling, *Internat. J. Math. Combin.*, **3** (2010) 22-40. https://vixra.org/pdf/1403.0419v1.pdf

[3685] R. Sridevi, S. Navaeethakrishnan, A. Nagarajan, and K. Nagarajan, Odd-even graceful graphs, *J. Appl. Math. Inform.,* **30**(5-6) (2012) 913-923. https://www.koreascience.kr/article/JAKO201219164030653.pdf

[3686] R. Sridevi, K. Nagarajan, A. Nellaimurugan, and S. Navaneethakrishnan, Fibonacci divisor cordial graphs, *Interat. J. Math. Soft Comput.,* **3**(3) (2013) 33-39.

[3687] R. Sridevi, K. Palani, S. Navanaeethakrishnan, and K. Nagarajan, Some results on Fibonacci divisor cordial graphs, https://www.researchgate.net/publication/334654897_Some_Results_on_Fibonacci_Divisor_Cordial_Graphs

[3688] R. Sridevi and S. Ragavi, Lucky-edge labeling of K_n and special types of graphs, *Internat. J. Math. Appl.*, **4**(1-C) (2016) 125–131. https://ijmaa.in/index.php/ijmaa/article/view/605/600

[3689] N. Sridharan and R. Umarani, E_k-cordial labellings of graphs, *Elixir Appl. Math.,* **38** (2011) 4564-4567. https://www.elixirpublishers.com/articles/1685940962_201109103.pdf

[3690] V. Srinivasan, N. Chidambaram, N. Devadoss, R. Pakkirisamy, and P. Krishnamoorthi, On the gracefulness of m-super subdivision of graphs, *J. Discrete Math. Sci. Crypt.*, **23**(6) (2020) 1359-1368.

[3691] S. Sriramr and R. Govindarajan, Permutation labeling of joins of kite graph, *Internat. J. Comput. Engin. Tech.,* (IJCET) **10**(3) (2019) 1–8, Article ID: IJCET_10_03_001. https://iaeme.com/MasterAdmin/Journal_uploads/IJCET/VOLUME_10_ISSUE_3/IJCET_10_03_001.pdf

[3692] S. Stalin Kumar and G. T. Marimuthu, $H-V$-super magic decomposition of complete bipartite graphs, *Commun. Korean Math. Soc.,* **30**(3) (2015) 313-325. https://www.researchgate.net/publication/280729062_H-V_-SUPER_MAGIC_DECOMPOSITION_OF_COMPLETE_BIPARTITE_GRAPHS

[3693] R. P. Stanley, Linear homogeneous Diophantine equations and magic labelings of graphs, *Duke Math. J.,* **40** (1973) 607-632. https://scispace.com/pdf/linear-homogeneous-diophantine-equations-and-magic-labelings-3bfoyed4dl.pdf

[3694] R. P. Stanley, Magic labelings of graphs, symmetric magic squares, systems of parameters and Cohen-Macaulay rings, *Duke Math. J.,* **43** (1976) 511-531. https://math.mit.edu/~rstan/pubs/pubfiles/31.pdf

[3695] R. Stanton and C. Zarnke, Labeling of balanced trees, *Proc. 4th Southeast Conf. Combin., Graph Theory, Comput.,* (1973) 479-495.

[3696] V. L. Stella Arputha Mary, S. Navaneethakrishnan, and A. Nagarajan, Strong Z_{4p}-magic labeling, *Ars Combin.,* **135** (2017) 197-211. https://combinatorialpress.com/article/ars/Volume%20135/volume-135-paper-16.pdf

[3697] B. M. Stewart, Magic graphs, *Canadian J. Math.,* **18** (1966) 1031-1059. https://www.cambridge.org/core/services/aop-cambridge-core/content/view/3FE1BA25EBC6130A8BC1544934A8D928/S0008414X00041006a.pdf/magic-graphs.pdf

[3698] B. M. Stewart, Supermagic complete graphs, *Canadian J. Math.,* **19** (1967) 427-438. https://www.cambridge.org/core/journals/canadian-journal-of-mathematics/article/supermagic-complete-graphs/B3910B84BD52F5ECB703F4EC446EB3DF

[3699] R. J. Stones, Do the windmills $K_5^{(3)}$ and $K_5^{(4)}$ admit graceful labellings?, https://mathoverflow.net/q/424682

[3700] A. Su, J. Buchanan, R. C. Bunge, E. Pelttari, G. Rasmuson, E. Sparks, and S. Tagaris, On decompositions of complete multipartite graphs

into the union of two even cycles, *J. Combin. Math. Combin. Comput., 96* (2016) 129-157. https://combinatorialpress.com/article/jcmcc/Volume%20096/vol-096-paper%209.pdf

[3701] J. Su, H. Wang, and B. Yao, On elegant labelling and magic labelling of large-scale graphs, *Discrete Dynamics in Nature and Society*, **2022** (2022) Article ID 6301674, 10 pages https://www.hindawi.com/journals/ddns/2022/6301674/

[3702] G. Subashini, K. Bhuvaneswari, and K. Manimekalai, Square difference labeling of theta graphs, *Internat. J. Engin. Res. Tech.,* (IJERT), **8**(9) (2019) 343-345 https://www.ijert.org/research/square-difference-labeling-of-theta-graphs-IJERTV8IS090122.pdf

[3703] M. Subbiah, A Study on Some Variations of Graph Labeling and its Applications in Various Fields, Ph. D. Thesis, Depart. Math., (2011), Bharathidasan University.

[3704] S. P. Subbiah and J. Pandimadevi, $H-E$-super magic decomposition of graphs, *Electron. J. Graph Theory Appl. (EJGTA)*, **2**(2) (2014) 115-128. https://www.ejgta.org/index.php/ejgta/article/view/51/23

[3705] S. P. Subbiah, J. Pandimadevi, and R. Chithra, Super total graceful graphs, *Elect. Notes Disc. Math.*, **48** (2015) 301-304. doi:DOI: 10.1016/j.endm.2015.05.045

[3706] G. Subhashini, K. Ramanathan and K. Manimekalai, Square sum labeling for pyramid graph and hanging pyramid graph, *Internat. Math. Appl.*, **6**(1) (2019) 97-102. https://ijmaa.in/index.php/ijmaa/article/view/1401/1359

[3707] I. W. Sudarsana, E. T. Baskova, D. Ismaimuza, and H. Assiyatun, Creating new super edge-magic total labelings from old ones, *J. Combin. Math. Combin. Comput.,* **55** (2005) 83-90. https://www.researchgate.net/publication/228940964_Creating_new_super_edge-magic_total_labelings_from_old_ones

[3708] I. W. Sudarsana, E. T. Baskoro, S. Uttunggadewa, and D. Ismaimuza, Expansion techniques on the super edge antimagic total graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 189-199. https://combinatorialpress.com/article/jcmcc/Volume%20071/vol-071-paper%2012.pdf

[3709] I. W. Sudarsana, A. Hendra, Adiwijaya, D. Y. Setyawan, On super edge anti magic total labeling for t-joint copies of wheel, *Far East J. Math. Sci.,* (FJMS) **69**(2) (2012) 275-283. https://www.researchgate.net/publication/267466292_On_super_edge_anti_magic_total_labeling_for_t-joint_copies_of_wheel

[3710] I. W. Sudarsana, D. Ismaimuza, E. Baskova, and H. Assiyatun, On super (a,d)-antimagic total labeling of disconnected graphs, *J. Combin. Math. Combin. Comput.,* **55** (2005) 149-158. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2014.pdf

[3711] I. W. Sudarsana, S. A. Suryanto, D. Lusianti, and N. Putri, An application of super mean and magic graphs labeling on cryptography system, *J. Physics Confer. Ser.* 1763(1):012052 (2020) https://www.researchgate.net/publication/348996477_An_application_of_super_mean_and_magic_graphs_labeling_on_cryptography_system/references#fullTextFileContent

[3712] G. Sudarvizhi and K. Balasangu, One modulo three mean labeling of graphs, *J. Phys.: Conf. Ser.,* (2021) 1770 012077, 1-12. https://iopscience.iop.org/article/10.1088/1742-6596/1770/1/012077

[3713] S. Sudha, Some chain graphs are graceful, *Math. Sci. Internat. Research J.,* **1**(3) (2012) 855-859.

[3714] S. Sudha, Gracefulness of joining complete bipartite graphs having a common vertex set, *Engin. Sci. Internat. Research J.,* **1**(1) (2013) 84-85.

[3715] M. Sudha and A. C. Babu, Even-even gracefulness of some families of graphs, *IOSR J. Math.,* (IOSR-JM), **8** (6) (2013) 7-11. https://www.iosrjournals.org/iosr-jm/papers/Vol8-issue6/B0860711.pdf

[3716] S. Sudha and V. Kanniga, Arbitrary supersubdivision of helms, centipedes and ladder graphs are graceful, *Math, Sci. Internat. Research J.*, **1**(3)(2012) 860-863.

[3717] S. Sudha and V. Kanniga, Gracefulness of joining isolated vertices to a path, *Global J. Math. Math. Sciences*, **2**(1) (2012) 91-94.

[3718] S. Sudha and V. Kanniga, Gracefulness of some new class of graphs, *Engin. Sci. Internat. Reser. J.*, **1**(1) (2013) 81-83.

[3719] S. Sudha and V. Kanniga, Graceful labeling on the combination of some graphs, *Math. Sci. Internat. Research J.,* **2**(2) (2013) 630-633.

[3720] S. Sudha and V. Kanniga, Superposition of stars on cycles and n-centipedes are graceful, *Math. Sci. Internat. Research J.,* **2**(2) (2013) 634-636. https://www.xisdxjxsu.asia/V16I11-8.pdf

[3721] V. Sudhakar, S. Leen, P. Anuradha, R. Indhumathi, N. Avinash, and V. Balaji, Super mean graph labeling: A novel cryptographic framework using five-Star graphs with applications in secure communications and epidemic modeling. *J. Comput. Math.*, **10**(2) (2026) 016-047. https://www.shcpub.in/index.php/cm/article/view/488/277

[3722] S. Sudhakar, T. Ranjani, V. Swathy, and V. Balaji, GMJ coding using even felicicitous labeling and two star graphs, July 2021 Journal of Physics Conference Series 1964(2):022025 8 pages. doi:10.1088/1742-6596/1964/2/022025

[3723] S. Suganya, V. Maheswari, and S. R. T. Mary, Verifications of harmonic mean labeling in degree splitting graphs, *Internat. J. Sci. Res. Sci., Engin. Tech.*, **10**(5) (2023) 10-13. https://www.academia.edu/109026162/Verifications_of_Harmonic_Mean_labeling_in_Degree_Splitting_Graphs

[3724] P. Sugapriya and A. Aishwariya, Lucky-edge mean labeling of corona product of graphs, *J. Manage. and Entrepreneurship*, **16**(2) (2022) 87–104. https://www.researchgate.net/publication/363009880_LUCKY_EDGE_MEAN_LABELING_OF_CORONA_PRODUCT_OF_GRAPHS

[3725] K. A. Sugeng, Z. Z. Barack, N. Hinding, and R. Simanjuntak, Modular irregular labeling on double-star and friendship graphs, *J. Math.*, (2021) Article ID 4746609, 6 pages. https://www.hindawi.com/journals/jmath/2021/4746609/

[3726] K. A. Sugeng and N. H. Bong, Vertex (a, d)-antimagic total labeling on circulant graph $C_n(1, 2, 3)$, *J. Indones. Math. Soc.,* Special Edition (2011) 79-88. https://jims-a.org/index.php/jimsa/article/view/21/pdf_1

[3727] K. A. Sugeng, D. Fronček, M. Miller, J. Ryan, and J. Walker, On distance magic labelings of graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 39-48. https://combinatorialpress.com/article/jcmcc/Volume%20071/vol-071-paper%203.pdf

[3728] K. A. Sugeng, B. N. Herawati, M. Miller, and M. Bača, On magicness and antimagicness of the union of 4-regular circulant graphs, *Austral. J. Combin.*, **50**, (2011) 141-153. https://ajc.maths.uq.edu.au/pdf/50/ajc_v50_p141.pdf

[3729] K. A. Sugeng, P. John, M. L. Lawrence, L. F. Anwar, M. Bača, and A. Semaničová-Feňovčíková, Modular irregularity strength on some flower graphs, *Elec. J. Graph Th. App.*, **11**(1) (2023) 27-38. https://www.ejgta.org/index.php/ejgta/article/view/1734/pdf_288

[3730] K. A. Sugeng and M. Miller, Relationship between adjacency matrices and super (a, d)-edge-antimagic-total labelings of graphs, *J. Combin. Math. Combin. Comput.,* **55** (2005) 71-82. https://www.researchgate.net/publication/267078657_Relationship_between_adjacency_matrices_and_super_ad-edge-antimagic-total_labeling_of_graphs

[3731] K. A. Sugeng and M. Miller, On consecutive edge magic total labelings of graphs, *J. Discrete Algorithms*, **6** (2008) 59-65. https://www.sciencedirect.com/science/article/pii/S1570866706000748?via%3Dihub

[3732] K. A. Sugeng, M. Miller, M. Bača, Super edge-antimagic total labelings, *Util. Math.,* **71** (2006) 131-141. https://www.researchgate.net/publication/262723143_Super_edge-antimagic_total_labelings

[3733] K. A. Sugeng, M. Miller, Y. Lin, and M. Bača, Super (a, d)-vertex-antimagic total labelings, *J. Combin. Math. Combin. Comput.,* **55** (2005) 91-102. https://www.researchgate.net/publication/262722984_Super_ad-vertex-_antimagic_total_labelings

[3734] K. A. Sugeng, M. Miller, Slamin, and M. Bača, (a, d)-edge-antimagic total labelings of caterpillars, *Lecture Notes Comput. Sci.,* **3330** (2005) 169-180. https://link.springer.com/content/pdf/10.1007/978-3-540-30540-8_19.pdf

[3735] K. A. Sugeng, M. Miller, Y. Lin, and M. Bača, Face antimagic labelings of prisms, *Util. Math.,* **71** (2006) 269-286. https://www.researchgate.net/publication/262722998_Face_antimagic_labelings_of_prisms

[3736] K. A. Sugeng and J. Ryan, On several classes of monographs, *Australas. J. Combin.,* **37** (2007) 277-284. https://www.researchgate.net/publication/266961369_On_several_classes_of_monographs

[3737] K. A. Sugeng, J. Ryan, and H. Fernau, A sum labelling for the flower $f_{p,q}$, unpublished. https://www.academia.edu/18724308/A_Sum_Labelling_for_the_Flower_fq_p

[3738] K. A. Sugeng and D. R. Silaban, Super (a, d)-vertex antimagic total labeling on disjoint union of regular graphs, *J. Combin. Math. Combin. Comput.,* **71** (2009) 217-225. https://www.researchgate.net/publication/266037854_Super_ad-vertex_antimagic_total_labeling__on_a_disjoint_union_of_regular_graphs

[3739] K. A. Sugeng, F. C. Sofyan, S. Sy, N. Hinding, and R. Simanjuntak, *Elect. J. of Graph Theory Appl.*, **14**(1) (2026) 47-70. doi:10.5614/ejgta.2026.14.1.4

[3740] K. A. Sugeng, Surip, and Rismayati, On odd harmonious labeling of m-shadow of cycle, gear with pendant, and Shuriken graphs, https://aip.scitation.org/doi/pdf/10.1063/1.5139141

[3741] K. A. Sugeng and W. Xie, Construction of super edge magic total graphs, in: *Proceed. Sixteenth Australasian Workshop oCombin. Algor. 2005*, Ballarat, Australia, (2005) 303-310/ https://core.ac.uk/download/pdf/213011717.pdf

[3742] M. Sugiyama, Generalization of magic graphs and their properties, *Inform. Process. Soc. Japan (IPSJ)*, **59**(6) (2017) 1394-1404. NOW Chat 10/12.26

[3743] A. Sugumaran and V. Mohan, Further results on prime cordial labeling, *Annals Pure and Appl. Math.*,, **14**(3) (2017) 489-496. http://www.researchmathsci.org/APAMart/apam-v14n3-16.pdf

[3744] A. Sugumaran and V. Mohan, Prime cordial labeling of some graphs related to $H-$graph, *Annals Pure and Appl. Math.*, **16**(1) (2018) 171-176. http://researchmathsci.org/APAMart/apam-v16n1-18.pdf

[3745] A. Sugumaran and V. Mohan, Difference cordial labeling of some special graphs and related to fan graphs, *Internat. J. Res. Engin. Appl. Management*, **4**(12) (2019) 57-60. http://ijream.org/papers/IJREAMV04I1248014.pdf

[3746] A. Sugumaran and K. Rajesh, Extended results on sum divisor cordial labeling, *Proyecc. J. Math.,* **38**(4) (2019) 653-663. https://www.revistaproyecciones.cl/index.php/proyecciones/article/view/3779/3239

[3747] A. Sugumaran and K. Suresh, Some results on divisor cordial labeling, *Internat. J. Applied Engin. Res,* **10**(72) (2015) 68-71.

[3748] A. Sugumaran and K. Suresh, Some results on vertex odd divisor cordial labeling of graphs, *Internat. J. Pure Appl. Math.*, **113**(11) (2017) 199-207. https://acadpubl.eu/jsi/2017-113-pp/articles/11/22.pdf

[3749] A. Sugumaran and K. Suresh, A special class of prime cordial graphs, *Internat. J. Pure Appl. Math.*,, **114**(6) (2017) 41-48. https://acadpubl.eu/jsi/2017-114-5/articles/2/5.pdf

[3750] A. Sugumaran and K. Suresh, Further results on vertex odd divisor cordial labeling of some graphs, *Annals Pure Appl. Math.*, **16**(2) (2018) 365-372. http://www.researchmathsci.org/APAMart/apam-v16n2-13.pdf

[3751] A. Sugumaran and K. Suresh, Some star related vertex odd divisor cordial labeling of graphs, *Annals Pure Appl. Math.*, **16**(2) (2018) 385-391. http://www.researchmathsci.org/APAMart/apam-v16n2-15.pdf

[3752] A. Sugumaran and K. Suresh, Cycle related vertex odd divisor cordial labeling of graphs, *J. Comput. Math. Sci.*, **9**(3) (2018) 204-211. http://www.compmath-journal.org/dnload/A-Sugumaran-and-K-Suresh-/CMJV09I03P0204.pdf

[3753] A. Sugumaran and P. Vishnu Prakash, Some new results of prime cordial labeling, *Internat. J. Pure Engin. Math., (IJPEM)*, **5**(1) (2017) 9-14. http://www.ascent-journals.com/IJPEM/Vol5No1/2-vishnuprakash.pdf

[3754] A. Sugumaran and P. Vishnu Prakash, Prime cordial labeling for theta graph, *Annals Pure Appl. Math.*, **14**(3) (2017) 379-386. https://pdfs.semanticscholar.org/38c3/54cd5af11aead326d7c7de6193719995481b.pdf

[3755] A. Sugumaran and P. Vishnu Prakash, Some new results on prime cordial labeling for theta graph, *J. Comput. Math. Sci.,* **8**(11) (2017) 630-634.

[3756] A. Sugumaran, and P. Vishnu Prakash, Prime cordial labeling of theta graphs, *Annals Pure Appl. Math.*, **14**(3) (2017) 379–386. http://www.researchmathsci.org/APAMart/apam-v14n3-4.pdf

[3757] A. Sugumaran and P. Vishnu Prakash, Some new families of 3-equitable prime cordial graphs, *Internat. J. Stat. Appl. Math.*, **3**(1) (2018) 45-49. http://www.mathsjournal.com/pdf/2018/vol3issue1/PartA/2-6-12-620.pdf

[3758] A. Sugumaran and P. Vishnu Prakash, E-cordial labeling for theta graph, *Research Reviews: Discrete Math. Structures*, **5**(1)(2018) 26-32.

[3759] A. Sugumaran and P. Vishnu Prakash, 3-equitable prime cordial labeling in theta graph, *J. Emerging Tech. Innovative Res.,* **5**(7) (2018) 332-338. https://www.jetir.org/papers/JETIR1807205.pdf

[3760] A. Sugumaran and K. Rajesh, Some new results on sum divisor cordial graphs, *Annals Pure Appl. Math.,* **14**(1) (2017) 45-52. https://pdfs.semanticscholar.org/eee0/93c3f3bb98122a7551b08f1e1a26db538370.pdf

[3761] A. Sugumaran and K. Rajesh, Sum divisor cordial labeling of theta graph, *Annals Pure Appl. Math.,* **14**(2) (2017) 313-320. https://pdfs.semanticscholar.org/a540/180f1aeaf801a1e9ba53e817dfade5afcfc1.pdf

[3762] A. Sugumaran and K. Rajesh, Sum divisor cordial labeling of Herschel graph, *Annals Pure Appl. Math.,* **14**(3) (2017) 465-472. https://pdfs.semanticscholar.org/6327/4d4d2ff4c6239308ac67cba007846e1688be.pdf

[3763] A. Sugumaran and K. Rajesh, Further results on sum divisor cordial labeling, *J. Comput. Math. Sci.*, **8**(12) (2017) 802-807. http://www.compmath-journal.org/dnload/A-Sugumaran-and-K-Rajesh/CMJV08I12P0802.pdf

[3764] A. Sugumaran and K. Rajesh, Some graph operations on sum divisor cordial labeling related to H-graph, *Internat. J. Stat. Appl. Math.*, **3**(1) (2018) 389-395. http://www.mathsjournal.com/archives/2018/vol3/issue1/PartE/3-1-15

[3765] A. Sugumaran, K. Rajesh, Sum divisor cordial labeling on some special graphs, *Internat. J. Math. Appl.*, **6**(2-B) (2018) 347-353. https://pdfs.semanticscholar.org/2105/7352842a3fb5d129c4e140c93775c65734a7.pdf

[3766] A. Sugumaran and K. Rajesh, Perrin graceful graphs, *Internat. J. Pure Appl. Math.*, **114**(6) (2017) 131-137. https://acadpubl.eu/jsi/2017-114-5/articles/2/14.pdf

[3767] A. Sugumaran and K. Rajesh, Sum divisor cordial labeling on some classes of graphs, *J. Comput. Math. Sci.*, **10**(2) (2019) 316-325. http://www.compmath-journal.org/dnload/A-Sugumaran-and-K-Rajesh/CMJV10I02P0316.pdf

[3768] A. Sugumaran and K. Rajesh, Lucas divisor cordial labeling, *Internat. J. Pure Appl. Math.*, **113**(6) (2017) 233-241. https://acadpubl.eu/jsi/2017-114-5/articles/2/14.pdf

[3769] J. E. Sulayman and A. C. Pedrano, On fibonacci cordial labeling of some snake graphs, *European J. Math. Stat.*, **4**(2) (2023) 29–35. https://eu-opensci.org/index.php/ejmath/article/view/14193/3204

[3770] L. H. Sullystiawati, M. Marsidi, E. D. Putra, and I. H. Agustin, The reflexive H-strength on some graphs, *CAUCHY J. Matematika Murni dan Aplikasi,* **9**(1) (2024) 36-49. DOI:10.18860/ca.v9i1.23172

[3771] P. Sumathi and P. Mala, (S, D) magic labeling of subdivision of some special trees, *Inter. J. Math. Comput. Sci.,* **12**(12) (2024) 4662-4668. https://ijmcr.in/index.php/ijmcr/article/view/845/668

[3772] P. Sumathi and P. Mala, (s, d) magic labeling in some trees, *Math. Stat. Eng. Appl.* , **72**(1) (2023) 1895-1904. https://www.philstat.org/index.php/MSEA/article/view/2608/2072

[3773] P. Sumathi and G. G. Raman, Arithmetic sequential graceful labeling on star related graphs, *Indian J. Sci. Tech.,* **15**(44) (2022) 2356-2362. https://indjst.org/articles/arithmetic-sequential-graceful-labeling-on-star-relate

[3774] P. Sumathi and A. Rathi Some more results on quotient labeling of special graphs, *Internat. J. Pure Appl. Math.*, **115**(9) (2017) 259-268. https://acadpubl.eu/jsi/2017-115-9/articles/9/22.pdf

[3775] P. Sumathi and A. Rathi, Quotient labeling of some ladder graphs, *American Engin. Res.*, (AJER), *7*(12) 38-42. https://ajer.org/papers/Vol-7-issue-12/E07123842.pdf

[3776] P. Sumathi and A. Rathi, Quotient labeling of snake related graphs, *IOSR J. Math.,* **14**(6) Ver. II (2018) 26-33. https://www.iosrjournals.org/iosr-jm/papers/Vol14-issue6/Version-2/D1406022633.pdf

[3777] P. Sumathi and J. Suresh Kumar, Fuzzy quotient-3 cordial labeling of star related graphs- Paper I, *Malaya J. Matematik*, **S**(1) (2019) 79-82. https://doi.org/10.26637/MJM0S01/0020

[3778] L. Sun, Radio number for middle graph of dandelion, *J. Discrete Math. Sci. Cryptogr,* **24**(4) (2021) 865-877.

[3779] R. G. Sun, Harmonious and sequential labelings of the book graphs B_m, *Gaoxiao Yingyong Shuxue Xuebao Ser. A*, **9** (1994) 335-337.

[3780] G.-Y. Sun, Z.-B. Gao, and S.-M. Lee, On full friendly index sets of twisted product of Möbius ladders, *Ars Combin.*, 128 (2016) 225–239. https://combinatorialpress.com/ars-articles/volume-128-ars-articles/on-full-friendly-index-sets-of-twisted-product-of-mobius-ladders/?utm_source=chatgpt.com

[3781] G. C. Sun, J. Guan, and S.-M. Lee, A labeling algorithm for magic graph, Proceedings of the Twenty-fifth Southeastern International Conference on Combinatorics, Graph Theory and Computing (Boca Raton, FL, 1994). *Congr. Numer.* **102** (1994) 129-137.

[3782] G. C. Sun and S. M. Lee, Construction of magic graphs, *Congr. Numer.*, **103** (1994) 243-251.

[3783] D. Q. Sun and D. Y. Wang, Even sequential harmonious graphs, personal communication (2015).

[3784] H. Sun, X. Zhang, and B. Yao, Construction of new graphical passwords with graceful-type labellings on trees, *Conference: 2018 2nd IEEE Adv. Infor. Management, Communicates, Elect. Automation Control Conf. (IMCEC)* https://www.researchgate.net/publication/328166886_Construction_of_New_Graphical_Passwords_with_Graceful-Type_Labellings_on_Trees

[3785] S. B. Sundar, R. Jeyabalan, R. Nishanthini, P. Swathi, and G. Rajchakit, Innovative perspectives on antimagic labeling in graphs, *Internat. J. Math. Comput. Sci.*, **20**(2) (2025) 591–602. https://www.researchgate.net/publication/389412370_Innovative_Perspectives_on_Antimagic_Labeling_in_Graphs

[3786] M. Sundaram, R. Ponraj and S. Somasundaram, Product cordial labeling of graphs, *Bull. Pure and Appl. Sci. (Math. & Stat.)*, **23E** (2004) 155-163. https://www.researchgate.net/publication/303486909_Product_cordial_labeling_of_graphs

[3787] M. Sundaram, R. Ponraj, and S. Somasundram, Prime cordial labeling of graphs, *J. Indian Acad. Math.*, **27** (2005) 373-390.

[3788] M. Sundaram, R. Ponraj, and S. Somasundram, Some results on total product cordial labeling of graphs, *J. Indian Acad. Math.*, **28** (2006) 309-320.

[3789] M. Sundaram, R. Ponraj, and S. Somasundaram, On a prime labeling conjecture, *Ars Combin.,* **80** (2006) 205-209. https://combinatorialpress.com/article/ars/Volume%20079/volume-79-paper-17.pdf

[3790] M. Sundaram, R. Ponraj, and S. Somasundram, Total product cordial labeling of graphs, *Bull. Pure Appl. Sci. Sect. E Math. Stat.,* **25** (2006) 199-203.

[3791] M. Sundaram, R. Ponraj and S. Somasundaram, On prime graphs of order ≤ 6, *Acta Ciencia Indica,* **32** (2006) 859-871.

[3792] M. Sundaram, R. Ponraj and S. Somasundaram, Mean number of a graph, *Pure Appl. Math. Sci.,* **57** (2007) 93-101.

[3793] M. Sundaram, R. Ponraj and S. Somasundaram, EP-cordial labeling of graphs, *Varahmihir J. Math. Sci.*, **7** (2007) 183-194. https://www.researchgate.net/publication/264966619_EP-cordial_labelings_of_graphs

[3794] M. Sundaram, R. Ponraj and S. Somasundaram, On graph labeling parameter, *J. Discrete Math. Sci. Cryp.,* **11** (2008) 219-229. https://www.connectjournals.com/file_full_text/54202H_jdmsc231_219-229.pdf

[3795] M. Sundaram, R. Ponraj, and S. Somasundaram, Some results on product cordial labeling, unpublished

[3796] K. Sunitha, Radio labeling of double triangular snake graphs, *Ann. Pure Appl. Math.,* **27**(2) (2023) 63-71. http://www.researchmathsci.org/apamart/APAM-v27n2-4.pdf

[3797] K. Sunitha, C. D. Raj, and A. Subramanian, Radio mean labeling of path and cycle related graphs, *Global J. Math. Sci.: Theory and Practical*, **9**(3) (2017) 337-345. https://www.ripublication.com/irph/gjms17/gjmsv9n3_10.pdf

[3798] K. Sunitha and T. Revathi, Vertex edge closed neighborhood (VECN) prime labeling of some m-fold snake related graphs, *Adv. Appl. Math. Sci.,* **21**(11) (2022) 6159-6165. https://www.researchgate.net/publication/375289679_VERTEX_EDGE_CLOSED_NEIGHBORHOOD_VECN_PRIME_LABELING_OF_SOME_m-FOLD_SNAKE_RELATED_GRAPHS

[3799] K. Sunitha and M. Sheriba, Gaussian tribonacci r-graceful labeling of some tree related graphs, *Ratio Mathematica*,**44** (2022) 188-196. https://eiris.it/ojs/index.php/ratiomathematica/article/view/906

[3800] K. Sunitha and K. Vimal Rani, Radio heronian mean labeling of some corona related graphs, *Malaya J. Matematik*, **09**(01) (2021) 1212-1215. https://www.researchgate.net/publication/375289606_Radio_Heronian_mean_labeling_of_some_Corona_related_graphs

[3801] K. Sunitha and K. Vimal Rani, Radio heronian mean k-graceful labeling on degree splitting of graphs, *Ratio Math.,* **45** (2023) 32-44. https://www.researchgate.net/publication/375289801_RADIO_HERONIAN_MEAN_GRACEFUL_LABELING_ON_DEGREE_SPLITTING_OF_GRAPHS

[3802] B. S. Sunoj and T. K. Mathew Varkey, ADCSS-Labeling for some middle graphs, *Annals Pure Applied Math.*, **12**(2) (2016) 161-167. http://www.researchmathsci.org/apamart/apam-v12n2-7.pdf

[3803] B. S. Sunoj and T. K. Mathew Varkey, A note on an absolute difference of cubic and square sum labeling, *Internat. J. Math. Sci. Engg. Appls.,* (IJMSEA) **11**(I) (2017) 1-10. http://www.ascent-journals.com/IJMSEA/Vol11No1/1-sunoj.pdf

[3804] B. S. Sunoj and T. K. Mathew Varkey, Square difference prime labeling for some snake graphs, *Global J. Pure Appl. Math.,* **13**3 (2017) 1083-1089. https://www.ripublication.com/gjpam17/gjpamv13n3_17.pdf

[3805] B. S. Sunoj and T. K. Mathew Varkey, Square difference prime labeling for some tree graphs, *Internat. J. Eng. Development Res.,* **5**(4) (2017 875-878. https://www.rjwave.org/ijedr/papers/IJEDR1704142.pdf

[3806] B. S. Sunoj and T. K. Mathew Varkey, Square difference prime labeling for some cycle related graphs, *Internat. J. Comput. Eng. Res.,* **7**(11) (2017) 22-25. https://mail.ijceronline.com/papers/Vol7_issue11/Version-2/D0711022225.pdf

[3807] B. S. Sunoj and T. K. Mathew Varkey, Square difference prime labeling of some star related graphs, *Internat. J. Math Trends Tech.*, **52**(3) (2017)

[3808] I. N. Suparta and I Dewa M. Agus Ariawan, Expanding graceful trees, *Electron. J. Graph Theory Appl,* (EJGTA) **8**(2) (2020) 217-232. https://www.ejgta.org/index.php/ejgta/article/view/936/pdf_137

[3809] N. Suparta and D. M. A. Ariawan, Some methods for constructing some classes of graceful uniform trees, *Indonesian J. Combin.,* **2**(2) (2018) 123-135. http://dx.doi.org/10.19184/ijc.2018.2.2.7

[3810] I. N. Suparta and I. P. Suharta, A note on edge irregularity strength of some graphs, *Indonesian J. Combin.*, **4**(1) (2020) 10-20. http://dx.doi.org/10.19184/ijc.2020.4.1.2

[3811] M. C. Superdock, The Graceful Tree Conjecture: A Class of Graceful Diameter-6 Trees, Senior Thesis, Department of Mathematics of Princeton University, 2013

[3812] Y. Susanti and M. A. Haq, On total edge irregularity strengths of odd and even staircase graphs, *Palestine J. Math.*, **13**(4)(2024) 1-18. https://www.preprints.org/frontend/manuscript/5985fc3deea52ae33dbcea3905684075/download_pub

[3813] Y. Susanti, H. Khotimah, M. Hidayati, and M. F. Wahyujati, On total edge irregularity strength of some graphs and the construction of some labeling, *Far East J. Math. Sci. (FJMS)*, **104**(1) (2018) 49-64. https://www.pphmj.com/abstract/11801.htm

[3814] Y. Susanti, I. Ernanto1, and B. Surodjo, On some new edge odd graceful graphs, *Proceed. 8th SEAMS-UGM Internat. Conf. Math. Appl. 2019* AIP Conf. Proc. 2192, 040016-1–040016-12. https://aip.scitation.org/doi/pdf/10.1063/1.5139142?class=pdf

[3815] Y. Susanti, Y. I. Puspitasari and H. Khotimah, On total edge irregularity strength of staircase graphs and related graphs, *Iran. J. Math. Sci. Inform.,* **15**(1) (2020) 1-13. http://ijmsi.ir/article-1-1121-en.html

[3816] Y. Susanti, S. Wahyuni, A. Sutjijana, S. Sutopo, and I. Ernanto, Generalized arithmetic staircase graphs and their total edge irregularity strengths, *Symmetry* **14**(9) (2022) :1853. https://www.researchgate.net/publication/363400767_Generalized_Arithmetic_Staircase_Graphs_and_Their_Total_Edge_Irregularity_Strengths

[3817] F. Susanto, On super (a, d)-C_n-antimagic total labeling of disjoint union of cycles, *AKCE Int. J. Graphs Comb.,* **15**(1) (2018) 22-26. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2018.01.017?scroll=top&needAccess=true

[3818] F. Susanto, On H-irregularity strengths of square chain graphs, *J. Ilmiah Matematika dan Terapan*, **23**(1) (2026) 1-10. https://bestjournal.untad.ac.id/index.php/JIMT/article/view/17989/12229?utm_source=chatgpt.com

[3819] F. Susanto, C. N. Betistiyan, I. Halikin, and K. Wijaya, On inclusive distance vertex irregularity strength of small identical copies of star graphs, *Journal of Physics: Conference Series* **1872** (1) (2021), 012005. https://www.researchgate.net/publication/351583634_On_inclusive_distance_vertex_irregularity_strength_of_small_identical_copies_of_star_graphs

[3820] F. Susanto, R. Simanjuntak, and E. T. Baskoro, Counterexamples to the total vertex irregularity strength's conjectures, *Discrete Math. Lett,* **12** (2023) 159-165. https://www.dmlett.com/archive/v12/DML23_v12_pp159-165.pdf

[3821] F. Susanto, R. Simanjuntak, and E. T. Baskoro, D-irregularity strength of a graph, *Util. Math.,* **121** (2024) 69-90. https://combinatorialpress.com/article/um/Volume%20121/d-irregularity-strength-of-a-graph.pdf

[3822] F. Susanto, R. Simanjuntak, and E. T. Baskoro, Further results on the total vertex irregularity strength of trees, *Elect. Graph Th. Appl.,* **13**(1) (2025) 123–140. https://www.ejgta.org/index.php/ejgta/article/view/2151

[3823] M. Ramalakshmi and K. M. Kathiresan, Total absolute difference edge irregularity strength of the graphs, *Kragujevac J. Math.*, **46**(6) (2022) 895–903. https://imi.pmf.kg.ac.rs/kjm/pub/kjom466/kjm_46_6-5.pdf

[3824] F. Susanto and K. Wijaya, Total absolute difference edge irregularity strength of several ladder-type graphs, *Euler : J. Ilmiah Matematika, Sains dan Teknologi*, **14**(2) (2026) 457–466.

[3825] F. Susanto, K. Wijaya, P. M. Purnama, and Slamin, On distance irregular labelings of disconnected graphs, *Kragujevac J. Math.,* **46**(4) (2020) 507-523. researchgate.net/publication/340593151_ON_DISTANCE_IRREGULAR_LABELING_OF_DISCONNECTED_GRAPHS

[3826] F. Susanto, K. Wijaya, and Slamin, A. Semaničová-Feňovčíková, Distance irregularity strength of graphs with pendant vertices, *Opuscula Math.* **42** (3) (2022), 439-458. https://www.researchgate.net/publication/360259244_Distance_irregularity_strength_of_graphs_with_pendant_vertices

[3827] F. Susanto, K. Wijaya, I. W. Sudarsana, and Slamin, Non-inclusive and inclusive distance irregularity strength for the join product of graphs, *Electron. J. Graph Theory Appl. (EJGTA)* **10** (1) (2022), 1-13. https://www.researchgate.net/publication/360299847_Non-inclusive_and_inclusive_distance_irregularity_strength_for_the_join_product_of_graphs

[3828] S. Susilawati, E. T. Baskoro, and R. Simanjuntak, Total vertex irregularity strength of trees with maximum degree five, *Elect. J. Graph Th. Appl.* **6** (2) (2018) 250-257. https://www.ejgta.org/index.php/ejgta/article/view/415/pdf_80

[3829] Susilawati, E. T. Baskaro, and R. Simanjuntak, On the vertex irregular clabeling for subdivision for trees, *Australas J. Comb.*, **71**(2) (2018) 293-302. https://ajc.maths.uq.edu.au/pdf/71/ajc_v71_p293.pdf

[3830] Susilawati, E. T. Baskoro and R. Simanjuntak, Total vertex irregularity labelings for subdivision of several classes of trees, *Procedia Comput. Sci.*, **74** (2015) 112-117. https://www.researchgate.net/publication/290021906_Total_Vertex-Irregularity_Labelings_for_Subdivision_of_Several_Classes_of_Trees

[3831] Susilawati, E. T. Baskoro and R. Simanjuntak, Total vertex irregularity strength of trees with maximum degree four, *AIP Conf. Proc.*, **1707** (2016) 1-7. https://www.researchgate.net/publication/283210408_Total_Vertex_Irregularity_Strength_of_Trees_with_Maximum_Degree_Four

[3832] Susilawati, E. T. Baskoro and R. Simanjuntak, Total vertex irregularity strength of trees with maximum degree five, *Elec. J. Graph Th. Appl.,* **6**(2) (2018) 250–257. https://www.researchgate.net/publication/328285434_Total_vertex_irregularity_strength_of_trees_with_maximum_degree_five

[3833] Susilawati, E. T. Baskoro and R. Simanjuntak, On the total vertex irregularity strength for trees with many vertices of degree 2, *Elec. J. Graph Th. Appl.*, **8**(2) (2020) 415–421. https://www.ejgta.org/index.php/ejgta/article/view/1058

[3834] L. Susilowati, T. Sania, and N. Estuningsih, Super (a, d)-C_n-antimagic total labeling of ladder graph, *Adv. Appl. Discrete Math.,* **10**(2) (2012) 77-93. https://www.researchgate.net/publication/268611699_Super_ad-C_n_-antimagic_total_labeling_of_ladder_graph

[3835] G. Suthakaran and R. Jeyabalan, Parity weighted totally antimagic total graphs, *Math. Stat.,* **13**(1) (2025) 62-67. https://www.hrpub.org/download/20250228/MS8-13440075.pdf

[3836] M. Sutton, Sumable Graphs Labellings and Their Applications, Ph. D. Thesis, Dept. Computer Science, The University of Newcastle, 2001.

[3837] M. Sutton, A. Draganova, and M. Miller, Mod sum numbers of wheels, *Ars Combin.*, **63** (2002) 273-287. https://combinatorialpress.com/article/ars/Volume%20063/volume-63-paper-22.pdf

[3838] M. Sutton and M. Miller, Mod sum graph labelling of $H_{n,n}$ and K_n, *Australas. J. Combin.,* **20** (1999) 233-240. https://ajc.maths.uq.edu.au/pdf/20/ocr-ajc-v20-p233.pdf

[3839] M. Sutton and M. Miller, On the sum number of wheels, *Discrete Math.,* **232** (2001) 185-188. https://www.sciencedirect.com/science/article/pii/S0012365X00003472?via%3Dihub

[3840] M. Sutton, M. Miller, J. Ryan, and Slamin, Connected graphs which are not mod sum graphs, *Discrete Math.,* **195** (1999) 287-293. https://www.sciencedirect.com/science/article/pii/S0012365X98001848?via%3Dihub

[3841] V. Swaminathan and P. Jeyanthi, Super vertex-magic labeling, *Indian J. Pure Appl. Math.,* **34**(6) (2003) 935-939. https://www.researchgate.net/publication/257305473_3_VSwaminathan_and_PJeyanthi_Super_Vertex-Magic_Labeling_Indian_Journal_of_Pure_and_Applied_Mathematics_34_6_2003

[3842] V. Swaminathan and P. Jeyanthi, On super vertex-magic labeling, *J. Discrete Math. Sci. & Cryptography,* **8** (2005) 217-224.

[3843] V. Swaminathan and P. Jeyanthi, Super edge-magic strength of fire crackers, banana trees and unicyclic graphs, *Discrete Math.,* **306** (2006) 1624-1636. https://reader.elsevier.com/reader/sd/pii/S0012365X06002172?token=A64AFC9690FA9E459EEA4506C7FD4BC3D30B4D2FEF4FA06418CB9802AEE5C6BB3AD0A20D863D001271originRegion=us-east-1&originCreation=20230311182022

[3844] V. Swaminathan and P. Jeyanthi, Super edge-magic strength of generalized theta graph, *Inter. J. Management and Systems,* **22** (2006) 203-220. https://www.researchgate.net/publication/257364394_5_VSwaminathan_and_PJeyanthi_Super_Edge-Magic_Strength_of_Generalised_Theta_Graphs_International_Journal_of_Management_and_SystemsVol22_No3_2006_203_-_220

[3845] V. Swaminathan and P. Jeyanthi, Super edge-magic strength of some trees, *Util. Math.,* **72** (2007) 199-210.

[3846] V. Swaminathan and P. Jeyanthi, (a, d)-1-vertex-antimagic vertex labeling, *Util. Math.*, **74** (2007) 179-186.

[3847] V. Swaminathan and P. Jeyanthi, Super edge-magic labeling of some new classes of graphs, *Math. Education*, **XLII**(2) (2008) 91-94. https://www.researchgate.net/publication/267677152_Super_edge-magic_labeling_of_some_new_classes_of_graphs

[3848] V. Swaminathan and P. Jeyanthi, Strong super edge-magic graphs, *Math. Education*, **XLII**(3) (2008) 156-160. https://www.researchgate.net/publication/257364590_6_VSwaminathan_and_PJeyanthi_Strong_Super_Edge-magic_Graphs_The_Mathematics_Education_VolXLIINo3_2008156-160

[3849] P. Swathi, G. Kumar, R. Jeyabalan, and R. Nishanthini, Perfectly antimagic total graphs, *J. Intell. Fuzzy Sys.,* **44**(1) (2023) 1517-1523.

[3850] P. Swathi and R. Jeyabalan, Perfectly antimagic total labelings of star-like trees and the corona product of two graphs, *Acta Math. Univ. Comenianae,* **XCIII**(2) (2024) 95-100. https://www.iam.fmph.uniba.sk/amuc/ojs/index.php/amuc/article/view/2099/1018

[3851] R. Sweetly and J. P. Joseph, The radio number of $(W_n : 2)$ graphs, *J. Discrete Math. Sci. Cryptogr.,* **12** (2009) 729-736.

[3852] B. Swita, R. Br Butar-Butar, D. C. Pramesti, N. Febriyan, D. Puspita, N. S. Saftri, L. Azhari, and M. Simanihuruk, A partial solution to an open problem on super edge-magic total labelings of (4, 3)-cycle books graphs, *Wiley Internat. J. Math. Math. Sci.*, **2025** Article ID 5582258, 9 pages. https://doi.org/10.1155/ijmm/5582258

[3853] B. Swita, U. Rafflesia, N. Henni Ms, D. S. Adji, and M. Simanihuruk, On edge magic total labeling of $(7, 3)$-cycle books, *Int. J. Math. Math. Sci.,* (2019) Art. ID 1801925, 8 pp. https://doi.org/10.1155/2019/1801925

[3854] Z. Szaniszló, k-equitable labellings of cycles and some other graphs, *Ars Combin.,* **37** (1994) 49-63. https://combinatorialpress.com/article/ars/Volume%20037/volume-37-paper-4.pdf

[3855] A. Tabraiz and M. Hussain, Magic and anti-magic total labeling on subdivision of grid graphs, *J. Graph Labeling,* **2**(1) (2016) 9-24.

[3856] Y. B. Tai, G. L. Chia, and P.-H. Ong, Multi-bridge graphs are anti-magic, *Electron. J. Graph Theory Appl.,* (EJGTA) **10**(1) (2022) 311-318. https://www.ejgta.org/index.php/ejgta/article/view/1369/pdf_222

[3857] A. Taimur, G. Ali, M. Numan, A. Aslam, and K. Anoh Yannick, Super H-antimagic total covering for generalized antiprism and toroidal octagonal map, *J. Math.,* (2021) Art. ID 9680137, 8 pp. https://www.hindawi.com/journals/jmath/2021/9680137

[3858] A. Taimur, M. Numan, A. Mumtaz, and A. Semaničová-Feňovčíková, Super (a,d) – H-antimagic labeling of subdivided graphs, *Open Math.,* **16** (2018) 688-697. https://www.researchgate.net/publication/326003389_Super_a_d-H-antimagic_labeling_of_subdivided_graphs

[3859] Y. Takahasi, F. A. Muntaner-Batle, and R. Ichishima, Some new results concerning the valences of (super) edge-magic graphs, *Bull. Math. Soc. Sci. Math. Roumanie,* **68** (116), No. 2, (2025) 185–200. https://ssmr.ro/bulletin/pdf/68-2/articol_6.pdf

[3860] Y. Takahashi, R. Ichishima, and F. A. Muntaner-Batle, On the strength and domination number of graphs, *Contrib. Math.,* **8** (2023) 11-15. https://shahindp.com/Contrib_Math/v8/CM23_v8_pp11-15.pdf

[3861] M. Tamilselvi, and K. Akilandeswari, k-Super Heronian mean labeling of some graphs, *J. Comput. Math. Sci.*, **9**(6) (2018) 657-664. https://www.ijser.in/archives/v5i9/IJSER171861.pdf

[3862] M. Tamilselvi, K. Akilandeswari, and V. Suguna, k-super mean labeling of some graphs, *Internat. J. Sci. and Res.,* (IJSR), **5**(6) 2016. https://bit.ly/2UEeMzu

[3863] M. Tamilselvi and N. Revathi, k–super harmonic mean labeling of some graphs, *Aryabhatta J. Math. Informatics,* **9**(1) (2017) 779-787. https://ijmr.net.in/current/2017/Jan-June,-2017/k3SYlZouzkxOAsd.pdf

[3864] D. Tanna, J. Ryan, and Semaničová-Feňovčíková, Edge irregular reflexive labeling of prisms and wheels, *Australasain J. Combin.*, **69**(3) (2017) 394-401. https://ajc.maths.uq.edu.au/pdf/69/ajc_v69_p394.pdf

[3865] R. Tao, On k-cordiality of cycles, crowns and wheels, *Systems Sci. Math. Sci.,* **11** (1998) 227-229. https://sysmath.cjoe.ac.cn/jssc/CN/Y1998/V11/I3?utm_source=chatgpt.com

[3866] I. Tarawneh, R. Hasni, and A. Ahmad, On the edge irregularity strength of corona product of graphs with paths, *Appl. Math. E-Notes,* **16** (2016) 80-87. https://www.academia.edu/36788611/On_The_Edge_Irregularity_Strength_Of_Corona_Product_Of_Graphs_With_Paths

[3867] I. Tarawneh, R. Hasni, and A. Ahmad, On the edge irregularity strength of corona product of cycle with isolated vertices, *AKCE Internat. J. Graphs Combin.,* **13** (2016) 213-217. https://www.researchgate.net/publication/305485534_

On_the_edge_irregularity_strength_of_corona_product_of_cycle_with_isolated_vertices

[3868] I. Tarawneh, R. Hasni, A. Ahmad, and M. A. Asim, On the edge irregularity strength for some classes of plane graphs, **6**(3) (2021) 2724-2731. http://www.aimspress.com/article/doi/10.3934/math.2021166

[3869] I. Tarawneh, R. Hasni, and A. Ahmad, On the edge irregularity strength of grid graphs, *AKCE Int. J. Graphs Comb.*, **17**(1) (2020) 414–418. https://doi.org/10.1016/j.akcej.2018.06.011

[3870] I. Tarawneh, R. Hasni, and M. A. Asim, On the edge irregularity strength of disjoint union of star graph and subdivison of star graph, *Ars Combin.,* **141** (2018) 93-100.

[3871] I. Tarawneh, R. Hasni, M. K. Siddiqui, and M. A. Asim, On the edge irregularity strength of disjoint union of graphs, *Ars Combin.,* **142** (2019) 239-249. https://www.researchgate.net/publication/328561737_On_the_edge_irregularity_strength_of_disjoint_union_of_graphs

[3872] M. Teffilia1 and J. Devaraj, On friendly index set of graphs, *Internat. J. Sci. Res. Math. Stat. Sci.*, **5**(4) (2018) 258-263. https://discovery.researcher.life/download/article/9454dd81f10c314295a46fa577feae70/full-text

[3873] H.-M. Teichert, The sum number of d-partite complete hypergraphs, *Discuss. Math. Graph Theory,* **19** (1999) 79-91. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1087

[3874] H.-M. Teichert, Classes of hypergraphs with sum number one, *Discuss. Math. Graph Theory,* **20** (2000) 93-103. https://eudml.org/doc/270220

[3875] H.-M. Teichert, Sum labellings of cycle hypergraphs, *Discuss. Math. Graph Theory*, **20** (2000) 255-265. https://eudml.org/doc/270630

[3876] S. Telang, private communication.

[3877] A. Tenguria and R. Verma, 3-Total edge sum cordial labeling for some graphs, *Internat. J. Comput. Appl.,* (0975 - 8887) **129**(8) (2015). https://www.ijcaonline.org/research/volume129/number8/tenguria-2015-ijca-906992.pdf

[3878] A. Tenguria and R. Verma. 3-Total super product cordial labeling for some graphs, *Internat. J. Science Res.,* **4**(2) (2015) 557-559.

[3879] A. Tenguria and R. Verma, 3-Total super sum cordial labeling for union of some graphs, *IJAIS*, Foundation of Computer Science, New York, USA, **8**(4) (2015) 25-30. https://scispace.com/pdf/3-total-super-sum-cordial-labeling-for-union-of-some-graphs-2jhnw1yxti.pdf

[3880] A. Tenguria and R. Verma, 3-total super sum cordial labeling by applying operations on some graphs, *Appl. Appl. Math.,* **11** (2016)(1) 444-446. https://www.pvamu.edu/mathematics/wp-content/uploads/sites/49/29_R827_Verma_AAM_.pdf

[3881] M. Tezer and I. Cahit, A note on (a, d)-vertex antimagic total labeling of paths and cycles, *Util. Math.,* **68** (2005) 217-221. https://www.researchgate.net/publication/269001467_A_note_on_a_d-Vertex_antimagic_total_labeling_of_paths_and_cycles

[3882] S. Thakare and M. Acharya, Exclusive sum labeling and exclusive sum number of prism graph (D_n), *Ars Combin.,* **154** (2021) 303-316.

[3883] A. P. Tharani and P. Saradha, Discrete labeling of graphs, *Indian J. Sci. a Tech.*, **17** Special Issue (2024) 93-102. https://indjst.org/articles/discrete-labeling-of-graphshttps://indjst.org/articles/discrete-labeling-of-graphs

[3884] T. Tharmaraj and P. B. Sarasija, Square difference labeling for certain graphs, *Internat. J. Math. Archive,* **4**(8) (2013) 183-186.

[3885] T. Tharmaraj and P. B. Sarasija, On beta combination labeling graphs, *Internat. J. Computer Applications,* **79**(13) (2013) 26-29. https://research.ijcaonline.org/volume79/number13/pxc3891807.pdf

[3886] T. Tharmaraj and P. B. Sarasija, Beta combination graphs, *Internat. J. Computer Applications,* **76**(14) (2013) 1-5. https://research.ijcaonline.org/volume76/number14/pxc3890589.pdf

[3887] K. G. Tharunraj and P. Ragukumar, Log-linear algorithm to generate prime trees. *J. Inform. Optim. Sci.*, (2016) 15 pp. https://www.researchgate.net/publication/404993414_C_Log-linear_algorithm_to_generate_prime_trees

[3888] K. G. Tharunraj and P. Ragukumar, Effective conversion of non-prime graphs to prime graphs, *Int. J. Anal. Appl.*, **23:254** (2025) 1-12. doi:DOI:10.28924/2291-8639-23-2025-254

[3889] K. Thejeshwi and S. Kirupa, Application of graceful graph in MPLS, *IJSRD - Internat. J. Sci. Res. Development,* **6**(06) (2018) 2321-0613. https://www.ijsrd.com/articles/IJSRDV6I60118.pdf

[3890] K. Thirugnanasambandam and G. Chitra, Fibonacci mean anti-magic labeling of graphs, *Inter. J. Res. Advent. Tech.*, **7**(4S) (2019) 7-10. http://www.ijrat.org/downloads/Vol-7/aprils-2019/PaperID-74S201905.pdf

[3891] K. Thirugnanasambandam, G. Chitra, and Y. Vishnupriya, Prime odd mean labeling of some special graphs, *J. Emerging Tech. Innov. Res.* (JETIR), **5**(6) (2018) 115-126. https://www.jetir.org/papers/JETIR1806795.pdf

[3892] K. Thirusangu and R. Bhrathiraja, Magic and antimagic labelings of Cayley digraphs over the groups $Z_{p^n} \oplus Z_{p^n}$ and $Z_m \oplus Z_n$, *Adv. Appl. Math. Sci.,* **20**(9) (2021) 1851-1864. https://www.mililink.com/upload/article/717844157aams_vol_209_july_2021_a19_p1851-1864_k._thirusangu_and_r._bharathiraja.pdf

[3893] K. Thirusangu and M. Madhu, Divisor cordial labeling in extended duplicate graph of star, bistar and double star, *JASC: J. Appl. Sci. Comput.,* **VI**(I) (2019) 583-596

[3894] K. Thirusangu, A. K. Nagar, and R. Rajeswari, Labelings in Cayley digraphs, *European J. Combin.,* **32** (2011) 133-139. https://www.sciencedirect.com/science/article/pii/S0195669810000946

[3895] K. Thirusangu and P. Nedumaran On 0-edge magic labeling in certain graphs, *Internat. J. Appl. Research,* **1**(12) (2015) 874-877. https://www.allresearchjournal.com/archives/2015/vol1issue12/PartM/1-11-70.pdf

[3896] K. Thirusangu and A. Selvaganapathy, Gaussian anti magic labeling in some graphs, *Internat. J. Math. Trends Tech.*, (2018) 81–84.

[3897] K. Thirusangu and A. Selvaganapathy, Gaussian anti magic labeling in sunlet and triangular snake graphs, *Malaya J. Mat.*, **S**(1) (2020) 148-152. https://www.malayajournal.org/articles/MJM0S200028.pdf

[3898] K. Thirusangu and A. Selvaganapathy, Gaussian anti magic labeling in some special graphs, *Malaya J. Mat.*, **S**(1) (2020) 231-235. https://www.malayajournal.org/articles/MJM0S200044.pdf

[3899] K. Thirusangu and A. Selvaganapathy, Gaussian anti magic labeling in some related graphs, *Adv. Appl. Math. Sci.*, **20**(9) (2021) 1767-1786. https://www.mililink.com/upload/article/2006614963aams_vol_209_july_2021_a11_p1767-1786_k._thirusangu_and_a._selvaganapathy.pdf

[3900] M. I. Tilukay, A. N. M. Salman, and E. R. Persulessy, On the total irregularity strength of fan, wheel, triangular book, and friendship graphs, *Procedia Compu. Sci,*, **74** (2015) 124-131. https://www.researchgate.net/publication/289994297_On_the_Total_Irregularity_Strength_of_Fan_Wheel_Triangular_Book_and_Friendship_Graphs

[3901] M. I. Tilukay, P. D. M. Taihuttu, A. N. M. Salman, F. Y. Rumlawang, and Z. A. Leleury, Complete bipartite graph is a totally irregular total graph, *Electron. J. Graph Theory Appl.,* (EJGTA) **9**(2) (2021) 387-395. https://www.ejgta.org/index.php/ejgta/article/view/453/pdf_186

[3902] M. I. Tilukay, A. Salman, V. Y. Ilwaru, and F. Y. Rumlawang, The total face irregularity strength of some plane graphs,*AKCE Int. J. Graphs Comb.,* **17**(1) (2020) 495-502. https://www.sciencedirect.com/science/article/pii/S0972860017302463

[3903] M. I. Tilukay and A. N. N. Salman, The total disjoint irregularity strength of some certain graphs, *Indonesian J. Combin.*, **4**(2) (2020) 91-97.

[3904] M. I. Tilukay, B. P. Tomasouw, F. Y. Rumlawang, and Z. A. Leleury, The total irregularity strength of complete graphs and complete bipartite graphs, *Far East J. Math. Sci.*, **102**(2) (2017) 317-327. http://dx.doi.org/10.17654/MS102020317

[3905] T. I. Titawanno, M. I. Tilukay, Z. A. Leleury, P. D. M. Taihuttu, L. Loves, The total disjoint irregularity strength of double and triple star graphs, *Tensor : Pure Appl. Math. J.*, **5**(2) (2024) 105-110. https://www.researchgate.net/publication/389411668_The_Total_Disjoint_Irregularity_Strength_of_a_Double_and_Triple_Star_Graphs

[3906] A. Tiwari and A. Tripathi, On the range of size of sum graphs & integral sum graphs of a given order, *Discrete Appl. Math.,* **161** (16-17) (2013) 2653-2661. https://www.sciencedirect.com/science/article/pii/S0166218X13002308?via%3Dihub

[3907] H. Todt, On edge-balance index sets of distance two, *Congr. Numer.,* **223** (2015) 105-117.

[3908] A. Tonapi and D. Paquin, On coprime-preserving transformations and dynamic coprime labeling, 13pp. https://doi.org/10.48550/arXiv.2603.22552

[3909] C. Tong, X. Lin, Y. Yang, and L. Wang, Irregular total labellings of $C_m \square C_n$, *Util. Math.,* **81** (2010) 3-13. https://www.researchgate.net/publication/267139738_Irregular_total_labellings_of_C_m_C_n

[3910] A. Tout, A. N. Dabboucy, and K. Howalla, Prime labeling of graphs, *Nat. Acad. Sci. Letters,* **11** (1982) 365-368.

[3911] T. Traetta, A complete solution to the two-table Oberwolfach problems, *J. Combin. Theory, Series A,* **120**(5) (2013) 984-997. https://www.sciencedirect.com/science/article/pii/S0097316513000149?via%3Dihub

[3912] M. Trenklér, Some results of magic graphs, graphs and other combinatorics topics, Teubner-Texte zur Mathematik - Band 59, Leipzig (1983) 328-332. https://www.researchgate.net/publication/236162416_Some_Results_of_Magic_Graphs

[3913] M. Trenklér, Numbers of vertices and edges of magic graphs, *Ars Combin.,* **55** (2000) 93-96. https://combinatorialpress.com/article/ars/Volume%20055/volume-55-paper-8.pdf?utm_source=chatgpt.com

[3914] M. Trenklér, Super-magic complete n-partite hypergraphs, *Graphs and Combin.,* **17** (2001) 171-175. https://www.researchgate.net/publication/220397559_Super-Magic_Complete_k-Partite_Hypergraphs

[3915] M. Trenklér, Magic p-dimensional cubes, *Acta Arithmetica,* **96** (2001) 361-364. https://www.researchgate.net/publication/236162460_Magic_p-dimensional_cubes

[3916] M. Trenklér and V. Vetchý, Magic powers of graphs, *Math. Bohemica,* **122** (1997) 121-124. https://dml.cz/bitstream/handle/10338.dmlcz/125922/MathBohem_122-1997-2_2.pdf

[3917] S. Triwahyuniti and D. Rahmadani, Pelabelan prima pada kelas graf hasil operasi perkalian tensor,*Jambura J. Math.*, **8**(1) (2026) 26-31. https://ejurnal.ung.ac.id/index.php/jjom/article/view/34596/12322

[3918] M. Truszczyński, Graceful unicyclic graphs, *Demonstatio Mathematica,* **17** (1984) 377-387. https://www.researchgate.net/publication/266239884_Graceful_unicyclic_graphs

[3919] Y. Tsao, A simple research of divisor graphs, *The 29th Workshop Combin. Math. Computation Th.*, National Taipei College of Business, Institute of Information and Decision Sciences, Taipei, Taiwan, April 27-28, 2012, 186-190.

[3920] M. Tuczyński, P. Wenus, and K. Węsek, On cordial labeling of hypertrees, *Discrete Math. Theor. Comput. Sci.,* **21**(4) (2019) Paper No. 1, 14 pp. https://dmtcs.episciences.org/5658/pdf

[3921] M. Tuga, M. Miller, J. Ryan, and Z. Ryjáček, Exclusive sum labelings of trees, *J. Combin. Math. Combin. Comput.,* **55** (2005) 109-121. https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2010.pdf

[3922] Z. Tuza, Graph labeling games, *Electronic Notes Disc. Math.,* **60** (2017) 61-68. https://doi.org/10.48550/arXiv.1609.03084

[3923] N. Ujwala, Applications of graceful graph, *Internat. J. Eng. Sci. Res. Tech.*, **4** (2015) 129-131. https://www.ijesrt.com/Old_IJESRT/issues%20pdf%20file/Archives-2015/August-2015/19_APPLICATIONS%20OF%20GRACEFUL%20GRAPHS.pdf

[3924] P. P. Ulaganathan, B. Selvam and P. V. Kumar, Signed product cordial labeling in duplicate graphs of bistar, double star and triangular ladder graph, *Internat. J. Math. Trends Tech.*, **33** (2016) 1-6. https://ijmttjournal.org/public/assets/volume-33/number-1/IJMTT-V33P505.pdf

[3925] R. Ullah, G. Rahmat, M. Numan, K. Anoh Yannick, and A. Aslam, 3-*total edge product cordial labeling for stellation of square grid graph*, *J. Math.,* (2021) Art. ID 1724687, 6 pp. https://downloads.hindawi.com/journals/jmath/2021/1724687.pdf

[3926] R. Ulfatimah, M. Roswitha, and T. A. Kusmayadi, H-supermagic labeling on shrubs graph and $L_m \cdot P_n$, *Internat. Confer. Math.: Edu., Th. Appl.,* IOP Conf. Series: Journal of Physics: Conf. Series **855** (2017) 012055 1-8.

[3927] J. Uma and A. Mazudha Shanofer, A note on edge–graceful labeling for corona and flower graph, *AIP Conf. Proceed. 2112, 020095,* (2019) 1-5. https://aip.scitation.org/doi/pdf/10.1063/1.5112280

[3928] L. Uma and G. Rajasekaran, On alpha labeling of tensor product of paths and cycles, *Heliyon,* **9**(11) E21430 (2023) 1-12. https://www.sciencedirect.com/science/article/pii/S2405844023086383

[3929] L. Uma and G. Rajasekaran, The vertex local antimagicness for Knödel and Fibonacci graphs, *Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Komp'yuternye Nauki*, **5**(2) (2025) 297–314. https://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=vuu&paperid=928&option_lang=eng

[3930] A. Uma Maheswari and S. Azhagarasi, New labeling for graphs-AUM block sum labeling, *Internat. J. Current Sci.* (IJCSPUB), **12**(1) (2022) 574-584. https://rjpn.org/ijcspub/papers/IJCSP22A1199.pdf

[3931] A. Uma Maheswari and S. Azhagasari, AUM block labeling for cycle cactus block graphs, *Compliance Engin. J.*, **13**(4) (2022) 84-96. https://www.researchgate.net/publication/361100048_AUM_BLOCK_LABELLING_FOR_CYCL_CACTUS_BLOC__GRAPHS

[3932] A. Uma Maheswari and S. Azhagarasi, AUM block sum labeling for some special graphs, *Internat. J. Mechanical Engin.* **7**(Special Issue 5) (2022) 102-110. https://www.researchgate.net/publication/361099607_AUM_BLOCK_SUM_LABELLING_FOR_SOME_SPECIAL_GRAPHS

[3933] A. Uma Maheswari and A. S. Purnalakshimi, Aum block labelling for friendship, tadpole and cactus graphs, *Neuro Quantology*, **20**(6) (2022) 7876-7884. https://www.neuroquantology.com/media/article_pdfs/7876-7884.pdf

[3934] A. Uma Maheswari and A. S. Purnalakshimi, AUM block labeling for kite, comb and diamond snake graphs, *J. North Eastern Univ.*, **25**(04).

[3935] A. Uma Maheswari and V. Sumathi, AUM block geometric mean labeling, *African J. Biological Sci.*, **6**(Si4) (2024) 4671-4682. https://www.afjbs.com/uploads/paper/40dc6615088dd6d13fc1ae730e5c691e.pdf

[3936] A. Uma Maheswari and V. Sumathi, AUM block geometric mean labeling for tadpole, kayak paddle and bull graphs, *Util. Math.*, **122**(2) (2025).

[3937] A. Uma Maheswari and V. Sumathi, Some characterisation of AUM block geometric mean labelled graphs, *Dandao Xuebao J. Ballistics*, **38**(2) (2026) 705-712. https://ballisticsjournal.com/index.php/journal/article/view/404/300

[3938] A. M. Uyesser and A. Pokrovskiy, On the Graham–Sloane harmonious labelling conjecture. arXiv:2509.05280v1 22 pages.

[3939] S. K. Vaidya and C. M. Barasara, Product cordial labeling for some new graphs, *J. Math. Research,* **3**(2) (2011) 206-211. http://www.ccsenet.org/journal/index.php/jmr/article/view/8703

[3940] U. Vaghela and D. Parmar, Difference perfect square cordial labeling, *J. Gujarat Res. Soc.,* **21**(15) (2019) 758-770. https://www.researchgate.net/publication/338503111_Difference_Perfect_Square_Cordial_Labeling

[3941] U. Vaghela and D. Parmar, Difference perfect square cordial labeling of some graphs, *J. Xidian Univer.,* **14**(2) (2020) 87-98. https://www.researchgate.net/publication/339676138_Difference_Perfect_Square_Cordial_Labeling_of_Some_Graphs

[3942] U. Vaghela and D. Parmar, Difference perfect square cordial labeling of snake graphs, *Zeichen J.*, **6**(3) 2020 54-73. https://www.researchgate.net/publication/340309173_Difference_Perfect_Square_Cordial_Labeling_of_Snake_Graphs/referencesW#fullTextFileContent

[3943] U. Vaghela and D. Parmar, Extra Skolem difference mean labeling of various graphs, *J. Math. Stat. Studies*, JMSS **2**(1) (2021) 26-39 https://al-kindipublisher.com/index.php/jmss/article/view/1465/1274

[3944] U. Vaghela and D. Parmar, Edge magic total labeling & edge trimagic total labeling of some graphs, *Annals of R.S.C.B.*, **25**(4) (2021) 5784-5809. https://www.annalsofrscb.ro/index.php/journal/article/view/3145/2616

[3945] U. Vaghela and D. Parmar, Tri sum perfect square cordial labeling of graphs, *Internat. J. Math. Stat. Invention.* (IJMSI) **12**(2) (2024) 38-47. https://www.researchgate.net/publication/380317221_Tri_Sum_Perfect_Square_Cordial_Labeling_of_Graphs

[3946] S. K. Vaidya and C. M. Barasara, Further results on product cordial labeling, *International J. Math. Combin.,* **3** (2012) 64-71. https://fs.unm.edu/IJMC/FurtherResultsOnProductCordialLabeling.pdf?utm_source=chatgpt.comh

[3947] S. K. Vaidya and C. M. Barasara, Product cordial graphs in the context of some graph operations, *Internat. J. Math. Sci. Comput.*, **1**(2) (2011) 1-6. https://www.researchgate.net/publication/265662133_Product_Cordial_Graphs_in_the_Context_of_Some_Graph_Operations

[3948] S. K. Vaidya and C. M. Barasara, Some product cordial graphs, *Elixir Discrete Math.*, **41** (2011) 5948-5952. https://www.researchgate.net/publication/258024450_Some_New_Product_Cordial_Graphs

[3949] S. K. Vaidya and C. M. Barasara, Harmonic mean labeling in the context of duplication of graph elements, *Elixir Discrete Math.*, **48** (2012) 9482-9485. https://www.researchgate.net/publication/374947045_Harmonic_mean_labeling_in_the_context_of_duplication_of_graph_elements

[3950] S. K. Vaidya and C. M. Barasara, Further results on product cordial graphs, *Internat. J. Math. Soft Comput.*, **2**(2) (2012) 67-74. https://www.researchgate.net/publication/277765650_Further_Results_on_Product_Cordial_Graphs

[3951] S. K. Vaidya and C. M. Barasara, Edge product cordial labeling of graphs, *J. Math.Comput. Science*, **2**(5) (2012) 1436-1450. http://scik.org/index.php/jmcs/article/view/420/189

[3952] S. K. Vaidya and C. M. Barasara, Geometric mean labeling in the context of duplication of graph elements, *Internat. J. Math. Sci. Engin. Appl.*, **6**(6) (2012) 311-319. http://www.ascent-journals.com/IJMSEA/Vol6No6/29-S.K.Vaidya.pdf

[3953] S. K. Vaidya and C. M. Barasara, Some new families of edge product cordial graphs, *Advanced Modeling Optimization*, **15**(1) (2013) 103-111. http://camo.ici.ro/journal/vol15/v15a9.pdf

[3954] S. K. Vaidya and C. M. Barasara, Some edge product cordial graphs, *Internat. J. Math. Soft Comput.*, **3**(3) (2013) 49-53. https://oaji.net/articles/2017/296-1504259589.pdf

[3955] S. K. Vaidya and C. M. Barasara, Edge product cordial labeling in the context of some graph operations, *Internat. J. Math. Scientific Comput.*, **3**(1) (2013) 4-7. https://veltech.edu.in/wp-content/uploads/2016/04/Paper-02-13.pdf

[3956] S. K. Vaidya and C. M. Barasara, Total edge product cordial labeling of graphs, *Malaya J. Matematik*, **3**(1) (2013) 55-63. http://www.malayajournal.org/articles/MJM039.pdf

[3957] S. K. Vaidya and C. M. Barasara, On total edge product cordial labeling, *Internat. J. Math. Scientific Comput.*, **3**(2) (2013) 12-16. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/Paper05-2013.pdf

[3958] S. K. Vaidya and C. M. Barasara, On edge product cordial labeling of some product related graphs, *Internat. J. Math. Appl.*, **2**(2) (2014) 15-22. https://oaji.net/articles/2014/1167-1408376924.pdf

[3959] S. K. Vaidya and C. M. Barasara, Product and edge product cordial labeling of degree splitting graph of some graphs, *Adv. Appl. Discrete Math.,* **15**(1) (2015) 61-74. https://www.researchgate.net/publication/276912945_PRODUCT_AND_EDGE_PRODUCT_CORDIAL_LABELING_OF_DEGREE_SPLITTING_GRAPH_OF_SOME_GRAPHS

[3960] S. K. Vaidya and C. M. Barasara, Embedding and NP-complete problems for some equitable labelings, *J. Appl. Comp. Sci. Math.*, **18**(8) (2014) 38-42. https://www.jacsm.ro/view/?pid=18_6

[3961] S. K. Vaidya and C. M. Barasara, On embedding and NP-complete problems of equitable labelings, *IOSR J. Math.*, **11**(1)(III) (2015) 80-85. http://iosrjournals.org/iosr-jm/papers/Vol11-issue1/Version-3/N011138085.pdf

[3962] S. K. Vaidya and C. M. Barasara, Product cordial labeling of line graph of some graphs, *Kragujevac J. Math.*, **40**(2) (2016) 290-297. https://pdfs.semanticscholar.org/0b62/22107afc5ad737e4c83bcb8862685d926724.pdf

[3963] S. K. Vaidya and L. Bijukumar, Odd graceful labeling of some new graphs, *Modern Appl. Sci.,* **4** (2010) 65-69. https://ccsenet.org/journal/index.php/mas/article/view/6684

[3964] S. K. Vaidya and L. Bijukumar, New mean graphs, *International J. Math. Combin.,* **3** (2011) 107-113. https://zenodo.org/records/9363

[3965] S. K. Vaidya and N. A. Dani, Some new star related graphs and their cordial as well as 3-equitable labeling, *J. Science* **1**(1) (2010) 111-114.

[3966] S. K. Vaidya and N. A. Dani, Cordial labeling and arbitrary super subdivision of some graphs, *Inter. J. Information Sci. Comput. Math.,* **2(1)** (2010) 51-60.

[3967] S. K. Vaidya and N. A. Dani, Some new product cordial graphs, *J. Applied Computer Science & Math.,* **8(4)** (2010) 62-65. http://jacs.usv.ro/index.php

[3968] S. K. Vaidya and N. A. Dani, Cordial and 3-equitable graphs induced by duplication of edge, *Math. Today,* **27** (2011) 71-82.

[3969] S. K. Vaidya, N. Dani, K. Kanani, and P. Vihol, Cordial and 3-equitable labeling for some star related graphs, *Internat. Mathematical Forum,* **4** (2009) 1543-1553. https://www.researchgate.net/publication/258026093_Cordial_and_3-Equitable_Labeling_for_Some_Star_Related_Graphs a

[3970] S. K. Vaidya, N. Dani, K. Kanani, and P. Vihol, Cordial and 3-equitable labeling for some shell related graphs, *J. Sci. Res.,* **1** (2009) 438-449. https://www.researchgate.net/publication/251013932_Cordial_And_3Equitable_Labeling_for_Some_Shell_Related_Graphs

[3971] S. K. Vaidya, N. Dani, K. Kanani, and P. Vihol, Some wheel related 3-equitable graphs in the context of vertex duplication, *Adv. and Appl. Disc. Math.,* **4** (2009) 71-85. https://www.researchgate.net/publication/258026468_Some_wheel_related_3-Equitable_Graphs_in_the_context_of_vertex_duplication

[3972] S. K. Vaidya, N. A. Dani, K. K. Kanani, and P. L. Vihol, Cordial and 3-equitable labeling for some wheel related graphs, *IAENG International J. Applied Math.* **41**(2) (2011) 99-105. https://www.iaeng.org/IJAM/issues_v41/issue_2/IJAM_41_2_03.pdf

[3973] S. K. Vaidya, N. A. Dani, P. L. Vihol, and K. K. Kanni, Strongly multiplicative labeling in the context of arbitrary supersubdivision, *J. Math. Research,* **2**(2) (2010) 28-33. https://pdfs.semanticscholar.org/4d0d/553257f337b63c04dc087b696fd0e293abc0.pdf

[3974] S. K. Vaidya, G. Ghodasara, S. Srivastav, and V. Kaneria, Cordial labeling for two cycle related graphs, *Math. Student,* **76** (2007) 237-246.

[3975] S. K. Vaidya, G. V. Ghodasara, S. Srivastav, and V. J. Kaneria, Cordial and 3-equitable labeling of star of a cycle, *Math. Today,* **24** (2008) 54-64.

[3976] S. K. Vaidya, G. Ghodasara, S. Srivastav, and V. Kaneria, Some new cordial graphs, *Internat. J. Scientific Computing,* **2** (2008) 81-92.

[3977] S. K. Vaidya and K. Kanani, Some new results on cordial labeling in the context of arbitrary supersubdivision of graph, *Applied Math. Sci.,* **4**(47) (2010) 2323-2329. https://www.m-hikari.com/ams/ams-2010/ams-45-48-2010/vaidyaAMS45-48-2010.pdf

[3978] S. K. Vaidya and K. K. Kanani, Some cycle related product cordial graphs, *Internat. J. Algor., Comput. and Math.,* **3**(1) (2010) 109-116. https://www.researchgate.net/publication/258026789_Some_Cycle_Related_Product_Cordial_Graphs

[3979] S. K. Vaidya and K. K. Kanani, Strongly multiplicative labeling for some cycle related graphs, *Modern Applied Science,* **4**(7) (2010) 82-88. https://ccsenet.org/journal/index.php/mas/article/view/6648

[3980] S. K. Vaidya and K. K. Kanani, Prime labeling for some cycle related graphs, *J. Math. Research,* **2**(2) (2010) 98-103. https://ccsenet.org/journal/index.php/jmr/article/view/4423

[3981] S. K. Vaidya and K. K. Kanani, Some new product cordial graphs, *Math. Today,* **27** (2011) 64-70. https://www.researchgate.net/publication/258024450_Some_New_Product_Cordial_Graphs

[3982] S. K. Vaidya and K. K. Kanani, Some strongly multiplicative graphs in the context of arbitrary supersubdivision, *Internat. J. Applied Math. and Comput.,* **3**(1) (2011) 60-64. https://www.researchgate.net/publication/258023265_Some_Strongly_Multiplicative_Graphs_in_the_Context_of_Arbitrary_Super_subdivision

[3983] S. K. Vaidya, V. J. Kaneria, S. Srivastav, and N. A. Dani, Gracefulness of union of two path graphs with grid graph and complete bipartite graph, *Proceedings of the First International Conference on Emerging Technologies and Applications in Engineering Technology and Sciences* (2008) 616-619.

[3984] S. K. Vaidya, K. Kanani, S. Srivastav, and G. Ghodasara, Baracentric subdivision and cordial labeling of some cycle related graphs, *Proceedings of the First International Conference on Emerging Technologies and Applications in Engineering, Technology and Sciences,* (2008) 1081-1083.

[3985] S. K. Vaidya, K. Kanani, P. Vihol, and N. Dani, Some cordial graphs in the context of barycentric subdivision, *Int. J. Comtemp. Math. Sciences,* **4** (2009) 1479-1492. https://www.m-hikari.com/ijcms-password2009/29-32-2009/vaidyaIJCMS29-32-2009.pdf

[3986] S. K. Vaidya and N. J. Kothari, Some new families of line graceful graphs, *Internat. J. Mathematics Sci. Comput.*, **1**(2) (2011) 26-28. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/IJMSC01-02-Paper05-2011.pdf

[3987] S. K. Vaidya and N. J. Kothari, Line gracefulness of some path related graphs, *Internat. J. Math. Sci. Comput.*, **4**(1) (2014) 15-18. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/Paper-05-14.pdf

[3988] S. K. Vaidya and B. Lekha, Odd graceful labeling of some new graphs, *Modern Appl. Sci.,* **4** (10) (2010) 65-70. https://ccsenet.org/journal/index.php/mas/article/view/6684

[3989] S. K. Vaidya and B. Lekha, Some new odd graceful graphs, *Advances Appl. Disc. Math.,* **6**(2). (2010) 101-108.

[3990] S. K. Vaidya and B. Lekha, New families of odd graceful graphs, *Internat. J. Open Problems Comp. Sci. Math.,* **3**(5) (2010) 166-171. www.ijopcm.org/Vol/10/IJOPCM(vol.3.5.18.D.10).pdf

[3991] S. K. Vaidya and B. Lekha, Some new graceful graphs, *Internat. J. Math. Soft Comp.,* **1**(1) (2011) 37-45. http://oaji.net/articles/2015/296-1433770065.pdf

[3992] S. K. Vaidya and B. Lekha, Some new results on E-cordial labeling, *Internat. J. Information Sci. Comput. Math.,* **3**(1) (2011) 21-29.

[3993] S. K. Vaidya and B. Lekha, Some new families of E-cordial graphs, *J. Math. Res.,* **3**(4) (2011) 105-111. https://ccsenet.org/journal/index.php/jmr/article/view/10600/8959

[3994] S. K. Vaidya and B. Lekha, Some new perspectives on odd sequential graphs, *Internat. J. Math. Soft Comput.*, **4**(1) (2014) 93-101. https://oaji.net/articles/2017/296-1504711954.pdf

[3995] S. K. Vaidya and U. M. Prajapati, Some results on prime and k-prime labeling, *J. Math. Research,* **3**(1) (2011) 66-75. http://www.ccsenet.org/journal/index.php/jmr/

[3996] S. K. Vaidya and U. M. Prajapati, Some switching invariant prime graphs, *Open J. Discrete Math.*, **2** (2012) 17-20. http://www.scirp.org/journal/PaperInformation.aspx?paperID=17155

[3997] S. K. Vaidya and U. M. Prajapati, Fibonacci and super Fibonacci graceful labelings of some cycle related graphs, *Inter. J. Math. Combin.*, **4** (2011) 59-69. https://fs.unm.edu/IJMC/FibonacciAndSuperFibonacciGracefulLabelings.pdf

[3998] S. K. Vaidya and U. M. Prajapati, Some new results on prime graphs, *Open J. Discrete Math.*, **2** (2012) 99-104. doi:10.4236/ojdm.2012.23019

[3999] S. K. Vaidya and U. M. Prajapati, Prime labeling in the context of duplication of graph elements, *Internat. J. Math. Soft Comput.*, **3**(1) (2013) 13-20.

[4000] S. K. Vaidya, U. M. Prajapati, and P. Vihol, Some important results on triangular sum graphs, *Appl. Math. Sciences,* **3** (2009) 1763-1772. https://www.m-hikari.com/ams/ams-password-2009/ams-password33-36-2009/vaidyaAMS33-36-2009.pdf

[4001] S. K. Vaidya and N. H. Shah, Some new families of prime cordial graphs, *J. Math. Research* **3**(4) (2011) 21-30. https://ccsenet.org/journal/index.php/jmr/article/view/9650

[4002] S. K. Vaidya and N. H. Shah, Some new odd harmonious graphs, *Internat. J. Math. and Soft Comput.,* **1**(1) (2011) 9-16. https://oaji.net/articles/2015/296-1433769758.pdf

[4003] S. K. Vaidya and N. H. Shah, Odd harmonious labeling of some graphs, *International J. Math. Combin.,* (3) (2012) 105-112. https://fs.unm.edu/IJMC/OddHarmoniousLabelingOfSomeGraphs.pdf

[4004] S. K. Vaidya and N. H. Shah, Prime cordial labeling of some graphs, *Open J. Discrete Math.*, **2** (2012) 11-16. http://www.scirp.org/journal/PaperInformation.aspx?paperID=17154

[4005] S. K. Vaidya and N. H. Shah, 3-Equitable labeling for some star and bistar related graphs, *Internat. J. Math. and Sci. Comput.*, **2**(1) (2012) 3-8. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/IJMSC02-01-Final-Paper-02.pdf

[4006] S. K. Vaidya and N. H. Shah, Graceful and odd graceful labeling of some graphs, *Internat. J. of Math. Soft Computing*, **3**(1) (2013) 61-68. https://oaji.net/articles/2017/296-1502694158.pdf

[4007] S. K. Vaidya and N. H. Shah, Some star and bistar related divisor cordial graphs, *Annals Pure Appl. Math.,* **3**(1) (2013) 67-77. http://www.researchmathsci.org/apamart/apam-v3n1-7.pdf

[4008] S. K. Vaidya and N. H. Shah, Prime cordial labeling of some wheel related graphs, *Malaya J. Mate.*, **4**(1) (2013) 148-156. https://www.malayajournal.org/articles/MJM057.pdf

[4009] S. K. Vaidya and N. H. Shah, Further results on divisor cordial labeling, *Annals Pure Appl. Math.*, **4**(2) (2013) 150-159. http://www.researchmathsci.org/apamart/apam-v4n2-5.pdf

[4010] S. K. Vaidya and N. H. Shah, Some new results on prime cordial labeling, *ISRN Combin.*, **2014** Article ID 607018, 9 pages, 2014. http://www.hindawi.com/journals/isrn.combinatorics/2014/607018/

[4011] S. K. Vaidya and N. H. Shah, Cordial labeling for some bistar related graphs, *Internat. J. Mathematics Soft Comput.*, **4**(2) (2014) 33-39. https://oaji.net/articles/2017/296-1504713044.pdf

[4012] S. K. Vaidya and N. H. Shah, Cordial labeling of snakes, *Internat. J. Math. Appl.*, **2**(3) (2014) 17-27. https://oaji.net/articles/2014/1167-1408373409.pdf

[4013] S. K. Vaidya and N. H. Shah, On square divisor cordial graphs, *J. Scientific Research*, **6**(3) (2014) 445-455. http://www.banglajol.info/index.php/JSR/article/view/16412

[4014] S. K. Vaidya and N. H. Shah, Cordial labeling of degree splitting graphs, *J. Graph Labeling,* **1**(2) (2015) 95-110.

[4015] S. K. Vaidya, S. Srivastav, G. Ghodasara, and V. Kaneria, Cordial labeling for cycle with one chord and its related graphs, *Indian J. Math. Sciences,* **4** (2008) 145-156.

[4016] S. K. Vaidya, S. Srivastav, V. J. Kaneria, and G. V. Ghodasara, Cordial and 3-equitable labeling of cycle with twin-chords, *Proceed. First International Conference on Emerging Technologies and Applications in Engineering, Technology and Sciences,* (2008) 905-907.

[4017] S. K. Vaidya, S. Srivastav, V. J. Kaneria, and K. K. Kanani, Some cycle related cordial graphs in the context of vertex switching, *Proceed. International Conf. Discrete Math. - 2008 RMS Lecturer Note Series,* No. **13** (2010) 243-252. https://www.researchgate.net/publication/258023343_Some_cycle_related_cordial_graphs_in_the_context_of_vertex_switching

[4018] S. K. Vaidya and P. L. Vihol, Total product cordial graphs induced by some graph operations on cycle related graphs, *Internat. J. Inform. Sci. and Comput. Math.,* **1**(2) (2010) 113-126.

[4019] S. K. Vaidya and P. L. Vihol, Cordial labeling for middle graph of some graphs, *Elixir Discrete Math.,* **34C** (2011) 2468-2475. https://www.elixirpublishers.com/articles/1686047788_201105056.pdf

[4020] S. K. Vaidya and P. L. Vihol, Prime cordial labeling for some graphs, *Modern Applied Science,* **4**(8) (2010) 119-126.

[4021] S. K. Vaidya and P. L. Vihol, Embedding and NP-complete problems for 3-equitable graphs, *Inter. J. Contemporary Advanced Math.,* **2**(1) (2011) 1-7. https://www.cscjournals.org/manuscript/Journals/IJCM/Volume2/Issue1/IJCM-15.pdf

[4022] S. K. Vaidya and P. L. Vihol, Prime cordial labeling for some cycle related graphs, *Internat. J. Open Problems in Comput. Sci. Math.,* **3**(5) (2010) 223-232. https://www.emis.de/journals/IJOPCM/Vol/10/IJOPCM(vol.3.5.24.D.10).pdf

[4023] S. K. Vaidya and P. L. Vihol, Fibonacci and super Fibonacci graceful labeling of some graphs, *Studies Math. Sciences,* **2**(2) (2011) 24-35. https://www.researchgate.net/publication/228972024_Fibonacci_and_Super_Fibonacci_Graceful_Labeling_of_Some_Graphs

[4024] S. K. Vaidya and P. L. Vihol, Radio labeling for some cycle related graphs, *Interna. J. Math. Soft Comput.,* **2**(2) (2012) 11-24. https://oaji.net/articles/2017/296-1502656338.pdf

[4025] S. K. Vaidya and N. B. Vyas, Product cordial labeling in the context of tensor product of graphs, *J. Math. Research,* **3**(3) (2011) 83-88. https://ccsenet.org/journal/index.php/jmr/article/view/9863

[4026] S. K. Vaidya and N. B. Vyas, *E*-cordial labeling of some mirror graphs, *Internat. Contemp. Adv. Math.,* **2(1)** (2011) 22-27. https://www.cscjournals.org/manuscript/Journals/IJCM/Volume2/Issue1/IJCM-19.pdf

[4027] S. K. Vaidya and N. B. Vyas, *E*-cordial labeling for Cartesian product of some graphs, *Studies Math. Sci.*, **3**(2) (2011) 11-15. https://www.researchgate.net/publication/277763305_E-cordial_Labeling_for_Cartesian_Product_of_Some_Graphs

[4028] S. K. Vaidya and N. B. Vyas, Further results on *E*-cordial labeling, *Internat. J. Informa. Sci. Comput. Math.*, **5**(1) (2012) 9-18. http://www.pphmj.com/abstract/6905.htm

[4029] S. K. Vaidya and N. B. Vyas, Antimagic labeling in the context of switching of a vertex, *Annals Pure Appl. Math.,* **2**(1) (2012) 33-39. http://www.researchmathsci.org/apamart/apam-v2n1-4.pdf

[4030] S. K. Vaidya and N. B. Vyas, Some results on *E*-cordial Llabeling, *Internat. J. Math. Sci. Comput.*, **2**(1) (2012) 9-13. https://demovtu.veltech.edu.in/wp-content/uploads/2016/04/IJMSC02-01-Final-Paper-03.pdf

[4031] S. K. Vaidya and N. B. Vyas, E-cordial labeling in the context of switching of a vertex, *Internat. J. Adv. Comput. Math. Sci.*, **3**(4) (2012) 446-452. http://bipublication.com/files/IJCMS-V3I4-2012-9.pdf

[4032] S. K. Vaidya and N. B. Vyas, Even mean labeling for path and bistar related graphs, *Internat. J. Graph Theory*, **1**(4) (2013) 122-130.

[4033] S. K. Vaidya and N. B. Vyas, Antimagic labeling of some path and cycle related graphs, *Annals Pure Appl. Math.,* **3**(2) (2013) 119-128. http://www.researchmathsci.org/apamart/apam-v3n2-4.pdf

[4034] S. K. Vaidya and N. B. Vyas, Product cordial labeling for alternate snake graphs, *Malaya Journal of Mathematik*, **2**(3) (2014) 188-196. http://www.malayajournal.org/articles/MJM082.pdf

[4035] S. K. Vaidya and N. B. Vyas, Product cordial labeling for some bistar related graphs, *Adv. Modeling Optim.*, **16**(2) (2014) 295-301. http://camo.ici.ro/journal/vol16/v16b4.pdf

[4036] K. Vaithilingam, Difference labeling of some graph families, *Internat. J. Math. Stat. Invention* (IJMSI), **2** (2014 37-43. https://www.ijmsi.org/Papers/Volume.2.Issue.6/E026037043.pdf

[4037] K. Vaithilingam and S. Meena, Prime labeling for some crown related graphs, *Internat. J. Sci. & Tech. Res.*, **2**(3) (2013) 92-95. https://www.ijstr.org/final-print/mar2013/Prime-Labeling-For-Some-Crown-Related-Graph

[4038] D. K. Valani and K. K. Kanani, Edge even graceful labeling of triangular belt graph, alternate triangular belt graph and $Z-P_n$ graph, *Indian J. Discrete Math.*, **9**(1) (2023) 31-50. https://www.researchgate.net/publication/382001535_

Edge_Even_Graceful_Labeling_Of_Triangular_Belt_Graph_Alternate_Triangular_Graph_And_Z-PnGraph

[4039] D. K. Valani and K. K. Kanani, Edge odd graceful lablinng of corona product of some graphs, *Bull. Cal. Math. Soc.*, **117**(1) (2025) 145–166. https://www.researchgate.net/publication/395382908_EDGE_ODD_GRACEFUL_LABELING_OF_CORONA_PRODUCT_OF_SOME_GRAPHS

[4040] D. K. Valani and K. K. Kanani, Edge odd garceful labelings of some snake graphs, *J. Indian Acad. Math.*, **47**(1) (2025) 133-152. https://www.researchgate.net/publication/395382587_EDGE_ODD_GRACEFUL_LABELING_OF_SOME_SNAKE_GRAPHS

[4041] A. Velankanni, A. Bernick Raj, and M. Sujasree, Odd graceful labeling of arbitrary supersubdivision of certain graphs, *Adv. Appl. Disc. Math.,* **34** (2022) 23-37. https://pphmjopenaccess.com/index.php/aadm/article/view/708/665

[4042] K. Valentin, Polychrome labelings of trees and cycles, *Ars Combin.,* **52** (1999) 272-284. https://combinatorialpress.com/article/ars/Volume%20052/volume-52-paper-24.pdf

[4043] L. S. Valencerina and A. C. Pedrano, On super heronian mean labeling of some subdivision graphs, *Comm. Math. Appl.,* **15**(2) (2024) 597-603. https://www.rgnpublications.com/journals/index.php/cma/article/view/2162/1696

[4044] F. Van Bussel, Relaxed graceful labellings of trees, *Electronic J. Combin.*, **9** (2002) #R4. (12 pages). https://www.combinatorics.org/ojs/index.php/eljc/article/view/v9i1r4/pdf

[4045] S. Vani Shree snd S. Dhanalakshmi, New results on face magic mean labeling of graphs, *Math. Stat.,* **11**(1) (2023) 199-205. https://www.researchgate.net/publication/371253652_New_Results_on_Face_Magic_Mean_Labeling_of_Graphs

[4046] V. Vanitha, A Study on Graph Labeling of Directed Graphs, PhD thesis, Mother Teresa Women's University, 2012.

[4047] R. Varatharajan, S. Navanaeethakrishnan, and K. Nagarajan, Divisor cordial graphs, *Internat. J. Math. Combin.,* **4** (2011) 15-25. https://fs.unm.edu/IJMC/DivisorCordialGraphs.pdf

[4048] R. Varatharajan, S. Navanaeethakrishnan, K. Nagarajan, Special classes of divisor cordial graphs, *Int. Math. Forum,* **7**(33-36) (2012) 1737-1749. https://www.m-hikari.com/imf/imf-2012/33-36-2012/nagarajanIMF33-36-2012.pdf

[4049] M. Vargheese and R. Arun, Even harmonious labelings of triangular book graphs, *Caribbean J. Sci.,* **53**(2) (2019) 2111-2117.

[4050] T. K. Mathew Varkey and B. S. Sunoj, Some new families of odd graceful graphs, *Internat. J. Math. Combin.,* **3** (2016) 158-161. https://fs.unm.edu/IJMC/SomeNewFamiliesOfOddGracefulGraphs.pd

[4051] T. K. Mathew Varkey and B. S. Sunoj, A note on absolute difference of cubic and square sum labeling of a class of trees, *Internat. J. Sci. Eng. Appl. Sci.*, **2**(8) (2016) 293-296. https://ijseas.com/volume2/v2i8/ijseas20160828.pdf

[4052] T. K. Mathew Varkey and B. S. Sunoj, An absolute difference of cubic and square sum labeling of splitting graphs, *Internat. Comput. Math. Sci.*, **5**(8) (2016) 16-18.

[4053] T. K. Mathew Varkey and B. S. Sunoj, ADCSS labeling of cycle related graphs, *Internat. J. Scientific Res. Edu.*, **4**(8) (2016) 5702-5705.

[4054] T. K. Mathew Varkey and B. S. Sunoj, An absolute difference of cubic and square sum labeling of certain class of graphs, *Internat. J. Math. Trends Tech.*, **36**(1) (2016) 77-79. https://www.ijmttjournal.org/2016/Volume-36/number-1/IJMTT-V36P509.pdf

[4055] T. K. Mathew Varkey and B. S. Sunoj, Some new results on absolute difference of cubic and square sum labeling of a class of graphs, *Internat. Sci. Res.*, **5**(8) (2016), 1465-1467. https://www.ijsr.net/archive/v5i8/ART20161261.pdf

[4056] T. K. Mathew Varkey and B. S. Sunoj, ADCSS-of product related graphs, *Internat. J. Math. Appl.,* **4**(2B) (2016) 145-149. https://ijmaa.in/index.php/ijmaa/article/view/1036/1021

[4057] T. K. Mathew Varkey and B. S. Suno, ADCSS-labeling of 2-tuple graphs of some graphs, *IOSR J. Math.*, **12**(5)(V) (2016) 12-15. https://www.iosrjournals.org/iosr-jm/papers/Vol12-issue5/Version-5/C1205051215.pdf

[4058] T. K. Mathew Varkey and B.S. Sunoj, ADCSS-labeling for some total graphs, *Internat. Math. Trends Tech.*, **38**(1) (2016) 1-4. https://www.ijmttjournal.org/2016/Volume-38/number-1/IJMTT-V38P501.pdf

[4059] T. K. Mathew Varkey and B. S. Sunoj, ADCSS-Labeling of shadow graph of some graphs, *J. Comput. Math. Sci.*, **7**(11) (2016) 593-598. http://www.compmath-journal.org/dnload/Mathew-Varkey-T-K-and-Sunoj-B-S-/CMJV07I11P0593.pdf

[4060] J. Vasanthi1 and N. Ramya, Prime cordial labelings of Heawood graph, pappus graph and Tutte-Coxeter graph, *J. Dynamics and Control,* **8**(11) (2024) 50-55. https://jodac.org/wp-content/uploads/2024/11/JoDaC-8-11-50-55.pdf

[4061] P. Vasoya and D. Bantva, Optimal radio labelings of the Cartesian product of the generalized Petersen graph and tree, arXiv:2304.10094.pdf https://www.researchgate.net/publication/370153207_Optimal_radio_labelings_of_the_Cartesian_product_of_the_generalized_Peterson_graph_and_tree

[4062] R. Vasuki, S. Arockiaraj, On super mean graphs, *Util. Math.,* **103** (2017) 335-351.

[4063] R. Vasuki and A. Nagarajan, Meanness of the graphs $P_{a,b}$ and P_a^b, *Internat. J. Appl. Math.*, **22**(4) (2009) 663-675.

[4064] R. Vasuki and A. Nagarajan, Some results on super mean graphs, *Internat. J. Math. Combin.* **3** (2009) 82-96.
https://zenodo.org/record/9459#.X61bqMhKiUk

[4065] R. Vasuki and A. Nagarajan, Further results on mean graphs, *Scientia Magna*, **6**(3) (2010) 1-14. https://fs.unm.edu/ScientiaMagna6no3.pdf?utm_source=chatgpt.com

[4066] R. Vasuki and A. Nagarajan, Odd mean labeling of the graphs $P_{a,b}, P_a^b$ and $P^b_{\langle 2a\rangle}$, *Kragujevac J. Math.* **36**(1) (2012) 141-150. https://imi.pmf.kg.ac.rs/kjm/pub/13476279057933_kjom3601-15.pdf

[4067] D. Vickrey, k-equitable labelings of complete bipartite and multipartite graphs, *Ars Combin.,* **54** (2000) 65-85. https://combinatorialpress.com/ars-articles/volume-054-ars-articles/k-equitable-labelings-of-complete-bipartite-and-multipartite-graphs/

[4068] A. Vietri, Graceful labellings for an infinite class of generalised Petersen graphs, *Ars. Combin.,* **81** (2006) 247-255. https://combinatorialpress.com/article/ars/Volume%20081/volume-81-paper-16.pdf
See also A. Vietri, Erratum: *Ars Combin.,* **83** (2007) 381. https://combinatorialpress.com/ars-articles/volume-083-ars-articles/erratum-a-little-emendation-to-the-graceful-labelling-of-the-generalised-petersen-8t3-when-t-5/

[4069] A. Vietri, On the graceful polynomials of a graph, *Austraasian J. Combin.,* **73**(2) (2019) 261-273. https://www.researchgate.net/publication/339676138_Difference_Perfect_Square_Cordial_Labeling_of_Some_Graphs

[4070] R. Vasuki, A. Nagarajan, and S. Arockiaraj, Even vertex odd mean labeling of graphs, *SUT J. Math.*, **49**(2) (2013) 79-92.https://projecteuclid.org/journals/sut-journal-of-mathematics/volume-49/issue-2/Even-vertex-odd-mean-labeling-of-graphs/10.55937/sut/1394108286.full

[4071] B. Vasuki, L. Shobana, and M. A. Ahmed, Face antimagic labeling for double duplication of barycentric and middle graphs, *Iraqi J. Sci.,* (2022) 3902-3908. doi:doi:10.24996/ijs.2022.63.9.22

[4072] B. Vasuki ·and L. Shobana, Face antimagic labeling for double duplication of crown related graphs, *AIP Conf. Proceed. 2516, 210048 (2022). 2nd Internat. Confer. Math. Tech. Appl.,* ICMTA2021.

[4073] R. Vasuki, P. Sugirtha, and J. Venkateswari, Super mean labeling of some subdivision graphs, *Kragujevac J. Math.,* **41**(2) (2017) 179-201. https://pdfs.semanticscholar.org/f225/53eefa3214addfe8300729d8dad5798dc076.pdf

[4074] R. Vasuki, L. Velmurugan, and P. Sugirtha, Some results on super pair sum graphs, *J. Discrete Math. Sci. Cryptogr.,* **20**(2) (2017) 463-476. https://www.researchgate.net/publication/317547428_Some_results_on_super_pair_sum_graphs

[4075] A. Villar, On the product cordial labeling for some crown graphs, U. Southeastern Philippines, Math. Stat. Depart., Davao City, Philippines, (2013) 19-86.

[4076] R. Vimal Kumar and D. Vijayalakshmi, Vertex magic total labeling of middle and total graph of cycle, *Commun. Fac. Sci. Univ. Ank. Ser. A1. Math. Stat.,* **68**(2) (2019) 1335-1340. https://www.researchgate.net/publication/334140866_Vertex_magic_total_labeling_of_middle_and_total_graph_of_cycle

[4077] S. Venkatesh, Open cyclic grid graphs are graceful, *IOSR J. Math.,* **10**(2) (2014) 65-67. http://iosrjournals.org/iosr-jm/papers/Vol10-issue2/Version-6/J010266567.pdf

[4078] S. Venkatesh and S. Bharathi, On generating graceful trees, *Int. J. Pure Appl. Math.,* (IJPAM) **118**(9) 899-904. https://www.researchgate.net/publication/323748246_On_Generating_Graceful_Trees

[4079] S. Venkatesh, B. Mahalakshmi and N. Amirthavahini, New results on some vertex labeling of graphs, *Inter. J. Pure Appl. Math.*, **118**(9) (2018) 891-898. https://www.researchgate.net/publication/323748420_New_Results_On_Some_Vertex_Labeling_of_Graphs

[4080] K. Venkatesan and K. Thirugnanasambandam, Super root square mean labeling of some cycle graphs, *Global J. Eng. Sci. Res.*, (GJESR), **5**(11) (2018) 221-234. https://www.gjesr.com/Issues%20PDF/Archive-2018/November-2018/23.pdf

[4081] S. Venkatesh and K. Balasubramanian, Some results on generating graceful trees,*Internat. J. Engin. Tech.,* 7(4.10) (2018) 570-572. https://www.researchgate.net/publication/332682658_Some_Results_on_Generating_Graceful_Trees

[4082] S. Venkatesh, J. Salah, and G. Sethuraman, Some results on E-cordial graphs, *Internat. J. Math. Trendsand Technology,* **7**(2) (2014) 121-125. https://ijmttjournal.org/public/assets/volume-7/number-2/IJMTT-V7P516.pdf

[4083] S. Venkatesh and S. Sivagurunathan, On the gracefulness of cycle related graphs, *Int. J. Pure Appl. Math.,* (IJPAM) **117**(15) (2017) 589-598.

https://www.researchgate.net/publication/317304532_Super_Root_Square_Mean_Labeling_of_Some_Graphs

[4084] E. Veronisha, A. Lourdusamy, and F. J. Beaula, Total 3-sum cordial labeling on zero divisor graphs, *AIP Conf. Proc.,* 2764, 030018 (2023) https://doi.org/10.1063/5.0144416

[4085] J. C. M. Vertudes and A. C. Pedrano, On topological cordial labeling of some graphs, *European J. Math. Stat.,* **6** (2025) 1-7. https://www.ej-math.org/index.php/ejmath/article/view/404

[4086] A. Vietri, A new infinite family of graceful generalised Petersen graphs, via "graceful collages" again, *Austral. J. Comb.,* **41** (2008) 273-282. https://ajc.maths.uq.edu.au/pdf/41/ajc_v41_p273.pdf

[4087] A. Vietri, Real-graceful labellings: a generalisation of graceful labellings, *Ars Combin.,* **102** (2011) 359-364. https://www.researchgate.net/publication/268019654_Real-graceful_labellings_A_generalisation_of_graceful_labellings

[4088] A. Vietri, Necessary conditions on graceful labels: a study case on trees and other examples, *Util. Math.,* **89** (2012) 275-287. https://www.researchgate.net/publication/266991516_Necessary_conditions_on_graceful_labels_A_study_case_on_trees_and_other_examples

[4089] A. Vietri, Sailing towards, and then against, the Graceful Tree Conjecture: some promiscuous results, *I.C.A. Bulletin*, **53** (2008) 31-46. https://www.researchgate.net/publication/266991516_Necessary_conditions_on_graceful_labels_A_study_case_on_trees_and_other_examples

[4090] M. Vieve, S. Gudin, and A. C. Pedrano, Tribonacci cordial labeling of some snake graphs, *Annals Communications Math.*, **8**(3) (2025) 393-405. https://www.technoskypub.com/journals/acm-2025-080306/

[4091] G. R. Vijayakumar, Arithmetic labelings and geometric labelings of finite graphs, *J. Combin. Math. Combin. Comput.,* **75** (2010) 229-231. https://combinatorialpress.com/article/jcmcc/Volume%20075/vol-075-paper%2020.pdf

[4092] G. R. Vijayakumar, Arithmetic labelings and geometric labelings of countable graphs, *Discuss. Math. Graph Theory,* **30** (2010) 539-544. https://www.dmgt.uz.zgora.pl/publish/pdf.php?doi=1511

[4093] G. R. Vijayakumar, A note on set graceful labeling, arXiv:1101.2729v1 [math.co] 14 Jan 2011.

[4094] G. Vijayalashmi and S. T. Arockiamary, Some results on k-prime labeling of graphs on graph operations, *Adv. Appl. Math. Sci.,* **21**(1) (2021) 283-298. https://www.researchgate.net/publication/357479219_SOME_RESULTS_ON_k-PRIME_LABELING_OF_GRAPHS_ON_GRAPH_OPERATIONS/references

[4095] P. Vijaya kumar, K. Thulukkanam, and T. Thirusangu, Skolem difference mean labeling in duplicate graphs of some path related graphs, *South East Asian J. Math. Math. Sci.*, **14**(3) (2018) 63-78. https://rsmams.org/journals/seajmams/article/322

[4096] P. Vijaya Kumar, P.P. Ulaganathan, and K. Thirusangu, Mean labeling in some duplicate graphs, *Internat. J. Pure App. Math.* **117**(11) (2017) 201-210. https://acadpubl.eu/jsi/2017-117-11-14/articles/11/24.pdf

[4097] K. Vilfred and T. Nicholas, The integral sum graph $_\Delta n$, *Graph Theory Notes N. Y.,* **57** (2009) 43-47. https://www.researchgate.net/publication/313561615_The_Integral_Sum_Graph_G_Delta_n

[4098] V. Vilfred, Families of graceful banana trees, *Internat. J. Management and Systems*, **15**(3) (1999) 275-286 https://www.researchgate.net/publication/342804270_FAMILIES_OF_GRACEFUL_BANANA_TREES

[4099] V. Vilfred, Sigma partitions and sigma labeled graphs, *J. Decision Math. Sci.,* **10**(1-3) (2005) 1-9. https://www.researchgate.net/publication/313561766_SIGMA_PARTITION_AND_SIGMA_LABELED_GRAPHS

[4100] V. Vilfred, Perfectly regular graphs or cyclic regular graphs and $\sum$-labeling and partitions, Srinivasa Ramanujan Centenary Celebrating International Conference in Mathematics, Anna University, Madras, India, Abstract A23 (1987).

[4101] V. Vilfred, $\sum$-labeled Graphs and Circulant Graphs, Ph.D., thesis University of Kerala, Trivandrum, India, (1994).

[4102] V. Vilfred Kamalappan, k-Distance magic labelings and long brush graphs, https://arxiv.org/pdf/2211.09666

[4103] V. Vilfred and M. Jinnah, On sigma labeled graphs, (2002) unpublished.

[4104] V. Vilfred and L. M. Florida, Partition technique in the study of (a, d)-antimagic graphs, unpublished

[4105] V. Vilfred and L. M. Florida, Odd antimagic and odd (a, d)-antimagic graphs, unpublished. 2010 https://www.researchgate.net/publication/313561570_ODD_ANTIMAGIC_AND_ODD_A_D-ANTIMAGIC_GRAPHS

[4106] V. Vilfred and L. M. Florida, Relation between sum, integer sum, chromatic and edge chromatic numbers of few graphs, unpublished.

[4107] V. Vilfred and L. M. Florida, Two problems on integral sum graphs, *Internat. J. Innovations Eng. Tech. (IJIET)*, **14**(4) 057-060. https://ijiet.com/wp-content/uploads/2019/12/8.pdf

[4108] V. Vilfred and L. M. Florida, Anti-sum and anti-integral sum graphs, unpublished.

[4109] V. Vilfred and L. M. Florida, Sum number and exclusiveness of graphs C_4, L_n and $P_3 \times P_3$, *Proceedings Inter. Conf. Math. Eng. Business Management* Stella Maris College, Chennai, India. (2012) 13-15.

[4110] V. Vilfred and T. Nicholas, On integral sum graphs G with $\Delta(G) = |V(G)| - 1$, unpublished.

[4111] V. Vilfred and T. Nicholas, Banana trees and unions of stars are integral sum graphs, *Ars Combin.,* **102** (2011) 79-85. https://combinatorialpress.com/article/ars/Volume%20102/volume-102-paper-7.pdf

[4112] V. Vilfred and T. Nicholas, Amalgamation of integral sum graphs, fans and Dutch M-windmills aredre integral sum graphs, National Seminar on Algebra and Discrete Mathematics held at Kerala Univ., Trivandrum, India, 2005, personal communication. https://www.researchgate.net/publication/313561463_Amalgamation_of_Integral_Sum_Graphs_Fan_and_Dutch_M-Windmill_are_Integral_Sum_Graphs

[4113] R. Vimal Kumar and D. Vijayalakshmi, V-super vertex magic labeling of some families of graphs, *Palest. J. Math.,* **10** (2021), Special Issue II, 191-196. https://pjm.ppu.edu/sites/default/files/papers/PJM_Special_Issue_July_2021_191to_196.pdf

[4114] Le Anh Vinh, Divisor graphs have arbitrary order and size, arXiv:0606483v1 [math.CO] 20 Jun 2006.

[4115] V. Vilfred, S. Somasundaram, and T. Nicholas, Classes of prime graphs, *International J. Management and Systems*, **18**(2) (2002).

[4116] V. Viola and T. Nicholas, A study on radio mean Dd-distance number of some basic graphs, *Internat. J. Appl. Engin. Res.,* **14**(8) (2019) (Special Issue) 77-80. https://www.ripublication.com/ijaerspl2019/ijaerv14n8spl_15.pdf

[4117] V. Vishnupriya, K. Manimekalai, and J. Baskar Babujee, Edge bimagic total labeling for some trees, *Acta Cienc. Indica Math.,* **35** (2009) 569-573.

[4118] M. F. Wahyujati and Y. Susanti, Computing the total H-irregularity strength of edge comb product of graphs, *Analele Stiintifice ale Universitatii Ovidius Constanta, Seria Matematica*, **13**(2), (2023) 177–189. https://www.researchgate.net/publication/362715105_Computing_the_Total_H-irregularity_Strength_of_Edge_Comb_Product_of_Graphs

[4119] C. D. Wallace, Mod Sum Numbers of Complete Bipartite Graphs, M. S. Thesis, East Tennessee State University, 1999.

[4120] W. D. Wallis, Magic Graphs, Birkhäuser, Boston, 2001.

[4121] W. D. Wallis, Vertex magic labelings of multiple graphs, *Congr. Numer.,* **152** (2001) 81-83.

[4122] W. D. Wallis, Two results of Kotzig on magic labelings, *Bull. Inst. Combin. Appl.,* **36** (2002) 23-28.

[4123] W. D. Wallis, Totally magic labellings and graphs, unpublished.

[4124] W. D. Wallis, E. T. Baskoro, M. Miller, and Slamin, Edge-magic total labelings, *Australas. J. Combin.,* **22** (2000) 177-190. https://ajc.maths.uq.edu.au/pdf/22/ocr-ajc-v22-p177.pdf

[4125] C. Wang, M. J. A. Khan, M. Ibrahim, E. Bonyah, M. K. Siddiqui, and S. Khalid, On edge irregular reflexive labeling for generalized prism, *J. Math.,* 2022, Art. ID 2886555, 7 pp. https://downloads.hindawi.com/journals/jmath/2022/2886555.pdf

[4126] G. Wang, On the number of graceful trees, *Internat. J. Math. Combin.,* **1** (2008) 91-101. https://core.ac.uk/download/pdf/144791728.pdf

[4127] J.-G. Wang, D. J. Jin, X.-G Lu, and D. Zhang, The gracefulness of a class of lobster trees, *Math. Comput. Modelling*, **20** (1994) 105-110. https://www.sciencedirect.com/science/article/pii/0895717794001677

[4128] H. Wang, The sum numbers and the integral sum numbers of the graph $K_{n+1}\setminus E(K_{1,r})$, *Discrete Math.,* **309** (2009) 4137-4143. https://www.sciencedirect.com/science/article/pii/S0012365X08007073

[4129] H. Wang and J. Gao, The sum numbers and the integral sum numbers of $\overline{P_n}$ and $\overline{F_n}$, *Ars Combin.,* **96** (2010) 9-31. https://combinatorialpress.com/article/ars/Volume%20096/volume-96-paper-2.pdf

[4130] H. Wang and J. Gao, The sum numbers and the integral sum numbers of $\overline{C_n}$ and $\overline{W_n}$, *Ars Combin.,* **96** (2010) 479-488. https://combinatorialpress.com/article/ars/Volume%20096/volume-96-paper-39.pdf

[4131] H. Wang and P. Li, Some results on exclusive sum graphs, *J. Appl. Math. Compu.,* **34** (2010) 343-351.

[4132] H. Wang and C. Li, New results on sum graph theory, *Ars Combin.,* **106** (2012) 509-519. https://combinatorialpress.com/article/ars/Volume%20106/volume-106-paper-45.pdf

[4133] H. Wang, C. Li, and B. Wei, Some results on integral sum graphs with no saturated vertices, *Util. Math.,* **97** (2015) 287-294.

[4134] H. Wang, Y. Ji, and P. Li, The sum numbers and integral sum numbers of the graph $K_n \backslash E(C_{n-1})$, *Ars Comb.,* **101** (2011) 15-26. https://combinatorialpress.com/article/ars/Volume%20101/volume-101-paper-2.pdf

[4135] H. Wang, J. Xu, M. Ma, and H. Zhang, A new type of graphical passwords based on odd-elegant labelled graphs, *Security and Communication Networks,* **2018**, Article ID 9482345, 11 pages, https://www.researchgate.net/publication/324557314_A_New_Type_of_Graphical_Passwords_Based_on_Odd-Elegant_Labelled_Graphs

[4136] H. Wang, J. Xu, and B. Yao. Twin odd-graceful trees towards information security, *Procedia Comput. Sci.,* **107** (2017) 15-20. doi:10.1016/j.procs.2017.03.050

[4137] H. Wang, J. Xu, and B. Yao, Exploring new cryptographical construction of complex network data, IEEE First International Conference on Data Science in Cyberspace, *IEEE Comput. Soc.,* (2016) 155-160.

[4138] H. Wang, J, Xu, and B. Yao, On generalized total graceful labelling of graphs, *Ars Combin.,* **140** (2018) 359-371.

[4139] H. Wang, X. Xu, Y. Yang, B. Zhang, M. Luo, and G. Wang, Radio number of ladder graphs, *Int. J. Comput. Math.,* **88** (2011) 2026-2034. https://www.researchgate.net/publication/220661587_Radio_number_of_ladder_graphs

[4140] H. Wang, B. Yao, and M. Yao, Generalized edge-magic total labellings of models from researching networks, *Inform. Sci.*, **279** (2014) 460-467. https://2024.sci-hub.se/2473/61edcb29db319cad8635dc31c7c77c6b/wang2014.pdf

[4141] J.-W. Wang, Vertex ranking of graphs, Master Thesis, Department of Applied Math., National Chiao-Tung University, June 1997.

[4142] S. Wang and Y. Gao, Product antimagic labeling of caterpillars, *J. Math.* (2021) Art. ID 3493941, 4 pp. https://www.hindawi.com/journals/jmath/2021/3493941/

[4143] T.-L. Wong and X. Zhu, Antimagic labelling of vertex weighted graphs, *J. Graph Theory,* **70**(3) (2012) 348-359.

[4144] T.-M. Wang, Toroidal grids are anti-magic, *Computing and combinatorics, Lecture Notes in Comput. Sci.*, **3595**, Springer, Berlin (2005) 671-679. https://www.researchgate.net/publication/221427243_Toroidal_Grids_Are_Anti-magic

[4145] T.-M. Wang and C.-C. Hsiao, On anti-magic labeling for graph products, *Discrete Math.*, **308** (2008) 3624-3633. https://www.researchgate.net/publication/220184812_On_anti-magic_labeling_for_graph_products

[4146] T.-M. Wang, C.-C. Hsiao, S.-M. Lee, A note on edge-graceful spectra of the square of paths, *Discrete Math.,* **308** (2008) 5878-5885. https://www.sciencedirect.com/science/article/pii/S0012365X07008552?via%3Dihub

[4147] T.-M. Wang, C.-C. Hsiao and S.-M. Lee, On anti-magic labeling of graphs products, *Discrete Math.*, **308** (16) 3624-3633. https://www.researchgate.net/publication/220184812_On_anti-magic_labeling_for_graph_products

[4148] T.-M. Wang and S. M. Lee, On edge-graceful spectrum of bicyclic graphs, *Congr. Numer.,* **205** (2010) 161-173.

[4149] T.-M. Wang, C.-M. Lin, and S.-M. Lee, On edge-balance index sets of regular graphs, The 26th Workshop on Combinatorial Mathematics and Computation Theory, 218-223. https://www.academia.edu/51218703/On_edge_balance_index_sets_of_regular_graphs

[4150] T.-M. Wang, C.-M. Lin, and S.-M. Lee, Edge-balance index sets of prisms and Möbius ladders, unpublished.

[4151] T. Wang and D. M. Li, Gracefulness of some special graph classes, *J. Wuhan Univ. Natur. Sci. Ed.,* **58**(5) (2012) 437-440.

[4152] T. Wang, D. Li, and Q. Wang, Some classes of antimagic graphs with regular subgraphs, *Ars Combin.,* **111** (2013) 241-250. https://combinatorialpress.com/article/ars/Volume%20111/volume-111-paper-19.pdf

[4153] T. Wang, H. S. Liu, and D. M. Li, Gracefulness of graphs related to wheels, *Acta Sci. Natur. Univ. Sunyatseni* **50**(6) (2011) 16-19.

[4154] T. Wang, M. J. Liu, and D. M. Li, Some classes of disconnected antimagic graphs and their joins, *Wuhan Univ. J. Nat. Sci.,* **17**(3) (2012) 195-199.

[4155] T. Wang, M. J. Liu, and D. M. Li, A class of antimagic join graphs, *Acta Math. Sin. (Engl. Ser.),* **29**(5) (2013) 1019-1026.

[4156] T. Wang, W. Miao, and D. Li, Deming, Antimagic graphs with even factors, *Wuhan Univ. J. Nat. Sci.,* **20**(3) (2015) 193-196. https://sci-hub.st/10.1007/s11859-015-1080-2

[4157] T. Wang, Q. Wang, and D. M. Li, Gracefulness of unconnected graphs $(P_3 + \overline{K_m} \cup G$ and $(C_3 + \overline{K_m} \cup G$, *Natur. Univ. Sunyatseni,* **51**(5) (2012) 54-57, 62. https://www.researchgate.net/publication/266958868_The_gracefulness_of_unconnected_graphs_P_3_K_m_G_and_C_3_K_m_G

[4158] T. Wang, Q. Wang, and D. M. Li, Gracefulness of disconnected graph $W_m^{(k)} \cup G$, *J. Hefei Univ. Technol. Nat. Sci.,* **35**(7) (2012) 987-990.

[4159] T. Wang, S.-F. Yan, D. M. Li, Graceful labeling of the union of some graphs, *Math. Pract. Theory,* **42**(16) (2012) 207-212.

[4160] T.-M. Wang, C.-C. Yang, L.-H. Hsu, and E. Cheng, Infinitely many equivalent versions of the graceful tree conjecture, *Appl. Anal. Discrete Math.,* **9**(1) (2015) 1-12. https://scispace.com/pdf/infinitely-many-equivalent-versions-of-the-graceful-tree-2dreue04w2.pdf

[4161] T.-M. Wang and G.-H. Zhang, Note on E-super vertex magic graphs, *Discrete Appl. Math.,* **178** (2014) 160-162. https://www.researchgate.net/publication/265603826_Note_on_E-super_vertex_magic_graphs

[4162] T.-M. Wang and G.-H. Zhang, On antimagic labeling of odd regular graphs, *Lecture Notes in Comput. Sci.,* S. Arumugam and B. Smyth (Eds.): IWOCA 2012, LNCS 7643 162-168, Springer-Verlag Berlin Heidelberg 2012. https://www.researchgate.net/publication/235942683_On_Antimagic_Labeling_of_Odd_Regular_Graphs

[4163] T.-M. Wang and G.-H. Zhang, On vertex magic total labeling of disjoint union of sun graphs, *Util. Math.,* **103** (2017) 289-298.

[4164] T.-M. Wang and G.-H. Zhang, On edge-graceful regular graphs with particular 3-factors, *Elect. Notes Discr. Math.,* **63** (2017) 33-40. https://www.academia.edu/81755260/On_Edge_Graceful_Regular_Graphs_with_Particular_3_Factors

[4165] T.-M. Wang and G.-H. Zhang, On edge-graceful labeling and deficiency for regular graphs, *AKCE Int. J. Graphs Comb.,* **15**(1) (2018) 105-111. https://www.tandfonline.com/doi/full/10.1016/j.akcej.2018.03.002#d1e198

[4166] W. Wang, M. Zhang, H. Yu, and D. Shi, The mod sum number of even fans and symmetric complete bipartite graphs, *Ars Combin.*, **94** (2010) 445-457. https://combinatorialpress.com/article/ars/Volume%20094/volume-94-paper-42.pdf

[4167] X. Wang and Y. Chang, Further results on $(v, 4, 1)$-perfect difference families, *Discrete Math.,* **310** (2010) 1995-2006. https://www.sciencedirect.com/science/article/pii/S0012365X10000968

[4168] Y. Wang and B. Liu, The sum number and integral sum number of complete bipartite graphs, *Discrete Math.,* **239** (2001) 69-82. https://www.sciencedirect.com/science/article/pii/S0012365X01001959

[4169] W. Wannasit and S. El-Zanati, On free α-labelings of cubic bipartite graphs, *24th MCCCC. J. Combin. Math. Combin. Comput.,* **82** (2012) 269-293. https://combinatorialpress.com/article/jcmcc/Volume%20082/vol-082-paper%2018.pdf

[4170] M. E. Watkins, A theorem on Tait colorings with an application to the generalized Petersen graphs, *J. Combin. Theory,* **6** (1969) 152-164. https://www.sciencedirect.com/science/article/pii/S002198006980116X?via%3Dihub

[4171] L. X. Wei and Z. Z. Jia, The gracefulness of unconnected graphs $G_1 \cup G_2$ and $G_1 \cup G_2 \cup K_2$, *Acta Math. Appl. Sin.,* **28** (2005) 689-694.

[4172] L. X. Wei, S. F. Yan, and K. L. Zhang, The researches on gracefulness of two kinds of unconnected graphs, *J. Shandong Univ. Nat. Sci.* **43** (2008) 90-96.

[4173] L. X. Wei and K. L. Zhang, Researches on graceful graphs $(P_1^{(1)} \vee P_n) \cup (P_1^{(2)} \vee P_{2n})$ and $(P_2 \vee \overline{K}_n) \cup G_{n-1}$. *J. Hefei Univ. Tech.*, **31** (2008) 276-279.

[4174] J. Wei, T, Wang, and C. Sun, Gracefulness of several unconnected graphs with wheel, *Wuhan Univ. J. Nat. Sci.*, **21**(3) (2016) 199-203.

[4175] L. X. Wei and K. L. Zhang, Graceful graphs on several kinds of union graphs, *Acta Sci. Natur. Univ. Sunyatseni,* **47** (2008) 10-13.

[4176] Y.-H. Wen, Necessary and sufficient condition for the class of $(P, P+1)$-graph to be balanced, *J. Lanzhou Univ. Nat. Sci.,* **43** (2007) 104-106.

[4177] Y.-H. Wen, Friendliness of generalized wheel, *J. Lanzhou Univ. Nat. Sci.,* **44** (2008) 103-108.

[4178] Y. Wen and S. M. Lee, On Eulerian graphs of odd sizes which are fully magic, unpublished.

[4179] Y. Wen, S.-M. Lee, H.-H. Su, On $Z_2 \oplus Z_2$-magic graphs, *Congr. Numer.*, **199** (2009) 13-31.

[4180] Y. Wen, S. M. Lee, and H. Sun, On the supermagic edge-splitting extension of graphs, *Ars Combin.,* **79** (2006) 115-128. https://combinatorialpress.com/article/ars/Volume%20079/volume-79-paper-10.pdf

[4181] R. W. Whitty, Rook polynomials on 2-dimensional surfaces and graceful labellings of graphs, *Discrete Math.,* **308** (2008) 674-683. https://www.sciencedirect.com/science/article/pii/S0012365X07005079

[4182] T. Wichianpaisarn and U. Mato, Star-supermagic decompositions of the complete bipartite graph minus a one-factor, *Internat. J. Math. Math. Sci.,* (2017) Article ID 5104701, 4 pp. https://onlinelibrary.wiley.com/doi/10.1155/2017/5104701

[4183] K. Wijaya, S. N. Aulia, I. Halikin, and K. Kusbudiono, An inclusive distance irregularity strength of n-ary tree, *JTAM (J. Teori dan Aplikasi Matematika),* **8**(2) (2024) 568-577. journal.ummat.ac.id/index.php/jtam/article/view/20549

[4184] K. Wijaya and E. Baskoro, Edge-magic labelings of a product of two graphs (in Indonesian), *Proc. Seminar MIPA*, ITB Bandung Indonesia, October 2000.

[4185] K. Wijaya, Slamin, Surahmat, and S. Jendroľ, Total vertex irregular labeling of complete bipartite graphs, *J. Combin. Math. Combin. Comput.,* **55** 129-136 https://combinatorialpress.com/article/jcmcc/Volume%20055/vol-055-paper%2012.pdf?utm_source=chatgpt.com

[4186] D. E. Wijayanti, N. Hidayat, D. Indriati, A. R. Alghofari, and S. Slamin, Total distance vertex irregularity strength of some corona product graphs, *Elect. J. Graph Th. Appl.*, **11**(1) (2023) 209. https://www.ejgta.org/index.php/ejgta/article/view/1473/pdf_273

[4187] D. E. Wijayanti, H. Noor, D. Indriati, A. R. Alghofari, and Slamin, On distance vertex irregular total k-labeling, *Sci. Tech. Indonesia*, **8**(3) (2023) 479–485. https://doi.org/10.26554/sti.2023.8.3.479-485

[4188] D. E. Wijayanti, H. Noor, D. Indriati, and A. R. Alghofari, The total distance vertex irregularity strength of fan and wheel graphs, AIP Conference Proceedings, 2326 (2021), 020043-1- 020043-14. https://aip.scitation.org/doi/pdf/10.1063/5.0039904

[4189] D. E. Wijayanti, H. Noor, D. Indriati, A. R. Alghofari, and Slamin, On d-distance vertex irregular total k-labeling, submitted (2021)

[4190] I. Wijayanti, D. E. Wijayanti, and S. Surono, Total distance vertex irregularity strength of hairy cycle C_m^n graph, *J. Data Science,* **2022:09** https://iuojs.intimal.edu.my/index.php/jods/article/view/48/45

[4191] H. S. Wilf and N. A. Yoshimura, Ranking rooted trees, and a graceful application, in *D*iscrete Algorithms and Complexity Proceedings of the Japan-US Joint Seminar, June 4–6, 1986, Kyoto, Japan, edited by D. S. Johnson, et al. 341-349.

[4192] A. William and C. R. Kenneth, Radio antipodal number of certain graphs, *J. Comp. Math. Sci.*, (2011) 868-872. https://www.researchgate.net/publication/350695685_Radio_Number_of_Some_Path_Related_Graph

[4193] K. J. Wilson and J. V. Vivik, Mean edge cordial labeling of path structured graphs with applications, *Panamerican Math. J.*, **35**(4s) (2025) 1894-1910. https://internationalpubls.com/index.php/pmj/article/view/6753/3840

[4194] K. J. Wilson K and J. V. Vivik, Mean edge cordial labeling of wheel related graphs, *Proceed. Internat. Confer. Environ. Sustainability Math. and Comput. Intell.*, (2026) https://www.academia.edu/166052339/MEAN_EDGE_CORDIAL_LABELING_OF_WHEEL_RELATED_GRAPHS

[4195] K. J. Wilson and J. V. Vivik, Mean edge cordial labeling of flower family graphs, *Internat. Conf. Environ. Sustainability Math. Comput. Intelligence*, – ICESMCI '26 https://doi.org/10.5281/zenodo.19626897

[4196] K. J. Wilson and J. V. Vivik, On mean edge cordial labeling of some cycle-related graphs, personal communication.

[4197] K. J. Wilson and J. V. Vivik, On Mean edge cordial labeling of path-derived graphs, personal communication.

[4198] K. J. Wilson and J. V. Vivik, Modeling biometric sensor arrays through mean edge cordial labeling of grid graphs, personal communication.

[4199] K. J. Wilson and J. V. Vivik, $m \times n$ grid graphs under mean edge cordial labeling for portrait pixel connectivity, personal communication.

[4200] K. J. Wilson and J. V. Vivik, Mean edge cordial labeling of ladder structured graphs, personal communication.

[4201] K. J. Wilson and J. V. Vivik, Mean edge cordial labeling in path structured irrigation networks, personal communication.

[4202] L. K. Wilson and V. M. Bebincy, Studies on some wheel related Zumkeller graphs, *AIP Conference Proceedings* (2022) 130049.

[4203] L. K. Wilson and V. M. Bebincy, Zumkeller labeling of some path related graphs, *Malaya J. Math.,* **9**(1) (2021) 519-523. https://www.malayajournal.org/articles/MJM09010087.pdf

[4204] L. K. Wilson and H. Jini, Prime labeling of torch graph, *Malaya J. Mat.,* **9**(1) (2021) 890–895. https://www.malayajournal.org/articles/MJM09010157.pdf

[4205] S. Wilson and A. Riskin, Edge-graceful labellings of odd cycles and their products, *Bulletin of the ICA,* **24** (1998) 57-64.

[4206] S. Winters, personal communication (circa 1990).

[4207] J. Wojciechowski, Long Induced Cycles in the Hypercube and Colourings of Graphs, Ph. D. Thesis, Cambridge University, England, 1990.

[4208] J. Wojciechowski, Equitable labelings of cycles, *J. Graph Theory*, **17** (1993) 531-547.

[4209] F. A. Winarti and D. Rahmadani, Odd harmonious labeling on prism snake graphs and generalized prism chain snake graphs, *J. Physics: Conf. Ser.*, 3276 (2026) 012002. https://iopscience.iop.org/article/10.1088/1742-6596/3276/1/012002

[4210] T. Wong and X. Zhu, Antimagic labelling of vertex weighted graphs, *J.* Graph Theory, **70**(3) (2012) 348-350.

[4211] D. Wood, On vertex-magic and edge-magic total injections of graphs, *Australas. J. Combin.,* **26** (2002) 49-63. https://ajc.maths.uq.edu.au/pdf/26/ajc_v26_p049.pdf

[4212] J. Wu, J. Mao, and D. Li, New types of integral sum graphs, *Discrete Math.,* **260** (2003) 163-176. https://www.sciencedirect.com/science/article/pii/S0012365X02004569

[4213] S.-L. Wu, Graceful labelings of graphs associated with vertex-saturated graphs, *Ars Combin.*, **62** (2002) 109-120. https://combinatorialpress.com/article/ars/Volume%20062/volume-62-paper-11.pdf

[4214] S.-L. Wu, New families of sequential graphs, *Ars Combin.,* **69** (2003) 9-17. https://combinatorialpress.com/article/ars/Volume%20069/volume-69-paper-2.pdf

[4215] S.-L. Wu, New graceful families on bipartite graphs, *Ars Combin.,* **73** (2004) 79-87. https://combinatorialpress.com/article/ars/Volume%20073/volume-73-paper-9.pdf

[4216] S.-L. Wu, A necessary condition for the existence of an α-labeling, unpublished.

[4217] S.-L. Wu, Graceful labelings of vertex-saturated graphs and related graphs, unpublished.

[4218] S.-L. Wu and H.-C. Lu, On the constructions of new families of graceful graphs, *Ars Combin.,* **106** (2012) 235-246. https://combinatorialpress.com/article/ars/Volume%20106/volume-106-paper-20.pdf

[4219] E. Wulandari and Y. Susanti, On total edge irregularity strength of arithmetic ladder graphs, *AIMS Math,* **11**(4) (2026) 10175–10190. https://www.aimspress.com/article/doi/10.3934/math.2026420

[4220] C. Wyels and M. Tomova, Radio Labeling Cartesian graph products, 9th Cologne-Twente Workshop on Graphs and Combinatorial Optimization, Cologne,Germany, May 25-27, 2010. Extended Abstracts (2010) 163–167. https://www.zpr.uni-koeln.de/AFS/conferences/CTW2010/CTW%202010%20Band/Kompilierfertiger%20CTW2010%20Band/39-(P163-167)-21.pdf

[4221] D. A. Xavier and R. Rathi, Proper lucky number of hexagonal mesh and honeycomb network, *Math. Internat. J. Math. Trends Tech.,* **48** (2017) 223-238. doi:10.14445/22315373/IJMTT-V48P532

[4222] D. A. Xavier and R. C. Thivyarathi, Radio antipodal mean number of certain graphs, *Internat. J. Math. Trends and Tech.*, **54**(6) (2018) Article Id. IJMTT-V54P556. https://ijmttjournal.org/public/assets/volume-54/number-6/IJMTT-V54P556.pdf

[4223] Y. Xi, Y. Yang, Mominul, and L. Wang, Super vertex-magic total labelings of $W_{3,n}$, *Ars Combin.,* **86** (2008) 121-128. https://combinatorialpress.com/article/ars/Volume%20086/volume-86-paper-12.pdf

[4224] Y. Xi, Y. Yang, and L. Wang, On harmonious labeling of the double triangular snake, *Indian J. Pure Appl. Math.,* **39** (2008) 177-184.

[4225] L. T. Xie and G. Z. Liu, A survey of the problem of graceful trees, *Qufu Shiyuan Xuebao,* (1984) 8-15.

[4226] J. Xie, T. Zhao, and B. Yao, The gracefulness of super sun-graphs, *2018 IEEE 3rd Adv. Infor. Tech., Elect. Automa. Control Conf.,* (IAEAC) (2018) 2098-2109. doi:10.1109/IAEAC.2018.8577870

[4227] B. Xu, On integral sum graphs, *Discrete Math.,* **194** (1999) 285-294. https://www.sciencedirect.com/science/article/pii/S0012365X98001502

[4228] S. D. Xu, Cycles with a chord are harmonious, *Mathematica Applicata*, **8** (1995) 31-37.

[4229] S. D. Xu, Harmonicity of triangular snakes, *J. Math. Res. Exposition*, **15** (1995) 475-476.

[4230] X. Xu, Jirimutu, L. Wang, and Z. Min, On the gracefulness of the digraphs n-$\vec{C}_m$, *Ars Combin.,* **92** (2009) 419-428. https://combinatorialpress.com/article/ars/Volume%20092/volume-92-paper-28.pdf

[4231] X. Xu, J.-M. Xu, M. Lü, Z. Baosheng, and C. Nan, On (a, d)-antimagic labelings of generalized Petersen graphs, *Ars Combin.,* **90** (2009) 411-421. https://scispace.com/pdf/on-a-d-antimagic-labelings-of-generalized-petersen-graphs-2qu47d3z6t.pdf

[4232] X. Xu, Y. Yang, H. Li, and Y. Xi, The graphs $C_{11}^{(t)}$ are graceful for $t \equiv 0, 1 \pmod 4$. *Ars Combin.*, **88** (2008) 429-435. https://combinatorialpress.com/article/ars/Volume%20088/volume-88-paper-35.pdf

[4233] X. Xu, Y. Yang, L. Han, and H. Li, The graphs $C_{13}^{(t)}$ are graceful for $t \equiv 0, 3 \pmod 4$. *Ars Combin.,* **90** (2009) 25-32. https://combinatorialpress.com/article/ars/Volume%20090/volume-90-paper-2.pdf

[4234] X. Xu, Y. Yang, Y. Xi, K. M. M. Haque, and L. Shen, Super edge-magic labelings of generalized Petersen graphs $P(n, 3)$, *Ars Combin.,* **85** (2007) 19-31. https://combinatorialpress.com/article/ars/Volume%20086/volume-86-paper-3.pdf

[4235] X. Xu, Y. Yang, Y. Xi, and H. Li, On (a, d)-antimagic labelings of generalized Petersen graphs $P(n, 3)$, *Ars Combin.,* **86** (2008) 23-31. https://combinatorialpress.com/article/ars/Volume%20086/volume-86-paper-3.pdf

[4236] L. Yahyaei and S. A. Katre, Representation number of a caterpillar, *J. Combin. Math. Combin. Comput.,* **101** (2017) 47-57. https://combinatorialpress.com/article/jcmcc/Volume%20101/vol-101-paper%205.pdf

[4237] Q. T. Yan, A proof of a conjecture related to of windmill graphs, *Math. Practice Theory,* **34** (2004) 116-117.

[4238] Q. T. Yan, The graceful labelings of $P_{2r,2m}$, *J. Systems Sci. Math. Sci.,* **26**(5) (2006) 513-517.

[4239] Q. T. Yan, Odd gracefulness and odd strongly harmoniousness of the product graphs $P_n \times P_m$, *J. Systems Sci. Math. Sci.,* **30** (2010) 341-348.

[4240] W. Yan and B. Liu, Some results on integral sum graphs, *Discrete Math.,* **240** (2001) 219-229. https://www.sciencedirect.com/science/article/pii/S0012365X00001187

[4241] W. Yan and B. Liu, The sum number and integral sum number of complete bipartite graphs, *Discrete Math.,* **239**(1-3) (2001) 69-82. https://www.sciencedirect.com/science/article/pii/S0012365X01001959

[4242] D. Yang, A note on antimagic orientations of even regular graphs, *Discrete Appl. Math.,* **267** (2019) 224-228. https://doi.org/10.48550/arXiv.1811.01904

[4243] X. Yang, H. Bian, H. Yu, and D. Liu, The local antimagic total chromatic number of some wheel-related graphs, *Axioms* **11**(3) (2022) 97 18 pp. https://www.mdpi.com/2075-1680/11/3/97

[4244] D. Yang, J. Carlson, A. Owens, K. E. Perry, I. Singgih, Z.-X. Song, F. Zhang, X. Zhang, Antimagic orientations of graphs with large maximum degree, *Discrete Math.,* **343**(12) (2020) 112-23. arXiv:1908.06072

[4245] H. Yang, M. A. Rashid, S. Ahmad, M. K. Siddiqui, and M. F. Hanif, Cycle super magic labeling of pumpkin, octagonal and hexagonal graphs, *J. Discrete Math. Sci. Cryptogr.,* **22**(7) (2019) 1165-1176.

[4246] H. Yang, M. K. Siddiqui, M. Ibrahim, S. Ahmad, and A. Ahmad, Computing the irregularity strength of planar graphs, *Mathematics,* (2018) 6, 150. https://www.mdpi.com/2227-7390/6/9/150

[4247] X.-W Yang and W. Pan, Gracefulness of the graph $\bigcup_{i=1}^{n} F_{m_i,4}$, *J. Jilin Univ. Sci.,* **41** (2003) 466-469.

[4248] Y. Yang, X. Lin, C. Yu, The graphs $C_5^{(t)}$ are graceful for $t \equiv 0, 3 \pmod 4$, *Ars Combin.* **74** (2005) 239-244. https://combinatorialpress.com/ars-articles/volume-074-ars-articles/the-graphs-c_5t-are-graceful-for-t-equiv-03-mod-4/

[4249] Y. Yang, W. Lu, and Q. Zeng, Harmonious graphs $C_{2k} \cup C_{2j+1}$, *Util. Math.,* **62** (2002) 191-198.

[4250] Y. S. Yang, Q. Rong, and X. R. Xu, A class of graceful graphs, *J. Math. Research and Exposition,* **24** (2004) 520-524.

[4251] Y. C. Yang and X. G. Wang, On the gracefulness of the product $C_n \times P_2$, *J. Math. Research and Exposition,* **1** (1992) 143-148.

[4252] Y. C. Yang and X. G. Wang, On the gracefulness of the union of two stars and three stars, *Combinatorics, Graph Theory, Algorithms and Applications (Beijing, 1993)*, 417-424, World Sci. Publishing, River Edge, NJ, 1994.

[4253] Y. C. Yang and X. G. Wang, On the gracefulness of product graph $C_{4n+2} \times P_{4m+3}$, *Combinatorics, Graph Theory, Algorithms and Applications (Beijing, 1993)*, 425-431, World Sci. Publishing, River Edge, NJ, 1994.

[4254] Y. Yang, X. Xu, Y. Xi, H. Li, and K. Haque, The graphs $C_7^{(t)}$ are graceful for $t \equiv 0, 1 \pmod 4$, *Ars Combin.,* **79** (2006) 295-301. https://combinatorialpress.com/article/ars/Volume%20079/volume-79-paper-26.pdf

[4255] Y. Yang, X. Xu, Y. Xi, and H. Huijun, The graphs $C_9^{(t)}$ are graceful for $t \equiv 0, 3 \pmod 4$, *Ars Combin.,* **85** (2007) 361-368. https://combinatorialpress.com/article/ars/Volume%20085/vol-85-paper%2030A.pdf

[4256] Y. Yang, X. Xu, Y. Xi, and J. Qian, On harmonious labelings of the balanced quadruple shells. *Ars Combin.,* **75** (2005) 289-296. https://combinatorialpress.com/article/ars/Volume%20075/volume-75-paper-25.pdf

[4257] B. Yao, X. Chen, M. Yao, and H. Cheng, On (k, λ)-magically total labeling of graphs, *JCMCC*, **87** (2013) 237-253. https://combinatorialpress.com/jcmcc-articles/volume-087/on-k-lambda-magically-total-labeling-of-graphs/

[4258] B. Yao, H. Cheng, M. Yao, and M. Zhao, Meimei, A note on strongly graceful trees, *Ars Combin.,* **92** (2009) 155-169. https://combinatorialpress.com/article/ars/Volume%20092/volume-92-paper-12.pdf

[4259] B. Yao, H. Cheng, Z. Zhongfu, and M. Yao, Labellings of trees with larger maximal degrees, unpublished.

[4260] B. Yao, X. Liu and M. Yao, Connections between labellings of trees, *Bull. Iranian Math. Soc.*, **43**(2) (2017) 275-283. http://bims.iranjournals.ir/article_930_493991d8c291f08d8bd8ff563e3af03f.pdf

[4261] B. Yao, M. Yao, and H. Cheng, On gracefulness of directed trees with short diameters, *Bull. Malays. Math. Sci. Soc.,* (2) **35**(1) (2012) 133-146. https://www.kurims.kyoto-u.ac.jp/EMIS/journals/BMMSS/pdf/v35n1/v35n1p12.pdf

[4262] B. Z. Yatin, Z. Y. Awanis, and I. G. A. W. Wardhana, Super $(a, d) - P_2 \odot P_n$-antimagie total labeling of corona product of two paths, *JTAM (J. Teori dan Aplikasi Matematika)*, **8**(2) (2024) 454-464. https://journal.ummat.ac.id/index.php/jtam/article/view/20065/pdf

[4263] V. Yegnanarayanan, On some additive analogues of graceful theme: cycle-related graphs, *Southeast Asian Bull. Math.,* **23** (1999) 317-333.

[4264] V. Yegnanarayanan, On magic graphs. *Util. Math.,* **59** (2001) 181-204.

[4265] V. Yegnanarayanan and P. Vaidhyanathan, On nice $(1, 1)$ edge-magic graphs, *Electronic Notes Discr. Math.,* **33** (2009) 115-122.

[4266] K. Yenoke, A. Jose, and P. Venugopal, On the radio antipodal mean number of certain types of ladder graphs, *Internat. J. Innov. Res. Sci. Engin. Tech. (IJIRSET)*, **9**(6) 2020. https://www.ijirset.com/upload/2020/june/92_On_NC.PDF

[4267] K. Yenoke, P. Selvagopal, K. M. B. Smitha, and R. Cranston, Bounds for the radio number of mesh derived architecture, *Int. J. of Innovat. Res. Sci., Eng. Tech.*, **9**(6) (2020) 4806-4810. https://www.researchgate.net/publication/342623957_Bounds_for_the_Radio_Number_of_Mesh_Derived_Architecture

[4268] Z. B. Yilma, Antimagic properties of graphs with large maximum degree, *J. Graph Th.,* **72**(4) (2013) 367-373. https://www.math.cmu.edu/~zyilma/docs/zby_antimagiclargemaxdegree.pdf?utm_source=chatgpt.com

[4269] R. Yilmaz and I. Cahit, E-cordial graphs, *Ars Combin.*, **46** (1997) 251-266. https://combinatorialpress.com/ars-articles/volume-046-ars-articles/e-cordial-graphs/

[4270] K.-K. Yoong, R. Hasni, G.-C. Lau, M. A. Asim, and A. Ahmad, Reflexive edge strength of convex polytopes and corona product of cycle with path, *AIMS Math.,* **7**(7) (2022) 11784-11800. https://www.aimspress.com/article/doi/10.3934/math.2022657?viewType=HTM

[4271] K. K. Yoong, R. Hasni, G. C. Lau, and M. J. Irfan, Edge irregular reflexive labeling for some classes of plane graphs, *Malaysian J. Math. Sci.,* **16**(1) (2022) 25-36. https://doi.org/10.47836/mjms.16.1.03

[4272] L. You, J. Yang and Z. You, The maximal total irregularity of unicyclic graphs, *Ars Combin.,* **114** (2014) 153-160. *NOW Chat 10/13/26

[4273] M. Z. Youssef, On Graceful, Harmonious and Prime Labelings of graphs, Ph. D. Thesis, Department of Mathematics, Ain Shams University, 2000. NOW 9/27/25

[4274] M. Z. Youssef, New families of graceful graphs, *Ars Combin.,* **67** (2003) 303-311. https://combinatorialpress.com/article/ars/Volume%20067/volume-67-paper-22.pdf

[4275] M. Z. Youssef, Two general results on harmonious labelings, *Ars Combin.*, **68** (2003) 225-230. https://combinatorialpress.com/article/ars/Volume%20068/volume-68-paper-22.pdf

[4276] M. Z. Youssef, A necessary condition on k-equitable labelings, *Util. Math.,* **64** (2003) 193-195. https://www.researchgate.net/publication/268864563_A_necessary_condition_on_k-equitable_labelings

[4277] M. Z. Youssef, Pseudograceful labelings of cycles, *Ars Combin.,* **76** (2005) 241-247. https://combinatorialpress.com/article/ars/Volume%20076/volume-76-paper-21.pdf

[4278] M. Z. Youssef, On Skolem-graceful and cordial graphs, *Ars Combin.,* **78** (2006) 167-177. https://combinatorialpress.com/article/ars/Volume%20078/volume-78-paper-16.pdf

[4279] M. Z. Youssef, On k-cordial labelling, *Australas. J. Combin.*, **43** (2009) 31-37. https://ajc.maths.uq.edu.au/pdf/43/ajc_v43_p031.pdf?utm_source=chatgpt.com

[4280] M. Z. Youssef, Graph operations and cordiality, *Ars Combin.*, **97** (2010) 161-174. https://www.academia.edu/66651229/Graph_operations_and_cordiality

[4281] M. Z. Youssef, On E_k-cordial labeling, *Ars Combin.,* **104** (2012) 271-279. https://combinatorialpress.com/article/ars/Volume%20104/volume-104-paper-23.pdf

[4282] M. Z. Youssef, On k-equitable and k-balanced labeling of graphs, *Ars Combin.,* **108** (2013) 51-64. https://combinatorialpress.com/article/ars/Volume%20108/volume-108-paper-5.pdf

[4283] M. Z. Youssef, On α-labelling of disconnected graphs, *Ars Comb.,* **123** (2015) 329-338. https://www.researchgate.net/publication/286559726_On_alpha-labeling_of_disconnected_graphs

[4284] M. Z. Youssef, Note on even harmonious labeling of cycles, *Util. Math.,* **100** (2016) 137-142.

[4285] M. Z. Youssef, personal communication (2014).

[4286] M. Z. Youssef, On even harmonious labelings, *Util. Math.,* **110** (2019) 243-253.

[4287] M. Z. Youssef and N. A. Al-Kuleab, 4-cordiality of some regular graphs and the complete 4-partite graph, *J. Combin. Math. Combin. Comput.,* **76** (2011) 111-136. https://combinatorialpress.com/article/jcmcc/Volume%20076/vol-076-paper%208.pdf

[4288] M. Z. Youssef and N. A. Al-Kuleab, Some new results on k-cordial labeling, *Ars Combin.,* **126** (2016) 311-322. https://combinatorialpress.com/article/ars/Volume%20126/volume-126-paper-27.pdf

[4289] M. Z. Youssef and N. A. Al-Kuleab, 3-equitable labelings of ladders and graphs related to cycles, *Utilitas Math.*, **92** (2013) 261-274.

[4290] M. Z. Youssef and Z. S. Almoreed, On odd prime labeling of graphs, *Open J. Discret. Appl. Math.,* **3**(3) (2020) 33-40. https://pisrt.org/psrpress/j/odam/2020/3/5/on-odd-prime-labeling-of-graphs.pdf

[4291] M. Z. Youssef and E.A. El Sakhawi, Some properties of prime graphs, *Ars Combin.,* **84** (2007) 129-140. https://combinatorialpress.com/article/ars/Volume%20084/volume-84-paper-10.pdf

[4292] P. Yu, A proof of a conjecture about arithmetic graphs, *J. Math. Res. Exposition,* **16** (1996) 594-598.

[4293] P. Yu, Strongly arithmetic graphs, *Chinese Quart. J. Math.,* **15** (2000) 22-27.

[4294] T.-K. Yu, D. T. Lee, and Y.-X. Chen, Graceful and harmonious labelings on 2-cube, 3-cube and 4-cube snakes, unpublished.

[4295] Y.-H. Yu, W.-X. Wang, and L.-x. Song, The gracefulness of a kind of unconnected graphs, *Adv. Intelligent Soft Comput.,* **128** (2012) 525-531.

[4296] X. Yue, Y. Yuan-sheng, and M. Xin-hong, Skolem-gracefulness of k-stars, *Ars Combin.,* **98** (2011) 149-160. https://combinatorialpress.com/article/ars/Volume%20098/volume-98-paper-13.pdf

[4297] J. Yuan and W. Zhu, Some results on harmonious labelings of graphs, *J. Zhengzhou Univ. Nat. Sci. Ed.,* **30** (1998) 7-12.

[4298] Y. Yuansheng, X. Yue, X. Xirong, and M. Xinhong, Super edge magic labelings of book graphs B_n, unpublished. Abstract at https://www.researchgate.net/publication/268023455_Super_edge-magic_labelings_of_book_graphs_B_n

[4299] X. Yue, Y. Yuansheng, and W. Liping, One edge union of k shell graphs is cordial, *Ars Combin.,* **86** (2008) 403-408. https://combinatorialpress.com/article/ars/Volume%20086/volume-86-paper-37.pdf

[4300] O. A. Zahraa, E. A. Nabeel, and F. A. Fawzi, Dividing graceful labeling of certain tree graphs, *Tikrit J. Pure Sci.*, **25**(4) (2020) 123-126. https://www.tjpsj.org/index.php/tjps/article/view/281

[4301] A. Zak, Harmonious orders of graphs, *D*iscrete Math., **309** (2009) 6055-6064.n

[4302] D. Zala, N. Chotaliya, and M. A. Chaurasiya, Even-odd harmonious labeling of some graphs, *Internat. J. Innov. Tech. Exploring Eng.*, — Special Issue, **10**(4) (2021) 149-151. doi:10.35940/ijitee.d8513.0210421

[4303] T. M. Zalsa, D. Indriati, and T. S. Martini, Edge irregular reflexive labeling of dumbbell graph and corona of open ladder and null graph, *BAREKENG: J. Math. App.*, **18**(4) (2024) 2767-2778. https://ojs3.unpatti.ac.id/index.php/barekeng/article/view/13892/8916

[4304] L. A. Zalzabila, D. Indriati, and T. Martini, Edge irregular reflexive labelings on alternate triangular snake and double alternate quadrilateral snake, https://ajc.maths.uq.edu.au/pdf/69/ajc_v69_p394.pdf

[4305] M. R. Zeen El Deen, Edge even graceful labeling of some graphs, *J. Egyptian Math. Soc.,* **27**(20) (2019) 1-15. https://www.researchgate.net/publication/338712966_Edge_even_graceful_labeling_of_some_graphs

[4306] M. R. Zeen El Deen, Edge δ-graceful labeling for some cyclic-related graphs, *Adv. Math. Phys.* (2020) 6273245. https://onlinelibrary.wiley.com/doi/epdf/10.1155/2020/6273245?utm_source=chatgpt.com

[4307] M. R. Zeen El Deen, Edge δ-graceful labeling of some snake and chain related networks, *J. Discr. Math. Sci. Cryp.*, **29**(7) (2026) 2641–2665. https://tarupublications.com/journals/jdmsc/volume/29/issue/7/edge-d-graceful-labeling-of-some-snake-and-chain-related-networks/pdf

[4308] M. R. Zeen El Deen, W. A. Aboamer, and H. M. El Sherbiny, Super edge magic harmonious labeling for certain graphs, *Frontiers Sci. Res. Tech.*, **8** (2024) 47-56.

[4309] M. R. Zeen El Deen and G. Elmahdy, New classes of graphs with edge $\delta-$ graceful labeling, *AIMS Math.*, **7**(3) (2022) 3554-3589. https://www.aimspress.com/article/doi/10.3934/math.2022197?viewType=HTML

[4310] M. R. Zeen El Deen, G. M. Elmahdy, E. M. Elkholy, and H. M. El-Sherbiny, Half-Zumkeller labeling for some Cartesian product graphs, *Frontiers Sci. Res. Tech.*, **4** (2022) 99-110. doi:10.21608/FSRT.2022.169109.1072

[4311] M. R. Zeen El Deen and N. A. Omar, Further results on edge even graceful labeling of the join of two graphs, *J. Egyptian Math. Soc.,* **28**, article 21, (2020) https://doi.org/10.1186/s42787-020-00077-5

[4312] D. Zhang, Y-S. Ho, S. M. Lee, and Y. Wen, On the balance index sets of trees with diameter at most four, unpublished.

[4313] D. Zhang, S. M. Lee, and L. Wen, On the balancedness of galaxies with at most four stars, *J. Combin. Math. Combin. Comput.,* **50** (2004) 3-15.

[4314] F. Zhang, S. Nazeer, M. Habib, T. J. Zia, and Z. Ren, Radio number for generalized Petersen graphs $P(n, 2)$, *IEEE Access* (2019) PP(99):1-9. https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=8848394

[4315] G.-H. Zhang and T.-M. Wang, E-super vertex magic regular graphs of odd degree, The Eighth International Workshop on Graph Labelings (IWOGL 2014), *Electron. Notes Discrete Math.*, **48** (2015) 49-56. https://www.academia.edu/116835395/E_Super_Vertex_Magic_Regular_Graphs_of_Odd_Degree

[4316] L. Zhang, L. Zhou, and J. Wang, Proofs of non-edge-magic for several kinds of composite graphs about K_n, *IEEE 6th Adv. Inform. Management, Communicates, Elect. Automation Control Confer.,* (IMCEC),

[4317] P. Zhang, Radio labelings of cycles, *Ars Combin.,* **65** (2002) 21-32. https://combinatorialpress.com/article/ars/Volume%20065/volume-65-paper-2.pdf

[4318] X. Zhang, M. Cancan, M. F. Nadeem and M. Imran, Edge irregularity strength of certain families of comb graph, *Proyecciones* (Antofagasta), **39**(4) (2020) 787-797. https://www.scielo.cl/pdf/proy/v39n4/0716-0917-proy-39-04-787.pdf

[4319] X. Zhang, M. Ibrahim, S. ul Haq Bokhary, and M. K. Siddiqui, Edge irregular reflexive labeling for the disjoint union of gear graphs and prism graphs, *Mathematics,* **6**(9):142 (2018), 10 pages. https://www.mdpi.com/2227-7390/6/9/142

[4320] X. Zhang, C. Ye, S. Zhang, and B. Yao, Graph colorings and labelings having multiple restrictive conditions in topological coding, *Mathematics,* **10** 1592 (2022) 1-9. https://www.mdpi.com/2227-7390/10/9/1592

[4321] X. Zhang, H. Sun, and B. Yao, Graph theory towards module-K odd-elegant labelling of graphical passwords, MATEC Web of Conferences 139, 00206 (2017) *ICMITE* 2017 doi:10.1051/matecconf/201713900206

[4322] Y. Zhang and H. Wang, Some new results on sum index and difference index, *AIMS Math.*, **8**(11) (2023) 26444–26458. https://www.aimspress.com/article/doi/10.3934/math.20231350

[4323] Z. Zhang, T. Mehmood, A. ur Rehman, M. Hussain, and X. Zhang, On valuation of edge irregularity strength of certain graphical families, *Hindawi J. Math.,* 2022, Article ID 3230932, 8 pages https://doi.org/10.1155/2022/3230932

[4324] Z. Zhang, M. Naeem, A. Tariq, and W. Zhao, On square sum labeling of two families of Petersen graphs, *J. Math.* (2022), Art. ID 1872695, 145 19-523. https://www.hindawi.com/journals/jmath/2022/1872695/

[4325] Z. Zhang, Q. Zhang, and C. Wang, On $(a, b; n)$-graceful labeling of path P_n, *Ars Combin.,* **132** (2017) 105-119. https://combinatorialpress.com/article/ars/Volume%20132/volume-132-paper-9.pdf

[4326] L. Zhao, W. Feng, and Jirimutu On the gracefulness of the digraphs $n - \vec{C}_m$, *Util. Math.,* **82** (2010) 129-134.

[4327] L. Zhao and Jirimutu, On the gracefulness of the digraph $n - \vec{C}_m$ for $m = 5, 7, 9, 11, 13$, *J. Prime Res. Math.,* **12**(4)Classes of prime graphs, (2008) 118-126.

[4328] L. Zhao, Siqintuya, and Jirimutu, On the gracefulness of the digraphs $n - \vec{C}_m$, *Ars Combin.,* **99** (2011) 421-428.https://combinatorialpress.com/article/ars/Volume%2099/volume-99-paper-36.pdf

[4329] S. L. Zhao, All trees of diameter four are graceful, Graph Theory and its Applications: East and West (Jinan, 1986), 700-706 *Ann. New York Acad. Sci.,* 576, New York Acad. Sci., New York, 1989.

[4330] X. Y. Zhao, J. Shao, B. Yao, and X. Wang, A class of trees having strongly graceful labellings, *J. Jilin Univ. Sci.*, **54**(2) (2016) 222-228. https://www.researchgate.net/publication/316049803_A_Class_of_Trees_Having_Strongly_Super_Total_Graceful_Labellings

[4331] Y. Zhao, personal communication (2010).

[4332] G. Zhenbin, The labelings of a variation of banana trees, *Ars Combin.*, **94** (2010) 175-181. https://combinatorialpress.com/article/ars/Volume%20094/volume-94-paper-17.pdf

[4333] G. Zhenbin, (a, d)-edge-antimagic total labelings of cycle, *Ars Combin.*, **101** (2011) 217-223. https://combinatorialpress.com/article/ars/Volume%20101/volume-101-paper-17.pdf

[4334] G. Zhenbin and F. Chongjin, Some discussions on super edge-magic labelings of $St(a_1, \ldots, a_n)$, *Ars Combin.*, **108** (2013) 187-192. https://combinatorialpress.com/article/ars/Volume%20108/volume-108-paper-15.pdf

[4335] X. Zeng, G. Deng, and C. Luo, Characterize group distance magic labeling of Cartesian product of two cycles, *Discrete Math.*, **346**(8):113407 (2023). doi:DOI:10.1016/j.disc.2023.113407

[4336] L. Zhihe, The balanced properties of bipartite graphs with applications, *Ars Combin.*, **48** (1998) 283-288. https://combinatorialpress.com/article/ars/Volume%20048/volume-48-paper-27.pdf

[4337] S. C. Zhou, Gracefulness of the graph $K_m \cup K_n$, *J. Lanzhou Railway Inst.*, **12** (1993) 70-72.

[4338] S. Zhou, Unifying approaches for constructing labeled graphs from known ones, *J. Combin. Inform. System Sci.*, **20** (1995) 305-319.

[4339] S. Zhou and J. Yuan, On constructions of sequential graphs, *Mathematica Applicata*, **6** (Supplement) (1993) 104-108.

[4340] X. Zhou, B. Yao, and X. Chen, Every lobster is odd-elegant, *Infor. Process. Letters,* **113** (2013) 30-33.

[4341] X. Zhou, B. Yao, X. Chen, and H. Tao, A proof to the odd-gracefulness of all lobsters, *Ars Combin.*, **103** (2012) 13-18. https://combinatorialpress.com/article/ars/Volume%20103/volume-103-paper-2.pdf

[4342] Z. Zhu and C. Liu, Characterizations and structure of sequential graphs, *Ars Combin.*, **116** (2014) 279-288. https://combinatorialpress.com/article/ars/Volume%20116/volume-116-paper-23.pdf

Index

Symbols

$(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial, 343
$(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial graph, 343
$(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial labeling, 343
(a, d)-F-antimagic, 251
(a, d)-1-vertex-antimagic vertex, 254
(a, d)-distance antimagic, 246
$(a, d) - D$ antimagic, 253
(a, r)-geometric, 421
(k, d)-Skolem graceful, 87
(k, d)-graceful labeling, 83
(k, d)-heronian mean, 396
(k, d)-hooked Skolem graceful, 41
(m, n)-gon star, 296
$< K_{m_1,n_1}, \ldots, K_{m_t,n_t} >$, 37
A-antimagic, 237
A-cordial graph, 105
A-magic, 221
$B(n, r, m)$, 27
$B^*_{n,n}$, 335
B_m, 25
$B_{n,n}$, 335
$C(G_1, G_2, \ldots, G_n)$, 37
$C(n \cdot G)$, 37
$CD_m(G)$, 419
$C_m * C_n$, 431
$C_m P_n$, 108
$C_m C_n$, 127
$C^+_{n,k}$, 18
$C^{(t)}_n$, 20
$Circ(n, s)$, 246
D-distance, 246
D-distance antimagic, 246
D-distance magic, 253
D-irregularity strength, 421
D-weight, 253
$DSt_n)$, 38
$D_2(G)$, 88, 120
$D_m(G)$, 92, 142, 419
E-super vertex magic, 197
E_k-cordial, 104
E_k-regular, 198
F-geometric mean, 388
F_n, 52
Fl_n, 313, 346
$G \odot H$, 22
$G \otimes H$, 85
$G \bigtriangledown H$, 56, 60
G^*, 100, 109
G', 37
$G_1 \oplus G_2$, 57
$G_1 \hat{o} G_2$, 302
$G_1[G_2]$, 26
H-E-super magic, 209
H-E-super magic decomposable, 209
H-cordial, 103
H-covering, 170, 261, 415
H-decomposition, 209
H-graph of P_n, 371
H-graph of a path, 382
H-magic, 204
H-supermagic strength, 208
H-union, 120
$H - V$-super magic decomposable, 209
H_n, 371
H_n-cordial, 103
H_n-graph, 127
JF_n, 431
J_n, 431
$KP(r, s, l)$, 72
$K^{(m)}_n$, 26
$M(G)$, 40, 100
$M_m(G)$, 142
M_n, 25
$P(G, f)$, 40
$P(n, k)$, 34
$P(n \cdot G)$, 37
P^t_n, 38
$P^t_n(tn \cdot H)$, 38
$P_n K_{1,n}$, 130

P_n^k, 33
$P_t(G)$, 101
$P_t(u, v)$, 101
$P_{a,b}$, 35, 367
Pl_n, 39
R-ring-magic, 225
$R_{\mathbf{m}}(G)$, 40
$S(G_1, G_2, \ldots, G_n)$, 37
$S(n \cdot G)$, 37
S_m, 25
S_n, 259
$S_{m,n}$, 121
$Spl_m(G)$, 142
$St(n)$, 31
$St(n_1, n_2, \ldots, n_k)$, 86
$T(G)$, 115
$T(P_n)$, 40
$T_n(G)$, 363
T_p-tree, 84
$W(t, n)$, 16
Γ-distance magic, 214
α-labeling
    eventually, 61
    free, 65
    near, 66
    strong, 65
    weakly, 64, 76
α-deficit, 62
α-labeling, 20, 53, 76, 88
α-mean labeling, 368
α-size, 64
α-valuation, 53
β-valuation, 5
$\chi_{slat}(G)$, 257
δ-optimal, 291
δ-optimal summable, 291
γ-labeling, 73
$\hat{\rho}$-labelings, 70
ρ-labeling, 71
ρ-valuation, 71
ρ^+-labeling, 72
$\rho^\star$, 72
θ-labeling, 72
$\tilde{\rho}$-labelings, 75
a-vertex consecutive bimagic labeling, 230
a-vertex consecutive magic labeling, 228
a-vertex multiple magic, 155
b-edge consecutive magic labeling, 229
b-edge multiple magic, 155
d-antimagic, 244
d-graceful, 63
d-lucky number, 448
f-permutation graph, 40
$grac(G)$, 43
k-cordial labeling, 105
k-difference cordial, 358
k-edge-weighting, 420
k-even mean graph, 380
k-even mean labeling, 381
k-even sequential harmonious, 130
k-fold, 187
k-graceful, 82
k-graceful digraph, 85
k-magic, 159
k-mean graph, 371
k-multilevel corona, 149
k-prime, 302
k-prime cordial, 363
k-product cordial, 338
k-ranking, 427
    minimal, 427
k-remainder cordial, 108
k-super mean, 371
k-total product cordial, 338
k-totally magic cordial, 226
k-ubiquitously graceful, 11
k-vertex amalgamation, 59
kC_n-snake, 21, 71
    linear, 21
m index, 114
m-gracefulness, 76
m-mirror graph, 142
m-shadow graph, 142
m-splitting graph, 142
mG, 28
n-cone, 16

n-cube, 25, 53
n-point suspension, 16
nth quadrilateral snake, 432
$n \cdot \vec{C}_m$, 43
q-labeling number, 454
r-distant irregular, 418
r-distant irregularity strength, 418
$rHs(G)$, 416
$rn \star (f)$, 326
$s(G)$, 402
$s_g(G)$, 412
t-fold, 59
t-join (m, n) kite, 441
t-ply graph, 101
$tdis(G)$, 418
$ts(G)$, 413
w-graph, 168
w-tree, 168
y-tree, 13
0-magic, 226
1-vertex bimagic, 220
2-link fence, 58
3-equitable prime cordial, 364
3-product cordial, 335
3-total super sum cordial graph, 342
3-total super sum cordial labeling, 342

A
abbreviated double tree of T, 160
absolutely harmonious graph, 128
additively (a, r)-geometric, 421
adjacency matrix, 71
almost graceful labeling, 70
almost-bipartite graph, 73
alpha-number, 183
alternate hexagonal snake, 21
alternate quadrilateral snake, 302, 333, 355
alternate shell, 100
alternate triangular snake, 302, 333, 355
amalgamation, 207
analytic mean graph, 395
antimagic edge chromatic, 243
antimagic orientation, 241
antipodal balanced, 218
antiprism, 208, 252, 276, 411
apex, 18, 119
arank number, 428
arbitrarily distance antimagic, 246
arbitrarily graceful, 82
arbitrary supersubdivision, 35, 100
arithmetic, 131
armed crown, 353

B
balance index set, 116
balanced cordial, 109
balanced distance graphs, 217
bamboo tree, 9, 89
banana tree, 13, 78, 89
barycentric subdivision, 38
bent ladder, 428
beta combination graph, 442
beta-number, 74
bi-odd sequential, 126
bicomposition, 73
bigraceful graph, 41
bipartite labeling, 64
bistar, 172, 178
block, 20, 182
block graph, 445
block-cut-vertex graph, 182
block-cutpoint, 57
block-cutpoint graph, 20
book, 6, 19, 25, 167
  generalized, 299
  stacked, 25
boundary value, 60
bow graph, 18
broom, 151

C
cactus
  k-angular, 94
  triangular, 20
Cartesian product, 23, 313
cartoon flower
  graph, 151

caterpillar, 9, 53, 66, 78, 124, 174
caterpillar cycle, 427
cells, 57
chain graph, 57, 182
chain of cycles, 19
chain tree, 58
chord, 17
chordal ring, 197, 253
circulant graph, 152
circular lobster, 428
closed m-shadow of a graph, 419
closed helm, 16
coalescence, 57
cocktail party graph, 142, 197, 289
coconut tree, 308
color number, 257
comb tree, 428
combination graph, 441
combs, 36
complete
    n-partite graph, 96, 282
    bipartite graph, 20, 26
    graph, 26
    tripartite graph, 26
complete mixed generalized sausage graph, 233
complete star, 407
component, 303
composition, 26, 94, 313
conjunction, 332
consecutive radio labeling, 326
consecutively super edge-magic, 178
consecutively super edge-magic deficiency, 178
contra harmonic mean, 387
converse skew product, 60
convex polytope, 210, 275
cordial graph, 95
cordial labeling, 93
corona, 22, 167
covering, 262
critical number, 60
crown, 21, 122, 124, 286, 424
cube, 24, 41
cube divisor cordial, 347
cubic graph, 185
cycle, 5, 289
cycle of a graph, 224
cycle of graphs, 37, 369
cycle with a P_k-chord, 17
cycle with parallel C_k- chord, 18
cycle with parallel P_k chords, 17
cyclic G-decomposition, 66
cyclic decomposition, 71
cylinders, 209

D

Dd-length, 330
decomposition, 5, 53, 66, 70, 74
deficiency
    edge-magic, 181
    super edge-magic, 181
degree splitting graph, 357
degree-magic, 149
difference cordial labeling, 355
difference graph, 435
difference index, 463
direct product, 215
directed edge-graceful, 321
directed graceful graph, 44
dis(G), 419
disjoint union, 28
distance k-antimagic, 245
distance antimagic, 245
    labeling, 234
distance magic labeling, 212
distance pair mean, 401
distance reflexive strength, 415
divisor cordial, 343
divisor graph, 444
dodecahedron, 41
double alternate hexagonal snake, 21
double alternate quadrilateral snake, 302, 333, 355
double alternate triangular snake, 302, 333, 355

double coconut trees, 308
double cone, 16
double fans, 36
double graph of G, 150
double hexagonal snake, 21
double path union, 85
double quadrilateral snake, 302, 333, 355
double star, 160
double step grid graph, 38
double tree, 159
double triangular snake, 302, 333, 355, 370
dragon, 19
duplication of a vertex, 37, 332
duplication of an edge, 37, 101, 333
Dutch t-windmill, 20
Dutch windmill, 156

E
EBI(G), 117
edge H-irregularity strength, 415
edge amalgamation, 299
edge comb product, 144
edge even graceful labeling, 77
edge irregular total labeling, 404
edge irregularity strength, 403, 416
edge linked cyclic snake, 367
edge magic graceful, 171
edge magic strength, 156
edge pair sum, 399
edge parity, 64
edge prime index, 464
edge product cordial labeling, 353
edge reduced
integral sum number, 289
sum number, 288
edge trimagic total labeling, 186
edge-antimagic graceful, 254
edge-antimagic total, 239
edge-balance index, 117
edge-bimagic total , 219
edge-coloring k-vertex weighting, 456
edge-covering, 415
edge-decomposition, 67
edge-friendly index, 116
edge-graceful deficiency, 315
edge-graceful spectrum, 316
edge-magic index, 187
edge-magic injection, 172
edge-magic total
labeling, 157
edge-odd graceful, 89, 92
edge-prime graph, 307
ehs(G, H), 415
elegant, 133
elegant labeling, 133
elem. parallel transformation, 84
elementary transformation, 55, 169
envelope graph, 118
EP-cordial graph, 338
EP-cordial labeling, 337
Eulerian graph, 119
even $2a$-sequential, 146
even 1-vertex bimagic, 220
even graceful, 59
even mean labeling, 380
even vertex equitable even, 433
even vertex magic total, 200
even vertex odd mean, 381
even-even, 93
even-odd harmonious, 147
exclusive sum labeling, 290
exclusive sum number, 290
extended w-tree, 168
extended edge vertex cordial labeling, 112
extended jewel graph, 393

F
face, 209, 275
face irregular total k-labeling, 413
fan, 52, 134, 148, 164, 167, 180, 197, 210, 294
fence, 58
FI(G), 113
Fibonacci graceful, 79
firecracker, 13
flag, 96, 130, 321

flower, 16, 195, 294
folded hypercube
graphs, 216
forest, 181
free α-labeling, 65
friendly index set, 113
friendship graph, 20, 94, 179, 195, 197, 209, 290
full r-ary tree, 13
full edge-friendly index, 113, 116
full friendly index set, 117
full hexagonal caterpillars, 59
full product-cordial index, 335
fully magic, 223
fully product-cordial, 334
functional extension, 160
fuzzy, 464

G

gamma-number, 42
gear graph, 16
generalized
book, 299
bundle, 102
fan, 102
wheel, 102
generalized kC_n-snake, 367
generalized antiprism, 268
generalized caterpillar, 36
generalized edge linked cyclic snake, 368
generalized helm, 195, 408
generalized Jahangir graph, 195
generalized prisms, 325
generalized sausage graph, 233
generalized shackle, 261
generalized spider, 36
generalized web, 16, 195
geometric mean 3-equitable, 402
geometric mean cordial, 402
glutting number, 350
Golomb ruler, 28
graceful
$H - E$ super graceful, 178
almost super Fibonacci, 80
graceful center, 66
graceful chromatic number, 425
graceful game, 77
graceful graph, 5
gracesize, 64
gracious k-labeling, 66
gracious labeling, 66
graph, 164, 357, 461
$(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial, 343
(ω, k)-antimagic, 242
(a, d)-F-antimagic, 251
(a, d)-antimagic, 249
(a, d)-distance antimagic, 246
(a, r)-geometric, 421
$(k + 1)$-equitable mean, 401
(k, λ)-magically total labeling, 221
(k, d)-Skolem graceful, 87
(k, d)-arithmetic, 131
(k, d)-balanced, 85
(k, d)-heronian mean, 396
(s, d) magic, 149
2-Skolem difference mean, 385
A-antimagic, 237
A-cordial, 105
A-vertex magic, 154
C-geometric, 389
D-distance, 246
D-distance antimagic, 246
E-cordial, 320
E-super arithmetic graceful, 459
E-super vertex magic, 197
E_k-cordial, 104
G-distance magic, 214
G-snake, 21
H-cordial, 103
H-elegant, 135
H-group magic, 153
H-harmonious, 135
H_k-cordial, 104
H_n-cordial, 103
L Fibonacci cordial graph, 341
S-magic, 153

V-cordial, 350
V-super vertex magic, 155
V_k super vertex in-magic, 230
V_k-super vertex magic, 199
Δ-optimum summable, 291
Γ irregular, 412
θ-Petersen, 317
a-vertex multiple magic, 155
b-edge multiple magic, 155
d-graceful, 63
f-permutation, 40
g-graph, 289
k odd-even prime graceful,, 122
k-antimagic, 242
k-balanced, 111, 121
k-difference cordial, 358
k-distance magic, 214
k-edge-magic, 158
k-enriched fan, 18
k-even edge-graceful, 316
k-magic, 159
k-modular multiplicative, 423
k-multilevel corona, 149
k-odd sequential harmonious, 129
k-prime cordial, 363
k-prime total, 304
k-product cordial, 338
k-shifted antimagic, 236
k-super cube root cube mean, 377
k-super graceful, 76
k-super root square mean, 376
k-ubiquitously, 11
kth Fibonacci prime, 309
$k(G)$ snake, 140
m-level wheel, 317
m-mirror, 142
m-shadow, 142
m-splitting, 142
m-star snake, 404
n-uniform, 408
n-uniform cactus chain, 408
p-distance magic, 215
t-m-Zumkeller, 447
t-uniform homeomorph, 101
w-graph, 168
w-tree, 168
(1,0,0)-F-face magic mean, 390
1-near mean cordial, 112
2-odd, 453
3-equitable prime cordial, 364
3-product cordial, 335
3-total super sum cordial labeling, 342
F-root square mean, 376
k-odd edge mean, 379
aboreale star, 356
absolutely k-harmonious graph, 128
absolutely antimagic, 236
absolutely harmonious, 128
actinia, 365
additively (a, r)-geometric, 421
additively $(a, r)*$-geometric, 421
additively graceful, 112
almost-bipartite, 73
alternate hexagonal snake, 21
alternate quadrilateral snake, 302, 333, 355, 443
alternate shell, 100
alternate triangular belt, 25
alternate triangular snake, 302, 333, 355, 443
analytic mean, 395
analytic odd mean, 395
anti skolem, 374
antimagic, 231
arbitrarily graceful, 82
arbitrary calendula, 382
arithmetic, 131, 422
arithmetic ladder, 404
armed crown, 353
armed helms, 97
balanced distance, 217
balloon, 127
barbell, 405
basket, 414
bent ladder, 428
beta combination, 442

bi-odd sequential, 126
bicomposition, 73
biconditional cordial, 112
bicyclic, 316, 331
bigraceful, 41
bijective product k-cordial, 336
bijective product square k-cordial, 336
blade, 103
block, 445
bow, 18, 344
braid, 186
braided star, 303, 356
branched-prism, 268
broken wheel, 115
broom, 151, 198
butterfly, 130, 316, 321
cactus, 408
calendula, 205
caterpillar cycle, 427
centered triangular difference mean, 385
centered triangular mean, 385
centroidal mean, 373
chain, 208, 417
cherry blossom, 461
chordal ring, 197, 253
circle, 141
circulant, 152
circular ladders, 103
circular lobster, 428
circulent, 246
clematis flower, 461
closed distance magic, 216
closed helm, 16
closed water wheel, 91
cocktail party, 142, 197, 289
coconut tree, 308
comb tree, 428
combs, 36
complete, 26
complete mixed generalized sausage graph, 233
composition, 26
cone, 336
conservative, 150
contra harmonic mean, 387
cordial, 180
countable infinite, 166
cube root cube mean, 377
cube sum, 438
cycle butterfly, 361
cycle with parallel chords, 29
cyclic snake, 379
dandelion, 325
deagon, 387
decomposable, 176
degree-magic, 149
diamond, 82
difference, 435
difference cordial, 355
difference perfect square cordial, 440
Diophantine prime, 465
directed, 7
directed Γ-distance magic, 217
directed edge-graceful, 321
disconnected, 28
distance, 330
distance k-antimagic, 245
distance antimagic, 245
distance irregular k, 415
distance pair mean, 401
divisor, 444
divisor 3-equitable, 446
double m-quadrilateral snake, 129
double alternate hexagonal snake, 21
double alternate quadrilateral snake, 302, 333, 355
double alternate triangular snake, 302, 333, 355
double arrow, 224
double coconut tree, 308
double divisor cordial, 348
double fans, 36
double graph of G, 150
double hexagonal snake, 21
double quadrilateral snake, 302, 333, 355

double squid, 144
double step grid, 38
double triangular, 21
double triangular snake, 302, 333, 355
double vertex wheel, 158
double wheels, 25
double-sided arrow, 398
dragon, 60, 399
dumbbell, 130, 316
Durer, 348
edg-bimagic harmonious, 186
edge corona path, 206, 252
edge edge-magic, 158
edge linked cyclic snake, 367
edge magic graceful, 171
edge pair sum, 400
edge product cordial, 353, 354
edge trimagic graceful, 158
edge vertex prime, 307
edge-friendly, 111
edge-magic, 157, 185
edge-odd graceful, 90
edge-prime, 307
EP-cordial, 338
equitable irregular, 420
even $2a$-sequential, 146
even edge-graceful, 90
even sum, 452
even vertex odd mean, 379, 381
even-multiple subdivision, 98
extended w-tree, 168
extended jewel, 393
extended vertex edge additive cordial, 112
extra Skolem difference mean, 383
face integer edge cordial, 351
face-magic toridal, 218
fan, 52
festoon, 130, 321
Fibonacci cordial, 340
Fibonacci divisor cordial, 342
Fibonacci graceful, 79
Fibonacci graceful antimagic, 460
Fibonacci mean antimagic, 240
Fibonacci product cordial, 341
Fibonacci range, 459
firecracker, 143
flag of C_n, 365
flower, 313, 346
flower snark, 361
fourth order of lemniscate, 94
fractional product cordial, 352
friendship, 20
fully product-cordial, 334
Gaussian antimagic, 247
Gaussian tribonacci r-graceful, 459
Gaussian tribonacci arbitrarily graceful, 459
generalize shackel, 261
generalized caterpillar, 36
generalized chain prism snake, 140
generalized cycle star, 308
generalized edge corona, 234
generalized edge linked cyclic snake, 368
generalized helm, 195, 405, 408
generalized Jahangir, 181, 195
generalized sausage, 233
generalized spider, 36
generalized web, 16, 195
generalized wheel, 334
globe, 114, 350
Goldberg snark) (a nontrivial, bridgeless, cubic graph that is 4-chromatic and cannot be edge colored with fewer than four colors, 97
graceful, 5
graceful antimagic, 254
gracefulness, 76
graph-block chain, 36
grid-like, 55
group S_3, 362
group S_3 cordial remainder, 450
group S_3 mean cordial, 391
hair-kC_4 graph, 140
half-Zumkeller, 447
Halin, 158

Hamming-graceful, 123
handicap distance d-antimagic, 235
Harary, 267
harmonic mean, 386
harmonious, 6, 30
hefty V_4-magic, 154
heronian mean, 396
hetro-cordial, 350
hexagonal snake, 21
highly irregular, 459
highly vertex prime, 304
Hilbert graceful, 123
holiday star, 356
homo-cordial, 350
hybrid quadrilateral snake, 139
hyper strongly multiplicative, 422
ice cream, 346
ideal magic, 173
indexable, 132
integer cordial, 109
integer edge cordial, 350
integral sum, 283
irregular dragon, 60
irregular quadrilateral snake, 355
irregular triangular snake, 355
Jahangir, 95
jelly fish, 171, 443
jewel, 369, 431
join, 32
join sum, 37, 141
kayak paddle, 19
king, 454
kite, 19, 179
Knödel, 197, 297
komodo dragon with many tails, 40
komodo dragons, 40
Kusadama, 356
ladder, 23
Lehmer-4 mean, 388
LH, 462
lict, 91
lilly, 437
lily, 110
line-graceful, 423
linear cactus, 137
litact, 91
local edge antimagic, 243
lollipop, 404
long brush, 214
lotus, 82, 340
Lucas divisor cordial, 347
lucky edge, 449
mean binary L-cordial, 365
mean edge cordial, 365
middle, 100
millipede, 85
minimally k-equitable, 121
mirror, 40
mixed generalized sausage, 233
modified arrow, 144
modified double arrow, 144
modified helm, 405
modular edge-graceful, 424
modular multiplicative, 423
multiple shell, 120
Mycielskian, 213
Mycieski, 119
Napier bridge, 336
nested triangle, 103
node-graceful, 86
north star, 340
nowhere zero modular edge-graceful, 424
odd (a, d)-antimagic, 253
odd antimagic, 253
odd Fibonacci mean, 380
odd prime, 308
odd sum, 127
odd vertex equitable even, 432
odd vertex magic, 152
odd-even prime graceful, 122
one modulo N graceful, 78
one modulo N difference mean, 394
one modulo three square mean, 394
one-sided arrow, 398
ordered, 239

orientable *Gamma*-distance magic, 217
pagoda, 389
pair difference cordial, 451
pair mean, 401
pair mean cordial, 451
pair sum, 399
pair sum modulo, 400
para-squares cactus, 408
para-squares cactus chain, 408
parity combination cordial, 349
path-block chain, 36
pentagonal sum, 444
perfect, 445
perfect super edge-magic, 171
perfectly antimagic total, 240
Perrin graceful, 80
plus, 39, 143
PMC, 451
polar grid, 59
prime, 154, 295, 296, 302
prime distance, 453
prime graceful, 43, 461
prime odd mean, 380
prime-graceful, 462
prism snake, 140
pronic graceful, 67
pseudo-magic, 155
pumpkin, 205
pyramid, 82, 143
radio mean, 391
reduction, 428
relatively prime edge, 464
relaxed mean, 372
relaxed Skolem mean, 372
remainder cordial, 108, 348
replicated, 40
restricted k-mean, 371
restricted triangular difference mean, 394
rigid ladders, 368
root square mean, 375
rose, 461
rose corona, 22
SD-divisor, 446
SD-harmonious, 457
SD-prime cordial, 301
semi Jahangir, 186, 224
semi-edge-prime, 307
semi-magic, 148
semismooth graceful, 84
semitotal-block, 436, 439
set graceful, 428
set sequential, 428
set-cordial, 350
shackel, 260
shackle, 207
shadow, 88, 120
sharp, 239
shell flower, 18
shell-butterfly, 18
shell-type, 18
shipping, 440
signed cordial, 455
signed product cordial, 360, 455
simply sequential, 426
Skolem difference Lucas mean, 383
Skolem difference mean, 382
Skolem even difference mean, 385
Skolem labeled, 87
Skolem-graceful, 86
slanting ladder, 127, 224
smooth graceful, 39
sparkler, 316
sparklers, 130
splitting, 36
square chain, 412
square difference, 436, 439
square harmonious, 130
square product, 458
square sum, 436
squid, 144
SSG(n), 91, 142
stalked subdivided shell, 60
star, 25
star extension, 135
star of, 100, 109

starlike, 30
stem-lotus, 137
step grid, 38, 369
step ladder, 142
strong edge-graceful, 315
strong face plane, 234
strong magic, 173
strong sum, 283
strong super edge-magic, 171
strongly c-elegant, 138
strongly k-indexable, 180
strongly 1-harmonious, 180
strongly felicitous, 137
strongly harmonious, 124
strongly indexable, 132
strongly multiplicative, 422
subdivided shell, 78, 91, 142
sum divisor cordial, 345
sun, 259
sunflower, 97
sunlet, 247
super (a, d)-F-antimagic, 251
super edge magic graceful, 171
super edge-graceful, 62
super Fibonacci graceful antimagic, 461
super graceful, 76
super Lehmer-3 mean, 387
super pair sum, 399
super root square mean, 376
super subdivision, 432
super totient, 463
super vertex mean, 370
supermagic, 148
supersubdivision, 34
swastik, 39
tadpoles, 19
theta, 134
theta graph, 36
Toeplitz, 270
topological cordial graph, 342
torch, 36, 297
tortoise, 368, 373
total, 40, 331
total edge Fibonacci irregular, 411
total mean cordial, 391, 393
total mean labeling, 382
total mixed, 417
total neighborhood magic, 208
total prime, 295, 303
totally antimagic total, 239
totally magic, 201
trangulated wheel, 158
tri sum perfect square cordial, 457
triangular, 25
triangular belt, 186
triangular difference mean, 385
triangular ladders, 335
triangular snake, 20
triangular tree, 53
triangulated prism, 158
triangulated water wheel, 91
tribonacci cordial, 341
trigraph of G, 364
trimagic harmonious, 186
triple star snake, 404
twing, 464
twisted cylinder, 117, 335
T–IASL signed graph, 430
umbrella, 114, 171
unicyclic, 17
uniform bow, 18
uniformly balanced, 111
uniformly cordial, 109
universal *alpha*-graceful, 66
universal graceful, 66
vanessa, 110
VECN prime, 306
vector basis S-cordial graph, 366
vertex k-prime, 330
vertex even mean, 379
vertex magic, 154
vertex odd divisor cordial, 347
vertex odd graceful, 93
vertex odd mean, 379
vertex product, 339
vertex switching, 37, 80, 100, 232

vertex-edge neighborhood-prime, 305
volcano, 399
water wheel, 90, 314
weak antimagic, 237
weak magic, 173
weak sum, 291
web graph without a center, 298
weighted-k-antimagic, 245
wheel gripped, 94
zero divisor, 346
zero-sum A-magic, 221
zig-zag triangle, 166
Zumkeller, 446, 447
graph labeling, 5
super edge-bimagic harmonious, 186
graph-block chain, 36
graphs
barbell wheel, 256
double-sided step, 398
dragon, 373
exponential, 397
firefly, 404
generalized friendship graph of H, 244
hurdle, 460
lilly, 225
long dumbbell, 456
middle, 376
multi-bridge, 236
octopus, 225
one-sided step, 397
Paley, 90
path union of n subdivided shell graphs, 438
radio heronian mean k-graceful, 460
radio mean graceful, 392
super-rooted, 55
superstar, 55
vanessa, 225
volcano, 307
weighted k-list-antimagic, 245
grid, 23, 83
grid-like graph, 55
group irregularity strength, 412

H
half-Zumkeller, 447
Halin graph, 158
Hamming-graceful graph, 123
handicap distance antimagic graphs, 235
handicap incomplete tournament, 235
harmonic labeling
mean cordial, 456
harmonic mean, 386
harmonic mean cordial, 456
harmonious graph, 6
harmonious number, 129
harmonious order, 42
Heawood graph, 42, 65
helm, 16, 294
closed, 97
generalized, 97
heronian mean, 396
Herschel graph, 42, 249
hexagonal lattice, 210
hexagonal snake, 21
Hilbert graceful
labeling, 123
Hilbert number, 123
holey α-labeling, 71
homeomorph, 116
honeycomb graph, 277
hooked Skolem sequence, 87
host graph, 61
hybrid quadrilateral snake, 139
hypercycle, 288
strong, 288
hypergraph, 153, 187, 242, 288
hyperwheel, 288

I
IC-coloring, 426
IC-index, 426
icicle graph, 428
icosahedron, 41
index of cordiality, 101
index of product cordiality, 341
integer-antimagic spectra, 238

integer-magic spectrum, 159, 223
integral radius, 288
integral sum
    number, 286
    tree, 286
integral sum-diameter, 285
irregular crown, 172
irregular fence, 58
irregular labeling, 402
irregular quadrilateral snake, 355
irregular triangle snake, 355
irregularity strength, 402

J
jewel graph, 369
join product, 182
join sum, 37

K
kayak paddle, 19, 72
kite, 19, 62, 179, 199
Kotzig's Conjecture, 72

L
L-cordial, 109
labeleing
    arithmetic cordial labeling modulo η, 353
labeling
    $(\alpha_1, \alpha_2, \ldots, \alpha_k)$-cordial, 343
    (ω, k)-antimagic, 242
    (a, b)-consecutive, 315
    $(a, b; n)$-graceful, 42
    (a, d)- vertex-antimagic edge, 249
    (a, d)-H-antimagic total labeling, 261
    (a, d)-1-vertex-antimagic vertex, 254
    (a, d)-distance antimagic, 246
    (a, d)-edge antimagic total, 250
    (a, d)-edge-antimagic total, 262
    (a, d)-edge-antimagic vertex, 262
    (a, d)-face antimagic, 275
    (a, d)-indexable, 262
    (a, d)-vertex-antimagic total, 259
    (a, r)-geometric, 421
    (k, λ)-magically total labeling, 221
    (k, d)-Skolem, 87
    (k, d)-arithmetic, 131
    (k, d)-even mean, 381
    (k, d)-graceful, 83
    (k, d)-heronian mean, 396
    (k, d)-hooked Skolem graceful, 41
    (k, d)-odd mean, 380
    (k, d)-super mean, 386
    (s, d) magic, 149
    2-Skolem difference mean, 384
    A-magic, 221
    A-vertex magic, 154
    C-geometric mean, 388
    D-irregular k-labeling, 421
    D-magic, 218
    E-cordial, 320
    E-sum cordial, 339
    F-centroidal mean, 373
    F-geometric, 388
    G-distance magic, 214
    H-E-super magic, 209
    H-group magic, 153
    H-irregular k, 415
    H-magic, 204
    L-Fibonacci cordial labeling, 341
    L-cordial, 364
    N-magic total, 155
    $P(a)Q(1)$-super vertex-graceful, 319
    $Q(a)P(b)$-super edge-graceful, 319
    R-ring-magic, 225
    V-cordial, 350
    V-super vertex in-antimagic total graceful labeling, 241
    V-super vertex magic, 155
    V_K-super vertex magic, 199
    V_k-super vertex in-magic labeling, 230
    Δ-exclusive sum labeling, 291
    Γ irregular, 412
    Θ-graceful, 81
    α-, 53
    α-mean, 368
    δ, 465

$\rho^{\star}$, 72
σ-, 72
a-vertex consec. edge-bimagic, 229
a-vertex-consecutive magic, 200
d-antimagic, 244
d-antimagic of type $(1, 1, 1)$, 275
d-graceful, 63
d-lucky, 448
k-DML, 214
k-antimagic, 242
k-balanced, 111
k-cordial, 105
k-distance magic, 214
k-distant edge total labeling, 462
k-edge graceful, 315
k-edge-magic, 158
k-equitable, 118, 121
k-even edge-graceful, 316
k-even mean, 380, 381
k-even sequential harmonious, 130
k-graceful, 85
k-indexable, 132
k-labeling, 330
k-mean, 371
k-odd edge mean, 379
k-odd mean, 380
k-odd sequential harmonious, 129
k-prime, 302
k-prime cordial, 363
k-prime cordial weak, 363
k-prime total, 304
k-product, 336
k-product cordial, 338
k-remainder cordial, 108
k-sequential, 425
k-sequentially additive, 435
k-super cube root mean labeling, 377
k-super harmonic, 396
k-super labeling, 76
k-super mean, 371, 386
k-total product cordial, 338
k-totally magic cordial, 226
kth Fibonacci prime, 309
m-bonacci graceful, 66
t-m-Zumkeller, 447
t-harmonious, 42
w-sum, 291
(1,0,0)-F-face magic, 390
1-near mean cordial labeling, 112
1-vertex bimagic, 220
1-vertex magic, 212
1-vertex magic vertex, 227
3-equitable prime cordial, 364
3-product cordial, 335
3-total super sum cordial labeling, 342
(k, d)-indexable, 132
fractional product cordial, 351
absolute differences of cubic and square difference, 454
absolute graceful, 398
absolutely k-harmonious, 128
absolutely harmonious, 128
additively (a, r)-geometric, 421
additively $(a, r)*$-geometric, 421
additively (k, d)-sequential, 435
additively graceful, 112, 131
almost graceful, 70
almost magic, 219
analytic mean, 395
analytic odd mean, 395
antimagic, 231, 241
antipodal, 328
arbitrarily graceful, 82
arithmetic cordial labeling modulo η, 352
arithmetic sequential graceful, 67
AUM block geomatric mean, 466
balanced, 53, 110
balanced cordial, 109
balanced mean cordial, 393
beta combination, 442
bi-odd sequential, 126
bigraceful, 67
bijective product k-cordial, 336
bijective product square k-cordial, 336
binary L-labeling, 365

binary magic total, 227
bipartite, 64
C-exponential mean, 397
centered triangular difference mean, 385
centered triangular mean, 385
closed distance magic, 215
complete k-equitable, 122
consecutive, 138
consecutive radio, 326
coprime, 308
coprime edge, 464
cordial, 93
cordial edge deficiency, 109
cordial vertex deficiency, 109
cube difference, 440
cube divisor cordial, 347
cube root mean labeling, 377
cubic roots cordial, 320
difference cordial, 355
difference perfect square cordial, 440
directed Γ-distance magic, 217
directed edge-graceful, 321
distance k-antimagic, 245
distance difference magic, 160
distance magic, 212, 227
distance pair mean, 401
distance vertex irregular total k-labeling, 418
dividing graceful, 457
divisor 3-equitable, 446
divisor cordial, 343
double divisor cordial, 348
double uniform $(t_1^{\ell-1}, t_2^{\ell_2})$-ply, 105
dynamic coprime labeling., 309
edge even graceful, 77
edge irregular k-labeling, 416
edge irregular reflexive k-labeling, 414
edge irregular total, 404
edge numbering, 458
edge pair sum, 399
edge product cordial, 353
edge trimagic total, 186
edge vertex prime, 307
edge-antimagic graceful, 254
edge-bimagic, 219
edge-bimagic total, 229
edge-friendly, 111
edge-graceful, 312
edge-magic, 157, 163, 185
edge-magic total, 163
edge-odd graceful, 89, 92
edge-prime, 307
elegant, 133
EP-cordial, 337
equitable, 219
equitably harmonious, 30
even $2a$-sequential, 146
even 1-vertex bimagic, 220
even felicitous, 138
even mean labeling, 380
even sequential harmonious, 129
even sum, 452
even vertex equitable even, 433
even vertex magic total, 200
even vertex odd mean, 381
even–odd cordial labeling, 139
even-even, 93
even-odd harmonious, 147
exponential, 397
extended edge vertex cordial labeling, 112
extended vertex odd mean, 379
F-face mean, 397
face integer cordial, 110
face integer edge cordial, 351
face irregular total k-labeling, 413
felicitous, 136
Fibonacci cordial, 340
Fibonacci divisor, 342
Fibonacci graceful, 79
Fibonacci mean antimagic, 240
Fibonacci product cordial, 341
fractional product cordial labeling, 351
friendly, 109
fuzzy quotient-3 cordial, 367
Gaussian antimagic, 247

Gaussian tribonacci r-graceful, 459
generalized geomertic mean, 389
geometric mean, 394
geometric mean 3-equitable, 402
geometric mean cordial, 402
graceful k-coloring, 425
graceful antimagic, 254
graceful data structure, 70
graceful difference, 44
graceful local antimagic, 243
graceful-harmonious, 43
gracefully consistent, 61
gracious, 66
group S_3, 362
group S_3 cordial, 450
group S_3 mean cordial, 391
half-Zumkeller, 447
handicap distance d-antimagic, 235
harmonious, 30
harmonious numbering, 129
heronian mean, 396
heronian mean Dd-distance, 460
hetro-cordial, 350
highly vertex prime, 304
homo-cordial, 350
i, 239
in-magic total, 230
inclusive distance vertex irregular, 418
indexable, 132
integer cordial, 109
integer cordial edge, 350
interlaced, 53
irregular, 402
L-cordial, 109
Legendre cordial labeling modulo p, 352
Legendre product cordial labeling modulo p, 352
Lehmer-4, 388
LH, 462
line-graceful, 423
local antimagic, 256
local distance antimagic, 255
local edge antimagic, 243
Lucas cordial, 383
Lucas divisor cordial, 347
luck edge mean, 449
luck-edge geometric mean, 450
lucky, 447
lucky edge, 448
magic, 148, 156
  consecutive, 209
  of type (0,1,1), 210
  of type (1,0,0), 210
  of type (1,1,0), 209
  of type (1,1,1), 209
magic valuation, 163
mean, 367
mean cordial, 390
mean edge cordial, 365
mean square cordial, 374
minimum coprime, 308
MMD, 446
modular irregular, 420
modular multiplicative divisor, 446
modulo antimagic, 244
near graceful, 70
near mean, 398
near-elegant, 134
nearly distance magic, 217
nearly graceful, 70
neighborhood-prime, 304
nice $(1, 1)$ edge-magic, 173
non-inclusive distance vertex irregular k-labeling, 419
numbering, 184, 458
odd 1-vertex bimagic, 220
odd Fibonacci mean, 380
odd harmonious, 139, 144
odd mean, 378
odd prime, 308
odd sum, 127
odd vertex equitable, 432
odd-elegant, 136
odd-even, 84, 93
odd-graceful, 70, 88
one modulo N graceful, 78

one modulo N-difference, 394
one modulo three graceful, 78
one modulo three mean, 393
one modulo three root square mean, 394
optimal, 465
optimal k-equitable, 122
optimal sum graph, 282
ordered, 239
orientable Γ-distance magic, 217
pair mean, 401, 451
pair sum, 399
parity combination cordial, 349
parity-weighted totally antimagic total labeling, 239
partial vertex, 111
partitional, 125
Pell graceful, 80
pentagonal sum, 444
perfect super edge-magic, 171
perfectly antimagic total labeling, 240
permuted cordial, 452
Perrin graceful, 80
polychrome, 136
prime, 294
prime cordial, 359
prime graceful, 461
prime-magic, 152
product antimagic, 280
product binary L-labeling, 365
product cordial labeling, 331
product edge-antimagic, 281
product edge-magic, 281
product integer cordial, 110
product magic, 280
product-irregular, 417
pronic graceful, 67
pronic Heron mean, 397
proper lucky, 447
properly even harmonious, 145
pseudo α, 74
pseudograceful, 74
quotient, 454
radial radio mean, 331
radio $\star$, 326
radio antipodal, 327, 328
radio antipodal geometric mean, 389
radio contra harmonic mean, 457
radio geometric mean, 389
radio heronian k-labeling, 460
radio mean, 391
radio mean D-distance, 392
radio mean Dd-distance, 330
radio mean graceful, 461
radio mean square, 377
rainbow antimagic vertex, 236
random walk, 466
range-relaxed graceful, 77
real-graceful, 42
relatively prime edge, 464
relaxed mean, 372
relaxed Skolem mean, 372
remainder cordial, 108, 348
restricted k-mean, 371
restricted super totient, 463
restricted triangular difference mean, 394
reverse edge magic, 160
reverse edge-bimagic, 177
reverse edge-trimagic, 186
reverse super edge magic, 160
reverse super edge-bimagic, 177
reverse super edge-trimagic, 187
rosy, 71
SD-divisor, 446
SD-prime cordial, 301
self-reverse, 213
semi (k, d)-arithmetic, 130
semi harmonious, 130
semi-elegant, 134
sequential, 124
set-ordered odd-graceful, 91
set-ordered strongly k-elegant, 134
sharp ordered, 239
sigma, 212
signed cordial, 455
signed product cordial, 360, 455

simply sequential, 425
Skolem difference Lucas mean, 383
Skolem difference mean, 382
Skolem even difference mean, 385
Skolem even vertex odd difference mean, 384
Skolem odd difference mean, 384
Skolem-graceful, 86
square difference, 439, 440
square difference prime, 440
square divisor cordial, 347
square harmonious, 130
square sum, 436
strength sum, 178
strong edge-graceful, 315
strong parity weighted totally antimagic total labeling, 239
strong rainbow antimagic, 236
strong super edge-magic, 171
strongly (k, d)-indexable, 132
strongly c-harmonious, 124
strongly k-elegant, 134
strongly balanced, 110
strongly edge-magic, 174
strongly even harmonious, 145
strongly graceful, 53, 65
strongly harmonious, 34, 124, 127
strongly indexable, 132
strongly odd harmonious, 139
strongly super edge-graceful, 319
strongly vertex-magic total, 199
successive vertex ordering, 466
sum divisor cordial, 345
sum graph, 282
sum perfect cube, 439
sum perfect square, 438
super (a, d)-F-antimagic, 251
super (a, d)-edge antimagic total, 250
super (a, d)-edge-antimagic graceful, 254
super (a, d)-vertex-antimagic total, 259
super arithmetic graceful, 459
super edge-antimagic total, 264
super edge-graceful, 316
super edge-magic, 174
super edge-magic total, 164
super Fibonacci graceful, 79
super geometric mean, 389
super graceful, 76
super heronian mean, 396
super Lehmer-3 mean, 387
super mean, 370
super pair sum, 399
super root mean, 376
super vertex in-out-antimagic total, 242
super vertex local antimagic total, 257
super vertex mean, 370
super vertex sum labeling, 465
super vertex-graceful, 319
super vertex-magic total, 196
super-edge bimagic cordial, 221
supermagic, 148, 180
topological cordial, 341
total, 238
total H-irregular α, 412
total cordial, 113
total edge product cordial, 354
total edge-cordial, 320
total irregular total k, 413
total magic cordial, 225
total mean, 382
total mean cordial, 391, 393
total neighborhood magic, 208
total neighborhood prime, 305
total prime, 303
total prime graceful, 461
total product cordial labeling, 337
totally antimagic total, 239
totally magic, 201
totally magic cordial, 228
totally vertex-magic cordial, 228
TPD-edge balanced, 352
tri sum perfect square cordial, 457
triangular difference mean, 385
triangular graceful, 77
triangular sum, 443

universal antimagic, 247
VECN prime, 306
vector basis S-cordial, 366
vertex k-prime, 330
vertex antimagic total, 270
vertex balanced cordial, 109
vertex equitable, 430
vertex even mean, 379
vertex in-out-antimagic total, 242
vertex irregular reflexive k-labeling, 415
vertex irregular total, 404
vertex magic total, 155
vertex odd divisor cordial, 347
vertex odd graceful, 93
vertex odd mean, 379
vertex prime, 303
vertex product cordial, 339
vertex-bimagic, 219
vertex-edge neighborhood-prime, 305
vertex-friendly, 117
vertex-graceful, 318
vertex-magic total, 193
vertex-relaxed graceful, 77
weak antimagic, 237
zero-sum A-magic, 221
Zumkeller, 446
Zumkeller cordial, 447
labeling number, 61
labelings
discrete, 352
edge irregular total absolute difference k-labeling, 405
labling
AUM block, 465
local super (a, d)-edge antimagic total, 250
ladder, 23, 124, 210
Langford sequence, 170
level joined planar grid, 133
lexicographic product, 157
linear cyclic snake, 21
lobster, 11, 71
local antimagic chromatic number, 254
local antimagic vertex coloring, 254
local distance antimagic chromatic number, 255
localantimagic total coloring, 244
lotus inside a circle, 210
Lucas divisor cordial, 347
luck edge odd mean number, 450
lucky number, 448

M

Möbius grid, 269
Möbius ladder, 25, 125, 148, 155, 210, 296, 314
magic b-edge consecutive, 200
magic constant, 163, 218
magic square, 148
magic strength, 156, 172
magic sum index, 154
mean cordial, 390
mean graph, 367
mean harmonic matrix, 464
mean number, 390
middle graph, 100
minimum coprime number, 308
mirror graph, 40
mixed generalized sausage graph, 233
mod difference digraph, 436
mod integral sum graph, 290
mod integral sum number, 290
mod sum graph, 289
mod sum number, 289
mod sum* graph, 292
mod sum* number, 292
modular edge-graceful, 424
modular irregularity strength, 420
modulo local antimagic chromatic number, 244
Mongolian tent, 23, 83
Mongolian village, 24, 83
MSG, 289
multigraph, 180, 187
multiple shell, 18
multiply divisor cordial, 351

mutation, 199
mutual duplication, 369

N
near α-labeling, 66
nearly distance magic, 217
nearly graceful labeling, 70
neighborhood-prime, 304
non-inclusive distance irregularity strength, 419
nowhere-zero, 424
nowhere-zero modular edge-gracefulness, 424
nullset, 154
numbering, 184

O
Oberwolfash Problem, 42
oblong numbers, 455
odd 1-vertex bimagic, 220
odd harmonious, 139, 144
odd mean graph, 378
odd mean labeling, 378
odd prime graph, 308
odd-elegant, 136
odd-even, 84, 93
odd-graceful labeling, 70, 88
olive tree, 9
one modulo N graceful, 78
one modulo three graceful labeling, 78
one-point union, 20, 27, 54, 88, 94, 136
open star of G, 224
optimal sum graph, 282

P
pair mean, 401
pair mean graph, 401
pair sum, 399
pair sum graph, 399
parachutes, 249
parallel chord, 115
path, 18, 134
path union, 37, 102
path-block chain, 36
pendent edge, 63
pentagonal number, 444
pentagonal sum labeling, 444
perfect Golomb ruler, 28
perfect system of difference sets, 83
permutation graph, 441
Perrin sequence, 80
Petersen graph, 41
  generalized, 33, 94, 167, 179, 194, 249, 259, 262
planar bipyramid, 210
planar graph, 209, 275
Platonic family, 209
plus graph, 39, 143
polar grid, 59
polyminoes, 83
polyominoes, 59
power graph, 41
prime cordial
  strongly, 362
prime cordial labeling, 359
prime graceful, 43
prime graph, 296, 302
prime labeling, 294
prism, 24, 209, 259, 275
product cordial, 331
product cordial labeling, 331
product graph, 282
product irregularity strength, 417
product-cordial index, 335
product-cordial set, 334
pronic number, 67, 397
properly even harmonious, 145
pseudo α-labeling, 74
pseudo-magic graph, 155
pseudograceful labeling, 74

Q
quadrilateral snakes, 21
quotient labeling number, 454

R
radial radio mean number, 331
radio k-chromatic number, 329

radio k-coloring, 329
radio k-number, 330
radio *star*-number, 326
radio antipodal geometric number, 389
radio antipodal labeling, 327
radio antipodal mean number, 328
radio antipodal number, 327
radio contra harmonic mean number, 457
radio geometric mean number, 389
radio graceful, 326
radio heronian Dd-distance number, 460
radio heronian mean number, 460
radio labeling, 324
radio mean D-distance number, 392
radio mean labeling, 391
radio mean number, 391
radio mean square number, 377
radio number, 324
rainbow antimagic connection number, 236
range-relaxed graceful game, 77
range-relaxed graceful labeling, 77
rank number, 427
real sum graph, 282
reflexive H strength, 416
reflexive edge strength, 414
reflexive vertex strength, 415
regular graph, 148, 152, 167, 195, 227
regular tree, 58
relaxed mean graph, 372
remainder cordial, 108
replicated graph, 40
representation, 433
representation number, 434
restricted triangular difference mean, 394
rigid ladders, 368
Ringel-Kotzig, 9
root, 96
root-union, 115

S

saturated vertex, 283
SD-divisor, 446
SD-prime cordial, 301
semi-edge-prime graph, 307
semismooth graceful, 84
separating set, 428
sequential join, 62
sequential number, 183
set-ordered odd-graceful, 91
shackle, 260
shadow graph, 88, 120
shadow of a graph G, 419
shell, 18, 97, 99, 119
  multiple, 18
shell graph, 107
Skolem labeled graph, 87
Skolem sequence, 12, 29
Skolem-graceful labelings, 86
SLAT, 257
smooth graceful, 39
snake, 20, 57
  n-polygonal, 78
  double triangular, 21
  edge linked cyclic, 367
  generalized edge linked cyclic, 368
  quadrilateral, 56
  triangular, 20, 70
snake polyomono, 57
sparse semi-magic square, 201
special super edge-magic, 175
spider, 9
split graph, 233
splitting graph, 36, 89, 357
spum, 282
square difference graph, 439
square divisor cordial, 347
SSG(n), 91
SSG(n), 142
stable set, 35, 40, 57
star, 31, 32, 195, 424
star of G, 337
star of a G, 100, 109
star of graphs, 37
star super edge-magic deficiency, 166
step grid graph, 38, 369
step ladder, 142

straight simple polymonial caterpillars, 59
strength
  edge magic, 156
  magic, 156, 172
  maximum magic, 173
strength sum, 178
strong product of graphs, 216
strong A-magic, 153
strong k-combination graph, 442
strong k-permutation graph, 442
strong beta-number, 74
strong edge-antimagic, 239
strong edge-graceful, 315
strong gamma-number, 42
strong harmonious number, 129
strong product, 417
strong rainbow antimagic connection number, 236
strong sequential number, 183
strong sum graph, 283
strong supersubdivision, 35
  arbitrary, 35
strong vertex-graceful, 318
strongly c-harmonious, 124
strongly *-graph, 443
strongly antimagic, 243
strongly even harmonious, 145
strongly graceful labeling, 65
strongly harmonious, 34, 124
strongly odd harmonious, 139
strongly prime cordial, 362
strs$G$, 178
stunted tree, 72
subdivided shell graph, 78, 91, 142
subdivision, 11, 23, 88, 210
Sudoku number, 461
sum graph, 282
  mod, 289
  mod integral, 290
  real, 282
sum index, 463
sum number, 282
sum perfect cube, 439
sum perfect square, 438
sum* graph, 291
sum* number, 291
sum-diameter, 284
sunflower, 97, 313
super (a, d)-F-antimagic, 251
super (a, d)-H-antimagic total labeling, 261
super (a, d)-edge-antimagic graceful, 254
super d-antimagic, 244
super edge magic graceful, 171
super edge-magic deficiency, 166
super edge-magic total labeling, 157
super Fibonacci graceful, 79
super geometric mean, 389
super graceful, 76
super labeling, 238
super magic frame, 152
super magic strength, 178
super mean, 370
super mean number, 377
super subdivision, 432
super totient, 462
super vertex local antimagic total chromatic number, 257
super vertex mean, 370
super vertex sum number, 465
super vertex-magic total, 196
super weak sumgraph, 291
supersubdivision, 34, 102
  arbitrary, 35
swastik graph, 39
switching invariant, 300
symmetric product, 26, 57

T

tadpoles, 19
tensor product, 85, 92, 111, 342
tes(G), 404
theta graph, 134
theta graphs, 36
toroidal polyhex, 268
torus grid, 24

total H-irregularity strength, 412
total k-distant chromatic number, 462
total absolute difference edge irregularity strength, 405
total edge (vertex) irregular strength, 404
total edge irregularity strength, 404
total edge product cordial labeling, 354
total graph, 40, 115, 331
total labeling, 238
total mean cordial, 391
total mixed, 417
total negative, 417
total negative edge, 417
total positive edge, 417
total product cordial, 337
  labeling, 339
total product cordial labeling, 337
total stable, 417
total stable edge, 417
totally magic cordial, 228
totally magic cordial deficiency, 227
totally vertex-magic cordial labeling, 228
$\mathrm{tr}(G)$, 326
tree, 5, 233, 289
  binary, 167
  path-like, 169
  symmetrical, 9
triameter, 326
triangular graceful labeling, 77
triangular snake, 20
$\mathrm{tvs}(G)$, 404

U

umbrella, 171
unicyclic graph, 19
uniform cycle snake, 296
uniform-distant tree, 13
union, 27, 163, 179, 181, 195, 286, 296, 303
universal antimagic, 247
unlabeled vertices, 111

V

vertex H-irregularity strength, 416
vertex balance index set, 117
vertex balanced cordial, 109
vertex equitable, 430
vertex irregular total labeling, 404
vertex parity, 64
vertex prime labeling, 303
vertex switching, 37, 80, 100, 232, 300
vertex weight, 257
vertex-antimagic total, 239
vertex-graceful, 318
vertex-relaxed graceful labeling, 77
$\mathrm{vhs}(G, H)$, 416

W

weak sum graph, 291
weak tensor product, 62, 66
weakly α-labeling, 64
web, 16
  generalized, 173
weight, 275
weight of vertex, 418
weight universal antimagic, 247
weighted-k-antimagic, 245
wheel, 16, 124, 148, 163, 197, 209, 231, 262
windmill, 26, 97
working vertex, 290
wreath product, 136

Y

Young tableau, 24, 83

Z

zero-sum A-magic, 221
zero-sum h-magic, 154