

Structure and Growth of \mathbb{R} -Bonacci Words

Sergey Dovgal^a

Sergey Kirgizov^b

Submitted: Jan 11, 2024; Accepted: Jun 21, 2025; Published: Aug 22, 2025

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

A binary word is called q -decreasing, for $q > 0$, if inside this word each of length-maximal (in the local sense) occurrences of a factor of the form $0^a 1^b$, $a > 0$, satisfies $q \cdot a > b$. We bijectively link q -decreasing words with certain prefixes of the cutting sequence of the line $y = qx$. We show that for any real positive q the number of q -decreasing words of length n grows as $C_q \cdot \Phi(q)^n$ for some constant C_q which depends on q but not on n . From previous works, it is already known that $\Phi(1)$ is the golden ratio, $\Phi(2)$ is equal to the tribonacci constant, $\Phi(k)$ is $(k+1)$ -bonacci constant. We prove that the function $\Phi(q)$ is strictly increasing, discontinuous at every positive rational point, and exhibits a fractal structure related to the Stern–Brocot tree and Minkowski’s question mark function.

Mathematics Subject Classifications: 05A05, 68R15, 11B39

1 Introduction

For any real $q > 0$, the *ray cutting word* $s(q)$ is defined as an intersection sequence of a straight half-line $y = qx$ for $x \in (0, \infty)$ with the lines of a square grid ($y = i$ or $x = i$ for $i \in \mathbb{N}^+$). Going along the half-line, starting from $(0, 0)$, we write 1 if the line intersects a horizontal edge and 0 in case of a vertical edge (see Figure 1), we write 01 (in this order) when crossing an intersection point of grid lines.

For any irrational slope q , the word $s(q)$ is aperiodic and Sturmian. In the general setting, Sturmian words are defined as cutting sequences of the line $y = ax + b$ for $x \in (0, \infty)$, irrational $a > 0$ and real $b \in [0, 1)$ or equivalently as binary words having exactly $n + 1$ factors (contiguous subwords) of length n . Sturmian words shine in several areas of mathematics: combinatorics, number theory, tilings, discrete dynamical systems. The structures similar to Sturmian words were already studied by Johann III Bernoulli [6] in 1771. Expositions of Sturmian words and related results can be found in Chapter 2 (written by Berstel and Séébold) of Lothaire’s œuvre [18] and in the book of Allouche and Shallit [1]. For a rational slope q , the word $s(q)$ is periodic, its shortest factor f such that

^aIMB UMR 5584, CNRS, Université Bourgogne-Europe, Dijon, France (dovgal.alea@gmail.com).

^bLIB UR 7534, Université Bourgogne-Europe, Dijon, France (sergey.kirgizov@u-bourgogne.fr).

$s(q) = f \cdot f \cdot f \dots$, where \cdot means concatenation, corresponds to the Christoffel word of slope q [9, 7].




Figure 1: Cutting sequences with slopes $\sqrt{2}$ and $\frac{2}{3}$.

One paradigmatic example of Sturmian words is the *Fibonacci word* $0100101001001\dots$ which is characterized by a cutting sequence of the line with a slope $1/\varphi$, where $\varphi = (1 + \sqrt{5})/2$. It can also be obtained either by a recursive simultaneous application of substitution rules $\{0 \mapsto 01, 1 \mapsto 0\}$ to an initial word 0 , or as a limit of recursive concatenations of words $S_n = S_{n-1}S_{n-2}$, where $S_0 = 0$ and $S_1 = 01$.

Now consider another Fibonacci object, (or, more generally, k -bonacci), which is an ensemble of binary words of length n avoiding k consecutive 1s. It seems that this object appears for the first time in Knuth's book [17, p. 286]. The set of such words is in bijection with tilings of stripes of length $(n+1) \times 1$ with tiles of size 1×1 (monomers), 2×1 (dimers), \dots , $k \times 1$ (k -mers), so it is convenient to call them *k-bonacci tilings*. The cardinality of the set of such words of length n is equal to n th k -bonacci number (see Feinberg [12] and Miles [19]), which is obtained by a recurrence relation $a_n = a_{n-1} + a_{n-2} + \dots + a_{n-k}$ with initial conditions $a_0 = a_{-1} = 1$ and $a_j = 0$ for any $j < -1$.

These two Fibonacci objects belong to two seemingly different worlds. In this paper we propose a link between these worlds: we show that certain subsets of prefixes of ray cutting words can be used as building blocks to construct generalized Fibonacci tilings. To demonstrate this link, we extend the family of q -decreasing words, defined in [5], to cover all positive real numbers as possible values of the parameter q .

Definition 1. For $q \in \mathbb{R}^+$, a binary word is called q -decreasing, if inside this word each of length-maximal occurrences of a factor of the form $0^a 1^b$, $a > 0$, satisfies $q \cdot a > b$. The length-maximality of the occurrences should be taken in the local sense: they are not preceded by a 0 or followed by a 1.

We denote by $\mathcal{W}_{q,n}$ the set of q -decreasing words of length n , Table 1 gives some examples. It is interesting to note that Egecioglu and Iršič [11] independently discovered

n	1	2	3	4	5	n	1	2	3	4	5
$\mathcal{W}_{\sqrt{2},n}$	0 1	00 01 10 11	000 001 010 100 101 110 111	0000	00000	$\mathcal{W}_{2/3,n}$	0 1	00 10 11	000 001 000 110 111	0000 0001 1000 1001 1100 1110 1111	00000
				0001	00001 10010						00001
				0010	00010 10011						00010
				0011	00011 10100						00100
				0100	00100 10101						10000
				0101	00101 11000						10001
				1000	00110 11001						10010
				1001	01000 11010						11000
				1010	01001 11100						11001
				1100	01010 11101						11100
				1101	10000 11110						11110
				1110	10001 11111						11111
				1111							
$ \mathcal{W}_{\sqrt{2},n} $	2	4	7	13	23	$ \mathcal{W}_{2/3,n} $	2	3	5	8	12

Table 1: q -decreasing words for $n \in [1, 5]$ and $q \in \{\sqrt{2}, \frac{2}{3}\}$.

and studied hypercube subgraphs associated with a subset of words from $\mathcal{W}_{1,n}$. By giving a Gray code for $\mathcal{W}_{1,n}$, Baril, Kirgizov and Vajnovszki [5] prove the Egecioğlu-Iršič conjecture [11] about the existence of a Hamiltonian path in such hypercube subgraphs. Recently, Wong, Liu, Lam and Im [25] found a 2-Gray code (where consecutive words differs in at most 2 positions) for $\mathcal{W}_{q,n}$ for any real positive q . The question whether there exists a 1-Gray code when q is a natural number greater than 1 remains open.

In the paper [5] it has been shown that q -decreasing words, for $q \in \mathbb{N}^+$, are in bijection with k -bonacci tilings, where $k = q + 1$, i.e. with the set of n -length binary words that avoid $q + 1$ consecutive 1s. Baril, Kirgizov and Vajnovszki are also proved that $\Phi(1)$ is the golden ratio, $\Phi(2)$ is equal to the tribonacci constant, $\Phi(k)$ is $(k + 1)$ -bonacci constant.

Intriguing bijective and enumerative connections between certain subsets of Dyck paths, integer compositions and q -decreasing words are studied by Barucci, Bernini, Bilotta and Pinzani [3, 4]. Hassler, Vajnovszki and Wong [15] provided, among other things, captivating Gray codes for the set of length n binary words of weight k (i.e., with exactly k 1s) with the property that any prefix contains at least p times as many 0s as 1s, for any $p \in \mathbb{R}^+$.

In Section 2 we decompose q -decreasing words into sequences of words corresponding to ray cutting prefixes ending on 1. The number $\Phi(q)$, called the exponential growth constant, is defined as the limit ratio of successive cardinalities $\Phi(q) = \lim_{n \rightarrow \infty} |\mathcal{W}_{q,n+1}|/|\mathcal{W}_{q,n}|$. In Sections 3 and 4 we show that this limit exists, and explore the structure of $\Phi(q)$ as a function of q . It turns out that the function $\Phi(q)$ is bounded, discontinuous at every positive rational point, strictly increasing over $(0, \infty)$, and also exhibits a nice fractal structure, shown in Figure 2, which bears visual resemblance to fractals arising from information-theoretical applications such as the (appropriately rescaled) number of coin tossings required to obtain a discrete uniform distribution on $[1, n]$ as n goes to infinity

(see a work [2, Fig. 6, right] by Bacher, Bodini, Hwang and Tsai). A characteristic trait of such fractals is that they demonstrate a sort of self-similarity, which is still quite tricky to explain, as the aforementioned similarity is only approximate. We also show that at the vicinity of each rational point q , $\Phi(q)$ converges locally to a piecewise linear function.




Figure 2: $\lim_{n \rightarrow \infty} |\mathcal{W}_{q,n+1}|/|\mathcal{W}_{q,n}|$ as a function of q in three different intervals. This function is jump discontinuous at every positive rational point.

2 Construction from ray cutting prefixes

Here we express q -decreasing words as sequences of ray cutting prefixes ending on 1. It is handy to use the Kleene star operator (it corresponds to **SEQ** operator in the Flajolet–Sedgewick book [13]), which constructs a disjoint union of finite concatenations from words of a given family. For instance, $(\{0, 10\})^*$ provides all binary words which are empty or end on 0 and do not contain two consecutive 1s. We also use the “.” symbol to denote all possible pairwise concatenations between the elements of two families.

Proposition 2. *For $q \in \mathbb{R}^+$, the set \mathcal{W}_q of q -decreasing words can be represented as*

$$\mathcal{W}_q = (\{1\})^* \cdot (\mathcal{S}_q)^*, \text{ where } \mathcal{S}_q = \cup_{i=0}^{\infty} \{0^{1+\lfloor i/q \rfloor} 1^i\}.$$

Proof. By Definition 1, a q -decreasing word is a concatenation of factors $0^a 1^b$ satisfying $a = 0$ or $qa > b$. If $a = 0$, the word starts with an arbitrary sequence of 1s, which is $(\{1\})^*$, otherwise the condition $qa > b$ can be rewritten as $a \geq 1 + \lfloor b/q \rfloor$. By grouping the extra zeros at the beginning of each factor $0^a 1^b$, we write it as $0^t 0^{1+\lfloor i/q \rfloor} 1^i$ with $i = b$ and $t = b - a$. Furthermore, since $0 \in \mathcal{S}_q$, the factor 0^t belongs to the family $(\mathcal{S}_q)^*$, and the remaining part $0^{1+\lfloor i/q \rfloor} 1^i$ belongs to \mathcal{S}_q itself. This procedure allows us to decompose the remainder into a sequence of words from \mathcal{S}_q , which finishes the proof. \square

For a binary word α containing n 0s and m 1s, we define a transformation $\kappa(\alpha) = 0^{n+1} 1^m$, so that the empty word ϵ is mapped to the word 0. We provide a decomposition of q -decreasing words into partitions of certain ray cutting prefixes by using the above transformation.

Proposition 3. For $q \in \mathbb{R}^+$, the transformation κ bijectively maps the set of prefixes ending with 1 of the ray cutting word $s(q)$ to the set \mathcal{S}_q , which is used in the construction of q -decreasing words.

Proof. Take a ray cutting word $s(q) = s_1s_2s_3s_4\ldots$ where every s_i is a binary digit. The index of i th 1 in this word is $i + \left\lfloor \frac{i}{q} \right\rfloor$. The prefix $s_1s_2\ldots s_{i+\lfloor i/q \rfloor}$ of $s(q)$ contains exactly i 1s and $\left\lfloor \frac{i}{q} \right\rfloor$ 0s. The word $\kappa(s_1s_2\ldots s_{i+\lfloor i/q \rfloor}) = 00^{\lfloor \frac{i}{q} \rfloor}1^i$ is a factor from the set \mathcal{S}_q , which completes the proof. \square

See Tables 2 and 3 for examples.

q	Ray cutting word	Factors from \mathcal{S}_q	Some q -decreasing words
$\sqrt{2}$	101011010110...	$\kappa(\epsilon) = 0$, $\kappa(1) = 01$, $\kappa(101) = 0011$, $\kappa(10101) = 000111, \dots$	111100011000111, 000111101000001, 100000101000001
$\frac{2}{3}$	010010100101...	$\kappa(\epsilon) = 0$, $\kappa(01) = 001$, $\kappa(01001) = 000011$, $\kappa(0100101) = 00000111, \dots$	111100001100001, 000011001000001, 001000000001111

Table 2: An illustration of the transformation κ . Prefixes ending with 1 of the ray cutting word $s(q)$ correspond to factors from the set \mathcal{S}_q .

q	Ray cutting word	Counting Sequence $ \mathcal{W}_{q,n} $	OEIS
$\frac{1}{2}$	0010010010010010...	1, 2, 3, 4, 6, 9, 13, 19, 28, 41, 60, 88, 129, 189, ...	Narayana's cows, A930
$1/\varphi$	0100101001001010... Fibonacci word	1, 2, 3, 5, 8, 12, 19, 30, 47, 74, 116, 182, 286, 448, ...	NEW
$\frac{2}{3}$	0100101001010010...	1, 2, 3, 5, 8, 12, 19, 30, 47, 74, 116, 182, 286, 449, ...	Comp. into 1s, 3s and 5s, A60961
1	0101010101010101...	1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, 377, 610, ...	Fibonacci, A45
2	1011011011011011...	1, 2, 4, 7, 13, 24, 44, 81, 149, 274, 504, 927, 1705, 3136, ...	Tribonacci, A73
$\frac{3}{2}$	1010110101101011...	1, 2, 4, 7, 13, 23, 42, 76, 138, 250, 453, 821, 1488, 2697, ...	NEW
$\sqrt{2}$	1010110101101010...	1, 2, 4, 7, 13, 23, 42, 76, 138, 250, 453, 821, 1488, 2697, ...	NEW
φ	1011010110110101-s	1, 2, 4, 7, 13, 24, 44, 81, 148, 272, 499, 916, 1681, 3085, ...	NEW
e	1101110111011011...	1, 2, 4, 8, 15, 29, 56, 108, 208, 401, 773, 1490, 2872, 5536, ...	NEW
π	1110111011101110...	1, 2, 4, 8, 16, 31, 61, 120, 236, 463, 910, 1788, 3513, 6901, ...	NEW

Table 3: Examples of ray cutting words and corresponding counting sequences for the cardinalities of q -decreasing words.

3 Rational discontinuity

In this section we study the function $\Phi(q) = \lim_{n \rightarrow \infty} \frac{|\mathcal{W}_{q,n+1}|}{|\mathcal{W}_{q,n}|}$, whose graph is shown on Figure 2. Using the previously mentioned SEQ operator, Proposition 2 yields the generating function $W_q(x) = \sum_{n=0}^{\infty} |\mathcal{W}_{q,n}| x^n$ of the family \mathcal{S}_q for any $q \in \mathbb{R}^+$:

$$W_q(x) = \frac{1}{(1-x) \left(1 - \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}\right)}. \quad (1)$$

The case where q is a positive *rational number* represented by an irreducible fraction $\frac{c}{d}$ is treated in [16] where the author expresses the generating function $W_q(x)$ as

$$W_{q=\frac{c}{d}}(x) = \frac{1 - x^{c+d}}{(1-x) \left(1 - x^{c+d} - \sum_{i=0}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor}\right)}. \quad (2)$$

In other words, Equation (1) holds for any positive real q , and is more general, although simpler, form of Equation (2) which is only valid for $q \in \mathbb{Q}^+$.

To prove the results about the asymptotic behaviour of coefficients $[x^n]W_q(x)$, we need the following lemma, which can be considered as a simpler variant of Daffodil Lemma from Flajolet–Sedgewick book [13, Lemma IV.1, p. 266].

Lemma 4 (“Little Narcissus Lemma”). *Let $f(x) = x + \sum_{k=2}^{\infty} a_k x^k$ be a power series with non-negative real coefficients $\{a_k\}_{k=2}^{\infty}$ and a positive radius of convergence ρ . The following hold:*

- 1) *there is a unique real root R of the equation $f(x) = 1$;*
- 2) *R has the multiplicity one;*
- 3) *there is no other root $x \in \mathbb{C}, x \neq R$ of the equation $f(x) = 1$ such that $|x| \leq R$.*




Figure 3: An element of the Little Narcissus Lemma proof.

Proof. Suppose that at least one of the coefficients $\{a_k\}_{k=2}^{\infty}$ is strictly positive, otherwise the result is trivially simple. The function $f(x)$ is monotonically increasing on $[0, \rho)$. We have $f(0) = 0$, $\lim_{x \rightarrow \rho} f(x) = +\infty$. So, there is a unique real value $R \in [0, \rho)$ such that

$f(R) = 1$. The multiplicity of R is one because $f'(R) > 0$. For any x such that $|x| < R$, we have, by the triangle inequality, $|f(x)| \leq f(|x|) < f(R) = 1$. Consider the case $|x| = R$. Again, by the triangle inequality we have $|\sum_{i=2}^{\infty} a_i x^i| \leq \sum_{i=2}^{\infty} a_i R^i$. If $x \neq R$, we have a strict inequality $|x + \sum_{i=2}^{\infty} a_i x^i| < R + \sum_{i=2}^{\infty} a_i R^i$, see Figure 3. \square

We have the following result about the asymptotic behaviour of $[x^n]W_q(x)$.

Proposition 5. *For any real $q > 0$, the number of q -decreasing words of length n grows as $C_q \cdot \Phi(q)^n$, where $1/\Phi(q)$ is the unique smallest in modulus root of $1 - \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}$, and*

$$C_q = -\frac{\Phi(q)}{\left((1-x)\left(1 - \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}\right)\right)'(1/\Phi(q))}.$$

Proof. Let $f_q(x) = \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor} = x + x^{2+\lfloor \frac{1}{q} \rfloor} + \dots$. For any $x \in [0, 1)$ we have $\sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor} \leq \sum_{i=1}^{\infty} x^i \leq \frac{x}{1-x}$. So, $f_q(x)$ evaluated at $x \in [0, 1)$ is bounded, thus convergent for any real $q \in (0, \infty)$. Apply Little Narcissus Lemma to see that $f_q(x) = 1$ has a unique smallest in modulus root $R < 1$ which has the multiplicity 1. Let $g_q(x) = (1-x)\left(1 - \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}\right)$. The function $W_q(x) = 1/g_q(x)$ is meromorphic in the unit disc. The root ρ_q of $f_q(x) = 1$ is the unique smallest in modulus pole of $W_q(x)$. The pole has the multiplicity 1. Let $\Phi(q) = 1/\rho_q$. Using classical asymptotic analysis of meromorphic functions (see Flajolet–Sedgewick book [13], Sedgewick’s online course [23] or Orlov’s paper [21]) we obtain the result, expressing C_q as $\Phi(q)/g'(1/\Phi(q))$. \square

The equation $1 = \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}$ shares the smallest in modulus root with

$$A_q := 1 - \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor+j},$$

a fact that has a nice geometrical interpretation. To see it, we have to decompose the set \mathcal{W}_q in another way, different from what was given in Section 2. Here we use a set \mathcal{F}_q of factors $0^a 1^b$ such that $qa > b$ and $b \geq 1$. With this, we decompose any word $w \in \mathcal{W}_q$ as a sequence of 1s, followed by a sequence of factors from \mathcal{F}_q , followed by a sequence of 0s. Any of these sequences can be empty. We have

$$w = \overbrace{1 \dots 1}^{\text{some ones}} \overbrace{f_1 f_2 \dots f_k}^{f_\ell \in \mathcal{F}_q} \overbrace{0 \dots 0}^{\text{some zeros}}, \quad \text{where } \mathcal{F}_q = \bigcup_{i=1}^{\infty} \bigcup_{j=0}^{\infty} \{ \overbrace{0 \dots 00}^{1+\lfloor \frac{i}{q} \rfloor+j \text{ zeros}} \underbrace{1 \dots 11}_i \}.$$

Now, we write the g.f. $W_q(x)$ as

$$W_q(x) = \frac{1}{1-x} \cdot \frac{1}{A_q} \cdot \frac{1}{1-x}.$$




Figure 4: The line $b = \frac{2}{3}a$ and the geometrical interpretation of factors $0^a 1^b$ where $\frac{2}{3}a > b$.

Consider the grid $\mathbb{Z}^+ \times \mathbb{Z}^+$, and make every point (a, b) correspond to a factor $0^a 1^b$. The power series A_q sums over all points with positive integer coordinates found under the line $b = qa$. Figure 4 gives some examples.

From Proposition 5 we see that the function $\Phi(q)$ is well-defined as $\lim_{n \rightarrow \infty} \frac{|\mathcal{W}_{q,n+1}|}{|\mathcal{W}_{q,n}|}$. In Proposition 6 we prove the basic properties of this strictly increasing and bounded function with a countable number of discontinuities, and then calculate the jump sizes in Proposition 8.

Proposition 6. *The function $\Phi(q) = \lim_{n \rightarrow \infty} \frac{|\mathcal{W}_{q,n+1}|}{|\mathcal{W}_{q,n}|}$ is*

- a) *strictly increasing over $q \in [0, \infty)$;*
- b) *bounded, $1 \leq \Phi(q) < 2$, with $\Phi(0) = 1$ and $\lim_{q \rightarrow \infty} \Phi(q) = 2$;*
- c) *left-continuous (and right-discontinuous) at every positive rational point;*
- d) *continuous at every positive irrational point.*

Proof. Let $\delta > 0$.

a) Note that $A_{q+\delta} = 1 - \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} x^{1+i+\lfloor \frac{i}{q+\delta} \rfloor + j}$ contains all terms of $A_q = 1 - \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor + j}$ together with terms not presented in A_q . This can be seen geometrically on Figure 4. The terms corresponding to points of positive integer coordinates, lying at the line $b = qa$, are included in $A_{q+\delta}$ for any $\delta > 0$, but none of these terms are in A_q . So, we necessarily have $\rho_{q+\delta} < \rho_q$, and thus $\Phi(q) < \Phi(q + \delta)$.

b) If $q = 0$, the only q -decreasing word of length n is 1^n , so $\Phi(0) = 1$. It is straightforward that $\lim_{q \rightarrow 0} \Phi(q) = 1$. As $q \rightarrow \infty$ we allow more and more binary words, and the functional limit of A_q can be expressed as 1 minus the sum over all integer points from the positive quadrant: $\lim_{q \rightarrow \infty} A_q(x) = 1 - \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} x^{i+j}$, so $\lim_{q \rightarrow \infty} \Phi(q) = 2$.

c) For a positive rational q represented by an irreducible fraction $\frac{c}{d}$ the line $b = qa$ contains points of integer coordinates (kd, kc) for $k \in [1, \infty)$. For any $\delta > 0$, these points are below the line $b = (q + \delta)a$. The corresponding terms of the form x^{kd+kc} are included in

$A_{q+\delta}$ but not in A_q . This, in turn, influences the smallest in modulus root of $A_{q+\delta}$ which is strictly less than the smallest in modulus root of A_q and we obtain $\Phi(q) < \lim_{\delta \rightarrow 0^+} \Phi(q + \delta)$. The difference between $\lim_{\delta \rightarrow 0^+}$ and $\Phi(q)$ is explicitly calculated in Proposition 8. In other case, no line of the form $b = (q - \delta)a$ can have the points (kd, kc) below it (even if $\delta = 0$). Any point lying under $b = qa$ also lies under $b = (q - \delta)a$ for a sufficiently small δ . We obtain $\lim_{\delta \rightarrow 0^+} \Phi(q - \delta) = \Phi(q)$.

d) For a positive irrational q , the line $b = qa$ contains no points of positive integer coordinates. There are therefore no terms included in A_q corresponding to this line. So, the smallest root ρ_q of A_q can be approached by smallest roots of $\{A_{r_i}\}_{i=1}^\infty$ where $\{r_i\}_{i=1}^\infty$ is a sequence of rational numbers such that $\lim_{i \rightarrow \infty} r_i = q$. \square

Assume that $q \in \mathbb{Q}^+$ is represented by an irreducible fraction c/d . From Equation (2) we see that the growth rate is dictated by the smallest in modulus root ρ_q , of the polynomial

$$\Pi_q := 1 - x^{c+d} - \sum_{i=0}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor}.$$

Comparing Equations (1) and (2) we see that Π_q shares the same smallest in modulus root with

$$1 - \sum_{i=0}^{\infty} x^{1+i+\lfloor \frac{i}{q} \rfloor}.$$

Lemma 7. *The smallest in modulus root ρ_q of Π_q is positive and real.*

Proof. By Little Narcissus Lemma 4. \square

Proposition 8. *For any irreducible $q = \frac{c}{d} \in \mathbb{Q}^+$, $\lim_{\delta \rightarrow 0^+} \Phi(q + \delta)$ equals the reciprocal of the smallest in modulus root, denoted by ρ_q^+ , of the polynomial*

$$\Pi_q^+ := 1 - (2 - x)x^{c+d} - \sum_{i=0}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor}.$$

We have

$$\lim_{\delta \rightarrow 0^+} \Phi(q + \delta) - \Phi(q) = \frac{1}{\rho_q^+} - \frac{1}{\rho_q},$$

where ρ_q is the smallest in modulus root of $\Pi_q := 1 - x^{c+d} - \sum_{i=0}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor}$.

Proof. Assume that q is represented by an irreducible fraction c/d . We consider a set of binary words $\mathcal{W}_{q,n}^+$ where every length-maximal (in the local sense) occurrence of a factor of the form $0^a 1^b$ respects $aq \geq b$. It differs from the definition of q -decreasing words which encloses a strict inequality. It is clear (see e.g. Figure 4) that the set $\mathcal{W}_{q+\delta,n}$ approaches $\mathcal{W}_{q,n}^+$ as $\delta \rightarrow 0$, factors $0^{kd} 1^{kc}$ corresponding to the points of the form (kd, kc) are included in both sets, no points located strictly above the line $b = qa$ are considered in both cases. The set \mathcal{W}_q^+ is constructed as $\mathcal{W}_q^+ = (\{1\})^* \cdot (\mathcal{S}_q^+)^*$, where $\mathcal{S}_q^+ = \{0\} \cup \bigcup_{i=1}^\infty \{0^{\lceil i/q \rceil} 1^i\}$ (c.f. Proposition 2).

Note that $\mathcal{S}_q^+ = \{0\} \cup \mathcal{B}$, where $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2 \cup \dots$, and $\mathcal{B}_0 = \bigcup_{i=1}^{c-1} \{0^{1+\lfloor i/q \rfloor} 1^i\} \cup \{0^d 1^c\}$, and \mathcal{B}_{j+1} is constructed by inserting the factor $0^d 1^c$ after the last 0 in words from \mathcal{B}_j . So, the g.f. $W_q^+(x)$ of the words in \mathcal{W}_q^+ is

$$\begin{aligned} W_{q=\frac{c}{d}}^+(x) &= \frac{1}{(1-x) \left(1 - \left(x + \frac{\sum_{i=1}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor} + x^{c+d}}{1-x^{c+d}} \right) \right)} = \\ &= \frac{1-x^{c+d}}{(1-x) \left(1 + x^{c+d+1} - 2x^{c+d} - \sum_{i=0}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor} \right)} = \frac{1-x^{c+d}}{(1-x) \cdot \Pi_q^+}. \end{aligned}$$

Consider $f_q(x) = x + \frac{\sum_{i=1}^{c-1} x^{1+i+\lfloor \frac{i}{q} \rfloor} + x^{c+d}}{1-x^{c+d}}$, from Little Narcissus Lemma 4 it follows that there are no complex roots of the equation $f_q(x) = 1$ smaller than or equal in modulus to ρ_q^+ that lies in $[0, 1)$. The claimed result is obtained by comparing this formula with Equation (2). \square

4 The fractal

As we can see on Figure 2 the graph of the function $\Phi(q)$ shows a certain amount of self-similarity. We explain some aspects of this fractality in this section. Firstly, we zoom into the intervals $q \in (\frac{k}{k+1}, 1]$ for $k \in [1, \infty)$, and then look into a more general setting using a rescaling based on Minkowski's question mark function and the Stern–Brocot tree. We use this tree to generate a sequence of nested intervals, narrower each time, around a given rational number. The summary of results and one open question are presented at the end of this section.

4.1 Around the point $q = 1$

Let us recall that the smallest in modulus root of $1 - x - x^2$ is $\rho_1 = 1/\varphi$, where $\varphi = (1 + \sqrt{5})/2$. The polynomials Π_q and Π_q^+ are defined in Proposition 8.

Proposition 9. *For natural $k \geq 1$, the smallest in modulus root $\rho_{k/(k+1)}$ of $\Pi_{k/(k+1)}$ is*

$$\rho_1 + C\rho_1^{2k}(1 + o(1)), \quad \text{as } k \rightarrow \infty,$$

where $C = \rho_1^3/(1 + 2\rho_1)$, and ρ_1 is the smallest in modulus root of $1 - x - x^2$.

Proof. After arithmetic transformations, the equation $\Pi_{k/(k+1)} = 0$, i.e. the equation

$$1 - x^{2k+1} - \sum_{i=0}^{k-1} x^{1+i+\lfloor \frac{i(k+1)}{k} \rfloor} = 0$$

turns into

$$1 - x^{2k+1} - \sum_{i=0}^{k-1} x^{1+2i} = 0 \quad \Leftrightarrow \quad 1 = x \cdot \frac{x^{2k+2} - 1}{x^2 - 1} \quad \Rightarrow \quad 1 = x + x^2 - x^{2k+3}.$$

We multiplied by $(x^2 - 1)$ both sides of equation, adding two new roots 1 and -1 . This does not change the overall picture of the asymptotics, because $0 < \rho_{k/(k+1)} < 1$. By Lemma 7 there is no complex roots smaller than or equal in modulus to $\rho_{k/(k+1)}$. The root can be represented as $\rho_{k/(k+1)} = \rho_1 + \varepsilon_k$ for some positive ε_k such that $0 < \rho_1 + \varepsilon_k < 1$. Note that $\varepsilon_k \rightarrow 0$ as k grows. Next, we substitute $\rho_1 + \varepsilon_k$ for x in $x + x^2 - 1 = x^{2k+3}$, use $1 = \rho_1 + \rho_1^2$ and obtain the following:

$$\begin{aligned}\rho_1 + \varepsilon_k + (\rho_1 + \varepsilon_k)^2 - 1 &= (\rho_1 + \varepsilon_k)^{2k+3}, \\ \rho_1 + \varepsilon_k + \rho_1^2 + 2\rho_1\varepsilon_k + \varepsilon_k^2 - 1 &= \rho_1^{2k+3}(1 + o(1)), \\ \varepsilon_k(1 + 2\rho_1 + \varepsilon_k) &= \rho_1^{2k+3}(1 + o(1)).\end{aligned}$$

The claimed result $\rho_{k/(k+1)} = \rho_1 + C\rho_1^{2k}(1 + o(1))$ follows, because $1 + 2\rho_1 + \varepsilon_k \rightarrow 1 + 2\rho_1$ as $k \rightarrow \infty$. \square

Using the Taylor expansion several times one can improve the root approximation and get $\rho_{k/(k+1)} = \rho_1 + C\rho_1^{2k} + O(k\rho_1^{4k})$. But in context of this paper, Proposition 9 is sufficient.

Proposition 10. *For natural $k \geq 2$, the smallest in modulus root $\rho_{(k-1)/k}^+$ of $\Pi_{(k-1)/k}^+$ is*

$$\rho_1 + C\rho_1^{2k}(1 + o(1)),$$

where $C = \rho_1^3/(1 + 2\rho_1)$, and ρ_1 is the smallest in modulus root of $1 - x - x^2$.

Proof. After arithmetic transformations similar to the ones in Proposition 9, the equation $\Pi_{(k-1)/k}^+ = 0$, i.e. the equation

$$1 - x^{2k-1} - \sum_{i=0}^{k-2} x^{1+i+\lfloor \frac{ik}{k-1} \rfloor} - x^{2k-1} + x^{2k} = 0$$

turns into

$$x + x^2 - 1 = x^{2(k-1)}(2x^3 - x - x^4 + x^2).$$

Note that, at some point, we multiply both sides of equation by $(1 - x^2)$, adding -1 and 1 as roots. This does not change the picture dramatically, because $0 < \rho_{(k-1)/k}^+ < 1$.

As in the proof of Proposition 9 the root is $\rho_1 + \varepsilon_k$, $\varepsilon_k \rightarrow 0$ when $k \rightarrow \infty$. Note that $2x^3 - x - x^4 + x^2 - x^5 = (x^3 - x)(1 - x - x^2)$, so we have $2\rho_1^3 - \rho_1 - \rho_1^4 + \rho_1^2 = \rho_1^5$, so $2(\rho_1 + \varepsilon_k)^3 - (\rho_1 + \varepsilon_k) - (\rho_1 + \varepsilon_k)^4 + (\rho_1 + \varepsilon_k)^2 = \rho_1^5(1 + o(1))$. Using the techniques from the previous proof we obtain the claimed result. \square

4.2 Minkowski rescaling and Stern–Brocot tree

Note that Propositions 9 and 10 already provide some insights into the fractal structure of $\Phi(q)$ displayed in Figure 5. On the left side of this figure, the horizontal axis is




Figure 5: Fractal structure of the function $\Phi(q) = \lim_{n \rightarrow \infty} \frac{|\mathcal{W}_{q,n+1}|}{|\mathcal{W}_{q,n}|}$, before and after rescaling on the intervals $(\frac{k-1}{k}, 1]$.

partitioned into intervals $(1/2, 2/3]$, $(2/3, 3/4]$, \dots , and the parts of the plot of $\Phi(\cdot)$ are grouped accordingly. In particular, we showed that

$$\lim_{k \rightarrow \infty} \frac{\Phi(1) - \lim_{\delta \rightarrow 0^+} \Phi(\frac{k-1}{k} + \delta)}{\Phi(1) - \Phi(\frac{k}{k+1})} = 1,$$

i.e. that the images of the intervals $(\frac{k-1}{k}, \frac{k}{k+1}]$ tend to straighten as $k \rightarrow \infty$.

To better observe the self-similarity, the intervals $q \in (k/(k+1), 1]$ and their images under $\Phi(\cdot)$ can be “normalized” using *simple rescaling* and *Minkowski’s question-mark function* [10, 20]. Simple rescaling takes a set of positive values V , containing at least 2 values, and maps every $v \in V$ to $\frac{v - \min V}{\max V - \min V}$, so the image lies in $[0, 1]$. Minkowski’s question-mark function is a little trickier, and we must first discuss mediants and the construction of the Stern–Brocot tree [24, 8].

For two irreducible fractions a/b and c/d their *mediant* is defined as $(a+c)/(b+d)$. The root of the Stern–Brocot tree is $1/1$, which is the mediant of two conventionally irreducible fractions $1/0$ and $0/1$. To determine the left (resp. right) child of a node x/y of the level i we need to find the greatest (resp. smallest) fraction $x'/y' < x/y$ (resp. $x'/y' > x/y$) that appears in set of values of first i levels together with $1/0$ and $0/1$, and compute the mediant $(x+x')/(y+y')$. For instance, the left child of $2/3$ is $3/5$, it is calculated as the mediant of $1/2$ and $2/3$. Figure 6 illustrates this process. Stern–Brocot tree contains all rationals once.

Minkowski’s question-mark function, denoted by $?(x)$, maps a positive rational value x to a positive dyadic rational $a/2^k$ with $a, k \in \mathbb{N}$. By definition, $?(0) = 0$ and $?(1) = 1$. Whenever $x \in (0, 1)$ is a rational number represented by an irreducible fraction a/b , such that in the Stern–Brocot tree it is constructed via taking a mediant of two fractions p/r and p'/r' , its image under Minkowski’s function is defined as

$$?(x) = ?\left(\frac{p+p'}{r+r'}\right) := \frac{1}{2} \left(?\left(\frac{p}{r}\right) + ?\left(\frac{p'}{r'}\right) \right).$$




Figure 6: The Stern–Brocot tree and Minkowski’s $?(x)$.

In other words, we descend the Stern–Brocot tree in search of the a/b , and “in parallel” construct a resulting value by applying the mean instead of the mediant. For $x > 1$, Minkowski’s function is defined as $?(x + 1) = ?(x) + 1$. In general, $?(x)$ is monotonically increasing, and can be defined on all \mathbb{R}^+ [10].

The right side of Figure 5 is obtained by applying the simple rescaling on the vertical axis and Minkowski’s question-mark function followed by the simple rescaling on the horizontal axis for intervals $(k/(k + 1), 1]$ and their images. The similar analysis can be done for intervals $(1, (k + 1)/k]$. The fractal structure of Φ presented in Figure 2 appears more regular in Figure 7 as we apply Minkowski’s question-mark function over the x -axis.




Figure 7: $\Phi(q) := \lim_{n \rightarrow \infty} |\mathcal{W}_{q,n+1}|/|\mathcal{W}_{q,n}|$ as a function of $?(q)$.

4.3 Around a positive rational number q

Now, we study the more general case, that is the fractal structure of $\Phi(q)$ on the intervals $(\frac{p+ck}{r+dk}, \frac{c}{d}]$ for $k \geq 1$, where fractions are irreducible, and $\frac{p+c}{r+d}$ is the left child of c/d in the

Stern–Brocot tree. Next proposition can be applied, for example, to the intervals $(\frac{3+5k}{2+3k}, \frac{5}{3}]$, where $8/5$ is the left child of $5/3$.

Proposition 11. *For natural $k \geq 1$, the smallest in modulus root of $\Pi_{\frac{p+ck}{r+dk}}$ is*

$$\rho_{\frac{p+ck}{r+dk}} = \rho_{c/d} + C\rho_{c/d}^{(c+d)k}(1 + o(1)),$$

where $k \geq 1$, $\frac{p+c}{r+d}$ is the left child of c/d in the Stern–Brocot tree, C is a constant depending only on p/r and c/d , and $\rho_{c/d}$ is the smallest in modulus root of $\Pi_{c/d}$.

Proof. Recall that from Proposition 8 we have

$$\Pi_{\frac{p+ck}{r+dk}} := 1 - x^{p+r+(c+d)k} - \sum_{i=0}^{p+ck-1} x^{1+i+\lfloor \frac{i(r+dk)}{p+ck} \rfloor}. \quad (3)$$

For $0 < i < p + ck$, $1 + i + \lfloor \frac{i(r+dk)}{p+ck} \rfloor$ equals the number of integer points with coordinates (i, y) lying between two diagonal lines intersecting at the origin with respective slopes $(-c)$ and $\frac{r+dk}{p+ck}$, see Figure 8 for an illustration.

Note that from the construction of Stern–Brocot tree, $\frac{p}{r} < \frac{c}{d}$. From the mediant inequality it follows that $\frac{p}{r} < \frac{p+ck}{r+dk} < \frac{c}{d}$, for any integer $k > 0$. Next, let us show that there are no integer points whose horizontal coordinate is equal to i , $0 < i < p + ck$, lying strictly between the lines with slopes $\frac{r+dk}{p+ck}$ and $\frac{d}{c}$. Consider the triangle ABC with following integer coordinates:


$$\begin{aligned} A &= (0, 0); \\ B &= (c(k+1), d(k+1)); \\ C &= (p+ck, r+dk). \end{aligned}$$

The area of ABC is

$$\frac{1}{2} \begin{vmatrix} c(k+1) & p+ck \\ d(k+1) & r+dk \end{vmatrix} = \frac{1}{2}(k+1) \begin{vmatrix} c & p \\ d & r \end{vmatrix} = \frac{k+1}{2}(cr - dp) = \frac{k+1}{2},$$

because $\frac{p}{r} < \frac{c}{d}$ and $(cr - dp) = 1$ by a joli property of the Stern–Brocot tree (see [14]). Pick’s Theorem [22] implies that the area of triangle ABC is equal to $I + B/2 - 1$, where I is the number of its interior points and exactly B points lie on the boundary. We have $B = k + 3$, because $\frac{p+ck}{r+dk}$ and $\frac{c}{d}$ are irreducible fractions. We conclude that $I = 0$ and there are no interior points inside ABC . This fact can be used to simplify $\Pi_{\frac{p+ck}{r+dk}}$ from Equation (3) by considering only points lying on or below the line of slope d/c and on or above the line of slope $(-c)$. Taking this into account, we rewrite the equation $\Pi_{\frac{p+ck}{r+dk}} = 0$, i.e. the equation

$$1 - x^{p+r+(c+d)k} - \sum_{i=0}^{p+ck-1} x^{1+i+\lfloor \frac{i(r+dk)}{p+ck} \rfloor} = 0$$



$$\begin{aligned}
1 - \sum_{j=0}^p x^{1+j+\lfloor \frac{jd}{c} \rfloor} - \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} \sum_{i=0}^{k-1} x^{(c+d)i} &= 0, \\
1 - \sum_{j=0}^p x^{1+j+\lfloor \frac{jd}{c} \rfloor} - \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} \frac{1 - x^{(c+d)k}}{1 - x^{c+d}} &= 0, \\
1 - \sum_{j=0}^p x^{1+j+\lfloor \frac{jd}{c} \rfloor} - x^{c+d} + x^{c+d} \sum_{j=0}^p x^{1+j+\lfloor \frac{jd}{c} \rfloor} - \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + x^{(c+d)k} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} &= 0,
\end{aligned}
\tag{4}$$

$$\begin{aligned}
1 - x^{c+d} - \sum_{j=0}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + \sum_{j=c}^{p+c} x^{1+j+d+\lfloor \frac{(j-c)d}{c} \rfloor} + x^{(c+d)k} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} &= 0, \\
1 - x^{c+d} - \sum_{j=0}^{c-1} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + x^{(c+d)k} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} &= 0.
\end{aligned} \tag{5}$$

At some point, we multiply both sides by $(1 - x^{c+d})$. All roots of Equation (4) are also the roots of Equation (5), but the latter is additionally satisfied by the roots of unity $1 = x^{c+d}$, the modulus of which is greater than $\rho_{\frac{p+ck}{r+dk}}$.

Finally, since the first three terms of the sum are equal to $\Pi_{c/d}(x)$, we conclude that for $x = \rho_{\frac{p+ck}{r+dk}}$ we have

$$-\Pi_{c/d}(x) = x^{(c+d)k} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor}.$$

Using the same method as in the proof of Proposition 9 and denoting by $\Pi'_{c/d}$ the derivative of $\Pi_{c/d}$ we obtain

$$\rho_{\frac{p+ck}{r+dk}} = \rho_{c/d} + C \rho_{c/d}^{(c+d)k} (1 + o(1)),$$

where

$$C = \frac{\sum_{j=p+1}^{p+c} \rho_{c/d}^{1+j+\lfloor \frac{jd}{c} \rfloor}}{-\Pi'_{c/d}(\rho_{c/d})}. \quad \square$$

Proposition 12. For natural $k \geq 2$, the smallest in modulus root $\rho_{\frac{p+c(k-1)}{r+d(k-1)}}^+$ of $\Pi_{\frac{p+c(k-1)}{r+d(k-1)}}^+$ is

$$\rho_{\frac{p+ck}{r+dk}} = \rho_{c/d} + C \rho_{c/d}^{(c+d)k} (1 + o(1)),$$

where $k \geq 1$, $\frac{p+c}{r+d}$ is the left child of c/d in the Stern–Brocot tree, C is the same constant as in Proposition 11 not depending on k , and $\rho_{c/d}$ is the smallest in modulus root of $\Pi_{c/d}$.

Proof. Recall that Proposition 8 defines $\Pi_{a/b}^+ := \Pi_{a/b} - x^{a+b} + x^{a+b+1}$. Having this in mind and adapting the equations from the proof of Proposition 11 by writing $k-1$ in place of k we obtain:

$$\begin{aligned}
1 - x^{c+d} - \sum_{j=0}^{c-1} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + x^{(c+d)(k-1)} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} - \\
-x^{p+r+(c+d)(k-1)}(1 - x^{c+d}) + x^{p+r+(c+d)(k-1)+1}(1 - x^{c+d}) &= 0,
\end{aligned}$$

$$1 - x^{c+d} - \sum_{j=0}^{c-1} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + x^{(c+d)(k-1)} \times \\ \times \left(\sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} - x^{p+r} + x^{p+r+(c+d)} + x^{p+r+1} - x^{p+r+(c+d)+1} \right) = 0. \quad (6)$$

Now, using the fact that $cr - dp = 1$ and a kind of geometrical argument (as in the previous proof), it can be shown that

$$\sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} - x^{p+r} + x^{p+r+(c+d)} + x^{p+r+1} - x^{p+r+(c+d)+1} - x^{(c+d)} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} = \\ = \left(1 - x^{c+d} - \sum_{j=0}^{c-1} x^{1+j+\lfloor \frac{jd}{c} \rfloor} \right) (x^{c+d+p+r} - x^{p+r}).$$

Adapting the techniques from the proofs of Propositions 9 and 10 we see that the smallest in modulus root of (6) is equal to the smallest in modulus root of

$$1 - x^{c+d} - \sum_{j=0}^{c-1} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + x^{(c+d)(k-1)} \left(x^{(c+d)} \sum_{j=p+1}^{p+c} x^{1+j+\lfloor \frac{jd}{c} \rfloor} + o(1) \right) = 0,$$

where $o(1)$ is considered as $k \rightarrow \infty$. Comparing it to Equation (5) from the proof of Proposition 11, we see that the claimed result follows. \square

4.4 Summary of fractal results and an open question

To summarize the results of the previous two propositions, let q denote $\frac{c}{d}$ and let q_k^* denote $\frac{p+ck}{r+dk}$, i.e. the k th left approximation of q in the Stern–Brocot tree. From Propositions 11 and 12 the following result follows.

Proposition 13. *The graph of the function $\Phi(q)$ rescaled on the intervals $(q_k^*, q]$ tends to a constant function on the semi-open intervals $(q_{k-1}^*, q_k^*]$:*

$$\lim_{k \rightarrow \infty} \frac{\Phi(q) - \lim_{\delta \rightarrow 0^+} \Phi(q_{k-1}^* + \delta)}{\Phi(q) - \Phi(q_k^*)} = 1.$$

The ratio between consecutive constants on the rescaled graph tends to ρ_q^{c+d} :

$$\lim_{k \rightarrow \infty} \frac{\Phi(q) - \Phi(q_k)}{\Phi(q) - \Phi(q_{k-1})} = \rho_q^{c+d}.$$

Using a similar technique, it is possible to demonstrate a resembling picture using the right child of 1 in the Stern–Brocot tree, generating intervals of the form $[1, \frac{k+1}{k})$, and, more generally, using the right approximation of any rational point $q = c/d$.

From Proposition 8 it follows that the function $\Phi(q)$ has the highest jump at point $q = 1$, the second highest jump is at $q = 1/2$, the third highest jump appears when $q = 2$, see Figure 2. The sequence of positive rational numbers ordered by corresponding jumps of the function $\Phi(q) = \lim_{n \rightarrow \infty} |\mathcal{W}_{q,n+1}|/|\mathcal{W}_{q,n}|$ starts with

$$1, \frac{1}{2}, 2, \frac{1}{3}, \frac{1}{4}, 3, \frac{2}{3}, \frac{1}{5}, \frac{1}{6}, \frac{3}{2}, \frac{1}{7}, 4, \frac{2}{5}, \frac{1}{8}, \frac{1}{9}, \frac{1}{10}, \frac{3}{4}, \frac{1}{11}, \frac{2}{7}, \frac{1}{12}, 5, \frac{3}{5}, \frac{1}{13}, \frac{4}{3}, \frac{1}{14}, \frac{2}{9}, \frac{1}{15}, \frac{1}{16}, \frac{5}{2}, \frac{1}{17}, \dots$$

Question 14. Is it possible to explain this sequence without polynomial root calculations?

Acknowledgements

Sergey Dovgal was supported by the EIPHI Graduate School (contract ANR-17-EURE-0002), FEDER Région Bourgogne Franche-Comté, and Sergey Kirgizov was supported in part by the project ANER ARTICO funded by Bourgogne-Franche-Comté region (France) and ANR-22-CE48-0002 funded by l'Agence Nationale de la Recherche. Special thanks to Michael A. Allen, Jean-Luc Baril, Khaydar Nurligareev, Nathanaël Hassler and Vincent Vajnovszki for inspiring and helpful discussions, to Vlady Ravelomanana for helping to connect the two authors, and to Natalia Kharchenko for helping to improve the current manuscript. We thank the anonymous reviewers who provided very good comments.

References

- [1] Allouche, J.-P., and Shallit, J. *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, 2003
- [2] Bacher, A., Bodini, O., Hwang, H. K., and Tsai, T. H. *Generating Random Permutations by Coin Tossing: Classical Algorithms, New Analysis, and Modern Implementation*. ACM Transactions on Algorithms, 13(2), 2017, pp. 1–43.
- [3] Barucci, E., Bernini, A., Bilotta, S., Pinzani R. *Dyck Paths Enumerated by the Q-bonacci Numbers*. GASCom 2024, in Electronic Proceedings in Theoretical Computer Science, 403, 2024, pp. 49–53.
- [4] Barucci, E., Bernini, A., Bilotta, S., Pinzani R. *Rational Dyck paths*. Journal of Integer Sequences, 28(3), 2025, Article 25.3.2.
- [5] Baril, J.-L., Kirgizov, S. and Vajnovszki, V. *Gray codes for Fibonacci q-decreasing words*. Theoretical Computer Science, 927, 2022, pp. 120–132.
- [6] Bernoulli, J. III. *Sur une nouvelle espece de calcul*. Recueil pour les Astronomes, 1, Berlin, 1771, pp. 255–284.
- [7] Berstel, J., Lauve, A., Reutenauer, C. and Franco, S. *Combinatorics on words: Christoffel words and repetitions in words*. CRM Monograph Series, 27, American Mathematical Society, 2009.
- [8] Brocot, A. *Calcul des rouages par approximation, nouvelle méthode*. Revue Chronométrique, 3, 1861, pp. 186–194.

- [9] Christoffel, E. B. *Observatio arithmetica*. Annali di Matematica Pura ed Applicata, 6, 1875, pp. 145–152.
- [10] Denjoy, A. *Sur une fonction réelle de Minkowski*. Journal de Mathématiques Pures et Appliquées. Neuvième Série, 17, 1938, pp 105–151.
- [11] Egecioğlu, Ö., and Iršič, V. *Fibonacci-run graphs I: Basic properties*. Discrete Applied Mathematics, 295, 2021, pp. 70–84.
- [12] Feinberg, M. *Fibonacci-Tribonacci*. Fibonacci Quarterly, 1(3), 1963, pp. 71–74.
- [13] Flajolet, P., and Sedgewick, R. *Analytic Combinatorics*. Cambridge University Press, 2009.
- [14] Graham, R.L., Knuth, D.E. and Patashnik, O. *Concrete Mathematics: A Foundation for Computer Science* 2nd ed., Addison-Wesley, 1994
- [15] Hassler, N., Vajnovszki, V. and Wong, D. *Greedy gray codes for some restricted classes of binary words*. In Brlek, S. and Ferrari, L. (eds) Proc. of the 13th ed. of the conf. on Random Generation of Combinatorial Structures (GASCom), Polyominoes and Tilings, Bordeaux, France, 24-28th June 2024, Electronic Proceedings in Theoretical Computer Science, 403, Open Publishing Association, 2024, pp. 108–112.
- [16] Kirgizov, S. *\mathbb{Q} -bonacci words and numbers*. Fibonacci Quarterly, 60(5), 2022, pp. 187–195.
- [17] Knuth, D. E. *The Art of Computer Programming, Volume 3: Sorting and Searching* 2nd ed., Addison-Wesley, 1998.
- [18] Lothaire, M. *Algebraic Combinatorics on Words*. Cambridge University Press, 2002.
- [19] Miles Jr., E. P. *Generalized Fibonacci numbers and associated matrices*. The American Mathematical Monthly, 67(8), 1960, pp. 745–752.
- [20] Minkowski, H. *Zur Geometrie der Zahlen*. Verhandlungen des III. internationalen Mathematiker-Kongresses in Heidelberg, Berlin, 1904, pp. 164–173.
- [21] Orlov, A. G. *On asymptotic behavior of the Taylor coefficients of algebraic functions*. Siberian Mathematical Journal, 25(5), 1994, pp. 1002–1013.
- [22] Pick, G. *Geometrisches zur Zahlenlehre*. Sitzungsberichte des deutschen naturwissenschaftlich-medicinischen Vereines für Böhmen "Lotos" in Prag. (Neue Folge), 19, 1899, pp. 311–319.
- [23] Sedgewick, R. *Analytic Combinatorics. Online Course. Chapter 4: Complex Analysis, Rational and Meromorphic Asymptotics*. <https://ac.cs.princeton.edu/online/slides/AC04-Poles.pdf>, Princeton University, 2024.
- [24] Stern, M. A. *Ueber eine zahlentheoretische Funktion*. Journal für die reine und angewandte Mathematik, 55, 1858, pp. 193–220.
- [25] Wong, D., Liu, B., Lam, C.-T., and Im, M. *Generating cyclic 2-Gray codes for Fibonacci q -decreasing words*. In Uehara, R., Yamanaka, K., Yen, H.-C. (eds) WALCOM: Algorithms and Computation. WALCOM 2024, Kanazawa, Japan, 18-20th March 2024, Lecture Notes in Computer Science, vol. 14549, Springer, Singapore, 2024, pp. 91–102.