

# Chromatically Unique Multibridge Graphs

F.M. Dong

Mathematics and Mathematics Education, National Institute of Education  
Nanyang Technological University, Singapore 637616  
fmdong@nie.edu.sg

K.L. Teo, C.H.C. Little\*, M. Hendy

Institute of Fundamental Sciences PN461, Massey University  
Palmerston North, New Zealand  
k.l.teo@massey.ac.nz, c.little@massey.ac.nz, m.hendy@massey.ac.nz

K.M. Koh

Department of Mathematics, National University of Singapore  
Singapore 117543  
matkohkm@nus.edu.sg

Submitted: Jul 28, 2003; Accepted: Dec 13, 2003; Published: Jan 23, 2004  
MR Subject Classification: 05C15

## Abstract

Let  $\theta(a_1, a_2, \dots, a_k)$  denote the graph obtained by connecting two distinct vertices with  $k$  independent paths of lengths  $a_1, a_2, \dots, a_k$  respectively. Assume that  $2 \leq a_1 \leq a_2 \leq \dots \leq a_k$ . We prove that the graph  $\theta(a_1, a_2, \dots, a_k)$  is chromatically unique if  $a_k < a_1 + a_2$ , and find examples showing that  $\theta(a_1, a_2, \dots, a_k)$  may not be chromatically unique if  $a_k = a_1 + a_2$ .

**Keywords:** Chromatic polynomials,  $\chi$ -unique,  $\chi$ -closed, polygon-tree

## 1 Introduction

All graphs considered here are simple graphs. For a graph  $G$ , let  $V(G)$ ,  $E(G)$ ,  $v(G)$ ,  $e(G)$ ,  $g(G)$ ,  $P(G, \lambda)$  respectively be the vertex set, edge set, order, size, girth and chromatic polynomial of  $G$ . Two graphs  $G$  and  $H$  are *chromatically equivalent* (or simply  *$\chi$ -equivalent*),

---

\*Corresponding author.

symbolically denoted by  $G \sim H$ , if  $P(G, \lambda) = P(H, \lambda)$ . Note that if  $H \sim G$ , then  $v(H) = v(G)$  and  $e(H) = e(G)$ . The *chromatic equivalence class* of  $G$ , denoted by  $[G]$ , is the set of graphs  $H$  such that  $H \sim G$ . A graph  $G$  is *chromatically unique* (or simply  $\chi$ -*unique*) if  $[G] = \{G\}$ . Whenever we talk about the chromaticity of a graph  $G$ , we are referring to questions about the chromatic equivalence class of  $G$ .

Let  $k$  be an integer with  $k \geq 2$  and let  $a_1, a_2, \dots, a_k$  be positive integers with  $a_i + a_j \geq 3$  for all  $i, j$  with  $1 \leq i < j \leq k$ . Let  $\theta(a_1, a_2, \dots, a_k)$  denote the graph obtained by connecting two distinct vertices with  $k$  independent (internally disjoint) paths of lengths  $a_1, a_2, \dots, a_k$  respectively. The graph  $\theta(a_1, a_2, \dots, a_k)$  is called a *multibridge* (more specifically  $k$ -*bridge*) graph (see Figure 1).




Figure 1

Given positive integers  $a_1, a_2, \dots, a_k$ , where  $k \geq 2$ , what is a necessary and sufficient condition on  $a_1, a_2, \dots, a_k$  for  $\theta(a_1, a_2, \dots, a_k)$  to be chromatically unique? Many papers [2, 4, 10, 6, 11, 12, 13, 14] have been published on this problem, but it is still far from being completely solved [8, 9]. In this paper, we shall solve this problem under the condition that  $\max_{1 \leq i \leq k} a_i \leq \min_{1 \leq i < j \leq k} (a_i + a_j)$ .

## 2 Known results

For two non-empty graphs  $G$  and  $H$ , an *edge-gluing* of  $G$  and  $H$  is a graph obtained from  $G$  and  $H$  by identifying one edge of  $G$  with one edge of  $H$ . For example, the graph  $K_4 - e$  (obtained from  $K_4$  by deleting one edge) is an edge-gluing of  $K_3$  and  $K_3$ . There are many edge-gluing of  $G$  and  $H$ . Let  $\mathcal{G}_2(G, H)$  denote the family of all edge-gluing of  $G$  and  $H$ . Zykov [15] showed that any member of  $\mathcal{G}_2(G, H)$  has chromatic polynomial

$$P(G, \lambda)P(H, \lambda)/(\lambda(\lambda - 1)). \tag{1}$$

Thus any two members in  $\mathcal{G}_2(G, H)$  are  $\chi$ -equivalent.

For any integer  $k \geq 2$  and non-empty graphs  $G_0, G_1, \dots, G_k$ , we can recursively define

$$\mathcal{G}_2(G_0, G_1, \dots, G_k) = \bigcup_{\substack{0 \leq i \leq k \\ G' \in \mathcal{G}_2(G_0, \dots, G_{i-1}, G_{i+1}, \dots, G_k)}} \mathcal{G}_2(G_i, G'). \tag{2}$$

Each graph in  $\mathcal{G}_2(G_0, G_1, \dots, G_k)$  is also called an *edge-gluing* of  $G_0, G_1, \dots, G_k$ . By (1), any two graphs in  $\mathcal{G}_2(G_0, G_1, \dots, G_k)$  are  $\chi$ -equivalent.

Let  $C_p$  denote the cycle of order  $p$ . It was shown independently in [12] and [13] that if  $G$  is  $\chi$ -equivalent to a graph in  $\mathcal{G}_2(C_{i_0}, C_{i_1}, \dots, C_{i_k})$ , then  $G \in \mathcal{G}_2(C_{i_0}, C_{i_1}, \dots, C_{i_k})$ . In other words, this family is a  $\chi$ -equivalence class.

For  $k = 2, 3$ , the graph  $\theta(a_1, a_2, \dots, a_k)$  is a cycle or a generalized  $\theta$ -graph respectively, and it is  $\chi$ -unique in both cases (see [10]). Assume therefore that  $k \geq 4$ . It is clear that if  $a_i = 1$  for some  $i$ , say  $i = 1$ , then  $\theta(a_1, a_2, \dots, a_k)$  is a member of  $\mathcal{G}_2(C_{a_2+1}, C_{a_3+1}, \dots, C_{a_k+1})$  and thus  $\theta(a_1, a_2, \dots, a_k)$  is not  $\chi$ -unique. Assume therefore that  $a_i \geq 2$  for all  $i$ . For  $k = 4$ , Chen, Bao and Ouyang [2] found that  $\theta(a_1, a_2, a_3, a_4)$  may not be  $\chi$ -unique.

**Theorem 2.1 ([2])** (a) *Let  $a_1, a_2, a_3, a_4$  be integers with  $2 \leq a_1 \leq a_2 \leq a_3 \leq a_4$ . Then  $\theta(a_1, a_2, a_3, a_4)$  is  $\chi$ -unique if and only if  $(a_1, a_2, a_3, a_4) \neq (2, b, b+1, b+2)$  for any integer  $b \geq 2$ .*

(b) *The  $\chi$ -equivalence class of  $\theta(2, b, b+1, b+2)$  is*

$$\{\theta(2, b, b+1, b+2)\} \cup \mathcal{G}_2(\theta(3, b, b+1), C_{b+2}).$$

□

Thus the problem of the chromaticity of  $\theta(a_1, a_2, \dots, a_k)$  has been completely settled for  $k \leq 4$ . For  $k \geq 5$ , we have

**Theorem 2.2 ([14])** *For  $k \geq 5$ ,  $\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique if  $a_i \geq k - 1$  for  $i = 1, 2, \dots, k$ .* □

**Theorem 2.3 ([11])** *Let  $h \geq s + 1 \geq 2$  or  $s = h + 1$ . Then for  $k \geq 5$ ,  $\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique if  $a_2 - 1 = a_1 = h$ ,  $a_j = h + s$  ( $j = 3, \dots, k - 1$ ),  $a_k \geq h + s$  and  $a_k \notin \{2h, 2h + s, 2h + s - 1\}$ .* □

Theorems 2.2 and 2.3 do not include the case where  $a_1 = a_2 = \dots = a_k < k - 1$ .

**Theorem 2.4 ([4], [6] and [13])**  *$\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique if  $k \geq 2$  and  $a_1 = a_2 = \dots = a_k \geq 2$ .* □

### 3 $\chi$ -closed families of g.p. trees

A *k-polygon tree* is a graph obtained by edge-gluing a collection of  $k$  cycles successively, i.e., a graph in  $\mathcal{G}_2(C_{i_1}, C_{i_2}, \dots, C_{i_k})$  for some integers  $i_1, i_2, \dots, i_k$  with  $i_j \geq 3$  for all  $j = 1, 2, \dots, k$ . A *polygon-tree* is a  $k$ -polygon tree for some integer  $k$  with  $k \geq 1$ . A graph is called a *generalized polygon tree* (g.p. tree) if it is a subdivision of some polygon tree.

Let  $\mathcal{GP}$  denote the set of all g.p. trees. Dirac [3] and Duffin [5] proved independently that a 2-connected graph is a g.p. tree if and only if it contains no subdivision of  $K_4$ .

A family  $\mathcal{S}$  of graphs is said to be *chromatically closed* (or simply  $\chi$ -closed) if  $\bigcup_{G \in \mathcal{S}} [G] = \mathcal{S}$ . By using Dirac's and Duffin's result, Chao and Zhao [1] obtained the following result.

**Theorem 3.1 ([1])** *The set  $\mathcal{GP}$  is  $\chi$ -closed.* □

The family  $\mathcal{GP}$  can be partitioned further into  $\chi$ -closed subfamilies. Let  $G \in \mathcal{GP}$ . A pair  $\{x, y\}$  of non-adjacent vertices of  $G$  is called a *communication pair* if there are at least three independent  $x - y$  paths in  $G$ . Let  $c(G)$  denote the number of communication pairs in  $G$ . For any integer  $r \geq 1$ , let  $\mathcal{GP}_r$  be the family of all g.p. trees  $G$  with  $c(G) = r$ .

**Theorem 3.2 ([13])** *The family  $\mathcal{GP}_r$  is  $\chi$ -closed for every integer  $r \geq 1$ .* □

Let  $G$  be a g.p. tree. We call a pair  $\{x, y\}$  of vertices in  $G$  a *pre-communication pair* of  $G$  if there are at least three independent  $x$ - $y$  paths in  $G$ . If  $x$  and  $y$  are non-adjacent, then  $\{x, y\}$  is a communication pair. Assume that  $c(G) = 1$ . Then  $G$  is a subdivision of a  $k$ -polygon tree  $H$  for some  $k \geq 2$ . It is clear that  $G$  and  $H$  have the same pre-communication pairs. But not every pre-communication pair in  $H$  is a communication pair. Since  $c(G) = 1$ , only one pre-communication pair in  $H$  is transformed into a communication pair in  $G$ . If  $G$  has only one pre-communication pair, then  $G$  is a multibridge graph. Otherwise,  $G$  is an edge-gluing of a multibridge graph and some cycles. Therefore

$$\mathcal{GP}_1 = \bigcup_{k \geq 3} \bigcup_{\substack{3 \leq t \leq k \\ b_1, b_2, \dots, b_k \geq 2}} \mathcal{G}_2(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}+1}, \dots, C_{b_k+1}). \quad (3)$$

Hence we have

**Lemma 3.1** *Let  $a_i \geq 2$  for  $i = 1, 2, \dots, k$ , where  $k \geq 3$ . If  $H \sim \theta(a_1, a_2, \dots, a_k)$ , then  $H$  is either a  $k$ -bridge graph  $\theta(b_1, \dots, b_k)$  with  $b_i \geq 2$  for all  $i$  or an edge-gluing of a  $t$ -bridge graph  $\theta(b_1, \dots, b_t)$  with  $b_i \geq 2$  for all  $i$  and  $k - t$  cycles for some integer  $t$  with  $3 \leq t \leq k - 1$ .* □

Note that for  $G \in \mathcal{G}_2(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}+1}, \dots, C_{b_k+1})$ ,

$$e(G) = v(G) + k - 2. \quad (4)$$

## 4 A graph function

For any graph  $G$  and real number  $\tau$ , write

$$\Psi(G, \tau) = (-1)^{1+e(G)} (1 - \tau)^{e(G)-v(G)+1} P(G, 1 - \tau). \quad (5)$$

Observe that  $\Psi(G, \tau) = \Psi(H, \tau)$  if  $G \sim H$ . However, the converse is not true. For example,  $\Psi(G, \tau) = \Psi(G \cup mK_1, \tau)$  but  $G \not\sim G \cup mK_1$  for any  $m \geq 1$ , where  $G \cup mK_1$  is the graph obtained from  $G$  by adding  $m$  isolated vertices. However, we have

**Lemma 4.1** For graphs  $G$  and  $H$ , if  $G \sim H$ , then  $\Psi(G, \tau) = \Psi(H, \tau)$ ; if  $v(G) = v(H)$  and  $\Psi(G, \tau) = \Psi(H, \tau)$ , then  $G \sim H$ .

*Proof.* We need to prove only the second assertion. Observe from (5) that  $\Psi(G, \tau)$  is a polynomial in  $\tau$  with degree  $e(G) + 1$ . Thus  $e(G) = e(H)$ . Since  $v(G) = v(H)$  and  $\Psi(G, \tau) = \Psi(H, \tau)$ , we have  $P(G, 1 - \tau) = P(H, 1 - \tau)$ . Therefore  $G \sim H$ .  $\square$

Thus, by Lemma 4.1, for any graph  $G$ ,  $[G]$  is the set of graphs  $H$  such that  $v(H) = v(G)$  and  $\Psi(H, \tau) = \Psi(G, \tau)$ . In this paper, we shall use this property to study the chromaticity of  $\theta(a_1, a_2, \dots, a_k)$ . We first derive an expression for  $\Psi(\theta(a_1, a_2, \dots, a_k), \tau)$ .

The following lemma is true even if  $k = 1$  or  $a_i = 1$  for some  $i$ .

**Lemma 4.2** For positive integers  $k, a_1, a_2, \dots, a_k$ ,

$$\Psi(\theta(a_1, a_2, \dots, a_k), \tau) = \tau \prod_{i=1}^k (\tau^{a_i} - 1) - \prod_{i=1}^k (\tau^{a_i} - \tau). \quad (6)$$

*Proof.* By the deletion-contraction formula for chromatic polynomials, it can be shown that

$$\begin{aligned} & P(\theta(a_1, a_2, \dots, a_k), \lambda) \\ &= \frac{1}{\lambda^{k-1}(\lambda - 1)^{k-1}} \prod_{i=1}^k ((\lambda - 1)^{a_i+1} + (-1)^{a_i+1}(\lambda - 1)) \\ & \quad + \frac{1}{\lambda^{k-1}} \prod_{i=1}^k ((\lambda - 1)^{a_i} + (-1)^{a_i}(\lambda - 1)). \end{aligned}$$

Let  $\tau = 1 - \lambda$ . Then

$$\begin{aligned} & (-1)^{1+a_1+a_2+\dots+a_k} (1 - \tau)^{k-1} P(\theta(a_1, a_2, \dots, a_k), 1 - \tau) \\ &= \tau \prod_{i=1}^k (\tau^{a_i} - 1) - \prod_{i=1}^k (\tau^{a_i} - \tau). \end{aligned}$$

Since  $v(G) = 2 - k + \sum_{i=1}^k a_i$  and  $e(G) = \sum_{i=1}^k a_i$ , by definition of  $\Psi(G, \tau)$ , (6) is obtained.  $\square$

**Corollary 4.1** For positive integers  $k, a_1, a_2, \dots, a_k$ ,

$$\begin{aligned} \Psi(\theta(a_1, a_2, \dots, a_k), \tau) &= (-1)^k (\tau - \tau^k) \\ & \quad + \sum_{\substack{1 \leq r \leq k \\ 1 \leq i_1 < i_2 < \dots < i_r \leq k}} (-1)^{k-r} (\tau - \tau^{k-r}) \tau^{a_{i_1} + a_{i_2} + \dots + a_{i_r}}. \end{aligned} \quad (7)$$

$\square$

We are now going to find an expression for  $\Psi(H, \tau)$  for any  $H$  in

$$\mathcal{G}_2(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}+1}, \dots, C_{b_k+1}).$$

**Lemma 4.3** *Let  $G$  and  $H$  be non-empty graphs, and  $M \in \mathcal{G}_2(G, H)$ . Then*

$$\Psi(M, \tau) = \Psi(G, \tau)\Psi(H, \tau)/((- \tau)(1 - \tau)). \quad (8)$$

*Proof.* Since  $v(M) = v(G) + v(H) - 2$ ,  $e(M) = e(G) + e(H) - 1$  and

$$P(M, \lambda) = P(G, \lambda)P(H, \lambda)/(\lambda(\lambda - 1)), \quad (9)$$

by (5), (8) is obtained.  $\square$

**Lemma 4.4** *Let  $k, t, b_1, b_2, \dots, b_k$  be integers with  $3 \leq t < k$  and  $b_i \geq 1$  for  $i = 1, 2, \dots, k$ . If  $H \in \mathcal{G}_2(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}+1}, \dots, C_{b_k+1})$ , then*

$$\Psi(H, \tau) = \tau \prod_{i=1}^k (\tau^{b_i} - 1) - \prod_{i=1}^t (\tau^{b_i} - \tau) \prod_{i=t+1}^k (\tau^{b_i} - 1). \quad (10)$$

*Proof.* By (5), we have  $\Psi(C_{b_{i+1}}, \tau) = (-\tau)(1 - \tau)(\tau^{b_i} - 1)$ . Thus by (6) and (8), (10) is obtained.  $\square$

## 5 $\chi$ -unique multibridge graphs

By Lemma 4.2, we can prove that  $\theta(a_1, a_2, \dots, a_k) \cong \theta(b_1, b_2, \dots, b_k)$  if  $\theta(a_1, a_2, \dots, a_k) \sim \theta(b_1, b_2, \dots, b_k)$ .

**Lemma 5.1** *Let  $a_i$  and  $b_i$  be integers with  $1 \leq a_1 \leq a_2 \leq \dots \leq a_k$  and  $1 \leq b_1 \leq b_2 \leq \dots \leq b_k$ , where  $k \geq 3$ . If*

$$\theta(a_1, a_2, \dots, a_k) \sim \theta(b_1, b_2, \dots, b_k), \quad (11)$$

then  $b_i = a_i$  for  $i = 1, 2, \dots, k$ .

*Proof.* By Lemma 4.1 and Corollary 4.1, we have

$$\begin{aligned} & \sum_{\substack{1 \leq r \leq k \\ 1 \leq i_1 < i_2 < \dots < i_r \leq k}} (-1)^{k-r} (\tau - \tau^{k-r}) \tau^{a_{i_1} + a_{i_2} + \dots + a_{i_r}} \\ = & \sum_{\substack{1 \leq r \leq k \\ 1 \leq i_1 < i_2 < \dots < i_r \leq k}} (-1)^{k-r} (\tau - \tau^{k-r}) \tau^{b_{i_1} + b_{i_2} + \dots + b_{i_r}}, \end{aligned} \quad (12)$$

after we cancel the terms  $(-1)^k(\tau - \tau^k)$  from both sides. The terms with lowest power in both sides have powers  $1 + a_1$  and  $1 + b_1$  respectively. Hence  $a_1 = b_1$ .

Suppose that  $a_i = b_i$  for  $i = 1, \dots, m$  but  $a_{m+1} \neq b_{m+1}$  for some integer  $m$  with  $1 \leq m \leq k - 1$ . Since  $a_i = b_i$  for  $i = 1, 2, \dots, m$ , by (12), we have

$$\begin{aligned} & \sum_{\substack{1 \leq r \leq k \\ 1 \leq i_1 < i_2 < \dots < i_r \leq k \\ i_r > m}} (-1)^{k-r} (\tau - \tau^{k-r}) \tau^{a_{i_1} + a_{i_2} + \dots + a_{i_r}} \\ = & \sum_{\substack{1 \leq r \leq k \\ 1 \leq i_1 < i_2 < \dots < i_r \leq k \\ i_r > m}} (-1)^{k-r} (\tau - \tau^{k-r}) \tau^{b_{i_1} + b_{i_2} + \dots + b_{i_r}}. \end{aligned} \quad (13)$$

The terms with lowest power in both sides of (13) have powers  $1 + a_{m+1}$  and  $1 + b_{m+1}$  respectively. Hence  $a_{m+1} = b_{m+1}$ , a contradiction. Therefore  $b_i = a_i$  for  $i = 1, 2, \dots, k$ .  $\square$

Let  $a_i$  be an integer with  $a_i \geq 2$  for  $i = 1, 2, \dots, k$  and suppose that  $a_1 \leq a_2 \leq \dots \leq a_k$ . We shall show that  $\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique if  $a_k < a_1 + a_2$ . It is well known (see [8]) that

**Lemma 5.2** *If  $G \sim H$ , then  $g(G) = g(H)$ .*  $\square$

**Theorem 5.1** *If  $2 \leq a_1 \leq a_2 \leq \dots \leq a_k < a_1 + a_2$ , where  $k \geq 3$ , then  $\theta(a_1, a_2, \dots, a_k)$  is chromatically unique.*

*Proof.* By Theorem 2.2, we may assume that  $a_1 \leq k - 2$ .

By Lemmas 3.1 and 5.1, it suffices to show that  $\theta(a_1, a_2, \dots, a_k) \not\sim H$  for any graph  $H \in \mathcal{G}_2(\theta(b_1, b_2, \dots, b_t), C_{b_{t+1}+1}, \dots, C_{b_k+1})$ , where  $t$  and  $b_i$  are integers with  $3 \leq t < k$  and  $b_i \geq 2$  for  $i = 1, 2, \dots, k$ . We may assume that  $b_1 \leq b_2 \leq \dots \leq b_t$  and  $b_{t+1} \leq \dots \leq b_k$ .

Suppose that  $H \sim \theta(a_1, a_2, \dots, a_k)$ . The girth of  $\theta(a_1, a_2, \dots, a_k)$  is  $a_1 + a_2$ . Since

$$g(H) = \min \left\{ \min_{1 \leq i < j \leq t} (b_i + b_j), \min_{t+1 \leq i \leq k} (b_i + 1) \right\}, \quad (14)$$

by Lemma 5.2, we have  $g(H) = a_1 + a_2$  and

$$\begin{cases} b_i + b_j \geq a_1 + a_2, & 1 \leq i < j \leq t, \\ b_i \geq a_1 + a_2 - 1, & t + 1 \leq i \leq k. \end{cases} \quad (15)$$

As  $e(H) = e(\theta(a_1, a_2, \dots, a_k))$ , we have

$$a_1 + a_2 + \dots + a_k = b_1 + b_2 + \dots + b_k. \quad (16)$$

By Lemma 4.1, (6) and (10), we have

$$\tau \prod_{i=1}^k (\tau^{a_i} - 1) - \prod_{i=1}^k (\tau^{a_i} - \tau) = \tau \prod_{i=1}^k (\tau^{b_i} - 1) - \prod_{i=1}^t (\tau^{b_i} - \tau) \prod_{i=t+1}^k (\tau^{b_i} - 1). \quad (17)$$

We expand both sides of (17), delete  $(-1)^k \tau$  from them and keep only the terms with powers at most  $a_1 + a_2$ . Since  $a_i + a_j \geq a_1 + a_2$  and  $b_i + b_j \geq a_1 + a_2$  for all  $i, j$  with  $1 \leq i < j \leq k$ , we have

$$\begin{aligned} & (-1)^{k-1} \sum_{i=1}^k \tau^{a_i+1} + (-1)^{k-1} \tau^k + (-1)^k \sum_{i=1}^k \tau^{k-1+a_i} \\ \equiv & (-1)^{k-1} \sum_{i=1}^k \tau^{b_i+1} + (-1)^{k-1} \tau^t + (-1)^k \sum_{i=1}^t \tau^{b_i+t-1} \\ & + (-1)^k \sum_{i=t+1}^k \tau^{b_i+t} \pmod{\tau^{a_1+a_2+1}}. \end{aligned} \tag{18}$$

Observe that  $b_i + t > a_1 + a_2$  for  $t + 1 \leq i \leq k$  and  $k - 1 + a_i > a_1 + a_2$  for  $2 \leq i \leq k$ . Thus

$$\begin{aligned} & (-1)^{k-1} \sum_{i=1}^k \tau^{a_i+1} + (-1)^{k-1} \tau^k + (-1)^k \tau^{k-1+a_1} \\ \equiv & (-1)^{k-1} \sum_{i=1}^k \tau^{b_i+1} + (-1)^{k-1} \tau^t + (-1)^k \sum_{i=1}^t \tau^{b_i+t-1} \pmod{\tau^{a_1+a_2+1}}. \end{aligned}$$

Hence

$$\sum_{i=1}^k \tau^{a_i+1} + \tau^k + \sum_{i=1}^t \tau^{b_i+t-1} \equiv \sum_{i=1}^k \tau^{b_i+1} + \tau^t + \tau^{k-1+a_1} \pmod{\tau^{a_1+a_2+1}}. \tag{19}$$

Since  $t \geq 3$ , we have  $b_i + t - 1 > a_1 + a_2$  for  $i \geq t + 1$ . If  $b_2 + t - 1 \leq a_1 + a_2$ , then since  $k - 1 + a_1 > k$ , the left side of (19) contains more terms with powers at most  $a_1 + a_2$  than does the right side, a contradiction. Hence  $b_i + t - 1 > a_1 + a_2$  for  $2 \leq i \leq t$ . Therefore

$$\sum_{i=1}^k \tau^{a_i+1} + \tau^k + \tau^{b_1+t-1} \equiv \sum_{i=1}^k \tau^{b_i+1} + \tau^t + \tau^{k-1+a_1} \pmod{\tau^{a_1+a_2+1}}. \tag{20}$$

Note that  $t \leq a_1 + a_2$ ; otherwise, since  $k > t$  and  $a_1, b_1 \geq 2$ , (20) becomes

$$\sum_{i=1}^k \tau^{a_i+1} = \sum_{i=1}^k \tau^{b_i+1},$$

which implies the equality of the multisets  $\{a_1, a_2, \dots, a_k\}$  and  $\{b_1, b_2, \dots, b_k\}$  in contradiction to (17).

**Claim 1:** There are no  $i, j$  such that

$$\{b_1, \dots, b_{i-1}, b_{i+1}, \dots, b_k\} = \{a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_k\}$$

as multisets.

Otherwise, by (16),  $\{b_1, \dots, b_k\} = \{a_1, \dots, a_k\}$  as multisets, which leads to a contradiction by (17).

**Claim 2:**  $a_2 \geq k - 1$ .

If  $a_2 < k - 1$ , then  $a_1 + k - 1 > a_1 + a_2$ . But  $a_i + 1 \leq a_1 + a_2$  for  $1 \leq i \leq k$ . So, by (20), the multiset  $\{a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_k\}$  is a subset of the multiset  $\{b_1, \dots, b_k\}$  for some  $j$  with  $1 \leq j \leq k$ , which contradicts Claim 1.

**Claim 3:**  $a_1 = t - 1$ .

Since  $\tau^t$  is a term of the right side of (20), the left side also contains  $\tau^t$ . But  $k > t$ ,  $b_1 + t - 1 > t$  and, by Claim 2,  $a_i + 1 \geq k > t$  for  $i \geq 2$ . Therefore  $a_1 + 1 = t$ .

By Claim 3, (20) is simplified to

$$\sum_{i=2}^k \tau^{a_i+1} + \tau^k + \tau^{b_1+t-1} \equiv \sum_{i=1}^k \tau^{b_i+1} + \tau^{k-1+a_1} \pmod{\tau^{a_1+a_2+1}}. \quad (21)$$

**Claim 4:**  $b_1 = k - 1$ .

Note that  $k < a_1 + a_2$ , by Claim 2. As  $\tau^k$  is a term of the left side of (21), the right side also contains this term. Thus  $b_i + 1 = k$  for some  $i$ . If  $i > t$ , then by (15) and Claims 2 and 3, we

$$b_i \geq a_1 + a_2 - 1 \geq k + t - 3 \geq k,$$

a contradiction. Thus  $i \leq t$  and  $b_1 \leq b_i = k - 1$ . If  $b_1 \leq k - 2$ , then the right side of (21) has a term with power at most  $k - 1$ . But the left side has no such term, a contradiction. Hence  $b_1 = k - 1$ .

By Claims 3 and 4, we have  $\tau^{b_1+t-1} = \tau^{k-1+a_1}$ . Thus (21) is further simplified to

$$\sum_{i=2}^k \tau^{a_i+1} + \tau^k \equiv \sum_{i=1}^k \tau^{b_i+1} \pmod{\tau^{a_1+a_2+1}}. \quad (22)$$

Therefore the multiset  $\{a_2, a_3, \dots, a_k\}$  is a subset of the multiset  $\{b_1, b_2, \dots, b_k\}$ , in contradiction to Claim 1.

Therefore  $H \not\sim \theta(a_1, a_2, \dots, a_k)$  and we conclude that  $\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique.  $\square$

## 6 $\chi$ -equivalent graphs

In Section 5, we proved that  $\theta(a_1, a_2, \dots, a_k)$  is  $\chi$ -unique if

$$\max_{1 \leq i \leq k} a_i < \min_{1 \leq i < j \leq k} (a_i + a_j). \quad (23)$$

Lemma 6.1 shows that, for any non-negative integer  $n$ , there exist examples where the graph  $\theta(a_1, a_2, \dots, a_k)$  is not  $\chi$ -unique and

$$\max_{1 \leq i \leq k} a_i - \min_{1 \leq i < j \leq k} (a_i + a_j) = n. \quad (24)$$

**Lemma 6.1** (i)  $\theta(2, 2, 2, 3, 4) \sim H$  for every  $H \in \mathcal{G}_2(\theta(2, 2, 3), C_4, C_4)$ .

(ii) For  $k \geq 4$  and  $a \geq 2$ ,  $\theta(k - 2, a, a + 1, \dots, a + k - 2) \sim H$  for every  $H \in \mathcal{G}_2(\theta(k - 1, a, a + 1, \dots, a + k - 3), C_{a+k-2})$ .

(iii) For  $k \geq 5$ ,  $\theta(2, 3, \dots, k-1, k, k-3) \sim H$  for every graph  $H$  in  $\mathcal{G}_2(\theta(2, 3, \dots, k-1), C_{k-1}, C_k)$ .  $\square$

It is straightforward to verify Lemma 6.1 by using Lemmas 4.1, 4.2 and 4.4.

It is natural to ask the following question: for which choices of  $(a_1, a_2, \dots, a_k)$  satisfying  $k \geq 5$  and

$$\max_{1 \leq i \leq k} a_i = \min_{1 \leq i < j \leq k} (a_i + a_j)$$

is the graph  $\theta(a_1, a_2, \dots, a_k)$  chromatically unique? If  $\theta(a_1, a_2, \dots, a_k)$  is not  $\chi$ -unique, what is its  $\chi$ -equivalence class? The solution to this question will be given in another paper.

## References

- [1] C.Y. Chao and L.C. Zhao, Chromatic polynomials of a family of graphs, *Ars. Combin.* **15** (1983) 111-129.
- [2] X.E. Chen, X.W. Bao and K.Z. Ouyang, Chromaticity of the graph  $\theta(a, b, c, d)$ , *J. Shaanxi Normal Univ.* **20** (1992) 75-79.
- [3] G.A. Dirac, A property of 4-chromatic graphs and some results on critical graphs, *J. London Math. Soc.* **27** (1952) 85-92.
- [4] F.M. Dong, On chromatic uniqueness of two infinite families of graphs, *J. Graph Theory* **17** (1993) 387-392.
- [5] R.J. Duffin, Topology of series-parallel networks, *J. Math. Anal. Appl.* **10** (1965) 303-318.
- [6] K.M. Koh and C.P. Teo, Some results on chromatically unique graphs, *Proc. Asian Math. Conf.* (World Scientific, Singapore, 1990) 258-262.
- [7] K.M. Koh and C.P. Teo, Chromaticity of series-parallel graphs, *Discrete Math.* **154** (1996) 289-295.
- [8] K.M. Koh and K.L. Teo, The search for chromatically unique graphs, *Graphs and Combinatorics* **6** (1990) 259-285.
- [9] K.M. Koh and K.L. Teo, The search for chromatically unique graphs -II, *Discrete Math.* **172** (1997) 59-78.
- [10] B. Loerinc, Chromatic uniqueness of the generalized  $\theta$ -graphs, *Discrete Math.* **23** (1978) 313-316.

- [11] Y.H. Peng, On the chromatic coefficients of a graph and chromatic uniqueness of certain  $n$ -partition graphs, in: *Combinatorics, Graph Theory, Algorithms and Applications* (Beijing , 1993) (World Scientific, River Edge, NJ, 1994) 307-316.
- [12] C.D. Wakelin and D.R. Woodall, Chromatic polynomials, polygon trees and outer-planar graphs, *J. Graph Theory* **16** (1992) 459-466.
- [13] S.J. Xu, Classes of chromatically equivalent graphs and polygon trees, *Discrete Math.* **133** (1994) 267-278.
- [14] S.J. Xu, J.J. Liu and Y.H. Peng, The chromaticity of  $s$ -bridge graphs and related graphs, *Discrete Math.* **135** (1994) 349-358.
- [15] A.A. Zykov, On some properties of linear complexes, *Amer. Math. Soc. Transl. No.* **79** (1952); translated from *Math. Sb.* **24**(66) (1949) 163-188.