

A note on an identity of Andrews

Zhizheng Zhang*

Department of Mathematics, Luoyang Teachers' College,
Luoyang 471022, P. R. China
zhzhzhang-yang@163.com

Submitted: Jan 26, 2005; Accepted: Feb 23, 2005; Published: Mar 7, 2005

Mathematics Subject Classifications: 33D15, 05A30

Abstract

In this note we use the q -exponential operator technique on an identity of Andrews.

1 Introduction

The following formula is equivalent to an identity of Andrews (see [3] or [1]):

$$\begin{aligned} & d \sum_{n=0}^{\infty} \frac{(q/bc, acdf; q)_n}{(ad, df; q)_{n+1}} (bd)^n - c \sum_{n=0}^{\infty} \frac{(q/bd, acdf; q)_n}{(ac, cf; q)_{n+1}} (bc)^n \\ &= d \frac{(q, qd/c, c/d, abcd, acdf, bcdf; q)_{\infty}}{(ac, ad, cf, df, bc, bd; q)_{\infty}}. \end{aligned} \quad (1)$$

Liu [3] showed it can be derived from the Ramanujan ${}_1\psi_1$ summation formula by the q -exponential operator techniques. In this short note, again using the q -exponential operator technique on it, we obtain a generalization of this identity. We have

Theorem 1.1. *Let $0 < |q| < 1$. Then*

$$\begin{aligned} & d \sum_{n=0}^{\infty} \frac{(q/bc, q/ce, acdf; q)_n}{(ad, df; q)_{n+1} (q^2/bcde; q)_n} q^n - c \sum_{n=0}^{\infty} \frac{(q/bd, q/de, acdf; q)_n}{(ac, cf; q)_{n+1} (q^2/bcde; q)_n} q^n \\ &= d \frac{(q, qd/c, c/d, abcd, acdf, bcdf, acde, cdef, bcde/q; q)_{\infty}}{(ac, ad, cf, df, bc, bd, ce, de, abc^2d^2ef/q; q)_{\infty}}. \end{aligned} \quad (2)$$

*This research is supported by the National Natural Science Foundation of China (Grant No. 10471016).

2 The proof of the Theorem

The q -difference operator and the q -shift operator η are defined by

$$D_q\{f(a)\} = \frac{1}{a}(f(a) - f(aq))$$

and

$$\eta\{f(a)\} = f(aq),$$

respectively. In [2] Chen and Liu construct the operator

$$\theta = \eta^{-1}D_q.$$

Based on these, they introduce a q -exponential operator:

$$E(b\theta) = \sum_{n=0}^{\infty} \frac{(b\theta)^n q^{\binom{n}{2}}}{(q; q)_n}.$$

For $E(b\theta)$, there hold the following operator identities.

$$E(b\theta) \{(at; q)_{\infty}\} = (at, bt; q)_{\infty}, \quad (3)$$

$$E(b\theta) \{(as, at; q)_{\infty}\} = \frac{(as, at, bs, bt; q)_{\infty}}{(abst/q; q)_{\infty}}. \quad (4)$$

Applying

$$(q/a; q)_n = (-a)^{-n} q^{\binom{n+1}{2}} \frac{(q^{-n}a; q)_{\infty}}{(a; q)_{\infty}}, \quad (5)$$

we rewrite (1) as

$$\begin{aligned} & d \sum_{n=0}^{\infty} \frac{(acdf; q)_n}{(ad, df; q)_{n+1}} \left(-\frac{d}{c}\right)^n q^{\binom{n+1}{2}} \cdot \{(q^{-n}bc, bd; q)_{\infty}\} \\ & - c \sum_{n=0}^{\infty} \frac{(acdf; q)_n}{(ac, cf; q)_{n+1}} \left(-\frac{c}{d}\right)^n q^{\binom{n+1}{2}} \cdot \{(q^{-n}bd, bc; q)_{\infty}\} \\ & = d \frac{(q, qd/c, c/d, acdf; q)_{\infty}}{(ac, ad, cf, df; q)_{\infty}} \cdot \{(abcd, bcdf; q)_{\infty}\}. \end{aligned} \quad (6)$$

Applying $E(e\theta)$ to both sides of the equation with respect to the variable b gives

$$\begin{aligned} & d \sum_{n=0}^{\infty} \frac{(acdf; q)_n}{(ad, df; q)_{n+1}} \left(-\frac{d}{c}\right)^n q^{\binom{n+1}{2}} \cdot E(e\theta) \{(q^{-n}bc, bd; q)_{\infty}\} \\ & - c \sum_{n=0}^{\infty} \frac{(acdf; q)_n}{(ac, cf; q)_{n+1}} \left(-\frac{c}{d}\right)^n q^{\binom{n+1}{2}} \cdot E(e\theta) \{(q^{-n}bd, bc; q)_{\infty}\} \\ & = d \frac{(q, qd/c, c/d, acdf; q)_{\infty}}{(ac, ad, cf, df; q)_{\infty}} \cdot E(e\theta) \{(abcd, bcdf; q)_{\infty}\}. \end{aligned} \quad (7)$$

Again, applying the results (3) and (4) of Chen and Liu, we have

$$E(e\theta) \{(q^{-n}bc, bd; q)_\infty\} = \frac{(q^{-n}bc, bd, q^{-n}ce, de; q)_\infty}{(q^{-n}bcde/q; q)_\infty}, \quad (8)$$

$$E(e\theta) \{(q^{-n}bd, bc; q)_\infty\} = \frac{(q^{-n}bd, bc, q^{-n}de, ce; q)_\infty}{(q^{-n}bcde/q; q)_\infty} \quad (9)$$

and

$$E(e\theta) \{(abcd, bcdf; q)_\infty\} = \frac{(abcd, bcdf, acde, cdef; q)_\infty}{(abc^2d^2ef/q; q)_\infty}. \quad (10)$$

Substituting these three identities into (7) and then using

$$(q^{-n}a; q)_\infty = (-a)^n q^{-\binom{n+1}{2}} (q/a; q)_n (a; q)_\infty, \quad (11)$$

we have

$$\begin{aligned} & d \frac{(bd, de, bc, ce; q)_\infty}{(bcde/q; q)_\infty} \sum_{n=0}^{\infty} \frac{(q/bc, q/ce, acdf; q)_n}{(ad, df; q)_{n+1} (q^2/bcde; q)_n} q^n \\ & - c \frac{(bc, ce, bd, de; q)_\infty}{(bcde/q; q)_\infty} \sum_{n=0}^{\infty} \frac{(q/bd, q/de, acdf; q)_n}{(ac, cf; q)_{n+1} (q^2/bcde; q)_n} q^n \\ & = d \frac{(q, qd/c, c/d, acdf, abcd, bcdf, acde, cdef; q)_\infty}{(ac, ad, cf, df, abc^2d^2ef/q; q)_\infty}. \end{aligned} \quad (12)$$

Hence we get

$$\begin{aligned} & d \sum_{n=0}^{\infty} \frac{(q/bc, q/ce, acdf; q)_n}{(ad, df; q)_{n+1} (q^2/bcde; q)_n} q^n - c \sum_{n=0}^{\infty} \frac{(q/bd, q/de, acdf; q)_n}{(ac, cf; q)_{n+1} (q^2/bcde; q)_n} q^n \\ & = d \frac{(q, qd/c, c/d, abcd, acdf, bcdf, acde, cdef, bcde/q; q)_\infty}{(ac, ad, cf, df, bc, bd, ce, de, abc^2d^2ef/q; q)_\infty}. \end{aligned} \quad (13)$$

The proof is completed.

References

- [1] G. E. Andrews, *Ramanujan's "lost" notebook. I. Partial θ -functions*, Adv. in Math. 41 (1981), 137-172 .
- [2] W. Y. C. Chen - Z. G. Liu, *Parameter augmentation for basic hypergeometric series, I*, in: B. E. Sagan, R. P. Stanley (Eds.), *Mathematical Essays in honor of Gian-Carlo Rota*, Birkhauser, Basel. 1998, pp. 111-129
- [3] Z. G. Liu, *Some operator identities and q -series transformation formulas*, Discrete Math. 265(2003), 119-139.