

# The Structure of Maximum Subsets of $\{1, \dots, n\}$ with No Solutions to $a + b = kc$

Andreas Baltz\*

Mathematisches Seminar  
University of Kiel, D-24098 Kiel, Germany  
aba@numerik.uni-kiel.de

Peter Hegarty, Jonas Knape, Urban Larsson

Department of Mathematics  
Chalmers University of Technology  
Göteborg, Sweden  
hegarty@math.chalmers.se {md9jonas,md0larur}@mdstud.chalmers.se

Tomasz Schoen \*\*

Wydział Matematyki i Informatyki  
Adam Mickiewicz University  
Poznań, Poland  
schoen@amu.edu.pl

Submitted: Nov 9, 2004; Accepted: Apr 22, 2005; Published: Apr 28, 2005

MR Subject Classifications: 05D05, 11P99

## Abstract

If  $k$  is a positive integer, we say that a set  $A$  of positive integers is  $k$ -sum-free if there do not exist  $a, b, c$  in  $A$  such that  $a + b = kc$ . In particular we give a precise characterization of the structure of maximum sized  $k$ -sum-free sets in  $\{1, \dots, n\}$  for  $k \geq 4$  and  $n$  large.

## 1 Introduction

A set of positive integers is called  $k$ -sum-free if it does not contain elements  $a, b, c$  such that

$$a + b = kc,$$

---

\*supported by DFG, Grant SR 7/9 – 2

\*\*research partially supported by KBN Grant 2 PO3A 007 24

where  $k$  is a positive integer. Denote by  $f(n, k)$  the maximum cardinality of a  $k$ -sum-free set in  $\{1, \dots, n\}$ . For  $k = 1$  these extremal sets are well-known: Deshouillers, Freiman, Sós, and Temkin [1] proved in particular that the maximum 1-sum-free sets in  $\{1, \dots, n\}$  are precisely the set of odd numbers and the “top half”  $\{\lceil \frac{n+1}{2} \rceil, \dots, n\}$ . For  $n > 8$  even  $\{\frac{n}{2}, \dots, n-1\}$  forms the only additional extremal set. The famous theorem of Roth [4] gives  $f(n, 2) = o(n)$ . Chung and Goldwasser [2] solved the case  $k = 3$  by showing that the set of odd integers is the unique extremal set for  $n > 22$ . For  $k \geq 4$  they gave an example of a  $k$ -sum-free set [3] of cardinality  $\frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n + \mathcal{O}(1)$ , which implies  $\lim_{n \rightarrow \infty} \frac{f(n, k)}{n} \geq \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}$ , and they conjectured that this lower bound is the actual value. Moreover they conjectured that extremal  $k$ -sum-free sets consist of three intervals of consecutive integers with slight modifications at the end-points if  $n$  is large.

In this paper we prove that the first conjecture is true, and we expose a structural result which is very close to the second. Our proof is elementary. In fact it is based on two simple observations:

Suppose we are given a  $k$ -sum-free set  $A$ . Then

- $kx - y \notin A$  for all  $x, y \in A$   
(Otherwise we could satisfy the equation  $kx = (kx - y) + y$  in  $A$ .)
- for all  $y \in A$  any interval centered around  $\frac{ky}{2}$  cannot share more than half of its elements with  $A$ .  
(Otherwise we would find a pair  $\lfloor \frac{ky}{2} \rfloor - d, \lceil \frac{ky}{2} \rceil + d$  in  $A$ , giving  $(\lfloor \frac{ky}{2} \rfloor - d) + (\lceil \frac{ky}{2} \rceil + d) = ky$ .)

## 2 Preparations

Let  $n \in \mathbb{N}$  be large and let  $k \in \mathbb{N}_{\geq 4}$ . We start by agreeing on some notations.

### Notations

Let  $A \subseteq \{1, \dots, n\}$  be a set of positive integers. Denote by

$$s_A := \min A \text{ and } m_A := \max A$$

the smallest and the largest elements of  $A$  respectively.

For  $l, r \in \mathbb{R}$  let

$$\begin{aligned} (l, r] &:= \{x \in \mathbb{N} \mid l < x \leq r\} \\ [l, r) &:= \{x \in \mathbb{N} \mid l \leq x < r\} \\ (l, r) &:= \{x \in \mathbb{N} \mid l < x < r\} \\ [l, r] &:= \{x \in \mathbb{N} \mid l \leq x \leq r\} \end{aligned}$$

abbreviate intervals of integers. Continuous intervals will be indicated by the subscript  $\mathbb{R}$ .

Furthermore for any  $y \in \mathbb{N}$  and  $d \in \mathbb{N}_0( := \mathbb{N} \cup \{0\} )$  put

$$I_y^d := \left[ \frac{ky - 1}{2} - d, \frac{ky + 1}{2} + d \right].$$

Note that if  $ky$  is even then  $I_y^d = \left\{ \frac{ky}{2} - d, \frac{ky}{2} - d + 1, \dots, \frac{ky}{2} + d \right\}$  and  $|I_y^d| = 2d + 1$ , while if  $ky$  is odd we have  $I_y^d = \left\{ \frac{ky-1}{2} - d, \dots, \frac{ky+1}{2} + d \right\}$  and  $|I_y^d| = 2d + 2$ .

The first Lemma restates our introductory observations.

**Lemma 1** *Let  $A \subseteq [1, n]$  be a  $k$ -sum-free set. If  $x, y \in A$  then  $kx - y \notin A$ . If  $y \in A$  and  $d \in \mathbb{N}_0$  then  $|I_y^d \setminus A| \geq d + 1$ .*

Suppose  $A'$  is a  $k$ -sum-free set consisting of intervals  $(l_i, r_i]$ . The interval  $(l_i, r_i]$  is  $k$ -sum-free if  $l_i \geq \frac{2r_i}{k}$ . Moreover we observe that reasonably large consecutive intervals  $(l_{i+1}, r_{i+1}], (l_i, r_i]$  (where we assume  $r_{i+1} < l_i$ ) should satisfy  $kr_{i+1} \leq l_i + s_{A'}$ . This leads to the following definition, describing a successive transformation of an arbitrary  $k$ -sum-free set  $A$  into a  $k$ -sum-free set of intervals.

**Definition 1** *Let  $n \in \mathbb{N}$  and let  $A \subseteq [1, n]$  be  $k$ -sum-free with smallest element  $s := s_A$ . Define sequences  $(r_i)$ ,  $(l_i)$ ,  $(A_i)$  by:*

$$\begin{aligned} A_0 &:= A, \quad r_1 := n, \\ l_i &:= \left\lfloor \frac{2r_i}{k} \right\rfloor, \quad r_{i+1} := \left\lfloor \frac{l_i + s}{k} \right\rfloor, \\ A_i &:= (A_{i-1} \setminus (r_{i+1}, l_i]) \cup (l_i, r_i] \cap [s, n] \text{ for } i \geq 1. \end{aligned}$$

The letter  $t = t_A$  will be reserved to denote the least integer such that  $r_{t+1} < s$ . Observe that, for all  $i \geq t$ ,

$$A_i = A_t = [\alpha, r_t] \cup \left( \bigcup_{j=1}^{t-1} (l_j, r_j] \right), \tag{1}$$

where  $\alpha = \alpha_A := \max\{l_t + 1, s\}$ .

### 3 The structure of maximum $k$ -sum-free sets

To obtain the structural result we consider the successive transformation of an arbitrary  $k$ -sum-free set  $A$  into a set  $A_t$  of intervals as in (1). Our plan is to show that each member of the transformation sequence  $(A_i)$  is  $k$ -sum-free and has size greater than or equal to  $|A|$ . For  $n$  sufficiently large, depending on  $k$ , and a maximum sized  $k$ -sum-free subset  $A$  of  $[1, n]$ , it will turn out that  $A_t$  consists of three intervals only, i.e.: that  $t = 3$ . This observation will do to determine  $f(n, k)$ , and we conclude our proof by showing that  $A$

could be enlarged if it did not contain (nearly) the whole interval  $(l_3, r_3]$  and consequently almost all elements from  $(l_2, r_2]$  and  $(l_1, r_1]$ , so that in fact almost nothing happens during the transformation of an extremal set.

**Lemma 2** *Let  $A \subseteq [1, n]$  be  $k$ -sum-free. Let  $i \in \mathbb{N}$ .*

a)  $A_i$  is  $k$ -sum-free.

b)  $|A_i| \geq |A_{i-1}|$ .

**Proof.** a) Clearly, it is enough to prove the claim for  $i \leq t$ , so we may assume that  $s \leq r_i$ . Suppose there are  $a, b, c \in A_i$  with  $a + b = kc$ .  $A_i$  is of the form

$$A_i = A_{i-1} \cap [s, r_{i+1}] \cup (l_i, r_i] \cap [s, n] \cup (l_{i-1}, r_{i-1}] \cup \dots \cup (l_1, r_1].$$

If  $c \in (l_1, r_1]$ , then  $kc > 2n$ , which is impossible. If  $i \geq 2$  and  $c \in (l_j, r_j]$  for some  $j \in [2, i]$ , then  $kc \in (2r_j, l_{j-1} + s]$  and the larger one of  $a, b$  must be in  $(r_j, l_{j-1}]$ . But  $(r_j, l_{j-1}] \cap A_i = \emptyset$  by construction. Hence  $c \in A_{i-1} \cap [s, r_{i+1}]$ . Now,  $kc \leq kr_{i+1} \leq l_i + s$ . Since  $(r_{i+1}, l_i] \cap A_i = \emptyset$ , both  $a$  and  $b$  have to be in  $A_{i-1} \cap [s, r_{i+1}] = A \cap [s, r_{i+1}]$ . But  $A$  is  $k$ -sum-free, a contradiction.

b) The inequality is trivial for  $i \geq t$ . For  $1 \leq i < t$  we have that  $l_i \geq s$  and hence

$$A_i = (A_{i-1} \cap [1, r_{i+1}]) \cup (l_i, r_i] \cup \left( \bigcup_{j=1}^{i-1} (l_j, r_j] \right).$$

Thus it suffices to prove that

$$|A_{i-1} \cap [1, r_i]| \leq |A_{i-1} \cap [1, r_{i+1}]| + \left\lceil \frac{(k-2)r_i}{k} \right\rceil.$$

Clearly, then, it suffices to prove the inequality for  $i = 1$ , i.e.: to prove that, for any  $n > 0$ , and any  $k$ -sum-free subset  $A$  of  $[1, n]$  with smallest element  $s_A$ , we have

$$|A| \leq |A \cap [1, r_{2,A}]| + \left\lceil \frac{(k-2)n}{k} \right\rceil, \tag{2}$$

where

$$r_{2,A} := \left\lceil \frac{\lfloor 2n/k \rfloor + s_A}{k} \right\rceil.$$

The proof is by induction on  $n$ . The result is trivial for  $n = 1$ . So suppose it holds for all  $1 \leq m < n$  and let  $A$  be a  $k$ -sum-free subset of  $[1, n]$ . Note that the result is again trivial if  $s_A > 2n/k$ , so we may assume that  $s_A \leq 2n/k$ , which implies that  $r_{2,A} \leq n/k$ , since  $k \geq 4$ .

First suppose that there exists  $x \in A \cap (n/k, 2n/k]$ . Then  $1 \leq kx - n \leq n$  and the

map  $f : y \mapsto kx - y$  is a 1-1 mapping from the interval  $[kx - n, n]$  to itself. For each  $y$  in this interval, at most one of the numbers  $y$  and  $f(y)$  can lie in  $A$ , since  $A$  is  $k$ -sum-free. To simplify notation, put  $w := kx - n - 1$ . Then our conclusion is that

$$|A \cap (w, n]| \leq \frac{1}{2}(n - w). \quad (3)$$

If  $w = 0$  or if  $A \cap [1, w] = \emptyset$ , then we are done (since  $k \geq 4$ ). Put  $B := A \cap [1, w]$ . Then we may assume  $B \neq \emptyset$ , hence  $s_B = s_A$ . Applying the induction hypothesis to  $B$ , we find that

$$|B| = |A \cap [1, w]| \leq |B \cap [1, r_{2,B}]| + \left\lceil \frac{(k-2)w}{k} \right\rceil. \quad (4)$$

But  $s_B = s_A$  implies that  $r_{2,B} \leq r_{2,A}$ , hence that  $B \cap [1, r_{2,B}] \subseteq A \cap [1, r_{2,A}]$ . Thus (3) and (4) yield the inequality

$$|A| \leq |A \cap [1, r_{2,A}]| + \left\lceil \frac{(k-2)w}{k} \right\rceil + \frac{1}{2}(n - w),$$

which in turn implies (2), since  $|A|$  is an integer. Thus we are reduced to completing the induction under the assumption that  $A \cap (n/k, 2n/k] = \emptyset$ . Suppose  $x \in A \cap (r_{2,A}, n/k]$ . Then  $\lfloor 2n/k \rfloor + s_A < kx \leq n$  and  $kx - s_A \notin A$ . In other words, we can pair off elements in  $A \cap (r_{2,A}, 2n/k]$  with elements in  $(2n/k, n] \setminus A$ . This immediately implies (2), and the proof of Lemma 2 is complete.  $\square$

We have seen so far that any  $k$ -sum-free set  $A$  can be turned into a  $k$ -sum-free set  $A_t$  having overall size at least  $|A|$ . The set  $A_t$  is a union of intervals, as given by (1), though note that the final interval  $[\alpha, r_t]$  may consist of a single point, since  $r_t = s$  is possible. The proof of the following Lemma uses a fact shown in [3] by Chung and Goldwasser, to prove that  $t$  must be equal to three if  $|A|$  is maximum.

**Lemma 3** *Let  $A$  be a maximum  $k$ -sum-free subset of  $[1, n]$ , where  $n > n_0(k)$  is sufficiently large. Let  $s := s_A$  and let  $t := \max\{i \in \mathbb{N} \mid r_i \geq s\}$ . Then  $t = 3$ .*

**Proof.** Let  $A_t$  be the set of positive integers given by (1). In a similar manner we now define a  $k$ -sum-free subset  $A'_t$  of  $(0, 1]_{\mathbb{R}}$ .

Put  $c := s/n$  and, for  $i = 1, \dots, t$  define real numbers  $R_i, L_i$  as follows :

$$R_1 := 1, \quad L_i := \frac{2R_i}{k}, \quad R_{i+1} := \frac{L_i + c}{k}.$$

Then we put

$$A'_t := [\alpha', R_t)_{\mathbb{R}} \cup \left( \bigcup_{j=1}^{t-1} [L_j, R_j)_{\mathbb{R}} \right),$$

where  $\alpha' := \max\{L_t, c\}$ . That  $A'_t$  is  $k$ -sum-free is shown in [3]. One sees easily that

$$|A_t| \leq n \cdot \mu(A'_t) + t, \tag{5}$$

where  $\mu$  denotes the Lebesgue-measure. Now suppose that  $t \neq 3$ . It is shown in [3] that there exists a constant  $c_k > 0$ , depending only on  $k$ , such that in this case

$$|\mu(A'_t)| \leq \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)} - c_k. \tag{6}$$

In fact, in the notation of page 8 of [3], an explicit value for  $c_k$  (which we will use later) is given by

$$c_k = \frac{2}{k}(R(3) - R(4)),$$

which by definition of  $R$  amounts to

$$c_k = \frac{8(k^4 - 4k^2 - 4)(k - 2)}{(k^6 - 2k^4 - 4k^2 - 8)(k^4 - 2k^2 - 4)k}. \tag{7}$$

Now (5) and (6) would imply that

$$|A| \leq \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n - c_k n + t.$$

But we have seen in the introduction that  $|A| \geq \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n + \mathcal{O}(1)$  and, since  $t = \mathcal{O}(\log_k n)$ , we thus have a contradiction for sufficiently large  $n$ . Hence  $t$  must equal three, for large enough  $n$ , as required.  $\square$

Now we are nearly in a position to determine  $f(n, k)$ . We want to calculate the cardinality of an extremal  $k$ -sum-free set  $A$  via computing  $|A_3|$ . Since  $|A_3|$  depends on  $s_A$ , the following lemma will be helpful :

**Lemma 4** *Let  $n > n_0(k)$  be sufficiently large. If  $A$  is a maximal  $k$ -sum-free subset of  $[1, n]$ , then  $S - 2k \leq s_A \leq S + 3$ , where  $S := \lfloor \frac{8n}{k^5 - 2k^3 - 4k} \rfloor$ .*

**Proof.** Set  $s := s_A$ . By Lemma 3, for  $n > n_0(k)$  we have  $r_4 < s$ . Since  $A$  is maximal we have  $|A| = |A_3|$ . Now, for a fixed  $n$ , the cardinality of  $A_3$  is a function of  $s \in [1, n]$  only. So we need to show that  $|A_3(s)|$  attains its maximum value only for some  $s \in [S - 2k, S + 3]$ . Define

$$s' := \min\{s \in [1, n] : l_3(s) < s\}.$$

A tedious computation (see the Appendix below) yields that  $s' = S + 1$  if  $k$  is even and  $s' = S$  or  $S + 1$  if  $k$  is odd. Hence

$$s' \in [S, S + 1]. \tag{8}$$

Clearly,

$$|A_3(s)| = \begin{cases} \lceil \frac{(k-2)n}{k} \rceil + r_2(s) - l_2(s) + r_3(s) - s + 1, & \text{if } s \geq s', \\ \lceil \frac{(k-2)n}{k} \rceil + r_2(s) - l_2(s) + r_3(s) - l_3(s), & \text{if } s < s'. \end{cases} \quad (9)$$

How does  $|A_3(s)|$  change (ignoring its maximality for a while) if we alter  $s$ ?

First suppose  $s \geq s'$ . If  $s$  increases by one, then  $|A_3|$  will decrease by one unless either  $r_2$  or  $r_3$  increases. Now  $r_2$  can only increase (by one) once in  $k(\geq 4)$  times. Almost the same is true of  $r_3$ , though its dependence on  $l_2$  makes things a little more complicated. However, it is not hard to see that we encounter an irreversible decrease in the cardinality of  $|A_3|$  after at most 3 steps of increment of  $s$ . Hence  $|A_3(s)| < |A_3(s')|$  if  $s \geq s' + 3$ .

Next suppose  $s < s'$ . If we decrease  $s$ , then  $|A_3|$  cannot increase at all, since  $l_i$  will not decrease unless  $r_i$  does. Moreover,  $|A_3|$  will become smaller if the size of any interval is diminished. So we can focus our attention on  $(l_2, r_2]$ . While  $r_2$  decreases once in  $k$  times,  $l_2$  does so no more than once in  $k \lfloor \frac{k}{2} \rfloor \geq 2k$  times. Thus  $|A_3(s)| < |A_3(s' - 1)|$  if  $s \leq s' - 1 - 2k$ .

We have now shown that, as a function of  $s \in [1, n]$ , the cardinality of  $A_3$  attains its maximum only for some  $s \in [s' - 2k, s' + 2]$ . This, together with (8), completes the proof of the lemma.  $\square$

Now we can prove the first conjecture of Chung and Goldwasser.

**Theorem 1**

$$\lim_{n \rightarrow \infty} \frac{f(n, k)}{n} = \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}.$$

**Proof.** Let  $A$  be a maximum  $k$ -sum-free set in  $[1, n]$ , with  $n$  sufficiently large. From Lemma 4 we have  $\frac{s_A}{n} = \frac{S^*}{n} + o(1)$ , where  $S^* = \frac{8n}{k^5-2k^3-4k}$ . Thus we can estimate

$$\begin{aligned} \frac{f(n, k)}{n} &= \frac{|A_3|}{n} = \frac{r_1 - l_1 + r_2 - l_2 + r_3 - S^* + 1}{n} + o(1) \\ &= \frac{1}{n} \left( n - \frac{2n}{k} + \frac{2n + kS^*}{k^2} - \frac{4n + 2kS^*}{k^3} + \frac{4n + 2kS^* + k^3S^*}{k^4} - S^* \right) + o(1) \\ &= \frac{k^4 - 2k^3 + 2k^2 - 4k + 4}{k^4} + \frac{S^*}{nk^3}(2k^2 - 2k + 2 - k^3) + o(1) \\ &= \frac{k^4 - 2k^3 + 2k^2 - 4k + 4}{k^4} + \frac{8(2k^2 - 2k + 2 - k^3)}{(k^5 - 2k^3 - 4k)k^3} + o(1) \\ &= \frac{k^5 - 2k^4 - 4k + 8}{(k^4 - 2k^2 - 4)k} + o(1) \\ &= \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)} + o(1), \end{aligned}$$

and the claim follows by taking the limit.  $\square$

We can now show the main result.

**Theorem 2** Let  $k \in \mathbb{N}_{\geq 4}$  and  $n > n_1(k)$ . Let  $S$  and  $s'$  be as in Lemma 4. Let  $A \subseteq \{1, \dots, n\}$  be a  $k$ -sum-free set of maximum cardinality, with smallest element  $s = s_A$ . Then  $s \in [S, S + 3]$  and  $A = \mathcal{I}_3 \cup \mathcal{I}_2 \cup \mathcal{I}_1$ , where

$$\begin{aligned} \mathcal{I}_3 &\in \begin{cases} \{[s, r_3], [s, r_3 + 1]\}, & \text{if } s \geq s' \\ \{[s, r_3], [s, r_3] \setminus \{r_3 - 1\}\}, & \text{if } s < s', \end{cases} \\ \mathcal{I}_2 &\in \begin{cases} \{[l_2 + 2, r_2], [l_2 + 2, r_2 + 1]\}, & \text{if } r_3 + 1 \in A \\ \{(l_2, r_2), (l_2, r_2 + 1), [l_2, r_2], [l_2, r_2] \setminus \{r_2 - 1\}\}, & \text{if } r_3 + 1 \notin A, \end{cases} \\ \mathcal{I}_1 &\in \begin{cases} \{[l_1 + 2, n]\}, & \text{if } r_2 + 1 \in A \\ \{[l_1, n], (l_1, n], [l_1, n] \setminus \{n - 1\}\}, & \text{if } r_2 + 1 \notin A, \end{cases} \end{aligned}$$

If  $k$  is even, then  $\mathcal{I}_i \neq [l_i, r_i] \setminus \{r_i - 1\}$  for  $1 \leq i \leq 3$ .

**Remark.** Note that Theorem 2 does not precisely determine the  $k$ -sum-free subsets of  $\{1, \dots, n\}$  of maximum size, for every  $n > n_1(k)$ . With  $n$  and  $k$  fixed, one first needs to determine for which value(s) of  $s \in [S, S + 3]$  the quantity  $|A_3(s)|$ , as given by (9), is maximized. The result will depend on  $n$  and  $k$ . Even then, for a fixed  $s$ , not all the possibilities for  $\mathcal{I}_3 \cup \mathcal{I}_2 \cup \mathcal{I}_1$  need be  $k$ -sum-free. See Section 4 below for further discussion.

**Proof.** We have already seen that  $|A_3| = |A|$ . Our first aim is to show by comparing  $A_3$  with  $A_2$  that almost the whole interval  $(l_3, r_3]$  must be in  $A$ . Having achieved this, we infer by Lemma 1 that  $(r_3, l_2] \cap A$  is nearly empty. Comparing  $A_2$  with  $A_1$  will then reveal that most of  $(l_2, r_2]$  is contained in  $A$ . Again Lemma 1 will help us to see that  $A$  cannot share many elements with  $(r_2, l_1]$  and a final comparison of  $A_1$  with  $A$  will conclude the proof.

(I) The first aim is easily reached if  $s := s_A \geq l_3 + 1$ . Simply note that

$$A_2 = (A \cap [s, r_3]) \cup (l_2, r_2] \cup (l_1, r_1] \subseteq [s, r_3] \cup (l_2, r_2] \cup (l_1, r_1] = A_3.$$

The maximality of  $|A_2|$  gives  $A_2 = A_3$  and hence  $[s, r_3] \subseteq A$ . Observe that  $s > l_3$  together with Lemma 4 and (8) give  $S \leq s \leq S + 3$ .

Assume now that  $s \leq l_3$ . We want to show that in this case  $s = l_3$ . Suppose  $s < l_3$  and let  $B = [S - 2k, l_3] \cap A$ . Define

$$C := I_{s_B}^1 \cup \bigcup_{b \in B \setminus \{s_B\}} I_b^0.$$

Clearly  $C \subseteq (l_3, r_3]$  for all  $n \gg 0$ . Then since  $C$  is the union of disjoint intervals, Lemma 1 gives that  $|C \setminus A| > |B|$ . Hence we get the contradiction  $|A_3| = |(A_2 \setminus B) \cup (l_3, r_3]| \geq |(A_2 \setminus B) \cup (C \setminus A)| > |A_2| - |B| + |B| = |A_2|$ . Therefore we are left with  $s = l_3$ , and this implies

$$|A_2| = |A_3| \iff |A \cap [s, r_3]| = |(l_3, r_3] \cap [s, r_3]| = |(s, r_3]|. \quad (10)$$



If  $r_3 \notin A$  we can infer from (10) that

$$A \cap [s, r_3] = [s, r_3 - 1] = [l_3, r_3 - 1].$$

If  $r_3 \in A$ , Lemma 1 gives  $kl_3 - r_3 \notin A$ , so  $-k + 1 \leq kl_3 - 2r_3 \leq -1$ . If  $kl_3 - 2r_3 \leq -2$  we get  $I_{l_3}^1 \subseteq (l_3, r_3]$  and  $|I_{l_3}^1 \setminus A| \geq 2$ , which is impossible since this would imply  $|A_3| > |A_2|$ . Hence  $kl_3 - 2r_3 = -1$  and  $k$  is odd. Using (10) one obtains

$$A \cap [s, r_3] = [l_3, r_3] \setminus \{r_3 - 1\}.$$

Suppose now that  $s = l_3$  and  $r_3 + 1 \in A$ . Then  $kl_3 - (r_3 + 1) \notin A$  and

$$r_3 - k \leq kl_3 - (r_3 + 1) \leq r_3 - 1.$$

This contradicts that  $[s, r_3 - 2] \subseteq A$  unless  $kl_3 - (r_3 + 1) = r_3 - 1$ , but then  $r_3 \notin A$  and  $|A \cap [s, r_3]| = |A \cap [s, r_3 - 2]|$  which contradicts (10). Hence  $r_3 + 1 \notin A$  if  $s = l_3$ .

Finally note that, if  $s = l_3$  and  $kl_3 \geq 2r_3 - 1$ , the latter being a requirement for either of the two possibilities for  $\mathcal{I}_3$  to be  $k$ -sum-free, then another computation similar to the one in the Appendix yields that  $s \geq S$ . Again, using Lemma 4 we obtain

$$S \leq s \leq S + 3, \tag{11}$$

as claimed in the statement of the theorem. This completes the first part of our proof.

**(II)** For the second part note that we have just shown

$$s \geq l_3. \tag{12}$$

Plugging (11) into the definition of  $l_3$  yields (after a further tedious computation similar to that in the Appendix)

$$S - 1 \leq l_3 \leq S + 1, \tag{13}$$

which implies in view of (12) and (11)

$$l_3 \leq s \leq l_3 + 4. \tag{14}$$

Moreover we have observed that  $[s, r_3 - 2] \subseteq A$ . Let  $\xi_1, \dots, \xi_5 \in \{0, \dots, k - 1\}$  be constants such that

$$kl_1 = 2r_1 - \xi_1 \tag{15}$$

$$kr_2 = l_1 + s - \xi_2 \tag{16}$$

$$kl_2 = 2r_2 - \xi_3 \tag{17}$$

$$kr_3 = l_2 + s - \xi_4 \tag{18}$$

$$kl_3 = 2r_3 - \xi_5. \tag{19}$$

We suppose that  $n$  is sufficiently large, so we can be sure that

$$[ks - (r_3 - 2), k(r_3 - 2) - s] \cap A = \emptyset.$$

By (14) we can infer that

$$\begin{aligned} \emptyset &= [k(l_3 + 4) - (r_3 - 2), k(r_3 - 2) - s] \cap A \\ &= [r_3 - \xi_5 + 4k + 2, l_2 - \xi_4 - 2k] \cap A. \end{aligned}$$

Let  $J = [r_3 + 2, r_3 - \xi_5 + 4k + 1] \cap A$  and  $K = \bigcup_{x \in J} \{kx - (s + 2), kx - (s + 1), kx - s\}$ . Then  $K \cap A = \emptyset$ ,  $|K| = 3|J|$  and by (18) and (19) we have

$$K \subseteq [l_2 - \xi_4 + 2k - 2, l_2 - \xi_4 - k\xi_5 + 4k^2 + k] \subseteq (l_2 + k - 2, l_2 + 4k^2 + k] \subseteq (l_2 + 2, r_2],$$

if  $n \gg 0$ . Let  $B = [l_2 - \xi_4 - 2k + 1, l_2] \cap A$ . If  $B \cup J \subseteq \{l_2\}$  then  $A \cap [r_3 + 2, l_2 - 1] = \emptyset$ . Otherwise, with  $C$  as in part (I) if  $|B| > 1$  we can verify that  $C \subseteq [r_2 - \frac{3k^2 - k + 2}{2}, r_2] \subseteq (l_2 + 1, r_2]$ , for  $n \gg 0$ , and  $|C \setminus A| > |B|$ . Put  $C := \emptyset$  if  $|B| \leq 1$ . For large  $n$ ,  $K$  and  $C$  are disjoint. Hence  $|B \cup J| < |(C \setminus A) \cup K|$  and we get

$$|A_2| = |[A_1 \setminus (J \cup B \cup \{r_3 + 1\})] \cup (l_2, r_2]| > |A_1 \setminus \{r_3 + 1\}|.$$

Thus if  $r_3 + 1 \notin A$  we get  $|A_2| > |A_1|$  so suppose  $r_3 + 1 \in A$ . Then neither  $l_2$  nor  $l_2 + 1$  can be in  $A_1$ . Otherwise, since  $(s - \xi_4 + k), s - \xi_4 + k - 1 \in [s, s + k] \subseteq [s, r_3 - 2] \subseteq A$  we get

$$k(r_3 + 1) = l_2 + (s - \xi_4 + k) = (l_2 + 1) + (s - \xi_4 + k - 1),$$

which is impossible. But  $l_2 + 1 \in A_2$ , so also in this case it follows that  $|A_2| > |A_1|$ , since  $l_2 + 1 \notin K \cup C$  for large  $n$ . Again we conclude that  $A \cap [r_3 + 2, l_2 - 1] = \emptyset$ . Consequently,

$$|A_2| = |A_1| \Leftrightarrow |A \cap ([l_2, r_2] \cup \{r_3 + 1\})| = |[l_2, r_2]|,$$

which gives  $A \cap [l_2, r_2] = [l_2 + 2, r_2]$  if  $r_3 + 1 \in A$ . If  $r_3 + 1 \notin A$  and either  $l_2 \notin A$  or  $r_2 \notin A$ , we get  $A \cap [l_2, r_2] = (l_2, r_2]$  or  $A \cap [l_2, r_2] = [l_2, r_2)$ , respectively. In case  $r_3 + 1 \notin A$  and both  $l_2, r_2 \in A$ , we see that  $kl_2 - r_2 = r_2 - \xi_3 \notin A$ . If  $\xi_3 \geq 2$  then  $I_{l_2}^1 \subseteq (l_2, r_2]$  and  $l_2$  could be profitably replaced. Hence  $\xi_3 = 1$ ,  $A \cap [l_2, r_2] = [l_2, r_2] \setminus \{r_2 - 1\}$  and  $k$  is odd.

**(III)** For the final interval  $(l_1, r_1]$  we use Lemma 1 to conclude from

$$[s, r_3 - 2] \subseteq A \text{ and } [l_2 + 2, r_2 - 2] \subseteq A$$

in view of (16) and (17) that, for  $n \gg 0$ ,

$$\begin{aligned} \emptyset &= A \cap [k(l_2 + 2) - (r_2 - 2), k(r_2 - 2) - (l_2 + 2)] \\ &= A \cap [r_2 - \xi_3 + 2k + 2, l_1 + s - \xi_2 - 2k - l_2 - 2], \text{ and} \\ \emptyset &= A \cap [k(l_2 + 2) - (r_3 - 2), k(r_2 - 2) - s] \\ &= A \cap [2r_2 - \xi_3 + 2k - r_3 + 2, l_1 - \xi_2 - 2k] \end{aligned}$$

Let  $J = [r_2 + 2, r_2 - \xi_3 + 2k + 1] \cap A$  and  $K = \cup_{x \in J} \{kx - s, kx - (s + 1), kx - (s + 2)\}$ . From (14) we have

$$K \subseteq [l_1 - \xi_2 + 2k - 2, l_1 - \xi_2 - k\xi_3 + 2k^2 + k] \subseteq (l_1 + k - 2, r_1], \text{ if } n \gg 0.$$

Let  $B = [l_1 - \xi_2 - 2k + 1, l_1] \cap A$ . If  $s_B < l_1$  with  $C$  as in (I) we can verify that, for sufficiently large  $n$ ,

$$C \subseteq \left[ \frac{2r_1 - \xi_1 - k\xi_2 - 2k^2 + k - 5}{2}, r_1 \right] \subseteq (l_1, r_1],$$

$|C \setminus A| > |B|$  and  $\max K < s_C$ . By analogy with part (II) we get  $A \cap [r_2 + 2, l_1 - 1] = \emptyset$  and the rest of the claim follows as before.  $\square$

## 4 Estimates and Periodicity

We first want to estimate values of  $n_i(k)$ ,  $i = 0, 1$ , for which Lemmas 3 and 4, and Theorem 2 respectively are valid. The estimates we shall arrive at can probably be improved upon. The example of a  $k$ -sum-free set  $A$  in [3], referred to in the proof of Lemma 3, satisfies

$$|A| > \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n - 3.$$

Hence the proof of Lemma 3 goes through provided  $n$  is sufficiently large so that

$$c_k n - t_0 \geq 3, \tag{20}$$

where  $t_0 = t_0(n, k)$  is the largest possible value for  $t$  in Definition 1. Now from Definition 1 we easily deduce that, if  $i < t$ , then  $r_{i+1} \leq \left(\frac{4}{k^2}\right)r_i$ , and hence that  $r_t \leq \left(\frac{4}{k^2}\right)^{t-1}n$ . Since  $r_t \geq 1$  a priori, we can thus estimate

$$t_0 \leq \frac{1}{2} \log_{k/2} n + 1. \tag{21}$$

Since, by (7),  $c_k = \mathcal{O}\left(\frac{1}{k^6}\right)$ , we thus deduce from (18) and (19) that one can take  $n_0(k) = \mathcal{O}(k^6)$ . It is then an easy and tedious exercise to go through the proof of Theorem 2 and check that one can also take  $n_1(k) = \mathcal{O}(k^6)$ .

Next, we explain what we mean by the word ‘periodicity’ in the title of this section. If  $k \geq 4$  is even then, for  $n > 0$ , we have  $s' = S + 1 = \lfloor \frac{8n}{k^5 - 2k^3 - 4k} \rfloor + 1$ . Hence for a fixed  $k$ , if we regard  $s'$  as a function of  $n$ , then  $s'(n) + 1 = s'(n + p_k)$ , where  $p_k := \frac{k^5 - 2k^3 - 4k}{8}$ . For odd  $k$ , we define  $p_k := k^5 - 2k^3 - 4k$  and in this case, a little more care is required to check that  $s'(n) + 8 = s'(n + p_k)$ .

Now for any  $k$  and  $n$ , let  $\mathcal{F}(k, n)$  denote the family of maximal  $k$ -sum-free subsets of  $\{1, \dots, n\}$ . Then for  $n$  sufficiently large, as estimated above, and  $k$  even (resp.  $k$  odd), the map  $s \mapsto s + 1$  (resp.  $s \mapsto s + 8$ ) clearly induces a 1-1 correspondence between the sets in  $\mathcal{F}(k, n)$  and  $\mathcal{F}(k, n + p_k)$ . This is what we mean by ‘periodicity’. This observation clearly reduces, for any fixed  $k$ , the full classification of all  $k$ -sum-free subsets of  $\{1, \dots, n\}$ , for all  $n$ , to a finite computation.

As an example, we now look at  $k = 4$ . By (7) we compute  $c_4 = \frac{47}{48290}$ . Then Lemma 3 is valid at least for all  $n$  satisfying

$$c_4 n - \frac{1}{2} \log_2 n - 1 \geq 3,$$

which reduces to  $n \geq 11008$ . One can then check that the proof of Theorem 2 also goes through for all such  $n$ . We have  $p_4 = 110$ . We now present the full classification of all 4-sum-free subsets of  $\{1, \dots, n\}$ , valid (at least) for all  $n \geq 11008$ . This was obtained with the help of a computer.

For each  $s, n \in \mathbf{N}$  we define the sets  $J_x(s)$ ,  $1 \leq x \leq 13$ , as follows (the  $l_i$  and  $r_i$  are functions of  $s$  and  $n$  as in Definition 1) :

$$\begin{aligned} J_1 &= [S, r_3 - 1] \cup [l_2, r_2 - 1] \cup [l_1, n - 1], \\ J_2 &= [S, r_3 - 1] \cup [l_2, r_2 - 1] \cup [l_1 + 1, n], \\ J_3 &= [S, r_3 - 1] \cup [l_2 + 1, r_2] \cup [l_1, n - 1], \\ J_4 &= [S, r_3 - 1] \cup [l_2 + 1, r_2] \cup [l_1 + 1, n], \\ J_5 &= [S, r_3 - 1] \cup [l_2 + 1, r_2 + 1] \cup [l_1 + 2, n], \\ J_6(s) &= [s, r_3] \cup [l_2, r_2 - 1] \cup [l_1, n - 1], \\ J_7(s) &= [s, r_3] \cup [l_2, r_2 - 1] \cup [l_1 + 1, n], \\ J_8(s) &= [s, r_3] \cup [l_2 + 1, r_2] \cup [l_1, n - 1], \\ J_9(s) &= [s, r_3] \cup [l_2 + 1, r_2] \cup [l_1 + 1, n], \\ J_{10}(s) &= [s, r_3] \cup [l_2 + 1, r_2 + 1] \cup [l_1 + 2, n], \\ J_{11}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2] \cup [l_1, n - 1], \\ J_{12}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2] \cup [l_1 + 1, n], \\ J_{13}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2 + 1] \cup [l_1 + 2, n]. \end{aligned}$$

Note that, by Theorem 2, for a given  $n \geq 11008$ , every maximal 4-sum-free subset of  $\{1, \dots, n\}$  is one of the sets  $J_x(s)$ , for some  $s \in [S, S + 3] = [s' - 1, s' + 2]$ . By the remarks above, for each  $i \in \{0, \dots, 109\}$ , there are natural 1-1 correspondences between the sets in the families  $\mathcal{F}(4, n)$  for all  $n \equiv i \pmod{110}$ . By slight abuse of notation, we denote any such family simply by  $\mathcal{F}_i$ . Our computer program yielded the following result :

If  $|\mathcal{F}_i| = 1$ , then  $i = 6, 7, 22, 23, 46, 47, 49, 51, 54, 55, 57, 59, 61, 70, 71, 73, 75, 77, 86, 87, 89$

or 91 and

$$\mathcal{F}_i = \{J_9(s')\},$$

or  $i = 36, 37, 100$  or  $101$  and

$$\mathcal{F}_i = \{J_9(s' + 1)\}.$$

If  $|\mathcal{F}_i| = 2$ , then  $\mathcal{F}_i$  is

$$\{J_9(s'), J_9(s' + 1)\} \quad \text{if } i = 93, 103, 105, 107,$$

$$\{J_4, J_9(s')\} \quad \text{if } i = 9, 11, 13, 25, 27,$$

$$\{J_8(s'), J_9(s')\} \quad \text{if } i = 48, 50, 56, 58, 60, 72, 74, 76, 88, 90$$

$$\{J_7(s'), J_9(s')\} \quad \text{if } i = 63, 65, 67, 79, 81.$$

If  $|\mathcal{F}_i| = 3$ :

$$\mathcal{F}_8 = \mathcal{F}_{24} = \{J_4, J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{15} = \{J_4, J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{29} = \{J_4, J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{39} = \{J_9(s'), J_{12}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{62} = \mathcal{F}_{78} = \{J_6(s'), J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{53} = \{J_9(s'), J_{10}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{83} = \{J_7(s'), J_9(s'), J_9(s' + 2)\},$$

$$\mathcal{F}_{92} = \{J_8(s'), J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{95} = \mathcal{F}_{97} = \{J_7(s'), J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{102} = \{J_9(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{109} = \{J_9(s'), J_7(s' + 1), J_9(s' + 1)\}.$$

If  $|\mathcal{F}_i| = 4$ :

$$\mathcal{F}_1 = \mathcal{F}_3 = \mathcal{F}_{17} = \{J_2, J_4, J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{10} = \mathcal{F}_{12} = \mathcal{F}_{26} = \{J_3, J_4, J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{38} = \{J_9(s'), J_{12}(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{41} = \mathcal{F}_{43} = \{J_4, J_9(s'), J_{12}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{52} = \{J_8(s'), J_9(s'), J_{10}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{64} = \mathcal{F}_{66} = \mathcal{F}_{80} = \{J_6(s'), J_7(s'), J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{104} = \mathcal{F}_{106} = \{J_8(s'), J_9(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{69} = \{J_7(s'), J_9(s'), J_{10}(s'), J_9(s' + 1)\}.$$

If  $|\mathcal{F}_i| = 5$  :

$$\begin{aligned}
\mathcal{F}_{14} &= \{J_3, J_4, J_6(s'), J_7(s'), J_9(s')\}, \\
\mathcal{F}_{19} &= \{J_2, J_4, J_7(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{28} &= \{J_3, J_4, J_8(s'), J_9(s'), J_9(s'+1)\}, \\
\mathcal{F}_{31} &= \{J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{82} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{94} &= \{J_6(s'), J_7(s'), J_9(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{99} &= \{J_7(s'), J_9(s'), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{108} &= \{J_8(s'), J_9(s'), J_6(s'+1), J_7(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 6$ :

$$\begin{aligned}
\mathcal{F}_5 &= \{J_2, J_4, J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{33} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{45} &= \{J_4, J_9(s'), J_{12}(s'), J_{13}(s'), J_7(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{68} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{85} &= \{J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{96} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_8(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 7$ :

$$\begin{aligned}
\mathcal{F}_0 = \mathcal{F}_{16} &= \{J_1, J_2, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s')\}, \\
\mathcal{F}_{40} &= \{J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 8$ :

$$\begin{aligned}
\mathcal{F}_2 &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s')\}, \\
\mathcal{F}_{21} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{30} &= \{J_3, J_4, J_6(s'), J_7(s'), J_9(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{35} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{42} &= \{J_3, J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{98} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_8(s'+1), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 9$ :

$$\begin{aligned}
\mathcal{F}_{18} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{84} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_8(s'+2), J_9(s'+2)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 10$ :

$$\begin{aligned}
\mathcal{F}_4 &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{44} &= \{J_3, J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_{13}(s'), J_6(s'+1), J_7(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If  $|\mathcal{F}_i| = 11, 13$  or  $14$ , we get precisely one family for each size:

$$\begin{aligned}\mathcal{F}_{32} &= \{J_1, J_2, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\ \mathcal{F}_{20} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), \\ &\quad J_9(s'+1), J_{12}(s'+1), J_8(s'+2), J_9(s'+2)\}, \\ \mathcal{F}_{34} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), \\ &\quad J_8(s'+1), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}.\end{aligned}$$

Note, in particular, that  $|\mathcal{F}(4, n)| \leq 14$  for all sufficiently large  $n$ . Computer simulations suggest the same may be true for any even  $k$ , with a similar result for odd  $k$ , but we leave the investigation of this possibility to a subsequent paper.

## Appendix

As a prototype for a type of calculation which appears in several places in the paper, we now show, in the notation of Lemma 4, that  $s' = S + 1$  when  $k$  is even.

We must investigate the condition  $l_3(s) < s$ . By definition of  $l_3$  this is just

$$\begin{aligned}\left\lfloor \frac{2r_3}{k} \right\rfloor < s &\Leftrightarrow \frac{2r_3}{k} < s \Leftrightarrow r_3 < \frac{ks}{2} \Leftrightarrow \left\lfloor \frac{l_2 + s}{k} \right\rfloor < \frac{ks}{2} \Leftrightarrow \frac{l_2 + s}{k} < \frac{ks}{2} \\ &\Leftrightarrow l_2 < \left(\frac{k^2}{2} - 1\right)s \Leftrightarrow \frac{2r_2}{k} < \left(\frac{k^2}{2} - 1\right)s \Leftrightarrow r_2 < \left(\frac{k^3}{4} - \frac{k}{2}\right)s \\ &\Leftrightarrow \frac{l_1 + s}{k} < \left(\frac{k^3}{4} - \frac{k}{2}\right)s \Leftrightarrow l_1 < \left(\frac{k^4}{4} - \frac{k^2}{2} - 1\right)s \\ &\Leftrightarrow \frac{2n}{k} < \left(\frac{k^4}{4} - \frac{k^2}{2} - 1\right)s \Leftrightarrow n < \left(\frac{k^5}{8} - \frac{k^3}{4} - \frac{k}{2}\right)s \Leftrightarrow s > \frac{8n}{k^5 - 2k^3 - 4k} \\ &\Leftrightarrow s > S.\end{aligned}$$

Thus  $s' = S + 1$ , as required.

## Acknowledgments

We would like to thank the anonymous referee whose detailed comments greatly helped us improving our paper.

## References

- [1] J. M. Deshoulliers, G. Freiman, V. Sós, and M. Temkin, On the structure of sum-free sets II, *Asterisque* **258** (1999), 149-161.

- [2] F. R. K. Chung and J. L. Goldwasser, Integer sets containing no solutions to  $x+y = 3k$ , in “The Mathematics of Paul Erdős”, R. L. Graham and J. Nešetřil eds., Springer Verlag, Heidelberg, (1996).
- [3] F. R. K. Chung and J. L. Goldwasser, Maximum subsets of  $(0, 1]$  with no solutions to  $x + y = kz$ , *Electron. J. Combin.* **3** (1996), R1.
- [4] K. Roth, On certain sets of integers, *J. London Math. Society* **28** (1953), 104-109.