

Rainbow H -factors

Raphael Yuster

Department of Mathematics
University of Haifa, Haifa 31905, Israel
raphy@research.haifa.ac.il

Submitted: Feb 21, 2005; Accepted: Feb 7, 2006; Published: Feb 15, 2006

Mathematics Subject Classifications: 05C35, 05C70, 05C15

Abstract

An H -factor of a graph G is a spanning subgraph of G whose connected components are isomorphic to H . Given a properly edge-colored graph G , a *rainbow H -subgraph* of G is an H -subgraph of G whose edges have distinct colors. A *rainbow H -factor* is an H -factor whose components are rainbow H -subgraphs. The following result is proved. If H is any fixed graph with h vertices then every properly edge-colored graph with hn vertices and minimum degree $(1 - 1/\chi(H))hn + o(n)$ has a rainbow H -factor.

1 Introduction

All the graphs considered here are finite, undirected and simple. For a graph G we let $v(G)$ and $e(G)$ denote the cardinality of the vertex set and edge set of G , respectively. Given two graphs G and H where $v(H)$ divides $v(G)$, we say that G has an H -factor if G contains $v(G)/v(H)$ vertex-disjoint subgraphs isomorphic to H . Thus, a K_2 -factor is simply a perfect matching. The study of H -factors is a major topic of research in extremal graph theory. A seminal result of Hajnal and Szemerédi [6] gave a sufficient condition for the existence of a K_k -factor. They proved that a graph with nk vertices and minimum degree at least $nk(1 - 1/k)$ has a K_k -factor, and this is best possible. Later, Alon and Yuster proved [3], using the Regularity Lemma [12], a general result guaranteeing the existence of H -factors. They showed that for every fixed graph H with chromatic number $\chi(H)$, any graph with $v(H)n$ vertices and minimum degree at least $v(H)n(1 - 1/\chi(H)) + o(n)$ has an H -factor, and this is asymptotically tight in terms of the chromatic number. Later, it was proved in [10] that the $o(n)$ term can be replaced with a constant $K = K(H)$.

An edge coloring of a graph is called *proper* if two edges sharing an endpoint receive distinct colors. Vizing's theorem asserts that there exists a proper edge coloring of a graph G which uses at most $\Delta(G) + 1$ colors. A *rainbow* subgraph of an edge-colored

graph is a subgraph all of whose edges receive distinct colors. Many graph theoretic parameters have corresponding rainbow variants. Erdős and Rado [5] were among the first to consider problems of this type. Jamison, Jiang and Ling [7], and Chen, Schelp and Wei [4] considered Ramsey type variants where an arbitrary number of colors can be used. Alon et. al. [1] studied the function $f(H)$ which is the minimum integer n such that any proper edge coloring of K_n has a rainbow copy of H . Keevash et. al. [8] considered the rainbow Turán number $ex^*(n, H)$ which is the largest integer m such that there exists a properly edge-colored graph with n vertices and m edges and which has no rainbow copy of H .

A *rainbow H -factor* of a properly edge-colored graph is an H -factor whose elements are rainbow copies of H . Our main result provides sufficient conditions for the existence of a rainbow H -factor. It turns out that the same asymptotic conditions that guarantee an H -factor also guarantee a rainbow H -factor.

Theorem 1.1 *Let H be a graph. There exists $K = K(H)$ such that every proper edge coloring of a graph with n vertices, where $v(H)$ divides n , and with minimum degree at least $(1 - 1/\chi(H))n + K$ has a rainbow H -factor.*

The result might seem a bit surprising as a rainbow version of the theorem of Hajnal and Szemerédi ceases to hold for small values of n . An example is provided in the final section. The proof of Theorem 1.1 is a consequence of a lemma that shows that if H is a complete r -partite graph then any proper edge coloring of some fixed (though much larger) complete r -partite graph has a rainbow H -factor. This lemma and the proof of Theorem 1.1 appear in Section 2. In Section 3 we consider the problem of finding an *almost rainbow H -factor*. Given $\epsilon > 0$, an (ϵ, H) -factor of a graph G is a set of vertex-disjoint copies of H that cover at least $(1 - \epsilon)v(G)$ vertices. Komlós [9] showed that the chromatic number in the main result of [2] can be replaced with another parameter, called the *critical chromatic number* (which, in many cases, is strictly smaller than the chromatic number) if one settles for an (ϵ, H) -factor. We prove a simple rainbow version of a strengthened version of his result due to Shokoufandeh and Zhao [11] where $\epsilon v(G)$ can be replaced by a constant depending only on H . The final section contains some concluding remarks.

2 Rainbow H -factors

Let $T_r(k)$ denote the complete r -partite graph with k vertices in each vertex class. Let H be a fixed graph with $v(H) = h$ and $\chi(H) = r$. Clearly, $T_r(h)$ has an H -factor. As $T_r(h)$ and H have the same chromatic number, this essentially means that it suffices to prove Theorem 1.1 for complete partite graphs. Now, if we can also show that for k sufficiently large, any proper edge coloring of $T_r(k)$ has a rainbow $T_r(h)$ -factor, we can use the results on (usual) H -factors in order to deduce a similar result for the rainbow analogue. We therefore need to prove the following lemma.

Lemma 2.1 *Let h and r be positive integers. There exists $k = k(h, r)$ such that any proper edge coloring of $T_r(k)$ has a rainbow $T_r(h)$ -factor.*

Proof: We shall prove a slightly stronger statement. For $0 \leq p \leq h$, Let $T_r(h, p)$ be the complete r -partite graph with h vertices in each vertex class, except the last vertex class which has only p vertices. Notice that $T_r(h, 0) = T_{r-1}(h, h)$. We prove that there exists $k = k(h, r, p)$ such that any proper edge coloring of $T_r(kh, kp)$ has a rainbow $T_r(h, p)$ -factor.

We fix h , and prove the result by induction on r , and for each r , by induction on $p \geq 1$. The base case $r = 2$ and $p = 1$ is trivial since every star subgraph of a proper edge-colored graph is rainbow. Given $r \geq 2$, assuming the result holds for r and $p - 1$, we prove it for r and p (if $p = 1$ then $p - 1 = 0$ so we use the induction on $T_{r-1}(h, h)$). Let $k = k(h, r, p - 1)$ and let t be sufficiently large (t will be chosen later). Consider a proper edge-coloring of $T = T_r(kth, ktp)$. We let $c(x, y)$ denote the color of the edge (x, y) . Denote the first $r - 1$ vertex classes of T by $V_1, \dots, V_{r-1}$ and denote the last vertex class by U_r . Let $V_r \subset U_r$ be an arbitrary subset of size $k(p - 1)t$ and let $W = U_r \setminus V_r$ be the remaining set with $|W| = kt$. For $i = 1, \dots, r$, we *randomly* partition V_i into t subsets $V_i(1), \dots, V_i(t)$, each of the same size. Each of the r random partitions is performed independently, and each partition is equally likely.

Let $S(j)$ be the subgraph of T induced by $V_1(j) \cup V_2(j) \cup \dots \cup V_r(j)$, for $j = 1, \dots, t$. Notice that $S(j)$ is a properly edge-colored $T_r(kh, k(p - 1))$ and hence, by the induction hypothesis $S(j)$ has a rainbow $T_r(h, p - 1)$ -factor.

Let $B = (X \cup W, F)$ be a bipartite graph where $X = \{S(j) : j = 1, \dots, t\}$ and there exists an edge $(S(j), v) \in F$ if for all $i = 1, \dots, r - 1$ and for all $x \in V_i(j)$, the color $c(x, v)$ does not appear at all in $S(j)$. If we can show that, with positive probability, B has a 1-to- k assignment in which each $S(j) \in X$ is assigned to precisely k elements of W and each $v \in W$ is assigned to a unique $S(j)$ then we can show that T has a rainbow $T_r(h, p)$ -factor. Indeed, consider $S(j)$ and the unique set X_j of k elements of W that are matched to $S(j)$. Since $S(j)$ has a rainbow $T_r(h, p - 1)$ -factor, we can arbitrarily assign a unique element of X_j to each element of this factor and obtain a $T_r(h, p)$ which is also rainbow because all the edges of this $T_r(h, p)$ incident with the assigned vertex from X_j have colors that did not appear at all in other edges of this $T_r(h, p)$.

In order to prove that B has the required 1-to- k assignment we shall use the 1-to- k extension of Hall's Theorem. Namely, we will show that, with positive probability, $|N(Y)| \geq k|Y|$ for each $Y \subset X$. (Hall's Theorem is simply the case $k = 1$. The 1-to- k generalization reduces to the 1-to-1 version by taking k vertex-disjoint copies of X .) To guarantee this condition, it suffices to prove that, with positive probability, each vertex of X has degree greater than $(k - 1/2)t$ in B and each vertex of W has degree greater than $t/2$ in B .

The second part is easy to guarantee, and randomness plays no role. Consider $S(j) \in X$. Let $C(j)$ be the set of all colors appearing in $S(j)$. As $S(j)$ is a $T_r(kh, k(p - 1))$ we have that $|C(j)| < k^2 h^2 \binom{r}{2}$. For each vertex x of $S(j)$, let $W_x \subset W$ be the set of vertices $v \in W$ such that $c(v, x) \in C(j)$. Clearly, $|W_x| < |C(j)|$ since no color appears more than once in edges incident with x . Let $W(j)$ be the union of all W_x taken over all vertices of $S(j)$. Hence, $|W(j)| < (khr)(k^2 h^2 \binom{r}{2})$. Each $v \in W \setminus W(j)$ is a neighbor of $S(j)$ in B . Thus, if we take $t > k^3 h^3 r^3$, we have that each $S(j)$ has more than $(k - 1/2)t$ neighbors

in B .

For the first part, fix some $v \in W$ and let $d_B(v)$ denote the degree of v in B . As $d_B(v)$ is a random variable, and since $|W| = kt$, it suffices to prove that

$$\Pr[d_B(v) \leq t/2] < 1/kt$$

which implies that

$$\Pr[\exists v : d_B(v) \leq t/2] < 1.$$

To simplify notation we let s_i be the size of the i 'th vertex class of each $S(j)$. Thus $s_i = kh$ for $i = 1, \dots, r-1$ and $s_r = k(p-1)$. Recall that the i 'th vertex class of $S(j)$ is formed by taking the j 'th block of a random partition of V_i into t blocks of equal size s_i . Alternatively, one can view the i 'th vertex class of $S(j)$ as the elements $s_i(j-1)+1, \dots, s_i j$ of a random permutation of V_i for $i = 1, \dots, r$. Let, therefore, π_i be a random permutation of V_i . Thus, for $i = 1, \dots, r$, $\pi_i(\ell) \in V_i$ for $\ell = 1, \dots, s_i t$. We define the ℓ 'th vertex of vertex class i of $S(j)$ to be $\pi_i(s_i(j-1) + \ell)$ for $i = 1, \dots, r$ and $\ell = 1, \dots, s_i$.

We define the following events. For three vertex classes $V_\alpha, V_\beta, V_\gamma$ with $1 \leq \alpha < \beta \leq r$, and $1 \leq \gamma \leq r-1$, for a block j where $1 \leq j \leq t$ and for three positive indices $\ell_1 \leq s_\alpha$, $\ell_2 \leq s_\beta$, $\ell_3 \leq s_\gamma$, let x be the ℓ_1 'th vertex of vertex class α in $S(j)$, let y be the ℓ_2 'th vertex of vertex class β in $S(j)$, and let z be the ℓ_3 'th vertex of vertex class γ in $S(j)$. Let $A(\alpha, \beta, \gamma, j, \ell_1, \ell_2, \ell_3)$ be the event that $c(x, y) = c(v, z)$. (Notice that if $\gamma = \alpha$ and $\ell_1 = \ell_3$ or $\gamma = \beta$ and $\ell_2 = \ell_3$ then the corresponding event never holds as our coloring is proper.) We now prove the following claim.

Claim 2.2 *If $d_B(v) \leq t/2$ then there exist $\alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3$ and there exists $J \subset \{1, \dots, t\}$ with $|J| > t/(khr)^3$ such that for each $j \in J$ the event $A(\alpha, \beta, \gamma, j, \ell_1, \ell_2, \ell_3)$ holds.*

Proof: If $d_B(v) \leq t/2$ then there exists $J' \subset \{1, \dots, t\}$ with $|J'| \geq t/2$ such that for each $j \in J'$ some event $A(., ., ., j, ., ., .)$ holds. There are $\binom{r}{2}$ choices for α and β . There are $r-1$ choices for γ . There are at most kh choices for each of ℓ_1, ℓ_2 and ℓ_3 . Hence for some $J \subset J'$ with

$$|J| \geq \frac{|J'|}{k^3 h^3 \binom{r}{2} (r-1)} > \frac{t}{(khr)^3}$$

the 6-tuple $(\alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3)$ is the same for all $j \in J$. ■

For each $\alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3$ where $\ell_1 \leq s_\alpha$, $\ell_2 \leq s_\beta$ and $\ell_3 \leq s_\gamma$ and for each subset $J \subset \{1, \dots, t\}$ of cardinality $|J| = \lceil \frac{t}{(khr)^3} \rceil$, let

$$A(J, \alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3) = \bigcap_{j \in J} A(\alpha, \beta, \gamma, j, \ell_1, \ell_2, \ell_3).$$

Claim 2.3 *If the probability of each of the events $A(J, \alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3)$ is smaller than $k^{-4} h^{-3} r^{-3} t^{-1} 2^{-t}$ then $\Pr[d_B(v) \leq t/2] < 1/kt$.*

Proof: The proof of the claim follows immediately from Claim 2.2 and from the fact that there are less than 2^t possible choices for J and less than $k^3 h^3 r^3$ possible choices for $\alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3$ where $\ell_1 \leq s_\alpha$, $\ell_2 \leq s_\beta$ and $\ell_3 \leq s_\gamma$. ■

By Claim 2.3, in order to complete the proof of Lemma 2.1 it suffices to prove the following claim.

Claim 2.4 *Let $1 \leq \alpha < \beta \leq r$, let $1 \leq \gamma \leq r - 1$, let $\ell_1 \leq s_\alpha$, $\ell_2 \leq s_\beta$, $\ell_3 \leq s_\gamma$ and let $J \subset \{1 \dots, t\}$ with $|J| = \lceil \frac{t}{(khr)^3} \rceil$. Then,*

$$\Pr[A(J, \alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3)] < \frac{1}{k^4 h^3 r^3 t 2^t}.$$

Proof: For convenience, let $A = A(J, \alpha, \beta, \gamma, \ell_1, \ell_2, \ell_3)$ and let $\Delta = \lceil \frac{t}{(khr)^3} \rceil$. We may assume, without loss of generality, that $J = \{1, \dots, \Delta\}$. For $j \in J$, let x_j be the ℓ_1 'th vertex of vertex class α in $S(j)$, let y_j be the ℓ_2 'th vertex of vertex class β in $S(j)$, and let z_j be the ℓ_3 'th vertex of vertex class γ in $S(j)$. Suppose that we are *given* the identity of the $3j - 2$ vertices $x_1, y_1, z_1, \dots, x_{j-1}, y_{j-1}, z_{j-1}$ and z_j (we assume here that all vertices are distinct since if $z_{j'}$ equals either $x_{j'}$ or $y_{j'}$ then $\Pr[A] = 0$ in this case, as our coloring is proper). If we can show that given this information, the probability that $c(x_j, y_j) = c(v, z_j)$ is less than q where q only depends on t, r, h then, by the product formula of conditional probabilities we have $\Pr[A] < q^\Delta$. Thus, assume that we are *given* the identity of the $3j - 2$ vertices $x_1, y_1, z_1, \dots, x_{j-1}, y_{j-1}, z_{j-1}$ and z_j . In particular, we know the color $c = c(v, z_j)$. What is the probability that $c(x_j, y_j) = c$? If $\alpha \neq \gamma$, let $X = V_\alpha \setminus \{x_1, \dots, x_{j-1}\}$ and if $\alpha = \gamma$ let $X = V_\alpha \setminus \{x_1, \dots, x_{j-1}, z_1, \dots, z_j\}$. If $\beta \neq \gamma$, let $Y = V_\beta \setminus \{y_1, \dots, y_{j-1}\}$ and if $\beta = \gamma$ let $Y = V_\beta \setminus \{y_1, \dots, y_{j-1}, z_1, \dots, z_j\}$. Each vertex of X has an equal chance of being x_j and each vertex of Y has an equal chance of being y_j . Thus, each edge of $X \times Y$ has an equal chance of being the edge (x_j, y_j) . Clearly $|X| \geq tkh - 2\Delta$ and $|Y| \geq tk(p - 1) - 2\Delta$ (if $\beta \neq r$ then, in fact, $|Y| \geq tkh - 2\Delta$ and if $p = 1$ then, trivially, $\beta \neq r$). Since our coloring is proper, the color c appears at most tkh times in $V_\alpha \times V_\beta$. Hence,

$$\Pr[c(x_j, y_j) = c] \leq \frac{tkh}{|X||Y|} \leq \frac{tkh}{(tk - 2\Delta)^2} < \frac{tkh}{(tk - tk/2)^2} = \frac{4h}{tk}.$$

It follows that for t sufficiently large as a function of k, h, r we have

$$\Pr[A] < \left(\frac{4h}{tk}\right)^\Delta \leq \left(\frac{4h}{tk}\right)^{t/(khr)^3} < \frac{1}{k^4 h^3 r^3 t 2^t}.$$

■

This completes the induction step and the proof of Lemma 2.1. ■

Proof of Theorem 1.1: Let H be a graph with $\chi(H) = r$ and $v(H) = h$. By Lemma 2.1 there exists $k = k(h, r)$ such that every proper edge coloring of $T_r(k)$ has a rainbow $K(h, r)$ -factor, and hence also a rainbow H -factor. By [10], there exists $K_0 = K_0(k, r)$ such that every graph with n vertices, where kr divides n , and with minimum degree at least $n(1 - 1/r) + K_0$ has a $T_r(k)$ -factor. Let $K = K_0 + kr$ and let G be a properly edge-colored graph with n vertices where h divides n , and with minimum degree at least

$n(1 - 1/r) + K$. Let $n^* \leq n$ be the largest integer which is a multiple of kr . Any graph obtained from G by deleting $n - n^*$ vertices has n^* vertices and minimum degree at least $n(1 - 1/r) + K_0 \geq n^*(1 - 1/r) + K_0$ and hence has a $T_r(k)$ -factor. In particular, we can greedily delete from G a set of $(n - n^*)/h$ vertex-disjoint rainbow copies of H , and the remaining graph has a $T_r(k)$ -factor. As each $T_r(k)$ in this factor is properly colored, each has a rainbow H -factor. Thus, G has a rainbow H -factor. ■

3 Rainbow “almost” H -factors

For an r -chromatic graph H on h vertices, let $u = u(H)$ be the smallest possible color-class size in any r -coloring of H . The *critical chromatic number* of H is $\chi_{cr}(H) = (r - 1)h/(h - u)$. It is easy to see that $\chi(H) - 1 < \chi_{cr}(H) \leq \chi(H)$ and $\chi_{cr}(H) = \chi(H)$ if and only if every r -coloring of H has equal color-class sizes. In [9], Komlós proved the following result.

Theorem 3.1 [Komlós [9]] *Let $\epsilon > 0$ and let H be a graph. There exists $n_0 = n_0(H, \epsilon)$ such that every graph with $n > n_0$ vertices and minimum degree at least $(1 - 1/\chi_{cr}(H))n$ has a set of vertex-disjoint copies of H that cover all but at most ϵn vertices.*

Solving a conjecture of Komlós, Shokoufandeh and Zhao proved the following strengthened version in [11].

Theorem 3.2 [Shokoufandeh and Zhao [11]] *For every graph H there exists $K_0 = K_0(H)$ such that every graph with n vertices and minimum degree at least $(1 - 1/\chi_{cr}(H))n$ has a set of vertex-disjoint copies of H that cover all but at most K_0 vertices.*

Let T be a complete r -partite graph with vertex class sizes $u_1 \leq u_2 \leq \dots \leq u_r$. For a positive integer k , let kT denote the complete r -partite graph with vertex class sizes $ku_1 \leq ku_2 \leq \dots \leq ku_r$. Clearly,

$$\chi_{cr}(kT) = \chi_{cr}(T) = \frac{(r - 1) \sum_{i=1}^r u_i}{\sum_{i=2}^r u_i}.$$

The following is a slight generalization of Lemma 2.1 whose proof is almost identical.

Lemma 3.3 *Let T be a complete r -partite graph with vertex class sizes $u_1 \leq u_2 \leq \dots \leq u_r$. There exists $k = k(T)$ such that any proper edge coloring of kT has a rainbow T -factor.* ■

Let H be a graph, and consider a coloring of H in which the smallest vertex class has size $u(H)$. Adding edges between any two vertices in distinct vertex classes we obtain a complete r -partite graph T with $\chi_{cr}(T) = \chi_{cr}(H)$. Thus, exactly as in the proof of Theorem 1.1 we can use Lemma 3.3 and Theorem 3.2 to obtain the following.

Proposition 3.4 *For every graph H there exists $K = K(H)$ such that every properly edge-colored graph with n vertices and minimum degree at least $(1 - 1/\chi_{cr}(H))n$ has a set of vertex-disjoint rainbow copies of H that cover all but at most K vertices.* ■

4 Concluding remarks

- The proof of Lemma 2.1 yields a *huge* constant $k = k(h, r)$. It is an interesting combinatorial problem to *determine* the minimum integer $k = k(h, r)$ which guarantees that a properly edge-colored $T_r(k)$ has a rainbow $T_r(h)$ -factor. Even for the case $h = 1$ (the case of complete graphs) we do not know the precise answer. Trivially $k(1, 2) = 1$ and $k(1, 3) = 1$. However, $k(1, 4) > 1$ since a proper edge coloring of K_4 need no be rainbow. The following example shows that $k(1, 4) > 2$. Assume the four vertex classes of $T_4(2)$ are $V_i = \{x_i, y_i\}$ for $i = 1, 2, 3, 4$. Color with 1 the edges $x_1x_2, y_1y_2, x_3x_4, y_3y_4$. Color with 2 the edges $x_1y_2, x_2y_1, x_3y_4, x_4y_3$. Color with 3 the edges $x_2x_3, y_2y_3, x_1y_4, y_1x_4$. Color with 4 the edges $x_2y_3, y_2x_3, x_1x_4, y_1y_4$. Color the remaining 8 edges in any way as to obtain a proper edge coloring. It is easily verified that any K_4 of this $T_4(2)$ is not rainbow. In particular, this example shows that the rainbow version of the theorem of Hajnal and Szemerédi ceases to hold for small values of n .
- An edge coloring of a graph is called m -good if each color appears at most m times at each vertex. A slightly more complicated version of Lemma 2.1 also holds in this setting. Namely, Let h, r and m be positive integers. There exists $k = k(h, r, m)$ such that any m -good edge coloring of $T_r(k)$ has a rainbow $T_r(h)$ -factor. We omit the details. Given this extended version of Lemma 2.1 it is straightforward to show that Theorem 1.1 also holds for m -good colored graphs.

References

- [1] N. Alon, T. Jiang, Z. Miller and D. Pritikin, *Properly colored subgraphs and rainbow subgraphs in edge-colorings with local constraints*, Random Struct. Algorithms 23 (2003), No. 4, 409–433.
- [2] N. Alon and R. Yuster, *Almost H -factors in dense graphs*, Graphs Combin. 8 (1992), no. 2, 95–102.
- [3] N. Alon and R. Yuster, *H -factors in dense graphs*, J. Combin. Theory Ser. B 66 (1996), no. 2, 269–282.
- [4] G. Chen, R. Schelp and B. Wei, *Monochromatic - rainbow Ramsey numbers*, 14th Cumberland Conference Abstracts.
Posted at “<http://www.msci.memphis.edu/~balistep/Abstracts.html>”.
- [5] P. Erdős and R. Rado, *A combinatorial theorem*, J. London Math. Soc. 25 (1950), 249–255.
- [6] A. Hajnal and E. Szemerédi, *Proof of a conjecture of Erdős*, in: *Combinatorial Theory and its Applications*, Vol. II (P. Erdős, A. Renyi and V. T. Sós eds.), Colloq. Math. Soc. J. Bolyai 4, North Holland, Amsterdam 1970, 601–623.

- [7] R. Jamison, T. Jiang and A. Ling, *Constrained Ramsey numbers*, J. Graph Theory 42 (2003), No. 1, 1–16.
- [8] P. Keevash, D. Mubayi, B. Sudakov and J. Verstraete, *Rainbow Turán Problems*, Combinatorics, Probability and Computing, to appear.
- [9] J. Komlós, *Tiling Turán Theorems*, Combinatorica 20 (2000), No. 2, 203–218.
- [10] J. Komlós, G. N. Sárközy and E. Szemerédi, *Proof of the Alon-Yuster conjecture*, Discrete Math. 235 (2001), No. 1-3, 255-269.
- [11] A. Shokoufandeh and Y. Zhao, *On a tiling conjecture of Komlós*, Random Struct. Algorithms 23 (2003), No. 2, 180–205.
- [12] E. Szemerédi, *Regular partitions of graphs*, in: *Proc. Colloque Inter. CNRS* (J. -C. Bermond, J. -C. Fournier, M. Las Vergnas and D. Sotteau eds.), 1978, 399–401.