

Adding layers to bumped-body polyforms with minimum perimeter preserves minimum perimeter

Winston C. Yang

winston@cs.wisc.edu

Submitted: Dec 1, 2004; Accepted: Dec 8, 2005; Published: Jan 25, 2006

Mathematics Subject Classifications: 05B50, 05B45

Abstract

In two dimensions, a polyform is a finite set of edge-connected cells on a square, triangular, or hexagonal grid. A layer is the set of grid cells that are vertex-adjacent to the polyform and not part of the polyform. A bumped-body polyform has two parts: a body and a bump. Adding a layer to a bumped-body polyform with minimum perimeter constructs a bumped-body polyform with min perimeter; the triangle case requires additional assumptions. A similar result holds for 3D polyominoes with minimum area.

1 Introduction

A *polyform* is a finite, edge-connected set of cells in a grid of any number of dimensions. In two dimensions, if the cell shape is a square, triangle, or hexagon, the polyform is a *polyomino*, *polyiamond*, or *polyhex*, respectively. In three dimensions, if the cell shape is a cube, the polyform is a *3D polyomino* or a *polycube*. See Figure 1 for examples of 2D polyforms.

In two dimensions, the *area* of a polyform is the number of polyform cells, and the *perimeter* is the number of edges belonging to only one cell. In three dimensions and higher, the *volume* of a polyform is the number of polyform cells, and the *area* is the number of faces belonging to only one cell. In all dimensions, a *layer* is the set of grid cells that are vertex-adjacent to the polyform and not part of the polyform.

Variations on the polyform theme exist; researchers sometimes consider rotations or reflections as distinct polyforms, or do not allow holes, or do not require edge-connectedness, or allow more than one cell shape. A revised classic reference on polyominoes is [Gol94]. Also see [Cla02].

Throughout the paper, “BB” means “bumped-body.” [BCC93] considers polyhexes in the context of hydrocarbons in chemistry, and it proves that if a polyhex can be circumscribed enough times, the resulting polyhex has min perimeter. (Circumscribing

is similar to, but not always equivalent to, adding a layer.) Two results in this paper are somewhat similar to this result. Theorem 23 states that adding a layer to a BB polyhex with min perimeter constructs a BB polyhex with min perimeter. Theorem 17 states that, in various cases, adding a layer to a BB polyiamond with min perimeter constructs a BB polyiamond with min perimeter. In one case, the number of layers must be large enough.

Some theorems are about the BB tuple, perimeter, or area of a polyform after the addition of a number of layers. The proofs depend on figures and induction and leave out the details.

2 Motivation from domain decomposition

In domain decomposition, we want to decompose a *domain* (a set of cells in a square, triangular, or hexagonal grid) into *subdomains* of equal area, as to minimize the total subdomain perimeter. Domain decomposition is a special case of *graph partitioning*: partition the nodes of a graph into sets of equal size as to minimize the *cut-arcs* (arcs between nodes in different sets). In domain decomposition, the nodes represent cells, and the arcs represent edge-connectivity.

Graph partitioning is difficult; it is NP-complete. Researchers have proposed various heuristics to solve it. [Chr96] uses genetic algorithms to try to solve domain decomposition. [KAS⁺02] is the web site of METIS, which gives code for graph partitioning. [GMT95] discusses experimental results with geometric bisection. [KL70] describes the Kernighan-Lin heuristic. [Mar98] uses stripes (rows of subdomains) for domain decomposition with rectangular domains.

Domain decomposition is essentially a tiling problem on a finite set. Finding polyhexes and polyiamonds with min perimeter that tile the plane was the original motivation for considering BB form. [Yan03] proves that all polyominoes constructed by the swirl algorithm in Theorem 3 tile the plane. Also, BB polyforms arise in the swirl algorithm.

3 Global concepts

Let p be a statement. A convenient notation from computer science is $[p]$, which is 1 if p is true, and 0 else. For example, $[i = j]$ is the Kronecker delta function δ_{ij} , and $[a \in A]$ is the characteristic function $\chi_A(a)$ of a set A evaluated at a point a .

Slices are an important concept. See Figure 1. Consider a square, triangular, or hexagonal grid. A *slice* is the set of cells between two parallel grid lines that are one cell apart. For squares, there are two types of slices, rows and columns, whereas for triangles and hexagons, there are three types of slices, rows, antidiagonals, and diagonals. Rotating the slices by $1/6$ turn results in columns, antidiagonals, and diagonals. Use rows instead of columns. Consider a polyform on the grid. A *subslice* is a maximal edge-connected set of cells in a slice. If a slice has one subslice, the term “slice” can also refer to that subslice. [Yan03] proves Theorem 1, which shows how subslices are related to perimeter.

Theorem 1. (*Perimeter slice formulas*) *The perimeter of a polyform with S subslices is: polyomino, $2S$; polyiamond, S ; polyhex, $2S$.*




Figure 1: Slices for polyominoes, polyiamonds, and polyhexes.

The following formulas are from [Yan03], and appeared earlier and in a slightly different form in [HH76]. The polyomino min perimeter formula appeared in [YMC97].

Theorem 2. (*Min perimeter formulas*) *The min perimeter of a polyform with area A is: polyomino, $2\lceil\sqrt{4A}\rceil$; polyiamond, $\lceil\sqrt{6A}\rceil + \lceil\sqrt{6A} \not\equiv A \pmod{2}\rceil$; polyhex, $2\lceil\sqrt{12A-3}\rceil$.*

[HH76], [BCC93], and [Yan03] discuss the swirl algorithm, which constructs polyominoes, polyiamonds, or polyhexes with fixed area and min perimeter. The constructed polyform is not always unique.

Theorem 3. (*Swirl algorithm*) *On a square, triangular, or hexagonal grid, to construct a polyform with area A and min perimeter, start at any cell, and follow a swirl (spiral) for A cells (including the starting cell).*

A *bumped-body (BB)* polyform has two parts: a body and a bump. The body depends on the polyform cell shape: for squares, it is a rectangle; for triangles, it is a hexagon; for hexagons, it is a polyhex whose boundary hexagons lie on a hexagon. The bump is a slice that is possibly empty, shorter than the top side, above the top side, and to the left. Represent a BB polyform by a tuple $(a, b, \dots|x)$, where a, b, \dots are the side lengths of the body, clockwise from the top, and x (for “extra”) is the number of cells in the bump. By rotation or reflection, a BB polyform might have more than one tuple. Use the one with the fewest number of extra cells.

4 Polyominoes

For BB polyominoes $(a, b|x)$, avoid the degenerate cases $a = 0$ or $b = 0$. (However, BB polyiamonds, in the next section, can have tuples in which some values are 0.)

Theorem 4. (*BB polyomino perimeter and area*) *A BB polyomino $(a, b|x)$ has perimeter $2(a + b + \lceil x > 0 \rceil)$ and area $x + ab$.*

Proof. See Figure 2. □




Figure 2: Adding one layer to a BB polyomino $(a, b|x)$.

Lemma 5. (*BB polyomino layers tuple*) Adding $\ell \geq 0$ layers to a BB polyomino $Q_0 = (a, b|x)$ constructs a BB polyomino $Q_\ell = (a + 2\ell, b + 2\ell[x > 0](x + 2\ell))$.

Proof. See Figure 2. Use induction. □

The results for BB polyominoes don't require full layers, just half layers. Consider a BB polyomino as a set of squares in an infinite square grid. A *half layer* of the BB polyomino is the set of grid squares that are edge-adjacent to the left or bottom of the polyomino, or edge-adjacent to at least two such squares.

Lemma 6. (*BB polyomino half layers tuple*) Adding $h \geq 0$ half layers to a BB polyomino $Q_0 = (a, b|x)$ constructs a BB polyomino $Q_h = (a + h, b + h[x > 0](x + h))$.

Proof. See Figure 2. Use induction. □

Theorem 7. (*BB polyomino layers and half layers*) Adding $\ell \geq 0$ layers to a BB polyomino is equivalent to adding 2ℓ half layers.

Proof. Use Lemmas 5 and 6. □

Theorem 8. (*BB polyomino half layers perimeter and area*) Adding $h \geq 0$ half layers to a BB polyomino with perimeter P_0 and area A_0 constructs a polyomino with perimeter $P_h = P_0 + 4h$ and area $A_h = A_0 + hP_0/2 + h^2$.

Proof. Use Theorem 4 and Lemma 6. □

Lemma 9 is arithmetical. It is motivated by BB polyominoes, but it does not require an interpretation in terms of them.

Lemma 9. (*BB polyomino arithmetic*) Let $A_0 \geq 1$, $P_0 = 2\lceil\sqrt{4A_0}\rceil$, $h \geq 0$, $P_h = P_0 + 4h$, and $A_h = A_0 + hP_0/2 + h^2$. Then $P_h = 2\lceil\sqrt{4A_h}\rceil$.

Proof. Note the following equivalences.

$$\begin{aligned} & 2\lceil\sqrt{4A_h}\rceil = P_h \\ \Leftrightarrow & (P_h/2 - 1)^2 < 4A_h \leq (P_h/2)^2 \\ \Leftrightarrow & -P_0 - 4h + 1 < 4A_0 - (P_0/2)^2 \leq 0. \end{aligned}$$

Consider the two inequalities after the last double implication arrow. The right inequality is true. By assumption, the left inequality is true for $h = 0$. For $h \geq 1$, the left side of the left inequality decreases, whereas the right side of the left inequality is constant. \square

Theorem 10. (*BB polyomino preservation of min perimeter through half layers*) Adding half layers to a BB polyomino with min perimeter constructs a BB polyomino with min perimeter.

Proof. Let the BB polyomino have min perimeter P_0 and area A_0 . By Theorem 2, $P_0 = 2\lceil\sqrt{4A_0}\rceil$. By Theorem 8, adding $h \geq 0$ half layers constructs a BB polyomino with perimeter $P_h = P_0 + 4h$ and area $A_h = A_0 + hP_0/2 + h^2$. By Lemma 9, $P_h = 2\lceil\sqrt{4A_h}\rceil$. By Theorem 2, the new polyomino has min perimeter. \square

5 Polyiamonds

Polyominoes use just half layers, but polyiamonds use full layers. The proofs of the polyiamond results are more complicated because the parities of the perimeters and areas are an issue. Some theorems add $\ell \geq 1$ instead of $\ell \geq 0$ layers; this simplifies some of the formulas.

Lemma 11. (*BB polyiamond conservation of slices*) A BB polyiamond $(a, b, c, d, e, f|x)$ satisfies $a + b = d + e$, $a + f = c + d$, $b + c = e + f$.

Proof. A row passes through sides b or c iff it passes through sides e or f . So $b + c = e + f$. The other equations follow from consideration of the antidiagonals and diagonals. \square

Theorem 12. (*BB polyiamond perimeter and area*) A BB polyiamond $(a, b, c, d, e, f|x)$ has perimeter $a + b + c + d + e + f + [x > 0](1 + [x \text{ even}])$. The area is $x + 2a(b + f) + 2bf + a^2 - d^2$.

Proof. See Figure 3. For the area formula, divide the polyiamond into four parts: extra, top, middle, and bottom. \square

Lemma 13. (*BB polyiamond layers tuple*) Adding $\ell \geq 1$ layers to a BB polyiamond $Q_0 = (a, \dots, f|x)$ constructs a BB polyiamond as follows:

- Case $x = 0$: $Q_\ell = (a + \ell, \dots, f + \ell|0)$.
- Case x odd: $Q_\ell = (a + \ell, \dots, f + \ell|x + 2\ell)$.




Figure 3: Calculating the area of a BB polyiamond $(a, b, c, d, e, f|x)$.




Figure 4: Adding one layer to a BB polyiamond $(a, b, c, d, e, f|x)$.

- Case $x > 0$, x even, $x < 2a - 2$: $Q_\ell = (a + \ell, \dots, f + \ell|x + 1 + 2\ell)$.
- Case $x = 2a - 2$:

$$Q_\ell = (a + \ell - 1, b + \ell + 1, c + \ell, d + \ell, e + \ell, f + \ell + 1|0).$$

Proof. See Figure 4. Use induction. □

Theorem 14. (*BB polyiamond layers perimeter and area*) Adding $\ell \geq 1$ layers to a BB polyiamond with x extra triangles, perimeter P_0 , and area A_0 constructs a BB polyiamond with perimeter $P_\ell = P_0 + 6\ell - [x > 0][x \text{ even}]$ and area $A_\ell = A_0 + 2\ell P_0 + 6\ell^2 - [x > 0][x \text{ even}](2\ell - 1)$.

Proof. Use Lemma 11, Theorem 12, and Lemma 13. See Figure 4. □

Lemma 15 is arithmetical. It is motivated by BB polyiamonds, but it does not require an interpretation in terms of them.

Lemma 15. (*BB polyiamond arithmetic*) Let $A_0 \geq 1$, $P_0 = \lceil \sqrt{6A_0} \rceil + \{0, 1\}$, $\ell \geq 0$, $A_\ell = A_0 + 2\ell P_0 + 6\ell^2$, and $P_\ell = P_0 + 6\ell$.

- Case $P_0 = \lceil \sqrt{6A_0} \rceil$: Then $P_\ell = \lceil \sqrt{6A_\ell} \rceil$.
- Case $P_0 = \lceil \sqrt{6A_0} \rceil + 1$: If $\ell > (\lceil \sqrt{6A_0} \rceil^2 - 6A_0)/12$, then $P_\ell = \lceil \sqrt{6A_\ell} \rceil$.

Proof. The proof is similar to the proof of Lemma 9. □

Lemma 16. (*BB polyiamond even criterion*) Consider a BB polyiamond with x extra triangles, $x > 0$, x even, area A , and min perimeter. Then adding one extra triangle constructs a BB polyiamond with area $A + 1$ and min perimeter, $\lceil \sqrt{6A} \rceil \not\equiv A \pmod{2}$, and $\lceil \sqrt{6(A+1)} \rceil \equiv A + 1 \pmod{2}$.

Proof. Let Q be the BB polyiamond, and let P be its perimeter. Adding one extra triangle constructs a BB polyiamond Q' with area $A + 1$ and perimeter $P - 1$.

By Theorem 2, $P = \lceil \sqrt{6A} \rceil + \{0, 1\}$, and the min perimeter of a polyiamond with area $A + 1$ is $\lceil \sqrt{6(A+1)} \rceil + \{0, 1\} \leq P - 1$. Of these four cases (two cases for P , and two cases for $P - 1$), the only consistent case is $P = \lceil \sqrt{6A} \rceil + 1$ and $P - 1 = \lceil \sqrt{6(A+1)} \rceil$. By Theorem 2, Q' has min perimeter, $\lceil \sqrt{6A} \rceil \not\equiv A \pmod{2}$, and $\lceil \sqrt{6(A+1)} \rceil \equiv A + 1 \pmod{2}$. □

Theorem 17. (*BB polyiamond preservation of min perimeter through layers*) Add $\ell \geq 1$ layers to a BB polyiamond Q_0 with x extra triangles, area A_0 , and min perimeter to construct a BB polyiamond Q_ℓ .

- Case $x = 0$ or x is odd:
 - Subcase $\lceil \sqrt{6A_0} \rceil \equiv A_0 \pmod{2}$: Then Q_ℓ has min perimeter.
 - Subcase $\lceil \sqrt{6A_0} \rceil \not\equiv A_0 \pmod{2}$: If $\ell > (\lceil \sqrt{6A_0} \rceil^2 - 6A_0)/12$, then Q_ℓ has min perimeter.
- Case $x > 0$ and x is even. Then Q_ℓ has min perimeter.

Proof. Let P_0 be the perimeter of Q_0 .

Case $x = 0$ or x odd: By Theorem 14, Q_ℓ has perimeter $P_\ell = P_0 + 6\ell$ and area $A_\ell = A_0 + 2\ell P_0 + 6\ell^2$. Subcase $\lceil \sqrt{6A_0} \rceil \equiv A_0 \pmod{2}$: By Theorem 2, $P_0 = \lceil \sqrt{6A_0} \rceil$. By Lemma 15, $P_\ell = \lceil \sqrt{6A_\ell} \rceil$. By Theorem 2, Q_ℓ has min perimeter. Subcase $\lceil \sqrt{6A_0} \rceil \not\equiv A_0 \pmod{2}$: By Theorem 2, $P_0 = \lceil \sqrt{6A_0} \rceil + 1$. By Lemma 15, $P_\ell = \lceil \sqrt{6A_\ell} \rceil$. By Theorem 2, Q_ℓ has min perimeter.

Case $x > 0$ and x even: Add one extra triangle to Q_0 to construct a BB polyiamond Q'_0 with area $A_0 + 1$. By Lemma 16, Q'_0 has min perimeter and $\lceil \sqrt{6(A_0+1)} \rceil \equiv A_0 + 1 \pmod{2}$. By Figure 4, adding ℓ layers to Q'_0 constructs Q_ℓ . Q'_0 has 0 or $x + 1$ (an odd number) of extra triangles. By Case $x = 0$ or x odd, Subcase $\lceil \sqrt{6A_0} \rceil \equiv A_0 \pmod{2}$, Q_ℓ has min perimeter. □

6 Polyhexes

The results for polyhexes are similar to those for polyiamonds, which is not too surprising, because a regular hexagon is made of six equilateral triangles. The proofs are omitted since they are similar to those in the previous paragraphs.

A *side* of a polyhex is a maximal set of boundary hexagons that are edge-connected and whose centers are collinear. The *length* of a side is the number of hexagons in the side. If a polyhex is nonempty, all sides have lengths at least 1.

Lemma 18. (*BB polyhex conservation of slices*) A BB polyhex $(a, b, c, d, e, f|x)$ satisfies $a + b = d + e$, $a + f = c + d$, $b + c = e + f$.

Theorem 19. (*BB polyhex perimeter and area*) A BB polyhex $(a, \dots, f|x)$ has perimeter $2(a+b+c+d+e+f+[x > 0]) - 6$ and area $x + (a+b-1)(a+f-1) - \frac{1}{2}a(a-1) - \frac{1}{2}d(d-1)$.

Lemma 20. (*BB polyhex layers tuple*) Adding $\ell \geq 0$ layers to a BB polyhex $Q_0 = (a, \dots, f|x)$ constructs a BB polyhex $Q_\ell = (a + \ell, \dots, f + \ell|[x > 0](x + \ell))$.

Theorem 21. (*BB polyhex layers perimeter and area*) Adding $\ell \geq 0$ layers to a BB polyhex with perimeter P_0 and area A_0 constructs a BB polyhex with perimeter $P_\ell = P_0 + 12\ell$ and area $A_\ell = A_0 + \ell P_0/2 + 3\ell^2$.

Lemma 22. (*BB polyhex arithmetic*) Let $A_0 \geq 1$, $P_0 = 2\lceil\sqrt{12A_0 - 3}\rceil$, $\ell \geq 0$, $P_\ell = P_0 + 12\ell$, and $A_\ell = A_0 + \ell P_0/2 + 3\ell^2$. Then $P_\ell = 2\lceil\sqrt{12A_\ell - 3}\rceil$.

Theorem 23. (*BB polyhex preservation of min perimeter through layers*) Adding layers to a BB polyhex with min perimeter constructs a BB polyhex with min perimeter.

7 3D polyominoes

This section is somewhat similar to the section on 2D polyominoes. A BB 3D polyomino has two parts: a body and a bump. The body is a box. The bump is a 2D BB polyomino, with its squares replaced by cubes, on the top face of the box, on the bottom left, arranged so that its extra squares (cubes) touch the left edge of the top face and face the back. Represent a BB 3D polyomino by a tuple $(c, d, e|a, b|x)$, where $(a, b|x)$ is the 2D BB polyomino, c is the height of the box, d is the width of the box, and e is the front-to-back distance of the box. See Figure 5.

A *cross-section* of a 3D polyomino is a 2D slice (a plane), parallel to any two of the three coordinate axes.

Theorem 24. (*BB 3D polyomino area, volume, and cross-sections*) A BB 3D polyomino $(c, d, e|a, b|x)$ has area $2(cd + ce + de + a + b + [x > 0])$, volume $x + ab + cde$, and $c + d + e + [ab > 0]$ cross-sections.

Proof. See Figure 5. □

As with 2D polyominoes, 3D polyominoes don't require full layers, just half layers. Consider a BB 3D polyomino in an infinite cubic grid. A *half layer* of the BB 3D polyomino is the set of grid cubes that are face-adjacent to the front, left, or bottom of the BB 3D polyomino, or face-adjacent to at least two such cubes. See Figure 5.




Figure 5: Adding one half layer to a 3D polyomino $(c, d, e|a, b|x)$.

Lemma 25. (*BB 3D polyomino half layers tuple*) Adding $h \geq 0$ half layers to a BB 3D polyomino $Q_0 = (c, d, e|a, b|x)$ constructs a BB 3D polyomino $Q_h = (c+h, d+h, e+h|[ab > 0](a+h), [ab > 0](b+h)|[x > 0](x+h))$.

Proof. See Figure 5. Use induction. □

Theorem 26. (*BB 3D polyomino half layers area, volume, and cross-sections*) Adding $h \geq 0$ half layers to a BB 3D polyomino with area A_0 , volume V_0 , and C_0 cross-sections constructs a BB 3D polyomino with area $A_h = 2(A_0/2 + 2hC_0 + 3h^2)$, volume $V_h = V_0 + hA_0/2 + h^2C_0 + h^3$, and $C_h = C_0 + 3h$ cross-sections.

Proof. See Figure 5. Use Theorem 24 and Lemma 25. □

Theorems 27 and 28 paraphrase two results in [AC96], which is about 3D polyominoes.

Theorem 27. (*Alonso-Cerf integer quasisquare decomposition*) Every positive integer has a unique “quasisquare decomposition:” a sum of a maximal quasicube, a maximal quasisquare, and a bar.

More precisely, if $V \geq 1$, there are unique nonnegative integers (c, d, e, a, b, x) such that $V = cde + ab + x$, $c \leq d \leq e \leq c + 1$, $b \leq a \leq b + 1$, $x < a$, $ab + x < de$.

Theorem 28. (*Alonso-Cerf 3D polyomino min area formula*) Let (c, d, e, a, b, x) be the unique quasisquare decomposition of $V \geq 1$. Then the min area of a 3D polyomino with volume V is $2((cd + ce + de) + (a + b) + [x > 0])$.

Theorem 28 motivates the following definition. A BB 3D polyomino $(c, d, e|a, b|x)$ is *quasisquare* iff (c, d, e, a, b, x) is a quasisquare decomposition of $cde + ab + x$.

Corollary 29. (*Quasisquare BB 3D polyomino min area*) A quasisquare BB 3D polyomino has min area.

Lemma 30. (*Quasisquare BB 3D polyomino half layers*) Adding half layers to a quasisquare BB 3D polyomino constructs a quasisquare BB 3D polyomino.

Proof. See Figure 5. Use Lemma 25. □

A yes answer to Problem 31 would make unnecessary the assumption about cross-sections in Theorem 32.

Problem 31. (*3D Polyomino equality of number of cross sections*) If two 3D polyominos have the same volume and min area, do they have the same number of cross-sections?

Theorem 32. (*BB 3D polyomino preservation of min area through half layers*) Let a BB 3D polyomino with min area have the same number of cross-sections as the quasisquare BB 3D polyomino with the same volume. Then adding half layers to the BB 3D polyomino with min area constructs a BB 3D polyomino with min area.

Proof. Let Q_0 be the BB 3D polyomino. Let Q_0 have area A_0 , volume V_0 , and C_0 cross-sections. By Theorem 26, adding $h \geq 0$ half layers to Q_0 constructs a BB 3D polyomino Q_h with area $A_h = 2(A_0/2 + 2hC_0 + 3h^2)$ and volume $V_h = V_0 + hA_0/2 + h^2C_0 + h^3$.

Let Q'_0 be the quasisquare BB 3D polyomino with volume $V'_0 = V_0$ and $C'_0 = C_0$ cross-sections. By Corollary 29, Q'_0 has min area $A'_0 = A_0$. By Theorem 26, adding h half layers to Q'_0 constructs a BB 3D polyomino Q'_h with area $A'_h = 2(A'_0/2 + 2hC'_0 + 3h^2) = A_h$ and volume $V'_h = V'_0 + hA'_0/2 + h^2C'_0 + h^3 = V_h$.

By Lemma 30, Q'_h is a quasisquare BB 3D polyomino. By Corollary 29, Q'_h has min area. Q'_h and Q_h have the same area and volume, so Q_h has min area. □

Consider earlier polyform area and perimeter formulas as continuous functions. Then $A_u = A_0 + c \int_0^u P_v dv + C$, where c is a constant and C is an extra term for polyiamonds.

8 Acknowledgments

I thank an anonymous referee for comments, for suggesting the assumption $\ell \geq 1$ layers for polyiamonds to simplify the presentation, and for helping to make the paper substantially shorter.

References

- [AC96] L. Alonso and R. Cerf. The three dimensional polyominoes of minimal area. *The Electronic Journal of Combinatorics*, 3, 1996.
- [BCC93] J. Brunvoll, B. N. Cyvin, and S. J. Cyvin. More about extremal animals. *Journal of Mathematical Chemistry*, 12:109–119, 1993.
- [Chr96] I. T. Christou. *Distributed genetic algorithms for partitioning uniform grids*. PhD thesis, University of Wisconsin, Madison, WI, 1996.
- [Cla02] A. L. Clarke. *The poly pages*. web, geocities.com/alclarke0/, 2002.
- [GMT95] J. R. Gilbert, G. L. Miller, and S.-H. Teng. Geometric mesh partitioning: implementation and experiments. *SIAM Journal of Scientific Computing*, 1995.
- [Gol94] S. W. Golomb. *Polyominoes: puzzles, patterns, problems, and packings*, 2nd ed. Princeton University Press, Princeton, NJ, 1994.
- [HH76] F. Harary and H. Harborth. Extremal animals. *Journal of Combinatorics, Information, & System Sciences*, 1:1–8, 1976.
- [KAS⁺02] G. Karypis, R. Aggarwal, K. Schloegel, V. Kumar, and S. Shekhar. Metis (set of programs for partitioning unstructured graphs and hypergraphs and computing fill-reducing orderings of sparse matrices), www-users.cs.umn.edu/~karypis/metis/. web, 2002.
- [KL70] B. W. Kernighan and S. Lin. An effective heuristic procedure for partitioning graphs. *Bell Systems Technical Journal*, pages 291–308, 1970.
- [Mar98] W. Martin. Fast equi-partitioning of rectangular domains using stripe decomposition. *Discrete Applied Mathematics*, 82:193–207, 1998.
- [Yan03] W. C. Yang. *Optimal polyform domain decomposition*. PhD thesis, University of Wisconsin, Madison, WI, 2003.
- [YMC97] J. Yackel, R. R. Meyer, and I. T. Christou. Minimum-perimeter domain assignment. *Mathematical Programming*, 78:283–303, 1997.