

Revisiting two classical results on graph spectra

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis,
Memphis TN 38152, USA
vnikifrv@memphis.edu.edu

Submitted: Sep 4, 2006; Accepted: Dec 18, 2006; Published: Jan 17, 2007

Mathematics Subject Classifications: 05C50

Abstract

Let $\mu(G)$ and $\mu_{\min}(G)$ be the largest and smallest eigenvalues of the adjacency matrix of a graph G . Our main results are:

(i) If H is a proper subgraph of a connected graph G of order n and diameter D , then

$$\mu(G) - \mu(H) > \frac{1}{\mu(G)^{2D} n}.$$

(ii) If G is a connected nonbipartite graph of order n and diameter D , then

$$\mu(G) + \mu_{\min}(G) > \frac{2}{\mu(G)^{2D} n}.$$

For large μ and D these bounds are close to the best possible ones.

Keywords: *smallest eigenvalue, largest eigenvalue, diameter, connected graph, bipartite graph*

1 Introduction

Our notation is standard (e.g., see [2], [3], and [5]). In particular, unless specified otherwise, all graphs are defined on the vertex set $[n] = \{1, \dots, n\}$ and $\mu(G)$ and $\mu_{\min}(G)$ stand for the largest and smallest eigenvalues of the adjacency matrix of a graph G .

The aim of this note is to refine quantitatively two well-known results on graph spectra. The first one, following from Frobenius's theorem on nonnegative matrices, asserts that if H is a proper subgraph of a connected graph G , then $\mu(G) > \mu(H)$. The second one, due to H. Sachs [7], asserts that if G is a connected nonbipartite graph, then $\mu(G) > -\mu_{\min}(G)$.

Our main result is the following theorem.

Theorem 1 *If H is a proper subgraph of a connected graph G of order n and diameter D , then*

$$\mu(G) - \mu(H) > \frac{1}{\mu(G)^{2D} n}. \quad (1)$$

It can be shown that, for large μ and D , the right-hand of (1) gives the correct order of magnitude; examples can be constructed as in the proofs of Theorems 2 and 3.

Theorem 2 *If G is a connected nonbipartite graph of order n and diameter D , then*

$$\mu(G) + \mu_{\min}(G) > \frac{2}{\mu(G)^{2D} n}. \quad (2)$$

Moreover, for all $k \geq 3$, $D \geq 4$, and $n = D + 2k - 1$, there exists a connected nonbipartite graph G of order n and diameter D with $\mu(G) > k$, and

$$\mu(G) + \mu_{\min}(G) < \frac{4}{(k-1)^{2D-4}}.$$

Theorem 2 shows that $\mu(G) + \mu_{\min}(G)$ can be extremely small, although G is nonbipartite and connected. Here is another viewpoint to this fact.

Theorem 3 *Let $0 < \varepsilon < 1/16$. For all sufficiently large n , there exists a connected graph G of order n with $\mu(G) + \mu_{\min}(G) < n^{-\varepsilon}$ such that, to make G bipartite, at least $(1/16 - \varepsilon)n^2$ edges must be removed.*

The picture is completely different for regular graphs. In [4] it is proved that if G is a connected nonregular graph of order n , size m , diameter D , and maximum degree Δ , then

$$\Delta - \mu(G) > \frac{n\Delta - 2m}{n(D(n\Delta - 2m) + 1)}.$$

This result and Theorem 1 imply the following theorems; we omit their straightforward proofs.

Theorem 4 *If H is a proper subgraph of a connected regular graph G of order n and diameter D , then*

$$\mu(G) - \mu(H) > \frac{1}{n(D+1)}.$$

Theorem 5 *If G is a connected regular nonbipartite graph of order n and diameter D , then*

$$\mu(G) + \mu_{\min}(G) > \frac{2}{n(2D+1)}.$$

Theorem 6 *If G is a connected, nonregular, nonbipartite graph of order n , diameter D , and maximum degree Δ , then*

$$\Delta + \mu_{\min}(G) > \frac{1}{n(D+1)} + \frac{1}{\mu(G)^{2D} n}.$$

Note that the last two theorems give some fine tuning of a result of Alon and Sudakov [1].

2 Proofs

Our proof of Theorem 1 stems from a result of Schneider [8] on eigenvectors of irreducible nonnegative matrices; for graphs it reads as: if G is a connected graph of order n and $x_{\min}, x_{\max}$ are minimal and maximal entries of an eigenvector to $\mu(G)$, then

$$\frac{x_{\min}}{x_{\max}} \geq \mu^{-n+1}(G).$$

We reprove this inequality in a more flexible form that sheds some extra light on the original matrix result of Schneider as well. Hereafter we write $\text{dist}(u, v)$ for the length of a shortest path joining the vertices u and v .

Proposition 7 *If G is a connected graph of order n and $(x_1, \dots, x_n)$ is an eigenvector to $\mu(G)$, then*

$$\frac{x_i}{x_j} \geq (\mu(G))^{-\text{dist}(i,j)} \quad (3)$$

for every two vertices $i, j \in V(G)$.

Proof Clearly we can assume that $i \neq j$. For convenience we also assume that $i = 1$ and the vertices $(1, \dots, j)$ form a path joining 1 to j . Then, for all $u = 1, \dots, j-1$, we have

$$\mu x_u = \sum_{uv \in E(G)} x_v \geq x_{u+1};$$

hence, (3) follows by multiplying all these inequalities. $\square$

Proof of Theorem 1 Since $\mu(H) \leq \mu(H')$ whenever $H \subset H'$, we may assume that H is a maximal proper subgraph of G , that is to say, $V(H) = V(G)$ and H differs from G in a single edge uv . Our proof is split into two cases: (a) H connected; (b) H disconnected.

Case (a): H is connected.

In this case we shall prove a stronger result than required, viz.

$$\mu(G) - \mu(H) > \frac{2}{\mu(G)^{2D} n}. \quad (4)$$

Our first goal is to prove that, for every $w \in V(H)$,

$$\text{dist}_H(w, u) + \text{dist}_H(w, v) \leq 2D. \quad (5)$$

Let $w \in V(H)$ and select in H shortest paths $P(u, w)$ and $P(v, w)$ joining u and v to w . Let $Q(u, x)$ and $Q(v, x)$ be the longest subpaths of $P(u, w)$ and $P(v, w)$ having no internal vertices in common. If $s \in Q(u, x)$ or $s \in Q(v, x)$, we obviously have

$$\text{dist}_H(w, s) = \text{dist}_H(w, x) + \text{dist}_H(s, x). \quad (6)$$

The paths $Q(u, x)$, $Q(v, x)$ and the edge uv form a cycle in G ; write k for its length. Assume that $\text{dist}(v, x) \geq \text{dist}(u, x)$ and select $y \in Q(v, x)$ with $\text{dist}_H(x, y) = \lfloor k/2 \rfloor$. Let $R(w, y)$ be a shortest path in G joining w to y ; clearly the length of $R(w, y)$ is at most D . If $R(w, y)$ does not contain the edge uv , it is a path in H and, using (6), we find that

$$\begin{aligned} D &\geq \text{dist}_G(w, y) = \text{dist}_H(w, y) = \text{dist}_H(w, x) + \lfloor k/2 \rfloor \\ &= \text{dist}_H(w, x) + \left\lfloor \frac{\text{dist}_H(x, u) + \text{dist}_H(x, v) + 1}{2} \right\rfloor \\ &\geq \text{dist}_H(w, x) + \frac{\text{dist}_H(x, u) + \text{dist}_H(x, v)}{2} = \frac{\text{dist}_H(w, u) + \text{dist}_H(w, v)}{2}, \end{aligned}$$

implying (5). Let now $R(w, y)$ contain the edge uv . Assume first that v occurs before u when traversing $R(w, y)$ from w to y . Then

$$\begin{aligned} \text{dist}_H(w, u) + \text{dist}_H(w, v) &\leq 2\text{dist}_H(w, x) + \text{dist}_H(x, u) + \text{dist}_H(x, v) \\ &\leq 2(\text{dist}_H(w, x) + \text{dist}_H(x, v)) < \text{dist}_G(w, y) \leq 2D, \end{aligned}$$

implying (5). Finally, if u occurs before v when traversing $R(w, y)$ from w to y , then

$$\begin{aligned} D &\geq \text{dist}_G(w, y) \geq \text{dist}_H(w, u) + 1 + \text{dist}_H(v, y) \\ &= \text{dist}_H(w, x) + \text{dist}_H(x, u) + 1 + \text{dist}_H(v, y) = \text{dist}_H(w, x) + \lceil k/2 \rceil \\ &\geq \text{dist}_H(w, x) + \frac{\text{dist}_H(x, u) + \text{dist}_H(x, v)}{2} = \frac{\text{dist}_H(w, u) + \text{dist}_H(w, v)}{2}, \end{aligned}$$

implying (5). Thus, inequality (5) is proved in full.

Let now $\mathbf{x} = (x_1, \dots, x_n)$ be a unit eigenvector to $\mu(H)$ and let x_w be a maximal entry of $\mathbf{x}$. In view of (3) and (5), we have

$$\frac{x_u x_v}{x_w^2} \geq \frac{1}{\mu^{\text{dist}(u, w) + \text{dist}(v, w)}(H)} \geq \frac{1}{\mu(H)^{2D}}.$$

Hence, in view of $x_w^2 \geq 1/n$, we see that

$$\mu(G) \geq 2 \sum_{ij \in E(G)} x_i x_j = 2x_u x_v + \mu(H) \geq \frac{2x_w^2}{\mu(H)^{2D}} + \mu(H) > \frac{2}{\mu(H)^{2D} n} + \mu(H),$$

completing the proof of (4) and thus of (1).

Case (b): H is disconnected.

Since G is connected, H is union of two connected graphs H_1 and H_2 such that $v \in H_1$, $u \in H_2$. Assume $\mu(H) = \mu(H_1)$, set $|H_1| = k$, $\mu = \mu(H_1)$, and let $\mathbf{x} = (x_1, \dots, x_k)$ be a unit eigenvector to μ . It is immediate to check that the desired inequality holds when $|H_1| = 2, 3$, so we shall assume that $k \geq 4$. Since the path of order 4 has the smallest maximal eigenvalue among all connected graphs of order at least 4, we may assume that $\mu \geq (\sqrt{5} + 1)/2$ and so $\mu^2 \geq \mu + 1$.

Since $\text{dist}(u, w) \leq \text{diam}G \leq D$ for every $w \in V(H_1)$, we see that $\text{dist}(v, w) \leq D - 1$ for every $w \in V(H_1)$. On the other hand, each maximal entry of $\mathbf{x}$ is at least $k^{-1/2}$; hence, Proposition 7 implies that $x_v \geq \mu^{-D+1}k^{-1/2}$. Setting

$$\mathbf{y} = (y_1, \dots, y_k, y_u) = \left(x_1, \dots, x_k, \frac{x_v}{\mu} \right),$$

we see that $\|\mathbf{y}\|^2 = 1 + (x_v/\mu)^2$; thus, letting B be the adjacency matrix of the graph $H_1 + u$, we have

$$\begin{aligned} \mu(G) &\geq \mu(H_1 + u) \geq \frac{\langle B\mathbf{y}, \mathbf{y} \rangle}{\|\mathbf{y}\|^2} \geq \frac{1}{1 + (x_v/\mu)^2} \left(2y_u y_v + 2 \sum_{ij \in E(H_1)} y_i y_j \right) \\ &= \frac{\mu^2}{\mu^2 + x_v^2} \left(\frac{2x_v^2}{\mu} + \mu \right) = \mu \frac{\mu^2 + 2x_v^2}{\mu^2 + x_v^2} > \mu + \mu \frac{x_v^2}{\mu^2 + \mu} = \mu + \frac{x_v^2}{\mu + 1}. \end{aligned}$$

To complete the proof of the theorem, observe that

$$\frac{x_v^2}{\mu + 1} \geq \frac{x_v^2}{\mu^2} = \frac{1}{k\mu^{2D}} > \frac{1}{n\mu^{2D}}.$$

□

Proof of Theorem 2 Let $\mathbf{x} = (x_1, \dots, x_n)$ be an eigenvector to $\mu_{\min}(G)$ and let $V_1 = \{u : x_u < 0\}$. Let H be the maximal bipartite subgraph of G , containing all edges with exactly one vertex in V_1 . It is not hard to see that H is connected proper subgraph of G , $V(H) = V(G)$, and $\mu_{\min}(H) < \mu_{\min}(G)$. Finally, let H' be a maximal proper subgraph of G containing H . We have

$$\mu(G) + \mu_{\min}(G) \geq \mu(G) + \mu_{\min}(H) = \mu(G) - \mu(H) \geq \mu(G) - \mu(H').$$

and (2) follows from case (a) of the proof of Theorem 1.

To construct the required example, set $G_1 = K_3$, $G_2 = K_{k,k}$, join G_1 to G_2 by a path P of length $n - 2k - 2$, and write G for the resulting graph; obviously G is of order n and diameter $n - 2k + 1$. Set $\mu = \mu(G)$ and note that $\mu(G) > k$. Let $V(G_1) = \{u_1, u_2, v_1\}$ and $P = (v_1, \dots, v_{n-2k-1})$, where $v_{n-2k-1} \in V(G_2)$. Let $\mathbf{x}$ be a unit eigenvector to $\mu(G)$ and assume that the entries $x_1, x_2, x_3, \dots, x_{n-2k+1}$ correspond to $u_1, u_2, v_1, \dots, v_{n-2k-1}$. Clearly $x_1 = x_2$, and so, from $\mu x_2 = x_2 + x_3$, we find that $x_1 = x_2 = x_3/(\mu - 1)$. Furthermore,

$$\mu x_3 = 2x_2 + x_4 = \frac{2x_3}{\mu - 1} + x_4 < x_3 + x_4,$$

and by induction we obtain $x_i < (\mu - 1)x_{i+1}$ for all $3 \leq i \leq n - 2k$. Therefore,

$$x_1 = x_2 \leq (\mu - 1)^{-n+2k+1} x_{n-2k+1} < (k - 1)^{-D+2},$$

and by Rayleigh's principle we deduce that

$$\mu(G) + \mu_{\min}(G) \leq 4x_1x_2 < \frac{4}{(k-1)^{2D-4}},$$

completing the proof. $\square$

Proof of Theorem 3 Set $r = \lceil n/4 \rceil + 1$, $s = \lceil (1/2 - \varepsilon)n \rceil$, select $G_1 = K_{r,r}$, $G_2 = K_s$, join G_1 to G_2 by a path P of length $n - 2r - s + 1$ and write G for the resulting graph. Note first that, to make G bipartite, we must remove at least

$$\binom{s}{2} - \left\lfloor \frac{s^2}{4} \right\rfloor \geq \frac{s^2}{4} - \frac{s}{2} > \frac{(1/2 - \varepsilon)^2 n^2}{4} - \frac{s}{2} \geq \left(\frac{1}{16} - \varepsilon \right) n^2$$

edges, for n large enough. Note also that

$$n - 2 \left\lceil \frac{n}{4} \right\rceil - 2 - \left\lceil \left(\frac{1}{2} - \varepsilon \right) n \right\rceil + 1 > n - \frac{n}{2} - \left(\frac{1}{2} - \varepsilon \right) n - 4 = \varepsilon n - 4.$$

so the length of P is greater than $\varepsilon n - 4$.

Let $\mathbf{x}$ be a unit eigenvector to $\mu(G)$. Clearly the entries of $\mathbf{x}$ corresponding to vertices from $V(G_1) \setminus V(P)$ have the same value α . Like in the proof of Theorem 2, we see that $\alpha < (n/4)^{-\varepsilon n + 5}$. Hence, by Rayleigh's principle, for n large enough, we deduce that

$$\mu(G) + \mu_{\min}(G) \leq 4\alpha^2 \binom{s}{2} < (n/4)^{-2\varepsilon n + 10} \frac{n^2}{2} < (n/4)^{-2\varepsilon n + 12} < n^{-\varepsilon n},$$

completing the proof. $\square$

Acknowledgment: The author is indebted to Béla Bollobás for his kind support and to Sebi Cioabă for interesting discussions. Finally, the referee suggested a number of improvements, in particular, a simplification of the proof of Theorem 1.

References

- [1] N. Alon, B. Sudakov, Bipartite subgraphs and the smallest eigenvalue, *Combin. Probab. Comput.* 9 (2000) 1-12.
- [2] B. Bollobás, *Modern Graph Theory*, Graduate Texts in Mathematics, 184, Springer-Verlag, New York (1998), xiv+394 pp.
- [3] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1980, 368 pp.
- [4] S. Cioabă, D. Gregory, V. Nikiforov, Extreme eigenvalues of nonregular graphs, to appear in *J. Combin. Theory Ser B*.

- [5] R. Horn, C. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985, xiii+561 pp.
- [6] V. Nikiforov, Walks and the spectral radius of graphs, *Linear Algebra Appl.* 418 (2006), 257-268
- [7] H. Sachs, Beziehungen zwischen den in einem Graphen enthaltenen Kreisen und seinem charakteristischen Polynom, *Publ. Math. Debrecen* 11 (1964) 119–134.
- [8] H. Schneider, Note on the fundamental theorem on irreducible non-negative matrices, *Proc. Edinburgh Math. Soc.* 11 (1958/1959) 127–130.