

Restricting supercharacters of the finite group of unipotent uppertriangular matrices

Nathaniel Thiem*

Department of Mathematics
University of Colorado at Boulder
thiemn@colorado.edu

Vidya Venkateswaran†

Department of Mathematics
California Institute of Technology
vidyav@caltech.edu

Submitted: Aug 22, 2008; Accepted: Feb 9, 2009; Published: Feb 20, 2009

Mathematics Subject Classification: 05E99, 20C33

Abstract

It is well-known that understanding the representation theory of the finite group of unipotent upper-triangular matrices U_n over a finite field is a wild problem. By instead considering approximately irreducible representations (supercharacters), one obtains a rich combinatorial theory analogous to that of the symmetric group, where we replace partition combinatorics with set-partitions. This paper studies the supercharacter theory of a family of subgroups that interpolate between U_{n-1} and U_n . We supply several combinatorial indexing sets for the supercharacters, supercharacter formulas for these indexing sets, and a combinatorial rule for restricting supercharacters from one group to another. A consequence of this analysis is a Pieri-like restriction rule from U_n to U_{n-1} that can be described on set-partitions (analogous to the corresponding symmetric group rule on partitions).

1 Introduction

The representation theory of the finite group of upper-triangular matrices U_n is a well-known wild problem. Therefore, it came as somewhat of a surprise when C. André was able to show that by merely “clumping” together some of the conjugacy classes and some of the irreducible representations one attains a workable approximation to the representation theory of U_n [1, 2, 3, 4]. In his Ph.D. thesis [14], N. Yan showed how the algebraic geometry of the original construction could be replaced by more elementary constructions. E. Arias-Castro, P. Diaconis, and R. Stanley [8] were then able to demonstrate that this theory can in fact be used to study random walks on U_n using techniques that traditionally required

*The authors would like to thank Diaconis and Marberg for many enlightening discussions regarding this work, and anonymous referees for their comments.

†Part of this work is Venkateswaran’s honors thesis at Stanford University.

knowledge of the full character theory [11]. Thus, the approximation is fine enough to be useful, but coarse enough to be computable.

André's approximate theory also has a remarkable combinatorial structure that recalls the classical connection between the representation theory of the symmetric group and partition combinatorics. In this case, we replace partition with set-partitions, so that

$$\left\{ \begin{array}{l} \text{Almost irreducible} \\ \text{representations of } U_n \end{array} \right\} \xleftrightarrow{1-1} \left\{ \begin{array}{l} \text{Set partitions} \\ \text{of } \{1, 2, \dots, n\} \end{array} \right\}.$$

In particular, the number of almost irreducible representations is a Bell number (or more generally a q -analogue of a Bell number). One of the main results of this paper is to extend the analogy with the symmetric group by giving a combinatorial Pieri-like formula for set-partitions that corresponds to restriction in U_n .

Our strategy is to study a family of groups – called pattern groups – that interpolate between U_n and U_{n-1} . A pattern group is a unipotent matrix group associated to a poset $\mathcal{P}$ of $\{1, 2, \dots, n\}$ subject to the condition that the (i, j) th can be nonzero only if $i \preceq j$ in $\mathcal{P}$ (a group version of the incidence algebra of $\mathcal{P}$). For example, U_n is the pattern group associated to the poset $1 \prec 2 \prec \dots \prec n$, and our interpolating pattern groups are associated to the posets $2 \prec 3 \prec \dots \prec n$ and $1 \prec m$ for some $1 < m \leq n$.

In [10], P. Diaconis and M. Isaacs generalized André's theory to the notion of a *supercharacter theory* for arbitrary finite groups, where irreducible characters are replaced by supercharacters and conjugacy classes are replaced by superclasses. In particular, their paper generalized André's original construction by giving a supercharacter theory for pattern groups (and even more generally algebra groups). The combinatorics of these supercharacter theories for general pattern groups is not yet understood: there seems to be a constant tension between the set partition combinatorics of U_n and the underlying poset $\mathcal{P}$ (see, for example, [12]). In particular, lengthy anti-chains seem to imply more complicated combinatorics. Another main result of this paper is to work out the combinatorics for the set of interpolating subgroups, demonstrating that while for these posets the combinatorics becomes more technical, it remains computable.

In [10], Diaconis and Isaacs also showed that the restriction of a supercharacter between pattern groups is a $\mathbb{Z}_{\geq 0}$ -linear combination of supercharacters in the subgroup. However, even for $U_m \subseteq U_n$, these coefficients are not well understood (and also depend on the particular embedding of U_m in U_n). This paper offers a first step in understanding this problem giving an algorithm for computing coefficients. In general, these will be polynomials in the size q of the underlying finite field, but it is unknown what these coefficients might count.

Section 2 reviews the basics of supercharacter theory and pattern groups. Section 3 defines the interpolating subgroups $U_{(m)}$, and finds two different sets of natural superclass and supercharacter representatives, which we call comb representatives and path representatives. Section 4 uses a general character formula from [12] to determine character formulas for both comb and path representatives. The character formula for comb representatives – Theorem 4.1 – is easier to compute directly, but the path representative character formula – Theorem 4.3 – has a more pleasing combinatorial structure. Section

5 uses the character formulas to derive a restriction rule for the interpolating subgroups given in Theorem 5.1. Corollary 5.1 iterates these restrictions to deduce a recursive decomposition formula for the restriction from U_n to U_{n-1} .

This paper is the companion paper to [13], which studies the superinduction of supercharacters. Other work related to supercharacter theory of unipotent groups, include C. André and A. Neto's exploration of supercharacter theories for unipotent groups of Lie types B , C , and D [5], C. André and A. Nicolás' analysis of supertheories over other rings [6], and an intriguing possible connection between supercharacter theories and Borovichenko and Drinfeld's work on L -packets [9].

2 Preliminaries

This section reviews several topics fundamental to our main results: Supercharacter theories, pattern groups, and a character formula for pattern groups.

2.1 Supertheories

Let G be a group. As defined in [10], a *supercharacter theory* for G is a partition $\mathcal{S}^\vee$ of the elements of G and a set of characters $\mathcal{S}$, such that

- (a) $|\mathcal{S}| = |\mathcal{S}^\vee|$,
- (b) Each $S \in \mathcal{S}^\vee$ is a union of conjugacy classes,
- (c) For each irreducible character γ of G , there exists a unique $\chi \in \mathcal{S}$ such that

$$\langle \gamma, \chi \rangle > 0,$$

where $\langle, \rangle$ is the usual innerproduct on class functions,

- (d) Every $\chi \in \mathcal{S}$ is constant on the elements of $\mathcal{S}^\vee$.

We call $\mathcal{S}^\vee$ the set of *superclasses* and $\mathcal{S}$ the set of *supercharacters*. Note that every group has two trivial supercharacter theories – the usual character theory and the supercharacter theory with $\mathcal{S}^\vee = \{\{1\}, G \setminus \{1\}\}$ and $\mathcal{S} = \{\mathbb{1}, \gamma_G - \mathbb{1}\}$, where $\mathbb{1}$ is the trivial character of G and γ_G is the regular character.

There are many ways to construct supercharacter theories, but this paper will study a particular version developed in [10] to generalize André's original construction to a larger family of groups called algebra groups.

2.2 Pattern groups

While many results can be stated in the generality of algebra groups, frequently statements become simpler if we restrict our attention to a subfamily called pattern groups. We follow the construction of [10] for the superclasses and supercharacters of pattern groups.

Let U_n denote the set of $n \times n$ unipotent upper-triangular matrices with entries in the finite field $\mathbb{F}_q$ of q elements. For any poset $\mathcal{P}$ on the set $\{1, 2, \dots, n\}$, the pattern group $U_{\mathcal{P}}$ is given by

$$U_{\mathcal{P}} = \{u \in U_n \mid u_{ij} \neq 0 \text{ implies } i \leq j \text{ in } \mathcal{P}\}.$$

This family of groups includes unipotent radicals of rational parabolic subgroups of the finite general linear groups $\mathrm{GL}_n(\mathbb{F}_q)$; the group U_n is the pattern group corresponding to the total order $1 < 2 < 3 < \dots < n$.

The group $U_{\mathcal{P}}$ acts on the $\mathbb{F}_q$ -algebra

$$\mathfrak{n}_{\mathcal{P}} = \{u - 1 \mid u \in U_{\mathcal{P}}\}$$

by left and right multiplication. Two elements $u, v \in U_{\mathcal{P}}$ are in the same *superclass* if $u - 1$ and $v - 1$ are in the same two-sided orbit of $\mathfrak{n}_{\mathcal{P}}$. Note that since every element of $U_{\mathcal{P}}$ can be decomposed as a product of elementary matrices, every element in the orbit containing $v - 1 \in \mathfrak{n}_{\mathcal{P}}$ can be obtained by applying a sequence of the following row and column operations.

- (a) A scalar multiple of row j may be added to row i if $j > i$ in $\mathcal{P}$,
- (b) A scalar multiple of column k may be added to column l if $k < l$ in $\mathcal{P}$.

There are also left and right actions of $U_{\mathcal{P}}$ on the dual space $\mathfrak{n}_{\mathcal{P}}^* = \mathrm{Hom}_{\mathbb{F}_q}(\mathfrak{n}_{\mathcal{P}}, \mathbb{F}_q)$ given by

$$(u\lambda v)(x - 1) = \lambda(u^{-1}(x - 1)v^{-1}), \quad \text{where } \lambda \in \mathfrak{n}_{\mathcal{P}}^*, u, v, x \in U_{\mathcal{P}}.$$

Fix a nontrivial group homomorphism $\theta : \mathbb{F}_q^+ \rightarrow \mathbb{C}^\times$. The *supercharacter* χ^λ with representative $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ is

$$\chi^\lambda = \frac{|U_{\mathcal{P}}\lambda|}{|U_{\mathcal{P}}\lambda U_{\mathcal{P}}|} \sum_{\mu \in U_{\mathcal{P}}\lambda U_{\mathcal{P}}} \theta \circ (-\mu).$$

We identify the functions $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ with matrices by the vector space isomorphism,

$$\begin{aligned} [\cdot] : \mathfrak{n}_{\mathcal{P}}^* &\longrightarrow M_n(\mathbb{F}_q)/\mathfrak{n}_{\mathcal{P}}^\perp \\ \lambda &\mapsto [\lambda] = \sum_{i < j \in \mathcal{P}} \lambda_{ij}(e_{ij} + \mathfrak{n}_{\mathcal{P}}^\perp), \end{aligned} \tag{1}$$

where $e_{ij} \in \mathfrak{n}_{\mathcal{P}}$ has (i, j) entry 1 and zeroes elsewhere, $\lambda_{ij} = \lambda(e_{ij})$, and

$$\begin{aligned} M_n(\mathbb{F}_q) &= \{n \times n \text{ matrices with entries in } \mathbb{F}_q\}, \\ \mathfrak{n}_{\mathcal{P}}^\perp &= \{y \in M_n(\mathbb{F}_q) \mid y_{ij} = 0 \text{ for all } i < j \text{ in } \mathcal{P}\}. \end{aligned}$$

We will typically choose the quotient representative to be in $\mathfrak{n}_{\mathcal{P}}$. Then, as with superclasses, every element in the orbit containing $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ can be obtained by applying a sequence of the following row and column operations to $[\lambda]$.

- (a) A scalar multiple of row i may be added to row j if $i < j$ in $\mathcal{P}$,
- (b) A scalar multiple of column l may be added to column k if $l > k$ in $\mathcal{P}$.

Note that since we are in the quotient space $M_n(\mathbb{F}_q)/\mathfrak{n}_{\mathcal{P}}^\perp$, we quotient by all nonzero entries that might occur through these operations that are not in allowable in $\mathfrak{n}_{\mathcal{P}}$.

Example. For U_n we have

$$\left\{ \begin{array}{c} \text{Superclasses} \\ \text{of } U_n \end{array} \right\} \longleftrightarrow \left\{ u \in U_n \mid \begin{array}{c} u - 1 \text{ has at most one nonzero} \\ \text{entry in every row and column} \end{array} \right\} \quad (2)$$

If $q = 2$, then

$$\left\{ u \in U_n \mid \begin{array}{c} u - 1 \text{ has at most one nonzero} \\ \text{entry in every row and column} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{Set partitions} \\ \text{of } \{1, 2, \dots, n\} \end{array} \right\}.$$

Similarly, if

$$\mathfrak{n}_n = U_n - 1,$$

then

$$\left\{ \begin{array}{c} \text{Supercharacters} \\ \text{of } U_n \end{array} \right\} \longleftrightarrow \left\{ \lambda \in \mathfrak{n}_n^* \mid \begin{array}{c} \text{The matrix } [\lambda] \text{ has at most one non-} \\ \text{zero entry in every row and column} \end{array} \right\}. \quad (3)$$

Let

$$\mathcal{S}_n(q) = \{ \lambda \in \mathfrak{n}_n^* \mid [\lambda] \text{ has at most one nonzero entry in every row and column} \}. \quad (4)$$

2.3 A supercharacter formula for pattern groups

Let $U_{\mathcal{P}}$ be a pattern group with corresponding nilpotent algebra $\mathfrak{n}_{\mathcal{P}}$. Let

$$J = \{(i, j) \mid i < j \text{ in } \mathcal{P}\}.$$

Given $u \in U_{\mathcal{P}}$ and $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$, define $a, b \in \mathbb{F}_q^{|J|}$ by

$$\begin{aligned} a_{ij} &= \sum_{j < k \text{ in } \mathcal{P}} u_{jk} \lambda_{ik}, & \text{for } (i, j) \in J, \\ b_{jk} &= \sum_{i < j \text{ in } \mathcal{P}} u_{ij} \lambda_{ik}, & \text{for } (j, k) \in J. \end{aligned}$$

Let M be the $|J| \times |J|$ matrix given by

$$M_{ij,kl} = \begin{cases} u_{jk} \lambda_{il}, & \text{if } i < j < k < l \text{ in } \mathcal{P}, \\ 0, & \text{otherwise.} \end{cases}, \quad \text{for } (i, j), (k, l) \in J.$$

Informally, if one superimposes the matrices u and $[\lambda]$, then

a	tracks occurrences of	$\begin{array}{ c } \hline \lambda_{jk} \\ \hline u_{ik} \\ \hline \end{array}$
b	tracks occurrences of	$\begin{array}{ c c } \hline u_{ij} & \lambda_{ik} \\ \hline \end{array}$
M	tracks occurrences of	$\begin{array}{ c c } \hline & \lambda_{il} \\ u_{jk} & \\ \hline \end{array}$

Remark. Each of a , b , and M depend on u , λ , and $\mathcal{P}$. However, to make the notation less heavy-handed, we leave this dependence out of the notation.

Let $\text{Null}(M)$ denote the nullspace of M and let $\cdot : \mathbb{F}_q^{|J|} \times \mathbb{F}_q^{|J|} \rightarrow \mathbb{F}_q$ be the usual inner product (dot product) on $\mathbb{F}_q^{|J|}$. The following theorem gives a general supercharacter formula for pattern groups. However, typical applications of the theorem make a particular choice of superclass and supercharacter representatives.

Theorem 2.1 ([12]). *Let $u \in U_{\mathcal{P}}$ and $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$. Then*

(a) *The character*

$$\chi^{\lambda}(u) = 0$$

unless there exists $x \in \mathbb{F}_q^{|J|}$ such that $Mx = -a$ and $b \cdot \text{Null}(M) = 0$,

(b) *If $\chi^{\lambda}(u)$ is not zero, then*

$$\chi^{\lambda}(u) = \frac{q^{|U_{\mathcal{P}}\lambda|}}{q^{\text{rank}(M)}} \theta(x \cdot b) \theta \circ \lambda(u - 1),$$

where $x \in \mathbb{F}_q^{|J|}$ is such that $Mx = -a$.

Remark. There are two natural choices for χ^{λ} , one of which is the conjugate of the other. Theorem 2.1 uses the convention of [10] rather than [12].

C. André proved the U_n -version of this supercharacter formula for large characteristic [3], and [8] extended it to all finite fields. Note that the following theorem follows from Theorem 2.1 by choosing appropriate representatives for the superclasses and supercharacters.

Theorem 2.2. *Let $\lambda \in \mathcal{S}_n(q)$, and let $u \in U_n$ be a superclass representative as in (2). Then*

(a) *The character degree*

$$\chi^{\lambda}(1) = \prod_{i < j, \lambda_{ij} \neq 0} q^{j-i-1}.$$

(b) The character

$$\chi^\lambda(u) = 0$$

unless whenever $u_{jk} \neq 0$ with $j < k$, we have $\lambda_{ik} = 0$ for all $i < j$ and $\lambda_{jl} = 0$ for all $l > k$.

(c) If $\chi^\lambda(u) \neq 0$, then

$$\chi^\lambda(u) = \frac{\chi^\lambda(1)\theta \circ \lambda(u-1)}{q^{|\{i < j < k < l \mid u_{jk}, \lambda_{il} \in \mathbb{F}_q^\times\}|}}.$$

3 Interpolating between U_{n-1} and U_n

Fix $n \geq 1$. For $2 \leq m \leq n$, let

$$U_{(m)} = \{u \in U_n \mid u_{1j} = 0, \text{ for } 1 < j \leq m\} = U_{\mathcal{P}_{(m)}},$$

$$\mathfrak{n}_{(m)} = \{u - 1 \mid u \in U_{(m)}\} = \mathfrak{n}_{\mathcal{P}_{(m)}},$$

where $\mathcal{P}_{(m)} = 1 \begin{array}{c} \nearrow m+1 \\ | m \\ \searrow m-1 \\ | \\ \vdots \\ | \\ 2 \end{array},$

and by convention, let $U_{(1)} = U_n$. Note that

$$U_{n-1} \cong U_{(n)} \triangleleft U_{(n-1)} \triangleleft \cdots \triangleleft U_{(1)} = U_n.$$

The goal of this section is to identify suitable orbit representatives for representatives for $U_{(m)} \backslash \mathfrak{n}_{(m)} / U_{(m)}$ and $U_{(m)} \backslash \mathfrak{n}_{(m)}^* / U_{(m)}$.

A matrix $A \in M_n(\mathbb{F}_q)$ has an underlying vertex-labeled graph structure G_A given by vertices

$$V_A = \{A_{ij} \mid 1 \leq i, j \leq n, A_{ij} \neq 0\}$$

and an edge from A_{ij} to A_{kl} if $i = k$ or $j = l$. We label each vertex by its location in the matrix, so A_{ij} has label (i, j) . For example, for $a, b, c, d, e, f, g, h \in \mathbb{F}_q^\times$,

$$A = \begin{pmatrix} 0 & 0 & a & 0 & b \\ c & 0 & 0 & 0 & d \\ 0 & e & 0 & f & 0 \\ 0 & 0 & 0 & 0 & g \\ 0 & 0 & 0 & h & 0 \end{pmatrix} \quad \text{implies} \quad G_A = \left(\begin{array}{ccc} & a & b \\ c & & d \\ & e & f \\ & & g \\ & & h \end{array} \right).$$

[illegible]
$$\text{bag}_{ij}(u) = \begin{cases} u_{ij}x_k(-x_{k-1})^{-1}x_{k-2}(-x_{k-3})^{-1}\dots((-1)^{k+1}x_1)^{(-1)^{k+1}}, & \text{if } S_u = (x_1, \dots, x_l), \\ 1, & u_{ij} = x_{k+1}, k < l, \\ & \text{otherwise.} \end{cases}$$
$$\begin{pmatrix} u_{1j_1} & & & & & \\ & & & & u_{i_{k+1}j_k} & \\ & & & u_{i_kj_{k-1}} & u_{i_kj_k} & \\ & & \ddots & u_{i_{k-1}j_{k-1}} & & \\ & u_{i_3j_2} & & & & \\ u_{i_2j_1} & u_{i_2j_2} & & & & \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} u_{1j_1} \text{bag}_{i_2j_2}(u) \cdots \text{bag}_{i_{k-1}j_{k-1}}(u) \text{bag}_{i_kj_k}(u) & & & & & \\ & & & & u_{i_{k+1}j_k} & \\ & & & & & u_{i_kj_{k-1}} \\ & & \ddots & & & \\ & u_{i_3j_2} & & & & \\ u_{i_2j_1} & u_{i_2j_2} & & & & \end{pmatrix} \quad (5)$$

Proposition 3.1. *Let $0 < m < n$. Then*

- Proof.* (a) Let $u \in \mathcal{T}_{(m)}^\vee$. Then $U_{(m)}(u-1)U_{(m)} \subseteq U_n(u-1)U_n$. In fact, if $v \in \mathcal{T}_{(m)}^\vee$, but $(v-1) \notin U_{(m)}(u-1)U_{(m)}$, then $(v-1) \notin U_n(u-1)U_n$. Thus, distinct elements of $\mathcal{T}_{(m)}^\vee$ correspond to distinct superclasses of $U_{(m)}$.

Let $u \in U_{(m)}$ and let $U_{n-1} \subseteq U_{(m)}$ be the subgroup of U_n obtained by taking the last $n - 1$ rows and columns. Then $U_{n-1}(u - 1)U_{n-1} \subseteq U_{(m)}(u - 1)U_{(m)}$. We may choose $(v - 1) \in U_{n-1}(u - 1)U_{n-1}$ such that

- (a) every row of $(v - 1)$ except row 1 has at most one nonzero entry,
- (b) every column of $(v - 1)$ has at most two nonzero entries,
- (c) if a column has two nonzero entries, then one of the entries must be in the first row.

We may now apply additional row operations allowable by $\mathcal{P}_{(m)}$ to obtain $(v' - 1) \in U_{(m)}(u - 1)U_{(m)}$, to replace (c) by

- (c') if a column has two nonzero entries, then one entry must be in the first row and the second in a row $\leq m$.

Therefore it suffices to show that if the rows of the second nonzero entries do not decrease as we move from left to right, we can convert them into an appropriate form. The following sequence of row and column operations effects such an adjustment.

$$\begin{aligned} \begin{pmatrix} u_{1k} & u_{1l} \\ u_{ik} & 0 \\ 0 & u_{jl} \end{pmatrix} &\xrightarrow{-u_{1k}^{-1}u_{1l}\text{Col}(k)+\text{Col}(l)} \begin{pmatrix} u_{1k} & 0 \\ u_{ik} & -u_{ik}u_{1k}^{-1}u_{1l} \\ 0 & u_{jl} \end{pmatrix} \\ &\xrightarrow{u_{jl}^{-1}u_{ik}u_{1k}^{-1}u_{1l}\text{Row}(j)+\text{Row}(i)} \begin{pmatrix} u_{1k} & 0 \\ u_{ik} & 0 \\ 0 & u_{jl} \end{pmatrix}. \end{aligned}$$

(b) follows from (a) and (5). □

3.2 Supercharacter representatives

Recall that we identify $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ with matrices $[\lambda] \in M_n(\mathbb{F}_q)/\mathfrak{n}_{\mathcal{P}}^\perp$ via the map (1). A function $\lambda \in \mathfrak{n}_{(m)}^*$ is a *comb representative* if

- (a) At most one connected component of $G_{[\lambda]}$ has more than one element,
- (b) If $G_{[\lambda]}$ has a connected component S with more than one element, then there exist $k_1 > k_2 > \cdots > k_r > m \geq i_{r'} > i_{r'-1} > \cdots > i_1 > 1$ with $r' \in \{r, r - 1\}$ such that

$$\left\{ \begin{pmatrix} & & & \lambda_{1k_1} \\ & & \lambda_{i_1k_2} & \lambda_{i_1k_1} \\ & \lambda_{i_2k_3} & & \lambda_{i_2k_1} \\ & \ddots & & \vdots \\ \lambda_{i_{r-1}k_r} & & & \lambda_{i_{r-1}k_1} \\ & & & \lambda_{i_rk_1} \end{pmatrix} \right\} \text{ or } \left\{ \begin{pmatrix} & & & \lambda_{1k_1} \\ & & \lambda_{i_1k_2} & \lambda_{i_1k_1} \\ & & \lambda_{i_2k_3} & \lambda_{i_2k_1} \\ & \ddots & & \vdots \\ \lambda_{i_{r-1}k_r} & & & \lambda_{i_{r-1}k_1} \end{pmatrix} \right\}$$

are the vertices of S .

A function $\lambda \in \mathfrak{n}_{(m)}^*$ is a *path representative* if

- (a) At most one connected component of $G_{[\lambda]}$ has more than one element,
- (b) If $G_{[\lambda]}$ contains a connected component S with more than one element, then there exist $k_1 > k_2 > \cdots > k_r > m \geq i_{r'} > i_{r'-1} > \cdots > i_1 > 1$ with $r' \in \{r, r-1\}$ such that

$$\left\{ \begin{array}{ccccccc} & & & & \lambda_{1k_1} & & \\ & & & & \lambda_{i_1k_2} & \lambda_{i_1k_1} & \\ & & & \lambda_{i_2k_3} & \lambda_{i_2k_2} & & \\ & & \ddots & & & & \\ \lambda_{i_{r-1}k_r} & \lambda_{i_{r-1}k_{r-1}} & & & & & \\ \lambda_{i_rk_r} & & & & & & \end{array} \right\} \text{ or } \left\{ \begin{array}{ccccccc} & & & & \lambda_{1k_1} & & \\ & & & & \lambda_{i_1k_2} & \lambda_{i_1k_1} & \\ & & & \lambda_{i_2k_3} & \lambda_{i_2k_2} & & \\ & & \ddots & & & & \\ \lambda_{i_{r-1}k_r} & \lambda_{i_{r-1}k_{r-1}} & & & & & \end{array} \right\}$$

are the vertices of S .

Let

$$\begin{aligned} \mathcal{T}_{(m)} &= \{\lambda \in \mathfrak{n}_{(m)}^* \mid \lambda \text{ a comb representative}\} \\ \mathcal{Z}_{(m)} &= \{\lambda \in \mathfrak{n}_{(m)}^* \mid \lambda \text{ a path representative}\}. \end{aligned}$$

Let $\lambda \in \mathcal{Z}_{(m)}$. If $G_{[\lambda]}$ has a connected component S_λ with a vertex in the first row, then we can order the vertices of S_λ by starting with the vertex in the first row and then numbering in order along the path. For example, if

$$S_\lambda = \left(\begin{array}{ccccccc} & & & & \lambda_{1j_1} & & \\ & & & & \lambda_{i_2j_2} - \lambda_{i_2j_1} & & \\ & & & & \lambda_{i_3j_2} & & \\ & & \ddots & & & & \\ & \lambda_{i_{k-1}j_{k-1}} & & & & & \\ \lambda_{i_kj_k} - \lambda_{i_kj_{k-1}} & & & & & & \end{array} \right)$$

then the order of the vertices is $S_\lambda = (\lambda_{1j_1}, \lambda_{i_2j_1}, \lambda_{i_2j_2}, \dots, \lambda_{i_kj_k})$. For $i < j$ in $\mathcal{P}$, define the baggage $\text{bag}_{ij} : \mathcal{Z}_{(m)} \rightarrow \mathbb{F}_q$ by the rule,

$$\text{bag}_{ij}(\lambda) = \begin{cases} \lambda_{ij} y_k (-y_{k-1})^{-1} y_{k-2} (-y_{k-3})^{-1} \cdots ((-1)^{k+1} y_1)^{(-1)^{k+1}}, & \text{if } S_\lambda = (y_1, \dots, y_l), \\ & \lambda_{ij} = y_{k+1}, k < l, \\ 1, & \text{otherwise.} \end{cases}$$

Note that the pairs

$$\begin{pmatrix} & & & \lambda_{1j_1} \\ & & \lambda_{i_2j_2}\lambda_{i_2j_1} & \\ & \ddots & \lambda_{i_3j_2} & \\ & & \lambda_{i_{k-1}j_{k-1}} & \\ \lambda_{i_kj_k} & \lambda_{i_kj_{k-1}} & & \\ \lambda_{i_{k+1}j_k} & & & \end{pmatrix} \text{ and } \begin{pmatrix} & & & \lambda_{1j_1} \\ & & \lambda_{i_2j_2} & -\lambda_{1j_1} \text{bag}_{i_2j_1}(\lambda) \\ & \ddots & & \vdots \\ & & \lambda_{i_{k-1}j_{k-1}} & -\lambda_{1j_1} \text{bag}_{i_{k-1}j_{k-2}}(\lambda) \\ \lambda_{i_kj_k} & & & -\lambda_{1j_1} \text{bag}_{i_kj_{k-1}}(\lambda) \\ & & & -\lambda_{1j_1} \text{bag}_{i_{k+1}j_k}(\lambda) \end{pmatrix} \quad (6)$$

$$\begin{pmatrix} & & & \lambda_{1j_1} \\ & & \lambda_{i_2j_2}\lambda_{i_2j_1} & \\ & \ddots & \lambda_{i_3j_2} & \\ & & \lambda_{i_{k-1}j_{k-1}} & \\ \lambda_{i_kj_k} & \lambda_{i_kj_{k-1}} & & \\ \lambda_{i_{k+1}j_k} & & & \end{pmatrix} \text{ and } \begin{pmatrix} & & & \lambda_{1j_1} \\ & & \lambda_{i_2j_2} & -\lambda_{1j_1} \text{bag}_{i_2j_1}(\lambda) \\ & \ddots & & \vdots \\ & & \lambda_{i_{k-1}j_{k-1}} & -\lambda_{1j_1} \text{bag}_{i_{k-1}j_{k-2}}(\lambda) \\ \lambda_{i_kj_k} & & & -\lambda_{1j_1} \text{bag}_{i_kj_{k-1}}(\lambda) \\ & & & \end{pmatrix}$$

are in the same two sided orbit in $\mathfrak{n}_{\mathcal{P}}^*$ according to the row and column operations given in Section 2.2.

Proposition 3.2. *Let $0 < m < n$. Then*

- (a) $\mathcal{T}_{(m)}$ is a set of supercharacter representatives,
- (b) $\mathcal{Z}_{(m)}$ is a set of supercharacter representatives.

Proof. (a) Let $\lambda \in \mathcal{T}_{(m)}$. Then $U_{(m)}\lambda U_{(m)} \subseteq U_n\lambda U_n$. In fact, if $\gamma \in \mathcal{T}_{(m)}$, but $\gamma \notin U_{(m)}\lambda U_{(m)}$, then $\gamma \notin U_n\lambda U_n$. Thus, distinct elements of $\mathcal{T}_{(m)}$ correspond to distinct two-sided orbits in $\mathfrak{n}_{\mathcal{P}}^*$.

Since 1 is incomparable to $j \in \{2, 3, \dots, m\}$ in $\mathcal{P}_{(m)}$, we may not add row 1 to row j if $j \leq m$ when computing two-sided orbits. Let $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ and let $U_{n-1} \subseteq U_{(m)}$ be the subgroup of U_n obtained by taking the last $n-1$ rows and columns. Then $U_{n-1}\lambda U_{n-1} \subseteq U_{(m)}\lambda U_{(m)}$. We may choose $\gamma \in U_{n-1}\lambda U_{n-1}$ such that

- (a) every row of γ except row 1 has at most one nonzero entry,
- (b) every column of γ has at most two nonzero entries,
- (c) if a column has two nonzero entries, then one of the entries must be in the first row.

We may now apply additional row operations allowable by $\mathcal{P}_{(m)}$ to obtain $\gamma' \in U_{(m)}\lambda U_{(m)}$, to replace (a),(b),(c) by

- (a') every column of γ except some column k has at most 1 nonzero entry,
- (b') every row of γ has at most two nonzero entries, and row 1 has at most 1,
- (c') if a row has two nonzero entries, then one entry must be in column k and the second in a column j such that $m < j < k$.

We can now readjust the nonzero entries to be in an appropriate arrangement as in the proof of Proposition 3.1.

(b) follows from (a) and (6). □

4 Supercharacter formulas for $U_{(m)}$

This section develops supercharacter formulas for both comb and path representatives. After developing tools that allow us to decompose characters as products of simpler characters, we prove a character formula for comb characters. We then use the translation between comb and path representatives of (5) and (6) to get a more combinatorial character formula for path representatives.

4.1 Multiplicativity of supercharacter formulas

In this section we begin with the general pattern group setting, so let $\mathcal{P}$ be a poset.

Let $u \in U_{\mathcal{P}}$. For a connected component S of G_{u-1} , let $[S] \in \mathfrak{n}_{\mathcal{P}}$ be given by

$$[S]_{jk} = \begin{cases} u_{jk}, & \text{if } u_{jk} \in V_S, \\ 0, & \text{otherwise.} \end{cases}$$

Similarly, let $\lambda \in \mathfrak{n}_{(m)}^*$. For a connected component T of $G_{[\lambda]}$, let $[T] \in M_n(\mathbb{F}_q)/\mathfrak{n}_{\mathcal{P}}^\perp$ be given by

$$[T]_{jk} = \begin{cases} \lambda_{jk}, & \text{if } \lambda_{jk} \in V_T, \\ 0, & \text{otherwise.} \end{cases}$$

The following lemma allows us to decompose the supercharacter formula of a pattern group $U_{\mathcal{P}}$ by connected components.

Lemma 4.1. *Let $u \in U_{\mathcal{P}}$ and $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$. Let $S_1, S_2, \dots, S_k$ be the connected components of G_{u-1} and $T_1, T_2, \dots, T_l$ be the connected components of $G_{[\lambda]}$. Then*

$$\chi^\lambda(u) = \prod_{j=1}^l \chi^{[T_j]}(1) \prod_{i=1}^k \frac{\chi^{[T_j]}([S_i] + 1)}{\chi^{[T_j]}(1)}.$$

Proof. Let $U = U_{\mathcal{P}}$. The proof follows from the following two claims:

- (1) If λ has two components T and T' , then

$$\chi^\lambda(u) = \chi^{[T]}(u) \chi^{[T']}(u).$$

- (2) If u has two components S and S' , then

$$\chi^\lambda(u) = \chi^\lambda(1) \frac{\chi^\lambda([S])}{\chi^\lambda(1)} \frac{\chi^\lambda([S'])}{\chi^\lambda(1)}.$$

- (1) Note that since T and T' involve distinct rows and columns, the left orbits of $[T]$ and $[T']$ are independent and involve distinct rows. Thus,

$$|U\lambda| = |U[T]| |U[T']|.$$

In fact, for $\lambda' \in U\lambda U$,

$$|\{(\gamma, \mu) \in (U[T]U) \times (U[T']U) \mid \lambda' = \gamma + \mu\}| = \frac{|U[T]U||U[T']U|}{|U\lambda U|}.$$

Thus, by definition

$$\begin{aligned} \chi^\lambda(u) &= \frac{|U\lambda|}{|U\lambda U|} \sum_{\lambda' \in U\lambda U} \theta(-\lambda'(u-1)) \\ &= \frac{|U\lambda|}{|U[T]U||U[T']U|} \sum_{\substack{\gamma \in U[T]U \\ \mu \in U[T']U}} \theta(-\gamma(u-1) - \mu(u-1)) \\ &= \frac{|U[T]||U[T']|}{|U[T]U||U[T']U|} \sum_{\substack{\gamma \in U[T]U \\ \mu \in U[T']U}} \theta(-\gamma(u-1))\theta(-\mu(u-1)) \\ &= \frac{|U[T]|}{|U[T]U|} \sum_{\gamma \in U[T]U} \theta(-\gamma(u-1)) \frac{|U[T']|}{|U[T']U|} \sum_{\mu \in U[T']U} \theta(-\mu(u-1)) \\ &= \chi^{[T]}(u) \chi^{[T']}(u). \end{aligned}$$

(2) For any $u' - 1 \in U(u-1)U$,

$$|\{(v-1, w-1) \in (U[S]U) \times (U[S']U) \mid u' - 1 = v - 1 + w - 1\}| = \frac{|U[S]U||U[S']U|}{|U(u-1)U|}.$$

We have that

$$\begin{aligned} \chi^\lambda(u) &= \frac{\chi^\lambda(1)}{|U(u-1)U|} \sum_{v-1 \in U(u-1)U} \theta(-\lambda(v-1)) \\ &= \frac{\chi^\lambda(1)}{|U[S]U||U[S']U|} \sum_{\substack{v-1 \in U[S]U \\ w-1 \in U[S']U}} \theta(-\lambda(v-1 + w-1)) \\ &= \frac{\chi^\lambda(1)}{\chi^\lambda(1)\chi^\lambda(1)} \frac{\chi^\lambda(1)}{|U[S]U|} \sum_{v-1 \in U[S]U} \theta(-\lambda(v-1)) \frac{\chi^\lambda(1)}{|U[S']U|} \sum_{w-1 \in U[S']U} \theta(-\lambda(w-1)) \\ &= \chi^\lambda(1) \frac{\chi^\lambda([S])}{\chi^\lambda(1)} \frac{\chi^\lambda([S'])}{\chi^\lambda(1)}, \end{aligned}$$

as desired. □

Corollary 4.1. *Let $u \in U_{\mathcal{P}}$ and $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$ with connected components $T_1, \dots, T_l$. Then*

$$\chi^\lambda(u) = \prod_{i=1}^l \chi^{[T_i]}(u).$$

To obtain character formulas for $U_{(m)}$ we will require a slightly more refined multiplicativity result that depends on the poset structure $\mathcal{P}_{(m)}$ and a choice of comb representatives.

For $u \in U_{(m)}$ and $1 \leq k \leq n$, let $u[k] \in U_{(m)}$ be given by

$$u[k]_{ij} = \begin{cases} u_{ij}, & \text{if } j \in \{i, k\}, \\ 0, & \text{otherwise.} \end{cases}$$

That is, $u[k]$ is the same as u on the diagonal and in the k th column, but has zeroes elsewhere. For $\lambda \in \mathfrak{n}_{\mathcal{P}}^*$, let $\lambda[u, k] \in \mathfrak{n}_{\mathcal{P}}^*$ be given by

$$[\lambda[u, k]]_{il} = \begin{cases} \lambda_{il}, & \text{if } k \leq l \text{ and } u_{jk} \neq 0 \text{ for some } i \leq j < k, \\ 0, & \text{otherwise.} \end{cases}$$

That is, $[\lambda[u, k]]$ is the same as $[\lambda]$ weakly NorthEast of the nonzero entries of u in the k th column, but has zeroes elsewhere.

The following lemma states that we can compute supercharacter formulas for $U_{(m)}$ column by column on the superclasses.

Lemma 4.2. *Let $u \in U_{(m)}$ with $u \in \mathcal{T}_{(m)}^\vee$ and let $\lambda \in \mathcal{T}_{(m)}$. Then*

- (a) *The character $\chi^\lambda(u) \neq 0$ if and only if $\chi^{\lambda[u, k]}(u[k]) \neq 0$ for all $2 \leq k \leq n$.*
- (b) *The character value*

$$\chi^\lambda(u) = \chi^\lambda(1) \prod_{k=2}^n \frac{\chi^{\lambda[u, k]}(u[k])}{\chi^{\lambda[u, k]}(1)}.$$

Proof. (a) Let M correspond to (λ, u) as in Theorem 2.1. Note that $M_{(i, j), (k, l)}, M_{(i, j), (k', l')} \in \mathbb{F}_q^\times$ implies $\lambda_{il}, u_{jk}, \lambda_{i'l'}, u_{j'k'} \in \mathbb{F}_q^\times$, so

$$u = \begin{matrix} & & k & k' \\ & & & \\ j & \begin{pmatrix} & & & \\ & u_{jk} & u_{jk'} & \end{pmatrix} \end{matrix} \quad \text{and} \quad [\lambda] = \begin{matrix} & & l & l' \\ i & \begin{pmatrix} & & & \\ & \lambda_{il} & \lambda_{i'l'} & \end{pmatrix} \end{matrix}.$$

However, since $u \in \mathcal{T}_{(m)}^\vee$, the only row of u which can have more than one nonzero entry is row 1. Since $i < j$, we have $k = k'$ and the nonzero entries of u contribute to distinct rows of M . Similiarly, if $M_{(i, j), (k, l)}, M_{(i', j'), (k, l)} \in \mathbb{F}_q^\times$ implies $\lambda_{il}, u_{jk}, \lambda_{i'l}, u_{j'k} \in \mathbb{F}_q^\times$, so

$$u = \begin{matrix} & & k \\ & & \\ j & \begin{pmatrix} & & \\ & u_{jk} & \\ j' & u_{j'k} \end{pmatrix} \end{matrix} \quad \text{and} \quad [\lambda] = \begin{matrix} & & l \\ i & \begin{pmatrix} & & \\ & \lambda_{il} & \\ i' & \lambda_{i'l} \end{pmatrix} \end{matrix}.$$

Thus, distinct columns of u contribute to distinct columns of M . For $1 \leq k \leq n$,

$$\begin{aligned} R_k &= \text{rows of } M \text{ that have nonzero entries corresponding} \\ &\quad \text{to the nonzero entries of } u \text{ in column } k \\ C_k &= \text{columns of } M \text{ that have nonzero entries corresponding} \\ &\quad \text{to the nonzero entries of } u \text{ in column } k. \end{aligned} \tag{7}$$

By choosing an appropriate order on the rows and columns of M ,

$$M = M_{R_1, C_1} \oplus M_{R_2, C_2} \oplus \cdots \oplus M_{R_n, C_n}, \quad (8)$$

where M_{R_k, C_k} is the submatrix of M using rows R_k and columns C_k .

Using (8), there exists a solution to $Mx = -a$ if and only if for each $1 \leq k \leq n$, there exist $x_k \in \mathbb{F}_q^{|C_k|}$ such that $M_{R_k, C_k} x_k = -a_{R_k}$.

If $a_{ij} \neq 0$, then there exist $\lambda_{ik}, u_{jk} \in \mathbb{F}_q^\times$ for some k . Since row j in u has at most one nonzero entry, $a_{ij} = u_{jk} \lambda_{ik}$. Thus, a_{R_k} only depends on the pair $(\lambda[u, k], u[k])$.

By (8), we have

$$\text{Null}(M) = \text{Null}(M_{R_1, C_1}) \oplus \text{Null}(M_{R_2, C_2}) \oplus \cdots \oplus \text{Null}(M_{R_n, C_n}),$$

so b is perpendicular to $\text{Null}(M)$ if and only if b_{C_k} is perpendicular to M_{R_k, C_k} for all k . The condition $(k, l) \in C_k$ implies $u_{jk} \neq 0$ for some j , so $b_{kl} \in \mathbb{F}_q^\times$ implies $b_{kl} = u_{1k} \lambda_{1l} + u_{jk} \lambda_{jl}$. Thus, b_{C_k} only depends on the pair $(\lambda[u, k], u[k])$, and (a) follows.

(b) Since $C_1 = R_1 = \emptyset$, it follows from (8) that

$$\text{rank}(M) = \sum_{k=1}^n \text{rank}(M_{R_k, C_k}) = \sum_{k=2}^n \text{rank}(M_{R_k, C_k}).$$

By (a),

$$\theta(x \cdot b) = \prod_{k=2}^n \theta(x_{C_k} \cdot b_{C_k}),$$

and by inspection

$$\theta \circ \lambda(u - 1) = \prod_{(j, k)} \theta(u_{jk} \lambda_{jk}) = \prod_{k=1}^n \prod_{j < k} \theta(u_{jk} \lambda_{jk}) = \prod_{k=1}^n \theta(\lambda[u, k](u[k] - 1)).$$

Now (b) follows from (a). □

Remark. This lemma depends on the choice of representatives. In particular, it is not true for path representatives.

4.2 A character formula for comb representatives

It follows from Lemmas 4.1 and 4.2 that to give the character value $\chi^\lambda(u)$, we may assume $u - 1$ has nonzero entries in one column and $G_{[\lambda]}$ has one connected component S .

Theorem 4.1. *Let $u \in U_{(m)}$ such that $u \in \mathcal{T}_{(m)}^\vee$ and $u - 1$ has support $\text{supp}(u - 1) \subseteq \{(1, k), (j, k)\}$. Let $\lambda \in \mathcal{T}_{(m)}$ be such that λ has one connected component S with $\text{Cols}(S) = \{l_1 < l_2 < \cdots < l_s\}$. Then*

(a) Let $i_1 > i_2 > \dots > i_{s-1}$ be such that $\lambda_{i_d l_d} \neq 0$ for $1 \leq d \leq s-1$. The character degree

$$\chi^\lambda(1) = \begin{cases} q^{l_s-m-2} \prod_{d=1}^{s-1} q^{l_d-i_d-1}, & \text{if } \lambda_{i l_s} = 0 \text{ for all } i > i_1, \\ q^{l_s-i-1} \prod_{d=1}^{s-1} q^{l_d-i_d-1}, & \text{if } \lambda_{i l_s} \neq 0 \text{ for some } i > i_1. \end{cases}$$

(b) The character

$$\chi^\lambda(u) = 0$$

unless at least one of the following occurs

- (1) $u_{jk}\lambda_{ik} \neq 0$ implies $i = 1$ with $j \leq m$ or $i > j$; and $u_{1k}\lambda_{1l} + u_{jk}\lambda_{jl} = 0$ for all $j < k < l$,
- (2) $u_{jk}\lambda_{ik} \neq 0$ for some $1 < i < j \leq m$, but $|R_k| = |C_k| > 0$ (R_k and C_k are as in (7)),
- (3) $u_{1k}\lambda_{1l_s} + u_{jk}\lambda_{jl} \neq 0$ for some $m < k < l$, but $\lambda_{ik} = 0$ for all i and $|R_k| \geq |C_k| > 0$,
- (4) $u_{jk}, \lambda_{j l_s}, \lambda_{i l_s} \in \mathbb{F}_q^\times$ with $i < j < k < l_s$ with $\lambda_{j k'} = 0$ for all $k < k' < l_s$.

(c) The character values are

$$\chi^\lambda(u) = \begin{cases} \frac{\chi^\lambda(1)}{q^{|C_k|-\delta_{RC}}} \theta(u_{jk}\lambda_{jk}), & \text{if (1)} \\ \frac{\chi^\lambda(1)}{q^{|C_k|}}, & \text{if (2) or (3) or (4)} \\ \frac{\chi^\lambda(1)}{q^{|C_k|}} \theta(-\lambda_{i l_s}^{-1} \lambda_{ik} (u_{1k}\lambda_{1l_s} + u_{jk}\lambda_{j l_s})), & \text{if (2) and (4),} \end{cases}$$

where $\delta_{RC} = 1$ if $|C_k| > |R_k|$ and $\delta_{RC} = 0$ if $|C_k| \leq |R_k|$.

Proof. (a) This is just a statement of the fact that

$$\chi^\lambda(1) = |U_{(m)}\lambda|,$$

combined with the structure of S .

(b) and (c). Note that by Lemma 4.2,

$$\chi^\lambda(u) = \frac{\chi^\lambda(1)}{\chi^{\lambda[u,k]}(1)} \chi^{\lambda[u,k]}(u[k]),$$

so we may assume $M = M_{R_k, C_k}$ (see (8)). Let $\text{Rows}(S) = \{i_1 > \dots > i_s\}$ or $\text{Rows}(S) = \{i_0 > i_1 > \dots > i_s\}$ be the rows with nonzero entries in S such that $\lambda_{i_d l_d}, \lambda_{i_d l_s} \in \mathbb{F}_q^\times$, and,

if $i_0 \in \text{Rows}(S)$, then $\lambda_{i_0 l_s} \neq 0$ (see the definition of comb representatives in Section 3.2). Let r and r' be minimal such that $l_r > k$ and $i_{r'} < j$. Then

$$M = \begin{pmatrix} & & & \delta' u_{jk} \lambda_{1l_s} \\ & & u_{jk} \lambda_{i_{s-1} l_{s-1}} & u_{jk} \lambda_{i_{s-1} l_s} \\ & \ddots & & \vdots \\ u_{jk} \lambda_{i_r l_r} & & & u_{jk} \lambda_{i_r l_s} \\ & & & u_{jk} \lambda_{i_{r-1} l_s} \\ & & & \vdots \\ & & & u_{jk} \lambda_{i_{r'} l_s} \end{pmatrix}, \text{ where } \delta' = \begin{cases} 1, & \text{if } j > m, \\ 0, & \text{if } j \leq m. \end{cases} \quad (9)$$

Thus, the rank of M is $q^{|C_k| - \delta_{RC}}$.

Furthermore, $a \in \mathbb{F}_q^{|R_k|}$ and $b \in \mathbb{F}_q^{|C_k|}$ are given by

$$\begin{aligned} a_{ij} &= u_{jk} \lambda_{ik}, & \text{for } (i, j) \in R_k, \\ b_{kl} &= \begin{cases} u_{1k} \lambda_{1l} + u_{jk} \lambda_{jl}, & \text{if } l \in \text{Cols}(S), \\ 0 & \text{otherwise.} \end{cases} & \text{for } (k, l) \in C_k. \end{aligned}$$

If $a = 0$ then $M \cdot 0 = -0$ is easily satisfied, and if $b = 0$ then $b \cdot \text{Null}(M) = 0$ is also trivially satisfied. Thus, $\chi^\lambda(u) \neq 0$ if $u_{jk} \lambda_{ik} = 0$ for all $i < j < k$ in $\mathcal{P}_{(m)}$ and $u_{1k} \lambda_{1l} + u_{jk} \lambda_{jl} = 0$ for all $1 < j < k < l$. Note that in the poset $\mathcal{P}_{(m)}$, $1 \not\leq j$ for $j \leq m$.

Suppose $a_{ij} \neq 0$. Note that $Mx = -a$ can only be satisfied if row (i, j) of M has a nonzero element. That is, there exists $i < j < k < l$ such that $\lambda_{il} \neq 0$. Consequently, we may assume $k < l_s$. If $j > m$, then $\delta = 1$, so $(Mx)_{1j} \neq 0$ if and only if $(Mx)_{ij} \neq 0$. However, $a_{1j} \neq 0$ and $(Mx)_{1j} \neq 0$ implies the first row of λ has two nonzero elements, contradicting the structure of S . Thus, if $a_{ij} \neq 0$ and $Mx = -a$ for some x , then $j \leq m$ and $k < l_s$.

Suppose $a_{ij} = u_{jk} \lambda_{ik} \neq 0$ with $j \leq m$ and $k < l_s$. By the definition of M and (9), $(i, k) = (i_{r-1}, l_{r-1})$. Note that $(Mx)_{ij} \neq 0$ if and only if $(Mx)_{i_{r'} j} \neq 0$. Since $u_{jk'} = 0$ for all $k' \neq k$, in this case $r' = r - 1$ or $|C_k| = |R_k|$. Then $Mx = -a$, where

$$x_{kl} = \begin{cases} -\lambda_{il_s}^{-1} \lambda_{ik}, & \text{if } l = l_s, \\ \lambda_{i_d l_d}^{-1} \lambda_{i_d l_s} \lambda_{il_s}^{-1} \lambda_{ik}, & \text{if } l = l_d, \\ 0, & \text{otherwise,} \end{cases} \quad \text{where } (k, l) \in C_k.$$

If $b_{kl} \neq 0$ and M has no nonzero entry in column (k, l) , then $b \cdot \text{Null}(M) \neq 0$. Thus, if $b_{kl} \neq 0$ we must have $\lambda_{jl}, \lambda_{il} \in \mathbb{F}_q^\times$ with $i < j$. In particular, $j \leq m$, and either $u_{1k} = 0$ or u has two nonzero elements. Since only the last column of S can have more than one nonzero entry, $l = l_s$, and $b_{kl_s} = u_{1k} \lambda_{1l_s} + u_{jk} \lambda_{jl_s}$. Note that

$$\dim(\text{Null}(M)) = \begin{cases} s - r, & \text{if } \delta' = 0, r' = r, \\ 0, & \text{otherwise.} \end{cases}$$

It follows that when $b \neq 0$, then b is perpendicular to $\text{Null}(M)$ if and only if $r' > r$ if and only if $|R_k| \geq |C_k|$ (if $\delta' = 1$, then $j > m$).

In the case that $j = i_{r-2}$ and $k = l_{r-1}$, we have

$$\theta(x \cdot b) = \theta(-\lambda_{il_s}^{-1} \lambda_{ik} (u_{1k} \lambda_{1l_s} + u_{jk} \lambda_{jl_s})), \quad \text{where } i = i_{r-1}.$$

Otherwise, $\theta(x \cdot b) = 1$. □

At this point, it may be helpful to give a more visual interpretation of the conditions in Theorem 4.1 by considering the configurations of superimposed graphs $G_{[\lambda]}$ and G_u . Recall, for $\lambda \in \mathcal{Z}_{(m)} \cup \mathcal{T}_{(m)}$ there is at most one connected component of $G_{[\lambda]}$ that can have more than one element (or can have a vertex in the first row of λ). Therefore, for $\lambda \in \mathcal{Z}_{(m)} \cup \mathcal{T}_{(m)}$, let

$$\begin{aligned} S_\lambda &= \text{the connected component of } G_{[\lambda]} \text{ that has a vertex in the first row} \\ \text{lc}(\lambda) &= \begin{cases} \min\{k \mid S_\lambda \text{ has a vertex in column } k\}, & \text{if } S_\lambda \neq \emptyset, \\ 0, & \text{otherwise,} \end{cases} \\ \text{br}(\lambda) &= \begin{cases} \max\{j \mid S_\lambda \text{ has a vertex in row } j\}, & \text{if } S_\lambda \neq \emptyset, \\ n, & \text{otherwise,} \end{cases} \\ \text{wt}(\lambda) &= \begin{cases} \#(\text{Nonzero entries in row } \text{br}(\lambda) \text{ of } \lambda) - 1, & \text{if } S_\lambda \neq \emptyset, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$


For example, if

$$\lambda = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & c & b \\ 0 & 0 & 0 & e & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & d \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \text{then } S_\lambda = \begin{array}{c} a \\ c-b \\ d \end{array}, \quad \begin{aligned} \text{lc}(\lambda) &= 5, \\ \text{br}(\lambda) &= 4, \\ \text{wt}(\lambda) &= 0. \end{aligned}$$

In the following discussion, we will suppress the values of the vertices and distinguish between G_{u-1} and $G_{[\lambda]}$ by the following conventions,

	G_{u-1}	$G_{[\lambda]}$
Vertices	•	■
Edges	•.....•	■—■

If $|S_\lambda| > 1$ and $\text{wt}(\lambda) = 0$, then add an edge to the non-zero vertex of row $\text{br}(\lambda)$ that extends West of this vertex,



thereby “completing” the comb.

Vertices of G_{u-1} see North in their column and East in their row, while vertices of $G_{[\lambda]}$ see South in their column and West in their row (in both cases they do not see the location they are in). That is,




Connected component S of G_{u-1} and T of $G_{[\lambda]}$ see *one-another* if when one superimposes their matrices, a vertex of S sees a vertex of T (and vice-versa).

The *tines* of S_λ are the pairs of horizontal edges with their leftmost vertices. For example, the tines of




Suppose $u \in U_{(m)}$ has at most two nonzero superdiagonal entries $u_{1k}, u_{jk} \in \mathbb{F}_q$, for some $1 \leq k \leq n$, and suppose $\lambda \in \mathfrak{n}_{(m)}^*$ such that $G_{[\lambda]}$ has exactly one connected component S . Then column k of u is *comb compatible* with S if the following conditions are satisfied.


(CC1) If column k of u has exactly one nonzero entry u_{jk} in column k and $S \neq S_\lambda$, then u_{jk} cannot see the vertex of S ,




(CC2) If $u_{1k}, u_{jk} \in \mathbb{F}_q^\times$, with $1 < j$ and $S \neq S_\lambda$, then the vertex of S cannot see u_{1k} or u_{jk} ,



(CC3) If column k of u has exactly one nonzero entry u_{jk} in column k and $S = S_\lambda$, then u_{jk} sees a vertex of S if and only if $j \leq m$, u_{jk} is South of the end of a tine and weakly North of the next tine to the South (if there is another tine),



(CC4) If $u_{1k}, u_{jk} \in \mathbb{F}_q^\times$, with $1 < j$ and $S = S_\lambda$, then a vertex of S sees u_{1k} or u_{jk} if and only if either u_{jk} is not South of the end of a tine but on a tine of S , or u_{jk} is South of the end of a tine and weakly North of the next tine to the South,



From this point of view, Theorem 4.1 translates to the following corollary.

Corollary 4.2. *Let $1 \leq m \leq n$. Suppose $u \in U_{(m)}$ has at most two nonzero superdiagonal entries $u_{1k}, u_{jk} \in \mathbb{F}_q$, for some $1 \leq k \leq n$. For $\lambda \in \mathcal{T}_{(m)}$, suppose $G_{[\lambda]}$ has one connected component S . Then*

(a) *The character degree*

$$\chi^\lambda(1) = \begin{cases} q^{|\{i < j \in \mathcal{P}_{(m)} \mid \lambda_{ik} \neq 0, \text{ for } k > j > i > 1, \lambda_{ik'} \neq 0 \text{ implies } k' \geq k\}|}, & \text{if } \text{wt}(\lambda) = 0, \\ q^{|\{i < j \in \mathcal{P}_{(m)} \mid \lambda_{ik} \neq 0, \text{ for } k > j > i, \lambda_{ik'} \neq 0 \text{ implies } k' \geq k\}|}, & \text{if } \text{wt}(\lambda) = 1. \end{cases}$$

(b) *The character*

$$\chi^\lambda(u) = 0$$

unless column k of u and S are comb compatible and in condition (CC4) if u_{1k} sees a vertex S and u_{jk} is not strictly South and weakly East of the end of a tine,

$$\begin{array}{c} u_{1k} \quad \lambda_{1l} \\ \blacksquare \quad \blacksquare \\ | \\ u_{jk} \quad \lambda_{jl} \\ \blacksquare \quad \blacksquare \\ | \\ \hline \end{array}, \quad \text{then} \quad u_{1k}\lambda_{1l} + u_{jk}\lambda_{jl} = 0. \quad (10)$$

(c) *If $\chi^\lambda(u) \neq 0$, then*

$$\chi^\lambda(u) = \chi^\lambda(1) \frac{\theta(u_{1k}\lambda_{1k} + u_{jk}\lambda_{jk})}{q^{c(u, \lambda)}} \prod_{\substack{i < j < l \\ \lambda_{il}, \lambda_{jl} \in \mathbb{F}_q^\times}} \theta(-\lambda_{il}^{-1} \lambda_{ik} (u_{1k}\lambda_{1l} + u_{jk}\lambda_{jl}))$$

where

$$c(u, \lambda) = \begin{cases} |\{l > k \mid \lambda_{il} \neq 0, \text{ for some } i < j\}|, & \text{if } u_{jk} \neq 0, j > m, \\ |\{l > k \mid \lambda_{il} \neq 0, \text{ for some } i < j\}|, & \text{if } u_{jk}, \lambda_{ij'}, \lambda_{il} \in \mathbb{F}_q^\times \text{ with } i < j' < j \leq m, k < l, \\ |\{l > k \mid \lambda_{il} \neq 0, \text{ for some } i < j\}| - 1, & \text{otherwise.} \end{cases}$$

will be of the form,

$$\begin{pmatrix} 0 & & & & \\ & \ddots & & & \\ & & 0 & & \\ \hline & & & 0 & \\ 0 & & & & \ddots \\ & & & & & 0 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 0 & & & & \\ & \ddots & & & \\ & & 0 & & \\ \hline & & & 0 & \\ 0 & & & & \ddots \\ & & & & & 0 \end{pmatrix}.$$

The *left corners* of u are the leftmost nonzero entries in the rows of $u - 1$. The *right corners* of u are the rightmost nonzero entries in the rows of $u - 1$.


Left and right corners *see* North in their column and East in their row, while top and bottom corners *see* South in their column and West in their row (in both cases they do not see the location they are in). That is,




Connected components S of G_u and T of $G_{[\lambda]}$ *see one-another* if when one superimposes their matrices, a corner of S sees a corner of T .

Fix $u \in \mathcal{Z}_{(m)}^\vee$ and $\lambda \in \mathcal{Z}_{(m)}$ with a connected component S of G_u and T of $G_{[\lambda]}$. The components S and T are *path compatible* if the following conditions are satisfied.


(PC1) If $S \neq S_u$ and $T \neq S_\lambda$, then S cannot see T ,




(PC2) If $S = S_u$ and $T \neq S_\lambda$, then S sees T if and only if T touches a vertical edge of S and no left corner of S sees T .



(PC3) If $S \neq S_u$ and $T = S_\lambda$, then S sees T if and only if S touches a vertical edge of T and no bottom corner of T sees S .



(PC4) If $S = S_u$ and $T = S_\lambda$, then S sees T if and only if T is never strictly South of S ; S ends weakly East of the beginning of T ; and left corners of S and bottom corners of T only see one-another horizontally.



Note that (PC1)-(PC4) are translations of (CC1)-(CC4) via the correspondence (6).
Define

$$\Theta : \mathcal{Z}_{(m)}^\vee \times \mathcal{Z}_{(m)} \longrightarrow \mathbb{C}$$

$$(u, \lambda) \mapsto \theta_{EB(u, \lambda)} \prod_{\substack{u_{jl}, u_{kl} \in \mathbb{F}_q^\times \\ \lambda_{il}, \lambda_{jl} \in \mathbb{F}_q^\times}} \theta(\text{bag}_{kl}(u) \text{bag}_{il}(\lambda)),$$

where

$$\theta_{EB(u, \lambda)} = \begin{cases} \theta(\text{bag}_{il}(u) \lambda_{il}), & \text{if } \lambda_{il} \neq 0, S_{u-1} \text{ ends with last} \\ & \text{vertex } u_{il} \text{ and a vertical edge,} \\ 1, & \text{otherwise.} \end{cases}$$

For a left corner u_{jk} in u , let

$$n_{jk}(\lambda) = \text{Number of bottom corners } \lambda_{il} \text{ with } i < j < k < l \text{ in } \mathcal{P}_{(m)}.$$

Corollary 4.3. *Let $u \in U_{(m)}$ be such that $u - 1 \in \mathcal{Z}_{(m)}^\vee$, and let $\lambda \in \mathcal{Z}_{(m)}$. Then*

(a) *The character*

$$\chi^\lambda(u) = 0$$

unless the connected components of G_{u-1} and $G_{[\lambda]}$ are pairwise path compatible, and in the superimposed matrices, every

$$\begin{array}{c} \leftarrow u_{ij} \text{---} \overset{j < k}{\text{---}} \lambda_{ik} \text{---} \rightarrow \\ \uparrow \\ \text{---} \end{array} \quad \text{implies} \quad \text{bag}_{ij}(u) \text{bag}_{ik}(\lambda) = 1. \quad (11)$$

(b) The character degree


$$\chi^\lambda(1) = \begin{cases} q^{|\{i < j \in \mathcal{P}_{(m)} \mid \lambda_{ik} \text{ is a bottom corner for some } k > j\}|}, & \text{if } \text{wt}(\lambda) = 0, \\ q^{|\{i < j \in \mathcal{P}_{(m)} \mid \lambda_{ik} \text{ is a top corner for some } k > j\}|}, & \text{if } \text{wt}(\lambda) = 1. \end{cases}$$

(c) If $\chi^\lambda(u) \neq 0$, then

$$\chi^\lambda(u) = \chi^\lambda(1) \frac{\theta(\lambda(u-1))\Theta(u, \lambda)}{q^{\text{wt}(\lambda)|\{m < j < k < \text{lc}(\lambda) \mid u_{jk} \neq 0\}|}} \prod_{\substack{\text{left corners} \\ u_{jk}}} q^{-n_{jk}(\lambda)}.$$

Proof. This corollary follows directly from Corollary 4.2 with the following observations, using (6).

(a) If a bottom corner of T sees a left corner of S horizontally, then we are in the situation of (11), so



so the comb representations of λ and u must satisfy condition (10). However, this is equivalent to $\text{bag}_{ij}(u)\text{bag}_{ik}(\lambda) = 1$. Thus, Corollary 4.2 (a) is satisfied if and only if Corollary 4.3 (a) is satisfied.

(b) is straight-forward translation of the combinatorics.

(c) Let $\tilde{\lambda} \in \mathcal{T}_{(m)}$ be the comb representative corresponding to $\lambda \in \mathcal{Z}_{(m)}$ and let $\tilde{u} \in \mathcal{T}_{(m)}^\vee$ be the comb representative in the same superclass as $u \in \mathcal{Z}_{(m)}^\vee$. If $\tilde{u}_{jk} \neq 0$, then

$$c(\tilde{u}[k], \tilde{\lambda}) = \begin{cases} n_{jk}(\lambda), & \text{if } \text{wt}(\lambda) = 0, j > m, \\ n_{jk}(\lambda) + 1, & \text{if } \text{wt}(\lambda) = 1, j > m, \\ n_{jk}(\lambda), & \text{if } \tilde{\lambda}_{ij'}, \tilde{\lambda}_{il} \in \mathbb{F}_q^\times, \text{ with } i < j' < j \leq m, k < l, \\ n_{jk}(\lambda), & \text{otherwise.} \end{cases}$$

Thus,

$$q^{\text{wt}(\lambda)|\{m < j < k < \text{lc}(\lambda) \mid u_{jk} \neq 0\}|} \prod_{\text{left corners } u_{jk}} q^{n_{jk}(\lambda)} = \prod_k q^{c(\tilde{u}[k], \tilde{\lambda})}.$$

If $\chi^\lambda(u) \neq 0$, and we have no configurations of the form

$$\begin{array}{c} \lambda_{1l} \\ \blacksquare \\ \vdots \\ u_{jl} \end{array} \longleftrightarrow \begin{array}{c} \lambda_{1l} \text{bag}_{jl}(u) \\ \blacksquare \end{array} \quad (12)$$

$$\begin{array}{c} \lambda_{1l} \\ \blacksquare \\ \vdots \\ \lambda_{ik} \text{---} \lambda_{ik'} \\ \vdots \\ u_{jk} \end{array} \longleftrightarrow \begin{array}{c} \text{bag}_{jk}(u) \text{---} \lambda_{1l} \\ \blacksquare \\ \lambda_{ik} \text{---} \lambda_{1l} \text{bag}_{ik'}(\lambda) \\ \vdots \\ u_{j'k} \text{---} \lambda_{1l} \text{bag}_{j'k}(\lambda) \end{array}, \quad (13)$$

then both character formulas are equal. If (12) occurs then both the path character formula and the comb character formula get a factor of

$$\theta(\lambda_{1l} \text{bag}_{jl}(u)).$$

If (13) occurs then the path character formula gets a factor

$$\theta(\lambda_{j'k} u_{j'k}) \theta(\text{bag}_{jk}(u) \text{bag}_{ik}(\lambda)),$$

and the comb character formula gets a factor of

$$\theta \left(- \left(- \lambda_{1l} \text{bag}_{ik'}(\lambda) \right)^{-1} \lambda_{ik} \left(\text{bag}_{jk}(u) \lambda_{1l} + u_{j'k} \left(- \lambda_{1l} \text{bag}_{j'k}(\lambda) \right) \right) \right),$$

However, a simple computation shows that these two quantities are equal. Thus, the character formulas for the two types of representatives are equal. $\square$

4.4 Example


Let $n = 7, m = 4$,

$$u = \begin{pmatrix} 1 & 0 & 0 & 0 & a & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & d & 0 \\ 0 & 0 & 1 & e & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & b & c & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \lambda = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & x \\ 0 & 0 & 0 & 0 & 0 & 0 & y \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & z & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

so the decorated superimposed matrix of $u - 1$ and λ is

$$\begin{pmatrix} 0 & 0 & 0 & 0 & a & 0 & x \\ 0 & 0 & 0 & 0 & 0 & d & y \\ 0 & 0 & 0 & e & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & bz & c & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

All the connected components are pairwise path compatible,



Thus,

$$\chi^\lambda(u) \neq 0$$

if and only if $\text{bag}(d)\text{bag}(y) = 1$ if and only if $d(-c^{-1})b(-a^{-1})y(-x^{-1}) = 1$. If $\chi^\lambda(u) \neq 0$, then

$$\chi^\lambda(u) = q^4 \theta(bz) \frac{1}{q^3} \cdot 1.$$

5 Interpolating from U_n to U_{n-1}

This section uses a restriction rule from $U_{(m-1)}$ to $U_{(m)}$ for supercharacters to deduce a restriction rule from U_n to U_{n-1} .

5.1 A restriction rule for $U_{(m)}$

Note that if $\lambda \in \mathcal{Z}_{(m-1)}$, then $\lambda \in \mathcal{Z}_{(m)}$ unless $\text{lc}(\lambda) = m$. Let $\lambda^\downarrow \in \mathcal{Z}_{(m)}$ be given by

$$\lambda^\downarrow = \begin{cases} \lambda, & \text{if } \text{lc}(\lambda) \neq m, \\ \lambda - \lambda_{im}e_{im} & \text{if } \text{lc}(\lambda) = m \text{ and } i \text{ is minimal so } \lambda_{im} \neq 0. \end{cases}$$

Then $\lambda^\downarrow$ is the path orbit representative of $U_{(m)}\lambda U_{(m)}$.

For $u \in \mathcal{Z}_{(m)}^\vee$, let

$$u^\uparrow = \begin{cases} u - u_{1j}e_{1j} - u_{mk}e_{mk} - u_{1j}u_{mj}^{-1}u_{mk}e_{1k}, & \text{if } u_{1j}, u_{mj}, u_{mk} \in \mathbb{F}_q^\times, j < k, \\ u - u_{1j}e_{1j}, & \text{if } u_{1j}, u_{mj} \in \mathbb{F}_q^\times, \text{wt}(u-1) = 1, \\ u, & \text{otherwise,} \end{cases}$$

so $u^\uparrow$ is the path orbit representative of $U_{(m-1)}uU_{(m-1)}$.

Theorem 5.1. *Let $\lambda \in \mathcal{Z}_{(m-1)}$. Then*

$$\text{Res}_{U_{(m)}}^{U_{(m-1)}}(\chi^\lambda) = \begin{cases} \chi^{\lambda^\downarrow}, & \text{if } \text{lc}(\lambda) = m \text{ or } \text{wt}(\lambda) = 0, \\ q\chi^{\lambda^\downarrow}, & \text{if } \text{lc}(\lambda) > m, \text{wt}(\lambda) = 1, \lambda_{mj} \neq 0 \text{ for some } j > \text{lc}(\lambda), \\ \sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + te_{m\text{lc}(\lambda)}}, & \text{if } \text{lc}(\lambda) > m, \text{wt}(\lambda) = 1, \lambda_{mj} = 0 \text{ for all } j > \text{lc}(\lambda). \end{cases}$$

Proof. Since at most one component of λ can satisfy case three of Theorem 5.1, it suffices to prove the statement for the case that $G_{[\lambda]}$ has one connected component. Note that Case 2 does not occur unless $G_{[\lambda]}$ has at least two components, and it follows directly from cases 1 and 3.

First, suppose that $S_\lambda = \emptyset$. Then $\text{wt}(\lambda) = \text{lc}(\lambda) = 0$, and $\lambda^\downarrow = \lambda$. Since the support of λ excludes the first row, the conditions implied by $\mathcal{P}_{(m)}$ are the same as the conditions implied by $\mathcal{P}_{(m-1)}$ in the character formulae and $\text{Res}_{U_{(m)}}^{U_{(m-1)}}(\chi^\lambda) = \chi^{\lambda^\downarrow}$.

Suppose that $\lambda = [S_\lambda]$. If $|S_\lambda| = 1$ with $\text{lc}(\lambda) = m$, then $\text{br}(\lambda) = 1$. In this case, $1 \not\leq m$ in $\mathcal{P}_{(m)}$ implies $\text{Res}_{U_{(m)}}^{U_{(m-1)}}(\chi^\lambda) = \mathbb{1}_{U_{(m)}} = \chi^{\lambda^\downarrow}$.

For the rest of the proof we will assume $\lambda = [S_\lambda]$ and $|S_\lambda| > 1$ or $\text{lc}(\lambda) > m$. Note that if $\text{lc}(\lambda) > m$, $\text{wt}(\lambda) = 1$, and $\lambda_{mj} = 0$ for all $j > k$, then

$$\sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + te_{mk}}(u) = \begin{cases} \chi^{\lambda^\downarrow}(u) \sum_{t \in \mathbb{F}_q} \theta(tu_{m,\text{lc}(\lambda)}), & \text{if } u_{mk} = 0, k < \text{lc}(\lambda), \\ \chi^{\lambda^\downarrow}(u), & \text{otherwise.} \end{cases} \quad (14)$$

Thus, all three cases of Theorem 5.1 require an understanding of the relationship between the values $\chi^{\lambda^\downarrow}(u)$ and $\chi^\lambda(u^\uparrow)$.

Since $\chi^{\lambda^\downarrow}$ is a supercharacter, by Lemma 4.1 WLOG we assume G_{u-1} has one connected component. It follows from the combinatorics that the two values $\chi^\lambda(u^\uparrow)$ and $\chi^{\lambda^\downarrow}(u)$ are simultaneously zero or nonzero unless $\text{wt}(\lambda) = 1$, $u_{m,\text{lc}(\lambda)} \neq 0$, and $u_{mk} = 0$ for $k > \text{lc}(\lambda)$. However, in this exceptional case,

$$\sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + t e_{mk}}(u) = \chi^{\lambda^\downarrow}(u) \sum_{t \in \mathbb{F}_q} \theta(t u_{m,\text{lc}(\lambda)}) = 0 = \chi^\lambda(u^\uparrow). \quad (15)$$

Suppose $\chi^\lambda(u^\uparrow)$ and $\chi^{\lambda^\downarrow}(u)$ are both nonzero. By inspection,

$$\frac{\chi^\lambda(1)}{\chi^{\lambda^\downarrow}(1)} = \begin{cases} 1, & \text{if } \text{lc}(\lambda) = m \text{ or } \text{wt}(\lambda) = 0, \\ q, & \text{otherwise.} \end{cases} \quad (16)$$

Since $\lambda^\downarrow$ has the same bottom corners as λ and $u^\uparrow$ has the same left corners as u ,

$$\frac{q^{\text{wt}(\lambda) \# \{m-1 < j < \text{lc}(\lambda) \mid u_{m-1,j}^\uparrow \neq 0\}} \prod_{\substack{\text{left corners } u_{jk}^\uparrow}} q^{n_{jk}(\lambda)}}{q^{\text{wt}(\lambda) \# \{m < j < \text{lc}(\lambda^\downarrow) \mid u_{mj} \neq 0\}} \prod_{\substack{\text{left corners } u_{jk}}} q^{n_{jk}(\lambda^\downarrow)}} = \frac{q^{\text{wt}(\lambda) \# \{m-1 < j < \text{lc}(\lambda) \mid u_{m-1,j}^\uparrow \neq 0\}}}{q^{\text{wt}(\lambda) \# \{m < j < \text{lc}(\lambda^\downarrow) \mid u_{mj} \neq 0\}}}. \quad (17)$$

Note that

$$\theta(\lambda(u-1)) = \begin{cases} \theta(\lambda^\downarrow(u-1))\theta(\lambda_{im}u_{im}), & \text{if } \text{lc}(\lambda) = m, i \text{ is minimal such that } \lambda_{im} \neq 0, \\ \theta(\lambda^\downarrow(u-1)), & \text{otherwise.} \end{cases}$$

However, in the first case, $u_{im} \neq 0$ implies $\chi^\lambda(u) = 0$, since $m < m+1$ and $u \in \mathcal{Z}_{(m)}$. Thus, for all $\text{lc}(\lambda)$ with $\chi^\lambda(u) \neq 0$,

$$\theta(\lambda^\downarrow(u-1)) = \theta(\lambda(u-1)).$$

In fact, if $S_{u-1} = \emptyset$ or $\text{br}(u-1) < m$, then $u^\uparrow = u$ and $\theta(\lambda(u-1)) = \theta(\lambda(u^\uparrow-1))$. In these cases we also have $\Theta(u^\uparrow, \lambda) = \Theta(u, \lambda^\downarrow)$.

Suppose $S_{u-1} \neq \emptyset$ and $\text{br}(u-1) = m$. If $S_{u-1} = \{u_{1k}, u_{mk}\}$, then $S_{u^\uparrow-1} = \{u_{mk}\}$, and

$$\theta(\lambda(u-1)) = \theta(\lambda_{1k}u_{1k})\theta(\lambda(u^\uparrow-1)).$$

However, since $\lambda \in \mathcal{Z}_{(m-1)}$, $\lambda_{1k} \neq 0$ implies $\chi^\lambda(u^\uparrow) = \chi^\lambda(u) = 0$.

If $\{u_{1j}, u_{mj}, u_{mk}\} \subseteq S_{u-1}$ with $j < k$, then $u_{1j}^\uparrow = u_{mk}^\uparrow = 0$, $u_{mj}^\uparrow = u_{mj}$, and $u_{1k}^\uparrow = -u_{1j}u_{mj}^{-1}u_{mk}$. Thus,

$$\theta(-\lambda_{1k}u_{1j}u_{mj}^{-1}u_{mk})\theta(\lambda(u-1)) = \theta(\lambda_{1j}u_{1j} + \lambda_{mk}u_{mk})\theta(\lambda(u^\uparrow-1)).$$

If $\lambda_{1j} \neq 0$ or $\lambda_{mk} \neq 0$, then $\chi^\lambda(u) = 0$, so

$$\theta(-\lambda_{1k}u_{1j}u_{mj}^{-1}u_{mk})\theta(\lambda(u-1)) = \theta(\lambda(u^\uparrow-1)).$$

If $\lambda_{1k} \neq 0$, then $\chi^\lambda(u) \neq 0$ only if

- (1) $S_{u-1} = \{u_{1j}, u_{mj}, u_{mk}\}$ and $\lambda_{ik} \neq 0$ for some $1 < i < k$,
- (2) there exists $1 < i < k$ such that $u_{ik}, \lambda_{ik} \in \mathbb{F}_q^\times$.

In both cases,

$$\Theta(u, \lambda)\theta(-\lambda_{1k}u_{1j}u_{mj}^{-1}u_{mk}) = \Theta(u^\uparrow, \lambda).$$

Note that in the first case, $\theta_{EB(u^\uparrow, \lambda)} = \theta(-\lambda_{1k}u_{1j}u_{mj}^{-1}u_{mk})$.

Thus, for all $u \in \mathcal{Z}_{(m)}$ and $\lambda \in \mathcal{Z}_{(m-1)}$ such that $\chi^\lambda(u) \neq 0$,

$$\Theta(u, \lambda^\downarrow)\theta(\lambda^\downarrow(u-1)) = \Theta(u, \lambda)\theta(\lambda(u-1)) = \Theta(u^\uparrow, \lambda)\theta(\lambda(u^\uparrow-1)). \quad (18)$$

We now use (16), (17), and (18) to check the cases of the theorem. Suppose $\text{lc}(\lambda) = m$ or $\text{wt}(\lambda) = 0$, then by (16), (17), and (18),

$$\chi^{\lambda^\downarrow}(u) = \chi^\lambda(u^\uparrow).$$

Suppose $\text{lc}(\lambda) > m$ and $\text{wt}(\lambda) = 1$. If $\lambda_{mj} = 0$ for all $j \geq \text{lc}(\lambda)$, then by (14),

$$\sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + te_{\text{mlc}(\lambda)}}(u) = \begin{cases} \chi^{\lambda^\downarrow}(u) \sum_{t \in \mathbb{F}_q} \theta(tu_{m, \text{lc}(\lambda)}), & \text{if } u_{mj} = 0, j < \text{lc}(\lambda), \\ \chi^{\lambda^\downarrow}(u), & \text{otherwise.} \end{cases}$$

In the first case, $\chi^\lambda(1) = q\chi^{\lambda^\downarrow}(1)$ and the ratio in (17) is 1. Thus, by (18) and (15) in this case,

$$\begin{aligned} \sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + te_{\text{mlc}(\lambda)}}(u) &= \begin{cases} q^{-1}\chi^{\lambda^\downarrow}(u^\uparrow)q, & \text{if } u_{\text{mlc}(\lambda)} = 0, \\ \chi^\lambda(u^\uparrow)(=0), & \text{otherwise.} \end{cases} \\ &= \chi^\lambda(u^\uparrow) \end{aligned}$$

In the second case of (14), $\chi^\lambda(1) = q\chi^{\lambda^\downarrow}(1)$ and the ratio in (17) is q . Thus,

$$\sum_{t \in \mathbb{F}_q} \chi^{\lambda^\downarrow + te_{\text{mlc}(\lambda)}}(u) = \chi^{\lambda^\downarrow}(u) = q^{-1}\chi^\lambda(u^\uparrow)q = \chi^\lambda(u^\uparrow),$$

completing the last case. □

5.2 A restriction rule for U_n

For $\lambda \in \mathcal{S}_n(q)$, and $i, k \in \mathbb{Z}_{\geq 1}$, let

$$\lambda *_{i} \{k\} = \begin{cases} \lambda, & \text{if } i = k, \\ q\lambda *_{i+1} \{k\}, & \text{if } \lambda_{il} \neq 0 \text{ for some } l > k, \\ \lambda|_{\lambda_{ik}=0} *_{i+1} \{k\}, & \text{if } \lambda_{ik} \neq 0, \\ \lambda *_{i+1} \{k\} + \sum_{\substack{t \in \mathbb{F}_q^\times \\ \lambda_{ij}=0}} \lambda|_{\lambda_{ik}=t} *_{i+1} \{j\}, & \text{if } \lambda_{ij} \neq 0 \text{ for some } i < j < k, \\ \lambda *_{i+1} \{k\} + \sum_{t \in \mathbb{F}_q^\times} \lambda|_{\lambda_{ik}=t}, & \text{if } \lambda_{ij} = 0 \text{ for all } j > i. \end{cases} \quad (19)$$

We will extend this product to supercharacters by the convention

$$\chi^\mu *_i \chi^{\{k\}} = \sum_{\lambda} c_{\mu k}^{\lambda} \chi^{\lambda}, \quad \text{if} \quad \mu *_i \{k\} = \sum_{\lambda} c_{\mu k}^{\lambda} \lambda.$$

Corollary 5.1. *For $\lambda \in \mathcal{S}_n(q)$,*

$$\text{Res}_{U_{n-1}}^{U_n}(\chi^{\lambda}) = \begin{cases} \chi^{\lambda} *_1 \chi^{\{k\}}, & \text{if } \lambda_{1k} \neq 0, \text{ for some } 1 < k, \\ \chi^{\lambda}, & \text{otherwise.} \end{cases}$$

Proof. Note that for $1 \leq m \leq n$, Theorem 5.1 implies

$$\text{Res}_{U_{(m)}}^{U_{(m-1)}}(\chi^{\lambda}) = \begin{cases} \chi^{\lambda}, & \text{if } \text{lc}(\lambda) \neq m \text{ and } \text{wt}(\lambda) = 0, \\ \chi^{\lambda - \lambda_{im} e_{im}}, & \text{if } \text{lc}(\lambda) = m, \\ q\chi^{\lambda}, & \text{if } \text{lc}(\lambda) > m, \text{ wt}(\lambda) = 1, \lambda_{mj} \neq 0 \text{ for some } j > \text{lc}(\lambda), \\ \sum_{t \in \mathbb{F}_q} \chi^{\lambda + t e_{m \text{lc}(\lambda)}}, & \text{if } \text{lc}(\lambda) > m, \text{ wt}(\lambda) = 1, \lambda_{mj} = 0, \text{ for all } j > \text{lc}(\lambda). \end{cases}$$

The multiplication given by (19) is an iterative version of the restrictions from $U_{(m-1)}$ to $U_{(m)}$, where the last two cases in (19) correspond to the last case in the restriction, depending on whether there exists $\lambda_{mj} \in \mathbb{F}_q^{\times}$ for some $j < \text{lc}(\lambda)$. $\square$

Remark. It follows that the coefficients of the restriction are polynomial in q .

5.3 Examples

Example 1. Consider the case $q = 2$. Then we may choose our representatives $\lambda \in \mathfrak{n}_n^*$ and $u \in U_n$ to correspond to set partitions of $\{1, 2, \dots, n\}$. For example, if

$$\lambda = \{1 \curvearrowright 5 \mid 2 \curvearrowright 6 \mid 3 \curvearrowright 4\},$$

then

$$\begin{aligned} \text{Res}_{U_5}^{U_6}(\chi^{\lambda}) &= \chi^{\{1-5|2-6|3-4\}} *_1 \{5\} \\ &= \chi^{\{1|5|2-6|3-4\}} *_2 \{5\} \\ &= 2\chi^{\{1|5|2-6|3-4\}} *_3 \{5\} \\ &= 2\chi^{\{1|5|2-6|3-4\}} *_4 \{5\} + 2\chi^{\{1|2-6|3-5|4\}} *_4 \{4\} \\ &= 2\chi^{\{1|5|2-6|3-4\}} *_5 \{5\} + 2\chi^{\{1|2-6|3-4-5\}} *_5 \{5\} + 2\chi^{\{1|2-6|3-5|4\}} \\ &= 2\chi^{\{1|5|2-6|3-4\}} + 2\chi^{\{1|2-6|3-4-5\}} + 2\chi^{\{1|2-6|3-5|4\}} \end{aligned}$$

Note that the final result gives the representative of U_{n-1} as a submatrix of U_n (since by construction the first row will always be zero). To obtain a set partition of $n - 1$, we would remove 1 and renumber the rest of the entries $j \mapsto j - 1$.

Alternatively, this algorithm may be viewed as a “bumping algorithm”, where we replace the 1 by all other “possibilities,” suitably defined.

Example 2. Linear characters of U_n are those characters whose superclass representative satisfies, i and j are in the same part only if $i + 1, i + 2, \dots, j - 1$ are also in the part. In this case,

$$\chi^\lambda *_1 \{j\} = \chi^\lambda.$$

Example 3. On the opposite extreme with have the case

$$\lambda = \{1 \curvearrowright n \mid 2 \mid 3 \mid \dots \mid n - 1\}.$$

In this case,

$$\chi^\lambda *_1 \{n\} = \sum_{\substack{\mu \in \mathcal{S}_n(q), \\ \lambda_{ij}=0, 1 \leq i < j < n}} \chi^\mu.$$

5.4 An alternate embedding of U_{n-1}

The paper [13] uses a different embedding of U_{n-1} into U_n (obtained by removing the last column rather than the first row). This alternate embedding gives a different restriction rule. For $\mu \in \mathcal{S}_n(q)$ and $j, l \in \mathbb{Z}_{\geq 1}$, let

$$\{j\} *_l \mu = \begin{cases} \mu, & \text{if } j = l, \\ q(\{j\} *_l \mu), & \text{if there is } i < j \text{ with } \mu_{il} \neq 0, \\ \{j\} *_l \mu|_{\mu_{jl}=0}, & \text{if } \mu_{jl} \neq 0, \\ \{j\} *_l \mu + \sum_{\substack{t \in \mathbb{F}_q^\times \\ \mu_{kl}=0 \\ \mu_{jl}=t}} \{k\} *_l \mu, & \text{if there is } k > j \text{ with } \mu_{kl} \neq 0, \\ \{j\} *_l \mu + \sum_{t \in \mathbb{F}_q^\times} \mu|_{\mu_{jl}=t}, & \text{otherwise.} \end{cases}$$

Then by symmetry arguments from [13], we obtain the following corollary for this alternate embedding of U_{n-1} into U_n .

Corollary 5.2. For $\lambda \in \mathcal{S}_n(q)$,

$$\text{Res}_{U_{n-1}}^{U_n}(\chi^\lambda) = \begin{cases} \chi^{\{j\}} *_n \chi^\lambda, & \text{if } \lambda_{jn} \neq 0, \text{ for some } j < n, \\ \chi^\lambda, & \text{otherwise.} \end{cases}$$

In particular, unlike in the symmetric group representation theory, the decomposition of restricted characters depends on the embedding of U_{n-1} into U_n . Other than the “anti-transpose” symmetry of this section, it is currently unknown what kind of combinatorial relationship might exist between the restriction coefficients of different embeddings.

Remark. The paper [13] studies the problem of induction from U_{n-1} to U_n . While this is done directly, one could obtain analogous results from this paper by Frobenius reciprocity.

References

- [1] André, C. “Basic characters of the unitriangular group,” *J. algebra* **175** (1995), 287–319.
- [2] André, C. “Irreducible characters of finite algebra groups,” *Matrices and group representations Coimbra, 1998* Textos Mat. Sér B **19** (1999), 65–80.
- [3] André, C. “The basic character table of the unitriangular group,” *J. algebra* **241** (2001), 437–471.
- [4] André, C. “Basic characters of the unitriangular group (for arbitrary primes),” *Proc. Amer. Math. Soc.* **130** (2002), 1934–1954.
- [5] André, C; Neto, A. “Super-characters of finite unipotent groups of types B_n , C_n and D_n ,” August 2006 preprint.
- [6] André, C; Nicolás, A. “Supercharacters of the adjoint group of a finite radical ring,” August 2006 preprint.
- [7] Arregi, J; Vera-Lopez, A. “Computing in unitriangular matrices over finite fields.” *Linear algebra and Appl.* **387** (2004), 193–219.
- [8] Arias-Castro E; Diaconis, P; Stanley, R. “A super-class walk on upper-triangular matrices,” *J. algebra* **278** (2004), 739–765.
- [9] Boyarchenko, M; Drinfeld, V. “A motivated introduction to character sheaves and the orbit method for unipotent groups in positive chracteristic.” Preprint (2006), arXiv.0609769.
- [10] Diaconis, P; Isaacs, M. “Supercharacters and superclasses for algebra groups,” To appear in *Trans. Amer. Math. Soc.*, 2006.
- [11] Diaconis, P; Saloff-Coste, L. “Comparison techniques for random walk on finite groups,” *Ann. Probab.* **21** (1993), 2131–2156.
- [12] Diaconis, P; Thiem, N. “Supercharacter formulas for pattern groups.” To appear in *Trans. Amer. Math. Soc.*, 2007.
- [13] Marberg, E; Thiem, N. “Superinduction for pattern groups.” A 2007 preprint.
- [14] Yan, N. *Representation theory of the finite unipotent linear groups*, Unpublished Ph.D. Thesis, Department of mathematics, University of Pennsylvania, 2001.
- [15] Yan, N. “Representations of finite unipotent linear groups by the method of clusters,” 2006 preprint.