

Tetrahedra on deformed spheres and integral group cohomology

Pavle V. M. Blagojević*

Mathematički Institut
Knez Michailova 35/1
11001 Beograd, Serbia

pavleb@mi.sanu.ac.rs

Günter M. Ziegler**

Inst. Mathematics, MA 6-2
TU Berlin
D-10623 Berlin, Germany

ziegler@math.tu-berlin.de

Submitted: Aug 28, 2008; Accepted: Jun 4, 2009; Published: Jun 10, 2009

Mathematics Subject Classification: 55S91, 55M20, 52C99

Dedicated to Anders Björner on the occasion of his 60th birthday

Abstract

We show that for every injective continuous map $f : S^2 \rightarrow \mathbb{R}^3$ there are four distinct points in the image of f such that the convex hull is a tetrahedron with the property that two opposite edges have the same length and the other four edges are also of equal length. This result represents a partial result for the topological Borsuk problem for $\mathbb{R}^3$. Our proof of the geometrical claim, via Fadell–Husseini index theory, provides an instance where arguments based on group cohomology with integer coefficients yield results that cannot be accessed using only field coefficients.

1 Introduction

The motivation for the study of the existence of particular types of tetrahedra on deformed 2-spheres is twofold. The topological Borsuk problem, as formulated by Soibelman in 1977 [6] (“estimate the minimal Borsuk partition number for the unit ball in $\mathbb{R}^n$ for general metrics!”), along with the square peg problem [5] first posed by Toeplitz 1911 (“does every Jordan curve contain the vertices of a square?”) inspire the search for possible polytopes with nice metric properties whose vertices lie on the continuous images of spheres. Beyond their intrinsic interest, these problems can be used as testing grounds for tools from equivariant topology, e.g. for comparing the strength of Fadell–Husseini index theory with ring resp. field coefficients.

*Supported by the grant 144018 of the Serbian Ministry of Science and Technological development

**Supported by the German Research Foundation DFG

The following theorem will be proved through the use of Fadell–Husseini index theory with coefficients in the ring $\mathbb{Z}$. It is also going to be demonstrated that Fadell–Husseini index theory with coefficients in field $\mathbb{F}_2$ has no power in this instance (Section 4.1).

Theorem 1.1. *Let $f : S^2 \rightarrow \mathbb{R}^3$ be an injective continuous map. Then its image contains vertices of a tetrahedron that has at least the symmetry of a square. That is, there are four distinct points ξ_1, ξ_2, ξ_3 and ξ_4 on S^2 such that*

$$d(f(\xi_1), f(\xi_2)) = d(f(\xi_2), f(\xi_3)) = d(f(\xi_3), f(\xi_4)) = d(f(\xi_4), f(\xi_1))$$

and

$$d(f(\xi_1), f(\xi_3)) = d(f(\xi_2), f(\xi_4)).$$

Thus the tetrahedron may even be regular and thus have symmetry group S_4 ; it may also degenerate to a (planar) square.

Remark 1.2. The proof is not going to use any properties of $\mathbb{R}^3$ except that it is a metric space. Thus in the statement of the theorem, $\mathbb{R}^3$ can be replaced by any metric space (M, d) .




Figure 1: A D_8 -invariant tetrahedron on a deformed 2-sphere

Let us try to relate this to the square peg problem and the topological Borsuk problem:

The square peg problem is settled for various classes of sufficiently piecewise-smooth Jordan curves, but open in general. Unfortunately, the methods used for the proof of Theorem 1.1 do not imply any conclusion when applied to the square peg problem (see Section 4.2). On the other hand, if the square peg problem could be solved for the continuous Jordan curves, then it would imply the result of Theorem 1.1.

The first open instance of the topological Borsuk problem considers the existence of a collection of four points with equal pairwise d -distances in a general metric space $(\mathbb{R}^3, d)$. The main result of the paper does not provide any new information concerning the topological Borsuk problem, as we work in the restricted parameter space $\{(x_1, x_2, x_3, x_4) \in (S^1)^4 \mid x_1 \neq x_3 \text{ or } x_2 \neq x_4\}$ whose dimension is much smaller compared to the one used in the topological Borsuk problem $\{(x_1, x_2, x_3, x_4) \in (\mathbb{R}^3)^4 \mid x_1 \neq x_3 \text{ or } x_2 \neq x_4\}$.

2 Introducing the equivariant question

Let $f : S^2 \rightarrow \mathbb{R}^3$ be an injective continuous map. Denote by D_8 the symmetry group of a square, that is, the 8-element dihedral group $D_8 = \langle \omega, j \mid \omega^4 = j^2 = 1, \omega j = j\omega^3 \rangle$.

A few D_8 -representations.

The real vector spaces

$$\begin{aligned} U_4 &= \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 \mid x_1 + x_2 + x_3 + x_4 = 0\}, \\ U_2 &= \{(x_1, x_2) \in \mathbb{R}^2 \mid x_1 + x_2 = 0\} \end{aligned}$$

are real D_8 -representations with actions given by

(a) for $(x_1, x_2, x_3, x_4) \in U_4$:

$$\omega \cdot (x_1, x_2, x_3, x_4) = (x_2, x_3, x_4, x_1), \quad j \cdot (x_1, x_2, x_3, x_4) = (x_3, x_2, x_1, x_4),$$

(b) for $(x_1, x_2) \in U_2$:

$$\omega \cdot (x_1, x_2) = (x_2, x_1), \quad j \cdot (x_1, x_2) = (x_2, x_1),$$

The configuration space.

Let $X = S^2 \times S^2 \times S^2 \times S^2$ and let Y be the subspace given by

$$Y = \{(x, y, x, y) \mid x, y \in S^2\} \approx S^2 \times S^2.$$

The configuration space to be considered is the space

$$\Omega := X \setminus Y.$$

Let a D_8 -action on X be induced by

$$\omega \cdot (\xi_1, \xi_2, \xi_3, \xi_4) = (\xi_2, \xi_3, \xi_4, \xi_1), \quad j \cdot (\xi_1, \xi_2, \xi_3, \xi_4) = (\xi_4, \xi_3, \xi_2, \xi_1),$$

for $(\xi_1, \xi_2, \xi_3, \xi_4) \in X$.

A test map.

Let $\tau : \Omega \rightarrow U_4 \times U_2$ be a map defined for $(\xi_1, \xi_2, \xi_3, \xi_4) \in X$ by

$$\tau(\xi_1, \xi_2, \xi_3, \xi_4) = (d_{12} - \frac{\Delta}{4}, d_{23} - \frac{\Delta}{4}, d_{34} - \frac{\Delta}{4}, d_{41} - \frac{\Delta}{4}) \times (d_{13} - \frac{\Phi}{2}, d_{24} - \frac{\Phi}{2}) \quad (1)$$

where $d_{ij} = d_{ji} := d(f(\xi_i), f(\xi_j))$ and

$$\Delta = d_{12} + d_{23} + d_{34} + d_{14}, \quad \Phi = d_{13} + d_{24}.$$

With the D_8 -actions introduced above the test map τ is D_8 -equivariant. Indeed,

$$\begin{aligned}\tau(\omega \cdot (\xi_1, \xi_2, \xi_3, \xi_4)) &= \tau(\xi_2, \xi_3, \xi_4, \xi_1) \\ &= (d_{23} - \frac{\Delta}{4}, d_{34} - \frac{\Delta}{4}, d_{41} - \frac{\Delta}{4}, d_{12} - \frac{\Delta}{4}) \times (d_{24} - \frac{\Phi}{2}, d_{13} - \frac{\Phi}{2}) \\ &= \omega \cdot ((d_{12} - \frac{\Delta}{4}, d_{23} - \frac{\Delta}{4}, d_{34} - \frac{\Delta}{4}, d_{41} - \frac{\Delta}{4}) \times (d_{13} - \frac{\Phi}{2}, d_{24} - \frac{\Phi}{2}))\end{aligned}$$

and

$$\begin{aligned}\tau(j \cdot (\xi_1, \xi_2, \xi_3, \xi_4)) &= \tau(\xi_4, \xi_3, \xi_2, \xi_1) \\ &= (d_{43} - \frac{\Delta}{4}, d_{32} - \frac{\Delta}{4}, d_{21} - \frac{\Delta}{4}, d_{14} - \frac{\Delta}{4}) \times (d_{42} - \frac{\Phi}{2}, d_{31} - \frac{\Phi}{2}) \\ &= j \cdot ((d_{12} - \frac{\Delta}{4}, d_{23} - \frac{\Delta}{4}, d_{34} - \frac{\Delta}{4}, d_{41} - \frac{\Delta}{4}) \times (d_{13} - \frac{\Phi}{2}, d_{24} - \frac{\Phi}{2})).\end{aligned}$$

The following proposition connects our set-up with the tetrahedron problem.

Proposition 2.1. *If there is no D_8 equivariant map*

$$\alpha : \Omega \rightarrow (U_4 \times U_2) \setminus (\{\mathbf{0}\} \times \{\mathbf{0}\}) \quad (2)$$

then Theorem 1.1 follows.

Proof. If there is no D_8 equivariant map $\Omega \rightarrow (U_4 \times U_2) \setminus (\{\mathbf{0}\} \times \{\mathbf{0}\})$, then for every continuous embedding $f : S^2 \rightarrow \mathbb{R}^3$ there is a point $\xi = (\xi_1, \xi_2, \xi_3, \xi_4) \in \Omega = X \setminus Y$ such that

$$\tau(\xi_1, \xi_2, \xi_3, \xi_4) = (\mathbf{0}, \mathbf{0}) \in U_4 \times U_2. \quad (3)$$

From (3) we conclude that

$$d_{12} = d_{23} = d_{34} = d_{14} = \frac{\Delta}{4} \quad \text{and} \quad d_{13} = d_{24} = \frac{\Phi}{2}. \quad (4)$$

It only remains to prove that all four points are different. Since $(\xi_1, \xi_2, \xi_3, \xi_4) \notin Y$ we have $\xi_1 \neq \xi_3$ or $\xi_2 \neq \xi_4$. By symmetry we may assume that $\xi_1 \neq \xi_3$. The map f is injective, therefore $f(\xi_1) \neq f(\xi_3)$ and consequently $d_{13} \neq 0$. Now

$$d_{13} \neq 0 \Rightarrow d_{24} \neq 0 \Rightarrow f(\xi_1) \neq f(\xi_3), f(\xi_2) \neq f(\xi_4) \Rightarrow \xi_1 \neq \xi_3, \xi_2 \neq \xi_4.$$

Let us assume, without loss of generality, that $\xi_1 = \xi_2$. Then $d_{12} = d_{23} = d_{34} = d_{14} = 0$, which implies that $d_{13} \leq d_{12} + d_{23} = 0$. This yield a contradiction to $d_{13} \neq 0$. Thus $\xi_1 \neq \xi_2$. $\square$

The unit sphere of the representation $U_4 \times U_2$ will be denoted by $S(U_4 \times U_2)$. Notice that there is a D_8 -equivariant deformation of $(U_4 \times U_2) \setminus (\{\mathbf{0}\} \times \{\mathbf{0}\})$ onto the sphere $S(U_4 \times U_2)$. Thus, there are D_8 -equivariant maps $(U_4 \times U_2) \setminus (\{\mathbf{0}\} \times \{\mathbf{0}\}) \rightarrow S(U_4 \times U_2)$ and $S(U_4 \times U_2) \rightarrow (U_4 \times U_2) \setminus (\{\mathbf{0}\} \times \{\mathbf{0}\})$. Hence by Proposition 2.1, Theorem 1.1 is a consequence of the following topological result.

Theorem 2.2. *There is no D_8 -equivariant map $\Omega \rightarrow S(U_4 \times U_2)$.*

Indeed, we will prove a stronger result: There is no $\mathbb{Z}_4$ -equivariant map $\Omega \rightarrow S(U_4 \times U_2)$.

3 Proof of Theorem 2.2

The proof is going to be conducted through a comparison of the Serre spectral sequences with $\mathbb{Z}$ -coefficients of the Borel constructions associated with the spaces Ω and $S(U_4 \times U_2)$ and the subgroup $\mathbb{Z}_4 = \langle \omega \rangle$ of D_8 . In other words, we determine the $\mathbb{Z}_4$ Fadell–Husseini index of these spaces living in $H^*(\mathbb{Z}_4; \mathbb{Z}) = \mathbb{Z}[U]/4U$, $\deg U = 2$.

The Fadell–Husseini index of a G -space X , $\text{Index}_{G, \mathbb{Z}} X$, is the kernel of the map

$$\pi_X^* : H^*(BG, \mathbb{Z}) \rightarrow H^*(X \times_G EG, \mathbb{Z})$$

induced by the projection $\pi_X : X \times_G EG \rightarrow BG$. Consider a G -equivariant map $f : X \rightarrow Y$ between two G -spaces. Then $\text{Index}_{G, \mathbb{Z}} X \supseteq \text{Index}_{G, \mathbb{Z}} Y$. Thus, the inclusion of indices of two G -spaces is a necessary condition for the existence of G -equivariant maps between these two spaces. If $E_*^{*,*}$ denotes the Serre spectral sequence of the Borel construction of X , then the homomorphism π_X^* can be presented as the composition

$$H^*(BG, \mathbb{Z}) \rightarrow E_2^{*,0} \rightarrow E_3^{*,0} \rightarrow E_4^{*,0} \rightarrow \dots \rightarrow E_\infty^{*,0} \subseteq H^*(X \times_G EG, \mathbb{Z}). \quad (5)$$

Since the E_2 -term of the spectral sequence is given by $E_2^{p,q} = H^p(BG, H^q(X, \mathbb{Z}))$ the first step in the computation of the index is study of the cohomology $H^*(X, \mathbb{Z})$ as a G -module (Section 3.2). The final step is explicit description of non-zero differentials in the spectral sequence and application of the presentation (5) of the homomorphism π_X^* (Section 3.3).

3.1 The Index of $S(U_4 \times U_2)$

Let V^1 be the 1-dimensional complex $\mathbb{Z}_4$ -representation, or 2-dimensional real $\mathbb{Z}_4$ -representation, induced by the correspondence $1 \mapsto e^{i\pi/2}$. Then the 3-dimensional real vector space $U_4 \subset \mathbb{R}^4$ seen as a real $\mathbb{Z}_4$ -representation decomposes into a sum of two irreducible real $\mathbb{Z}_4$ -representations

$$U_4 = \text{span}\{(1, 0, -1, 0), (0, 1, 0, -1)\} \oplus \text{span}\{(1, -1, 1, -1)\} \cong V^1 \oplus U_2.$$

Here “span” stands for all $\mathbb{R}$ -linear combinations of the given vectors. It can be also seen that there is an isomorphism of real $\mathbb{Z}_4$ -representations

$$U_4 \times U_2 \cong V^1 \oplus U_2 \oplus U_2 \cong V^1 \oplus (V^1 \otimes_{\mathbb{C}} V^1).$$

Here $V^1 \otimes_{\mathbb{C}} V^1$ is a tensor product of complex representations and therefore a 1-dimensional complex $\mathbb{Z}_4$ -representation or a 2-dimensional real $\mathbb{Z}_4$ -representation. Following [1, Section 8, p. 271 and Appendix, page 285] we deduce the total Chern class of the $\mathbb{Z}_4$ -representation $U_4 \times U_2$

$$c(U_4 \times U_2) = c(V^1) \cdot c(V^1 \otimes V^1).$$

Therefore the top Chern class, or the Euler class of the underlying real representation, is

$$c_2(U_4 \times U_2) = c_1(V^1) \cdot c_1(V^1 \otimes V^1) = c_1(V^1) \cdot (c_1(V^1) + c_1(V^1)) = 2U^2 \in H^*(\mathbb{Z}_4; \mathbb{Z}).$$

The $\mathbb{Z}_4$ -index of the sphere $S(U_4 \times U_2)$ is generated by the Euler class [2, Proposition 3.11], and so

$$\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2) = \langle 2U^2 \rangle. \quad (6)$$

3.2 The cohomology $H^*(\Omega; \mathbb{Z})$ as a $\mathbb{Z}_4$ -module

The cohomology is going to be determined via Poincaré–Lefschetz duality and an explicit study of cell structures for the spaces X and Y .

Poincaré–Lefschetz duality [4, Theorem 70.2, page 415] implies that

$$H^*(\Omega; \mathbb{Z}) = H^*(X \setminus Y; \mathbb{Z}) \cong H_{8-*}(X, Y; \mathbb{Z}) \quad (7)$$

and therefore we analyze the homology of the pair (X, Y) . The inclusion $Y \hookrightarrow X$ induces a map in homology. In particular, we consider this map in dimensions 2 and 4,

$$\Phi : H_2(Y; \mathbb{Z}) \rightarrow H_2(X; \mathbb{Z}) \quad \text{and} \quad \Psi : H_4(Y; \mathbb{Z}) \rightarrow H_4(X; \mathbb{Z}).$$

The long exact sequence in homology of the pair (X, Y) yields that the possibly non-zero homology groups of the pair (X, Y) with $\mathbb{Z}$ -coefficients are

$$H_i(X, Y; \mathbb{Z}) = \begin{cases} \mathbb{Z}[\mathbb{Z}_4]/\text{im}\Phi, & i = 2 \\ \ker \Phi, & i = 3 \\ \mathbb{Z}[\mathbb{Z}_4] \oplus \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2]/\text{im}\Psi, & i = 4 \\ \ker \Psi, & i = 5 \\ \mathbb{Z}[\mathbb{Z}_4], & i = 6 \\ \mathbb{Z}, & i = 8 \end{cases}$$

Thus explicit formulas for the maps Φ and Ψ are needed in order to determine the homology $H_*(X, Y; \mathbb{Z})$ and its exact $\mathbb{Z}_4$ -module structure.

Let $x_1, x_2, x_3, x_4 \in H_2(X; \mathbb{Z})$ be generators carried by individual copies of S^2 in the product $X = S^2 \times S^2 \times S^2 \times S^2$. The generator of the group $\mathbb{Z}_4 = \langle \omega \rangle$ acts on this basis of $H_2(X; \mathbb{Z})$ by $\omega \cdot x_i = x_{i+1}$ where $x_5 = x_1$. Then by $x_i x_j \in H_4(X; \mathbb{Z})$, $i \neq j$, we denote the generator carried by the product of i -th and j -th copy of S^2 in X . The action of ω on $H_4(X; \mathbb{Z})$ is described by

$$x_1 x_2 \xrightarrow{\omega} x_2 x_3 \xrightarrow{\omega} x_3 x_4 \xrightarrow{\omega} x_1 x_4 \quad \text{and} \quad x_1 x_3 \xrightarrow{\omega} x_2 x_4.$$

Let similarly $y_1, y_2 \in H_2(Y; \mathbb{Z})$ be generators carried by individual copies of S^2 in the product $Y = S^2 \times S^2$. Then $\omega \cdot y_1 = y_2$ and $\omega \cdot y_2 = y_1$. Again $y_1 y_2$ denotes the generator of $H_4(Y; \mathbb{Z})$ and $\omega \cdot y_1 y_2 = y_1 y_2$. Note that ω preserves the orientations of X and Y and therefore acts trivially on $H_8(X; \mathbb{Z})$ and on $H_4(Y; \mathbb{Z})$.

The inclusion $Y \subset X$ induces a map in homology $H_*(X; \mathbb{Z}) \subset H_*(Y; \mathbb{Z})$, which in dimensions 2 and 4 is given by

$$y_1 \longmapsto x_1 + x_3, \quad y_2 \longmapsto x_2 + x_4,$$

$$y_1 y_2 \longmapsto x_1 x_2 + x_2 x_3 + x_3 x_4 + x_1 x_4.$$

This can be seen from the dual cohomology picture: An element is mapped to a sum of generators intersecting its image, with appropriately attached intersection numbers.

Thus Φ and Ψ are injective and

$$\operatorname{im}\Phi = \langle x_1 + x_3, x_2 + x_4 \rangle, \quad \operatorname{im}\Psi = \langle x_1x_2 + x_2x_3 + x_3x_4 + x_1x_4 \rangle.$$

Let $N = \mathbb{Z} \oplus \mathbb{Z}$ be the $\mathbb{Z}_4$ -representation given by $\omega \cdot (a, b) = (b, -a)$, while M denotes the representation $\mathbb{Z}[\mathbb{Z}_4]/(1+\omega+\omega^2+\omega^3)\mathbb{Z}$. Then the non-trivial cohomology of the space $X \setminus Y$, as a $\mathbb{Z}_4$ -module via the isomorphism (7), is given by

$$H^i(\Omega; \mathbb{Z}) = \begin{cases} N, & i = 6 \\ M \oplus \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2], & i = 4 \\ \mathbb{Z}[\mathbb{Z}_4], & i = 2 \\ \mathbb{Z}, & i = 0 \end{cases} \quad (8)$$

3.3 The Serre spectral sequence of $\Omega \times_{\mathbb{Z}_4} \mathbb{E}\mathbb{Z}_4$

The Serre spectral sequence associated to the fibration $\Omega \rightarrow \Omega \times_{\mathbb{Z}_4} \mathbb{E}\mathbb{Z}_4 \rightarrow \mathbb{B}\mathbb{Z}_4$ is a spectral sequence with non-trivial local coefficients, since $\pi_1(\mathbb{B}\mathbb{Z}_4) = \mathbb{Z}_4$ acts non-trivially (8) on the cohomology $H^*(\Omega; \mathbb{Z})$. The first step in the study of such a spectral sequence is to understand the $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module structure on the rows of its E_2 -term.

The E_2 -term of the sequence is given by

$$E_2^{p,q} = \begin{cases} H^p(\mathbb{Z}_4, N), & q = 6 \\ H^p(\mathbb{Z}_4, M) \oplus H^p(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2]), & q = 4 \\ H^p(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]), & q = 2 \\ H^p(\mathbb{Z}_4; \mathbb{Z}), & q = 0 \\ 0, & \text{otherwise.} \end{cases} \quad (9)$$

Lemma 3.1. $H^p(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) = \begin{cases} \mathbb{Z}, & p = 0 \\ 0, & p > 0 \end{cases}$

and multiplication by $U \in H^2(\mathbb{Z}_4; \mathbb{Z})$ is trivial, $U \cdot H^p(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) = 0$.

For the proof one can consult [3, Example 2, page 58].

Lemma 3.2. $H^*(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2]) \cong H^*(\mathbb{Z}_2; \mathbb{Z})$, where the module structure is given by the restriction homomorphism $\operatorname{res}_{\mathbb{Z}_2}^{\mathbb{Z}_4} : H^*(\mathbb{Z}_4; \mathbb{Z}) \rightarrow H^*(\mathbb{Z}_2; \mathbb{Z})$.

In other words, if we denote $H^*(\mathbb{Z}_2; \mathbb{Z}) = \mathbb{Z}[T]/2T$, $\deg T = 2$, then $\operatorname{res}_{\mathbb{Z}_2}^{\mathbb{Z}_4}(U) = T$ and consequently:

- (A) $H^*(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2])$ is generated by one element of degree 0 as a $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module, and
- (B) multiplication by U in $H^*(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2])$ is an isomorphism, while multiplication by $2U$ is zero.

The proof is a direct application of Shapiro's lemma [3, (6.3), page 73] and a small part of the restriction diagram [2, Section 4.5.2].

Lemma 3.3. There exists an element $\Lambda \in H^*(\mathbb{Z}_4, M)$ of degree 1 such that $4\Lambda = 0$ and $H^*(\mathbb{Z}_4, M) \cong H^*(\mathbb{Z}_4; \mathbb{Z}) \cdot \Lambda$ as an $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module.

Proof. The short exact sequence of $\mathbb{Z}_4$ -modules

$$0 \longrightarrow \mathbb{Z} \xrightarrow{1+\omega+\omega^2+\omega^3} \mathbb{Z}[\mathbb{Z}_4] \longrightarrow M \longrightarrow 0$$

induces a long exact sequence in cohomology [3, Proposition 6.1, page 71], which is natural with respect to $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module multiplication. Since $\mathbb{Z}[\mathbb{Z}_4]$ is a free module we get enough zeros to recover the information we need:

$$\begin{array}{ccccccc} 0 \longrightarrow & H^0(\mathbb{Z}_4; \mathbb{Z}) & \xrightarrow{\xi} & H^0(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) & \longrightarrow & H^0(\mathbb{Z}_4, M) & \longrightarrow & H^1(\mathbb{Z}_4; \mathbb{Z}) & \longrightarrow \\ & \mathbb{Z} & & \mathbb{Z} & & & & 0 & \\ & & & & & & & & \\ & \longrightarrow & H^1(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) & \longrightarrow & H^1(\mathbb{Z}_4, M) & \longrightarrow & H^2(\mathbb{Z}_4; \mathbb{Z}) & \longrightarrow \\ & & 0 & & & & \mathbb{Z}_4 & \\ & & & & & & & \\ & \longrightarrow & H^2(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) & \longrightarrow & \dots & & & \\ & & 0 & & & & & \\ & & & & & & & \\ & \longrightarrow & H^i(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) & \longrightarrow & H^i(\mathbb{Z}_4, M) & \longrightarrow & H^{i+1}(\mathbb{Z}_4; \mathbb{Z}) & \longrightarrow \\ & & 0 & & & & & \\ & \longrightarrow & H^{i+1}(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) & \longrightarrow & \dots & & & \\ & & 0 & & & & & \end{array}$$

The map $\xi : H^0(\mathbb{Z}_4; \mathbb{Z}) \cong \mathbb{Z}^{\mathbb{Z}_4} \rightarrow H^0(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4]) \cong \mathbb{Z}[\mathbb{Z}_4]^{\mathbb{Z}_4}$ is a surjection. Indeed, ξ is induced by the map $\mathbb{Z} \xrightarrow[1-1 \text{ and onto}]{1+\omega+\omega^2+\omega^3} \mathbb{Z}[\mathbb{Z}_4]^{\mathbb{Z}_4} \hookrightarrow \mathbb{Z}[\mathbb{Z}_4]$ which bijectively factorizes through the invariants of $\mathbb{Z}[\mathbb{Z}_4]$. $\square$

Lemma 3.4. *There exists an element $\Upsilon \in H^*(\mathbb{Z}_4, N)$ of degree 1 such that $2\Upsilon = 0$ and $H^*(\mathbb{Z}_4, N) \cong H^*(\mathbb{Z}_4; \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2]) \cdot \Upsilon$ as an $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module.*

Proof. There is a short exact sequence of $\mathbb{Z}_4$ -modules

$$0 \rightarrow N \xrightarrow{\alpha} \mathbb{Z}[\mathbb{Z}_4] \xrightarrow{\beta} L \rightarrow 0$$

where $L = \mathbb{Z}[\mathbb{Z}_4]/N$ and $\alpha(p, q) = (p, q, -p, -q)$. The map α is well defined because the following diagram commutes

$$\begin{array}{ccccc} N =_{ab} \mathbb{Z} \oplus \mathbb{Z} \ni & (p, q) & \xrightarrow{\alpha} & (p, q, -p, -q) & \in \mathbb{Z}[\mathbb{Z}_4] \\ & \downarrow \cdot \omega & & \downarrow \cdot \omega & \\ N =_{ab} \mathbb{Z} \oplus \mathbb{Z} \ni & (q, -p) & \xrightarrow{\alpha} & (q, -p, -q, p) & \in \mathbb{Z}[\mathbb{Z}_4]. \end{array}$$

The $\mathbb{Z}_4$ -module L is isomorphism to $\mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2] \cong \mathbb{Z} \oplus \mathbb{Z}$ and the map β is given, on generators, by

$$(1, 0, 0, 0) \mapsto (1, 0), (0, 1, 0, 0) \mapsto (0, 1), (0, 0, 1, 0) \mapsto (1, 0), (0, 0, 0, 1) \mapsto (1, 0).$$

Therefore, the induced map of invariants $\mathbb{Z} \cong \mathbb{Z}[\mathbb{Z}_4]^{\mathbb{Z}_4} \xrightarrow{\beta} \mathbb{Z}[\mathbb{Z}_4/\mathbb{Z}_2] \cong \mathbb{Z}$ is a multiplication by 2. Now, the long exact sequence in group cohomology [3, Prop. 6.1, p 71] implies the result. $\square$

6	0		$\Upsilon_{\mathbb{Z}_2}$	0		$T\Upsilon_{\mathbb{Z}_2}$		0		$T^2\Upsilon_{\mathbb{Z}_2}$		0		$T^3\Upsilon_{\mathbb{Z}_2}$																			
5	0		0		0		0		0		0		0		0																		
4	$I_{\mathbb{Z}_2}$		0		0		$\Lambda_{\mathbb{Z}_4}$		$T_{\mathbb{Z}_2}$		0		0		$\Lambda U_{\mathbb{Z}_4}$		$T^2_{\mathbb{Z}_2}$		0		0		$\Lambda U^2_{\mathbb{Z}_4}$		$T^3_{\mathbb{Z}_2}$		0		0		$\Lambda U^3_{\mathbb{Z}_4}$		
3	0		0		0		0		0		0		0		0		0		0														
2	$\mathbb{Z}$		0		0		0		0		0		0		0		0		0														
1	0		0		0		0		0		0		0		0		0		0														
0	$1_{\mathbb{Z}}$		0		$U_{\mathbb{Z}_4}$		0		$U^2_{\mathbb{Z}_4}$		0		$U^3_{\mathbb{Z}_4}$		0																		
		0		1		2		3		4		5		6		7																	

Figure 2: The E_2 -term

The E_2 -term of the Borel construction $(X \setminus Y) \times_{\mathbb{Z}_4} E\mathbb{Z}_4$, with the $H^*(\mathbb{Z}_4; \mathbb{Z})$ -module structure, is presented in Figure 2.

The differentials of the spectral sequence are retrieved from the fact that the $\mathbb{Z}_4$ action on Ω is free. Therefore $H_{\mathbb{Z}_4}^i(\Omega; \mathbb{Z}) = 0$ for all $i > 8$. Since the spectral sequence is converging to the graded group associated with $H_{\mathbb{Z}_4}^i(\Omega; \mathbb{Z})$ this means that for $p + q > 8$ nothing survives. Thus the only non-zero second differentials are $d_2 : E_2^{2i+1,6} \rightarrow E_2^{2i+4,4}$, $d_2(T^i\Upsilon) = T^{i+1}$, $i > 0$, as displayed in Figure 3.

The diagram illustrates a sequence of transformations on a grid of mathematical symbols. The grid is 8 rows by 8 columns, with rows indexed 0 to 7 and columns indexed 0 to 7. The symbols are arranged in a pattern that suggests a sequence of operations, with some symbols circled in red and arrows indicating a path through them.

Left Grid (Annotated):

- Row 6: 0 , $\Upsilon_{\mathbb{Z}_2}$ (circled), 0 , $T\Upsilon_{\mathbb{Z}_2}$ (circled), 0 , $T^2\Upsilon_{\mathbb{Z}_2}$ (circled), 0 , $T^3\Upsilon_{\mathbb{Z}_2}$ (circled)
- Row 5: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 4: $I_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda_{\mathbb{Z}_4}$, $T_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda U_{\mathbb{Z}_4}$, $T^2_{\mathbb{Z}_2}$ (circled), 0 , 0 , $\Lambda U^2_{\mathbb{Z}_4}$, $T^3_{\mathbb{Z}_2}$ (circled), 0 , 0 , $\Lambda U^3_{\mathbb{Z}_4}$
- Row 3: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 2: $\mathbb{Z}$, 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 1: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 0: $I_{\mathbb{Z}}$, 0 , $U_{\mathbb{Z}_4}$, 0 , $U^2_{\mathbb{Z}_4}$, 0 , $U^3_{\mathbb{Z}_4}$, 0

Right Grid (Unannotated):

- Row 6: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 5: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 4: $I_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda_{\mathbb{Z}_4}$, $T_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda U_{\mathbb{Z}_4}$, $T^2_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda U^2_{\mathbb{Z}_4}$, $T^3_{\mathbb{Z}_2}$, 0 , 0 , $\Lambda U^3_{\mathbb{Z}_4}$
- Row 3: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 2: $\mathbb{Z}$, 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 1: 0 , 0 , 0 , 0 , 0 , 0 , 0 , 0
- Row 0: $I_{\mathbb{Z}}$, 0 , $U_{\mathbb{Z}_4}$, 0 , $U^2_{\mathbb{Z}_4}$, 0 , $U^3_{\mathbb{Z}_4}$, 0

Figure 3: Differentials in E_2 and E_3 -terms

The last remaining non-zero differentials are $d_4 : E_4^{2i+1,4} \rightarrow E_4^{2i+6,0}$, $d_6(U^i\Lambda) = U^{i+3}$, $i > 0$. Then $E_5 = E_\infty$, cf. Figure 4.

3.4 The index of Ω

The conclusion $d_6(\Lambda) = U^3$ implies that

$$\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} \Omega = \langle U^3 \rangle.$$

Since the generator $2U^2$ of the $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2)$ is not contained in $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} \Omega$ it follows that **there is no $\mathbb{Z}_4$ -equivariant map $\Omega \rightarrow S(U_4 \times U_2)$** . This concludes the proof of Theorem 2.2.




Figure 4: Differentials in E_4 and E_5 -terms

Remark 3.5. As one of the referees observed, in order to prove Theorem 2.2 there was no need to compute the $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} \Omega$. The structure of E_2 -term (9) of the spectral sequence of the fibration $\Omega \times_{\mathbb{Z}_4} \mathbb{E}\mathbb{Z}_4$ and Lemma 3.1 guarantee that the elements U^2 and $2U^2$ survive to E_∞ -term. This provides the contradiction since $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2) = \langle 2U^2 \rangle$.

4 Concluding remarks

4.1 The $\mathbb{F}_2$ -index

Let $H^*(\mathbb{Z}_4, \mathbb{F}_2) = \mathbb{F}_2[e, u]/e^2$, $\deg(e) = 1$, $\deg(u) = 2$. The homomorphism of coefficients $j : \mathbb{Z} \rightarrow \mathbb{F}_2$, $j(1) = 1$, induces a homomorphism in group cohomology $j^* : H^*(\mathbb{Z}_4; \mathbb{Z}) \rightarrow H^*(\mathbb{Z}_4, \mathbb{F}_2)$ given by $j^*(U) = u$ (compare [2, Section 4.5.2]).

The $\mathbb{F}_2$ -index of the configuration space Ω is

$$\text{Index}_{\mathbb{Z}_4, \mathbb{F}_2} \Omega = \langle eu^2, u^3 \rangle.$$

This can be obtained in a similar fashion as we obtained the index with $\mathbb{Z}$ -coefficients in Section 3.3. The relevant E_2 -term of the Serre spectral sequence of the fibration $\Omega \rightarrow \Omega \times_{\mathbb{Z}_4} \mathbb{E}\mathbb{Z}_4 \rightarrow \mathbb{B}\mathbb{Z}_4$ is described in Figure 5.

The $\mathbb{F}_2$ -index of the sphere $S(U_4 \times U_2)$ is generated by the j^* image of the generator $2U^2$ of the index with $\mathbb{Z}$ -coefficients $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2)$. Since $j^*(2U^2) = 0$ the index $\text{Index}_{\mathbb{Z}_4, \mathbb{F}_2} S(U_4 \times U_2)$ is trivial. Therefore, for our problem **no conclusion can be obtained from the study of the $\mathbb{F}_2$ -index**. The same observation holds even when the complete group D_8 is used. The $\mathbb{F}_2$ -index of the sphere $S(U_4 \times U_2)$ would be generated by $xyw = 0 \in H^*(D_8; \mathbb{F}_2)$, in the notation of [2].

4.2 The square peg problem

The method of configuration spaces can also be set up for to the continuous square peg problem. Following the ideas presented in Section 2, taking for X the product $S^1 \times S^1 \times S^1 \times S^1$, for Y the subspace $Y = \{(x, y, x, y) \mid x, y \in S^1\}$ and for the configuration space

6	1	t	t^2	t^3	t^4	t^5	t^6	t^7									
5	0	0	0	0	0	0	0	0									
4	1	e	t	u	t^2	eu	t^3	u^2	t^4	eu^2	t^5	u^3	t^6	eu^3	t^7	u^4	
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
2	$\mathbf{F}_2$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
0	1	e	u	eu	u^2	eu^2	u^3	eu^3									
	0	1	2	3	4	5	6	7									

Figure 5: E_2 -term with $\mathbb{F}_2$ -coefficients

$\Omega = X \setminus Y$, the square peg problem can be related to the question of the existence of a D_8 -equivariant map $\Omega \rightarrow S(U_4 \times U_2)$. The Fadell–Husseini indexes can be retrieved:

$$\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} \Omega = \langle U^2 \rangle \quad \text{and} \quad \text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2) = \langle 2U^2 \rangle,$$

but since $\text{Index}_{\mathbb{Z}_4, \mathbb{Z}} \Omega \supseteq \text{Index}_{\mathbb{Z}_4, \mathbb{Z}} S(U_4 \times U_2)$ the result does not yield any conclusion. The same can be done for the complete symmetry group D_8 , explicitly $\text{Index}_{D_8, \mathbb{Z}} S(U_4 \times U_2) = \langle 2\mathcal{W} \rangle$ and $\mathcal{W} \in \text{Index}_{D_8, \mathbb{Z}} \Omega$.

Acknowledgements.

Thanks to Anton Dochterman and to the referees for many interesting and useful comments.

References

- [1] M. F. ATIYAH, *Characters and cohomology of finite groups*, Inst. Hautes Études Sci. Publ. Math. No. 9 (1961), 23–64.
- [2] P. V. M. BLAGOJEVIĆ, G. M. ZIEGLER, *The ideal-valued index for a dihedral group action, and mass partition by two hyperplanes*, preprint, revised version [arXiv:0704.1943v2](https://arxiv.org/abs/0704.1943v2), July 2008, 42 pages.
- [3] K. S. BROWN, *Cohomology of Groups*, Graduate Texts in Math. 87, Springer-Verlag, New York, Berlin, 1982.
- [4] J. R. MUNKRES, *Elements of Algebraic Topology*, Addison-Wesley, Menlo Park CA, 1984.
- [5] L. G. SHNIRELMAN, *On certain geometrical properties of closed curves (in Russian)*, Uspehi Matem. Nauk 10 (1944), 34–44, <http://tinyurl.com/28gsy3>.
- [6] Y. SOIBELMAN, *Topological Borsuk problem*, Preprint 2002, 4 pages, <http://arxiv.org/abs/math/0208221v2>.