

# Invariant and coinvariant spaces for the algebra of symmetric polynomials in non-commuting variables

François Bergeron\*

LaCIM  
Université du Québec à Montréal  
Montréal (Québec) H3C 3P8, CANADA  
bergeron.francois@uqam.ca

Aaron Lauve

Department of Mathematics  
Texas A&M University  
College Station, TX 77843, USA  
lauve@math.tamu.edu

Submitted: Oct 2, 2009; Accepted: Nov 26, 2010; Published: Dec 10, 2010

Mathematics Subject Classification: 05E05

## Abstract

We analyze the structure of the algebra  $\mathbb{K}\langle \mathbf{x} \rangle^{\mathfrak{S}_n}$  of symmetric polynomials in non-commuting variables in so far as it relates to  $\mathbb{K}[\mathbf{x}]^{\mathfrak{S}_n}$ , its commutative counterpart. Using the “place-action” of the symmetric group, we are able to realize the latter as the invariant polynomials inside the former. We discover a tensor product decomposition of  $\mathbb{K}\langle \mathbf{x} \rangle^{\mathfrak{S}_n}$  analogous to the classical theorems of Chevalley, Shephard-Todd on finite reflection groups.

**Résumé.** Nous analysons la structure de l’algèbre  $\mathbb{K}\langle \mathbf{x} \rangle^{\mathfrak{S}_n}$  des polynômes symétriques en des variables non-commutatives pour obtenir des analogues des résultats classiques concernant la structure de l’anneau  $\mathbb{K}[\mathbf{x}]^{\mathfrak{S}_n}$  des polynômes symétriques en des variables commutatives. Plus précisément, au moyen de “l’action par positions”, on réalise  $\mathbb{K}[\mathbf{x}]^{\mathfrak{S}_n}$  comme sous-module de  $\mathbb{K}\langle \mathbf{x} \rangle^{\mathfrak{S}_n}$ . On découvre alors une nouvelle décomposition de  $\mathbb{K}\langle \mathbf{x} \rangle^{\mathfrak{S}_n}$  comme produit tensoriel, obtenant ainsi un analogues des théorèmes classiques de Chevalley et Shephard-Todd.

## 1 Introduction

One of the more striking results of invariant theory is certainly the following: if  $W$  is a finite group of  $n \times n$  matrices (over some field  $\mathbb{K}$  containing  $\mathbb{Q}$ ), then there is a  $W$ -module decomposition of the polynomial ring  $S = \mathbb{K}[\mathbf{x}]$ , in variables  $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ , as a tensor product

$$S \simeq S_W \otimes S^W \tag{1}$$

---

\*F. Bergeron is supported by NSERC-Canada and FQRNT-Québec.

if and only if  $W$  is a group generated by (pseudo) reflections. As usual,  $S$  is afforded a natural  $W$ -module structure by considering it as the symmetric space on the defining vector space  $X^*$  for  $W$ , e.g.,  $w \cdot f(\mathbf{x}) = f(\mathbf{x} \cdot w)$ . It is customary to denote by  $S^W$  the ring of  $W$ -invariant polynomials for this action. To finish parsing (1), recall that  $S_W$  stands for the **coinvariant space**, i.e., the  $W$ -module

$$S_W := S / \langle S_+^W \rangle \tag{2}$$

defined as the quotient of  $S$  by the ideal generated by constant-term free  $W$ -invariant polynomials. We give  $S$  an  $\mathbb{N}$ -grading by degree in the variables  $\mathbf{x}$ . Since the  $W$ -action on  $S$  preserves degrees, both  $S^W$  and  $S_W$  inherit a grading from the one on  $S$ , and (1) is an isomorphism of graded  $W$ -modules. One of the motivations behind the quotient in (2) is to eliminate trivially redundant copies of irreducible  $W$ -modules inside  $S$ . Indeed, if  $\mathcal{V}$  is such a module and  $f$  is any  $W$ -invariant polynomial with no constant term, then  $\mathcal{V}f$  is an isomorphic copy of  $\mathcal{V}$  living within  $\langle S_+^W \rangle$ . Thus, the coinvariant space  $S_W$  is the more interesting part of the story.

The context for the present paper is the algebra  $T = \mathbb{K}\langle \mathbf{x} \rangle$  of noncommutative polynomials, with  $W$ -module structure on  $T$  obtained by considering it as the tensor space on the defining space  $X^*$  for  $W$ . In the special case when  $W$  is the symmetric group  $\mathfrak{S}_n$ , we elucidate a relationship between the space  $S^W$  and the subalgebra  $T^W$  of  $W$ -invariants in  $T$ . The subalgebra  $T^W$  was first studied in [4, 20] with the aim of obtaining noncommutative analogs of classical results concerning symmetric function theory. Recent work in [2, 15] has extended a large part of the story surrounding (1) to this noncommutative context. In particular, there is an explicit  $\mathfrak{S}_n$ -module decomposition of the form  $T \simeq T_{\mathfrak{S}_n} \otimes T^{\mathfrak{S}_n}$  [2, Theorem 8.7]. See [7] for a survey of other results in noncommutative invariant theory.

By contrast, our work proceeds in a somewhat complementary direction. We consider  $\mathcal{N} = T^{\mathfrak{S}_n}$  as a tower of  $\mathfrak{S}_d$ -modules under the “place-action” and realize  $S^{\mathfrak{S}_n}$  inside  $\mathcal{N}$  as a subspace  $\Lambda$  of invariants for this action. This leads to a decomposition of  $\mathcal{N}$  analogous to (1). More explicitly, our main result is as follows.

**Theorem 1.** *There is an explicitly constructed subspace  $\mathcal{C}$  of  $\mathcal{N}$  so that  $\mathcal{C}$  and the place-action invariants  $\Lambda$  exhibit a graded vector space isomorphism*

$$\mathcal{N} \simeq \mathcal{C} \otimes \Lambda. \tag{3}$$

An analogous result holds in the case  $|\mathbf{x}| = \infty$ . An immediate corollary in either case is the Hilbert series formula

$$\text{Hilb}_t(\mathcal{C}) = \text{Hilb}_t(\mathcal{N}) \prod_{i=1}^{|\mathbf{x}|} (1 - t^i). \tag{4}$$

Here, the **Hilbert series** of a graded space  $\mathcal{V} = \bigoplus_{d \geq 0} \mathcal{V}_d$  is the formal power series defined as

$$\text{Hilb}_t(\mathcal{V}) = \sum_{d \geq 0} \dim \mathcal{V}_d t^d,$$

where  $\mathcal{V}_d$  is the **homogeneous degree  $d$  component** of  $\mathcal{V}$ . The fact that (4) expands as a series in  $\mathbb{N}[[t]]$  is not at all obvious, as one may check that the Hilbert series of  $\mathcal{N}$  is

$$\text{Hilb}_t(\mathcal{N}) = 1 + \sum_{k=1}^{|\mathbf{x}|} \frac{t^k}{(1-t)(1-2t)\cdots(1-kt)}. \quad (5)$$

In Sections 2 and 3, we recall the relevant structural features of  $S$  and  $T$ . Section 4 describes the place-action structure of  $T$  and the original motivation for our work. Our main results are proven in Sections 5 and 6. We underline that the harder part of our work lies in working out the case  $|\mathbf{x}| < \infty$ . This is accomplished in Section 6. If we restrict ourselves to the case  $|\mathbf{x}| = \infty$ , both  $\mathcal{N}$  and  $\Lambda$  become Hopf algebras and our results are then consequences of a general theorem of Blattner, Cohen and Montgomery. As we will see in Section 5, stronger results hold in this simpler context. For example, (4) may be refined to a statement about “shape” enumeration.

## 2 The algebra $S^\mathfrak{S}$ of symmetric functions

### 2.1 Vector space structure of $S^\mathfrak{S}$

We specialize our introductory discussion to the group  $W = \mathfrak{S}_n$  of permutation matrices (writing  $|\mathbf{x}| = n$ ). The action on  $S = \mathbb{K}[\mathbf{x}]$  is simply the **permutation action**  $\sigma \cdot x_i = x_{\sigma(i)}$  and  $S^{\mathfrak{S}_n}$  comprises the familiar symmetric polynomials. We suppress  $n$  in the notation and denote the subring of symmetric polynomials by  $S^\mathfrak{S}$ . (Note that upon sending  $n$  to  $\infty$ , the elements of  $S^\mathfrak{S}$  become formal series in  $\mathbb{K}[[\mathbf{x}]]$  of bounded degree; we call both finite and infinite versions “functions” in what follows to affect a uniform discussion.) A monomial in  $S$  of degree  $d$  may be written as follows: given an  $r$ -subset  $\mathbf{y} = \{y_1, y_2, \dots, y_r\}$  of  $\mathbf{x}$  and a **composition** of  $d$  into  $r$  parts,  $\mathbf{a} = (a_1, a_2, \dots, a_r)$  ( $a_i > 0$ ), we write  $\mathbf{y}^\mathbf{a}$  for  $y_1^{a_1} y_2^{a_2} \cdots y_r^{a_r}$ . We assume that the variables  $y_i$  are naturally ordered, so that whenever  $y_i = x_j$  and  $y_{i+1} = x_k$  we have  $j < k$ . Reordering the entries of a composition  $\mathbf{a}$  in decreasing order results in a partition  $\lambda(\mathbf{a})$  called the **shape** of  $\mathbf{a}$ . Summing over monomials  $\mathbf{y}^\mathbf{a}$  with the same shape leads to the monomial symmetric function

$$m_\mu = m_\mu(\mathbf{x}) := \sum_{\lambda(\mathbf{a})=\mu, \mathbf{y} \subseteq \mathbf{x}} \mathbf{y}^\mathbf{a}.$$

Letting  $\mu = (\mu_1, \mu_2, \dots, \mu_r)$  run over all partitions of  $d = |\mu| = \mu_1 + \mu_2 + \cdots + \mu_r$  gives a basis for  $S_d^\mathfrak{S}$ . As usual, we set  $m_0 := 1$  and agree that  $m_\mu = 0$  if  $\mu$  has too many parts (i.e.,  $n < r$ ).

### 2.2 Dimension enumeration

A fundamental result in the invariant theory of  $\mathfrak{S}_n$  is that  $S^\mathfrak{S}$  is generated by a family  $\{f_k\}_{1 \leq k \leq n}$  of algebraically independent symmetric functions, having respective degrees

$\deg f_k = k$ . (One may choose  $\{m_k\}_{1 \leq k \leq n}$  for such a family.) It follows that the Hilbert series of  $S^\mathfrak{S}$  is

$$\text{Hilb}_t(S^\mathfrak{S}) = \prod_{i=1}^n \frac{1}{1-t^i}. \quad (6)$$

Recalling that the Hilbert series of  $S$  is  $(1-t)^{-n}$ , we see from (1) and (6) that the Hilbert series for the coinvariant space  $S_\mathfrak{S}$  is the well-known  $t$ -analog of  $n!$ :

$$\prod_{i=1}^n \frac{1-t^i}{1-t} = \prod_{i=1}^n (1+t+\cdots+t^{i-1}). \quad (7)$$

In particular, contrary to the situation in (4), the series  $\text{Hilb}_t(S)/\text{Hilb}_t(S^\mathfrak{S})$  in  $\mathbb{Q}[[t]]$  *obviously* belongs to  $\mathbb{N}[[t]]$ .

### 2.3 Algebra and coalgebra structures of $S^\mathfrak{S}$

Given partitions  $\mu$  and  $\nu$ , there is an explicit multiplication rule for computing the product  $m_\mu \cdot m_\nu$ . In lieu of giving the formula, see [2, §4.1], we simply give an example

$$m_{21} \cdot m_{11} = 3m_{2111} + 2m_{221} + 2m_{311} + m_{32} \quad (8)$$

and highlight two features relevant to the coming discussion.

First, we note that if  $n < 4$ , then the first term is equal to zero. However, if  $n$  is sufficiently large then analogs of this term always appear with positive integer coefficients. If  $\mu = (\mu_1, \mu_2, \dots, \mu_r)$  and  $\nu = (\nu_1, \nu_2, \dots, \nu_s)$  with  $r \leq s$ , then the partition indexing the left-most term in  $m_\mu m_\nu$  is denoted by  $\mu \cup \nu$  and is given by sorting the list  $(\mu_1, \dots, \mu_r, \nu_1, \dots, \nu_s)$  in increasing order; the right-most term is indexed by  $\mu + \nu := (\mu_1 + \nu_1, \dots, \mu_r + \nu_r, \nu_{r+1}, \dots, \nu_s)$ . Taking  $\mu = 31$  and  $\nu = 221$ , we would have  $\mu \cup \nu = 32211$  and  $\mu + \nu = 531$ .

Second, we point out that the leftmost term (indexed by  $\mu \cup \nu$ ) is indeed a *leading term* in the following sense. An important partial order on partitions takes

$$\lambda \leq \mu \quad \text{iff} \quad \sum_{i=1}^k \lambda_i \leq \sum_{i=1}^k \mu_i \quad \text{for all } k.$$

With this ordering,  $\mu \cup \nu$  is the least partition occurring with nonzero coefficient in the product of  $m_\mu m_\nu$ . That is,  $S^\mathfrak{S}$  is **shape-filtered**:  $(S^\mathfrak{S})_\lambda \cdot (S^\mathfrak{S})_\mu \subseteq \bigoplus_{\nu \geq \lambda \cup \mu} (S^\mathfrak{S})_\nu$ . Here  $(S^\mathfrak{S})_\lambda$  denotes the subspace of  $S^\mathfrak{S}$  indexed by partitions of shape  $\lambda$  (the linear span of  $m_\lambda$ ), which we point out in preparation for the noncommutative analog.

The ring  $S^\mathfrak{S}$  is afforded a coalgebra structure with counit  $\varepsilon : S^\mathfrak{S} \rightarrow \mathbb{K}$  and coproduct  $\Delta : S_d^\mathfrak{S} \rightarrow \bigoplus_{k=0}^d S_k^\mathfrak{S} \otimes S_{d-k}^\mathfrak{S}$  given, respectively, by

$$\varepsilon(m_\mu) = \delta_{\mu,0} \quad \text{and} \quad \Delta(m_\nu) = \sum_{\lambda \cup \mu = \nu} m_\lambda \otimes m_\mu.$$

If  $|\mathbf{x}| = \infty$ ,  $\Delta$  and  $\varepsilon$  are algebra maps, making  $S^\mathfrak{S}$  a graded connected Hopf algebra.

### 3 The algebra $\mathcal{N}$ of noncommutative symmetric functions

#### 3.1 Vector space structure of $\mathcal{N}$

Suppose now that  $\mathbf{x}$  denotes a set of non-commuting variables. The algebra  $T = \mathbb{K}\langle \mathbf{x} \rangle$  of noncommutative polynomials is graded by degree. A degree  $d$  **noncommutative monomial**  $\mathbf{z} \in T_d$  is simply a length  $d$  “word”:

$$\mathbf{z} = z_1 z_2 \cdots z_d, \quad \text{with each } z_i \in \mathbf{x}.$$

In other terms,  $\mathbf{z}$  is a function  $\mathbf{z} : [d] \rightarrow \mathbf{x}$ , with  $[d]$  denoting the set  $\{1, 2, \dots, d\}$ . The permutation-action on  $\mathbf{x}$  clearly extends to  $T$ , giving rise to the subspace  $\mathcal{N} = T^{\mathfrak{S}}$  of noncommutative  $\mathfrak{S}$ -invariants. With the aim of describing a linear basis for the homogeneous component  $\mathcal{N}_d$ , we next introduce set partitions of  $[d]$  and the type of a monomial  $\mathbf{z} : [d] \rightarrow \mathbf{x}$ . Let  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$  be a set of subsets of  $[d]$ . Say  $\mathbf{A}$  is a **set partition** of  $[d]$ , written  $\mathbf{A} \vdash [d]$ , iff  $A_1 \cup A_2 \cup \dots \cup A_r = [d]$ ,  $A_i \neq \emptyset$  ( $\forall i$ ), and  $A_i \cap A_j = \emptyset$  ( $\forall i \neq j$ ). The **type**  $\tau(\mathbf{z})$  of a degree  $d$  monomial  $\mathbf{z} : [d] \rightarrow \mathbf{x}$  is the set partition

$$\tau(\mathbf{z}) := \{\mathbf{z}^{-1}(x) : x \in \mathbf{x}\} \setminus \{\emptyset\} \quad \text{of } [d],$$

whose parts are the non-empty fibers of the function  $\mathbf{z}$ . For instance,

$$\tau(x_1 x_8 x_1 x_5 x_8) = \{\{1, 3\}, \{2, 5\}, \{4\}\}.$$

Note that the type of a monomial is a set partition with at most  $n$  parts. In what follows, we lighten the heavy notation for set partitions, writing, e.g., the set partition  $\{\{1, 3\}, \{2, 5\}, \{4\}\}$  as 13.25.4. We also always order the parts in increasing order of their minimum elements. The **shape**  $\lambda(\mathbf{A})$  of a set partition  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$  is the (integer) partition  $\lambda(|A_1|, |A_2|, \dots, |A_r|)$  obtained by sorting the part sizes of  $\mathbf{A}$  in increasing order, and its **length**  $\ell(\mathbf{A})$  is its number of parts ( $r$ ). Observing that the permutation-action is *type preserving*, we are led to index the **monomial** linear basis for the space  $\mathcal{N}_d$  by set partitions:

$$m_{\mathbf{A}} = m_{\mathbf{A}}(\mathbf{x}) := \sum_{\tau(\mathbf{z})=\mathbf{A}, \mathbf{z} \in \mathbf{x}^{[d]}} \mathbf{z}$$

For example, with  $n = 2$ , we have  $m_1 = x_1 + x_2$ ,  $m_{12} = x_1^2 + x_2^2$ ,  $m_{1.2} = x_1 x_2 + x_2 x_1$ ,  $m_{123} = x_1^3 + x_2^3$ ,  $m_{12.3} = x_1^2 x_2 + x_2^2 x_1$ ,  $m_{13.2} = x_1 x_2 x_1 + x_2 x_1 x_2$ ,  $m_{1.2.3} = 0$ , and so on. (We set  $m_{\emptyset} := 1$ , taking  $\emptyset$  as the unique set partition of the empty set, and we agree that  $m_{\mathbf{A}} = 0$  if  $\mathbf{A}$  is a set partition with more than  $n$  parts.)

#### 3.2 Dimension enumeration and shape grading

Above, we determined that  $\dim \mathcal{N}_d$  is the number of set partitions of  $d$  into at most  $n$  parts. These are counted by the (length restricted) **Bell numbers**  $B_d^{(n)}$ . Consequently,

(5) follows from the fact that its right-hand side is the ordinary generating function for length restricted Bell numbers. See [10, §2]. We next highlight a finer enumeration, where we grade  $\mathcal{N}$  by shape rather than degree.

For each partition  $\mu$ , we may consider the subspace  $\mathcal{N}_\mu$  spanned by those  $m_{\mathbf{A}}$  for which  $\lambda(\mathbf{A}) = \mu$ . This results in a direct sum decomposition  $\mathcal{N}_d = \bigoplus_{\mu \vdash d} \mathcal{N}_\mu$ . A simple dimension description for  $\mathcal{N}_d$  takes the form of a **shape Hilbert series** in the following manner. View commuting variables  $q_i$  as marking parts of size  $i$  and set  $\mathbf{q}_\mu := q_{\mu_1} q_{\mu_2} \cdots q_{\mu_r}$ . Then

$$\text{Hilb}_{\mathbf{q}}(\mathcal{N}_d) = \sum_{\mu \vdash d} \dim \mathcal{N}_\mu \mathbf{q}_\mu, = \sum_{\mathbf{A} \vdash [d]} q_{\lambda(\mathbf{A})}. \quad (9)$$

Here,  $\mathbf{q}_\mu$  is a marker for set partitions of shape  $\lambda(\mathbf{A}) = \mu$  and the sum is over all partitions into at most  $n$  parts. Such a shape grading also makes sense for  $S_d^{\mathfrak{S}}$ . Summing over all  $d \geq 0$  and all  $\mu$ , we get

$$\text{Hilb}_{\mathbf{q}}(S^{\mathfrak{S}}) = \sum_{\mu} \mathbf{q}_\mu = \prod_{i \geq 1} \frac{1}{1 - q_i}. \quad (10)$$

Using classical combinatorial arguments, one finds the enumerator polynomials  $\text{Hilb}_{\mathbf{q}}(\mathcal{N}_d)$  are naturally collected in the **exponential generating function**

$$\sum_{d=0}^{\infty} \text{Hilb}_{\mathbf{q}}(\mathcal{N}_d) \frac{t^d}{d!} = \sum_{m=0}^n \frac{1}{m!} \left( \sum_{k=1}^{\infty} q_k \frac{t^k}{k!} \right)^m. \quad (11)$$

See [1, Chap. 2.3], Example 13(a). For instance, with  $n = 3$ , we have

$$\text{Hilb}_{\mathbf{q}}(\mathcal{N}_6) = q_6 + 6 q_5 q_1 + 15 q_4 q_2 + 15 q_4 q_1^2 + 10 q_3^2 + 60 q_3 q_2 q_1 + 15 q_2^3,$$

thus  $\dim \mathcal{N}_{222} = 15$  when  $n \geq 3$ . Evidently, the  $\mathbf{q}$ -polynomials  $\text{Hilb}_{\mathbf{q}}(\mathcal{N}_d)$  specialize to the length restricted Bell numbers  $B_d^{(n)}$  when we set all  $q_k$  equal to 1.

In view of (10), (11), and Theorem 1, we claim the following refinement of (4).

**Corollary 2.** *Sending  $n$  to  $\infty$ , the shape Hilbert series of the space  $\mathcal{C}$  is given by*

$$\text{Hilb}_{\mathbf{q}}(\mathcal{C}) = \sum_{d \geq 0} d! \exp \left( \sum_{k=1}^{\infty} q_k \frac{t^k}{k!} \right) \Big|_{t^d} \prod_{i \geq 1} (1 - q_i), \quad (12)$$

with  $(-)|_{t^d}$  standing for the operation of taking the coefficient of  $t^d$ .

This refinement of (4) will follow immediately from the isomorphism  $\mathcal{C} \otimes \Lambda \rightarrow \mathcal{N}$  in Section 5, which is shape-preserving in an appropriate sense. Thus we have the expansion

$$\begin{aligned} \text{Hilb}_{\mathbf{q}}(\mathcal{C}) &= 1 + 2 q_2 q_1 + (3 q_3 q_1 + 2 q_2^2 + 3 q_2 q_1^2) \\ &\quad + (4 q_4 q_1 + 9 q_3 q_2 + 6 q_3 q_1^2 + 10 q_2^2 q_1 + 4 q_2 q_1^3) + \cdots \end{aligned}$$

### 3.3 Algebra and coalgebra structures of $\mathcal{N}$

Since the action of  $\mathfrak{S}$  on  $T$  is multiplicative, it is straightforward to see that  $\mathcal{N}$  is a subalgebra of  $T$ . The *multiplication rule* in  $\mathcal{N}$ , expressing a product  $m_{\mathbf{A}} \cdot m_{\mathbf{B}}$  as a sum of basis vectors  $\sum_{\mathbf{C}} m_{\mathbf{C}}$ , is easy to describe. Since we make heavy use of the rule later, we develop it carefully here. We begin with an example (digits corresponding to  $\mathbf{B} = \mathbf{1.2}$  appear in bold):

$$\begin{aligned} m_{\mathbf{13.2}} \cdot m_{\mathbf{1.2}} &= m_{\mathbf{13.2.4.5}} + m_{\mathbf{134.2.5}} + m_{\mathbf{135.2.4}} \\ &\quad + m_{\mathbf{13.24.5}} + m_{\mathbf{13.25.4}} + m_{\mathbf{135.24}} + m_{\mathbf{134.25}} \end{aligned} \tag{13}$$

Notice that the shapes indexing the first and last terms in (13) are the partitions  $\lambda(13.2) \cup \lambda(1.2)$  and  $\lambda(13.2) + \lambda(1.2)$ . As was the case in  $S^{\mathfrak{S}}$ , one of these shapes, namely  $\lambda(\mathbf{A}) + \lambda(\mathbf{B})$ , will always appear in the product, while appearance of the shape  $\lambda(\mathbf{A}) \cup \lambda(\mathbf{B})$  depends on the cardinality of  $\mathbf{x}$ .

Let us now describe the multiplication rule. Given any  $D \subseteq \mathbb{N}$  and  $k \in \mathbb{N}$ , we write  $D^{+k}$  for the set

$$D^{+k} := \{a + k : a \in D\}.$$

By extension, for any set partition  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$  we set  $\mathbf{A}^{+k} := \{A_1^{+k}, A_2^{+k}, \dots, A_r^{+k}\}$ . Also, we set  $\mathbf{A}_{\hat{i}} := \mathbf{A} \setminus \{A_i\}$ . Next, if  $\mathcal{X}$  is a collection of set partitions of  $D$ , and  $A$  is a set disjoint from  $D$ , we extend  $\mathcal{X}$  to partitions of  $A \cup D$  by the rule

$$A \diamond \mathcal{X} := \bigcup_{\mathbf{B} \in \mathcal{X}} \{A\} \cup \mathbf{B}.$$

Finally, given partitions  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$  of  $C$  and  $\mathbf{B} = \{B_1, B_2, \dots, B_s\}$  of  $D$  (disjoint from  $C$ ), their **quasi-shuffles**  $\mathbf{A} \omega \mathbf{B}$  are the set partitions of  $C \cup D$  recursively defined by the rules:

- $\mathbf{A} \omega \emptyset = \emptyset \omega \mathbf{A} := \mathbf{A}$ , where  $\emptyset$  is the unique set partition of the empty set;
- $\mathbf{A} \omega \mathbf{B} := \bigcup_{i=0}^s (A_1 \cup B_i) \diamond (\mathbf{A}_{\hat{1}} \omega (\mathbf{B}_{\hat{i}}))$ , taking  $B_0$  to be the empty set.

If  $\mathbf{A} \vdash [c]$  and  $\mathbf{B} \vdash [d]$ , we abuse notation and write  $\mathbf{A} \omega \mathbf{B}$  for  $\mathbf{A} \omega \mathbf{B}^{+c}$ . As shown in [2, Prop. 3.2], the multiplication rule for  $m_{\mathbf{A}}$  and  $m_{\mathbf{B}}$  in  $\mathcal{N}$  is

$$m_{\mathbf{A}} \cdot m_{\mathbf{B}} = \sum_{\mathbf{C} \in \mathbf{A} \omega \mathbf{B}} m_{\mathbf{C}}. \tag{14}$$

The subalgebra  $\mathcal{N}$ , like its commutative analog, is freely generated by certain monomial symmetric functions  $\{m_{\mathbf{A}}\}_{\mathbf{A} \in \mathcal{A}}$ , where  $\mathcal{A}$  is some carefully chosen collection of set partitions. This is the main theorem of Wolf [20]. We use two such collections later, our choice depending on whether or not  $|\mathbf{x}| < \infty$ .

The operation  $(-)^{+k}$  has a left inverse called the **standardization** operator and denoted by “ $(-)^{\downarrow}$ ”. It maps set partitions  $\mathbf{A}$  of any cardinality  $d$  subset  $D \subseteq \mathbb{N}$  to set

partitions of  $[d]$ , by defining  $\mathbf{A}^\downarrow$  as the pullback of  $\mathbf{A}$  along the unique increasing bijection from  $[d]$  to  $D$ . For example,  $(18.4)^\downarrow = 13.2$  and  $(18.4.67)^\downarrow = 15.2.34$ . The coproduct  $\Delta$  and counit  $\varepsilon$  on  $\mathcal{N}$  are given, respectively, by

$$\Delta(m_{\mathbf{A}}) = \sum_{\mathbf{B} \cup \mathbf{C} = \mathbf{A}} m_{\mathbf{B}^\downarrow} \otimes m_{\mathbf{C}^\downarrow} \quad \text{and} \quad \varepsilon(m_{\mathbf{A}}) = \delta_{\mathbf{A}, \emptyset},$$

where  $\mathbf{B} \cup \mathbf{C} = \mathbf{A}$  means that  $\mathbf{B}$  and  $\mathbf{C}$  form complementary subsets of  $\mathbf{A}$ . In the case  $|\mathbf{x}| = \infty$ , the maps  $\Delta$  and  $\varepsilon$  are algebra maps, making  $\mathcal{N}$  a graded connected Hopf algebra.

## 4 The place-action of $\mathfrak{S}$ on $\mathcal{N}$

### 4.1 Swapping places in $T_d$ and $\mathcal{N}_d$

On top of the permutation-action of the symmetric group  $\mathfrak{S}_{\mathbf{x}}$  on  $T$ , we also consider the “place-action” of  $\mathfrak{S}_d$  on the degree  $d$  homogeneous component  $T_d$ . Observe that the permutation-action of  $\sigma \in \mathfrak{S}_{\mathbf{x}}$  on a monomial  $\mathbf{z}$  corresponds to the functional composition

$$\sigma \circ \mathbf{z} : [d] \xrightarrow{\mathbf{z}} \mathbf{x} \xrightarrow{\sigma} \mathbf{x}$$

(notation as in Section 3.1). By contrast, the **place-action** of  $\rho \in \mathfrak{S}_d$  on  $\mathbf{z}$  gives the monomial

$$\mathbf{z} \circ \rho : [d] \xrightarrow{\rho} [d] \xrightarrow{\mathbf{z}} \mathbf{x},$$

composing  $\rho$  on the right with  $\mathbf{z}$ . In the linear extension of this action to all of  $T_d$ , it is easily seen that  $\mathcal{N}_d$  (even each  $\mathcal{N}_\mu$ ) is an invariant subspace of  $T_d$ . Indeed, for any set partition  $\mathbf{A} = \{A_1, A_2, \dots, A_r\} \vdash [d]$  and any  $\rho \in \mathfrak{S}_d$ , one has

$$m_{\mathbf{A}} \cdot \rho = m_{\rho^{-1} \cdot \mathbf{A}} \tag{15}$$

(see [15, §2]), where as usual  $\rho^{-1} \cdot \mathbf{A} := \{\rho^{-1}(A_1), \rho^{-1}(A_2), \dots, \rho^{-1}(A_r)\}$ .

### 4.2 The place-action structure of $\mathcal{N}$

Notice that the action in (15) is shape-preserving and transitive on set partitions of a given shape (i.e.,  $\mathcal{N}_\mu$  is an  $\mathfrak{S}_d$ -submodule of  $\mathcal{N}_d$  for each  $\mu \vdash d$ ). It follows that there is exactly one copy of the trivial  $\mathfrak{S}_d$ -module inside  $\mathcal{N}_\mu$  for each  $\mu \vdash d$ , that is, a basis for the place-action invariants in  $\mathcal{N}_d$  is indexed by partitions. We choose as basis the functions

$$\mathbf{m}_\mu := \frac{1}{(\dim \mathcal{N}_\mu) \mu^\dagger} \sum_{\lambda(\mathbf{A}) = \mu} m_{\mathbf{A}}, \tag{16}$$

with  $\mu^\dagger = a_1! a_2! \dots$  whenever  $\mu = 1^{a_1} 2^{a_2} \dots$ . The rationale for choosing this normalizing coefficient will be revealed in (20).

To simplify our discussion of the structure of  $\mathcal{N}$  in this context, we will say that  $\mathfrak{S}$  acts on  $\mathcal{N}$  rather than being fastidious about underlying in each situation that individual

$\mathcal{N}_d$ 's are being acted upon on the right by the corresponding group  $\mathfrak{S}_d$ . We denote the set  $\mathcal{N}^{\mathfrak{S}}$  of **place-invariants** by  $\Lambda$  in what follows. To summarize,

$$\Lambda = \text{span}\{\mathbf{m}_\mu : \mu \text{ a partition of } d, d \in \mathbb{N}\}. \quad (17)$$

The pair  $(\mathcal{N}, \Lambda)$  begins to look like the pair  $(S, S^{\mathfrak{S}})$  from the introduction. This was the observation that originally motivated our search for Theorem 1.

We next decompose  $\mathcal{N}$  into irreducible place-action representations. Although this can be worked out for any value of  $n$ , the results are more elegant when we send  $n$  to infinity. Recall that the **Frobenius characteristic** of a  $\mathfrak{S}_d$ -module  $\mathcal{V}$  is a symmetric function

$$\text{Frob}(\mathcal{V}) = \sum_{\mu \vdash d} v_\mu s_\mu,$$

where  $s_\mu$  is a Schur function (the character of “the” irreducible  $\mathfrak{S}_d$  representation  $\mathcal{V}_\mu$  indexed by  $\mu$ ) and  $v_\mu$  is the multiplicity of  $\mathcal{V}_\mu$  in  $\mathcal{V}$ . To reveal the  $\mathfrak{S}_d$ -module structure of  $\mathcal{N}_\mu$ , we use (15) and techniques from the theory of combinatorial species.

**Proposition 3.** *For a partition  $\mu = 1^{a_1} 2^{a_2} \cdots k^{a_k}$ , having  $a_i$  parts of size  $i$ , we have*

$$\text{Frob}(\mathcal{N}_\mu) = h_{a_1}[h_1] h_{a_2}[h_2] \cdots h_{a_k}[h_k], \quad (18)$$

with  $f[g]$  denoting plethysm of  $f$  and  $g$ , and  $h_i$  denoting the  $i^{\text{th}}$  homogeneous symmetric function.

Recall that the **plethysm**  $f[g]$  of two symmetric functions is obtained by linear and multiplicative extension of the rule  $p_k[p_\ell] := p_{k\ell}$ , where the  $p_k$ 's denote the usual power sum symmetric functions (see [12, I.8] for notation and details).

Let  $\text{Par}$  denote the combinatorial species of set partitions. So  $\text{Par}[n]$  denotes the set partitions of  $[n]$  and permutations  $\sigma: [n] \rightarrow [n]$  are transferred in a natural way to permutations  $\text{Par}[\sigma]: \text{Par}[n] \rightarrow \text{Par}[n]$ . The number  $\text{fix Par}[\sigma]$  of fixed points of this permutation is the same as the character  $\chi_{\text{Par}[n]}(\sigma)$  of the  $\mathfrak{S}_n$ -representation given by  $\text{Par}[n]$ . Given a partition  $\mu = 1^{a_1} 2^{a_2} \cdots k^{a_k}$ , put  $z_\mu := 1^{a_1} a_1! 2^{a_2} a_2! \cdots k^{a_k} a_k!$ . (There are  $n!/z_\mu$  permutations in  $\mathfrak{S}_n$  of cycle type  $\mu$ .) The **cycle index series** for  $\text{Par}$  is defined by

$$Z_{\text{Par}} = \sum_{n \geq 0} \sum_{\mu \vdash n} \text{fix Par}[\sigma_\mu] \frac{p_\mu}{z_\mu},$$

where  $\sigma_\mu$  is any permutation with cycle type  $\mu$  and  $p_\mu := p_1^{a_1} p_2^{a_2} \cdots p_k^{a_k}$  (taking  $p_i$  as the  $i$ -th power sum symmetric function).

*Proof.* Recall that the Schur and power sum symmetric functions are related by

$$s_\lambda = \sum_{\mu \vdash |\lambda|} \chi_\lambda(\sigma_\mu) \frac{p_\mu}{z_\mu},$$

so  $Z_{\text{Par}} = \text{Frob}(\text{Par})$ . Because  $\text{Par}$  is the composition  $\mathbf{E} \circ \mathbf{E}_+$  of the species of sets and nonempty sets, we also know that its cycle index series is given by plethystic substitution:  $Z_{\mathbf{E} \circ \mathbf{E}_+} = Z_{\mathbf{E}}[Z_{\mathbf{E}_+}]$ . See Theorem 2 and (12) in [1, I.4]. Combining these two results will give the proof.

First, we are only interested in that piece of  $\text{Frob}(\text{Par})$  coming from set partitions of shape  $\mu$ . For this we need weighted combinatorial species. If a set partition has shape  $\mu$ , give it the weight  $q_1^{a_1} q_2^{a_2} \cdots q_k^{a_k}$  in the cycle index series enumeration. The relevant identity is

$$Z_{\mathbf{P}}(\mathbf{q}) = \exp \sum_{k \geq 1} \frac{1}{k} \left( \exp \left( \sum_{j \geq 1} q_j^k \frac{p_{jk}}{j} \right) - 1 \right)$$

(cf. Example 13(c) of Chapter 2.3 in [1]). Collecting the terms of weight  $\mathbf{q}_\mu$  gives  $\text{Frob}(\mathcal{N}_\mu)$ . We get

$$\text{coeff}_{\mathbf{q}_\mu} [Z_{\text{Par}}(\mathbf{q})] = \prod_{i=1}^k \left( \sum_{\lambda \vdash a_i} \frac{p_\lambda}{z_\lambda} \right) \left[ \sum_{\nu \vdash i} \frac{p_\nu}{z_\nu} \right].$$

Standard identities [12, (2.14')] in I.2] between the  $h_i$ 's and  $p_j$ 's finish the proof.  $\square$

As an example, we consider  $\mu = 222 = 2^3$ . Since

$$h_2 = \frac{p_1^2}{2} + \frac{p_2}{2} \quad \text{and} \quad h_3 = \frac{p_1^3}{6} + \frac{p_1 p_2}{2} + \frac{p_3}{3},$$

a plethysm computation (and a change of basis) gives

$$\begin{aligned} h_3[h_2] &= \frac{p_1^3}{6} \left[ \frac{p_1^2}{2} + \frac{p_2}{2} \right] + \frac{p_1 p_2}{2} \left[ \frac{p_1^2}{2} + \frac{p_2}{2} \right] + \frac{p_3}{3} \left[ \frac{p_1^2}{2} + \frac{p_2}{2} \right] \\ &= \frac{1}{6} \left( \frac{p_1^2}{2} + \frac{p_2}{2} \right)^3 + \frac{1}{2} \left( \frac{p_1^2}{2} + \frac{p_2}{2} \right) \left( \frac{p_2^2}{2} + \frac{p_4}{2} \right) + \frac{1}{3} \left( \frac{p_3^2}{2} + \frac{p_6}{2} \right) \\ &= s_6 + s_{42} + s_{222}. \end{aligned}$$

That is,  $\mathcal{N}_{222}$  decomposes into three irreducible components, with the trivial representation  $s_6$  being the span of  $\mathbf{m}_{222}$  inside  $\Lambda$ .

### 4.3 $\Lambda$ meets $S^{\mathfrak{S}}$

We begin by explaining the choice of normalizing coefficient in (16). Analyzing the **abelianization** map  $\mathbf{ab} : T \rightarrow S$  (the map making the variables  $\mathbf{x}$  commute), Rosas and Sagan [15, Thm. 2.1] show that  $\mathbf{ab}|_{\mathcal{N}}$  satisfies:

$$\mathbf{ab}(m_{\mathbf{A}}) = \lambda(\mathbf{A})! m_{\lambda(\mathbf{A})}. \tag{19}$$

In particular,  $\mathbf{ab}$  maps onto  $S^{\mathfrak{S}}$  and

$$\mathbf{ab}(\mathbf{m}_\mu) = m_\mu. \tag{20}$$

Note that  $\mathbf{ab}$  is also an algebra map. The reader may wish to use (19) to compare (8) and (13). Formula (20) suggests that a natural right-inverse to  $\mathbf{ab}|_{\mathcal{N}}$  is given by

$$\iota : S^{\mathfrak{S}} \hookrightarrow \mathcal{N}, \quad \text{with} \quad \iota(m_\mu) := \mathbf{m}_\mu \quad \text{and} \quad \iota(1) = 1. \quad (21)$$

This fact, combined with the observation that  $\iota(S^{\mathfrak{S}}) = \Lambda$ , affords a quick proof of Theorem 1 when  $|\mathbf{x}| = \infty$ . We explain this now.

## 5 The coinvariant space of $\mathcal{N}$ (Case: $|\mathbf{x}| = \infty$ )

### 5.1 Quick proof of main result

When  $|\mathbf{x}| = \infty$ , the pair of maps  $(\mathbf{ab}, \iota)$  have further properties: the former is a Hopf algebra map and the latter is a coalgebra map [2, Props. 4.3 & 4.5]. Together with (20) and (21), these properties make  $\iota$  a **coalgebra splitting** of  $\mathbf{ab} : \mathcal{N} \rightarrow S^{\mathfrak{S}} \rightarrow 0$ . A theorem of Blattner, Cohen, and Montgomery immediately gives our main result in this case.

**Theorem 4** ([5], Thm. 4.14). *If  $H \xrightarrow{\pi} \overline{H} \rightarrow 0$  is an exact sequence of Hopf algebras that is split as a coalgebra sequence, and the splitting map  $\iota$  satisfies  $\iota(\overline{1}) = 1$ , then  $H$  is isomorphic to a crossed product  $A \# \overline{H}$ , where  $A$  is the left Hopf kernel of  $\pi$ . In particular,  $H \simeq A \otimes \overline{H}$  as vector spaces.*

For the technical definition of crossed products, we refer the reader to [5, §4]. We mention only that: (i) the crossed product  $A \# \overline{H}$  is a certain algebra structure placed on the tensor product  $A \otimes \overline{H}$ ; and (ii) the **left Hopf kernel** is the subalgebra

$$A := \{h \in H : (\text{id} \otimes \pi) \circ \Delta(h) = h \otimes \overline{1}\}.$$

We take  $H = \mathcal{N}$ ,  $\overline{H} = S^{\mathfrak{S}}$ , and  $\pi = \mathbf{ab}$ . Since our  $\iota$  is a coalgebra splitting, the coinvariant space  $\mathcal{C}$  we seek seems to be the left Hopf kernel of  $\mathbf{ab}$ . Before setting off to describe  $\mathcal{C}$  more explicitly, we point out that the left Hopf kernel is graded: the maps  $\Delta$ ,  $\text{id}$ , and  $\mathbf{ab}$  are graded, as is the map  $\mathcal{C} \# \Lambda \xrightarrow{\simeq} \mathcal{N}$  used in the proof of Theorem 4 (which is simply  $a \otimes \overline{h} \mapsto a \cdot \iota(\overline{h})$ ). Theorem 1 follows immediately from this result.

### 5.2 Atomic set partitions.

Recall the main result of Wolf [20] that  $\mathcal{N}$  is freely generated by some collection of functions. We announce our first choice for this collection now, following the terminology of [3]. Let  $\Pi$  denote the set of all set partitions (of  $[d]$ ,  $\forall d \geq 0$ ). The **atomic set partitions**  $\Pi$  are defined as follows. A set partition  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$  of  $[d]$  is *atomic* if there does not exist a pair  $(s, c)$  ( $1 \leq s < r$ ,  $1 \leq c < d$ ) such that  $\{A_1, A_2, \dots, A_s\}$  is a set partition of  $[c]$ . Conversely,  $\mathbf{A}$  is not atomic if there are set partitions  $\mathbf{B}$  of  $[d']$  and  $\mathbf{C}$  of  $[d'']$  splitting  $\mathbf{A}$  in two:  $\mathbf{A} = \mathbf{B} \cup \mathbf{C}^{+d'}$ . We write  $\mathbf{A} = \mathbf{B}|\mathbf{C}$  in this situation. A **maximal splitting**  $\mathbf{A} = \mathbf{A}'|\mathbf{A}'' \cdots |\mathbf{A}^{(t)}$  of  $\mathbf{A}$  is one where each  $\mathbf{A}^{(i)}$  is atomic. For example, the partition 17.235.4.68 is atomic, while 12.346.57.8 is not. The maximal splitting of

the latter would be  $12|124.35|1$ , but we abuse notation and write  $12|346.57|8$  to improve legibility.

It follows from [3, Corollary 9] that  $\mathcal{N}$  is freely generated by the atomic monomial functions  $\{m_{\mathbf{A}} : \mathbf{A} \in \dot{\Pi}\}$ . We now introduce an order on  $\Pi$  that will make this explicit. First we introduce the *restricted growth function* associated to a set partition (see Section 6.1): if  $\mathbf{A} = \{A_1, A_2, \dots, A_r\} \vdash [d]$ , define  $w(\mathbf{A}) \in \mathbb{N}^d$  by

$$w(\mathbf{A}) = w_1 w_2 \cdots w_d, \quad \text{with} \quad w_i := k \iff i \in A_k. \quad (22)$$

For example,  $w(\mathbf{13.24}) = 1212$  and  $w(\mathbf{17.235.4.68}) = 12232414$ . Now, given two atomic set partitions  $\mathbf{A} \vdash [c]$  and  $\mathbf{B} \vdash [d]$ , we put:

- $\mathbf{A} \succ \mathbf{B}$  when  $c > d$ ; or
- $\mathbf{A} \succ \mathbf{B}$  when  $c = d$  and  $w(\mathbf{A}) >_{\text{lex}} w(\mathbf{B})$ .

Finally, given two set partitions  $\mathbf{A}$  and  $\mathbf{B}$ , put  $\mathbf{A} > \mathbf{B}$  if  $\lambda(\mathbf{A}) <_{\text{lex}} \lambda(\mathbf{B})$  in the usual lexicographic order on integer partitions. If  $\lambda(\mathbf{A}) = \lambda(\mathbf{B})$ , then determine maximal splittings of  $\mathbf{A}$  and  $\mathbf{B}$ , view them as words in the atomic set partitions and use the lexicographic order induced by  $\succ$ . The following chain of set partitions of shape 3221 illustrates our total ordering on  $\Pi$ :

$$1|23|45|678 < 13.2|456|78 < 13.24|568.7 < 13.24|578.6 < 17.235.4.68 < 17.236.4.58.$$

In fact,  $1|23|45|678$  is the unique minimal element of  $\Pi$  of shape 3221.

Define the **leading term** of a sum  $\sum_{\mathbf{C}} \alpha_{\mathbf{C}} m_{\mathbf{C}}$  to be the monomial  $m_{\mathbf{C}_0}$  such that  $\mathbf{C}_0$  is greatest (according to  $>$  above) among all  $\mathbf{C}$  with  $\alpha_{\mathbf{C}} \neq 0$ . Combined with (14), our definition of  $>$  makes it clear that the leading term of  $m_{\mathbf{A}} \cdot m_{\mathbf{B}}$  is  $m_{\mathbf{A|B}}$  and that  $\mathcal{N}$  is freely generated by the atomic monomial functions. Moreover, it is clear that multiplication in  $\mathcal{N}$  is shape-filtered. Since the left Hopf kernel  $\mathcal{C}$  is a subalgebra,  $\mathcal{C}$  is shape-filtered as well. Finally, the isomorphism  $\mathcal{C} \# \Lambda \xrightarrow{\simeq} \mathcal{N}$  constructed in the proof of Theorem 4 is also shape-filtered. These facts give Corollary 2 immediately.

### 5.3 Explicit description of the Hopf algebra structure of $\mathcal{C}$

We begin by partitioning  $\dot{\Pi}$  into two sets according to length,

$$\dot{\Pi}_{(1)} := \{ \mathbf{A} \in \dot{\Pi} : \ell(\mathbf{A}) = 1 \} \quad \text{and} \quad \dot{\Pi}_{(>1)} := \{ \mathbf{A} \in \dot{\Pi} : \ell(\mathbf{A}) > 1 \}.$$

It is easy to find elements of the left Hopf kernel  $\mathcal{C}$ . For instance, if  $\mathbf{A}$  and  $\mathbf{B}$  belong to  $\dot{\Pi}_{(1)}$ , then the Lie bracket  $[m_{\mathbf{A}}, m_{\mathbf{B}}]$  belongs to  $\mathcal{C}$ . Indeed,

$$\begin{aligned} \Delta([m_{\mathbf{A}}, m_{\mathbf{B}}]) &= \Delta(m_{\mathbf{A|B}} - m_{\mathbf{B|A}}) \\ &= m_{\mathbf{A|B}} \otimes 1 + m_{\mathbf{A}} \otimes m_{\mathbf{B}} + m_{\mathbf{B}} \otimes m_{\mathbf{A}} + 1 \otimes m_{\mathbf{A|B}} \\ &\quad - m_{\mathbf{B|A}} \otimes 1 - m_{\mathbf{B}} \otimes m_{\mathbf{A}} - m_{\mathbf{A}} \otimes m_{\mathbf{B}} - 1 \otimes m_{\mathbf{B|A}} \\ &= (m_{\mathbf{A|B}} - m_{\mathbf{B|A}}) \otimes 1 + 1 \otimes (m_{\mathbf{A|B}} - m_{\mathbf{B|A}}). \end{aligned}$$

Since  $\mathbf{ab}(m_{\mathbf{A}|\mathbf{B}}) = \mathbf{ab}(m_{\mathbf{B}|\mathbf{A}})$ , we have

$$(\mathrm{id} \otimes \mathbf{ab}) \circ \Delta([m_{\mathbf{A}}, m_{\mathbf{B}}]) = [m_{\mathbf{A}}, m_{\mathbf{B}}] \otimes 1$$

as desired. Similarly, the difference of monomial functions  $m_{13.2} - m_{12.3}$  belongs to  $\mathcal{C}$ . The leading term here is indexed by  $13.2 \in \dot{\Pi}_{(>1)}$ . These two simple examples essentially exhaust the different ways in which an element can belong to  $\mathcal{C}$ . The following discussion makes this precise.

From [3, Theorem 15], we learn that  $\mathcal{N}$  is cofree cocommutative with minimal cogenerated set indexed by the Lyndon words in  $\dot{\Pi}$ . (This result and the previously mentioned freeness result may also be deduced from the techniques developed in [9].) Since single letters are Lyndon words, we know there are primitive elements associated to each atomic set partition. Recall that an element  $h$  in a Hopf algebra is **primitive** if  $\Delta(h) = h \otimes 1 + 1 \otimes h$ . Let  $\mathrm{Prim}(\mathcal{N})$  denote the set of primitive elements in  $\mathcal{N}$ —a Lie algebra under the commutator bracket.

Bearing the free and cofree cocommutative results in mind, a classical theorem of Milnor and Moore [13] guarantees that  $\mathcal{N}$  is isomorphic to the universal enveloping algebra  $\mathfrak{U}(\mathfrak{L}(\dot{\Pi}))$  of the free Lie algebra  $\mathfrak{L}(\dot{\Pi})$  on the set  $\dot{\Pi}$ . In the isomorphism  $\mathfrak{L}(\dot{\Pi}) \xrightarrow{\cong} \mathrm{Prim}(\mathcal{N})$ , one may map  $\mathbf{A} \in \dot{\Pi}_{(1)}$  to  $m_{\mathbf{A}}$  since these monomial functions are already primitive. The choice of where to send  $\mathbf{A} \in \dot{\Pi}_{(>1)}$  is the subject of the next proposition.

**Proposition 5.** *For each  $\mathbf{A} \in \dot{\Pi}_{(>1)}$ , there is a primitive element  $\tilde{m}_{\mathbf{A}}$  of  $\mathcal{N}$ ,*

$$\tilde{m}_{\mathbf{A}} = m_{\mathbf{A}} - \sum_{\mathbf{B} \in \dot{\Pi}} \alpha_{\mathbf{B}} m_{\mathbf{B}},$$

satisfying: (i) if  $\mathbf{B} \in \dot{\Pi}$  or  $\lambda(\mathbf{B}) \neq \lambda(\mathbf{A})$ , then  $\alpha_{\mathbf{B}} = 0$ ; and (ii)  $\sum_{\mathbf{B}} \alpha_{\mathbf{B}} = 1$ .

*Proof.* Suppose  $\mathbf{A} \in \dot{\Pi}_{(>1)}$ . A primitive  $\tilde{m}_{\mathbf{A}}$  exists by the Milnor–Moore theorem, as explained above.

(i). Since  $\mathcal{N} = \bigoplus_{\mu} \mathcal{N}_{\mu}$  is a coalgebra grading by shape, we may assume  $\lambda(\mathbf{B}) = \lambda(\mathbf{A})$  for any nonzero coefficients  $\alpha_{\mathbf{B}}$ . Now, since there are linearly independent primitive elements in  $\mathcal{N}$  associated to every atomic set partition, we may use Gaussian elimination and our ordering on  $\dot{\Pi}$  to ensure that  $\alpha_{\mathbf{B}} = 0$  for any  $\mathbf{B} \in \dot{\Pi}$ .

(ii). Define linear maps  $\Delta_+^j : \mathcal{N}_+ \rightarrow \mathcal{N} \otimes \mathcal{N}$  recursively by

$$\begin{aligned} \Delta_+(h)^1 &:= \Delta(h) - h \otimes 1 - 1 \otimes h, \\ \Delta_+^{j+1}(h) &:= (\Delta_+ \otimes \mathrm{id}^{\otimes j}) \circ \Delta_+^j(h) \quad \text{for } j > 0. \end{aligned}$$

Assume that (i) is satisfied for  $\tilde{m}_{\mathbf{A}}$  and that  $\mathbf{A} = \{A_1, A_2, \dots, A_r\}$ . Since  $\Delta_+(\tilde{m}_{\mathbf{A}}) = 0$ , we have  $\Delta_+^j(m_{\mathbf{A}}) = \Delta_+^j(\sum_{\mathbf{B}} \alpha_{\mathbf{B}} m_{\mathbf{B}})$  for all  $j > 1$ . Now,

$$\Delta_+^r(m_{\mathbf{A}}) = \sum_{\sigma \in \mathfrak{S}_r} m_{A_{\sigma 1} \downarrow} \otimes m_{A_{\sigma 2} \downarrow} \otimes \cdots \otimes m_{A_{\sigma r} \downarrow}.$$

Indeed, the same holds for any  $\mathbf{B}$  with  $\lambda(\mathbf{B}) = \lambda(\mathbf{A})$ :

$$\Delta_+^r \left( \sum_{\mathbf{B}} \alpha_{\mathbf{B}} m_{\mathbf{B}} \right) = \left( \sum_{\mathbf{B}} \alpha_{\mathbf{B}} \right) \sum_{\sigma \in \mathfrak{S}_r} m_{A_{\sigma_1} \downarrow} \otimes m_{A_{\sigma_2} \downarrow} \otimes \cdots \otimes m_{A_{\sigma_r} \downarrow}.$$

Conclude that  $\sum_{\mathbf{B}} \alpha_{\mathbf{B}} = 1$ . □

We say an element  $h \in \mathcal{N}_\mu$  has the “zero-sum” property if it satisfies (ii) from the proposition. Put  $\tilde{m}_{\mathbf{A}} := m_{\mathbf{A}}$  for  $\mathbf{A} \in \dot{\Pi}_{(1)}$ . We next describe the coinvariant space  $\mathcal{C}$ .

**Corollary 6.** *Let  $\mathcal{C}$  be the Lie ideal in  $\mathfrak{L}(\dot{\Pi})$  given by  $\mathcal{C} = [\mathfrak{L}(\dot{\Pi}), \mathfrak{L}(\dot{\Pi})] \oplus \dot{\Pi}_{(>1)}$ . If  $\varphi : \mathfrak{U}(\mathfrak{L}(\dot{\Pi})) \rightarrow \mathcal{N}$  is the Milnor–Moore isomorphism given by putting  $\varphi(\mathbf{A}) := \tilde{m}_{\mathbf{A}}$  for all  $\mathbf{A} \in \dot{\Pi}$  and extending multiplicatively, then the left Hopf kernel  $\mathcal{C}$  is the Hopf subalgebra  $\varphi(\mathfrak{U}(\mathcal{C}))$ .*

*Proof.* We first show that  $\varphi(\mathfrak{U}(\mathcal{C})) \subseteq \mathcal{C}$ . We certainly have  $\tilde{m}_{\mathbf{A}} \in \mathcal{C}$  for all  $\mathbf{A} \in \dot{\Pi}_{(>1)}$ , since the zero-sum property means  $\mathbf{ab}(\tilde{m}_{\mathbf{A}}) = 0$ . Next suppose  $f \in [\mathfrak{L}(\dot{\Pi}), \mathfrak{L}(\dot{\Pi})]$  is a sum of Lie brackets  $[\mathbf{A}] = [[\dots [\mathbf{A}', \mathbf{A}''], \dots], \mathbf{A}^{(t)}]$ . In this case,  $\varphi(f) \in \mathcal{C}$  because each  $\varphi([\mathbf{A}])$  is primitive and  $\mathbf{ab}$  is an algebra map. Indeed,  $\mathbf{ab}([\tilde{m}_{\mathbf{A}'}, \tilde{m}_{\mathbf{A}''}]) = 0$ . The inclusion follows, since  $\mathfrak{U}(\mathcal{C})$  is generated by elements of these two types.

It remains to show that  $\mathcal{C} \subseteq \varphi(\mathfrak{U}(\mathcal{C}))$ . To begin, note that  $\mathfrak{L}(\dot{\Pi})/\mathcal{C}$  is isomorphic to the abelian Lie algebra generated by  $\dot{\Pi}_{(1)}$ . The universal enveloping algebra of this latter object is evidently isomorphic to  $S^{\mathfrak{S}}$ . (Send  $\mathbf{A} = \{[d]\}$  to  $m_d$ .) The Poincaré–Birkhoff–Witt theorem guarantees that the map  $\varphi(\mathfrak{U}(\mathcal{C})) \otimes S^{\mathfrak{S}} \rightarrow \mathcal{N}$  given by  $a \otimes b \mapsto a \cdot \iota(b)$  is onto  $\mathcal{N}$ . Conclude that  $\mathcal{C} \subseteq \varphi(\mathfrak{U}(\mathcal{C}))$ , as needed. □

Before turning to the case  $|\mathbf{x}| < \infty$ , we remark that we have left unanswered the question of finding a systematic procedure (e.g., a closed formula in the spirit of Möbius inversion) that constructs a primitive element  $\tilde{m}_{\mathbf{A}}$  for each  $\mathbf{A} \in \dot{\Pi}_{(>1)}$ . This is accomplished in [11].

## 6 The coinvariant space of $\mathcal{N}$ (Case: $|\mathbf{x}| \leq \infty$ )

### 6.1 Restricted growth functions

We repeat our example of Section 3.3 in the case  $n = 3$ . The leading term with respect to our previous order would be  $m_{13.2.4.5}$ , except that this term does not appear because 13.2.4.5 has more than  $n = 3$  parts:

$$m_{13.2} \cdot m_{1.2} = 0 + m_{134.2.5} + m_{135.2.4} + m_{13.24.5} + m_{13.25.4} + m_{135.24} + m_{134.25}.$$

Fortunately, the map  $w$  from set partitions to words on the alphabet  $\mathbb{N}_{>0}$  reveals a more useful leading term, underlined below:

$$m_{121} \cdot m_{12} = 0 + m_{12113} + m_{12131} + m_{12123} + m_{12132} + m_{12121} + \underline{m_{12112}}. \quad (23)$$

Notice that the words appearing on the right in (23) all begin by 121 and that the concatenation  $\underline{121}\underline{12}$  is the lexicographically smallest word appearing there. This is generally true and easy to see: if  $w(\mathbf{A}) = u$  and  $w(\mathbf{B}) = v$ , then  $uv$  is the lexicographically smallest element of  $w(\mathbf{A} \cup \mathbf{B})$ .

The map  $w$  maps set partitions to **restricted growth functions**, i.e., the words  $w = w_1 w_2 \cdots w_d$  satisfying  $w_1 = 1$  and  $w_i \leq 1 + \max\{w_1, w_2, \dots, w_{i-1}\}$  for all  $2 \leq i \leq d$ . We call them restricted growth words here. See [16, 17, 19] and [6, 8] for some of their combinatorial properties and applications. These words are also known as “*rhyme scheme words*” in the literature; see [14] and [18, A000110]. Before looking for a coinvariant space  $\mathcal{C}$  within  $\mathcal{N}$ , we first fix the representatives of  $\Lambda$ . Consider the partition  $\mu = 3221$ . Of course,  $\mathbf{m}_\mu$  is the sum of all set partitions of shape  $\mu$ , but it will be nice to have a single one in mind when we speak of  $\mathbf{m}_\mu$ . A convenient choice turns out to be 123.45.67.8: if we use the length plus lexicographic order on  $w(\Pi)$ , then it is easy to see that  $w(123.45.67.8) = 11122334$  is the minimal element of  $\Pi$  of shape 3221. We are led to introduce the words

$$w(\mu) := 1^{\mu_1} 2^{\mu_2} \cdots k^{\mu_k}$$

associated to partitions  $\mu = (\mu_1, \mu_2, \dots, \mu_k)$ ; we call such restricted growth words **convex words** since  $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_k$ .

## 6.2 Proof of main theorem

We say that a restricted growth word is **non-splittable** if  $w_i \cdots w_{n-1} w_n$  is not a restricted growth word for any  $i > 1$ . The **maximal splitting** of a restricted growth word  $w$  is the maximal deconcatenation  $w = w' | w'' | \cdots | w^{(r)}$  of  $w$  into non-splittable words  $w^{(i)}$ . For example, 12314 is non-splittable while 11232411 is a string of four non-splittable words  $1 | 12324 | 1 | 1$ .

It is easy to see that if  $a, b, c$ , and  $d$  are non-splittable, then  $ac = bd$  if and only if  $a = b$  and  $c = d$ . Together with the remarks on  $\mathbf{A} \cup \mathbf{B}$  following (23), this implies that if  $\{u_1, u_2, \dots, u_r\}$  and  $\{v_1, v_2, \dots, v_s\}$  are two sets of non-splittable words, then

$$m_{u_1} m_{u_2} \cdots m_{u_r} \quad \text{and} \quad m_{v_1} m_{v_2} \cdots m_{v_s}$$

share the same leading term (namely,  $m_{u_1 | u_2 | \cdots | u_r}$ ) if and only if  $r = s$  and  $u_i = v_i$  for all  $i$ . In other words, our algebra  $\mathcal{N}$  is *non-splittable word-filtered* and freely generated by the monomial functions  $\{m_{w(\mathbf{A})} : w(\mathbf{A}) \text{ is non-splittable}\}$ . This is one of the collections of monomial functions originally chosen by Wolf [20].

We aim to index  $\mathcal{C}$  by the restricted growth words that don't end in a convex word. Toward that end, we introduce the notion of **bimodal words**. These are words with a maximal (but possibly empty) convex prefix, followed by one non-splittable word. The **bimodal decomposition** of a restricted growth word  $w$  is the expression of  $w$  as a product  $w = w' | w'' | \cdots | w^{(r)} | w^{(r+1)}$ , where  $w', w'', \dots, w^{(r)}$  are bimodal and  $w^{(r+1)}$  is a possibly empty convex word (which we call a **tail**). For a given word  $w$ , this decomposition is accomplished by first splitting  $w$  into non-splittable words, then recombining, from

left to right, consecutive non-splittable words to form bimodal words. For instance, the maximal splitting of 1122212 into non-splittable words is  $1|1222|12$ . The first two factors combine to make one bimodal word; the last factor is a convex tail:  $1122212 \mapsto \widehat{1|1222} \widehat{1}2$ . Similarly,

$$1231231411122311 \mapsto 123|12314|1|1|1223|1|1 \mapsto \widehat{123|12314} \widehat{1|1|1223} \widehat{1} \widehat{1}.$$

Suppose now that  $u$  and  $v$  are restricted growth words and that the bimodal decomposition of  $u$  is tail-free. Then by construction, the bimodal decomposition of  $uv$  is the concatenation of the respective bimodal decompositions of  $u$  and  $v$ . We are ready to identify  $\mathcal{C}$  as a subalgebra of  $\mathcal{N}$ .

**Theorem 7.** *Let  $\mathcal{C}$  be the subalgebra of  $\mathcal{N}$  generated by  $\{m_v : v \text{ is bimodal}\}$ . Then  $\mathcal{C}$  has a basis indexed by restricted growth words  $w$  whose bimodal decompositions are tail-free. Moreover, the map  $\varphi : \mathcal{C} \otimes \Lambda \rightarrow \mathcal{N}$  given by  $m_w m_{w''} \cdots m_{w^{(r)}} \otimes \mathbf{m}_\mu \mapsto m_{w'|w''|\dots|w^{(r)}}|w(\mu)$  is a vector space isomorphism.*

*Proof.* The advertised map is certainly onto, since  $\{m_w : w \in w(\Pi)\}$  is a basis for  $\mathcal{N}$  and every restricted growth word has a bimodal decomposition  $w'|w''|\dots|w^{(r)}|w(\mu)$ . It remains to show that the map is one-to-one.

Note that the monomial functions  $\{m_v : v \text{ is bimodal}\}$  are algebraically independent: certainly, the leading term in a product  $m_{v_1} m_{v_2} \cdots m_{v_s}$  (with  $v_i$  bimodal) is  $m_{v_1|v_2|\dots|v_s}$ ; now, since every word has a unique bimodal decomposition, no (nontrivial) linear combination of products of this form can be zero. Finally, the leading term in the simple tensor  $m_w m_{w''} \cdots m_{w^{(r)}} \otimes \mathbf{m}_\mu$  is the basis vector  $m_{w'|w''|\dots|w^{(r)}} \otimes m_{w(\mu)}$ , so no (nontrivial) linear combination of these will vanish under the map  $\varphi$ .  $\square$

## References

- [1] François Bergeron, Gilbert Labelle, and Pierre Leroux. *Combinatorial species and tree-like structures*, volume 67 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1998. Translated from the 1994 French original by Margaret Readdy, With a foreword by Gian-Carlo Rota.
- [2] Nantel Bergeron, Christophe Reutenauer, Mercedes Rosas, and Mike Zabrocki. Invariants and coinvariants of the symmetric groups in noncommuting variables. *Canad. J. Math.*, 60(2):266–296, 2008.
- [3] Nantel Bergeron and Mike Zabrocki. The Hopf algebras of symmetric functions and quasi-symmetric functions in non-commutative variables are free and co-free. *J. Algebra Appl.*, 8(4):581–600, 2009.
- [4] George M. Bergman and Paul M. Cohn. Symmetric elements in free powers of rings. *J. London Math. Soc. (2)*, 1:525–534, 1969.
- [5] Robert J. Blattner, Miriam Cohen, and Susan Montgomery. Crossed products and inner actions of Hopf algebras. *Trans. Amer. Math. Soc.*, 298(2):671–711, 1986.

- [6] David Bremner and Lars Schewe. Edge-graph diameter bounds for convex polytopes with few facets. preprint, arXiv:0809.0915v2.
- [7] Edward Formanek. Noncommutative invariant theory. In *Group actions on rings (Brunswick, Maine, 1984)*, volume 43 of *Contemp. Math.*, pages 87–119. Amer. Math. Soc., Providence, RI, 1985.
- [8] Ira Gessel, Jonathan Weinstein, and Herbert S. Wilf. Lattice walks in  $\mathbf{Z}^d$  and permutations with no long ascending subsequences. *Electron. J. Combin.*, 5:Research Paper 2, 11 pp. (electronic), 1998.
- [9] Florent Hivert, Jean-Christophe Novelli, and Jean-Yves Thibon. Commutative combinatorial Hopf algebras. *J. Algebraic Combin.*, 28(1):65–95, 2008.
- [10] Martin Klazar. Bell numbers, their relatives, and algebraic differential equations. *J. Combin. Theory Ser. A*, 102(1):63–87, 2003.
- [11] Aaron Lauve and Mitja Mastnak. The primitives and antipode in the Hopf algebra of symmetric functions in noncommuting variables. preprint, arXiv:1006.0367v3.
- [12] Ian G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [13] John W. Milnor and John C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965.
- [14] John Riordan. A budget of rhyme scheme counts. In *Second International Conference on Combinatorial Mathematics (New York, 1978)*, volume 319 of *Ann. New York Acad. Sci.*, pages 455–465. New York Acad. Sci., New York, 1979.
- [15] Mercedes H. Rosas and Bruce E. Sagan. Symmetric functions in noncommuting variables. *Trans. Amer. Math. Soc.*, 358(1):215–232 (electronic), 2006.
- [16] Bruce E. Sagan. Pattern avoidance in set partitions. *Ars Combin.*, 94:79–96, 2010.
- [17] Rodica Simion. Combinatorial statistics on noncrossing partitions. *J. Combin. Theory Ser. A*, 66(2):270–301, 1994.
- [18] Niel J. A. Sloane. The on-line encyclopedia of integer sequences. published electronically at <http://www.research.att.com/~njas/sequences/>.
- [19] Michelle Wachs and Dennis White.  $p, q$ -Stirling numbers and set partition statistics. *J. Combin. Theory Ser. A*, 56(1):27–46, 1991.
- [20] Margarete C. Wolf. Symmetric functions of non-commutative elements. *Duke Math. J.*, 2(4):626–637, 1936.