

Optimal lower bound for 2-identifying codes in the hexagonal grid*

Ville Junnila[†]

Turku Centre for Computer Science TUCS
Department of Mathematics
University of Turku, FI-20014 Turku, Finland
`viljun@utu.fi`

Tero Laihonen

Department of Mathematics
University of Turku, FI-20014 Turku, Finland
`terolai@utu.fi`

Submitted: Oct 26, 2010; Accepted: May 28, 2012; Published: Jun 13, 2012
Mathematics Subject Classifications: 05C70, 68R05, 94B65, 94C12

Abstract

An r -identifying code in a graph $G = (V, E)$ is a subset $C \subseteq V$ such that for each $u \in V$ the intersection of C and the ball of radius r centered at u is non-empty and unique. Previously, r -identifying codes have been studied in various grids. In particular, it has been shown that there exists a 2-identifying code in the hexagonal grid with density $4/19$ and that there are no 2-identifying codes with density smaller than $2/11$. Recently, the lower bound has been improved to $1/5$ by Martin and Stanton (2010). In this paper, we prove that the 2-identifying code with density $4/19$ is optimal, i.e. that there does not exist a 2-identifying code in the hexagonal grid with smaller density.

Keywords: Identifying code; optimal code; hexagonal grid

1 Introduction

Let $G = (V, E)$ be a simple connected and undirected graph with V as the set of vertices and E as the set of edges. Let u and v be vertices in V . If u and v are adjacent to

*This paper has been presented in part at the International Workshop on Coding and Cryptography 2011, WCC 2011.

[†]Research supported by the Academy of Finland under grant 210280.

each other, then the edge between u and v is denoted by $\{u, v\}$ (or in short by uv). The *distance* between u and v is denoted by $d(u, v)$ and is defined as the number of edges in any shortest path between u and v . Let r be a positive integer. We say that u *r -covers* v if the distance $d(u, v)$ is at most r . The *ball of radius r centered at u* is defined as

$$B_r(u) = \{x \in V \mid d(u, x) \leq r\}.$$

A non-empty subset of V is called a *code* in G , and its elements are called *codewords*. Let $C \subseteq V$ be a code and u be a vertex in V . An *I -set* (or an *identifying set*) of the vertex u with respect to the code C is defined as

$$I_r(C; u) = I_r(u) = B_r(u) \cap C.$$

The following definition is due to Karpovsky *et al.* [5].

Definition 1.1. Let r be a positive integer. A code $C \subseteq V$ is said to be *r -identifying* in G if for all $u, v \in V$ ($u \neq v$) the set $I_r(C; u)$ is non-empty and

$$I_r(C; u) \neq I_r(C; v).$$

Let X and Y be subsets of V . The *symmetric difference* of X and Y is defined as $X \triangle Y = (X \setminus Y) \cup (Y \setminus X)$. We say that the vertices u and v are *r -separated* by a code $C \subseteq V$ (or by a codeword of C) if the symmetric difference $I_r(C; u) \triangle I_r(C; v)$ is non-empty. The definition of r -identifying codes can now be reformulated as follows: $C \subseteq V$ is an r -identifying code in G if and only if for all $u, v \in V$ ($u \neq v$) the vertex u is r -covered by a codeword of C and

$$I_r(C; u) \triangle I_r(C; v) \neq \emptyset.$$

In this paper, we study identifying codes in the hexagonal grid. We define the hexagonal grid $G_H = (V_H, E_H)$ using the brick wall representation as follows: the set of vertices $V_H = \mathbb{Z}^2$ and the set of edges

$$E_H = \{\{\mathbf{u} = (i, j), \mathbf{v}\} \mid \mathbf{u}, \mathbf{v} \in \mathbb{Z}^2, \mathbf{u} - \mathbf{v} \in \{(0, (-1)^{i+j+1}), (\pm 1, 0)\}\}.$$

This definition is illustrated in Figure 1(a). The hexagonal grid can also be illustrated using the honeycomb representation as in Figure 1(b). In both illustrations, lines represent the edges and intersections of the lines represent the vertices of G_H . The labeling of the vertices in the brick wall representation is self-explanatory. This labeling can also be applied to the honeycomb representation, if we visualize the honeycomb to be obtained from the brick wall by squeezing it from left and right. For an example of the labeling of the vertices, we refer to Figure 1.

To measure the size of an identifying code in the infinite hexagonal grid, we introduce the notion of density. For the formal definition, we first define

$$Q_n = \{(x, y) \in V_H \mid |x| \leq n, |y| \leq n\}.$$




Figure 1: The brick wall and the honeycomb representations illustrated.

Then the *density* of a code $C \subseteq V_H$ is defined as

$$D(C) = \limsup_{n \rightarrow \infty} \frac{|C \cap Q_n|}{|Q_n|}.$$

Naturally, we try to construct identifying codes with as small density as possible. Moreover, we say that an r -identifying code is *optimal*, if there do not exist any r -identifying codes with smaller density.

Previously, r -identifying codes in G_H have been studied in various papers. The first results concerning r -identifying codes in G_H have been presented in the seminal paper [5] in the case $r = 1$. Later these results have been improved by showing that there exists a 1-identifying code with density $3/7$ (see Cohen *et al.* [3]) and that there do not exist 1-identifying codes in G_H with density smaller than $12/29$ (see Cranston and Yu [4]). For general $r \geq 2$, Charon *et al.* [1] showed that each r -identifying code C in G_H has $D(C) \geq 2/(5r + 3)$ if r is even and $D(C) \geq 2/(5r + 2)$ if r is odd. They also presented a construction for each $r \geq 2$ giving an r -identifying code $C \subseteq V_H$ with $D(C) \sim 8/(9r)$.

For small values of r , the previous constructions have been improved in [2] by Charon *et al.* In particular, it is shown that there exists a 2-identifying code in G_H with density $4/19$. In the case $r = 2$, the aforementioned general lower bound is improved in Martin and Stanton [6] by showing that the density of any 2-identifying code in G_H is at least $1/5$. In this paper, we further improve this lower bound to $4/19$. In other words, we show that the previously presented 2-identifying code with density $4/19$ is optimal.

The organization of the paper is as follows. In Section 2, we first introduce some preliminary definitions and results. Then, in Section 3, we proceed with the actual proof of the lower bound.

2 Basics

Let $G = (V, E)$ be a simple connected and undirected graph. Assume also that C is a code in G . The following concept of the share of a codeword has been introduced by Slater in [7]. The *share* of a codeword $c \in C$ is denoted by $s_r(c)$ and defined as

$$s_r(C; c) = s_r(c) = \sum_{u \in B_r(c)} \frac{1}{|I_r(C; u)|}.$$

The notion of share proves to be useful in determining lower bounds of r -identifying codes (as explained in the following paragraph).

Assume that $G = (V, E)$ is a finite graph and D is a code in G such that $B_r(u) \cap D$ is non-empty for all $u \in V$. Then it is easy to conclude that $\sum_{c \in D} s_r(D; c) = |V|$. Assume further that $s_r(D; c) \leq \alpha$ for all $c \in D$. Then we have $|V| \leq \alpha|D|$, which immediately implies

$$|D| \geq \frac{1}{\alpha}|V|.$$

Assume then that for any r -identifying code C in G we have $s_r(C; c) \leq \alpha$ for all $c \in C$. By the aforementioned observation, we then obtain the lower bound $|V|/\alpha$ for the size of an r -identifying code in G . In other words, by determining the maximum share for any r -identifying code, we obtain a lower bound for the minimum size of an r -identifying code.

The previous reasoning can also be generalized to the case when an infinite graph is considered. In particular, if for any r -identifying code in G_H we have $s_r(C; c) \leq \alpha$ for all $c \in C$, then it can be shown that the density of an r -identifying code in G_H is at least $1/\alpha$ (compare to Theorem 3.5). The main idea behind the proof of the lower bound (in Section 3) is based on this observation, although we use a more sophisticated method by showing that for any 2-identifying code the share is on *average* at most $19/4$. In Theorem 3.5, we present a formal proof to verify that this method is indeed valid.

In the proof of the lower bound, we need to determine upper bounds for shares of codewords. To formally present a way to estimate shares, we first need to introduce some notations. Let $D \subseteq V$ be a code and c be a codeword of D . Consider then the I -sets $I_r(D; u)$ when u goes through all the vertices in $B_r(c)$. (Notice that all of these I -sets do not have to be different.) Denote the different identifying sets by $I_1, I_2, \dots, I_k$, where k is a positive integer. Furthermore, denote the number of identifying sets equal to I_j by i_j ($j = 1, 2, \dots, k$). Now we are ready to present the following lemma, which provides a method to estimate the shares of the codewords.

Lemma 2.1. *Let C be an r -identifying code in G and let D be a non-empty subset of C . For $c \in D$, using the previous notations, we have*

$$s_r(C; c) \leq \sum_{j=1}^k \left(\frac{1}{|I_j|} + (i_j - 1) \frac{1}{|I_j| + 1} \right).$$

Proof. Assume that $c \in D$. Then, for each $j = 1, 2, \dots, k$, define $\mathcal{I}_j = \{u \in B_r(c) \mid I_j = I_r(D; u)\}$. Now it is obvious that for at most one vertex $u \in \mathcal{I}_j$ we have $I_j = I_r(C; u)$ and the other vertices of $\mathcal{I}_j$ are r -covered by at least $|I_j| + 1$ codewords of C . Hence, the claim immediately follows. $\square$

The previous lemma will be used numerous times in the following presentation. The computations needed in applying this lemma may sometimes be a little bit tedious, but always very straightforward. It is also quite easy to implement an algorithm to compute the upper bound given by the lemma. Furthermore, the use of the previous lemma is illustrated in the following example.




Figure 2: The cases of Example 2.2 illustrated. The black dots represent codewords of C .

Example 2.2. Let C be a 2-identifying code in the hexagonal grid G_H . For the first case (see Figure 2(a)), we assume that $D = \{(0, 0), (0, 1), (1, -1)\}$ is a subset of C . Now we have the following facts:

- $I_2(D; \mathbf{u}) = \{(0, 0), (1, -1)\}$ for $\mathbf{u} = (-1, -1), (1, -1)$ or $(2, 0)$ (the vertices labeled with 1 in the figure),
- $I_2(D; \mathbf{u}) = \{(0, 1), (0, 0), (1, -1)\}$ for $\mathbf{u} = (0, 0)$ or $(1, 0)$ (the vertices labeled with 2 in the figure),
- $I_2(D; \mathbf{u}) = \{(0, 0), (0, 1)\}$ for $\mathbf{u} = (-1, 1), (0, 1), (1, 1)$ or $(-1, 0)$ (the vertices labeled with 3 in the figure),
- $I_2(D; \mathbf{u}) = \{(0, 1)\}$ for $\mathbf{u} = (-2, 1), (-1, 2), (1, 2)$ or $(2, 1)$ (the vertices labeled with 4 in the figure), and
- $I_2(D; (-2, 0)) = \{(0, 0)\}$ (the vertex labeled with 5 in the figure).

Thus, by Lemma 2.1, we obtain that

$$s_2(C; (0, 0)) \leq \left(\frac{1}{2} + 2 \cdot \frac{1}{3}\right) + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{2} + 3 \cdot \frac{1}{3}\right) + 1 = \frac{17}{4}.$$

Similarly, we also have

$$s_2(C; (0, 1)) \leq 1 + 4 \cdot \frac{1}{2} + 4 \cdot \frac{1}{3} + \frac{1}{4} = \frac{55}{12}.$$

For the second case (see Figure 2(b)), we assume that $D = \{(0, 0), (0, 1), (2, 0)\}$ is a subset of C . As in the previous case (with the aid of the figure), it can be concluded that

$$s_2(C; (0, 0)) \leq \frac{13}{3} \text{ and } s_2(C; (0, 1)) \leq \frac{9}{2}.$$

It should be noted that the results of this example will be later used in the proof of Lemma 3.3.

3 The proof of the lower bound

For the rest of the section, assume that C is a 2-identifying code in G_H . It can be shown that $s_2(\mathbf{c}) \leq 5$ for all $\mathbf{c} \in C$ (see the proof of Lemma 3.4). This provides another approach to obtain the lower bound $D(C) \geq 1/5$, which was previously shown in [6]. In order to improve this lower bound, we need to consider the shares of codewords on average. Indeed, we can show that on average the share of a codeword is at most $19/4$. Therefore, as shown in Theorem 3.5, we prove that the density $D(C) \geq 4/19$.

The averaging process is done by introducing a *shifting scheme* designed to even out the shares among the codewords of C . (Notice that the shifting scheme can also be understood as a discharging method.) The rules of the shifting scheme are defined in Section 3.1. In Section 3.2, we introduce three lemmas, which state the following results:

- If $s_2(\mathbf{c}) > 19/4$ for some $\mathbf{c} \in C$, then at least $s_2(\mathbf{c}) - 19/4$ units of share is shifted from $\mathbf{c}$ to other codewords. (Lemma 3.4)
- If share is shifted to a codeword $\mathbf{c} \in C$, then $s_2(\mathbf{c}) \leq 19/4$ and the codeword $\mathbf{c}$ receives at most $19/4 - s_2(\mathbf{c})$ units of share. (Lemmas 3.2 and 3.3)

In other words, after the shifting is done, the share of each codeword is at most $19/4$. Using this fact, we are able to prove the main theorem (Theorem 3.5) of the paper according to which $D(C) \geq 4/19$. Finally, in Section 3.3, we provide the proofs of the lemmas.

3.1 The rules of the shifting scheme

The *rules* of the shifting scheme are illustrated in Figure 3. Translations, rotations and reflections (over the line passing vertically through $\mathbf{u}$) can be applied to each rule in such a way that the structure of the underlying graph G_H is preserved. In the rules, share is shifted as follows:

- In the rules 1, 2, 4 and 7, we shift $1/4$ units of share from $\mathbf{u}$ to $\mathbf{v}$.
- In the rule 3, we shift $1/6$ and $1/12$ units of share from $\mathbf{u}$ to $\mathbf{v}$ and $\mathbf{v}'$, respectively.
- In the rule 5, we shift $1/6$ units of share from $\mathbf{u}$ to $\mathbf{v}$.
- In the rules 6, 8, 9 and 10, we shift $1/12$ units of share from $\mathbf{u}$ to $\mathbf{v}$.

We also have the following modifications to the previous rules:

- If in the rules 1, 2 and 7 we have $\mathbf{u} + (0, -1) \in C$, then we only shift $1/12$ units of share from $\mathbf{u}$ to $\mathbf{v}$ and denote these new rules (respectively) by 1.1, 2.1 and 7.1. Moreover, in the rule 1.1, we shift $1/12$ units of share to $\mathbf{v}$ whether $(-3, 2)$ belongs to C or not.
- If in the rule 1 we have $\mathbf{u} + (-3, 2) \in C$, then then we shift $1/4$ units of share from $\mathbf{u}$ to $\mathbf{u} + (-1, 2)$ (no share is shifted to $\mathbf{v}$) and denote this new rule by 1.2.

- If in the rule 2 we have $\mathbf{u} + (-3, 1) \in C$, then we shift $1/6$ units of share from $\mathbf{u}$ to $\mathbf{v}$ and denote this new rule by 2.2.
- If in the rule 2 we have $\mathbf{u} + (1, 2) \in C$, then we shift $1/12$ units of share from $\mathbf{u}$ to $\mathbf{v}$ and denote this new rule by 2.3.




Figure 3: The rules of the shifting scheme illustrated. The black dots represent codewords and the white dots represent non-codewords. In the rules 7 and 8, at least one of the vertices marked with a white square is a codeword.

The modified share of a codeword $\mathbf{c} \in C$, which is obtained after the shifting scheme is applied, is denoted by $\bar{s}_2(\mathbf{c})$. The use of the rules is illustrated in the following example.

Example 3.1. Consider the codeword $\mathbf{c}$ with the surroundings as illustrated in Figure 4. The share of the codeword $\mathbf{c}$ is equal to 5. The rules 6, 8 and 9 apply to the codeword $\mathbf{c}$ and according to the rules $1/12$ units of share is shifted from $\mathbf{c}$ to $\mathbf{u}_1$, $\mathbf{u}_2$ and $\mathbf{u}_3$, respectively. (Recall that reflections and rotations can be applied to the constellations in Figure 3.) Hence, after the shifting scheme is applied, we have $\bar{s}_2(\mathbf{c}) = 19/4$ for the modified share. In order to ensure that also $\bar{s}_2(\mathbf{u}_i) \leq 19/4$ for any $i = 1, 2, 3$, we refer to the proofs of Lemmas 3.2 and 3.3.




Figure 4: An example of the use of the shifting rules.

3.2 The main theorem

The following three lemmas show that $\bar{s}_2(\mathbf{c}) \leq 19/4$ for all $\mathbf{c} \in C$. The proofs of the lemmas are postponed to Section 3.3.

Lemma 3.2. *Let $\mathbf{c} \in C$ be a codeword such that $\mathbf{c}$ is not adjacent to another codeword and share is shifted to $\mathbf{c}$ according to the previous rules. Then we have $\bar{s}_2(\mathbf{c}) \leq 19/4$.*

Lemma 3.3. *Let $\mathbf{c} \in C$ be a codeword such that $\mathbf{c}$ is adjacent to another codeword and share is shifted to $\mathbf{c}$ according to the previous rules. Then we have $\bar{s}_2(\mathbf{c}) \leq 19/4$.*

Lemma 3.4. *Let $\mathbf{c} \in C$ be a codeword such that no share is shifted to $\mathbf{c}$ according to the previous rules. Then we have $\bar{s}_2(\mathbf{c}) \leq 19/4$.*

As stated in the previous lemmas, we have $\bar{s}_2(\mathbf{c}) \leq 19/4$ for any $\mathbf{c} \in C$. Now we are ready to prove the main theorem of the paper.

Theorem 3.5. *If C is a 2-identifying code in the hexagonal grid G_H , then the density*

$$D(C) \geq \frac{4}{19}.$$

Proof. Assume that C is a 2-identifying code in G_H . Since each vertex $\mathbf{u} \in Q_{n-2}$ with $|I_2(\mathbf{u})| = i$ contributes the summand $1/i$ to $s_2(\mathbf{c})$ for each of the i codewords $\mathbf{c} \in B_2(\mathbf{u})$, we have

$$\sum_{\mathbf{c} \in C \cap Q_n} s_2(\mathbf{c}) \geq |Q_{n-2}|. \quad (1)$$

Furthermore, we have

$$\sum_{\mathbf{c} \in C \cap Q_n} s_2(\mathbf{c}) \leq \sum_{\mathbf{c} \in C \cap Q_n} \bar{s}_2(\mathbf{c}) + \frac{19}{4} |Q_{n+6} \setminus Q_n|, \quad (2)$$

because shifting shares inside Q_n does not affect the sum and each codeword in $Q_{n+6} \setminus Q_n$ can receive at most $19/4$ units of share (by Lemmas 3.2 and 3.3). Notice also that

codewords in Q_n cannot shift share to codewords outside Q_{n+6} . Therefore, combining the equations (1) and (2) with the fact that $\bar{s}_2(\mathbf{c}) \leq 19/4$ for any $\mathbf{c} \in C$, we obtain

$$\frac{|C \cap Q_n|}{|Q_n|} \geq \frac{4}{19} \cdot \frac{|Q_{n-2}|}{|Q_n|} - \frac{|Q_{n+6} \setminus Q_n|}{|Q_n|}.$$

Since $|Q_k| = (2k+1)^2$ for any positive integer k , it is straightforward to conclude from the previous inequality that the density $D(C) \geq 4/19$. $\square$

3.3 The proofs of the lemmas

In what follows, we provide the proofs of Lemmas 3.2, 3.3 and 3.4.

Proof of Lemma 3.2. Notice first that share can be shifted to the codeword $\mathbf{c}$ only according to the rules 3, 7, 7.1, 8, 9 and 10 since $\mathbf{c}$ is not adjacent to another codeword. The main idea of the following proof is to show that $\mathbf{c}$ cannot receive share according to two different rules. In each case, this observation then quite straightforwardly implies the claim.

Assume first that $\mathbf{c}$ receives share according to the rule 10. Without loss of generality, we may assume that the rule is applied as in Figure 3(j) (when $\mathbf{c} = \mathbf{v}$). Now the vertices $\mathbf{c} + (-2, -1)$, $\mathbf{c} + (0, 1)$ and $\mathbf{c} + (2, -1)$ belong to the code C . Since this is not the case with the other rules (see Figure 3), they cannot be applied to $\mathbf{c}$. Moreover, choosing $D = \{\mathbf{c}, \mathbf{c} + (-2, -1), \mathbf{c} + (0, 1), \mathbf{c} + (2, -1)\}$ in Lemma 2.1, we have $s_2(\mathbf{c}) \leq 9/2$. Thus, since according to the rule 10 share can be shifted to $\mathbf{c}$ only from $\mathbf{c} + (-2, -1)$, $\mathbf{c} + (0, 1)$ and $\mathbf{c} + (2, -1)$ and at most once for each of these codewords, we obtain that $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 3 \cdot 1/12 \leq 19/4$.

Assume then that share is shifted to $\mathbf{c}$ according to the rule 9. First of all, by the previous paragraph, the rule 10 cannot be applied to $\mathbf{c}$. Since now we have $I_2(\mathbf{c}) = \{\mathbf{c}\}$, it is immediate that $\mathbf{c}$ cannot receive share according to the rules 3, 7, 7.1 or 8. Moreover, it is easy to see that the rule 9 can be applied only once. Therefore, since we have $s_2(\mathbf{c}) \leq 14/3$ by Lemma 2.1, we obtain that $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/12 \leq 19/4$.

Assume that $\mathbf{c}$ receives share according to the rule 8 and that the rule is used as in Figure 3(h). Now, if the rule 7 or 7.1 was used, then there would exist two codewords in $B_2(\mathbf{c})$ such that the distance between them is equal to 4. By the constellation of the rule 8, this is impossible. Let us then show that neither the rule 3 can be used. Assume to the contrary that share is shifted to $\mathbf{c}$ according to the rule 3. Since there is a pair of adjacent codewords in the constellation of the rule 3, the vertex $\mathbf{c} + (0, 1)$ belongs to C . Furthermore, we have either $\mathbf{c} + (-1, 1) \in C$ or $\mathbf{c} + (1, 1) \in C$ (but not both). If $\mathbf{c} + (-1, 1) \in C$, then we have a contradiction since share is shifted from $\mathbf{c} + (3, 0)$, which is not a codeword in the constellation of the rule 8. On the other hand, if $\mathbf{c} + (1, 1) \in C$, then share is received from $\mathbf{c} + (-3, 0)$. This again leads to a contradiction since $\mathbf{c} + (-2, -1) \in C$. In conclusion, only the rule 8 can be applied to $\mathbf{c}$. Moreover, the rule 8 can be used at most once. If $\mathbf{c} + (-1, 1) \in C$ or $\mathbf{c} + (1, 1) \in C$, then $s_2(\mathbf{c}) \leq 55/12$ or $s_2(\mathbf{c}) \leq 23/6$ by Lemma 2.1, respectively. Hence, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/12 \leq 56/12 \leq 19/4$.




Figure 5: The cases of the proof of Lemma 3.3 illustrated.

Assume that $\mathbf{c}$ receives share according to the rule 3 and that the rule is used as in Figure 3(c) (when $\mathbf{c} = \mathbf{v}'$). By the previous considerations, we know that share cannot be shifted to $\mathbf{c}$ according to the rules 8, 9 and 10. Let us then show that neither the rules 7 or 7.1 can be applied to $\mathbf{c}$. Assume to the contrary that $\mathbf{c}$ receives share according to the rule 7. Now $\mathbf{c}$ can receive share only from the vertices $\mathbf{c} + (-1, -2)$, $\mathbf{c} + (1, -2)$, $\mathbf{c} + (3, 0)$, $\mathbf{c} + (2, 1)$, $\mathbf{c} + (-3, 0)$ and $\mathbf{c} + (-2, 1)$. Now we have the following observations:

- Since $\mathbf{c} + (-1, -2) \in C$ share cannot be shifted from $\mathbf{c} + (-1, -2)$ and $\mathbf{c} + (1, -2)$.
- Since $\mathbf{c} + (-2, 0) \in C$ share cannot be shifted from $\mathbf{c} + (-3, 0)$ and $\mathbf{c} + (-2, 1)$.
- Since $\mathbf{c} + (-1, -1) \notin C$ and $\mathbf{c} + (1, -1) \notin C$ share cannot be shifted from $\mathbf{c} + (3, 0)$ and $\mathbf{c} + (2, 1)$.

A contradiction now follows from these facts. Hence, only the rule 3 can be applied to $\mathbf{c}$. Moreover, it is easy to see that share can be shifted to $\mathbf{c}$ at most twice according to the rule 3. Therefore, since we have $s_2(\mathbf{c}) \leq 9/2$ by Lemma 2.1, we obtain that $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 2 \cdot 1/12 \leq 19/4$.

Finally, assume that $\mathbf{c}$ receives share according to the rule 7 or 7.1 and that the rule is used as in Figure 3(g). As shown above, other rules cannot be applied to $\mathbf{c}$. Moreover, the rules 7 and 7.1 can be applied to $\mathbf{c}$ only once. Furthermore, if $\mathbf{c} + (2, 0) \in C$ or $\mathbf{c} + (1, 1) \in C$, then we have $s_2(\mathbf{c}) \leq 23/6$ or $s_2(\mathbf{c}) \leq 53/12$ by Lemma 2.1, respectively. Thus, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/4 \leq 14/3 \leq 19/4$. $\square$

Proof of Lemma 3.3. Since $\mathbf{c}$ is adjacent to another codeword, it is immediate that $\mathbf{c}$ can receive share only according to the rules 1, 1.1, 1.2, 2, 2.1, 2.2, 2.3, 3, 4, 5 and 6. The proof of the lemma is now divided into three cases depending on the number of codewords adjacent to $\mathbf{c}$.

1) Assume first that $\mathbf{c}$ is adjacent to exactly one codeword. Without loss of generality, we may assume that $\mathbf{c} = (0, 0)$ and the adjacent codeword is $(0, 1)$. (Notice that $(-1, 0) \notin C$ and $(1, 0) \notin C$.) Now the only possibilities for $\mathbf{c}$ to receive share is from $(-3, 0)$ or $(3, 0)$

(the rule 1.2) and from the vertices that belong to $S_1 = \{(5, 1), (4, 1), (3, 1), (3, 2), (2, 2)\}$ and $S_2 = \{(-5, 1), (-4, 1), (-3, 1), (-3, 2), (-2, 2)\}$. These observations are illustrated in Figure 5(a).

Consider then more closely the set S_1 . In what follows, we show that at most $1/4$ units of share is shifted from the vertices of S_1 to $\mathbf{c}$. Notice that if share is shifted only from one vertex of S_1 , then we are immediately done. Observe then that if $\mathbf{u} \in S_1$ shifts share to $\mathbf{c}$ according to the rules 1–6 or their modifications, then we have $I_2(\mathbf{u}) = \{\mathbf{u}\}$. Therefore, if two vertices of S_1 shift share, then one of these vertices is $(4, 1)$ or $(5, 1)$. Assume that share is shifted from $(4, 1)$ according to the rule 5 (see Figure 3(e)). Now we have $C \cap S_1 = \{(2, 2), (4, 1)\}$ and $(4, 3) \in C$. Therefore, since at most $1/12$ units of share is shifted from $(4, 1)$ according to the rule 2.1, we obtain that no more than $1/6 + 1/12 = 1/4$ units of share can be shifted from S_1 to $\mathbf{c}$. Similarly, if share is shifted from $(5, 1)$ according to the rule 6, then it can be shown that $\mathbf{c}$ receives at most $1/4$ units of share from S_1 . In conclusion, at most $1/4$ units of share is shifted from the vertices of S_1 to $\mathbf{c}$. Analogously, this statement also holds for the vertices of S_2 .

Assume that the rule 1.2 is used. (Clearly, this rule can be used only once.) Without loss of generality, we may assume that $(-3, 0) \in C$ and $(1, -1) \in C$. Therefore, choosing $D = \{\mathbf{c}, (0, 1), (-3, 0), (1, -1)\}$ in Lemma 2.1, we obtain that $s_2(\mathbf{c}) \leq 15/4$. Thus, since the codewords in each of the sets S_1 and S_2 can shift at most $1/4$ units of share to $\mathbf{c}$, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 3 \cdot 1/4 \leq 9/2$. Assume then that the rule 1.2 cannot be applied to $\mathbf{c}$. Since $\mathbf{c}$ and $(0, 1)$ are 2-separated by C , there exists at least one codeword in the symmetric difference $B_2(\mathbf{c}) \triangle B_2(0, 1)$. Thus, without loss of generality, we may assume that $(1, -1) \in C$, $(1, 2) \in C$, $(2, 0) \in C$ or $(2, 1) \in C$. The first part of the proof is then concluded by the following four cases:

- Assume that $(1, -1) \in C$. Now, by the first case of Example 2.2, we know that $s_2(\mathbf{c}) \leq 17/4$. Therefore, since share is shifted to $\mathbf{c}$ only from the vertices of the sets S_1 and S_2 , we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 2 \cdot 1/4 \leq 19/4$.
- Assume that $(1, 2) \in C$. It is straightforward to verify that share can be shifted to $\mathbf{c}$ only from $(-3, 1)$, $(-2, 2)$, $(-3, 2)$, $(-4, 1)$ and $(-5, 1)$ according to the rules 1.1, 2.3, 3, 5 and 6, respectively. Moreover, it is easy to see that at most one of these rules can be used (and only once). Thus, the codeword $\mathbf{c}$ receives at most $1/6$ units of share. Hence, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/6 \leq 19/4$ since $s_2(\mathbf{c}) \leq 55/12$ by the first case of Example 2.2 (and obvious symmetrical argument).
- Assume that $(2, 0) \in C$. Consider then more closely the vertices of S_1 . Then it is easy to conclude that the vertices $(5, 1)$, $(4, 1)$ and $(3, 1)$ cannot shift share to $\mathbf{c}$. Hence, only either $(3, 2)$ according to the rule 3 or $(2, 2)$ according to the rule 2.2 (but not both) is capable of shifting share to $\mathbf{c}$. In both cases, $\mathbf{c}$ receives at most $1/6$ units of share. By the second case of Example 2.2, we know that $s_2(\mathbf{c}) \leq 13/3$. Therefore, since at most $1/4$ units of share is received from S_2 , we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/4 + 1/6 \leq 19/4$.

- Assume that $(2, 1) \in C$. By the second case of Example 2.2 (and symmetry), we obtain that $s_2(\mathbf{c}) \leq 9/2$. Since now share can be shifted to $\mathbf{c}$ only from S_2 , we obtain that $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 1/4 \leq 19/4$.

2) Assume that $\mathbf{c}$ is adjacent to exactly two codewords. Without loss of generality, we may assume that $\mathbf{c} = (0, 0)$ and that the adjacent codewords are $(-1, 0)$ and $(0, 1)$. Let then S_3 and S_4 be sets which are obtained by rotating respectively the sets S_1 and S_2 by $2\pi/3$ (counter-clockwise in the honeycomb representation) around the origin. Again the codewords in each of these sets S_i can shift at most $1/4$ units of share to $\mathbf{c}$. In addition to the previous ones, at most $1/4$ units of share can also be shifted to $\mathbf{c}$ from either $(2, -1)$ or $(3, 0)$ (but not both) according to the rule 1.2. These observations are illustrated in Figure 5(b).

Assume first that share is shifted to $\mathbf{c}$ according to the rule 1.2. Without loss of generality, we may assume that $\mathbf{c}$ receives share from the vertex $(3, 0)$. Then we immediately have $(3, 0) \in C$ and $(-1, -1) \in C$. Therefore, as above, we have $s_2(\mathbf{c}) \leq 15/4$ by Lemma 2.1. Furthermore, since $(-1, -1) \in C$, the codeword $\mathbf{c}$ does not receive share from the set S_4 . Thus, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 4 \cdot 1/4 \leq 19/4$.

Assume then that the rule 1.2 is not used. Since the vertices $\mathbf{c}$ and $(0, 1)$ are 2-separated by C , there exists a codeword in $B_r(\mathbf{c}) \triangle B_r(0, 1)$. If $(1, -1) \in C$ or $(2, 0) \in C$, then $s_2(\mathbf{c}) \leq 53/15$ (by Lemma 2.1) and we are immediately done since at most 1 unit of share can be shifted to $\mathbf{c}$ from the union of the sets S_i . If $(-2, 1) \in C$, then $s_2(\mathbf{c}) \leq 21/5$ and we are done since share cannot be shifted to $\mathbf{c}$ from S_2 and S_3 . If $(-1, -1) \in C$ or $(-2, 0) \in C$, then $s_2(\mathbf{c}) \leq 79/20$ (by Lemma 2.1) and we are again done (since share is not shifted from S_3). Hence, we may assume that $(-1, 2) \in C$, $(1, 2) \in C$ or $(2, 1) \in C$. Analogously, it can also be assumed that $(-3, 0) \in C$, $(-2, -1) \in C$ or $(0, -1) \in C$ since $\mathbf{c}$ and $(-1, 0)$ are 2-separated by C . Thus, at most two of the sets S_i can shift share to $\mathbf{c}$. Therefore, since $s_2(\mathbf{c}) \leq 17/4$ (choose $D = \{\mathbf{c}, (-1, 0), (0, 1)\}$ in Lemma 2.1), we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 2 \cdot 1/4 \leq 19/4$.

3) Finally, assume that all the vertices adjacent to $\mathbf{c}$ are codewords, i.e. $(-1, 0) \in C$, $(0, 1) \in C$ and $(1, 0) \in C$ (see Figure 5(c)). Notice that now the rule 1.2 cannot be used. Since there again exists a codeword in $B_r(\mathbf{c}) \triangle B_r(0, 1)$, it is easy to conclude as above that at most $5 \cdot 1/4$ units of share is shifted to $\mathbf{c}$. Therefore, since $s_2(\mathbf{c}) \leq 67/20$ by Lemma 2.1, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) + 5/4 \leq 19/4$. This completes the proof of the lemma. $\square$

Proof of Lemma 3.4. Without loss of generality, we may assume that $\mathbf{c} = (0, 0)$. Assume first that $|I_2(\mathbf{c})| \geq 2$. If now $\mathbf{c}$ is adjacent to another codeword, then $\bar{s}_2(\mathbf{c}) \leq 19/4$ (by Lemma 2.1). Hence, we may assume that $(-1, 0), (1, 0), (0, 1) \notin C$. Let then $(2, 0)$ be a codeword of C . Since the vertices $\mathbf{c}$ and $(1, 0)$ are 2-separated by C , there is at least one codeword in the symmetric difference $B_2(\mathbf{c}) \triangle B_2(1, 0)$ (see Figure 6). Therefore, by Lemma 2.1, it is straightforward (albeit tedious) to verify that in all possible cases $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) \leq 19/4$. Indeed, we can choose in Lemma 2.1 the set D to consist of $\mathbf{c}$, $(2, 0)$ and a codeword in the symmetric difference $B_2(\mathbf{c}) \triangle B_2(1, 0)$.




Figure 6: The symmetric difference $B_2(\mathbf{c}) \triangle B_2(1, 0)$ consists of the squared vertices. One of these vertices is a codeword.




Figure 7: The sets A_1 , A_2 , A_3 , A'_1 , A'_2 and A'_3 illustrated.

From now on, we may assume that $I_2(\mathbf{c}) = \{\mathbf{c}\}$. In what follows, we denote $A_1 = \{(-1, 1), (0, 1), (1, 1)\}$, $A_2 = \{(-2, 0), (-1, 0), (-1, -1)\}$, $A_3 = \{(1, -1), (1, 0), (2, 0)\}$, $A'_1 = \{(-1, 2), (1, 2)\}$, $A'_2 = \{(-3, 0), (-2, -1)\}$ and $A'_3 = \{(2, -1), (3, 0)\}$. These sets are illustrated in Figure 7. The proof of the lemma now divides into three cases depending on the number of codewords in the set $\{(-2, 1), (2, 1), (0, -1)\}$

1) Assume first that $(-2, 1), (2, 1), (0, -1) \notin C$. Since the vertices $\mathbf{c}$, $(-1, 0)$, $(1, 0)$ and $(0, 1)$ are 2-separated by C , each one of the sets A'_1 , A'_2 and A'_3 contains at least one codeword. Hence, each of the sets A_1 , A_2 and A_3 contains a vertex whose I -set contains at least three codewords. Indeed, if for example $(-1, 2) \in C$, then $(-1, 1)$ or $(0, 1)$ is such a vertex in A_1 . Therefore, we obtain that $s_2(\mathbf{c}) \leq 1 + 6 \cdot 1/2 + 3 \cdot 1/3 = 5$.

Consider then the set A'_1 that contains at least one codeword as stated above. Assume first that both $(-1, 2) \in C$ and $(1, 2) \in C$. If the vertex $(0, 2)$ also belongs to C , then the I -sets of all the vertices in A_1 have size at least 3. Then there are at least 5 vertices in $B_2(\mathbf{c})$ which are 2-covered by at least 3 codewords. Therefore, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) \leq 1 + 4 \cdot 1/2 + 5 \cdot 1/3 = 14/3$ (and we are done). Hence, suppose that $(0, 2) \notin C$. Furthermore, assume first that $(-2, 2) \in C$. If also $(-3, 1) \in C$, then $|I_2(0, 1)| \geq 3$ and $|I_2(-1, 1)| \geq 4$. Thus, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) \leq 1 + 5 \cdot 1/2 + 3 \cdot 1/3 + 1/4 = 19/4$. On the other hand, if $(-3, 1) \notin C$, then the rule 2.3 can be applied to $\mathbf{c}$ and we obtain that $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) - 1/12 \leq 29/6 - 1/12 \leq 19/4$. Hence, we may assume that $(-2, 2) \notin C$ and




Figure 8: The surroundings of the codeword $\mathbf{c}$ illustrated in (a) the first and (b) the second part of the proof of Lemma 3.4.

$(-2, 2) \notin C$ (by symmetry). Since the vertices $(-1, 2)$ and $(1, 2)$ are 2-separated by C , at least one of the vertices $(-3, 2)$, $(-2, 3)$, $(2, 3)$ and $(3, 2)$ belongs to C . Hence, we can shift at least $1/4$ units of share from $\mathbf{c}$ according to the rule 7. Therefore, we have $\bar{s}_2(\mathbf{c}) \leq s_2(\mathbf{c}) - 1/4 \leq 19/4$.

By the considerations above, we may without loss of generality assume that $(-1, 2) \in C$ and $(1, 2) \notin C$. If $(0, 2) \in C$, then $1/4$ units of share can be shifted from $\mathbf{c}$ according to the rule 1 or 1.2, and we are done. Thus, suppose that $(0, 2) \notin C$. Assume then that $(-2, 2) \in C$. If the rule 2 applies to $\mathbf{c}$, then we are immediately done ($1/4$ units of share is shifted from $\mathbf{c}$). On the other hand, if $(-3, 1) \in C$, then by the fact that $|I_2(-1, 1)| \geq 4$ we have $s_2(\mathbf{c}) \leq 59/12$ and we are again done since at least $1/6$ units of share is shifted from $\mathbf{c}$ according to the rule 2.2. Hence, assume that $(-2, 2) \notin C$. Since $(0, 1)$ and $(-1, 1)$ are 2-separated by C , the vertex $(-3, 1)$ belongs to C . Now, if $(-3, 2) \in C$, then $1/4$ units of share can be shifted from $\mathbf{c}$ according to the rule 3 and we are done. Thus, we may assume that $(-3, 2) \notin C$. If $(-4, 1) \in C$ and $(-3, 0) \notin C$, then the rule 4 can be applied to $\mathbf{c}$ and we are done. Furthermore, if $(-4, 1) \in C$, and $(-3, 0) \in C$, then instead of the set A'_1 consider the set A'_3 . Repeating the previous arguments for the set A'_3 , we obtain that $\bar{s}_2(\mathbf{c}) \leq 19/4$ also in this case. Thus, we may assume that $(-4, 1) \notin C$. The surroundings of the codeword $\mathbf{c}$ (obtained above) is illustrated in Figure 8(a).

The previous reasoning also applies when we consider the sets A'_2 and A'_3 instead of A'_1 . This leads straightforwardly to the observation that we have only two different neighbourhoods of $\mathbf{c}$ (up to rotations and reflections). These neighbourhoods are illustrated in Figure 9.

Consider first the case in Figure 9(a). In what follows, we show that $\mathbf{c}$ shifts $1/12$ units of share to $(-1, 2)$ or $(1, 3)$, or that we originally have at least 2 vertices in A_1 such that their I -sets are at least of size 3. In both cases, the (maximum) share of $\mathbf{c}$ is reduced by at least $1/12$. (Actually, in the latter case, the share is reduced by $1/6$.) This observation can then be generalized to the (other two) symmetrical cases implying that $\bar{s}_2(\mathbf{c}) \leq 5 - 3 \cdot 1/12 = 19/4$. If $(2, 2) \in C$, then $(-1, 1) \in A_1$ and $(1, 1) \in A_1$ are 2-covered by three codewords. Hence, we may assume that $(2, 2) \notin C$. If we can shift $1/12$ units of




Figure 9: Two cases of the proof of Lemma 3.4 illustrated.

share from $\mathbf{c}$ to $(-1, 2)$ according to the rule 8, then we are immediately done. Therefore, we may assume that $(-2, 3) \notin C$ and $(0, 3) \notin C$. Since $(-1, 2)$ and $(1, 2)$ are 2-separated by C , the vertex $(2, 3)$ belongs to C . Furthermore, since $(-1, 2)$ and $(0, 2)$ are 2-separated by C , at least one of the vertices $(-1, 3)$ and $(1, 3)$ is a codeword. Therefore, at least $1/12$ units of share can be shifted from $\mathbf{c}$ according to the rule 6 or 9.

Consider then the case in Figure 9(b). Let us now show that $\mathbf{c}$ shifts at least $1/6$ units of share to $(-1, -2)$ or $(1, -2)$, or that we originally have more than 2 vertices in $A_2 \cup A_3$ such that their I -sets are at least of size 3. In both cases, the (maximum) share of $\mathbf{c}$ is reduced by at least $1/6$. This result together with the observation in the previous paragraph, then implies that $\bar{s}_2(\mathbf{c}) \leq 5 - 1/6 - 1/12 = 19/4$. If now $(0, -2) \in C$, then we know that $(-2, 0) \in A_2$, $(2, 0) \in A_3$, and at least one of the vertices $(-1, -1) \in A_2$ and $(1, -1) \in A_3$ are such that their I -sets are at least of size 3. Hence, we may assume that $(0, -2) \notin C$. Since $\mathbf{c}$, $(-1, -1)$ and $(1, -1)$ are 2-separated by C , the vertices $(-2, -2)$ and $(2, -2)$ belong to C . Furthermore, since $(0, -1)$ is 2-covered by a codeword of C , we have $(-1, -2) \in C$ or $(1, -2) \in C$. Therefore, we can shift at least $1/6$ units of share to $(-1, -2)$ or $(1, -2)$ according to the rule 5. In conclusion, if $\mathbf{c}$ is a codeword such that $I_2(\mathbf{c}) = \{\mathbf{c}\}$ and $(-2, 1), (2, 1), (0, -1) \notin C$, then we have $\bar{s}_2(\mathbf{c}) \leq 19/4$.

2) Assume then that $\mathbf{c}$ is a codeword such that $I_2(\mathbf{c}) = \{\mathbf{c}\}$, $(-2, 1) \notin C$, $(2, 1) \notin C$ and $(0, -1) \in C$. Now, by considering the vertices $(-1, -1) \in A_2$, $(-1, 0) \in A_2$, $(1, -1) \in A_3$ and $(1, 0) \in A_3$, we obtain that there are at least 3 vertices in $A_2 \cup A_3$ which are 2-covered by at least 3 codewords. Thus, since there is also one such vertex in A_1 , we have $s_2(\mathbf{c}) \leq 1 + 5 \cdot 1/2 + 4 \cdot 1/3 = 29/6$. Assume first that both $(-1, 2)$ and $(1, 2)$ belong to C . If $(-2, 2) \in C$, $(0, 2) \in C$ or $(2, 2) \in C$, there are at least 2 vertices in A_1 with the I -set of size at least 3 and, therefore, we have $s_2(\mathbf{c}) \leq 19/4$. Hence, according to the rule 7.1, $1/12$ units of share can be shifted from $\mathbf{c}$ (since $(-1, 2)$ and $(1, 2)$ are 2-separated by C) and we are done. Thus, without loss of generality, we may assume that $(-1, 2) \in C$ and $(1, 2) \notin C$. If now $(-3, 1) \in C$ and $(-2, 2) \in C$, then $|I_2(-1, 1)| \geq 4$ and there is no problem since $s_2(\mathbf{c}) \leq 1 + 5 \cdot 1/2 + 3 \cdot 1/3 + 1/4 = 19/4$. Furthermore, if the rules 1.1 or 2.1 can be used, then we are again done. Thus, we may assume that $(-2, 2) \notin C$ and $(0, 2) \notin C$. Therefore, since $(-1, 1)$ and $(0, 1)$ are 2-separated by C , we obtain that

$(-3, 1) \in C$. We have now arrive at the constellation illustrated in Figure 8(b). If now one of the vertices $(-4, 0)$, $(-3, 0)$, $(-3, -1)$, $(-2, -1)$, $(0, -2)$ and $(2, -1)$ belongs to C , then by Lemma 2.1 we immediately have $s_2(\mathbf{c}) \leq 19/4$. Hence, we may assume that none of these vertices belongs to C . Thus, since $(-1, 0)$, $(-1, -1)$ and $(1, 0)$ are 2-separated by C , we have $(-2, -2) \in C$ and $(3, 0) \in C$. If now $(3, -1) \in C$ or $(4, 0) \in C$, then again $s_2(\mathbf{c}) \leq 19/4$ and we are done. Therefore, since $(-1, 0)$ and $(1, -1)$ are 2-separated by C , we have $(2, -2) \in C$ and $1/12$ units of share can be shifted from $\mathbf{c}$ to $(0, -1)$ according to the rule 10. Hence, we have $\bar{s}_2(\mathbf{c}) \leq 19/4$.

3) Finally, assume that $\mathbf{c}$ is a codeword such that $I_2(\mathbf{c}) = \{\mathbf{c}\}$, and that at least two of the vertices $(-2, 1)$, $(2, 1)$ and $(0, -1)$ belong to C . Then, by Lemma 2.1, we have $s_2(\mathbf{c}) \leq 14/3$. This observation completes the proof of the lemma. $\square$

Acknowledgements

We would like to thank the anonymous referee for providing constructive comments to improve the paper.

References

- [1] I. Charon, I. Honkala, O. Hudry, and A. Lobstein. General bounds for identifying codes in some infinite regular graphs. *Electron. J. Combin.*, 8(1):#R39, 21 pp., 2001.
- [2] I. Charon, O. Hudry, and A. Lobstein. Identifying codes with small radius in some infinite regular graphs. *Electron. J. Combin.*, 9(1):#R11, 25 pp., 2002.
- [3] G. Cohen, I. Honkala, A. Lobstein, and G. Zémor. Bounds for codes identifying vertices in the hexagonal grid. *SIAM J. Discrete Math.*, 13(14):492–504, 2000.
- [4] D. W. Cranston and G. Yu. A new lower bound on the density of vertex identifying codes for the infinite hexagonal grid. *Electron. J. Combin.*, 16(1):#R113, 16 pp., 2009.
- [5] M. G. Karpovsky, K. Chakrabarty, and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Trans. Inform. Theory*, 44(2):599–611, 1998.
- [6] R. Martin and B. Stanton. Lower bounds for identifying codes in some infinite grids. *Electron. J. Combin.*, 17(1):#R122, 16 pp., 2010.
- [7] P. J. Slater. Fault-tolerant locating-dominating sets. *Discrete Math.*, 249(1–3):179–189, 2002.