

Integral Cayley multigraphs over Abelian and Hamiltonian groups

Matt DeVos*

Department of Mathematics
Simon Fraser University
Burnaby, B.C. V5A 1S6

mdevos@sfu.ca

Roi Krakovski†

Department of Mathematics
Simon Fraser University
Burnaby, B.C. V5A 1S6

roikr@cs.bgu.ac.il

Bojan Mohar‡§

Department of Mathematics
Simon Fraser University
Burnaby, B.C. V5A 1S6

mohar@sfu.ca

Azhvan Sheikh Ahmady

Department of Mathematics
Simon Fraser University
Burnaby, B.C. V5A 1S6

ssheikha@sfu.ca

Submitted: Sep 24, 2012; Accepted: Jun 23, 2013; Published: Jun 30, 2013

Mathematics Subject Classifications: 05C50, 05E15

Abstract

It is shown that a Cayley multigraph over a group G with generating multiset S is integral (i.e., all of its eigenvalues are integers) if S lies in the integral cone over the boolean algebra generated by the normal subgroups of G . The converse holds in the case when G is abelian. This in particular gives an alternative, character-theoretic proof of a theorem of Bridges and Mena (1982). We extend this result by providing a necessary and sufficient condition for a Cayley multigraph over a hamiltonian group to be integral, in terms of character sums and the structure of the generating set.

1 Introduction

A graph X is said to be *integral* if all eigenvalues of the adjacency matrix of X are integers. This property was first defined by Harary and Schwenk [6] who suggested the problem of

*Supported in part by an NSERC Discovery Grant (Canada) and a Sloan Fellowship.

†Supported in part by postdoctoral support at the Simon Fraser University.

‡Supported in part by an NSERC Discovery Grant (Canada), by the Canada Research Chair program, and by the Research Grant P1-0297 of ARRS (Slovenia).

§On leave from: IMFM & FMF, Department of Mathematics, University of Ljubljana, Ljubljana, Slovenia.

determining which graphs satisfy it. Although this problem appears far too complicated in general, there is a broad literature on it with numerous interesting successes.

Throughout we shall assume that G is a finite group written multiplicatively. For a subset $S \subseteq G$ which is inverse-closed (i.e., $S = S^{-1}$), we define the *Cayley graph* of G over S , denoted $\text{Cay}(G, S)$, to be the graph with vertex set G and $x, y \in G$ adjacent if $xy^{-1} \in S$. In what follows, we shall consider the multigraph version. Recall that a *multiset* is a set S together with multiplicity function $\mu_S : S \rightarrow \mathbb{N}$, where $\mu_S(x)$ is a positive integer for every $x \in S$ (counting “how many times x occurs in the multiset”). We set $\mu_S(x) = 0$ for $x \notin S$. A multiset S of group elements is *inverse-closed* if $\mu_S(s) = \mu_S(s^{-1})$ for every $s \in S$. If S is an inverse-closed multiset of elements of a group G , the *Cayley multigraph* $\text{Cay}(G, S)$ is defined as above except that it is a multigraph and the number of edges joining x, y in $\text{Cay}(G, S)$ is equal to $\mu_S(xy^{-1})$.

Any collection \mathcal{S} of subsets of G naturally generates a boolean algebra of sets by taking all sets of elements of G that are obtained from the sets in \mathcal{S} by repeatedly taking unions, intersections and the symmetric difference. We let $\mathcal{B}(G)$ denote the boolean algebra generated by the normal subgroups of G . When extending this notion to multisets, we include multiset operations. Formally, we take all atoms of the Boolean algebra $\mathcal{B}(G)$ and take all multisets that can be expressed as non-negative integer combinations of these atoms. This defines the collection $\mathcal{C}(G)$ of multisets that is called the *integral cone* over $\mathcal{B}(G)$.

The following theorem gives a complete characterization of which Cayley multigraphs over abelian groups are integral.

Theorem 1 (Bridges, Mena [1]). *If G is an abelian group, then $\text{Cay}(G, S)$ is integral if and only if $S \in \mathcal{C}(G)$.*

Although the result of Bridges and Mena was originally stated for simple Cayley graphs, these authors proved the multigraph version. For each group element $g \in G$, let A_g denote the permutation matrix (indexed by $G \times G$) associated with g and for a set $S \subseteq G$ let $A_S = \sum_{s \in S} A_s$. Bridges and Mena proved that for an abelian group G , a complex linear combination of the matrices $\{A_g : g \in G\}$ is a rational matrix with rational eigenvalues if and only if it is a rational combination of the matrices $\{A_Q : Q \text{ is an atom of } \mathcal{B}(G)\}$.

There has been some recent interest in finding a new proof of this result. Notably, So [9] found a new proof in the special case when G is a cyclic group, and Klotz and Sander [8] found a new proof of the “if” direction for all abelian groups. Our initial goal was to give a new proof of Theorem 1. Our proof is based on characters, and is a fairly direct generalization of that given by So. However, our approach generalizes to non-abelian groups and enables us to consider more general classes of groups.

First, we obtain a sufficient condition for integrality of Cayley multigraphs over arbitrary groups.

Theorem 2. *For every finite group G and every $S \in \mathcal{C}(G)$, the Cayley multigraph $\text{Cay}(G, S)$ is integral.*

In the course of proving Theorem 2, we obtain a new proof of Theorem 1 based on character theory.

It is worth mentioning that having used normal subgroups and not all subgroups in the definition of $\mathcal{B}(G)$ and $\mathcal{C}(G)$ is necessary for Theorem 2 to hold. There are examples (e.g. dihedral groups) where using a subgroup in the generating set does not yield integral Cayley graphs.

By Theorem 1, the converse statement to Theorem 2 holds in abelian groups. In the second part of the paper we investigate to what extent the converse would hold in some other groups. As a natural candidate, we have decided to consider *dedekind groups*, i.e. groups whose every subgroup is normal. Every abelian group is dedekind; non-abelian dedekind groups are also called *hamiltonian groups*, and they have a simple characterization that is due to Baer, cf. [5, Theorem 12.5.4].

Theorem 3. *A finite group is hamiltonian if and only if it can be written as a direct product $Q_8 \times A$, where Q_8 is the group of quaternions and A is an abelian group without elements of order 4.*

We provide sufficient and necessary conditions for integrality of the spectra of Cayley multigraphs over such groups (Theorem 11). By using this characterization, we show that integrality of Cayley graphs over hamiltonian groups is easy to decide in certain special cases, while it leads to challenging combinatorial problems in some other special cases. See Sections 5 and 6 for more details.

2 B -integrality

If \mathcal{G} is a collection of graphs on the common vertex set V and B is an orthogonal basis of \mathbb{C}^V , then we say that \mathcal{G} is *B -integral* if for every X in \mathcal{G} , B is a set of eigenvectors for X and all eigenvalues of X are integral. Equivalently, if $A(X)$ denotes the adjacency matrix of X , then $A(X)B = B\Lambda$, where Λ is a diagonal matrix with integer entries (and B is viewed as a matrix whose columns are the vectors from B). If X and Y are (simple) graphs on the same vertex set V , then we denote by $X \cup Y$ the simple graph on V in which vertices u, v are adjacent if and only if they are adjacent in X or in Y (or in both). For any family of graphs \mathcal{G} on a common vertex set, we let $U(\mathcal{G})$ be the closure of \mathcal{G} under the operation \cup . We begin with an easy lemma.

- Lemma 4.** (a) *If X is B -integral and $\mathbb{1} \in B$, then \overline{X} is B -integral.*
 (b) *If X and $X \cap Y$ are B -integral, then $X \cap \overline{Y}$ is B -integral.*
 (c) *If \mathcal{G} is an intersection-closed family of B -integral graphs, then $U(\mathcal{G})$ is B -integral.*

Proof. To prove (b), observe that $A(X \cap \overline{Y}) = A(X) - A(X \cap Y)$, thus

$$A(X \cap \overline{Y})B = A(X)B - A(X \cap Y)B = B\Lambda_1 - B\Lambda_2 = B(\Lambda_1 - \Lambda_2).$$

Since Λ_1 and Λ_2 are integral, so is their difference, hence $X \cap \overline{Y}$ is B -integral.

If $\mathbb{1} \in B$, then the complete graph K_n is B -integral since $A(K_n)$ has eigenvalue $n-1$ with eigenvector $\mathbb{1}$ and eigenvalue -1 with eigenspace $\mathbb{1}^\perp$. Thus (a) follows from (b) by taking K_n and X playing the role of X and Y , respectively.

By observing that $A(X \cup Y) = A(X) + A(Y) - A(X \cap Y)$, a proof similar to the above proof of (b) shows that (c) holds. \square

For any set of graphs \mathcal{G} on a common vertex set V , we let $\mathcal{B}(\mathcal{G})$ denote the set of all graphs on V which may be expressed using members of \mathcal{G} and the operations \cap , \cup , and complement.

Lemma 5. *Let \mathcal{G} be an intersection-closed family of B -integral graphs and assume that $\mathbb{1} \in \mathcal{B}$. Then $\mathcal{B}(\mathcal{G})$ is B -integral.*

Proof. Set $\mathcal{G}_0 = \mathcal{G}$ and for every $k \in \mathbb{N}$ recursively define

$$\mathcal{G}_{k+1} = \{X_1 \cap \cdots \cap X_n : \text{either } X_i \in \mathcal{G}_k \text{ or } \overline{X_i} \in \mathcal{G}_k \text{ for every } 1 \leq i \leq n\}.$$

It is immediate that each \mathcal{G}_k is intersection-closed and it follows from De Morgan's law that $\mathcal{B}(\mathcal{G}) = \cup_{k=0}^{\infty} \mathcal{G}_k$. To complete the proof, we shall show, by induction on k , that every graph in \mathcal{G}_k is B -integral. As a base, we observe that this holds for $k = 0$ by assumption. For the inductive step, let X be a graph in \mathcal{G}_{k+1} and suppose that $X = X_1 \cap \cdots \cap X_\ell \cap \overline{Y_1} \cap \cdots \cap \overline{Y_m}$ where $X_1, \dots, X_\ell, Y_1, \dots, Y_m \in \mathcal{G}_k$. Then we have

$$X = (\cap_{i=1}^{\ell} X_i) \cap \overline{(\cup_{j=1}^m Y_j)}.$$

Since \mathcal{G}_k is intersection-closed and $X' = X_1 \cap \cdots \cap X_\ell \in \mathcal{G}_k$ and $X' \cap (\cup_{j=1}^m Y_j) = (\cup_{j=1}^m (X' \cap Y_j)) \in \mathcal{U}(\mathcal{G}_k)$, it follows from Lemma 4 that X is B -integral, as desired. \square

Lemma 6. *Let S and T be inverse-closed multisets of a group G . If $gT = Tg$ (equality holding as multisets) for every $g \in G$, then the adjacency matrices of Cayley multigraphs $\text{Cay}(G, S)$ and $\text{Cay}(G, T)$ commute.*

Proof. Let A_S and A_T be the adjacency matrices of both Cayley multigraphs, and let $g, h \in G$. Since S and T are inverse-closed, we have

$$\begin{aligned} (A_S A_T)_{g,h} &= \sum_{x \in G} \mu_S(gx^{-1}) \mu_T(xh^{-1}) = \sum_{x \in G} \mu_{Sg}(x) \mu_{Th}(x) \\ &= \sum_{x \in G} \mu_{Sg}(xg) \mu_{Th}(xg) = \sum_{x \in G} \mu_S(x) \mu_{Thg^{-1}}(x). \end{aligned}$$

Taking a similar expression for $A_T A_S$ and using the fact that S and T are inverse-closed, we derive

$$\begin{aligned} (A_T A_S)_{g,h} &= \sum_{x \in G} \mu_{Tgh^{-1}}(x) \mu_S(x) = \sum_{x \in G} \mu_{Tgh^{-1}}(x^{-1}) \mu_S(x^{-1}) \\ &= \sum_{x \in G} \mu_{hg^{-1}T}(x) \mu_S(x). \end{aligned}$$

To obtain equality for every g and h , it suffices to see that $\mu_{Tgh^{-1}}(x) = \mu_{hg^{-1}T}(x)$ for every $g, h, x \in G$; equivalently, $\mu_{Tk}(x) = \mu_{kT}(x)$ for every $k, x \in G$. But this is precisely our assumption that $Tk = kT$. \square

Lemma 6 implies that the adjacency matrices of all Cayley multigraphs $\text{Cay}(G, T)$, where T is any normal subgroup of G (or any other union of conjugacy classes), commute. Therefore, they have a common set of eigenvectors.

Proof of Theorem 2. Let $\mathcal{G} = \{\text{Cay}(G, H) : H \triangleleft G\}$. By Lemma 6, the adjacency matrices of all graphs in \mathcal{G} commute and hence they have a common orthogonal set B of eigenvectors. For every $X \in \mathcal{G}$ we have that X has B as a basis of eigenvectors, and X is a disjoint union of cliques (with loops at every vertex), so X is B -integral. It now follows from Lemma 5 that $\mathcal{B}(\mathcal{G}) = \{\text{Cay}(G, S) : S \in \mathcal{B}(G)\}$ is B -integral. Since the adjacency matrix A_T of each multigraph $\text{Cay}(G, T)$, $T \in \mathcal{C}(G)$, is an integral linear combination of adjacency matrices of $\text{Cay}(G, S)$, $S \in \mathcal{B}(G)$, also A_T is B -integral. This completes the proof. \square

3 The boolean algebra of normal subgroups

In this short section, we shall describe the atoms of $\mathcal{B}(G)$. For an element $x \in G$, let $N(x)$ be the intersection of all normal subgroups containing x . For $x, y \in G$, define $x \sim y$ if $N(x) = N(y)$. This is equivalent to the condition that $y \in N(x)$ and $x \in N(y)$. Clearly, \sim is an equivalence relation in G .

Proposition 7. *The equivalence classes of \sim are precisely the atoms of the boolean algebra $\mathcal{B}(G)$, and therefore $S \in \mathcal{B}(G)$ if and only if S is the union of such equivalence classes, and a multiset is in $\mathcal{C}(G)$ if and only if it is a multiset generated by the \sim -classes.*

Proof. Suppose that $x \sim y$. Then $x \in N(y)$ and $y \in N(x)$. Thus, every normal subgroup of G either contains both or neither of these two elements. This property is preserved by taking unions, intersections and complements. Thus, every $S \in \mathcal{B}(G)$ is the union of equivalence classes of \sim .

Suppose now that $x \not\sim y$. If $x \in N(y)$ and $y \in N(x)$, then $N(x) = N(y)$. Thus, we may assume that $x \notin N(y)$. Since $N(y)$ is in $\mathcal{B}(G)$, the atom of $\mathcal{B}(G)$ containing y does not contain x . This shows that each atom is an equivalence class of \sim . \square

4 Abelian groups

Throughout this section, G will always be a (finite) abelian group. Let G^* denote the dual group of G , consisting of all complex characters, i.e., homomorphisms from G to (the multiplicative group of) \mathbb{C} . It is well known that G^* is a group under pointwise multiplication, and that $G^* \cong G$. We define F to be the matrix indexed by $G^* \times G$ and given by the rule that for $\alpha \in G^*$ and $x \in G$ we have $F_{\alpha, x} = \alpha(x)$. Note that each row of F is a character. Furthermore, it follows from the fact that the characters form an orthogonal basis that $FF^* = nI$, where F^* is the conjugate transpose of F and $n = |G|$. Finally, observe that if r is the exponent of G , then every element of F is an r^{th} root of unity.

In the remainder, for any vector $v \in \mathbb{C}^I$ (where I is a non-empty index set) and any $n \in \mathbb{Z}$, we let v^n denote the vector in \mathbb{C}^I given by coordinate-wise exponentiation, i.e., $(v^n)_i = (v_i)^n$ for each $i \in I$.

Observation 8. *Let $x, y \in G$ and let F_x, F_y denote the column vectors of F indexed by x and y , respectively. If $x \sim y$, then there exist integers $j, k \in \mathbb{Z}$ so that $(F_x)^j = F_y$ and $(F_y)^k = F_x$.*

Proof. Since $x \sim y$, we may choose $j, k \in \mathbb{Z}$ so that $x^j = y$ and $y^k = x$. Now, for any character $\alpha \in G^*$ we have $\alpha(y) = \alpha(x^j) = (\alpha(x))^j$ and it follows that $F_y = (F_x)^j$. A similar argument shows that $F_x = (F_y)^k$. \square

Lemma 9. *Let $v \in \mathbb{Q}^G$. If $Fv \in \mathbb{Q}^{G^*}$, then for every $x, y \in G$ with $x \sim y$, we have $v_x = v_y$.*

Proof. Let F_x and F_y denote the column vectors of F indexed by x and y and let ℓ (m) be the smallest integer so that every term of F_x (F_y) is a ℓ^{th} (m^{th}) root of unity. It follows from Observation 8 that $\ell = m$. Now, fix a primitive ℓ^{th} root of unity ω and express each entry of F_x and F_y in the form ω^i for some $i \in \{0, 1, \dots, \ell - 1\}$. Using this interpretation, and recalling that $u := Fv$ is rational, we obtain an expression for the inner product of F_x and u as

$$F_x \cdot u = (F^*u)_x = \sum_{i=0}^{\ell-1} a_i \omega^i$$

where each $a_i \in \mathbb{Q}$. Note that $F_x \cdot u = (F^*u)_x = nv_x$. Now, let $P(z) \in \mathbb{C}[z]$ denote the polynomial $P(z) = \sum_{i=0}^{\ell-1} a_i z^i - nv_x$. Observe that $P(\omega) = 0$. Next, choose $j \in \{0, 1, \dots, \ell - 1\}$ so that $F_y = (F_x)^j$. Note that $\gcd(j, \ell) = 1$. We may assume $x \neq y$, as otherwise there is nothing to prove. It follows that $j \geq 2$, so $\ell \geq 3$. The polynomial P has rational coefficients and has ω as a root. It follows from this and the fact that the polynomial

$$\Phi_\ell(z) = \prod_{i \in \{1.. \ell\}: \gcd(i, \ell) = 1} (z - \omega^i)$$

is irreducible over \mathbb{Q} , that ω^j is also a root of P . But then we have

$$0 = P(\omega^j) = \sum_{i=0}^{\ell-1} a_i \omega^{ij} - nv_x = F_y \cdot u - nv_x$$

which implies that $v_y = \frac{1}{n} F_y \cdot u = v_x$ as desired. \square

We are ready to give a proof of Theorem 1. Observe that sufficiency part follows from Theorem 2.

Proof of Theorem 1, necessity. It is well known that for every $S \subseteq G$ the Cayley graph $\text{Cay}(G, S)$ has each character $\alpha \in G^*$ as an eigenvector with eigenvalue $\alpha(S) = \sum_{g \in S} \alpha(g)$. Alternately, if we view α as a vector and let $\mathbb{1}_S \in \mathbb{C}^G$ denote the characteristic vector of S , this eigenvalue may be written as $\alpha \cdot \mathbb{1}_S$. Suppose that $\text{Cay}(G, S)$ is integral. Then we have $\alpha \cdot \mathbb{1}_S \in \mathbb{Q}$ for every $\alpha \in G^*$. Equivalently, $F\mathbb{1}_S$ is rational-valued. But then, it follows from the previous lemma that whenever $x, y \in G$ satisfy $x \sim y$, we have $(\mathbb{1}_S)_x = (\mathbb{1}_S)_y$. By Proposition 7, this implies that $S \in \mathcal{B}(G)$, as desired. \square

5 Hamiltonian groups

Let \mathcal{H} be the family of groups of the form $Q_8 \times A$ where A is a finite abelian group and Q_8 is the quaternion group represented as follows:

$$Q_8 = \langle -1, i, j, k \mid (-1)^2 = 1, i^2 = j^2 = k^2 = ijk = -1 \rangle.$$

Let us recall that a finite group G is *hamiltonian* if it is non-abelian and every subgroup of G is normal. By a well-known result of Baer (Theorem 3), every hamiltonian group is in \mathcal{H} .

In this section, we obtain a necessary and sufficient condition for a multigraph $\text{Cay}(G, S)$ to be integral, where $G \in \mathcal{H}$ and $S \subseteq G$ is an inverse-closed multiset of elements of G .

We require some definitions and notation from representation theory. For a more detailed account the reader is referred to [3, 7].

In what follows, let G be a finite group. We denote by $\mathbb{C}G$ its group algebra. That is, $\mathbb{C}G$ is the vector space over \mathbb{C} with basis G and multiplication defined by extending the group multiplication linearly. Identifying $\sum_{g \in G} a_g g$ with the function $g \mapsto a_g$, we can view the vector space $\mathbb{C}G$ as the space of all \mathbb{C} -valued functions on G . We sometimes identify a subset or a multiset M of G with the element $\sum_{g \in G} \mu_M(g)g$ of the group algebra $\mathbb{C}G$.

Let V be an n -dimensional vector space over \mathbb{C} . A *complex representation* (or simply a *representation*) of G on V is a group homomorphism $\rho : G \rightarrow GL(V)$, where $GL(V)$ denotes the group of invertible endomorphism of V . The *degree* of representation is the dimension of V . Two representations ρ_1 and ρ_2 of G on V_1 and V_2 , respectively, are *equivalent* (written $\rho_1 \cong \rho_2$) if there is a linear isomorphism $T : V_1 \rightarrow V_2$ such that for every $g \in G$ we have $T\rho_1(g) = \rho_2(g)T$.

If ρ is a representation of G then the *character* χ_ρ induced by ρ , is the linear functional $\chi_\rho : \mathbb{C}G \rightarrow \mathbb{C}$ defined by

$$\chi_\rho(g) = \text{tr}(\rho(g)), \quad g \in G,$$

and extended by linearity to $\mathbb{C}G$. (The trace $\text{tr}(\alpha)$ of a linear map α is the trace of any matrix representing α according to some basis.) The *degree* of the character χ_ρ is the degree of ρ , and is equal to $\chi_\rho(1)$. A character of degree one is called a *linear character*. The character χ which assigns 1 to every element of a group G is called the *principal character* of G and is denoted by $\mathbb{1}_G$.

The $|G|$ -dimensional representation $\rho_{reg} : G \rightarrow GL(\mathbb{C}G)$ defined by $\rho_{reg}(g)(x) = gx$ ($g \in G, x \in \mathbb{C}G$) is called the *left-regular representation*. Choosing G as a basis for $\mathbb{C}G$, we see that for every $g \in G$, $[\rho_{reg}(g)]_G$ (the matrix representing the linear map $\rho_{reg}(g)$ according to the basis G) is the $(|G| \times |G|)$ -matrix, indexed with the elements of G , such that for every $k, h \in G$

$$([\rho_{reg}(g)]_G)_{h,k} = \begin{cases} 1 & \text{if } h = gk \\ 0 & \text{otherwise.} \end{cases}$$

This gives a natural link to Cayley graphs since the adjacency matrix of a Cayley multi-graph $\text{Cay}(G, S)$ can be written as

$$A = \sum_{s \in S} \mu_S(s) [\rho_{\text{reg}}(s)]_G. \quad (1)$$

Let $\rho : G \rightarrow GL(V)$ be a representation. A subspace W of V is said to be ρ -invariant, if $\rho(g)w \in W$ for every $g \in G$ and $w \in W$. If W is a ρ -invariant subspace of V , then the restriction of ρ to W , that is $\rho|_W : G \rightarrow GL(W)$, is a representation of G on W . If V has no ρ -invariant subspaces, then ρ is said to be an *irreducible representation* for G and the corresponding character χ_ρ an *irreducible character* for G .

For a group G , we denote by $\text{IRR}(G)$ and $\text{Irr}(G)$ a complete set of non-equivalent irreducible representations of G and the complete set of non-equivalent irreducible characters of G , respectively. Note that $\text{IRR}(G)$ is not necessarily unique, but $\text{Irr}(G)$ is unique. If G is abelian, then every irreducible representation $\rho \in \text{IRR}(G)$ is 1-dimensional and thus it can be identified with its character $\chi_\rho \in \text{Irr}(G)$.

The following result has been observed by many authors (see for example [2]).

Lemma 10. *Let G be a group and let $S \subseteq G$ be a multiset of elements of G . Let $\text{IRR}(G) = \{\rho_1, \dots, \rho_k\}$. For $t = 1, \dots, k$, let d_t be the degree of ρ_t , and let Λ_t be the multiset of eigenvalues of the matrix $\sum_{g \in S} \mu_S(g) \rho_t(g)$. Then the following holds:*

- (1) *The set of eigenvalues of $\text{Cay}(G, S)$ equals $\cup_{t=1}^k \Lambda_t$.*
- (2) *If the eigenvalue λ occurs with multiplicity $m_t(\lambda)$ in $\sum_{g \in S} \mu_S(g) \rho_t(g)$ ($1 \leq t \leq k$), then the multiplicity of λ in $\text{Cay}(G, S)$ is $\sum_{t=1}^k d_t m_t(\lambda)$.*

The table below is the character table of Q_8 .

$g \in Q_8$	1	-1	i	j	k
$cl(g)$	{1}	{-1}	{ $i, -i$ }	{ $j, -j$ }	{ $k, -k$ }
$\mathbb{1}_{Q_8}$	1	1	1	1	1
λ_i	1	1	1	-1	-1
λ_j	1	1	-1	1	-1
λ_k	1	1	-1	-1	1
ε	2	-2	0	0	0

where ε is the character afforded by the representation ρ_ε defined below:

$$\rho_\varepsilon(1) = I, \quad \rho_\varepsilon(i) = \begin{pmatrix} \mathbf{i} & 0 \\ 0 & -\mathbf{i} \end{pmatrix}, \quad \rho_\varepsilon(j) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \rho_\varepsilon(k) = \begin{pmatrix} 0 & \mathbf{i} \\ \mathbf{i} & 0 \end{pmatrix}$$

where the value \mathbf{i} appearing in the matrices is the complex imaginary unit $\sqrt{-1}$. Also note that $\rho_\varepsilon(-g) = -\rho_\varepsilon(g)$ for every $g \in Q_8$ and that $\text{IRR}(Q_8) = \{\mathbb{1}_{Q_8}, \lambda_i, \lambda_j, \lambda_k, \rho_\varepsilon\}$.

Let $G = Q_8 \times A$, where A is an abelian group, and let $S \subseteq G$ be an inverse-closed multiset of elements of G . For every $q \in Q_8$, let B_q be the multiset

$$B_q = \{a \in A \mid (q, a) \in S\} \tag{2}$$

in which the multiplicity of $a \in B_q$ is equal to the multiplicity of (q, a) in S .

Since S is inverse-closed, we have $B_1 = B_1^{-1}$, $B_{-1} = B_{-1}^{-1}$, and $B_{-q} = B_q^{-1}$ for every $q \in Q_8 \setminus \{1, -1\}$. In particular, this implies that $\lambda(B_{-q}) = \overline{\lambda(B_q)}$, for every $\lambda \in \text{Irr}(A)$. For every multiset D of elements of A , we define

$$\widehat{\lambda}(D) = \lambda(D) - \lambda(D^{-1}) = \sum_{g \in D} \mu_D(g)(\lambda(g) - \lambda(g^{-1})) = \sum_{g \in D} \mu_D(g)(\lambda(g) - \overline{\lambda(g)}).$$

In particular, for every $q \in Q_8$, $\widehat{\lambda}(B_q) = \lambda(B_q) - \lambda(B_{-q})$. The following is the main result of this section.

Theorem 11. *Let $G = Q_8 \times A$, where A is an abelian group, and let S be an inverse-closed multiset of elements of G . Then $\text{Cay}(G, S)$ is integral if and only if the following holds:*

- (i) $B_1, B_{-1} \in \mathcal{C}(A)$.
- (ii) *The multiset union $B_q \cup B_{-q} \in \mathcal{C}(A)$, for every $q \in Q_8 \setminus \{-1, 1\}$.*
- (iii) $\widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2$ *is a negative square of an integer, for every $\lambda \in \text{Irr}(A)$.*

Proof. Lemma 10 shows that $\text{Cay}(G, S)$ is integral if and only if the matrix $\sum_{s \in S} \mu_S(s)\phi(s)$ is integral for every $\phi \in \text{IRR}(G)$. Since ϕ is an irreducible representation of the direct product $Q_8 \times A$, it can be written in the form $\phi = \rho \times \lambda$ for some $\rho \in \text{IRR}(Q_8)$ and $\lambda \in \text{Irr}(A)$ (where we identify $\text{Irr}(A)$ and $\text{IRR}(A)$ since all irreducible representations of A are 1-dimensional). In other words, $\phi(q, a) = \lambda(a)\rho(q)$ for every $(q, a) \in Q_8 \times A$. Consequently, $\text{Cay}(G, S)$ is integral if and only if the matrices

$$A^{(\rho, \lambda)} = \sum_{(q, a) \in S} \mu_S((q, a))(\rho \times \lambda)(q, a) = \sum_{(q, a) \in S} \mu_S((q, a))\lambda(a)\rho(q)$$

are integral for every $\rho \in \text{IRR}(Q_8)$ and every $\lambda \in \text{Irr}(A)$. By definition of B_q we can write the matrix $A^{(\rho, \lambda)}$ in the following form:

$$A^{(\rho, \lambda)} = \sum_{(q, a) \in S} \mu_S((q, a))\lambda(a)\rho(q) = \sum_{q \in Q_8} \lambda(B_q)\rho(q). \tag{3}$$

Integrality of the matrix in (3) (with $\rho = \rho_\varepsilon$ and $\lambda \in \text{Irr}(A)$ arbitrary) together with the fact that the trace of a matrix is equal to the sum of its eigenvalues implies that

$$\text{tr}\left(\sum_{q \in Q_8} \lambda(B_q)\rho_\varepsilon(q)\right) = \sum_{q \in Q_8} \lambda(B_q)\varepsilon(q) = 2(\lambda(B_1) - \lambda(B_{-1})) \in \mathbb{Z}.$$

It follows that

$$\lambda(B_1) - \lambda(B_{-1}) \in \mathbb{Q}. \tag{4}$$

Let $\rho \in \{\mathbb{1}_{Q_8}, \lambda_i, \lambda_j, \lambda_k\}$ be a degree-one representation of Q_8 and let $\lambda \in \text{Irr}(A)$. Define $\lambda^+(B_q) = \lambda(B_q) + \lambda(B_{-q})$. Observe that $\rho(q) = \rho(-q)$ for every $q \in Q_8$. Therefore, integrality of the matrices $A^{(\rho, \lambda)}$ in (3) implies by the same argument as above that

$$\rho(1)\lambda^+(B_1) + \rho(i)\lambda^+(B_i) + \rho(j)\lambda^+(B_j) + \rho(k)\lambda^+(B_k) \in \mathbb{Z}.$$

This yields the following four conditions (one for each $\rho \in \{1, \lambda_i, \lambda_j, \lambda_k\}$):

$$\begin{aligned} \lambda^+(B_1) + \lambda^+(B_i) + \lambda^+(B_j) + \lambda^+(B_k) &\in \mathbb{Z} \\ \lambda^+(B_1) + \lambda^+(B_i) - \lambda^+(B_j) - \lambda^+(B_k) &\in \mathbb{Z} \\ \lambda^+(B_1) - \lambda^+(B_i) + \lambda^+(B_j) - \lambda^+(B_k) &\in \mathbb{Z} \\ \lambda^+(B_1) - \lambda^+(B_i) - \lambda^+(B_j) + \lambda^+(B_k) &\in \mathbb{Z} \end{aligned} \tag{5}$$

Since the matrix of coefficients of the linear system (5) is invertible, this implies that $\lambda^+(B_q) \in \mathbb{Q}$ for every $q \in Q_8$. In particular, since $\lambda^+(B_1) = \lambda(B_1) + \lambda(B_{-1}) \in \mathbb{Q}$, we conclude by using (4) that $\lambda(B_1) \in \mathbb{Q}$ and $\lambda(B_{-1}) \in \mathbb{Q}$, while for $q \in Q_8 \setminus \{1, -1\}$, we have $\lambda(B_q) + \lambda(B_{-q}) \in \mathbb{Q}$.

The rationality of $\lambda(X)$ for every $\lambda \in \text{Irr}(A)$ has been discussed in the proof of Lemma 9, where it was proved that this is equivalent to the condition that $X \in \mathcal{C}(A)$. Therefore, the conclusions stated in the previous paragraph imply (i) and (ii).

Conversely, notice that by Theorem 1, (i) and (ii) imply integrality of the matrices in (3), where ρ is any degree-one representation of Q_8 and $\lambda \in \text{Irr}(A)$.

For (iii), we consider the degree-two representation ρ_ε . As observed above, $\rho_\varepsilon(-q) = -\rho_\varepsilon(q)$ for every $q \in Q_8$, and hence

$$\sum_{q \in Q_8} \lambda(B_q)\rho_\varepsilon(q) = \widehat{\lambda}(B_1)I + \widehat{\lambda}(B_i) \begin{pmatrix} \mathbf{i} & 0 \\ 0 & -\mathbf{i} \end{pmatrix} + \widehat{\lambda}(B_j) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} + \widehat{\lambda}(B_k) \begin{pmatrix} 0 & \mathbf{i} \\ \mathbf{i} & 0 \end{pmatrix}.$$

As mentioned above, (4) implies that $\widehat{\lambda}(B_1) \in \mathbb{Z}$. Therefore, $\sum_{q \in Q_8} \lambda(B_q)\rho_\varepsilon(q)$ is integral if and only if the matrix

$$M = \begin{pmatrix} \mathbf{i}\widehat{\lambda}(B_i) & \widehat{\lambda}(B_j) + \mathbf{i}\widehat{\lambda}(B_k) \\ -\widehat{\lambda}(B_j) + \mathbf{i}\widehat{\lambda}(B_k) & -\mathbf{i}\widehat{\lambda}(B_i) \end{pmatrix}.$$

is integral. By considering the characteristic polynomial of M , it is easy to see that M is integral if and only if $\widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2$ is the negative square of an integer. Hence, integrality of $\text{Cay}(G, S)$ implies (iii), and conversely, (iii) implies integrality of the matrices $A^{(\rho, \lambda)}$. This completes the proof. \square

6 Some special cases

In this section we consider some special cases of hamiltonian groups by applying Theorem 11. This result gives a simple characterization in some cases, and leads to interesting combinatorial problems in some other cases.

6.1 Simple Cayley graphs of $Q_8 \times C_p$, where $p \neq 3$

As the first special case of using Theorem 11 we consider hamiltonian groups $G = Q_8 \times C_p$, where $p \neq 3$ is a prime and C_p is the cyclic group of order p . In analogy with the abelian case, we obtain the following complete characterization for integrality of *simple* Cayley graphs over this group. The multigraph version is different and is treated in a separate section.

Theorem 12. *Let $p \neq 3$ be a prime and let $S \subseteq Q_8 \times C_p$ be an inverse-closed subset of $Q_8 \times C_p$. The Cayley graph $\text{Cay}(Q_8 \times C_p, S)$ is integral if and only if $S \in \mathcal{B}(Q_8 \times C_p)$.*

This result is a direct consequence of the following:

Theorem 13. *Let $G = Q_8 \times C_p$, for a prime $p \neq 3$. Let $S \subseteq G$ be an inverse-closed subset of G , and let B_q ($q \in Q_8$) be defined as in (2). Then $\text{Cay}(G, S)$ is integral if and only if the following conditions hold:*

(P1) $B_1, B_{-1} \in \mathcal{B}(C_p)$.

(P2) For every $q \in Q_8 \setminus \{1, -1\}$, $B_q = B_{-q} \in \mathcal{B}(C_p)$.

Proof. By Theorem 11, it suffices to show that (P2) holds if and only if conditions (ii) and (iii) in Theorem 11 hold. The “only if” part is trivial, since (P2) implies that $\widehat{\lambda}(B_i) = \widehat{\lambda}(B_j) = \widehat{\lambda}(B_k) = 0$. For the “if” part suppose that conditions (ii) and (iii) of Theorem 11 hold. By condition (iii), for every $\lambda \in \text{Irr}(C_p)$ there is an integer α_λ so that

$$\widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2 = -\alpha_\lambda^2. \quad (6)$$

Let e be the unit in C_p and let $E_1 = \{e\}$ and $E_2 = C_p \setminus E_1$ be the two equivalence classes of C_p . Let $q \in Q_8 \setminus \{1, -1\}$. Recall that since S is inverse-closed we have $B_q^{-1} = B_{-q}$.

If $B_q \in \mathcal{B}(C_p)$ then $B_{-q} = B_q$. This is true because $B_q^{-1} = B_{-q}$, and the sets E_1 and E_2 are inverse-closed. Hence in this case $B_q = B_{-q}$ and $\widehat{\lambda}(B_q) = 0$. If $p = 2$, then every subset of C_p is in $\mathcal{B}(C_p)$, so (P2) holds in this case, and we may henceforth assume that $p \geq 5$.

If $B_q \notin \mathcal{B}(C_p)$, then by condition (ii) and the fact that $B_q^{-1} = B_{-q}$, we conclude that $E_2 \subseteq B_q \Delta B_{-q}$, thus the support of $B_q - B_{-q}$, viewed as an element of the group algebra $\mathbb{C}C_p$, contains $p - 1$ distinct elements (that is, the whole class E_2), where an element and its inverse appear with opposite signs. In particular, the sum of coefficients of elements of $B_q - B_{-q}$ is 0.

Let us write $B'_q = B_q \setminus B_{-q}$, and observe that for every $q \in Q_8$, either $B'_q = \emptyset$ or $|B'_q| = \frac{1}{2}(p-1)$. Now, (6) can be written as follows

$$\begin{aligned}
 -\alpha_\lambda^2 &= \widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2 \\
 &= \widehat{\lambda}(B'_i)^2 + \widehat{\lambda}(B'_j)^2 + \widehat{\lambda}(B'_k)^2 \\
 &= \lambda((B'_i - B'_{-i})^2) + \lambda((B'_j - B'_{-j})^2) + \lambda((B'_k - B'_{-k})^2) \\
 &= \lambda((B'_i - B'_{-i})^2 + (B'_j - B'_{-j})^2 + (B'_k - B'_{-k})^2) \\
 &= \lambda\left(-2(|B'_i| + |B'_j| + |B'_k|)e + \sum_{g \in E_2} a_g g\right) \\
 &= -2(|B'_i| + |B'_j| + |B'_k|) + \lambda\left(\sum_{g \in E_2} a_g g\right) \tag{7}
 \end{aligned}$$

where $a_g \in \mathbb{Z}$ for every $g \in E_2$. Since the sum of coefficients in $B_q - B_{-q}$ is zero, it follows that the sum of coefficients in $(B_q - B_{-q})^2$ is also zero. Thus, (7) implies that

$$\sum_{g \in E_2} a_g = 2(|B'_i| + |B'_j| + |B'_k|). \tag{8}$$

By (7), $\lambda(\sum_{g \in E_2} a_g g) \in \mathbb{Q}$ for every $\lambda \in \text{Irr}(C_p)$. It follows by Lemma 9, that all coefficients a_g are equal, and from (8) we conclude that for every $g \in E_2$:

$$a_g = \frac{2(|B'_i| + |B'_j| + |B'_k|)}{p-1}.$$

We also know that for each non-principal character $\lambda \in \text{Irr}(C_p)$ we have $\sum_{g \in C_p} \lambda(g) = 0$. Thus, $\sum_{g \in E_2} \lambda(g) = -1$, and we can rewrite (7) as follows:

$$-\alpha_\lambda^2 = -2(|B'_i| + |B'_j| + |B'_k|) - \frac{2(|B'_i| + |B'_j| + |B'_k|)}{p-1}.$$

This gives the following conclusion:

$$\alpha_\lambda^2 = 2(|B'_i| + |B'_j| + |B'_k|) \frac{p}{p-1}. \tag{9}$$

We know that for every $q \in Q_8$, $|B'_q|$ is either 0 or $\frac{1}{2}(p-1)$. By (9), $p-1$ divides $2(|B'_i| + |B'_j| + |B'_k|)$. Let β denote the number of elements $q \in \{i, j, k\}$ such that $|B'_q| = \frac{p-1}{2}$. Then we conclude from (9) that $\alpha_\lambda^2 = \beta p$. Since $0 \leq \beta \leq 3$ and $p \geq 5$, this is possible only when $\alpha = 0$. However, in that case (P2) holds. \square

6.2 $Q_8 \times C_3$

The conclusion of Theorem 13 does not hold for $p = 3$. An example is provided in the next observation.

Observation 14. Let $G = Q_8 \times C_3$, and $S = \{(i, 1), (-i, 2), (j, 1), (-j, 2), (k, 1), (-k, 2)\}$. Then $\text{Cay}(G, S)$ is integral but $S \notin \mathcal{B}(G)$.

To see this, we verify conditions (i)–(iii) of Theorem 11. Conditions (i) and (ii) are obvious; (iii) is left to the reader.

This graph is indeed a very interesting vertex-transitive graph whose properties are discussed below. Let us remark at this point that the proof of the Theorem 13 shows that the example in Observation 14 is the only integral simple Cayley graph of $Q_8 \times C_3$ (up to Cayley graph isomorphisms and up to choice of $B_1, B_{-1} \in \mathcal{B}(C_3)$) that fails to satisfy the conclusion of Theorem 13.

This graph has a natural tripartition according to the first coordinate, and the bipartite graphs obtained from it by removing one of these tripartite classes is the Möbius-Kantor graph. The Möbius-Kantor graph is the unique double-cover of the cube of girth 6 and it sits naturally as a subgraph of the 4-cube. The graph of the 24-cell is also tripartite with classes of size 8, and deleting any one yields a 4-cube.

6.3 Cayley multigraphs of $Q_8 \times C_p$

Theorem 13 does not hold for the multigraph case. In this section we shall provide infinitely many examples confirming this. We let $C_p = \{a^t \mid 0 \leq t < p\}$, the cyclic group of order p generated by a . We consider the multisets B_q ($q \in Q_8 \setminus \{1, -1\}$) defined as in (2), and we set $B_1 = B_{-1} = \emptyset$. In order to satisfy conditions (i)–(iii) of Theorem 11, we need that $B_q \cup B_{-q} \in \mathcal{C}(C_p)$ and $\widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2$ is a negative square of an integer, for every $\lambda \in \text{Irr}(C_p)$. As before, for every $q \in Q_8 \setminus \{1, -1\}$ we define $B'_q = B_q \setminus (B_q \cap B_{-q})$, where $B_q \cap B_{-q}$ is the multiset in which the multiplicity of any $x \in C_p$ is equal to the minimum of multiplicities of x in B_q and in B_{-q} . Thus, in particular, $\widehat{\lambda}(B_q) = \widehat{\lambda}(B'_q)$. Note that B'_q and B'_{-q} are disjoint and the condition that generating multiset is inverse-closed is equivalent to the requirement that the multiplicity of a^t ($0 \leq t < p$) in B'_q is the same as the multiplicity of a^{-t} in B'_{-q} . The following is a well-known result from number theory.

Lemma 15. If p is a prime number and $p \equiv 1 \pmod{4}$, then the diophantine equation $x^2 + y^2 = pz^2$ has infinitely many solutions satisfying $\gcd(x, y, z) = 1$.

A solution of the diophantine equation $x^2 + y^2 = pz^2$ is *primitive* if $\gcd(x, y, z) = 1$. Clearly, every integral multiple of (x, y, z) is also a solution. The solution $(0, 0, 0)$ is called the *trivial solution*.

Lemma 16. Let (r, s, t) be a non-trivial solution for the diophantine equation $x^2 + y^2 = 5z^2$. Let $D_1 = ra + sa^2$, $D_2 = ra + sa^3$ and $D_3 = 0$, be elements of $\mathbb{C}C_5$, where a is a generator of C_5 . Then

$$\widehat{\lambda}(D_1)^2 + \widehat{\lambda}(D_2)^2 + \widehat{\lambda}(D_3)^2 = -(5t)^2.$$

Proof. For any x and y in \mathbb{R} , we have the following equation in $\mathbb{C}C_5$:

$$(x(a - a^4) + y(a^2 - a^3))^2 = -2(x^2 + y^2) + (y^2 - 2xy)(a + a^4) + (x^2 + 2xy)(a^2 + a^3).$$

Thus,

$$\begin{aligned}\widehat{\lambda}(D_1)^2 &= -2(r^2 + s^2) + (s^2 - 2rs)\lambda(a + a^4) + (r^2 + 2rs)\lambda(a^2 + a^3), \\ \widehat{\lambda}(D_2)^2 &= -2(r^2 + s^2) + (r^2 + 2rs)\lambda(a + a^4) + (s^2 - 2rs)\lambda(a^2 + a^3).\end{aligned}$$

Clearly, $\widehat{\lambda}(D_3)^2 = 0$. We notice also that for each non-principal character $\lambda \in \text{Irr}(C_5)$ we have $\sum_{i=1}^4 \lambda(a^i) = -1$. Therefore, $\widehat{\lambda}(D_1)^2 + \widehat{\lambda}(D_2)^2 + \widehat{\lambda}(D_3)^2 = -5(r^2 + s^2) = -(5t)^2$. \square

Corollary 17. *There are infinitely many multisets S (none of which is a multiple of another) such that $\text{Cay}(Q_8 \times C_5, S)$ is integral but $S \notin \mathcal{C}(G)$.*

Proof. Let us start with a primitive solution (m, n, α) of the diophantine equation $x^2 + y^2 = 5z^2$. Since $(2m, 2n, 2\alpha)$ is a solution of the diophantine equation $x^2 + y^2 = 5z^2$, we can construct D_1, D_2 and D_3 as in the previous lemma, i.e., $D_1 = 2ma + 2na^2, D_2 = 2na + 2ma^3$ and $D_3 = 0$. Suppose without loss of generality that $n \leq m$. Let us take $B_i = \{2ma, (m+n)a^2, (m-n)a^3\}, B_j = \{(m+n)a, (m-n)a^4, 2ma^3\}, B_{-i} = B_i^{-1}, B_{-j} = B_j^{-1}$, and $B_k = B_{-k} = \emptyset$ (where the coefficients of a^t in the set notation denote multiplicities). Then $B_i + B_{-i} \in \mathcal{C}(C_5), B_j + B_{-j} \in \mathcal{C}(C_5)$, and $B_k + B_{-k} \in \mathcal{C}(C_5)$. We also have $B'_i = D_1, B'_j = D_2$ and $B'_k = D_3$. From the previous lemma we get

$$\widehat{\lambda}(B_i)^2 + \widehat{\lambda}(B_j)^2 + \widehat{\lambda}(B_k)^2 = \widehat{\lambda}(B'_i)^2 + \widehat{\lambda}(B'_j)^2 + \widehat{\lambda}(B'_k)^2 = -(10\alpha)^2.$$

So, clearly conditions (i)–(iii) of Theorem 11 are satisfied for the generating multiset S arising from B_i, B_j and B_k , but $B_i, B_j \notin \mathcal{C}(C_5)$. Thus $S \notin \mathcal{C}(Q_8 \times C_5)$, while according to Theorem 11 the Cayley graph $\text{Cay}(Q_8 \times C_5, S)$ is integral. \square

The case $p = 7$ is similar. First, we observe that there are infinitely many primitive solutions for diophantine equation $x^2 + y^2 + z^2 = 7\alpha^2$. If we assume (m, n, l, α) is one of these solutions, then we can define

$$B'_i = ma + na^2 + la^3, \quad B'_j = la + ma^2 + na^3, \quad B'_k = na + la^2 + ma^3.$$

It is easy to see that condition (iii) of Theorem 11 holds. As in Corollary 17, we can define B_i, B_j and B_k using correspondence with B'_i, B'_j and B'_k such that conditions (i)–(iii) of Theorem 11 are satisfied. The details are left to the reader. This gives rise to integral Cayley multigraphs of $Q_8 \times C_7$ whose generating multiset is not in the lattice $\mathcal{C}(Q_8 \times A)$.

6.4 Simple Cayley graphs of $Q_8 \times C_p^d$

As the last special case we consider the group $G = Q_8 \times C_p^d$, where p is a prime and $d \geq 2$. Here the abelian direct factor of G is an elementary abelian p -group, thus every non-identity element has order p . If $[a]$ denotes the equivalence class containing a with respect to the relation \sim in C_p^d and if $a \neq e$ (where e is the identity element of C_p^d), then $[a] = \{a^t \mid 1 \leq t \leq p-1\}$. We also know that $[e] = \{e\}$; we call this the trivial equivalence class. Since each non-identity element in C_p^d has order p , each non-trivial class is of order

$p - 1$, and the number of non-trivial classes is equal to $n_d = \frac{p^d - 1}{p - 1}$. Label these classes as A_r for $1 \leq r \leq n_d$. If λ is a non-principal character of C_p^d , then $|\text{Im}(\lambda)| = p$ and therefore $\ker(\lambda)$ is a subgroup of order p^{d-1} .

Let us assume that $\text{Cay}(G, S)$ is integral. Then we derive in the same way as in the case of $Q_8 \times C_p$ that there is an integer α_λ such that

$$-\alpha_\lambda^2 = \widehat{\lambda}(B'_i)^2 + \widehat{\lambda}(B'_j)^2 + \widehat{\lambda}(B'_k)^2 = -2(|B'_i| + |B'_j| + |B'_k|) + \lambda\left(\sum_{g \in C_p^d \setminus \{e\}} a_g g\right). \quad (10)$$

Since (10) holds for every $\lambda \in \text{Irr}(C_p^d)$, we conclude by Lemma 9 that the coefficients a_g are constant on each equivalence class A_r . Let b_r be the common value for a_g , $g \in A_r$. Then

$$\begin{aligned} -\alpha_\lambda^2 &= -2(|B'_i| + |B'_j| + |B'_k|) + \lambda\left(\sum_{r=1}^{n_d} \sum_{g \in A_r} b_r g\right) \\ &= -2(|B'_i| + |B'_j| + |B'_k|) + \sum_{r=1}^{n_d} b_r \lambda(A_r). \end{aligned} \quad (11)$$

Since each $A_r \cup \{e\}$ is a subgroup of order p , we have

$$\lambda(A_r) = \sum_{g \in A_r} \lambda(g) = \begin{cases} p - 1, & A_r \subseteq \ker(\lambda) \\ -1, & A_r \not\subseteq \ker(\lambda). \end{cases} \quad (12)$$

We also notice that for $q \in \{i, j, k\}$ the element $B_q - B_{-q}$ of the group algebra has the sum of the coefficients equal to zero. By using this fact in combination with (11) and (12) for the case when λ is the principal character and noting that $\alpha_\lambda = 0$ in that case, we obtain the following analogue of (9):

$$2(|B'_i| + |B'_j| + |B'_k|) = (p - 1) \sum_{r=1}^{n_d} b_r. \quad (13)$$

Using (13), we have for every non-principal character λ :

$$\alpha_\lambda^2 = 2(|B'_i| + |B'_j| + |B'_k|) - \sum_{r=1}^{n_d} b_r \lambda(A_r) = \sum_{r=1}^{n_d} b_r (p - 1 - \lambda(A_r)). \quad (14)$$

The equality (12) shows that a non-zero contribution in the sum on the right side of (14) arises only when $A_r \not\subseteq \ker(\lambda)$. Let $I_\lambda \subseteq \{1, \dots, n_d\}$ be the set of values r for which $A_r \not\subseteq \ker(\lambda)$. Then we have:

$$\alpha_\lambda^2 = \sum_{r \in I_\lambda} b_r (p - 1 - \lambda(A_r)) = p \sum_{r \in I_\lambda} b_r. \quad (15)$$

There is a natural geometric setting for these equations. View C_p^d as a vector space over C_p and consider the projective geometry $PG(d - 1, p)$ consisting of all subspaces of C_p^d .

The points in our projective geometry are the 1-dimensional subspaces of C_p^d which are in correspondence with A_1, A_2, \dots, A_{n_d} , and we label the point associated with A_i by b_i . The kernels of the non-principal characters of C_p^d correspond to the hyperplanes in our projective geometry (i.e. subspaces of dimension $d-1$ of C_p^d). So, equation (15) implies that the sum of the labels on the complement of every hyperplane must be a square divided by p . Although this is a meaningful constraint, it is not difficult to find labelings of the points in a projective geometry which satisfy this property, so a more complicated analysis will be required to understand the integrality of such Cayley graphs.

References

- [1] Bridges, W. G., and Mena, R. A. Rational G-matrices with rational eigenvalues. *J. Combin. Theory, Ser. A* 32 (1982), 264–280.
- [2] Diaconis, P., and Shahshahani, M. Generating a random permutation with random transpositions. *Z. Wahrsch. Verw. Gebiete* 57 (1981) 159–179.
- [3] Fulton, W., Harris, J. *Representation theory. A first course*. Springer-Verlag, 1991.
- [4] Godsil, C., and Royle, G. *Algebraic graph theory*. Graduate Texts in Mathematics. Vol. 207. Springer, 2001.
- [5] Hall, M. *The theory of groups*. Macmillan, New York, 1959.
- [6] Harary, F., and Schwenk, A. J. Which graphs have integral spectra? In “Graphs and Combinatorics (Proc. Capital Conf., George Washington Univ., Washington, D.C., 1973)”. Lecture Notes in Mathematics 406. Springer, Berlin, 1974, pp. 45–51.
- [7] James, G., Liebeck, M. *Representations and characters of groups*. Second edition. Cambridge University Press, 2001.
- [8] Klotz, W., and Sander, T. Integral Cayley graphs over abelian groups. *Electronic J. Combinatorics* 17 (2010), #R81.
- [9] So, W. Integral circulant graphs. *Discrete Math.* 306 (2005), 153–158.