

General Restriction of (s,t) -Wythoff's Game

Wen An Liu*

College of Mathematics and Information Science
Henan Normal University
Xinxiang, P. R. China
liuwenan@126.com

Haiyan Li

College of Mathematics and Information Science
Henan Normal University
Xinxiang, P. R. China
lihaiyan1107@126.com

Submitted: Dec 11, 2012; Accepted: ; Published: XX
Mathematics Subject Classifications: 91A46, 68R05

Abstract

A.S. Fraenkel introduced a new (s,t) -Wythoff's game which is a generalization of both Wythoff's game and α -Wythoff's game. Four new models of a restricted version of (s,t) -Wythoff's game, Odd-Odd (s,t) -Wythoff's Game, Even-Even (s,t) -Wythoff's Game, Odd-Even (s,t) -Wythoff's Game and Even-Odd (s,t) -Wythoff's Game, are investigated. Under normal or misère play convention, all P -positions of these four models are given for arbitrary integers $s, t \geq 1$. For Even-Even (s,t) -Wythoff's Game, the structure of P -positions is given by recursive characterizations in term of mex function. For other models, the structures of P -positions are of algebraic form, which permit to decide in polynomial time whether or not a given game position (a,b) is a P -position.

Keywords: impartial combinatorial game; normal play convention; misère play convention; P -position; (s,t) -Wythoff's game

1 Introduction

By game we mean a combinatorial game; we restrict our attention to classical impartial games. There are two conventions: in *normal play convention*, the player first unable to

*This work is supported by the National Natural Science Foundation of China under Grants 11171368 and 11171094.

move is the loser (his opponent is the winner); in *misère play convention*, the player first unable to move is the winner (his opponent is the loser). The positions from which the previous player can win regardless of the opponent's moves are called *P-positions* and those from which the next player can win regardless of the opponent's moves are called *N-positions*. The theory of such games can be found in [1, 2, 4, 8].

Throughout this paper, we use the following notations.

- (1) By $Z^{\geq m}$ we denote the set of all integers not less than m , i.e., $Z^{\geq m} = \{x \geq m \mid x \text{ is an integer}\}$. Let $Z^{\text{even}} = \{2n \mid n \in Z^{\geq 0}\}$, $Z^{\text{odd}} = \{2n + 1 \mid n \in Z^{\geq 0}\}$.
- (2) For any set $U \subseteq Z^{\geq 0}$, by $\text{mex}(U)$ we denote the *Minimum EXcluded value* of U , i.e., the smallest nonnegative integer not in U . In particular, $\text{mex}(\emptyset) = 0$.
- (3) By $\lfloor x \rfloor$ we denote the largest integer $\leq x$.
- (4) We use the notation $(x_1, y_1) \rightarrow (x_2, y_2)$ if there is a legal move from (x_1, y_1) to (x_2, y_2) .

1.1 Wythoff's game

Wythoff's game is played with two heaps of tokens. Each player can either remove any number of tokens from a single heap (*Nim rule*) or remove the same number of tokens from both heaps (*Wythoff's rule*). All *P-positions* of Wythoff's game under normal play convention were given in [19]. All *P-positions* of Wythoff's game under misère play convention were determined in [13].

1.2 Extension of Wythoff's game

In many papers devoted to variations of Wythoff's game, new rules are adjoined to the original ones. Such variations are called *extensions*.

As an example, *a-Wythoff's game* was investigated in [9]: Given an integer $a \geq 1$ and two heaps of finitely many tokens. Two rules of moves are allowed.

(*Nim Rule*) Take any positive number of tokens from a single heap, possibly the entire heap.

(*General Wythoff's Rule*) Take tokens from both heaps, $k > 0$ tokens from one heap, and $\ell > 0$ tokens from the other, and $|k - \ell| < a$, where $a > 0$ is a fixed integer parameter.

A.S. Fraenkel [11] introduced a new (s, t) -*Wythoff's game*. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. There are two types of moves:

(*Nim Rule*) Take any positive number of tokens from a single heap, possibly the entire heap.

(*More General Wythoff's Rule*) Take tokens from both heaps, $k > 0$ from one heap and $\ell > 0$ from the other, and

$$0 < k \leq \ell < sk + t. \quad (1)$$

In [11], the author gave the following results: Denote by $\mathcal{P}_{s,t}$ the set of all *P-positions* of (s, t) -Wythoff's game under normal play convention. Then $\mathcal{P}_{s,t} = \bigcup_{n=0}^{\infty} \{(A_n, B_n)\}$,

where for $n \geq 0$,

$$\begin{cases} A_n = \text{mex}\{A_i, B_i | 0 \leq i < n\}, \\ B_n = sA_n + tn. \end{cases} \quad (2)$$

It is worth to mention that Wythoff's game is a special case $s = t = 1$ in (s, t) -Wythoff's game, and a -Wythoff's game is a special case $s = 1$ and $t = a$ in (s, t) -Wythoff's game. Thus (s, t) -Wythoff's game is a generalization of both Wythoff's game and a -Wythoff's game.

Under normal play convention, the set $\mathcal{P}_{1,a}$ of all P -positions of a -Wythoff's game and the set $\mathcal{P}_{1,1}$ of all P -positions of Wythoff's game are given by letting $(s = 1$ and $t = a)$ and $s = t = 1$ in Eq. (2), respectively (see [9, 19]).

Under misère play convention, all P -positions of a -Wythoff's game were given in [13]. All P -positions of (s, t) -Wythoff's game were determined in [17], for all integers $s, t \geq 1$.

Other examples of extensions of Wythoff's game were given in [3, 12, 14, 15, 18].

1.3 Restriction of (s, t) -Wythoff's game

There are a few papers where only subsets of Wythoff's moves are allowed (see [5, 6, 10]). Such variations are called *restrictions* of Wythoff's game.

We now introduce a new *General Restriction of (s, t) -Wythoff's Game*. Let S_h, S_v, D_1 and D_2 be subsets of $Z^{\geq 0}$. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. One of the heaps is designated as "first heap" and the other as "second heap" throughout this game. By (x, y) we denote a position of present game, where x and y denote the numbers of tokens in the first and the second heaps, respectively. There are three types of moves.

(*Horizontal Move*) A player chooses the first heap and takes $k \in (\{w | 0 < w \leq x\} \cap S_h)$ tokens, i.e.,

$$(x, y) \rightarrow (x - k, y) \text{ and } k \in (\{w | 0 < w \leq x\} \cap S_h). \quad (3)$$

In this case, we call that (x, y) is moved to $(x - k, y)$ in *horizontal direction*, and k is called *horizontal distance*.

(*Vertical Move*) A player chooses the second heap and takes $\ell \in (\{z | 0 < z \leq y\} \cap S_v)$ tokens, i.e.,

$$(x, y) \rightarrow (x, y - \ell) \text{ and } \ell \in (\{z | 0 < z \leq y\} \cap S_v). \quad (4)$$

In this case, we call that (x, y) is moved to $(x, y - \ell)$ in *vertical direction*, and ℓ is called *vertical distance*.

(*Extended Diagonal Move*) A player takes tokens from both heaps, $k \in (\{w | 0 < w \leq x\} \cap D_1)$ from the first heap and $\ell \in (\{z | 0 < z \leq y\} \cap D_2)$ from the second heap, and

$$0 \leq |\ell - k| < (s - 1)\lambda + t, \quad \lambda = \min\{k, \ell\} \in Z^{\geq 1}. \quad (5)$$

In this case, we call that (x, y) is moved to $(x - k, y - \ell)$ in *extended diagonal direction*, k and ℓ are called *extended diagonal distance*.

Remark 1. Note that Eq. (1) is equivalent to

$$0 \leq \ell - k < (s - 1)k + t, \quad k \in Z^{\geq 1}, \quad (6)$$

and Eq. (6) is equivalent to Eq. (5).

Remark 2. (s, t) -Wythoff's game introduced by A.S. Fraenkel in [11] is equivalent to $S_h = S_v = D_1 = D_2 = Z^{\geq 0}$ in General Restriction of (s, t) -Wythoff's Game. Also a -Wythoff's game investigated in [9] is equivalent to $S_h = S_v = D_1 = D_2 = Z^{\geq 0}$, $s = 1$ and $t = a$ in General Restriction of (s, t) -Wythoff's Game. Wythoff's game investigated in [19] is equivalent to $S_h = S_v = D_1 = D_2 = Z^{\geq 0}$, $s = 1$ and $t = 1$ in General Restriction of (s, t) -Wythoff's Game.

Remark 3. In [5], the authors investigated the case of Wythoff's game, where “horizontal distance”, “vertical distance” and “diagonal distance” are bounded by a given positive integer R . This problem is equivalent to $S_h = S_v = D_1 = D_2 = \{n \leq R | n \in Z^{\geq 0}\}$ and $s = t = 1$ in General Restriction of (s, t) -Wythoff's Game. The set of all P -positions of this game under normal play convention was determined in [5].

Remark 4. In [5], the authors presented the following problems:

(1) One can investigate the case of Wythoff's game, where only “diagonal distance” is bounded. This problem is equivalent to $S_h = S_v = Z^{\geq 0}$, $D_1 = D_2 = \{n \leq R | n \in Z^{\geq 0}\}$ (R is a fixed positive integer) and $s = t = 1$ in General Restriction of (s, t) -Wythoff's Game.

(2) One can investigate the case of Wythoff's game, where “horizontal distance” and “vertical distance” are bounded, but “diagonal distance” is infinite. This problem is equivalent to $S_h = S_v = \{n \leq R | n \in Z^{\geq 0}\}$ (R is a fixed positive integer), $D_1 = D_2 = Z^{\geq 0}$ and $s = t = 1$ in General Restriction of (s, t) -Wythoff's Game. Under normal play convention, the set of all P -positions of this game was given in [16].

(3) One can investigate the bounded version of a -Wythoff's game, where “horizontal distance” and “vertical distance” are bounded, but “diagonal distance” is infinite. This problem is equivalent to $S_h = S_v = \{n \leq R | n \in Z^{\geq 0}\}$ (R is a fixed positive integer), $D_1 = D_2 = Z^{\geq 0}$ and $s = 1$, $t = a$ in General Restriction of (s, t) -Wythoff's Game; If “horizontal distance”, “vertical distance” and “diagonal distance” are bounded, then this problem is equivalent to $S_h = S_v = D_1 = D_2 = \{n \leq R | n \in Z^{\geq 0}\}$ (R is a fixed positive integer), $s = 1$ and $t = a$ in General Restriction of (s, t) -Wythoff's Game.

1.4 Our results

For all extensions and restrictions of Wythoff's game, our main goal is to find characterizations of P -positions, which almost always differs from the original Wythoff's sequence (see [7, 16]). In this paper, four models of General Restriction of (s, t) -Wythoff's Game are investigated. Let us now briefly present the content of this paper.

In Section 3, we define the first model, *Odd-Odd (s, t) -Wythoff's Game*, which is equivalent to $S_h = S_v = D_1 = D_2 = Z^{odd}$ in General Restriction of (s, t) -Wythoff's Game. Under normal play convention and for all $s, t \in Z^{\geq 1}$, the set of all P -positions is

given by $\bigcup_{m=0}^{\infty} \bigcup_{n=0}^{\infty} \{(2m, 2n)\}$; Under misère play convention and for all $s, t \in \mathbb{Z}^{\geq 1}$, the set of all P -positions is given by

$$\{(0, 2p+1), (2p+1, 0) | p \in \mathbb{Z}^{\geq 0}\} \cup \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{(2m, 2n)\}.$$

The structures of P -positions are of algebraic form (Theorems 7 and 9), which permit to decide in polynomial time whether or not a given game position (a, b) is a P -position.

In Section 4, we define the second model, *Even-Even (s, t) -Wythoff's Game*, which is equivalent to $S_h = S_v = D_1 = D_2 = \mathbb{Z}^{even}$ in General Restriction of (s, t) -Wythoff's Game. Under normal or misère play convention, and for all $s, t \geq 1$, the sets of all P -positions are given by recursive characterizations in term of mex function (Theorems 13 and 14).

In Section 5, we define the third model, *Odd-Even (s, t) -Wythoff's Game*, which is equivalent to $(S_h = D_1 = \mathbb{Z}^{odd} \text{ and } S_v = D_2 = \mathbb{Z}^{even})$ in General Restriction of (s, t) -Wythoff's Game. Under normal or misère play convention, and for all $s, t \in \mathbb{Z}^{\geq 1}$, the sets of all P -positions are given by algebraic characterizations (Theorems 17, 18, 19 and 20), which provide polynomial time procedures.

In Section 6, we define the fourth model, *Even-Odd (s, t) -Wythoff's Game*, which is equivalent to $(S_h = D_1 = \mathbb{Z}^{even} \text{ and } S_v = D_2 = \mathbb{Z}^{odd})$ in General Restriction of (s, t) -Wythoff's Game. Under normal or misère play convention, and for all $s, t \in \mathbb{Z}^{\geq 1}$, the sets of all P -positions are given by explicit formulas (Corollaries 21, 22, 23 and 24), which provide polynomial time procedures.

2 Preliminaries

Given any game Γ , we say informally that a P -position is any position u of Γ from which the *Previous* player can force a win, that is, the opponent of the player moving from u . An N -position is any position v of Γ from which the *Next* player can force a win, that is, the player who moves from v . The set of all P -positions of Γ is denoted by \mathcal{P} , and the set of all N -positions of Γ is denoted by \mathcal{N} . Denote by $Option(u)$ all options of u , i.e., the set of all positions that can be reached in one move from u . It follows from Fraenkel [8] that

$$\begin{aligned} u \in \mathcal{P} &\iff Option(u) \subseteq \mathcal{N}, \\ u \in \mathcal{N} &\iff Option(u) \cap \mathcal{P} \neq \emptyset. \end{aligned} \tag{7}$$

In order to better understand the legal moves of General Restriction of (s, t) -Wythoff's Game, we define the following notations.

By $Option_h(x, y)$ we denote the set of all positions that can be reached in one move in “horizontal direction” from a position (x, y) ;

By $Option_v(x, y)$ we denote the set of all positions that can be reached in one move in “vertical direction” from a position (x, y) ;

By $Option_e(x, y)$ we denote the set of all positions that can be reached in one move in “extended diagonal direction” from a position (x, y) .

It is obvious that for any position (x, y) ,

- (I) $Option(x, y) = Option_h(x, y) \cup Option_v(x, y) \cup Option_e(x, y)$;
- (II) $Option_h(x, y)$, $Option_v(x, y)$ and $Option_e(x, y)$ are pairwise disjoint.

Example 5. Eq. (7) can be used to check whether or not a given game position (a, b) is a P -position. We consider General Restriction of (s, t) -Wythoff’s Game under normal play convention and $s = t = 2$, where $S_h = S_v = D_1 = D_2 = Z^{\geq 0}$. Then $u = (1, 4)$ is a P -position.

Proof. By \mathcal{P} and \mathcal{N} we denote the sets of all P -positions and all N -positions, respectively. It is obvious that $(0, 0)$ is a P -position, i.e., $(0, 0) \in \mathcal{P}$.

(1) The positions $(0, 1), (0, 2), (0, 3), (0, 4)$ are N -positions. In fact, fix $m \in \{1, 2, 3, 4\}$ and let $w = (0, m)$, one can move $(0, m) \rightarrow (0, 0)$ by taking m tokens in the vertical direction. Thus $(0, 0) \in Option_v(w)$, i.e., $Option_v(w) \cap \mathcal{P} \neq \emptyset$. By Eq. (7), $(0, m)$ is an N -position.

(2) The position $(1, 0)$ is an N -position. For $w = (1, 0)$, one can move $(1, 0) \rightarrow (0, 0)$ by taking 1 tokens in the horizontal direction. Thus $(0, 0) \in Option_h(w)$, i.e., $Option_h(w) \cap \mathcal{P} \neq \emptyset$. By Eq. (7), $(1, 0)$ is an N -position.

(3) The positions $(1, 1), (1, 2), (1, 3)$ are N -positions. In fact, fix $m \in \{1, 2, 3\}$ and let $w = (1, m)$. For $w = (1, m)$, one can move $(1, m) \rightarrow (0, 0)$ by taking $k = 1$ token from the first heap and $\ell = m$ token from the second heap. Note that Eq. (5) is true:

$$|\ell - k| = m - 1 < 1 + 2 = (s - 1)\lambda + t, \lambda = k = 1.$$

(4) $Option_e(1, 4) = \{(0, 1), (0, 2), (0, 3)\}$. For $w = (1, 4)$, one can move $(1, 4) \rightarrow (0, m)$ with $1 \leq m \leq 3$ by taking $k = 1$ token from the first heap and $\ell = m$ tokens from the second heap, and $|\ell - k| = m - 1 < 1 + 2 = (s - 1)\lambda + t, \lambda = k = 1$.

(5) It is obvious that $Option_h(1, 4) = \{(0, 4)\}$, $Option_v(1, 4) = \{(1, m) | 0 \leq m \leq 3\}$. Thus

$$\begin{aligned} Option(1, 4) &= Option_h(1, 4) \cup Option_v(1, 4) \cup Option_e(1, 4) \\ &= \{(0, 4)\} \cup \{(1, m) | 0 \leq m \leq 3\} \cup \{(0, 1), (0, 2), (0, 3)\}. \end{aligned}$$

It follows from (1), (2) and (3) that $Option(1, 4) \subseteq \mathcal{N}$. By Eq. (7), the position $(1, 4)$ is a P -position. \square

Proposition 6. ([7], *Characterization of the P-positions of an impartial acyclic game*)
The sets of P - and N -positions of any impartial acyclic game (like Wythoff’s game) are uniquely determined by the following two properties:

- Any move from a P -position leads to an N -position (stability property of the P -positions).
- From any N -position, there exists a move leading to a P -position (absorbing property of the P -positions).

Proof. See Proposition 1 in [7]. \square

3 Odd-Odd (s, t) -Wythoff's Game

In this section, we introduce a new *Odd-Odd (s, t) -Wythoff's Game* (Denoted by OOW). Let S_h, S_v, D_1 and D_2 be subsets of $Z^{\geq 0}$. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. One of the heaps is designated as “first heap” and the other as “second heap” throughout this game. By (x, y) we denote a position of present game, where x and y denote the numbers of tokens in the first and the second heaps, respectively. Two rules of moves are allowed:

(*Odd-Odd Nim Rule*) A player chooses one heap and takes an arbitrary *odd* number k of tokens.

(*Odd-Odd More General Wythoff's Rule*) A player takes tokens from both heaps, *odd* $k > 0$ tokens from the first heap, *odd* $\ell > 0$ tokens from the second heap, and

$$0 \leq |\ell - k| < (s - 1)\lambda + t, \quad \lambda = \min\{k, \ell\} \in Z^{\geq 1}. \quad (8)$$

Obviously, OOW is equivalent to $S_h = S_v = D_1 = D_2 = Z^{odd}$ in General Restriction of (s, t) -Wythoff's Game.

By the definition of OOW, the positions (x, y) and (y, x) are equivalent, i.e., both (x, y) and (y, x) are P -positions, or are N -positions. Theorems 7 and 9 will give the sets of all P -positions of OOW under normal or misère play convention, respectively. The corresponding winning strategies are also presented.

We define a function δ_n for $n \in Z^{\geq 0}$:

$$\delta_n = \begin{cases} 0, & \text{if } n \text{ is even,} \\ 1, & \text{if } n \text{ is odd.} \end{cases}$$

Theorem 7. By \mathcal{P}_1 we denote the set of all P -positions of OOW. Then for all $s, t \in Z^{\geq 1}$,

$$\mathcal{P}_1 = \bigcup_{m=0}^{\infty} \bigcup_{n=0}^{\infty} \{(2m, 2n)\}.$$

Proof. Let $\mathcal{M}_1 = \bigcup_{m=0}^{\infty} \bigcup_{n=0}^{\infty} \{(2m, 2n)\}$. It suffices to show two things:

Fact A. No options of a position in \mathcal{M}_1 can be in \mathcal{M}_1 .

Fact B. Any position not in \mathcal{M}_1 can land in a position in \mathcal{M}_1 .

Proof of Fact A. Let $(x, y) \in \mathcal{M}_1$ be a position. Suppose that $(x, y) \rightarrow (x', y') \in \mathcal{M}_1$. By the definition of \mathcal{M}_1 , x, y, x' and y' are even. Thus both $x - x'$ and $y - y'$ are even. This contradicts the rules of moves of OOW.

Proof of Fact B. Let $(x, y) \notin \mathcal{M}_1$ be a position. In this case, at least one of x and y is odd, i.e., $(\delta_x, \delta_y) = (0, 1)$ or $(1, 0)$ or $(1, 1)$. Thus we can move $(x, y) \rightarrow (x - \delta_x, y - \delta_y) \in \mathcal{M}_1$ by taking one token from an odd-size heap.

The proof is completed. □

Remark 8. Given a game Γ . Let \mathcal{M} be the set of all P -positions of game Γ . The following facts are true:

Fact 1. No options of a position in \mathcal{M} can be in \mathcal{M} .

Fact 2. Any position not in \mathcal{M} can land in a position in \mathcal{M} by a legal move.

We will determine the sets of all P -positions of the games investigated in this paper, respectively. In all proofs, the validity of *Fact 1* and *Fact 2* will be proved. The method of the proofs is the same, though the proofs themselves vary greatly.

Theorem 9. *By \mathcal{P}_2 we denote the set of all P -positions of OOW under misère play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$,*

$$\mathcal{P}_2 = \{(0, 2p+1), (2p+1, 0) | p \in \mathbb{Z}^{\geq 0}\} \cup \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{(2m, 2n)\}.$$

Proof. Let

$$\begin{aligned} \mathcal{M}_2 &= \{(0, 2p+1), (2p+1, 0) | p \in \mathbb{Z}^{\geq 0}\} \cup \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{(2m, 2n)\}, \\ \mathcal{M}'_2 &= \{(0, 2p+1), (2p+1, 0) | p \in \mathbb{Z}^{\geq 0}\}, \\ \mathcal{M}''_2 &= \bigcup_{m=1}^{\infty} \bigcup_{n=1}^{\infty} \{(2m, 2n)\}. \end{aligned}$$

Proof of Fact 1. Let (x, y) be a position in \mathcal{M}_2 . For $(0, 2p+1) \in \mathcal{M}'_2$, $(0, 2p+1) \rightarrow (x', y') \in \mathcal{M}''_2$ is impossible since $x' > 0$; $(0, 2p+1) \rightarrow (0, 2q+1) (q < p)$ is also impossible, since $2(p-q)$ is even.

For $(2m, 2n) \in \mathcal{M}''_2$, $(2m, 2n) \rightarrow (0, 2p+1)$ (or $(2p+1, 0)$) is impossible, since $2m$ (or $2n$) is even; $(2m, 2n) \rightarrow (2m', 2n') \in \mathcal{M}''_2$ is also impossible, since $2(m-m')$ is even, which contradicts the rules of moves of OOW.

Proof of Fact 2. Let (x, y) with $x \leq y$ be a position not in \mathcal{M}_2 .

If $(x, y) = (0, 2v)$ for some $v \in \mathbb{Z}^{\geq 1}$, we move $(x, y) = (0, 2v) \rightarrow (0, 2v-1)$ by taking one token from the heap of size $2v$. If $(x, y) = (0, 0)$ then next player wins without doing any thing.

If $(x, y) = (2m, 2n+1)$ for some $m, n \in \mathbb{Z}^{\geq 1}$ and $n \geq m$, we move $(2m, 2n+1) \rightarrow (2m, 2n)$, by taking one token from the heap of size $2n+1$.

If $(x, y) = (2u+1, w)$ for some $u \in \mathbb{Z}^{\geq 0}$ and $w \geq 2u+1$. We need to consider two subcases:

(i) $u = 0$. In this case, $w \geq 1$. If w is odd, we move $(2u+1, w) = (1, w) \rightarrow (0, w) \in \mathcal{M}'_2$; If w is even, we move $(2u+1, w) = (1, w) \rightarrow (0, w-1) \in \mathcal{M}'_2$, by taking one token from each heap.

(ii) $u > 0$. In this case, $w \geq 2u+1 \geq 3$. If w is odd, we move $(2u+1, w) \rightarrow (2u, w-1) \in \mathcal{M}''_2$. If w is even, thus we move $(2u+1, w) \rightarrow (2u, w) \in \mathcal{M}''_2$.

The proof is completed. \square

4 Even-Even (s, t) -Wythoff's Game

In this section, we introduce a new *Even-Even (s, t) -Wythoff's Game* (Denoted by EEW). Let S_h, S_v, D_1 and D_2 be subsets of $Z^{\geq 0}$. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. One of the heaps is designated as “first heap” and the other as “second heap” throughout this game. By (x, y) we denote a position of present game, where x and y denote the numbers of tokens in the first and the second heaps, respectively. Two rules of moves are allowed:

(*Even-Even Nim Rule*) A player chooses one heap and takes an arbitrary *even* number $k > 0$ of tokens.

(*Even-Even More General Wythoff's Rule*) A player takes tokens from both heaps, *even* $k > 0$ tokens from the first heap, *even* $\ell > 0$ tokens from the second heap and

$$0 \leq |\ell - k| < (s - 1)\lambda + t, \quad \lambda = \min\{k, \ell\} \in Z^{\geq 1}. \quad (9)$$

Obviously, EEW is equivalent to $S_h = S_v = D_1 = D_2 = Z^{\text{even}}$ in General Restriction of (s, t) -Wythoff's Game.

Remark 10. The “symmetric” notation $\{x, y\}$ for unordered pairs of non-negative integers is used whenever the positions (x, y) and (y, x) are equivalent, i.e., both (x, y) and (y, x) are P -positions, or are N -positions.

Example 11. We consider EEW under normal play convention. Fix two integers $s = t = 1$. It is obvious that $(0, 1)$ is a P -position, $(1, 0)$ is also a P -position. Thus we use $\{0, 1\}$ to denote two positions $(0, 1)$ and $(1, 0)$, i.e., $\{0, 1\} = \{(0, 1), (1, 0)\}$. Generally, by the definition of EEW, two positions (x, y) and (y, x) are equivalent, i.e., both (x, y) and (y, x) are P -positions, or are N -positions. Thus we use $\{x, y\}$ to denote two positions (x, y) and (y, x) , i.e., $\{x, y\} = \{(x, y), (y, x)\}$.

Theorems 13 and 14 will give the sets of all P -positions of EEW under normal or misère play convention, respectively. The corresponding winning strategies are also presented. Before the main results, we define two sequences and give some properties in Lemma 12.

We define two sequences A_n and B_n for $n \in Z^{\geq 0}$ and all $s, t \in Z^{\geq 1}$:

$$\begin{cases} A_n = \text{mex}\{A_i, A_i + 1, B_i, B_i + 1 | 0 \leq i < n\}, \\ B_n = sA_n + (t + \delta_t)n. \end{cases} \quad (10)$$

Tables 1 and 2 list the first few values of A_n and B_n for $s = t = 1$ and $s = t = 2$, respectively.

Table 1. The first few values of A_n and B_n for $s = t = 1$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	2	6	8	12	16	18	22	24	28	32	34	38	42	44
B_n	0	4	10	14	20	26	30	36	40	46	52	56	62	68	72

Table 2. The first few values of A_n and B_n for $s = t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	2	4	8	10	14	16	18	20	24	26	30	32	34	36
B_n	0	6	12	22	28	38	44	50	56	66	72	82	88	94	100

Lemma 12. Let $\{A_n\}_{n=0}^\infty$ and $\{B_n\}_{n=0}^\infty$ be defined by Eq. (10). We have the following properties:

- (I) Both A_n and B_n are even for $n \geq 0$.
- (II) Both A_n and B_n are strictly increasing sequences for $n \geq 0$.
- (III) $B_n > A_n + 1 > A_n$ for $n \geq 1$.
- (IV) Let

$$\begin{cases} A = \bigcup_{n=1}^\infty \{A_n\} \cup \bigcup_{n=1}^\infty \{A_n + 1\}, \\ B = \bigcup_{n=1}^\infty \{B_n\} \cup \bigcup_{n=1}^\infty \{B_n + 1\}. \end{cases} \quad (11)$$

Then A, B are complementary with respect to $Z^{\geq 2}$, i.e., $A \cup B = Z^{\geq 2}$ and $A \cap B = \emptyset$.

Proof. (I) Note that $t + \delta_t$ is even for $t \in Z^{\geq 1}$. We proceed by induction on n . Obviously, $A_0 = B_0 = 0$, $A_1 = 2$ and $B_1 = sA_1 + (t + \delta_t)$ are even. Suppose $m < n$, both A_m and B_m are even. We now show that A_n is even, and then $B_n = sA_n + (t + \delta_t)n$ is even.

Indeed, suppose that A_n is odd. Let $k = A_n$ and $S = \{A_i, A_i + 1, B_i, B_i + 1 \mid 0 \leq i < n\}$. By Eq. (10), the fact $k = \text{mex}(S)$ implies that $k \notin S$ and $k - 1 \in S$.

By the hypothesis of induction, A_i and B_i are even for all $i \in \{0, 1, \dots, n - 1\}$. Note that the facts $k - 1 \in S$ and $k - 1$ is even imply that $k - 1 \neq A_i + 1$ or $B_i + 1$. If there exists an integer $i_0 \in \{0, 1, \dots, n - 1\}$ such that $k - 1 = A_{i_0}$ or B_{i_0} , then $k \in S$. This contradicts $k \notin S$.

(II) By the definition of A_n and mex property, A_n is strictly increasing sequence. Also B_n is strictly increasing sequence. Indeed, for $m > n$,

$$B_m - B_n = s(A_m - A_n) + (t + \delta_t)(m - n) > 0.$$

(III) Note that $t + \delta_t \geq 2$ for $t \in Z^{\geq 1}$. By Eq. (10), we have $B_n = sA_n + (t + \delta_t)n \geq A_n + 2n > A_n + 1 > A_n$, for $n \in Z^{\geq 1}$.

(IV) In fact, $A \cup B = Z^{\geq 2}$ follows from the mex property and $A_0 = 0$, $B_0 = 0$, $A_1 = 2$. Suppose $A \cap B \neq \emptyset$. It follows from (I) that $A_m + 1 \neq B_n$ and $A_m \neq B_n + 1$, thus the only possibility is $A_m = B_n$ for two integers $m, n \in Z^{\geq 1}$. If $m > n$, then A_m is mex of a set containing $B_n = A_m$, a contradiction. If $m \leq n$, then by (II) we have $B_n = sA_n + (t + \delta_t)n \geq sA_m + (t + \delta_t)m > A_m$, another contradiction.

The proof is completed. \square

Theorem 13. By \mathcal{P}_3 we denote the set of all P -positions of EEW under normal play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$,

$$\mathcal{P}_3 = \bigcup_{i=0}^{\infty} \left\{ \begin{array}{l} \{A_i, B_i\}, \{A_i, B_i + 1\}, \\ \{A_i + 1, B_i\}, \{A_i + 1, B_i + 1\} \end{array} \right\},$$

where A_n and B_n are defined by Eq. (10).

Proof. Before we give the proof of Theorem 13, Tables 3 and 4 list the first few values of A_n and B_n , which show us how to determine \mathcal{P}_3 by using Theorem 13:

Table 3. The first few values of A_n and B_n for $s = t = 1$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	2	6	8	12	16	18	22	24	28	32	34	38	42	44
$A_n + 1$	1	3	7	9	13	17	19	23	25	29	33	35	39	43	45
B_n	0	4	10	14	20	26	30	36	40	46	52	56	62	68	72
$B_n + 1$	1	5	11	15	21	27	31	37	41	47	53	57	63	69	73

For $s = t = 1$, it follows from Table 3 that

$$\mathcal{P}_3 = \left\{ \begin{array}{l} (0, 0), (0, 1), (1, 0), (1, 1); \\ (2, 4), (2, 5), (3, 4), (3, 5); (4, 2), (5, 2), (4, 3), (5, 3); \\ (6, 10), (6, 11), (7, 10), (7, 11); (10, 6), (11, 6), (10, 7), (11, 7); \dots \end{array} \right\}$$

Table 4. The first few values of A_n and B_n for $s = t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	2	4	8	10	14	16	18	20	24	26	30	32	34	36
$A_n + 1$	1	3	5	9	11	15	17	19	21	25	27	31	33	35	37
B_n	0	6	12	22	28	38	44	50	56	66	72	82	88	94	100
$B_n + 1$	1	7	13	23	29	39	45	51	57	67	73	83	89	95	101

For $s = t = 2$, it follows from Table 4 that

$$\mathcal{P}_3 = \left\{ \begin{array}{l} (0, 0), (0, 1), (1, 0), (1, 1); \\ (2, 6), (2, 7), (3, 6), (3, 7); (6, 2), (7, 2), (6, 3), (7, 3); \\ (4, 12), (4, 13), (5, 12), (5, 13); (12, 4), (13, 4), (12, 5), (13, 5); \dots \end{array} \right\}$$

We now give the proof of Theorem 13. Let

$$\mathcal{M}_3 = \bigcup_{i=0}^{\infty} \left\{ \begin{array}{l} \{A_i, B_i\}, \{A_i, B_i + 1\}, \\ \{A_i + 1, B_i\}, \{A_i + 1, B_i + 1\} \end{array} \right\}.$$

Proof of Fact 1. Let (x, y) with $x \leq y$ be a position in \mathcal{M}_3 . It follows from (III) of Lemma 12 that there exists an integer $n \geq 0$ such that $x = A_n$ or $A_n + 1$, and $y = B_n$ or $B_n + 1$.

Suppose that $(x, y) \rightarrow (x', y) \in \mathcal{M}_3$ with $x' = A_m$ or $A_m + 1$, by Even-Even Nim Rule. Then the fact $x' \leq x - 2 < A_n$ implies that $m < n$. Thus $(x', y) \notin \mathcal{M}_3$, a contradiction.

Suppose that $(x, y) \rightarrow (x, y') \in \mathcal{M}_3$ with $y' = B_m$ or $B_m + 1$, by Even-Even Nim Rule. Then $y' \leq y - 2 < B_n$ implies that $m < n$. Thus $(x, y') \notin \mathcal{M}_3$, another contradiction.

Suppose that $(x, y) \rightarrow (x', y') \in \mathcal{M}_3$ by Even-Even More General Wythoff's Rule. Then $x - x' > 0$ is even and $y - y' > 0$ is even. It follows from (I) and (III) of Lemma 12 that $k = x - x' = A_n - A_m$, $\ell = y - y' = B_n - B_m$ and $m < n$. Thus

$$0 < k \leq \ell = s(A_n - A_m) + (t + \delta_t)(m - n) \geq sk + t,$$

which contradicts Eq. (9).

Proof of Fact 2. Let (x, y) be a position not in \mathcal{M}_3 . Without loss of generality, assume that $0 \leq x \leq y$.

If $x = 0$ or 1 , we move $(x, y) \rightarrow (x, \delta_y) \in \mathcal{M}_3$. This move is legal, since $y - \delta_y$ is even and $(x, y) \notin \mathcal{M}_3$ implies that $y > 1 \geq \delta_y$.

If $x \geq 2$ then the integer x appears exactly once in exactly one of A and B , since A and B are complementary with respect to $Z^{\geq 2}$ (Lemma 12, (IV)). Therefore, we have one of the following two cases: (i) $x = B_n$ or $x = B_n + 1$, (ii) $x = A_n$ or $x = A_n + 1$, for some $n \geq 1$.

Case (i): $x = B_n$ or $x = B_n + 1$, $n \geq 1$. We move

$$(x, y) \rightarrow (x, A_n + \delta_y) \in \mathcal{M}_3,$$

i.e., we take $y - A_n - \delta_y$ tokens from the heap of y tokens. It follows from (I) and (III) of Lemma 12 that

$$y \geq x \geq B_n > A_n + 1 \geq A_n + \delta_y$$

and $y - A_n - \delta_y$ is even. Thus the above move is legal.

Case (ii): $x = A_n$ or $x = A_n + 1$, $n \geq 1$. In this case, we have $y > B_n + 1$ or $x \leq y < B_n$.

(1) $y > B_n + 1$. We move

$$(x, y) \rightarrow (x, B_n + \delta_y) \in \mathcal{M}_3,$$

i.e., we take $y - B_n - \delta_y$ tokens from the heap of y tokens. This is a legal move, since $y > B_n + 1 \geq B_n + \delta_y$ and $y - B_n - \delta_y$ is even.

(2) $x \leq y < B_n$. We distinguish the following two subcases: $x \leq y < sA_n + t + \delta_t$ or $sA_n + t + \delta_t \leq y < B_n$:

(2.1) $x \leq y < sA_n + t + \delta_t$. We move

$$(x, y) \rightarrow (x - A_n, \delta_y) \in \mathcal{M}_3,$$

since $x - A_n = 0$ or 1 , and $\delta_y = 0$ or 1 . This move is legal:

- 1) $k = A_n$ is even;
- 2) $\ell = y - \delta_y$ is even;

3) $y \geq x$ implies that $\ell = y - \delta_y \geq A_n = k$. Note that $y < sA_n + t + \delta_t$ and $sA_n + t + \delta_t$ is even, so $y \leq sA_n + t + \delta_t - 2 + \delta_y$. Hence,

$$\begin{aligned} |\ell - k| &= y - \delta_y - A_n \\ &\leq (s-1)A_n + t + \delta_t - 2 \\ &< (s-1)A_n + t. \end{aligned}$$

(2.2) $sA_n + t + \delta_t \leq y < B_n$. Put $m = \lfloor \frac{y-sA_n-\delta_y}{t+\delta_t} \rfloor$. We move

$$(x, y) \rightarrow (x - A_n + A_m, B_m + \delta_y) \in \mathcal{M}_3,$$

since $x - A_n = 0$ or 1 , and $\delta_y = 0$ or 1 . This move is legal:

(a) $k = A_n - A_m > 0$ and k is even. Firstly, we show that $0 \leq m < n$. Note that $y - sA_n \geq t + \delta_t \geq 2 > \delta_y$, so $\frac{y-sA_n-\delta_y}{t+\delta_t} > 0$. Thus $m = \lfloor \frac{y-sA_n-\delta_y}{t+\delta_t} \rfloor \geq 0$. On the other hand, the facts $y < B_n$ and B_n is even imply that $y - \delta_y < B_n$. Thus

$$y - sA_n - \delta_y < B_n - sA_n = (t + \delta_t)n,$$

and

$$m = \lfloor \frac{y - sA_n - \delta_y}{t + \delta_t} \rfloor \leq \frac{y - sA_n - \delta_y}{t + \delta_t} < n.$$

It follows from (I) and (II) of Lemma 12 that $k = A_n - A_m > 0$ and k is even.

(b) $\ell = y - B_m - \delta_y > 0$ and ℓ is even. By the definition of m , we have $m \leq \frac{y-sA_n-\delta_y}{t+\delta_t}$ and

$$\begin{aligned} y &\geq (t + \delta_t)m + sA_n + \delta_y \\ &= B_m + \delta_y + s(A_n - A_m) \\ &> B_m + \delta_y. \end{aligned}$$

Thus $\ell = y - B_m - \delta_y > 0$ and ℓ is even.

(c) $|\ell - k| < (s-1)k + t$. By the definition of m , we have $m > \frac{y-sA_n-\delta_y}{t+\delta_t} - 1$ and $y < (t + \delta_t)(m + 1) + sA_n + \delta_y$. Thus

$$\begin{aligned} \ell &= y - B_m - \delta_y \\ &< (t + \delta_t)(m + 1) + sA_n - sA_m - (t + \delta_t)m \\ &= s(A_n - A_m) + t + \delta_t. \end{aligned}$$

We note that $y - B_m - \delta_y$ and $s(A_n - A_m) + t + \delta_t$ are even, so

$$\begin{aligned} \ell &= y - B_m - \delta_y \\ &\leq s(A_n - A_m) + t + \delta_t - 2 \\ &< s(A_n - A_m) + t; \end{aligned}$$

On the other hand, $y - B_m - \delta_y \geq s(A_n - A_m) \geq A_n - A_m$ by virtue of (b). Therefore, $|\ell - k| < (s-1)k + t$.

The proof is completed. □

Theorem 14. By \mathcal{P}_4 we denote the set of all P -positions of EEW under misère play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$,

$$\mathcal{P}_4 = \bigcup_{i=0}^{\infty} \left\{ \begin{array}{l} \{E_i, H_i\}, \{E_i, H_i + 1\}, \\ \{E_i + 1, H_i\}, \{E_i + 1, H_i + 1\} \end{array} \right\},$$

where E_n and H_n are given by the following two cases:

(A) If $s \neq 1$ or $t > 2$, then for $n \geq 0$,

$$\begin{cases} E_n = \text{mex}\{E_i, E_i + 1, H_i, H_i + 1 | 0 \leq i < n\}, \\ H_n = sE_n + (t + \delta_t)n + 2. \end{cases} \quad (12)$$

(B) If $s = 1$ and $t \in \{1, 2\}$, then $E_0 = H_0 = 4$ and for $n \geq 1$,

$$\begin{cases} E_n = \text{mex}\{E_i, E_i + 1, H_i, H_i + 1 | 0 \leq i < n\}, \\ H_n = E_n + 2n. \end{cases} \quad (13)$$

Proof. Before we give the proof of Theorem 14, Tables 5 and 6 list the first few values of E_n and H_n , which show us how to determine \mathcal{P}_4 by using Theorem 14:

Table 5. The first few values of E_n and H_n for $s = t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
E_n	0	4	6	8	10	14	16	20	22	26	28	32	34	36	38
H_n	2	12	18	24	30	40	46	56	62	72	78	88	94	100	106

For $s = t = 2$, it follows from Table 5 that

$$\mathcal{P}_4 = \left\{ \begin{array}{l} (0, 2), (0, 3), (1, 2), (1, 3); (2, 0), (3, 0), (2, 1), (3, 1); \\ (4, 12), (4, 13), (5, 12), (5, 13); (12, 4), (13, 4), (12, 5), (13, 5); \\ (6, 18), (6, 19), (7, 18), (7, 19); (18, 6), (19, 6), (18, 7), (19, 7); \dots \end{array} \right\}$$

Table 6. The first few values of E_n and H_n for $s = t = 1$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
E_n	4	0	6	8	12	16	18	22	24	28	32	34	38	42	44
H_n	4	2	10	14	20	26	30	36	40	46	52	56	62	68	72

For $s = t = 1$, it follows from Table 6 that

$$\mathcal{P}_4 = \left\{ \begin{array}{l} (4, 4), (4, 5), (5, 4), (5, 5); \\ (0, 2), (0, 3), (1, 2), (1, 3); (2, 0), (3, 0), (2, 1), (3, 1); \\ (6, 10), (6, 11), (7, 10), (7, 11); (10, 6), (11, 6), (10, 7), (11, 7); \dots \end{array} \right\}$$

We now give the proof of Theorem 14. Let

$$\mathcal{M}_4 = \bigcup_{i=0}^{\infty} \left\{ \begin{array}{l} \{E_i, H_i\}, \{E_i, H_i + 1\}, \\ \{E_i + 1, H_i\}, \{E_i + 1, H_i + 1\} \end{array} \right\},$$

and $E = \bigcup_{n=0}^{\infty} \{E_n\} \cup \bigcup_{n=0}^{\infty} \{E_n + 1\}$, $H = \bigcup_{n=0}^{\infty} \{H_n\} \cup \bigcup_{n=0}^{\infty} \{H_n + 1\}$. Then we claim that:

Fact A.1 If $s \neq 1$ or $t > 2$, then both E_n and H_n are even for $n \in \mathbb{Z}^{\geq 0}$, and both E_n and H_n are strictly increasing sequences for $n \geq 0$. The proofs are similar to ones of (I) and (II) of Lemma 12.

Fact A.2 If $s \neq 1$ or $t > 2$, then $E \cup H = \mathbb{Z}^{\geq 0}$ and $E \cap H = \emptyset$. In fact, $E \cup H = \mathbb{Z}^{\geq 0}$ follows from the definition of mex . Suppose $E \cap H \neq \emptyset$. It follows from Fact A.1 that $E_m + 1 = H_n$ and $E_m = H_n + 1$ are impossible, thus there exist two integers $m, n \in \mathbb{Z}^{\geq 0}$ such that $E_m = H_n$. If $m > n$ then $E_m = \text{mex}\{E_i, E_i + 1, H_i, H_i + 1 \mid 0 \leq i < m\}$, which contradicts $E_m = H_n$; If $m \leq n$ then

$$H_n = sE_n + (t + \delta_t)n + 2 \geq sE_m + (t + \delta_t)m + 2 > E_m,$$

which also contradicts $E_m = H_n$.

Fact B.1 If $s = 1$ and $t \in \{1, 2\}$, then both E_n and H_n are even for $n \in \mathbb{Z}^{\geq 0}$, both E_n and H_n are strictly increasing sequences for $n \geq 1$. Indeed,

$$E_1 = 0 < 6 = E_2 < 8 = E_3 < 12 = E_4 < 16 = E_5 < \dots$$

and

$$H_1 = 2 < 10 = H_2 < 14 = H_3 < 20 = E_4 < 26 = E_5 < \dots$$

Fact B.2 If $s = 1$ and $t \in \{1, 2\}$, then $E \cup H = \mathbb{Z}^{\geq 0}$ and $E \cap H = \{4\}$. Its proof is similar to those of Fact A.2.

Proof of Fact 1. Let (x, y) with $x \leq y$ be a position in \mathcal{M}_4 . By the definition of \mathcal{M}_4 , there exists an integer $n \in \mathbb{Z}^{\geq 0}$ such that $x = E_n$ or $E_n + 1$, and $y = H_n$ or $H_n + 1$.

Suppose that $(x, y) \rightarrow (x', y) \in \mathcal{M}_4$ with $x' = E_m$ or $E_m + 1$, by Even-Even Nim Rule. Then the fact $x' \leq x - 2 < E_n$ implies that $m < n$. Thus $(x', y) \notin \mathcal{M}_4$, a contradiction. Similarly, $(x, y) \rightarrow (x, y') \in \mathcal{M}_4$ is also impossible.

Suppose that $(x, y) \rightarrow (x', y') \in \mathcal{M}_4$ by Even-Even More General Wythoff's Rule. Then $x - x' > 0$ is even and $y - y' > 0$ is even. It follows from Facts A.1 and B.1 that $k = x - x' = E_n - E_m$, $\ell = y - y' = H_n - H_m$ and $m < n$. Thus

$$0 < k \leq \ell = s(E_n - E_m) + (t + \delta_t)(m - n) \geq sk + t,$$

which contradicts Eq. (9).

Proof of Fact 2. Let (x, y) be a position not in \mathcal{M}_4 . Without loss of generality, assume that $0 \leq x \leq y$. By Facts A.2 and B.2, we have one of the following two cases: (i) $x = H_n$ or $x = H_n + 1$, (ii) $x = E_n$ or $x = E_n + 1$, for some $n \geq 0$.

Case (i): $x = H_n$ or $x = H_n + 1$, $n \geq 0$. In this case, the fact $(x, y) \notin \mathcal{M}_4$ implies that $y > E_n + 1$. In fact, if $n = 0$, then $y \geq x \geq H_0 \geq E_0$, thus $y > E_0 + 1$; For $n \geq 1$, if $y \leq E_n + 1$, then it follows from Eqs (12) and (13) that

$$x \geq H_n \geq E_n + 2 > E_n + 1 \geq y,$$

a contradiction.

We move

$$(x, y) \rightarrow (x, E_n + \delta_y) \in \mathcal{M}_4,$$

i.e., we take $y - E_n - \delta_y$ tokens from the heap of y tokens. Note that $y - E_n - \delta_y > 1 - \delta_y \geq 0$ and $y - E_n - \delta_y$ is even, thus the above move is legal.

Case (ii): $x = E_n$ or $x = E_n + 1$, $n \geq 0$. The fact $(x, y) \notin \mathcal{M}_4$ implies that $y > H_n + 1$ or $x \leq y < H_n$.

(1) $y > H_n + 1$. We move

$$(x, y) \rightarrow (x, H_n + \delta_y) \in \mathcal{M}_4,$$

i.e., we take $y - H_n - \delta_y$ tokens from the heap of y tokens. This is a legal move, since $y - H_n - \delta_y > 1 - \delta_y \geq 0$ and $y - H_n - \delta_y$ is even.

(2) $x \leq y < H_n$. We need to consider the situations for (A) and for (B), respectively.

(2-A) $s \neq 1$ or $t > 2$. If $n = 0$, then $0 \leq x \leq y < 2$ and hence $(x, y) = (0, 0)$ or $(0, 1)$ or $(1, 1)$. The next player wins without doing anything. It remains to consider $n \geq 1$. We proceed by distinguishing the following two subcases: (2-A-1) $x \leq y < sE_n + t + \delta_t + 2$; (2-A-2) $sE_n + t + \delta_t + 2 \leq y < H_n$.

(2-A-1) $x \leq y < sE_n + t + \delta_t + 2$. In this subcase, we move

$$(x, y) \rightarrow (x - E_n, 2 + \delta_y) \in \mathcal{M}_4,$$

since $x - E_n \in \{0, 1\} = \{E_0, E_0 + 1\}$ and $2 + \delta_y \in \{H_0, H_0 + 1\}$. Note that $sE_n + t + \delta_t$ is even, so $y < sE_n + t + \delta_t + 1 + \delta_y$. This move is legal:

1) $k = E_n > 0$ is even;

2) $\ell = y - 2 - \delta_y$ is even. Note that $y \geq x \geq E_n$ and E_n is even, thus $y - \delta_y \geq E_n$ and $\ell = y - 2 - \delta_y \geq E_n - 2 \geq E_1 - 2 > 0$;

3) It is easy to see that

$$\begin{aligned} |\ell - k| &= |y - 2 - \delta_y - E_n| \\ &< (s - 1)E_n + t + \delta_t - 1 \\ &\leq (s - 1)E_n + t. \end{aligned}$$

(2-A-2) $sE_n + t + \delta_t + 2 \leq y < H_n$. Put

$$m = \left\lfloor \frac{y - sE_n - 2 - \delta_y}{t + \delta_t} \right\rfloor. \quad (14)$$

We move

$$(x, y) \rightarrow (x - E_n + E_m, H_m + \delta_y) \in \mathcal{M}_4, \quad (15)$$

since $x - E_n + E_m \in \{E_m, E_m + 1\}$ and $H_m + \delta_y \in \{H_m, H_m + 1\}$. This move is legal:

(A-a) $k = E_n - E_m > 0$ and k is even. Firstly, we show that $0 \leq m < n$. Note that

$$y - sE_n - 2 \geq t + \delta_t \geq 2 > \delta_y,$$

so $\frac{y-sE_n-2-\delta_y}{t+\delta_t} > 0$. Thus $m = \lfloor \frac{y-sE_n-2-\delta_y}{t+\delta_t} \rfloor \geq 0$; On the other hand, the fact $y < H_n$ implies that $y - \delta_y < H_n$. Thus

$$y - sE_n - 2 - \delta_y < H_n - sE_n = (t + \delta_t)n,$$

and

$$m = \lfloor \frac{y - sE_n - 2 - \delta_y}{t + \delta_t} \rfloor \leq \frac{y - sE_n - 2 - \delta_y}{t + \delta_t} < n.$$

By Facts A.1 and A.2, $k = E_n - E_m > 0$ and k is even.

(A-b) $\ell = y - H_m - \delta_y > 0$ and ℓ is even. By the definition of m , we have $m \leq \frac{y-sE_n-2-\delta_y}{t+\delta_t}$ and

$$\begin{aligned} y &\geq (t + \delta_t)m + sE_n + 2 + \delta_y \\ &= H_m + \delta_y + s(E_n - E_m) \\ &> H_m + \delta_y. \end{aligned}$$

Thus $\ell = y - H_m - \delta_y > 0$ and ℓ is even.

(A-c) $0 \leq \ell - k < (s - 1)k + t$. By the definition of m , we have

$$m > \frac{y - sE_n - 2 - \delta_y}{t + \delta_t} - 1,$$

i.e.,

$$y < (t + \delta_t)(m + 1) + sE_n + 2 + \delta_y.$$

By Eq. (12), we have $H_m = sE_m + (t + \delta_t)m + 2$. Thus

$$\begin{aligned} \ell &= y - H_m - \delta_y \\ &< (t + \delta_t)(m + 1) + sE_n - sE_m - (t + \delta_t)m \\ &= s(E_n - E_m) + t + \delta_t. \end{aligned}$$

Note that $y - H_m - \delta_y$ and $s(E_n - E_m) + t + \delta_t$ are even, so

$$y - H_m - \delta_y \leq s(E_n - E_m) + t + \delta_t - 2 < s(E_n - E_m) + t;$$

On the other hand,

$$\ell = y - H_m - \delta_y \geq s(E_n - E_m) \geq E_n - E_m = k$$

by virtue of (A-b). Therefore, $0 \leq \ell - k < (s - 1)k + t$.

(2-B) $s = 1$ and $t \in \{1, 2\}$. Note that $x \leq y < H_n$. In this case, $n = 0$ is impossible; If $n = 1$ then $0 = E_1 \leq x \leq y < H_1 = 2$, i.e., $(x, y) = (0, 0)$ or $(0, 1)$, or $(1, 1)$. The next player wins without doing anything. It remains to consider $n \geq 2$:

Put

$$m = \lfloor \frac{y - E_n - \delta_y}{2} \rfloor. \tag{16}$$

We move

$$(x, y) \rightarrow (x - E_n + E_m, H_m + \delta_y) \in \mathcal{M}_4, \tag{17}$$

since $x - E_n = 0$ or 1 , and $\delta_y = 0$ or 1 . This move is legal:

(B-a) $k = E_n - E_m > 0$ and k is even. Firstly, we show that $0 \leq m < n$. Note that if y is even, we have $y \geq x \geq E_n = E_n + \delta_y$; if y is odd, we have $y \geq E_n + 1 = E_n + \delta_y$. Thus $y \geq E_n + \delta_y$, i.e., $m = \lfloor \frac{y - E_n - \delta_y}{2} \rfloor \geq 0$; On the other hand, the fact $y < H_n$ implies that $y - \delta_y < H_n$. Thus

$$y - E_n - \delta_y < H_n - E_n = 2n,$$

and

$$m = \lfloor \frac{y - E_n - \delta_y}{2} \rfloor \leq \frac{y - E_n - \delta_y}{2} < n.$$

By Fact B.1, $k = E_n - E_m > 0$ and k is even.

(B-b) $\ell = y - H_m - \delta_y > 0$ and ℓ is even. By the definition of m , we have $m \leq \frac{y - E_n - \delta_y}{2}$. It follows from Eq. (13) and $E_0 = H_0 = 4$ that

$$H_n = E_n + 2n \text{ for } n \geq 0. \quad (18)$$

Thus

$$\begin{aligned} y &\geq 2m + E_n + \delta_y \\ &= H_m + \delta_y + E_n - E_m \\ &> H_m + \delta_y. \end{aligned}$$

Hence $\ell = y - H_m - \delta_y > 0$ and ℓ is even.

(B-c) $0 \leq \ell - k < t$. By the definition of m , we have $m > \frac{y - E_n - \delta_y}{2} - 1$, i.e., $y < 2(m + 1) + E_n + \delta_y$. By Eq. (18), we have

$$\begin{aligned} \ell &= y - H_m - \delta_y \\ &< 2(m + 1) + E_n - E_m - 2m \\ &= E_n - E_m + 2. \end{aligned}$$

Note that $y - H_m - \delta_y$ and $E_n - E_m + 2$ are even, so $\ell = y - H_m - \delta_y \leq E_n - E_m < E_n - E_m + t$; On the other hand, $\ell = y - H_m - \delta_y \geq E_n - E_m = k$ by virtue of (B-b). Therefore, $0 \leq \ell - k < t$.

The proof is completed. \square

5 Odd-Even (s, t) -Wythoff's Game

In this section, we introduce a new *Odd-Even (s, t) -Wythoff's Game* (denoted by OEW). Let S_h , S_v , D_1 and D_2 be subsets of $Z^{\geq 0}$. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. One of the heaps is designated as “first heap” and the other as “second heap” throughout the game. By (x, y) we denote a position of present game, where x and y denote the numbers of tokens in the first and the second heaps, respectively. Two rules of moves are allowed:

(*Odd-Even Nim Rule*) A player chooses the first heap and takes *odd* $k > 0$ tokens, or chooses the second heap and takes *even* $\ell > 0$ tokens.

(*Odd-Even More General Wythoff's Rule*) A player takes tokens from both heaps, *odd* $k > 0$ tokens from the first heap, *even* $\ell > 0$ tokens from the second heap, and

$$0 \leq |\ell - k| < (s - 1)\lambda + t, \quad \lambda = \min\{k, \ell\} \in \mathbb{Z}^{\geq 1}. \quad (19)$$

Obviously, OEW is equivalent to $S_h = D_1 = Z^{\text{odd}}$ and $S_v = D_2 = Z^{\text{even}}$ in General Restriction of (s, t) -Wythoff's Game. Therefore OEW is a restricted version of (s, t) -Wythoff's game.

Remark 15. OEW has no “symmetry”, i.e., two positions (x, y) and (y, x) maybe not equivalent (see Remark 10). As an example, we consider OEW under normal play convention. Obviously, $(0, 0)$ is a P -position. The position $(0, 1)$ is a P -position, as the only possible move is by taking 1 token from the second heap. But this is not a legal move. The position $(1, 0)$ is an N -position as one can move $(1, 0)$ to $(0, 0)$ by taking 1 tokens from the first heap. Thus two positions $(0, 1)$ and $(1, 0)$ are not equivalent.

The position $(3, 8)$ is an option of position $(10, 8)$, as one can move $(10, 8)$ to $(3, 8)$ by taking 7 tokens from the first heap. But the position $(8, 3)$ is not an option of position $(8, 10)$, as the move of taking 7 tokens from the second heap is not a legal move.

Remark 16. In OEW, two parameters $s \geq 1$ and $t \geq 1$ are positive integers. If $s = t = 1$, then Eq. (19) can not hold:

$$|\ell - k| \geq 1 = (s - 1)\lambda + t, \lambda = \min\{k, \ell\} \in \mathbb{Z}^{\geq 1},$$

i.e., Odd-Even More General Wythoff's Rule is invalid. For $(s = 1 \text{ and } t > 1)$ or $(s > 1 \text{ and } t \geq 1)$, Odd-Even More General Wythoff's Rule is valid. Therefore, we will give the results on $s = t = 1$ and $s + t > 2$, respectively. All P -positions of OEW under normal or misère play convention are given, and the corresponding winning strategies are also presented (Theorems 17, 18, 19 and 20).

5.1 All P -positions of OEW: $s = t = 1$

Theorem 17. *Given two parameters $s = t = 1$. By \mathcal{P}_5 we denote the set of all P -positions of OEW under normal play convention. Then*

$$\mathcal{P}_5 = \bigcup_{n=0}^{\infty} \{(2n, 0), (2n, 1), (2n + 1, 2), (2n + 1, 3)\}.$$

Proof. Let

$$\mathcal{W} = \bigcup_{n=0}^{\infty} \{(2n, 0), (2n, 1), (2n + 1, 2), (2n + 1, 3)\}.$$

Proof of Fact 1. Let (a, b) and (a', b') are two distinct positions of \mathcal{W} . It is easy to see that there exists no legal move such that $(a, b) \rightarrow (a', b')$ or $(a', b') \rightarrow (a, b)$.

Proof of Fact 2. Let (a, b) be a position not in \mathcal{W} . It suffices to show that there exists a legal move such that $(a, b) \rightarrow (a', b') \in \mathcal{W}$.

(2.1) $a = 2n$ for some integer $n \in \mathbb{Z}^{\geq 0}$. In this case, the fact $(a, b) \notin \mathcal{W}$ implies that $b \geq 2$. We move

$$(a, b) \rightarrow (2n, \delta_b) \in \mathcal{W}.$$

(2.2) $a = 2n + 1$ for some integer $n \in \mathbb{Z}^{\geq 0}$. In this case, the fact $(a, b) \notin \mathcal{W}$ implies that $b = 0$, $b = 1$ or $b \geq 4$:

- $b \in \{0, 1\}$. We move $(a, b) = (2n + 1, b) \rightarrow (2n, b) \in \mathcal{W}$.
- $b \geq 4$. We move $(a, b) = (2n + 1, b) \rightarrow (2n + 1, 2 + \delta_b) \in \mathcal{W}$.

The proof is completed. \square

Theorem 18. *Given two parameters $s = t = 1$. By \mathcal{P}'_5 we denote the set of all P -positions of OEWS under misère play convention. Then*

$$\mathcal{P}'_5 = \bigcup_{n=0}^{\infty} \{(2n, 2), (2n, 3), (2n + 1, 0), (2n + 1, 1)\}.$$

Proof. Let

$$\mathcal{M} = \bigcup_{n=0}^{\infty} \{(2n, 2), (2n, 3), (2n + 1, 0), (2n + 1, 1)\}.$$

Proof of Fact 1. Let (a, b) and (a', b') are two distinct positions of \mathcal{M} . It is easy to see that there exists no legal move such that $(a, b) \rightarrow (a', b')$ or $(a', b') \rightarrow (a, b)$.

Proof of Fact 2. Let (a, b) be a position not in \mathcal{M} . It suffices to show that there exists a legal move such that $(a, b) \rightarrow (a', b') \in \mathcal{M}$.

(2.1) $a = 2n + 1$ for some integer $n \in \mathbb{Z}^{\geq 0}$. In this case, the fact $(a, b) \notin \mathcal{M}$ implies that $b \geq 2$. We move

$$(a, b) \rightarrow (2n + 1, \delta_b) \in \mathcal{M}.$$

(2.2) $a = 2n$ for some integer $n \in \mathbb{Z}^{\geq 0}$. We distinguish the following two subcases: $n = 0$ or $n \geq 1$.

(2.2.1) $n = 0$. In this subcase, the fact $(a, b) \notin \mathcal{M}$ implies that $b = 0$ or $b = 1$ or $b \geq 4$. It is obvious that $(0, 0)$ and $(0, 1)$ are N -positions. If $b \geq 4$, we move

$$(a, b) = (0, b) \rightarrow (0, 2 + \delta_b) \in \mathcal{M}.$$

(2.2.2) $n \geq 1$. In this subcase, the fact $(a, b) \notin \mathcal{M}$ implies that $b = 0$ or $b = 1$ or $b \geq 4$.

- $b \in \{0, 1\}$. We move $(a, b) = (2n, b) \rightarrow (2n - 1, b) \in \mathcal{M}$.
- $b \geq 4$. We move $(a, b) = (2n, b) \rightarrow (2n, 2 + \delta_b) \in \mathcal{M}$.

The proof is completed. \square

5.2 All P -positions of OEW: $s + t > 2$

Theorem 19. By \mathcal{P}_6 we denote the set of all P -positions of OEW under normal play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$ with $s + t > 2$,

$$\mathcal{P}_6 = \bigcup_{n=0}^{\infty} \{(A_n, B_n), (A_n, B'_n)\},$$

where for $n \geq 0$,

$$\begin{cases} A_n = n, \\ B_n = \delta_n(sA_n + t + \delta_{s+t}), \\ B'_n = B_n + 1. \end{cases} \quad (20)$$

Proof. Before we give the proof of Theorem 19, Tables 7 and 8 list the first few values of A_n and B_n for $s = t = 2$, $s = 2$ and $t = 3$, respectively, which show us how to determine \mathcal{P}_6 by using Theorem 19.

Table 7. The first few values of A_n and B_n for $s = t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
B_n	0	4	0	8	0	12	0	16	0	20	0	24	0	28	0
B'_n	1	5	1	9	1	13	1	17	1	21	1	25	1	29	1

For $s = t = 2$, it follows from Table 7 that

$$\mathcal{P}_6 = \left\{ \begin{array}{l} (0, 0), (0, 1), (1, 4), (1, 5), (2, 0), (2, 1), (3, 8), (3, 9), (4, 0), (4, 1), \\ (5, 12), (5, 13), (6, 0), (6, 1), (7, 16), (7, 17), (8, 0), (8, 1), \dots \end{array} \right\}$$

Table 8. The first few values of A_n and B_n for $s = 2$ and $t = 3$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
A_n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
B_n	0	6	0	10	0	14	0	18	0	22	0	26	0	30	0
B'_n	1	7	1	11	1	15	1	19	1	23	1	27	1	31	1

For $s = 2$ and $t = 3$, it follows from Table 8 that

$$\mathcal{P}_6 = \left\{ \begin{array}{l} (0, 0), (0, 1), (1, 6), (1, 7), (2, 0), (2, 1), (3, 10), (3, 11), (4, 0), (4, 1), \\ (5, 14), (5, 15), (6, 0), (6, 1), (7, 18), (7, 19), (8, 0), (8, 1), \dots \end{array} \right\}$$

We now give the proof of Theorem 19. Let $\mathcal{W} = \bigcup_{n=0}^{\infty} \{(A_n, B_n), (A_n, B'_n)\}$.

Proof of Fact 1. By the definition of function δ_n , we have

$$B_n = \delta_n(sA_n + t + \delta_{s+t}) = \delta_n(sn + t + \delta_{s+t})$$

is even and $B'_n = B_n + 1$ is odd.

Given $(A_n, B_n) \in \mathcal{W}$. Note that $(A_n, B_n) \rightarrow (A_m, B'_m) \in \mathcal{W}$ is not a legal move, since $\ell = B_n - B'_m$ is odd. Similarly, $(A_n, B'_n) \rightarrow (A_m, B_m) \in \mathcal{W}$ is also impossible.

Given $(A_n, B_n) \in \mathcal{W}$ or $(A_n, B'_n) \in \mathcal{W}$. Suppose that $(A_n, B_n) \rightarrow (A_m, B_m) \in \mathcal{W}$ or $(A_n, B'_n) \rightarrow (A_m, B'_m) \in \mathcal{W}$. In both cases, we have $m < n$, $k = A_n - A_m = n - m$ is odd, and $\ell = B_n - B_m$.

If n is even and m is odd then

$$\ell = B_n - B_m = 0 - (sm + t + \delta_{s+t}) < 0,$$

which is impossible; If n is odd and m is even then

$$\ell = B_n - B_m = (sA_n + t + \delta_{s+t}) - 0 \geq sn \geq s(n - m) \geq k > 0,$$

and

$$\begin{aligned} \ell &= B_n - B_m = sA_n + t + \delta_{s+t} \\ &\geq sn + t \geq s(n - m) + t = sk + t, \end{aligned}$$

which contradicts Eq. (19).

Proof of Fact 2. Let (x, y) be a position not in \mathcal{W} . We will show that there exists a legal move such that $(x, y) \rightarrow (A_n, B_n) \in \mathcal{W}$ or $(x, y) \rightarrow (A_n, B'_n) \in \mathcal{W}$.

Put $x = n = A_n$ for some integer $n \in \mathbb{Z}^{\geq 0}$. We distinguish two cases:

(2.1) $x = n = A_n$ is even. In this case, $B_n = 0$, $B'_n = 1$. The fact $(x, y) \notin \mathcal{W}$ implies that $y \geq 2$. We move $(x, y) \rightarrow (A_n, B_n + \delta_y) \in \mathcal{W}$ by virtue of $\delta_y = 0$ or 1 .

(2.2) $x = n = A_n$ is odd. In this case,

$$B_n = sx + t + \delta_{s+t} \text{ is even,} \tag{21}$$

and

$$B'_n = B_n + 1 \text{ is odd.} \tag{22}$$

The fact $(x, y) \notin \mathcal{W}$ implies that $y \geq B_n + 2$ or $0 \leq y \leq B_n - 1$.

(2.2.1) $y \geq B_n + 2$. Now $y \geq B_n + 2 > B_n + \delta_y$ and $y - B_n - \delta_y$ is even. We move

$$(x, y) \rightarrow (A_n, B_n + \delta_y) \in \mathcal{W},$$

by taking $y - B_n - \delta_y$ tokens from the second heap.

(2.2.2) $0 \leq y \leq B_n - 1$. We distinguish the following three subcases: $y \in \{0, 1\}$ or $2 \leq y \leq x$ or $x + 1 \leq y \leq B_n - 1$.

• $y \in \{0, 1\}$. We move

$$(x, y) \rightarrow (x - 1, \delta_y) = (A_{n-1}, B_{n-1} + \delta_y) \in \mathcal{W},$$

since $n - 1$ is odd, $B_{n-1} = 0$, and $y = B_{n-1} + \delta_y$.

• $2 \leq y \leq x$. We move

$$(x, y) \rightarrow (x - y + 1 + \delta_y, \delta_y),$$

by taking $y - 1 - \delta_y$ tokens from the first heap and $y - \delta_y$ tokens from the second heap. Let $x - y + 1 + \delta_y = m$. We note that $m = A_m$ is even and $B_m = 0$. Note that $\delta_y = B_m$ if y is even, $\delta_y = B_m + 1 = B'_m$ if y is odd. Thus

$$(x - y + 1 + \delta_y, \delta_y) = (A_m, B_m + \delta_y) \in \mathcal{W}.$$

This move is legal. Indeed,

- 1) $k = y - 1 - \delta_y > 0$ is odd;
 - 2) $\ell = y - \delta_y > 0$ is even;
 - 3) $0 \leq |\ell - k| = 1 < (s - 1)k + t$.
- $x + 1 \leq y \leq B_n - 1$. We move $(x, y) \rightarrow (0, \delta_y) \in \mathcal{W}$, by taking x tokens from the first heap and $y - \delta_y$ tokens from the second heap. This move is legal. Indeed,
- 1) $k = x$ is odd;
 - 2) $\ell = y - \delta_y$ is even;
 - 3) We note that $y \geq x + 1$ implies that $\ell = y - \delta_y \geq x = k$. By Eq. (21), $B_n - 1$ is odd. The fact $y \leq B_n - 1$ implies that $y \leq B_n - 2 + \delta_y$. Hence,

$$\begin{aligned} 0 \leq |\ell - k| &= y - \delta_y - x \\ &\leq sx + t + \delta_{s+t} - 2 - x \\ &= (s - 1)k + t + \delta_{s+t} - 2 \\ &< (s - 1)k + t, \end{aligned}$$

by virtue of $k \geq 1$.

The proof is completed. \square

Theorem 20. By \mathcal{P}'_6 we denote the set of all P -positions of OEWS under misère play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$ with $s + t > 2$,

$$\mathcal{P}'_6 = \bigcup_{n=0}^{\infty} \{(E_n, H_n), (E_n, H'_n)\},$$

where $E_0 = 0$, $H_0 = 2$, $H'_0 = 3$ and for $n \geq 1$,

$$\begin{cases} E_n = n, \\ H_n = (1 - \delta_n)(sE_n - s + t + \delta_{s+t}), \\ H'_n = H_n + 1. \end{cases} \quad (23)$$

Proof. Before we give the proof of Theorem 20, Tables 9 and 10 list the first few values of E_n , H_n and H'_n for $s = t = 2$, $s = 1$ and $t = 2$, respectively, which show us how to determine \mathcal{P}'_6 by using Theorem 20.

Table 9. The first few values of E_n , H_n and H'_n for $s = t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
E_n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
H_n	2	0	4	0	8	0	12	0	16	0	20	0	24	0	28
H'_n	3	1	5	1	9	1	13	1	17	1	21	1	25	1	29

Table 10. The first few values of E_n , H_n and H'_n for $s = 1, t = 2$.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
E_n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
H_n	2	0	5	0	9	0	13	0	17	0	21	0	25	0	29
H'_n	3	1	6	1	10	1	14	1	18	1	22	1	26	1	30

We now give the proof of Theorem 20. Let

$$\mathcal{G} = \{(0, 2), (0, 3)\} \cup \bigcup_{n=1}^{\infty} \{(E_n, H_n), (E_n, H'_n)\}.$$

Proof of Fact 1. By the definition of function δ_n , if n is odd then $H_n = 0$, if $n \geq 1$ is even then $H_n = s(n-1) + t + \delta_{s+t}$ is even. Thus for $n \in \mathbb{Z}^{\geq 1}$, H_n is even and $H'_n = H_n + 1$ is odd.

Note that $(0, 2)$ or $(0, 3)$ can not move to (E_n, H_n) or (E_n, H'_n) as $E_n > 0$. If n is odd, $H_n = 0$ implies that (E_n, H_n) or (E_n, H'_n) can not move to $(0, 2)$ or $(0, 3)$; If n is even, $E_n = n$ is even implies that (E_n, H_n) or (E_n, H'_n) can not move to $(0, 2)$ or $(0, 3)$.

Given $(E_n, H_n) \in \mathcal{G}$. Note that $(E_n, H_n) \rightarrow (E_m, H'_m) \in \mathcal{G}$ is not a legal move, since $\ell = H_n - H'_m$ is odd. Similarly, $(E_n, H'_n) \rightarrow (E_m, H_m) \in \mathcal{G}$ is also impossible.

Given $(E_n, H_n) \in \mathcal{G}$ or $(E_n, H'_n) \in \mathcal{G}$. Suppose that $(E_n, H_n) \rightarrow (E_m, H_m) \in \mathcal{G}$ or $(E_n, H'_n) \rightarrow (E_m, H'_m) \in \mathcal{G}$. In both cases, we have $1 \leq m < n$, $k = E_n - E_m = n - m$ is odd, and $\ell = H_n - H_m$.

If n is odd and m is even then

$$\ell = H_n - H_m = 0 - (s(m-1) + t + \delta_{s+t}) < 0,$$

which is impossible; If n is even and m is odd, then

$$\ell = H_n - H_m = s(n-1) + t + \delta_{s+t} - 0 \geq s(n-m) + t > sk \geq k > 0,$$

and

$$\begin{aligned} \ell &= H_n - H_m \\ &= sE_n - s + t + \delta_{s+t} \\ &\geq s(n-1) + t \\ &= s(n-m) + s(m-1) + t \\ &\geq s(n-m) + t = sk + t, \end{aligned}$$

by virtue of $m \geq 1$, which contradicts Eq. (19).

Proof of Fact 2. Let (x, y) be a position not in \mathcal{G} . It suffices to show that there exists a legal move such that $(x, y) \rightarrow (E_n, H_n) \in \mathcal{G}$ or $(x, y) \rightarrow (E_n, H'_n) \in \mathcal{G}$.

Put $x = n = E_n$ for some integer $n \in \mathbb{Z}^{\geq 0}$.

(2.1) $x = n = 0$. In this case, we have $y = 0$ or $y = 1$ or $y \geq 4$. It is obvious that $(0, 0)$ and $(0, 1)$ are N -positions. If $y \geq 4$, then we move $(0, y) \rightarrow (0, 2 + \delta_y) \in \mathcal{G}$ by virtue of $\delta_y = 0$ or 1 .

(2.2) $x = n > 0$ is odd. In this case, $H_n = 0$, $H'_n = 1$. The fact $(x, y) \notin \mathcal{G}$ implies that $y \geq 2$. We move $(x, y) \rightarrow (E_n, H_n + \delta_y) \in \mathcal{G}$.

(2.3) $x = n > 0$ is even. In this case,

$$H_n = sx - s + t + \delta_{s+t} \text{ is even,} \quad (24)$$

and

$$H'_n = H_n + 1 \text{ is odd.} \quad (25)$$

The fact $(x, y) \notin \mathcal{G}$ implies that $y \geq H_n + 2$ or $0 \leq y \leq H_n - 1$.

(2.3.1) $y \geq H_n + 2$, $n \in Z^{\geq 1}$. We move

$$(x, y) \rightarrow (E_n, H_n + \delta_y) \in \mathcal{G},$$

which is legal, since $y - H_n - \delta_y > 0$ is even.

(2.3.2) $0 \leq y \leq H_n - 1$, $n \in Z^{\geq 1}$. We distinguish the following three subcases: $y \in \{0, 1\}$ or $2 \leq y \leq x - 1$ or $x \leq y \leq H_n - 1$.

• $y \in \{0, 1\}$. We move

$$(x, y) \rightarrow (x - 1, \delta_y) = (E_{n-1}, H_{n-1} + \delta_y) \in \mathcal{G},$$

since $n - 1$ is odd, $H_{n-1} = 0$, and $\delta_y = 0$ or 1 .

• $2 \leq y \leq x - 1$. We move

$$(x, y) \rightarrow (x - y - 1 + \delta_y, \delta_y),$$

by taking $y + 1 - \delta_y$ tokens from the first heap and $y - \delta_y$ tokens from the second heap. Let $x - y - 1 + \delta_y = m$. Note that m is odd, $H_m = 0$, and $\delta_y = H_m + \delta_y$. Obviously, $\delta_y = H_m$ if y is even, $\delta_y = H_m + 1 = H'_m$ if y is odd. Thus

$$(x - y - 1 + \delta_y, \delta_y) = (E_m, H_m + \delta_y) \in \mathcal{G}.$$

This move is legal. Indeed,

- 1) $k = y - \delta_y + 1 > 0$ is odd;
 - 2) $\ell = y - \delta_y > 0$ is even and $\ell \geq k$;
 - 3) $0 < |\ell - k| = 1 < (s - 1)k + t$.
- $x \leq y \leq H_n - 1$, $n \in Z^{\geq 1}$. We move

$$(x, y) \rightarrow (1, \delta_y) \in \mathcal{G},$$

by taking $x - 1$ tokens from the first heap and $y - \delta_y$ tokens from the second heap. If y is even, $\delta_y = 0 = H_1$; If y is odd, $\delta_y = 1 = H_1 + 1 = H'_1$. Thus $(1, \delta_y) = (E_1, H_1 + \delta_y) \in \mathcal{G}$.

This move is legal. Indeed,

- 1) $k = x - 1$ is odd;
- 2) $\ell = y - \delta_y$ is even;
- 3) Note that $y \geq x$ implies that

$$\ell = y - \delta_y \geq x > x - 1 = k.$$

By Eq. (24), $H_n - 1$ is odd, so the fact $y \leq H_n - 1$ implies that $y \leq H_n - 2 + \delta_y$. Hence,

$$\begin{aligned}
0 \leq |\ell - k| &= y - \delta_y - x + 1 \\
&\leq sx - s + t + \delta_{s+t} - 2 - x + 1 \\
&= (s-1)(k+1) - s + t + \delta_{s+t} - 1 \\
&= (s-1)k + t + \delta_{s+t} - 2 \\
&< (s-1)k + t,
\end{aligned}$$

by virtue of $k \geq 1$.

The proof is completed. \square

6 Conclusion

Three new models, Odd-Odd (s, t) -Wythoff's Game, Even-Even (s, t) -Wythoff's Game, Odd-Even (s, t) -Wythoff's Game, are investigated. Under normal or misère play convention, all P -positions of these three models are given for all integers $s, t \geq 1$.

Similar to Odd-Even (s, t) -Wythoff's Game, we can define the fourth model, *Even-Odd (s, t) -Wythoff's Game* (Denoted by EOW): Let S_h, S_v, D_1 and D_2 be subsets of $Z^{\geq 0}$. Given two parameters $s, t \in Z^{\geq 1}$ and two heaps of finitely many tokens. One of the heaps is designated as “first heap” and the other as “second heap” throughout the game. By (x, y) we denote a position of present game, where x and y denote the numbers of tokens in the first and the second heaps, respectively. Two rules of moves are allowed:

(*Even-Odd Nim Rule*) A player chooses the first heap and takes *even* $k > 0$ tokens, or chooses the second heap and takes *odd* $\ell > 0$ tokens.

(*Even-Odd More General Wythoff's Rule*) A player takes tokens from both heaps, *even* $k > 0$ tokens from the first heap, *odd* $\ell > 0$ tokens from the second heap, and

$$0 \leq |\ell - k| < (s-1)\lambda + t, \quad \lambda = \min\{k, \ell\} \in Z^{\geq 1}. \quad (26)$$

Obviously, EOW is equivalent to $S_h = D_1 = Z^{\text{even}}$ and $S_v = D_2 = Z^{\text{odd}}$ in General Restriction of (s, t) -Wythoff's Game.

If (x, y) is a P -position of OEOW, then (y, x) is a P -position of EOW. Thus we have

Corollary 21. *Given two parameters $s = t = 1$. By \mathcal{P}_7 we denote the set of all P -positions of EOW under normal play convention. Then*

$$\mathcal{P}_7 = \bigcup_{n=0}^{\infty} \{(0, 2n), (1, 2n), (2, 2n+1), (3, 2n+1)\}.$$

Corollary 22. *Given two parameters $s = t = 1$. By \mathcal{P}'_7 we denote the set of all P -positions of EOW under misère play convention. Then*

$$\mathcal{P}'_7 = \bigcup_{n=0}^{\infty} \{(2, 2n), (3, 2n), (0, 2n+1), (1, 2n+1)\}.$$

Corollary 23. By \mathcal{P}_8 we denote the set of all P -positions of EOW under normal play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$ with $s + t > 2$,

$$\mathcal{P}_8 = \bigcup_{n=0}^{\infty} \{(B_n, A_n), (B'_n, A_n)\},$$

where for $n \geq 0$,

$$\begin{cases} A_n = n, \\ B_n = \delta_n(sA_n + t + \delta_{s+t}), \\ B'_n = B_n + 1. \end{cases} \quad (27)$$

Corollary 24. By \mathcal{P}'_8 we denote the set of all P -positions of EOW under misère play convention. Then for all $s, t \in \mathbb{Z}^{\geq 1}$ with $s + t > 2$,

$$\mathcal{P}'_8 = \{(2, 0), (3, 0)\} \cup \bigcup_{n=1}^{\infty} \{(H_n, E_n), (H'_n, E_n)\},$$

where for $n \geq 1$,

$$\begin{cases} E_n = n, \\ H_n = (1 - \delta_n)(sE_n - s + t + \delta_{s+t}), \\ H'_n = H_n + 1. \end{cases} \quad (28)$$

Recall that Wythoff's game is a special case $s = t = 1$ in (s, t) -Wythoff's game, and a -Wythoff's game is a special case $s = 1$ and $t = a$ in (s, t) -Wythoff's game. Thus (s, t) -Wythoff's game is a generalization of both Wythoff's game and a -Wythoff's game. Under normal play convention, the set of all P -positions of a -Wythoff's game and the set of all P -positions of Wythoff's game can be obtained by letting $(s = 1 \text{ and } t = a)$ and $s = t = 1$ in Eq. (2), respectively (see [19],[9]).

Our results on OOW, EEW, OEW and EOW are given for all integers $s \geq 1$ and $t \geq 1$. Thus the corresponding results on $(s = 1 \text{ and } t = a)$ or $s = t = 1$ have been obtained.

Given two integer $K \geq 1$ and $r \in \{0, 1, 2, \dots, K - 1\}$. We use notation $Z_K^{(r)} = \{Kn + r | n \in \mathbb{Z}^{\geq 0}\}$.

Open Problem 25. Let $S_h = S_v = D_1 = D_2 = Z_K^{(0)}$ in General Restriction of (s, t) -Wythoff's Game. How to determine all P -positions under normal or misère play convention?

Note that $Z^{\text{even}} = Z_2^{(0)}$, thus Theorems 13 and 14 have settled the special case $K = 2$ of this problem. Can we generalize Theorems 13 and 14 from $K = 2$ to an arbitrary integer $K \geq 3$?

Open Problem 26. Let $S_h = S_v = D_1 = D_2 = Z_K^{(r)}$ in General Restriction of (s, t) -Wythoff's Game, where $r \in \{0, 1, 2, \dots, K - 1\}$ be a fixed integer. How to determine all P -positions under normal or misère play convention?

Note that $Z^{\text{odd}} = Z_2^{(1)}$, thus Theorems 7 and 9 have settled the special case $K = 2$ and $r = 1$ of this problem. Can we generalize Theorems 7 and 9 from $K = 2$ and $r = 1$ to arbitrary integers $K \geq 3$ and $r \in \{0, 1, 2, \dots, K - 1\}$?

Open Problem 27. Let $S_h = D_1 = Z_K^{(r_1)}$ and $S_v = D_2 = Z_K^{(r_2)}$ in General Restriction of (s, t) -Wythoff's Game, where $r_1, r_2 \in \{0, 1, 2, \dots, K-1\}$ and $r_1 \neq r_2$ be fixed integers. How to determine all P -positions under normal or misère play convention?

Note that $Z^{even} = Z_2^{(0)}$ and $Z^{odd} = Z_2^{(1)}$, thus Theorems 17, 18, 19 and 20 have settled the special case $K = 2$ of this problem. Can we generalize these results from $(K = 2, r_1 = 1 \text{ and } r_2 = 0)$ to arbitrary integers $K \geq 3$ and $r_1 \neq r_2 \in \{0, 1, 2, \dots, K-1\}$?

Open Problem 28. One can investigate the case of (s, t) -Wythoff's game which is only restricted on Extended Diagonal Moves. As an example, let $S_h = S_v = Z^{\geq 0}$ and $D_1 = D_2 = Z_K^{(r)}$ (K is a fixed positive integer, $r \in \{0, 1, 2, \dots, K-1\}$) in General Restriction of (s, t) -Wythoff's Game. Can we obtain the corresponding results?

Acknowledgements The authors are grateful to the responsible editor and the anonymous referees for their valuable comments and suggestions, which have greatly improved the earlier version of this paper. The research is supported by the National Natural Science Foundation of China under Grants 11171368 and 11171094.

References

- [1] E.R. Berlekamp, J.H. Conway and R.K. Guy. Winning Ways for Your Mathematical Plays. Volume 1-4, A K Peters, Wellesley, MA, 2001-2004. 2nd edition: vol. 1 (2001), vols. 2, 3 (2003), vol. 4 (2004).
- [2] C.L. Bouton. Nim, a game with a complete mathematical theory. *Annals of Mathematics*, 3:35-39, 1905.
- [3] I.G. Connell. A generalization of Wythoff's game. *Canad. Math. Bull.*, 2:181-190, 1959.
- [4] J.H. Conway, *On Numbers and Games*, Academic Press, London, 1976.
- [5] E. Duchêne and S. Gravier. Geometrical extensions of Wythoff's game. *Discrete Math.*, 309(11):3595-3608, 2009.
- [6] E. Duchêne, A.S. Fraenkel, S. Gravier and R.J. Nowakowski. Another bridge between Nim and Wythoff. *Australasian J. of Combinatorics*, 44:43-56, 2009.
- [7] E. Duchêne, A.S. Fraenkel, R.J. Nowakowski and M. Rigo. Extensions and restrictions of Wythoff's game preserving its P-positions. *J. Combinatorial Theory Ser. A*, 117(5):545-567, 2010.
- [8] A.S. Fraenkel. New games related to old and new sequences. *Integers: Electr. J of Combinat. Number Theory*, 4:#G6, 18 pp., 2004.
- [9] A.S. Fraenkel. How to beat your Wythoff games' opponent on three fronts. *Amer. Math. Monthly*, 89:353-361, 1982.
- [10] A.S. Fraenkel and M. Lorberbom. Nimhoff games. *J. Combinatorial Theory Ser. A*, 58(1):1-25, 1991.
- [11] A.S. Fraenkel. Heap games, Numeration systems and Sequences. *Annals of Combinatorics*, 2(3):197-210, 1998.

- [12] A.S. Fraenkel and M. Ozery. Adjoining to Wythoff's game its P-positions as moves. *Theoretical Computer Science*, 205:283-296, 1998.
- [13] A.S. Fraenkel. Wythoff games, continued fractions, cedar trees and fibonacci searches. *Theoretical Computer Science*, 29(1-2):49C73, 1984.
- [14] A.S. Fraenkel. Euclid and Wythoff games. *Discrete Math.*, 304:65-68, 2005.
- [15] A.S. Fraenkel and E. Reisner. The game of End-Wythoff. *Games of No Chance III*, Proc. BIRS Workshop on Combinatorial Games, July 2005, Banff, Alberta, Canada, MSRI Publ., M.H. Albert and R.J. Nowakowski, Eds., Vol. 56, Cambridge University Press, Cambridge, (2009) 329-347.
- [16] W.A. Liu, H.F. Li and B. Li. A restricted version of Wythoff's game. *Electronic Journal of Combinatorics*, 18:#P207, 17pp, 2011.
- [17] W.A. Liu, Y. Wang and N. Li. The (s, t) -Wythoff's game under misère play convention. *Fifth International Joint Conference on Computational Sciences and Optimization*, Harbin, China, 419-422, 2012.
- [18] X. Sun and D. Zeilberger. On Fraenkel's N-heap Wythoff's conjectures. *Annals of Combinatorics*, 8:225-238, 2004.
- [19] W. Wythoff. A modification of the game of Nim. *Nieuw Arch. Wisk.*, 7:199-202, 1907.