

# Closing the gap on path-kipas Ramsey numbers

*We dedicate this paper to the memory of Ralph Faudree,  
one of the exponents of Ramsey theory who died on January 13, 2015*

Binlong Li\*

Department of Mathematics  
University of West Bohemia  
European Centre of Excellence NTIS -  
New Technologies for the Inf. Society  
306 14 Pilsen, Czech Rep.

Department of Applied Mathematics  
Northwestern Polytechnical University  
Xi'an, Shaanxi 710072, P.R. China

libinlong@mail.nwpu.edu.cn

Yanbo Zhang

Faculty of Electrical Engineering  
University of Twente  
7500 AE Enschede, The Netherlands

Department of Mathematics  
Nanjing University  
Nanjing 210093, P.R. China

y.zhang@utwente.nl

Halina Bielak

Institute of Mathematics  
Maria Curie-Skłodowska University  
20-031 Lublin, Poland  
hbiel@hektor.umcs.lublin.pl

Hajo Broersma

Faculty of Electrical Engineering,  
Mathematics and Computer Science  
University of Twente  
P.O. Box 217, 7500 AE Enschede, The Netherlands  
h.j.broersma@utwente.nl

Přemysl Holub<sup>†</sup>

Department of Mathematics  
University of West Bohemia  
P.O. Box 314, 306 14 Pilsen, Czech Republic  
holubpre@kma.zcu.cz

Submitted: Jan 30, 2015; Accepted: Jul 26, 2015; Published: Aug 14, 2015

Mathematics Subject Classifications: 05C55, 05D10

---

## Abstract

\*The first author is partly supported by the Doctorate Foundation of Northwestern Polytechnical University (No. cx201202) and by the project NEXLIZ-CZ.1.07/2.3.00/30.0038, which is co-financed by the European Social Fund and the state budget of the Czech Republic.

<sup>†</sup>Research partly supported by project P202/12/G061 of the Czech Science Foundation.

Given two graphs  $G_1$  and  $G_2$ , the Ramsey number  $R(G_1, G_2)$  is the smallest integer  $N$  such that, for any graph  $G$  of order  $N$ , either  $G_1$  is a subgraph of  $G$ , or  $G_2$  is a subgraph of the complement of  $G$ . Let  $P_n$  denote a path of order  $n$  and  $\widehat{K}_m$  a kipas of order  $m + 1$ , i.e., the graph obtained from a  $P_m$  by adding one new vertex  $v$  and edges from  $v$  to all vertices of the  $P_m$ . We close the gap in existing knowledge on exact values of the Ramsey numbers  $R(P_n, \widehat{K}_m)$  by determining the exact values for the remaining open cases.

**Keywords:** Ramsey number; path; kipas

## 1 Introduction

We only consider finite simple graphs. A cycle, a path and a complete graph of order  $n$  are denoted by  $C_n$ ,  $P_n$  and  $K_n$ , respectively. A complete  $k$ -partite graph with classes of cardinalities  $n_1, n_2, \dots, n_k$  is denoted by  $K_{n_1, n_2, \dots, n_k}$ . For a nonempty proper subset  $S \subseteq V(G)$ , let  $G[S]$  and  $G - S$  denote the subgraph induced by  $S$  and  $V(G) - S$ , respectively. For a vertex  $v \in V(G)$ , we let  $N_S(v)$  denote the set of neighbors of  $v$  that are contained in  $S$ . For two vertex-disjoint graphs  $H_1, H_2$ , we define  $H_1 + H_2$  to be the graph with vertex set  $V(H_1) \cup V(H_2)$  and edge set  $E(H_1) \cup E(H_2) \cup \{uv \mid u \in V(H_1) \text{ and } v \in V(H_2)\}$ . For two disjoint vertex sets  $X, Y$ ,  $e(X, Y)$  denotes the number of edges with one end in  $X$  and one end in  $Y$ . We use  $mG$  to denote  $m$  vertex-disjoint copies of  $G$ . A star  $K_{1, n} = K_1 + nK_1$ , a kipas  $\widehat{K}_n = K_1 + P_n$  and a wheel  $W_n = K_1 + C_n$ . The term kipas and its notation were adopted from [8]. Kipas is the Malay word for fan; the motivation for the term kipas is that the graph  $K_1 + P_n$  looks like a hand fan (especially if the path  $P_n$  is drawn as part of a circle) but the term fan was already in use for the graphs  $K_1 + nK_2$ .

We use  $\delta(G)$  and  $\Delta(G)$  to denote the minimum and maximum degree of  $G$ , respectively.

Given two graphs  $G_1$  and  $G_2$ , the Ramsey number  $R(G_1, G_2)$  is the smallest integer  $N$  such that, for any graph  $G$  of order  $N$ , either  $G$  contains  $G_1$  or  $\overline{G}$  contains  $G_2$ , where  $\overline{G}$  is the complement of  $G$ . It is easy to check that  $R(G_1, G_2) = R(G_2, G_1)$ , and, if  $G_1$  is a subgraph of  $G_3$ , then  $R(G_1, G_2) \leq R(G_3, G_2)$ . Thus,  $R(P_n, K_{1, m}) \leq R(P_n, \widehat{K}_m) \leq R(P_n, W_m)$ . In [7], an explicit formula for  $R(P_n, K_{1, m})$  is given, while in [5], the Ramsey numbers  $R(P_n, W_m)$  for all  $m, n$  have been obtained. It follows from these results that  $R(P_n, K_{1, m}) = R(P_n, W_m)$  for  $m \geq 2n$ . Therefore,  $R(P_n, \widehat{K}_m) = R(P_n, K_{1, m}) = R(P_n, W_m)$  for  $m \geq 2n$ , and the exact values of these Ramsey numbers can be found in both [5] and [7].

It is trivial that  $R(P_1, \widehat{K}_m) = 1$  and  $R(P_n, \widehat{K}_1) = n$ . Many nontrivial exact values for  $R(P_n, \widehat{K}_m)$  have been obtained by Salman and Broersma in [8]. Here we completely solve the case by determining all the remaining path-kipas Ramsey numbers.  $R(P_n, \widehat{K}_m)$  can easily be determined for  $m \geq 2n$  (and follows directly from earlier results, as indicated above). In this note we close the gap by proving the following theorem.

**Theorem 1.**  $R(P_n, \widehat{K}_m) = \max\{2n - 1, \lceil 3m/2 \rceil - 1, 2\lfloor m/2 \rfloor + n - 2\}$  for  $m \leq 2n - 1$  and  $m, n \geq 2$ .

## 2 Proof of Theorem 1

We first list the following eight useful results that we will use in our proof of Theorem 1, as separate lemmas.

**Lemma 2.** (*Gerencsér and Gyárfás [4]*). For  $m \geq n \geq 2$ ,  $R(P_m, P_n) = m + \lfloor n/2 \rfloor - 1$ .

**Lemma 3.** (*Faudree et al. [3]*). For  $n \geq 2$  and even  $m \geq 4$ ,  $R(C_m, P_n) = \max\{m + \lfloor n/2 \rfloor - 1, n + m/2 - 1\}$ .

**Lemma 4.** (*Parsons [6]*). For  $n \geq m \geq 2$ ,  $R(K_{1,m}, P_n) = \max\{2m - 1, n\}$ .

**Lemma 5.** (*Salman and Broersma [8]*).  $R(P_4, \widehat{K}_6) = 8$ .

**Lemma 6.** (*Dirac [2]*). If  $G$  is a connected graph, then  $G$  contains a path of order at least  $\min\{2\delta(G) + 1, |V(G)|\}$ .

**Lemma 7.** (*Bondy [1]*). If  $\delta(G) \geq |V(G)|/2$ , then  $G$  contains cycles of every length between 3 and  $|V(G)|$ , or  $r = |V(G)|/2$  and  $G = K_{r,r}$ .

**Lemma 8.** (*Zhang et al. [9]*). Let  $C$  be a longest cycle of a graph  $G$  and  $v_1, v_2 \in V(G) - V(C)$ . Then  $|N_{V(C)}(v_1) \cup N_{V(C)}(v_2)| \leq \lfloor |V(C)|/2 \rfloor + 1$ .

**Lemma 9.** Let  $G$  be a graph with  $|V(G)| \geq 6$  and  $\delta(G) \geq 2$ . Then  $G$  contains two vertex-disjoint paths, one with order three and one with order two.

*Proof.* If  $G$  is connected, by Lemma 6,  $G$  contains a path of order at least 5. Let  $x_1x_2x_3x_4x_5$  be a path in  $G$ . Then  $G$  contains two vertex-disjoint paths  $x_1x_2x_3$  and  $x_4x_5$ . If  $G$  is disconnected, then each component of  $G$  contains a path of order three. This completes the proof of Lemma 9.  $\square$

We proceed to prove Theorem 1. Let  $N = \max\{2n - 1, \lceil 3m/2 \rceil - 1, 2\lfloor m/2 \rfloor + n - 2\}$ , and let  $m \leq 2n - 1$  and  $m, n \geq 2$ . It suffices to show that  $R(P_n, \widehat{K}_m) = N$ .

If  $n = 2$ , then  $m \leq 2n - 1$  and  $m, n \geq 2$  imply  $m = 2$  or  $m = 3$ . It is obvious that  $R(P_2, \widehat{K}_m) = m + 1$ , and one easily checks that  $m + 1 = N$  for these values of  $m$  and  $n$ . Next we assume that  $n \geq 3$ . We first show that  $R(P_n, \widehat{K}_m) \geq N$ . For this purpose, we note that it is straightforward to check that any of the graphs  $G \in \{K_{n-1, n-1}, K_{\lfloor m/2 \rfloor, \lfloor m/2 \rfloor - 1, \lfloor m/2 \rfloor - 1}, K_{n-1, \lfloor m/2 \rfloor - 1, \lfloor m/2 \rfloor - 1}\}$  contains no  $\widehat{K}_m$ , whereas  $\overline{G}$  contains no  $P_n$ . Thus,  $R(P_n, \widehat{K}_m) \geq \max\{2n - 1, \lceil 3m/2 \rceil - 1, 2\lfloor m/2 \rfloor + n - 2\} = N$ .

It remains to prove  $R(P_n, \widehat{K}_m) \leq N$ . To the contrary, we assume there exists a graph  $G$  of order  $N$  such that neither  $G$  contains a  $\widehat{K}_m$ , nor  $\overline{G}$  contains a  $P_n$ .

We first claim that  $\Delta(G) \geq N - \lfloor n/2 \rfloor$ . To prove this claim, assume to the contrary that  $\Delta(G) \leq N - \lfloor n/2 \rfloor - 1$ . Then  $\delta(\overline{G}) \geq \lfloor n/2 \rfloor$ . Let  $H$  be a largest component of  $\overline{G}$ . If  $|V(H)| \geq n$ , then, since  $\delta(H) \geq \lfloor n/2 \rfloor$ ,  $H$  contains a  $P_n$  by Lemma 6, a contradiction. Thus,  $|V(H)| \leq n - 1$  and  $|V(G)| - |V(H)| \geq N - n + 1$ . Since  $m \leq 2n - 1$ , we have  $n \geq \lfloor m/2 \rfloor$ . From the definition of  $N$  we get that  $N - n + 1 \geq n$  and  $N - n + 1 \geq 2\lfloor m/2 \rfloor - 1$ ,

so  $N - n + 1 \geq \max\{2\lfloor m/2 \rfloor - 1, n\}$ . Since  $\overline{G} - V(H)$  contains no  $P_n$ , by Lemma 4,  $G - V(H)$  contains a  $K_{1, \lfloor m/2 \rfloor}$ . If  $|V(H)| \geq \lfloor m/2 \rfloor$ , since every vertex of  $V(H)$  is adjacent to every vertex of  $V(G) - V(H)$  in  $G$ , then  $G$  contains a  $\widehat{K}_m$ , a contradiction. This implies that  $|V(H)| \leq \lfloor m/2 \rfloor - 1$ . Recall that  $H$  is a largest component of  $\overline{G}$ . Thus  $\overline{G}$  contains at least four components; otherwise  $|V(\overline{G})| \leq 3(\lfloor m/2 \rfloor - 1) < \lfloor 3m/2 \rfloor - 1 \leq N$ , a contradiction. Let  $H'$  be a smallest component of  $\overline{G}$ . Then  $|V(H')| \leq N/4$  and  $|V(G)| - |V(H')| \geq 3N/4 \geq 3/4(\lfloor 3m/2 \rfloor - 1) \geq 9m/8 - 3/4 > m - 3/4$ . That is,  $|V(G)| - |V(H')| \geq m$ . Since every component in  $\overline{G} - V(H')$  is of order at most  $\lfloor m/2 \rfloor - 1$ , then every vertex in  $\overline{G} - V(H')$  is of degree at most  $\lfloor m/2 \rfloor - 2$ . Thus, we have  $\delta(G - V(H')) > (|V(G)| - |V(H')|)/2$ . By Lemma 7,  $G - V(H')$  contains a  $P_m$ , which together with any vertex of  $V(H')$  forms a  $\widehat{K}_m$  in  $G$ , a contradiction. This proves our claim that  $\Delta(G) \geq N - \lfloor n/2 \rfloor$ .

Let  $u$  be a vertex of  $G$  with  $d(u) = d = \Delta(G)$ , let  $F = G[N(u)]$  and  $Z = V(G) - V(F) - \{u\}$ . Then  $|V(F)| = d \geq N - \lfloor n/2 \rfloor = \max\{n + \lfloor n/2 \rfloor - 1, \lfloor 3m/2 \rfloor - \lfloor n/2 \rfloor - 1, 2\lfloor m/2 \rfloor + \lfloor n/2 \rfloor - 2\}$ . We claim that  $R(P_m, P_n) > d$ ; otherwise  $R(P_m, P_n) \leq d$ , and either  $F$  contains a  $P_m$ , which together with  $u$  forms a  $\widehat{K}_m$ , a contradiction; or  $\overline{F}$  contains a  $P_n$ , also a contradiction. If  $m \leq n$ , or if  $m = n + 1$  and  $m$  is even, then by Lemma 2,  $R(P_m, P_n) = \max\{n + \lfloor m/2 \rfloor - 1, m + \lfloor n/2 \rfloor - 1\} \leq n + \lfloor n/2 \rfloor - 1 \leq d$ , a contradiction. Therefore, it remains to deal with the cases that  $m \geq n + 2$ , and that  $m = n + 1$  and  $m$  is odd. We first deal with the latter case.

Let  $m = n + 1$  and  $m$  is odd. Then  $n$  is even, hence  $n \geq 4$ . We claim that  $|Z| \geq 1$ ; otherwise  $d = N - 1 = 2n - 2$ , and then  $R(P_m, P_n) = m + n/2 - 1 \leq 2n - 2 = d$  by Lemma 2, a contradiction. By Lemma 3,  $R(C_{m-1}, P_n) = m - 1 + n/2 - 1 = n + n/2 - 1 \leq d$ . Since  $\overline{F}$  contains no  $P_n$ , then  $F$  contains a  $C_{m-1}$ . Let  $C_{m-1} = x_1x_2 \dots x_{m-1}x_1$ ,  $Y = V(F) - V(C_{m-1}) = \{y_1, y_2, \dots, y_k\}$ . Then  $k \geq n/2 - 1$ . If  $e(V(C_{m-1}), Y) \geq 1$ , say  $x_1y_1 \in E(G)$ , then  $y_1x_1x_2 \dots x_{m-1}$  is a path in  $G$ , which together with  $u$  forms a  $\widehat{K}_m$ , a contradiction. Thus,  $e(V(C_{m-1}), Y) = 0$ . If there is an edge in  $\overline{G}[V(C_{m-1})]$ , say  $x_ix_j \in E(\overline{G})$  ( $1 \leq i < j \leq m - 1$ ), then  $x_ix_jy_1x'_1y_2x'_2 \dots y_{n/2-1}x'_{n/2-1}$  with  $\{x'_k : 1 \leq k \leq n/2 - 1\} \subseteq V(C_{m-1}) - \{x_i, x_j\}$  is a path of order  $n$  in  $\overline{G}$ , a contradiction. Thus,  $G[V(C_{m-1})]$  is a complete graph. Set  $z \in Z$ . If  $e(\{z\}, V(C_{m-1})) \geq 1$  in  $\overline{G}$ , say  $zx_1 \in E(\overline{G})$ , then  $uzx_1y_1 \dots x_{n/2-1}y_{n/2-1}$  is a path of order  $n$  in  $\overline{G}$ , a contradiction. Thus,  $e(\{z\}, V(C_{m-1})) = 0$  in  $\overline{G}$ , and  $G$  contains a path  $ux_1zx_2x_3 \dots x_{m-2}$ , which together with  $x_{m-1}$  forms a  $\widehat{K}_m$ , another contradiction. This completes the case that  $m = n + 1$  and  $m$  is odd. We proceed with the case that  $n + 2 \leq m \leq 2n - 1$ , and first consider the small values of  $n$ .

For  $n = 3$  and  $m = 5$ , or  $n = 4$  and  $m = 7$ , or  $n = 5$  and  $7 \leq m \leq 9$ , we get that  $R(P_m, P_n) = m + \lfloor n/2 \rfloor - 1 \leq \lfloor 3m/2 \rfloor - \lfloor n/2 \rfloor - 1 \leq d$ , a contradiction. By Lemma 5,  $R(P_4, \widehat{K}_6) = 8 = N$ . Hence it remains to consider the case that  $m \geq n + 2 \geq 8$ .

We first claim that  $|Z| \geq 2$ . If not,  $|Z| \leq 1$  and  $d = N - 1 - |Z| \geq N - 2$ . By Lemma 2,  $R(P_m, P_n) = m + \lfloor n/2 \rfloor - 1$ . If  $m \geq n + 3$ , then  $m + \lfloor n/2 \rfloor - 1 \leq \lfloor 3m/2 \rfloor - 3 \leq N - 2 \leq d$ , a contradiction; if  $n \geq 7$  or  $(n, m) = (6, 8)$ , then  $m + \lfloor n/2 \rfloor - 1 \leq 2\lfloor m/2 \rfloor + n - 4 \leq N - 2 \leq d$ , also a contradiction. Thus, for  $m \geq n + 2 \geq 8$ , we have  $|Z| \geq 2$ .

Since  $m \geq n + 2 \geq 8$ , by Lemma 3,  $R(C_{2\lfloor m/2 \rfloor - 2}, P_n) = \max\{2\lfloor m/2 \rfloor + \lfloor n/2 \rfloor -$

$3, n + \lfloor m/2 \rfloor - 2\} < 2\lfloor m/2 \rfloor + \lfloor n/2 \rfloor - 2 \leq d$ . Since  $\overline{F}$  contains no  $P_n$ ,  $F$  contains a  $C_{2\lfloor m/2 \rfloor - 2}$ . Let  $C$  be a longest cycle in  $F$ . Then  $|V(C)| \geq m - 3$ . If  $|V(C)| \geq m$ , then  $F$  contains a  $P_m$ , which together with  $u$  forms a  $\widehat{K}_m$  in  $G$ , a contradiction. Thus,  $m - 3 \leq |V(C)| \leq m - 1$ . We complete the proof by distinguishing the three cases that  $|V(C)| = m - 1$ ,  $|V(C)| = m - 2$  or  $|V(C)| = m - 3$ . In each case, let  $C = x_1x_2 \dots x_{|V(C)|}x_1$  and  $Y = V(F) - V(C) = \{y_1, y_2, \dots, y_k\}$ .

**Case 1:**  $|V(C)| = m - 1$ .

We have  $k = d - (m - 1) \geq \lfloor n/2 \rfloor - 2$ . If  $e(V(C), Y) \geq 1$ , say  $x_1y_1 \in E(G)$ , then  $y_1x_1x_2 \dots x_{m-1}$  is a path in  $G$ , which together with  $u$  forms a  $\widehat{K}_m$ , a contradiction. Thus,  $e(V(C), Y) = 0$ . Let  $z_1, z_2 \in Z$ . If  $e(\{z_1\}, V(C)) \geq 1$  in  $\overline{G}$ , say  $z_1x_1 \in E(\overline{G})$ , then  $z_2uz_1x_1y_1 \dots x_{\lfloor n/2 \rfloor - 2}y_{\lfloor n/2 \rfloor - 2}x_{\lfloor n/2 \rfloor - 1}$  is a path of order at least  $n$  in  $\overline{G}$ , a contradiction. This implies that  $e(\{z_1\}, V(C)) = 0$  in  $\overline{G}$ . For the same reason,  $e(\{z_2\}, V(C)) = 0$  in  $\overline{G}$ .

We claim that  $\delta(\overline{G}[V(C)]) \leq 1$ . If not,  $\delta(\overline{G}[V(C)]) \geq 2$ . Since  $m \geq 8$ , by Lemma 9, there are two vertex-disjoint paths in  $\overline{G}[V(C)]$ , one with order three and one with order two. Without loss of generality, let  $x'_1x'_2x'_3$  and  $x'_4x'_5$  be the two paths in  $\overline{G}[V(C)]$ . Because  $m - 1 \geq \lfloor n/2 \rfloor + 2$ , we may assume that  $x'_6, \dots, x'_{\lfloor n/2 \rfloor + 2} \in V(C) - \{x'_1, \dots, x'_5\}$ . Then  $x'_1x'_2x'_3y_1x'_4x'_5y_2x'_6y_3 \dots x'_{\lfloor n/2 \rfloor + 1}y_{\lfloor n/2 \rfloor - 2}x'_{\lfloor n/2 \rfloor + 2}$  is a path of order at least  $n$  in  $\overline{G}$ , a contradiction. This proves our claim that  $\delta(\overline{G}[V(C)]) \leq 1$ . That is, there exists a vertex of  $V(C)$  which is adjacent to at least  $|V(C)| - 2$  vertices of  $V(C)$ . Without loss of generality, let  $x_1$  be a vertex with maximum degree in  $G[V(C)]$ , and let  $x_3$  be the possible vertex that is nonadjacent to  $x_1$ . Then  $ux_2z_1x_4z_2x_5x_6 \dots x_{m-1}$  is a path of order  $m$ , which together with  $x_1$  forms a  $\widehat{K}_m$  in  $G$ , our final contradiction in Case 1.

**Case 2:**  $|V(C)| = m - 2$ .

We have  $k = d - (m - 2)$ . Note that  $k \geq \lfloor n/2 \rfloor - 1$  for odd  $m$ , and  $k \geq \lfloor n/2 \rfloor$  for even  $m$ . Let  $X$  be the set of all vertices of  $V(C)$  that are nonadjacent to  $Y$  in  $G$ . For  $1 \leq i \leq m - 2$ , either  $x_i \in X$ , or  $x_{i+1} \in X$ . Here,  $x_{m-1} = x_1$ . This is because, if  $x_i$  and  $x_{i+1}$  have a common neighbor in  $Y$ , say  $y_1$ , then by replacing  $x_ix_{i+1}$  by  $x_iy_1x_{i+1}$  in  $C$ , we obtain a cycle longer than  $C$ , a contradiction; if  $x_i$  and  $x_{i+1}$  are adjacent to different vertices of  $Y$ , say  $x_iy_1, x_{i+1}y_2 \in E(G)$ , then  $y_2x_{i+1}x_{i+2} \dots x_{m-2}x_1 \dots x_iy_1$  is a path of length  $m$ , which together with  $u$  forms a  $\widehat{K}_m$  in  $G$ , also a contradiction. Thus, at least one end of each edge of  $C$  is nonadjacent to  $Y$  in  $G$ . Note that  $|X| \geq \lfloor n/2 \rfloor$  and  $|Y| \geq \lfloor n/2 \rfloor - 1$  for odd  $m$  and  $|Y| \geq \lfloor n/2 \rfloor$  for even  $m$ . If  $m$  is even or  $n$  is odd, then we get a path  $P_n$  in  $\overline{G}[X \cup Y]$ . This implies it remains to consider the case that  $n$  is even and  $m$  is odd, with  $m \geq n + 3$ .

If  $|V(C) - X| \geq 2$ , say  $x_i, x_j \notin X$ , then  $x_{i+1}, x_{j+1} \in X$ . Moreover,  $x_{i+1}x_{j+1} \notin E(G)$ ; otherwise we may obtain either a cycle longer than  $C$  in  $F$ , or a path of length  $m$  in  $F$ , which together with  $u$  forms a  $\widehat{K}_m$  in  $G$ , both of which are contradictions. Now let  $x'_1, x'_2, \dots, x'_{|X|-2} \in X - \{x_{i+1}, x_{j+1}\}$ . Since  $|X| - 2 \geq \lfloor |V(C)|/2 \rfloor - 2 \geq n/2 - 1$ , let  $P = x_{i+1}x_{j+1}y_1x'_1y_2x'_2 \dots y_{n/2-1}x'_{n/2-1}$ . Note that  $P$  is a path of order  $n$  in  $\overline{G}$ , a contradiction. Thus,  $m - 3 \leq |X| \leq m - 2$  and there exists a vertex in  $V(C)$ , say  $x_1$ , such that  $e(V(C) - \{x_1\}, Y) = 0$ .

Since  $m \geq n+3 \geq 9$ , we have  $m-3 \geq \lceil n/2 \rceil + 2$ . If there is an edge in  $\overline{G}[V(C) - \{x_1\}]$ , say  $x_i x_j \in E(\overline{G})$ , then  $\overline{G}[X \cup Y]$  contains a path  $P_n$ , a contradiction. Thus,  $G[V(C) - \{x_1\}]$  is a complete graph of order  $m-3$ .

Let  $z_1, z_2 \in Z$ . We claim that  $e(\{z_1\}, V(C) - \{x_1\}) = 0$  in  $\overline{G}$ ; otherwise, say for  $z_1 x_2 \in E(\overline{G})$ ,  $z_2 u z_1 x_2 y_2 x_3 y_3 \dots x_{n/2-1} y_{n/2-1} x_{n/2}$  is a path of order  $n$  in  $\overline{G}$ , a contradiction. For the same reason,  $e(\{z_2\}, V(C) - \{x_1\}) = 0$  in  $\overline{G}$ .

It is easy to check that  $x_1 u x_3 z_1 x_4 z_2 x_5 \dots x_{m-2}$  is a path of order  $m$ , which together with  $x_2$  forms a  $\widehat{K}_m$  in  $G$ , our final contradiction in Case 2.

**Case 3:**  $|V(C)| = m-3$ .

If  $m = n+2 \geq 8$ , then  $m$  and  $n$  have the same parity. In that case,  $R(C_{2\lfloor (m-1)/2 \rfloor}, P_n) = 2\lfloor (m-1)/2 \rfloor + \lfloor n/2 \rfloor - 1 \leq 2\lfloor m/2 \rfloor + \lfloor n/2 \rfloor - 2 \leq d$ . Since  $\overline{F}$  contains no  $P_n$ ,  $F$  contains a  $C_{2\lfloor (m-1)/2 \rfloor}$ . This contradicts the fact that  $C$  with  $|V(C)| = m-3$  is a longest cycle in  $F$ . It remains to consider the case that  $m \geq n+3 \geq 9$ .

We have  $k = d - (m-3) \geq \lceil n/2 \rceil$ . By Lemma 8, any two vertices of  $Y$  have at least  $\lceil (m-3)/2 \rceil - 1 \geq \lceil n/2 \rceil - 1$  common nonadjacent vertices of  $V(C)$  in  $G$ . Since  $C$  is a longest cycle in  $G$ , any vertex of  $Y$  has at least  $\lceil (m-3)/2 \rceil \geq \lceil n/2 \rceil$  nonadjacent vertices of  $V(C)$  in  $G$ . By these observations,  $y_1$  and  $y_2$  have a common nonadjacent vertex in  $V(C)$ , say  $x_1$ ; for  $2 \leq i \leq \lceil n/2 \rceil - 1$ ,  $y_i$  and  $y_{i+1}$  have a common nonadjacent vertex in  $V(C) - \{x_1, x_2, \dots, x_{i-1}\}$ , say  $x_i$ ;  $y_{\lceil n/2 \rceil}$  have a nonadjacent vertex in  $X - \{x_1, x_2, \dots, x_{\lceil n/2 \rceil - 1}\}$ , say  $x_{\lceil n/2 \rceil}$ . Then  $y_1 x_1 y_2 x_2 \dots y_{\lceil n/2 \rceil} x_{\lceil n/2 \rceil}$  is a path of order at least  $n$  in  $\overline{G}$ . This final contradiction completes the proof of Case 3 and of Theorem 1.  $\square$

## References

- [1] J.A. Bondy, Pancyclic graphs I, *Journal of Combinatorial Theory, Series B* 11 (1971), 80–84.
- [2] G.A. Dirac, Some theorems on abstract graphs, *Proceedings of the London Mathematical Society* 2 (1952), 69–81.
- [3] R.J. Faudree, S.L. Lawrence, T.D. Parsons, and R.H. Schelp, Path-cycle Ramsey numbers, *Discrete Mathematics* 10 (1974), 269–277.
- [4] L. Gerencsér and A. Gyárfás, On Ramsey-type problems, *Annales Universitatis Scientiarum Budapestinensis, Eötvös Sect. Math.* 10 (1967), 167–170.
- [5] B. Li and B. Ning, The Ramsey numbers of paths versus wheels: a complete solution, *The Electronic Journal of Combinatorics* 21 (2014), #P4.41.
- [6] T.D. Parsons, Path-star Ramsey numbers, *Journal of Combinatorial Theory, Series B* 17 (1974), 51–58.
- [7] C.C. Rousseau and J. Sheehan, A class of Ramsey problems involving trees, *Journal of the London Mathematical Society* 2 (1978), 392–396.
- [8] A.N.M. Salman and H.J. Broersma, Path-kipas Ramsey numbers, *Discrete Applied Mathematics* 155 (2007), 1878–1884.

- [9] Y.B. Zhang, Y.Q. Zhang, and Y.J. Chen, The Ramsey numbers of wheels versus odd cycles, *Discrete Mathematics* 323 (2014), 76–80.