

Extremal graph for intersecting odd cycles

Xinmin Hou^{a*} Yu Qiu^b Boyuan Liu^b

Key Laboratory of Wu Wen-Tsun Mathematics
Chinese Academy of Sciences
School of Mathematical Sciences
University of Science and Technology of China
Hefei, Anhui 230026, China.

^axmhou@ustc.edu.cn, ^byuqiu@mail.ustc.edu.cn

^clby1055@mail.ustc.edu.cn

Submitted: Jan 8, 2016; Accepted: Apr 24, 2016; Published: May 13, 2016

Mathematics Subject Classifications: 05C35, 05C38

Abstract

An extremal graph for a graph H on n vertices is a graph on n vertices with maximum number of edges that does not contain H as a subgraph. Let $T_{n,r}$ be the Turán graph, which is the complete r -partite graph on n vertices with part sizes that differ by at most one. The well-known Turán Theorem states that $T_{n,r}$ is the only extremal graph for complete graph K_{r+1} . Erdős et al. (1995) determined the extremal graphs for intersecting triangles and Chen et al. (2003) determined the maximum number of edges of the extremal graphs for intersecting cliques. In this paper, we determine the extremal graphs for intersecting odd cycles.

Keywords: extremal graph; Turán graph; intersecting odd cycle

1 Introduction

In this paper, all graphs considered are simple and finite. For a graph G and a vertex $x \in V(G)$, the neighborhood of x in G is denoted by $N_G(x)$. Let $N_G[x] = \{x\} \cup N_G(x)$. The *degree* of x , denoted by $\deg_G(x)$, is $|N_G(x)|$. Let $\delta(G)$ and $\Delta(G)$ denote the minimum and maximum degrees of G , respectively. A *matching* M in G is a subgraph of G with $\delta(M) = \Delta(M) = 1$. The *matching number* of G , denoted by $\nu(G)$, is the maximum number of edges in a matching in G . Let $e(G)$ be the number of edges of G . For a graph G and $S, T \subset V(G)$, let $e_G(S, T)$ be the number of edges $e = xy \in E(G)$ such that $x \in S$

*Supported by NNSF of China (No. 11271348) and the Fundamental Research Funds for the Central Universities.

and $y \in T$, if $S = T$, we use $e_G(S)$ instead of $e_G(S, S)$; and we use $e_G(u, T)$ instead of $e_G(\{u\}, T)$ for convenience, the index G will be omitted if no confusion from the context. For a subset $X \subseteq V(G)$ or $X \subseteq E(G)$, let $G[S]$ be the subgraph of G induced by X , that is $G[X] = (X, E(X))$ if $X \subseteq V(G)$, or $G[X] = (V(X), X)$ if $X \subseteq E(G)$. A cycle of length q is called a q -cycle.

Given two graphs G and H , we say that G is H -free if G does not contain an H as a subgraph. The Turán function, denoted by $ex(n, H)$, is the largest possible number of edges of an H -free graph on n vertices. That is,

$$ex(n, H) = \max\{e(G) : |V(G)| = n, G \text{ is } H\text{-free}\}.$$

And for positive integers $n \geq r$, the Turán graph, denoted by $T_{n,r}$, is the complete r -partite graph on n vertices with part sizes that differ by at most one (also called the complete balanced r -partite graph). The well-known Turán Theorem states that $ex(n, K_{r+1}) = e(T_{n,r})$ and $T_{n,r}$ is the only extremal graph for complete graph K_{r+1} .

A k -fan, denoted by F_k , is a graph on $2k + 1$ vertices consisting of k triangles which intersect in exactly one common vertex. In 1995, Erdős et al. [4] gave the value of $ex(n, F_k)$ and determined the extremal graphs for F_k as follows.

Theorem 1 ([4]). *For $k \geq 1$ and $n \geq 50k^2$,*

$$ex(n, F_k) = e(T_{n,2}) + g(k),$$

where

$$g(k) = \begin{cases} k^2 - k & \text{if } k \text{ is odd,} \\ k^2 - \frac{3}{2}k & \text{if } k \text{ is even.} \end{cases}$$

Moreover, when k is odd, the extremal graph must be a $T_{n,2}$ with two vertex disjoint copies of K_k embedding in one partite set. When k is even, the extremal graph must be a $T_{n,2}$ with a graph having $2k - 1$ vertices, $k^2 - \frac{3}{2}k$ edges with maximum degree $k - 1$ embedded in one partite set.

In 2003, Chen et al. [2] proved that $ex(n, F_{k,r}) = e(T_{n,r-1}) + g(k)$, where $F_{k,r}$ is a graph consisting of k complete graphs of order r which intersect in exactly one common vertex and $g(k)$ is the same as in Theorem 1. Recently, Glebov [5] and Liu [6] gave the extremal graphs for blow-ups of paths [5], cycles and a large class of trees [6] (A blow-up of a graph G is a graph obtained from G by replacing each edge of G by a clique, two cliques share a common vertex if and only if their corresponding edges have a common end).

In this paper, motivated by the results in [2, 4, 5, 6], we generalize Theorem 1 in another way. For a positive integer k and an odd integer q with $q \geq 5$, let $C_{k,q}$ be the graph consisting of k q -cycles which intersect exactly in one common vertex, called the center of it. For $n \geq 4(k - 1)^2$, let $\mathcal{F}_{n,k}$ be the family of graphs where each member is a Turán graph $T_{n,2}$ with a complete bipartite graph $K_{k-1,k-1}$ embedded into one class. Our main result is as follows.

Theorem 2. For an integer $k \geq 2$ and an odd integer $q \geq 5$, there exists $n_1(k, q) \in \mathbb{N}$ such that for all $n \geq n_1(k, q)$, we have

$$\text{ex}(n, C_{k,q}) = e(T_{n,2}) + (k - 1)^2,$$

and the only extremal graphs for $C_{k,q}$ are members of $\mathcal{F}_{n,k}$.

The remainder of the paper is arranged as follows. Section 2 gives some lemmas. Section 3 gives the proof of Theorem 2.

2 Lemmas

The following two lemmas are useful to estimate the number of edges of a graph with restricted degrees and matching number.

Lemma 3 (Chavátal [1]). For any graph G with maximum degree $\Delta \geq 1$ and matching number $\nu \geq 1$, then $e(G) \leq f(\nu, \Delta) = \nu\Delta + \lfloor \frac{\Delta}{2} \rfloor \lfloor \frac{\nu}{\lceil \Delta/2 \rceil} \rfloor \leq \nu(\Delta + 1)$.

The following stability result due to Erdős [3] and Simonovits [7] gives the rough structure of the extremal graphs for a graph H with $\chi(H) = r \geq 3$ and $H \neq K_r$.

Lemma 4 ([3, 7]). Let H be a graph with $\chi(H) = r \geq 3$ and $H \neq K_r$. Then, for every $\gamma > 0$, there exists $\delta > 0$ and $n_0 = n_0(H, \gamma) \in \mathbb{N}$ such that the following holds. If G is an H -free graph on $n \geq n_0$ vertices with $e(G) \geq \text{ex}(n, H) - \delta n^2$, then there exists a partition of $V(G) = V_1 \dot{\cup} \dots \dot{\cup} V_{r-1}$ such that $\sum_{i=1}^{r-1} e(V_i) < \gamma n^2$.

The following is a simple observation.

Observation 5. Let G be a graph with no isolated vertex. If $\Delta(G) \leq 2$, then

$$\nu(G) \geq \frac{|V(G)| - \omega(G)}{2},$$

where $\omega(G)$ is the number of components of G .

Proof. Since $\Delta(G) \leq 2$, each component of G is a path or a cycle. Hence each component C of G has matching number at least $\frac{|V(C)|-1}{2}$. This implies the desired result. \square

Lemma 6. Let G be a graph with no isolated vertex. If for all $x \in V(G)$, $\deg(x) + \nu(G - N[x]) \leq r$, then $e(G) \leq r^2$. Moreover, equality holds if and only if $G = K_{r,r}$.

Proof. Clearly, $\Delta(G) \leq r$. We claim that $\nu(G) \leq r$. Let $u_1 v_1, \dots, u_\ell v_\ell$ be a matching in G . Without loss of generality, assume that

$$\begin{aligned} \{u_1, v_1, \dots, u_s, v_s, u_{s+1}, \dots, u_{s+t}\} &\subseteq N[u_1], \\ \{v_{s+1}, \dots, v_{s+t}, u_{s+t+1}, v_{s+t+1}, u_\ell, v_\ell\} &\subseteq V(G) \setminus N[u_1]. \end{aligned}$$

Then $s \geq 1$, $2s + t - 1 \leq \deg(u_1)$ and $\ell - (s + t) \leq \nu(G - N[u_1])$. Thus,

$$\ell \leq \nu(G - N[u_1]) + s + t \leq \deg(u_1) + \nu(G - N[u_1]) \leq r,$$

the claim is true. Now we prove the result according to the following two cases.

Case 1. Suppose $\Delta(G) < r$. Then, by Lemma 3, we have $e(G) \leq f(\nu, \Delta) \leq r(r - 1 + 1) = r^2$, and the equality holds only if $\nu = r$ and $\Delta = r - 1$. We claim that the equality does not hold in this case. For $r \geq 4$, $e(G) \leq f(r, r - 1) = r(r - 1) + \lfloor \frac{r-1}{2} \rfloor \lfloor \frac{r}{\lceil (r-1)/2 \rceil} \rfloor < r^2$. For $r = 3$, $e(G) \leq f(3, 2) = 3^2 = 9$. Suppose to the contrary that $e(G) = 9$. Since $\Delta(G) \leq 2$ and G has no isolated vertex, $|V(G)| \geq e(G) = 9$ (equality holds if and only if G is 2-regular) and $\omega(G) \leq \nu(G) = 3$. By Observation 5,

$$3 = \nu(G) \geq \frac{|V(G)| - \omega(G)}{2} \geq \frac{|V(G)| - 3}{2}.$$

Hence $|V(G)| \geq 9$. Thus $|V(G)| = 9$ (and so G is 2-regular) and $\omega(G) = 3$. Therefore, G consists of three vertex-disjoint triangles. But this contradicts the assumption that $\deg(x) + \nu(G - N[x]) \leq r$ for all $x \in V(G)$.

Case 2. Suppose $\Delta(G) = r$. Choose $x \in V(G)$ such that $\deg(x) = r$, then $\nu(G - N[x]) = 0$. Hence $e(G - N[x]) = 0$ and so each vertex in $G - N[x]$ must be adjacent to vertices in $N(x)$. Let $N(x) = \{x_1, \dots, x_r\}$. For each $i \in [1, r]$, let $d_i = \deg(x_i)$ and $\tilde{d}_i = \deg_{G[N(x)]}(x_i)$. Then

$$\begin{aligned} e(G) &= e(G[N(x)]) + e(N(x), V(G) \setminus N(x)) = \frac{1}{2} \sum_{i=1}^r \tilde{d}_i + \sum_{i=1}^r (d_i - \tilde{d}_i) \\ &= \sum_{i=1}^r d_i - \frac{1}{2} \sum_{i=1}^r \tilde{d}_i \leq r^2 - \frac{1}{2} \sum_{i=1}^r \tilde{d}_i \leq r^2. \end{aligned}$$

Moreover, equality holds if and only if $d_i = r$ and $\tilde{d}_i = 0$ for each $i \in [1, r]$, that is, G is a bipartite graph with partite sets $N(x) = \{x_1, \dots, x_r\}$ and $V(G) \setminus N(x)$. To show that $G = K_{r,r}$, it suffices to prove that $|V(G) \setminus N(x)| = r$. If not, then $|V(G) \setminus N(x)| > r$. Since $\deg(x_1) = d_1 = r$, there must exist a vertex $y \in (V(G) \setminus N(x)) \setminus N(x_1)$. Since G has no isolated vertex, y must be adjacent to some vertex x_j with $j \neq 1$. This implies that $\nu(G - N[x_1]) \geq 1$, a contradiction with $\deg(x_1) + \nu(G - N[x_1]) \leq r$. \square

The following lemma states that the members $\mathcal{F}_{n,k}$ are $C_{k,q}$ -free.

Lemma 7. *Each member of $\mathcal{F}_{n,k}$ is $C_{k,q}$ -free for all $k \geq 2$, $n \geq 4(k - 1)^2$, and odd integer $q \geq 5$.*

Proof. Suppose to the contrary that there is a graph $G \in \mathcal{F}_{n,k}$ containing a copy of $C_{k,q}$. Let K be the copy of $K_{k-1,k-1}$ in G . Then each odd cycle of $C_{k,q}$ must contain odd number of the edges of K . Let $A = E(C_{k,q}) \cap E(K)$. Then $|A| \geq k$. We claim that the center of $C_{k,q}$ must lie in K . If not, then $G[A]$ contains a matching of order at least k by the

structure of $C_{k,q}$, a contradiction with $\nu(K) = k - 1$. Let $x \in V(K)$ be the center of $C_{k,q}$. Assume that $\deg_{G[A]}(x) = s$ and let $E_A(x)$ be the set of edges incident with x in $G[A]$. Then at most s cycles of $C_{k,q}$ intersect $E_A(x)$, that is $A - E_A(x)$ contains a matching of K of order at least $k - s$. This is impossible since $\nu(K - N_{G[A]}(x)) \leq k - s - 1$. \square

Lemma 8. *Let n_0 be an integer and let G be a graph on $n \geq n_0 + \binom{n_0}{2}$ vertices with $e(G) = e(T_{n,2}) + j$ for some integer $j > 0$. Then G contains a subgraph G' on $n' > n_0$ vertices such that $\delta(G') \geq \delta(T_{n',2})$ and $e(G') \geq e(T_{n',2}) + j + n - n'$.*

Proof. If $\delta(G) \geq \lfloor \frac{n}{2} \rfloor$, then G is the desired graph and we have nothing to do. So assume that $\delta(G) < \lfloor \frac{n}{2} \rfloor$. Choose $v \in V(G)$ with $\deg_G(v) < \lfloor \frac{n}{2} \rfloor$. Let $G_1 = G - v$. Then $e(G_1) \geq e(G) - \deg_G(v) \geq e(T_{n,2}) + j - \lfloor \frac{n}{2} \rfloor + 1 = e(T_{n-1,2}) + j + 1$, since $e(T_{n,2}) - e(T_{n-1,2}) = \lfloor \frac{n}{2} \rfloor$. We may continue this procedure until we get G' on $n - i$ vertices with $\delta(G') \geq \delta(T_{n-i,2})$ for some $i < n - n_0$, or until $i = n - n_0$. For the latter case, G' has n_0 vertices but $e(G') \geq e(T_{n_0,2}) + j + i > n - n_0 \geq \binom{n_0}{2}$, which is impossible. \square

3 Proof of Theorem 2

Let G be an extremal graph for $C_{k,q}$ on $n \geq n_1(k, q)$ ($n_1(k, q)$ is given below) vertices. By Lemma 7, $e(G) \geq e(T_{n,2}) + (k - 1)^2$. We will show that $e(G) = e(T_{n,2}) + (k - 1)^2$ and G is a member of $\mathcal{F}_{n,k}$. Let

$$\begin{aligned} \gamma &= \frac{1}{1600}, \\ n_0 &= n_0(C_{k,q}, \gamma) \text{ (which is determined by } C_{k,q} \text{ and } \gamma \text{ by applying Lemma 4),} \\ n_1 &= n_1(k, q) = n_0 + 20k^2q + \binom{n_0 + 20k^2q}{2}. \end{aligned}$$

By the choice of n_1 and Lemma 8, we may assume $\delta(G) \geq \delta(T_{n,2}) = \lfloor \frac{n}{2} \rfloor$, otherwise, we consider a subgraph G' with the desired minimum degree instead of G . Let $V_0 \dot{\cup} V_1$ be a partition of $V(G)$ such that $e(V_0, V_1)$ is maximized. Lemma 4 implies that $m = e(V_0) + e(V_1) < \gamma n^2$. The following claim asserts that the partition is close to being balanced. The following inequality holds:

Claim 9.

$$\frac{n}{2} - \sqrt{\gamma}n < |V_i| < \frac{n}{2} + \sqrt{\gamma}n \text{ for } i = 0, 1.$$

Furthermore, $m = e(V_0) + e(V_1) \geq (k - 1)^2$ and if the equality holds then G contains a complete balanced bipartite graph with classes V_0 and V_1 .

Proof. Let $|V_0| = \frac{n}{2} + a$. Then $|V_1| = \frac{n}{2} - a$. Since

$$\left\lfloor \frac{n^2}{4} \right\rfloor + (k - 1)^2 = e(T_{n,2}) + (k - 1)^2 \leq e(G) \leq |V_0||V_1| + m = \frac{n^2}{4} - a^2 + m,$$

we have $m \geq (k - 1)^2$ and $m \geq a^2$. Since $m < \gamma n^2$, $a^2 < \gamma n^2$. Hence $|a| < \sqrt{\gamma}n$.

If $m = (k - 1)^2$, then

$$e(T_{n,2}) + (k - 1)^2 \leq e(G) = e(V_0, V_1) + (k - 1)^2.$$

Hence $e(V_0, V_1) = e(T_{n,2})$, that is, V_0, V_1 are balanced and so G contains a complete balanced bipartite graph with classes V_0 and V_1 . \square

In the following, let $G_i = G[V_i]$, $\Delta_i = \Delta(G_i)$ and $\nu_i = \nu(G_i)$, $i = 0, 1$. For a vertex $x \in V_i$, let $E_{1-i}(x) = \{e \in E(G_{1-i}) \mid V(e) \cap N_G(x) \neq \emptyset\}$.

Claim 10. For any vertex $x \in V_i$,

$$\deg_{G_i}(x) + \nu(G_i - N_{G_i}[x]) + \nu(G[E_{1-i}(x)]) \leq k - 1.$$

Proof. We prove this claim by contradiction. Without loss of generality, assume that there is an $x \in V_0$ such that $\deg_{G_0}(x) + \nu(G_0 - N_0[x]) + \nu(G[E_1(x)]) \geq k$. Let $xx_1, xx_2, \dots, xx_s \in E(G_0)$, and let $M_0 = \{u_{s+1}v_{s+1}, \dots, u_tv_t\}$ be a matching of $G[V_0 \setminus \{x, x_1, \dots, x_s\}]$ and $M_1 = \{u_{t+1}v_{t+1}, \dots, u_kv_k\}$ a matching of $G[E_1(x)]$ such that $xu_{t+1}, \dots, xu_k \in E_G(x, V_1)$, where $s, t \in [0, k]$. Define

$$\xi(j) = \begin{cases} 1 & \text{if } j \text{ is odd,} \\ 0 & \text{if } j \text{ is even.} \end{cases}$$

We say that v is *bad* if $\deg_{G[V_i]}(v) > t_1$, otherwise v is said to be *good*, where $t_1 = 6\sqrt{\gamma}n$.

Then the number of bad vertices in G is at most $\frac{2m}{t_1} < \frac{2\gamma n^2}{t_1} = \frac{1}{3}\sqrt{\gamma}n$.

For any vertex $u \in V_i$ ($i = 0, 1$), by the maximality of $e(V_0, V_1)$, we have

$$e_G(u, V_{1-i}) \geq \max\{e_G(u, V_i), \lfloor \frac{n}{2} \rfloor - e_G(u, V_i)\} \geq \frac{1}{2} \lfloor \frac{n}{2} \rfloor.$$

Particularly, if $u \in V_i$ is good, then we have

$$e_G(u, V_{1-i}) \geq \lfloor \frac{n}{2} \rfloor - e_G(u, V_i) \geq \lfloor \frac{n}{2} \rfloor - t_1 \left(\geq \frac{1}{2} \lfloor \frac{n}{2} \rfloor \right).$$

Let $A = V(\{xx_1, xx_2, \dots, xx_s\} \cup M_i \cup M_{1-i})$. We find a copy of $C_{k,q}$ passing through all the vertices of A to get a contradiction.

For each $\ell \in [1, s]$, we find a sequence of vertices $w_{1\ell}^1, w_{0\ell}^2, \dots, w_{0\ell}^{q-3}, w_{1\ell}^{q-2}$ with $w_{\xi(j)\ell}^j \in V_{\xi(j)} \setminus A$ for $1 \leq j \leq q - 2$, such that $w_{0\ell}^{q-3}$ is good and $xw_{1\ell}^1 \dots w_{0\ell}^{q-3} w_{1\ell}^{q-2} x_\ell x$ is a q -cycle. Furthermore, we require that $w_{\xi(j)\ell}^j$ ($\ell \in [1, s]$, $j \in [1, q - 2]$) are pairwise different. This is possible since together with all vertices in A , the total number of good vertices which we have found is at most $|V(C_{k,q})| = k(q - 1) + 1$ and each vertex $u \in V_i$ has at least

$$e_G(u, V_{1-i}) - \frac{1}{3}\sqrt{\gamma}n \geq \frac{1}{2} \lfloor \frac{n}{2} \rfloor - \frac{1}{3}\sqrt{\gamma}n \geq |V(C_{k,q})| \text{ (since } n \geq 20k^2q)$$

good neighbors in V_{1-i} and the number of common neighbors of $w_{0\ell}^{q-3}$ and x_ℓ in V_1 is at least (since $n \geq 20k^2q$)

$$e_G(w_{0\ell}^{q-3}, V_1) + e_G(x_\ell, V_1) - |V_1| \geq \lfloor \frac{n}{2} \rfloor - t_1 + \frac{1}{2} \lfloor \frac{n}{2} \rfloor - \left(\frac{n}{2} + \sqrt{\gamma}n \right) \geq |V(C_{k,q})|.$$

Thus we have found a copy of $C_{s,q}$ centered at x and passing through the edges of $\{xx_1, xx_2, \dots, xx_s\}$. Particularly, since G is $C_{k,q}$ -free, we have $s \leq k - 1$. Thus

$$\Delta_i = \Delta(G_i) \leq k - 1 < t_1, \quad i = 0, 1.$$

Consequently, all the vertices of G are good, and for each vertex $u \in V_i$,

$$e_G(u, V_{1-i}) \geq \left\lfloor \frac{n}{2} \right\rfloor - (k - 1) \geq \frac{n}{2} - k$$

and thereby

$$\frac{n}{2} - k \leq |V_i| \leq \frac{n}{2} + k, \quad i = 0, 1.$$

Next we will find a copy of $C_{k-s,q}$ centered at x disjoint from the copy of $C_{s,q}$. For every u_ℓ ($\ell \in [s + 1, t]$), choose a common neighbor of x and u_ℓ , say $w_{1\ell}^1$, in V_1 such that $w_{1\ell}^1 \neq w_{1\ell'}^1$ if $\ell \neq \ell'$. We can do this because the number of common neighbors of x and u_ℓ in V_1 is at least

$$e_G(x, V_1) + e_G(u_\ell, V_1) - |V_1| \geq 2\left(\frac{n}{2} - k\right) - \left(\frac{n}{2} + k\right) \geq |V(C_{k,q})| \quad (\text{since } n \geq 20k^2q).$$

For each $\ell \in [s + 1, t]$, with the same reason as above, we find vertices $w_{1\ell}^3, w_{0\ell}^4, \dots, w_{0\ell}^{q-3}$ one by one, then a common neighbor of $w_{0\ell}^{q-3}$ and v_ℓ , say $w_{1\ell}^{q-2}$, in V_1 such that

$$u_\ell w_{1\ell}^1 x w_{1\ell}^3 \cdots w_{1\ell}^{q-2} v_\ell u_\ell$$

is a q -cycle. And for every $l \in [t, k]$, begin with x , we can find vertices $w_{1l}^2, w_{0l}^3, \dots, w_{1l}^{q-3}$ one by one, then a common neighbor, say w_{0l}^{q-2} , of w_{1l}^{q-3} and v_l in V_0 such that

$$u_l x w_{1l}^2 w_{0l}^3 \cdots w_{0l}^{q-2} v_l u_l$$

is a q -cycle. Also, we may require that $w_{\xi(j)\ell}^j$ ($\ell \in [s + 1, k], j \in [1, q - 2]$) are pairwise different. Thus the $k - s$ q -cycles form a copy of $C_{k-s,q}$ centered at x and passing through all of the edges of $M_0 \cup M_1$. Therefore, $G[\{x, x_1, \dots, x_t\} \cup \{w_{\xi(j),l}^j : j \in [1, q - 2], \ell \in [1, k]\}]$ contains a desired copy of $C_{k,q}$ with center x . This completes the proof. \square

Claim 11. *We have that $\nu_0 + \nu_1 \leq k - 1$.*

Proof. If not, suppose that $M_0 = \{u_1v_1, \dots, u_s v_s\}$ and $M_1 = \{u_{s+1}v_{s+1}, \dots, u_k v_k\}$ are matchings in G_0 and G_1 , respectively. Without loss of generality, assume that $s \geq 1$. First we find a common neighbor, say x , of u_{s+1}, \dots, u_k in $V_0 \setminus V(M_0)$. This is possible since the number of such neighbors is at least

$$\begin{aligned} & \sum_{l=s+1}^k e_G(u_l, V_0) - (k - s - 1)|V_0| - 2s \\ & \geq (k - s)\left(\frac{n}{2} - k\right) - (k - s - 1)\left(\frac{n}{2} + k\right) - 2k \\ & = \frac{n}{2} - (2(k - s) + 1)k \\ & \geq \frac{n}{2} - 2k^2 > 0 \quad (\text{since } n \geq 20k^2q). \end{aligned}$$

Let M'_0 be the maximal subset of M_0 such that M'_0 is a matching of $G_0 - N_{G_0}[x]$. By Claim 10, $\deg_{G_0}(x) + |M'_0| + |M_1| \leq k - 1$. Clearly, $\deg_{G_0}(x) + |M'_0| \geq s$. Hence $\deg_{G_0}(x) + |M'_0| + |M_1| \geq k$, a contradiction. \square

Claim 12. *We have that $\max\{\Delta_0, \Delta_1\} = k - 1$.*

Proof. If not, then by Claim 10, $\max\{\Delta_0, \Delta_1\} \leq k - 2$. Thus by Lemma 3 and Claim 11,

$$\begin{aligned} m &= e(V_0) + e(V_1) \leq f(\nu_0, k - 2) + f(\nu_1, k - 2) \\ &\leq f(\nu_0 + \nu_1, k - 2) \leq f(k - 1, k - 2). \end{aligned}$$

If $k \neq 4$, $m \leq f(k - 1, k - 2) = (k - 1)^2 - 1$, contradicts to $m \geq (k - 1)^2$ (by Claim 9).

If $k = 4$, then $m \leq f(3, 2) = (k - 1)^2 = 9$. By Claim 9, $m = (k - 1)^2 = 9$ and G contains a complete balanced bipartite subgraph with classes V_0 and V_1 . Let H be the subgraph consisting of nonempty components of $G_0 \cup G_1$. Then H is a graph with $e(H) = 9$, $\Delta(H) = 2$ and $\nu(H) = 3$. By Observation 5 and with a similar discussion as in Case 1 of the proof of Lemma 6, H consists of three vertex-disjoint triangles. Then we can easily find a vertex x in H with $\deg_H(x) = 2$ and a matching of order 2 in $H - N_H[x]$. That is $\deg_{G_i}(x) + \nu(G_i - N_{G_i}[x]) + \nu(G[E_{1-i}(x)]) \geq 4 = k$, a contradiction Claim 10. \square

Claim 13. *We have that $e(V_0) \cdot e(V_1) = 0$.*

Proof. At first, by Claim 12 and Claim 11, we have

$$\begin{aligned} m &\leq f(\nu_0, k - 1) + f(\nu_1, k - 1) \leq f(\nu_0 + \nu_1, k - 1) \\ &\leq f(k - 1, k - 1) \leq k(k - 1). \end{aligned}$$

Next, by Claim 12, $\max\{\Delta_0, \Delta_1\} = k - 1$. Without loss of generality, assume $\Delta_0 = k - 1$. Let $x \in V_0$ with $\deg_{G_0}(x) = k - 1$. We show that $e(V_1) = 0$. If $e(V_1) > 0$, then $\nu_1 \geq 1$. By Claim 11, $\nu_0 \leq k - 1 - \nu_1 = k - 2$. Let $A_1 = \{u \in V_1 : \deg_{G_1}(u) > 0\}$. By Claim 10, we have $A_1 \cap N_G(x) = \emptyset$. So $e(V_0, V_1) \leq |V_0||V_1| - |A_1| \leq e(T_{n,2}) - |A_1|$. Thus we have

$$e(T_{n,2}) + (k - 1)^2 \leq e(G) \leq e(T_{n,2}) - |A_1| + m.$$

Therefore, $|A_1| \leq m - (k - 1)^2 (\leq k - 1)$. Again by Lemma 3, we have

$$\begin{aligned} m &\leq \nu_0(\Delta_0 + 1) + \nu_1(\Delta_1 + 1) \\ &\leq k\nu_0 + (k - 1 - \nu_0)|A_1| \quad (\text{since } \Delta_1 + 1 \leq |A_1|) \\ &= \nu_0(k - |A_1|) + (k - 1)|A_1| \\ &\leq (k - 2)(k - |A_1|) + (k - 1)|A_1| \quad (\text{since } |A_1| \leq k - 1 \text{ and } \nu_0 \leq k - 2) \\ &= (k - 1)^2 + |A_1| - 1 \leq (k - 1)^2 + m - (k - 1)^2 - 1 \\ &= m - 1, \text{ a contradiction.} \end{aligned} \quad \square$$

By Claim 13, Without loss of generality, we may assume $e(V_1) = 0$, so $m = e(V_0)$. Let A_0 be the set of non-isolated vertices in G_0 . By Claim 10 and Lemma 6, $m = e(G[A_0]) \leq (k - 1)^2$. By Claim 9, we must have $m = (k - 1)^2$ and therefore G contains a complete balanced bipartite subgraph with classes V_0 and V_1 . Again by Claim 10 and Lemma 6, since $e(G[A_0]) = (k - 1)^2$, $G[A_0]$ must be a copy of $K_{k-1, k-1}$. This completes the proof of Theorem 2. \square

References

- [1] V. Chavátal, D. Hanson, Degrees and matchings, *J. Combin. Theory Ser. B*, 20(2): 128–138, 1976.
- [2] G. Chen, R. J. Gould, F. Pfender, and B. Wei, Extremal graphs for intersecting cliques, *J. Combin. Theory Ser. B* 89: 159–171, 2003.
- [3] P. Erdős, On some new inequalities concerning extremal properties of graphs, In *Theory of Graphs, Proc. Colloq.*, Tihany, 1966, Academic Press, New York, pp. 77-81, 1968.
- [4] P. Erdős, Z. Füredi, R. J. Gould, D. S. Gunderson, Extremal graphs for intersecting triangles, *J. Combin. Theory Ser. B*, 64(1): 89–100, 1995.
- [5] R. Glebov, Extremal graphs for clique-paths, [arXiv:1111.7029v1](https://arxiv.org/abs/1111.7029v1).
- [6] H. Liu, Extremal graphs for blow-ups of cycles and trees, *The Electron. J. of Combin.*, 20(1): P65, 2013.
- [7] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, In *Theory of Graphs, Proc. Colloq.*, Tihany, 1966, Academic Press, New York, pp. 279–319, 1968.