

Queens in exile: non-attacking queens on infinite chess boards

F. Michel Dekking

Applied Mathematics (DIAM), Delft University of Technology
2600 GA Delft, The Netherlands

F.M.Dekking@math.tudelft.nl

Jeffrey Shallit

School of Computer Science, University of Waterloo
Waterloo, ON N2L 3G1, Canada

shallit@waterloo.ca

N. J. A. Sloane*

The OEIS Foundation Inc.
11 So. Adelaide Ave., Highland Park, NJ 08904, U.S.A.

njasloane@gmail.com

Submitted: Jul 28, 2019; Accepted: Feb 24, 2020; Published: Mar 6, 2020

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

Number the cells of a (possibly infinite) chessboard in some way with the numbers $0, 1, 2, \dots$. Consider the cells in order, placing a queen in a cell if and only if it would not attack any earlier queen. The problem is to determine the positions of the queens. We study the problem for a doubly-infinite chessboard of size $\mathbb{Z} \times \mathbb{Z}$ numbered along a square spiral, and an infinite single-quadrant chessboard (of size $\mathbb{N} \times \mathbb{N}$) numbered along antidiagonals. We give a fairly complete solution in the first case, based on the Tribonacci word. There are connections with combinatorial games.

Keywords: Tribonacci word, Tribonacci representation, Greedy Queens, Wythoff Nim, combinatorial games, Sprague-Grundy function

Mathematics Subject Classifications: 91A46

*Corresponding author

1 Queens in exile




Figure 1: The great plain of Attakía. [John Tenniel, Illustration for Lewis Carroll, *Through the looking-glass and what Alice found there* (1871).]

The rival queens in the mythical country of Attakía have been quarreling, and have agreed to go into exile. The great plain has been divided into squares, which have been numbered in a square spiral (Figs. 1, 2). The first queen settles at square 0. The next queen proceeds along the square spiral and settles at the first square she reaches from which she cannot attack the first queen: this is square 9. The process is repeated for all the queens. Each queen settles at the first square along the spiral from which she cannot attack any queen who is already settled. The positions of the first nine queens are indicated by circles in Fig. 2. The squares along the spiral where they settle form the sequence

$$0, 9, 13, 17, 21, 82, 92, 102, 112, 228, 244, 260, 276, 445, 467, 489, 511, 630, \dots \quad (1)$$

([A273059](#)¹ in [15]).

Figure 3 shows the positions of the first 1409 queens. At this scale one can see that the points lie essentially on four straight lines, and that the configuration has cyclic four-fold symmetry. The main goals of the first part of the paper are to determine the positions

¹Six-digit numbers prefixed by A refer to entries in the *On-Line Encyclopedia of Integer Sequences* [15].




Figure 2: Squares of a doubly-infinite chessboard numbered along a square spiral. Positions of the first nine exiled queens are circled. [Figure courtesy of Jessica Gonzalez.]

of the queens, to establish the cyclic symmetry, and to show that the slopes of the four lines are $\pm\psi$ and $\pm 1/\psi$, where $\psi \approx 1.8393$ is the Tribonacci constant, the real root of $x^3 = x^2 + x + 1$. These results are established in Sections 2–6.

In Section 2 we first show that the positions of the queens are determined by certain recursively defined quadruples of integers $X_n, Y_n, M_n, P_n, n \geq 0$ (see (2), Tables 1, 2, and Theorem 1). A study of the first differences ΔX_n , etc., of the X_n, Y_n, M_n, P_n sequences suggests that all four can be defined in terms of a certain three-letter sequence that we call the “theme song”, and denote by $\Theta(a, b, c)$ (Section 3).

Rather surprisingly, the theme song turns out to be a disguised version of the classic three-letter Tribonacci word $\mathbf{T}(a, b, c)$ (see Theorem 2 in Section 4). Section 5 contains a number of properties of the Tribonacci word that will be used later. Some of these properties appear to be new (Theorems 13–17, for example), although it is difficult to be certain because so much has already been published about the Tribonacci word.

In Section 6 we establish our main theorem, Theorem 20, which shows that the rows of the $XYMP$ table are in one-to-one correspondence with the terms of the theme song Θ (or, if we ignore the initial $n = 0$ term, with the terms of the Tribonacci word \mathbf{T}). Corollary 21 establishes some unexpected connections between the X_n, Y_n, M_n, P_n sequences and the A_n, B_n, C_n sequences studied in [2, 4, 8] and in §5.4. Remark (iv) following Theorem 20 shows that the slopes of the lines containing the queens are as claimed.




Figure 3: Positions of the first 1409 queens (those with maximum coordinate in the range -1000 to 1000). At this scale the points lie essentially on four straight lines. [Figure courtesy of Alois Heinz.]

The reader may wonder why we use both the theme song $\Theta(a, b, c)$ and the Tribonacci word $\mathbf{T}(a, b, c)$, when these sequences are so similar. The answer is that we need $\mathbf{T}(a, b, c)$ because so much is known about its properties (see Section 5), whereas $\Theta(a, b, c)$ is more in tune with the *XYMP* table, since the lengths 6, 5, 4 of the images $\theta(a)$, $\theta(b)$, $\theta(c)$ (see (6)) match the block structure of the table, as can be seen by comparing (4) and (5) with (7).

In Section 7 we consider the same problem in the setting of combinatorial games. The positions of the queens are the \mathcal{P} -positions in a certain game, and so correspond to the 0 entries in the table of Sprague-Grundy values for the game (see Fig. 4 below). Although we have been able to determine the positions of the queens, we have not been able to answer a natural question about the Sprague-Grundy values: are all the rows, columns, and diagonals of Fig. 4 permutations of the nonnegative integers (Conjecture 22)?

Similar questions can be asked for chessboards of other shapes. The general setting for the problem is that the cells of the board are numbered in some way with the numbers $0, 1, 2, 3, \dots$. We consider the cells in order, placing a chess queen in cell n if and only if it would not attack any earlier queen. The problem is to determine the positions of the queens.

In Section 8 we consider the case of an infinite $\mathbb{N} \times \mathbb{N}$ board (that is, a single-quadrant board), where the squares are numbered along successive antidiagonals, as shown in Table 5. For this version of the problem, the data shows overwhelmingly that the queens lie essentially on two straight lines, of slopes ϕ and $1/\phi$, where ϕ is the golden ratio. It is

regrettable that we have not been able to prove this. On the other hand, we have been able to prove that all the rows and columns of the Sprague-Grundy table are permutations of the nonnegative integers (although not that the diagonals are). So for this problem, our results are both weaker and stronger than for the queens-on-a-square-spiral problem.

Many other examples can be found in [15]. These include:

- the board formed from a 45-degree sector of a single quadrant (cells $\{(x, y) \in \mathbb{N} \times \mathbb{N} : x \geq y\}$) ([A274650](#)),
- finite boards of size $n \times n$ ([A308880](#), [A308881](#)),
- boards with hexagonal cells ([A274820](#), [A296339](#)),
- one may also ask similar questions using other chess pieces instead of queens: kings ([A275609](#)), knights ([A308884](#)), rooks ([A308896](#)), or Maharajas (pieces that combine the moves of a queen and a knight [12]: [A307282](#)).

Much is known about these examples, but there is no space to discuss them here. There are many open questions.

Historical remarks. To the best of our knowledge, the first mention of any of these queens-in-exile problems was in [15], in October 2001, when Antti Karttunen contributed [A065188](#), a version of the single-quadrant sequence [A275895](#). He called it a “Greedy Queens” sequence (referring to the fact that the queens are placed using the greedy algorithm—no disrespect to the queens was intended). The problem on the $\mathbb{Z} \times \mathbb{Z}$ board stated at the beginning of this article was introduced (using somewhat different language) by Paul D. Hanna in June 2008 when he submitted [A140100](#)–[A140103](#) to [15] (these are the X_n, Y_n, M_n, P_n sequences), and implicitly stated what is now Theorem 1 below. The Sprague-Grundy values for the single-quadrant version were contributed by Alec Jones in April 2016, in [A269526](#). The connections between the exiled queens problems and combinatorial games were pointed out by Allan C. Wechsler in a comment on [A274528](#). The Sprague-Grundy values for the $\mathbb{Z} \times \mathbb{Z}$ board numbered along a square spiral originated in [A274640](#), contributed in June 2016 by Zak Seidov and Kerry Mitchell. Since then, a large number of other authors (too many to mention here) have added further sequences of this type, or contributed comments, computer programs, additional terms, etc.

Very recently, Fokkink and Rust introduced in [9] a two-pile combinatorial game they call *Splythoff*, where the \mathcal{P} -positions are given by the queen positions (X_n, Y_n) . This is another variant of Wythoff’s Nim. It is different from the game we discuss in Section 7, since their piles contain only nonnegative numbers of tokens.

Notation. The ternary Tribonacci word will be denoted by $\mathbf{T} = t_1 t_2 t_3 \dots$, or by $\mathbf{T}(a, b, c)$ when we wish to emphasize which three-letter alphabet is being used. $\psi = 1.839286755214\dots$ is the Tribonacci constant, the real root of $x^3 - x^2 - x - 1$, and ϕ is the golden ratio. $|S|$ denotes the cardinality of a set, the length of a word, or the absolute value of a complex number. For a set S , $\text{mex}(S)$ is the *minimum excluded value*, that

is, the smallest nonnegative number not in S [10]. For a sequence $\{s_n\}$, the difference operator is defined by $\Delta s_n := s_{n+1} - s_n$. We use a centered dot (\cdot) to indicate concatenation of words (or, rarely, the product of two numbers). \mathbb{Z} and \mathbb{N} are the integers and nonnegative integers, respectively. For any undefined terms from combinatorial games or combinatorics on words, see [1, 10, 13].

2 Queens on a square spiral

In this section we study the problem on a doubly-infinite chessboard. The cells are unit squares centered at the points of a $\mathbb{Z} \times \mathbb{Z}$ grid. We construct a “square spiral” by starting at the central square and proceeding counter-clockwise, moving successively East, North, West, South, East, North, \dots . The cells are numbered $0, 1, 2, 3, \dots$ (see Fig. 2). The exiled queens are placed according to the rule specified in the opening paragraph of the previous section.

We take the x -axis to point South and the y -axis to point East, as shown in the coordinate axes in Fig. 2. This puts the main line of queens (the queens in cells $0, 9, 82, 228, 445, 630, \dots$ in the first quadrant, and is also consistent with having the origin for the single-quadrant version of the problem (Section 8) in the top left corner of the board, as in the discussions of the related games Wyt Queens and Wythoff’s Nim in [3].

We consider the square spiral as being built up from a series of square “shells”. Shell 0 is the starting cell at the center. Shell k ($k = 1, 2, \dots$) consists of the $8k$ cells labeled $(2k - 1)^2$ to $4k(k + 1)$. The spiral traverses shells $0, 1, 2, \dots$ in order. Shell k has four edges, each containing $2k$ cells. Edge 1 (on the right) consists of cells $(2k - 1)^2$ through $4k^2 - 2k$, edge 2 (at top): cells $4k^2 - 2k + 1$ through $4k^2$, edge 3 (on left): cells $4k^2 + 1$ through $4k^2 + 2k$, and edge 4 (at bottom): cells $4k^2 + 2k + 1$ through $4k^2 + 4k$. The spiral traverses shell k along successive edges 1, 2, 3, 4.

We see that the cyclic group of order 4 generated by $(x, y) \mapsto (-y, x)$ preserves the points in each shell.

As Paul Hanna realized in 2008, the positions of the queens in the spiral are determined by certain quadruples of nonnegative integers X_n, Y_n, M_n, P_n ($n \geq 0$), defined by $X_0 = Y_0 = M_0 = P_0 = 0$ and, for $n > 0$,

$$\begin{aligned} X_n &= \text{mex}\{X_i, Y_i : i < n\}, \\ M_n &= \text{mex}\{M_i, P_i : i < n\}, \\ Y_n &= X_n + M_n, \\ P_n &= X_n + Y_n, \end{aligned} \tag{2}$$

where mex denotes “minimum excluded value” as defined above. The initial values of these quadruples are shown in Table 1 (the “XYMP table”), and a more extensive list is given in Table 2 below. These are Paul Hanna’s sequences [A140100–A140103](#).

The following properties are immediate consequences of the definition: $\{X_n\}$ and $\{Y_n\}$ are a pair of complementary sequences, as are $\{M_n\}$ and $\{P_n\}$. All four sequences are monotonically increasing, so $\Delta X_n \geq 1$, $\Delta M_n \geq 1$, $\Delta Y_n \geq 2$, $\Delta P_n \geq 3$. Also $\Delta X_n \leq 2$

Table 1: Initial values of X_n, Y_n, M_n, P_n .

n	X_n	Y_n	M_n	P_n
0	0	0	0	0
1	1	2	1	3
2	3	5	2	8
3	4	8	4	12
4	6	11	5	17
5	7	13	6	20
6	9	16	7	25
7	10	19	9	29
8	12	22	10	34
9	14	25	11	39
10	15	28	13	43
11	17	31	14	48

(if $\Delta X_n = 3$ there would be a pair of adjacent Y values differing by 1, contradicting $\Delta Y_n \geq 2$). Similarly $\Delta M_n \leq 2$, $\Delta Y_n \leq 4$, $\Delta P_n \leq 6$. (In fact ΔY_n is never 4 and ΔP_n is never 6, although we will not prove this until Section 6.) Also $Y_n \geq X_n + 1$ and $P_n \geq M_n + 2$ for $n > 0$.

Let q_n ($n \geq 0$) denote the (x, y) coordinates of the n th queen in the spiral. We saw in Fig. 2 that $q_0 = (0, 0)$, $q_1 = (1, 2)$, $q_2 = (-2, 1)$, $q_3 = (-1, -2)$, $q_4 = (2, -1)$, $q_5 = (3, 5)$, \dots .

The following theorem is implicit in Paul Hanna's remarks in [A140100](#)–[A140103](#).

Theorem 1. *After the initial queen is placed at $q_0 = (X_0, Y_0)$, the subsequent queens are placed at*

$$q_{4k+1} = (X_k, Y_k), \quad q_{4k+2} = (-Y_k, X_k), \quad q_{4k+3} = (-X_k, -Y_k), \quad q_{4k+4} = (-Y_k, X_k), \quad (3)$$

for $k = 0, 1, 2, \dots$.

Proof. Note that the points (3) lie on shell Y_k of the spiral, and this set of four points is preserved by the cyclic group of order 4. We establish (3) by induction on k . The result is true for $k = 1$.

Suppose that the hypothesis holds for $k = 0, 1, \dots, n$. Call a square “free” if a queen at that square would not attack any existing queen. After q_0, \dots, q_{4n+4} have been placed, a square (x, y) is not free if any of the following hold:

- x is equal to $\pm X_i$ or $\pm Y_i$ for some $0 \leq i \leq n$,
- y is equal to $\pm X_i$ or $\pm Y_i$ for some $0 \leq i \leq n$,
- $y - x$ is equal to $\pm M_i$ or $\pm P_i$ for some $0 \leq i \leq n$,

- $y + x$ is equal to $\pm M_i$ or $\pm P_i$ for some $0 \leq i \leq n$,

because (x, y) would then be on the same row, column, or diagonal as one of the existing queens.

When we move along the spiral after placing q_{4n+4} , the first square we reach that does not satisfy any of these conditions is (by (2)) (X_{n+1}, Y_{n+1}) , which is therefore q_{4n+5} . Since $0 < X_{n+1} < Y_{n+1}$, this lies on edge 1 of shell Y_{n+1} .

As we continue around the spiral, we next reach edge 2 of the same shell. Since the configuration of existing queens is preserved by the cyclic group, we would have $q_{4n+6} = (-Y_{n+1}, X_{n+1})$, except we must check that this square does not attack the queen q_{4n+5} we just placed. However, the line from $(-Y_{n+1}, X_{n+1})$ to (X_{n+1}, Y_{n+1}) has slope $(Y_{n+1} - X_{n+1})/(Y_{n+1} + X_{n+1})$, which is not ± 1 , since neither X_{n+1} nor Y_{n+1} is 0. So $q_{4n+6} = (-Y_{n+1}, X_{n+1})$.

Similar arguments show that $q_{4n+7} = (-X_{n+1}, -Y_{n+1})$ and $q_{4n+8} = (Y_{n+1}, -X_{n+1})$. Thus (3) holds for $k = n + 1$. \square

3 The “theme song”

Although it is not immediately apparent, all four columns of the XYMP table are variations on a single sequence. This “theme song” is most visible when we examine the differences $\{\Delta P_n\}$ of the P_n column of the table, keeping in mind the observations about these differences that were made following (2). The differences $\{\Delta P_n\}$ begin

$$3, 5, 4, 5, \quad 3, 5, 4, 5, 5, 4, 5, \quad 3, 5, 4, 5, 4, 5, \quad 3, 5, 4, 5, 5, 4, 5, \quad 3, 5, 4, 5, \quad 3, \dots, \quad (4)$$

where we have inserted spaces to highlight the block structure. The differences of the other columns show a similar, although less obvious, structure. For example, the differences $\{\Delta X_n\}$ begin

$$1, 2, 1, 2, \quad 1, 2, 1, 2, 2, 1, 2, \quad 1, 2, 1, 2, 1, 2, \quad 1, 2, 1, 2, 2, 1, 2, \quad 1, 2, 1, 2, \quad 1, \dots, \quad (5)$$

where we have used the same block lengths. As we will prove in Theorem 20, all four column differences are instances of the sequence $\Theta = \Theta(a, b, c)$ (the “theme song”), the fixed point of the morphism θ defined over the alphabet $\{a, b, c\}$ by

$$\theta : \quad a \rightarrow cabaaba, \quad b \rightarrow cababa, \quad c \rightarrow caba. \quad (6)$$

$\Theta(a, b, c)$ begins

$$c, a, b, a, \quad c, a, b, a, a, b, a, \quad c, a, b, a, b, a, \quad c, a, b, a, a, b, a, \quad c, \dots, \quad (7)$$

and, as we will see, $\{\Delta P_n\} = \Theta(5, 4, 3)$ and $\{\Delta X_n\} = \Theta(2, 1, 1)$.

Table 2: The sequences X_n, Y_n, M_n, P_n and their differences, the identification of the rows with the “theme song” $\Theta(a, b, c) = \{t_n : n \geq 0\}$, and the sequences A_n, B_n, C_n .

n	t_n	X_n	Y_n	M_n	P_n	ΔX_n	ΔY_n	ΔM_n	ΔP_n	A_n	B_n	C_n
0	c	0	0	0	0	1	2	1	3	0	0	0
1	a	1	2	1	3	2	3	1	5	1	2	4
2	b	3	5	2	8	1	3	2	4	3	6	11
3	a	4	8	4	12	2	3	1	5	5	9	17
4	c	6	11	5	17	1	2	1	3	7	13	24
5	a	7	13	6	20	2	3	1	5	8	15	28
6	b	9	16	7	25	1	3	2	4	10	19	35
7	a	10	19	9	29	2	3	1	5	12	22	41
8	a	12	22	10	34	2	3	1	5	14	26	48
9	b	14	25	11	39	1	3	2	4	16	30	55
10	a	15	28	13	43	2	3	1	5	18	33	61
11	c	17	31	14	48	1	2	1	3	20	37	68
12	a	18	33	15	51	2	3	1	5	21	39	72
13	b	20	36	16	56	1	3	2	4	23	43	79
14	a	21	39	18	60	2	3	1	5	25	46	85
15	b	23	42	19	65	1	3	2	4	27	50	92
16	a	24	45	21	69	2	3	1	5	29	53	98
17	c	26	48	22	74	1	2	1	3	31	57	105
18	a	27	50	23	77	2	3	1	5	32	59	109
19	b	29	53	24	82	1	3	2	4	34	63	116
20	a	30	56	26	86	2	3	1	5	36	66	122
21	a	32	59	27	91	2	3	1	5	38	70	129
22	b	34	62	28	96	1	3	2	4	40	74	136
23	a	35	65	30	100	2	3	1	5	42	77	142
24	c	37	68	31	105	1	2	1	3	44	81	149
25	a	38	70	32	108	2	3	1	5	45	83	153
26	b	40	73	33	113	1	3	2	4	47	87	160
27	a	41	76	35	117	2	3	1	5	49	90	166
28	c	43	79	36	122	1	2	1	3	51	94	173

4 The Tribonacci word

It was a further surprise to discover that $\Theta(a, b, c)$ is itself a lightly disguised version of the classic Tribonacci word. The *Tribonacci word* $\mathbf{T} = \mathbf{T}(a, b, c) = \{t_n : n \geq 1\}$ is the fixed point of the *Tribonacci morphism*

$$\tau : \quad a \rightarrow ab, \quad b \rightarrow ac, \quad c \rightarrow a. \quad (8)$$

$\mathbf{T}(a, b, c)$ begins

$$a, b, a, c, a, b, a, a, b, a, c, a, b, a, b, a, c, a, b, a, a, b, a, c, a, \dots \quad (9)$$

There is an extensive literature—see for example [2, 4, 7, 8, 14, 16], as well as the references cited in those papers.

Theorem 2. *We have*

$$\Theta(a, b, c) = c \cdot \mathbf{T}(a, b, c). \quad (10)$$

Proof. The morphism $\alpha := \tau^3$ maps

$$a \rightarrow abacaba, \quad b \rightarrow abacab, \quad c \rightarrow abac.$$

If the prefix aba in these three images is changed to a suffix, α becomes θ ; that is, for single letters x , $\alpha(x)aba = aba\theta(x)$. So $\alpha(x) = aba\theta(x)(aba)^{-1}$, and since α and θ are morphisms, $\alpha(w) = aba\theta(w)(aba)^{-1}$ for any word w , that is,

$$\alpha(w)aba = aba\theta(w). \quad (11)$$

We prove (10) by showing that, for all $k \geq 0$,

$$\theta^k(c)c = c\alpha^k(c). \quad (12)$$

We use induction on k . The result is certainly true for $k = 0$ and 1. Suppose it holds for k , and set $w = \alpha^k(c)$ in (11). We have

$$\begin{aligned} c\alpha^{k+1}(c)aba &= caba\theta(\alpha^k(c)), \quad \text{by (11),} \\ &= \theta(c)\theta(\alpha^k(c)) \\ &= \theta(c\alpha^k(c)) \\ &= \theta(\theta^k(c)c) \quad (\text{by the induction hypothesis}) \\ &= \theta^{k+1}(c)caba, \end{aligned}$$

and canceling aba from both sides we obtain $c\alpha^{k+1}(c) = \theta^{k+1}(c)c$, as required. Letting $k \rightarrow \infty$ in (12) completes the proof of (10). \square

In view of Theorem 2 we define $t_0 = c$, so that $\Theta(a, b, c) = \{t_n : n \geq 0\}$.

5 Properties of the Tribonacci word

The Tribonacci word $\mathbf{T} = \{t_n : n \geq 1\}$ is an analog for a three-letter alphabet of the even more classic two-letter Fibonacci word (see [A003849](#) for an extensive bibliography). In this section we discuss various properties of \mathbf{T} for use later in the paper. Most of the properties are analogs of similar properties of the Fibonacci word.

5.1 The Tribonacci representation of numbers

Define the Tribonacci numbers $\{T_n : n \geq 0\}$ as follows: $T_{-3} = 0$, $T_{-2} = 0$, $T_{-1} = 1$, and $T_n = T_{n-1} + T_{n-2} + T_{n-3}$ for $n \geq 0$ ([A000073](#)).

For $E = e_1e_2 \cdots e_i$ any finite string of 0's and 1's, let $[E]_T$ be the number $n = \sum_{1 \leq j \leq i} e_j T_{i-j}$; we call this a *Tribonacci representation* for n . Among all such representations, one is *canonical*, that obtained by the greedy algorithm (repeatedly subtract the

largest possible Tribonacci number). As is well known (cf. [4]), the canonical representation is uniquely characterized by not containing three consecutive 1's. For integers $n \geq 0$, let $(n)_T$ be this canonical representation for n , written with the least significant digit on the right. For example, 111 and 1000 are both representations of the number 7, but only the latter is canonical. Note that we are distinguishing between $[E]_T$, which is a number, and $(n)_T$, which is a binary string. The notations are combined in the formulas in Theorems 14 and 15.

Lemma 3. *Let x, y be binary strings. Then $[x]_T = [y]_T$ if and only if $[x0]_T = [y0]_T$.*

Proof. Let z be the canonical Tribonacci representation of the integer $[x]_T$. Then $[z]_T = [x]_T$, so it suffices to prove that $[x]_T = [z]_T$ if and only if $[x0]_T = [z0]_T$, where z is canonical.

Suppose $[x]_T = [z]_T$. Consider obtaining z by applying the following replacement rule repeatedly, until no more replacements can be done: 0111 is replaced by 1000. (If 111 appears as a prefix, treat it as if it were 0111.) Each replacement reduces the number of 1's, so the procedure must eventually halt. For example, if $x = 1011011101$, then the following transformations take place (underlining highlights the block that is replaced):

$$1011011101 \rightarrow 1011100001 \rightarrow 1100000001.$$

This normalization procedure transforms every non-canonical binary Tribonacci representation into a canonical one.

Now observe that if we carry out the same process starting instead with $x0$, then the rightmost 0 cannot participate in any of these replacements, and so we end up with $z0$. Hence $[x0]_T = [z0]_T$.

On the other hand, if z is canonical, then so is $z0$. So if $[x0]_T = [z0]_T$, we can obtain $z0$ by processing the representation of $x0$ as above. If we perform these replacements, then the last 0 of both representations never participates in a replacement, and so omitting the last 0 gives exactly the same sequence of replacements. Hence $[x]_T = [z]_T$. \square

Note that the hypothesis that x and y are binary strings is necessary. For example, if we write 4 in a non-canonical way as $[20]_T = [100]_T$, then $8 = [200]_T \neq [1000]_T = 7$.

Corollary 4. *Suppose x, y, w_1, w_2 are binary strings such that $|w_1| = |w_2|$, $[xw_1]_T = [yw_2]_T$, and $[w_1]_T = [w_2]_T$. Then $[x]_T = [y]_T$.*

Proof. Take the equality $[xw_1]_T = [yw_2]_T$ and subtract the equality $[w_1]_T = [w_2]_T$ from it. The result is $[x0^i]_T = [y0^i]_T$, where $i = |w_1| = |w_2|$. Then by applying Lemma 3 i times, we get $[x]_T = [y]_T$. \square

Corollary 5. *Let $e_1 \cdots e_i$ be a binary Tribonacci representation for n . Then the quantity $[e_1 \cdots e_{i-1}]_T + e_i$ does not depend on the particular representation $e_1 \cdots e_i$ chosen for n .*

Proof. Let $E = e_1 \cdots e_i$ and $F = f_1 \cdots f_j$ be two binary Tribonacci representations for n . Without loss of generality, we can assume that one of the representations is canonical.

If $e_i = f_j$, then by Corollary 4 we get $[e_1 \cdots e_{i-1}]_T = [f_1 \cdots f_{j-1}]_T$, and hence

$$[e_1 \cdots e_{i-1}]_T + e_i = [f_1 \cdots f_{j-1}]_T + f_j.$$

Otherwise assume that $e_i = 0$ and $f_j = 1$. If F is the canonical representation for n , then in carrying out the normalization procedure to convert E to F (as we did in the proof of Lemma 3), we evidently cannot change E 's last bit, so $f_j = 0$, a contradiction. So E must be the canonical representation for n . Now consider carrying out the normalization procedure to convert F to E . To change $f_j = 1$ into the 0 corresponding to e_i , the only possibility is that the rightmost four bits of F are 0111 and the rightmost four bits of E are 1000. Write $E = E'1000$ and $F = F'0111$. Then $[E'1000]_T = [F'0111]_T$, and by Corollary 4 we get $[E']_T = [F']_T$. By applying Lemma 3 three times, we get $[E'000]_T = [F'000]_T$. Adding $[100]_T + 0 = [011]_T + 1$ to both sides gives $[E'100] + 0 = [F'011]_T + 1$, as desired. \square

5.2 The Tribonacci morphism

Using the Tribonacci morphism (8), we define a sequence of finite binary words by $\mathbf{T}_n := \tau^n(a)$ for $n \geq 0$. Then the Tribonacci word is $\mathbf{T} = \lim_{n \rightarrow \infty} \mathbf{T}_n$.

Theorem 6. *We have*

$$\begin{aligned}\tau^n(a) &= \mathbf{T}_n, \quad n \geq 0, \\ \tau^n(b) &= \mathbf{T}_{n-1} \cdot \mathbf{T}_{n-2}, \quad n \geq 1, \\ \tau^n(c) &= \mathbf{T}_{n-1}, \quad n \geq 0.\end{aligned}$$

Proof. An easy induction on n . \square

Lemma 7. *We have $\mathbf{T}_n = \mathbf{T}_{n-1} \cdot \mathbf{T}_{n-2} \cdot \mathbf{T}_{n-3}$ for $n \geq 3$.*

Proof. We have

$$\begin{aligned}\mathbf{T}_n &= \tau^n(a) = \tau^{n-3}(\tau^3(a)) = \tau^{n-3}(abacaba) = \tau^{n-3}(abac)\tau^{n-3}(ab)\tau^{n-3}(a) \\ &= \tau^{n-1}(a)\tau^{n-2}(a)\tau^{n-3}(a) = \mathbf{T}_{n-1} \cdot \mathbf{T}_{n-2} \cdot \mathbf{T}_{n-3}.\end{aligned}$$

\square

The next two lemmas are also easily established by induction:

Lemma 8. *We have $|\mathbf{T}_n| = T_n$ for $n \geq 0$.*

Lemma 9. *For $n \geq 0$, \mathbf{T}_n contains T_{n-1} a's, T_{n-2} b's, and T_{n-3} c's.*

5.3 The Tribonacci word

Lemma 10. *Let U_m be the set of length- m binary strings consisting of the Tribonacci representations (padded with leading zeros, if necessary) of the numbers from 0 to $T_m - 1$. Then*

$$U_m = 0U_{m-1} \sqcup 10U_{m-2} \sqcup 110U_{m-3},$$

for $m \geq 3$, where \sqcup denotes disjoint union.

Proof. Again an easy induction on m . \square

Theorem 11. ([8]). *Let the Tribonacci representation of $n - 1$ be $e_1 e_2 \cdots e_i 0 1^j$, where $j \in \{0, 1, 2\}$. Then*

$$t_n = \begin{cases} a, & \text{if } j = 0, \\ b, & \text{if } j = 1, \\ c, & \text{if } j = 2. \end{cases}$$

Proof. An easy proof by induction on n , where $T_m + 1 \leq n \leq T_{m+1}$, using Lemma 9. \square

For $n \geq 1$ let $N_a(n), N_b(n), N_c(n)$ be the number of a 's, b 's, c 's respectively in $t_1 \cdots t_n$.

Theorem 12. *For $n \geq 1$ let the Tribonacci representation of n be $e_1 e_2 \cdots e_i$. Then*

$$\begin{aligned} N_a(n) &= [e_1 \cdots e_{i-1}]_T + e_i, \\ N_b(n) &= [e_1 \cdots e_{i-2}]_T + e_{i-1}, \\ N_c(n) &= [e_1 \cdots e_{i-3}]_T + e_{i-2}. \end{aligned}$$

Proof. We know from Theorem 11 that t_n depends on the Tribonacci representation of $n - 1$ rather than n , so for this proof we set $\nu = n - 1$. The proof is by induction on ν . Consider a ν in the range $T_m \leq \nu < T_{m+1}$. We prove the result for $N_a(n)$, with the other results being exactly analogous. The base cases are easy.

For the induction step, there are two cases to consider: (a) $T_m \leq \nu < T_m + T_{m-1}$ and (b) $T_m + T_{m-1} \leq \nu < T_{m+1}$.

If (a) holds, then write $\nu = T_m + \nu'$ with $0 \leq \nu' < T_{m-1}$, and consider the length- m Tribonacci representations of the numbers from 0 to $\nu - 1$. Using Lemma 10, we see that the length- m representations of the numbers from 0 to $T_m - 1$ start with 0, while the numbers from T_m to $T_m + \nu' - 1$ have representations that start with 10. For these latter numbers, subtracting T_m removes a leading 1 from the Tribonacci representation, and using Lemma 7 we see that the Tribonacci representations of the numbers from T_m to $T_m + \nu' - 1$ are those of the numbers from 0 to $\nu' - 1$, except for this leading 1. By Theorem 11 it follows that the number of a 's in $t_{T_m+1} \cdots t_{T_m+\nu'}$ is equal to $N_a(\nu')$.

Hence

$$\begin{aligned} N_a(n) &= N_a(T_m) + N_a(\nu') \\ &= T_{m-1} + N_a(\nu') \quad (\text{by Lemma 9}) \\ &= T_{m-1} + [e_3 \cdots e_{i-1}]_T + e_i \quad (\text{by the induction hypothesis}) \\ &= [e_1 \cdots e_{i-1}]_T + e_i. \end{aligned}$$

Case (b) is similar. \square

Theorem 13. *Let the Tribonacci representation of $n - 1$ be $e_1 \cdots e_i$, the representation of $n - 2$ be $f_1 \cdots f_j$, and the representation of $n - 4$ be $g_1 \cdots g_k$. Then*

$$\begin{aligned} N_a(n) &= [e_1 \cdots e_{i-1}]_T + 1, \\ N_b(n) &= [f_1 \cdots f_{j-2}]_T + 1, \\ N_c(n) &= [g_1 \cdots g_{k-3}]_T + 1. \end{aligned}$$

Proof. We prove the result for $N_a(n)$, the others being similar. There are three cases, depending on whether $(n-1)_T = x0$, $x01$, or $x011$. If $(n-1)_T = x0$, then $x1$ is a representation of n ; if $(n-1)_T = x01$, $x10$ is a representation of n ; and if $(n-1)_T = x011$, $x100$ is a representation of n . The result then follows by combining Corollary 5 with Theorem 12 \square

An analogous property to Theorem 13 for the Fibonacci word was established in [6, Section 5].

5.4 The indexing sequences A_n , B_n , C_n .

Let A_n ($n \geq 1$) denote the index of the n th occurrence of the letter a in \mathbf{T} , with similar definitions for B_n and C_n . We also set $A_0 = B_0 = C_0 = 0$. The initial values of A_n , B_n , C_n are

n	0	1	2	3	4	5	6	7	8	9	10	11	12	...
A_n	0	1	3	5	7	8	10	12	14	16	18	20	21	...
B_n	0	2	6	9	13	15	19	22	26	30	33	37	39	...
C_n	0	4	11	17	24	28	35	41	48	55	61	68	72	...

(13)

For further terms see Table 2 or [A003144](#), [A003145](#), [A003146](#). These sequences are inverses to the sequences $N_a(n)$, $N_b(n)$, $N_c(n)$ defined in §5.3. For example, $A_5 = 8$, while $N_a(8) = 5$. They are studied in many references ([2, 4, 8]).

Theorem 14. *We have*

- $A_n = [(n-1)_T 0]_T + 1$,
- $B_n = [(n-1)_T 01]_T + 1$,
- $C_n = [(n-1)_T 011]_T + 1$.

Proof. We use Theorem 12, which tells us how many a 's (resp., b 's, c 's) occur in a prefix of \mathbf{T} of a given length. We prove the result for A_n , with the other results being proved analogously. Let us count how many a 's there are in a prefix $[(n-1)_T 0]_T$. By Theorem 12, there are $n-1$ of them. Similarly, Theorem 12 says that there are n a 's in the prefix of length $[(n-1)_T 1]_T$. Since we index \mathbf{T} starting at position 1, it now follows that the symbol at position $1 + [(n-1)_T 0]_T$ must be an a . \square

We also record some further properties of A_n , B_n , C_n established in [8]. For $n \geq 1$, we have

$$A_n = \text{mex}\{A_i, B_i, C_i : 0 \leq i < n\}, \quad (14)$$

$$B_n = A_n + \text{mex}\{B_i - A_i, C_i - B_i : 0 \leq i < n\}, \quad (15)$$

$$C_n = A_n + B_n + n. \quad (16)$$

There are some similarities with (2), and in particular (15) is consistent with equation (41) of Corollary 21 below, although we will not prove this observation is correct until Section 6.

Furthermore, \mathbf{T} is the unique ternary sequence satisfying (14)–(16). Also from [8] (see Remarks 2.1–2.3), we know that

$$\{\Delta A\} = \Theta(2, 2, 1), \{\Delta B\} = \Theta(4, 3, 2), \{\Delta C\} = \Theta(7, 6, 4). \quad (17)$$

The next three properties are easy consequences of the definitions.

$$A_n = n + N_a(n-1) + N_b(n-1), \quad (18)$$

$$B_n = A_n + N_a(A_n) + N_b(A_n), \quad (19)$$

$$C_n = B_n + N_a(B_n) + N_b(B_n). \quad (20)$$

We prove (18), since the same argument will be used later. Since \mathbf{T} is the fixed point of τ , $\mathbf{T} = \tau(\mathbf{T})$. Writing $\tau(t_n)$ underneath t_n , we see:

$$\begin{array}{cccccccccccccccc} n : & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & \dots \\ t_n : & a & b & a & c & a & b & a & a & b & a & c & a & b & \dots \\ \tau(t_n) : & ab & ac & ab & a & ab & ac & ab & ab & ac & ab & a & ab & ac & \dots \end{array}$$

Each letter t_n in \mathbf{T} produces an a in $\tau(t_n)$, and this a is at position p , where p equals n plus the total number of a 's and b 's before t_n . This is exactly the assertion (18). Properties (19) and (20) have similar proofs.

Many other properties are known, such as ([2, 8])

$$A_{A_n} + 1 = B_n, \quad A_{B_n} = B_{A_n} + 1, \quad A_{B_n} + 1 = C_n. \quad (21)$$

5.5 Numerical bounds.

Appending a 0 to the Tribonacci representation of a number (as in the formula for A_n given in Theorem 14) has about the same effect as multiplying the number by ψ . To get precise estimates we must study the Tribonacci numbers T_n themselves.

From the theory of linear recurrences we know that if ψ , ψ_2 and $\psi_3 := \overline{\psi_2}$ denote the roots of $x^3 = x^2 + x + 1$ then there are constants c_1 , c_2 , and $c_3 := \overline{c_2}$ such that

$$T_n = c_1 \psi^n + c_2 \psi_2^n + c_3 \psi_3^n, \quad \text{for } n \geq 0. \quad (22)$$

The numerical values of these constants are $\psi_2 = -0.419643\dots + 0.606291\dots i$, $c_1 = 0.336228\dots$, $c_2 = -0.168114\dots - 0.198324\dots i$.

From (22) it is straightforward to show that, for $n \geq 0$, we have

$$|T_n - c_1 \psi^n| \leq 0.283 (0.738)^n \quad (23)$$

and

$$|T_{n+1} - \psi T_n| \leq 0.731 (0.738)^n, \quad (24)$$

$$|T_{n+2} - \psi^2 T_n| \leq 1.113 (0.738)^n, \quad (25)$$

$$|T_{n+3} - \psi^3 T_n| \leq 1.877 (0.738)^n. \quad (26)$$

Theorem 15. For all $n \geq 0$ we have

$$-0.596 < [(n)_T 0]_T - \psi n < 0.856, \quad (27)$$

$$-0.883 < [(n)_T 00]_T - \psi^2 n < 1.460, \quad (28)$$

$$-1.461 < [(n)_T 000]_T - \psi^3 n < 2.298. \quad (29)$$

Proof. We prove (27), the arguments in the other two cases being similar. We write n in its canonical Tribonacci representation, say $n = T_{e_1} + T_{e_2} + \cdots + T_{e_s}$, where $e_1 > e_2 > \cdots > e_s$. Then $[(n)_T 0]_T = T_{e_1+1} + T_{e_2+1} + \cdots + T_{e_s+1}$, so

$$[(n)_T 0]_T - \psi n = \sum_{1 \leq j \leq s} (T_{e_j+1} - \psi T_{e_j}).$$

We break up this sum into two parts, one where $2 \leq e_j \leq 18$, and one where $e_j > 18$. From (24), the latter sum is bounded in absolute value by $\sum_{j \geq 19} 0.731 \cdot 0.738^j \leq .009$. The former sum can be bounded by actually computing it for all $n < T_{19} = 121415$. The minimum is achieved at $n = 65915$ and is, rounded down, equal to -0.587 . The maximum is achieved at $n = 78748$ and is, rounded up, equal to 0.847 . Hence $-0.596 < [(n)_T 0]_T - \alpha_1 n < 0.856$. \square

Theorem 16. For all $n \geq 0$,

$$\lfloor \psi n \rfloor - 1 \leq A_n \leq \lfloor \psi n \rfloor + 1, \quad (30)$$

$$\lfloor \psi^2 n \rfloor - 2 \leq B_n \leq \lfloor \psi^2 n \rfloor + 1, \quad (31)$$

$$\lfloor \psi^3 n \rfloor - 3 \leq C_n \leq \lfloor \psi^3 n \rfloor + 1. \quad (32)$$

Proof. From Theorem 15 we get $-.596 < [(n-1)_T 0]_T - \psi(n-1) < .856$. Since $A_n = [(n-1)_T 0]_T + 1$, we have $-.596 < A_n - 1 - \psi(n-1) < .856$. Hence $-.017 < \psi n - A_n < 1.436$, so $A_n - .017 < \psi n < A_n + 1.436$. Taking floors gives

$$A_n - 1 \leq \lfloor \psi n \rfloor \leq A_n + 1,$$

which proves the first assertion. The other two have similar proofs, which we omit. \square

Values of n for which the “+ 1” on the right-hand sides of (30)–(32) are actually needed seem quite rare, the first instances occurring at $n = 12737$, 329 , and 2047 , respectively (see [A275158](#), [A278352](#), [A278353](#)).

For use in the proof of Theorem 20 we note that

$$\psi C_n > C_{n+1} \text{ for } n \geq 2. \quad (33)$$

Bounds on the inverse quantities are easier to derive, and we just state the result. Deleting the least significant bit of the Tribonacci representation of n has about the same effect as dividing n by ψ . From Theorem 12 we obtain:

Theorem 17. For all $n \geq 1$,

$$\left\lfloor \frac{n}{\psi} \right\rfloor \leq \frac{n}{\psi} \leq N_a(n) \leq \left\lceil \frac{n}{\psi} \right\rceil + 1, \quad (34)$$

$$\left\lfloor \frac{n}{\psi^2} \right\rfloor \leq \frac{n}{\psi^2} \leq N_b(n) \leq \left\lceil \frac{n}{\psi^2} \right\rceil + 1, \quad (35)$$

$$\left\lfloor \frac{n}{\psi^3} \right\rfloor \leq \frac{n}{\psi^3} \leq N_c(n) \leq \left\lceil \frac{n}{\psi^3} \right\rceil + 1. \quad (36)$$

6 The main theorem

We can now establish our main theorem, Theorem 20, which shows that the rows of the *XYMP* table are in one-to-one correspondence with the terms of the theme song Θ (or, if we ignore the initial $n = 0$ term, with the terms of the Tribonacci word \mathbf{T}). The bijection can be seen in the second column of Table 2.

Lemma 18. Let τ' be the morphism $\tau' : a \rightarrow ba, b \rightarrow ca, c \rightarrow a$. Then

$$\tau'(c \cdot \mathbf{T}(a, b, c)) = \mathbf{T}(a, b, c). \quad (37)$$

Proof. Let A'_n ($n \geq 1$) denote the index of the n th occurrence of the letter a in the image $\tau'(c \cdot \mathbf{T}(a, b, c))$, with similar definitions for B'_n and C'_n . Write the terms of $\tau'(c \cdot \mathbf{T}(a, b, c))$ under the corresponding terms of $c \cdot \mathbf{T}(a, b, c)$, as we did in the proof of (18)–(20) above. As in that proof, we observe that each term of $c \cdot \mathbf{T}(a, b, c)$ produces an a in the image, only now the initial c produces an extra a at the start, and the a produced by $t_n \in \mathbf{T}(a, b, c)$ is displaced from $n + 1$ by the number of copies of a and b in $\mathbf{T}(a, b, c)$ at or before t_n . This implies that

$$A'_{n+1} = n + 1 + N_a(n) + N_b(n).$$

So $A'_n = n + N_a(n - 1) + N_b(n - 1) = A_n$. Similar arguments show that $B'_n = B_n$ and $C'_n = C_n$, and so (37) holds. \square

Lemma 19. Let τ'' be the morphism $\tau'' : a \rightarrow acab, b \rightarrow aab, c \rightarrow ab$. Then

$$\tau''(c \cdot \mathbf{T}(a, b, c)) = \mathbf{T}(a, b, c). \quad (38)$$

Proof. We have $\tau'' = \tau \circ \tau'$ (first apply τ' then τ), and τ fixes $\mathbf{T}(a, b, c)$, so the result follows from Lemma 18. \square

Theorem 20. We have

$$\begin{aligned} \{\Delta X\} &= \Theta(2, 1, 1), \\ \{\Delta Y\} &= \Theta(3, 3, 2), \\ \{\Delta M\} &= \Theta(1, 2, 1), \\ \{\Delta P\} &= \Theta(5, 4, 3). \end{aligned} \quad (39)$$

Proof. There are three parts to the proof. In the first part we show that, for any $i \geq 1$, if terms 0 through $C_i - 1$ of $\{\Delta Y\}$ (that is, the terms until just before the i th occurrence of $c = 2$ in $\{\Delta Y\}$) agree with terms 0 through $C_i - 1$ of $\Theta(3, 3, 2)$, then terms 0 through $\lfloor \psi(C_i - 1) \rfloor$ of $\{\Delta X\}$ agree with terms 0 through $\lfloor \psi(C_i - 1) \rfloor$ of $\Theta(2, 1, 1)$.

We form the partial sums of the $\{\Delta Y\}$ to get C_i terms of $\{Y\}$, compute the complement to get a certain number of initial terms of $\{X\}$, and take the differences to get the initial terms of $\{\Delta X\}$.

A 3 in the $\{\Delta Y\}$ sequence corresponds to a succession $\dots, x, x + 2, x + 3, x + 5, \dots$ of terms in $\{X\}$, with differences $\dots, 2, 1, 2, \dots$. The initial 2 was already present in $\{\Delta X\}$, so the 3 in $\{\Delta Y\}$ produces a pair 1, 2 in $\{\Delta X\}$. Similarly, a term $\Delta Y = 2$ produces a single 2 in $\{\Delta X\}$. The initial C_i terms of $\{\Delta Y\}$ are transformed by the action of the map $3 \rightarrow 1, 2; 2 \rightarrow 2$ into a sequence over the alphabet $\{1, 2\}$. We can state this assertion in an equivalent way. We identify the initial terms of $\{\Delta Y\}$ with the initial terms of $\Theta(a, b, c) = \Theta(3, 3, 2)$, and the initial terms of $\{\Delta X\}$ with the initial terms of $\Theta(a, b, c) = \Theta(2, 1, 1)$. The map can be described as $\tau' : a \rightarrow ba, b \rightarrow ca, c \rightarrow a$, and Lemma 18 guarantees that the image is indeed a prefix of $\Theta(2, 1, 1)$.

Table 3: Showing how the difference sequence $\{\Delta Y\}$ generates a larger number of terms of the difference sequence $\{\Delta X\}$. The first four lines refer to $\{\Delta Y\}$ and $\{Y\}$, the second four to $\{X\}$ and $\{\Delta X\}$.

$n:$	0	1		2			3		4		5		6		7				
$\Theta:$	c	a		b			a		c		a		b		a				
$\Delta Y_n:$	2	3		3			3		2		3		3		3				
$Y_n:$	0	2		5			8		11		13		16		19				
$X_n:$	0	1		3	4		6	7		9	10		12	14	15		17	18	
$\Delta X_n:$	1		2	1		2	1		2	1		2	1		2	1		2	
$\Theta:$	c		a	b		a	c		a	b		a	a	b		a	c	a	
$n:$	0		1	2		3	4		5	6		7		8	9		10	11	12

$n:$			8			9			10			11			12			13			14
$\Theta:$			a			b			a			c			a			b			a
$\Delta Y_n:$			3			3			3			2			3			3			3
$Y_n:$			22			25			28			31			33			36			39
$X_n:$	20	21		23	24		26	27		29	30		32		34	35			37	38	
$\Delta X_n:$	1		2	1		2	1		2	1		2		2	1		2	1		2	
$\Theta:$	b		a	b		a	c		a	b		a		a	b		a	c		a	
$n:$	13		14	15		16	17		18	19		20		21	22		23	24		25	

This process is illustrated in Table 3. The first four rows show $\{\Delta Y\}$ as $\Theta(a, b, c) = \Theta(3, 3, 2)$, and its partial sums $\{Y\}$. The second four rows show $\{X\}$ being formed as the complement of $\{Y\}$, and the identification of $\{\Delta X\}$ with $\Theta(a, b, c) = \Theta(2, 1, 1)$.

From Theorem 2, in terms 1 through $k := C_i - 1$ of $\Theta(a, b, c)$ there are $N_a(k)$, $N_b(k)$, $N_c(k)$ copies of a , b , c , respectively. After applying the map τ' , and taking into account the slightly irregular behavior at the start of these sequences, we obtain $2 + 2N_a(k) + 2N_b(k) + N_c(k) = 2 + 2k - N_c(k)$ terms of $\{\Delta X\}$. For illustration, in Table 3, we may take $i = 2$, $C_2 = 11$, $k = 10$, and using $N_a(10) = 6$, $N_b(10) = 3$, $N_c(10) = 1$, we see that we obtain $19 + 2 = 21$ terms of $\{\Delta X\}$, that is, terms 0 through 20.

Using (36), the number of terms we obtain is at least $1 + k(2 - 1/\psi^3) = \psi k + 1 = \psi(C_i - 1) + 1$. This completes the first part of the proof of the theorem.

In the second part of the proof we show that, for any $i \geq 1$, if terms 0 through $C_i - 1$ of $\{\Delta P\}$ agree with terms 0 through $C_i - 1$ of $\Theta(5, 4, 3)$, then terms 0 through $\lfloor \psi^2(C_i - 1) \rfloor$ of $\{\Delta M\}$ agree with terms 0 through $\lfloor \psi^2(C_i - 1) \rfloor$ of $\Theta(1, 2, 1)$.

The argument is parallel to that for the first part: we proceed from $\{\Delta P\}$ to $\{P\}$ to its complement, $\{M\}$, and then to $\{\Delta M\}$ (see Table 4). Every 5 in $\{\Delta P\}$ produces a subsequence 1, 1, 1, 2 in $\{\Delta M\}$, every 4 produces 1, 1, 2, and every 3 produces 1, 2.

Table 4: Showing how the difference sequence $\{\Delta P\}$ generates a larger number of terms of the difference sequence $\{\Delta M\}$. The first four lines refer to $\{\Delta P\}$ and $\{P\}$, the second four to $\{M\}$ and $\{\Delta M\}$.

$n:$	0			1					2					3					4	
$\Theta:$	c			a					b					a					c	
$\Delta P_n:$	3			5					4					5					3	
$P_n:$	0			3					8					12					17	
$M_n:$	0	1	2		4	5	6	7		9	10	11		13	14	15	16		18	19
$\Delta M_n:$	1	1		2	1	1	1		2	1	1		2	1	1	1		2	1	1
$\Theta:$	c	a		b	a	c	a		b	a	a		b	a	c	a		b	a	
$n:$	0	1		2	3	4	5		6	7	8		9	10	11	12		13	14	

$n:$	5				6				7					8					9
$\Theta:$	a				b				a					a					b
$\Delta P_n:$	5				4				5					5					4
$P_n:$	20				25				29					34					39
$M_n:$		21	22	23	24		26	27	28		30	31	32	33		35	36	37	38
$\Delta M_n:$	2	1	1	1		2	1	1		2	1	1	1		2	1	1	1	2
$\Theta:$	b	a	c	a		b	a	a		b	a	c	a		b	a	c	a	b
$n:$	15	16	17	18		19	20	21		22	23	24	25		26	27	28	29	30

We identify the initial terms of $\{\Delta P\}$ with the initial terms of $\Theta(a, b, c) = \Theta(5, 4, 3)$, and the initial terms of $\{\Delta M\}$ with the initial terms of $\Theta(a, b, c) = \Theta(1, 2, 1)$. The map can be described as $\tau'' : a \rightarrow acab, b \rightarrow aab, c \rightarrow ab$, and Lemma 19 guarantees that the image is indeed a prefix of $\Theta(1, 2, 1)$.

When τ'' is applied to terms 0 through $k := C_i - 1$ of $\Theta(a, b, c)$ we obtain $3 + 4N_a(k) +$

$3N_b(k) + 2N_c(k)$ terms of $\{\Delta M\}$. From (34)–(36), this quantity is at least $3 + k(4/\psi + 3/\psi^2 + 2/\psi^3) = \psi^2 k + 3$, which is enough to complete the second part of the proof.

For the third part of the proof we use induction. The induction hypothesis is that for some $i \geq 2$, terms 0 through $C_i - 1$ of $\{\Delta Y\}$ agree with terms 0 through $C_i - 1$ of $\Theta(3, 3, 2)$, and terms 0 through $C_i - 1$ of $\{\Delta P\}$ agree with terms 0 through $C_i - 1$ of $\Theta(5, 4, 3)$. From Table 2 we can verify that these assertions are true for all $i \leq 3$.

From the first two parts of the proof the induction hypothesis implies that $\{\Delta X\}$ agrees with $\Theta(2, 1, 1)$ for $\psi C(i)$ terms, and $\{\Delta M\}$ agrees with $\Theta(1, 2, 1)$ for at least the same number of terms. Since $\{Y\} = \{X\} + \{M\}$, $\{\Delta Y\}$ agrees with $\Theta(2, 1, 1) + \Theta(1, 2, 1) = \Theta(3, 3, 2)$ for ψC_i terms. Since $\{P\} = \{X\} + \{Y\}$, $\{\Delta P\}$ agrees with $\Theta(2, 1, 1) + \Theta(3, 3, 2) = \Theta(5, 4, 3)$ for ψC_i terms. But by (33), $\psi C_i > C_{i+1}$ (this is where we need $i \geq 2$), so the induction hypothesis holds for $i + 1$. This completes the proof. \square

Remarks.

(i) The identification of $\{\Delta P\}$ with $\Theta(5, 4, 3)$ establishes the bijection between the rows of the XYMP table and the terms of the sequence $\Theta(a, b, c)$ (see column 2 of Table 2). The label for row $n \geq 0$ is determined by the value of the quadruple $(\Delta X_n, \Delta Y_n, \Delta M_n, \Delta P_n)$: if this is

$$\begin{aligned} (2, 3, 1, 5) & \text{ then row } n \text{ is labeled } a, \\ (1, 3, 2, 4) & \text{ then row } n \text{ is labeled } b, \\ (1, 2, 1, 3) & \text{ then row } n \text{ is labeled } c. \end{aligned} \tag{40}$$

(ii) Since $\Delta X_n = 2$ only in the first case, and $\Delta M_n = 2$ only in the second case, we conclude that:

row n is labeled a if and only if $X_n + 1 \in \{Y\}$,
row n is labeled b if and only if $M_n + 1 \in \{P\}$, and
otherwise row n is labeled c .

(iii) We can now connect the X_n, Y_n, M_n, P_n sequences with the A_n, B_n, C_n sequences.

Corollary 21. For $n \geq 0$,

$$X_n = B_n - A_n, \tag{41}$$

$$Y_n = C_n - B_n, \tag{42}$$

$$M_n = C_n - 2B_n + A_n, \tag{43}$$

$$P_n = C_n - A_n. \tag{44}$$

Proof. These formulas follow from (17). The first one, for example, follows because $\{\Delta B\} - \{\Delta A\} = \Theta(4, 3, 2) - \Theta(2, 2, 1) = \Theta(2, 1, 1) = \{\Delta X\}$, by (39), and so $X_n = B_n - A_n$. \square

These formulas can be confirmed by looking at the columns of Table 2.

(iv) The queens in the first quadrant (cf. Fig. 3) have coordinates (X_n, Y_n) , $n \geq 0$, and we can now determine the slope of the line that they (approximately) lie on. For

$$\frac{Y_n}{X_n} = \frac{C_n - B_n}{B_n - A_n}, \quad (45)$$

which from Theorem 16 converges to $(\psi^3 - \psi^2)/(\psi^2 - \psi) = \psi$ as n increases.

(v) We can also answer a question left over from Section 2. From Theorem 20, we see that ΔY_n is never 4, and ΔP_n is never 6. So there is no run of three consecutive X_n all differing by 1, and no run of five consecutive M_n all differing by 1.




Figure 4: Sprague-Grundy values ([A274641](#)) for game based on the queens-in-exile problem on a square spiral. The cells with value 0 are both the \mathcal{P} -positions for the game and the locations of the exiled queens. [Figure courtesy of Jessica Gonzalez.]

7 The Sprague-Grundy values

The queens-in-exile problems can be described in terms of two-person, impartial, combinatorial games. We fix a numbering of the cells (such as along a square spiral, or by antidiagonals). A queen is placed anywhere on the board, and the players take turns, moving the queen to any lower-numbered cell which is a queen’s move away. The first player who is unable to move loses. The Sprague-Grundy value of a cell is the “mex” of

the values of the cells that are a legal move away [3, 10]. The \mathcal{P} -positions are the cells with Sprague-Grundy value 0, and are also the locations of the exiled queens.

For the $\mathbb{Z} \times \mathbb{Z}$ board numbered along a square spiral, the Sprague-Grundy values are shown in Fig. 4. By construction, the Sprague-Grundy values along any row are distinct, and similarly for any column, and any diagonal of slope ± 1 . The following conjecture is very plausible but unsolved.

Conjecture 22. Every row, column, and diagonal of slope ± 1 in the array of Sprague-Grundy values is a permutation of \mathbb{N} .

We do not even know, for example, that the numbers along the horizontal axis,

$$\dots, 24, 12, 16, 9, 14, 6, 5, 4, 2, 0, 1, 3, 7, 10, 11, 15, 8, 18, 23, 21, \dots, \quad (46)$$

([A324778](#), [A324774](#)) include every nonnegative number.

Table 5: Single-quadrant board numbered along upwards antidiagonals; circles indicate positions of exiled queens.

①	2	5	9	14	20	27	35	44	...
1	4	8	⑬	19	26	34	43	...	
3	⑦	12	18	25	33	42	...		
6	11	17	24	③②	41	...			
10	16	②③	31	40	...				
15	22	30	39	...					
21	29	38	...						
28	37	...							
36	...								
...									

8 The single-quadrant board

We have fewer results about the positions of the queens in this version of the problem, so we will start right away with the combinatorial game. This is played on an infinite $\mathbb{N} \times \mathbb{N}$ board where the squares are numbered along successive upward antidiagonals, as shown in Table 5. In the game, a queen is placed anywhere on the board, and the players take turns moving it to a lower-numbered square that is a queen's-move away. As usual the first player unable to move loses. A small portion of the table of Sprague-Grundy values is shown in Table 6, and a color-coded illustration of the top 500×500 corner of the table is given in Fig. 5.

Table 6: Sprague-Grundy values for single-quadrant board numbered along upwards antidiagonals ([A274528](#)). The indexing of the rows and columns in this table start at 0.

0	2	1	5	3	4	9	10	12	...
1	3	4	0	7	2	5	11	...	
2	0	5	1	8	6	4	...		
3	1	2	4	0	7	...			
4	6	0	3	1	...				
5	7	8	6	...					
6	4	3	...						
7	5	...							
8	...								
...									




Figure 5: The top 500×500 corner of the table of Sprague-Grundy values. The color ranges from red (for values near 0) to blue (for values near 1000). [Figure courtesy of Remy Sigrist.]

The squares where the exiled queens settle in this version of the problem (i.e., the 0's in the table) have indices, reading along the successive antidiagonals,

$$0, 7, 13, 23, 32, 96, 114, 142, 163, 183, 197, 261, 290, 446, 484, 581, \dots \quad (47)$$

([A275897](#)). The first five of these values are indicated by the circles in Fig. 5.

An alternative way to specify the positions of the exiled queens is by the sequence $\{S_c : c \geq 0\}$ ([A275895](#)), which indicates which row contains the queen in column c (this is well-defined, thanks to Theorem 23 below). The initial values of S_c are

$$0, 2, 4, 1, 3, 8, 10, 12, 14, 5, 7, 18, 6, 21, 9, 24, 26, 28, 30, 11, 13, 34, \dots \quad (48)$$

The only theorem we have for this version of the problem is:

Theorem 23. *Every column and every row of the table of Sprague-Grundy values is a permutation of \mathbb{N} .*

Proof. (Based on arguments in [A269526](#) given by Rob Pratt, Bob Selcoe, and N.J.A.S. in June 2016.) There can be no repeated terms in any column, row, or diagonal, by construction, so we must just show that there are no missing terms. Consider column $c \geq 0$. Since the Sprague-Grundy values are calculated moving upwards along the antidiagonals, a number k will appear in column c unless it is blocked by the presence of a k in an earlier column. But the first c columns contain at most c copies of k , so eventually every k will appear in column c .

Consider row $r \geq 0$, and suppose a number k never appears. There are at most r copies of k in the earlier rows, and these can affect only a bounded portion of row r . Consider a square (r, n) , where $n \geq 0$ is large. If k is not to appear in that cell, there must be a copy of k in the antidiagonal to the South-West. So in the right triangle bounded by row r , column 0, and the antidiagonal through (r, n) , there must be at least $n + 1 - r$ copies of k (allowing for the $\leq r$ copies of k in the first r rows). Imagine these k 's replaced by chess queens. By construction they are mutually non-attacking. But it is known ([17, Problem 252], or [A274616](#)) that on a right triangular half-chessboard of side n , there can be at most $2n/3 + 1$ mutually non-attacking queens. Since $2n/3 + 1 < n + 1 - r$ for large n , a k must eventually appear in that row. \square

As to the diagonals, although they *appear* to be permutations, there is no proof.

Conjecture 24. *Every diagonal of the table of Sprague-Grundy values is a permutation of \mathbb{N} .*

The argument using non-attacking queens breaks down here because the diagonal of the half-chessboard contains only half as many squares as the sides. (The antidiagonals are certainly not permutations of \mathbb{N} , since they have finite length.)

We return to the discussion of the positions of the exiled queens. We observed in 2016 that the first 50000 queens appear to lie almost exactly on two straight lines, of slopes ϕ and $1/\phi$, where ϕ is the golden ratio. Figure 6 shows a plot of the first 10000 queens. This is similar to what we saw in Fig. 3, only now we do not have a proof that the points lie on these lines, nor do we have a proof that the slopes are what they appear to be.

Donald Knuth (see Exercise 7.2.2.1–38 in [11]) investigated this question in 2018, although his version of the problem is superficially different. In his problem, the queens are placed column-by-column. A queen is placed in column $c \geq 0$ in the first square




Figure 6: Positions of the first 10000 queens on the single-quadrant board. The points appear to lie essentially on two straight lines. The tick-marks on the horizontal axis (the column indices) are at 0, 2000, 4000, ..., 10000, and on the vertical axis (the row indices) at 0, 5000, 10000, 15000.

from which it cannot attack any earlier queen. However, it is easy to prove that it makes no difference whether the board is scanned by successive antidiagonals or by successive columns. The same sequence $\{S_c\}$ is obtained in both cases.

Knuth has written an efficient program for computing this sequence, and finds that the points are extremely close to the two lines: for $c < 10^9$, he finds that

$$-2 < S_c - c\phi < 1, \quad \text{if } S_c > c, \quad (49)$$

$$-3 < S_c - \frac{c}{\phi} < 5, \quad \text{if } S_c < c. \quad (50)$$

The evidence for the next conjecture is therefore overwhelming:

Conjecture 25. There are constants ϵ_1, ϵ_2 such that

$$|S_c - c\phi| < \epsilon_1, \quad \text{if } S_c > c, \quad (51)$$

$$\left| S_c - \frac{c}{\phi} \right| < \epsilon_2, \quad \text{if } S_c < c. \quad (52)$$

This combinatorial game can be described in a different way, which suggests a possible attack on the conjecture. If the queen being moved is located at square (x, y) , we represent its position by two piles of tokens, of sizes x and y . The legal moves are to remove any positive number of tokens from one pile, or to remove an equal positive number of tokens from both piles, or to move a positive number of tokens from the y pile to the x pile (the latter move corresponds to moving the queen down the antidiagonal). The first person unable to move loses.

A simpler two-pile game is the classic Wythoff's Nim (also called Wyt Queens) [3, 18], which has the same moves except that moving tokens from one pile to another is not

allowed. The array of Sprague-Grundy values can be found in [3, First edition, Chap. 3, Table 3], [5], and [A004481](#). The sequence $\{W_c\}$ specifying which rows the queens are in (the analog of $\{S_c\}$) is [A002251](#):

$$0, 2, 1, 5, 7, 3, 10, 4, 13, 15, 6, 18, 20, 8, 23, 9, 26, 28, 11, 31, 12, 34, \dots \quad (53)$$

Wythoff [18] himself showed in 1907 that these points lie on two lines ([A000201](#), [A001950](#)):

$$W_c = \lfloor c\phi \rfloor, \quad \text{if } W_c > c, \quad (54)$$

$$W_c = \left\lfloor \frac{c}{\phi} \right\rfloor, \quad \text{if } W_c < c. \quad (55)$$

The reason we mention this is that Larsson and Wästlund [12] were able to analyze the two lines of queens in Maharaja Nim, a variant of Wythoff's Nim. Can their method be adapted to our problem?

Although the array of Sprague-Grundy values for our problem looks irregular, it appears that the columns eventually become quasi-periodic. Column 1 is 2, 3, 0, 1, 6, 7, 4, ... ([A004482](#)), which is the Nim-sum $r \oplus 2$, and has generating function

$$\frac{2 - x - 2x^2 + 3x^3}{(1 - x)^2(1 + x^2)} = \frac{(1 + x)(2 - x - 2x^2 + 3x^3)}{(1 - x)(1 - x^4)}. \quad (56)$$

All subsequent columns appear to have a generating function with denominator $(1 - x)(1 - x^{16})$. Column 2, for example, appears to have generating function $g(x)/((1 - x)(1 - x^{16}))$, where $g(x)$ is

$$1 + 3x + x^2 - 3x^3 - 2x^4 + 8x^5 - 5x^6 + 3x^7 + x^8 + 5x^9 + \\ + x^{10} - 3x^{11} + x^{12} - 2x^{13} + 8x^{14} - 3x^{15} + 2x^{17}.$$

Conjecture 26. Every column of the array of Sprague-Grundy values for the single-quadrant problem has a generating function with denominator $(1 - x)(1 - x^{16})$.

(The conjecture also holds for columns 0 and 1.) If true, this would mean that every column eventually becomes quasi-periodic with period 16, something that we find surprising. No such property seems to hold for the rows of the table. The quasi-periodicity of the Sprague-Grundy values for Wythoff's Nim and certain other combinatorial games was studied by Dress, Flammenkamp, and Pink in [5]. However, it does not seem that Conjecture 26 follows from their work.

Acknowledgments

Alois Heinz and Rémy Sigrist were always ready to design efficient computer programs for these sequences, to generate extensive tables, and to provide spectacular illustrations (such as Fig. 5). Many of their programs and tables can be found in the entries in [15].

We thank Alois Heinz for providing Fig. 3, and Jessica Gonzalez for drawing Figures 2 and 4. We also thank Donald E. Knuth for telling us about his investigations of the single-quadrant problem. Achim Flammenkamp kindly provided a copy of [5].

References

- [1] J.-P. Allouche and J. Shallit, *Automatic sequences*. Cambridge University Press, 2003.
- [2] E. Barucci, L. Belanger, and S. Brlek, On Tribonacci sequences. *Fibonacci Quarterly*, 42.4:314–320, 2004.
- [3] E. R. Berlekamp, J. H. Conway, and R. K. Guy, *Winning ways for your mathematical plays*, 2nd ed., 4 vols. A. K. Peters, Boston, 2004.
- [4] L. Carlitz, R. Scoville, and V. E. Hoggatt, Jr., Fibonacci representations of higher order. *Fibonacci Quarterly*, 10.1:43–69, 1972.
- [5] A. Dress, A. Flammenkamp, and N. Pink, Additive periodicity of the Sprague-Grundy function of certain Nim games. *Adv. Appl. Math.*, 22:249–270, 1999.
- [6] C. F. Du, H. Mousavi, L. Schaeffer, and J. Shallit, Decision algorithms for Fibonacci-automatic words, III: Enumeration and abelian properties. *International Journal of Foundations of Computer Science*, 27.08:943–963, 2016.
- [7] E. Duchêne, A. S. Fraenkel, V. Gurvich, N. B. Ho, C. Kimberling, and U. Larsson, Wythoff visions. In U. Larsson, ed., *Games of no chance*, Vol. 5, pp. 101–153. Cambridge University Press, 2017.
- [8] E. Duchêne and M. Rigo, A morphic approach to combinatorial games: the Tribonacci case. *RAIRO—Theoretical Informatics and Applications*, 42.2:375–383, 2008.
- [9] R. Fokkink and D. Rust, A modification of Wythoff’s Nim. [arXiv:1904.08339v1](https://arxiv.org/abs/1904.08339v1).
- [10] R. K. Guy, ed., *Combinatorial games*. American Mathematical Society, Proceedings of Symposia in Applied Mathematics, Vol. 43, 1991.
- [11] D. E. Knuth, *The art of computer programming*. Addison-Wesley, Boston, Vol. 4B, Fascicle 5c, In preparation, 2019. (See <http://www-cs-faculty.stanford.edu/~knuth/fasc5c.ps.gz>.)
- [12] U. Larsson and J. Wästlund, Maharaja Nim: Wythoff’s queen meets the knight. *Integers: Electronic Journal of Combinatorial Number Theory*, 14#G05, 2014.
- [13] M. Lothaire, *Combinatorics on words*. Cambridge University Press, Encyclopedia of mathematics and its applications, Vol. 17, 1983.
- [14] H. Mousavi and J. Shallit, Mechanical proofs of properties of the Tribonacci word. In F. Manea and D. Nowotka, eds., *Combinatorics on Words: WORDS 2015. Lecture Notes in Computer Science*, Vol. 9304. Springer, 2015, pp. 170–190.
- [15] The OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <https://oeis.org>.
- [16] B. Tan and Z.-Y. Wen, Some properties of the Tribonacci sequence. *European Journal of Combinatorics* 28.6:1703–1719, 2007.
- [17] P. Vanderlind, R. K. Guy, and L. C. Larson, *The inquisitive problem solver*. The Mathematical Association of America, 2002.
- [18] W. A. Wythoff, A modification of the game of Nim. *Nieuw Arch. Wisk.*, 7:199–202, 1907.