

Extremal trees with fixed degree sequence

Eric O. D. Andriantiana*

Department of Mathematics (Pure and Applied)
Rhodes University
PO Box 94, 6140 Grahamstown, South Africa
`E.Andriantiana@ru.ac.za`

Valisoa Razanajatovo Misanantenaina

Department of Mathematical Sciences
Stellenbosch University
Private Bag X1, Matieland 7602, South Africa
`valisoa@aims.ac.za`

Stephan Wagner†

Department of Mathematics
Uppsala Universitet
Box 480, 751 06 Uppsala, Sweden
`stephan.wagner@math.uu.se`

and

Department of Mathematical Sciences
Stellenbosch University
Private Bag X1, Matieland 7602, South Africa
`swagner@sun.ac.za`

Submitted: Aug 7, 2020; Accepted: Dec 15, 2020; Published: Jan 15, 2021

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

The greedy tree $\mathcal{G}(D)$ and the \mathcal{M} -tree $\mathcal{M}(D)$ are known to be extremal among trees with degree sequence D with respect to various graph invariants. This paper provides a general theorem that covers a large family of invariants for which $\mathcal{G}(D)$ or $\mathcal{M}(D)$ is extremal. Many known results, for example on the Wiener index, the number of subtrees, the number of independent subsets and the number of

*Supported by the National Research Foundation of South Africa, grant 96310.

†Supported by the National Research Foundation of South Africa, grant 96236, and the Knut and Alice Wallenberg Foundation.

matchings follow as corollaries, as do some new results on invariants such as the number of rooted spanning forests, the incidence energy and the solvability. We also extend our results on trees with fixed degree sequence D to the set of trees whose degree sequence is majorised by a given sequence D , which also has a number of applications.

Mathematics Subject Classifications: 05C05, 05C07, 05C09, 05C35, 05C92

1 Introduction

In the context of chemical graph theory, (molecular) graphs are used to model molecules: the vertices of the graph represent the atoms of the molecule, and the edges correspond to the chemical bonds between atoms. Thus vertex degrees amount to valencies of atoms. If the entries of $D = (d_1, d_2, \dots, d_n)$ are possible valencies of atoms, then the set of all graphs with degree sequence D contains the molecular graphs of all possible isomers of a certain molecule. This is one of the motivations to study graphs with a given degree sequence. In this paper, we will be particularly concerned with trees whose degree sequence is given.

The *greedy tree* $\mathcal{G}(D)$ with degree sequence D , formally defined in the next section, is a tree that can be constructed by starting with the largest degree vertex and always assigning the largest available degree to a neighbour of the vertex with the largest degree whose neighbour degrees are not yet fully specified. On the other hand, large degrees and small degrees alternate in the \mathcal{M} -tree $\mathcal{M}(D)$ with degree sequence D , see Definition 27.

The greedy tree $\mathcal{G}(D)$ and the \mathcal{M} -tree $\mathcal{M}(D)$ are known to be extremal for a number of graph invariants. Let us list a few examples.

- The sum $W(G)$ of the distances $d(u, v)$ between all (unordered) pairs of vertices $\{u, v\}$, better known as the Wiener index, is among the most popular graph invariants. It was shown by Wang in [33] and by Zhang et al. in [41] that, among all trees with degree sequence D , $\mathcal{G}(D)$ has the minimum Wiener index.
- Schmuck et al. [29] showed a more general statement: if $W(G)$ is replaced by a Wiener-like invariant

$$W_f(G) = \sum_{\{u,v\} \subseteq V(G)} f(d(u,v))$$

for a nonnegative and nondecreasing function f , then $\mathcal{G}(D)$ still attains the minimum value among trees with degree sequence D . Likewise, if f is a nonnegative and nonincreasing function, then $\mathcal{G}(D)$ attains the maximum value. In particular, this includes a well-studied invariant known as the Harary index, which corresponds to $f(x) = 1/x$ (see [32]).

- The greedy tree $\mathcal{G}(D)$ is extremal with respect to several other invariants including:
 - the number of subtrees (and the number of antichains when rooted trees are considered) [3, 42],
 - the spectral radius [5] and the Laplacian spectral radius [40],

– the spectral moments [2, 24]. See also [7] for a generalisation.

- The total number of independent subsets is also known as the Merrifield-Simmons index of a graph. It is proven in [1] that the Merrifield-Simmons index is maximised by $\mathcal{M}(D)$. The total number of matchings is also called Hosoya index. It is minimised by $\mathcal{M}(D)$ (see [1]).
- The energy of a graph G is defined as $\text{En}(G) = \sum_i |\lambda_i|$, where the λ_i s are the eigenvalues of the adjacency matrix of G . For trees, the energy is connected to matchings by the well-known Coulson formula [14]: let $m(G, k)$ be the number of matchings of cardinality k in a graph G . For every tree T ,

$$\text{En}(T) = \frac{2}{\pi} \int_0^\infty \frac{dx}{x^2} \log \sum_{k \geq 0} m(T, k) x^{2k}. \quad (1)$$

For a positive real number x , define

$$M(G, x) = \sum_{k \geq 0} m(G, k) x^k.$$

It is shown in [1] that, among all trees with degree sequence D , $\mathcal{M}(D)$ is the unique tree with minimum $M(\cdot, x)$ for any $x > 0$. Thus, it also has the minimum energy.

The long list of graph invariants for which $\mathcal{G}(D)$ or $\mathcal{M}(D)$ is extremal leads us to the following natural question: what condition(s) does a graph invariant need to satisfy for $\mathcal{G}(D)$ or $\mathcal{M}(D)$ to be extremal among trees with degree sequence D ? In this paper, we provide sufficient conditions that cover a number of examples, both old and new. The idea is inspired by an exchange lemma used to study bounded degree trees in [17], and trees with given degree sequence in [1].

A sequence $B = (b_1, b_2, \dots, b_n)$ is said to majorise D if $b_1 + \dots + b_n = d_1 + \dots + d_n$ and $d_1 + \dots + d_k \leq b_1 + \dots + b_k$ for every k with $1 \leq k < n$. We then write $B \triangleright D$. In Section 4, we will consider the set of all trees whose degree sequence is majorised by D . Several natural sets of trees satisfy majorisation properties. This includes the set of all n -vertex trees, n -vertex trees with bounded degrees, and trees of order n with fixed number of leaves. There are several instances of graph invariants for which the extremal trees with degree sequence D are also extremal among those whose degree sequence is majorised by D . We will see many such examples at the end of this paper.

We need to introduce some notation to describe our results. Let T be a rooted tree. The root of T is denoted by $r(T)$. We write $T = [T_1, T_2, \dots, T_k]$ if the rooted trees T_1, T_2, \dots, T_k are the connected components of $T - r(T)$, where the roots of T_1, T_2, \dots, T_k are the neighbours of $r(T)$.

Let v and w be two different leaves of a tree H . The tree obtained by merging the root of $[L_1, L_2, \dots, L_k]$ with v and the root of $[R_1, R_2, \dots, R_\ell]$ with w , as seen in Figure 1, is denoted by $[L_1, L_2, \dots, L_k]vHw[R_1, R_2, \dots, R_\ell]$.




Figure 1: The tree $[L_1, L_2, \dots, L_k]vHw[R_1, R_2, \dots, R_\ell]$.

Let $\rho(T)$ be a quantity associated to a rooted tree T , which satisfies a recursive relation of the form

$$\rho([T_1, T_2, \dots, T_k]) = f_\rho(\rho(T_1), \rho(T_2), \dots, \rho(T_k)) \quad (2)$$

for a symmetric function f_ρ , i.e., the value of f_ρ is invariant under permutation of the branches. We call f_ρ the *recurrence rule* for ρ . In our examples, the recurrence rule can always be expressed as a function of sums or products over all branches.

Definition 1. We say that a tree T is ρ -*exchange-extremal* if, whenever we can decompose T as

$$T = [L_1, L_2, \dots, L_k]vHw[R_1, R_2, \dots, R_\ell]$$

for some H , then we have $k \geq \ell$ and

$$\min\{\rho(L_1), \rho(L_2), \dots, \rho(L_k)\} \geq \max\{\rho(R_1), \rho(R_2), \dots, \rho(R_\ell)\}$$

or $k \leq \ell$ and

$$\max\{\rho(L_1), \rho(L_2), \dots, \rho(L_k)\} \leq \min\{\rho(R_1), \rho(R_2), \dots, \rho(R_\ell)\}.$$

We will prove the following theorem:

Theorem 2. *Let ρ be an invariant of rooted trees that satisfies the recurrence relation (2). Suppose that a tree T with degree sequence D is ρ -exchange-extremal.*

- *If f_ρ is increasing (in each variable and under addition of further variables) and the minimum of ρ is attained by the single-vertex tree, then T has to be $\mathcal{G}(D)$.*
- *If f_ρ is decreasing (in each variable and under addition of further variables) and the maximum of ρ is attained by the single-vertex tree, then T has to be $\mathcal{M}(D)$.*

It turns out that extremality with respect to many different invariants implies ρ -exchange-extremality for a suitable choice of ρ . This includes some of the previously mentioned instances (Wiener index, number of subtrees, Merrifield-Simmons index, Hosoya index, etc.), but we will also discuss examples that have not been considered before.

The rest of this paper is organised as follows. Section 2 treats the case of increasing f_ρ , where the greedy tree $\mathcal{G}(D)$ is extremal. The case of decreasing f_ρ is considered in Section 3, it leads to the situation where the \mathcal{M} -tree $\mathcal{M}(D)$ is extremal. Various special cases are listed at the end of each section. This includes many known results, but we also obtain several new results, specifically for the number of rooted spanning forests (related to the coefficients of the Laplacian characteristic polynomial), the incidence energy, and an invariant called the solvability. We also settle an open question on the Steiner Wiener index. Furthermore, in Section 4, we compare trees with different degree sequences and prove extremality results for the set of all trees whose degree sequence is majorised by a fixed sequence D .

The following technical terms will be needed. The *height* $h(T)$ of a rooted tree T is the greatest distance of a vertex to $r(T)$. A subgraph B of a tree T is called a *complete branch* of T if there is an edge e such that B is one of the components of $T - e$, see Figure 2. We denote by $\text{rd}(B)$ the degree of the root $r(B)$ (the end of e that belongs to B) as a vertex of B .




Figure 2: Complete branches B and $T - B$ of T .

2 Increasing recurrence rule f_ρ

Definition 3. [29] Given a degree sequence of a tree D , the greedy tree, denoted $\mathcal{G}(D)$, is constructed by the following “greedy algorithm”:

- (i) Label the vertex with the largest degree v (the root).
- (ii) Label the neighbours of v as $v_1, v_2, \dots, v_{d(v)}$, and assign the largest degrees available to them so that $d(v_1) \geq d(v_2) \geq \dots \geq d(v_{d(v)})$.
- (iii) Label the neighbours of v_1 (except v) as $v_{1,1}, v_{1,2}, \dots, v_{1,d(v_1)-1}$, and assign the largest degrees available to them so that $d(v_{1,1}) \geq d(v_{1,2}) \geq \dots \geq d(v_{1,d(v_1)-1})$. Then do the same for $v_2, v_3, \dots, v_{d(v)}$.
- (iv) Repeat (ii) and (iii) for all the newly labelled vertices. Always start with the neighbours of the labelled vertex with the largest degree whose neighbours are not labelled yet.

Figure 3 shows the greedy tree $\mathcal{G}(D)$ for $D = (4, 4, 3, 3, 3, 3, 3, 3, 2, 2, 2, 1, 1, \dots, 1)$.




Figure 3: The greedy tree $\mathcal{G}(4, 4, 3, 3, 3, 3, 3, 3, 2, 2, 2, 1, 1, \dots, 1)$.

We will see in this section that the greedy tree is the unique ρ -exchange-extremal tree if ρ satisfies the following conditions:

- I.1** the recurrence relation (2) holds, for some symmetric recurrence rule f_ρ ,
- I.2** the function f_ρ is strictly increasing (strictly increasing in each single variable and strictly increasing under addition of further variables),
- I.3** $\rho(\bullet) < \rho(B)$, for all rooted trees B with $|V(B)| > 1$, where \bullet denotes a single vertex tree.

2.1 Special case: the Wiener index

Let us first consider the special case where $\rho(T)$ is the number of vertices of T , i.e., $\rho(T) = \rho_0(T) = |V(T)|$. We have

$$\rho_0([T_1, \dots, T_k]) = f_{\rho_0}(\rho_0(T_1), \dots, \rho_0(T_k)) = 1 + \sum_{i=1}^k \rho_0(T_i).$$

The recurrence rule f_{ρ_0} is indeed symmetric and increasing with respect to each of its variables and under addition of further variables. Clearly $\rho_0(\bullet) = 1$ is the minimum among all rooted trees. So conditions **I.1** to **I.3** are all satisfied.

Now recall that the Wiener index is the sum of all distances between pairs of vertices:

$$W(T) = \sum_{u,v \in V(T)} d(u, v).$$

An important alternative formula for the Wiener index for trees is (see [9])

$$W(T) = \sum_{uv \in E(T)} \rho_0(T_u) \rho_0(T_v), \tag{3}$$

where T_u and T_v are the components of $T - uv$ containing u and v respectively. We use this formula to show that minimality with respect to the Wiener index implies ρ_0 -exchange-extremality.

Lemma 4 (cf. [31]). *Let T be a tree with degree sequence D for which $W(T)$ attains its minimum. Then, for any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_0(A_1), \dots, \rho_0(A_k)\} \geq \max\{\rho_0(B_1), \dots, \rho_0(B_\ell)\}$,
- or $k \leq \ell$ and $\max\{\rho_0(A_1), \dots, \rho_0(A_k)\} \leq \min\{\rho_0(B_1), \dots, \rho_0(B_\ell)\}$.




Figure 4: Decomposition of T in the proof of Lemma 4.

Proof. Let $P = r(A)u_1 \dots u_t r(B)$ be the unique path between $r(A)$ and $r(B)$. To simplify notation, we put $\alpha = \rho_0(A) = 1 + \sum_{i=1}^k \rho_0(A_i)$ and $\beta = \rho_0(B) = 1 + \sum_{i=1}^\ell \rho_0(B_i)$.

For $1 \leq j \leq t$, let U_j be the component containing u_j when we remove all the edges of the path P , and set $z_j = \rho_0(U_j)$, $p_j = z_1 + \dots + z_j$, $q_j = z_t + \dots + z_{t-j+1}$ and $p_0 = q_0 = 0$.

Using the formula for the Wiener index in (3), we have:

$$W(T) = \sum_{j=0}^t (\alpha + p_j)(q_{t-j} + \beta) + C_T,$$

where C_T is the contribution of edges in U_1, \dots, U_t, A and B .

With $q = q_1 + \dots + q_t$ and $p = p_1 + \dots + p_t$, this can be rewritten as

$$W(T) = \alpha q + \beta p + \alpha \beta (t + 1) + \sum_{j=0}^t p_j q_{t-j} + C_T.$$

The last two terms $\sum_{j=0}^t p_j q_{t-j} + C_T$ are invariant under any rearrangements of the A_i s and B_i s. Note that the degree sequence remains the same if all the A_i s are switched with all the B_i s (interchanging the degrees of $r(A)$ and $r(B)$), if $A_1, A_2, \dots, A_k, B_1, B_2, \dots, B_\ell$ are permuted in an arbitrary way, or both. Hence, if $W(T)$ is minimal, then the expression

$\alpha q + \beta p + \alpha\beta(t + 1)$ must have its minimum value under these operations. Note that the sum $\Sigma = \alpha + \beta$ remains constant under any of these rearrangements, and we have

$$\alpha q + \beta p + \alpha\beta(t + 1) = \alpha q + (\Sigma - \alpha)p + \alpha(\Sigma - \alpha)(t + 1),$$

which is a strictly concave function of α . Thus the minimum is attained if α is either as small or as large as possible. For this we must have

$$k \leq \ell \quad \text{and} \quad \max\{\rho_0(A_i) : 1 \leq i \leq k\} \leq \min\{\rho_0(B_i) : 1 \leq i \leq \ell\}$$

or

$$k \geq \ell \quad \text{and} \quad \min\{\rho_0(A_i) : 1 \leq i \leq k\} \geq \max\{\rho_0(B_i) : 1 \leq i \leq \ell\}.$$

In words, the largest possible branches (and as many of them as possible) have to be gathered in one place. This is precisely the statement of the lemma. \square

We know now that the tree which minimises the Wiener index is ρ_0 -exchange-extremal. Moreover, ρ_0 -exchange-extremality is equivalent to the semi-regular property defined in Definition 4 of [29], where it is proved that such a tree is greedy. Thus we have the following observation, which will be the basis for the proof of the main theorem of this section.

Theorem 5 (cf. [29, 31]). *If T is a ρ_0 -exchange-extremal tree, then T is a greedy tree.*

2.2 Main result

We require one more lemma before stating the main result. This lemma establishes a connection between ρ_0 and other invariants ρ .

Lemma 6. *Let ρ be an invariant of rooted trees that satisfies **I.1**, **I.2** and **I.3**, and let T be a ρ -exchange-extremal tree. For any two disjoint complete branches A and B in T , $\rho(A) \geq \rho(B)$ if and only if $\rho_0(A) \geq \rho_0(B)$. In particular, $\rho(A) = \rho(B)$ if and only if $\rho_0(A) = \rho_0(B)$.*

Proof. We reason by induction on the heights of A and B , specifically $\max\{h(A), h(B)\}$. If $\max\{h(A), h(B)\} = 0$, then $h(A) = h(B) = 0$, so A and B are both single vertices and the claim holds trivially. Assume that the statement holds whenever $\max\{h(A), h(B)\} \leq t$, for some $t \geq 0$. Now, consider two complete branches A and B in T such that $\max\{h(A), h(B)\} = t + 1$ and $\rho(A) \geq \rho(B)$. If A only has a single vertex, then so does B by condition **I.3**. If B only has a single vertex, then $\rho_0(A) \geq \rho_0(B)$ is trivially satisfied. In both cases, there is nothing left to show. So we can now assume that A and B have more than one vertex and can thus be decomposed. Let $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ for some $k, \ell \geq 1$. Since T is a ρ -exchange-extremal tree, we are left with two possibilities. If $k \geq \ell$, and $\min\{\rho(A_i) : 1 \leq i \leq k\} \geq \max\{\rho(B_i) : 1 \leq i \leq \ell\}$, then by the induction hypothesis, we have $\min\{\rho_0(A_i) : 1 \leq i \leq k\} \geq \max\{\rho_0(B_i) : 1 \leq i \leq \ell\}$. Thus $\rho_0(A) \geq \rho_0(B)$, because f_{ρ_0} is increasing.

On the other hand, if $k \leq \ell$ and $\max\{\rho(A_i) : 1 \leq i \leq k\} \leq \min\{\rho(B_i) : 1 \leq i \leq \ell\}$, then $\rho(A) \leq \rho(B)$ since f_ρ is increasing, so $\rho(A) = \rho(B)$. For this to hold, we must have $k = \ell$ and $\rho(A_1) = \dots = \rho(A_k) = \rho(B_1) = \dots = \rho(B_\ell)$. By the induction hypothesis, this implies $\rho_0(A_1) = \dots = \rho_0(A_k) = \rho_0(B_1) = \dots = \rho_0(B_\ell)$, so $\rho_0(A) = \rho_0(B)$.

Thus we have shown that $\rho(A) \geq \rho(B)$ implies $\rho_0(A) \geq \rho_0(B)$, and the proof of the converse is analogous. \square

Now we are ready for the main theorem.

Theorem 7. *Let ρ be an invariant of rooted trees that satisfies **I.1**, **I.2** and **I.3**. If a tree T is ρ -exchange-extremal, then T is a greedy tree.*

Proof. By Lemma 6, the tree T is also ρ_0 -exchange-extremal. Theorem 5 gives us the desired result. \square

In the remainder of this section, we describe different applications of Theorem 7.

2.3 The number of subtrees

As a first example, we consider the number of subtrees (here, a subtree is any induced subgraph that is connected and thus again a tree). We will be able to show that there is a suitable function ρ such that a tree with the greatest possible number of subtrees among all trees with the same degree sequence is necessarily ρ -exchange-extremal. To this end, we need a technical lemma, which will also be useful later.

Lemma 8. *Let $x_1 \geq \dots \geq x_n \geq 1$ and $a \geq b$ be positive real numbers, $n \geq k \geq n/2$ and \mathbb{S}_n the set of permutations of $1, \dots, n$. Then*

$$a \prod_{i=1}^k x_i + b \prod_{i=k+1}^n x_i = \max \left\{ a \prod_{i=1}^{\ell} x_{\sigma(i)} + b \prod_{i=\ell+1}^n x_{\sigma(i)} : \sigma \in \mathbb{S}_n \text{ and } \ell \in \{n-k, k\} \right\}.$$

Moreover, we have uniqueness: if

$$a \prod_{i=1}^k x_i + b \prod_{i=k+1}^n x_i = a \prod_{i=1}^{\ell} x_{\sigma(i)} + b \prod_{i=\ell+1}^n x_{\sigma(i)},$$

$a > b$ and the x_i s are not all equal to 1, then $\ell = k$ and $x_{\sigma(1)}, \dots, x_{\sigma(k)}$ are a permutation of x_1, \dots, x_k . If $a = b$, then either the same holds, or $\ell = n - k$ and $x_{\sigma(1)}, \dots, x_{\sigma(n-k)}$ are a permutation of x_{k+1}, \dots, x_n .

In words, Lemma 8 says that the maximum value is reached when we assemble the largest elements together in the same product.

Proof. Set $g(x) = ax + bx^{-1} \prod_{i=1}^n x_i$, and note that

$$a \prod_{i=1}^{\ell} x_{\sigma(i)} + b \prod_{i=\ell+1}^n x_{\sigma(i)} = g\left(\prod_{i=1}^{\ell} x_{\sigma(i)}\right).$$

Since $g''(x) = 2bx^{-3} \prod_{i=1}^n x_i > 0$, g is strictly convex, so it can only attain its maximum value when $x = \prod_{i=1}^{i=\ell} x_{\sigma(i)}$ is either largest or smallest. The largest possible value of x is clearly $L = \prod_{i=1}^k x_i$, the smallest $S = \prod_{i=k+1}^n x_i$. Comparing the two possibilities, we see that

$$g(L) - g(S) = a \prod_{i=1}^k x_i + b \prod_{i=k+1}^n x_i - a \prod_{i=k+1}^n x_i - b \prod_{i=1}^k x_i = (a - b) \left(\prod_{i=1}^k x_i - \prod_{i=k+1}^n x_i \right) \geq 0$$

since $a \geq b$ and $k \geq \frac{n}{2}$. We have strict inequality if $a > b$ unless $k = \frac{n}{2}$ and all x_i s are equal (in which case $x_{\sigma(1)}, \dots, x_{\sigma(k)}$ are still a permutation of x_1, \dots, x_k) or all x_i s are equal to 1. The statement of the lemma follows. \square

The number of subtrees of a tree T is denoted by $\eta(T)$, and the number of subtrees of T containing the vertices v_1, \dots, v_ℓ is denoted by $\eta(T, v_1, \dots, v_\ell)$. In the following lemma, we establish the necessary property to apply our main theorem.

Lemma 9. *Let T be a tree with degree sequence D for which $\eta(T)$ attains its maximum. For any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and

$$\min\{\eta(A_1, r(A_1)), \dots, \eta(A_k, r(A_k))\} \geq \max\{\eta(B_1, r(B_1)), \dots, \eta(B_\ell, r(B_\ell))\},$$

- or $k \leq \ell$ and

$$\max\{\eta(A_1, r(A_1)), \dots, \eta(A_k, r(A_k))\} \leq \min\{\eta(B_1, r(B_1)), \dots, \eta(B_\ell, r(B_\ell))\}.$$

In other words, T is ρ -exchange-extremal with $\rho(T) = \eta(T, r(T))$.

Proof. Consider a decomposition of T as $[A_1, \dots, A_k]vHw[B_1, \dots, B_\ell]$, for some tree H . Each subtree of T has to belong to one of the following types:

- Subtrees which contain neither v nor w . Each subtree of this type is either in $H - v - w$ or in one of the A_i s or B_i s. Thus, there are $\eta(H - v - w) + \sum_{i=1}^k \eta(A_i) + \sum_{i=1}^\ell \eta(B_i)$ of them.
- Subtrees which contain w , but not v . A subtree of this type consists of a part in $H - v$ and another part in $[B_1, \dots, B_\ell]$, each containing w . Their number is

$$\eta(H - v, w) \prod_{i=1}^\ell (1 + \eta(B_i, r(B_i))).$$

- Subtrees which contain v , but not w . By similar reasoning as in the second case, the number of these subtrees is

$$\eta(H - w, v) \prod_{i=1}^k (1 + \eta(A_i, r(A_i))).$$

- Subtrees which contain both v and w . This corresponds to subtrees of H which contain both v and w , to which induced subtrees of the A_i s and B_i s containing the respective roots can be attached. Thus, there are

$$\eta(H, v, w) \prod_{i=1}^{\ell} (1 + \eta(B_i, r(B_i))) \prod_{i=1}^k (1 + \eta(A_i, r(A_i)))$$

of them.

In total we have

$$\eta(T) = \alpha_T + \beta_T,$$

where

$$\begin{aligned} \alpha_T &= \eta(H - v, w) \prod_{i=1}^{\ell} (1 + \eta(B_i, r(B_i))) + \eta(H - w, v) \prod_{i=1}^k (1 + \eta(A_i, r(A_i))), \\ \beta_T &= \eta(H - v - w) + \sum_{i=1}^k \eta(A_i) + \sum_{i=1}^{\ell} \eta(B_i) \\ &\quad + \eta(H, v, w) \prod_{i=1}^{\ell} (1 + \eta(B_i, r(B_i))) \prod_{i=1}^k (1 + \eta(A_i, r(A_i))). \end{aligned}$$

As in the proof of Lemma 4, this must be maximal under all permutations of the A_i s and B_i s, possibly also switching the degrees of v and w . Note here that β_T is not affected by any of these rearrangements. So the maximality of $\eta(T)$ depends only on α_T .

Since all the quantities involved are positive and the factors $1 + \eta(A_i, r(A_i))$ and $1 + \eta(B_i, r(B_i))$ are even greater than 1, Lemma 8 applies. Thus we obtain the maximum of α_T if either $\eta(H - w, v) \geq \eta(H - v, w)$, $k \geq \ell$ and

$$\min\{\eta(A_1, r(A_1)), \dots, \eta(A_k, r(A_k))\} \geq \max\{\eta(B_1, r(B_1)), \dots, \eta(B_\ell, r(B_\ell))\},$$

or $\eta(H - w, v) \leq \eta(H - v, w)$, $k \leq \ell$ and

$$\max\{\eta(A_1, r(A_1)), \dots, \eta(A_k, r(A_k))\} \leq \min\{\eta(B_1, r(B_1)), \dots, \eta(B_\ell, r(B_\ell))\}. \quad \square$$

So we have established ρ -exchange-extremality with respect to $\rho(T) = \eta(T, r(T))$, the number of root-containing subtrees. Moreover, if T can be decomposed as $[T_1, \dots, T_k]$, then

$$\eta(T, r(T)) = \prod_{i=1}^k (1 + \eta(T_i, r(T_i))), \quad (4)$$

which is increasing in all of its variables and under addition of further variables, so **I.1** and **I.2** hold. Moreover, the minimum of $\eta(T, r(T))$, which is trivially equal to 1, is only reached if $T = \bullet$. Thus, **I.3** holds as well, and Theorem 7 implies the following theorem, which was already established in [3, 42].

Theorem 10 ([3, 42]). *Given a degree sequence, the corresponding greedy tree is the unique tree that maximises the number of subtrees.*

2.4 Rooted spanning forests, incidence energy, Laplacian-energy-like invariant

Let T be an n -vertex tree and $A(T)$ its adjacency matrix. The *Laplacian matrix* of T is $L(T) = A(T) - D(T)$, where $D(T)$ is the diagonal matrix whose diagonal entry d_{ii} is the degree of the i -th vertex v_i . The *Laplacian characteristic polynomial* $L(T, x)$ is the characteristic polynomial of the Laplacian matrix. In 1967, Kel'mans [22] gave a combinatorial interpretation for the coefficients of the Laplacian characteristic polynomial as follows.

Theorem 11. *If $L(T, x) = \det(xI_n - L(T)) = \sum_{k=1}^n (-1)^{n-k} c_k(T) x^k$, then*

$$c_k(T) = \sum_{F \in \mathcal{F}(T, k)} \gamma(F),$$

where $\mathcal{F}(T, k)$ is the set of all spanning forests of the tree T containing exactly k components and $\gamma(F)$ is the product of the number of vertices in each component of F .

The quantity $\gamma(F)$ can be interpreted as the number of ways to assign roots to the components of a forest F , and therefore $c_k(T)$ is the number of k -rooted spanning forests of T (spanning forests with k components, where each component is rooted at one of its vertices).

Let us consider the polynomial in which we associate to every rooted spanning forest F a weight $x^{\lambda(F)}$, where $\lambda(F)$ is the number of components of F . Replacing x by $-x$ in Theorem 11, we see that this polynomial is connected to the Laplacian matrix as follows:

$$\text{rf}(T, x) = \sum_{k=1}^n c_k(T) x^k = \det(L(T) + xI_n). \quad (5)$$

Note that the total number of rooted spanning forests in a tree T is equal to $\text{rf}(T, 1)$. To avoid confusion as we also consider rooted spanning forests within rooted trees, we will refer to “marked” spanning forests rather than rooted spanning forests in the following, and call the components’ roots “markers”. We show that the invariant $\text{rf}(\cdot, x)$ fits our general scheme. This generalises the approach taken in [20], where trees with given maximum degree are considered.

We define an auxiliary quantity for rooted trees T , which is denoted by $f(T, x)$. It counts marked spanning forests F weighted with $x^{\gamma(F)-1}$, in which the root of T is also a marker of one of the forest’s components. Note that $f(T, x)$ also counts (weighted) spanning forests of T where all components, except the one containing the root of T , have a marker. Finally, we set

$$\rho_1(T, x) = \frac{\text{rf}(T, x)}{\text{rf}(T, x) + f(T, x)}.$$

Lemma 12. *Let $T = [A_1, \dots, A_k]vHw[B_1, \dots, B_\ell]$ for some tree H , and let $x > 0$. Suppose that $\text{rf}(T, x) \leq \text{rf}(T', x)$ for every tree T' with the same degree sequence as T . Then*

- either $k \geq \ell$ and

$$\min\{\rho_1(A_1, x), \dots, \rho_1(A_k, x)\} \geq \max\{\rho_1(B_1, x), \dots, \rho_1(B_\ell, x)\},$$

- or $k \leq \ell$ and

$$\max\{\rho_1(A_1, x), \dots, \rho_1(A_k, x)\} \leq \min\{\rho_1(B_1, x), \dots, \rho_1(B_\ell, x)\}.$$

Proof. Let $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$. To obtain an expression for $\text{rf}(T, x)$, we consider the following cases for a marked spanning forest of T :

- $r(A)$ and $r(B)$ belong to components that have a marker in H (possibly the same component). The parts that form spanning forests of $A_1 \cup \dots \cup A_k \cup B_1 \cup \dots \cup B_\ell$ are either fully marked or marked except for the root's component (which is joined to the component of $r(A)$ or $r(B)$ in H). Let $c_{11}(H, r)$ be the number of r -component marked spanning forests of H , and set $a = \sum_{r \geq 1} c_{11}(H, r)x^r$. The contribution of this case to $\text{rf}(T, x)$ is

$$a \cdot \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x)).$$

- $r(A)$ and $r(B)$ belong to the same component, but this component does not have a marker in H . In this case, the marker of the component that contains $r(A)$ and $r(B)$ lies in one of the A_i s or B_j s. So we have to choose exactly one of them and replace the factor $\text{rf}(\cdot, x) + f(\cdot, x)$ by $\text{rf}(\cdot, x)$. Let $c_{00}(H, r)$ be the number of r -component spanning forests of H , where $r(A)$ and $r(B)$ lie in the same component and all components except the one containing those are marked. Set $b = \sum_{r \geq 1} c_{00}(H, r)x^{r-1}$. This gives a contribution of

$$b \cdot \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x)) \\ \left(\sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} + \sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)} \right).$$

- $r(A)$ and $r(B)$ lie in different components, both vertices have markers outside of H . Now, one of the A_i s has to contain the marker of the component of $r(A)$, and one of the B_j s the marker of the component of $r(B)$. Let $c'_{00}(H, r)$ be the number of r -component spanning forests of H where $r(A)$ and $r(B)$ belong to different components and all but those two components are marked. Set $c = \sum_{r \geq 2} c'_{00}(H, r)x^{r-2}$. For this case we get

$$c \cdot \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x))$$

$$\left(\sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} \right) \left(\sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)} \right).$$

- $r(A)$ and $r(B)$ lie in different components, one has a marker in H , the other does not. Let $c_{10}(H, r)$ be the number of r -component spanning forests of H such that $r(A)$ and $r(B)$ lie in different components and all but the one containing $r(B)$ are marked. Define $c_{01}(H, r)$ analogously, with the roles of $r(A)$ and $r(B)$ reversed. Now set $d_1 = \sum_{r \geq 2} c_{10}(H, r)x^{r-1}$ and $d_2 = \sum_{r \geq 2} c_{01}(H, r)x^{r-1}$. Using a similar reasoning as before, we get a contribution of

$$\begin{aligned} & d_1 \cdot \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x)) \cdot \sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} \\ & + d_2 \cdot \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x)) \cdot \sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)}. \end{aligned}$$

Hence, we finally obtain

$$\begin{aligned} \text{rf}(T, x) &= \prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x)) \\ & \left[a + (b + d_1) \sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} + (b + d_2) \sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)} \right. \\ & \left. + c \left(\sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} \right) \left(\sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)} \right) \right]. \end{aligned}$$

The product $\prod_{i=1}^k (\text{rf}(A_i, x) + f(A_i, x)) \prod_{j=1}^{\ell} (\text{rf}(B_j, x) + f(B_j, x))$ remains constant when the A_i s and B_j s are rearranged, as does the sum

$$\begin{aligned} \Sigma &= \sum_{i=1}^k \frac{\text{rf}(A_i, x)}{\text{rf}(A_i, x) + f(A_i, x)} + \sum_{j=1}^{\ell} \frac{\text{rf}(B_j, x)}{\text{rf}(B_j, x) + f(B_j, x)} \\ &= \sum_{i=1}^k \rho_1(A_i, x) + \sum_{j=1}^{\ell} \rho_1(B_j, x). \end{aligned}$$

We can now argue exactly as in the proof of Lemma 4. Write $y = \sum_{i=1}^k \rho_1(A_i, x)$. For $\text{rf}(T, x)$ to attain its minimum under all possible permutations of A_i s and B_i s, the function

$$y \mapsto a + (b + d_1)y + (b + d_2)(\Sigma - y) + cy(\Sigma - y),$$

which is strictly concave in y , has to attain its minimum. This occurs when y is either as large or as small as possible. That is,

- $k \geq \ell$ and $\min\{\rho_1(A_1, x), \dots, \rho_1(A_k, x)\} \geq \max\{\rho_1(B_1, x), \dots, \rho_1(B_\ell, x)\}$, or
- $k \leq \ell$ and $\max\{\rho_1(A_1, x), \dots, \rho_1(A_k, x)\} \leq \min\{\rho_1(B_1, x), \dots, \rho_1(B_\ell, x)\}$. □

We have established now that minimality with respect to $\text{rf}(\cdot, x)$ implies ρ_1 -exchange-extremality. Moreover, the quantity ρ_1 can be determined recursively as follows. If T can be decomposed as $[T_1, \dots, T_k]$, then

$$\text{rf}(T, x) = \prod_{i=1}^k (\text{rf}(T_i, x) + f(T_i, x)) \left(x + \sum_{i=1}^k \frac{\text{rf}(T_i, x)}{\text{rf}(T_i, x) + f(T_i, x)} \right),$$

and

$$f(T, x) = \prod_{i=1}^k (\text{rf}(T_i, x) + f(T_i, x))$$

using similar arguments as before. Thus,

$$\rho_1(T, x) = \frac{x + \sum_{i=1}^k \frac{\text{rf}(T_i, x)}{\text{rf}(T_i, x) + f(T_i, x)}}{1 + x + \sum_{i=1}^k \frac{\text{rf}(T_i, x)}{\text{rf}(T_i, x) + f(T_i, x)}} = \frac{x + \sum_{i=1}^k \rho_1(T_i, x)}{1 + x + \sum_{i=1}^k \rho_1(T_i, x)}. \quad (6)$$

The recurrence rule f_{ρ_1} corresponding to (6) is increasing in all of its variables and under addition of further variables, moreover $\rho_1(\bullet, x) = \frac{x}{1+x}$ is the unique minimum. So, we may use Theorem 7 to obtain the following result.

Theorem 13. *For every tree T with degree sequence D and every $x > 0$,*

$$\text{rf}(T, x) \geq \text{rf}(\mathcal{G}(D), x),$$

with equality if and only if T is isomorphic to $\mathcal{G}(D)$. In particular, $\mathcal{G}(D)$ has the smallest total number of marked spanning forests among trees with degree sequence D .

Now, let us consider other quantities related to the polynomial $\text{rf}(T, x)$.

Definition 14. A *subdivision graph*, denoted $S(G)$, is a graph obtained by inserting a new vertex of degree 2 on each edge of G .

Lemma 15 ([43]). *Let T be a tree of order n and $S(T)$ its corresponding subdivision graph. Then*

$$c_k(T) = m(S(T), k), \quad k = 0, \dots, n,$$

where $m(S(T), k)$ is the number of k -matchings of $S(T)$.

Let $M(T, x) = \sum_{k \geq 0} m(T, k)x^k$ be the matching generating polynomial of T ; then Lemma 15 implies that $\text{rf}(T, x) = M(S(T), x)$. Thus, we obtain the following corollary of Theorem 13:

Corollary 16. *For every tree T with degree sequence D , and for every $x > 0$,*

$$M(S(T), x) \geq M(S(\mathcal{G}(D)), x),$$

with equality if and only if T is isomorphic to $\mathcal{G}(D)$.

For a tree T , let $\mu_1, \mu_2, \dots, \mu_n$ be the eigenvalues of $L(T)$. They are also the eigenvalues of the signless Laplacian matrix $L^+(T)$ (this is in fact true for every bipartite graph, see [8]). Recall that the energy of a graph is the sum of the absolute values of the eigenvalues of its adjacency matrix. A variant of the energy, known as the Laplacian-energy-like invariant (LEL for short, see [27]), is defined by

$$\text{LEL}(T) = \sum_{i=1}^n \sqrt{\mu_i}.$$

It is closely related to the incidence energy IE of a graph, defined in [21] as the sum of the singular values of its (vertex-edge) incidence matrix. For every tree T , one has $\text{LEL}(T) = \text{IE}(T)$. Furthermore, it is known that (see [13])

$$\text{LEL}(T) = \text{IE}(T) = \frac{1}{2} \text{En}(S(T)), \quad (7)$$

where $S(T)$ is the subdivision graph of T . This allows us to prove the following corollary.

Corollary 17. *Given a degree sequence of a tree, the incidence energy IE , or equivalently the Laplacian-energy-like invariant LEL , is minimised by the greedy tree.*

Proof. Let T be a tree and $S(T)$ its subdivision graph. Using the Coulson formula (1) for the energy and the relation (7), we obtain:

$$\text{LEL}(T) = \text{IE}(T) = \frac{2}{\pi} \int_0^\infty \frac{1}{x^2} \ln \left(\sum_k m(S(T), k) x^{2k} \right) dx = \frac{2}{\pi} \int_0^\infty \frac{1}{x^2} \ln M(S(T), x^2) dx.$$

Thus, the claim readily follows from Corollary 16. □

2.5 A common generalisation of Wiener index and terminal Wiener index

In analogy to the Wiener index, the terminal Wiener index [12] is defined as the sum of all distances between pairs of leaves, and the spinal Wiener index [4] is the sum of all distances between pairs of non-leaves. It is known that both are minimised by greedy trees [31, 4]. We consider a common generalisation defined as follows: for two fixed positive numbers a and b , we set

$$W_{a,b}(T) = \sum_{\{v,w\} \subseteq V(T)} \omega(v)\omega(w) d(v,w) = \frac{1}{2} \sum_{v \in V(T)} \sum_{w \in V(T)} \omega(v)\omega(w) d(v,w),$$

where

$$\omega(u) = \begin{cases} a & \text{if } u \text{ is a leaf,} \\ b & \text{otherwise.} \end{cases}$$

It is easy to see that $W_{1,1}$ is the Wiener index, $W_{1,0}$ is the terminal Wiener index, and $W_{0,1}$ is the spinal Wiener index. Let us prove an equivalent representation for $W_{a,b}$ that generalises (3).

Proposition 18. *We have*

$$\sum_{\{v,w\} \subseteq V(T)} \omega(v)\omega(w) d(v,w) = \sum_{vw \in E(T)} \left(\sum_{u \in V(T_v)} \omega(u) \right) \left(\sum_{u \in V(T_w)} \omega(u) \right),$$

where T_v and T_w are the components of $T - vw$ containing v and w respectively.

Proof. A pair of two vertices v' and w' occurs on the right side of the equation with weight $\omega(v')\omega(w')$ for every edge vw such that $v' \in T_v$ and $w' \in T_w$ (or the other way around). Equivalently, whenever the unique path $P(v', w')$ from v' to w' contains the edge vw . Every pair of two vertices v' and w' is counted $d(v', w')$ times in this way and therefore contributes $\omega(v')\omega(w') d(v', w')$. The sum over all pairs gives the left side of the equation. \square

Let $\mathcal{L}(T)$ be the set of leaves of a tree T . If T is rooted, the root is only counted as a leaf if it is the only vertex. In view of Proposition 18, we can write

$$W_{a,b}(T) = \sum_{vw \in E(T)} \rho_2(T_v)\rho_2(T_w),$$

where $\rho_2(T) = a|\mathcal{L}(T)| + b|V(T) - \mathcal{L}(T)|$. Here, T_v and T_w are regarded as rooted at v and w , respectively.

Lemma 19. *Let T be a tree with degree sequence D for which $W_{a,b}(T)$ attains its minimum. For any pair of disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_2(A_1), \dots, \rho_2(A_k)\} \geq \max\{\rho_2(B_1), \dots, \rho_2(B_\ell)\}$,
- or $k \leq \ell$ and $\max\{\rho_2(A_1), \dots, \rho_2(A_k)\} \leq \min\{\rho_2(B_1), \dots, \rho_2(B_\ell)\}$.

Proof. The proof is analogous to the one for the Wiener index (Lemma 4), using ρ_2 instead of ρ_0 . \square

In other words, a tree that minimises $W_{a,b}$ is ρ_2 -exchange-extremal. Moreover, if $H = [H_1, \dots, H_k]$, then

$$\rho_2(H) = \sum_{i=1}^k \rho_2(H_i) + b. \tag{8}$$

We see that conditions **I.1**, **I.2** and **I.3** are satisfied for every fixed pair of positive numbers a, b . Hence, we have the following theorem:

Theorem 20. Given a degree sequence D , $W_{a,b}$ is minimised by $\mathcal{G}(D)$ for every fixed pair of positive numbers a, b .

If we take the limits $a \rightarrow 0$ or $b \rightarrow 0$, we find that the greedy tree minimises $W_{0,b}$ and $W_{a,0}$, thus in particular the terminal and the spinal Wiener index. However, it may not be the unique optimal tree, since strict inequalities may become non-strict in the limit. Let us, for example, exhibit this phenomenon for the terminal Wiener index $W_{1,0}$. Consider the trees with degree sequence $(3, 2, 2, 2, 2, 2, 1, 1, 1)$ shown in Figure 5. Note




Figure 5: Optimal trees for the terminal Wiener index.

that $W_{1,0}(T^*) = 6 + 5 + 5 = 16$ and $W_{1,0}(T) = 6 + 6 + 4 = 16$, so the greedy tree T^* and the tree T have the same terminal Wiener index and are thus both extremal. Both trees satisfy Lemma 19.

2.6 The Steiner Wiener index

The Steiner distance of a graph, introduced by Chartrand et al. [6] in 1989, is a natural and nice generalisation of the classical graph distance. For $r \geq 2$, let $\{v_1, v_2, \dots, v_r\}$ be a set of vertices of a graph G . We denote by $S(v_1, v_2, \dots, v_r)$ the smallest subtree of G which contains all the vertices v_1, v_2, \dots, v_r . Then the *Steiner distance* of $\{v_1, \dots, v_r\}$, denoted by $sd(v_1, v_2, \dots, v_r)$, is the number of edges in $S(v_1, v_2, \dots, v_r)$. For $r = 2$, the Steiner distance clearly coincides with the classical distance, i.e., $sd(v_1, v_2) = d(v_1, v_2)$.

In [25], the authors define the *Steiner r -Wiener index* $SW_r(G)$ of a graph G as a generalisation of the Wiener index in the following way:

$$SW_r(G) = \sum_{\{v_1, \dots, v_r\} \subseteq V(G)} sd(v_1, \dots, v_r).$$

It is straightforward that the case $r = 2$ corresponds to the classical Wiener index. We have $SW_1(G) = 0$ for every graph G and $SW_n(G) = n - 1$ for every n -vertex graph G , so it is natural to restrict r to the set $\{2, 3, \dots, n - 1\}$. There is also an alternative formula for the Steiner Wiener index of trees generalising (3), see [25]:

Proposition 21. Let T be an n -vertex tree, and r a positive integer. We have

$$SW_r(T) = \sum_{uv \in E(T)} \sum_{i=1}^{r-1} \binom{|V(T_u)|}{i} \binom{|V(T_v)|}{r-i} \quad (9)$$

$$= \sum_{uv \in E(T)} \left(\binom{n}{r} - \binom{|V(T_u)|}{r} - \binom{|V(T_v)|}{r} \right), \quad (10)$$

where T_u and T_v are the components of $T - uv$ containing u and v respectively.

Proof. Equation (9) is already proven in [25], and Equation (10) follows directly from the Vandermonde identity. There is also a direct combinatorial argument: the edge uv is contained in $S(v_1, v_2, \dots, v_r)$ if and only if the set $\{v_1, v_2, \dots, v_r\}$ contains vertices of both T_u and T_v . Equivalently, $\{v_1, v_2, \dots, v_r\}$ can be any set of r vertices that is not a subset of either $V(T_u)$ or $V(T_v)$. Our formula follows immediately. \square

Zhang et al. [39, Question 1.1] asked in a recent paper whether the greedy tree minimises the Steiner r -Wiener index for every r . In the following, we will answer this question affirmatively. Recall for the following lemma that $\rho_0(T)$ is simply defined to be the number of vertices of T . As for the Wiener index, we will show ρ_0 -exchange-extremality. To avoid degeneracies (see the example below), we consider $\text{SW}_r(T) + \varepsilon W(T)$ for some fixed $\varepsilon > 0$ and later let ε go to 0.

Lemma 22. *Let ε be a fixed positive real number, and let T be a tree for which $\text{SW}_r(T) + \varepsilon W(T)$ attains its minimum among trees with degree sequence D . Then, for any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_0(A_1), \dots, \rho_0(A_k)\} \geq \max\{\rho_0(B_1), \dots, \rho_0(B_\ell)\}$,
- or $k \leq \ell$ and $\max\{\rho_0(A_1), \dots, \rho_0(A_k)\} \leq \min\{\rho_0(B_1), \dots, \rho_0(B_\ell)\}$.

In other words, T is ρ_0 -exchange-extremal.

Proof. We use the same notation as in the proof of Lemma 4, see also Figure 4 again. We have

$$\begin{aligned} \text{SW}_r(T) + \varepsilon W(T) &= \sum_{j=0}^t \left(\binom{n}{r} - \binom{\alpha + p_j}{r} - \binom{q_{t-j} + \beta}{r} + \varepsilon(\alpha + p_j)(q_{t-j} + \beta) \right) + C_T, \\ &= \sum_{j=0}^t \left(\binom{n}{r} - \binom{\alpha + p_j}{r} - \binom{n - \alpha - p_j}{r} \right. \\ &\quad \left. + \varepsilon(\alpha + p_j)(n - \alpha - p_j) \right) + C_T, \end{aligned}$$

where C_T is invariant under permutations of the A_i s and B_i s. This is a strictly concave function of α , so it can only attain its minimum when α is either at its largest or smallest value. As in the proof of Lemma 4, this implies the statement. \square

We thus have the following theorem.

Theorem 23. *Given a degree sequence D , a positive integer r and a positive real number ε , $\text{SW}(T) + \varepsilon W(T)$ attains its minimum if and only if T is the greedy tree $\mathcal{G}(D)$.*

As mentioned before, we now take the limit $\varepsilon \rightarrow 0$ to obtain the following theorem.

Theorem 24. *Given a degree sequence D , the greedy tree attains the minimum of the Steiner r -Wiener index SW_r .*

As was also the case previously for the terminal Wiener index and the spinal Wiener index, the greedy tree might not be unique with the minimum Steiner r -Wiener index. This can happen when r is quite large (compared to the total number of vertices), as in the following example: for the degree sequence $D = (3, 2, 2, 1, 1, 1)$, both the greedy tree and the only other tree with the same degree sequence have the same Steiner 5-Wiener index of 27.




Figure 6: Both trees with degree sequence $(3, 2, 2, 1, 1, 1)$ have the same Steiner 5-Wiener index.

3 Decreasing recurrence rule f_ρ

Let us first mention some definitions, which are needed to describe an \mathcal{M} -tree. We use similar terminology and notation as in [1]. Let us remark that, even though large and small degrees alternate, the concept is slightly different from that of an “alternating greedy tree”, as introduced in [35].

Definition 25. A complete branch $B = [B_1, \dots, B_k]$ of a tree T is called a *pseudo-leaf branch* and its root a *pseudo-leaf* if $|V(B_1)| = |V(B_2)| = \dots = |V(B_k)| = 1$; i.e., all vertices in B , except for the root $r(B)$, are leaves.

We simply write $[d]$ for a pseudo-leaf branch with d vertices (a root and $d - 1$ leaves). In particular, $[1]$ stands for a single vertex.

Definition 26. Let $(d_1, \dots, d_t, 1, \dots, 1)$ be the degree sequence of a tree T , where $d_j \geq 2$ for $1 \leq j \leq t$. The t -tuple (d_1, \dots, d_t) is called the *reduced degree sequence* of T . We assume that the d_i s are in non-increasing order, i.e., $d_1 \geq d_2 \geq \dots \geq d_t$.

By the handshake lemma, the sum of the number of leaves of T and the degrees of the non-leaf vertices, i.e., $|\mathcal{L}(T)| + \sum_{i=1}^t d_i$, where (d_1, \dots, d_t) is the reduced degree sequence of T , is equal to $2(|\mathcal{L}(T)| + t - 1)$. Thus, the number of leaves is $|\mathcal{L}(T)| = 2 - 2t + \sum_{i=1}^t d_i$. This implies that two trees with the same reduced degree sequence have the same number of leaves, therefore they have the same degree sequence.

Next, we give an explicit construction for the \mathcal{M} -tree $\mathcal{M}(D)$ with degree sequence D . Following the previous remark, we also write $\mathcal{M}(D')$ instead of $\mathcal{M}(D)$ for the same \mathcal{M} -tree, where D' is the reduced degree sequence corresponding to D .

Definition 27. Let (d_1, \dots, d_t) be a reduced degree sequence of a tree. If $t \leq d_t + 1$, then $\mathcal{M}(d_1, \dots, d_t)$ is the tree obtained by merging $t - 1$ leaves of a star $[1 + d_t]$ with the roots of the $t - 1$ stars $[d_1], [d_2], \dots, [d_{t-1}]$. In our formal notation, it can be written as $[[d_1], [d_2], \dots, [d_{t-1}], [1], \dots, [1]]$. We label selected vertices as shown in Figure 7, in such a way that

$$d(v_i) \leq d(v_j) \quad \text{if } i < j. \tag{11}$$

At this point all non-leaf vertices are labelled.




Figure 7: Labelling of the vertices of $\mathcal{M}(d_1, \dots, d_t)$ when $t \leq d_t + 1$.

On the other hand, if $t \geq d_t + 2$, we construct $\mathcal{M}(d_1, \dots, d_t)$ recursively. Let ℓ be the largest integer such that a vertex labelled v_ℓ occurs in $\mathcal{M}(d_t, \dots, d_{t-1})$, and let s be the smallest integer such that v_s is adjacent to a leaf in $\mathcal{M}(d_t, \dots, d_{t-1})$. Let $R_{d_t} = [[d_1], \dots, [d_{t-1}]]$, where the pseudo-leaves are labelled $v_{\ell+1}, \dots, v_{\ell+d_t-1}$, still respecting (11). $\mathcal{M}(d_1, \dots, d_t)$ is the tree obtained by merging the root of R_{d_t} with a leaf adjacent to v_s .




Figure 8: Step-by-step construction of $\mathcal{M}(5, 4, 4, 4, 3, 3, 3, 2)$.

We will show in this section that the \mathcal{M} -tree $\mathcal{M}(D)$ is the unique ρ -exchange-extremal tree with degree sequence D if ρ satisfies the following conditions:

II.1 the quantity ρ satisfies (2), where the function f_ρ is symmetric,

II.2 the function f_ρ is strictly decreasing (strictly decreasing in each single variable and strictly decreasing under addition of further variables),

II.3 the quantity ρ satisfies $\rho(\bullet) > \rho(B)$, for all rooted trees B with $|V(B)| > 1$.

3.1 Special case: the Hosoya index

Let T be a rooted tree, and recall that the Hosoya index, denoted by $z(T)$, is the total number of matchings of T . The number of matchings in T that do not cover the root $r(T)$ will be denoted by $z_0(T)$. We consider the following ratio, which can be interpreted as the probability that a random matching does not cover the root:

$$\rho_3(T) = \frac{z_0(T)}{z(T)}.$$

If $T = [T_1, \dots, T_k]$, then it is not hard to see that

$$z_0(T) = \prod_{j=1}^k z(T_j)$$

and

$$z(T) = z_0(T) + \sum_{j=1}^k z_0(T_j) \prod_{\substack{i=1 \\ i \neq j}}^k z(T_i).$$

It follows that

$$\rho_3(T) = \rho_3([T_1, T_2, \dots, T_k]) = \frac{1}{1 + \sum_{j=1}^k \rho_3(T_j)}. \quad (12)$$

Note that this recurrence rule f_{ρ_3} is symmetric and decreasing with respect to each of its variables and under addition of further variables. In addition, we obviously have $\rho_3(\bullet) = 1 > \rho_3(B)$ for every rooted tree B with more than one vertex. Thus **II.1**, **II.2** and **II.3** are all satisfied.

Lemma 28 (cf. [17]). *Let T be a tree with degree sequence D for which $z(T)$ attains its maximum. For any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_3(A_1), \dots, \rho_3(A_k)\} \geq \max\{\rho_3(B_1), \dots, \rho_3(B_\ell)\}$,
- or $k \leq \ell$ and $\max\{\rho_3(A_1), \dots, \rho_3(A_k)\} \leq \min\{\rho_3(B_1), \dots, \rho_3(B_\ell)\}$.

In other words, the tree with degree sequence D that minimises the Hosoya index is ρ_3 -exchange-extremal. In addition, as a special case of Theorem 22 in [1], the following holds.

Theorem 29 ([1]). *If T is a ρ_3 -exchange-extremal tree, then T is an \mathcal{M} -tree.*

3.2 Main result

In analogy to Theorem 7, we obtain the following result.

Theorem 30. *Let ρ be an invariant of rooted trees that satisfies conditions **II.1**, **II.2** and **II.3**. If a tree T is ρ -exchange-extremal, then T is an \mathcal{M} -tree.*

The proof of Theorem 30 is essentially identical to that of Theorem 7, with minor modifications to take into account that f_ρ and f_{ρ_3} are decreasing and $\rho(\bullet)$ and $\rho_3(\bullet)$ are maxima. The rest of this section will be devoted to applications of Theorem 30.

3.3 Matching polynomial and energy

Let T be a rooted tree. Denote by $m(T, k)$ the number of k -matchings in T and by $m_0(T, k)$ the number of k -matchings not containing the root. We write $M(T, x) = \sum_{k \geq 0} m(T, k)x^k$ and $M_0(T, x) = \sum_{k \geq 0} m_0(T, k)x^k$ for the generating polynomials corresponding to m and m_0 respectively. Moreover, we define the following ratio for $x > 0$:

$$\tau(T, x) = \frac{M_0(T, x)}{M(T, x)}.$$

Lemma 31 ([1]). *Let $x > 0$, and let T be a tree such that $M(T, x) \leq M(T', x)$ for every tree T' with the same degree sequence. For any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\tau(A_j, x) : 1 \leq j \leq k\} \geq \max\{\tau(B_j, x) : 1 \leq j \leq \ell\}$,
- or $k \leq \ell$ and $\max\{\tau(A_j, x) : 1 \leq j \leq k\} \leq \min\{\tau(B_j, x) : 1 \leq j \leq \ell\}$.

The tree that minimises the matching polynomial $M(T, x)$ for some $x > 0$ is therefore $\tau(\cdot, x)$ -exchange-extremal. In addition, we have the following generalisation of (12): if $T = [T_1, T_2, \dots, T_k]$, then

$$\tau(T, x) = \frac{1}{1 + x \sum_{j=1}^k \tau(T_j, x)}.$$

The recurrence rule f_ρ , with $\rho(T) = \tau(T, x)$, is thus symmetric and decreasing with respect to each of its variables and under addition of further variables. In addition, $\tau(\bullet, x) = 1$ is the unique maximum among all rooted trees for every x . Thus the following result from [1] can be seen as a special case of Theorem 30:

Theorem 32 ([1]). *Let x be a fixed positive real number. If T is a tree with degree sequence D , then*

$$M(T, x) \geq M(\mathcal{M}(D), x),$$

and equality holds only if T is isomorphic to $\mathcal{M}(D)$.

In view of the connection between the energy and the matching polynomial (namely the Coulson formula (1)), we also get the following corollary.

Corollary 33 ([1]). *Given the degree sequence D of a tree, the energy is minimised by the \mathcal{M} -tree $\mathcal{M}(D)$.*

3.4 Merrifield-Simmons index

Recall that the number of independent sets in T is also called the Merrifield-Simmons index and denoted by $\sigma(T)$. Let $\sigma_0(T)$ be the number of independent sets of T that do not contain the root, and define $\rho_4(T) = \sigma(T)/\sigma_0(T)$. The following lemma from [1] states that the tree that maximises the number of independent sets among all trees with a given degree sequence D has to be ρ_4 -exchange-extremal.

Lemma 34 ([1]). *Let T be a tree with degree sequence D for which $\sigma(T)$ attains its maximum. Then, for any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_4(A_j) : 1 \leq j \leq k\} \geq \max\{\rho_4(B_j) : 1 \leq j \leq \ell\}$,
- or $k \leq \ell$ and $\max\{\rho_4(A_j) : 1 \leq j \leq k\} \leq \min\{\rho_4(B_j) : 1 \leq j \leq \ell\}$.

The recurrence rule f_{ρ_4} corresponding to ρ_4 is found in a similar way to (12), and is given by

$$\rho_4([T_1, T_2, \dots, T_k]) = 1 + \prod_{j=1}^k \rho_4(T_j)^{-1}.$$

Note that it is decreasing with respect to each of its variables (and under addition of further variables), and $\rho_4(\bullet) = 2$ is easily seen to be maximal among all trees. Hence we can deduce the following theorem as a consequence of Theorem 30.

Theorem 35 ([1]). *Given a degree sequence D , the Merrifield-Simmons index is maximised by $\mathcal{M}(D)$.*

3.5 Solvability

Let G be a graph and v a vertex in G . The open neighbourhood of v , denoted $N(v)$, is the set $\{u \in V(G) : uv \in E(G)\}$, and the closed neighbourhood of v is $N[v] = N(v) \cup \{v\}$.

Let \mathbb{F}_2 be the field with two elements. The *solvability* of G , introduced in [15] and denoted $s(G)$, is the number of pairs $(a, b) \in \mathbb{F}_2^{V(G)} \times \mathbb{F}_2^{V(G)}$ such that there exists a vector $x \in \mathbb{F}_2^{V(G)}$ that satisfies

$$(A + \text{diag}(a))x = b,$$

where A is the adjacency matrix of G and $\text{diag}(a)$ is the diagonal matrix whose diagonal is a .

This can be interpreted as a domination problem with parity constraints. We are looking for a set S of vertices satisfying, for each of the vertices of the graph, one of four possible conditions: the open/closed neighbourhood has to contain an even/odd number of vertices in S . Here, a encodes the open/closed neighbourhood condition and b encodes the required parity. The solvability of a graph measures how many instances of the problem have solutions. It turns out that the solvability of trees can be calculated by a recursion involving a second auxiliary quantity t associated with rooted trees, as stated in the following lemma.

Lemma 36 ([15]). *Let T be a rooted tree. If $T = [T_1, T_2, \dots, T_k]$, then*

$$s(T) = 8 \prod_{i=1}^k s(T_i) - 5 \prod_{i=1}^k t(T_i),$$

$$t(T) = 8 \prod_{i=1}^k s(T_i) - 6 \prod_{i=1}^k t(T_i),$$

with initial values $s(\bullet) = 3$ and $t(\bullet) = 2$.

We remark that the value of the parameter t generally depends on the choice of root, unlike s . Note also that $s(T)$ and $t(T)$ are both positive, and that $s(T) > t(T)$ holds for every rooted tree T .

It was shown in [15] that the path has the greatest solvability among all trees (in fact all graphs) of a given order, while the star has the least solvability, but no other extremal results for the solvability of trees are available. Let us show that our general results apply to the solvability as well. To this end, define $\rho_5(T)$ to be the ratio $\rho_5(T) = \frac{s(T)}{t(T)} > 1$. If T can be decomposed as $[T_1, \dots, T_k]$, then by Lemma 36 we have:

$$\rho_5(T) = \frac{8 \prod_{i=1}^k \rho_5(T_i) - 5}{8 \prod_{i=1}^k \rho_5(T_i) - 6}. \quad (13)$$

Note that this is a symmetric and decreasing recurrence rule, so ρ_5 satisfies conditions **II.1** and **II.2**. Moreover, **II.3** is also satisfied by (13), since

$$\rho_5(T) = 1 + \frac{1}{8 \prod_{i=1}^k \rho_5(T_i) - 6} < 1 + \frac{1}{8 - 6} = \frac{3}{2} = \rho_5(\bullet)$$

for every tree rooted tree T with more than one vertex. Let us now prove the main lemma of this subsection.

Lemma 37. *Let T be a tree with degree sequence D for which $s(T)$ attains its minimum. Then, for any two disjoint complete branches $A = [A_1, \dots, A_k]$ and $B = [B_1, \dots, B_\ell]$ in T , we have*

- either $k \geq \ell$ and $\min\{\rho_5(A_j) : 1 \leq j \leq k\} \geq \max\{\rho_5(B_j) : 1 \leq j \leq \ell\}$,
- or $k \leq \ell$ and $\max\{\rho_5(A_j) : 1 \leq j \leq k\} \leq \min\{\rho_5(B_j) : 1 \leq j \leq \ell\}$.

Proof. Let $P = u_0 u_1 \dots u_m$ be the path between $r(A) = u_0$ and $r(B) = u_m$. For $1 \leq j < m$, let $U_j = [U_j^1, \dots, U_j^{\text{rd}(U_j)}]$ be the component containing u_j (rooted at u_j) when we remove all the edges of the path P . We also denote by X_j the component containing u_j in $T - u_{j-1}u_j$, rooted at u_j . We consider the tree T as rooted at $r(A)$. Using Lemma 36, we can write $s(T)$ and $t(T)$ in terms of matrices as follows:




Figure 9: Decomposition of T in the proof of Lemma 37.

$$\begin{pmatrix} s(T) \\ t(T) \end{pmatrix} = \begin{pmatrix} 8 \prod_{i=1}^k s(A_i) & -5 \prod_{i=1}^k t(A_i) \\ 8 \prod_{i=1}^k s(A_i) & -6 \prod_{i=1}^k t(A_i) \end{pmatrix} \begin{pmatrix} s(X_1) \\ t(X_1) \end{pmatrix}.$$

Moreover, for $1 \leq j < m$, we have

$$\begin{pmatrix} s(X_j) \\ t(X_j) \end{pmatrix} = \begin{pmatrix} 8 \prod_{i=1}^{\text{rd}(U_j)} s(U_j^i) & -5 \prod_{i=1}^{\text{rd}(U_j)} t(U_j^i) \\ 8 \prod_{i=1}^{\text{rd}(U_j)} s(U_j^i) & -6 \prod_{i=1}^{\text{rd}(U_j)} t(U_j^i) \end{pmatrix} \begin{pmatrix} s(X_{j+1}) \\ t(X_{j+1}) \end{pmatrix}.$$

Let us denote the 2×2 -matrix on the right side of this equation by M_j . Empty products (if $\text{rd}(U_j) = 0$) are defined to be 1, which is consistent with the recursion. Then we have

$$\begin{pmatrix} s(T) \\ t(T) \end{pmatrix} = \begin{pmatrix} 8 \prod_{i=1}^k s(A_i) & -5 \prod_{i=1}^k t(A_i) \\ 8 \prod_{i=1}^k s(A_i) & -6 \prod_{i=1}^k t(A_i) \end{pmatrix} M \begin{pmatrix} s(B) \\ t(B) \end{pmatrix},$$

where $M = M_0 \times M_1 \times M_2 \times \cdots \times M_{m-1}$, M_0 being the identity matrix. If we set $M = \begin{pmatrix} M_{00} & M_{01} \\ M_{10} & M_{11} \end{pmatrix}$, then we can write the solvability of T as

$$\begin{aligned} s(T) &= \prod_{i=1}^k s(A_i) \prod_{i=1}^{\ell} s(B_i) (64M_{00} + 64M_{01}) - \prod_{i=1}^k s(A_i) \prod_{i=1}^{\ell} t(B_i) (40M_{00} + 48M_{01}) \\ &\quad - \prod_{i=1}^k t(A_i) \prod_{i=1}^{\ell} s(B_i) (40M_{10} + 40M_{11}) + \prod_{i=1}^k t(A_i) \prod_{i=1}^{\ell} t(B_i) (25M_{10} + 30M_{11}) \end{aligned}$$

$$\begin{aligned}
&= \prod_{i=1}^k t(A_i) \prod_{i=1}^{\ell} t(B_i) \left[(64M_{00} + 64M_{01}) \prod_{i=1}^k \rho_5(A_i) \prod_{i=1}^{\ell} \rho_5(B_i) \right. \\
&\quad \left. - (40M_{00} + 48M_{01}) \prod_{i=1}^k \rho_5(A_i) - (40M_{10} + 40M_{11}) \prod_{i=1}^{\ell} \rho_5(B_i) + (25M_{10} + 30M_{11}) \right].
\end{aligned}$$

Since $\prod_{i=1}^k t(A_i) \prod_{i=1}^{\ell} t(B_i)$ and $(64M_{00} + 64M_{01}) \prod_{i=1}^k \rho_5(A_i) \prod_{i=1}^{\ell} \rho_5(B_i) + (25M_{10} + 30M_{11})$ remain invariant under any rearrangements of the A_i s and B_i s, $s(T)$ attains its minimum under permutations of the A_i s and B_i s if and only if

$$\alpha \prod_{i=1}^k \rho_5(A_i) + \beta \prod_{i=1}^{\ell} \rho_5(B_i)$$

attains its maximum, where $\alpha = 40M_{00} + 48M_{01}$ and $\beta = 40M_{10} + 40M_{11}$.

Next we show that α and β are positive. For $m = 1$, M is equal to the identity matrix. So $\alpha = \beta = 40 > 0$.

Now, for $m > 1$, let us prove by induction on m that $5M_{00} + 6M_{01} > 0$, $5M_{10} + 6M_{11} > 0$ and $M_{01}, M_{11} \leq 0$. Note that the positivity of $5M_{00} + 6M_{01}$ implies the positivity of α since $\alpha = 8(5M_{00} + 6M_{01})$, and the positivity of $5M_{10} + 6M_{11}$, combined with the inequality $M_{11} \leq 0$, implies the positivity of β since $\beta = 8(5M_{10} + 5M_{11}) \geq 8(5M_{10} + 6M_{11})$. Let us write

$$\begin{bmatrix} M_{00}(m) & M_{01}(m) \\ M_{10}(m) & M_{11}(m) \end{bmatrix} = M(m) = M_0 \times M_1 \times \cdots \times M_{m-1}.$$

For $m = 2$, we have, since $s(U_1^i) > t(U_1^i) > 0$ for all i ,

$$\begin{aligned}
M_{01}(2) &= -5 \prod_{i=1}^{\text{rd}(U_1)} t(U_1^i) \leq 0, & M_{11}(2) &= -6 \prod_{i=1}^{\text{rd}(U_1)} t(U_1^i) \leq 0, \\
5M_{00}(2) + 6M_{01}(2) &= 40 \prod_{i=1}^{\text{rd}(U_1)} s(U_1^i) - 30 \prod_{i=1}^{\text{rd}(U_1)} t(U_1^i) > 0, \\
5M_{10}(2) + 6M_{11}(2) &= 40 \prod_{i=1}^{\text{rd}(U_1)} s(U_1^i) - 36 \prod_{i=1}^{\text{rd}(U_1)} t(U_1^i) > 0.
\end{aligned}$$

Suppose now that the statement is true for $m = d$. For the induction step, we take $m = d + 1$, and $M(d + 1) = M_0 \times M_1 \times \cdots \times M_d = M(d) \times M_d$. We have

$$\begin{aligned}
M_{00}(d + 1) &= (8M_{00}(d) + 8M_{01}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i), \\
M_{01}(d + 1) &= -(5M_{00}(d) + 6M_{01}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i),
\end{aligned}$$

$$M_{10}(d+1) = (8M_{10}(d) + 8M_{11}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i),$$

$$M_{11}(d+1) = -(5M_{10}(d) + 6M_{11}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i).$$

By the induction hypothesis, we have $M_{01}(d), M_{11}(d) \leq 0$, $5M_{00}(d) + 6M_{01}(d) > 0$ and $5M_{10}(d) + 6M_{11}(d) > 0$. For all i , $s(U_{d+1}^i) > t(U_{d+1}^i) > 0$, so

$$M_{01}(d+1) \leq 0, \text{ and } M_{11}(d+1) \leq 0.$$

Moreover, we get

$$\begin{aligned} & 5M_{00}(d+1) + 6M_{01}(d+1) \\ &= 8 \cdot (5M_{00}(d) + 5M_{01}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i) - 6 \cdot (5M_{00}(d) + 6M_{01}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i) \\ &\geq (5M_{00}(d) + 6M_{01}(d)) \left(8 \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i) - 6 \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i) \right) > 0 \end{aligned}$$

and

$$\begin{aligned} & 5M_{10}(d+1) + 6M_{11}(d+1) \\ &= 8 \cdot (5M_{10}(d) + 5M_{11}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i) - 6 \cdot (5M_{10}(d) + 6M_{11}(d)) \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i) \\ &\geq (5M_{10}(d) + 6M_{11}(d)) \left(8 \prod_{i=1}^{\text{rd}(U_{d+1})} s(U_{d+1}^i) - 6 \prod_{i=1}^{\text{rd}(U_{d+1})} t(U_{d+1}^i) \right) > 0. \end{aligned}$$

This completes our induction. Since we know now that α and β are positive, and that $\rho_5(A_1), \dots, \rho_5(A_k), \rho_5(B_1), \dots, \rho_5(B_\ell)$ are all greater than 1 by definition, Lemma 8 applies and shows that $s(T)$ can only be minimal if

- $\alpha \geq \beta$, $k \geq \ell$ and $\min\{\rho_5(A_j) : 1 \leq j \leq k\} \geq \max\{\rho_5(B_j) : 1 \leq j \leq \ell\}$,
- or $\alpha \leq \beta$, $k \leq \ell$ and $\max\{\rho_5(A_j) : 1 \leq j \leq k\} \leq \min\{\rho_5(B_j) : 1 \leq j \leq \ell\}$.

This completes the proof. □

So a tree that minimises the solvability, given the degree sequence, must be ρ_5 -exchange-extremal. Applying Theorem 30, we immediately get the following theorem.

Theorem 38. *Given a degree sequence of a tree D , the solvability is minimised by $\mathcal{M}(D)$.*

4 Majorisation of degree sequences

Recall that a sequence $B = (b_1, b_2, \dots, b_n)$ is said to majorise D if $b_1 + \dots + b_n = d_1 + \dots + d_n$ and $d_1 + \dots + d_k \leq b_1 + \dots + b_k$ for every k with $1 \leq k < n$. In this case we write $B \triangleright D$. Let \mathbb{T}_n be the set of n -vertex trees. In this section we study the class of trees whose degree sequence is majorised by a fixed degree sequence B of a tree, i.e.,

$$\mathbb{T}_{\triangleleft B} = \{T \in \mathbb{T}_n : B \triangleright D, \text{ where } D \text{ is the degree sequence of } T\}.$$

Definition 39. Let ρ be an invariant of rooted trees which satisfies a recursive relation as in Equation (2), and let \mathbb{S}_n denote the set of permutations of $1, 2, \dots, n$. We say that the tree invariant I is maximum- ρ -compatible if for every tree H and every choice of two of its leaves v and w , every collection of rooted trees T_1, T_2, \dots, T_r with $\rho(T_1) \geq \rho(T_2) \geq \dots \geq \rho(T_r)$, and every integer s with $r/2 \leq s \leq r$, the maximum value of I among all trees in the set

$$\mathbb{K}_s = \{[T_{\sigma(1)}, \dots, T_{\sigma(k)}]vHw[T_{\sigma(k+1)}, \dots, T_{\sigma(r)}] : \sigma \in \mathbb{S}_r \text{ and } r - s \leq k \leq s\}$$

is attained by (at least) one of the following two trees:

$$[T_1, \dots, T_s]vHw[T_{s+1}, \dots, T_r] \text{ or } [T_{s+1}, \dots, T_r]vHw[T_1, \dots, T_s].$$

In words, among all possibilities to attach T_1, T_2, \dots, T_r to v and w in such a way that their degrees are not greater than some fixed bound $s + 1$, the maximum value of I is reached when the s trees T_i with largest ρ -values are attached to one of the two vertices v and w , so that its degree is $s + 1$. Note here that if $s \leq s'$ then $\mathbb{K}_s \subseteq \mathbb{K}_{s'}$. In the same way, one defines an invariant I to be minimum- ρ -compatible (replacing “maximum value” by “minimum value” in the definition).

Several examples from the previous sections fit this scheme. In each case, the proof is essentially the same as the proof that extremality with respect to I implies ρ -exchange-extremality.

- The Wiener index is minimum- ρ_0 -compatible, and this extends to the generalisations discussed in Subsections 2.5 and 2.6.
- The number of subtrees is maximum- ρ -compatible with respect to the invariant $\rho(T) = \eta(T, r(T))$ defined in Subsection 2.3.
- The weighted number of rooted spanning forests $\text{rf}(T, x)$ discussed in Subsection 2.4 is minimum- ρ_1 -compatible with respect to the invariant ρ_1 defined there.
- The Hosoya index is minimum- ρ_3 -compatible (see Subsection 3.1), and this extends to the matching generating polynomial $M(T, x)$.
- The Merrifield-Simmons index is maximum- ρ_4 -compatible (Subsection 3.4).

- The solvability is minimum- ρ_5 -compatible (Subsection 3.5).

We note that for any maximum- ρ -compatible invariant I , a tree with degree sequence D that attains the maximum value of I is necessarily ρ -exchange-extremal, since the compatibility condition is an extension of ρ -exchange-extremality.

Theorem 40. *Let I be a maximum- ρ -compatible tree invariant, for some ρ that satisfies (2), and let B be a degree sequence of a tree. Then*

- i) $\max\{I(T) : T \in \mathbb{T}_{\triangleleft B}\} = I(\mathcal{G}(B))$ if ρ satisfies properties **I.1**, **I.2** and **I.3**,
- ii) $\max\{I(T) : T \in \mathbb{T}_{\triangleleft B}\} = I(\mathcal{M}(B))$ if ρ satisfies properties **II.1**, **II.2** and **II.3**,

Proof. We only prove part i), the proof of part ii) is analogous. As mentioned earlier, by definition of maximum- ρ -compatibility of I , $\max\{I(T) : T \in \mathbb{T}_{\triangleleft B}\}$ is reached by a ρ -exchange-extremal tree, say E . It is only left to prove that E can be chosen to have degree sequence B . Let $T \in \mathbb{T}_{\triangleleft B}$, with a degree sequence D (thus $D \triangleleft B$), such that $D = (d_1, d_2, \dots, d_n) \neq B = (b_1, b_2, \dots, b_n)$. Let r be the smallest index for which $d_r > b_r$ (such an index must exist, since B and D have the same sum), and let ℓ be the largest index less than r such that $d_\ell < b_\ell$ (such an index must exist, since $b_1 + b_2 + \dots + b_r \geq d_1 + d_2 + \dots + d_r$ by majorisation). Observe that $d_i = b_i$ for $\ell < i < r$. Now define $B_1 = (b'_1, b'_2, \dots, b'_n)$ by $b'_i = b_i$ for $i \notin \{\ell, r\}$, $b'_\ell = b_\ell - 1$ and $b'_r = b_r + 1$. We note that B_1 is still non-increasing, thus a valid degree sequence: $b'_\ell = b_\ell - 1 \geq d_\ell \geq d_{\ell+1} \geq b'_{\ell+1}$ and $b'_r = b_r + 1 \leq d_r \leq d_{r-1} \leq b'_{r-1}$. Moreover, it is easy to verify that $D \triangleleft B_1$, since $b'_i \geq d_i$ for $i < r$, while $b'_1 + b'_2 + \dots + b'_j = b_1 + b_2 + \dots + b_j$ for $j \geq r$. Clearly, we also have $B_1 \triangleleft B$. We can repeat this procedure with B_1 and D to obtain a degree sequence B_2 with $D \triangleleft B_2 \triangleleft B_1$, and so on. This yields a sequence of degree sequences $B = B_0, B_1, B_2, \dots, B_p = D$ such that

$$B = B_0 \triangleright B_1 \triangleright \dots \triangleright B_p = D,$$

and B_i and B_{i+1} always only differ in two entries (by exactly 1 each). We therefore restrict ourselves first to the case that B and D only differ in two entries: $b_\ell = d_\ell + 1$, $b_r = d_r - 1$ and $b_i = d_i$ for $i \notin \{\ell, r\}$.

Let v and w be distinct vertices in T such that $d(v) = d_\ell \geq d_r = d(w)$. Then T can be decomposed as $[T_1, \dots, T_{d_\ell-1}]vHw[T_{d_\ell}, \dots, T_{d_\ell+d_r-2}]$ for some tree H and branches $T_1, T_2, \dots, T_{d_\ell+d_r-2}$. Let $\sigma \in \mathbb{S}_{d_\ell+d_r-2}$ such that $\rho(T_{\sigma(1)}) \geq \rho(T_{\sigma(2)}) \geq \dots \geq \rho(T_{\sigma(d_\ell+d_r-2)})$.

By the definition of maximum- ρ -compatibility, we can find a tree T' , which is either

$$\begin{aligned} & [T_{\sigma(1)}, \dots, T_{\sigma(d_\ell)}]vHw[T_{\sigma(d_\ell+1)}, \dots, T_{\sigma(d_\ell+d_r-2)}] \quad \text{or} \\ & [T_{\sigma(d_\ell+1)}, \dots, T_{\sigma(d_\ell+d_r-2)}]vHw[T_{\sigma(1)}, \dots, T_{\sigma(d_\ell)}], \end{aligned}$$

such that its degree sequence is B and $I(T) \leq I(T')$. In the general case, we can find a sequence of trees $T_0, T_1, \dots, T_p = T$ whose degree sequences are $B = B_0, B_1, \dots, B_p = D$ respectively, such that

$$I(T_0) \geq I(T_1) \geq \dots \geq I(T_p) = I(T).$$

This shows that the extremal tree E mentioned earlier can indeed be chosen to have degree sequence B . The statement of the theorem follows. \square

Corollary 41. *Let $\mathbb{A} \subseteq \mathbb{T}_n$ be a class of trees with n vertices and B a degree sequence such that for every tree $T \in \mathbb{A}$, its degree sequence D satisfies $D \triangleleft B$. If I is a maximum- ρ -compatible tree invariant, then*

- $\max\{I(T) : T \in \mathbb{A}\} = I(\mathcal{G}(B))$ if ρ satisfies **I.1**, **I.2** and **I.3** and $\mathcal{G}(B) \in \mathbb{A}$,
- $\max\{I(T) : T \in \mathbb{A}\} = I(\mathcal{M}(B))$ if ρ satisfies **II.1**, **II.2** and **II.3** and $\mathcal{M}(B) \in \mathbb{A}$.

Again, an analogous statement holds if I is minimum- ρ -compatible (replacing max by min).

Proof. The degree sequence condition implies that $\mathbb{A} \subseteq \mathbb{T}_{\triangleleft B}$, so the statement is immediate from Theorem 40. □

We conclude this paper by discussing different applications of Corollary 41.

- The first example is the class of all trees of order n , $\mathbb{A} = \mathbb{T}_n$, where we can take $B = (n-1, 1, \dots, 1)$. The only tree with this degree sequence is the star $S_n = \mathcal{G}(B) = \mathcal{M}(B)$. Therefore, we obtain the following known results as immediate corollaries: among all n -vertex trees, the star minimises the Wiener index [9], the Hosoya index and the energy [11], the quantity $\text{rf}(T, x)$ [43] for every $x > 0$ (thus also the incidence energy) and the solvability [15], while it maximises the number of subtrees [30] and the Merrifield-Simmons index [28].
- Next, we take \mathbb{A} to be the set of all n -vertex trees whose vertex degrees are at most equal to d . Here, we can take $B = (d, \dots, d, r, 1, \dots, 1)$, where $1 \leq r < d$ and $r \equiv n-1 \pmod{d-1}$. The greedy tree $\mathcal{G}(B)$ is in this case called a Volkmann tree. Again, several extremality results now follow as direct applications of Corollary 41. For example, the Volkmann tree minimises the Wiener index [10] and the polynomial $\text{rf}(T, x)$ for every $x > 0$ [20] (thus also the incidence energy), while it maximises the number of subtrees [23]. On the other hand, the \mathcal{M} -tree $\mathcal{M}(B)$ is known to be extremal for the Hosoya index, the energy of a graph and the Merrifield-Simmons index [18, 16]. We also obtain new results for the Steiner Wiener index (which is minimised by $\mathcal{G}(B)$) and the solvability (which is minimised by $\mathcal{M}(B)$).
- Next, let \mathbb{A} be the set of n -vertex trees with exactly ℓ leaves. Now we can take $B = (\ell, 2, \dots, 2, 1, \dots, 1)$ ($n-\ell-1$ copies of 2, ℓ copies of 1) and find that either $\mathcal{G}(B)$ or $\mathcal{M}(B)$ is extremal for each of the invariants we considered in the previous sections, thus recovering several known results (and adding some new ones). For instance, the greedy tree $\mathcal{G}(B)$ is known to minimise the Wiener index [9, 34] and the polynomial $\text{rf}(T, x)$ as well as the incidence energy [19]. On the other hand, it maximises the number of subtrees [3]. Similarly, the extremal tree for the Hosoya index, the Merrifield-Simmons index and the energy is known to be $\mathcal{M}(B)$ [38, 36, 37]. We find now also that $\mathcal{G}(B)$ minimises the Steiner Wiener index, while $\mathcal{M}(B)$ minimises the solvability.

- Finally, let \mathbb{A} be the set of n -vertex trees having r branching vertices, i.e., vertices of degree greater than or equal to 3. Here, we can take $B = (n-2r+1, 3, \dots, 3, 1, \dots, 1)$ ($r-1$ copies of 3, $n-r$ copies of 1). It was shown in [26] that the greedy tree $\mathcal{G}(B)$ minimises the Wiener index in this class of trees if $r \leq \frac{n+2}{3}$, while the case $r > \frac{n+2}{3}$ was left as an open problem. Minimality of $\mathcal{G}(B)$ with respect to the Wiener index is a consequence of our results for arbitrary r now, and we also obtain extremality of $\mathcal{G}(B)$ or $\mathcal{M}(B)$ for all the other invariants mentioned in this paper.

References

- [1] E. O. D. Andriantiana. Energy, Hosoya index and Merrifield-Simmons index of trees with prescribed degree sequence. *Discrete Appl. Math.*, 161:724–741, 2013.
- [2] E. O. D. Andriantiana and S. Wagner. Spectral moments of trees with given degree sequence. *Linear Algebra Appl.*, 439(12):3980–4002, 2013.
- [3] E. O. D. Andriantiana, S. Wagner, and H. Wang. Greedy trees, subtrees and antichains. *Electron. J. Combin.*, 20(3):#P28, 2013.
- [4] M. Bartlett, E. Krop, C. Magnant, F. Mutiso, and H. Wang. Variations of distance-based invariants of trees. *J. Combin. Math. Combin. Comput.*, 91:19–29, 2014.
- [5] T. Bıyıkoglu and J. Leydold. Graphs with given degree sequence and maximal spectral radius. *Electron. J. Combin.*, 15:#R119, 2008.
- [6] G. Chartrand, O. R. Oellermann, S. Tian, and H. Zou. Steiner distance in graphs. *Časopis Pest. Mat.*, 114:399–410, 1989.
- [7] Y.-H. Chen, D. Gray, Y.-L. Jin, and X.-D. Zhang. On majorization of closed walk vectors of trees with given degree sequences. *Appl. Math. Comput.*, 336:326–337, 2018.
- [8] D. Cvetković, P. Rowlinson, and S. Simić. Signless Laplacians of finite graphs. *Lin. Algebra Appl.*, 423:155–171, 2007.
- [9] A. A. Dobrynin, R. Entringer, and I. Gutman. Wiener index of trees: theory and applications. *Acta Appl. Math.*, 66(3):211–249, 2001.
- [10] M. Fischermann, A. Hoffmann, D. Rautenbach, L. Székely, and L. Volkmann. Wiener index versus maximum degree in trees. *Discrete Appl. Math.*, 122(1):127–137, 2002.
- [11] I. Gutman. Acyclic systems with extremal Hückel π -electron energy. *Theor. Chim. Acta*, 45:79–87, 1977.
- [12] I. Gutman, B. Furtula, and M. Petrović. Terminal Wiener index. *J. Math. Chem.*, 46(2):522–531, 2009.
- [13] I. Gutman, D. Kiani, and M. Mirzakhah. On incidence energy of graphs. *MATCH Commun. Math. Comput. Chem.*, 62:573–580, 2009.
- [14] I. Gutman and O. E. Polansky. *Mathematical Concepts in Organic Chemistry*. Springer, Berlin, 1986.

- [15] J. Hatzl and S. Wagner. Combinatorial properties of a general domination problem with parity constraints. *Discrete Math.*, 308:6355–6367, 2008.
- [16] C. Heuberger and S. Wagner. Maximizing the number of independent subsets over trees with bounded degree. *J. Graph Theory*, 58(1):49–68, 2008.
- [17] C. Heuberger and S. Wagner. Chemical trees minimizing energy and Hosoya index. *J. Math. Chem.*, 46:214–230, 2009.
- [18] C. Heuberger and S. Wagner. On a class of extremal trees for various indices. *MATCH Commun. Math. Comput. Chem.*, 62:437–464, 2009.
- [19] A. Ilić and M. Ilić. Laplacian coefficients of trees with given number of leaves or vertices of degree two. *Linear Algebra Appl.*, 431(11):2195–2202, 2009.
- [20] Y.-L. Jin, Y.-N. Yeh, and X.-D. Zhang. Laplacian coefficient, matching polynomial and incidence energy of trees with described maximum degree. *J. Comb. Optim.*, 31:1345–1372, 2016.
- [21] M. R. Jooyandeh, D. Kiani, and M. Mirzakhah. Incidence energy of a graph. *MATCH Commun. Math. Comput. Chem.*, 62:561–572, 2009.
- [22] A. K. Kel'mans. On properties of the characteristic polynomial of a graph. *Cybernetics in the Service of Communism [In Russian]*, 4:27–41, 1967.
- [23] R. Kirk and H. Wang. Largest number of subtrees of trees with a given maximum degree. *SIAM J. Discrete Math.*, 22(3):985–995, 2008.
- [24] S. Li and Y. Song. On the spectral moment of trees with given degree sequences. *arXiv:math/1209.2188*, 2012.
- [25] X. Li, Y. Mao, and I. Gutman. The Steiner Wiener index of a graph. *Disc. Math. Graph Theory*, 36(2):455–465, 2016.
- [26] H. Lin. On the Wiener Index of Trees with Given Number of Branching Vertices. *MATCH Commun. Math. Comput. Chem.*, 72:301–310, 2014.
- [27] J. Liu and B. Liu. A Laplacian-energy like invariant of a graph. *MATCH Commun. Math. Comput. Chem.*, 59:355–372, 2008.
- [28] H. Prodinger and R. F. Tichy. Fibonacci numbers of graphs. *Fibonacci Quart.*, 20:16–21, 1982.
- [29] N. S. Schmuck, S. G. Wagner, and H. Wang. Greedy Trees, Caterpillars, and Wiener-type Graph Invariants. *MATCH Commun. Math. Comput. Chem.*, 68:273–292, 2012.
- [30] L. A. Székely and H. Wang. On subtrees of trees. *Adv. Appl. Math.*, 34(1):138–155, 2005.
- [31] L. A. Székely, H. Wang, and T. Wu. The sum of the distances between the leaves of a tree and the semi-regular property. *Discrete Math.*, 311(13):1197–1203, 2011.
- [32] S. Wagner, H. Wang, and X.-D. Zhang. Distance-based graph invariants of trees and the Harary index. *Filomat*, 27(1):41–50, 2013.
- [33] H. Wang. The extremal values of the Wiener index of a tree with given degree sequence. *Discrete Appl. Math.*, 156:2647–2654, 2009.

- [34] H. Wang. On trees with given number of pendant edges and their Wiener indices. *Adv. Appl. Math. Sci.*, 2(1):167–175, 2010.
- [35] H. Wang. Functions on adjacent vertex degrees of trees with given degree sequence. *Cent. Eur. J. Math.*, 12(11):1656–1663, 2014.
- [36] L. Ye and X. Yuan. On the minimal energy of trees with a given number of pendent vertices. *MATCH Commun. Math. Comput. Chem.*, 57(1):193–201, 2007.
- [37] A. Yu and X. Lv. Minimum energy on trees with k pendent vertices. *Linear Algebra Appl.*, 418(2-3):625–633, 2006.
- [38] A. Yu and X. Lv. The Merrifield–Simmons indices and Hosoya indices of trees with k pendant vertices. *J. Math. Chem*, 41:33–43, 2007.
- [39] J. Zhang, G.-J. Zhang, H. Wang, and X.-D. Zhang. Extremal trees with respect to the Steiner Wiener index. *Discrete Math. Algorithms Appl.*, 11(6):article 1950067, 16pp., 2019.
- [40] X.-D. Zhang. The Laplacian spectral radii of trees with given degree sequences. *Discrete Math.*, 308(15):3143–3150, 2008.
- [41] X.-D. Zhang, Q.-Y. Xiang, L.-Q. Xu, and R.-Y. Pan. The Wiener index of trees with given degree sequences. *MATCH Commun. Math. Comput. Chem*, 60:623–644, 2008.
- [42] X.-M. Zhang, X.-D. Zhang, D. Gray, and H. Wang. The number of subtrees of trees with given degree sequence. *J. Graph Theory*, 73(3):280–295, 2013.
- [43] B. Zhou and I. Gutman. A connection between ordinary and Laplacian spectra of bipartite graphs. *Linear Multilinear Algebra*, 56:305–310, 2008.