

# On well-connected sets of strings

Peter Frankl\*      János Pach†

Rényi Institute

POB 127 Budapest, 1364 Hungary

`peter.frankl@gmail.com`

`pach@cims.nyu.edu`

Submitted: Mar 8, 2021; Accepted: Feb 7, 2022; Published: Mar 25, 2022

© The authors. Released under the CC BY-ND license (International 4.0).

## Abstract

Given  $n$  sets  $X_1, \dots, X_n$ , we call the elements of  $S = X_1 \times \dots \times X_n$  *strings*. A nonempty set of strings  $W \subseteq S$  is said to be *well-connected* if for every  $v \in W$  and for every  $i$  ( $1 \leq i \leq n$ ), there is another element  $v' \in W$  which differs from  $v$  only in its  $i$ th coordinate. We prove a conjecture of Yaokun Wu and Yanzhen Xiong by showing that every set of more than  $\prod_{i=1}^n |X_i| - \prod_{i=1}^n (|X_i| - 1)$  strings has a well-connected subset. This bound is tight.

**Mathematics Subject Classifications:** 05C88, 05C89

## 1 Introduction

Let  $X_1, \dots, X_n$  be pairwise disjoint sets with  $|X_i| = d_i > 1$  for  $1 \leq i \leq n$ . Let

$$S = X_1 \times \dots \times X_n = \{(x_1, \dots, x_n) : x_i \in X_i \text{ for every } i \in [n]\}$$

be the set of *strings*  $x = (x_1, \dots, x_n)$ , where  $x_i$  is called the  $i$ th coordinate of  $x$  and  $[n] = \{1, \dots, n\}$ .

A subset  $W \subseteq S$  is called *well-connected* if for every  $x \in W$  and for every  $i \in [n]$ , there is another element  $x' \in W$  which differs from  $x$  only in its  $i$ th coordinate. That is,  $x'_j \neq x_j$  if and only if  $j = i$ .

The following statement was conjectured by Yaokun Wu and Yanzhen Xiong [4].

---

\*Partially supported by the Ministry of Education and Science of the Russian Federation in the framework of MegaGrant 075-15-2019-1926.

†Partially supported by ERC Advanced Grant GeoScope, NKFIH (Hungarian National Research, Development and Innovation Office) grant K-131529, and by the Ministry of Education and Science of the Russian Federation in the framework of MegaGrant 075-15-2019-1926.

**Theorem 1.** Let  $T$  be a subset of  $S = X_1 \times \cdots \times X_n$  with  $|X_i| = d_i > 1$  for every  $i \in [n]$ . If

$$|T| > \prod_{i=1}^n d_i - \prod_{i=1}^n (d_i - 1),$$

then  $T$  has a nonempty well-connected subset. This bound cannot be improved.

To see the tightness of the theorem, fix an element  $y_i$  in each  $X_i$  and let  $X'_i = X_i \setminus \{y_i\}$ . We claim that the set of strings

$$T_0 = (X_1 \times \cdots \times X_n) \setminus (X'_1 \times \cdots \times X'_n) \tag{1}$$

does not have any nonempty well-connected subset. Suppose for contradiction that there is such a subset  $W \subseteq T_0$ , and let  $x = (x_1, \dots, x_n)$  be an element of  $W$  with the minimum number of coordinates  $i$  for which  $x_i = y_i$  holds. Obviously, this minimum is positive, otherwise  $x \notin T_0$ . Pick an integer  $k$  with  $x_k = y_k$ . Using the assumption that  $W$  is well-connected, we obtain that there exists  $x' \in W$  that differs from  $x$  only in its  $k$ th coordinate. However, then  $x'$  would have one fewer coordinates with  $x_i = y_i$  than  $x$  does, contradicting the minimality of  $x$ .

In the next section, we establish a result somewhat stronger than Theorem 1: we prove that under the conditions of Theorem 1,  $T$  also has a subset  $W$  such that for every  $x \in W$  and  $i \in [n]$ , the number of elements  $x' \in W$  which differ from  $x$  only in its  $i$ th coordinate is *odd* (see Theorem 6). In Section 3, we present a self-contained argument which proves this stronger statement.

Shortly after learning about our proof of the conjecture of Wu and Xiong, another proof was found by Chengyang Qian.

## 2 Exact sequence of maps

In this section, we introduce the necessary definitions and terminology, and we apply a basic topological property of simplicial complexes to establish Theorem 1. We will assume throughout, without loss of generality, that the sets  $X_i$  are pairwise disjoint.

For every  $k$  ( $0 \leq k \leq n$ ), let

$$S_k = \{A \subseteq X_1 \cup \dots \cup X_n : |A| = k \text{ and } |A \cap X_i| \leq 1 \text{ for every } i\}.$$

Clearly, we have  $|S_n| = |S| = \prod_{i=1}^n |X_i|$ . With a slight abuse of notation, we identify  $S_n$  with  $S$ . The set system  $\cup_{k=0}^n S_k$  is an *abstract simplicial complex*, that is, for each of its elements  $A$ , every subset of  $A$  also belongs to  $\cup_{k=0}^n S_k$ . This simplicial complex has a geometric realization in  $\mathbb{R}^{2n-1}$ , where every element  $A$  is represented by an  $(|A| - 1)$ -dimensional simplex. (See [1], part II, Section 9 or [3], Section 1.5. Note that not all textbooks consider the empty set a  $-1$ -dimensional simplex, but we do.)

Assign to each  $A \in S_k$  a different symbol  $v_A$ , and define  $V_k$  as the family of all formal sums of these symbols with coefficients 0 or 1. Then

$$V_k = \left\{ \sum_{A \in S_k} \lambda_A v_A : \lambda_A = 0 \text{ or } 1 \right\}$$

can be regarded as a vector space over  $\text{GF}(2)$  whose dimension satisfies

$$\dim V_k = |S_k| = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} d_{j_1} d_{j_2} \cdot \dots \cdot d_{j_k}. \quad (2)$$

We use the standard definition of the *boundary operations*  $\partial_k$ . (See [2], Section 2.1.) Informally, the boundary of each  $(k - 1)$ -dimensional simplex that corresponds to a member  $A \in S_k$  consists of all  $(k - 2)$ -dimensional simplices corresponding to  $(k - 1)$ -element subsets  $B \subset A$ . This definition naturally extends to any collection (“chain”) of  $(k - 1)$ -dimensional simplices that correspond to members of  $S_k$ , with multiplicities taken modulo 2.

**Definition 2.** Let  $\partial_0 : V_0 \rightarrow 0$ . For every  $k \in [n]$  and every  $A \in S_k$ , let

$$\partial_k(v_A) = \sum_{\substack{B \subset A \\ |B|=k-1}} v_B.$$

Extend this map to a homomorphism  $\partial_k : V_k \rightarrow V_{k-1}$  by setting

$$\partial_k\left(\sum_{A \in S_k} \lambda_A v_A\right) = \sum_{A \in S_k} \lambda_A \partial_k(v_A),$$

where the sum is taken over  $\text{GF}(2)$ .

Let  $\ker(\partial_k) \subseteq V_k$  and  $\text{im}(\partial_k) \subseteq V_{k-1}$  denote the *kernel* and the *image* of  $\partial_k$ , respectively.

Our proof is based on the following lemma.

**Lemma 3.** *The sequence of homomorphisms  $V_n \xrightarrow{\partial_n} V_{n-1} \xrightarrow{\partial_{n-1}} \dots \xrightarrow{\partial_1} V_0 \xrightarrow{\partial_0} 0$  is an exact sequence, i.e.,  $\text{im}(\partial_k) = \ker(\partial_{k-1})$  holds for every  $k \in [n]$ .*

**Proof.** Before proving the statement, we show that  $\text{im}(\partial_k) \subseteq \ker(\partial_{k-1})$  for every  $k \in [n]$ . The statement is obviously true for  $k = 1$ . If  $k \geq 2$ , then for every  $A \in S_k$ , we have

$$\partial_{k-1} \partial_k v_A = \sum_{\substack{B \subset A \\ |B|=k-1}} \sum_{\substack{C \subset B \\ |C|=k-2}} v_C = \sum_{\substack{C \subset A \\ |C|=k-2}} 2v_C = 0.$$

Thus,  $\partial_{k-1} \partial_k(v) = 0$  for every  $v \in V_k$ , as claimed. In fact, the containment  $\text{im}(\partial_k) \subseteq \ker(\partial_{k-1})$  holds for *every* simplicial complex.

We prove that in our case, all the above containments hold with equality. For every  $i \in [n]$ , let  $K_i$  denote the 0-dimensional abstract simplicial complex consisting of the 1-element subsets of  $X_i$  and the empty set. Consider now their *join*  $K = K_1 * \dots * K_n$ ; see [2], Chapter 0. By definition,  $K$  is the same as the simplicial complex  $\cup_{i=0}^n S_i$ .

Let  $j \geq -1$  be an integer. We need three well-known properties of the notion of *j-connectedness* of complexes; see Proposition 4.4.3 in [3].

- (i) A complex is  $-1$ -connected if and only if it contains a nonempty simplex.

(ii) If  $K_1$  is  $a$ -connected and  $K_2$  is  $b$ -connected, then their join  $K_1 * K_2$  is  $(a + b - 2)$ -connected.

(iii) If a complex is  $j$ -connected, then  $\text{im}(\partial_k) = \ker(\partial_{k-1})$  holds for every  $k$ ,  $1 \leq k \leq j+2$ .

In our case, each  $X_i$  is nonempty, hence, by property (i), each  $K_i$  is  $-1$ -connected. By repeated application of (ii), we obtain that  $K = K_1 * \dots * K_n$  is  $(n - 2)$ -connected. In view of (iii), this implies that  $\text{im}(\partial_k) = \ker(\partial_{k-1})$  for every  $k \in [n]$ , as required.  $\square$

**Corollary 4.** For every  $k$  ( $0 \leq k \leq n$ ), we have  $\dim \ker(\partial_k) = \sum_{i=0}^k (-1)^{k-i} \dim V_i$ .

**Proof.** By induction on  $k$ . According to the Rank Nullity Theorem, we have

$$\dim V_i = \dim \ker(\partial_i) + \dim \text{im}(\partial_i), \tag{3}$$

for every  $i \leq n$ . Since  $\dim V_0 = 1$  and  $\dim \text{im}(\partial_0) = \dim 0 = 0$ , the corollary is true for  $k = 0$ .

Assume we have already verified it for some  $k < n$ . To show that it is also true for  $k + 1$ , we use that  $\dim \text{im}(\partial_{k+1}) = \dim \ker(\partial_k)$ , by Lemma 3. Plugging this into (3) with  $i = k + 1$ , we obtain

$$\dim V_{k+1} = \dim \ker(\partial_{k+1}) + \dim \ker(\partial_k).$$

Hence, using the induction hypothesis, we have

$$\begin{aligned} \dim \ker(\partial_{k+1}) &= \dim V_{k+1} - \dim \ker(\partial_k) \\ &= \dim V_{k+1} - \sum_{i=0}^k (-1)^{k-i} \dim V_i = \sum_{i=0}^{k+1} (-1)^{k+1-i} \dim V_i, \end{aligned}$$

as required.  $\square$

By (2), we know the value of  $\dim V_i$  for every  $i$ . Therefore, Corollary 4 enables us to compute  $\dim \ker(\partial_n)$  and, hence,  $\dim V_n - \dim \ker(\partial_n)$ .

**Corollary 5.** We have

$$\dim V_n - \dim \ker(\partial_n) = \prod_{i=1}^n d_i - \prod_{i=1}^n (d_i - 1).$$

**Proof.** From Corollary 4, we get

$$\dim V_n - \dim \ker(\partial_n) = \sum_{i=0}^{n-1} (-1)^{n-1-i} \dim V_i.$$

Using (2) and the fact that  $\dim V_0 = 1$ , this is further equal to

$$\sum_{i=1}^{n-1} (-1)^{n-1-i} \sum_{1 \leq j_1 < j_2 < \dots < j_i \leq n} d_{j_1} d_{j_2} \dots d_{j_i} + (-1)^{n-1} = \prod_{i=1}^n d_i - \prod_{i=1}^n (d_i - 1). \quad \square$$

Now we are in a position to establish the following statement, which is somewhat stronger than Theorem 1.

**Theorem 6.** *Let  $T$  be a subset of  $S = X_1 \times \cdots \times X_n$  with  $|X_i| = d_i > 1$  for every  $i \in [n]$ . If*

$$|T| > \prod_{i=1}^n d_i - \prod_{i=1}^n (d_i - 1),$$

*then there is a nonempty subset  $W \subseteq T$  with the property that for every  $x \in W$  and  $i \in [n]$ , the number of elements  $x' \in W$  which differ from  $x$  only in their  $i$ th coordinate is odd. This bound cannot be improved.*

**Proof.** The tightness of the bound follows from the tightness of Theorem 1 shown at the end of the Introduction.

Let  $T$  be a system of strings of length  $n$  satisfying the conditions of the theorem. Using the notation introduced at the beginning of this section, let

$$V(T) = \left\{ \sum_{A \in T} \lambda_A v_A : \lambda_A = 0 \text{ or } 1 \right\}.$$

Then  $V(T)$  can be regarded as a linear subspace of  $V_n$  with  $\dim V(T) = |T|$ . Comparing the size of  $T$  with the value of  $\dim V_n - \dim \ker(\partial_n)$  given by Corollary 5, we obtain that there is a nonzero vector  $v = \sum_{A \in T} \lambda_A v_A$  that belongs to  $V(T) \cap \ker(\partial_n)$ . Let  $W = \{A \in T : \lambda_A = 1\}$ . Then we have

$$0 = \partial_n(v) = \sum_{A \in W} \partial_n(v_A) = \sum_{A \in W} \sum_{\substack{B \subset A \\ |B|=n-1}} v_B = \sum_{\substack{B \subset [n] \\ |B|=n-1}} |\{A \in W : A \supseteq B\}| v_B.$$

Thus, for each  $B$ , the coefficient of  $v_B$  is *even*. This means that the set of strings  $W \subset T$  meets the requirements of the theorem.  $\square$

### 3 Direct proof of Theorem 6

In this section, we prove Corollary 5 and, hence, Theorem 6 directly, without using Lemma 3.

As in the Introduction, fix an element  $y_i \in X_i$  and let  $X'_i = X_i \setminus \{y_i\}$ , for every  $i \in [n]$ . Defining  $T_0$  as in (1), we have that  $|T_0| = \prod_{i=1}^n d_i - \prod_{i=1}^n (d_i - 1)$ .

Suppose that  $|T| > |T_0|$ . To prove Corollary 5, it is sufficient to show that there exists a nonzero vector  $v = \sum_{A \in T} \lambda_A v_A$  with suitable coefficients  $\lambda_A \in \{0, 1\}$  such that  $v \in \ker(\partial_n)$ , *i.e.*, we have  $\partial_n v = \sum_{A \in T} \lambda_A (\partial_n v_A) = 0$ . Thus, it is enough to establish the following statement.

**Lemma 7.** *Let  $T$  be a subset of  $S = X_1 \times \cdots \times X_n$  with  $|X_i| > 1$  for every  $i \in [n]$ . If  $|T| > |T_0|$ , then the set of vectors  $\{\partial_n v_A : A \in T\}$  is linearly dependent over  $\text{GF}(2)$ .*

**Proof.** First, we show that the set of vectors  $\{\partial_n v_A : A \in T_0\}$  is linearly independent. Suppose, for a contradiction, that there is a nonempty subset  $W \subset T_0$  such that  $\sum_{A \in W} \partial_n v_A = 0$ . Pick an element  $A = \{x_1, \dots, x_n\}$  of  $W$  for which the number of coordinates  $i$  with  $x_i = y_i$  is as small as possible. By the definition of  $T_0$ , there is at least one such coordinate  $x_k = y_k$ . In view of Definition 2, one of the terms of the formal sum  $\partial_n v_A$  is  $v_B$  with  $B = A \setminus \{y_k\}$ , and this term cannot be canceled out by a term of  $\partial_n v_{A'}$  for any other  $A' \in W$ , because in this case  $A'$  would have fewer coordinates that are equal to some  $y_i$  than  $A$  does. Hence,  $\sum_{A \in W} \partial_n v_A \neq 0$ , contradicting our assumption.

It remains to prove that  $\{\partial_n v_A : A \in T_0\}$  is a *base* of  $\text{im}(\partial_n)$ , that is, there exists no set of strings  $T \supset T_0$  with  $|T| > |T_0|$  such that the set of vectors  $\{\partial_n v_A : A \in T\}$  is linearly independent over  $\text{GF}(2)$ .

To see this, consider any string  $C = \{z_1, \dots, z_n\} \in S \setminus T_0$ . Since  $C \notin T_0$ , we have  $z_i \neq y_i$  for every  $i$ . Define  $T(C)$  as the set of all strings  $A = \{x_1, \dots, x_n\} \in S$  whose every coordinate  $x_i$  is either  $y_i$  or  $z_i$ . Then we have  $\sum_{A \in T(C)} \partial_n v_A = 0$ . As we have  $T(C) \subseteq T_0 \cup \{C\}$ , this means that the set of vectors  $\{\partial_n v_A : A \in T_0 \cup \{C\}\}$  is linearly dependent over  $\text{GF}(2)$ . This completes the proof of the lemma and, hence, of Theorem 6.  $\square$

## Acknowledgements

We thank Gábor Tardos and an anonymous referee for several helpful suggestions.

## References

- [1] A. Björner. Topological Methods. *Handbook of Combinatorics*, R. Graham, M. Grötschel, L. Lovász (eds.), Vol. 2, 1995, 1819–1872.
- [2] A. Hatcher. Algebraic Topology. Cambridge University Press, Cambridge, 2001.
- [3] J. Matoušek. Using the Borsuk-Ulam Theorem. Springer, Berlin, 2003.
- [4] Y. Wu and Y. Xiong. Sparse  $(0, 1)$  arrays and tree-like partition systems. Manuscript, 2020.