

# Every Steiner Triple System Contains an Almost Spanning $d$ -Ary Hypertree

Andrii Arman

Department of Mathematics  
Emory University  
Atlanta, GA, 30322, U.S.A.  
andrew0arman@gmail.com

Vojtěch Rödl\*

Department of Mathematics  
Emory University  
Atlanta, GA, 30322, U.S.A.  
rodl@mathcs.emory.edu

Marcelo Tadeu Sales

Department of Mathematics  
Emory University  
Atlanta, GA, 30322, U.S.A.

marcelo.tadeu.sales@emory.edu

Submitted: May 23, 2021; Accepted: May 30, 2022; Published: Jul 15, 2022

© The authors. Released under the CC BY-ND license (International 4.0).

## Abstract

In this paper we make a partial progress on the following conjecture: for every  $\mu > 0$  and a large enough  $n$ , every Steiner triple system  $S$  on at least  $(1 + \mu)n$  vertices contains every hypertree  $T$  on  $n$  vertices. We prove that the conjecture holds if  $T$  is a perfect  $d$ -ary hypertree.

**Mathematics Subject Classifications:** 05B07, 05C65

## 1 Introduction

In this paper we study the following conjecture, raised by the second author and Bradley Elliot [4].

**Conjecture 1.** Given  $\mu > 0$  there is  $n_0$ , such that for any  $n \geq n_0$ , any hypertree  $T$  on  $n$  vertices and any Steiner triple system  $S$  on at least  $(1 + \mu)n$  vertices,  $S$  contains  $T$  as a subhypergraph.

---

\*The second author was supported by NSF grant DMS 1764385

Note that any hypertree  $T$  can be embedded into any Steiner triple system  $S$ , provided  $|V(T)| \leq \frac{1}{2}(|V(S)| + 3)$ . The problem becomes more interesting if the size of the tree is larger. In [3] (see also [4] Section 5) Conjecture 1 was verified for some special classes of hypertrees. In this paper we verify the conjecture for another class of hypertrees – perfect  $d$ -ary hypertrees.

**Definition 2.** A perfect  $d$ -ary hypertree  $T$  of height  $h$  is a hypertree  $T$  with  $V(T) = \bigcup_{i=0}^h V_i$ , where  $|V_i| = (2d)^i$  for all  $i \in [0, h]$ , such that for every  $i \in [0, h - 1]$  and  $v \in V_i$  there are  $2d$  vertices  $\{u_1, \dots, u_{2d}\} \subseteq V_{i+1}$  such that  $\{v, u_{2j-1}, u_{2j}\}$  is a hyperedge of  $T$  for all  $j \in [d]$ .

In other words,  $T$  is a perfect  $d$ -ary hypertree if every non-leaf vertex has  $2d$  children (or a forward degree  $d$ ). The main result of this paper is the following theorem.

**Theorem 3.** *For any real  $\mu > 0$  there is  $n_0$  such that the following holds for all  $n \geq n_0$  and any positive integer  $d$ . If  $S$  is a Steiner triple system with at least  $(1 + \mu)n$  vertices and  $T$  is a perfect  $d$ -ary hypertree on at most  $n$  vertices, then  $T \subseteq S$ .*

## 2 Preliminaries

For a positive integer  $k$  let  $[k] = \{1, \dots, k\}$  and for positive integers  $k < \ell$  let  $[k, \ell] = \{k, k + 1, \dots, \ell\}$ . We write  $x = y \pm z$  if  $x \in [y - z, y + z]$ . We write  $A = B \sqcup C$  if  $A$  is a union of disjoint sets  $B$  and  $C$ .

A hypertree is a connected, simple (linear) 3-uniform hypergraph in which every two vertices are joined by a unique path. A hyperstar  $S$  of size  $a$  centered at  $v$  is a hypertree on the vertex set  $v, v_1, v_2, \dots, v_{2a}$  with the edge set  $E(S) = \{\{v, v_{2i-1}, v_{2i}\} : i \in [a]\}$ . A Steiner triple system (STS) is a 3-uniform hypergraph in which every pair of vertices is contained in exactly one edge.

If  $H$  is a hypergraph and  $v \in V(H)$ , then  $d_H(v)$  (or  $d(v)$  when the context is clear) is the degree of a vertex  $v$  in  $H$ .

For  $V(H) = X \sqcup Y$  we denote by  $H[X, Y]$  the spanning subhypergraph of  $H$  with

$$E(H[X, Y]) = \{e \in E(H) : |e \cap X| = 1, |e \cap Y| = 2\}.$$

The proof of Theorem 3 relies on the application of an existence of an almost perfect matching in an almost regular 3-uniform simple hypergraph. We will use two versions of such results. In the first version the degrees of a small proportion of vertices are allowed to deviate from the average degree. We will use Theorem 4.7.1 from [2] (see [5] and [8] for earlier versions).

**Theorem 4.** *For any  $\delta > 0$  and  $k > 0$  there exists  $\varepsilon$  and  $D_0$  such that the following holds. Let  $H$  be a 3-uniform simple hypergraph on  $N$  vertices and  $D \geq D_0$  be such that*

(i) *for all but at most  $\varepsilon N$  vertices  $x$  of  $H$  the degree of  $x$*

$$d(x) = (1 \pm \varepsilon)D.$$

(ii) for all  $x \in V(H)$  we have

$$d(x) \leq kD.$$

Then  $H$  contains a matching on at least  $N(1 - \delta)$  vertices.

A second version is a result by Alon, Kim and Spencer [1], where under the assumption that all degrees are concentrated near the average, a stronger conclusion may be drawn. We use a version of this result as stated in [7]\*.

**Theorem 5.** *For any  $K > 0$  there exists  $D_0$  such that the following holds. Let  $H$  be a 3-uniform simple hypergraph on  $N$  vertices and  $D \geq D_0$  be such that  $\deg(x) = D \pm K\sqrt{D \ln D}$  for all  $x \in V(H)$ . Then  $H$  contains a matching on  $N - O(ND^{-1/2} \ln^{3/2} D)$  vertices.*

Here the constant in  $O()$ -notation is depending on  $K$  only and is independent of  $N$  and  $D$ .

In Lemma 9 we consider a random partition of the vertex set of Steiner triples system  $S$  and heavily use the following version of Chernoff's bound (this is Corollary 2.3 of Janson, Luczak, Rucinski [6]).

**Theorem 6.** *Let  $X \sim \text{Bi}(n, p)$  be a binomial random variable with the expectation  $\mu$ , then for  $t \leq \frac{3}{2}\mu$*

$$\mathbb{P}(|X - \mu| > t) \leq 2e^{-t^2/(3\mu)}.$$

In particular for  $K \leq \frac{3}{2}\sqrt{\frac{\mu}{\ln \mu}}$  and  $t = K\sqrt{\mu \ln \mu}$

$$\mathbb{P}(|X - \mu| > K\sqrt{\mu \ln \mu}) \leq 2(\mu)^{-K^2/3}. \tag{1}$$

If  $\varepsilon > 0$  is fixed and  $\mu > \mu(\varepsilon)$ , then

$$\mathbb{P}(X = (1 \pm \varepsilon)\mu) = 1 - e^{-\Omega(\mu)}. \tag{2}$$

### 3 Proof of Theorem 3

#### 3.1 Proof Idea

Assume that  $S$  is an STS on at least  $(1 + \mu)n$  vertices. We will choose a small constant  $\varepsilon \ll \mu$ .

Let  $T$  be a perfect  $d$ -ary tree on at most  $n$  vertices with levels  $V_i$ ,  $i \in [0, h]$  and let  $i_0 = \max\{i, |V_i| \leq \varepsilon n\}$ . Let  $T_0$  be a subhypertree of  $T$  induced on  $\bigcup_{i=0}^{i_0} V_i$ . To simplify our notation we set  $t = h - i_0$  and for all  $i \in [0, t]$  we set  $L_i = V_{i_0+i}$ . Our goal is to find  $L \subset V(S)$  with  $|L| = |V(T)|$  such that  $S[L]$ , the subhypergraph induced by  $L$ , contains a spanning copy of  $T$ . In particular we would find such an  $L$ , level by level, first embedding  $T_0$ , and then  $L_1, \dots, L_t$ .

To start we consider a partition  $\mathcal{P} = \{C_0, \dots, C_t, R\}$  of  $V(S)$  with "random-like" properties (see Lemma 9 for the description of  $\mathcal{P}$ ) in the following way:

---

\*We refer to Theorem 3 from that paper. There is a typo in the conclusion part of that theorem, where instead of  $O(ND^{1/2} \ln^{3/2} D)$  there should be  $O(ND^{-1/2} \ln^{3/2} D)$

- The first few levels of  $T$ , constituting  $T_0$  with the subset of leafs  $L_0$ , will be embedded greedily into  $S[C_0]$ .
- Then Lemma 7 and Lemma 8 will be used to find star forests in  $S[L_0, C_1]$  and  $S[C_{i-1}, C_i]$  for  $i \in [2, t]$ . The union of these star forests will establish embedding of almost all vertices of  $T$  (see Claim 12).
- Finally, reservoir vertices  $R$  will be used to complete the embedding (see Claim 13).

### 3.2 Auxiliary Lemmas

We start our proof with the following Lemma that will later allow us to verify that  $S[L_0, C_1]$  contains an almost perfect packing of hyperstars, each of size at most  $d$ , centered at the vertices of  $L_0$

**Lemma 7.** *For any positive real  $\delta$ ,  $k > 1$  there are  $\varepsilon > 0$  and  $D_0$  such that the following holds for all  $D \geq D_0$  and all positive integers  $d$ . Let  $G = (V, E)$  be a 3-uniform simple hypergraph on  $N$  vertices such that  $V = X \sqcup Y$  and*

(i) *for all  $e \in E$ ,  $|e \cap X| = 1$  and  $|e \cap Y| = 2$ .*

(ii) *for all vertices  $v \in X$  we have*

$$d(v) = dD(1 \pm \varepsilon),$$

*and for all but at most  $\varepsilon N$  vertices  $v \in Y$  we have*

$$d(v) = D(1 \pm \varepsilon).$$

(iii)  *$d(v) \leq kD$  for all  $v \in Y$ .*

*Then  $G$  contains a packing of hyperstars of size at most  $d$  centered at vertices of  $X$  that covers all but at most  $\delta N$  vertices.*

*Proof.* For given  $\delta$  and  $k$  set  $\delta_{2.1} = 2\delta/3$  and  $k_{2.1} = k$ . With these parameters as an input, Theorem 4 yields  $\varepsilon_{2.1}$  and  $D_0$ . Set  $\varepsilon = \varepsilon_{2.1}/2$  and note that if Theorem 4 holds for some  $\varepsilon_{2.1}$  and  $D_0$  then it also holds for smaller values of  $\varepsilon_{2.1}$  and larger values of  $D_0$ . Therefore we may assume that  $\varepsilon$  is sufficiently small with respect to  $\delta$  and  $k$ .

Let  $G$  that satisfies conditions (i)–(iii) be given. We start with constructing an auxiliary hypergraph  $H$  that is obtained from  $G$  by repeating the following splitting procedure for each vertex  $v \in X$ : split the hyperedges incident to  $v$  into  $d$  disjoint groups, each of size  $D(1 \pm 2\varepsilon) = D(1 \pm \varepsilon_{2.1})$ , and then replace  $v$  with new vertices  $v_1, \dots, v_d$  and each hyperedge  $\{v, u, w\}$  that belongs to the group  $j$  with a hyperedge  $\{v_j, u, w\}$ .

First, we show that  $|V(H)| \leq \frac{3}{2}N$ . Note that due to conditions (i)–(iii)

$$|E(G)| \sim |X|dD \sim \frac{1}{2}|Y|D.$$

Provided  $\varepsilon$  is small enough and  $N$  is large enough compared to  $\delta$  we can guarantee  $|X| \leq \frac{N}{2d}$ . Finally,  $|V(H)| \leq d|X| + |Y|$  by the construction of  $H$ , so

$$N < |V(H)| \leq (d-1)|X| + |X| + |Y| \leq (d-1)|X| + N \leq \frac{3}{2}N.$$

Hypergraph  $H$  satisfies the assumptions of Theorem 4 with parameters  $\delta_4 = \frac{2}{3}\delta$ ,  $k_{2,1} = k$ ,  $\varepsilon_{2,1} = 2\varepsilon$ ,  $D_{2,1} = D$  and  $N_{2,1} = |V(H)|$ . Indeed, all of the vertices in  $H$  still have degrees at most  $k_{2,1}D_{2,1}$ , and for all but at most  $\varepsilon_{2,1}N_{2,1}$  vertices we have  $d_H(v) = D_{2,1}(1 \pm \varepsilon_{2,1})$ . Therefore, there is a matching  $M$  in  $H$  that omits at most  $\delta_4 N_{2,1} \leq \delta N$  vertices.

Now, the matching  $M$  in  $H$  corresponds to a collection of hyperstars  $S_1, \dots, S_k$  in  $G$  with centers at vertices of  $X$ , and such that the size of each  $S_i$  is at most  $d$ . Indeed, recall that during the construction of  $H$  some vertices  $v \in X$  were replaced by  $d$  vertices  $v_1, \dots, v_d$ , hyperedges incident to  $v$  were split into  $d$  almost equal in size disjoint groups, and then each hyperedge  $\{v, u, w\}$  in  $j$ -th group was replaced with  $\{v_j, u, w\}$ . Consequently, a matching in  $H$  that covers some vertices  $v_i$  gives a rise to a hyperstar centered at  $v$  of size at most  $d$  in  $G$ .

Moreover since  $M$  in  $H$  omits at most  $\delta N$  vertices, the union of hyperstars  $S_1, \dots, S_k$  also omits at most  $\delta N$  vertices.  $\square$

The following Lemma will later allow us to verify that for all  $i \in [t-1]$  the subhypergraph  $S[C_i, C_{i+1}]$  contains an almost perfect packing of hyperstars, each of size at most  $d$ , centered at the vertices of  $C_i$ .

**Lemma 8.** *For any positive real  $K$  there is  $D_0$  such that the following holds for all  $D \geq D_0$ ,  $\Delta = K\sqrt{D \ln D}$  and any positive integer  $d$ . Let  $G = (V, E)$  be a 3-uniform simple hypergraph on  $N$  vertices such that  $V = X \sqcup Y$  and*

(i) *for all  $e \in E$ ,  $|e \cap X| = 1$  and  $|e \cap Y| = 2$ .*

(ii)  *$d(v) = d(D \pm \Delta)$  for all  $v \in X$  and  $d(v) = D \pm \Delta$  for all  $v \in Y$ .*

*Then  $G$  contains a packing of hyperstars of size at most  $d$  centered at the vertices of  $X$  that covers all but at most  $O(ND^{-1/2} \ln^{3/2} D)$  vertices.*

Here the constant in  $O()$ -notation depends on  $K$  only.

*Proof.* The proof is almost identical to the proof of Lemma 7. For a given  $K$  let  $D_0$  be the number guaranteed by Theorem 5 with  $2K$  as input.

Let  $G$  that satisfies conditions (i),(ii) be given. We start with constructing an auxiliary hypergraph  $H$  that is obtained from  $G$  by splitting every vertex  $v \in X$  into  $d$  new vertices  $v_1, \dots, v_d$  that have degrees  $D \pm 2\Delta$ .

First, we will show that  $|V(H)| = \Theta(N)$ . Note that due to conditions (i) and (ii), we have

$$\frac{|Y|(D \pm \Delta)}{2} = |E(G)| = |X|d \left( D \pm \frac{\Delta}{2} \right).$$

In particular,

$$|Y| = |X|d \left( \frac{2D \pm \Delta}{D \pm \Delta} \right).$$

As  $|X| + |Y| = N$  we have that  $|Y| = \Theta(N)$  and hence  $d|X| = \Theta(N)$ . Then  $|V(H)| = d|X| + |Y|$  by construction of  $H$ , so  $|V(H)| = \Theta(N)$  as well.

Hypergraph  $H$  satisfies the assumptions of the Theorem 5 with parameters  $2K$  and  $D \geq D_0$ . Therefore, there is a matching  $M$  in  $H$  that omits at most  $O(ND^{-1/2} \ln^{3/2} D)$  vertices.

Now, matching  $M$  in  $H$  corresponds to a collection of hyperstars  $S_1, \dots, S_k$  in  $G$  with centers at the vertices of  $X$ . Each hyperstar  $S_i$  contains at most  $d$  hyperedges and hyperstars  $S_1, \dots, S_k$  cover all but at most  $O(ND^{-1/2} \ln^{3/2} D)$  vertices of  $G$ , which finishes the proof.  $\square$

### 3.3 Formal Proof

We start by defining constants, proving some useful inequalities and proving Lemma 9.

Let  $S$  be a Steiner triple system on  $m \geq (1 + \mu)n$  vertices and let  $T$  be the largest perfect  $d$ -ary hypertree with at most  $n$  vertices. Our goal is to show that  $T \subset S$ .

We make few trivial observations. First, if  $d > \sqrt{n}$  and  $T$  is perfect  $d$ -ary hypertree with  $|V(T)| \leq n$ , then  $T$  is just a hyperstar which  $S$  clearly contains. Second, if  $m > 2n$ , then  $T$  can be found in  $S$  greedily. Finally, if Theorem 3 holds for some value of  $\mu$ , then Theorem 3 holds for larger values of  $\mu$ . Hence we may assume without loss of generality that  $d \leq \sqrt{n}$ ,  $m \leq 2n$  and  $\mu \leq \frac{1}{4}$ .

**Constants.** We will choose new constant  $\varepsilon < \delta < \rho < \mu$  independent of  $m, n$ :

$$\rho = \left( \frac{3\mu - \mu^2}{8(1 + \mu)} \right)^2, \quad \delta = \frac{(1 + \mu)\rho}{20}. \quad (3)$$

Let  $\varepsilon_{3.1}$  be a constant guaranteed by Lemma 7 with  $\delta$  and  $k = 2$  as an input. We choose  $\varepsilon$  to be small enough, in particular we want

$$\varepsilon < \min\{\delta^2, (\mu/16)^2, (\varepsilon_{3.1})^{10}, 1/10^{100}\}. \quad (4)$$

**Properties of  $T$ .** Here we define the levels of  $T$  and prove some useful inequalities. Recall that  $V_i$ ,  $i \in [0, h]$  denoted the levels of  $T$ . For  $i_0 = \max\{i, |V_i| \leq \varepsilon n\}$  let  $T_0$  be a subhypertree of  $T$  induced on  $\bigcup_{i=0}^{i_0} V_i$ . To simplify our notation we also set  $t = h - i_0$  and for all  $i \in [0, t]$ ,  $L_i = V_{i_0+i}$  and  $\ell_i = |L_i|$ . Then we have for  $i \in [t]$

$$\ell_i = (2d)^i \ell_0, \quad \varepsilon n \geq \ell_0 > \frac{\varepsilon}{2d} n. \quad (5)$$

Since  $n \geq \ell_t$ , (5) implies  $n \geq (2d)^{t-1} \varepsilon n$ , and consequently

$$t \leq 1 + \frac{\log \frac{1}{\varepsilon}}{\log(2d)} \leq 1 + \log \frac{1}{\varepsilon}. \quad (6)$$

Finally,  $T_0 = \bigcup_{i=0}^{i_0} V_i$ , where  $|V_{i_0}| = (2d)^{i_0} = \ell_0$ , so

$$|V(T_0)| = \frac{(2d)^{i_0+1} - 1}{2d - 1} \leq \frac{(2d)\ell_0}{2d - 1} \stackrel{(5)}{\leq} 2\varepsilon n. \quad (7)$$

**Partition Lemma.**

For a given Steiner triple system  $S$  with  $m$  vertices our goal will be to find a partition  $\mathcal{P} = \{C_1, \dots, C_t, R\}$  of  $V(S)$  so that  $S[C_0]$  contains a copy of  $T_0$  (and  $L_0$ ), sets  $C_1, \dots, C_t$  are the ‘‘candidates’’ for levels  $L_1, \dots, L_t$  of  $T$ , and  $R$  is a reservoir. Such a partition will be guaranteed by Lemma 9.

In the proof we will consider a random partition  $\mathcal{P}$ , where each vertex  $v \in V(S)$  ends up in  $C_i$  with probability  $p_i$  and in  $R$  with probability  $\gamma$  independently of other vertices.

To that end set

$$p_0 = 4\sqrt{\varepsilon}, \quad (8)$$

then by (7) and (4)

$$\frac{p_0^2}{4}(m - 1) \geq |V(T_0)|, \quad p_0 \leq \frac{\mu}{4}. \quad (9)$$

Now, for all  $i \in [t]$  define

$$p_i = \frac{\ell_i}{m} \stackrel{(5)}{\geq} \frac{\varepsilon n}{m} \geq \frac{\varepsilon}{2}. \quad (10)$$

Finally let  $\gamma = 1 - \sum_{i=0}^t p_i$ . Then

$$\gamma \geq 1 - \frac{\mu}{4} - \sum_{i=1}^t p_i = 1 - \frac{\mu}{4} - \frac{\sum_{i=1}^t \ell_i}{m} \geq 1 - \frac{\mu}{4} - \frac{|V(T)|}{m},$$

and so

$$\gamma \geq 1 - \frac{\mu}{4} - \frac{n}{m} \geq 1 - \frac{\mu}{4} - \frac{1}{1 + \mu} = \frac{3\mu - \mu^2}{4(1 + \mu)} \stackrel{(3)}{=} 2\sqrt{\rho}. \quad (11)$$

Hence  $\gamma \in (0, 1)$ .

**Lemma 9.** *Let  $\varepsilon, \ell_0, \dots, \ell_t, p_0, \dots, p_t, \gamma$  and  $\rho$  be defined as above. Then for some  $m_0 = m_0(\varepsilon)$  and  $K = 8$  the following is true for any  $m \geq m_0$ . If  $S$  is a STS on  $m$  vertices, then there is a partition  $\mathcal{P} = C_0 \sqcup C_1 \cdots \sqcup C_t \sqcup R$  of  $V(S)$  with the following properties:*

(a)  $|C_i| = \ell_i \pm K\sqrt{\ell_i \ln \ell_i}$  for all  $i \in [t]$ .

(b) for all  $i \in [t]$  and all  $v \in C_{i-1}$

$$d_{S[C_{i-1}, C_i]}(v) = d\left(p_i \ell_{i-1} \pm K\sqrt{p_i \ell_{i-1} \ln p_i \ell_{i-1}}\right).$$

(c) for all  $i \in [2, t]$  and all  $v \in C_i$

$$d_{S[C_{i-1}, C_i]}(v) = p_i \ell_{i-1} \pm K\sqrt{p_i \ell_{i-1} \ln p_i \ell_{i-1}}.$$

(d) for all  $v \in V(S)$ ,  $d_{S[v \cup R]}(v) \geq \rho m$ .

(e)  $|C_0| = p_0 m \pm K\sqrt{p_0 m \ln p_0 m}$  and  $S[C_0]$  contains a copy of a hypertree  $T_0$  with  $L_0$  as its last level. Moreover for all but at most  $\varepsilon^{0.1}|C_1|$  vertices  $v \in C_1$

$$d_{S[L_0, C_1]}(v) = (1 \pm \varepsilon^{0.1})p_1 \ell_0,$$

and for all vertices  $v \in C_1$

$$d_{S[L_0, C_1]}(v) \leq 2p_1 \ell_0.$$

*Proof.* Recall that  $\sum_{i=0}^t p_i + \gamma = 1$ . Consider a random partition  $\mathcal{P} = \{C_0, \dots, C_t, R\}$ , where vertices  $v \in V(S)$  are chosen into partition classes independently so that  $\mathbb{P}[v \in C_i] = p_i$  for  $i \in [0, t]$  while  $\mathbb{P}[v \in R] = \gamma$ . For  $j \in \{a, b, c, d, e\}$  let  $X^{(j)}$  be the event that the corresponding part of Lemma 9 fails. We will prove that  $\mathbb{P}[X^{(j)}] = o(1)$  for each  $j \in \{a, b, c, d, e\}$ .

**Proof of Property (a).** For all  $i \in [t]$  let  $X_i^{(a)}$  be the event that

$$||C_i| - p_i m| > K\sqrt{p_i m \ln p_i m}.$$

Then since  $|C_i| \sim \text{Bi}(m, p_i)$  and  $\mathbb{E}(|C_i|) = p_i m \stackrel{(10)}{=} \ell_i \stackrel{(5)}{=} \Omega(m)$ , Theorem 6 implies that

$$\mathbb{P}[X_i^{(a)}] \leq 2(\ell_i)^{-K^2/3} = o(m^{-20}).$$

Since by (6),  $t \leq 1 + \log \frac{1}{\varepsilon} \ll m$  we infer that

$$\mathbb{P}[X^{(a)}] = \mathbb{P}\left[\bigcup_{i=1}^t X_i^{(a)}\right] \leq \sum_{i=1}^t \mathbb{P}[X_i^{(a)}] = o(1).$$

**Proof of Property (b).** For all  $i \in [t]$  and  $v \in V(G)$  let  $X_{i,v}^{(b)}$  be the event

$$|d_{S[C_{i-1}, C_i]}(v) - dp_i \ell_{i-1}| > Kd\sqrt{p_i \ell_{i-1} \ln p_i \ell_{i-1}},$$

and  $Y_{i,v}^{(b)}$  be the event

$$|d_{S[C_{i-1}, C_i]}(v) - (m-1)p_i^2/2| > \frac{K}{2}\sqrt{(m-1)p_i^2/2 \ln(m-1)p_i^2/2}.$$

Then for  $i \in [t]$  and  $v \in C_{i-1}$  we have  $d_{S[C_{i-1}, C_i]}(v) \sim \text{Bi}(\frac{m-1}{2}, p_i^2)$  and

$$\mathbb{E}(d_{S[C_{i-1}, C_i]}(v)) = (m-1)p_i^2/2 \stackrel{(10),(5)}{=} dp_i \ell_{i-1} \pm 1.$$

Therefore  $X_{i,v}^{(b)} \subseteq Y_{i,v}^{(b)}$  for a large enough  $m$ . Moreover, Theorem 6 implies that

$$\mathbb{P}[X_{i,v}^{(b)}] \leq \mathbb{P}[Y_{i,v}^{(b)}] \leq 2((m-1)p_i^2/2)^{-K^2/12} \stackrel{(10)}{=} O(m^{-2}).$$



Finally, the union bound yields

$$\mathbb{P}[X^{(b)}] \leq \sum_{i \in [t], v \in C_i} \mathbb{P}[X_{i,v}^{(b)}] = o(1).$$

**Proof of Property (c).** Proof follows the lines of the proof of part (b), since for  $i \in [2, t]$  and  $v \in C_i$  we have  $d_{S[C_{i-1}, C_i]}(v) \sim \text{Bi}(\frac{m-1}{2}, 2p_i p_{i-1})$  and  $\mathbb{E}(d_{S[C_{i-1}, C_i]}(v)) = (m-1)p_i p_{i-1} = p_i \ell_{i-1} \pm 1$ . Hence we have  $\mathbb{P}[X^{(c)}] = o(1)$

**Proof of Property (d).** Proof follows the lines of the proof of part (b), since for all  $v \in V(S)$  we have  $d_{S[v \cup R]}(v) \sim \text{Bi}(\frac{m-1}{2}, \gamma^2)$  and  $\mathbb{E}(d_{S[v \cup R]}(v)) = \frac{m-1}{2} \gamma^2 \stackrel{(11)}{\geq} 2\rho(m-1)$ . Hence we have  $\mathbb{P}[X^{(d)}] = o(1)$

**Proof of Property (e)**

We say that a set  $C \subseteq V(S)$  is *typical* if  $|C| = p_0 m \pm K \sqrt{p_0 m \ln p_0 m}$  and  $S[C]$  contains a copy of  $T_0$ . For a partition  $\mathcal{P} = \{C_0, \dots, C_t, R\}$  set  $C_i(\mathcal{P}) = C_i$  for all  $i \in [0, t]$ .

Next we will show that the first statement of (e), namely that  $C_0(\mathcal{P})$  is typical, holds asymptotically almost surely.

**Claim 10.**

$$\mathbb{P}[C_0(\mathcal{P}) \text{ is typical}] = 1 - o(1).$$

*Proof.* Let  $X$  be the event that  $||C_0| - p_0 m| \leq K \sqrt{p_0 m \ln p_0 m}$ , and  $Y$  be the event that  $S[C_0(\mathcal{P})]$  contains a copy of  $T_0$ . Since  $|C_0| \sim \text{Bi}(m, p_0)$  and  $\mathbb{E}(|C_0|) = p_0 m \stackrel{(5)}{=} \Omega(m)$ , Theorem 6 implies  $\mathbb{P}[X] = 1 - o(1)$ .

For  $v \in V(S)$  let  $Z_v$  denote the event that  $d_{S[C_0]}(v) \geq |V(T_0)|$ , then  $\bigcap_{v \in V(S)} Z_v \subseteq Y$ . Indeed, if every vertex has degree at least  $|V(T_0)|$  in  $S[C_0]$ , then  $T_0$  can be found in  $S[C_0]$  greedily, adding one hyperedge at a time.

Following the lines of proof of (d), we have  $d_{S[C_0]}(v) \sim \text{Bi}(\frac{m-1}{2}, p_0^2)$ , and

$$\mathbb{E}(d_{S[C_0]}(v)) = \frac{m-1}{2} p_0^2 \stackrel{(9)}{\geq} 2|V(T_0)|,$$

so by Theorem 6 for all  $v \in V(S)$  we have  $\mathbb{P}[Z_v] \geq 1 - o(m^{-20})$ . Finally,

$$\mathbb{P}[Y] \geq \mathbb{P}\left[\bigcap_{v \in V(S)} Z_v \geq 1 - m \cdot o(m^{-20}) \geq 1 - o(1)\right].$$

Therefore  $\mathbb{P}[X] = \mathbb{P}[Y] = 1 - o(1)$  and hence  $\mathbb{P}[X \cap Y] = \mathbb{P}[C_0(\mathcal{P}) \text{ is typical}] = 1 - o(1)$ . □

Now, for every typical set  $C$ , we fix one copy of  $T_0$  in  $S[C]$ .

We first show that there are not many vertices in  $\bar{C} = V(S) \setminus C$  that have low degree in  $S[L_0, \bar{C}]$ .

**Claim 11.** For any  $\alpha > 0$  and any typical set  $C$  all but at most  $|C|/\alpha$  vertices in  $v \in \bar{C}$  satisfy

$$d_{S[L_0, \bar{C}]}(v) = (1 \pm \alpha)\ell_0.$$

*Proof of Claim.* For  $v \in \overline{C}$  and  $x \in L_0$  there is a unique  $w \in V(S)$  such that  $\{v, x, w\} \in E(S)$ . Consequently,  $d_{S[L_0, \overline{C}]}(v) \leq \ell_0$  holds for any  $v \in \overline{C}$ .

Let  $A = \{\{x, v, w\} : x \in L_0, v \in \overline{C}, w \in \overline{C}\}$ . Since for every  $x \in L_0$  there are at most  $|C|$  edges  $\{v, x, w\}$  with  $v \in \overline{C}$  and  $w \in C$

$$|A| \geq \ell_0(|\overline{C}| - |C|). \tag{12}$$

On the other hand, let  $b$  be the number of “bad” vertices  $v \in \overline{C}$ , i.e., vertices  $v$  with  $d_{S[L_0, \overline{C}]}(v) < (1 - \alpha)\ell_0$ . Then we have

$$|A| \leq b(1 - \alpha)\ell_0 + (|\overline{C}| - b)\ell_0. \tag{13}$$

Comparing (12) and (13) yields that  $b \leq |C|/\alpha$ . □

Let  $E$  be the event that property (e) holds. Next we will show that

$$\mathbb{P}[E|C_0(\mathcal{P}) = C] = 1 - o(1) \text{ for every typical } C. \tag{14}$$

This implies that  $E$  holds with probability  $1 - o(1)$ .

Indeed, by Claim 10,  $\mathbb{P}[C_0(\mathcal{P}) \text{ is typical}] = \sum_{C \text{ is typical}} \mathbb{P}[C_0(\mathcal{P}) = C] = (1 - o(1))$  and so

$$\begin{aligned} \mathbb{P}[E] &\geq \sum_{C \text{ is typical}} \mathbb{P}(C_0(\mathcal{P}) = C) \mathbb{P}[E|C_0(\mathcal{P}) = C] \\ &\stackrel{(14)}{\geq} (1 - o(1)) \sum_{C \text{ is typical}} \mathbb{P}(C_0(\mathcal{P}) = C) \geq 1 - o(1). \end{aligned}$$

It remains to prove (14).

Denote by  $(\Omega, \mathcal{F}, \mathbb{P})$  the space of all partitions of  $V(S)$  with  $\mathbb{P}[v \in C_i] = p_i$  for  $i \in [0, t]$  and  $\mathbb{P}[v \in R] = \gamma$ , and for fixed  $C$  let  $(\Omega, \mathcal{F}, \mathbb{P}_C)$  to be the space of all partitions of  $V(S)$  with the probability function  $\mathbb{P}_C(A) = \mathbb{P}(A|C_0(\mathcal{P}) = C)$ .

With this notation we need to show that  $\mathbb{P}_C(E) = 1 - o(1)$  for every typical  $C$ .

Recall that  $\overline{C} = V(S)/C$  and for all  $v \in \overline{C}$  let

$$\chi(v) = \begin{cases} 1, & \text{if } v \in C_1 \\ 0, & \text{otherwise.} \end{cases}$$

Note that for all  $v \in \overline{C}$

$$\begin{aligned} \mathbb{P}_C(\chi(v) = 1) &= \mathbb{P}_C(v \in C_1) = \mathbb{P}(v \in C_1 | C_0(\mathcal{P}) = C) = \frac{\mathbb{P}(v \in C_1 \wedge C_0(\mathcal{P}) = C)}{\mathbb{P}(C_0(\mathcal{P}) = C)} \\ &= \frac{p_1 \cdot p_0^{|C|} (1 - p_0)^{m - |C| - 1}}{p_0^{|C|} (1 - p_0)^{m - |C|}} = \frac{p_1}{1 - p_0} = q. \end{aligned}$$

Then by (8)

$$q = (1 \pm \varepsilon^{0.3})p_1. \quad (15)$$

Moreover, since for fixed  $v \in V(S)$  the event  $\{v \in C_1\}$  was, in the “initial” space  $(\Omega, \mathcal{F}, \mathbb{P})$ , independent of the outcome of a random experiment for the remaining vertices  $w \in V(S) \setminus \{v\}$ , we infer that the random variables  $\{\chi(v) : v \in \overline{C}\}$  are mutually independent.

Therefore for the rest of the proof we assume that typical  $C$  with  $L_0 \subset C$  is fixed and all events and random variables are considered in the space  $(\Omega, \mathcal{F}, \mathbb{P}_C)$ .

For a typical  $C$  define

$$M = M(C) = \{v \in \overline{C} : d_{S[L_0, \overline{C}]}(v) = (1 \pm \varepsilon^{0.2})\ell_0\}. \quad (16)$$

Recall that since  $C$  is typical we have

$$|C| = (1 + o(1))p_0m, \text{ and } |\overline{C}| = (1 - o(1))(1 - p_0)m. \quad (17)$$

Then by Claim 11 with  $\alpha = \varepsilon^{0.2}$

$$|M| \geq \overline{C} - \frac{|C|}{\varepsilon^{0.2}} \stackrel{(17)}{=} |\overline{C}| - \frac{|\overline{C}|p_0}{\varepsilon^{0.2}(1 - p_0)}(1 - o(1)) \stackrel{(8)}{\geq} (1 - \varepsilon^{0.2})|\overline{C}|. \quad (18)$$

Note that  $M$  is independent of the choice of  $C_1$  and is fully determined by  $C$  and  $S$ .

Next we verify that certain events  $E^{(1)}$ ,  $E^{(2)}$ ,  $E^{(3)}$  hold asymptotically almost surely and that  $E^{(1)} \wedge E^{(2)} \wedge E^{(3)} \subseteq E$ . Let event  $E^{(1)}$  be defined as

$$E^{(1)} : |M \cap C_1| \geq (1 - \varepsilon^{0.1})|C_1|.$$

Since  $|C_1| \sim \text{Bi}(|\overline{C}|, q)$  and  $|M \cap C_1| \sim \text{Bi}(|M|, q)$ , we have

$$\mathbb{E}(|C_1|) = |\overline{C}|q \text{ and } \mathbb{E}(|M \cap C_1|) \stackrel{(18)}{\geq} (1 - \varepsilon^{0.2})|\overline{C}|q.$$

Hence Theorem 6 implies that with probability  $1 - o(1)$  we have  $|M \cap C_1|/|C_1| \geq 1 - \varepsilon^{0.1}$  and so  $\mathbb{P}_C[E^{(1)}] = 1 - o(1)$ .

Now for every  $v \in \overline{C}$  let  $N(v)$  be the random variable that equals to the number of hyperedges  $\{v, x, w\}$ , where  $x \in L_0$  and  $w \in C_1$ . Then  $N(v) \sim \text{Bi}(d_{S[L_0, \overline{C}]}(v), q)$  for all  $v \in \overline{C}$ .

Let  $E^{(2)}$  be the event

$$E^{(2)} : N(v) = (1 \pm \varepsilon^{0.1})\ell_0p_1 \text{ for all } v \in M.$$

For every  $v \in M$ , we have  $N(v) \sim \text{Bi}(d_{S[L_0, \overline{C}]}(v), q)$  and so  $\mathbb{E}(N(v)) = (1 \pm 2\varepsilon^{0.2})\ell_0p_1$  by (16) and (15). Then Theorem 6 combined with the union bound implies  $\mathbb{P}_C[E^{(2)}] = 1 - o(1)$ .

Let  $E^{(3)}$  be the event

$$E^{(3)} : N(v) \leq 2\ell_0p_1 \text{ for all } v \in \overline{C}.$$

For every  $v \in \overline{C}$  we have  $N(v) \sim \text{Bi}(d_{S[L_0, \overline{C}]}(v), q)$  and  $d_{S[L_0, \overline{C}]} \leq \ell_0$ , hence we always have  $\mathbb{E}(N(v)) \stackrel{(15)}{\leq} (1 + \varepsilon^{0.3})\ell_0 p_1$ . Therefore by Theorem 6 and the union bound we have  $\mathbb{P}_C[E^{(3)}] = 1 - o(1)$ .

It remains to notice that for  $v \in C_1$  we have  $d_{S[L_0, C_1]}(v) = N(v)$  and so  $E^{(1)} \wedge E^{(2)} \wedge E^{(3)} \subseteq E$ . Therefore,  $\mathbb{P}_C[E] \geq 1 - o(1)$ , finishing the proof of (14).  $\square$

**Embedding of  $T$ .** We start with applying Lemma 9 to  $S$  obtaining a partition  $\mathcal{P} = \{C_0, \dots, C_t, R\}$  of  $V(S)$  that satisfies properties (a)-(e) of Lemma 9. To simplify our notation we set  $G_1 = S[L_0, C_1]$  and for  $i \in [2, t]$   $G_i = S[C_{i-1}, C_i]$ .

- 1) We first verify that Lemma 9 guarantees that the assumptions of Lemma 7 and Lemma 8 are satisfied. These Lemmas then yield systems of stars  $\mathcal{S}_i = \{S_i^1, \dots, S_i^{p_i}\}$  for  $i \in [2, t]$ , such that each  $\mathcal{S}_i$  covers almost all vertices of the respective  $G_i$ . (See Figure 1, where each star  $S_i^j$  is represented by a single grey edge.)
- 2) Let  $F$  be the union of  $T_0$  with  $\mathcal{S}_i$ 's. The ‘‘almost cover’’ property of  $\mathcal{S}_i$ 's then allows us to show that hyperforest  $F$  contains a large connected component  $T_1$  which contains almost all vertices of  $T$ . (See Figure 1, green and grey edges form  $T_1$ .)
- 3) Finally, we extend  $T_1$  into a full copy of  $T$  in a greedy procedure using the vertices of  $R$ . (See Figure 1, vertices of  $R$  are blue.)




Figure 1: Case  $d = 1$  and  $t = 2$ . Green edges form  $T_0$ , grey edges are hyperstars  $S_i^j$ , blue edges are constructed by using vertices in reservoir  $R$ .

**Step 1.** Construction of the hyperforest  $F$ .

We start with applying Lemma 9 to  $S$  and obtaining a partition  $\mathcal{P} = \{C_0, \dots, C_t, R\}$  of  $V(S)$  that satisfies properties (a)-(e) of Lemma 9. Recall that  $G_1 = S[L_0, C_1]$  and for  $i \in [2, t]$  let  $G_i = S[C_{i-1}, C_i]$ .

Let  $N_i = |V(G_i)|$ , then  $N_1 = \ell_0 + |C_1|$  and  $N_i = |C_{i-1}| + |C_i|$  for  $i \in [2, t]$ . Due to property (a) of Lemma 9 we have that for a sufficiently large  $m$  and for all  $i \in [t]$

$$N_i = (1 \pm \varepsilon)(\ell_{i-1} + \ell_i) \leq (1 \pm \varepsilon) \left( \frac{\ell_i}{2d} + \ell_i \right) \leq 2\ell_i. \quad (19)$$

In what follows we will show that  $G_1$  satisfies the assumptions of Lemma 7 and for  $i \in [2, t]$ ,  $G_i$  satisfies the assumptions of Lemma 8.

We start with  $G_1$ . Recall that for a given  $\mu$  we have defined  $\delta$  (see (3)), and  $\varepsilon_{3.1}$  was a constant guaranteed by Lemma 7 with  $\delta$  and  $k = 2$  as input. We then defined  $\varepsilon$  (see (4)) to be small enough, in particular such that  $\varepsilon_{3.1} \geq \varepsilon^{0.1}$ . We will now show that  $G_1$  satisfies conditions of Lemma 7 with  $D = D_1 = p_1\ell_0$  and  $N = N_1$ .

To verify condition (i) of Lemma 7 it is enough to recall that  $G_1 = S[L_0, C_1]$  and so (i) holds with  $X = L_0$  and  $Y = C_1$ .

Note that condition (iii) is guaranteed by property (e), since for all  $v \in C_1 = Y$  we have  $d_{G_1}(v) \leq 2D$ .

We now verify condition (ii) of Lemma 7. Property (b) with  $i = 1$  guarantees that for all  $v \in X = L_0$

$$d_{G_1}(v) = d \left( D \pm K\sqrt{D \ln D} \right).$$

Since  $D = p_1\ell_0 \stackrel{(5),(10)}{=} \Omega(\sqrt{n})$ , for a large enough  $n$  we have for all  $v \in X = L_0$

$$d_{G_1}(v) = dD(1 \pm \varepsilon_{3.1}). \quad (20)$$

Property (e) in turn guarantees that for all but at most  $\varepsilon^{0.1}|C_1| \leq \varepsilon_{3.1}N_1$  vertices  $v \in Y = C_1$  we have

$$d_{G_1}(v) = (1 \pm \varepsilon^{0.1})p_1\ell_0 = D(1 \pm \varepsilon_{3.1}). \quad (21)$$

Now, (20) and (21) imply that  $G_1$  satisfies condition (ii) of Lemma 7.

Lemma 7 produces a collection  $\{S_1^1, \dots, S_{p_1}^1\}$  of disjoint hyperstars of  $G_1$  centered at vertices of  $L_0$ , each  $S_j^1$  has at most  $d$  hyperedges, and

$$\text{the star forest } \mathcal{S}_1 = \bigcup_{j=1}^{p_1} S_j^1 \text{ covers all but at most } \delta N_1 \text{ vertices of } G_1. \quad (22)$$

Similarly for all  $i \in [2, t]$ , the hypergraph  $G_i$  satisfies the assumptions of Lemma 8 with the parameters  $K_{3.2} = 8, d_{3.2} = d$  and  $D = D_i = p_i\ell_{i-1}$ . Indeed, condition (i) of Lemma 8 is guaranteed by taking  $X = C_{i-1}$  and  $Y = C_i$ , and condition (ii) is guaranteed by properties (b) and (c) of Lemma 9.

Then for  $i \in [2, t]$ , Lemma 8 when applied to  $G_i$  yields a collection  $\{S_1^i, \dots, S_{p_i}^i\}$  of disjoint hyperstars centered at vertices of  $C_{i-1}$ , each  $S_j^i$  has at most  $d$  hyperedges, and the

star forest  $\mathcal{S}_i = \bigcup_{j=1}^{p_i} S_j$  covers all but at most  $O(N_i D_i^{-1/2} \ln^{3/2} D_i)$  vertices of  $G_i$ . Once again recall that for  $i \in [2, t]$  we have  $D_i = p_i \ell_{i-1} \stackrel{(5),(10)}{=} \Omega(n)$  and so for a large enough  $n$ ,

$$\mathcal{S}_i \text{ covers all but at most } \varepsilon N_i \text{ vertices of } G_i. \quad (23)$$

Now, let  $F = T_0 \cup \bigcup_{i=1}^t \mathcal{S}_i$ , then  $F$  is a hyperforest and we will now find a hypertree  $T_1 \subseteq F$  such that  $T_1$  is an almost spanning subhypertree of  $T$ .

We first will estimate  $|V(F)|$ . Since for  $i \in [2, t]$  a star forest  $\mathcal{S}_i$  misses at most  $\varepsilon N_i$  vertices of  $C_i$  and  $\mathcal{S}_1$  misses at most  $\delta N_1$  vertices of  $C_1$  we have:

$$|V(F)| \geq |V(T_0)| + |C_1| - \delta N_1 + \sum_{i=2}^t (|C_i| - \varepsilon N_i).$$

Now, by property (a) of Lemma 9, for large enough  $m$  and for any  $i \in [t]$ , we have  $|C_i| = (1 \pm \varepsilon)\ell_i$ . Therefore

$$|V(F)| \geq |V(T_0)| + \sum_{i=1}^t (1 - \varepsilon)\ell_i - \delta N_1 - \varepsilon \sum_{i=2}^t N_i \stackrel{(19)}{\geq} |V(T_0)| + \sum_{i=1}^t \ell_i - (\varepsilon + 2\delta)\ell_1 - 3\varepsilon \sum_{i=2}^t \ell_i.$$

Since  $|V(T_0)| + \sum_{i=1}^t \ell_i = |V(T)|$  and  $\varepsilon < \delta$  we have

$$|V(F)| \geq |V(T)| - 3\delta \sum_{i=1}^t \ell_i \geq (1 - 3\delta)|V(T)|. \quad (24)$$

**Step 2.** Embedding the most of  $T$ .

**Claim 12.**  $S$  contains a hypertree  $T_1 \subseteq F$  such that  $T_1$  is a subhypertree of  $T$  and  $|E(T_1)| \geq (1 - 20\delta)|E(T)|$ .

*Proof.* Recall that while  $V(T) = \bigcup_{i=0}^h V_i$ , the forest  $F$  has the vertex set  $V(F) = \bigcup_{i=0}^{i_0} V_i \cup \bigcup_{i=1}^t C_i$ , where we have set  $t = h - i_0$  and  $V_{i_0} = L_0$ . Also recall that  $V_0 = \{v_0\}$  was a root of  $T$  (and  $F$ ).

For a non-root vertex  $v \in V_i$  (or  $v \in C_i$ ) a parent of  $v$  is a unique vertex  $u \in V_{i-1}$  (or  $u \in C_{i-1}$ ) such that  $\{u, v, w\} \in F$ . If a vertex  $v$  has a parent  $u$  we will write  $p(v) = u$ , if a vertex  $v$  has no parent we will say that  $v$  is an orphan.

For each  $v \in V(F) \setminus \{v_0\}$  consider “a path of ancestors”  $v = a_i, a_{i-1}, \dots, a_1 = a^*$ , i.e., a path satisfying  $p(a_j) = a_{j-1}$ ,  $j \in [2, i]$  and such that  $a^* = a^*(v)$  is an orphan in  $F$ .

Let  $T_1 \subseteq F$  be a subtree of  $F$  induced on a set  $\{v \in V(F) : a^*(v) = v_0\}$ . Note that if for some  $v \in V(F)$  we have  $a^*(v) \neq v_0$ , then  $a^*(v) \notin V(T_0)$ .

For  $i \in [t]$  let  $U_i \subseteq C_i$  be the set of vertices not covered by  $\mathcal{S}_i$ . Then

$$|U_1| \stackrel{(22)}{\leq} \delta N_1 \text{ and } |U_i| \stackrel{(23)}{\leq} \varepsilon N_i \text{ for all } i \in [2, t]. \quad (25)$$

Note that all orphan vertices, except of  $v_0$ , belong to  $\bigcup_{i=1}^t U_i$  as they were not covered by some  $\mathcal{S}_i$ . In particular, for every  $v \notin V(T_1)$  its ancestor  $a^*(v)$  is an orphan and hence belongs to  $\bigcup_{i=1}^t U_i$

For an orphan vertex  $a^*$  let  $T(a^*)$  be a subtree of  $F$  rooted at  $a^*$ . Then for every orphan vertex  $a^* \in U_i$  we have

$$|V(T(a^*))| \leq 1 + 2d + \dots + (2d)^{t-i} \leq 3(2d)^{t-i}.$$

Finally, every  $v \notin V(T_1)$  is in  $T(a^*(v))$ , where  $a^*(v) \in U_i$  for some  $i \in [t]$ , and therefore we have

$$|V(T_1)| \geq |V(F)| - \sum_{i=1}^t |U_i| \cdot 3(2d)^{t-i} \stackrel{(24),(25)}{\geq} (1-3\delta)|V(T)| - 3\delta N_1(2d)^{t-1} - 3\varepsilon \sum_{i=2}^t N_i(2d)^{t-i}.$$

Now, by (19), we have  $N_i \leq 2\ell_i$  and, by (5),  $\ell_i(2d)^{t-i} = \ell_t$  for all  $i \in [t]$ , and so

$$|V(T_1)| \geq (1-3\delta)|V(T)| - 6\delta\ell_t - 6\varepsilon(t-1)\ell_t.$$

Recall that  $\ell_t$  is the size of the last level of  $T$ , so  $\ell_t \leq |V(T)|$ . Also  $t \leq 1 + \log(\frac{1}{\varepsilon})$ , and since  $\varepsilon$  is sufficiently small  $6\varepsilon(t-1) \leq \sqrt{\varepsilon} \leq \delta$  and so

$$|V(T_1)| \geq (1-10\delta)|V(T)|.$$

For every hypertree  $T'$  we have  $|V(T')| = 2|E(T')| + 1$ , and so

$$|E(T_1)| \geq (1-20\delta)|E(T)|,$$

finishing the proof of the Claim. □

**Step 3.** Finally, we complete  $T_1$  to a full copy of  $T$  by using the reservoir  $R$ .

**Claim 13.** *Assume that  $\mathcal{P}$  is a partition guaranteed by Lemma 9 and  $T_1$  be a hypertree guaranteed by Claim 12. Then  $T_1$  can be extended to a copy of  $T$  in  $S$ .*

*Proof.* Since  $T_1$  is a subhypertree of  $T$ , there is a sequence of hyperedges  $\{e_1, \dots, e_{p-1}\}$  such that each  $T_i = T_1 \cup_{j=1}^{i-1} e_j$  for  $i \in [p]$  is a hypertree an  $T_p \cong T$ . Every vertex  $v \in V(S)$  has degree at least  $\rho m$  in  $R$  (by Lemma 9) and

$$p-1 = |E(T)| - |E(T_1)| \stackrel{\text{Claim 12}}{\leq} 20\delta|E(T)| \leq 10\delta n \stackrel{(3)}{\leq} \frac{\rho m}{2}.$$

Hence we can greedily embed edges  $e_1, \dots, e_{p-1}$ . Indeed, having embedded the edges  $e_1, \dots, e_{i-1}$  for some  $i \in [p-1]$ , for set  $R_i = R \setminus \cup_{j=1}^{i-1} e_j$  and all  $v \in V(S)$  we have

$$d_{S[v \cup R_i]} \geq \rho m - 2(i-1) > 0$$

allowing the greedy embedding to continue. Then the last hypertree  $T_p$  is, by the construction, isomorphic to  $T$ . □

## 4 Concluding remarks

We notice that with a similar proof one can verify Conjecture 1 for some other types of hypertrees.

Let  $D = \{d_1, \dots, d_k\}$  be a sequence of integers. Let  $T$  be a tree rooted at  $v_0$  and let  $V(T) = V_0 \sqcup V_1 \sqcup \dots \sqcup V_h$  be a partition of  $V(T)$  into levels, so that  $V_i$  consist of vertices distance  $i$  from  $v_0$ . We say that  $T$  is  $D$ -ary hypertree if for every  $i \in [0, h - 1]$  there is  $j \in [k]$  such that for every  $v \in V_i$  the forward degree of  $v$  is  $d_j$ . In other words, forward degree of every non-leaf vertex of  $T$  is in  $D$ , and depends only on the height of a vertex in  $T$ .

Following the lines of the proof of Theorem 3 one can conclude that for any finite set  $D \subset \mathbb{N}$ , and any  $\mu$ , any large enough STS  $S$  contains any  $D$ -ary hypertree  $T$ , provided  $|V(T)| \leq |V(S)|/(1 + \mu)$ .

Another type of hypertrees for which Conjecture 1 holds are truncated  $d$ -ary hypertrees. In a perfect  $d$ -ary hypertree label children of every vertex with numbers  $\{1, \dots, 2d\}$ . Then every leaf can be identified with a sequence  $\{a_1, \dots, a_h\} \in [2d]^h$  based on the way that leaf is reached from the root, and all leaves can be ordered by a lexicographic order. We say that  $T$  is a *truncated perfect  $d$ -ary hypertree* if, for some integer  $t$ , the hypertree  $T$  is obtained from a perfect  $d$ -ary hypertree by removing the smallest  $2t$  leaves (according to the lexicographic order).

With an essentially same proof as of Theorem 3, for every  $\mu > 0$  and  $d$ , any sufficiently large Steiner triple system  $S$  contains any truncated  $d$ -ary hypertree  $T$  with  $|V(T)| \leq |V(S)|/(1 + \mu)$ .

## References

- [1] N. Alon, J.-H. Kim, J. H. Spencer. Nearly perfect matchings in regular simple hypergraphs. *Israel J. Math.* 100 (1997), 171–187.
- [2] N. Alon, J. H. Spencer. *The probabilistic method*. Wiley Series in Discrete Mathematics and Optimization, John Wiley & Sons, Inc., Hoboken, NJ, 2016.
- [3] A. Arman, V. Rödl, Marcelo Tadeu Sales. Colorful matchings. [arXiv:2102.09633](https://arxiv.org/abs/2102.09633).
- [4] B. Elliott, V. Rödl. Embedding hypertrees into Steiner triple systems. *J. Combin. Des.* 27(2) (2019), 82–105.
- [5] P. Frankl, V. Rödl, Near perfect coverings in graphs and hypergraphs. *European J. Combin.* 6(4) (1985), 317–326.
- [6] S. Janson, T. Łuczak, A. Ruciński. *Random graphs*. Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000.
- [7] A. V. Kostochka, V. Rödl. Partial Steiner systems and matchings in hypergraphs. *Random Structures Algorithms* 13(3-4) (1998), 335–347.
- [8] N. Pippenger, J. H. Spencer. Asymptotic behavior of the chromatic index for hypergraphs. *J. Combin. Theory Ser. A* 5(1) (1989), 24–42.