

# Majority Edge-Colorings of Graphs

Felix Bock

Institute of Optimization and OR  
Ulm University  
Ulm, Germany

`felix.bock@uni-ulm.de`

Rafał Kalinowski

Department of Discrete Mathematics  
AGH University  
Krakow, Poland

`kalinows@agh.edu.pl`

Johannes Pardey

Institute of Optimization and OR  
Ulm University  
Ulm, Germany

`johannes.pardey@uni-ulm.de`

Monika Piłśniak

Department of Discrete Mathematics  
AGH University  
Krakow, Poland

`pilsniak@agh.edu.pl`

Dieter Rautenbach

Institute of Optimization and OR  
Ulm University  
Ulm, Germany

`dieter.rautenbach@uni-ulm.de`

Mariusz Woźniak

Department of Discrete Mathematics  
AGH University  
Krakow, Poland

`mwozniak@agh.edu.pl`

Submitted: May 30, 2022; Accepted: Feb 10, 2023; Published: Mar 10, 2023

© The authors. Released under the CC BY-ND license (International 4.0).

## Abstract

We propose the notion of a majority  $k$ -edge-coloring of a graph  $G$ , which is an edge-coloring of  $G$  with  $k$  colors such that, for every vertex  $u$  of  $G$ , at most half the edges of  $G$  incident with  $u$  have the same color. We show the best possible results that every graph of minimum degree at least 2 has a majority 4-edge-coloring, and that every graph of minimum degree at least 4 has a majority 3-edge-coloring. Furthermore, we discuss a natural variation of majority edge-colorings and some related open problems.

**Mathematics Subject Classifications:** 05C15

## 1 Introduction

Motivated by similar notions considered for vertex-colorings, we propose and study *majority edge-colorings* of graphs: For a (finite, simple, and undirected) graph  $G$ , an edge-coloring  $c : E(G) \rightarrow [k]$  is a *majority  $k$ -edge-coloring* if, for every vertex  $u$  of  $G$  and every color  $\alpha$  in  $[k]$ , at most half the edges incident with  $u$  have the color  $\alpha$ .

Before we present our results, we discuss some related research. Lovász [9] showed that every graph  $G$  has a 2-vertex-coloring such that, for every vertex  $u$  of  $G$ , at most half the neighbors of  $u$  have the same color as  $u$ . For infinite graphs, this leads to the *Unfriendly Partition Conjecture* [2]. Kreutzer, Oum, Seymour, van der Zypen, and Wood [8] showed that every digraph  $D$  has a 4-vertex-coloring such that, for every vertex  $u$  of  $D$ , at most half the out-neighbors of  $u$  have the same color as  $u$ , and they conjecture that 3 colors suffice. Anholcer, Bosek, and Grytczuk [4] studied a choosability version for digraphs. It follows from a result of Wood [13] that every digraph  $D$  has a 4-arc-coloring such that, for every vertex  $u$  of  $D$ , at most half the arcs leaving  $u$  have the same color. Further related research concerns *defective* or *frugal* edge-colorings [1, 3, 7], where maximum degree conditions are imposed on the subgraphs formed by edges having the same color.

Our first result is that 2 colors almost suffice for a majority edge-coloring.

**Theorem 1.** *Let  $G$  be a connected graph.*

- (i) *If  $G$  has an even number of edges or  $G$  contains vertices of odd degree, then  $G$  has a 2-edge-coloring such that, for every vertex  $u$  of  $G$ , at most  $\left\lceil \frac{d_G(u)}{2} \right\rceil$  of the edges incident with  $u$  have the same color.*
- (ii) *If  $G$  has an odd number of edges, all vertices of  $G$  have even degree, and  $u_G$  is any vertex of  $G$ , then  $G$  has a 2-edge-coloring such that, for every vertex  $u$  of  $G$  distinct from  $u_G$ , exactly  $\frac{d_G(u)}{2}$  of the edges incident with  $u$  have the same color, and exactly  $\frac{d_G(u_G)}{2} + 1$  of the edges incident with  $u_G$  have the same color.*

Using Vizing's bound [12] on the chromatic index leads to our second result.

**Theorem 2.** *Every graph of minimum degree at least 2 has a majority 4-edge-coloring.*

Clearly, a graph containing a vertex of degree 1 does not have a majority edge-coloring, which motivates the minimum degree condition in Theorem 2. Furthermore, since graphs of minimum degree at least 2, maximum degree 3, and chromatic index 4 have no majority 3-edge-coloring, the number of colors in Theorem 2 is best possible under this minimum degree condition. In fact, if a graph  $G$  of minimum degree at least 2 has an induced subgraph  $H$  such that  $H$  is a graph of maximum degree 3 and chromatic index 4 such that all vertices of  $H$  have degree 2 or 3 in  $G$ , then  $G$  has no majority 3-edge-coloring. We conjecture that all graphs for which 4 colors are needed contain an induced subgraph of maximum degree 3 and chromatic index 4.

Our third result supports this conjecture.

**Theorem 3.** *Every graph of minimum degree at least 4 has a majority 3-edge-coloring.*

Since a graph containing a vertex of odd degree at least 3 does not have a majority 2-edge-coloring, the number of colors in Theorem 3 is best possible under the minimum degree condition in that result. In Section 2 we prove our results, and in a conclusion we discuss a variation of majority edge-colorings.

## 2 Proofs

Theorem 1 is a consequence of *Euler's Theorem* [6].

*Proof of Theorem 1.*

- (i) Let the multigraph  $G'$  arise from  $G$  by adding the edges of a perfect matching  $M$  on the possibly empty set of vertices of odd degree. Clearly, the multigraph  $G'$  is connected and every vertex has even degree in  $G'$ . Let  $e_0e_1 \cdots e_{m-1}$  be an *Euler tour* of  $G'$ , where, provided that  $M$  is not empty, we may assume that  $e_{m-1} \in M$ . Setting  $c(e_i) = (i \bmod 2) + 1$  for every index  $i$  such that  $e_i$  belongs to  $G$ , yields the desired 2-edge-coloring of  $G$ .
- (ii) Let  $e_0e_1 \cdots e_{m-1}$  be an Euler tour of  $G$  such that  $e_0$  is incident with  $u_G$ . Now, setting  $c(e_i) = (i \bmod 2) + 1$  for every index  $i$ , yields the desired 2-edge-coloring of  $G$ .  $\square$

Theorem 2 is a consequence of *Vizing's Theorem* [12].

*Proof of Theorem 2.* Let  $G$  be a graph of minimum degree at least 2. If  $u$  is a vertex of degree  $d$ , and  $d = d_1 + \cdots + d_k$  is a partition of  $d$  into positive integers  $d_i$ , then the graph  $H$  arises from  $G$  by splitting  $u$  into vertices of degrees  $d_1, \dots, d_k$  if there is a partition  $N_G(u) = N_1 \cup \cdots \cup N_k$  of  $N_G(u)$  with  $|N_i| = d_i$  for  $i \in [k]$ ,  $V(H) = (V(G) \setminus \{u\}) \cup \{u_1, \dots, u_k\}$  for  $u_1, \dots, u_k \notin V(G)$ , and  $E(H) = E(G - u) \cup \bigcup_{i \in [k]} \{u_i v : v \in N_i\}$ . See Figure 1 for an illustration.




Figure 1: Splitting a vertex  $u$  of degree 7 into vertices of degrees 2, 2, and 3.

Now, let  $G^*$  arise from  $G$  by splitting every vertex of degree  $d > 3$  into vertices of degrees

- $3, \dots, 3$ , if  $d \equiv 0 \bmod 3$ ,
- $2, 2, 3, \dots, 3$ , if  $d \equiv 1 \bmod 3$ , and
- $2, 3, \dots, 3$ , if  $d \equiv 2 \bmod 3$ .

Note that there is a natural bijection between the edges of  $G$  and those of  $G^*$ . By Vizing's Theorem [12], the graph  $G^*$  has a proper 4-edge-coloring, which yields a majority 4-edge-coloring of  $G$ . In fact, we obtain an edge-coloring of  $G$  such that, for every vertex of degree  $d$  at least 4, at most  $(d + 2)/3$  of the incident edges have the same color.  $\square$

We proceed to the proof of Theorem 3.

*Proof of Theorem 3.* Let  $G$  be a graph of minimum degree  $\delta$  at least 4. Let  $V(G) = D \cup A \cup C$  be the *Gallai-Edmonds decomposition* of  $G$ , that is,  $D$  is the set of all vertices of  $G$  that are missed by some maximum matching,  $A$  is the set of all vertices of  $G$  outside of  $D$  that have a neighbor in  $D$ , and  $C$  contains the remaining vertices, cf. [10].

Let  $D'$  be the set of isolated vertices in  $G[D]$ .

**Claim 4.** *It is possible to select, for every vertex  $u$  in  $D'$ , exactly one edge incident with  $u$  in such a way that every vertex  $v$  in  $A$  is incident with at most  $\left\lfloor \frac{d_G(v)}{2} \right\rfloor$  of the selected edges.*

*Proof of Claim 4.* Let  $H_0$  be the bipartite subgraph of  $G$  with partite sets  $D'$  and  $A$  whose edges are exactly all edges of  $G$  between  $D'$  and  $A$ . Let  $H$  arise from  $H_0$  by replacing each vertex  $v$  in  $A$  by  $\left\lfloor \frac{d_G(v)}{2} \right\rfloor$  copies having the same neighbors in  $D'$  as  $v$ . Clearly, the desired statement follows if  $H$  has a matching saturating all vertices in  $D'$ . Suppose, for a contradiction, that such a matching does not exist. By *Hall's Theorem* [5], there is a subset  $S$  of  $D'$  with  $|S| > |N_H(S)|$ . By the definition of  $D'$  and the construction of  $H$ , we have  $|N_H(S)| = \sum_{v \in N_G(S)} \left\lfloor \frac{d_G(v)}{2} \right\rfloor$ . Let  $m$  denote the number of edges of  $G$  between  $S$  and  $N_G(S)$ . Since every vertex in  $D'$  has all its neighbors in  $A$ , we have  $m \geq \delta|S|$ . Furthermore,  $m \leq \sum_{v \in N_G(S)} d_G(v)$ . Combining these estimates, we obtain

$$\sum_{v \in N_G(S)} \delta \left\lfloor \frac{d_G(v)}{2} \right\rfloor = \delta|N_H(S)| < \delta|S| \leq m \leq \sum_{v \in N_G(S)} d_G(v). \quad (1)$$

For integers  $\delta$  and  $d$  with  $3 \leq \delta \leq d$ , it is easy to verify that  $\delta \left\lfloor \frac{d}{2} \right\rfloor \geq d$ , which yields a contradiction to (1). This completes the proof of Claim 4.  $\square$

The properties of the *Gallai-Edmonds decomposition* imply that  $G[C]$  has a perfect matching  $M_C$ , that there is a matching  $M_A$  using edges between  $A$  and  $D$  that connects each vertex from  $A$  to a distinct component of  $G[D]$ , and that every component of  $G[D]$  is *factor-critical*; recall that a graph  $H$  is factor-critical if  $H - u$  has a perfect matching for every vertex  $u$  of  $H$ .

We now construct a subset  $E_1$  of the edge set  $E(G)$  of  $G$  as follows, starting with the empty set:

- We add to  $E_1$  all  $|D'|$  selected edges as in Claim 4.
- We add  $M_C$  to  $E_1$ .
- For every vertex  $v$  from  $A$  that is not incident with a selected edge, we add to  $E_1$  the unique edge from  $M_A$  incident with  $v$ . Let  $M'_A$  be the subset of  $M_A$  added to  $E_1$ .

- For every component  $K$  of  $G[D]$  of order at least 3 such that some vertex  $x$  of  $K$  is incident with an edge from  $M'_A$ , we add to  $E_1$  a perfect matching of  $K - x$ .
- For every component  $K$  of  $G[D]$  of order at least 3 such that no vertex of  $K$  is incident with an edge from  $M'_A$ , we add to  $E_1$  a perfect matching of  $K - x$  for some vertex  $x$  of  $K$  as well as one further edge of  $K$  incident with  $x$ .

Up to some small modifications explained below, this completes the description of  $E_1$ .

By construction, the spanning subgraph  $G_1$  of  $G$  with edge set  $E_1$  satisfies

$$1 \leq d_{G_1}(u) \leq \left\lfloor \frac{d_G(u)}{2} \right\rfloor \text{ for every vertex } u \text{ of } G. \quad (2)$$

Let  $G_2$  be the spanning subgraph of  $G$  with edge set  $E(G) \setminus E_1$ .

For every component  $K$  of  $G_2$  such that all vertices of  $K$  have even degree in  $G_2$ ,  $K$  has an odd number of edges, and all vertices from  $V(K)$  have degree 1 in  $G_1$ , we select any edge  $e_K$  from  $K$  and move it from  $G_2$  to  $G_1$ . Note that  $K - e_K$  contains exactly two vertices of odd degree, and, hence, is still connected. Furthermore, since  $G$  has minimum degree at least 4, it follows that (2) still holds after each such modification. Having performed these modifications for each such component of  $G_2$ , every component  $K$  of (the modified)  $G_2$  now

- either contains at least one vertex of odd degree in  $K$ ,
- or all vertices of  $K$  have even degrees in  $K$ , and the number of edges of  $K$  is even,
- or all vertices of  $K$  have even degrees in  $K$ , the number of edges of  $K$  is odd, and  $K$  contains a vertex  $u_K$  such that the degree of  $u_K$  in  $G_1$  is at least 2.

The components of  $G_2$  as in the final point are called *type 2* components, and the remaining components of  $G_2$  are called *type 1* components.

We are now in a position to describe a majority 3-edge-coloring  $c : E(G) \rightarrow [3]$ .

- For all edges  $e$  of  $G_1$ , let  $c(e) = 3$ .
- For every component  $K$  of  $G_2$  that is of type 1, let  $c : E(K) \rightarrow [2]$  be as in Theorem 1(i) (applied to  $K$  as  $G$ ).
- For every component  $K$  of  $G_2$  that is of type 2, let  $c : E(K) \rightarrow [2]$  be as in Theorem 1(ii) (applied to  $K$  and  $u_K$  as  $G$  and  $u_G$ ).

It is now easy to verify that  $c$  is a majority 3-edge-coloring of  $G$ , which completes the proof.  $\square$

### 3 Conclusion

The most natural question motivated by our results is which graphs of minimum degree at least 2 do not have a majority 3-edge-coloring.

As a variation of majority edge-colorings, we propose the study of  $\alpha$ -majority edge-colorings for  $\alpha \in (0, 1)$ , where at most an  $\alpha$ -fraction of the edges incident with each vertex are allowed to have the same color. If  $k$  is a positive integer at least 2, then every positive integer at least  $k(k-1)$  can be written as a non-negative integral linear combination of  $k$  and  $k+1$ . Using this fact, a straightforward adaptation of the proof of Theorem 2 yields the following statement: *If a graph  $G$  has minimum degree at least  $k(k-1)$ , then  $G$  has a  $\frac{1}{k}$ -majority  $(k+2)$ -edge-coloring.* A probabilistic argument implies that, for a sufficiently large minimum degree, one color less suffices.

**Theorem 5.** *For every integer  $k$  at least 2, there is a positive integer  $\delta_k$  such that every graph of minimum degree at least  $\delta_k$  has a  $\frac{1}{k}$ -majority  $(k+1)$ -edge-coloring.*

*Proof.* Let  $G$  be a graph of minimum degree  $\delta$  at least  $\delta_k$ , where we specify  $\delta_k$  later. Let  $c : E(G) \rightarrow [k+1]$  be a random  $(k+1)$ -edge-coloring, where we choose the color of each edge uniformly and independently at random. For every vertex  $u$  of  $G$ , we consider the bad event  $A_u$  that more than  $\frac{1}{k}d_G(u)$  of the edges incident with  $u$  have the same color.

For  $d = d_G(u)$ , the union bound and the Chernoff inequality, cf. [11], imply

$$\begin{aligned} \mathbb{P}[A_u] &\leq (k+1)\mathbb{P}\left[\text{Bin}\left(d, \frac{1}{k+1}\right) > \frac{d}{k}\right] && \text{(union bound)} \\ &= (k+1)\mathbb{P}\left[\text{Bin}\left(d, \frac{1}{k+1}\right) > \left(1 + \frac{1}{k}\right) \frac{d}{k+1}\right] \\ &\leq (k+1)e^{-\frac{d}{3k^2(k+1)}}. && \text{(Chernoff inequality)} \end{aligned}$$

For every vertex  $u$  of  $G$ , the event  $A_u$  is determined only by the colors of the edges incident with  $u$ , which are chosen uniformly and independently at random. Therefore, the event  $A_u$  is mutually independent of all events  $A_v$  with  $v \in V(G) \setminus (\{u\} \cup N_G(u))$ . In order to complete the proof, we use the *weighted Lovász Local Lemma*, cf. [11], which states that with positive probability none of the bad events  $A_u$  occurs provided that there is a positive integer  $t_u$  for every vertex  $u$  of  $G$  and there is some real  $p$  with  $0 \leq p \leq \frac{1}{4}$  such that

- $\mathbb{P}[A_u] \leq p^{t_u}$  for every vertex  $u$  of  $G$  and
- $\sum_{v \in N_G(u)} (2p)^{t_v} \leq \frac{t_u}{2}$  for every vertex  $u$  of  $G$ .

Let  $p = (k+1)e^{-\frac{\delta}{3k^2(k+1)}}$  and, for every vertex  $u$  of  $G$ , let  $t_u = \left\lfloor \frac{d_G(u)}{\delta} \right\rfloor$ . Note that  $d_G(u) \geq \delta$  implies that  $t_u$  is a positive integer, and that  $2t_u = 2 \left\lfloor \frac{d_G(u)}{\delta} \right\rfloor \geq \frac{d_G(u)}{\delta}$ .

Choosing  $\delta_k$  sufficiently large, we may ensure that  $p \leq \frac{1}{4}$ , and, hence,  $\mathbb{P}[A_u] \leq p^{\frac{d_G(u)}{\delta}} \leq p^{t_u}$ . Furthermore, we obtain

$$\sum_{v \in N_G(u)} (2p)^{t_v} \leq 2pd_G(u) \leq 4p\delta t_u = \underbrace{\left(4(k+1)e^{-\frac{\delta}{3k^2(k+1)}}\delta\right)}_{\rightarrow 0 \text{ for } \delta \rightarrow \infty} t_u,$$

which is at most  $t_u/2$  for  $\delta_k$  sufficiently large.

Altogether, choosing  $\delta_k$  sufficiently large, the weighted Lovász Local Lemma implies that with positive probability none of the bad events  $A_u$  occurs, which implies the existence of a  $\frac{1}{k}$ -majority  $(k+1)$ -edge-coloring and completes the proof.  $\square$

The estimates in the above proof allow to show that  $\delta_k$  can be chosen to be  $O(k^3 \log k)$ . Our Theorem 3 implies that 4 is the smallest possible value for  $\delta_2$ .

## Acknowledgements

The research reported in this paper was carried out at the *25th C5 Graph Theory Workshop* organized by Ingo Schiermeyer in Rathen, May 2022. After a pause due to the COVID-19 pandemic, we very strongly felt the value of the creative, collaborative, and enjoyable atmosphere and the direct personal exchange at that workshop. We express our gratitude to Ingo, not just for this year but also for the long tradition of this wonderful meeting.

## References

- [1] P. Aboulker, G. Aubian, and C.-C. Huang. Vizing's and Shannon's Theorems for defective edge colouring. [arXiv:2201.11548](#), 2022.
- [2] R. Aharoni, E.C. Milner, and K. Prikry. Unfriendly partitions of a graph. *J. Combinatorial Theory, Ser. B*, 50:1–10, 1990.
- [3] O. Amini, L. Esperet, and J. van den Heuvel. Frugal Colouring of Graphs. [arXiv:0705.0422](#), 2007.
- [4] M. Anholcer, B. Bosek, and J. Grytczuk. Majority choosability of digraphs. *Electron. J. Combin.*, 24:#P3.57, 2017.
- [5] P. Hall. On representatives of subsets. *J. London Math. Soc.*, 10:26–30, 1935.
- [6] C. Hierholzer. Ueber die Möglichkeit, einen Linienzug ohne Wiederholung und ohne Unterbrechung zu umfahren. *Math. Ann.*, 6:30–32, 1873.
- [7] A.J.W. Hilton, T. Slivnik, and D.S.G. Stirling. Aspects of edge list-colourings. *Discrete Math.*, 231:253–264, 2001.
- [8] S. Kreutzer, S. Oum, P. Seymour, D. van der Zypen, and D.R. Wood. Majority colourings of digraphs. *Electron. J. Combin.*, 24:#P2.25, 2017.
- [9] L. Lovász. On decomposition of graphs. *Studia Sci. Math. Hungar.*, 1:237–238, 1966.

- [10] L. Lovász and M.D. Plummer. Matching theory. Annals of Discrete Mathematics, 29, North-Holland Publishing Co., Amsterdam, 1986.
- [11] M. Molloy and B. Reed. Graph colouring and the probabilistic method. Springer-Verlag, Berlin, 2002.
- [12] V.G. Vizing. On an estimate of the chromatic class of a  $p$ -graph. *Diskret. Analiz*, 3:25–30, 1964.
- [13] D.R. Wood. Bounded degree acyclic decompositions of digraphs. *J. Combinatorial Theory, Ser. B*, 90:309–313, 2004.