

The Lexicographically Least Square-Free Word with a Given Prefix

Siddharth Berera^a Andrés Gómez-Colunga^b Joey Lakerdas-Gayle^c
John López^d Mauditra Matin^e Daniel Roebuck^f Eric Rowland^g
Noam Scully^h Juliet Whiddenⁱ

Submitted: Nov 5, 2022; Accepted: May 12, 2023; Published: Jul 28, 2023

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

The lexicographically least square-free infinite word on the alphabet of non-negative integers with a given prefix p is denoted $L(p)$. When p is the empty word, this word was shown by Guay-Paquet and Shallit to be the ruler sequence. For other prefixes, the structure is significantly more complicated. In this paper, we show that $L(p)$ reflects the structure of the ruler sequence for several words p . We provide morphisms that generate $L(n)$ for letters $n = 1$ and $n \geq 3$, and $L(p)$ for most families of two-letter words p .

Mathematics Subject Classifications: 68R15

1 Introduction

A word is *square-free* if it contains no block of letters that occurs twice consecutively. In 2009, Guay-Paquet and Shallit [4] established the structure of the lexicographically least square-free infinite word on the alphabet $\mathbb{N} := \{0, 1, 2, \dots\}$. This word's letters comprise the ruler sequence 0102010301020104 \dots [9, A007814], and it is the fixed point $\rho^\infty(0)$ of the *ruler morphism* ρ defined by $\rho(n) = 0(n+1)$.

However, this result is not robust; a minor variation produces words that are quite different. Given a word w , let $L(w)$ denote the lexicographically least infinite word on \mathbb{N}

^aUniversity of Edinburgh, Edinburgh, U.K. (s2020550@ed.ac.uk).

^bPennsylvania State University, University Park, U.S.A. (azg6165@psu.edu).

^cUniversity of Waterloo, Waterloo, Canada (ja29991a@uwaterloo.ca).

^dTulane University, New Orleans, U.S.A. (jlopez12@tulane.edu).

^eDelft University of Technology, Delft, The Netherlands (a.mauditra@gmail.com).

^fUniversity of St Andrews, St Andrews, U.K. (dr211@st-andrews.ac.uk).

^gHofstra University, Hempstead, U.S.A. (eric.rowland@hofstra.edu).

^hYale University, New Haven, U.S.A. (noamscully166@gmail.com).

ⁱVassar College, Poughkeepsie, U.S.A. (jwhidden@vassar.edu).

beginning with w whose only square factors are contained in the prefix w . In particular, if w is square-free, then so is $L(w)$. If w is infinite, then $L(w) = w$. For example,

$$L(1) = 10120102012021012010201203010201 \dots \quad [9, \text{A356677}],$$

$$L(2) = 20102012021012010201202102010210 \dots \quad [9, \text{A356678}],$$

and

$$L(33) = 33010201030102012021012010201202 \dots \quad [9, \text{A356679}].$$

Unlike $L(\varepsilon) = \rho^\infty(0)$, the letters in these words do not alternate between 0s and positive integers. Moreover, the letters 3, 4, 5, ... take much longer to appear, and there is no clear pattern for these words, as in the case for the ruler sequence.

In this paper we determine the structure of $L(w)$ for certain simple words w . Our main results are that $L(1)$ and $L(n)$ for $n \geq 3$ reflect the structure of $L(\varepsilon)$ as follows.

Theorem 1. *There exists a morphism α and a word Y_1 with length 5177 such that $L(1) = Y_1 \alpha(L(\varepsilon))$.*

Theorem 2. *For each $n \geq 3$, there exists a finite word Y_n such that $L(n) = Y_n \rho(\alpha(L(\varepsilon)))$, where α is the morphism in Theorem 1.*

These results imply that the suffix $\alpha(L(\varepsilon))$ and the related suffix $\rho(\alpha(L(\varepsilon)))$ are two attractors for infinite square-free extensions of words on \mathbb{N} . We have

$$\alpha(L(\varepsilon)) = 01020301201020120210120102012023 \dots \quad [9, \text{A356676}].$$

The structure of $L(2)$ appears to also reflect the structure of $L(\varepsilon)$ but, surprisingly, appears not to be related to the morphism α . Instead, we give a morphism γ in Section 4.3 for which we conjecture the following.

Conjecture 3. *There exists a morphism γ such that $L(2) = 2\gamma(L(\varepsilon))$.*

In Corollary 44, we show that $L(012) = 01201 \lim_{n \rightarrow \infty} \rho^{-1}(\alpha(n))$ where α is the morphism described above. We also describe $L(p)$ for many two-letter words. For example, we show that $L(nn) = nL(n)$ for all letters n . On the other hand, we do not have conjectures for the structures of $L(1n)$ when $n > 1$ and $L(2n)$ when $n \notin \{0, 2\}$. For example we have

$$L(12) = 12010201202101201020120212010201 \dots \quad [9, \text{A356680}],$$

$$L(13) = 13010201030102012021012010201202 \dots \quad [9, \text{A356681}],$$

$$L(21) = 21012010201202101201020121012010 \dots \quad [9, \text{A356682}].$$

Section 2 contains the main definitions and some preliminary results we will use to prove Theorems 1 and 2. Section 3 establishes the structure of finite words $T(n)$ for which $Y_n = nT(n)A$ for a constant word A when $n \geq 3$. We see in Section 4.2 that $T(n)$ is the only component of $L(n)$ that depends on n . In Section 4 we give explicit constructions of α and the words Y_n , and we prove Theorems 1 and 2. Unsurprisingly, the proofs are fairly technical.

In Section 5 we give two conditions under which $L(uv) = uL(v)$. For example, we show that $L(n_1n_2) = n_1L(n_2)$ for all $n_1 \geq 3$ and $n_2 \geq 3$. We use this to describe the structure of $L(p)$ for several families of two-letter words p . Finally, in Section 6, we study the inverse problem of finding a prefix p that induces a given finite square-free word w under L . We show that this can always be done and present an algorithm that computes a prefix p such that pw is a prefix of $L(p)$. Section 7 is a glossary of all of the functions, morphisms, and constants defined in the paper.

This work was motivated by results of several papers [7, 6, 8] studying the lexicographically least word on \mathbb{N} that avoids $\frac{a}{b}$ -powers, for various rational numbers in the interval $1 < \frac{a}{b} < 2$. These words exhibit a remarkable diversity of behaviors. Some of these words (for example, when $\frac{a}{b} = \frac{24}{17}$ [6, Figure 4 on page 36]) alternate between two different modes before settling into their long-term behavior, suggesting that there may be multiple attractors in the dynamical systems that generate them. The current paper is the first exploration of the set of attractors for the alphabet \mathbb{N} . By varying the prefix, we show there are multiple attractors for square-free words. In contrast, on a binary alphabet there are patterns with only one attractor; Allouche, Currie, and Shallit [2] showed that the lexicographically least overlap-free word on $\{0, 1\}$ with a fixed prefix, if it exists, always has a suffix that is a suffix of $10010110 \dots$ (the complement of the Thue–Morse word).

Frequently in this paper, we use computations to show that a particular finite word possesses a certain property. Often, this involves verifying that the word is square-free. As an example, in the proof of Lemma 7, we verify that the word $\psi_1(010)$ is square-free. These words can easily be computed from their definitions. All computations in this paper that verify that a word is square-free can be done quickly using the Main–Lorentz algorithm described in their 1985 paper [5].

We sometimes require the computation of finite prefixes of words $L(w)$. For example, the word Y_1 is defined as the 5177-letter prefix of $L(1)$. These computations can be completed quickly using the following simple greedy algorithm: To find the next letter of $L(w)$ after the prefix wv , check whether wvn has a square suffix for non-negative integers n , increasing from zero. Then the least n for which wvn has no square suffix is the next letter of $L(w)$. If w contains a square, we test all even-length suffixes of wvn . If w is square-free, then we can use a more efficient variation of the Main–Lorentz algorithm [5] that only searches for squares that are suffixes. This variation is easiest to implement by first reversing the string and then using the algorithm described in their paper with “Step 3” altered so that only the first block of length l is examined.

2 Definitions and preliminary results

We assume the reader is familiar with basic definitions and notation regarding words and morphisms. See the survey by Allouche [1] for a short introduction. All words in the paper are words on the alphabet \mathbb{N} .

We say that v is a *factor* of w if $w = xvy$ for some words x, y . If v is a factor of w , we also say that v *occurs in* w and w *contains* v . A *square* is a nonempty word of the form

yy. A word is *square-free* if it contains no square factors.

We index letters in a word beginning with 0, and we use Python notation to extract factors: Suppose a word w has length ℓ and is written as a sequence of letters $w = w_0w_1w_2 \cdots w_{\ell-2}w_{\ell-1}$. If $0 \leq i \leq j \leq \ell - 1$, then $w[i] := w_i$ and $w[i : j] := w_iw_{i+1} \cdots w_{j-1}$, with “default values” for i and j being 0 and ℓ , respectively. If $i = j$, then $w[i : j] = \varepsilon$. Negative values index letters from the end of the word. For example, if $w = \text{hydrant}$, then

$$\begin{aligned} w[3 : 6] &= \text{ran} = w[3 : -1] \\ w[4 :] &= \text{ant} = w[-3 :] \\ w[: 5] &= \text{hydra} = w[: -2]. \end{aligned}$$

Note also that $w[: 0] = \varepsilon$ for any word w .

Definition. A word w is *even-grounded* if $w_i = 0$ for even i and $w_i \neq 0$ for odd i . A word w is *odd-grounded* if $w_i = 0$ for odd i and $w_i \neq 0$ for even i . A word is *grounded* if it is even-grounded or odd-grounded.

For example, the words 010 and 0102 are even-grounded, 301 and 3010 are odd-grounded, and the words 0120 and 0100 are not grounded.

Definition. The *ruler morphism* $\rho: \mathbb{N}^* \cup \mathbb{N}^\omega \rightarrow \mathbb{N}^* \cup \mathbb{N}^\omega$ is defined by $\rho(n) = 0(n + 1)$ for letters $n \in \mathbb{N}$.

Notation. We will denote by R_n the prefix of the ruler sequence up to the first occurrence of the letter n , i.e. $R_n = \rho^n(0) = L(\varepsilon)[: 2^n]$. For example, $R_0 = 0$, $R_1 = 01$, $R_2 = 0102$, and so on.

Notation. For a nonempty finite word w , we define w^+ , the *successor* of w , to be the word that is identical to w except for the last letter, which is increased by 1. For example, if $w = 0102$ then $w^+ = 0103$. Formally, $w^+ = w[: -1](w[-1] + 1)$.

Definition. Let ϕ be a morphism.

- ϕ is *non-erasing* if $\phi(k) \neq \varepsilon$ for all letters k .
- Let Δ be a set of words. We say that ϕ is *square-free over* Δ if ϕ is non-erasing and $\phi(w)$ is square-free for all square-free words $w \in \Delta$. We say that ϕ is *square-free* if ϕ is square-free over $\mathbb{N}^* \cup \mathbb{N}^\omega$.
- ϕ is *letter-injective* if, given letters k and ℓ , $\phi(k) = \phi(\ell)$ implies $k = \ell$.

Definition. Given two words u and v , we say that u is *lexicographically less than* v and we write $u \prec v$ if there is an index i such that $u[: i] = v[: i]$ and $u[i] < v[i]$ as letters. It can be seen that \prec is a partial ordering on $\mathbb{N}^* \cup \mathbb{N}^\omega$. The only case when u and v are not comparable by \prec is when one word is a prefix of the other. We can also see that if w is a nonempty word of finite length and $v \prec w^+$, then either $v \prec w$ or w is a prefix of v .

Definition. Given words w and u , we say that u is *irreducible in wu* if for all words $s \prec u$, s introduces a square in ws . That is, there is a square in ws that ends at a letter in s .

Notice that if u is irreducible in wu , it can still introduce a square. In this case, u^+ is irreducible in wu^+ .

Example. Consider $w = 0102010$, $u = 23$, and $v = 301$. Any word y beginning with a 0, 1, or 2 introduces a square in wy , so u is irreducible and also introduces a square in wu . Thus, $u^+ = 24$ is also irreducible. The word $v = 301$ does not introduce a square, but every word lexicographically less than v either begins with 0, 1, or 2, or includes a square. So v is irreducible, but $v^+ = 302$ is not.

Definition. We say that p *generates ps* if $L(p) = L(ps)$. In other words, $L(p)$ starts with ps .

The main results of this paper, that $L(1) = Y_1\alpha(L(\varepsilon))$ and $L(n) = Y_n\rho(\alpha(L(\varepsilon)))$ for $n \geq 3$ can be restated as 1 generates $Y_1\alpha(L(\varepsilon))$, and n generates $Y_n\rho(\alpha(L(\varepsilon)))$ for $n \geq 3$.

Remark 4. To prove that a square-free word p generates another word $w = ps$, we must show that w is square-free and that s is irreducible in $w = ps$.

In particular, we have the useful property that if p generates $w = ps$ and uw is square-free for another word u , then up generates uw . The square-free condition is known by assumption, and the fact that s is irreducible in $w = ps$ implies that s is irreducible in $uw = ups$.

One common application of this property is when $w = L(p)$. Then if $uL(p)$ is square-free, we get $L(up) = L(uL(p)) = uL(p)$. Or in other words, up generates $uL(p)$.

Example. The above remark is often used implicitly in this paper. For example in Lemma 11, we prove that for $n \geq 1$, $\psi_1(n)$ generates $\psi_1(n)202101$. To do this, we first show that $\psi_1(n)202101$ is square-free and that 2 is irreducible in $\psi_1(n)2$. This proves that $\psi_1(n)$ generates $\psi_1(n)2$.

We then show that $\psi_1(n)2$ has suffix R_2R_12 , and via computation show that R_2R_12 generates $R_2R_1202101$. Letting $p = R_2R_12$, $s = 02101$, and u be the word for which $up = \psi_1(n)2$, we have that p generates ps , and $ups = \psi_1(n)202101$ is square-free. So Remark 4 says that $up = \psi_1(n)2$ generates $ups = \psi_1(n)202101$.

We now have that $\psi_1(n)$ generates $\psi_1(n)2$, and $\psi_1(n)2$ generates $\psi_1(n)202101$. By definition, this means that $L(\psi_1(n)) = L(\psi_1(n)2) = L(\psi_1(n)202101)$, so $\psi_1(n)$ generates $\psi_1(n)202101$.

Notation. For any word w , $\max(w)$ denotes the maximum letter value in w if it exists.

The remainder of this section introduces the notion of *chunks* which will be key in Section 3 to prove Theorem 24 regarding the structure of a certain prefix of $L(n)$ and in Section 4 to prove Theorem 28 related to the square-freeness of the morphism α , determining the structure of $L(1)$.

Given a morphism ϕ and a word $w = w_0w_1w_2 \cdots$ we can write $\phi(w)$ as

$$\phi(w) = [\phi(w_0)] [\phi(w_1)] [\phi(w_2)] \cdots,$$

where we use square brackets to delineate the contributions of each individual letter of w . A factor of $\phi(w)$ that arises as the image $\phi(k)$ of the letter k under ϕ is a *chunk* or, more specifically, a *k-chunk*. For example, consider the ruler morphism ρ and the word 0121. We would break $\rho(0121)$ into chunks as

$$\rho(0121) = [01] [02] [03] [02].$$

Each factor 02 is a 1-chunk.

Sometimes, when we take a particular occurrence of a factor of $\phi(w)$, we find that the factor starts or ends partway through a chunk. For example $\rho(0121)[1 : 6]$ can be written as

$$\rho(0121)[1 : 6] = 1] [02] [03] [0.$$

We refer to 1] and [0 as *partial chunks*. 1] is the *initial* partial chunk and [0 is the *final* partial chunk.

In this paper, we frequently consider words of the form $\phi(w)$, and then characterize the possible locations of certain factors of $\phi(w)$ with respect to its chunks. For example, in Lemma 30, we show that for any grounded square-free word w , the constant word E can only occur in $\alpha(w)$ as a prefix or a suffix of a 0-chunk. This is related to the morphism property that ϕ *locates words of length ℓ* introduced by Pudwell and Rowland [6], which restricts the possible starting index for a word of length ℓ , relative to the chunks in $\phi(w)$. It is also related to the *synchronization point* of a word x introduced by Cassaigne [3] which describes a point in a word x with respect to a morphism ϕ that must occur at a chunk boundary whenever x occurs in $\phi(w)$.

Definition. Given a word w and a morphism ϕ , we say that two (possibly partial) chunks in $\phi(w)[i : j]$ *come from the same letter* to mean that both chunks arise as images of the same letter in w . For example, in $\rho(0121)[1 : 6] = 1][02][03][0$, the first whole chunk, [02], and the final partial chunk, [0, come from the same letter, 1. If a word v occurs twice in $\phi(w)$, we say that the occurrences have the same *chunk decomposition* if every whole chunk in one occurrence corresponds to a whole chunk in the other occurrence and both chunks come from the same letter.

Example. Consider the morphism $\phi: \mathbb{N}^* \cup \mathbb{N}^\omega \rightarrow \mathbb{N}^* \cup \mathbb{N}^\omega$ defined by

$$\phi(n) = \begin{cases} 0 & \text{if } n = 0 \\ 01 & \text{if } n = 1 \\ n - 1 & \text{if } n \geq 2. \end{cases}$$

The single occurrence of $v = 103$ in $\phi(2041) = [1] [0] [3] 01$ has the same chunk decomposition as its occurrence in $\phi(2204) = 1 [1] [0] [3]$. On the other hand, $w = 01$ in $\phi(01) = 0 [01]$ does not have the same chunk decomposition as its occurrence in $\phi(024) = [0] [1] 3$.

In $\rho(01012)$, we consider the square 1020 1020:

$$\rho(01012) = [0 (1)[02][0] (1)[02][0] 3].$$

The two halves of the square have the same chunk decomposition because in both, 02 is a whole 1-chunk and there are no other whole chunks. The two initial partial chunks come from the same letter because they are both part of 0-chunks. The two final partial chunks do not come from the same letter because the final partial chunk of the first half is part of a 0-chunk, and the final partial chunk of the second half is part of a 2-chunk.

Consider a morphism ϕ where $\phi(0) = 123$, and $\phi(1) = 13$. Then, $\phi(010) = [12(3)[1](3)[1]23]$. In the square 3131, both halves vacuously have the same chunk decomposition since there are no whole chunks. But neither their initial nor final partial chunks come from the same letter.

The following theorem is used frequently in this paper to show that a morphism is square-free.

Theorem 5. *Let ϕ be a letter-injective morphism such that $\phi(\ell)$ is square-free for all letters ℓ . Suppose w is a word such that $\phi(w)$ contains a square yy for which the following three properties hold.*

1. *Each half of the square contains at least one whole chunk.*
2. *The two halves of the square have the same chunk decomposition.*
3. *If either half has a partial chunk, then either their initial partial chunks or their final partial chunks come from the same letter in w .*

Then w contains a square.

Proof. By the first and second conditions, the corresponding whole chunks in both halves come from the same letters in w . So there is a nonempty factor u of w that yields all whole chunks in both halves. That is, we can write the square as

$$yy = [b] [\phi(u)] [c] [b] [\phi(u)] [c]$$

where $b]$ and $[c$ are possibly empty partial chunks.

If there are no partial chunks ($b = c = \varepsilon$), then $yy = [\phi(u)][\phi(u)]$ and uu is a square factor of w .

Now, suppose there is a partial chunk in either half. Then since the halves have the same chunk decomposition, there must be an initial and final partial chunk in both halves (neither b nor c is empty). Let a and d be the smallest words that complete the partial chunks of yy . That is, $\phi(w)$ has the factor $[a(b] [\phi(u)] [c](b] [\phi(u)] [c)d] = ayyd$. Since ϕ is letter-injective, ϕ^{-1} is well defined on chunks. The third condition says that either $\phi^{-1}(ab) = \phi^{-1}(cb)$ or $\phi^{-1}(cb) = \phi^{-1}(cd)$. In the first case, $\phi^{-1}(ab)u\phi^{-1}(cb)u$ is a square factor of w . In the second case, $u\phi^{-1}(cb)u\phi^{-1}(cd)$ is a square factor of w . \square

As a corollary to this theorem, suppose Γ is a family of square-free words, and ϕ is a letter-injective morphism with $\phi(\ell)$ square-free for all letters ℓ . For any word $w \in \Gamma$, $\phi(w)$ cannot contain a square that possesses all three properties in Theorem 5, or else w would contain a square. So we can prove that ϕ is square-free over Γ by showing that for all $w \in \Gamma$, if $\phi(w)$ contains a square, then it contains a square with these three properties. In this paper, we use this technique to show that the morphisms ψ_1 , ψ_2 , and α are square-free over grounded words.

3 A certain prefix of $L(n)$

For all $n \geq 1$, it is clear that $L(n)$ always starts with the letter n followed by a prefix of the ruler sequence $L(\varepsilon)$. In this section, we show that for $n \geq 3$, $L(n)$ has prefix $nT(n)$ which has length exponential in n . In Section 4.2 we will show that this is the only part of $L(n)$ that depends on n .

3.1 The morphism ψ_1

We begin by showing that $L(n)$ has a shorter prefix, $nP_0(n)P_1(n)$, which is proved in Theorem 14. Next we define the words $P_0(n)$, $P_1(n)$ and the morphism ψ_1 .

Definition. For $n \geq 0$, let $P_0(n)$ be the maximum prefix of the ruler sequence such that $nP_0(n)$ is a prefix of $L(n)$. Define the morphism $\psi_1: \mathbb{N}^* \rightarrow \mathbb{N}^*$ by

$$\psi_1(n) = \begin{cases} 202101 & \text{if } n = 0, \\ (n+1)P_0(n+1) & \text{otherwise,} \end{cases}$$

and for $n \geq 3$ define

$$P_1(n) = \psi_1(P_0(n-1)).$$

For $n \geq 3$, after the word $L(n)$ deviates from the ruler sequence prefix, we will show that it continues with the word $P_1(n)$. These definitions can be referenced in the glossary, Section 7.

Remark 6. Note that $P_0(0) = \varepsilon$. Also, it is not hard to see that the length of $P_0(n)$ is $2^{n+1} - 2$, hence for all n we have

$$P_0(n) = R_{n+1}[: -2] = R_n R_n[: -2] = R_n R_{n-1} \cdots R_3 R_2 R_1.$$

And for $n \geq 1$,

$$P_0(n) = R_n P_0(n-1).$$

By repeated application of this argument, for $1 \leq k \leq n$, we have that $P_0(n)$ has suffix $P_0(k)$.

We would like to show that ψ_1 is square-free over grounded words, since that will imply that $P_1(n)$ is square-free. The next three lemmas describe some of the important behavior of ψ_1 over grounded square-free words.

Lemma 7. *Let $n \geq 1$. Then $\psi_1(0n0)$ is square-free.*

Proof. We can verify computationally that $\psi_1(010)$ is square-free. So let $n \geq 2$ and suppose there is a square yy in

$$\psi_1(0n0) = [202101] [(n+1)P_0(n+1)] [202101].$$

Since $(n+1)P_0(n+1)$ is a prefix of $L(n+1)$, $\psi_1(n)$ is square-free for all n . So yy lies over at least one of the chunk boundaries.

In the first chunk boundary we find the word $1(n+1)$, and in the second the word 12 . Since $n+1 \geq 3$ and $\psi_1(n)$ is grounded, these words appear nowhere else in $\psi_1(0n0)$. Therefore, neither chunk boundary can be completely contained in y . This means that yy must be a factor of $\psi_1(0n)$ or $\psi_1(n0)$ and each occurrence of y must be completely contained in a different chunk. Every prefix of $\psi_1(n)$ begins with $n+1 \geq 3$ but no suffix of $\psi_1(0)$ does, and every suffix of $\psi_1(n)$ is grounded and ends with a 1 which is not true for any prefix of $\psi_1(0)$. Therefore, such a square cannot exist. \square

Lemma 8. *Let $n > k \geq 0$. Then neither of $\psi_1(n)$ and $\psi_1(k)$ is a factor of the other.*

Proof. From the definition, it is clear that since $n > k$, $|\psi_1(n)| > |\psi_1(k)|$, so $\psi_1(n)$ cannot be a factor of $\psi_1(k)$. If $k = 0$, then $\psi_1(k)$ is not grounded but $\psi_1(n)$ is. Thus, assume $n > k > 0$. Then since $\psi_1(n) = (n+1)P_0(n+1) = (n+1)R_{n+2}[: -2]$, it is sufficient to show that $\psi_1(k) = (k+1)R_{k+2}[: -2]$ cannot occur in R_{n+2} .

Since $R_{n+1} = R_n R_n^+$ for all n , we have that

$$\begin{aligned} R_{n+2} &= R_{k+2} R_{k+2}^+ R_{k+2} R_{k+2}^{++} R_{k+2} R_{k+2}^+ \cdots R_{k+2} R_{k+2}^{++++} \\ &= R_{k+2}^* R_{k+2}^* \cdots R_{k+2}^*, \end{aligned}$$

where $*$ represents the application of zero or more $^+$'s according to the ruler sequence pattern.

The largest letter in $R_{k+2}[: -2]$ is $k+1$ and the last letter of each R_{k+2}^* is at least $k+2$. So every occurrence of $R_{k+2}[: -2]$ in $R_{k+2}^* R_{k+2}^* \cdots R_{k+2}^*$ must be as a prefix of an R_{k+2}^* . This means that any occurrence of $R_{k+2}[: -2]$ in R_{n+2} is either a prefix of R_{n+2} or is preceded by a letter that is at least $k+2$. Therefore, $(k+1)R_{k+2}[: -2]$ can never occur in R_{n+2} , and $\psi_1(k)$ cannot occur in $\psi_1(n)$ which proves the result. \square

Lemma 9. *Let w be a grounded square-free word. Then for $\ell \geq 0$, every occurrence of $\psi_1(\ell)$ in $\psi_1(w)$ is a ℓ -chunk.*

Proof. It is not hard to see that the word 21 can only occur in $\psi_1(w)$ in the middle of 0-chunks. Thus, 202101 occurs in $\psi_1(w)$ only as a 0-chunk.

Suppose $\ell \geq 1$. Then $\psi_1(\ell) = (\ell+1)P_0(\ell+1) = (\ell+1)R_{\ell+1}R_{\ell+1}[: -2]$ which is odd-grounded. For $n \geq 0$, $\psi_1(n)$ begins and ends with a non-zero letter, so any word lying over a chunk boundary in $\psi_1(w)$ cannot be grounded. Therefore, every occurrence of $\psi_1(\ell)$ in $\psi_1(w)$ must be totally contained in some k -chunk where $k \geq \ell$. By Lemma 8 $\psi_1(\ell)$ cannot be a factor of $\psi_1(k)$ when $k > \ell$, so $\psi_1(\ell)$ can only occur as an ℓ -chunk. \square

Proposition 10. ψ_1 is square-free over grounded words.

Proof. Suppose w is a grounded square-free word and that $\psi_1(w)$ contains a square yy . From its definition, ψ_1 is letter injective and $\psi_1(n)$ is square-free for all n . Also, Lemma 9 implies that both halves of yy have the same chunk decomposition. We will show that each half of the square contains a whole chunk, and that if either half contains a partial chunk, then the final partial chunk of both halves comes from the same letter. Then Theorem 5 will imply that w contains a square, which is a contradiction.

Suppose neither half of the square contains a whole chunk. Then the whole square contains no more than one whole chunk. By Lemma 7, $\psi_1(0n0)$ is square-free, so yy must be a proper factor of $\psi_1(n0k)$ for $n, k > 0, n \neq k$. The factor 21 only occurs in the middle of 0-chunks, so the square in $\psi_1(n0k)$ has its center at the 21 in the 0-chunk. The square cannot be totally contained in the 0-chunk, so the second half of the square begins with 101 which cannot occur in the first half because $\psi_1(n) = (n + 1)R_{n+2}[: -2]$. This is a contradiction so at least one half of the square contains a whole chunk. Since both halves have the same chunk decomposition, both halves contain a whole chunk.

Suppose either half contains a partial chunk. Then since the halves have the same chunk decomposition, they must both end with a partial chunk. The final partial chunks begin with the same letter, so they must be equal chunks or one of them must be a 0-chunk. But since the halves have the same chunk decomposition and contain whole chunks, their last whole chunk is equal. So the final partial chunks are either both 0-chunks or both equal nonzero chunks. \square

The next three lemmas are used to prove the irreducibility condition in Theorem 14.

Lemma 11. Let $n \geq 1$. Then $\psi_1(n)$ generates $\psi_1(n0)$.

Proof. Since $n0$ is square-free and grounded, $\psi_1(n0)$ is square-free by Proposition 10, so we only need to show that $\psi_1(0)$ is irreducible in $\psi_1(n0) = \psi_1(n)202101$. We will first show that 2 is an irreducible suffix of $\psi_1(n)2$, and then that 02101 is an irreducible suffix of $\psi_1(n)202101$.

Recall that $\psi_1(n) = (n + 1)P_0(n + 1) = (n + 1)R_{n+1}[: -1](n + 1)R_{n+1}[: -2]$. From the structure of R_n , we know that R_{n+1} has suffix $010(n + 1)$ when $n \geq 1$. So $\psi_1(n)0 = (n + 1)R_{n+1}[: -1](n + 1)R_{n+1}[: -1]$ which is a square. The last letter of $\psi_1(n)$ is a 1, so $\psi_1(n)1$ has square suffix 11. Therefore, 2 is irreducible at the end of $\psi_1(n)2$. Since $\psi_1(n)$ ends with R_2R_1 , then $\psi_1(n)2$ ends with R_2R_12 . We can computationally verify that R_2R_12 generates $R_2R_1\psi_1(0)$, which implies the result by using Remark 4. \square

Lemma 12. For $0 \leq k \leq n$, $k + 1$ is irreducible at the end of $\psi_1(nR_k[: -1])(k + 1)$.

Proof. Let $0 \leq m \leq k$, we will show that $w_m := \psi_1(nR_k[: -1])m$ has a square suffix. If $k = 0$, then $m = 0$ and $\psi_1(nR_k[: -1]) = \psi_1(n)$. Then $\psi_1(nR_k[: -1])m = \psi_1(n)0$ which has a square suffix by Lemma 11.

Now note that for $k \geq 1$, the last letter of $R_k[: -1]$ is 0 and $\psi_1(0) = 202101$, so w_m has a square suffix for $m < 2$.

Now assume $2 \leq m \leq k$. By definition $\psi_1(n) = (n+1)P_0(n+1)$, and by Remark 6, $P_0(n+1)$ has suffix $P_0(m)$. Hence, if $m = k$, w_m has suffix $P_0(m)\psi_1(R_{m-1}[: -1])m$.

On the other hand, if $m < k$ it is easy to see that $nR_k[: -1]$ has suffix $mR_m[: -1]$. By definition $\psi_1(m) = (m+1)P_0(m+1)$, and since $m \geq 2$, Remark 6 implies that $P_0(m+1)$ has suffix $P_0(m)$. So w_m has suffix $P_0(m)\psi_1(R_m[: -1])m$ for any $2 \leq m \leq k$. Finally note that

$$\begin{aligned} P_0(m)\psi_1(R_m[: -1])m &= P_0(m)\psi_1(R_{m-1}[: -1])\psi_1(m-1)\psi_1(R_{m-1}[: -1])m \\ &= P_0(m)\psi_1(R_{m-1}[: -1])mP_0(m)\psi_1(R_{m-1}[: -1])m, \end{aligned}$$

which is a square. □

Lemma 13. For $n \geq 1$ and $0 \leq k \leq n$, $\psi_1(n)$ generates $\psi_1(nR_k)$.

Proof. Since $k \leq n$ we know that nR_k is square-free and grounded, so Proposition 10 implies that $\psi_1(nR_k)$ is square-free. It is now sufficient to show that $\psi_1(R_k)$ is irreducible in $\psi_1(nR_k)$.

We prove this inductively over n . The base case $n = 1$ implies that $k = 0$ or $k = 1$. We can computationally verify that $\psi_1(1)$ generates $\psi_1(1R_0) = \psi_1(10)$ and that $\psi_1(1)$ generates $\psi_1(1R_1) = \psi_1(101)$.

Fix $n > 1$ and suppose the result holds for all $1 \leq n_0 < n$. That is,

$$\psi_1(R_k) \text{ is irreducible in } \psi_1(n_0R_k) \text{ for all } 0 \leq k \leq n_0 \text{ and } 1 \leq n_0 < n. \quad (\text{i})$$

We will show that the result holds for $n_0 = n$. That is, $\psi_1(R_k)$ is irreducible in $\psi_1(nR_k)$ for all $0 \leq k \leq n$.

We can prove this intermediate step by a second induction, now over k . The base case is $k = 0$, which holds by Lemma 11 since $R_0 = 0$. Now fix $k > 0$ and suppose

$$\psi_1(R_{k_0}) \text{ is irreducible in } \psi_1(nR_{k_0}) \text{ for all } 0 \leq k_0 < k \quad (\text{ii})$$

We will show that the result holds for $k_0 = k$. That is, $\psi_1(R_k)$ is irreducible in $\psi_1(nR_k)$.

We have that $\psi_1(nR_k) = \psi_1(nR_{k-1}R_{k-2} \cdots R_2R_1R_0k)$. By the second inductive hypothesis (ii), $\psi_1(R_{k-1})$ is irreducible in $\psi_1(nR_{k-1})$. The last letter of R_{k-1} is $k-1$, so the first inductive hypothesis (i) says that $\psi_1(R_{k-2})$ is irreducible in $\psi_1((k-1)R_{k-2})$ and so $\psi_1(R_{k-1}R_{k-2})$ is irreducible in $\psi_1(nR_{k-1}R_{k-2})$. Repeating this argument shows that $\psi_1(R_{k-1}R_{k-2} \cdots R_2R_1R_0) = \psi_1(R_k[: -1])$ is irreducible in $\psi_1(nR_k[: -1])$. Lemma 12 implies that $k+1$ is irreducible in $\psi_1(nR_k[: -1])(k+1)$. And $\psi_1(k) = (k+1)P_0(k+1)$ is a prefix of $L(k+1)$ by the definition of P_0 , so $k+1$ generates $\psi_1(k)$, meaning that $\psi_1(k)$ is irreducible in $\psi_1(nR_k[: -1])\psi_1(k) = \psi_1(nR_k)$, which proves the result. □

In particular, this lemma implies that for all $n \geq 1$, $\psi_1(n)$ generates $\psi_1(nR_n)$.

Theorem 14. For all $n \geq 3$, let $P_1(n) = \psi_1(P_0(n-1))$. Then for $n \geq 3$, $L(n)$ has prefix $nP_0(n)P_1(n)$.

Proof. Note that $nP_0(n)P_1(n) = \psi_1((n-1)P_0(n-1))$. Then, since $(n-1)P_0(n-1)$ is square-free and grounded, Proposition 10 implies that $nP_0(n)P_1(n)$ is square-free. It remains to show that $P_1(n) = \psi_1(P_0(n-1)) = \psi_1(R_{n-1}R_{n-1}[: -2])$ is irreducible in $nP_0(n)P_1(n)$. Indeed, note that

$$nP_0(n)P_1(n) = \psi_1((n-1)P_0(n-1)) = \psi_1((n-1)R_{n-1}R_{n-1}[: -2]).$$

Lemma 13 implies that $\psi_1(n-1)$ generates $\psi_1((n-1)R_{n-1})$, which has suffix $\psi_1(n-1)$. Hence, by applying Lemma 13 a second time we obtain in particular that $\psi_1(R_{n-1}[: -2])$ is irreducible in $\psi_1((n-1)R_{n-1}[: -2])$, and so it is also irreducible in

$$nP_0(n)P_1(n) = \psi_1((n-1)R_{n-1}R_{n-1}[: -2]).$$

Therefore, the whole factor $P_1(n)$ is an irreducible suffix of $nP_0(n)P_1(n)$. □

Remark 15. Using Remark 6, for all $n \geq 4$ we have

$$\begin{aligned} P_1(n) &= \psi_1(P_0(n-1)) \\ &= \psi_1(R_{n-1}[: -1](n-1)R_{n-1}[: -2]) \\ &= \psi_1(R_{n-1}[: -1])\psi_1(n-1)\psi_1(P_0(n-2)) \\ &= \psi_1(R_{n-1}[: -1])nP_0(n)P_1(n-1) \\ &= \psi_1(R_{n-1}[: -1])nR_nP_0(n-1)P_1(n-1), \end{aligned}$$

so $P_1(n)$ has $nP_0(n-1)P_1(n-1)$ as suffix.

Repeated application of this argument implies that for all $2 \leq k < n$, the word $P_1(n)$ has the suffix $(k+1)P_0(k)P_1(k)$.

3.2 The morphism ψ_2

For all $n \geq 3$ after the prefix given by the previous result, $L(n)$ continues with another sequence, $P_2(n)$ defined as follows.

Definition. Define the morphism $\psi_2: \mathbb{N}^* \rightarrow \mathbb{N}^*$ by

$$\begin{aligned} \psi_2(0) &= 2021020102101201020120210120102013010201030102012021012010201 \\ &\quad 2021013010201030102012021012010201202301020103010201202101201 \\ &\quad 0201203010201030102030103020102030102010301020301030201202101 \\ &\quad 2010201202101202, \\ \psi_2(n) &= (n+2)P_0(n+2)P_1(n+2), \text{ if } n > 0 \end{aligned}$$

and let

$$P_2(n) = \psi_2(P_0(n-2))$$

which can be referenced in Section 7.

In particular, we note that for $n \geq 1$, $\psi_2(n) = \psi_1((n+1)P_0(n+1)) = \psi_1^2(n)$, but $\psi_2(0) \neq \psi_1^2(0)$. Similarly to ψ_1 , we would like to show that ψ_2 is square-free over grounded words because that will imply that $nT(n) = \psi_2((n-2)P_0(n-2))$ is square-free. The next four lemmas prove some properties of ψ_2 that are used to prove this condition.

Lemma 16. *Let $n \geq 1$. Then $\psi_2(n0)$ is square-free.*

Proof. We can verify computationally that $\psi_2(10)$ is square-free. So let $n \geq 2$ and suppose there is a square yy in $\psi_2(n0)$. Since $\psi_2(n) = \psi_1((n+1)P_0(n+1))$, $\psi_2(n)$ is square-free by Proposition 10. This implies that yy overlaps both chunks. We have that $\psi_2(n)$ ends with

$$P_1(n+2) = \psi_1(P_0(n+1)) = \psi_1(R_{n+2}[: -2]) = \psi_1(R_{n+1}R_n \cdots R_3R_2R_1).$$

Since $n \geq 2$, this ends with $\psi_1(3R_2R_1)$. One verifies computationally that $\psi_1(3R_2R_1)\psi_2(0)$ is square-free, which means that any square in $\psi_2(n0)$ must contain all of $\psi_1(3R_2R_1)[2$ at the chunk boundary. Since $\psi_1(3)$ contains 4's, the square must contain 4's.

Let k be the largest letter in the square yy . We know that $4 \leq k \leq n+2$ since $\max(\psi_2(n0)) = n+2$. Since $\max(\psi_2(0)) = 3$, all occurrences of k in $\psi_2(n0)$ are in the n -chunk. From above, $\psi_2(n)$ ends with $\psi_1(R_m[: -2])$ for $m \leq n+2$. We consider two cases:

Case 1: $k < n+2$. Since $k+1 \leq n+2$, $\psi_2(n)$ ends with

$$\begin{aligned} \psi_1(R_{k+1}[: -2]) &= \psi_1(R_k[: -1]) \quad \psi_1(k) && \psi_1(R_k[: -2]) \\ &= \psi_1(R_k[: -1]) \quad \underline{(k+1)R_{k+1}[: -1]} \underline{(k+1)R_{k+1}[: -2]} && \psi_1(R_k[: -2]) \end{aligned}$$

which contains the last two occurrences of $k+1$ in $\psi_2(n)$. Since k is the largest letter in the square, yy must be contained in the suffix of $\psi_2(n0)$ after the last occurrence of $k+1$ which is

$$\begin{aligned} &R_{k+1}[: -2] && \psi_1(R_k[: -2]) && \psi_2(0) \\ = &R_kR_k[: -2] && \psi_1(R_{k-1}R_{k-1}[: -2]) && \psi_2(0) \\ = &R_k[: -1] \underline{k} R_k[: -2] && \psi_1(R_{k-1}[: -1])\psi_1(k-1) && \psi_1(R_{k-1}[: -2]) \quad \psi_2(0) \\ = &R_k[: -1] \underline{k} R_k[: -2] && \psi_1(R_{k-1}[: -1]) \underline{k} R_k[: -1] \underline{k} R_k[: -2] && \psi_1(R_{k-1}[: -2]) \quad \psi_2(0) \end{aligned}$$

These are the only three occurrences of k that can occur in yy . The square must contain an even number of k occurrences, so since yy overlaps part of $\psi_2(0)$, it cannot include the first occurrence of k . Therefore, the square contains only the last two occurrences of k . Since the last letter of $R_m[: -1]$ is 0 for all m , the second last occurrence of k is preceded by $\psi_1(0)$ which ends in 1, and the last occurrence of k is preceded by a 0. This means that k must be the first letter of y in the square. It can be seen from the above equation that this square would not reach $\psi_2(0)$ which is a contradiction.

Case 2: $k = n + 2$. We have that

$$\begin{aligned}\psi_2(n) &= \psi_1(n + 1) \cdot \psi_1(P_0(n + 1)) \\ &= \psi_1(n + 1) \cdot \psi_1(R_{n+1}R_{n+1}[: -2]) \\ &= \psi_1(n + 1) \cdot \psi_1(R_{n+1}[: -1]) \cdot \psi_1(n + 1) \cdot \psi_1(R_{n+1}[: -2]) \\ &= (n + 2)P_0(n + 2) \cdot \psi_1(R_{n+1}[: -1]) \cdot (n + 2)P_0(n + 2) \cdot \psi_1(R_{n+1}[: -2]).\end{aligned}$$

And so $\psi_2(n)$ is the “almost square”

$$\psi_2(n) = \frac{(n + 2)}{(n + 2)} \frac{R_{n+2}}{R_{n+2}} R_{n+2}[: -2]\psi_1(R_{n+1}[: -1]) \psi_1(R_{n+1}[: -2]). \quad (1)$$

and these are all four of the occurrences of $n + 2$ in $\psi_2(n0)$. The square must contain an even number of occurrences of $n + 2$.

The square cannot contain all four occurrences of $n + 2$ since then $\psi_2(0)$ would need to begin with $\psi_1(R_{n+1}[: -2]) = \psi_1(0)$ which it does not.

So the square only contains the last two occurrences of $n + 2$. The second last occurrence is preceded by $\psi_1(0)$ which ends in a 1, and the last occurrence is preceded by a 0. This means that $n + 2$ must be the first letter of y in the square. It can be seen from the equation for $\psi_2(n)$ that this square would not reach $\psi_2(0)$ which is a contradiction. \square

Lemma 17. *Let $n \geq 1$. Then $\psi_2(0n0)$ is square-free.*

Proof. We can verify computationally that $\psi_2(010)$ is square-free. So let $n \geq 2$ and suppose there is a square yy in $\psi_2(0n0)$. We have

$$\psi_2(0n0) = [\psi_2(0)][(n + 2)P_0(n + 2)P_1(n + 2)][\psi_2(0)]$$

Since $\psi_2(n)$ is square-free, yy lies over at least one of the chunk boundaries.

The largest letter of $\psi_2(0)$ is 3, so the largest letter of $\psi_2(0n0)$ is $n + 2$ which occurs exactly four times:

$$\psi_2(n) = \frac{(n + 2)}{(n + 2)} \frac{R_{n+2}}{R_{n+2}} R_{n+2}[: -2]\psi_1(R_{n+1}[: -1])(n + 2) \frac{R_{n+2}}{R_{n+2}} R_{n+2}[: -2]\psi_1(R_{n+1}[: -2])$$

The second and fourth occurrences of $n + 2$ are the last letter of an R_{n+2} , so they are preceded by a 0. The third occurrence is after $\psi_1(R_{n+1}[: -1])$ which ends with $\psi_1(0) = 202101$.

The first chunk boundary in $\psi_2(0n0)$ has the letters $2(n + 2)$. The second, third, and fourth occurrences of $n + 2$ all immediately follow a 0 or a 1, so this is the only occurrence of $2(n + 2)$ in $\psi_2(0n0)$. This means that it cannot be contained in either half of the square. Since $\max(\psi_2(0)) = 3$ and $n + 2 \geq 4$, $2(n + 2)$ cannot be the middle of the square either. So the square is contained in $\psi_2(n0)$ which contradicts Lemma 16. \square

Lemma 18. *For $n > k > 0$, neither $\psi_2(n)$ nor $\psi_2(k)$ is a suffix of the other.*

Proof. Since $n > k$, $|\psi_2(n)| > |\psi_2(k)|$, so $\psi_2(n)$ cannot be a suffix of $\psi_2(k)$.

From the definition of ψ_2 , $\psi_2(k) = (k+2)P_0(k+2)P_1(k+2)$ and $\psi_2(n)$ ends with $P_1(n+2)$. By Remark 15, $P_1(n+2)$ has suffix $(k+3)P_0(k+2)P_1(k+2)$. Therefore, $\psi_2(k)$ cannot be a suffix of $\psi_2(n)$. \square

Lemma 19. *Let w be a grounded square-free word. Then in $\psi_2(w)$, every occurrence of $\psi_2(0)[:6]$ is a prefix of a 0-chunk, and every occurrence of $\psi_2(0)[6:]$ is a suffix of a 0-chunk.*

Proof. Let $p := \psi_2(0)[:6] = 202102$ and $s := \psi_2(0)[6:] = 202101202$. Since s is a suffix of $\psi_2(0)[6:]$, proving the result for p and s is sufficient.

For $n \geq 1$, $\psi_2(n)$ ends with 1 and $\psi_2(0)$ begins with 2 and 12 does not occur in $p = 202102$, so p cannot lie over a $\psi_2(n0)$ chunk boundary. The first letter of $\psi_2(n)$ is $n+2 \geq 3$ which is larger than any letter in p , so p cannot lie over a $\psi_2(0n)$ chunk boundary. Therefore, any occurrence of p in $\psi_2(w)$ must be contained in a single chunk.

Since $\psi_1(n) = (n+1)R_{n+2}[: -2]$ does not contain 202, p does not occur in any $\psi_1(n)$. Also, $\psi_1(n)$ ends with 201, $\psi_2(n) = \psi_1((n+1)P_0(n+1))$ and $(n+1)P_0(n+1)$ is grounded, 202 only occurs in $\psi_2(n)$ at the beginning of instances of $\psi_1(0) = 202101$. This means that $p = 202102$ can never occur in $\psi_2(n)$. We can verify computationally that 202102 only occurs in $\psi_2(0)$ as a prefix. Therefore, p only occurs in $\psi_2(w)$ as a prefix of 0-chunks.

If $s = 202101202$ lies over a $\psi_2(n0)$ chunk boundary, then the 12 would need to be at the boundary. But this cannot happen since no $\psi_2(n)$ ends with 202101. It also cannot lie over a $\psi_2(0n)$ boundary since the first letter of $\psi_2(n)$ is $n+2 \geq 3$ which does not occur in s .

By Lemma 9, $\psi_1(0) = 202101$ only occurs in $\psi_2(n) = \psi_1((n+1)P_0(n+1))$ as an 0-chunk. So since no $\psi_1(n)$ begins with 202, $s = 202101202$ does not occur in $\psi_2(n)$. We can verify computationally that s only occurs in $\psi_2(0)$ as a suffix. Therefore, s only occurs in $\psi_2(w)$ as a suffix of 0-chunks. \square

We can easily see that this lemma implies that any occurrence of $\psi_2(0)$ in $\psi_2(w)$ is a 0-chunk when w is square-free and grounded.

Proposition 20. *ψ_2 is square-free over grounded words.*

Proof. Suppose w is a grounded square-free word and that $\psi_2(w)$ contains a square yy . We will first show that there is a whole 0-chunk in both halves of the square, and then use that to show that both halves have the same chunk decomposition. We then show that if the square contains any partial chunks, then the final partial chunks of the two halves of the square come from the same letter in w . We know that ψ_2 is letter injective and $\psi_2(n)$ is square-free for all n . Hence, Theorem 5 will imply that w must contain a square, which is a contradiction.

Suppose that there are no whole 0-chunks in either half. The square must contain a whole 0-chunk, or else it would be a factor of $\psi_2(0n0)$, contradicting Lemma 17. Then the whole 0-chunk in the square must be split between the two halves and the square yy is a proper factor of $\psi_2(0n0k0)$ for some $n, k \geq 1$ with the center of the square lying

in the middle 0-chunk. Consider the prefix $p := \psi_2(0)[: 6]$ and suffix $s := \psi_2(0)[6 :]$ of $\psi_2(0) = ps$. Then either p is totally contained in the first half of the square, or s is totally contained in the second half.

If s is contained in the second half, then it must also occur in the first half and by Lemma 19, the only place for this is as a suffix of the first 0-chunk in $\psi_2(0n0k0)$. The two occurrences of s in the square are followed by the first letters of $\psi_2(n)$ and $\psi_2(k)$ in each half respectively, so the first letter of $\psi_2(n)$, $n + 2$ and of $\psi_2(k)$, $k + 2$ must be equal. This means that $n = k$ which is a contradiction since w is square-free and cannot contain $0n0n0$.

If p is contained in the first half, then it must also occur in the second half and by Lemma 19, the only place for this is as a prefix of the third 0-chunk in $\psi_2(0n0k0) = [ps][\psi_2(n)][ps][\psi_2(k)][ps]$. The center of the square lies in the suffix s of the middle 0-chunk, so the boundary is formed by words s_1, s_2 such that $s = s_1s_2$. We then get from the first half of the square that y is a proper suffix of $ps\psi_2(n)ps_1$ and from the second half that y is a prefix of $s_2\psi_2(k)ps$. Since p occurs exactly once in each half, we can see that the second half is $y = s_2\psi_2(k)ps_1$. Then $s_2\psi_2(k)$ is a suffix of $ps\psi_2(n)$ implying that one of $\psi_2(n)$ or $\psi_2(k)$ must be a suffix of the other, so by Lemma 18, $n = k$ which is a contradiction since w is square-free and cannot contain $0n0n0$. Therefore, there must be a whole 0-chunk in one of the halves of the square. Then $\psi_2(0)$ occurs in both halves and by Lemma 19, both halves contain a whole 0-chunk.

We now show that both halves of the square have the same chunk decomposition. Let $[\psi_2(\ell)]$, $0 \leq \ell$ be any whole chunk in either half of the square. If $\ell = 0$, then by Lemma 19, this is a whole 0-chunk in both halves. If $\ell > 0$, then since there is a whole 0-chunk in each half and w is grounded, there must be a whole 0-chunk adjacent to this chunk. Thus, either $[ps]\psi_2(\ell)$ or $\psi_2(\ell)[ps]$ is a factor of y . If $[ps]\psi_2(\ell)$ is a factor of y , then $\psi_2(\ell)$ must be a whole ℓ -chunk in both halves since any other chunk would start with a different letter. If $\psi_2(\ell)[ps]$ is a factor, then $\psi_2(\ell)$ must be a whole ℓ -chunk in both halves since no other chunk can be a suffix of $\psi_2(\ell)$ or have $\psi_2(\ell)$ as a suffix by Lemma 18. Thus, $\psi_2(\ell)$ is a whole chunk in both halves of the square, so both halves have the same chunk decomposition.

Suppose either half contains a partial chunk. Then since the halves have the same chunk decomposition, they must both end with a partial chunk and share the first letter of their final partial chunk, say ℓ . From the definition of ψ_2 , this means that both final partial chunks are partial $(\ell - 2)$ -chunks.

This verifies the conditions of Theorem 5 which implies that w contains a square, a contradiction. \square

The next three lemmas are used to prove the irreducibility condition in Theorem 24. They are analogous to Lemmas 11, 12, and 13 about ψ_1 .

Lemma 21. *Let $n \geq 1$. Then $\psi_2(n)$ generates $\psi_2(n0)$.*

Proof. Since $n0$ is square-free and grounded, $\psi_2(n0)$ is square-free by Proposition 20, so we only need to show that $\psi_2(0)$ is irreducible in $\psi_2(n0)$. We can computationally verify that $\psi_2(1)$ generates $\psi_2(10)$ and assume that $n \geq 2$.

Since $\psi_2(n)$ has suffix $P_1(n+2)$ and $n+2 \geq 4$, Remark 15 implies that $\psi_2(n)$ has suffix $P_0(3)P_1(3)$. We can computationally verify that $P_0(3)P_1(3)$ generates $P_0(3)P_1(3)20210$. The next letter of $\psi_2(0)$ is a 2. Hence, we need to show that 2 is irreducible at the end of $\psi_2(n)202102$. Clearly, it cannot be a 0, so we will show that $\psi_2(n)202101$ contains a square. Using Equation (1) we have

$$\begin{aligned} & \psi_2(n) 202101 \\ &= \psi_2(n) \psi_1(0) \\ &= (n+2)R_{n+2}R_{n+2}[: -2]\psi_1(R_{n+1}[: -1])(n+2)R_{n+2}R_{n+2}[: -2]\psi_1(R_{n+1}[: -2]) \psi_1(0) \\ &= (n+2)R_{n+2}R_{n+2}[: -2]\psi_1(R_{n+1}[: -1])(n+2)R_{n+2}R_{n+2}[: -2]\psi_1(R_{n+1}[: -1]), \end{aligned}$$

which is a square, so 202102 is irreducible in $\psi_2(n)202102$. We can then computationally verify that $P_0(3)P_1(3)202102$ generates $P_0(3)P_1(3)\psi_2(0)$, which implies the desired result. \square

Lemma 22. For $0 \leq k \leq n$, $k+2$ is irreducible at the end of $\psi_2(nR_k[: -1])(k+2)$.

Proof. Let $0 \leq m \leq k+1$, we will show that $w_m = \psi_2(nR_k[: -1])m$ has a square suffix. If $k=0$, then $m \leq 1$. For $m=0$ we can use Remark 6 as in Lemma 12 to show that $\psi_2(nR_k[: -1])0 = \psi_2(n)0$ is a square, and for $m=1$, $\psi_2(n)1$ has the square suffix 11. Hence, $k+2=2$ is irreducible.

Note that for $k \geq 1$ the last letter of $R_k[: -1]$ is 0 and $\psi_2(0)$ ends with 01202101202, so w_m has a square suffix for $m < 3$. Assume $m \geq 3$, by definition $\psi_2(n) = (n+2)P_0(n+2)P_1(n+2)$, and according to Remark 15 $P_1(n+2)$ has suffix $P_0(m)P_1(m)$.

If $m-1 = k$, then w_m has suffix $P_0(m)P_1(m)\psi_2(R_{m-1}[: -1])m$. Also, if $m-1 < k$ it is easy to see that $nR_k[: -1]$ has suffix $(m-1)R_{m-1}[: -1]$. By definition $\psi_2(m-1) = (m+1)P_0(m+1)P_1(m+1)$, and again Remark 15 implies that $P_1(m+1)$ has suffix $P_0(m)P_1(m)$. Hence, w_m has suffix $P_0(m)P_1(m)\psi_2(R_{m-1}[: -1])m$ for all $3 \leq m \leq k+1$. Finally, we have that

$$\begin{aligned} & P_0(m)P_1(m)\psi_2(R_{m-1}[: -1])m \\ &= P_0(m)P_1(m)\psi_2(R_{m-2}[: -1])\psi_2(m-2)\psi_2(R_{m-2}[: -1])m \\ &= P_0(m)P_1(m)\psi_2(R_{m-2}[: -1])m P_0(m)P_1(m)\psi_2(R_{m-2}[: -1])m, \end{aligned}$$

which is a square. \square

Lemma 23. If $n \geq 1$ and $0 \leq k \leq n$, then $\psi_2(n)$ generates $\psi_2(nR_k)$.

Proof. Since nR_k is square-free and grounded, Proposition 20 implies that $\psi_2(nR_k)$ is square-free. It is now sufficient to show that $\psi_2(R_k)$ is irreducible in $\psi_2(nR_k)$.

We prove this inductively over n . The base case $n=1$ implies that $k=0$ or $k=1$. We can computationally verify that $\psi_2(1)$ generates $\psi_2(1R_0) = \psi_2(10)$ and that $\psi_2(1)$ generates $\psi_2(1R_1) = \psi_2(101)$.

Fix $n > 1$ and suppose the result holds for all $1 \leq n_0 < n$. That is,

$$\psi_2(R_k) \text{ is irreducible in } \psi_2(n_0R_k) \text{ for all } 0 \leq k \leq n_0 \text{ and } 1 \leq n_0 < n. \quad (i)$$

We will show that the result holds for $n_0 = n$. That is, $\psi_2(R_k)$ is irreducible in $\psi_2(nR_k)$ for all $0 \leq k \leq n$.

We can prove this intermediate step by a second induction, now over k . The base case is $k = 0$, which holds by Lemma 21 since $R_0 = 0$. Now fix k and suppose

$$\psi_2(R_{k_0}) \text{ is irreducible in } \psi_2(nR_{k_0}) \text{ for all } 0 \leq k_0 < k. \tag{ii}$$

We will show that the result holds for $k_0 = k$. That is, $\psi_2(R_k)$ is irreducible in $\psi_2(nR_k)$.

We have that $\psi_2(nR_k) = \psi_2(nR_{k-1}R_{k-2} \cdots R_2R_1R_0k)$. By the second inductive hypothesis (ii), $\psi_2(R_{k-1})$ is irreducible in $\psi_2(nR_{k-1})$. The last letter of R_{k-1} is $k-1$, so the first inductive hypothesis (i) says that $\psi_2(R_{k-2})$ is irreducible in $\psi_2((k-1)R_{k-2})$ and so $\psi_2(R_{k-1}R_{k-2})$ is irreducible in $\psi_2(nR_{k-1}R_{k-2})$. Repeating this argument shows that $\psi_2(R_{k-1}R_{k-2} \cdots R_2R_1R_0) = \psi_2(R_k[: -1])$ is irreducible in $\psi_2(nR_k[: -1])$. Lemma 22 implies that $k+2$ is irreducible in $\psi_2(nR_k[: -1])(k+2)$. And $\psi_2(k) = (k+2)P_0(k+2)P_1(k+2)$ is a prefix of $L(k+2)$ by Theorem 14, so $k+2$ generates $\psi_2(k)$, meaning that $\psi_2(k)$ is irreducible in $\psi_2(nR_k[: -1])\psi_2(k) = \psi_2(nR_k)$, which proves the result. \square

We now prove the main theorem of this section. For $n \geq 3$, define

$$T(n) = P_0(n)P_1(n)P_2(n).$$

Theorem 24. For $n \geq 3$, $L(n)$ has prefix $nT(n)$.

Proof. Note that $nT(n) = nP_0(n)P_1(n)P_2(n) = \psi_2((n-2)P_0(n-2))$. Hence, since $(n-2)P_0(n-2)$ is square-free and grounded, Proposition 20 implies that $nT(n)$ is square-free. It remains to show that $P_2(n) = \psi_2(P_0(n-2)) = \psi_2(R_{n-2}R_{n-2}[: -2])$ is irreducible in $nP_0(n)P_1(n)P_2(n)$.

We know from Theorem 14 that n generates $nP_0(n)P_1(n)$. The fact that $P_2(n) = \psi_2(R_{n-2}R_{n-2}[: -2])$ is irreducible in $nP_0(n)P_1(n)P_2(n)$ follows from Lemma 23 by the same argument used in the proof of Theorem 14. \square

Remark 25. Again, using Remark 6 we have that for $n \geq 4$

$$\begin{aligned} P_2(n) &= \psi_2(P_0(n-2)) \\ &= \psi_2(R_{n-2}[: -1](n-1)R_{n-2}[: -2]) \\ &= \psi_2(R_{n-2}[: -1]) \psi_2(n-2) \psi_2(P_0(n-3)) \\ &= \psi_2(R_{n-2}[: -1]) n P_0(n) P_1(n) P_2(n-1), \end{aligned}$$

Hence, by Remark 15 we see that $P_2(n)$ has $T(n-1) = P_0(n-1)P_1(n-1)P_2(n-1)$ as a suffix. This means that $T(n)$ has suffix $T(n-1)$, and applying the same argument repeatedly we have that $T(n)$ has $T(3)$ as a suffix.

4 The structure of $L(n)$

In this section we prove that the word $L(n)$ reflects the structure of the ruler sequence for $n = 1$ and $n \geq 3$. Namely, $L(n) = Y_n \phi(L(\varepsilon))$ for a finite prefix Y_n and a morphism ϕ .

Definition. We say a morphism ϕ is *L-commuting* over a set of words $\Delta \subset \mathbb{N}^* \cup \mathbb{N}^\omega$ if $L(\phi(w)) = \phi(L(w))$ for all $w \in \Delta$.

Let Σ be the set of all nonempty even-grounded square-free words. If ϕ is *L-commuting* over Σ , we get in particular

$$L(\phi(0)) = \phi(L(0)) = \phi(L(\varepsilon)),$$

which lets us determine the lexicographically least square-free word with prefix $\phi(0)$, as the result of applying the morphism ϕ to the ruler sequence.

In Section 4.1 we introduce a morphism α and prove that it is *L-commuting* over the set Σ of even-grounded square-free words. In Section 4.2 we use Remark 4 and the *L-commuting* property of α to find the general structure of $L(1)$ and $L(n)$ for $n \geq 3$. In Section 4.3 we state a conjecture about the structure of $L(2)$ being given by a morphism γ .

We start by showing that the ruler morphism ρ is *L-commuting* over the set of square-free words, and then use this fact to prove a result that establishes properties that are sufficient for a morphism to satisfy the *L-commuting* property over the set Σ .

Theorem 26. *The ruler morphism ρ is L-commuting over the set of all nonempty square-free words.*

Proof. Let x be any nonempty square-free word. If $|x| = \infty$, then $\rho(L(x)) = \rho(x) = L(\rho(x))$.

On the other hand, if $|x| = n < \infty$, let $w = \rho(L(x))$ and $v = L(\rho(x))$. We proceed to prove that $w = v$ by induction, proving that if w and v agree on the first $2k$ letters, then they will also agree on the next two letters. For the Base case, it is easy to see that $w[: 2n] = v[: 2n] = \rho(x)$.

For the inductive step, assume that w and v agree on the first $2k$ letters, $k \geq n$. It is clear from the definition of these words that w and v are both square-free, and if $v \neq w$, then $v \prec w$.

This implies that $v[2k] \leq w[2k] = \rho(L(x))[2k] = 0$, so w and v agree at position $2k$. Now suppose toward a contradiction that $v[2k + 1] < w[2k + 1]$ and let $l = v[2k + 1]$ and $y = L(x)[: k](l - 1)$. We have

$$\rho(y) = \rho(L(x)[: k](l - 1)) = \rho(L(x)[: k]) 0 l = L(\rho(x))[: 2k + 2] = v[: 2k + 2],$$

so y is square-free. However

$$y[k] = l - 1 = v[2k + 1] - 1 < w[2k + 1] - 1 = \rho(L(x))[2k + 1] - 1 = L(x)[k].$$

Since $y[: k] = L(x)[: k]$, this implies that y is a square-free word beginning with x that is smaller than $L(x)$. This is a contradiction and so $w[2k + 1] = v[2k + 1]$. Therefore w and v agree at position $2k + 1$, which proves the inductive step. \square

For example, Theorem 26 implies that for $n \geq 1$, $\rho(L(n-1)) = L(\rho(n-1)) = L(0n)$. So if we determine the structure of the word $L(n-1)$, this Theorem gives us the structure of the word $L(0n)$ as the ruler morphism applied to $L(n-1)$. In particular for $n = 1$ we have

$$\rho(L(0)) = \rho(\rho^\infty(0)) = \rho^\infty(0) = L(01) = L(\rho(0)).$$

The ruler morphism is not L -commuting over the set of all words on \mathbb{N} . For example,

$$L(\rho(00)) = L(0101) = 01012010 \cdots \neq 01010201 \cdots = \rho(0010 \cdots) = \rho(L(00)).$$

Theorem 27. *Let Σ be the set of all nonempty even-grounded square-free words. Let ϕ be a non-erasing morphism satisfying the following conditions.*

1. For all $w \in \Sigma$, $\phi(w)$ is square-free.
2. $\phi(0)$ generates $\phi(01)$.
3. $\phi(0n)$ generates $\phi(0n0)$ for all $n > 0$.
4. $\phi(0n)^+$ generates $\phi(0(n+1))$ for all $n > 0$.

Then ϕ is L -commuting over Σ .

Proof. Let $w \in \Sigma$. We first show that $L(w) \in \Sigma$. If $w = 0$, then $L(w)$ is the ruler sequence which is even-grounded and square-free. If $w \neq 0$, $|w| \geq 2$ and any even-length prefix of w is the image of a nonempty square-free word under ρ . So there is a nonempty square-free word w_0 such that if $|w|$ is even, $w = \rho(w_0)$, and if $|w|$ is odd, $w[: -1] = \rho(w_0)$. If $|w|$ is odd, its last letter is 0 which is irreducible, so $L(w) = L(w[: -1])$. Thus in either case, $L(w) = L(\rho(w_0)) = \rho(L(w_0))$ which is even-grounded and square-free. Here, the second equality is because ρ is L -commuting over square-free words by Theorem 26. Since $L(w) \in \Sigma$, it follows from Condition 1 that $\phi(L(w))$ is square-free.

Now we need to show that $\phi(L(w))$ is irreducible. We proceed by induction, assume that for some number $k \geq |w|$, $\phi(L(w)[: k+1])$ is a prefix of $L(\phi(w))$. We break this next bit down into cases, letting $m = L(w)[k]$.

Case 1: If $m \neq 0$ then $L(w)[k-1] = 0$ and so $L(w)[: k+1]$ ends in $0m$. In this case we have from Condition 3 that $\phi(L(w)[: k+1])$ generates $\phi(L(w)[: k+1]0)$ by Remark 4. As $L(w)[: k+2] = L(w)[: k+1]0$ this would mean that $\phi(L(w)[: k+2])$ is a prefix of $L(\phi(w))$, demonstrating the inductive step.

Case 2: In the case that $m = 0$ it must follow that $l := w[k+1] \neq 0$. We have that $\phi(L(w)[: k+1])$ ends with $\phi(0)$ which by Condition 2 we get that $\phi(1)$ at the end of $\phi(L(w)[: k+1]1)$ is irreducible. If $l = 1$ then we're done. If not then we enter the following argument.

Let n be such that $\phi(n)$ at the end of $\phi(L(w)[: k+1]n)$ is irreducible and $0 < n < l$. $L(w)[: k+1]n$ has w as prefix and is lexicographically less than $L(w)$ so $L(w)[: k+1]n$ contains a square and therefore $\phi(L(w)[: k+1]n)$ contains a square. We have that $\phi(0n)$ at the end of $\phi(L(w)[: k+1]n)$ is irreducible meaning that all words lexicographically less than

$\phi(0n)$ would introduce a square. But $\phi(0n)$ also introduces a square. This means that all words less than $\phi(0n)^+$ introduce a square, so $\phi(0n)^+$ is irreducible in $\phi(L(w)[: k + 1]n)^+$. Then from Condition 4 we get that $\phi(0(n + 1))$ at the end of $\phi(L(w)[: k + 1](n + 1))$ is irreducible. Then by induction on n , $\phi(L(w)[: k + 1]l)$ is irreducible.

This argument allows us to show that $L(\phi(w))$ and $\phi(L(w))$ agree on their $(k + 1)$ th chunk and so provides the inductive step. The base case is simply when $k = |w|$ which is trivial since $\phi(w)$ is a prefix of both words. \square

4.1 The morphism α

The morphism α is defined as follows.

Definition. For all $n \geq 0$, let

$$\alpha(n) = \begin{cases} EFE & \text{if } n = 0 \\ B_1 R_4 C B_1 R_4 & \text{if } n = 1 \\ \alpha(n - 1)^+ R_{n+3} C \alpha(n - 1)^+ R_{n+3} & \text{if } n \geq 2, \end{cases}$$

where

$$\begin{aligned} C &= 0102030102, \\ B_0 &= 0301 \psi_1(1010)[: -3] \psi_2(1010)[: -6] \psi_2(10)[: -12] 301020, \\ B_1 &= \rho(B_0[7 : -5]), \\ E &= 0102B_01B_0[: -9], \\ F &= B_0[-9 :]3010302C0103C^+02, \text{ and} \\ G &= 010203012. \end{aligned}$$

These definitions can be referenced in Section 7. The lengths of the auxiliary words are $|C| = 10$, $|B_0| = 798$, $|B_1| = 1572$, $|E| = 1592$, $|F| = 42$ and $|G| = 9$. We can computationally verify that E is the largest word that is both a prefix and a suffix of $\alpha(0)$. It is of interest to note that for $n > 0$, $\alpha(n)$ has prefix B_1 and that B_1 has prefix F^{++} , so F cannot be a prefix of any $\alpha(n)$. Also, the word G is useful since it is the shortest word that generates $\alpha(0)$.

Over the next two subsections, we prove that α satisfies the conditions of Theorem 27, which will imply that it is L -commuting over nonempty even-grounded square-free words.

4.1.1 Condition 1: α is square-free over grounded words

Condition 1 of Theorem 27 says that α is square-free over even-grounded words. In this section, we prove the following stronger property.

Theorem 28. α is square-free over grounded words.

Theorem 28 will be shown to follow from Theorem 5. This requires the results about α shown in the following lemmas.

Lemma 29. *Let $n > k > 0$. Then $\alpha(n)$ and $\alpha(k)$ end with different letters and neither is a prefix of the other.*

Proof. By definition, $\alpha(n)$ ends with R_{n+3} which has $n + 3$ as its last letter, so $\alpha(n)$ and $\alpha(k)$ end with different letters. Consider that since $n > k$, $|\alpha(n)| > |\alpha(k)|$ so $\alpha(n)$ cannot be a prefix of $\alpha(k)$. Also, $\alpha(n)$ begins with $\alpha(k)^+$ so $\alpha(k)$ cannot be a prefix of $\alpha(n)$. \square

The next three results show how E can be used to restrict the placement of chunks throughout a word $\alpha(w)$ where w is square-free and grounded.

Lemma 30. *If w is a grounded square-free word, then every occurrence of E in $\alpha(w)$ is a prefix or suffix of a 0-chunk.*

Proof. We can verify computationally that E only occurs in $\alpha(0)$ as a prefix and a suffix. For $n > 0$, $\alpha(n)$ is even-grounded, but E is not grounded so it cannot be a factor of $\alpha(n)$. So any other occurrence of E in $\alpha(w)$ must lie over the chunk boundary in $\alpha(0n)$ or $\alpha(n0)$ for some $n > 0$.

If E lies over the chunk boundary in $\alpha(0n)$, then there must be a nonempty suffix of E that is also a prefix of $\alpha(n)$. But every prefix of $\alpha(n)$ is even-grounded and E has no nonempty even-grounded suffix since it ends with 1 2.

If E lies over the chunk boundary in $\alpha(n0)$, then there must be a nonempty prefix of E that is also a suffix of $\alpha(n)$. But $\max(E) = 3$ and the last letter of $\alpha(n)$ is $n + 3$ which is greater than 3. \square

Corollary 31. *If w is a grounded square-free word, then every occurrence of $\alpha(0)$ in $\alpha(w)$ is a 0-chunk.*

Proof. Any occurrence of $\alpha(0) = EFE$ in $\alpha(w)$ begins and ends with E . There are only two occurrences of E in $\alpha(0)$ so by Lemma 30, one E must be a prefix of a 0-chunk and the other must be a suffix of a 0-chunk. F is shorter than every chunk, so no other chunk can be contained in it. Therefore, this must be a whole 0-chunk. \square

Corollary 32. *Let w be a grounded square-free word and $l > 0$. If $E\alpha(l)$ or $\alpha(l)E$ is a factor of $\alpha(w)$, then that occurrence of $\alpha(l)$ is an l -chunk.*

Proof. If $E\alpha(l)$ occurs in $\alpha(w)$, E is followed by the prefix F^{++} of $\alpha(l)$, so E cannot be followed by F and this cannot be the prefix of a 0-chunk. Thus, E must be a suffix of a 0-chunk by Lemma 30. So a nonzero chunk begins at the start of $\alpha(l)$. By Lemma 29, this must be an l -chunk.

An analogous argument uses the fact that F cannot be the suffix of any $\alpha(l)$ to show the result for $\alpha(l)E$. \square

The following is the final result that we need for proving Theorem 28, that α is square-free over grounded words.

Proposition 33. $\alpha(0n0)$ is square-free for all $n > 0$.

The proof requires Lemmas 34 to 40. Lemmas 34 to 36 show some results about the structure of $\alpha(n)$. Lemmas 37 and 38 show that $\alpha(n)$ is square-free for all n . Finally, Lemmas 39 and 40 show that $\alpha(0n)$ and $\alpha(n0)$ are square-free, which is then used to prove Proposition 33.

Lemma 34. For all $n \geq 1$, $0203R_3^+R_4^+\cdots R_{n+2}^+R_{n+3}$ is a suffix of $\alpha(n)$.

Proof. We proceed by induction. For $n = 1$ we can check directly that $0203R_3^+R_4$ is a suffix of $\alpha(1)$. For the inductive step, assume that for some $k \geq 1$, we have that $0203R_3^+\cdots R_{k+2}^+R_{k+3}$ is a suffix of $\alpha(k)$ and recall that

$$\alpha(k+1) = \alpha(k)^+ R_{k+4} C \alpha(k)^+ R_{k+4}.$$

Since $\alpha(k)$ ends with $0203R_3^+\cdots R_{k+2}^+R_{k+3}$, we have that $\alpha(k+1)$ must end with $0203R_3^+\cdots R_{k+2}^+R_{k+3}^+R_{k+4}$, which concludes the proof. \square

Lemma 35. For all $n \geq 1$, $R_{n+4}[i :]$ is a suffix of $\alpha(n)^+$ if and only if $i \geq 6$.

Proof. We can write R_{n+4} as

$$\begin{aligned} R_{n+4} &= R_3R_3^+R_4^+\cdots R_{n+2}^+R_{n+3}^+ \\ &= 01020103R_3^+R_4^+\cdots R_{n+2}^+R_{n+3}^+. \end{aligned}$$

Also, it follows from Lemma 34 that $\alpha(n)^+$ ends with $0203R_3^+\cdots R_{n+2}^+R_{n+3}^+$. These two words are identical starting with the $03R_3^+$, but not including any letters before. Therefore, $R_{n+4}[i :]$ is a suffix of $\alpha(n)^+$ if and only if $i \geq 6$. \square

Lemma 36. For all $n \geq 1$, $\alpha(n)$ ends in $n+3$, and does not contain any letter greater than $n+3$. For $n \geq 2$, $\alpha(n)$ contains exactly four occurrences of $n+3$.

Proof. This can be proved using induction. For the base case, it can be checked by direct computation that $\alpha(1)$ and $\alpha(2)$ satisfy the lemma.

For the inductive step, assume that for some $k \geq 2$, $\alpha(k)$ satisfies the lemma. Since C does not contain any occurrence of $k+4$ or higher letters and

$$\alpha(k+1) = \alpha(k)^+ R_{k+4} C \alpha(k)^+ R_{k+4},$$

it is clear that our assumption implies that $\alpha(k+1)$ contains exactly four occurrences of $k+4$, no higher letters, and ends with $k+4$. \square

Lemma 37. For all $n \geq 1$, $C\alpha(n)^+$ is square-free.

Proof. We proceed by induction. We can check computationally that the claim holds for $n = 1$.

For the inductive step assume that for some $k \geq 1$, $C\alpha(k)^+$ is square-free and suppose that the word $w = C\alpha(k+1)^+$ contains a square. We can think of w as the concatenation of 6 factors:

$$w = w_1w_2w_3w_4w_5w_6 := [C][\alpha(k)^+][R_{k+4}][C][\alpha(k)^+][R_{k+4}^+].$$

By the inductive hypothesis we have that $w_1w_2 = w_4w_5$ is square-free. We divide the rest of the proof in four steps.

Step 1: Proving that $w_1w_2w_3$ is square-free. Suppose that $w_1w_2w_3$ contains a square. We know that the square must cross the boundary between w_2 and w_3 , and hence it includes the last letter of $w_2 = \alpha(k)^+$, which by Lemma 36 is $k + 3 + 1 = k + 4$. The only other occurrence of $k + 4$ in $w_1w_2w_3$ is the last letter of $w_3 = R_{k+4}$, so the square ends in $k + 4$. Hence, the second half of the square is the entire R_{k+4} , and so R_{k+4} is a suffix of $\alpha(k)^+$, which contradicts Lemma 35. This completes the proof of our first claim.

Step 2: Proving that $w_1w_2w_3w_4$ is square-free. Suppose that $w_1w_2w_3w_4$ contains a square. By the previous step, the square must cross the boundary between w_3 and w_4 , and hence it includes the letter $w_3[-1] = k + 4$. Since $k + 4$ does not occur in $w_4 = C$, and the only other occurrence of $k + 4$ in this word is as the last letter of $w_2 = \alpha(k)^+$, each half of the square must have length equal to $|w_3| = 2^{k+4}$. The largest common prefix between the ruler sequence and C is $C[: 5]$. So at most these five letters can appear in each half of the square after the occurrences of $k + 4$. On the other hand, Lemma 35 implies that at most a suffix of $w_3 = R_{k+4}$ with length $2^{k+4} - 6$ can appear as a suffix of $w_2 = \alpha(k)^+$, and so at most this many letters can appear in each half of the square up to the occurrences of $k + 4$.

Therefore, each half of the square contains at most $2^{k+4} - 6 + 5 = 2^{k+4} - 1$ letters, which just falls short of the number required. Therefore no square can exist in $w_1w_2w_3w_4$.

Step 3: Proving that $w_1w_2w_3w_4w_5$ is square-free. Suppose that it contains a square. By the previous step the square can not be contained in $w_1w_2w_3w_4$. Also, by the inductive hypothesis the square can not be contained in w_4w_5 . So the square must contain a nonempty suffix of $w_3 = R_{k+4}$, all of $w_4 = C$, and a nonempty prefix of $w_5 = \alpha(k)^+$. In particular it includes $k + 4$, the last letter of w_3 . Also, note that this word includes only three occurrences of $k + 4$ ($w_2[-1]$, $w_3[-1]$ and $w_5[-1]$), hence the square must contain only two of them.

If the square contains $w_2[-1]$ and $w_3[-1]$, then the second half of the square contains the whole factor $w_4 = C$ right after $w_3[-1]$. This implies that C must also appear right after $w_2[-1]$, and so C must be a prefix of $w_3 = R_{k+4}$, which is a contradiction. On the other hand, if the square contains $w_3[-1]$ and $w_5[-1]$, then the second half of the square must be $w_4w_5 = C\alpha(n)^+$ and the first half must be contained in $w_3 = R_{k+4}$. This is impossible since C is not contained in R_{k+4} .

Step 4: Proving that $w_1w_2w_3w_4w_5w_6$ is square-free. Suppose that it contains a square.

The square must contain $w_5[-1] = k + 4$ and since $w_6 = R_{k+4}^+$ does not contain $k + 4$, the occurrence of $k + 4$ in the first half of the square must be at $w_3[-1]$. Note that the first half of the square cannot contain the whole $w_4 = C$, because this is not a prefix of $w_6 = R_{n+4}$. This implies that $w_5 = \alpha(n)^+$ is totally contained in the second half of the square. Hence, since $|R_{k+4}| < |\alpha(n)^+|$ we have that $w_3 = R_{n+4}$ is a suffix of $w_5 = \alpha(n)^+$, which contradicts Lemma 35.

Therefore we conclude that $w_1w_2w_3w_4w_5w_6$ is square-free, which completes the proof of the lemma. \square

Lemma 38. $\alpha(n)$ is square-free for all $n \geq 0$.

Proof. We proceed by induction. We can check computationally that $\alpha(0)$ and $\alpha(1)$ are square-free.

For the inductive step assume that $\alpha(k)$ is square-free for some $k \geq 1$. We can think of $w = \alpha(k + 1)$ as the concatenation of 5 factors:

$$w = w_1w_2w_3w_4w_5 := [\alpha(k)^+][R_{k+4}][C][\alpha(k)^+][R_{k+4}].$$

Suppose w contains a square. By Lemma 37, $w^+ = \alpha(k + 1)^+$ is square-free. So $w[: -1]$ is square-free and the square in w is a suffix, containing $w_5[-1] = k + 4$. Also, the square must contain $w_4[-1] = k + 4$, since $w_5 = R_{k+4}$ is square-free. Now, since w contains exactly four occurrences of $k + 4$ we have the following cases:

Case 1: the square contains only $w_4[-1]$ and $w_5[-1]$. In this case the second half of the square would have to be the whole factor $w_5 = R_{k+4}$. Hence the first half of the square would have R_{k+4} as a suffix of $w_4 = \alpha(n)^+$ which contradicts Lemma 36.

Case 2: the square contains all four occurrences of $k + 4$ which are $w_1[-1]$, $w_2[-1]$, $w_4[-1]$, and $w_5[-1]$. In this case $k + 4$ must be the final letter in each half of the square, and so the first half of the square would be a suffix of w_1w_2 . This is not possible since the second half of the square would be $w_3w_4w_5$ which is longer than w_1w_2 .

Therefore we conclude that $w = \alpha(k + 1)$ is square-free, which completes the inductive step. \square

Lemma 39. $\alpha(0n)$ is square-free for all $n \geq 1$.

Proof. From Lemma 38 we have that any square in $\alpha(0n)$ must cross into both chunks. We consider two cases based on the length of the square.

Suppose there is a square yy in $\alpha(0n)$ such that $|y| \leq |\alpha(0)|$. Then the square's total length would be at most $|\alpha(0)| \times 2 = 6452$ letters. Computationally, we can see that $\alpha(3)$ and $\alpha(4)$ share their first 13029 letters. Since $\alpha(n)$ is a prefix of $\alpha(n + 1)$, all $\alpha(n)$ for $n \geq 3$ have the same first 13029 letters. Thus, $\alpha(0n)$ has the same first $|\alpha(0)| + 13029 = 16255$ letters for all $n \geq 3$. Checking with a computer, we find that $\alpha(01)$, $\alpha(02)$, and $\alpha(03)$ are square-free, so the first 16255 letters of $\alpha(0n)$ are square-free for all n . Since the square must intersect the 0-chunk and has length at most $|\alpha(0)| \times 2$, the square must be contained in the first $|\alpha(0)| \times 3 = 9678$ letters. This is a contradiction so no square yy with length $|y| \leq |\alpha(0)|$ can occur in $\alpha(0n)$.

Now suppose $|y| > |\alpha(0)|$. Then the second half of the square is entirely contained in $\alpha(n)$. However, $\alpha(n)$ is grounded and $\alpha(0)$ ends with 12. This means the first half of the square can't contain more than the last letter of $\alpha(0)$ so it remains to show that $2\alpha(n)$ is square-free.

Since

$$C[-1]\alpha(n)^+ = 2\alpha(n)^+$$

is a factor of $\alpha(n+1)$, it must be square-free. So we can show that decreasing the last letter by one does not introduce a square. We know that $2\alpha(n)[1:]$ and $2\alpha(n)[: -1]$ are square-free, so a square would need to be the entire word. However,

$$|2\alpha(n)| = 1 + 2|\alpha(n-1)| + 2|R_{n+3}| + 10,$$

which is odd so it is impossible for the entire word to be a square. □

Lemma 40. $\alpha(n0)$ is square-free for all $n \geq 1$.

Proof. We can check by computer that $\alpha(10)$ is square-free. Assume $n \geq 2$ from now on.

Since $\alpha(0)$ and $\alpha(n)$ are square-free, any square in $\alpha(n0)$ must cross into both chunks and include the $n+3$ at the end of $\alpha(n)$. Since $\max(\alpha(0)) = 3$, the $n+3$ in each half must come from $\alpha(n)$. As a result, we know the entire first half of the square is in $\alpha(n)$. Additionally, $\alpha(0)$ becomes ungrounded at the ninth letter and $\alpha(n)$ is grounded, so the square can't extend past the eighth letter of $\alpha(0)$. Thus this proof simplifies to proving

$$\alpha(n)01020301$$

is square-free. From Lemma 36, $\alpha(n)$ contains 4 occurrences of $n+3$, so the full square must either contain the last two or all four.

Case 1: The square contains only the third and fourth occurrences of $n+3$, so the square is contained in

$$C\alpha(n-1)^+R_{n+3}01020301 = C\alpha(n-1)^+R_{n+3}C[:8],$$

which is a prefix of $C\alpha(n)^+$, which is square-free by Lemma 37.

Case 2: The square contains all occurrences of $n+3$, so it appears in

$$\alpha(n-1)^+R_{n+3}C\alpha(n-1)^+R_{n+3}C[:8].$$

The second and fourth occurrences of $n+3$ are separated by a distance of

$$|C| + |\alpha(n-1)^+| + |R_{n+3}| = 10 + |\alpha(n-1)^+| + |R_{n+3}|,$$

so the entire square would need to have twice this length. However, the whole word has length

$$|\alpha(n-1)^+| + |R_{n+3}| + |C| + |\alpha(n-1)^+| + |R_{n+3}| + |C[:8]| = 18 + 2|\alpha(n-1)^+| + 2|R_{n+3}|,$$

which is less than the required length of square.

Since both cases are ruled out, no square can exist in $\alpha(n0)$. □

We can now prove that $\alpha(0n0)$ is square-free:

Proof of Proposition 33. We can prove that $\alpha(010)$ and $\alpha(020)$ are square-free via computation. Assume $n \geq 3$ from here.

Using Lemmas 39 and 40, it follows that if a square exists in $\alpha(0n0)$, then it includes the entire n -chunk as well as a part of each 0-chunk. We can check using a computer that $|\alpha(0)| = 3226$ and $|\alpha(3)| = 13030$. For any $p > q > 1$, $|\alpha(p)| > |\alpha(q)|$. Therefore, $|\alpha(n)| \geq |\alpha(3)| > |\alpha(0)|$.

Case 1: Suppose the boundary between the halves of the square appears either in a 0-chunk or between chunks. Then the half of the square entirely in a 0-chunk would have a length at most 3226. The other half would also have the same length, but we know this half needs to contain the entire n -chunk, which has length greater than 3226. Thus this case is impossible.

Case 2: Now suppose the boundary lies within the n -chunk. Let x be the nonempty suffix of the first 0-chunk contained in the square, and z be the nonempty prefix of the last 0-chunk contained in the square. The occurrence of x in the second half of the square begins in $\alpha(n)$. If it extends into the last 0-chunk, then the whole square has length less than

$$|xzxz| \leq 4 \times |\alpha(0)| = 12904 < 13030 = |\alpha(3)| \leq |\alpha(n)|.$$

This is not possible since the square contains all of $\alpha(n)$. Thus, there are nonempty words y such that the square can be written as

$$x]yzxy][z,$$

where $yzxy = \alpha(n)$. Since all of $\alpha(n)$ is grounded, x and z are both grounded. We can check by computer that the longest grounded prefix of $\alpha(0)$ is 01020301, and its longest grounded suffix is 2. It follows that $x = 2$ and z is a prefix of 01020301. This means that $|zx| \leq 9$ and zx must appear in the exact center of $\alpha(n)$. The middle 10 letters of $\alpha(n)$ are $C = 0102030102$, so zx is located at the center of C , meaning that it must be grounded. Thus, the only possible values for zx are 02, 0102, 010202, and 01020302. Clearly none of these appear at the center of C , so this square cannot exist. \square

We can now use the above results to prove that α is square-free over grounded words.

Proof of Theorem 28. Suppose w is a square-free grounded word and $\alpha(w)$ contains a square yy . We will first show that E must be a factor of y , and then use that to show that both halves of the square have the same chunk decomposition. We then show that each half of the square contains a whole chunk, and that if the square contains any partial chunks, then the initial partial chunks of the two halves of the square come from the same letter. It is clear from its definition that α is letter-injective, and we know from Lemma 38 that $\alpha(n)$ is square-free for all letters n . Hence, Theorem 5 will imply that w contains a square, which is a contradiction.

The square yy contains a whole 0-chunk, since otherwise it would be a factor of $\alpha(0n0)$, contradicting Proposition 33. Since $\alpha(0) = EFE$, there are at least two whole occurrences

of E in yy . At least one of these occurrences must be completely contained in one half of the square. Thus, E is a factor of y .

Let $[\alpha(l)]$, $0 \leq l$ be any whole chunk in either half of the square. We will show that the corresponding occurrence of $\alpha(l)$ in the other half is also a whole l -chunk. If $l = 0$, then by Corollary 31, $\alpha(l)$ must be a whole 0-chunk in both halves. If $l > 0$, then since w is grounded and E is a factor of y , there must be a whole occurrence of E adjacent to this chunk and entirely contained in this half of the square. Thus, either $E\alpha(l)$ or $\alpha(l)E$ is a factor of y . Then by Corollary 32, $\alpha(l)$ is a whole l -chunk in both halves of the square, so both halves have the same chunk decomposition.

Suppose neither half of the square contains a whole chunk. Then yy cannot span over more than three chunks. Since $\alpha(0n0)$ is square-free by Proposition 33, yy must be a proper factor of $\alpha(n0k)$. Then yy overlaps all three chunks because $\alpha(n0)$ and $\alpha(0k)$ are square-free. By Lemma 30, there are exactly two occurrences of E in $\alpha(n0k)$ and each must be in a different half of the square since E is a factor of y . But the letter before the E in the first half is the last letter of $\alpha(n)$, which is $n + 3$, and the letter before the E in the second half is the last letter of F , which is 2. We cannot have E as a prefix of y , or else the square would not overlap the first of the three chunks. This is a contradiction so one half of the square must contain a whole chunk. Since both halves have the same chunk decomposition, both halves contain a whole chunk.

Suppose either half of the square contains a partial chunk. Then since the halves have the same chunk decomposition, they must both begin with a partial chunk. We will show that the halves share their initial partial chunk. The initial partial chunks end with the same letter, so by Lemma 29, they must be equal chunks or one of them must be a 0-chunk. But since the halves of the square have the same chunk decomposition and contain a whole chunk, their first whole chunks are equal. So the final partial chunks are either both 0-chunks or equal nonzero chunks.

This verifies the conditions of Theorem 5 which implies that w contains a square, a contradiction. \square

4.1.2 Conditions 2, 3, and 4

In this section we prove that α satisfies the remaining conditions of Theorem 27. Condition 2 ($\alpha(0)$ generates $\alpha(01)$) can be verified via direct computation. In the following result we prove that α satisfies Condition 3.

Theorem 41. *For all $n > 0$, $\alpha(0n)$ generates $\alpha(0n0)$.*

Proof. The case $n = 1$ can be verified via direct computation, so we assume $n \geq 2$. By Theorem 33 $\alpha(0n0)$ is square-free, hence since the word G generates $\alpha(0)$, it is enough to show that G is irreducible in $\alpha(0n)G$. Indeed, consider

$$\alpha(0n)G = \alpha(0n)01020\underline{3}01\underline{2}.$$

It is clear that the only letters in G that could potentially be reduced are the underlined

ones. The 3 could only be reduced to 1, in which case note that

$$\begin{aligned}
 \alpha(0n)010201 &= \cdots \alpha(n)010201 \\
 &= \cdots \alpha(n-1)^+ R_{n+3} C \alpha(n-1)^+ R_{n+3} 010201 \\
 &= \cdots \alpha(n-1)^+ R_{n+3} 010201 \\
 &= \cdots R_{n+3}[6:]R_{n+3}010201 && \text{(by Lemma 35)} \\
 &= \cdots R_{n+3}[6:]010201R_{n+3}[6:]010201,
 \end{aligned}$$

which contains a square, so the 3 in G is irreducible.

Now, the last 2 in G could only be reduced to 0, and in this case note that

$$\begin{aligned}
 \alpha(0n)010203010 &= \cdots 2\alpha(n)010203010 \\
 &= \cdots 2\alpha(n-1)^+ R_{n+3} C \alpha(n-1)^+ R_{n+3} 010203010 \\
 &= \cdots 2\alpha(n-1)^+ R_{n+3} 0102030102\alpha(n-1)^+ R_{n+3} 010203010,
 \end{aligned}$$

which contains a square, so the last letter of G is irreducible. \square

Before proving that α satisfies Condition 4 of Theorem 27 we need to establish the following lemmas.

Lemma 42. C is irreducible in $\alpha(n)^+ R_{n+4} C$ for all $n > 0$.

Proof. Recall that $C = 0102030102$. Clearly, the only letter that is reducible within C is the 3, which could only be made a 1. In this case we would have $\alpha(n)^+ R_{n+4} 010201$. Since $n \geq 1$, Lemma 35 says that $R_{n+4}[6:]$ is a suffix of $\alpha(n)^+$. Also, $R_{n+4}[:6] = 010201$ for all n . Therefore

$$\alpha(n)^+ R_{n+4} 010201 = \cdots R_{n+4}[6:]010201R_{n+4}[6:]010201$$

which contains a square. Hence C is irreducible. \square

Lemma 43. C generates $C\alpha(n)^+$ for all $n > 0$.

Proof. We proceed by induction. We can check in the case $n = 1$, that $C\alpha(1)^+$ is a prefix of $L(C)$ by direct computation.

For the inductive step, assume that C generates $C\alpha(k)^+$ for some $k \geq 1$. First note that

$$C\alpha(k+1)^+ = C\alpha(k)^+ R_{k+4} C\alpha(k)^+ R_{k+4}^+$$

is square-free, since it is a factor of $\alpha(k+2)$. Then we just need to show that $\alpha(k+1)^+$ is irreducible.

From the inductive hypothesis $\alpha(k)^+$ is irreducible after C and since R_{k+4} is a prefix of the ruler sequence, it is also irreducible. Now, Lemma 42 implies that C is irreducible after $C\alpha(k)^+ R_{k+4}$ and from the inductive hypothesis again we conclude that $\alpha(k)^+$ is irreducible after $C\alpha(k)^+ R_{k+4} C$. Finally, the last letter in R_{k+4}^+ cannot be reduced by 1 because it would create the square $(C\alpha(k)^+ R_{k+4})^2$, and cannot be reduced by more than 1 because R_{k+4} is irreducible. Therefore $\alpha(k)^+ R_{k+4} C\alpha(k)^+ R_{k+4}^+ = \alpha(k+1)^+$ is irreducible in $C\alpha(k+1)^+$, which concludes the proof. \square

We note that Lemma 43 immediately describes the structure of $L(012)$.

Corollary 44. $L(012) = 01201 \lim_{n \rightarrow \infty} \rho^{-1}(\alpha(n))$.

Proof. First note that ρ^{-1} is well defined on $\alpha(n)$ since it is even-grounded. Since 010203 generates C , Lemma 43 implies that $L(\rho(012)) = L(010203) = L(C) = C \lim_{n \rightarrow \infty} \alpha(n)$. Since ρ is L -commuting over square-free words by Theorem 26, $L(\rho(012)) = \rho(L(012))$. Thus, $L(012) = \rho^{-1}(C)\rho^{-1}(\lim_{n \rightarrow \infty} \alpha(n)) = 01201 \lim_{n \rightarrow \infty} \rho^{-1}(\alpha(n))$. \square

Finally, we prove that α satisfies Condition 4.

Theorem 45. $\alpha(0n)^+$ generates $\alpha(0(n+1))$ for all $n > 0$.

Proof. From Lemma 39 we know that $\alpha(0(n+1))$ is square-free, so it is enough to show that $\alpha(n)^+$ generates $\alpha(n+1)$. To show this, recall that

$$\alpha(n+1) = \alpha(n)^+ R_{n+4} C \alpha(n)^+ R_{n+4}.$$

Consider that $\alpha(n)^+$ generates $\alpha(n)^+ R_{n+4} C$, because R_{n+4} is a prefix of the ruler sequence and C is irreducible by Lemma 42. Similarly, Lemma 43 implies that $\alpha(n)^+ R_{n+4}$ is irreducible after C . Therefore $\alpha(n)^+$ generates $\alpha(n+1)$. \square

4.1.3 Conclusion

We have proved that the morphism α satisfies all the requirements of Theorem 27, hence we have the following result.

Theorem 46. α is L -commuting over Σ , the set of all nonempty even-grounded square-free words.

Corollary 47. $L(G) = L(\alpha(0)) = \alpha(L(\varepsilon))$.

Proof. Recall that G generates $\alpha(0)$, so $L(G) = L(\alpha(0))$. The other equality follows directly from Theorem 46, since $0 \in \Sigma$. \square

4.2 Structure of $L(1)$ and $L(n)$ for $n \geq 3$

The following result will reduce the task of proving the square-freeness of a word formed by a finite prefix followed by $\alpha(L(\varepsilon))$ to a finite computation.

Lemma 48. Let w be a finite square-free word. If $w\alpha(L(\varepsilon))$ contains a square, then that square contains no letter greater than $\max(w\alpha(0))$.

Proof. Suppose toward a contradiction that there is a square yy with a letter greater than $\max(w\alpha(0))$. Since w and $\alpha(L(\varepsilon))$ are square-free, yy must cross the boundary between these two factors. Choose n to be some letter such that $\max(\alpha(n))$ is greater than any letter in y . Then yy must be contained in $w\alpha(R_n[: -1]n) = w\alpha(R_n)$.

Let $\ell := \max(y) > \max(w\alpha(0))$ and choose some occurrence of ℓ in the first half of the square. Since ℓ is neither contained in w nor $\alpha(0)$, and since R_n is even-grounded,

this occurrence of ℓ is from an i -chunk which is after a 0-chunk. Also, this 0-chunk must be totally contained within the first half of the square, since the square involves w .

Let s be the suffix of the first half of the square starting right after w . By our previous reasoning, s contains the whole first 0-chunk. We claim that the occurrences of s at the end of each half of the square yy have the same chunk decomposition. Indeed, let s_1 and s_2 be the occurrences of s in the first and second half of the square respectively. Consider the first whole occurrence of $\alpha(0)$ in s_1 and s_2 , Lemma 31 implies that these occurrences of $\alpha(0)$ are indeed 0-chunks.

Now recall that by Lemma 29, for $n > k > 0$, we have that $\alpha(n)$ and $\alpha(k)$ end with different letters and neither is a prefix of the other. Hence, it is not hard to see that if for some $n > 0$, there is a whole n -chunk in s_1 either right before or immediately after this 0-chunk, then s_2 must also have this n -chunk in the same position. Inductively we have that s_1 and s_2 have the same chunk decomposition.

We claim that ℓ occurs in a full chunk in both halves of the square. Indeed, ℓ cannot occur in an initial partial chunk of the first half of the square, since it occurs after $w\alpha(0)$. Suppose now that ℓ occurs in an i -chunk, $i > 0$, which splits over the two halves. By Lemma 36, the last letter of the chunk is its largest letter, which must be at least ℓ , and so cannot be in w or in a 0-chunk. So this i -chunk ends after the suffix of w and the $\alpha(0)$ at the beginning of the second half. This is a contradiction since an i -chunk cannot contain a 0-chunk by Corollary 31. Hence ℓ must occur in a whole i -chunk, and then the claim follows from the fact that s_1 and s_2 have the same chunk decomposition.

From our knowledge of the ruler sequence, for any two occurrences of i within R_n there exists an $i + 1$ between them, and so $\alpha(i + 1)[-1]$ is contained within yy . Finally, from Lemma 36 we have that $\alpha(i + 1)[-1] > \alpha(i)[-1] \geq \ell$, which contradicts our choice of ℓ . \square

Remark 49. We can see from the properties of the ruler sequence that Lemmas 31, 36, and 29 apply analogously to the morphism $\rho \circ \alpha$. Then the proof of Lemma 48 can be easily adapted to show that if w is square-free, then any square in $w\rho(\alpha(L(\varepsilon)))$ contains no letter greater than $\max(w\rho(\alpha(0)))$. This will be used in Lemma 51 to prove that $A\rho(\alpha(L(\varepsilon)))$ is square-free.

Now we can prove Theorem 1.

Theorem 1. *Let Y_1 be the 5177-letter prefix of $L(1)$. Then $L(1) = Y_1 \alpha(L(\varepsilon))$.*

Proof. We first show that $Y_1\alpha(L(\varepsilon))$ is square-free. Indeed, suppose that $Y_1\alpha(L(\varepsilon))$ contains a square. Since Y_1 and $\alpha(L(\varepsilon))$ are square-free, the square must start in Y_1 . We can verify by computation that $Y_1\alpha(R_2)$ is square-free, so the square must end after $\alpha(R_2)$. Hence it contains $\alpha(2)[-1] = 5$. But by Lemma 48, the square cannot contain any letter larger than $\max(Y_1\alpha(0)) = 4$. This is a contradiction so $Y_1\alpha(L(\varepsilon))$ is square-free.

We can check by direct computation that $L(1) = L(Y_1G)$. Then using Remark 4 with $p = G$, $w = L(G)$, and $u = Y_1$, and Corollary 47, we obtain that

$$L(1) = L(Y_1G) = Y_1 L(G) = Y_1\alpha(L(\varepsilon)). \quad \square$$

The structure of $L(n)$ for $n \geq 3$ is similar to that of $L(1)$, although the prefix is different and the morphism α is replaced with the composition $\rho \circ \alpha$. From Theorem 24, we know that $L(n)$ has prefix $nT(n)$ which has length exponential in n . This is followed by A , a constant word of length 13747 which can be easily found computationally. It is noteworthy that A has prefix $\psi_2(0)^+$.

Theorem 2. For all $n \geq 3$, $L(n) = Y_n \rho(\alpha(L(\varepsilon)))$, where $Y_n = nT(n)A$.

In order to prove that n generates $nT(n)A\rho(\alpha(\varepsilon))$, we need first to show that it is square-free and then show that $A\rho(\alpha(\varepsilon))$ is irreducible. We begin with some lemmas used to prove the square-free condition.

Lemma 50. For $n \geq 3$, $nT(n)A$ is square-free.

Proof. From Theorem 24 we know that $nT(n)$ is square-free and we can verify that A is square-free computationally, so any square would have to overlap both factors. We can also computationally check the cases $n = 3, 4, 5$, so assume $n \geq 6$ and suppose that $nT(n)A$ contains a square yy .

From Remark 25 we have that $T(n)$ has suffix $T(6)$. We can computationally verify that $T(6)A$ is square-free, so the square contains $T(6)$. Since $\max(T(6)A) = 6$, then let k be the largest letter in the square, we have that $k \geq 6$. Also, $\max(A) = 5$, so all occurrences of k are in $nT(n)$. Since both halves contain at least one letter k , then the center of the square lies in $nT(n)$.

Recall that A begins with $\psi_2(0)^+$ which never occurs in $nT(n) = \psi_2((n-2)P_0(n-2))$ by Lemma 19. Since the first half is contained in $nT(n)$, y cannot contain $\psi_2(0)^+$. The second half of the square starts in $nT(n)$, but it cannot contain all of $\psi_2(0)^+$ at the beginning of A . Therefore, the square is a factor of

$$nT(n)\psi_2(0)[:-1] = \psi_2((n-2)R_{n-2}[:-1](n-2)R_{n-2}[:-1])[:-1].$$

We first consider that $R_{n-2}[:-1](n-2)R_{n-2}[:-1] = R_{n-1}[:-1]$ is square-free, so by Proposition 20, $\psi_2(R_{n-2}R_{n-2}[:-1])$ is square-free meaning that the square intersects the first chunk, $\psi_2(n-2)$.

This means that the square contains all of the middle occurrence of $\psi_2(n-2)$ which contains four occurrences of n . Since yy contains n and $\max(nT(n)) = n$ we have $n = k$. Also, there are exactly 8 occurrences of n in $nT(n)A$, 4 in each occurrence of $\psi_2(n-2)$. The square yy must include the last 4 occurrences of n and either none of the earlier ones, just the last 6 n 's, or all 8 n 's. Recall that

$$nT(n)\psi_2(0)[:-1] = \underline{\psi_2(n-2)} \psi_2(R_{n-2}[:-2]) \psi_2(0) \underline{\psi_2(n-2)} \psi_2(R_{n-2}[:-2]) \psi_2(0)[:-1]$$

and from Equation (1) in Section 3,

$$\psi_2(n-2) = \underline{n} \underline{R_n} R_n[:-2] \psi_1(R_{n-1}[:-1]) \underline{n} \underline{R_n} R_n[:-2] \psi_1(R_{n-1}[:-2]),$$

which shows the locations of the 4 occurrences of n in $\psi_2(n - 2)$.

If yy contains only the last 4 occurrences of n , then $|y| = |nR_nR_n[: -2]\psi_1(R_{n-1}[: -1])|$. But yy contains $\psi_2(0)\psi_2(n - 2)$ which has length more than twice the length of $nR_nR_n[: -2]\psi_1(R_{n-1}[: -1])$. This is a contradiction.

If yy contains only the last 6 occurrences of n , we consider the first and second n in each half. The first two n 's in the first half occur together as nR_n . But the first two n 's in the second half occur in $R_n R_n[: -2] \psi_1(R_{n-1}[: -1]) \underline{n}$. Clearly, $nR_n[: -2]\psi_1(R_{n-1}[: -1])n \neq nR_n$, so this is a contradiction.

If yy contains all 8 occurrences of n , then it starts at the first letter of $nT(n)$. Then $y = \psi_2(n - 2)\psi_2(R_{n-2}[: -2])\psi_2(0)$, but then $yy = nT(n)\psi_2(0)$ is not a factor of $nT(n)\psi_2(0)[: -1]$. So this is also a contradiction. \square

Lemma 51. $A\rho(\alpha(L(\varepsilon)))$ is square-free.

Proof. Suppose $A\rho(\alpha(L(\varepsilon)))$ contains a square. Since A is square-free, we can use Remark 49 to see that the square contains no letter greater than $\max(A\rho(\alpha(0))) = 5$. Since $\rho(\alpha(L(\varepsilon)))$ is square-free, the square overlaps A . For all letters $n \geq 0$, $\max(\rho(\alpha(n))) = n + 4$, so the square would need to be contained in $A\rho(\alpha(R_2))$ which is a prefix of $A\rho(\alpha(L(\varepsilon)))$ that contains 6. We can computationally verify that $A\rho(\alpha(R_2))$ is square-free which is a contradiction. \square

Theorem 52. For all $n \geq 3$, $nT(n)A\rho(\alpha(L(\varepsilon)))$ is square-free.

Proof. Suppose that there is a square yy in $nT(n)A\rho(\alpha(L(\varepsilon)))$. We have from Lemmas 50 and 51 that $nT(n)A$ and $A\rho(\alpha(L(\varepsilon)))$ are both square-free. So the square yy must contain all of A and overlap some nonempty suffix of $nT(n)$ and some nonempty prefix of $\rho(\alpha(L(\varepsilon)))$.

Consider the prefix $p := A[: 254]$ and the suffix $s := A[-88 :]$, and define w such that $A = pws$. Since A is totally contained in yy , then at least one of p or s must be totally contained in y . We will show that p and s each occur exactly once in $nT(n)A\rho(\alpha(L(\varepsilon))) = nT(n)pws\rho(\alpha(L(\varepsilon)))$, which leads to a contradiction, since at least one of p or s must appear in both halves of the square.

First we show that p appears exactly once in $nT(n)A\rho(\alpha(L(\varepsilon)))$. We can verify computationally that p occurs exactly once in A . Also, since p begins with $\psi_2(0)^+$ which by Lemma 19 never occurs in $nT(n)$, then p cannot occur in $nT(n)$. Moreover, if p occurred over the boundary between $nT(n)$ and A , then since p is a prefix of A , there would be a square in $nT(n)A$ which is not true according to Lemma 50. Finally, since p ends with 12 and $\rho(\alpha(L(\varepsilon)))$ is even-grounded we conclude that p occurs neither in $\rho(\alpha(L(\varepsilon)))$ nor on the boundary between A and $\rho(\alpha(L(\varepsilon)))$.

Secondly, we show that s appears exactly once in $nT(n)A\rho(\alpha(L(\varepsilon)))$. It can be checked that s occurs exactly once in A . To show that s is not contained in $nT(n) = \psi_2((n - 2)P_0(n - 2))$ we use the following properties of s , which can be verified computationally: s is even-grounded, $\psi_2(0)$ does not contain s , $\max(s) = 4$ and s contains 7 occurrences of 4. Since s is even-grounded and for all $n \geq 0$, $\psi_2(n)$ begins and ends with nonzero letters, s cannot lie over a ψ_2 chunk boundary. So if s occurs in $nT(n)$, it is within $\psi_2(k)$ for

some $k > 0$. We can computationally verify that s does not occur in $\psi_2(1)$ or $\psi_2(2)$, so assume $k > 2$. Consider that $\psi_2(k) = \psi_1((k+1)P_0(k+1)) = \psi_1((k+1)R_{k+2}[: -2])$ and that $\psi_1(\ell)$ contains no 4's when $\ell < 3$, two 4's and no 5's when $\ell = 3$, and contains 5's when $\ell > 3$. So if s is contained in $\psi_2(k)$, it must contain at least two whole occurrences of $\psi_1(3)$, and no whole occurrence of $\psi_1(4)$. But since $k > 2$, any two occurrences of 3 in $(k+1)R_{k+2}[: -2]$ have an occurrence of 4 between them. So s cannot be contained in $nT(n)$.

Note that A begins with 2 and recall from Remark 25 that $T(n)$ has suffix $T(3)$ which ends with a 1. Hence, since s is grounded, s cannot lie over the boundary between $nT(n)$ and A . Also, if s occurred over the boundary between A and $\rho(\alpha(L(\varepsilon)))$, then since s is a suffix of A , there would be a square in $A\rho(\alpha(L(\varepsilon)))$, which is not true according to Lemma 51.

Finally we show that s cannot occur in $\rho(\alpha(L(\varepsilon)))$. Since s is even-grounded and has even length, $\rho^{-1}(s)$ is well-defined. Hence, it is enough to show that $\rho^{-1}(s)$ does not occur in $\alpha(L(\varepsilon))$. The first two letters of $\rho^{-1}(s)$ are 13. For $n \geq 0$, $\alpha(n)$ begins with zero, so 13 cannot occur over a chunk boundary in $\alpha(L(\varepsilon))$. Also, for $n \geq 1$, $\alpha(n)$ is grounded so it does not contain 13. We can verify directly that $\rho^{-1}(s)$ does not occur in $\alpha(0)$, so s does not occur in $\rho(\alpha(L(\varepsilon)))$. \square

Proposition 53. *For $n \geq 3$, $L(n)$ has prefix $nT(n)A\rho(G)$.*

Proof. We know from Theorem 24 that $L(n)$ has prefix $nT(n)$. First we show that $L(n)$ has prefix $nT(n)\psi_2(0)^+$. Indeed, using Remark 25, $T(n)$ has suffix $T(3)$ which has suffix $\psi_2(1)$, which according to Lemma 21 generates $\psi_2(10)$. This means that $\psi_2(0)$ at the end of $nT(n)\psi_2(0)$ is irreducible. However, $\psi_2(0)$ also introduces a square:

$$\begin{aligned} nT(n)\psi_2(0) &= nP_0(n)P_1(n)\psi_2(R_{n-2}R_{n-2}[: -2])\psi_2(0) \\ &= \psi_2(n-2)\psi_2(R_{n-2}[: -1])\psi_2(n-2)\psi_2(R_{n-2}[: -2])\psi_2(0) \\ &= \psi_2(n-2)\psi_2(R_{n-2}[: -1])\psi_2(n-2)\psi_2(R_{n-2}[: -1]) \\ &= \psi_2((n-2)R_{n-2}[: -1])^2. \end{aligned}$$

Therefore $nT(n)\psi_2(0)^+$ is irreducible. It is also square-free by Proposition 52, since A has prefix $\psi_2(0)^+$.

Now, from Remark 25 we know that $nT(n)\psi_2(0)^+$ has suffix $T(3)\psi_2(0)^+$. A computer can then verify that this generates $T(3)A\rho(G)$. Since G is a prefix of $\alpha(0)$ we obtain from Proposition 52 that $nT(n)A\rho(G)$ is square-free, which concludes the proof. \square

We can now prove that $L(n) = Y_n\rho(\alpha(L(\varepsilon)))$.

Proof of Theorem 2. From Proposition 53 we get $L(n) = L(nT(n)A\rho(G)) = L(Y_n\rho(G))$. Also, Theorem 26 implies that $L(\rho(G)) = \rho(L(G))$, so by Corollary 47 we have that

$$Y_nL(\rho(G)) = Y_n\rho(L(G)) = Y_n\rho(\alpha(L(\varepsilon))),$$

which is square-free by Proposition 52. Hence, Theorem 2 follows from Remark 4 with $p = \rho(G)$, $u = Y_n$ and $w = L(\rho(G))$. \square

4.3 The structure of $L(2)$

In this section, we briefly describe a conjectured structure for $L(2)$ that is similar to the structures of $L(n)$ in the previous sections.

First note that R_n can be written recursively as

$$\begin{aligned} R_1 &= 01, \\ R_n &= R_{n-1}R_{n-1}^+. \end{aligned}$$

Define

$$\begin{aligned} b_2 &= 0102012021012, \\ b_n &= b_{n-1}b_{n-1}^+R_{n-1}R_{n-1}^+ = b_{n-1}b_{n-1}^+R_n. \end{aligned}$$

Also, let c_3 be the 261-letter word:

$$\begin{aligned} c_3 &= 0102012021012010201202102010210120102012021012010201301020103 \\ &0102012021012010201202101301020103010201202101201020120230102 \\ &0103010201202101201020120301020103010203010302010203010201030 \\ &1020301030201202101201020120210120230102010301020120210120102 \\ &01202101301020103, \\ c_n &= c_{n-1}c_{n-1}^+R_{n-1}R_{n-1}^+b_{n-1}b_{n-1}^+R_{n-1}R_{n-1}^+ \\ &= c_{n-1}c_{n-1}^+R_nb_n. \end{aligned}$$

Notice that for all n , c_n has c_{n-1} as a prefix. Thus, c_n has all previous c_k ($3 \leq k < n$) as prefixes and we have

$$\begin{aligned} \lim_{n \rightarrow \infty} c_n &= c_3 c_3^+ R_4 b_4 c_4^+ R_5 b_5 c_5^+ R_6 b_6 \cdots \\ &= c_3 c_3^+ R_4 b_4 c_3 c_3^+ R_4 b_4^+ R_5 b_5 c_4 c_4^+ R_5 b_5^+ R_6 b_6 \cdots \\ &= \underline{c_3 c_3^+} R_4 b_4 \underline{c_3 c_3^+} R_4 b_4^+ R_5 b_5 \underline{c_3 c_3^+} R_4 b_4 \underline{c_3 c_3^+} R_4 b_4^+ R_5 b_5^+ R_6 b_6 \cdots, \end{aligned}$$

which gives rise to the following morphism:

Definition. For all $n \geq 0$, $\gamma(n)$ is the morphism defined by

$$\begin{aligned} \gamma(0) &= c_3 c_3^+ \\ \gamma(n) &= R_4 b_4^+ R_5 b_5^+ \cdots R_{n+2} b_{n+2}^+ R_{n+3} b_{n+3}. \end{aligned}$$

From the structure of c_n and γ , we can see that

$$\lim_{n \rightarrow \infty} c_n = \gamma(L(\varepsilon)).$$

Conjecture 3. $L(2) = 2 \lim_{n \rightarrow \infty} c_n = 2\gamma(L(\varepsilon)).$

5 Extending from known words

In this section we give two results establishing conditions for when $L(uv) = uL(v)$ for words u and v . If uv is square-free, then so is $L(uv)$. Thus, by Remark 4, $uL(v)$ being square-free is a necessary and sufficient condition for $L(uv) = uL(v)$ when uv is square-free. The following result lets us use our knowledge of $L(n)$ to show that we can omit the square-free condition when u and v are letters ≥ 3 , and Theorem 59 demonstrates a test for the case when uv is not square-free and has a particular structure. The proof of the latter theorem does not use Theorem 2.

Lemma 54. *For all $n_1, n_2 \geq 0$, if the word n_1n_2 is not a factor of $L(n_2)$, then $L(n_1n_2) = n_1L(n_2)$.*

Proof. Since n_1n_2 never occurs in $L(n_2)$ and $L(n_2)$ is square-free, $n_1L(n_2)$ is square-free unless $n_1 = n_2$ in which case its only square factor is the prefix n_1n_1 . Thus, $L(n_1n_2)$ and $n_1L(n_2)$ are both infinite words beginning with n_1n_2 whose only square factors are contained in the prefix n_1n_2 . So by the definition of L , $L(n_1n_2) \preceq n_1L(n_2)$. If $L(n_1n_2) \prec n_1L(n_2)$, then $L(n_1n_2)[1:] \prec L(n_2)$, which is a contradiction since these are both infinite square-free words beginning with n_2 . Therefore, $L(n_1n_2) = n_1L(n_2)$. \square

This immediately describes all words of the form $L(nn)$:

Theorem 55. *For all $n \geq 0$, $L(nn) = nL(n)$.*

Proof. Since $L(n)$ is square-free, it does not contain nn . The result then follows from Lemma 54. \square

Theorem 56. *For all $n_1 \geq 3$ and $n_2 \geq 3$, we have $L(n_1n_2) = n_1L(n_2)$.*

Proof. By Lemma 54, it is sufficient to show that the word n_1n_2 never appears in $L(n_2)$.

By Theorem 2, $L(n_2) = Y_{n_2}\rho(\alpha(L(\varepsilon)))$. Since $\rho(\alpha(L(\varepsilon)))$ is grounded, it cannot contain n_1n_2 . Also, the first letter of $\rho(\alpha(L(\varepsilon)))$ is 0, so n_1n_2 cannot lie over the boundary. For the prefix we have

$$Y_{n_2} = n_2T(n_2) = \psi_2((n_2 - 2)P_0(n_2 - 2)),$$

hence it is enough to show that n_1n_2 does not occur in any ℓ -chunk $\psi_2(\ell)$, nor over any chunk boundary.

Since $\max(\psi_2(0)) = 3$, n_1n_2 could only occur in a 0-chunk if $n_1n_2 = 33$, which is not a factor of $\psi_2(0)$. For the case $\ell \geq 1$, $\psi_2(\ell) = \psi_1((\ell + 1)P_0(\ell + 1))$. Hence, it is sufficient to show that n_1n_2 cannot occur in any $\psi_1(\ell)$ nor over the chunk boundary of any $\psi_1(0\ell)$ or $\psi_1(\ell 0)$. Indeed, since $\psi_1(0) = 202101$, and all other $\psi_1(\ell)$ are grounded, then n_1n_2 cannot occur in any $\psi_1(\ell)$. Also, $\psi_1(\ell)$ ends with a 1 for all $\ell \geq 0$, so n_1n_2 cannot occur in $\psi_1(0\ell)$ or in $\psi_1(\ell 0)$.

Finally, to show that n_1n_2 does not occur over any chunk boundary of ψ_2 recall that for $\ell \geq 1$, $\psi_2(\ell)$ has suffix

$$P_1(\ell + 2) = \psi_1(P_0(\ell + 1)) = \psi_1(R_{\ell+2}[: -2]),$$

which has suffix $\psi_1(1)$, which ends with a 1, so n_1n_2 cannot lie over a $\psi_2(\ell 0)$ chunk boundary. Also, $\psi_2(0)$ ends with 2 so n_1n_2 cannot lie over a $\psi_2(0\ell)$ chunk boundary.

Therefore, n_1n_2 cannot occur anywhere in $n_1L(n_2)$, except as a prefix. And so $L(n_1n_2) = n_1L(n_2)$. \square

Experiments suggest the following related result.

Conjecture 57. For all $n \geq 3$, we have $L(n1) = nL(1)$ and $L(n2) = nL(2)$.

For example, it appears that $L(31) = 3L(1)$ and $L(32) = 3L(2)$. Since Theorem 26 implies that $L(0n) = \rho(L(n-1))$ for all $n > 0$, we have a proven or conjectural description of $L(w)$ for all 2-letter words w except for $L(1n)$ when $n > 1$ and $L(2n)$ when $n \notin \{0, 2\}$. However, it does appear that these words also have structures related to the ruler sequence and to the other words discussed in this paper.

The rest of this section deals with the case when $w = uv$ is not square-free and is a particular decomposition of w . The next lemma describes this decomposition.

Lemma 58. *Let w be any nonempty finite word containing a square. Then there is a unique decomposition $w = psq$ such that sq is the maximal square-free suffix of w , and $p[-1]s$ is the maximal square prefix of $p[-1]sq$.*

Proof. Any single letter is square-free, so w is guaranteed to have some square-free suffix. Each suffix of w has a different length, so the maximal square-free suffix, sq is unique. Since w contains a square, sq is a proper suffix of w , so p is nonempty and unique.

If $p[-1]sq$ is square-free, then it would be a square-free suffix larger than sq which is a contradiction. Any square in $p[-1]sq$ cannot be contained in sq which is square-free, so $p[-1]sq$ has a square prefix. No two distinct prefixes of have the same length, so the maximal square prefix of $p[-1]sq$ is unique. \square

Remark. In the $w = psq$ decomposition in Lemma 58, p and s are always nonempty, while q can be empty. The last letter of w is always square-free, so sq is always nonempty. Since w contains a square, we cannot have $sq = w$ so p is nonempty. Since $p[-1]$ is a single letter, it cannot be a square, so s must be nonempty.

Example. For $w = 012323045$, we have that $p = 012$, $s = 323$, $q = 045$.

For $w = 1121123210$, we have that $p = 1121$, $s = 1$, and $q = 23210$.

For $w = 11011$, we have that $p = 1101$, $s = 1$, and $q = \varepsilon$.

Theorem 59. *Let w be any nonempty finite word containing a square. Write $w = psq$ such that sq is the maximal square-free suffix of w , and $p[-1]s$ is the maximal square prefix of $p[-1]sq$. Then $L(psq) = pL(sq)$ if and only if $L(psq)[: 2|ps|] = (pL(sq))[[: 2|ps|]]$.*

In other words, to verify that $L(psq) = pL(sq)$, it is sufficient to verify that they match for their first $2|ps|$ letters. Note that this is potentially useful because $L(sq)$ is the maximal square-free tail of $pL(sq)$. Before proving Theorem 59, we look at a few examples.

Example. For $w = 012323045$, since $ps = 012323$ the theorem implies that $L(w) = 012L(323045)$ if and only if their first $2|ps| = 12$ letters match. Since $|w| = 9$, and both words have w as a prefix, it is sufficient to compute the next three letters of each. In this case, both have 010 as their next letters and so we conclude that $L(012323045) = 012L(323045)$.

For $w = 1121123210$, since $ps = 11211$ we obtain that $L(w) = 1121L(123210)$ are equal if and only if they match for the first $2|ps| = 10$ letters. Since $|w| = 10$, no further computations are necessary.

For $w = 11011$, since $ps = 11011$ we have that $L(w) = 1101L(1)$ if and only if they match for the first $2|ps| = 10$ letters. However, in this case $L(11011)[: 10] = 1101120102 \neq 1101101201 = 1101L(1)[: 10]$.

Proof of Theorem 59. The forward direction is trivial. We prove the other direction by induction. The base case is our supposition that $L(w)[: 2|ps|] = (pL(sq))[: 2|ps|]$. Now suppose $L(w)[: n] = (pL(sq))[: n]$ for some $n \geq 2|ps|$. We will prove that $L(w)[: n+1] = (pL(sq))[: n+1]$. We let $a = L(w)[n]$ and $b = (pL(sq))[n] = L(sq)[n - |p|]$, then we need to show that $a = b$.

Suppose that $a < b$, which implies that $(pL(sq))[: n]a$ has a square suffix. By the inductive hypothesis, $(pL(sq))[: n]a = L(w)[: n]a = L(w)[: n+1]$. By definition of L , squares in $L(w)$ are contained in the prefix $w = L(w)[: |w|]$. Thus, $n < |w|$. But then $a = L(w)[n] = w[n] = (pL(sq))[n] = b$. Therefore we cannot have $a < b$.

Now suppose that $a > b$, which analogously to the previous case implies that $L(w)[: n]b$ has a square suffix, say yy . By the inductive hypothesis, $L(w)[: n]b = (pL(sq))[: n]b = (pL(sq))[: n+1]$. Since sq is square-free, $L(sq)$ is square-free, so the square must start in the prefix p . Let k be the length of the suffix of p contained in the square. Then $1 \leq k \leq |p|$ and we have

$$(pL(sq))[: n]b = p(L(sq)[: n - |p|])b$$

so the square has length

$$|yy| = k + n - |p| + 1 \geq k + 2|ps| - |p| + 1 = k + |p| + 2|s| + 1.$$

Note that the first half of the square cannot contain all of s , otherwise it would contain $p[-1]s$ which is a square. This in turn would imply that the second half contains $p[-1]s$ and is a factor of $L(sq)$ which is square-free. Therefore, the first half of the square ends within $ps[: -1]$, and so $|y| < k + |s|$.

This implies that $|yy| = 2|y| < 2k + 2|s| \leq k + |p| + 2|s|$ which is a contradiction since $|yy| = k + |p| + 2|s| + 1$. Therefore, $a > b$ is not possible either.

This proves the inductive step, hence $L(w)[: n] = (pL(sq))[: n]$ for all n , as wanted. \square

6 Inducing factors

In this section we consider the following problem. Given a finite square-free word w , find a word p (not necessarily square-free) such that p generates pw , i.e. $L(p) = L(pw)$.

Definition. Let w be a finite square-free word. For $0 \leq j < |w|$ and $0 \leq k < w[j]$, we call a nonempty word of the form

$$r_{j,k}(w) = w[:j]k$$

a *restriction* of w . That is, $r_{j,k}(w)$ is obtained by starting with $w[:j+1]$ and decreasing the last letter by some amount. Let $m(w)$ be the total number of square-free restrictions of w , and relabel the square-free restrictions in the lexicographic order as $r_0(w), \dots, r_{m-1}(w)$, this is called the *restriction sequence* of w .

Example. For the word $w = 2021$, we have

$$r_{0,0} = 0, r_{0,1} = 1, r_{2,0} = 200, r_{2,1} = 201, \text{ and } r_{3,0} = 2020.$$

Taking only the square-free $r_{j,k}$ and sorting them lexicographically, we obtain that the restriction sequence of w is $r_0 = 0, r_1 = 1, r_2 = 201$.

From the definition of lexicographic order, the set of restrictions is always totally ordered by \prec . For the rest of this section, we will just write $r_{j,k}$, r_i , and m where the dependence on w is inferred by context.

Intuitively, the restriction sequence of w does the following: Whenever we extend a word with L , we follow the ruler sequence until we can't anymore due to a square. So, to generate w with L , we need to have exactly the right squares coming up at the right positions to deviate from the ruler sequence and spell out w instead. The restriction sequence for w is composed of all the words that will provide those necessary squares.

In other words, the restriction sequence is the collection of all square-free words that are not longer than w , and are lexicographically less than w . We then design p so that pr_i contains a square for all i . Thus, p will generate the lexicographically least word that is greater than all r_i , which is w .

Definition. Let w be a finite square-free word, and let $v_i(w) = \max(w) + i + 1$ and $V_i(w) = v_i \cdots v_1 v_0$. For $i \leq m = m(w)$, we define words $x_i(w)$ by

$$\begin{aligned} x_0 &= v_0, \\ x_i &= v_i x_{i-1} r_{i-1} x_{i-1}, \end{aligned}$$

where $v_i = v_i(w)$, $V_i = V_i(w)$, and $x_i = x_i(w)$. This dependence on w will be inferred by context.

Note that for all $j \leq i$, x_i has suffix x_j and so x_m has all x_j 's as suffixes. Also, for all i , $\max(r_i) \leq \max(w) < v_0 < v_1 < v_2 < \dots$ and $\max(x_i) = v_i$.

Example. Continuing with the example $w = 2021$, since $\max(w) = 2$, we have $v_i = i + 3$ for $0 \leq i \leq m = 3$. Then

$$\begin{aligned} x_0 &= v_0 &&= 3, \\ x_1 &= v_1 x_0 r_0 x_0 &&= 4303, \\ x_2 &= v_2 x_1 r_1 x_1 &&= 5 \ 4303 \ 1 \ 4303, \\ x_3 &= v_3 x_2 r_2 x_2 &&= 6 \ 5430314303 \ 201 \ 5 \ 4303 \ 1 \ 4303. \end{aligned}$$

We can write x_3 with spacing suggestive of the next lemma:

$$\begin{aligned} x_3 &= 6543 \cdot 03 \cdot 143 \cdot 03 \cdot 201543 \cdot 03 \cdot 143 \cdot 03 \\ &= V_3 \cdot r_0V_0 \cdot r_1V_1 \cdot r_0V_0 \cdot r_2V_2 \cdot r_0V_0 \cdot r_1V_1 \cdot r_0V_0. \end{aligned}$$

Lemma 60. *Let w be a finite square-free word, and let ϕ_w be the morphism defined on the alphabet $\{0, 1, \dots, m\}$ by $\phi_w(k) = r_kV_k$ for letters k . Then for $0 \leq i \leq m$, $x_i = V_i\phi_w(R_i[: -1])$.*

Proof. We proceed by induction. For the base case,

$$x_0 = v_0 = V_0 = V_0\phi_w(\varepsilon) = V_0\phi_w(R_0[: -1]).$$

For the inductive step, suppose $x_i = V_i\phi_w(R_i[: -1])$ for some $i < m$. Then

$$\begin{aligned} x_{i+1} &= v_{i+1}x_i r_i x_i \\ &= v_{i+1}V_i\phi_w(R_i[: -1]) r_i V_i\phi_w(R_i[: -1]) \\ &= V_{i+1}\phi_w(R_i[: -1]) \phi_w(i) \phi_w(R_i[: -1]) \\ &= V_{i+1}\phi_w(R_i[: -1]iR_i[: -1]) \\ &= V_{i+1}\phi_w(R_{i+1}[: -1]). \quad \square \end{aligned}$$

Finally we present the main result of this section, which implies that for every finite square-free word w , there exists a prefix p that generates pw . Indeed, it states that such prefix is given by $p = x_m$.

Theorem 61. *Let w be a finite square-free word. Then x_m generates x_mw .*

Proof. We will show that for all $0 \leq j < |w|$, $x_mw[: j]$ generates $x_mw[: j + 1]$. This means that $L(x_mw[: j]) = L(x_mw[: j + 1])$, which proves the desired result:

$$L(x_m) = L(x_mw[: 0]) = L(x_mw[: 1]) = \dots = L(x_mw[: |w|]) = L(x_mw).$$

To show that $L(x_mw[: j]) = L(x_mw[: j + 1])$ we need to prove the last letter of $x_mw[: j + 1]$, which is $w[j]$, is irreducible and does not introduce a square.

First we prove the irreducibility condition. Let $0 \leq j < |w|$ and $\ell < w[j]$, we need to show that $x_mw[: j]\ell$ has a square suffix. Indeed, since $0 \leq j < |w|$ and $0 \leq \ell < w[j]$, we have that $w[: j]\ell = r_{j,\ell}(w)$. If $w[: j]\ell$ contains a square, since w is square-free, this must be a square suffix, and we are done. Otherwise, if $w[: j]\ell$ is square-free, then w has a restriction $r_i = w[: j]\ell$ for some $i < m$. Hence x_m has suffix x_{i+1} , which means that $x_mw[: j]\ell$ has suffix

$$x_{i+1} w[: j]\ell = v_{i+1}x_i r_i x_i r_i,$$

which also has a square suffix. Therefore $w[j]$ is irreducible in $x_mw[: j + 1]$.

Now we prove that $w[j]$ does not introduce a square in $x_mw[: j]$. Suppose toward a contradiction that $x_mw[: j + 1]$ has square suffix yy . Since w is square-free, the square must start in the prefix x_m , and so y contains the last letter of x_m which is $x_0 = v_0$.

Hence, since $v_0 > \max(w)$, y cannot be completely contained in w . Therefore, the second half of the square starts in x_m and contains all of $w[: j + 1]$. This implies that y has suffix $v_0w[: j + 1]$.

By Lemma 60,

$$x_m = V_m\phi_w(R_m[: -1]) = V_m \cdot r_0V_0 \cdot r_1V_1 \cdot r_0V_0 \cdot r_2V_2 \cdots r_0V_0,$$

where $\phi_w(k) = r_kV_k$ for letters $k < |w|$. Since $y[-1] = w[j] \leq \max(w)$ and all letters in any V_k are greater than $\max(w)$, we have that the last letter of the first half of the square is in a factor r_i for some i . Therefore, there are no partial V_k 's in the second half.

Since $\max(\phi_w(k)) = v_k > \max(w)$, the largest letter in y is some letter $V_k[0] = v_k$ occurring k -chunk. Let $v_\ell = \max(y)$, then every occurrence of v_ℓ in the second half of the square is as the first letter of V_ℓ in an ℓ -chunk. In order to contain v_ℓ , the first half of the square must overlap a k -chunk with $k \geq \ell$. From the structure of the ruler sequence, we know that the latest k -chunk with $k \geq \ell$ before the first ℓ -chunk in the second half is an $(\ell + 1)$ -chunk. The first half cannot contain the letter $v_{\ell+1}$ in this $(\ell + 1)$ -chunk, but it must include the letter v_ℓ . Therefore, v_ℓ is the first letter of the first half and since there are no partial V_k 's in the second half, then y has prefix V_ℓ . The second half then begins in the middle of an ℓ -chunk at the beginning of the factor V_ℓ and so the first half has suffix r_ℓ from the same ℓ -chunk. Before this factor r_ℓ there is another chunk, which always ends with v_0 and so y has suffix v_0r_ℓ .

We have proved that the words $v_0w[: j + 1]$ and v_0r_ℓ are both suffixes of y . Since $v_0 > \max(w[: j + 1])$ and $v_0 > \max(r_\ell)$, we have that $r_\ell = w[: j + 1]$. But this is a contradiction since the restriction r_ℓ cannot be a prefix of w . This proves that no such square exists and the result follows. \square

Example. Again using $w = 2021$, we can verify that

$$x_3 = 654303143032015430314303 \text{ generates } x_3w,$$

for example the suffix 303 prevents a 0, the suffix 430314303 prevents a 1, so since 2 does not introduce a square, it can be located after this suffix. Similarly, we can continue checking that the whole word w is the lexicographically least extension of x_3 .

7 Glossary

A list of all the important mathematical objects in the paper, along with their definitions.

7.1 Sequences of Words

R_n is defined for all letters $n \geq 0$, and is the ruler sequence up to the first appearance of n . We can also define R_n by $R_n = \rho^n(0)$, or inductively by $R_0 = 0$ and $R_n = R_{n-1}R_{n-1}^+$. R_n is always even-grounded, the length of R_n is 2^n , and $\max(R_n) = n$.

b_n is defined for letters $n \geq 2$, and is defined inductively by $b_2 = 0102012021012$ and $b_n = b_{n-1}b_{n-1}^+R_n$. b_n is never grounded, the length of b_n is $2^{n-2}(4n + 5)$, and $\max(b_n) = n$.

c_n is defined for letters $n \geq 3$, and is defined inductively by $c_n = c_{n-1}c_{n-1}^+R_nb_n$ with a base case c_3 of length 261. c_n is never grounded, the length of c_n is $2^{n-3}(4n^2 + 22n + 159)$, and $\max(c_n) = n$.

7.2 Functions on Letters

$P_0(n)$ is defined for all letters $n \geq 0$, and is the largest prefix of the ruler sequence such that $nP_0(n)$ is a prefix of $L(n)$. The length of $P_0(n)$ is $2^{n+1} - 2$, so $P_0(n) = R_{n+1}[: -2]$.

$P_1(n)$ is defined for letters $n \geq 3$, and is equal to $\psi_1(P_0(n - 1))$. The length of $P_1(n)$ is $(4n + 1)2^{n-1} - 5$.

$P_2(n)$ is defined for letters $n \geq 3$, and is equal to $\psi_2(P_0(n - 2))$. The length of $P_2(n)$ is $(4n^2 + 14n + 149)2^{n-2} - 193$.

$T(n)$ is defined for letters $n \geq 3$, and is equal to $P_0(n)P_1(n)P_2(n)$. The length of $T(n)$ is $(4n^2 + 22n + 159)2^{n-2} - 200$.

7.3 Morphisms

The ruler morphism ρ is defined by $\rho(n) = 0(n + 1)$.

ψ_1 is defined by $\psi_1(0) = 202101$ and $\psi_1(n) = (n + 1)P_0(n + 1)$ for $n > 0$. The length of $\psi_1(n)$ is $2^{n+2} - 1$ for $n > 0$.

ψ_2 is defined by $\psi_2(n) = (n + 2)P_0(n + 2)P_1(n + 2)$ for $n > 0$, with $\psi_2(0)$ a specific word of length 199. For $n > 0$, the length of $\psi_2(n)$ is $(4n + 13)2^{n+1} - 6$.

α is defined by

$$\alpha(n) = \begin{cases} EFE & \text{if } n = 0 \\ B_1 R_4 C B_1 R_4 & \text{if } n = 1 \\ \alpha(n - 1)^+ R_{n+3} C \alpha(n - 1)^+ R_{n+3} & \text{if } n \geq 2 \end{cases}$$

for constants C , B_0 , B_1 , E , and F .

7.4 Constants

ε is the empty word.

$C = 0102030102$ is a grounded word of length 10 which is used in the inductive definition of α . While not mathematically relevant, we would be remiss not to note that coding C into letters yields the word **abacadabac**, which is amusingly similar to **abracadabra**.

$B_0 = 0301 \psi_1(1010)[: -3] \psi_2(1010)[: -6] \psi_2(10)[: -12] 301020$ is a non-grounded word of length 798 which is used in the definition of $\alpha(0)$.

$B_1 = \rho(B_0[7 : -5])$ is a grounded word of length 1572 which is used in the definition of $\alpha(1)$, which is the base case for the inductive definition of α .

$E = 0102B_01B_0[: -9]$ and $F = B_0[-9 :]3010302C0103C^+02$. These are useful in the structure of α because $EFE = \alpha(0)$.

$G = 010203012$ is a word of length 9, which is the shortest prefix that generates $\alpha(0)$ and appears in various proofs.

A is a word of length 13747 with the property that $nT(n)A$ is a prefix of $L(n)$ for all $n \geq 3$.

Acknowledgements

This work was done in the 2021 Polymath Jr. program and was partially supported by NSF award DMS-2113535. We thank the organizers of the Polymath Jr. program 2021 for creating the environment that allowed this collaboration. We also thank our colleagues Alycia Doucette, Bridget Duah, Bill Feng, Mordechai Goldberger, Luke Hammer, Ziqi He, Amanda Lamphere, Mary Olivia Liebig, Jacob Micheletti, Adil Oryspayev, Sara Salazar, Shiyao Shen, Wangsheng Song, and Thomas Sottosanti, who contributed in coding, generating data, and presenting results.

References

- [1] J.-P. Allouche, A short walk across combinatorics of words, *Da Pitagora a Schützenberger Symposium* (2018), University of Calabria, in Cosenza and Crotona. <http://www-igm.univ-mlv.fr/~berstel/Colloque-Pitagore/word-survey-march2020.pdf>.
- [2] J.-P. Allouche, James Currie, and Jeffrey Shallit, Extremal infinite overlap-free binary words, *The Electronic Journal of Combinatorics* **5** (1998) #R27, 11 pages.
- [3] J. Cassaigne, An algorithm to test if a given circular HDOL-language avoids a pattern, *IFIP World Computer Congress'94* Elsevier (North-Holland) **1** (1994) 459–464.
- [4] M. Guay-Paquet and J. Shallit, Avoiding squares and overlaps over the natural numbers, *Discrete Mathematics* **309** (2009) 6245–6254.
- [5] M.G. Main, and R.J. Lorentz, Linear time recognition of squarefree strings, *Combinatorial Algorithms on Words* NATO ASI Series, vol. F 12 (1985) 271–278. Springer, Heidelberg.
- [6] L. Pudwell and E. Rowland, Avoiding fractional powers over the natural numbers, *The Electronic Journal of Combinatorics* **25** (2018) #P2.27, 46 pages.
- [7] E. Rowland and J. Shallit, Avoiding $3/2$ -powers over the natural numbers, *Discrete Mathematics* **312** (2012) 1282–1288.
- [8] E. Rowland and M. Stipulanti, Avoiding $5/4$ -powers on the alphabet of nonnegative integers, *The Electronic Journal of Combinatorics* **27** (2020) #P3.42, 39 pages.
- [9] N. Sloane et al., The On-Line Encyclopedia of Integer Sequences, <https://oeis.org>.