

# A characterization of one-element commutation classes

Ricardo Mamede

José Luis Santos

Diogo Soares

Submitted: Jan 26, 2024; Accepted: Oct 15, 2024; Published: Dec 17, 2024

© The authors. Released under the CC BY-ND license (International 4.0).

## Abstract

A reduced word for a permutation of the symmetric group is its own commutation class if it has no commutation moves available. These words have the property that every factor of length 2 is formed by consecutive integers, but in general words of this form may not be reduced. In this paper we give a necessary and sufficient condition for a word with the previous property to be reduced. In the case of involutions, we give an explicit construction of their one-element commutation classes and relate their existence with pattern avoidance problems.

**Mathematics Subject Classifications:** 05A05, 20F55

## 1 Introduction

Given a positive integer  $n \geq 1$ , let  $\mathfrak{S}_{n+1}$  denote the *symmetric group* on  $n + 1$  elements formed by all permutations of the set  $[n + 1] := \{1, 2, \dots, n + 1\}$ , with composition (read from the right) as group operation. We usually write a permutation  $\sigma$  using the *one-line notation*  $\sigma = [\sigma(1), \dots, \sigma(n + 1)]$ . In some cases, we will also use the *cyclic notation* of a permutation, using parenthesis to represent the cycles with commas to separate the images. For example, the permutation  $\sigma = [2, 5, 7, 1, 4, 3, 6]$  can be written in cyclic notation as  $\sigma = (1, 2, 5, 4)(3, 7, 6)$ .

The group  $\mathfrak{S}_{n+1}$  is generated by the involutions  $\{s_1, \dots, s_n\}$ , also known as *simple reflections*, where  $s_i = [1, \dots, i + 1, i, \dots, n + 1] = (i, i + 1)$ , for all  $i \in [n]$ . This fact can be easily understood by noticing that multiplying a permutation  $\sigma$  on the right by  $s_i$  interchanges the values in positions  $i$  and  $i + 1$  in the one-line notation of  $\sigma$ , that is  $\sigma s_i = [\sigma(1), \dots, \sigma(i + 1), \sigma(i), \dots, \sigma(n + 1)]$ . Since  $\mathfrak{S}_{n+1}$  is generated by the involutions  $s_i$ , any permutation  $\sigma$  can be written as a product  $s_{i_1} s_{i_2} \cdots s_{i_l}$ , with  $i_j \in [n]$ , for all  $j \in [l]$ . When  $l$  is minimal, we say that the product is a *reduced decomposition* and  $i_1 \cdots i_l$  a *reduced word* of  $\sigma$ . The integer  $l(\sigma) := l$  is the *length* of  $\sigma$ . Let  $R(\sigma)$  be the set of all reduced words of  $\sigma$ .

---

University of Coimbra, CMUC, Department of Mathematics, 3000-143 Coimbra, Portugal  
(mamede@mat.uc.pt, zeluis@mat.uc.pt, soaresdiogo@hotmail.com).

Reduced words of permutations are widely studied in combinatorics (see [1, 11, 12, 13, 15]). Perhaps, one of the most important facts concerning reduced words is a well-known result of Tits [15], which says that two reduced words for the same permutation differ by a sequence of the following two types of moves:

$$ij \leftrightarrow ji, \text{ if } |i - j| > 1, \quad (1)$$

$$i(i + 1)i \leftrightarrow (i + 1)i(i + 1), \quad (2)$$

where (1) is called a *commutation move*, and (2) is called a *braid move*.

We can define an equivalence relation on the set  $R(\sigma)$  by setting  $s \sim t$  if  $s$  and  $t$  differ by a sequence of commutation moves. The equivalence classes generated by this relation are called *commutation classes*. These structures were already considered by some authors (see [3, 6, 5, 7, 8]), but there is still much to understand about commutation classes. For instance, there is no known formula for the number of commutation classes of a given permutation. Recently, Tenner [14] studied the commutation classes that have only one reduced word, giving a necessary condition for a reduced word to be its own commutation class in terms of pinnacles and vales. A nice consequence of this result is that the number of one-element commutation classes of  $w_0$ , the longest element in  $\mathfrak{S}_{n+1}$ , is exactly 4, a result previously obtained in [5]. The goal of this paper is to extend Tenner's work on this topic. We start Section 2 by introducing the terminology used in [14] in order to define what we have called a *segment* in a word. This notion will be crucial to prove the main result of Section 3, which is a necessary and sufficient condition for a word to be a one-element commutation class of some permutation. An application of this characterization will be done in Section 4, where we give an explicit construction of the one-element commutation classes for involutions and relate their existence with pattern avoidance problems.

## 2 Definitions and background

Let  $[n]^*$  be the set of all words with finite length over the alphabet  $[n]$ . A *subword* of a word  $s = i_1 \cdots i_l$  in  $[n]^*$  is a word obtained from  $s$  by deleting some of its letters, and a *factor* of  $s$  is a subword of  $s$  of the form  $s_{i_j} s_{i_{j+1}} \cdots s_{i_k}$ , with  $1 \leq j \leq k \leq l$ . When  $j = 1$ , we call it a *left factor* of  $s$ . Given a permutation  $\sigma \in \mathfrak{S}_{n+1}$ , we denote by  $R_\bullet(\sigma) \subseteq R(\sigma)$  the set of reduced words of  $\sigma$  that are their own commutation class. By definition, a word in  $R_\bullet(\sigma)$  has no commutation moves available, which means that all of its factors of length 2 are formed by consecutive integers. These words can be described in terms of its pinnacles and vales, notions used in [14] to study one-element commutation classes, previously introduced in [2, 4].

**Definition 1** ([14]). Let  $s \in [n]^*$  be a word. The endpoints of  $s$  are its leftmost and rightmost letters. A *pinnacle* of  $s$  is a letter that is larger than its immediate neighbor(s), and a *vale* is a letter that is smaller than its immediate neighbor(s). We call pinnacles and vales the *spikes* of  $s$ . Write  $\mathbf{p}(s)$  for the subword of pinnacles of  $s$ , and  $\mathbf{v}(s)$  for the subword of vales. The subword of pinnacles and vales will be written as  $\mathbf{pv}(s)$ . If every

factor of length 2 of  $s$  is formed by consecutive integers, then we say that  $s$  is a word *formed by consecutive integers*.

As an example, if  $s = 23454321234$  we have  $\mathbf{p}(s) = 54$ ,  $\mathbf{v}(s) = 21$  and  $\mathbf{pv}(s) = 2514$ . When  $s$  is a word formed by consecutive integers, each factor  $ij$  of  $\mathbf{pv}(s)$  corresponds in  $s$  to the factor  $i(i+1)\cdots(j-1)j$  if  $i < j$ , or  $i(i-1)\cdots(j+1)j$  if  $i > j$ . Sometimes, instead of writing all of its letters, it will be more useful to write only the endpoints of those factors, and for that we use the notation  $\underline{i}$  and  $\bar{j}$  to denote a vale  $i$  or to denote a pinnacle  $j$ , respectively. When using this identification to represent the entire word, we write  $s \equiv \mathbf{pv}(s)$ . In the example above, we have  $23454321234 \equiv \underline{2}\bar{5}\underline{1}\bar{4}$ . A graphical representation for these type of words can be given using line diagrams.

**Definition 2.** Let  $s \in [n]^*$  be a word formed by consecutive integers with  $\mathbf{pv}(s) = i_1 \cdots i_l$ . The *line diagram* of  $s$  is formed by the set of points  $(j, i_j) \in [l] \times [n]$ , where there is a line segment connecting each pair  $(j, i_j)$  and  $(j+1, i_{j+1})$ , for all  $j \in [l-1]$ .

The line diagram of the word in the previous example is represented in Figure 1. Each




Figure 1: Line diagram of  $\underline{2}\bar{5}\underline{1}\bar{4}$ .

factor of length 2 of  $\mathbf{pv}(s)$  is encoded by a line segment in the line diagram of  $s$ . Thereby, we say that  $\underline{i}\bar{j}$  or  $\bar{j}\underline{i}$  are *segments* of  $s$  if  $ij$  or  $ji$  are factors of  $\mathbf{pv}(s)$ , with  $i < j$ . For our running example, its segments are  $\underline{2}\bar{5}$ ,  $\bar{5}\underline{1}$  and  $\underline{1}\bar{4}$ , which we can see clearly from its diagram in Figure 1. Notice that multiplying a permutation  $\sigma \in \mathfrak{S}_{n+1}$  on the right by the permutation associated to the segment  $\underline{i}\bar{j}$  (resp.  $\bar{j}\underline{i}$ ) has the effect of “moving” the integer  $\sigma(i)$  to position  $j+1$  (resp. the integer  $\sigma(j+1)$  to position  $i$ ) in the one-line notation of  $\sigma$ . In this sense, we say that the segment *moves* an integer. In the above example, the segment  $\underline{1}\bar{4}$  acts in the permutation  $[1, 6, 3, 4, 5, 2]$  (the permutation associated to the left factor  $2345432$  of  $s$ ) by moving the integer in the first position of the one-line notation to position 5, obtaining the permutation  $[6, 3, 4, 5, 1, 2]$ . Not every word formed by consecutive integers is a reduced word. For instance, the word  $1232123 \equiv \underline{1}\bar{3}\underline{1}\bar{3}$  is not reduced. A set of necessary conditions for a word  $s$  to be a one-element commutation class was given in [14] using properties of the words  $\mathbf{p}(s)$ ,  $\mathbf{v}(s)$  and  $\mathbf{pv}(s)$ .

**Definition 3.** Let  $s = i_1 \cdots i_l \in [n]^*$  be a word. If there exist  $j$  and  $k$  such that  $1 \leq j \leq k \leq l$  and

$$i_1 < \cdots < i_j = i_k > \cdots > i_l,$$

then  $s$  is a *wedge*. If

$$i_1 > \cdots > i_j = i_k < \cdots < i_l,$$

then  $s$  is a *vee*. If  $j = k$ , then that wedge or vee is strict.

For example, the word 24731 is a strict wedge and the word 245521 is a wedge that is not strict. The words 42157 and 632245 are examples of a strict vee and a non-strict vee, respectively.

**Proposition 4** ([14, Theorem 3.1]). *For any  $\sigma \in \mathfrak{S}_{n+1}$ , if  $s \in R_\bullet(\sigma)$  then:*

1.  $\mathbf{p}(s)$  is a wedge,
2.  $\mathbf{v}(s)$  is a vee,
3.  $\mathbf{p}(s)$  and/or  $\mathbf{v}(s)$  is strict,
4. the minimum and maximum values of  $\mathbf{pv}(s)$  appear consecutively and,
5. if  $\mathbf{p}(s)$  (or  $\mathbf{v}(s)$ ) has more than one integer  $i$ , then one of those  $i$ 's is an endpoint of  $s$ .

As a consequence, if  $s$  is a word formed by consecutive integers that does not satisfy one of the previous 5 conditions, then  $s$  cannot be a reduced word. It follows directly from condition 3 of the previous theorem that the word  $s \equiv \underline{1}\bar{3}\underline{1}\bar{3}$  is not reduced because neither  $\mathbf{p}(s) = 33$  nor  $\mathbf{v}(s) = 11$  are strict.

The “converse” of this theorem is not true, *i.e.* a word  $s$  formed by consecutive integers that satisfies all the conditions of the previous theorem is not necessarily a reduced word. Consider for instance the word  $s = 2343212345654345 \equiv \underline{2}\bar{4}\underline{1}\bar{6}\underline{3}\bar{5}$ , which satisfies all the conditions of Proposition 4 but is not a reduced word. The reason is that  $s$  contains the factor  $t = 34321234565434 \equiv \underline{3}\bar{4}\underline{1}\bar{6}\underline{3}\bar{4}$  which is not reduced (the permutation associated to  $t$  is  $[5, 2, 7, 3, 4, 6, 1]$  which has length 12, but  $t$  has 14 letters). Notice that  $t$  contains two occurrences of the segment  $\underline{3}\bar{4}$ . This is not a coincidence, as we are going to see in the next section. The line diagrams of  $s$  and  $t$  are depicted in Figure 2.




Figure 2: Line diagrams of  $s \equiv \underline{2}\bar{4}\underline{1}\bar{6}\underline{3}\bar{5}$  and  $t \equiv \underline{3}\bar{4}\underline{1}\bar{6}\underline{3}\bar{4}$ .

We end this section with the following lemma which will be useful more ahead.

**Lemma 5.** *Let  $s \in [n]^*$  be a word formed by consecutive integers. Suppose that  $t$  is a factor of  $s$ . Then:*

1. Every spike of  $t$  that is not an endpoint of  $t$  is also a spike of  $s$ .
2. If  $\underline{i}\bar{j}$  (resp.  $\bar{j}\underline{i}$ ) is a segment of  $t$  that does not contain any endpoint of  $t$ , then  $\underline{i}\bar{j}$  (resp.  $\bar{j}\underline{i}$ ) is also a segment of  $s$ .

*Proof.* If  $i$  is a vale (resp. pinnacle) of  $t$  that is not one of its endpoints, then it is between two letters that are larger (resp. smaller) in the word  $t$ . Since  $t$  is a factor of  $s$ , that letter  $i$  is also between the same letters in the word  $s$ , implying that  $i$  is also a vale (resp. pinnacle) of  $s$ . To prove condition 2, if  $\underline{i}\bar{j}$  or  $\bar{j}\underline{i}$  is a segment of  $t$  that does not contain any endpoint, then  $i$  and  $j$  are also spikes of  $s$ , by condition 1, which appear consecutively in  $\mathbf{pv}(s)$ . Therefore, it is also a segment of  $s$ .  $\square$

In other words, if we have a word  $s$  formed by consecutive integers and  $t$  a factor of  $s$ , then the only segments of  $t$  that may not be segments of  $s$  are its leftmost and rightmost ones. The reason is that endpoints are always considered spikes, and so the endpoints of  $t$  will be always spikes of  $t$ , but not necessarily spikes of  $s$ . Considering  $s$  and  $t$  as in Figure 2, the only segments of  $t$  that are segments of  $s$  are  $\bar{4}\underline{1}$ ,  $\underline{1}\bar{6}$  and  $\bar{6}\underline{3}$ . If we consider the word  $u = 123456543 \equiv \underline{1}\bar{6}\underline{3}$  which is also a factor of  $s$ , then every segment of  $u$  is a segment of  $s$ , because the endpoints of  $u$  are also spikes in  $s$ . The word  $v = 1234565434 \equiv \underline{1}\bar{6}\underline{3}\bar{4}$  is a factor of  $s$  where only one of its endpoints is a spike of  $s$ . The line diagrams of  $u$  and  $v$  are depicted in Figure 3.




Figure 3: Line diagrams of  $u \equiv \underline{1}\bar{6}\underline{3}$  and  $v \equiv \underline{1}\bar{6}\underline{3}\bar{4}$ .

### 3 A characterization of one-element commutation classes

As we saw in the previous section, the conditions stated in Proposition 4 are not enough to completely characterize one-element commutation classes, as there are words formed by consecutive integers satisfying all five conditions of the theorem which are not reduced. In this section, we give a necessary and sufficient condition for a word formed by consecutive integers to be reduced.

Given  $s = i_1 \cdots i_l \in [n]^*$ , define the words

- $s^r = i_l \cdots i_1$ , called the *reverse word* of  $s$ ,
- $s^c = i'_1 \cdots i'_l$  with  $i'_j = n + 1 - i_j$ , called the *complement word* of  $s$ ,
- $s^{rc} = (s^r)^c = (s^c)^r$ , called the *reverse complement word* of  $s$ .

These words are called the *symmetries* of  $s$ . It is easy to check that if  $s \in R_\bullet(\sigma)$ , then  $s^r \in R_\bullet(\sigma^{-1})$ ,  $s^c \in R_\bullet(w_0\sigma w_0)$  and  $s^{rc} \in R_\bullet(w_0\sigma^{-1}w_0)$ , since all permutations  $\sigma, \sigma^{-1}$  and  $w_0\sigma w_0$  have the same length. (see [1]). The following result is a well-known property about reduced words of permutations, which we will use in a moment.

**Lemma 6** ([1]). Let  $\sigma \in \mathfrak{S}_{n+1}$  and  $i \in [n]$ . Then

$$l(\sigma s_i) = \begin{cases} l(\sigma) + 1, & \text{if } \sigma(i) < \sigma(i+1) \\ l(\sigma) - 1, & \text{if } \sigma(i) > \sigma(i+1) \end{cases}.$$

As a consequence we have the following.

**Corollary 7.** Let  $s = t \cdot i \in R(\sigma)$  be a reduced word for some  $\sigma \in \mathfrak{S}_{n+1}$ , with  $t \in [n]^*$  and  $i \in [n]$ . Then,  $\sigma(i) > \sigma(i+1)$ .

**Proposition 8.** Let  $s = \underline{i}\bar{j} \ t \ \underline{i}\bar{j} \in [n]^*$  be a word formed by consecutive integers, with  $1 \leq i < j \leq n$  and  $t \in [n]^*$ . Then,  $s$  is not reduced.

*Proof.* Suppose that  $s$  is reduced, and let  $\sigma$  be its associated permutation so that  $s \in R_\bullet(\sigma)$ . We know that  $t$  must contain a spike of  $s$ , otherwise  $s \equiv \underline{i}\bar{j}\underline{i}\bar{j}$  which is not reduced by condition 3 of Proposition 4. Let  $\mathbf{v}(s) = i \ v_1 \cdots v_l \ i$  and  $\mathbf{p}(s) = j \ p_1 \cdots p_l \ j$  be the words of vales and pinnacles of  $s$ , respectively, for some integer  $l$ . From Proposition 4, since  $\mathbf{v}(s)$  is a vee (resp.  $\mathbf{p}(s)$  is a wedge) we have  $v_k < i$  (resp.  $p_k > j$ ), for all  $k \in [l]$ . This allow us to write  $s$  as

$$s = (\mathbf{i} \cdots \mathbf{j} \cdots \mathbf{i})(i-1 \cdots v_1 \cdots i-1) \cdots (\mathbf{i} \cdots \mathbf{p}_k \cdots \mathbf{i})(i+1 \cdots j).$$

(See Figure 4) The permutation associated to the bold left factor of  $s$  is

$$\pi = (i, j+1)(i, v_1) \cdots (i, p_k+1) = (i, p_k+1, \cdots, v_1, j+1).$$

Notice that the permutation associated to  $(i+1 \cdots j)$  moves the integer  $\pi(i+1)$  to position  $j+1$ . Since  $\pi(i+1) = i+1$  and  $\pi(j+1) = i$ , we have that  $\sigma(j) = i$  and  $\sigma(j+1) = i+1$ , which contradicts Corollary 7. Therefore,  $s$  cannot be reduced.  $\square$




Figure 4: Diagram of a word of the form  $\underline{i}\bar{j}t\underline{i}\bar{j}$

It follows from the previous proposition and the definition of reverse word that a word of the form  $\bar{j}\underline{i}t\bar{j}\underline{i}$ , with  $i < j$  that is formed by consecutive integers is also not reduced. These words contain at least two occurrences of the same segment, a fact that motivated the following definition.

**Definition 9.** A word formed by consecutive integers is said to have a factor with repeated segments if it contains a factor  $\underline{i}\bar{j}t\underline{i}\bar{j}$  or  $\bar{j}\underline{i}t\bar{j}\underline{i}$ , for some  $i < j$  and  $t \in [n]^*$ .

For example, the word  $s \equiv \overline{241635}$  has a factor with repeated segments since  $u = \overline{34t34}$ , with  $t = 3212345654$ , is a factor of  $s$ . As a consequence of Proposition 8, we have a new necessary condition for a word to be a one-element commutation class of some permutation.

**Theorem 10.** *Let  $\sigma \in \mathfrak{S}_{n+1}$  and  $s \in R_\bullet(\sigma)$ . Then  $s$  does not have a factor with repeated segments.*

*Proof.* If  $s$  has a factor with repeated segments, then it contains a factor  $\underline{i}\overline{j}t\underline{i}\overline{j}$  or  $\overline{j}\underline{i}t\overline{j}\underline{i}$  for some word  $t \in [n]^*$ , which by the previous proposition is not reduced.  $\square$

A natural question that one may ask is whether this new condition plus the ones stated in Proposition 4 are sufficient to completely characterize one-element commutation classes. As we are going to see, we just need this new one and condition 5 of Proposition 4 to complete this characterization. We will start by giving a criterion to identify words that contain factors with repeated segments just by looking at its line diagram.

**Proposition 11.** *Let  $s \in [n]^*$  be a word formed by consecutive integers. The following statements are equivalent:*

1. *There is no factor of  $s$  with repeated segments.*
2. *The word  $s$  does not contain a factor  $\underline{x}\overline{j}t\underline{i}\overline{y}$  with  $x \leq i < j \leq y$ , or a factor  $\overline{y}\underline{i}t\overline{j}\underline{x}$  with  $y \geq j > i \geq x$ , for some word  $t \in [n]^*$ , where  $x$  and  $y$  are spikes of  $s$ .*

*Proof.* We prove the contrapositive assertions. A word  $s$  contains a factor with repeated segments if and only if  $s$  contains a factor  $\underline{i}\overline{j}t\underline{i}\overline{j}$  or  $\overline{j}\underline{i}t\overline{j}\underline{i}$ . Consider the first case (the second is analogous). From Lemma 5, the only spikes of  $\underline{i}\overline{j}t\underline{i}\overline{j}$  that may not be spikes of  $s$  are its endpoints. Therefore  $s$  contains a factor  $\underline{x}\overline{j}t\underline{i}\overline{y}$  with  $x \leq i < j \leq y$ , with  $x$  and  $y$  spikes of  $s$ .

Reciprocally, suppose that  $s$  contains a factor  $\underline{x}\overline{j}t\underline{i}\overline{y}$  with  $x \leq i < j \leq y$  (the other case is analogous). Then,  $\underline{i}\overline{j}t\underline{i}\overline{j}$  is a factor of  $s$ , and we have the result.  $\square$

It follows that if  $s$  is a one-element commutation class, then its line diagram must avoid the two shapes depicted in Figure 5.




Figure 5: Avoiding shapes for one-element commutation classes.

The following definition will allow us to translate condition 5 of Proposition 4 into the language of segments.

**Definition 12.** A word formed by consecutive integers is said to have a factor with symmetric segments if it has a factor  $\underline{i}\bar{j}t\bar{j}\underline{i}$  or  $\bar{j}\underline{i}t\underline{i}\bar{j}$ , for some  $i < j$  where  $t \in [n]^*$  contains at least one spike of  $s$ .

For instance, the word  $s \equiv \underline{2}\bar{4}\underline{1}\bar{4}\underline{2}$ , depicted in Figure 6, has symmetric segments. The word  $s \equiv \bar{4}\underline{2}\bar{4}$  does not contain symmetric segments.




Figure 6: Diagram of the word  $s = \underline{2}\bar{4}\underline{1}\bar{4}\underline{2}$ .

**Proposition 13.** Let  $s \in [n]^*$  be a word formed by consecutive integers. The following statements are equivalent:

1. There is no factor of  $s$  with symmetric segments.
2. If  $\mathbf{p}(s)$  (or  $\mathbf{v}(s)$ ) has more than one integer  $i$ , then one of those  $i$ 's is an endpoint of  $s$ .

*Proof.* We prove the contrapositive assertions. If  $s$  contains a factor with symmetric segments, then  $s$  must contain a factor  $u = \underline{i}\bar{j}t\bar{j}\underline{i}$ , or  $u = \bar{j}\underline{i}t\underline{i}\bar{j}$ , with  $i, j \in [n]$  and  $t \in [n]^*$  a word which contains some spike of  $s$ . Considering the first case (the other is analogous), by Lemma 5 we have that both  $\bar{j}$  in  $u$  are pinnacles of  $s$ , and neither of them is an endpoint. Therefore, condition 2 does not hold.

Now suppose that  $\mathbf{v}(s)$  contains two letters  $i$  such that neither of them is an endpoint (the  $\mathbf{p}(s)$  case is analogous). Then  $s$  will contain a factor  $u$  of the form  $u = \bar{j}\underline{i}t\underline{i}\bar{k}$ , with  $j, k \in [n]$  and  $t$  a word which contains some spike of  $s$ . If  $j \leq k$ , then  $s$  will contain the factor  $\bar{j}\underline{i}t\underline{i}\bar{j}$ . If  $j > k$ , then  $s$  will contain the factor  $\bar{k}\underline{i}t\underline{i}\bar{k}$  (see Figure 7). In either case, we have a factor with symmetric segments.  $\square$




Figure 7: Diagram of the word  $u = \bar{j}\underline{i}t\underline{i}\bar{k}$  with  $j \leq k$  and  $j > k$ .

Given  $i < j$ , let  $[i, j] := \{i, i + 1, \dots, j\}$  and  $[i, j]^*$  the set of words with finite length in the alphabet  $[i, j]$ . The following lemma will be useful more ahead.

**Lemma 14.** Let  $s \in R(\sigma)$  a reduced word for some  $\sigma \in \mathfrak{S}_{n+1}$ . If  $s \in [i, j]^*$ , then  $\sigma(k) = k$  for all  $k \in [n + 1] \setminus [i, j + 1]$ .



*Proof.* Suppose that there is  $k \in [n+1] \setminus [i, j+1]$  such that  $\sigma(k) \neq k$ . Then, every reduced word for  $\sigma$  must contain a letter  $k$  or a letter  $k-1$ . But this is a contradiction because  $k$  and  $k-1$  belong to  $[n+1] \setminus [i, j+1]$  and  $s \in [i, j]^*$ .  $\square$

Before stating the main result of this section, we need a better understanding of how segments behave. Suppose that  $s = \underline{i}\bar{j}t \in R_\bullet(\sigma)$ , for some word  $t \in [n]^*$ . The segment  $\underline{i}\bar{j}$  moves the integer  $i$  to position  $j+1$ . If  $s$  contains another segment that moves the integer  $i$ , then the following segment to move this integer is of the form  $\underline{j}\bar{y}$ , with  $j < y$ , or  $\bar{j}\underline{x}$  with  $j > x$ . Notice that the second case cannot hold, otherwise we would have two pinnacles  $j$  in  $s$  where neither is an endpoint, contradicting condition 5 of Proposition 4. We can do a similar reasoning if  $s = \bar{j}\underline{i}t$  and concluding the following results:

**Lemma 15.** *Let  $s \in R_\bullet(\sigma)$  for some  $\sigma \in \mathfrak{S}_{n+1}$ . If a segment  $\underline{i}\bar{j}$  (resp.  $\bar{j}\underline{i}$ ) of  $s$  moves an integer  $k \in [n+1]$ , then every segment of  $s$  that moves  $k$  is of the form  $\underline{x}\bar{y}$  (resp.  $\bar{y}\underline{x}$ ).*

**Lemma 16.** *Let  $s \in R_\bullet(\sigma)$  for some  $\sigma \in \mathfrak{S}_{n+1}$  and  $i \in [n]$ .*

1. *If  $\sigma(i+1) < i$ , then there is a segment  $\underline{x}\bar{y}$  of  $s$  that moves the integer  $\sigma(i+1)$ .*
2. *If  $i+1 < \sigma(i)$ , then there is a segment  $\bar{y}\underline{x}$  of  $s$  that moves the integer  $\sigma(i)$ .*

There is also a restriction on the integers that are moved by segments.

**Lemma 17.** *Let  $s \in R_\bullet(\sigma)$  for some  $\sigma \in \mathfrak{S}_{n+1}$ . If  $\underline{i}\bar{j}$  (resp.  $\bar{j}\underline{i}$ ) is a segment of  $s$ , then it moves an integer  $k \in [i+1]$  (resp.  $k \in [j, n+1]$ ).*

*Proof.* Assume that  $s$  contains a segment  $\underline{i}\bar{j}$  (the other case is similar) which moves an integer  $k$ . Then, we can write  $s = t_1\underline{i}\bar{j}t_2$ , for some words  $t_1, t_2 \in [n]^*$ . Suppose by contradiction that  $i+1 < k$ . Since  $s$  is a reduced word, the word  $t_1$  is also a reduced word for some permutation  $\pi \in \mathfrak{S}_{n+1}$  with  $\pi(i) = k$  (because  $\underline{i}\bar{j}$  moves the integer  $k$ ). But  $i+1 < k$ , which from the previous lemma implies that  $t$  contains a segment  $\bar{y}\underline{x}$  that moves  $k$ , contradicting Lemma 15. Therefore  $k \leq i+1$ .  $\square$

**Proposition 18.** *Let  $s \in [n]^*$  be a word formed by consecutive integers such that  $s = t \cdot j$  with  $t \in R_\bullet(\sigma)$  for some  $\sigma \in \mathfrak{S}_{n+1}$  and  $j \in [n]$ . If  $s$  is not reduced, then  $s$  contains a factor with repeated or symmetric segments.*

*Proof.* Since  $s$  is not reduced, from Lemma 6 we have  $\sigma(j) > \sigma(j+1)$ . Assume that  $j$  is a pinnacle of  $s$  (the case when  $j$  is a vale follows from the application of complement word). Then, we can write  $s = u\underline{i}\bar{j}$  for some  $i < j$ , and we need to consider two cases.

**Case 1:**  $j > i+1$

If  $j > i+1$ , we have  $t = u\underline{i}\bar{j} - 1$  and from Lemma 17, since  $t$  is a one-element commutation class, the segment  $\underline{i}\bar{j} - 1$  moves an integer  $k \in [i+1]$  to position  $j$ . But then,  $\sigma(j) = k$  and we have  $j > i+1 \geq k > \sigma(j+1)$  which implies, from Lemma 16, that  $t$  contains

a segment  $\underline{x}\bar{y}$  that moves the integer  $\sigma(j+1)$  (call it  $l$ ), for some  $x < y$ . The rightmost such segment  $\underline{x}\bar{y}$  must have  $y = j$ , so we can write

$$t = u' \underline{x} \bar{j} u'' \bar{i} \bar{j} - \bar{1},$$

for some words  $u', u'' \in [n]^*$ . Our goal is to prove that  $x \leq i$ . If  $x > i$ , the fact that  $\mathbf{v}(t)$  is a vee implies that  $u'$  is a word in the alphabet  $[x+1, n]$ , because  $u'$  cannot contain vales of  $t$  that are smaller than or equal to  $x$ . Considering  $\pi$  the permutation associated to  $u'$ , from Lemma 14 we have  $\pi(x) = x$ . Since the segment  $\underline{x}\bar{j}$  moves the integer  $l$ , we have  $x = l$ . But this is a contradiction because  $x \geq i+1 > l$ . Therefore  $x \leq i$  and we have  $s = u' \underline{x} \bar{j} u'' \bar{i} \bar{j}$ , which, by Proposition 11, contains a factor with repeated segments (the factor  $\underline{x} \bar{j} u'' \bar{i} \bar{j}$ ).

**Case 2:  $j = i+1$**

In this case we have  $s = u \bar{i} \bar{i} + \bar{1}$ , with  $t = u \ i$ . If  $u$  does not contain any letter  $i$ , then  $u$  is a word in the alphabet  $[i+1, n]$ . But then, its associated permutation fixes  $i$ , so  $\sigma(i+1) = i$ . Since  $\sigma(i+1) > \sigma(i+2)$ , we have  $i+1 > i > \sigma(i+2)$ , which from Lemma 16 implies that  $t$  contains a segment  $\underline{x}\bar{y}$  that moves the integer  $\sigma(i+2)$ , for some  $x < y$ . The rightmost such segment  $\underline{x}\bar{y}$  must have  $y = i+1$ , and so we can write

$$s = u' \underline{x} \bar{i} + \bar{1} u'' \bar{i} \bar{i} + \bar{1},$$

for some words  $u', u'' \in [n]^*$ . Since  $x \leq i$ , we have that  $s$  contains a factor with repeated segments (the factor  $\underline{x} \bar{i} + \bar{1} u'' \bar{i} \bar{i} + \bar{1}$ ).

For the case where  $u$  contains letters  $i$ , we need to consider two sub-cases:

**Sub-case 1:  $u$  does not contain vales  $i$**

Our goal is to prove that  $i+1 > \sigma(i+2)$  in order to use the previous argument. If the first appearance of a letter  $i$  in  $t$  is preceded by a letter  $i-1$ , then  $\sigma(i+1) = i+1$ . If it's preceded by a letter  $i+1$ , then  $\sigma(i+1) = i$ . Since  $\sigma(i+1) > \sigma(i+2)$  we have the result.

**Sub-case 2:  $u$  contains vales  $i$**

We can write  $s = u' \bar{i} \bar{y} u'' \bar{i} \bar{i} + \bar{1}$ . If  $u'$  is not empty, then  $s$  contains two vales  $i$  where neither is an endpoint, which from Proposition 13 implies that  $s$  contains symmetric segments. If  $u'$  is empty, then  $t = \bar{i} \bar{y} u'' \bar{i}$ , which from the proof of Proposition 8 we have  $\sigma(i+1) = i+1$ . Using the same argument as in Sub-case 1, we have that  $s$  contains a factor with repeated segments.  $\square$

We are now in condition to state and prove the main result of this section.

**Theorem 19.** *Let  $s \in [n]^*$  be a word formed by consecutive integers and let  $\sigma \in \mathfrak{S}_{n+1}$  be the corresponding permutation. Then  $s \in R_{\bullet}(\sigma)$  if and only if there is no factor of  $s$  with repeated or symmetric segments.*

*Proof.* From Theorem 10, if  $s \in R_{\bullet}(\sigma)$ , then  $s$  cannot contain factors with repeated segments. Moreover,  $s$  must satisfy condition 5 of Proposition 4, which is equivalent to say that  $s$  does not have a factor with symmetric segments by Proposition 13. Reciprocally, suppose that  $s \notin R_{\bullet}(\sigma)$  for all  $\sigma \in \mathfrak{S}_{n+1}$  (we want to prove the contrapositive assertion).

Then  $s$  cannot be a reduced word and so it must contain a left factor  $s' = t \cdot j$  where  $t$  is reduced but  $s'$  is not, for some  $j \in [n]$ . From the previous proposition,  $s'$  contains a factor with repeated or symmetric segments and we have the result.  $\square$

## 4 One-element classes for involutions

In this section we give an explicit construction of one-element commutation classes for involutions and relate their existence with pattern avoidance problems. We start by recalling the following result proved in [5].

**Lemma 20.** *The word  $\underline{1}\bar{n}$  (resp.  $\bar{n}\underline{1}$ ) is the only reduced word formed by consecutive integers with length  $\geq n$  over the alphabet  $[n]$ , having left (resp. right) endpoint the letter 1 and right (resp. left) endpoint the letter  $n$ .*

In other words, there are no spikes between letters 1 and  $n$  in a one-element commutation class. As a consequence, we have the following.

**Lemma 21.** *Let  $\sigma$  be a permutation in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n+1$ . If  $|R_\bullet(\sigma)| > 0$ , then  $\sigma(n+1) = 1$  or  $\sigma(1) = n+1$ . Moreover, if  $\sigma(n+1) = 1$  (resp.  $\sigma(1) = n+1$ ) every one-element commutation class of  $\sigma$  contains a segment  $\underline{1}\bar{n}$  (resp.  $\bar{n}\underline{1}$ ).*

*Proof.* Since 1 and  $n+1$  are not fixed points of  $\sigma$ , every reduced word for  $\sigma$  must contain at least a letter 1 and a letter  $n$ . Let  $s \in R_\bullet(\sigma)$  and suppose that there is a letter 1 preceded by a letter  $n$  in  $s$ . By Lemma 20, we can write  $s$  as

$$s = u\underline{1}\bar{n}v,$$

where  $u$  is a word that does not contain letters 1 and  $v$  is a word that does not contain letters  $n$ . But then, the permutation associated to  $u$  fixes the integer 1, and so the segment  $\underline{1}\bar{n}$  will move the integer 1 to position  $n+1$ . Since the permutation associated to  $v$  fixes the integer  $n+1$ , we have  $\sigma(n+1) = 1$ . With analogous arguments one can prove that if there is a letter  $n$  preceded by a letter 1, then  $\sigma(1) = n+1$  and every one-element commutation class contains a segment  $\bar{n}\underline{1}$ .  $\square$

The previous lemma gives us a necessary condition for a permutation that does not fix 1 nor  $n+1$  to contain one-element commutation classes. It is not, however, a sufficient condition; consider for instance the permutation  $\sigma = [3, 4, 5, 2, 1] \in \mathfrak{S}_5$ , which does not fix 1 nor 5 and  $\sigma(5) = 1$ . One can check that this permutation contains 4 commutation classes and neither of them is a one-element commutation class. In the case of involutions, one can get more information. Before that, let's recall that the symmetries of a word  $s$  are the words  $s, s^r, s^c$  and  $s^{cr}$ .

**Lemma 22.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n+1$ . If  $|R_\bullet(\sigma)| > 0$ , then  $\sigma(n+1) = 1$  and  $\sigma(1) = n+1$ . Moreover, if  $s \in R_\bullet(\sigma)$ , then  $s$  is a symmetry of  $\underline{1}\bar{n}\underline{1}t$ , for some word  $t \in [2, n-1]^*$ .*

*Proof.* The equalities  $\sigma(1) = n + 1$  and  $\sigma(n + 1) = 1$  follows from the previous lemma and from the fact that  $\sigma$  is an involution. We have also from the previous lemma that, if  $s \in R_\bullet(\sigma)$ , then  $s$  contains a segment  $\underline{1}\bar{n}$  (because  $\sigma(n + 1) = 1$ ) and a segment  $\bar{n}\underline{1}$  (because  $\sigma(1) = n + 1$ ). The only possibility for  $s$  to contain those segments at the same time is to have the factor  $\underline{1}\bar{n}\underline{1}$  or  $\bar{n}\underline{1}\bar{n}$ , which must contain an endpoint, by Theorem 19.  $\square$

The previous two lemmas can be generalized for any permutation  $\sigma$  by replacing 1 and  $n + 1$  with the minimum and maximum non-fixed points of  $\sigma$ , respectively.

Notice that there is always an endpoint of a one-element commutation class of an involution that does not fix 1 nor  $n + 1$  that is the letter 1 or  $n$ . We have the following.

**Proposition 23.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$ . Then:*

1.  *$\sigma$  has a one-element commutation class that contains a letter 1 as endpoint if and only if  $\sigma = \prod_{k=1}^l (k, j_k + 1)$ , with  $j_1 = n > j_2 > \dots > j_l$  for some integer  $l$ , and  $k < j_k + 1$  for all  $k \in [l]$ .*
2.  *$\sigma$  has a one-element commutation class that contains a letter  $n$  as endpoint if and only if  $\sigma = \prod_{k=1}^l (i_k, n + 2 - k)$ , with  $i_1 = 1 < i_2 < \dots < i_l$  for some integer  $l$ , and  $i_k < n + 2 - k$  for all  $k \in [l]$ .*

*Proof.* We prove only statement 1 (the proof of 2 is analogous). Since  $\sigma$  is an involution, if there is a one-element commutation class of  $\sigma$  that contains a letter 1 as endpoint, there must be  $s \in R_\bullet(\sigma)$  such that  $s = \underline{1}\bar{n}\underline{1}t$ , for some word  $t \in [2, n - 1]^*$ . Note that the permutation associated to  $t$  (call it  $\pi$ ) is also an involution because the left factor  $\underline{1}\bar{n}\underline{1}$  is a reduced word for the cycle  $(1, n + 1)$  and  $\pi$  fixes 1 and  $n + 1$ . Since the left-endpoint of  $t$  is a letter 2, we must have  $t = \underline{2}\bar{j}\underline{2}'t'$  for some  $j < n$  and  $t' \in [3, j - 1]^*$ . Continuing this procedure, we conclude that

$$s = (1 \dots j_1 \dots 1)(2 \dots j_2 \dots 2) \dots (l \dots j_l \dots l), \quad (3)$$

with  $n = j_1 > \dots > j_l$ . Notice that  $l$  can be equal to  $j_l$  if the right endpoint of  $s$  is a pinnacle. In that case  $(l \dots j_l \dots l)$  would be just the letter  $l$ . Each factor  $(k \dots j_k \dots k)$  encodes the involution  $(k, j_k + 1)$ , so  $\sigma = \prod_{k=1}^l (k, j_k + 1)$ . Moreover,  $s$  does not contain any factor with repeated or symmetric segments (see Figure 8). Therefore, from Theorem 19,  $s$  is a reduced word of  $\sigma$ .

For the converse, just consider  $s$  as in (3), which is a reduced word for  $\sigma$  and is a word formed by consecutives integers.  $\square$

If  $s = \underline{1}\bar{n}\underline{1}t$ , then its associated permutation is completely determined by its pinnacles, as we saw in the previous proof. Hence we can conclude the following.




Figure 8: Possible diagrams for a one-element commutation class of an involution that fixes neither 1 nor  $n + 1$

**Lemma 24.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$ . Then, there is at most one word in  $R_{\bullet}(\sigma)$  of the form  $\underline{1}\bar{n}\underline{1}t$  (the same is true for its symmetries).*

Before stating one of the main results of this section, we need two auxiliar lemmas.

**Lemma 25.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$  and  $s \in R_{\bullet}(\sigma)$ . If  $\mathbf{p}(s)$  or  $\mathbf{v}(s)$  is a symmetry of the word  $1\ 2\ \cdots\ l$  for some  $l$ , then  $s^c \in R_{\bullet}(\sigma)$ .*

*Proof.* Suppose that  $s = \underline{1}\bar{n}\underline{1}t$  for some  $t \in [2, n - 1]^*$  (the other cases are analogous). Since  $\mathbf{v}(s)$  is not strict we must have that  $\mathbf{p}(s)$  is the word that belong to the symmetries of  $1\ 2\ \cdots\ l$  for some  $l$ . From the proof of the previous lemma  $\sigma = \prod_{k=1}^l (k, n + 2 - k)$ , and since  $w_0\sigma w_0 = \sigma$ , we have  $s^c \in R_{\bullet}(\sigma)$ .  $\square$

**Lemma 26.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$  and  $s \in R_{\bullet}(\sigma)$ . If  $s = s^r$ , then  $s \equiv \underline{1}\bar{n}\underline{1}$  or  $s \equiv \bar{n}\underline{1}\bar{n}$ .*

*Proof.* The assumption that  $s = s^r$  implies that its endpoints are equal. Therefore, both of its endpoints are the letter 1 or  $n$ , which from Lemma 24 we have the result.  $\square$

**Theorem 27.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$ . Suppose that  $s \in R_{\bullet}(\sigma)$ .*

1. *If  $n = 1$ , then  $R_{\bullet}(\sigma) = \{1\}$ .*
2. *If  $\sigma = (1, n + 1)$ , then  $R_{\bullet}(\sigma) = \{\underline{1}\bar{n}\underline{1}, \bar{n}\underline{1}\bar{n}\}$ .*
3. *If  $n > 1$  and neither  $\mathbf{p}(s)$  nor  $\mathbf{v}(s)$  is a symmetry of the word  $1\ 2\ \cdots\ l$  for some integer  $l$ , then  $R_{\bullet}(\sigma) = \{s, s^r\}$ .*
4. *If  $n > 1$ ,  $\sigma \neq (1, n + 1)$  and one of the words  $\mathbf{p}(s)$  or  $\mathbf{v}(s)$  is a symmetry of the word  $1\ 2\ \cdots\ l$ , then  $R_{\bullet}(\sigma) = \{s, s^r, s^c, s^{cr}\}$*

*Proof.* Condition 1 is trivial. From Lemma 24, we have at most one word  $s \in R_{\bullet}(\sigma)$  with left (resp. right) endpoint the letter 1 or with left (resp. right) endpoint the letter  $n$ . If  $\sigma = (1, n + 1)$ , then  $\underline{1}\bar{n}\underline{1}$  and  $\bar{n}\underline{1}\bar{n}$  are one-element commutation classes and they must be the only ones. To prove conditions 3 and 4, we are going to assume that  $s = \underline{1}\bar{n}\underline{1}t$  (the other cases are analogous). We need to consider two cases:

**Case 1:**  $\mathbf{p}(s)$  is not a symmetry of  $1\ 2\ \cdots\ l$

From Lemma 23 we have  $\sigma = \prod_{k=1}^l (k, j_k + 1)$ , where  $j_k$  is the  $k$ -th pinnacle of  $s$  and  $k < j_k$ .

Our goal is to show that  $\sigma$  cannot contain one-element commutation classes with endpoint the letter  $n$ . If it contains such class, from Lemma 23 we have  $\sigma = \prod_{k=1}^l (i_k, n+2-k)$ . Since  $i_k < n+2-k$  for all  $k \in [l]$ , and since  $\sigma$  has a unique decomposition into disjoint cycles, we must have  $\mathbf{p}(s) = j_1 j_2 \cdots j_l = n \ n \cdots n + 1 - l$ , which is a symmetry of the word  $1 \ 2 \cdots l$ , contradicting our assumption. Since  $s^r \in R_\bullet(\sigma)$ , we have  $R_\bullet(\sigma) = \{s, s^r\}$ .

**Case 2:**  $\mathbf{p}(s)$  is a symmetry of  $1 \ 2 \cdots l$

From Lemma 25 we have  $s^c \in R_\bullet(\sigma)$ , and since  $\sigma$  is an involution,  $s^{cr} \in R_\bullet(\sigma)$ . The fact that  $\sigma \neq (1, n+1)$  implies that  $s \neq s^r$ . Therefore, we have  $R_\bullet(\sigma) = \{s, s^r, s^c, s^{cr}\}$ .  $\square$

For an arbitrary involution  $\sigma \in \mathfrak{S}_{n+1}$ , the previous theorem can be generalized by defining the complementary word of  $s = i_1 i_2 \cdots i_l$  as

$$(M + m - 1 - i_1)(M + m - 1 - i_2) \cdots (M + m - 1 - i_l),$$

where  $m$  and  $M$  are the minimum and maximum non-fixed points of  $\sigma$ , respectively.

We end this section with a relation between involutions that contain one-element commutation classes and pattern avoidance problems.

**Definition 28.** Let  $\sigma \in \mathfrak{S}_{n+1}$  and  $p \in \mathfrak{S}_k$  with  $k \leq n+1$ . We say that  $\sigma$  *contains* the pattern  $p$  if there is a subword of the one-line notation of  $\sigma$  order isomorphic to  $p$ . If not, we say that  $\sigma$  is *p-avoiding*.

When writing permutation as patterns, we drop the brackets and commas. For instance, the permutation  $[4, 1, 2, 5, 3]$  is 321-avoiding, but contains two patterns 123, namely the subwords  $1 \ 2 \ 5$  and  $1 \ 2 \ 3$ .

**Proposition 29.** Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n+1$ . If  $\sigma$  is 132 and 3412-avoiding, then  $\sigma(1) = n+1$  and  $\sigma(n+1) = 1$ .

*Proof.* Assume by contradiction that  $\sigma(1) \neq n+1$  and  $\sigma(1) \neq 1$ . Since  $\sigma$  does not fix 1 nor  $n+1$ , we must have

$$\sigma = [i, \dots, 1, \dots, n+1, \dots, j]$$

or

$$\sigma = [i, \dots, n+1, \dots, 1, \dots, j],$$

for some integers  $i, j \in [2, n]$ . The first case cannot hold because we have the subword  $1 \ n+1 \ j$ , which is a 132-pattern. In the second case, if  $i < j$ , then  $i \ n+1 \ j$  is a 132-pattern. If  $i > j$ , then  $\sigma$  contains the subword  $i \ n+1 \ 1 \ j$ , which is a 3412-pattern. Therefore, we must have  $\sigma(1) = n+1$  or  $\sigma(n+1) = 1$ . The fact that  $\sigma$  is an involution implies  $\sigma(1) = n+1$  and  $\sigma(n+1) = 1$ .  $\square$

Before stating the main result of this section, we need to recall two known results about pattern avoidance.

**Proposition 30** ([10]). A permutation  $\sigma$  is 2143 and 3412-avoiding if and only if  $\sigma$  can be partitioned into an increasing and decreasing sequence.

For instance, the permutation  $\sigma = [3, 6, 4, 7, 5, 2, 1]$  is 2143 and 3412-avoiding because we can partitioned  $\sigma$  into the sequences 3 4 7 and 6 5 2 1.

**Proposition 31** ([9]). *Let  $\sigma \in \mathfrak{S}_{n+1}$  and  $p \in \mathfrak{S}_k$  with  $k \leq n + 1$ . Then  $\sigma$  contains the pattern  $p$  if and only if  $w_0\sigma w_0$  contains the patterns  $w_0pw_0$ .*

We have the following.

**Theorem 32.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n + 1$ . Then,  $|R_\bullet(\sigma)| > 0$  if and only if  $\sigma$  avoids the patterns 132 and 3412 or the patterns 213 and 3412.*

*Proof.* Assume that  $|R_\bullet(\sigma)| > 0$ . Using Lemma 23, we start by consider  $\sigma = \prod_{k=1}^l (k, j_k + 1)$ , with  $j_1 = n > j_2 > \dots > j_l$  for some integer  $l$ . To prove that  $\sigma$  is 3412-avoiding, just notice that the non-fixed points forms a decreasing sequence and the fixed points forms an increasing sequence. Then, by Proposition 30 we have the result. It remains to prove that  $\sigma$  is 132-avoiding. By way of contradiction, assume that  $\sigma$  contains a pattern 132. Then  $\sigma$  contains a subword  $xzy$  such that  $x < y$ ,  $x < z$  and  $z > y$ . We then have two cases.

**Case 1:**  $x$  is not a fixed point.

Since the sequence of non-fixed points of  $\sigma$  is decreasing, we have that  $y$  and  $z$  are fixed points. But that cannot happen because  $z > y$  and the sequence of fixed points is increasing.

**Case 2:**  $x$  is a fixed point.

If  $x$  is a fixed point of  $\sigma$ , then  $x > l$ . We have that all non-fixed points of  $\sigma$  that are to the right of  $x$  are smaller than  $x$ . Since  $x < y$  and  $x < z$ , the integers  $y$  and  $z$  are also fixed points of  $\sigma$ . But that cannot be possible because  $z > y$ . Therefore,  $\sigma$  must be 132-avoiding.

Now considering the case where  $\sigma = \prod_{k=1}^l (i_k, n + 2 - k)$ , with  $i_1 = 1 < i_2 < \dots < i_l$  for some integer  $l$ , we have that  $w_0\sigma w_0 = \prod_{k=1}^l (k, n + 2 - i_k)$ , which we already proved is 132 and 3412-avoiding. From Proposition 31,  $\sigma$  is 213 and 3412-avoiding.

For the converse, suppose that  $\sigma$  avoids the patterns 132 and 3412. From Lemma 29, we have  $\sigma(1) = n + 1$  and  $\sigma(n + 1) = 1$ , so  $\sigma$  contains the cycle  $(1, n + 1)$  in its disjoint cycle decomposition. Without loss of generality, the disjoint decomposition of  $\sigma$  can be written as

$$\sigma = (i_1, j_1)(i_2, j_2) \cdots (i_l, j_l),$$

with  $1 = i_1 < i_2 < i_3 < \dots < i_l$ ,  $j_1 = n + 1$  and  $i_k < j_k$  for all  $k \in [l]$ . Notice that  $i_2 = 2$ , otherwise  $2 < i_2 < j_2$  and 2 would be a fixed point of  $\sigma$  implying that  $2 j_2 i_2$  would be a subword of  $\sigma$ , which is a 132-pattern. We also have that  $n + 1 = j_1 > j_2$ . Now suppose that

$$\sigma = (1 \ n + 1)(2 \ j_2) \cdots (m \ j_m)(i_{m+1} \ j_{m+1}) \cdots (i_l, j_l),$$

with  $n + 1 > j_2 > j_3 > \dots > j_m$  and  $i_k < j_k$  for all  $k \in [m + 1, l]$ . Our goal is to show that  $i_{m+1} = m + 1$  and  $j_m > j_{m+1}$ . If  $i_{m+1} \neq m + 1$ , then  $m + 1 < i_{m+1} < j_{m+1}$  and we have two cases. If  $m + 1$  is a fixed point, then  $\sigma$  will contain the subword  $m + 1 \ j_{m+1} \ i_{m+1}$ ,

which is a 132-pattern. If  $m+1$  is not a fixed point, then  $(m, j_m) = (m, m+1)$ . But then,  $\sigma$  will contain the subword  $m j_{m+1} i_{m+1}$ , which is a 132-pattern. Therefore  $i_{m+1} = m+1$ . If  $j_m < j_{m+1}$ , then  $\sigma$  will contain the subword  $j_m j_{m+1} m m+1$ , which is a 3412-pattern. Using an inductive argument we show that  $\sigma = (1, j_1)(2, j_2) \cdots (l, j_l)$ , with  $n+1 = j_1 > j_2 > \cdots > j_l$ . From Lemma 23, we have that  $\sigma$  contains a one-element commutation class with some endpoint the letter 1. If  $\sigma$  avoids the patterns 213 and 3412, then  $w_0 \sigma w_0$  avoids the patterns 132 and 3412 from Proposition 31, which we proved to have a one-element commutation class  $s$ . Hence  $s^c$  is a one-element commutation class for  $\sigma$ , proving that  $R_\bullet(\sigma) > 0$ .  $\square$

Notice that this result is not true in general for any permutation that does not fix 1 nor  $n+1$ . For instance, the permutation  $[3, 4, 5, 2, 1]$  is 132, 213 and 3412-avoiding and does not contain one-element commutation classes.

As a corollary, we have a necessary and sufficient condition for an involution to contain 4 one-element commutation classes.

**Corollary 33.** *Let  $\sigma$  be an involution in  $\mathfrak{S}_{n+1}$  that fixes neither 1 nor  $n+1$ . Assume that  $\sigma \neq (1, n+1)$ . The following are equivalent:*

1.  $|R_\bullet(\sigma)| = 4$ .
2.  $\sigma$  is 132, 213 and 3412-avoiding.

*Proof.* From Theorem 27, if  $|R_\bullet(\sigma)| = 4$ , then  $R_\bullet(\sigma) = \{s, s^r, s^c, s^{cr}\}$  for some word  $s \in [n]^*$  formed by consecutive integers. We can assume that  $s = \underline{1}\bar{n}\underline{1}t$  for some word  $t \in [2, n-1]^*$ . From the proof of the previous theorem, its associated permutation is 132 and 3412 avoiding. Since  $s^c \in R_\bullet(\sigma)$ , we have  $\sigma = w_0 \sigma w_0$  and from Proposition 31,  $\sigma$  will also avoid the pattern 213. Reciprocally, since  $\sigma$  avoids the patterns 132 and 3412, from the proof of the previous theorem we have that  $\sigma$  contains a one-element commutation class  $s$  with some endpoint the letter 1. We also have that  $\sigma$  is 213 and 3412-avoiding, so  $\sigma$  contains a one-element commutation class  $t$  with some endpoint the letter  $n$ . Since  $\sigma \neq (1, n+1)$ , from Lemma 24 we have  $R_\bullet(\sigma) = \{s, s^r, t, t^r\}$ .  $\square$

Corollary 33 allow us to recover the result from [5] and [14] which states that for  $n > 1$  the longest permutation  $w_0$  contains 4 one-element commutation classes, since  $w_0$  is an involution that fix neither 1 nor  $n+1$  and the only patterns of length 3 and 4 that  $w_0$  contains are 321 and 4321, respectively. One can generalize the previous results to any involution.

**Corollary 34.** *Let  $\sigma \in \mathfrak{S}_{n+1}$  an involution such that  $m$  and  $M$  are the minimum and maximum non-fixed points of  $\sigma$ . Consider the permutation  $\pi = [\sigma(m), \sigma(m+1), \dots, \sigma(M)]$ . Then:*

1.  $|R_\bullet(\sigma)| > 0$  if and only if  $\pi$  avoids the patterns 132 and 3412 or the patterns 213 and 3412.
2. Suppose that  $\sigma \neq (m, M)$ . Then  $|R_\bullet(\sigma)| = 4$  if and only if  $\pi$  avoids the patterns 132, 213 and 3412.



## 5 Acknowledgements

This work was partially supported by the Centre for Mathematics of the University of Coimbra (funded by the Portuguese Government through FCT/MCTES, DOI 10.54499/UIDB/00324/2020).

## References

- [1] A. Björner and F. Brenti. *Combinatorics of Coxeter Groups*. Springer, 2005. doi:10.1007/3-540-27596-7.
- [2] R. Davis, S. Nelson, T. Petersen and B. Tenner, The pinnacle set of a permutation, *Discrete Mathematics*, 341(11), 3249–3270, 2018. doi:10.1016/j.disc.2018.08.011.
- [3] S. Elnitsky. Rhombic tilings of polygons and classes of reduced words in Coxeter groups. *Journal of Combinatorial Theory, Series A*, 77(2):193–221, 1997. doi:10.1006/jcta.1997.2723.
- [4] N. González, P. Harris, G. Kirby, M. Garcia and B. Tenner. Pinnacle sets of signed permutations. *Discrete Mathematics*, 346(7): 113–439, 2023. doi:10.1016/j.disc.2023.113439.
- [5] G. Gutierrez, R. Mamede, and J. Santos. Commutation classes of the reduced words for the longest element of  $\mathfrak{S}_n$ . *The Electronic Journal of Combinatorics*, 27(2):#P2.21, 2020. doi:10.37236/9481
- [6] G. Gutierrez, R. Mamede and J. Santos. Diameter of the commutation classes graph of a permutation. *European Journal of Combinatorics*, 103, 2022. doi:10.1016/j.ejc.2022.103525.
- [7] D. Knuth. Axioms and hulls. In *Lecture Notes in Computer Science*, 1992. doi:10.1007/3-540-55611-7.
- [8] R. Mamede, J. Santos, D. Soares. The commutation graph for the longest signed permutation. *Discrete Mathematics*, 345(11):113055, 2022. doi:10.1016/j.disc.2022.113055 .
- [9] F. Negassi, L. Pudwell, C. Cratty and S. Erickson. Pattern avoidance in double lists. *Involve, a Journal of Mathematics*, 10:379–398, 2017. doi:10.2140/involve.2017.10.379.
- [10] Z. Stankova. Forbidden subsequences. *Discrete Mathematics*, 132(1):291–316, 1994. doi:10.1016/0012-365X(94)90242-9.
- [11] R. Stanley. On the number of reduced decompositions of elements of Coxeter groups. *European Journal of Combinatorics*, 5(4):359–372, 1984. doi:10.1016/S0195-6698(84)80039-6.
- [12] B. Tenner. Reduced decompositions and permutation patterns. *Journal of Algebraic Combinatorics*, 24:263–284, 2005. doi:10.1007/s10801-006-0015-6.

- [13] B. Tenner. Reduced word manipulation: patterns and enumeration. *Journal of Algebraic Combinatorics*, 46:189–217, 2015. [doi:10.1007/s10801-017-0752-8](https://doi.org/10.1007/s10801-017-0752-8).
- [14] B. Tenner. One-element commutation classes. *Journal of Combinatorics*, 15(3): 401–408, 2024. [doi:10.4310/JOC.240907014318](https://doi.org/10.4310/JOC.240907014318).
- [15] J. Tits. Le problème des mots dans les groupes de coxeter. *In: Symposia Mathematica*, pages 175–185, 1969. [doi:10.1016/B978-1-4832-2995-9.50013-1](https://doi.org/10.1016/B978-1-4832-2995-9.50013-1).