

The Latin Tableau Conjecture

Timothy Y. Chow^a Mark G. Tiefenbruck^b

Submitted: Jul 31, 2024; Accepted: Apr 5, 2025; Published: Jun 20, 2025

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

A *Latin tableau of shape λ and type μ* is a Young diagram of shape λ in which each box contains a single positive integer, with no repeated integers in any row or column, and the i th most common integer appearing μ_i times. Over twenty years ago, Chow et al., in their study of a generalization of Rota's basis conjecture that they called the wide partition conjecture, conjectured a necessary and sufficient condition for the existence of a Latin tableau of shape λ and type μ . We report some computational evidence for this conjecture, and prove that the conjecture correctly characterizes, for any given λ , at least the first four parts of μ .

Mathematics Subject Classifications: 05C15, 05A17, 05B30

1 Introduction

Over twenty years ago, Chow et al. [3] proposed the *wide partition conjecture*, which generalized Gian-Carlo Rota's famous *basis conjecture* [10]. They hoped that enlarging the context of the basis conjecture would suggest new lines of attack (e.g., proofs by induction on the number of boxes in the Young diagram of a partition). Unfortunately, even the special case of the wide partition conjecture that Chow et al. called the *wide partition conjecture for free matroids* has shown itself to be difficult, and remains open to this day. The only significant progress we are aware of is a recent preprint [1] that formulates and proves a dual version of the wide partition conjecture for free matroids.

Chow et al. mentioned in passing something they called the *Latin Tableau Question*; a positive answer to the question would imply the wide partition conjecture for free matroids. We believe that the Latin Tableau Question, in addition to having interesting connections with Rota's basis conjecture, is interesting in its own right. In the present paper, we give a self-contained presentation of the Latin Tableau Question (that requires no knowledge of Rota's basis conjecture or the wide partition conjecture), and we prove some partial results.

^aCenter for Communications Research—Princeton, NJ (tchow@alum.mit.edu).

^bCenter for Communications Research—La Jolla, CA (mgtiefe@ccr-lajolla.org).

We begin by defining a *Latin tableau*. A *partition* λ is a weakly decreasing finite sequence $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell$ of positive integers. The individual λ_i are called the *parts* of the partition, and the sum of the parts is denoted $|\lambda|$. The *Young diagram* or simply *diagram* of a partition λ is a left-justified grid of boxes whose i th row contains λ_i boxes. If we color each box of the diagram of λ in such a way that there are no repeated colors in any row or column, then we call the colored diagram T a *Latin tableau of shape* λ . The *type* of T is the partition μ where μ_1 is the number of times that the most frequently occurring color appears, μ_2 is the number of times that the next most frequently occurring color appears, and so on. See Figure 1 for an example.




Figure 1: Latin tableau of shape $(4, 2, 1)$ and type $(2, 2, 2, 1)$

The word *Latin* is inspired by the well-known notion of a *Latin square*, but we caution the reader that even if λ is a square shape, a Latin tableau of shape λ is not required to have $\text{type}(\mu) = \lambda$, and hence is not necessarily a Latin square in the usual sense. We can now state the following natural question.

Given λ and μ , what are necessary and sufficient conditions for there to exist a Latin tableau of shape λ and type μ ?

A conjectural answer to the above question was proposed in [3, Section 6]. To state this *Latin Tableau Conjecture*, we need a few more definitions.

Given a partition λ and a nonnegative integer n , we let α_n denote the maximum number of boxes of the diagram of λ that we can color using n colors without repeating a color in any row or column. (Technically, we should write $\alpha_n(\lambda)$ to emphasize the dependence of α_n on λ , but to avoid cluttering the notation, we just write α_n when there is no danger of confusion.) We define the *chromatic difference sequence* $\delta = (\delta_1, \delta_2, \dots)$ by letting

$$\delta_n := \alpha_n - \alpha_{n-1}$$

for each positive integer n . It is clear from the definitions that if there exists a Latin tableau of shape λ and type μ , then necessarily

$$\begin{aligned} \delta_1 &= \alpha_1 \geq \mu_1 \\ \delta_2 + \delta_1 &= \alpha_2 \geq \mu_1 + \mu_2 \\ \delta_3 + \delta_2 + \delta_1 &= \alpha_3 \geq \mu_1 + \mu_2 + \mu_3 \\ &\vdots \end{aligned}$$

i.e., δ *dominates* or *majorizes* μ (written $\delta \succeq \mu$). The Latin Tableau Conjecture asserts that this necessary condition is sufficient.

Latin Tableau Conjecture. A Latin tableau of shape λ and type μ exists if and only if $\delta \succeq \mu$.

In fact, Proposition 5 below reduces the conjecture to showing that there always exists a Latin tableau of shape λ and type δ .

More than once, when we have shown the Latin Tableau Conjecture to someone, the reaction has been that such a natural statement must be either known or false. We are confident that it is not known; Chow et al. [3] surveyed a lot of closely related literature, and in the present paper, we establish further connections with the literature on matching theory (see the proof of Theorem 14) and chromatic difference sequences. If anything, the prior work in these related areas makes the Latin Tableau Conjecture seem unexpected, since even slight generalizations are known to be false (see the discussion of Figure 3 below).

The conjecture could of course be false, but we have checked that it is true for all partitions λ that fit inside a 12×12 square. Moreover, the main result of this paper is that there always exists a Latin tableau of shape λ and type μ such that $\mu_i = \delta_i$ for $1 \leq i \leq 4$. There is some hope that our methods might generalize to prove the full conjecture, but we currently rely on case analyses that become increasingly complicated for higher terms of the CDS.

2 Graph-Theoretic Perspective

In this section, we describe two different ways to think of a Young diagram as a graph. Each way has its own merits.

All our graphs $G = \{V, E\}$ will be finite, simple, and undirected. A *coloring* of G is a map $\kappa: V \rightarrow \mathbb{N}$ such that $\kappa(u) \neq \kappa(v)$ whenever u and v are adjacent. (In the literature, the term *proper coloring* is sometimes used, but we will have no occasion to refer to any other type of coloring, so we drop the word *proper*.) A *stable set* or *independent set* of G is a subset $S \subseteq V$ such that whenever $u \in S$ and $v \in S$ then there is no edge between u and v . Observe that each *color class* of a coloring κ (meaning the inverse image of some $n \in \mathbb{N}$) is a stable set. Given a coloring κ , we define $\text{shape}(\kappa)$ to be the sequence of cardinalities of the color classes of κ , arranged in weakly decreasing order.

A very important concept for us is the chromatic difference sequence of a graph [2].

Definition 1. Given a graph G , we let $\alpha_i(G)$ (or simply α_i , if the graph under consideration is understood) denote the maximum cardinality of a union of i (disjoint) stable sets of G . The *chromatic difference sequence* or *CDS* of G is the sequence $(\delta_1, \delta_2, \delta_3, \dots)$ defined by

$$\delta_i := \alpha_i - \alpha_{i-1}, \quad \text{for } i \geq 1.$$

We say that G is *CDS-colorable* if there exists a coloring κ of G such that $\text{shape}(\kappa)$ equals the CDS of G .

The first of our two ways of associating a graph with a partition is given by the following definition, which establishes a connection between the above definition of a CDS and the definition given in the Introduction.

Definition 2. Given a partition λ , the *partition graph* G_λ is the graph whose vertices are the boxes of the diagram of λ , and in which two boxes are adjacent if and only if they lie in the same row or column.

The CDS of an arbitrary graph is not necessarily weakly decreasing (see for example [5, Remark 2] or [2, Figure 1]). However, we have the following result.

Proposition 3. *The CDS of a partition graph is itself a partition; i.e., it is weakly decreasing.*

Proof. This was proved by de Werra [5, Lemma 2.1], who actually showed that the result holds for the line graph of any bipartite graph. \square

In particular, the CDS of G_λ coincides with the CDS of λ as defined in the Introduction, and the Latin Tableau Conjecture implies (and in fact, by Proposition 5 below, is equivalent to) the CDS-colorability of G_λ for all partitions λ .

It is well known that not every graph is CDS-colorable; indeed, despite the considerable literature on chromatic difference sequences, very few graphs are known to be CDS-colorable. In Figure 2, we regard the picture on the left as a graph whose vertices are the boxes and in which two boxes are adjacent if and only if they lie in the same row or column. The other two pictures in Figure 2 demonstrate that $\alpha_1 = 5$ and $\alpha_2 = 4 + 4 = 8$. So the CDS is $(5, 3, 1)$, but it is easy to check that there is no coloring κ such that $\text{shape}(\kappa) = (5, 3, 1)$. A slightly larger graph (Figure 3, reproduced from [3]) shows that that Latin Tableau Conjecture cannot be generalized even to skew Young diagrams; the CDS is $(6, 6, 3, 1)$ but there is no coloring κ such that $\text{shape}(\kappa) = (6, 6, 3, 1)$.




Figure 2: A non-CDS-colorable graph

There is a second way to think of a Young diagram as a graph.

Definition 4. Given a partition λ , we let B_λ be the bipartite graph whose vertices are the rows and columns of the diagram of λ , and in which row i is adjacent to column j if and only if the diagram of λ contains a box in row i and column j .

The bipartite graph B_λ is a useful tool for proving the following Proposition, which, together with Proposition 3, implies that if we can prove the Latin Tableau Conjecture in the case that $\mu = \delta$, then the full conjecture follows.




Figure 3: A non-CDS-colorable skew shape

Proposition 5. *If there exists a Latin tableau with shape λ and type μ , and $\mu \succeq \nu$, then there exists a Latin tableau with shape λ and type ν .*

Proof. We need the following fact about the dominance order on partitions [8, Chapter I, (1.16)]. If $\mu \succeq \nu$, then we can transform μ to ν by performing a (carefully chosen) sequence of operations of the following type: take two parts of the current partition that differ in size—call them a and b , with $a < b$ —and replace them with $a + 1$ and $b - 1$ (reordering the parts into weakly decreasing order if necessary) to get a new partition. Given this fact, it suffices to show that if T is a Latin tableau with shape λ and type μ , and colors i and j occur with multiplicities μ_i and μ_j respectively, where $\mu_i < \mu_j$, then we can recolor those boxes in such a way that color i occurs with multiplicity $\mu_i + 1$ and color j occurs with multiplicity $\mu_j - 1$.

Now consider the bipartite graph B_λ , and identify the coloring of the boxes of T with an edge coloring of B_λ . In B_λ , the edges with color i form a *matching* (i.e., a set of edges, no two of which touch each other), and similarly the edges with color j form a matching that is disjoint from the first. The union of these two matchings form a subgraph of B_λ in which every vertex has degree 1 or 2, and hence each connected component must be a cycle or a path with edges of alternating colors. Since $\mu_i < \mu_j$, at least one of these connected components must be a path with an odd number of edges and with color j edges in the majority. Now we can simply reverse the colors of the edges in this path, and observe that this decreases the number of color j edges by 1, increases the number of color i edges by 1, and preserves the Latin tableau property. \square

Corollary 6. *Assume that for every λ , there exists a Latin tableau of shape λ and type δ . Then the Latin Tableau Conjecture follows.*

3 The CDS of a Partition Graph

It will be convenient to coordinatize a Young diagram; box (i, j) denotes the box in row i and column j , where we number the rows and columns starting with 0 (i.e., the top left box is $(0, 0)$).

Definition 7. For $k \geq 0$, *antidiagonal k* is the set of boxes $\{(0, k), (1, k - 1), (2, k - 2), \dots, (k, 0)\}$ whose coordinates sum to k .

The portion of a Young diagram lying (weakly) below and to the right of a box (i, j) plays an important role, so we introduce the following notation for it.

Definition 8. For a partition λ and nonnegative integers i and j , we define

$$\lambda[\overline{i, j}] := \{(i', j') \in \lambda \mid i' \geq i \text{ and } j' \geq j\},$$

where “ $(i', j') \in \lambda$ ” means that (i', j') are the coordinates of a box in the diagram of λ . The notation $\lambda[\overline{i, j}]$ is pronounced, “ λ corner i, j .”

The following Lemma states a simple but important fact about $\lambda[\overline{i, j}]$.

Lemma 9. A stable set in a partition graph G_λ contains at most $i + j$ boxes outside $\lambda[\overline{i, j}]$.

Proof. The portion of the diagram of λ outside $\lambda[\overline{i, j}]$ is contained in the union of i rows and j columns. But at most i elements of the stable set can lie in i rows and at most j elements of the stable set can lie in j columns. \square

Lemma 9 implies a useful characterization of δ_1 , the first term of the CDS.

Proposition 10. Let k be a positive integer. Then the CDS of a partition graph G_λ begins with k if and only if

1. the diagram of λ contains the entirety of antidiagonal $k - 1$, and
2. the diagram of λ omits some box from antidiagonal k .

Proof. Suppose that conditions 1 and 2 hold. Condition 1 implies that there exists a stable set with k boxes—simply take antidiagonal $k - 1$. Let (i, j) be the coordinates of a missing box in antidiagonal k (whose existence is guaranteed by condition 2). Then Lemma 9 tells us that any stable set contains at most k boxes outside $\lambda[\overline{i, j}]$, and since all of λ lies outside $\lambda[\overline{i, j}]$, this means that the maximum size of a stable set is k . That is, the CDS begins with k .

Conversely, suppose that the CDS begins with k . If there were some box in antidiagonal $k - 1$ missing from the diagram of λ , then a similar argument to the one in the previous paragraph would imply that a stable set could have size at most $k - 1$, which is a contradiction. So Condition 1 holds. Condition 2 must also hold, because if the diagram of λ contained the entirety of antidiagonal k , then there would be a stable set of size at least $k + 1$, again a contradiction. \square

We use the term *main antidiagonal* to refer to antidiagonal $\delta_1 - 1$. Proposition 10 tells us a lot about δ_2 as well. If $\delta_1 = k$, then in the worst case, we can color antidiagonal $k - 1$ one color and antidiagonal $k - 2$ another color. Therefore, $\delta_2 \geq k - 1$. A slightly more careful argument yields the following result.

Proposition 11. If δ is the CDS of a partition graph G_λ and $\delta_1 = k$, then $\delta_r \geq k - r + 1$ for $1 \leq r \leq k$.

Proof. Suppose that α_{r-1} boxes of the diagram of λ have been colored using $r-1$ colors. It suffices to show that at least $k-r+1$ uncolored boxes can be colored using an r th color. Proposition 10 tells us that the diagram of λ contains the entirety of antidiagonal $k-1$. That is, for $0 \leq i \leq k-1$, the diagram of λ contains at least $k-i$ boxes in row i . In particular, in row $k-r$, there are at least r boxes, and therefore at least one uncolored box, since only $r-1$ colors have been used so far. Let us color an uncolored box in row $k-r$ with color r . By the same kind of reasoning, there are at least two uncolored boxes in row $k-r-1$, and at least one of these can be colored with color r , because at most one of them is rendered unavailable by the box in row $k-r$ that we colored with color r . Continuing in this way with rows $k-r-2$, $k-r-3$, etc., we find that we can color one box in each of the rows $0, 1, \dots, k-r$ with color r , for a total of $k-r+1$ boxes. This is what we set out to prove. \square

Combining Propositions 3 and 11, we see that if we fix $\delta_1 := k$, then there are at most five possibilities for the first three terms of a CDS of a partition graph:

$$(k, k, k), \quad (k, k, k-1), \quad (k, k, k-2), \quad (k, k-1, k-1), \quad (k, k-1, k-2).$$

Similarly, there are at most C_r possibilities for the first r terms, where C_r is the r th Catalan number. It turns out, however, that not all of these possibilities are realizable. But before proving that, we need to develop a bit of general theory.

3.1 Corner Constraints

Lemma 9 can be used to derive what we call *corner constraints* on the CDS of a partition graph. First, we need a definition.

Definition 12. Let λ and μ be partitions, and let d be a nonnegative integer. Then λ *includes* μ at d if there exist nonnegative integers i and j such that

1. $i + j = d$, and
2. $\lambda[\overline{i, j}] \simeq \mu$ (i.e., the diagram of μ is isomorphic to $\lambda[\overline{i, j}]$).

If λ does not include μ at d , then we say that λ *excludes* μ at d .

As an example, Definition 12 allows us to rephrase Proposition 10 as follows: $\delta_1 = k$ if and only if λ excludes the empty partition at $d-1$ and includes the empty partition at d .

Lemma 13 (corner constraints). *If λ includes μ at d , then for all $r \geq 1$,*

$$\alpha_r(G_\lambda) \leq rd + |\mu|. \tag{1}$$

Proof. By Lemma 9, every stable set contains at most d elements outside $\lambda[\overline{i, j}]$, so the union of r stable sets contains at most rd elements *outside* $\lambda[\overline{i, j}]$. On the other hand, since $\lambda[\overline{i, j}] \simeq \mu$, the union of r stable sets contains at most $|\mu|$ elements *inside* $\lambda[\overline{i, j}]$. Combining these two facts yield Inequality (1). \square

A remarkable consequence of matching theory is that corner constraints actually determine the values of α_r , in the following sense.

Theorem 14. *The value of $\alpha_r(G_\lambda)$ is the minimum value of $rd + |\mu|$ as i and j range over all nonnegative integers, where $d := i + j$ and $\mu := \lambda[\overline{i, j}]$.*

Proof. It is a standard fact from matching theory [11, Corollary 21.4b] that if B is a bipartite graph with vertex set V , then the maximum size of the union of r matchings is the minimum value of

$$r \cdot |X| + |E[V \setminus X]|$$

as X ranges over all subsets of V , and $|E[V \setminus X]|$ is the number of edges, neither of whose endvertices lie in X . In the context of B_λ , this means that we should consider all subsets of rows and columns—let i denote the number of rows and let j denote the number of columns—and minimize over $r \cdot (i + j)$ plus the number of boxes not in any of the chosen rows or columns. The key observation is that since λ is a partition, we can safely restrict attention to the case in which the chosen rows and columns are the *first* i rows and the *first* j columns, because choosing any other i rows or j columns can only increase the value of $|E[V \setminus X]|$ (while keeping the value of $|X| = i + j$ the same), thereby yielding a weaker upper bound on the value of α_r . But if we choose the first i rows and the first j columns, then the inequality we obtain is just a corner constraint. \square

3.2 Combinatorial Characterization of the CDS

From Theorem 14, we can infer an algorithm for computing the CDS of a partition graph G_λ . At first glance, it seems that we must consider infinitely many nonnegative integers i and j , but it turns out that we need only consider finitely many pairs (i, j) . To see this, first find (i_0, j_0) not in the diagram of λ such that $i_0 + j_0$ is as small as possible; by Proposition 10, $\delta_1 = i_0 + j_0$, so the associated corner constraints are $\alpha_r \leq r\delta_1$. No other corner constraints with $\mu = \emptyset$ need to be considered, because they all involve larger values of d and are therefore weaker. So to compute the CDS of G_λ , the only other pairs (i, j) we need to consider are those that are inside the diagram of λ .

The above algorithm is nice, but we can say more.

Definition 15. If δ is the CDS of λ , then the *normalized CDS* $\bar{\delta} = (\bar{\delta}_1, \bar{\delta}_2, \dots)$ is defined by $\bar{\delta}_i := \delta_1 - \delta_i$ for all i .

Knowing the CDS is equivalent to knowing the normalized CDS plus the value of δ_1 . This may seem like a trivial change of variables, but the advantage of the normalized CDS is that for a fixed r , there are only finitely many possible values that the sequence $(\bar{\delta}_1, \bar{\delta}_2, \dots, \bar{\delta}_r)$ can take, and as we now show, we can give a combinatorial characterization of each such sequence.

We begin by rephrasing the corner constraints in terms of the normalized CDS. Since $\alpha_r = \delta_1 + \dots + \delta_r$, we can subtract both sides of the corner constraint $\alpha_r \leq rd + |\mu|$ from $r\delta_1$ to obtain the equivalent inequality

$$\bar{\delta}_1 + \bar{\delta}_2 + \dots + \bar{\delta}_r \geq r(\delta_1 - d) - |\mu|. \quad (2)$$

We can now rephrase the algorithm in the first paragraph of this subsection in terms of the normalized CDS. First we compute δ_1 , and we also note that $\bar{\delta}_1 = 0$ always. We can then compute $\bar{\delta}_r$ by induction; if we have already computed $\bar{\delta}_1, \dots, \bar{\delta}_{r-1}$, then Theorem 14 tells us that $\bar{\delta}_r$ is the smallest nonnegative integer consistent with all constraints of the form of Inequality (2). Notice that for this calculation, we just need to know the ordered pairs $(\lambda[\bar{i}, \bar{j}], \delta_1 - i - j)$ as (i, j) ranges over the diagram. This observation is important enough that we introduce the following definition.

Definition 16. The *inclusion set* of a partition λ is the set of all pairs $(\mu, \delta_1 - d)$ such that λ includes μ at d and $\mu \neq \emptyset$.

If we forget λ itself but remember the inclusion set of λ and the value of δ_1 , then we can recover the normalized CDS. In fact, to compute the normalized CDS, we do not even need the entire inclusion set. Any $(\mu, \delta_1 - d)$ with $\delta_1 - d \leq 0$ can be safely omitted from the inclusion set, because then Inequality (2) is already implied by the nonnegativity of the $\bar{\delta}_i$.

Now let us fix r and think about how to compute the first r terms of the normalized CDS. We can omit any $(\mu, \delta_1 - d)$ with $|\mu| \geq r(\delta_1 - d)$, because again Inequality (2) is then implied by the nonnegativity of the $\bar{\delta}_i$. We also have the following result.

Lemma 17. *If we delete every $(\mu, \delta_1 - d)$ with $\delta_1 - d \geq r$ from the inclusion set of λ , then the remaining elements still suffice to recover $\bar{\delta}_1, \dots, \bar{\delta}_r$.*

Proof. We may assume that $r \geq 2$ since $\bar{\delta}_1 = 0$ always. If $(\mu, \delta_1 - d)$ is in the inclusion set, then there exists (i, j) such that $i + j = d$ and $\lambda[\bar{i}, \bar{j}] \simeq \mu$. Proposition 10 tells us that the diagram of λ contains all of antidiagonal $\delta_1 - 1$. By assumption, $i + j = d \leq \delta_1 - r$, so $\lambda[\bar{i}, \bar{j}]$ must contain all the boxes in row i extending from (i, j) out to antidiagonal $\delta_1 - 1$, namely the boxes $(i, j), (i, j + 1), \dots, (i, j + r - 1)$. Therefore, the first part of μ is at least r .

Let ν be the partition obtained from μ by deleting the first part of μ , so that $\nu \simeq \lambda[\bar{i} + 1, \bar{j}]$. Since $r \geq 2$ by assumption, $\nu \neq \emptyset$. As we just noted, the first part of μ is at least r , so

$$|\nu| \leq |\mu| - r. \quad (3)$$

Equations (2) and (3), together with the fact that λ includes ν at $d + 1$, imply that

$$\bar{\delta}_1 + \dots + \bar{\delta}_r \geq r(\delta_1 - d - 1) - |\nu| \geq r(\delta_1 - d) - |\mu|,$$

which is the constraint associated with $(\mu, \delta_1 - d)$. So omitting $(\mu, \delta_1 - d)$ from the inclusion set does no harm, since the inclusion of ν at $d + 1$ implies a constraint that is at least as strong. \square

In summary, to compute $\bar{\delta}_1, \dots, \bar{\delta}_r$, the only $(\mu, \delta_1 - d)$ in the inclusion set that we need are those satisfying $0 < \delta_1 - d < r$, and we need only consider μ such that

$$|\mu| < r(\delta_1 - d) < r^2.$$

These bounds have an important consequence: If we let S_r be the set of all pairs (μ, s) where $0 < s < r$ and $|\mu| < r^2$, then S_r is finite, and the first r terms of the normalized CDS of any λ , *no matter how large*, can be computed from a restriction of the inclusion set of λ to a subset of S_r . In particular, suppose we want to characterize all the partition graphs whose normalized CDS begins with a specific sequence of r numbers. Even though this is an infinite set of partition graphs, there are only finitely many possibilities for how the inclusion set of λ can restrict to a subset of S_r , and this restriction determines the first r terms of the normalized CDS. Therefore, we can prove results such as Propositions 18 and 19 below just by carrying out a finite computation (exhausting over all possible subsets of S_r). Since $\bar{\delta}_1 = 0$ always, we usually suppress mention of $\bar{\delta}_1$.

Proposition 18. *If λ is a partition, then*

- (a) $\bar{\delta}_2 = 1$ if and only if λ includes (1) at $\delta_1 - 1$, and
- (b) $\bar{\delta}_2 = 0$ if and only if λ excludes (1) at $\delta_1 - 1$.

Proposition 19. *If λ is a partition, then*

- (a) $(\bar{\delta}_2, \bar{\delta}_3) = (1, 2)$ if and only if λ includes (1) at $\delta_1 - 1$ and (2, 1) at $\delta_1 - 2$;
- (b) $(\bar{\delta}_2, \bar{\delta}_3) = (1, 1)$ if and only if λ includes (1) at $\delta_1 - 1$ and excludes (2, 1) at $\delta_1 - 2$;
- (c) $(\bar{\delta}_2, \bar{\delta}_3) = (0, 2)$ if and only if λ excludes (1) at $\delta_1 - 1$ and includes (2, 2) at $\delta_1 - 2$;
- (d) $(\bar{\delta}_2, \bar{\delta}_3) = (0, 1)$ if and only if λ excludes (1) at $\delta_1 - 1$ and (2, 2) at $\delta_1 - 2$ and includes (2) or (1, 1) at $\delta_1 - 1$;
- (e) $(\bar{\delta}_2, \bar{\delta}_3) = (0, 0)$ if and only if λ excludes (1) and (2) and (1, 1) at $\delta_1 - 1$ and (2, 2) at $\delta_1 - 2$.

We have computed similar characterizations of $(\bar{\delta}_2, \dots, \bar{\delta}_r)$ up to $r = 7$, but the corresponding Propositions become increasingly long and complicated, so we have chosen not to state them explicitly. But there is one remark we should make. As we noted earlier, there are at most Catalan-many possible values for the first r terms of the normalized CDS of a partition. The first surprise is that for $r = 4$, there are only 13, and not 14, possibilities; it turns out that for no partition graph do we have $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 2, 3)$. That is not all; for example, $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4, \bar{\delta}_5, \bar{\delta}_6)$ cannot assume any of the following values:

$$\begin{aligned} &(0, 0, 0, 2, 5), \quad (0, 0, 0, 3, 5), \quad (0, 0, 0, 4, 5), \quad (0, 0, 1, 3, 5), \quad (0, 0, 1, 4, 5), \\ &(0, 0, 2, 3, 5), \quad (0, 0, 2, 4, 4), \quad (0, 0, 2, 4, 5), \quad (0, 0, 3, 3, 4), \quad (0, 0, 3, 3, 5), \\ &(0, 0, 3, 4, 4), \quad (0, 0, 3, 4, 5), \quad (0, 1, 1, 4, 5), \quad (0, 1, 2, 4, 5), \quad (0, 1, 3, 3, 5), \\ &(0, 1, 3, 4, 4), \quad (0, 1, 3, 4, 5), \quad (0, 2, 2, 4, 5), \quad (1, 1, 3, 4, 5), \end{aligned}$$

as well as of course anything starting with $(0, 2, 3)$. The number of possible values for the first r terms of the normalized CDS of a partition is an integer sequence that begins

$$1, 2, 5, 13, 37, 108, 334, \dots$$

The closest match to this sequence in the Online Encyclopedia of Integer Sequences (oeis.org) as of this writing is A036249, the number of rooted trees of nonempty sets with n points, which begins

$$1, 2, 5, 13, 37, 108, 332, \dots$$

4 Partial CDS-Colorability

We mentioned in the Introduction that we have been able to prove the existence of a coloring that matches the first four terms of the CDS. We give the proof in this section.

Definition 20. Let \mathcal{L}_n denote the statement that for every partition graph G , there exists a coloring κ such that the first n parts of $\text{shape}(\kappa)$ coincide with the first n terms of the CDS of G .

The scrupulous reader may worry about what Definition 20 means if the length of $\text{shape}(\kappa)$ or of the CDS is less than n . For the purposes of Definition 20, in such cases we simply “pad out” the sequences with trailing zeros. So in particular, if \mathcal{L}_n is true for all n , then the Latin Tableau Conjecture is true.

The statement \mathcal{L}_1 becomes obvious as soon as one unwinds the definitions. The statements \mathcal{L}_2 and \mathcal{L}_3 are not too difficult either, as we now show.

Theorem 21. \mathcal{L}_2 and \mathcal{L}_3 are true.

Proof. We know from Propositions 3 and 11 that the only possibilities for δ_2 are $\delta_2 = \delta_1$ and $\delta_2 = \delta_1 - 1$. Suppose first that $\delta_2 = \delta_1$. That means that $\alpha_2 = 2\delta_1$, which means that there exists a disjoint union of two stable sets whose cardinality is $2\delta_1$. But by definition of δ_1 , each stable set has cardinality at most δ_1 . So each of the two stable sets has cardinality δ_1 , and we have exhibited a coloring with the desired properties. Similarly, if $\delta_2 = \delta_1 - 1$, then there must exist a union of two disjoint stable sets whose cardinality is $2\delta_1 - 1$. But the only way this can be realized is if one stable set has size δ_1 and the other has size $\delta_1 - 1$. The proof of \mathcal{L}_2 is complete.

There are five possibilities for the first three terms of the CDS. A similar argument to the one in the previous paragraph handles the cases $\delta_1 = \delta_2 = \delta_3$ and $\delta_1 = \delta_2$, $\delta_3 = \delta_1 - 1$. If $\delta_3 = \delta_2 = \delta_1 - 1$, then there exists a union of three disjoint stable sets whose cardinality is $3\delta_1 - 2$, but because $\alpha_2 = \delta_1 + \delta_2 = 2\delta_1 - 1$, at most one of these stable sets can have cardinality δ_1 . So the only possibility is that we have one stable set of size δ_1 and two stable sets of size $\delta_1 - 1$, which is what we need.

In the remaining two cases, $\delta_3 = \delta_1 - 2$. Since \mathcal{L}_2 is true, we know that there exist two stable sets with cardinalities δ_1 and δ_2 respectively. We can now establish the existence of a third stable set, disjoint from the first two and having cardinality $\delta_1 - 2$, via the same kind of argument that we used in the proof of Proposition 11. Namely, the diagram of λ contains the entirety of antidiagonal $\delta_1 - 1$, and so there are at least 3 boxes in row $\delta_1 - 3$, one of which we can use for our third stable set. We can work our way up the diagram of λ one row at a time, adding one box to our stable set in each row. This completes the proof of \mathcal{L}_3 . \square

The proof of \mathcal{L}_4 is of course more involved. We can handle all but five cases using ideas similar to those in the proof of Theorem 21. For convenience, we state this partial result as a separate Proposition.

Proposition 22. \mathcal{L}_4 is true except possibly when $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4)$ equals $(0, 0, 2)$ or $(0, 1, 2)$ or $(0, 2, 2)$ or $(1, 1, 2)$ or $(1, 2, 2)$.

Proof. Let us first dispense with the case $\bar{\delta}_4 = 3$. Theorem 21 tells us that \mathcal{L}_3 is true, so there exist three disjoint stable sets with cardinalities $\delta_1, \delta_2, \delta_3$ respectively. The diagram of λ contains the entire main antidiagonal, so there are at least 4 boxes in row $\delta_1 - 4$, one of which we can use for our fourth stable set. We can work our way up the diagram of λ one row at a time, adding one box to our fourth stable set in each row. This gives us a stable set of size $\delta_1 - 3$, so indeed $\bar{\delta}_4 = 3$.

If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4)$ is $(0, 0, 0)$ (respectively, $(0, 0, 1)$), then the only way to cover $4\delta_1$ (respectively, $4\delta_1 - 1$) boxes with four stable sets, each of size at most δ_1 , is for the stable sets to have sizes $\delta_1, \delta_1, \delta_1, \delta_1$ (respectively, $\delta_1, \delta_1, \delta_1, \delta_1 - 1$), which is precisely what we want.

Suppose $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 1, 1)$. Then at most one color can occur δ_1 times, so the only way to cover $4\delta_1 - 3$ boxes with four stable sets is with stable sets of sizes $\delta_1, \delta_1 - 1, \delta_1 - 1, \delta_1 - 1$. Finally, if $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 1, 1)$, then at most two colors can occur δ_1 times, so the only way to cover $4\delta_1 - 2$ boxes with four stable sets is with stable sets of sizes $\delta_1, \delta_1, \delta_1 - 1, \delta_1 - 1$. \square

For the five outstanding cases, we can deduce some necessary consequences.

Proposition 23. Let λ be a partition.

- (a) If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 0, 2)$ then λ excludes (1) , $(1, 1)$, and (2) at $\delta_1 - 1$ and excludes $(3, 3, 3)$ at $\delta_1 - 3$.
- (b) If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 1, 2)$ then λ excludes (1) at $\delta_1 - 1$, excludes $(2, 2)$ at $\delta_1 - 2$, and excludes $(3, 3, 2)$ at $\delta_1 - 3$.
- (c) If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 2, 2)$ then λ excludes (1) at $\delta_1 - 1$.
- (d) If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 1, 2)$ then λ excludes $(2, 1)$ at $\delta_1 - 2$, and excludes $(3, 3, 1)$ and $(3, 2, 2)$ at $\delta_1 - 3$.
- (e) If $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 2, 2)$ then λ excludes $(3, 2, 1)$ at $\delta_1 - 3$.

Proof. The proof is a routine verification. For example, if $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 0, 2)$ then λ must exclude $(3, 3, 3)$ at $\delta_1 - 3$, because otherwise Equation (2) with $r = 4$ would imply

$$2 = \bar{\delta}_2 + \bar{\delta}_3 + \bar{\delta}_4 \geq 4(\delta_1 - (\delta_1 - 3)) - 9 = 3,$$

which is false. All the other assertions are easily confirmed using the same kind of argument; we omit the details. \square

The way they are currently stated, the exclusion conditions in Proposition 23 are not so easy to grasp, but fortunately, in several cases, there is an easier way to think about them. For the remainder of this section, we shall use the notation $(i_1, j_1), (i_2, j_2), \dots, (i_s, j_s)$ (where $i_1 < i_2 < \dots < i_s$) to denote the coordinates of all the boxes along antidiagonal δ_1 that are *not* in the diagram of λ (at least one such box must exist, by Proposition 10).

As an illustration of why this notation is useful, suppose that λ excludes (1) at $\delta_1 - 1$. Then $j_{p+1} \leq j_p - 2$ (for $1 \leq p < s$); i.e., there cannot be two consecutive missing boxes in antidiagonal δ_1 . See Figure 4, where for clarity, we have put dots in the boxes on the main antidiagonal; the two consecutive missing boxes (1, 3) and (2, 2) in antidiagonal 4 force $\lambda[1, 2] \simeq (1)$, as indicated by the red box, so λ includes (1) at 3. It is intuitively easier to grasp “no two consecutive missing boxes in antidiagonal δ_1 ” than “ λ excludes (1) at $\delta_1 - 1$.”




Figure 4: Two consecutive missing boxes in antidiagonal 4

This type of reasoning will play an important role in our proof of Theorem 24.

Theorem 24. \mathcal{L}_4 is true.

Proof. Proposition 22 leaves five remaining cases, which we prove in approximately increasing order of complexity. In each case, we let λ be a partition whose normalized CDS begins with the specified numbers, and we construct disjoint stable sets S_1, S_2, S_3, S_4 of cardinalities $\delta_1, \delta_2, \delta_3, \delta_4$ respectively. We sometimes refer to the inclusion of a box in S_i as *coloring* the box with color i . In our figures below, the presence of i in a box means that the box is in S_i .

Case $(\delta_2, \delta_3, \delta_4) = (1, 2, 2)$. We claim that there cannot be four consecutive missing boxes in antidiagonal δ_1 ; for if there were, then by drawing a diagram analogous to Figure 4, we would deduce that λ includes $(3, 2, 1)$ at $\delta_1 - 3$, but this is ruled out by Proposition 23(e). For $i = 1, 2, 3$, we let S_i comprise all the boxes on antidiagonal $\delta_1 - i$. We let S_4 comprise any and all boxes on antidiagonal δ_1 in the diagram of λ , as well as all boxes with coordinates (i_p, j_{p+3}) for $1 \leq p \leq s - 3$ (by Proposition 19, λ includes $(2, 1)$ at $\delta_1 - 2$, so $s \geq 3$). See Figure 5 for an example.

There are two things to check: first, that S_4 has size $\delta_1 - 2$, and second, that S_4 really is a stable set that is disjoint from the other three. There are $\delta_1 + 1$ boxes on antidiagonal δ_1 , of which s are not in the diagram of λ , and $\delta_1 + 1 - s$ are in S_4 . So the cardinality of S_4 is $(\delta_1 + 1 - s) + (s - 3) = \delta_1 - 2$, as claimed. The row and column coordinates of the boxes of S_4 are all distinct by construction, so S_4 is a stable set. The crucial point is that

				3	2	1	4
			4	3	2	1	
	4		3	2	1		
4		3	2	1			
	3	2	1	4			
3	2	1					
2	1	4					
1							

Figure 5: Example for Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 2, 2)$

$j_{p+3} \leq j_p - 4$ (for $1 \leq p \leq s - 3$) because there cannot be four consecutive missing boxes in antidiagonal δ_1 . Therefore,

$$i_p + j_{p+3} \leq i_p + j_p - 4 = \delta_1 - 4,$$

and thus (i_p, j_{p+3}) does not lie on antidiagonals $\delta_1 - 1$, $\delta_1 - 2$, or $\delta_1 - 3$. So S_4 is indeed disjoint from the other three stable sets.

Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 1, 2)$. By Proposition 19, λ includes (1) at $\delta_1 - 1$, so $s \geq 2$. We claim that $j_{p+2} \leq j_p - 4$ for $1 \leq p \leq s - 2$. For if $j_{p+2} \geq j_p - 3$, then three out of four consecutive boxes on antidiagonal δ_1 must be missing, but by Proposition 23(d), λ excludes (2, 1) at $\delta_1 - 2$, so no three consecutive boxes on antidiagonal δ_1 are missing. So somewhere in λ , one of the configurations in Figure 6 must exist (where we have put dots in the boxes along the main antidiagonal). That is, λ must include either (3, 3, 1) or (3, 2, 2) at $\delta_1 - 3$, but this contradicts Proposition 23(d).




Figure 6: Missing 3 out of 4 consecutive boxes in antidiagonal δ_1

For $i = 1, 2$, we let S_i comprise all the boxes on antidiagonal $\delta_1 - i$, and we let S_4 comprise all the boxes on antidiagonal $\delta_1 - 3$. We let S_3 comprise any and all boxes on antidiagonal δ_1 , as well as all boxes with coordinates (i_p, j_{p+2}) for $1 \leq p \leq s - 2$. The rest of the argument is similar to the previous case; we readily verify that the cardinality of S_3 is $\delta_1 - 1$, and because $j_{p+2} \leq j_p - 4$ (for $1 \leq p \leq s - 2$), it follows that the boxes with coordinates (i_p, j_{p+2}) lie in antidiagonal $\delta_1 - 4$ or lower, and hence are disjoint from S_1 , S_2 , and S_4 . See Figure 7 for an example. This completes the proof of Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 1, 2)$.

The remaining three cases rely on the decomposition of λ into what we call *building blocks*. We regard the missing boxes $(i_1, j_1), \dots, (i_s, j_s)$ as dividing up λ into smaller (nonempty) pieces. More precisely, building block 0 is the set of boxes (i, j) in λ with $i < i_1$ and $j_1 \leq j$ (as long as this set is not empty), building block 1 comprises the boxes (i, j) in λ with $i_1 \leq i < i_2$ and $j_2 \leq j < j_1$, and so on. See Figure 8 for an example, where

				3	4	2	1
				4	2	1	3
			4	2	1	3	
3		4	2	1			
	4	2	1				
4	2	1	3				
2	1	3					
1	3						

Figure 7: Example for Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (1, 1, 2)$

the building blocks are colored green (and dots are placed along the main antidiagonal for reference). By construction, along antidiagonal δ_1 , a building block omits no boxes in its range, except possibly the bottom leftmost box, the top rightmost box, or both; we call these three kinds of building blocks *top*, *bottom*, and *interior* blocks respectively. In Figure 8, there are three interior blocks, and a bottom block, but no top block.




Figure 8: Building Blocks

The construction of S_1, \dots, S_4 proceeds in two phases. In Phase 1, we show how to color individual building blocks. Actually, for each kind of building block (top, bottom, interior), we show only how to color the *minimal* building blocks of a given height; minimality means that any other building block (for partitions with the specified normalized CDS values) of the same kind and height contains a minimal one. It suffices to show how to color minimal building blocks; if a building block is not minimal, we can simply ignore the extra boxes when forming S_1, \dots, S_4 . In Phase 2, we color λ by coloring each building block the way it was colored in Phase 1, and then coloring additional boxes as needed.

There is one important subtlety. Let $\Lambda(a, b, c)$ denote the set of partitions such that $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (a, b, c)$. A building block for $\Lambda(a, b, c)$ is often, but not always, itself a member of $\Lambda(a, b, c)$. For example, the partition $(1, 1, 1)$ is a bottom block for $\Lambda(0, 0, 2)$ (e.g., it is a building block for the partition $(6, 6, 6, 4, 4, 1, 1, 1) \in \Lambda(0, 0, 2)$) but $(1, 1, 1) \notin \Lambda(0, 0, 2)$ since $(1, 1, 1) \in \Lambda(0, 0, 1)$. To check that our colorings are valid, we must verify that the building blocks that are themselves in $\Lambda(a, b, c)$ are correctly colored in Phase 1, and we must also verify that Phase 2 produces correct colorings whether or not the constituent building blocks are themselves in $\Lambda(a, b, c)$.

Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 0, 2)$. Proposition 23(a) says that λ excludes (1) at $\delta_1 - 1$ (which as we noted above means that there are no two consecutive missing boxes in antidiagonal δ_1), and that λ excludes (1, 1) and (2) at $\delta_1 - 1$. Therefore, $j_{p+1} \leq j_p - 3$ (for

$1 \leq p \leq s-1$); i.e., the missing boxes in antidiagonal δ_1 must be at least three apart, and an interior building block must have a box at the start and end of antidiagonal $\delta_1 + 1$. Figures 9 and 10 illustrate the minimal interior building blocks of heights 4–12, along with a Phase 1 coloring (there is no interior building block of height 3 because Proposition 23(a) says that λ excludes $(3, 3, 3)$ at $\delta_1 - 3$). It should be clear from Figure 10 how the pattern extends to minimal interior building blocks with larger height.




Figure 9: Minimal Interior Building Blocks of Heights 4, 5, 6




Figure 10: Minimal Interior Building Blocks of Heights 7–12

The minimal top (respectively, bottom) building blocks, together with their colorings, may be obtained by deleting the top three rows (respectively, left three columns) from the minimal interior building blocks. All these building blocks, except (3) and $(1, 1, 1)$ and $(4, 4, 4, 3)$, are in $\Lambda(0, 0, 2)$, and it is readily verified that the given colorings are valid.

In Phase 2, after coloring each building block as in Phase 1, S_1 will always already have the correct cardinality. Moreover, in each building block, S_2 and S_3 are the same size as S_1 , so they also already have the correct cardinality. Thus we just need to show that S_4 can be suitably enlarged. To see this, first note that if (3) and $(1, 1, 1)$ and $(4, 4, 4, 3)$ are not involved, then between any two consecutive building blocks, we can put two more 4's

in antidiagonal $\delta_1 - 3$, and this will make S_4 the right size without putting any 4's in a row or column with an existing 4. See Figure 11 for an example.




Figure 11: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 0, 2)$: Phase 2

Finally, if any of the building blocks (3) or $(1, 1, 1)$ or $(4, 4, 4, 3)$ are involved, we put a 4 in antidiagonal $\delta_1 - 2$ between building blocks, and if $(4, 4, 4, 3)$ is involved, we also put a 4 in the one as-yet-uncolored box of $(4, 4, 4, 3)$ if necessary. It is easy to check that this construction works. See Figure 12.




Figure 12: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 0, 2)$: Phase 2

Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 1, 2)$. Proposition 23(b) says that λ excludes (1) at $\delta_1 - 1$ and excludes $(2, 2)$ at $\delta_1 - 2$; by arguments similar to those we have seen before, this implies that $j_{p+1} \leq j_p - 3$ for $1 \leq p \leq s - 1$. Moreover, Proposition 23(b) says that λ excludes $(3, 3, 2)$ at $\delta_1 - 3$, so if $j_{p+1} = j_p - 3$ (where $1 \leq p \leq s - 1$) then $\lambda|_{\overline{i_p, j_{p+1}}} \simeq (3, 3, 3)$. Therefore:

1. There is no restriction on the heights of minimal top and bottom building blocks, and they contain no boxes in antidiagonals $\delta_1 + 1$ or higher.
2. The shape $(3, 3, 3)$ is a minimal interior building block of height 3.
3. The remaining minimal interior building blocks have height at least 4, and they contain no boxes in antidiagonals $\delta_1 + 1$ or higher.

For Phase 1, in both Cases 1 and 3, we put all boxes (if any) in the diagram of λ on antidiagonal $\delta_1 - 1$ (respectively, $\delta_1, \delta_1 - 2, \delta_1 - 3$) in S_1 (respectively, S_2, S_3, S_4). In Case 3, we also put the top left box $(0, 0)$ in S_2 . See Figure 13, which also shows how we color $(3, 3, 3)$.




Figure 13: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 1, 2)$: Building Blocks

For Phase 2, if $(3, 3, 3)$ is not one of the building blocks, we simply fill antidiagonal $\delta_1 - 2$ with 3's and antidiagonal $\delta_1 - 3$ with 4's. If $(3, 3, 3)$ is involved, then see Figure 14. Again, the verification that this construction is correct is straightforward and we omit the details.




Figure 14: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 1, 2)$: Phase 2

Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 2, 2)$. Proposition 23(c) says that λ excludes (1) at $\delta_1 - 1$. Therefore, the partition (1) is not a valid building block; the minimal building blocks of height at most 5 are those shown in Figure 15. The larger minimal building blocks are interior building blocks that contain all the boxes in antidiagonal δ_1 except the box in row 0 and the box in column 0, and no other boxes. (We need not separately consider top and bottom building blocks of height at least 4 because they contain a valid interior building block as a proper subset.) To color one of these larger building blocks, we let S_1 and S_4 comprise all of antidiagonals $\delta_1 - 1$ and $\delta_1 - 3$ respectively; we let S_3 comprise all of antidiagonal $\delta_1 - 2$ except for the box in column 0, and we let S_2 comprise all of antidiagonal δ_1 plus the box with coordinates $(0, 0)$.




Figure 15: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 2, 2)$: Building Blocks

In Phase 2, if the building blocks are sufficiently large, then when we concatenate them, we simply add two more 3's along antidiagonal $\delta_1 - 2$ and two more 4's along antidiagonal $\delta_1 - 3$. The smaller building blocks are handled as shown in Figure 16. \square




Figure 16: Case $(\bar{\delta}_2, \bar{\delta}_3, \bar{\delta}_4) = (0, 2, 2)$: Phase 2

5 Concluding Remarks

We believe that there is some hope that the coloring algorithms in the proof of Theorem 24 can be generalized to prove the full conjecture, though at the moment, there remain several annoying ad hoc steps.

Another possible approach to proving the conjecture, which we have not investigated, is to examine closely what happens if one takes B_λ and applies the known general algorithms for constructing maximum-cardinality unions of i matchings in a bipartite graph, for $i = 1, 2, 3, \dots$. For an arbitrary bipartite graph, these unions of i matchings will not “nest” nicely, but maybe one can see how to make them nest in the case of B_λ .

It is also natural to ask more generally which classes of graphs are CDS-colorable. We suspect that CDS-colorability is NP-complete, but perhaps there are interesting classes of CDS-colorable graphs. Griggs [7] showed that incomparability graphs of symmetric chain orders are CDS-colorable. It is an unpublished result of the first author (proved independently by others [6, 9]) that indifference graphs (also known as unit interval graphs) are CDS-colorable, and that the coloring can be constructed by scanning the vertices from left to right, assigning the lowest available color at each step. Karen Collins (personal communication) has also informed us that *unbalanced split graphs* [4], which are graphs whose vertex set can be partitioned into a clique and an independent set in more than one way, are CDS-colorable. Finally, it may also be possible to extract some examples from the literature on integer multiflows [12]; there is no general “max-flow min-cut” theorem in this setting, but some partial results are known, which may be relevant.

For those with an interest in Rota’s basis conjecture, here is an ambitious generalization to matroids of the Latin Tableau Conjecture. Let λ be a partition, let δ denote the CDS of G_λ , and let $\mu = \delta'$ be the conjugate partition (i.e., the diagram of μ is obtained from the diagram of δ by transposing rows and columns). Let M be any matroid that is a disjoint union of independent sets of cardinalities μ_1, μ_2, \dots . Then we may conjecture that there is a way to place the elements of M into the diagram of λ , one element per box, such that the rows and the columns are all independent sets of M . We have not made any serious attempt to disprove this conjecture, so maybe it is false, but even if it is false, perhaps the counterexamples will help illuminate the Latin Tableau Conjecture and/or Rota’s basis conjecture.

Acknowledgments

We thank Katharine C. Walker for helpful conversations and ideas in the early stages of this research project.

References

- [1] Ron Aharoni, Eli Berger, He Guo, and Daniel Kotlar, “2-covers of wide Young diagrams,” [arXiv:2311.17670v2](#), 2023.
- [2] Michael O. Albertson and David M. Berman, “The chromatic difference sequence of a graph,” *Journal of Combinatorial Theory, Series B* **29** (1980), 1–12.
- [3] Timothy Y. Chow, C. Kenneth Fan, Michel X. Goemans, and Jan Vondrak, “Wide partitions, Latin tableaux, and Rota’s basis conjecture,” *Advances in Applied Mathematics* **31** (2003), 334–358.
- [4] Karen L. Collins and Ann N. Trenk, “Split Graphs,” in *Topics in Algorithmic Graph Theory*, by Lowell W. Beineke, Martin Charles Golumbic, and Robin J. Wilson (eds.), Cambridge University Press, 2021, Chapter 9, pp. 189–206.
- [5] D. de Werra, “Investigations on an edge coloring problem,” *Discrete Mathematics* **1** (1971), 167–179.
- [6] Molly Fenn and Eric Sommers, “A transitivity result for ad-nilpotent ideals in type A ,” *Indagationes Mathematicae* **32** (2021), 1175–1189.
- [7] Jerrold R. Griggs, “Problems on chain partitions,” *Discrete Mathematics* **72** (1988), 157–162.
- [8] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd edition, Oxford University Press, 1995.
- [9] Jacob P. Matherne, Alejandro H. Morales, and Jesse Selover, “The Newton polytope and Lorentzian property of chromatic symmetric functions,” *Selecta Mathematica New Series* **30** (2024), 42.
- [10] Gian-Carlo Rota, “Ten mathematics problems I will never solve,” *Mitteilungen der Deutschen Mathematiker-Vereinigung* **6** (1998), 45–52.
- [11] Alexander Schrijver, *Combinatorial Optimization: Polyhedra and Efficiency, Vol. A*, Springer, 2003.
- [12] Alexander Schrijver, *Combinatorial Optimization: Polyhedra and Efficiency, Vol. C*, Springer, 2003.