

Temperatures of Robin Hood

Niranjan Balachandran^a Ankita Dargad^a Urban Larsson^b

Submitted: Mar 19, 2025; Accepted: Jan 31, 2026; Published: May 8, 2026

© The authors. Released under the CC BY license (International 4.0).

Abstract

Cumulative Games were introduced by Larsson, Meir, and Zick (2020) to bridge some conceptual and technical gaps between Combinatorial Game Theory (CGT) and Economic Game Theory. The partizan ruleset ROBIN HOOD is an instance of a Cumulative Game, namely, WEALTH NIM. It is played on multiple heaps, each associated with a pair of cumululations, interpreted here as wealth. Each player chooses one of the heaps, removes tokens from that heap not exceeding their own wealth, while simultaneously diminishing the other player's wealth by the same amount. In CGT, the *temperature* of a *disjunctive sum* game component is an estimate of the urgency of moving first in that component. It turns out that most of the positions of ROBIN HOOD are *hot*. The temperature of ROBIN HOOD on a single large heap shows a dichotomy in behavior depending on the ratio of the wealths of the players. Interestingly, this bifurcation is related to Pingala (Fibonacci) sequences and the Golden Ratio ϕ : when the ratio of the wealths lies in the interval (ϕ^{-1}, ϕ) , the temperature increases linearly with the heap size, and otherwise it remains constant, and the mean values have a reciprocal property. It turns out that despite ROBIN HOOD displaying high temperatures, playing in the hottest component might be a sub-optimal strategy.

Mathematics Subject Classifications: 91A46, 91A05, 11B39

1 Introduction

In the Era of Pingala, two wetland tribes engage in a dispute over land pieces on various islands for farming, which escalates into a war. Being honorable tribes, their chiefs agree to certain rules for fighting: as a preparation, each tribe allocates a number of soldiers to each island, and the tribe with the smaller number of soldiers gets to start. On day one, this tribe gets to challenge their opponent on an island of their choice, and by the principle of “first player advantage” they get to weaken the opponent's strength on that

^aDepartment of Mathematics, Indian Institute of Technology, Bombay, India
(niranj@math.iitb.ac.in, ankitadargad.iitb@gmail.com).

^bDepartment of Industrial Engineering and Operations Research, Indian Institute of Technology, Bombay, India (larsson@iitb.ac.in).

island. The next day, the other tribe retaliates and attacks any island of their choice. This continues each day, and as a result of the fighting, some land pieces get ruined. The war concludes when only one tribe is left on each island. At that stage, the tribes get to collect the remaining fertile lands that are under their control in their respective conquered islands, and return home to their respective villages to celebrate the end of the war.¹

How can the tribes maximize their gains in terms of conquered land pieces? And, which island should they target first?




Figure 1: A three days’ war that is initiated by the red tribe. The war ends on the 3rd day as only one tribe is left on each island. Ruined land pieces and beaten soldiers are colored gray. The colored arrows specify the attacking tribe.

This scenario describes a game of a ruleset dubbed **ROBIN HOOD**, which is an instance of **WEALTH NIM**, a **CUMULATIVE GAME** [10]. This is a two-player alternating play combinatorial game [15, 2] that is played on a finite number of heaps of finite sizes. To each heap, each player has an associated wealth, an integer that we shall refer to as the *heap wealth*, which determines their strength on that heap. On their turn, a player

- chooses one of the heaps,
- removes a certain number of tokens from that heap not exceeding their own heap wealth, and
- reduces the opponent’s wealth on that heap by the same amount.

¹This has some resemblance with the classical “Colonel Blotto” war game [3, 5]. The main difference is that they use simultaneous play, and their game concerns the assignment of the forces to the islands, while in our setting, we will regard the assignments of soldiers as given, and the main challenge is the sequential selections for ‘the next fight’. The story, of course, is entirely fictional.

A player who cannot move loses (normal play). Let $\mathbb{N} = \{1, 2, \dots\}$, and let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. We denote by $(n; a, b)$ a ROBIN HOOD instance comprising of a single heap of size $n \in \mathbb{N}_0$, Left's wealth equal to $a \in \mathbb{N}_0$ and Right's wealth equal to $b \in \mathbb{N}_0$. In case the player with greater wealth removes more tokens than the opponent's wealth, their wealth drops below zero. If so, by convention, we set the opponent's wealth to zero, since they cannot play further. In the scenario described above, each island represents a heap, the number of land pieces represents the heap size and the tribes' strengths represent the players' wealths. The blue (resp. red) tribe represents Left (resp. Right).²

Figure 1 illustrates a play on a ROBIN HOOD instance with two heaps of sizes 9 and 4 (land pieces). Left (blue tribe) has heap wealths of 3 and 2, and Right (red tribe) has heap wealths of 2 and 1, respectively. The '+' sign denotes the disjunctive sum, meaning that on their turn, a player chooses exactly one heap (island) to play on, while the other heap remains the same. The war ends when only one tribe remains on each island, but ROBIN HOOD continues due to the normal play convention. It ends when the current player has no token to remove. In this example, after the war, Left has at most three rounds of play before she runs out of tokens, while Right's resources can still be abundant. Thus, Right wins the game. (A careful scrutiny ought to leave Left somewhat disappointed; if she had instead continued responding in the first component, then Right would have won by only two tokens.)

By separating an instance of ROBIN HOOD into a war phase and a conclusion phase, we make a first observation. After the war phase, for each player there is a total number of remaining tokens in all heaps, where they still have non-zero heap wealth. This number can be considered as 'their number of free moves', as those tokens can be removed one at a time, without interference from their opponent.

Intuitively, both players desire to start as they get to diminish the opponent's strength as early as possible by as much as possible. Additionally, it turns out that, in the Figure 1 example, playing in the first component is preferential for either player.

To elucidate: If Right starts in the first component, then as discussed before, Left can respond in the first component, and Right can win by $(5 - 3) = 2$ moves, whereas, if Right starts in the second component, Left can respond in the first component and gain 7 moves, while Right receives only 2 moves from the second component. Therefore, Right loses by $(7 - 2) = 5$ moves. Thus, the total loss for Right by starting in the second component instead of the first one is $2 - (-5) = 7$ moves.

If Left starts and plays in the first component, she gains 7 land pieces in this component and also receives 2 land pieces from the second one, leading to a win by 9 moves. However, if Left starts in the second component, Right can play in the first component on his turn. In the end, Left gains 3 land pieces from the second component, while Right gets 5 land pieces from the first component. Therefore, Left loses by 2 moves. Thus, the total loss for Left by starting in the second component instead of the first is $9 - (-2) = 11$.

This also highlights the urgency to start the game, as a player forfeits $(2 + 9) = 11$ moves by not starting. The notion of *temperature* [15, 2] (see Definition 15) is an estimate,

²Rumors, from the Sherwood forest, tell that Robin Hood got inspired by folktales about the wetland tribes' wars, in the spirit: "I take, and you pay!".

which attempts to capture this sense of urgency. In many cases, a higher temperature indicates greater urgency, while a lower temperature indicates less urgency. In keeping with this terminology, a game with positive temperature is called a “hot” game. The computation of the temperature is often facilitated by considering a pictorial gadget called the *thermograph* [15] (see Definition 16). We will discuss this in greater detail in Section 3. Another related concept is the *mean value* of a game. It is an estimate of how good a game is for the respective player; a larger mean value is usually better for Left and vice versa.

Based on this intuition our main question is:

What are the temperatures and mean values of ROBIN HOOD?

CGsuite [16] guides us on the temperatures and the mean values of the two ROBIN HOOD games, $(n; 5, 4)$ and $(n; 5, 3)$, with varying n . For any game G , let $t(G)$ and $m(G)$ denote the temperature and mean of G , respectively. See Table 1.

Table 1: The temperatures and mean values of the games $(n; 5, 4)$ and $(n; 5, 3)$ for a few initial heap sizes n . We indicate with italics and bold text when patterns emerge.

n	$t(n; 5, 4)$	$m(n; 5, 4)$	$t(n; 5, 3)$	$m(n; 5, 3)$
3	0	0	0	0
4	0	0	1/2	1/2
5	1/2	1/2	1	1
6	5/4	3/4	7/4	5/4
7	17/8	7/8	19/8	13/8
8	49/16	15/16	3	2
9	4	<i>1</i>	7/2	5/2
10	5	<i>1</i>	4	<i>3</i>
11	6	<i>1</i>	4	<i>4</i>
12	7	<i>1</i>	4	<i>5</i>
13	8	<i>1</i>	4	<i>6</i>

Consistency can be observed in the values of the temperature and the mean of the two games in Table 1 for large heap sizes. It also shows a reciprocal behavior between the temperature and the mean values: The temperature (mean value) of the game $(n; 5, 4)$ increases (stabilizes) as the heap size grows, whereas the temperature (mean value) of the game $(n; 5, 3)$ stabilizes (increases) with increasing heap size.

Next, we observe the temperatures for different wealth pairs in Table 2, which illustrates whether the temperature of $(n; a, b)$ is stabilizing or increasing with n for different pairs (a, b) .

When one tribe is significantly stronger than the other and the number of land pieces is large, the stronger tribe has (intuitively speaking) a better chance of securing a huge loot of war, regardless of the actions of the weaker tribe. As a result, neither tribe has a strong desire to start the war, irrespective of the number of land pieces. The low desire to

Table 2: Temperatures of the games $(n; a, b)$ for different pairs (a, b) . The second column specifies the behavior of the temperature with increasing n . The last column is the lower bound of n for which the property holds. Cases where the temperature increases with heap size are indicated in bold.

$(n; a, b)$	<i>Property</i>	$t(n; a, b)$	<i>Bound</i>
$(n; 1, 1)$	Increasing	$n - 1$	1
$(n; 1, 2)$	Stabilizing	1	3
$(n; 1, 3)$	Stabilizing	1	4
$(n; 2, 3)$	Increasing	$n - 3$	8
$(n; 2, 4)$	Stabilizing	2	6
$(n; 2, 9)$	Stabilizing	2	11
$(n; 3, 4)$	Increasing	$n - 4$	7
$(n; 3, 5)$	Stabilizing	4	10
$(n; 3, 6)$	Stabilizing	3	9
$(n; 5, 8)$	Increasing	$n - 8.5$	15
$(n; 5, 9)$	Stabilizing	6	16
$(n; 7, 11)$	Increasing	$n - 11.5$	20
$(n; 7, 12)$	Stabilizing	9	15

start a war shows that the heat in this situation should not increase with an increase in the number of land pieces. However, when the players' wealths are relatively equal, both players benefit from starting the game. This increases their desire to begin, leading to a rise in temperatures as the heap size increases.

This indicates that the two different patterns of the temperature values are related to the ratio of the players' wealth. It is clear from Table 2 that whenever the ratio of wealth (larger to smaller) is bigger than or equal to $5/3$, the temperature stabilizes, and it keeps increasing when the ratio is smaller than or equal to $8/5$.

Our main result gives a complete description of the temperatures and mean values of ROBIN HOOD when played on a single large heap. Let $\phi = \frac{1+\sqrt{5}}{2}$ denote the *golden ratio*, and given two positive integers a and b , let $(U_k)_{k \geq 0}$ be the sequence defined by

1. $U_0 \geq U_1$,
2. $U_{k+2} = U_{k+1} + U_k$ for all $k \geq 0$, and
3. for some $q \geq 0$, $U_q = \min \{a, b\}$ and $U_{q+1} = \max \{a, b\}$.

Theorem 1 (Main Theorem). *For fixed positive integers a and b , let $G_n = (n; a, b)$ be instances of ROBIN HOOD and $(U_k)_{k \geq 0}$ be defined as above. If $n \geq a + b$, then G_n is a*

hot game. Moreover, for all sufficiently large n ,

$$t(G_n) = \begin{cases} b - U_0 & \text{if } \frac{a}{b} < \phi^{-1}; \\ n - a + \frac{U_0 - b}{2} & \text{if } \phi^{-1} < \frac{a}{b} < 1; \\ n - a & \text{if } \frac{a}{b} = 1; \\ n - b + \frac{U_0 - a}{2} & \text{if } 1 < \frac{a}{b} < \phi; \\ a - U_0 & \text{if } \phi < \frac{a}{b}, \end{cases}$$

and

$$m(G_n) = \begin{cases} -(n - (a + b) + U_0) & \text{if } \frac{a}{b} < \phi^{-1}; \\ \frac{U_0 - b}{2} & \text{if } \phi^{-1} < \frac{a}{b} < 1; \\ 0 & \text{if } \frac{a}{b} = 1; \\ \frac{a - U_0}{2} & \text{if } 1 < \frac{a}{b} < \phi; \\ n - (a + b) + U_0 & \text{if } \phi < \frac{a}{b}. \end{cases}$$

The notions of ‘urgency’ and temperature are not always the same in the sense that playing on the hottest component is not always an optimal strategy. Here, by optimal strategy, we mean the alternating play strategy that maximizes the earning of the players and minimizes the loss in terms of ‘number of moves’. A playing strategy that implies ‘playing in the hottest component in a disjunctive sum of games is an optimal strategy for either player’ is known as *Hotstrat* [15].³ If a ruleset satisfies hotstrat, then knowing the temperature and the thermograph of a game reveals all the facets of the game. Many ROBIN HOOD positions satisfy hotstrat, for instance the position in Figure 1. However, ROBIN HOOD is even more interesting, as the following theorem shows.

Theorem 2 (No Hotstrat). *There exists a ROBIN HOOD disjunctive sum game, with components of distinct temperatures, such that the unique winning move is in the coolest component.*

We prove this result at the end of Section 4. The rest of the paper is organized as follows.

- In Section 2 we mention several papers that inspired this work.
- In Section 3 we review the notion of temperature and mean value using Left and Right stops, along with thermographs.
- In Section 4 we define ROBIN HOOD formally and analyze some basic properties and positions.
- In Section 5, we see the connection between Pingala sequences and ROBIN HOOD.
- Section 6 collects some relevant facts about Pingala sequences and its generalizations.

³Hotstrat also specifies the move to be made in the hottest component [15].

- In Section 7, we study a simpler game that we dub LITTLE JOHN.
- In Section 8 we show that for large heaps, the Left and Right stops for ROBIN HOOD and LITTLE JOHN are the same, and we also study the thermographs of LITTLE JOHN.
- In Section 9, we prove the main result, Theorem 1, by first justifying that the thermographs of LITTLE JOHN and ROBIN HOOD are the same for large heap sizes.

2 Literature review

Curiously, our ruleset has a resemblance to the classical *impartial* ruleset EUCLID, which is played on two non-empty heaps of pebbles. A player must remove a multiple of the size of the smaller heap from the larger heap. A position is represented by a pair of positive integers (x, y) , where say $x \leq y$. Note that if $x = y$, then the position is terminal. Example: $(2, 7) \rightarrow (2, 3) \rightarrow (1, 2) \rightarrow (1, 1)$. Since we put the requirement that (both) heaps remain non-empty, no further move is possible. Note that the losing moves are forced. Optimal play reduces to minimizing the relative distance of the heaps.

Theorem 3 ([4]). *A player wins EUCLID if and only if they can remove a multiple of the smaller heap such that the ratio of the heap sizes (x, y) satisfies $1 \leq y/x < \phi$.*

In ROBIN HOOD, the player with lesser wealth cannot remove a non-trivial multiple of their own wealth. Similarly, in EUCLID, for positions (x, y) where $y/x < \phi$, players are restricted from removing non-trivial multiples. Consequently, the removal recurrences are identical in both scenarios.

Cumulative Games were defined in a broad sense in [10]. The purpose of that monograph is to explore an intersection of classical game theory with combinatorial game theory. The ruleset WEALTH NIM a.k.a. WEALTH PEBBLES is introduced as an example where player cumulations are part of the rules of how to move, but do not contribute to the payoffs that the players gain when the game ends. Since the purpose of [10] is to provide a broad framework for further study, no efforts were made to solve proposed individual games and rulesets. The current paper is among the first ones to do so.

The problem of bounding the temperature of combinatorial games has been extensively studied for specific games. A central example is DOMINEERING, where it is a folklore conjecture, generally attributed to Berlekamp, that the temperature is at most two. Shankar and Sridharan [14] performed an extensive computational search over billions of positions of DOMINEERING, discovering many new temperature values, but none exceeding two. Subsequently, Uiterwijk and Barton [17] provide further supporting evidence for the same bound by constructing and analyzing endgame databases for all single-component positions up to 15 squares, finding one position with temperature two and no positions with temperature greater than two. Very recently, Huntemann and Maciosowski [8] conducted a parallel exhaustive search, identifying many more DOMINEERING positions with temperature two.

Berlekamp [1] explored the notion of temperature across a variety of combinatorial games, including GO, DOMINEERING and AMAZONS, and ordered them according to the temperatures of their hottest known positions.

Huntemann, Nowakowski, and dos Santos [9] contributed significantly toward general techniques for bounding temperature. They show that if the lengths of the confusion intervals of all subpositions in a game are bounded by a constant K , then the temperature of the game is bounded by $3K/2$.

Later, Huntemann and Maciosowski [7] disprove the long-held conjecture that the temperature of SNORT is bounded by the maximum degree of the graph, and further conjecture that it is bounded by a polynomial in the maximum n^{th} degree of the graph (the n^{th} degree of a vertex is the number of vertices at distance exactly n from it, and the maximum n^{th} degree of a graph is the maximum n^{th} degree over all vertices).

Interestingly, the temperature of ROBIN HOOD for fixed wealths shows both bounded and unbounded behavior (in terms of the heap size) depending on the relationship between the wealth ratio and the golden ratio.

3 Preliminaries

In order to establish the foundation for proving our main results regarding temperatures and mean values, we revisit the concept of a ‘thermograph’. Understanding thermographs requires familiarity with several key terminologies from CGT. We will not discuss standard CGT concepts, such as game comparison and canonical forms, as they are less critical in this context, and are well-covered in the existing literature (e.g., [2, 15]). We begin by defining the pivotal concept of a Number game.

3.1 Numbers, mean values and stops

Intuitively, a game is called a ‘Number’ if each player prefers that the other player starts. Recall that a *sub-position* of a game can be the game itself or any option of the game or any option of options, etc.

Definition 4 (Numbers). A short game x is a Number if, in the canonical form of x , every sub-position y satisfies $y^L < y^R$ for all y^L and y^R .⁴

As usual, \emptyset denotes the empty set (of options). Every game is associated with a Number game in the following sense.

Definition 5 (Mean Value). Consider a short game G . Its mean value $m(G)$ is the Number such that, for any positive dyadic ϵ , for all sufficiently large n , $n \cdot m(G) - \epsilon \leq n \cdot G \leq n \cdot m(G) + \epsilon$.

In [15, Theorem 3.23] it is proved that every game has a mean value.

The most basic Number games are as follows: For all $k \in \mathbb{Z}_{>0}$, we define the *integer games* k and $-k$ recursively as:

⁴The definition of a Number in [15] does not hold for $\{* | *\}$ and many other literal form games that equal some canonical form Number.

- $k = \{k - 1 \mid \emptyset\}$;
- $-k = \{\emptyset \mid -k + 1\}$,

where $0 = \{\emptyset \mid \emptyset\}$.

For all odd $k \in \mathbb{Z}$ and $n \in \mathbb{N}$, we define the *dyadic rational games* recursively as:

$$\frac{k}{2^n} = \left\{ \left[\frac{k-1}{2^n} \right] \mid \left[\frac{k+1}{2^n} \right] \right\},$$

where the brackets denote the reduction of the fraction such that the numerator is odd, unless 0. For example, with $k = 1$ and $n = 3$, the game $1/8 = \{0 \mid 1/4\}$.

By [15, Proposition 3.5], integer and dyadic rational games follow the standard arithmetic properties. For instance, the disjunctive sum of the games 1 and $\frac{1}{2}$ equals the game $1 + \frac{1}{2} = \frac{3}{2}$.

The games 0 and 1 are vacuously Numbers and similarly, all integers are also Numbers. Moreover, $\frac{1}{2} = \{0 \mid 1\}$ is a Number, since $0 < 1$. The game $\{* \mid *\}$ is also a Number since its canonical form is 0, which is a Number.

Henceforth, we shall call all integers and dyadic rationals simply as *dyadics*. Let the set of dyadics be $\mathbb{D} = \left\{ \frac{k}{2^n} : k \in \mathbb{Z}, n \in \mathbb{N}_0 \right\}$, where, with some abuse of notation, the cases for $n = 0$ correspond to the integers. Since our definition of “Number” is not the same as in most textbooks, we include a proof of the consistency of terminology.

Theorem 6. *The set of dyadics has a bijective relation with the set of canonical form Numbers.*

Proof. All integers are vacuously Numbers, and they are in canonical form. Next, we show that every non-integer dyadic $\frac{k}{2^n}$ is a canonical form Number. First note that $\frac{k-1}{2^n} < \frac{k+1}{2^n}$, and so it is a Number. Moreover, this game cannot be reduced (indeed, domination does not apply, and it does not reverse out).

Now, we will show that any literal form Number x equals a dyadic. Let y be the canonical form of x . By induction, every y^L and y^R is dyadic. Now, by domination, y will have only one left and one right option, (say) y^L and y^R . By the simplicity theorem, y is the simplest dyadic between y^L and y^R . \square

A Number game can also be interpreted as the number of ‘free moves’ available for Left (if positive) or Right (if negative). This raises the question of the maximum number of free moves a player is guaranteed in alternating play of a game. In CGT, the maximum number of free moves Left is guaranteed in alternating play, when starting a game, is called the Left stop of the game (this may be negative; if so, it is the minimum guaranteed loss for Left). Similarly, the negative of the maximum number of free moves Right is guaranteed when starting the game is called the Right stop of the game (this may be positive).

Definition 7 (Stops). For a game G , the Left stop $\ell(G)$ and the Right stop $r(G)$ are defined as:

$$\ell(G) = \begin{cases} x & \text{if } G \text{ equals a dyadic } x; \\ \max_{G^L} (r(G^L)) & \text{otherwise;} \end{cases}$$

$$r(G) = \begin{cases} x & \text{if } G \text{ equals a dyadic } x; \\ \min_{GR} (\ell(G^R)) & \text{otherwise.} \end{cases}$$

The stops of a game G is the ordered pair $s(G) = (\ell(G), r(G))$.

Remark 8. If the Left and Right stops of a game G are not the same, then G does not equal a dyadic.

We shall recall a few results about the stops from [15] that will be of use to us later.

Proposition 9 ([15, Proposition 3.17]). *Let G be a game and let x be a number. Then,*

1. $\ell(-G) = -r(G)$ and $r(-G) = -\ell(G)$;
2. if $G \geq x$ then, $\ell(G) \geq r(G) \geq x$. Likewise if $G \leq x$ then, $r(G) \leq \ell(G) \leq x$.

Proposition 10 ([15, Proposition 3.18]). *Let G be a short game and let x be any dyadic. Then,*

1. $\ell(G) \geq r(G)$;
2. $\ell(G + x) = \ell(G) + x$ and $r(G + x) = r(G) + x$.

When we analyze games in terms of the stops, we momentarily stop thinking about winning, while rather emphasizing the stops. Sometimes we abuse language and instead of “stops”, we say *scores* (the loot of war). This terminology would be consistent with Milnor’s positional games [13], where his ‘scoring functions’ correspond to normal play stops.⁵

In a disjunctive sum of games, if the first player can guarantee a higher total score by playing on a particular component, in comparison to the other components, then that component is considered ‘urgent’. For example, in the sum $\{4 \mid -5\} + \{1 \mid -2\}$ the game $\{4 \mid -5\}$ is urgent compared to $\{1 \mid -2\}$.

3.2 Penalized positions and temperature

To numerically estimate this notion of urgency, we recursively apply equal penalties to both players. The minimum penalty at which the Left and Right stops of a game become equal (i.e., the game is no longer urgent) provides an estimate of the urgency. Note that this estimate does not depend on any other components. Let \mathbb{D}^+ denote the set of non-negative dyadics. Let X be a set of games and y be a Number, then $X + y$ denotes the set $\{x + y : x \in X\}$.

Definition 11 (Penalized Position). Let G be a short game in canonical form and let $p \in \mathbb{D}^+$. Then, G *penalized by p* , denoted by G_p , is recursively defined as

⁵It turns out that the normal play reduced canonical forms [6] correspond to Milnor’s positional games [13].

- $G_p = \{G_p^{\mathcal{L}} - p \mid G_p^{\mathcal{R}} + p\}$ for all $0 \leq p \leq t$ where t is the minimum p for which the Left and Right stops of G_p are equal to a dyadic, say x ,
- $G_p = x$ for all $p > t$.

Here $G_p^{\mathcal{L}}$ denotes the set of penalized Left option of G , and similarly for Right.

Example 12. Let $G = \{9 \mid 7\}$. Then, $G_p = \{9 - p \mid 7 + p\}$, for all $p \leq 1$. At $p = 1$, G_p becomes $\{8 \mid 8\}$, and hence, for all $p > 1$, $G_p = 8$. In summary,

$$G_p = \begin{cases} \{9 - p \mid 7 + p\} & \text{if } p \leq 1; \\ 8 & \text{if } 1 < p. \end{cases}$$

The concept of *cooling by t* defined in [15] is similar but not identical to the concept of *penalized by p* . The next observation helps to understand the similarity and the differences between the two.

Observation 13. Let G_p^C and G_p denote the game G cooled by p and G penalized by p , respectively. Then, $G_p^C = G_p$ for all but finitely many p at which they differ infinitesimally.

Using Observation 13, the stops of G_p and G_p^C are the same for all p . Therefore, we can use the results on the stops of G cooled by t for that of G_p penalized by p .

Proposition 14 ([15, Theorem 5.11(b)]). For any game G , for all $p \in \mathbb{D}^+$, $\ell(G) \geq \ell(G_p) \geq r(G_p) \geq r(G)$.

Proof. The middle inequality follows by Proposition 10 and the other two inequalities follow because the penalty reduces the benefit for both players. \square

The minimum penalty, for which the penalized game remains no longer urgent, is the *temperature* of the original game.

Definition 15 (Temperature). The temperature of a dyadic $G = k/2^n$ is $t(G) = -1/2^n$, where $k \in \mathbb{Z}$ and $n \in \mathbb{N} \cup \{0\}$ and if $n > 0$, k is an odd integer. The temperature $t(G)$ of a non-dyadic G is the smallest $p \in \mathbb{D}^+$ such that $\ell(G_p) = r(G_p)$.

When the game G is unambiguous from the context, we sometimes simply write $t = t(G)$ and $m = m(G)$, and similarly $\ell = \ell(G)$ and $r = r(G)$. In Example 12, $t(G) = 1$.

Combinatorial games are divided into three categories depending on the temperature:

- *Hot*: A game G is hot if $t(G) > 0$;
- *Tepid*: A game G is tepid if $t(G) = 0$;
- *Cold*: A game G is cold if $t(G) < 0$.

3.3 Thermographs and their walls

One of the issues with the definition of temperature is that it is somewhat unwieldy from a computational point of view. A more intuitive and appealing way of understanding (and computing) the temperature of a game comes from a more pictorial device, the *thermograph*.

Definition 16 (Thermograph). Let G be a short game. Then, the thermograph of G , $\text{Therm}(G)$, is a plot of the Left and Right stops of G_p (on the X -axis) with respect to p (on the Y -axis).

For a given (hot) game G , $\ell(G_p)$ and $r(G_p)$ are bounded functions on the penalty p . Sometimes we think of them as the walls of $\text{Therm}(G)$.

Definition 17 (Walls). The sets

$$\begin{aligned} \text{LW}(G) &= \{(\ell(G_p), p) : p \in \mathbb{D}^+\} \text{ and} \\ \text{RW}(G) &= \{(r(G_p), p) : p \in \mathbb{D}^+\} \end{aligned}$$

are called the large left and large right walls of $\text{Therm}(G)$, respectively. The sets

$$\begin{aligned} \text{lw}(G) &= \{(\ell(G_p), p) : p \in \mathbb{D}^+, p \leq t\} \text{ and} \\ \text{rw}(G) &= \{(r(G_p), p) : p \in \mathbb{D}^+, p \leq t\} \end{aligned}$$

are called the small left and small right walls of $\text{Therm}(G)$, respectively. A wall is either a small or a large wall.

At times, we may refer to the ‘walls of the game’ rather than explicitly stating the ‘walls of the thermograph of the game.’ However, in both cases, we are referring to the same concept.

The large left (right) wall can be seen as an extension of the small left (right) wall, continuing indefinitely. In Figure 2, we depict a thermograph where $ABCD\infty$ and $ED\infty$ is the large left and large right wall of the thermograph, respectively, while $ABCD$ and ED is the small left and small right wall, respectively.

The temperature is the y -coordinate of the point where the small left and right walls merge. In Figure 2, the y -coordinate of D is the temperature. Moreover, in [15, Theorem 5.17], there is a proof that the x -coordinate of the same point equals the mean value of the game. Thus, given a game, the thermograph gives us a computational means to find both the temperature and the mean value of the game. While computing the walls of a game G , in general, the large walls of the options need to be considered.

3.4 An example of a wall computation

For any game G , the left and right walls of $\text{Therm}(G)$ correspond to the graphs of $\ell(G_p)$ and $r(G_p)$ as a function of p , respectively.




Figure 2: The thermograph of $G = \{6, \{10 \mid \{5 \mid 3\}\} \mid -5\}$.

In the case where $\ell(G) = r(G)$, we have $G_p = G$ when $p = 0$, and $G_p = \ell(G)$ for $p > 0$. Consequently, $\ell(G_p) = \ell(G) = r(G) = r(G_p)$ for all $p \geq 0$, which implies that $\text{Therm}(G)$ is a vertical line at $\ell(G)$.

Now, consider the case when $\ell(G) > r(G)$. By Definition 15, the smallest p for which $\ell(G_p) = r(G_p)$ is $t(G)$, denoted by t in short. For all $p \leq t$, we have

$$\begin{aligned} \ell(G_p) &= \max_{G^L} r(G_p^L - p) \\ &= \max_{G^L} r(G_p^L) - p \end{aligned} \tag{1}$$

$$\begin{aligned} r(G_p) &= \min_{G^R} \ell(G_p^R + p) \\ &= \max_{G^R} \ell(G_p^R) + p \end{aligned} \tag{2}$$


Equations (1) and (2) follow using Proposition 10. Geometrically, $r(G_p^L)$ represents the large right wall of $\text{Therm}(G_p^L)$, denoted as $\text{RW}(G_p^L)$. Moreover, the function ‘max’ in Equation (1) corresponds to the function ‘leftmost’ in a geometric sense, as the orientation of the number line is reversed. Consequently, $\max_{G^L} r(G_p^L)$ represents the leftmost $\text{RW}(G_p^L)$. This leftmost $\text{RW}(G_p^L)$ may be a combination of right walls from different options, as the right walls of G_p^L ’s could intersect. The term $-p$ in Equation (1) indicates a 45° rightward tilt of the wall $\max_{G^L} r(G_p^L)$. We also refer to this transformation as a *right tilt*.

Similarly, the right wall of G below $p = t$ is the left-tilted rightmost large left wall of right options. This aligns with Equation (2) as the left tilt captures the $+p$ shift, the rightmost aspect corresponds to the min, and the large left wall of right options corresponds to $\ell(G_p^R)$.

The two tilted walls meet when $\ell(G_p) = r(G_p)$, and which occurs at $p = t$. In other words, the value of p at which the two walls intersect is the temperature of G . Furthermore, $G_p = \ell(G_t)$ for all $p > t$. Consequently, the thermograph above this point is simply a vertical line at $\ell(G_t)$.

For instance, consider the game $H = \{\{2 \mid 0\}, \{3 \mid -3\}, 1/2 \mid -4\}$. The thermographs of the left options of H are shown in Figure 3a. The solid edges in Figure 3b represents the

leftmost $\text{RW}(H_p^L)$. Note that it is a combination of the right walls of different options. The 45° rightward tilt of this wall is depicted by the solid edges in Figure 3c. Similarly, the rightmost $\text{LW}(H_p^R)$ is represented by purple dashed edge in Figure 4 and the $\text{Therm}(H)$ is given by solid edges in the same figure.



(a) Thermographs of Left options of H . The thermographs of $\{3 \mid -3\}$, $\{2 \mid 0\}$ and $1/2$ are represented in red, blue and green, respectively. (b) Leftmost large right wall of thermographs of Left options of H (solid line). (c) Tilted leftmost large right wall of thermographs of Left options H (solid line).

Figure 3: Thermographs corresponding to $H = \{\{2 \mid 0\}, \{3 \mid -3\}, 1/2 \mid -4\}$.




Figure 4: Thermograph of $H = \{\{2 \mid 0\}, \{3 \mid -3\}, 1/2 \mid -4\}$.

3.5 Masts and tents

Sometimes, we view a thermograph as the two functions that define it, but other times, it is convenient to view it as a vertical structure. In this spirit, we define some particularly simple structures. In our proofs to come, we will assume such structures of the options by induction, and prove that they survive in the inductive step.

Let $A \subset \mathbb{D} \times \mathbb{D}$. If, for all $(x, y) \in A$, $y = kx + c$, for some constant c , then we say that A has slope k .

Definition 18 (Masts and Tents). Consider a (hot or tepid) game G . Then:

- $G \in \text{Mast}$, if $\text{LW}(G) = \text{RW}(G)$;
- $G \in \mathcal{DT}$ (double tent), if $\text{lw}(G)$ has slope -1 and $\text{rw}(G)$ has slope $+1$;
- $G \in \mathcal{LT}$ (left single tent), if $\text{lw}(G)$ has slope -1 and $\text{rw}(G)$ has slope 0 ;

- $G \in \mathcal{RT}$ (right single tent), if $\text{lw}(G)$ has slope 0 and $\text{rw}(G)$ has slope +1.

The terms double tent, left single tent, and right single tent refer to the shapes of a game's thermograph. These thermograph shapes can be seen in Figure 5. The three categories – double tent, left single tent, and right single tent – are collectively referred to as *tents*.




Figure 5: Mast and Tents.

In more detail we get the following (by keeping track of the constant c in the definition of slope).

Observation 19. *We have:*

- $G \in \mathcal{DT}$, if for all $p \leq t(G)$, $\ell(G_p) = t - p + m$ and $r(G_p) = p - t + m$;
- $G \in \mathcal{LT}$, if for all $p \leq t(G)$, $\ell(G_p) = t - p + m$ and $r(G_p) = m$;
- $G \in \mathcal{RT}$, if for all $p \leq t(G)$, $\ell(G_p) = m$ and $r(G_p) = p - t + m$,

where, as usual $t = t(G)$ and $m = m(G)$.

The following lemma helps in determining the temperatures and mean values of games with tent-shaped thermographs. Let us reformulate the case of $p = 0$, how it applies to the proof of our main theorem.

Lemma 20 (Temperature and Mean of Tents). *Let G be a game. Then,*

1. if $G \in \mathcal{DT}$, then $t = (\ell - r)/2$ and $m = (\ell + r)/2$;
2. if $G \in \mathcal{LT}$, then $t = \ell - r$ and $m = r$;
3. if $G \in \mathcal{RT}$, then $t = \ell - r$ and $m = \ell$,

where, as usual $t = t(G)$, $m = m(G)$, $\ell = \ell(G)$ and $r = r(G)$.

Proof. Apply Observation 19 with $p = 0$. □

In this spirit of preparing for the main proofs to come, let us present a general lemma concerning the simplest of thermographs.

Lemma 21. Consider a game G and let H denote the Left (Right) option of G with the largest (smallest) Left (Right) stop. If $H \in \mathbf{Mast}$, then $\text{LW}(G)$ ($\text{RW}(G)$) does not depend on any other Left (Right) option than H .

Proof. It suffices to prove that the leftmost large right wall of Left options of G is $\text{LW}(H)$. Recall that for a game G , G_p denotes the game G penalized by p . We get, for all $p \geq 0$ and all $G^L \in G^{\mathcal{L}}$,

$$r(H_p) = r(H) = \ell(H) \tag{3}$$

$$\geq \ell(G^L) \tag{4}$$

$$\geq \ell(G_p^L) \quad (\text{by Prop 14})$$

$$\geq r(G_p^L). \quad (\text{by Prop 10})$$

Equation (3) follows by Definition 16 as $H \in \mathbf{Mast}$. Equation (4) follows as H is the option with the largest Left stop. A similar proof works for the Right options. \square

4 Sherwood Organization

Let us formalize the ruleset ROBIN HOOD. We denote by $[a]$ the set $\{1, 2, \dots, a\}$.

Definition 22 (Robin Hood). Let $n, a, b \in \mathbb{N}_0$. A single-heap ROBIN HOOD game $(n; a, b)$, where n is the heap size, a and b are the wealths of Left and Right players, respectively, has the following options:

1. the Left options are $(n - i; a, b - i)$ where $i \in [\min \{n, a\}]$;
2. the Right options are $(n - j; a - j, b)$ where $j \in [\min \{n, b\}]$.

By convention, non-positive wealth is deemed to be 0 because a player with 0 or negative wealth cannot make a move. Recall that this game can also be played on multiple heaps. A ROBIN HOOD game on multiple heaps is the same as the disjunctive sum of single heap ROBIN HOOD games.

To understand a multiple heap game, it suffices to know the game values of single heap games. For this reason, from now onward in this paper, we will only consider the single heap games. However, the simplest game values, the canonical forms, often quickly become intractable. Let us give some intuition why.

The canonical form of the game $(4; 2, 2)$ is $G = \pm\{2, \{2 \mid \pm 1\}\}$. Suppose that we play the sum $(4; 2, 2) + (3; 1, 2) = G + \{\pm 1 \mid -2\}$. Left loses if she plays to $2 + \{\pm 1 \mid -2\}$, but she wins if she plays to $\{\pm 1 \mid -2\} + \{2 \mid \pm 1\}$. In a sense, the most likely ‘best’ move can fail depending on the surrounding context. The standard abstract way to explain this type of situation is that, indeed the game 2 is incomparable with the game $\{2 \mid \pm 1\}$, and neither option reverses out (which has to be checked). Similar arguments show that generic games of the form $(n; b, b)$ have b canonical options for each player (all options are sensible depending on situation). Thus, the complexity of canonical form games quickly becomes intractable. However, there are some very obvious options that never come into play.

Proposition 23. Consider the ROBIN HOOD game $(n; a, b)$, with $n > b$. If $a > b$, then the Left options $(n - i; a, 0)$, $b < i \leq \min\{n, a\}$, are dominated by the Left option $(n - b; a, 0)$.

Proof. For $i > b$, $(n - i; a, 0) = n - i < n - b = (n - b; a, 0)$. □

Thus, from now onward, we only consider the non-dominated options (with $i \leq \min\{a, b\}$).

Proposition 24. Consider $n, a, b \in \mathbb{N}_0$ and let $G = (n; a, b)$. Then

1. $-n \leq G \leq n$;
2. $-n \leq \ell(G) \leq n$ and $-n \leq r(G) \leq n$;
3. $-G = (n; b, a)$.

Proof. Left can win $n - (n; a, b)$ and $(n; a, b) + n$ playing second, by playing on the number in every turn. The 2nd item follows using the 1st item along with Proposition 9. At last, the negative of a game is swapping the roles of the players. □

Theorem 25 (Robin Hood Positions). Let $G = (n; a, b)$ be a ROBIN HOOD position. We have the following facts:

1. $G = 0$ if $a = b = 0$;
2. $G = n$ if $a > b = 0$, and $G = -n$ if $b > a = 0$;
3. $G = *n$ if $n \leq \min\{a, b\}$;
4. G is hot, if $n > \min\{a, b\} > 0$;
5. $\ell(G) = \max_{GL} r(G^L)$ and $r(G) = \min_{GR} \ell(G^R)$, if $n > \max\{a, b\}$.

Proof. Starting with the first item, neither player has moves, and the result is trivial.

Regarding the second item, it suffices to check that $(n; a, 0) - n$ is a \mathcal{P} -position. If Left starts and moves to $(n - j; a, 0) - n$, Right wins by responding with $(n - j; a, 0) - (n - 1)$. This happens because, by induction, $(n - j; a, 0) = (n - j) \leq (n - 1)$. Similarly, if Right starts and moves to $(n; a, 0) - (n - 1)$, Left wins by responding with $(n - 1; a, 0) - (n - 1)$. By the same argument, $(n, 0, b) = -n$.

In the third item, the Left options are $(n - j; a, b - j)$ where $0 < j \leq n$. Since $n - j \leq \min\{a, b - j\}$, by induction, the Left options are $0, *, *2, \dots, *(n - 1)$. By the same argument, the Right options are also the same. Hence, the game G is $\{0, *, \dots, *(n - 1) \mid 0, *, \dots, *(n - 1)\}$, which is $*n$.

For the fourth item, suppose without loss of generality that $a \geq b$. Then $n > b$. Thus, $\ell(G) \geq r(n - b; a, 0) = n - b > 0$.

On the other hand $r(G) \leq \ell(n - b; a - b, b)$. If $a = b$, then $\ell(n - b; a - b, b) = -(n - b) < 0$, and otherwise, since the heap size $n - b$ will decrease further in computing $\max\{r(n - b - i; a - b, b - i)\}$, $r(G) < n - b$.

Therefore, $r(G) \leq \ell(n - b; a - b, b) < n - b < \ell(G)$ and consequently, G is hot.

The last item is a consequence of item 4. □

As promised in the introduction, let us prove that ROBIN HOOD does not belong to hotstrat.

Proof of Theorem 2. Let $G_1 = (11; 1, 1)$ and $G_2 = (12; 2, 1)$ be two ROBIN HOOD games. Thus, $G_1 = \{10 \mid -10\}$ and $G_2 = \{11 \mid \{10 \mid -10\}\}$. Hence $t(G_1) = 10$ and $t(G_2) = 1$.

Let $G = G_1 + G_2$. If Right starts the game G by playing in the hottest component, G_1 , to $(10; 0, 1) + G_2$, then Left can respond by playing in G_2 to $(10; 0, 1) + (11; 2, 0)$. This game is Left winning (by one land piece). Whereas, if Right starts the game by playing in the cooler component, G_2 , to the game $(11; 1, 1) + (11; 1, 1)$, Right wins by henceforth mimicking Left's moves. \square

5 Robin Hood meets Little John

By Theorem 25, we know that the ROBIN HOOD positions with sufficiently large heap sizes are hot; both players benefit from starting the game. Is the advantage the same regardless of the initial move each player makes, or is there a specific move that offers the greatest benefit? Since reduced wealth is disadvantageous for a player, the move that maximizes the reduction of the opponent's wealth is likely to yield the highest advantage.


Definition 26 (Little John Move). Consider a ROBIN HOOD game $(n; a, b)$, where $n > 0$. For $a > 0$, if $b > 0$, Left's Little John move is $(n - \min\{n, a, b\}; a, b - \min\{n, a, b\})$, and if $b = 0$, it is $(n - 1; a, 0)$. Right's Little John move is analogously defined.⁶

Definition 27 (Little John Path). A Little John path is a sequence of alternating play Little John moves.

After a Little John path, there will usually be a number of free Little John moves for either player, depending on the size of n . In Figure 6, we illustrate the Little John path on $(n; a, b)$ if $a/b = 1.4$.

If Left and Right have wealths of 27 and 17 respectively, the sequence of pairs (Left's wealth, Right's wealth) along the Little John path when Right starts the game is $(27, 17)$, $(10, 17)$, $(10, 7)$, $(3, 7)$, $(3, 4)$, $(0, 4)$. Observe that the wealths of Left and Right changes alternatively until one of the players' wealth reaches 0. Thus, we can simplify and reduce this sequence to 27, 17, 10, 7, 3, 4, 0, where the first value belongs to Left, the second to Right, the third to Left, and so on. If we omit the 0th term from the reversed sequence of players' wealths on the Little John path, the resulting sequence $(4, 3, 7, 10, 17, 27)$ follows the pattern where each term is the sum of the two preceding terms. The next section explores the properties of such sequences.

⁶The term 'Little John move' is inspired by Little John, a companion of Robin Hood in English folklore, known for his great physical strength. The name reflects that this move yields the greatest advantage in the game.



(a) The Little John Path when Left starts. (b) The Little John Path when Right starts.

Figure 6: The Little John Paths of $(n; a, b)$, for $a/b = 1.4$ and sufficiently large n .

6 Pingala sequences and their properties

The previous section points at classical number sequences.

Definition 28 (Pingala Sequence). The Pingala Sequence $(P_k)_{k \geq 0}$ is given by $P_0 = 0$, $P_1 = 1$ and, for all $k \geq 0$, $P_{k+2} = P_{k+1} + P_k$.⁷

The wealths of the players may not lie in this sequence, so we define a modified version of it.

Definition 29 (Modified Pingala Sequence). A sequence $(U_k)_{k \geq 0}$ is a Modified Pingala Sequence (MP-sequence) if U_0 and U_1 are positive integers such that $U_0 \geq U_1$, and $U_{k+2} = U_{k+1} + U_k$ for all $k \geq 0$.

Remark 30. The Pingala sequence includes 0 as its 0th term. However, as noted earlier, the reversed sequence of players' wealths follows the Pingala sequence pattern only when 0 is excluded. The condition $U_0, U_1 \in \mathbb{Z}_{>0}$ in the definition of the Modified Pingala sequence ensures that 0 and any negative numbers are omitted from the sequence.

Next, we see some sequences derived from the Pingala sequence and their properties.

Definition 31 (Ratio Sequences). The ratio sequences $(O_k)_{k \geq 0}$ and $(E_k)_{k \geq 0}$ are given by, for all $k \geq 0$, $O_k = P_{2k+2}/P_{2k+1}$ and $E_k = P_{2k+3}/P_{2k+2}$.

We refer to $(O_k)_{k \geq 0}$ and $(E_k)_{k \geq 0}$ as the Odd and Even Ratio sequences, respectively. It is well known that Even and Odd ratio sequences are strictly decreasing and increasing, respectively, and both converge to the golden ratio.

The following proposition is routine.

⁷This is commonly known as the Fibonacci sequence. We attribute the sequence instead to Acharya Pingala, an ancient (3rd-2nd century BCE) Indian mathematician and poet who used this sequence in poetry long before Fibonacci lived.

Proposition 32. Consider two positive integers a and b such that $a \geq b$. Then, there exists

1. a unique MP-sequence $(U_k)_{k \geq 0}$, such that $U_\mu = b$ and $U_{\mu+1} = a$, for some $\mu \geq 0$;
2. a unique MP-sequence $(V_k)_{k \geq 0}$, such that $V_\nu = a$ and $V_{\nu+1} = b$, for some $\nu \geq 0$.

Proof. The proof follows simply by generating the sequence by using the MP-sequence rules that every term is sum of the last two terms and the starting term in the sequence is greater than equal to the next term. \square

As an illustrative example, consider $a = 31$ and $b = 20$. Then $(U_k) = (7, 2, 9, 11, 20, 31, 51, 82, \dots)$, with $\mu = 4$, and $(V_k) = (31, 20, 51, 71, 122, \dots)$, with $\nu = 0$.

Let us list a few elementary properties of MP-sequences.

Observation 33. If $(U_k)_{k \geq 0}$ is a MP-sequence, then the following hold:

1. $U_1 \leq U_0 < U_2 < U_3 < \dots < U_n < \dots$;
2. $U_k \in \mathbb{Z}_{>0}$ for all $k \geq 0$.

The next proposition determines the μ and ν from Proposition 32, given a and b . First, we relate the ratio of consecutive terms of a MP-sequence with those of the Pingala sequence.

Lemma 34. Let $(U_k)_{k \geq 0}$ be an MP-sequence. Then, for all $k \geq 1$, for all i such that $0 \leq 2i \leq k - 1$,

1. $U_{k-2i}/U_{k-2i-1} \leq O_j \iff U_{k+1}/U_k \geq E_{j+i}$;
2. $U_{k-2i+1}/U_{k-2i} \leq O_j \iff U_{k+1}/U_k \leq O_{j+i}$.

Proof. Let k be fixed. We will prove both statements together by induction on i .

Base Case: Let $i = 0$.

1. Then the following statements are equivalent.

$$\frac{U_k}{U_{k-1}} \leq O_j \left(= \frac{P_{2j+2}}{P_{2j+1}} \right) \tag{5}$$

$$\frac{U_{k+1} - U_k}{U_k} \geq \frac{P_{2j+1}}{P_{2j+2}} \tag{6}$$

$$\frac{U_{k+1}}{U_k} \geq \frac{P_{2j+3}}{P_{2j+2}} (= E_{j+0}). \tag{7}$$

Equation (6) follows by inverting the Equation (5).

2. For $i = 0$, both sides are the same.

Suppose that the statement is true for i and let $2(i + 1) \leq k - 1$. Then, for part (1), the following statements are equivalent by induction on i .

$$\frac{U_{k-2(i+1)}}{U_{k-2(i+1)-1}} \leq \frac{P_{2j+2}}{P_{2j+1}} \quad (8)$$

$$\frac{U_{k-2i-1} - U_{k-2i-2}}{U_{k-2i-2}} \geq \frac{P_{2j+1}}{P_{2j+2}} \quad (9)$$

$$\frac{U_{k-2i-1}}{U_{k-2i-2}} \geq \frac{P_{2j+3}}{P_{2j+2}} \quad (10)$$

$$\frac{U_{k-2i}}{U_{k-2i-1}} \leq \frac{P_{2(j+1)+2}}{P_{2(j+1)+1}} = O_{j+1} \quad (11)$$

$$\frac{U_{k+1}}{U_k} \geq E_{j+1+i}. \quad (\text{by induction})$$

Equation (9) follows by inverting Equation (8) and Equation (11) follows by inverting Equation (10) and adding 1. This completes the induction. We skip the proof of part (2) as it is similar to part (1). \square

Proposition 35. *Let $a, b > 0$ be integers and let $(U_k)_{k \geq 0}$ be the unique MP-sequence such that $U_\mu = b$, $U_{\mu+1} = a$ for some $\mu \geq 0$. Then μ is determined as follows:*

1. If $\frac{a}{b} < \phi$, then $\mu = 2\hat{\mu}$, where $\hat{\mu} = \min\{k \geq 0 : \frac{a}{b} \leq O_k\}$;
2. If $\frac{a}{b} > \phi$, then $\mu = 2\tilde{\mu} + 1$, where $\tilde{\mu} = \min\{k \geq 0 : \frac{a}{b} \geq E_k\}$.

Proof. The indexes $\hat{\mu}$ and $\tilde{\mu}$ always exist because the sequences $(O_k)_{k \geq 0}$ and $(E_k)_{k \geq 0}$ are increasing and decreasing respectively, and both converge to ϕ .

1. The following inequalities are equivalent:

$$\frac{a}{b} \leq O_{\hat{\mu}} \quad (12)$$

$$\frac{U_{\mu+1}}{U_\mu} \leq O_{\hat{\mu}}$$

$$\frac{U_{\mu-2\hat{\mu}+1}}{U_{\mu-2\hat{\mu}}} \leq O_0 (= 1) \quad (13)$$

$$U_{\mu-2\hat{\mu}+1} \leq U_{\mu-2\hat{\mu}} \quad (14)$$

$$\mu - 2\hat{\mu} = 0. \quad (\text{Obs 33(1)})$$

Equation (12) follows by the definition of $\hat{\mu}$ and Equation (13) follows by Proposition 34(2) for $i = \hat{\mu}$ and $j = 0$.

2. The following inequalities are equivalent:

$$\frac{a}{b} \geq E_{\tilde{\mu}} \quad (15)$$

$$\begin{aligned} \frac{U_{\mu+1}}{U_{\mu}} &\geq E_{\tilde{\mu}} \\ \frac{U_{\mu-2\tilde{\mu}}}{U_{\mu-2\tilde{\mu}-1}} &\leq O_0 (= 1) \\ U_{\mu-2\tilde{\mu}} &\leq U_{\mu-2\tilde{\mu}-1} \\ \mu - 2\tilde{\mu} - 1 &= 0. \end{aligned} \tag{16}$$

(Obs 33(1))

Equation (15) follows by the definition of $\tilde{\mu}$ and Equation (16) follows by Proposition 34(1) for $i = \tilde{\mu}$ and $j = 0$.

This completes the proofs of both statements. □

We now introduce the Pingala sequence with alternating signs, which will be used to establish a relation between two MP-sequences and hence two games.

Definition 36. The alternating Pingala sequence $(\bar{P}_k)_{k \geq 0}$ is given by $\bar{P}_k = (-1)^{k+1} P_k$.

Proposition 37. *The following equalities hold.*

1. $\bar{P}_{k+2} = \bar{P}_k - \bar{P}_{k+1}$ for all $k \geq 0$;
2. $\sum_{i=1}^k P_i = P_{k+2} - 1$ for all $k \geq 1$;
3. $\sum_{i=1}^k \bar{P}_i = 1 + (-1)^{k-1} P_{k-1}$ for all $k \geq 1$;
4. For any MP-sequence $(U_k)_{k \geq 0}$, $\sum_{i=1}^k U_i = U_{k+2} - U_1 - U_0$ for all $k \geq 1$.

Proof. The proofs of all the statements follow by standard induction arguments. □

Later we will consider two ROBIN HOOD games for which the wealth of one of the players differ by one, while the other player's wealth remains the same. To compare them, we here give a relation between the MP-sequences generated by their respective pair of wealths.

Proposition 38. *Let $(U_n)_{n \geq 0}$ and $(V_n)_{n \geq 0}$ be two MP-sequences. Suppose there exist constants α and β such that:*

1. $V_{\alpha} = U_{\beta}$ and $V_{\alpha+1} = U_{\beta+1} + 1$. Then, for all $0 \leq k \leq \min\{\alpha, \beta\}$, $V_{\alpha-k} = U_{\beta-k} + \bar{P}_k$;
2. $V_{\alpha} = U_{\beta} + 1$ and $V_{\alpha+1} = U_{\beta+1}$. Then, for all $0 \leq k \leq \min\{\alpha, \beta\}$, $V_{\alpha-k} = U_{\beta-k} + \bar{P}_{k+1}$.

Proof. We only prove part 1, as the proof of part 2 is similar. We induct on k . For $k = 0$, the statement follows by the assumption, and for $k = 1$ (assume $\min\{\alpha, \beta\} \geq 1$), we have

$$\begin{aligned} V_{\alpha-1} &= V_{\alpha+1} - V_{\alpha} \\ &= U_{\beta+1} + 1 - U_{\beta} \end{aligned}$$

$$= U_{\beta-1} + \bar{P}_1,$$

where the first and last equalities follow by Definition 29.

Suppose the statement holds for all $k \leq n$ where n satisfies $1 \leq n+1 \leq \min\{\alpha, \beta\}$. Then, by the induction hypothesis $V_{\alpha-(n-1)} = U_{\beta-(n-1)} + \bar{P}_{n-1}$ and $V_{\alpha-n} = U_{\beta-n} + \bar{P}_n$, and hence,

$$\begin{aligned} V_{\alpha-(n+1)} &= V_{\alpha-(n-1)} - V_{\alpha-n} \\ &= U_{\beta-(n-1)} - U_{\beta-n} + \bar{P}_{n-1} - \bar{P}_n \\ &= U_{\beta-(n+1)} + \bar{P}_{n+1}, \end{aligned}$$

where the first and last equalities follow by Definition 29. □

7 Little John's Ruleset

In this section, we study a simpler ruleset derived from ROBIN HOOD. It will turn out that the stops of this new ruleset are the same as the stops of ROBIN HOOD, which we prove in the next section.

Intuitively, in ROBIN HOOD, players gain by minimizing their opponent's wealth because it reduces the opponent's wealth-reducing power. So, we digress along the Little John Path (Definition 27) to understand the stops and we do so by playing a variation of ROBIN HOOD where players are only allowed to make Little John moves (Definition 26).

Definition 39 (LITTLE JOHN). Let $(n; a, b)^*$ denotes a position of the ruleset LITTLE JOHN on a heap of size n .

1. if $n = 0$ or $a = 0 = b$, neither player has any option, and otherwise:
2. if $a, b > 0$, the only Left option is $(n - \gamma; a, b - \gamma)^*$ and the only Right option is $(n - \gamma; a - \gamma, b)^*$ where $\gamma = \min\{n, a, b\}$;
3. if $a > 0 = b$, the only Left option is $(n - 1; a, 0)^*$ and Right has no option. Similarly, if $b > 0 = a$, the only Right option is $(n - 1; 0, b)^*$ and Left has no option.

The next propositions and lemma allow us to compute the stops of LITTLE JOHN. For simplicity of notations, we remove the extra set of bracket from $\ell((n; a, b)^*)$ and write $\ell(n; a, b)^*$ and follow similar notion for Right stop.

Proposition 40. Consider $n, a, b \in \mathbb{N}_0$ and let $G = (n; a, b)^*$. Then

1. $-n \leq G \leq n$;
2. $-n \leq \ell(G) \leq n$ and $-n \leq r(G) \leq n$.

Proof. The proof is same as that of Proposition 24. □

Lemma 41. Consider $n, a, b \in \mathbb{N}_0$ such that $n > \min\{a, b\}$ and $a \geq b > 0$. Then

1. $(n; a, b)^*$ is hot;
2. $\ell(n; a, b)^* = r(n - b; a, 0)^*$ and $r(n; a, b)^* = \ell(n - b; a - b, b)^*$.

Proof. The proof is similar to that of Theorem 25. □

The Left and Right stops of a game G are the same as the game value if and only if G is a Number. Since Numbers are cold, and Lemma 41 establishes that LITTLE JOHN positions with large n and positive wealths are hot, these positions cannot be Numbers. Consequently, the Left and Right stops of such LITTLE JOHN positions differ from their game value. The next lemma computes these stops.

Lemma 42 (Little John Stops). *Consider $n, a, b \in \mathbb{N}_0$ and suppose $n \geq a + b$.*

1. If $a = b = 0$, then $r(n; a, b)^* = 0$.
2. If $b > 0 = a$, then $r(n; a, b)^* = -n$ and if $a > 0 = b$, then $r(n; a, b)^* = n$.
3. If $b \geq a > 0$, then $r(n; a, b)^* = -(n - a)$.
4. If $a > b > 0$, let $(U_i)_{i \geq 0}$ denote the unique MP-sequence such that $U_\mu = b$ and $U_{\mu+1} = a$.
 - (a) If $\frac{a}{b} < \phi$, then $r(n; a, b)^* = a + b - n - U_0$.
 - (b) If $\frac{a}{b} > \phi$, then $r(n; a, b)^* = n - (a + b) + U_0$.

Proof. To calculate $r(n; a, b)^*$, we observe the following:

1. For all $n \geq 0$, $r(n; 0, 0)^* = 0$ as $(n; 0, 0)^* = 0$.
2. If $b > 0 = a$, then it follows easily that $r(n; 0, b)^* = r(-n) = -n$ for all $n \geq 0$. Similarly, if $a > 0 = b$, then we have $r(n; a, 0)^* = r(n) = n$ for all $n \geq 0$.
3. if $0 < \frac{a}{b} \leq 1$, then $r(n; a, b)^* = \ell(n - a; 0, b)^* = -(n - a)$. The first equality holds for all $n \geq a + b$, by Lemma 41. The second equality holds as $(n - a; 0, b)^* = -(n - a)$ by Theorem 25(2).
4. (a) Case $1 < \frac{a}{b} < \phi$. From Lemma 35, $\mu = 2\hat{\mu}$ where $\hat{\mu} = \min\{k \geq 0 : \frac{a}{b} \leq O_k\}$. Then, for all $n \geq a + b$,

$$\begin{aligned} r(n; a, b)^* &= r(n; U_{2\hat{\mu}+1}, U_{2\hat{\mu}})^* \\ &= \ell(n - U_{2\hat{\mu}}; U_{2\hat{\mu}-1}, U_{2\hat{\mu}})^* \end{aligned} \tag{17}$$

$$= r\left(n - \sum_{i=2\hat{\mu}-1}^{2\hat{\mu}} U_i; U_{2\hat{\mu}-1}, U_{2\hat{\mu}-2}\right)^* \tag{18}$$

$$= r\left(n - \sum_{i=1}^{2\hat{\mu}} U_i; U_1, U_0\right)^* \tag{19}$$

$$= \ell(n - \sum_{i=1}^{2\tilde{\mu}} U_i - U_1; 0, U_0)^* \quad (20)$$

$$= -(n - \sum_{i=1}^{\mu} U_i - U_1) \quad (21)$$

$$= -(n - U_{\mu+2} + U_0) \quad (22)$$

$$= -(n - (a + b) + U_0).$$

Equation (17) follows using Observation 33(1), Lemma 41(2), and the fact that, at each step, the heap size is at least the sum of the players' wealths, as $n \geq a + b$, and both decrease equally in each iteration. By repeating this process, we get Equations (18), (19) and (20). Equation (21) holds using item (2) of this proof and Proposition 9(1). Equation (22) is obtained using Proposition 37(4) as $\frac{a}{b} > 1 \implies \mu > 0$. The final equality holds as $U_{\mu+1} = a$ and $U_{\mu} = b$.

- (b) Case $\frac{a}{b} > \phi$. From Lemma 35, $\mu = 2\tilde{\mu} + 1$ where $\tilde{\mu} = \min \{k \geq 0 : \frac{a}{b} \geq E_k\}$. Then, for all $n \geq a + b$,

$$r(n; a, b)^* = r(n; U_{2\tilde{\mu}+2}, U_{2\tilde{\mu}+1})^* \\ = \ell(n - U_{2\tilde{\mu}+1}; U_{2\tilde{\mu}}, U_{2\tilde{\mu}+1})^* \quad (23)$$

$$= r(n - \sum_{i=2\tilde{\mu}}^{2\tilde{\mu}+1} U_i; U_{2\tilde{\mu}}, U_{2\tilde{\mu}-1})^* \quad (24)$$

$$= \ell(n - \sum_{i=1}^{2\tilde{\mu}+1} U_i; U_0, U_1)^* \quad (25)$$

$$= r(n - \sum_{i=1}^{2\tilde{\mu}+1} U_i - U_1; U_0, 0)^* \quad (26)$$

$$= (n - \sum_{i=1}^{\mu} U_i - U_1) \quad (27)$$

$$= n - U_{\mu+2} + U_0 \quad (28)$$

$$= n - (a + b) + U_0.$$

Equation (23) follows using Observation 33(1) and Lemma 41(2) and the fact that, at each step, the heap size is at least the sum of the players' wealths, as $n \geq a + b$, and both decrease equally in each iteration. By repeating this process, we get Equations (24), (25) and (26). Equation (27) holds using item (2) of this proof and Equation (28) follows using Proposition 37(4). The last equality holds as $U_{\mu+1} = a$ and $U_{\mu} = b$.

This concludes the proof. □

Intuitively, having more wealth does not disadvantage a LITTLE JOHN player. The next theorem establishes this by comparing the stops of two LITTLE JOHN games that

differ only in the wealth of one of the players; we refer to this property as *stop monotonicity*. Later, in Lemma 47, we prove an analogous monotonicity result for ROBIN HOOD. The key difference is that, in the LITTLE JOHN setting, the two games being compared need not be related as options of the same game and differ only in one player's wealth, whereas in the ROBIN HOOD setting, the comparison is between two options of the same game, which therefore differ in both the heap size and the wealth of one of the players.

Theorem 43 (Little John Stop Monotonicity). *Consider $n, a, b \in \mathbb{N} \cup \{0\}$. Then, for all $n \geq a + b + 1$,*

1. $r(n; a, b)^* \leq r(n; a + 1, b)^*$;
2. $r(n; a, b + 1)^* \leq r(n; a, b)^*$;
3. $\ell(n; a, b)^* \leq \ell(n; a + 1, b)^*$;
4. $\ell(n; a, b + 1)^* \leq \ell(n; a, b)^*$.

Proof. We prove the first item and the others are similar.

Let $(U_i)_{i \geq 0}$ be the unique MP-sequence such that for some $\mu \geq 0$, $U_\mu = \min\{a, b\}$ and $U_{\mu+1} = \max\{a, b\}$. The uniqueness follows by Proposition 32.

Similarly, let $(V_i)_{i \geq 0}$ be the unique MP-sequence such that for some $\nu \geq 0$, $V_\nu = \min\{a + 1, b\}$ and $V_{\nu+1} = \max\{a + 1, b\}$. Let

$$R_U = r(n; a, b)^*, \quad R_V = r(n; a + 1, b)^*.$$

We need to prove $R_U \leq R_V$. There are 5 cases based on the relative values of a and b with respect to the golden ratio. We will use the Stop Values Lemma 42 several times here.

- (a) If $a = b = 0$, then, by Lemma 42(1, 2), $R_U = 0 \leq n = R_V$ for all $n \geq 0$.
- (b) If $a > 0 = b$, then, by Lemma 42(2), $R_U = n = R_V$ for all $n \geq 0$.
- (c) If $b > 0 = a$, then, $R_U = -n \leq R_V$ for all $n \geq 0$, where the first equality holds using Lemma 42(2) and second inequality holds using Proposition 40(2).
- (d) Case $\phi < \frac{a}{b}$. By Lemma 42, we have,

$$R_U = n - (a + b) + U_0, \tag{29}$$

$$R_V = n - (a + 1 + b) + V_0, \tag{30}$$

for all $n \geq a + b + 1$. Now, to compare R_U and R_V , we compare U_0 and V_0 . Recall that $V_{\nu+1} = a + 1 = U_{\mu+1} + 1$ and $V_\nu = b = U_\mu$, as $\frac{a}{b} > \phi$. Thus, by Proposition 38, we have, for all $0 \leq k \leq \min\{\mu, \nu\}$,

$$V_{\nu-k} = U_{\mu-k} + \bar{P}_k. \tag{31}$$

Now, by Lemma 35, $\mu = 2\tilde{\mu} + 1$ and $\nu = 2\tilde{\nu} + 1$ where

$$\tilde{\mu} = \min \left\{ i \geq 0 : \frac{a}{b} \geq E_i \right\} \text{ and } \tilde{\nu} = \min \left\{ i \geq 0 : \frac{a+1}{b} \geq E_i \right\}.$$

Recall that the sequence $(E_i)_{i \geq 0}$, where $E_i = P_{2i+3}/P_{2i+2}$, is decreasing. Therefore, $\tilde{\nu} \leq \tilde{\mu}$ and hence, $\nu \leq \mu$. Now,

$$\begin{aligned} R_V - R_U &= V_0 - 1 - U_0 && \text{(by Eq (29)-(30))} \\ &= U_{\mu-\nu} + \bar{P}_\nu - 1 - U_0 && \text{(by Eq (31))} \\ &= U_{2(\tilde{\mu}-\tilde{\nu})} + \bar{P}_{2\tilde{\nu}+1} - 1 - U_0 && (\nu = 2\tilde{\nu} + 1, \mu = 2\tilde{\mu} + 1) \\ &\geq U_0 + P_{2\tilde{\nu}+1} - 1 - U_0 && (32) \\ &\geq 0. && (P_{2\tilde{\nu}+1} \geq 1) \end{aligned}$$

Equation (32) follows using the following facts:

- if $\tilde{\mu} = \tilde{\nu}$, then $U_{2(\tilde{\mu}-\tilde{\nu})} = U_0$;
- if $\tilde{\mu} > \tilde{\nu}$, then by Observation 33, $U_{2(\tilde{\mu}-\tilde{\nu})} \geq U_2 > U_0$;
- $\bar{P}_{2\tilde{\nu}+1} = P_{2\tilde{\nu}+1}$ by Definition 36.

(e) Case $\frac{a}{b} < \phi < \frac{a+1}{b}$. By Lemma 42, we have,

$$\begin{aligned} R_U &= \begin{cases} a - n & \text{if } a \leq b ; \\ a + b - n - U_0 & \text{if } a > b , \end{cases} \\ R_V &= n - (a + 1 + b) + V_0, \end{aligned}$$

for all $n \geq a + b + 1$. Hence $R_U < 0 < R_V$.

(f) Case $\frac{a+1}{b} < \phi$. By Lemma 42, we have,

$$R_U = \begin{cases} a - n & \text{if } a \leq b ; \\ a + b - n - U_0 & \text{if } a > b , \end{cases} \quad (33)$$

$$R_V = \begin{cases} a + 1 - n & \text{if } a + 1 \leq b ; \\ a + 1 + b - n - V_0 & \text{if } a + 1 > b , \end{cases} \quad (34)$$

for all $n \geq a + b + 1$. The following subcases arise depending on the relative values of a and b .

- (i) If $a + 1 \leq b$, then $R_U = a - n < a + 1 - n = R_V$.
- (ii) If $a = b (\geq 1)$, then $V_{\nu+1} = a + 1$ and $V_\nu = b$ and consequently, $V_{\nu-1} = 1$. In the case where $b = 1$, we have $V_{\nu-1} \geq V_\nu$, and by Observation 33, it follows that

$\nu = 1$. Otherwise, $V_{\nu-2} = b - 1 \geq 1 = V_{\nu-1}$ and which implies $\nu = 2$. In both cases, we have $V_0 \leq b$ and therefore,

$$\begin{aligned} R_V &= a + b + 1 - n - V_0 \\ &\geq a - n + b + 1 - b \\ &\geq a - n = R_U. \end{aligned}$$

(iii) If $a > b$, then we must compare U_0 and V_0 in order to compare R_U and R_V . We know $V_{\nu+1} = a + 1 = U_{\mu+1} + 1$ and $V_\nu = b = U_\mu$. Thus, by Proposition 38, for all $0 \leq k \leq \min\{\nu, \mu\}$ we have,

$$V_{\nu-k} = U_{\mu-k} + \bar{P}_k. \tag{35}$$

Now, by Lemma 35, $\nu = 2\hat{\nu}$ and $\mu = 2\hat{\mu}$ where,

$$\hat{\mu} = \min \left\{ i \geq 0 : \frac{a}{b} \leq O_i \right\}, \quad \hat{\nu} = \min \left\{ i \geq 0 : \frac{a+1}{b} \leq O_i \right\}.$$

Recall that the sequence $(O_i)_{i \geq 0}$, where $O_i = P_{2i+2}/P_{2i+1}$, is increasing. Therefore, $\hat{\mu} \leq \hat{\nu}$ and consequently, $\mu \leq \nu$. Thus, we have,

$$\begin{aligned} R_U - R_V &= V_0 - 1 - U_0 && \text{(by Eq (33)-(34))} \\ &= V_0 - 1 - (V_{\nu-\mu} - \bar{P}_\mu) && \text{(by Eq (35))} \\ &= V_0 - V_{2(\hat{\nu}-\hat{\mu})} + \bar{P}_{2\hat{\mu}} - 1 && (\mu = 2\hat{\mu}, \nu = 2\hat{\nu}) \\ &\leq V_0 - V_0 - P_{2\hat{\mu}} - 1 && \text{(36)} \\ &< 0. && (P_{2\hat{\mu}} \geq 0) \end{aligned}$$

Equation (36) follows using the following facts:

- if $\hat{\nu} = \hat{\mu}$, then $V_{2(\hat{\nu}-\hat{\mu})} = V_0$;
- if $\hat{\nu} > \hat{\mu}$, then by Observation 33(1), $V_{2(\hat{\nu}-\hat{\mu})} \geq V_2 > V_0$;
- $\bar{P}_{2\hat{\mu}} = -P_{2\hat{\mu}}$ by Definition 36.

This concludes the proof. □

Let us restate this result as we often will use it.

Corollary 44. *Consider $n, a, b \in \mathbb{N}$. Then, for all $n \geq a+b$, $r(n; a, b)^* \leq r(n; a+1, b-1)^*$.*

Proof. By Theorem 43, we have $r(n; a, b)^* \leq r(n; a+1, b)^* \leq r(n; a+1, b-1)^*$. □

We understand the stops of LITTLE JOHN. Next, we will show that the stops of LITTLE JOHN and ROBIN HOOD are the same.

8 Little John guides Robin Hood

For large heap sizes, ROBIN HOOD resembles LITTLE JOHN. Robin Hood is wise when he listens to Little John.

Theorem 45 (Wise Robin Hood). *Consider $n, a, b \in \mathbb{N}_0$. Then, for all $n \geq a + b$, the stops of $(n; a, b)$ are the same as the stops of $(n; a, b)^*$.*

Proof. We prove the statement only for the Right stops, as, for any game G , $\ell(G) = -r(-G)$.

We prove this using induction. Before initiating the induction steps, we first verify the statement for the cases where at least one of a or b is zero. Suppose $n \geq 0$.

1. If $a = 0 = b$, then $(n; a, b) = 0 = (n; a, b)^*$;
2. If $a > 0 = b$, then $(n; a, b) = n = (n; a, b)^*$;
3. If $b > 0 = a$, then $(n; a, b) = -n = (n; a, b)^*$.

In all these cases, the stops of both games are equal as their game values are equal. Since the choice of n was arbitrary, the statement holds true for all $n \geq 0$ in all these cases.

We now proceed to the induction on $a + b$. The base case of induction, $a + b = 1$, is already proven.

Suppose $n \geq a + b$. To proceed, we consider three cases based on the relative values of a and b . Note that the scenarios where $a = 0$, or $b = 0$, or both $a = b = 0$ have already been resolved. Therefore, we now focus solely on cases where $a, b > 0$.

1. If $a \leq b$, then, $r(n; a, b)^* = -(n - a)$ by Lemma 42. Next, we compute $r(n; a, b)$ to verify the equivalence. By Theorem 25(5) and Proposition 23,

$$r(n; a, b) = \min_{i \in [a]} \ell(n - i; a - i, b)$$

We know, by Lemma 42(2) and Proposition 9(1), $\ell(n - a; 0, b) = -(n - a)$. Moreover, for all $1 \leq i \leq a - 1$, we have,

$$\begin{aligned} \ell(n - i; a - i, b) &= \ell(n - i; a - i, b)^* && \text{(by induction)} \\ &= r(n - a; a - i, b - (a - i))^* \\ &\geq -(n - a). && \text{(by Prop 40(2))} \end{aligned}$$

Hence, $\min_{i \in [a]} \ell(n - i; a - i, b) = -(n - a)$. This completes the proof of this case.

2. If $b \leq \frac{a}{2}$, then

$$\begin{aligned} r(n; a, b) &= \min_{i \in [b]} \ell(n - i; a - i, b) && \text{(by Thm 25(5))} \\ &= \min_{i \in [b]} \ell(n - i; a - i, b)^* && \text{(by induction)} \end{aligned}$$

$$\begin{aligned}
&= \min_{i \in [b]} -r(n - i; b, a - i)^* && \text{(by Prop 9(1))} \\
&= \min_{i \in [b]} (n - i - b) && (37) \\
&= n - 2b.
\end{aligned}$$

Equation (37) follows using Lemma 42(3) as $a - i \geq b$ for all $i \in [b]$ and $n \geq a + b$.

Now, we compute $r(n; a, b)^*$.

$$\begin{aligned}
r(n; a, b)^* &= \ell(n - b; a - b, b)^* && \text{(by Lem 41(2))} \\
&= -r(n - b; b, a - b)^* && \text{(by Prop 9(1))} \\
&= n - 2b. && (38)
\end{aligned}$$

Equation (38) holds using Lemma 42(3) as $a - b \geq b$ and $n \geq a + b$. This concludes the proof for this case.

3. If $\frac{a}{2} < b < a$, then,

$$\begin{aligned}
r(n; a, b) &= \min_{i \in [b]} \ell(n - i; a - i, b) \\
&= \min_{i \in [b]} -r(n - i; b, a - i)^*. && \text{(by induction and Prop 9(1))}
\end{aligned}$$

Now, we divide the range of i in two parts, in one, $a - i \geq b$ and in the other, $a - i < b$. Let $A = \{i \in [b] : a - i \geq b\}$ and $B = [b] \setminus A$. Note that B cannot be empty by the assumption. Thus, $A = \{1, \dots, a - b\}$ and $B = \{a - b + 1, \dots, b\}$. Then,

$$\begin{aligned}
r(n; a, b) &= \min \left\{ \min_{i \in A} (-r(n - i; b, a - i)^*), \min_{i \in B} (-r(n - i; b, a - i)^*) \right\} \\
&= \min \left\{ \min_{i \in A} (n - i - b), \min_{i \in B} (-\ell(n - a; b - (a - i), a - i)^*) \right\} \\
&= \min \left\{ n - (a - b) - b, \min_{i \in B} r(n - a; a - i, b - (a - i))^* \right\} \\
& && \text{(by definition of } A) \\
&= \min \{n - a, r(n - a; a - b, b - (a - b))^*\} && \text{(by Cor 44)} \\
&= r(n - a; a - b, b - (a - b))^* && \text{(by Prop 40(2))} \\
&= \ell(n - b; a - b, b)^* && (39) \\
&= r(n; a, b)^*. && (40)
\end{aligned}$$

Equation (39) holds because, for $a - b < b$, we have $\ell(n - b; a - b, b)^* = r(n - a; a - b, b - (a - b))^*$ by Lemma 41(2). Similarly, Equation (40) holds since, for $b < a$, we have $r(n; a, b)^* = \ell(n - b; a - b, b)^*$.

Since, the choice of n was arbitrary from the set $\{a + b, a + b + 1, \dots\}$, all these cases hold for all $n \geq a + b$. This completes the proof. \square

The next results focus on the geometric aspects of $\text{Therm}(n; a, b)^*$. As we indicated in the Introduction, the typical behavior will depend of the ‘wealth ratio’ a/b . By convention we choose $a \geq b$. Therefore the Left option will (trivially) be a Number, and so all of our efforts will concern the Right options. Let the *wealth ratio* of a Right option be $w_b := (a - b)/b$.

Theorem 46 (Little John Thermographs). *For fixed integers $a, b > 0$, let $G = (n; a, b)^*$. Then, for n sufficiently large,*

1. $G \in \mathcal{LT}$, if $\frac{a}{b} > \phi$;
2. $G \in \mathcal{RT}$, if $\frac{a}{b} < \phi^{-1}$;
3. $G \in \mathcal{DT}$, otherwise.

Proof. We induct on $a + b$. Without loss of generality, consider $a \geq b$, as the thermograph of $(n; a, b)^*$ is the mirror image of thermograph of $(n; b, a)^*$.

Base case 1: If $a + b = 2$, then $a/b = 1$ and $G = (n; 1, 1)^* = \{n - 1 \mid 1 - n\}$. Hence, $G \in \mathcal{DT}$ for all $n \geq 2$, as in Figure 7a.



(a) The thermograph of $(n; 1, 1)^*$.

(b) The thermograph of $(n; 2, 1)^*$.

Figure 7: Thermographs of $(n; a, b)^*$ for small a and b .

Base case 2: If $a + b = 3$, then $a/b = 2$ and $G = (n; 2, 1)^* = \{n - 1 \mid \{n - 2 \mid 2 - n\}\}$. Thus, for all $n \geq 3$, $G \in \mathcal{LT}$, as in Figure 7b.

Since $a \geq b$, for all $n \geq b$, the Left option of $G = (n; a, b)^*$ is the Number $n - b$, and thus its thermograph is a mast at $n - b$. Therefore, the contribution of $\text{Therm}(G^L)$ to $\text{Therm}(G)$ is a small left wall of slope -1 .

Next we analyze the contributions of the Right option for all sufficiently large n . This will require an induction argument, and we have analyzed the base cases above. Suppose that, for sufficiently large n , the statement holds for the thermographs of the options of $G = (n; a, b)^*$. To prove that the statement holds for the thermograph of G , we take different cases based on the ratio of the players’ wealths.

(1) $a/b > \phi$: In this case $G^R = (n - b; a - b, b)^*$. We must prove that, for sufficiently large n , $G \in \mathcal{LT}$. The thermograph of G^R depends on the wealth ratio $w_b = (a - b)/b$. Since $a/b > \phi$, then $(a - b)/b > \phi^{-1}$. Hence, by induction, $G^R \in \mathcal{DT} \cup \mathcal{LT}$. Thus, $\text{lw}(G^R)$ has slope -1 . In either case, since we must prove that $\text{RW}(G)$ is a vertical line, we must verify that the contribution to $\text{Therm}(G)$ from $\text{Therm}(G^L)$ meets the contribution from small left wall of $\text{Therm}(G^R)$.

(1A) $w_b > \phi$: In this case, $G^R \in \mathcal{LT}$ (see Figure 8a). The small left wall of $\text{Therm}(G^R)$ ends at the temperature $t(G^R) = \ell(G^R) - r(G^R)$. Recall that the contribution from $\text{Therm}(G^L)$ to $\text{Therm}(G)$ is a small left wall with slope -1 . It intersects the left tilted $\text{lw}(G^R)$ if and only if

$$\begin{aligned} \ell(G) - r(G) &= \ell(G) - \ell(G^R) \\ &\leq \ell(G^R) - r(G^R), \end{aligned} \tag{41}$$



(a) The wealth ratio of G^R is $\frac{a-b}{b} > \phi$. (b) The wealth ratio of G^R is $\phi^{-1} < \frac{a-b}{b} < \phi$.

Figure 8: Thermographs of $G = (n; a, b)^*$ and its options when $\frac{a}{b} > \phi$. In each figure, the Red dashed line represents $\text{Therm}(G^L)$, blue dashed lines represent $\text{Therm}(G^R)$, black dotted line indicates the temperature of G^R and $\text{Therm}(G)$ is given by solid purple lines.

where the first equality holds since there is only one Right option. For the second inequality, we need to find $r(G^R)$ and $\ell(G^R)$. For this purpose, let $(U_i)_{i \geq 0}$ be an MP-sequence with $U_{\mu+1} = a - b$ and $U_\mu = b$ for some $\mu \geq 0$. Then, by Lemma 42, we have $\ell(G) = n - b$, $\ell(G^R) = n - 2b$, and $r(G^R) = n - b - (a - b + b) + U_0$ which implies

$$\ell(G) - \ell(G^R) = b \text{ and } \ell(G^R) - r(G^R) = a - b - U_0.$$

Hence, by Equation (41), it suffices to prove that $U_0 \leq a - 2b$. Note that $a - 2b > 0$ and therefore $U_{\mu-1} = a - 2b$ and $\mu \geq 1$. Thus, the problem reduced to show that $U_0 \leq U_{\mu-1}$, for all $\mu \geq 1$.

If $\mu \geq 3$ or $\mu = 1$, then, by Observation 33(1), $U_0 \leq U_{\mu-1}$. If $\mu = 2$, then $U_0 = 3b - a$ and $U_{\mu-1} = U_1 = a - 2b$. Now, by applying Observation 33(1), we get $3b - a \geq a - 2b$. This implies $w_b \leq 1.5$, which is a contradiction to our assumption.

(1B) $w_b < \phi$: In this case, $G^R \in \mathcal{DT}$ (see Figure 8b). Thus, $\text{lw}(G^R)$ ends at the temperature $t(G^R) = \frac{1}{2}(\ell(G^R) - r(G^R))$. Recall that the contribution from $\text{Therm}(G^L)$ to $\text{Therm}(G)$ is a small left wall of slope -1 , and it meets the left rotated $\text{lw}(G^R)$ if and only if

$$\ell(G) - \ell(G^R) \leq \frac{1}{2}(\ell(G^R) - r(G^R)). \tag{42}$$

By Lemma 42, $\ell(G^R) - r(G^R) = 2n + c$ and $\ell(G) - \ell(G^R) = d$ for some constants c and d , with respect to n . Hence, for all sufficiently large n , $\ell(G) - \ell(G^R) \leq \frac{1}{2}(\ell(G^R) - r(G^R))$.

(2) $1 < \mathbf{a/b} < \phi$: In this case, we must prove that, for all sufficiently large n , $G \in \mathcal{DT}$. Here $G = \{n-b \mid (n-b; a-b, b)^*\}$, and thus, by $\frac{a-b}{b} < \phi^{-1}$, by induction, for n sufficiently large, $G^R \in \mathcal{RT}$ (see Figure 9a). Hence, the contribution to $\text{Therm}(G)$ from $\text{Therm}(G^R)$ is a small right wall with slope $+1$ and the contribution from $\text{Therm}(G^L)$ is a small left wall with slope -1 . Hence, for all sufficiently large n , $G \in \mathcal{DT}$ (as in Figure 9a). In this case, the lower bound for n is $a + b$, by using Lemma 42.




Figure 9: Thermographs of G when $\frac{a}{b} < \phi$. In each figure, the Red dashed line represents $\text{Therm}(G^L)$, blue dashed lines represent $\text{Therm}(G^R)$, and $\text{Therm}(G)$ is given by purple solid lines.

(3) $\mathbf{a/b} = 1$: If $n > a = b$, obviously $G \in \mathcal{DT}$.

Thus LITTLE JOHN's tent structures have been established. \square

The term ‘‘orthodox option’’ is often used in the context of thermograph plots. Such options contribute to the thermograph. The key to the ROBIN HOOD thermographs depends on its orthodox link with LITTLE JOHN. Later in Theorem 49 we will exploit this connection between the thermographs of ROBIN HOOD and LITTLE JOHN. The next lemma gives the order in which the thermographs of the options of ROBIN HOOD are placed on the graph, by providing the stop monotonicity for these options.

Lemma 47 (Options Stop Monotonicity). *Let $n, a, b > 0$ be integers and consider the ROBIN HOOD game $(n; a, b)$, with Left and Right options $L_i = (n - i; a, b - i)$ and $R_i = (n - i; a - i, b)$, respectively, where $i \in [b]$. For all sufficiently large heap sizes n ,*

1. $\ell(L_1) = \ell(L_2) = \dots = \ell(L_b) = n - b$;
2. $r(L_1) \leq r(L_2) \leq \dots \leq r(L_b) = n - b$;
3. $\ell(R_1) \geq \ell(R_2) \geq \dots \geq \ell(R_b)$;
4. $r(R_1) \geq r(R_2) \geq \dots \geq r(R_b)$.

Proof. Without loss of generality, consider $a \geq b$. Now, we compare the stops of the Left options of G using LITTLE JOHN Stop Monotonicity, Theorem 43 and Corollary 44. We use the following results several times in the proof:

- By Theorem 45, the stops of Robin Hood and Little John are the same;
- Lemma 42 gives the exact stop values.

For any fixed $i \in [b]$, we have:

$$\begin{aligned} \ell(L_i) &= \ell(n - i; a, b - i)^* \\ &= r(n - b; a, 0)^* \end{aligned} \tag{43}$$

$$= n - b \tag{44}$$

$$\begin{aligned} r(L_i) &= r(n - i; a, b - i)^* \\ &= \begin{cases} n - b & \text{if } i = b; \\ \ell(n - b; a - b + i, b - i)^* & \text{otherwise.} \end{cases} \end{aligned} \tag{45}$$

We have used that $b - i < a$ for all $i \in [b]$. Therefore, by Corollary 44,

$$r(L_1) \leq r(L_2) \leq \dots \leq r(L_{b-1}). \tag{46}$$

And, by Proposition 40,

$$r(L_{b-1}) \leq n - b. \tag{47}$$

Thus, equations (44)-(47) together prove items (1) and (2).

For item (3), we compare the Left stops of the Right options. By Theorem 45, for all $i \in [b]$, we have

$$\ell(R_i) = \ell(n - i; a - i, b)^* \tag{48}$$

$$= \begin{cases} n - b - i, & \text{if } a - i \geq b; \\ r(n - a; a - i, b - a + i)^*, & \text{otherwise.} \end{cases} \tag{49}$$

In Equation (49), the behavior of $\ell(R_i)$ changes when i increases such that $a - i < b$. So, we define α as the minimum natural number i such that $a - i < b$, i.e., $\alpha := \min \{i \in \mathbb{N} : a - i < b\}$. Note that $\alpha = a - b + 1$. Put differently, α is the fewest tokens Right needs to remove in $(n; a, b)$ to make Left's wealth less than Right's wealth. But, if $\alpha > b$, then Right cannot do this. Therefore, let us first assume $\alpha \leq b$. Now, we rewrite Equation (49) as follows:

$$\ell(R_i) = \begin{cases} n - b - i, & \text{if } i < \alpha; \\ r(n - a; a - i, b - a + i)^*, & \text{if } i \geq \alpha. \end{cases} \tag{50}$$

Then, by Corollary 44, we have

$$\ell(R_\alpha) \geq \ell(R_{\alpha+1}) \geq \dots \geq \ell(R_b). \tag{51}$$

Now, we compare $\ell(R_{\alpha-1})$ with $\ell(R_\alpha)$,

$$\ell(R_{\alpha-1}) = n - b - \alpha + 1 \tag{by Eq (50)}$$

$$\begin{aligned}
&= n - a \\
&\geq r(n - a; a - \alpha, b - a + \alpha)^* && \text{(by Prop 40)} \\
&= \ell(R_\alpha). && (52)
\end{aligned}$$

Equations (50)-(52) prove item (3) when $\alpha \leq b$. If $\alpha > b$, then the proof of item 3 is complete because $\ell(R_i) = n - b - i$ for all $i \in [b]$.

For item (4), we compare the Right stops of the Right options of $(n; a, b)$.

$$r(R_i) = r(n - i; a - i, b)^* \tag{53}$$

$$= \begin{cases} \ell(n - i - b; a - i - b, b)^*, & \text{if } a - i > b; \\ -(n - a), & \text{otherwise.} \end{cases} \tag{54}$$

$$= \begin{cases} n - i - 2b, & \text{if } a - i \geq 2b; \\ r(n - a; a - i - b, 2b - a + i)^*, & \text{if } b < a - i < 2b; \\ -(n - a), & \text{if } a - i \leq b. \end{cases} \tag{55}$$

Define $\beta := \min \{i \in \mathbb{N} : a - i < 2b\}$ and $\gamma := \min \{i \in \mathbb{N} : a - i \leq b\}$. Note that $\beta = a - 2b + 1$ and $\gamma = a - b$. If $\gamma \leq b$, then we rewrite Equation (55) as:

$$r(R_i) = \begin{cases} n - i - 2b, & \text{if } i < \beta; \\ r(n - a; a - i - b, 2b - a + i)^*, & \text{if } \beta \leq i < \gamma; \\ -(n - a), & \text{if } \gamma \leq i \leq b. \end{cases} \tag{56}$$

Thus, we have

$$r(R_1) \geq r(R_2) \geq \dots \geq r(R_{\beta-1}), \tag{57}$$

$$r(R_\beta) \geq r(R_{\beta+1}) \geq \dots \geq r(R_{\gamma-1}), \tag{58}$$

$$r(R_\gamma) \geq r(R_{\gamma+1}) \geq \dots \geq r(R_b), \tag{59}$$

where Equation (58) follows by Corollary 44. Now, by Proposition 24,

$$r(R_{\beta-1}) = (n - a) \geq r(R_\beta), \tag{60}$$

$$r(R_{\gamma-1}) \geq -(n - a) = r(R_\gamma). \tag{61}$$

Equations (57)-(61) complete the proof when $\gamma \leq b$. If $\gamma > b$ and $\beta \leq b$, then Equations (57),(58) and (60) complete the proof. The proof of case when $\beta > b$ follows by Equation (57). \square

9 A solution for the Pingala Era wetland tribes

The toolbox is now complete, and we arrive at the main theorem, in terms of thermographs. At last, in this section, we revisit Theorem 1 by including a short proof,

interpreting ROBIN HOOD's thermographs in terms of mean values and temperatures. To simplify reading the proof we locally abbreviate some of our standard notation.

Notation. Consider a ROBIN HOOD game $G = (n; a, b)$. For $i \in [b]$, let $L^i = (n - i; a, b - i)$ represent the Left options and let $R_i = (n - i; a - i, b)$ represent the Right options. The reason for the super- and sub-scripts is the following short hand notation for the stops of these options:

- Let r^i and r_i denote the Right stops of L^i and R_i , respectively, and let ℓ^i and ℓ_i denote the Left stops of L^i and R_i , respectively;
- Let RW^i and RW_i denote the large right walls of L^i and R_i , respectively, and let LW^i and LW_i denote the large left walls of L^i and R_i , respectively;
- Similarly, let rw^i , rw_i , lw^i and lw_i denote the small walls.

This use of sub- and super-scripts can be generalized to any function on R_i and L^i respectively; for example $m(R_b) = m_b$ and $m(L^a) = m^a$, etc. Moreover, when the options are penalized by p , we write $r^i(p)$ and $\ell^i(p)$ for the Right and Left stops of L^i penalized by p , respectively, and $r_i(p)$ and $\ell_i(p)$ for the Right and Left stops of R_i penalized by p , respectively. Aligning with these notations, denote a typical $G^R \in \mathcal{DT}$ by R_δ , with Left and Right stops ℓ_δ and r_δ , respectively, and denote a typical $G^L \in \mathcal{DT}$ by L^δ , with Left and Right stops ℓ^δ and r^δ , respectively. Similarly, denote $G^R \in \mathcal{ST}$ by R_σ , with stops ℓ_σ and r_σ , and denote $G^L \in \mathcal{ST}$ by L^σ , with stops ℓ^σ and r^σ .

We make use of a partial order of large left and right walls.

Definition 48 (Wall Partial Order). Let $f, g : \mathbb{D}^+ \rightarrow \mathbb{D}$, and let $F = \{(f(y), y) \mid y \in \mathbb{D}^+\}$ and $G = \{(g(y), y) : y \in \mathbb{D}^+\}$. Then $F \geq G$ if, for all $y \in \mathbb{D}^+$, $f(y) \geq g(y)$.

Thus, for example $LW_i \geq LW_j$ if, for all $p \in \mathbb{D}^+$, $\ell_i(p) \geq \ell_j(p)$. We will see that LITTLE JOHN and ROBIN HOOD have the same mean values and temperatures for large heaps, and the reason for that is that they have the same thermographs.

Theorem 49 (Robin Hood Thermographs). *Let $a, b \geq 0$ be integers. Then, for any sufficiently large heap size n , $\text{Therm}(n; a, b) = \text{Therm}(n; a, b)^*$.*

Proof. Let $G = (n; a, b)$ and let $H = (n, a, b)^*$. Our goal is to demonstrate that, for n sufficiently large, the thermographs of H and G are identical. Thus, it suffices to show that, for large n , the thermograph of G solely depends on the Little John options.

We induct on $a + b$. Without loss of generality, consider $a \geq b$, as $\text{Therm}(n; a, b)$ is the mirror image of $\text{Therm}(n, b, a)$.

If $a = b = 0$, then $G = 0 = H$ and if $a > b = 0$, then $G = n = H$. Thus, the result holds in these cases, so suppose both a and b are non-zero.

If $a + b = 2$, then $(n; 1, 1) = \{(n - 1) \mid -(n - 1)\} = (n; 1, 1)^*$. Hence, the thermographs of G and H are the same.

Now, suppose that the statement holds for the options of G , for all sufficiently large heap sizes n .

We begin by examining the Left options of G and their corresponding thermographs. By Option Stop Monotonicity, Lemma 47, we have

$$\ell^1 = \ell^2 = \dots = \ell^b = n - b = r^b \geq r^{b-1} \geq \dots \geq r^1. \quad (62)$$

This implies that L^b is the option with the Largest Left stop and the thermograph of L^b is a mast at $n - b$. Hence, by Lemma 21, the left wall of $\text{Therm}(G)$ does not depend on any Left option other than L^b , which is the Little John option.

For the rest of the proof, we demonstrate that, for large n , $\text{RW}(G)$ does not depend on any Right option other than the Little John Right option R_b . That is, we prove that, for large n , for all i , $\text{LW}_b \leq \text{LW}_i$. By induction, the thermographs of the Right options are the same as those of the corresponding LITTLE JOHN thermographs, and consequently they depend on the option's wealth ratio $w_i = \frac{a-i}{b}$, for $i \in [b]$. Recalling Theorem 46, we have $R_i \in \mathcal{RT}$ if $w_i < \phi^{-1}$, $R_i \in \mathcal{DT}$ if $\phi^{-1} < w_i < \phi$, and $R_i \in \mathcal{LT}$, otherwise.

We take different cases based on G 's wealth ratio $\frac{a}{b}$.

(1) $\frac{a}{b} > \phi$: As $i \in [b]$, the wealth ratios, $w_i = \frac{a-i}{b}$, of the Right options fall within the interval $(\phi^{-1}, \frac{a}{b})$. Hence, by induction, their thermographs are either \mathcal{DT} or \mathcal{LT} . When the ratio w_i drops (as i increases) below ϕ , the Right options' thermographs change from \mathcal{LT} to \mathcal{DT} . Therefore, we define α as the smallest Right removal (i) for which $R_i \in \mathcal{DT}$, i.e.,

$$\alpha = \min \{i \geq 1 : w_i < \phi\}.$$

By this definition, if $\alpha > b$, then the thermographs of all Right options are left single tents. Otherwise, at least one of the thermographs is a double tent. So, we take sub-cases based on α .

Recall, by Lemma 47,

$$\ell_1 \geq \ell_2 \geq \dots \geq \ell_b, \text{ and} \quad (63)$$

$$r_1 \geq r_2 \geq \dots \geq r_b. \quad (64)$$

(1A) $\alpha > b$: Let $R_\sigma \in \mathcal{LT}$ be any Right option other than $R_b \in \mathcal{LT}$. We must prove that $\text{LW}_b \leq \text{LW}_\sigma$. Clearly $\ell_b \leq \ell_\sigma$, by the monotonicity property in Equation (63), so $\text{LW}_b \not\leq \text{LW}_\sigma$.

Assume, for a contradiction, that $\text{LW}_b \not\leq \text{LW}_\sigma$. Since $\ell_b \leq \ell_\sigma$, this assumption implies that LW_σ crosses LW_b at some point, that is, there is a penalty p such that $\ell_b(p) > \ell_\sigma(p)$. The only way this can occur is if lw_σ intersects LW_b above t_b (see Figure 10c). In this case, since both $R_b, R_\sigma \in \mathcal{LT}$, we must have $\text{RW}_\sigma < \text{RW}_b$. This would imply $r_\sigma < r_b$, which contradicts the monotonicity property in Equation (64). Hence, we are left with the situations in Figures 10a and 10b that both confirm that $\text{LW}_b \leq \text{LW}_\sigma$.

(1B) $\alpha \leq b$: If $\alpha \leq i \leq b$, then by induction and definition of α , $R_i \in \mathcal{DT}$. Otherwise, if $i < \alpha$, then $R_i \in \mathcal{LT}$. Therefore, in this case, we must compare $R_b \in \mathcal{DT}$ with both $R_\sigma \in \mathcal{LT}$ and $R_\delta \in \mathcal{DT}$. We start with $R_b \in \mathcal{DT}$ and $R_\sigma \in \mathcal{LT}$.

Claim: If $R_b \in \mathcal{DT}$ and $R_\sigma \in \mathcal{LT}$, then, for sufficiently large heap sizes, $m_b \leq m_\sigma$.




Figure 10: Thermographs of Right options of G in case (1A) when both $R_b, R_\delta \in \mathcal{LT}$.

Proof of Claim. By Lemma 42 and Theorem 45, we have

$$\ell_b = n + c_1, \tag{65}$$

$$r_b = -n + c_2, \tag{66}$$

$$r_\sigma = n + c_3, \tag{67}$$

where c_1, c_2 and c_3 are constants with respect to n . Since $R_b \in \mathcal{DT}$, Equations (65)-(66) and Lemma 20 together imply $m_b = (\ell_b + r_b)/2 = (c_1 + c_2)/2$. Similarly, Equation (67) imply $m_\sigma = r_\sigma = n + c_3$ as $R_\sigma \in \mathcal{LT}$. Therefore, since we are taking n to be large, we have

$$m_b \leq m_\sigma. \tag{68}$$




Figure 11: Thermographs of Left walls of G , in case (1B).

Since both small left walls, lw_b and lw_σ , have slope -1 , and $\ell_b \leq \ell_\sigma$ as well as $m_b \leq m_\sigma$, the proof of $LW_b \leq LW_\sigma$ follows similarly to Case (1A). Therefore, we omit the details. This argument does not depend on other properties of the thermographs, such as the relation between temperatures; see Figure 11.

We omit the analysis of $R_b \in \mathcal{DT}$ and $R_\delta \in \mathcal{DT}$, as it parallels that of $R_b \in \mathcal{LT}$ with $R_\sigma \in \mathcal{LT}$ in Case (1A), with the only distinction being the argument that $m(R_b) \leq m(R_\delta)$ due to the shape of the thermographs and the monotonicity of the stops. This completes the proof in this case.

(2) $1 < a/b < \phi$: In this case, the wealth ratios $w_i = (a - i)/b$ for the Right options fall within the interval $(0, \phi)$. By the induction hypothesis, $R_i \in \mathcal{DT} \cup \mathcal{RT}$. Define β as the smallest right removal index for which $R_i \in \mathcal{RT}$, i.e.,

$$\beta = \min \{i \geq 1 : w_i < \phi^{-1}\}.$$

Observe that $\beta \leq b$, since $(a - b)/b = a/b - 1 < \phi^{-1}$. Consequently, $R_\delta \in \mathcal{DT}$ for $\delta < \beta$, while $R_\sigma \in \mathcal{RT}$, if $\sigma \geq \beta$. Hence, we must compare the large left wall of $R_b \in \mathcal{RT}$ with those of both $R_\delta \in \mathcal{DT}$ and $R_\sigma \in \mathcal{RT}$.

We begin by verifying that $LW_b \leq LW_\delta$, for large n .



(a) The case $m_b > m_\delta$ does not appear for large n . (b) The case $m_b \leq m_\delta$ appears for large n .

Figure 12: Thermographs of the Right options R_b and R_δ of G in case (2).

Similar to the claim in case (1B), for large n , we get $m_b \leq m_\delta$. We indicate with a red arrow in Figure 12a that $\text{Therm}(R_b)$ shifts to the right with increasing n , which instead creates a situation as in (b). Indeed, since $R_b \in \mathcal{RT}$, LW_b is a vertical line, and thus $LW_\delta \geq LW_b$ by stop monotonicity and $m_b \leq m_\delta$ (see Figure 12b).

Next we compare $R_b \in \mathcal{RT}$ with $R_\sigma \in \mathcal{RT}$. In this case, the Left walls of both R_b and R_σ are vertical lines determined by their Left stops, i.e., for all $p \geq 0$, $\ell_b(p) = \ell_b$ and $\ell_\sigma(p) = \ell_\sigma$. By Lemma 47, we have $\ell_b \leq \ell_\sigma$. Hence, $LW_b \leq LW_\sigma$.

(3) $a/b = 1$: In this case, $R_b = (n - a; 0, b) = -(n - a)$. Thus, $R_b \in \text{Mast}$. Now, by Lemma 47, R_b is the Right option of G with the smallest Right stop and hence, by applying Lemma 21, $\text{RW}(G)$ does not depend on any other Right option than R_b .

Thus the ROBIN HOOD thermographs are the same as those of LITTLE JOHN. □

Remark 50. The claim in Case (1B) in the proof of Theorem 49, might not hold for smaller heap sizes, and we explain what can happen in terms of thermographs in Figure 13b; our “tent structures” may not survive for small heap sizes.

We can now compute the temperatures and mean values of ROBIN HOOD as stated in the main theorem from the Introduction, Theorem 1.

Theorem 1 (Main Theorem) For fixed positive integers a and b , let $G_n = (n; a, b)$ be instances of ROBIN HOOD. Let $(U_k)_{k \geq 0}$ be the MP-sequence such that $U_q = \min \{a, b\}$




Figure 13: Thermographs of Right options, as in case (1B), for smaller heap sizes.

and $U_{q+1} = \max\{a, b\}$ for some $q \geq 0$. If $n \geq a + b$, then G is a hot game. Moreover, for all sufficiently large heap sizes n ,

$$t(G_n) = \begin{cases} b - U_0 & \text{if } \frac{a}{b} < \phi^{-1}, \\ n - a + \frac{U_0 - b}{2} & \text{if } \phi^{-1} < \frac{a}{b} < 1, \\ n - a & \text{if } \frac{a}{b} = 1, \\ n - b + \frac{U_0 - a}{2} & \text{if } 1 < \frac{a}{b} < \phi, \\ a - U_0 & \text{if } \phi < \frac{a}{b}, \end{cases}$$

and

$$m(G_n) = \begin{cases} a + b - n - U_0 & \text{if } \frac{a}{b} < \phi^{-1}, \\ \frac{U_0 - b}{2} & \text{if } \phi^{-1} < \frac{a}{b} < 1, \\ 0 & \text{if } \frac{a}{b} = 1, \\ \frac{a - U_0}{2} & \text{if } 1 < \frac{a}{b} < \phi, \\ n - (a + b) + U_0 & \text{if } \phi < \frac{a}{b}. \end{cases}$$

Proof. Let G_n^* represent the LITTLE JOHN game associated with G_n , i.e., $G_n^* = (n; a, b)^*$. According to Theorem 49, for sufficiently large n , $\text{Therm}(G_n)$ equals $\text{Therm}(G_n^*)$, implying:

$$t(G_n) = t(G_n^*) \quad \text{and} \quad m(G_n) = m(G_n^*).$$

Thus, the problem reduces to determining the temperature and mean value of G_n^* . As shown in Theorem 46, $\text{Therm}(G_n^*)$ depends on the wealth ratio a/b for sufficiently large n . Therefore, we consider different cases based on the wealth ratio a/b and take n to be large enough that both Theorems 46 and 49 are applicable.

- (1) $a/b < \phi^{-1}$: In this case, by Theorem 46, $G_n^* \in \mathcal{RT}$, and by Lemma 42, the stops are $s(G_n^*) = (a + b - n - U_0, a - n)$. Hence, we get:

$$m(G_n^*) = a + b - n - U_0 \quad \text{and} \quad t(G_n^*) = b - U_0,$$

as per Lemma 20. A similar argument applies if $a/b > \phi$.

(2) $\phi^{-1} < a/b < 1$: In this case, by Theorem 46, $G_n^* \in \mathcal{DT}$, and by Lemma 42, the stops are $s(G_n^*) = (n - (a + b) + U_0, a - n)$. Therefore, we have:

$$m(G_n^*) = \frac{U_0 - b}{2} \quad \text{and} \quad t(G_n^*) = n - a + \frac{U_0 - b}{2},$$

as shown in Lemma 20. A symmetric argument holds if $1 < a/b < \phi$.

(3) $a/b = 1$: In this case, by Theorem 46, $G_n^* \in \mathcal{DT}$, and by Lemma 42, the stops are $s(G_n^*) = (n - a, a - n)$. The desired result follows by applying Lemma 20.

This concludes the proof. □

Open problems

1. Solve the ‘middle region’ of ROBIN HOOD, in terms of thermographs, whenever $0 < \min\{a, b\} < n \leq a + b$.
2. Find explicit bounds on “sufficiently large” heap sizes for the main theorem to apply.
3. Study the Canonical Forms of ROBIN HOOD, and in cases where this is hard, study instead the so-called Reduced Canonical Form [15].
4. Study more instances of WEALTH NIM.
5. Study the outcomes of a ruleset where there is only one global pair of wealths for the entire compound, governing how many tokens may be removed from any single heap. The reduction of the opponent’s wealth would be the same as here. For example if the wealth pair is $(a, b) = (2, 3)$ and there are two heaps of sizes one and two, then, if Left starts, she can win if she removes one token from the second heap and reduce Right’s wealth by one rupee. However, if she plays the Little John move and reduces Right’s wealth by two rupees, and removes the second heap, then she loses. In this perspective, ROBIN HOOD is a local variation, where each heap has an individual wealth pair. Compare with two papers on *local* vs. *global* FIBONACCI NIM [12, 11].
6. Given a number of heaps, and a global wealth for each player, solve the simultaneous play ‘optimal’ assignments of wealth per heap, in the sense of Colonel Blotto [3]. In cases where hotstrat applies, the main results of this paper should guide such assignments, and otherwise one would need better understanding of (reduced) canonical forms.
7. Study the notion of ‘thermograph convergence’, similar to ROBIN HOOD, for other ruleset families.

Acknowledgements

We would like to express our heartfelt gratitude to Prof. Carlos Pereira dos Santos for his invaluable assistance and insightful discussions; in particular concerning Theorem 25.

References

- [1] E. R. Berlekamp, Temperatures of games and coupons. (Ed. U. Larsson) *Games of No Chance 5*, MSRI CUP, Volume 70, 2019: 21–33. [doi:10.1017/9781108641999.002](https://doi.org/10.1017/9781108641999.002).
- [2] E. R. Berlekamp, J. H. Conway and R. K. Guy, *Winning Ways for Your Mathematical Plays*. AK Peters/CRC Press, 2004.
- [3] É. Borel, The theory of play and integral equations with skew symmetric kernels. *Econometrica*, 21(1), (1953), 97–100. (The paper was originally published in French in 1921.) [doi:10.2307/1906946](https://doi.org/10.2307/1906946).
- [4] A. J. Cole and A. J. T. Davie, A game based on the Euclidean algorithm and a winning strategy for it. *The Mathematical Gazette*, volume 53, (1969), 354–357. [doi:10.2307/3612461](https://doi.org/10.2307/3612461).
- [5] O. Gross and R. Wagner, A continuous Colonel Blotto game. *RAND Corporation*, RM-408, Santa Monica, CA, 1950.
- [6] J. P. Grossman and A. N. Siegel, Reductions of partizan games. in (Eds. M. H. Albert and R. J. Nowakowski) *Games of No Chance 3*, MSRI CUP, Volume 56, 2009. [doi:10.1017/CB09780511807251.022](https://doi.org/10.1017/CB09780511807251.022)
- [7] S. Huntemann, and T. Maciosowski, Degrees are useless in SNORT when measuring temperature. *Theoretical Computer Science* (2025): 115597. [doi:10.1016/j.tcs.2025.115597](https://doi.org/10.1016/j.tcs.2025.115597).
- [8] S. Huntemann, and T. Maciosowski, High temperature Domineering positions. *Integers*, vol. 26, (2026). [doi:10.5281/zenodo.18154236](https://doi.org/10.5281/zenodo.18154236).
- [9] S. Huntemann, R. J. Nowakowski, and C. P. Dos Santos. Bounding game temperature using confusion intervals. *Theoretical Computer Science 855* (2021): 43-60. [doi:10.1016/j.tcs.2020.11.015](https://doi.org/10.1016/j.tcs.2020.11.015).
- [10] U. Larsson, R. Meir and Y. Zick, Cumulative Games: Who is the current player? [arXiv:2005.06326](https://arxiv.org/abs/2005.06326).
- [11] U. Larsson, S. Rubinstein-Salzedo, Global Fibonacci Nim. *Int. J. Game Theory* 47, Special Issue on Combinatorial Games, 2018. Invited paper from Combinatorial Game Theory Colloquium, Lisbon 2015 (2018), 595–611. [doi:10.1007/s00182-017-0574-x](https://doi.org/10.1007/s00182-017-0574-x).
- [12] U. Larsson, S. Rubinstein-Salzedo, Grundy values of Fibonacci Nim. *Int. J. Game Theory*, (2015), 45: 617. [doi:10.1007/s00182-015-0473-y](https://doi.org/10.1007/s00182-015-0473-y).
- [13] J. Milnor, Sums of positional games. *Ann. of Math. Stud.* (Contributions to the Theory of Games, H. W. Kuhn and A. W. Tucker, eds.), Princeton University Press (1953), 291–301. [doi:10.1515/9781400881970-017](https://doi.org/10.1515/9781400881970-017).

- [14] A. Shankar and M. Sridharan, New temperatures in Domineering. *Integers* 5 (2005), G04. doi:10.5281/zenodo.7727185.
- [15] A. N. Siegel, *Combinatorial Game Theory*. American Mathematical Society, 2013.
- [16] A. N. Siegel, *CGSuite*, A tool for computational combinatorial game theory, <https://www.cgsuite.org/>.
- [17] J. W. H. M. Uiterwijk and M. Barton, New results for Domineering from combinatorial game theory endgame databases. *Theoretical Computer Science* 592 (2015): 72-86. doi:10.1016/j.tcs.2015.05.017.