

The Number of Solutions of $X^2 = 0$ in Triangular Matrices Over $GF(q)$

Shalosh B. EKHAD¹ and Doron ZEILBERGER¹

Abstract: We prove an explicit formula for the number of $n \times n$ upper triangular matrices, over $GF(q)$, whose square is the zero matrix. This formula was recently conjectured by Sasha Kirillov and Anna Melnikov [KM].

Theorem. *The number of $n \times n$ upper-triangular matrices over $GF(q)$ (the finite field with q elements), whose square is the zero matrix, is given by the polynomial $C_n(q)$, where,*

$$C_{2n}(q) = \sum_j \left[\binom{2n}{n-3j} - \binom{2n}{n-3j-1} \right] \cdot q^{n^2-3j^2-j},$$

$$C_{2n+1}(q) = \sum_j \left[\binom{2n+1}{n-3j} - \binom{2n+1}{n-3j-1} \right] \cdot q^{n^2+n-3j^2-2j}.$$

Proof. In [K] it was shown that the quantity of interest is given by the polynomial $A_n(q) = \sum_{r \geq 0} A_n^r(q)$, where the polynomials $A_n^r(q)$ are defined recursively by

$$A_{n+1}^{r+1}(q) = q^{r+1} \cdot A_n^{r+1}(q) + (q^{n-r} - q^r) \cdot A_n^r(q) \quad ; \quad A_{n+1}^0(q) = 1. \quad (Sasha)$$

For any Laurent formal power series $P(w)$, let $CT_w P(w)$ denote the coefficient of w^0 . Recall that the q -binomial coefficients are defined by

$$\binom{m}{n}_q := \frac{(1-q^m)(1-q^{m-1}) \cdots (1-q^{m-n+1})}{(1-q)(1-q^2) \cdots (1-q^n)}, \quad (Carl)$$

whenever $0 \leq n \leq m$, and 0 otherwise.

Lemma 1. *We have*

$$A_n^r(q) = CT_w \left[\frac{(1-w)(1+w)^n q^{r(n-r)}}{w^r} \sum_{i=0}^{\infty} (-1)^i q^{-(i+1)i/2-i(n-2r)} \binom{i+n-2r}{i}_q w^i \right]. \quad (Anna)$$

Proof. Call the right side of Eq. (Anna), $S_n^r(q)$. Since $S_{n+1}^0(q) = 1$, the lemma would follow by induction if we could show that

$$S_{n+1}^{r+1}(q) - q^{r+1} \cdot S_n^{r+1}(q) - (q^{n-r} - q^r) \cdot S_n^r(q) = 0. \quad (Sasha')$$

¹ Department of Mathematics, Temple University, Philadelphia, PA 19122, USA. [ekhad,zeilberg]@math.temple.edu, [http://www.math.temple.edu/~\[ekhad,zeilberg\]](http://www.math.temple.edu/~[ekhad,zeilberg]) [ftp://ftp.math.temple.edu/pub/\[ekhad,zeilberg\]](ftp://ftp.math.temple.edu/pub/[ekhad,zeilberg]) .

Supported in part by the NSF. Nov. 28, 1995.

Using the linearity of CT_w , manipulating the series, using the definition (*Carl*) of the q -binomial coefficients, and simplifying, brings the left side of (*Sasha'*) to be $CT_w \Phi_n^r(q, w)$, where Φ_n^r is zero except when n is odd and $r = (n-1)/2$, in which case it is a monomial in q times $\frac{(1-w)(1+w)^n}{w^{r+1}}$, and applying CT_w kills it all the same, thanks to the symmetry of the Chu-Pascal triangle. $\square$

Summing the expression proved for $A_n^r(q)$, yields that

$$A_n(q) = CT_w \left[(1-w)(1+w)^n \cdot \sum_{r=0}^{\infty} \sum_{i=0}^r (-1)^i q^{r(n-r)-(i+1)i/2-i(n-2r)} \binom{i+n-2r}{n-2r}_q w^{i-r} \right].$$

Letting $l = r - i$, and changing the order of summation, yields

$$A_n(q) = CT_w \left[(1-w)(1+w)^n \cdot \sum_{\ell=0}^{\lfloor n/2 \rfloor} w^{-\ell} \cdot q^{\ell n - \ell^2} \sum_{i=0}^{\lfloor (n-2\ell)/2 \rfloor} (-1)^i q^{i(i-1)/2} \binom{n-2\ell-i}{i}_q \right].$$

(*SumAnna*)

Luckily, the inner sum evaluates nicely thanks to the following.

Lemma 2. *We have*

$$\sum_{i=0}^{\lfloor m/2 \rfloor} (-1)^i q^{i(i-1)/2} \binom{m-i}{i}_q = (-1)^{\lfloor m/3 \rfloor} q^{m(m-1)/6} \cdot \chi(m \not\equiv 2 \pmod{3}),$$

where $\chi(\dots)$ is the truth value (=1 or 0) of the proposition “ $\dots$ ”.

Proof. While this is unlikely to be new², it is also irrelevant whether or not it is new, since such things are *now* routine, thanks to the package **qEKHAD**, accompanying [PWZ]. Let's call the left side divided by $q^{m(m-1)/6}$, $Z(m)$. Then we have to prove that $Z_0(m) := Z(3m)$ equals $(-1)^m$, $Z_1(m) := Z(3m+1)$ equals $(-1)^m$, and $Z_2(m) := Z(3m+2)$ equals 0. It is directly verified that these are true for $m = 0, 1$, and the general result follows from the second order recurrences produced by **qEKHAD**. The input files **inZ0**, **inZ1**, **inZ2** as well as the corresponding output files, **outZ0**, **outZ1**, **outZ2** can be obtained by anonymous **ftp** to **ftp.math.temple.edu**, directory **pub/ekhad/sasha**. The package **qEKHAD** can be downloaded from <http://www.math.temple.edu/~zeilberg>. $\square$

To complete the proof of the theorem, we use lemma 2 to evaluate the inner sum of (*SumAnna*), then to get $A_{2n}(q)$, we replace n by $2n$, and then replace l by $l + n$, and finally use the binomial theorem. Similarly for $A_{2n+1}(q)$. $\square$

References

[K] A.A. Kirillov, *On the number of solutions to the equation $X^2 = 0$ in triangular matrices over a finite field*, *Funct. Anal. and Appl.* **29** (1995), no. 1.

[KM] A.A. Kirillov and A. Melnikov, *On a remarkable sequence of polynomials*, preprint.

[PWZ] M. Petkovšek, H.S. Wilf, and D. Zeilberger, “ $A = B$ ”, A.K.Peters, 1996.

² For a ‘classical’ proof, see Christian Krattenthaler’s message, at <ftp://ftp.math.temple.edu/pub/ekhad/sasha>.