

On the function “sandwiched” between $\alpha(G)$ and $\bar{\chi}(G)$

V. Y. Dobrynin¹

Submitted: July 18, 1997; Accepted: September 2, 1997

Abstract

A new function of a graph G is presented. Say that a matrix B that is indexed by vertices of G is feasible for G if it is real, symmetric and $I \leq B \leq I + A(G)$, where I is the identity matrix and $A(G)$ is the adjacency matrix of G . Let $\mathcal{B}(G)$ be the set of all feasible matrices for G , and let $\bar{\chi}(G)$ be the smallest number of cliques that cover the vertices of G . We show that

$$\alpha(G) \leq \min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\} \leq \bar{\chi}(G)$$

and that $\alpha(G) = \min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\}$ implies $\alpha(G) = \bar{\chi}(G)$.

The well known Lovász number $\vartheta(G)$ of a graph G [1] is “sandwiched” between the size of the largest stable set in G and the smallest number of cliques that cover the vertices of G

$$\alpha(G) \leq \vartheta(G) \leq \bar{\chi}(G).$$

Some alternative definitions of $\vartheta(G)$ are introduced in [2][3]. For example,

$$\begin{aligned} \vartheta(G) = \max\{ \Lambda(B) \mid & B \text{ is a real positive semidefinite matrix} \\ & \text{indexed by vertices of } G, \\ & B_{vv} = 1 \text{ for all } v \in V(G), \\ & B_{uv} = 0 \text{ whenever } u - v \text{ in } G \}, \end{aligned}$$

where $\Lambda(B)$ is the maximum eigenvalue of B , $V(G)$ — the set of vertices of G , $u - v$ denotes the adjacency of vertices u and v .

Call the matrix B indexed by vertices of G feasible for G if

$$\begin{aligned} & B \text{ is real and symmetric,} \\ & B_{vv} = 1 \text{ for all } v \in V(G), \\ & B_{uv} = 0 \text{ whenever } u \not\sim v \text{ in } G, \\ & 0 \leq B_{uv} \leq 1 \text{ whenever } u \sim v \text{ in } G. \end{aligned}$$

Let $\mathcal{B}(G)$ be the set of all feasible matrices for G . Then [4]

$$\bar{\chi}(G) = \min\{\text{rank}(B) \mid B \in \mathcal{B}(G), B = C^T C, C \geq 0\},$$

where the inequality denotes componentwise inequality.

The aim of this paper is to study a new function of graph G

Theorem. For all graphs G

$$\alpha(G) = \min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\}$$

implies $\alpha(G) = \bar{\chi}(G)$.

Proof. Let $S = \{v_1, v_2, \dots, v_{\alpha(G)}\}$ be the stable set of G , $\bar{S} = V(G) \setminus S$ and $B \in \mathcal{B}(G)$ is a matrix such that $\text{rank}(B) = \alpha(G)$. We can assume that

$$B = \begin{pmatrix} I_{\alpha(G)} & X \\ X^T & Y \end{pmatrix},$$

where $I_{\alpha(G)}$ is the identity $\alpha(G) \times \alpha(G)$ -matrix.

Applying block Gauss elimination, B reduces to the matrix

$$B' = \begin{pmatrix} I_{\alpha(G)} & X \\ 0 & Y - X^T X \end{pmatrix}.$$

We have

$$Y - X^T X = 0$$

or

$$Y_{uv} - \sum_{w \in S} X_{wu} X_{wv} = 0, \quad u, v \in \bar{S} \quad (1)$$

since $\text{rank}(B') = \text{rank}(B) = \alpha(G)$.

Equation (1) gives us further information about the graph G .

- (i) If $v \in \bar{S}$ then exists $u \in S$ such that $u - v$. Indeed, $Y_{vv} = 1$ and $X_{wv} \geq 0$ for all $w \in S$. Hence, $\sum_{w \in S} X_{wv}^2 = 1$ and $X_{uv} > 0$ for some $u \in S$.
- (ii) If $v', v'' \in \bar{S}$ and $X_{uv'} X_{uv''} > 0$ for some $u \in S$ then $v' - v''$. Indeed, if $\sum_{w \in S} X_{wv'} X_{wv''} > 0$ then $Y_{v'v''} > 0$.

Let

$$V_u = \{u\} \cup \{v \mid v \in \bar{S}, X_{uv} > 0\}$$

for all $u \in S$ and $G(V_u)$ be the subgraph induced from G by leaving out all vertices except vertices from V_u . We know from (i) and (ii) that $G(V_u)$ is a clique and $V(G) = \cup_{u \in S} V_u$. Hence, $\bar{\chi}(G) = \alpha(G)$. $\square$

Corollary. If $\bar{\chi}(G) \leq \alpha(G) + 1$ then

$$\bar{\chi}(G) = \min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\}.$$

For example, consider the Petersen graph G . We have $\alpha(G) = \vartheta(G) = 4$, $\bar{\chi}(G) = 5$. Hence, $\min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\} = 5$.

There is a graph G such that

$$\alpha(G) < \min\{\text{rank}(B) \mid B \in \mathcal{B}(G)\} < \bar{\chi}(G).$$

Let $V(G) = 2^{\{1,2,3,4,5,6\}}$ and $u - v$ iff $2 \leq |(u \setminus v) \cup (v \setminus u)| \leq 5$ for all $u, v \in V(G)$. Then [5] $\chi(G) = 32$ and $\text{rank}(A(G)) = 29$ where $A(G)$ is the adjacency matrix of G . Then $\bar{\chi}(G) = 32$ and

and

$$\alpha(\overline{G}) < \min\{\text{rank}(B) \mid B \in \mathcal{B}(\overline{G})\}$$

by theorem.

Acknowledgement.

The author is grateful to the anonymous referee for valuable remarks.

References

1. L. Lovász, “On the Shannon capacity of a graph”, *IEEE Transactions on Information Theory* **IT-25** (1979), 1–7.
2. Martin Grötschel, László Lovász, and Alexander Schrijver, *Geometric Algorithms and Combinatorial Optimization* (Springer-Verlag, 1988).
3. Donald E.Knuth, “The Sandwich Theorem”, *Electronic J. Combinatorics* **1**, A1 (1994), 48 pp.
4. V.Y.Dobrynin, “The chromatic number of a graph and rank of matrix associated with it”, *Vestnik Sankt-Peterburgskogo Universiteta*, Ser.1, Issue 2, Vol.15 (1995), 120–122.
5. N.Alon, P.D.Seymour. “A counterexample to the rank-coloring conjecture”, *Journal of Graph Theory* **13** (1989), 523–525.