

# Hook lengths in a skew Young diagram

Svante Janson

Department of Mathematics, Uppsala University

PO Box 480, S-751 06 Uppsala, Sweden

`svante.janson@math.uu.se`

Submitted: October 15, 1997; Accepted: October 20, 1997

## Abstract

Regev and Vershik (Electronic J. Combinatorics **4** (1997), #R22) have obtained some properties of the set of hook lengths for certain skew Young diagrams, using asymptotic calculations of character degrees. They also conjectured a stronger form of one of their results.

We give a simple inductive proof of this conjecture.

Very recently, Regev and Zeilberger (Annals of Combinatorics, to appear) have independently proved this conjecture.

## 1 Introduction

Regev and Vershik [1] have recently obtained some properties of the set of hook lengths for certain skew Young diagrams. They prove the results using asymptotic calculations of the degrees of certain sequences of characters of the symmetric group, and note that they do not know a direct “finite” proof of their results.

The purpose of the present note is to present such a proof for one of their results, viz. their Theorem 1.2.2. Moreover, Regev and Vershik’s Theorem 1.2.2 states that two different sets of hook lengths have the same product, and the authors conjecture that in fact these two sets are equal. (More precisely, the sets in question should be regarded as multisets, i.e. the elements may have multiplicities.) We prove this conjecture.

Very recently, Regev and Zeilberger [2] have independently proved this conjecture.

## 2 Notation

If  $n_1 \geq n_2 \geq \dots \geq n_m \geq 0$  are integers, with  $m \geq 1$ , let  $D = D(n_1, \dots, n_m)$  be the Young diagram with rows of lengths  $n_1, \dots, n_m$ . Following (and slightly extending) Regev and Vershik [1], we introduce the following definitions, see the examples in Figures 1 and 2.

$R = R(m, n)$  is an  $m \times n$  rectangle, i.e. a Young diagram with  $m$  rows of equal length  $n$ .

We assume that  $n \geq n_1$  and that  $R$  and  $D$  are positioned such that their top left corners coincide. Then  $D \subseteq R$ . (Regev and Vershik consider only the case when  $R$  is the smallest rectangle containing  $D$ , i.e. when  $n = n_1$  and  $n_m \geq 1$ . We find it convenient to treat a slightly more general situation.)

$D^*$  is obtained by rotating  $D$  a half turn about the center of  $R$ ; thus  $D^* \subseteq R$  and  $R \setminus D^*$  is a Young diagram with rows of lengths  $n - n_m, n - n_{m-1}, \dots, n - n_1$ .

$SQ = SQ(n_1, \dots, n_m; n)$  is obtained from  $R \setminus D^*$  by adding two copies of  $D^*$ , one along the left edge and one along the top edge of  $R$ .

We will use a coordinate system with the  $x$ -axis directed upwards and the  $y$ -axis directed to the left. (Note that thus rows are numbered from the bottom to the top and columns from the right to the left.) We may then describe the skew diagrams as follows.

$$\begin{aligned}
 R(m, n) &= \{(i, j) : 1 \leq i \leq m, 1 \leq j \leq n\} \\
 D^*(n_1, \dots, n_m) &= \{(i, j) : 1 \leq i \leq m, 1 \leq j \leq n_i\} \\
 SQ(n_1, \dots, n_m; n) &= \{(i, j) : 1 \leq i \leq m, n_i + 1 \leq j \leq n_i + n\} \\
 &\quad \cup \{(i, j) : m + 1 \leq i \leq 2m, 1 \leq j \leq n_{i-m}\}.
 \end{aligned}$$

The hook length  $h_A(x)$  of an element  $x$  of a (skew) diagram  $A$  is, as usual, the number of elements of  $A$  directly below or to the right of  $x$ , including  $x$  itself. We define  $H(A)$  to be the multiset  $\{h_A(x) : x \in A\}$ .




Figure 1:  $D^*(4, 2, 1)$  marked with 0;  $SQ(4, 2, 1; 4)$  marked with  $\times$




Figure 2:  $SQ(4, 2, 1; 4)$ ,  $R(3, 4)$  and  $D(4, 2, 1)$  with hook lengths

### 3 Result

**Theorem.** Let  $n \geq n_1 \geq n_2 \geq \dots \geq n_m \geq 0$  ( $m \geq 1$ ) be integers. We then have the equality of multisets of hook lengths

$$H(SQ(n_1, \dots, n_m; n)) = H(R(m, n)) \cup H(D(n_1, \dots, n_m)). \tag{1}$$

*Proof.* We say that  $(n, n_1, \dots, n_m)$  is good if (1) holds. The result follows by double induction, in  $m$  and  $n_m$ , from the following three claims.

- (i) If  $n \geq n_1 \geq 0$ , then  $(n, n_1)$  is good.
- (ii) If  $(n, n_1, \dots, n_m)$  is good, then  $(n, n_1, \dots, n_m, 0)$  is good.
- (iii) If  $(n, n_1, \dots, n_m)$  is good and  $n_{m-1} > n_m$ , then  $(n, n_1, \dots, n_{m-1}, n_m + 1)$  is good.

(i) is trivial:  $SQ(n_1; n)$  consists of two rows with  $n$  and  $n_1$  elements, positioned such that their hook lengths are  $1, \dots, n$  and  $1, \dots, n_1$ , respectively, corresponding to the hook lengths of  $R(1, n)$  and  $D(n_1)$ .

For (ii), note that  $SQ' = SQ(n_1, \dots, n_m, 0; n)$  is obtained from  $SQ = SQ(n_1, \dots, n_m; n)$  by inserting a new row  $(m + 1, 1), \dots, (m + 1, n)$ , moving up all elements  $(i, j)$  with  $i \geq m$ ; equivalently,  $SQ'$  is obtained from  $SQ$  by adding a new element on top of each column  $1, \dots, n$ . Each of these new top elements has  $m$  elements beneath it, and thus their hook lengths are  $m + 1, \dots, m + n$ . Moreover, all elements in  $SQ$  keep the same hook length in  $SQ'$ ; consequently  $H(SQ') = H(SQ) \cup \{m + 1, \dots, m + n\}$ , see Figure 3.

For the right hand side of (1), we observe that adding a new row of length 0 does not change  $D$ , while  $R = R(m, n)$  is changed to  $R' = R(m + 1, n)$ , which equals  $R$  with an additional top row having hook lengths  $m + 1, \dots, m + n$ . Thus  $H(R') = H(R) \cup \{m + 1, \dots, m + n\}$ . Consequently, if  $H(SQ) = H(R) \cup H(D)$ , then  $H(SQ') = H(R') \cup H(D)$  too, which proves (ii).

For (iii) we let  $SQ' = SQ(n_1, \dots, n_m + 1; n)$  and  $D' = D(n_1, \dots, n_m + 1)$  (this time  $R = R(m, n)$  stays the same), and argue similarly.  $SQ'$  differs from  $SQ$  in three places: the



For the right hand side of (1), we observe that  $D'$  differs from  $D$  in that a new element is added to the last row. The hook lengths in this row in  $D$  are  $1, \dots, n_m$ , while in  $D'$  they are  $1, \dots, n_m + 1$ , a net addition of  $n_m + 1$ .

The element above the new element in the  $k$ th row from top has hook length in  $D$   $n_k - n_m + m - k - 1$ , while its hook length in  $D'$  is increased by 1 to  $n_k - n_m + m - k$ .

No other hook lengths are affected, and consequently,

$$H(D') = H(D) \cup \{n_m + 1\} \cup \{n_k - n_m + m - k\}_{k=1}^{m-1} \setminus \{n_k - n_m + m - k - 1\}_{k=1}^{m-1}.$$

Comparing this with (2), we see that if  $H(SQ) = H(R) \cup H(D)$ , then  $H(SQ') = H(R) \cup H(D')$  too. This completes the proof of (iii), and thus of the theorem.  $\square$

## References

- [1] A. Regev and A. Vershik, Asymptotics of Young diagrams and hook numbers. *Electron. J. Combin.* **4** (1997), #R22, 12pp.
- [2] A. Regev and D. Zeilberger, Proof of a conjecture on multisets of hook numbers. *Ann. Combin.*, to appear.