

TIGHT UPPER BOUNDS FOR THE DOMINATION NUMBERS OF GRAPHS WITH GIVEN ORDER AND MINIMUM DEGREE

W. EDWIN CLARK

UNIVERSITY OF SOUTH FLORIDA

TAMPA, FL 33620-5700

eclark@math.usf.edu

AND

LARRY A. DUNNING

BOWLING GREEN STATE UNIVERSITY

BOWLING GREEN, OH 43403-0214

dunning@cs.bgsu.edu

Submitted: June 2, 1997; Accepted: October 27, 1997

ABSTRACT. Let $\gamma(n, \delta)$ denote the maximum possible domination number of a graph with n vertices and minimum degree δ . Using known results we determine $\gamma(n, \delta)$ for $\delta = 0, 1, 2, 3$, $n \geq \delta + 1$ and for all n, δ where $\delta = n - k$ and n is sufficiently large relative to k . We also obtain $\gamma(n, \delta)$ for all remaining values of (n, δ) when $n \leq 14$ and all but 6 values of (n, δ) when $n = 15$ or 16 .

1. INTRODUCTION

We denote the domination number of a graph G by $\gamma(G)$. By an (n, δ) -graph we mean a graph with n vertices and minimum degree δ . Let $\gamma(n, \delta)$ be the maximum of $\gamma(G)$ where G is an (n, δ) -graph. Using known results [3], [7], [8], [9]

one easily finds the exact values of $\gamma(n, \delta)$ when $\delta = 0, 1, 2, 3$. It is also fairly easy to obtain $\gamma(n, \delta)$ when $\delta = n - k$ for n sufficiently large relative to k . By various methods we also find $\gamma(n, \delta)$ for all remaining values of (n, δ) when $n \leq 14$ and all but 6 values of (n, δ) when $n = 15$ and 16 . Many values can be established using the upper bounds in [2] together with examples found by computer search or *ad hoc* techniques. In a number of cases we have used Brendan McKay's program makeg to

1991 *Mathematics Subject Classification.* 05C35.

Key words and phrases. graph, domination number, upper bounds, minimum degree.

generate all nonisomorphic (n, δ) -graphs [6] while checking the domination numbers using a simple recursive, depth-first search.

Our results give support to the the natural conjecture that $\gamma(n, \delta)$ is attained by an (n, δ) -graph with the minimum number of edges, that is, by a regular graph if $n\delta$ is even or by a graph with degree sequence $(\delta + 1, \delta, \delta, \dots, \delta)$ if $n\delta$ is odd. We are able to verify the conjecture for all the cases mentioned above where we are able to determine $\gamma(n, \delta)$ and in at least one case where we are not (see Proposition 4.11). However, see Section 5 for some evidence in opposition to the conjecture.

To simplify discussion we say that a graph is *almost δ -regular* if its degree sequence has the form $(\delta + 1, \delta, \delta, \dots, \delta)$ and we define $\gamma_r(n, \delta)$ to be the maximum of $\gamma(G)$ where G is an (n, δ) -graph which is regular if $n\delta$ is even and almost regular if $n\delta$ is odd. In this notation the above mentioned conjecture becomes $\gamma(n, \delta) = \gamma_r(n, \delta)$ for all n and δ .

We use the following standard notation. Let $G = (V, E)$ be a graph with vertex set V and edge set E . We write $x \sim y$ to indicate that the vertices x and y are adjacent. For $v \in V$ the open neighborhood $N(v)$ is the set of all vertices adjacent to v and the closed neighborhood of v is the set $N[v] = N(v) \cup \{v\}$. $S \subseteq V$ is a dominating set for G if $\bigcup_{x \in S} N[x] = V$. The domination number, $\gamma(G)$, is the cardinality of a smallest dominating set for G . If $x \sim y$ or $x = y$ we say that x covers y . If $A \subseteq V$ then we let $\langle A \rangle$ denote the subgraph of G induced by A .

Table 1 contains the value of $\gamma(n, \delta)$ for $n \leq 16$, $0 \leq \delta \leq n - 1$ if the value is known, otherwise upper and lower bounds for $\gamma(n, \delta)$. We establish these values in Sections 2, 3 and 4.

Table 1. Values of $\gamma(n, \delta)$ for $n \leq 16$, $0 \leq \delta \leq n - 1$.

$n \setminus \delta$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	1															
2	2	1														
3	3	1	1													
4	4	2	2	1												
5	5	2	2	1	1											
6	6	3	2	2	2	1										
7	7	3	3	2	2	1	1									
8	8	4	4	3	2	2	2	1								
9	9	4	4	3	3	2	2	1	1							
10	10	5	4	3	3	2	2	2	2	1						
11	11	5	5	4	3	3	3	2	2	1	1					
12	12	6	6	4	4	3	3	2	2	2	2	1				
13	13	6	6	4	4	3	3	3	2	2	2	1	1			
14	14	7	6	5	4	4	3	3	3	2	2	2	2	1		
15	15	7	7	5	5	4	3-4	3	3	2-3	2	2	2	1	1	
16	16	8	8	6	5	4-5	4	3-4	3	2-3	2-3	2	2	2	2	1

2. THE CASES $\delta = 0, 1, 2, 3$.

Lemma 2.1. *If $n \geq m > \delta \geq 1$ then*

$$(2.1) \quad \gamma(n, \delta) + \gamma(m, \delta) \leq \gamma(n + m, \delta)$$

and if $n\delta$ or $m\delta$ is even

$$(2.2) \quad \gamma_r(n, \delta) + \gamma_r(m, \delta) \leq \gamma_r(n + m, \delta).$$

Proof. The lemma is immediate from the fact that if G is an (n, δ) -graph and H is an (m, δ) -graph then the disjoint union $G \cup H$ is an $(n + m, \delta)$ -graph with domination number $\gamma(G \cup H) = \gamma(G) + \gamma(H)$. (2.2) follows from the fact that if G is regular and H is regular (resp., almost regular) then $G \cup H$ is regular (resp., almost regular). $\square$

Corollary 2.2. *For every positive integer k we have*

$$(2.2) \quad k\gamma(n, \delta) \leq \gamma(kn, \delta)$$

and provided that $n\delta$ is even,

$$(2.3) \quad k\gamma_r(n, \delta) \leq \gamma_r(kn, \delta). \quad \square$$

We will need the following two theorems:

Ore's Theorem [7] . *If G is an (n, δ) -graph with $\delta \geq 1$ then $\gamma(G) \leq n/2$. $\square$*

Reed's Theorem [9] . *If G is an (n, δ) -graph with $\delta \geq 3$ then $\gamma(G) \leq 3n/8$. $\square$*

Proposition 2.3. *For $n \geq 1$*

$$\gamma(n, 0) = \gamma_r(n, 0) = n$$

and for $n \geq 2$

$$\gamma(n, 1) = \gamma_r(n, 1) = \lfloor n/2 \rfloor. \quad \square$$

Proof. The case $\delta = 0$ is trivial and the case $\delta = 1$ is immediate from Ore's Theorem. $\square$

Proposition 2.4. *For $n \geq 3$,*

$$\gamma(n, 2) = \gamma_r(n, 2) = \begin{cases} \lfloor n/2 \rfloor - 1, & \text{if } n \equiv 2 \pmod{4} \\ \lfloor n/2 \rfloor, & \text{otherwise.} \end{cases}$$

Proof. From Ore's Theorem we have $\gamma(n, 2) \leq \lfloor n/2 \rfloor$. Consider the four cases:

$$(1) \quad n = 4k, \quad k \geq 1.$$

$$(2) \quad n = 4k + 1, \quad k \geq 1.$$

$$(3) \quad n = 4k + 2, \quad k \geq 1.$$

$$(4) \quad n = 4k + 3, \quad k \geq 0.$$

In cases (1) and (2) $\lfloor n/2 \rfloor = 2k$ so in these cases it suffices to exhibit a 2-regular graph G with $\gamma(G) = 2k$. In case (1) we can take G to be the disjoint union of k

4-cycles. In case (2) we can take G to be the disjoint union of $k-1$ 4-cycles and

one 5-cycle. In case (4) we can take G to be the union of k 4-cycles and one 3-cycle. Then $\gamma(G) = 2k + 1 = \lfloor n/2 \rfloor$.

For case (3) we first note that by [8] or [3] if a graph G has no isolated vertices and if $\gamma(G) = n/2$ then each connected component of G is either a 4-cycle or has a vertex of degree 1. Since we are interested here only in graphs with $\delta = 2$ it follows that such a graph cannot have $\gamma(G) = n/2$ unless it has order $4k$. So in case (3) we have $\gamma(n, 2) \leq n/2 - 1$. To see that this upper bound can be attained one must only consider the disjoint union of $k - 1$ 4-cycles and one 6-cycle. $\square$

Proposition 2.5. *If $n \geq 4$ then*

$$\gamma(n, 3) = \gamma_r(n, 3) = \lfloor 3n/8 \rfloor.$$

Proof. From Reed's Theorem $\gamma(n, 3) \leq 3n/8$. Thus it suffices to exhibit for each $n \geq 4$ an $(n, 3)$ -graph G_n which is 3-regular if n is even and almost regular if n is odd such that $\gamma(G_n) = \lfloor 3n/8 \rfloor$.

We first note that it suffices to find the graphs G_n for $4 \leq n \leq 11$: For $12 \leq n \leq 15$ we can take

$$G_{15} = G_8 \cup G_7, \quad G_{14} = G_8 \cup G_6, \quad G_{13} = G_8 \cup G_5, \quad G_{12} = G_8 \cup G_4.$$

If $n \geq 16$ we can write $n = 8k + r$ where $k \geq 1$ and $r \in \{8, 9, 10, 11, 12, 13, 14, 15\}$. Then $\lfloor 3n/8 \rfloor = 3k + \lfloor 3r/8 \rfloor$. So if we let G_n be the disjoint union of k copies of G_8 and one copy of G_r we have

$$\gamma(G_n) = k\gamma(G_8) + \gamma(G_r) = 3k + \lfloor 3r/8 \rfloor = \lfloor 3n/8 \rfloor.$$

Moreover since G_8 is 3-regular, G_n will be regular if r is even and almost regular if r is odd.

This leaves the cases $4 \leq n \leq 11$. It is easy to see that we may take $G_4 = K_4$, $G_5 = K_5 - \{\text{two disjoint edges}\}$, $G_6 = \text{any regular } (6, 3)\text{-graph}$, $G_7 = \text{any almost regular } (7, 3)\text{-graph}$, G_8 can be taken to be the 8-cycle with 4 diameters added, and $G_{10} = G_4 \cup G_6$. This leaves only G_9 and G_{11} . See appendix B for these graphs, namely the graphs listed there as $G(9, 3, 3)$ and $G(11, 3, 4)$. $\square$

3. THE CASES $\delta = n - k$ FOR k SMALL.

We first describe an upper bound for $\gamma(n, \delta)$ which is a variation of Theorem 2 in [2]. This result plays a major role in almost all of our arguments.

Proposition 3.1. *Let G be an (n, δ) -graph containing a set S of s vertices which covers at least λ vertices. Define the sequence $g_s, g_{s+1}, \dots, g_{n-\delta}$ by $g_s = n - \lambda$ and*

$$g_{t+1} = \left\lfloor g_t \left(1 - \frac{\delta + 1}{n - t} \right) \right\rfloor = g_t - \left\lceil \frac{g_t(\delta + 1)}{n - t} \right\rceil, \quad t \geq s.$$

Then for any t , $s \leq t \leq n - \delta$, there is a set of t vertices that covers at least $n - g_t$ vertices. Thus if we set

$$M(n, \delta, \lambda, s) = \min \{ t \mid g_t = 0 \}$$

we have

$$\gamma(G) \leq M(n, \delta, \lambda, s).$$

In particular,

$$\gamma(n, \delta) \leq M(n, \delta, \delta + 1, 1)$$

and if $n\delta$ is odd,

$$\gamma(n, \delta) \leq M(n, \delta, \delta + 2, 1).$$

Proof. Starting with the set $S = \{v_1, v_2, \dots, v_s\}$ we select successively and greedily the elements $v_{s+1}, v_{s+2}, \dots, v_t$. Let u_t , $t \geq s$, be the number of vertices left uncovered after the vertices $v_1, v_2, \dots, v_t$ have been chosen. It clearly suffices to prove that $u_t \leq g_t$ for $t \geq s$. We prove this by induction on t . If $t = s$ then since S covers at least λ vertices $u_s \leq n - \lambda = g_s$.

Assume that $u_t \leq g_t$ holds for some $t \geq s$. Let

$$U_t = \{x_1, x_2, \dots, x_{u_t}\}$$

be the set of vertices not covered by $v_1, v_2, \dots, v_t$ and let

Now define the $u_t \times (n - t)$ matrix M whose (i, j) -th entry is given by

$$M_{i,j} = \begin{cases} 1, & \text{if } x_i \text{ covers } y_j \\ 0, & \text{otherwise.} \end{cases}$$

Since none of the x_i 's are covered by any of the vertices $v_1, v_2, \dots, v_t$ chosen so far and $\deg(x_i) \geq \delta$ there are at least $\delta + 1$ ones in each row of M . This gives at least $u_t(\delta + 1)$ ones in the entire matrix. Since there are $n - t$ columns at least one column must contain at least

$$N = \left\lceil \frac{u_t(\delta + 1)}{n - t} \right\rceil$$

ones. Say it is the j -th column. This means that y_j covers at least N of the x_i 's. So if we select v_{t+1} to cover the maximum number N_{max} of the x_i 's we have $N \leq N_{max}$ and the number of vertices now left uncovered is

$$\begin{aligned} u_{t+1} &= u_t - N_{max} \\ &\leq u_t - N \\ &= u_t - \left\lceil \frac{u_t(\delta + 1)}{n - t} \right\rceil \\ &= u_t + \left\lfloor -\frac{u_t(\delta + 1)}{n - t} \right\rfloor \\ &= \left\lfloor u_t \left(1 - \frac{\delta + 1}{n - t} \right) \right\rfloor \\ &\leq \left\lfloor g_t \left(1 - \frac{\delta + 1}{n - t} \right) \right\rfloor \\ &= g_{t+1}. \end{aligned}$$

This completes the proof. $\square$

Proposition 3.2.

- (1) $\gamma(n, n - 1) = \gamma_r(n, n - 1) = 1$ for any $n \geq 1$.
- (2) $\gamma(n, n - 2) = \gamma_r(n, n - 2) = 1$ for odd $n \geq 2$.
- (3) $\gamma(n, n - 2) = \gamma_r(n, n - 2) = 2$ for even $n \geq 2$.

Proof. For each of the cases (1) and (2) there is a vertex of degree $n - 1$ which by itself forms a dominating set. For case (3) any regular graph with $\delta = n - 2$ clearly has domination number 2. $\square$

Lemma 3.3. *Let $n \geq 2$ and let G be an (n, δ) -graph with a vertex of degree Δ . Then $\gamma(G) \leq 2$ if*

$$(3.1) \quad (n - \Delta - 1)(n - \delta - 2) < n - 1$$

.

Proof. By Proposition 3.1 $\gamma(G) \leq M(n, \delta, \Delta + 1, 1)$ and $M(n, \delta, \Delta + 1, 1) = 2$ if and only if $g_2 = 0$. Now $g_1 = n - \Delta - 1$ and

$$g_2 = \left\lfloor g_1 \left(1 - \frac{\delta + 1}{n - 1} \right) \right\rfloor.$$

So $g_2 = 0$ if and only if

$$(n - \Delta - 1) \left(1 - \frac{\delta + 1}{n - 1} \right) < 1$$

which is equivalent to (3.1). $\square$

Proposition 3.4. *Let $3 \leq k \leq n - 1$.*

(1) *If $(k - 1)(k - 2) + 1 < n$ then*

$$\gamma(n, n - k) = \gamma_r(n, n - k) = 2.$$

(2) *If n is odd, k is even and $(k - 2)^2 + 1 < n$ then*

$$\gamma(n, n - k) = \gamma_r(n, n - k) = 2$$

Proof. If $k \geq 3$ then $\gamma(n, n - k) \neq 1$ since a regular or almost regular graph with $\delta = n - k$ has vertices of degree at most $n - k + 1$ so a single vertex cannot cover all n vertices. Hence whenever $\gamma(n, n - k) \leq 2$ we have $\gamma(n, n - k) = \gamma_r(n, n - k) = 2$. By Lemma 3.3 $\gamma(n, n - k) \leq 2$ if $(k - 1)(k - 2) + 1 < n$. This proves (1). To prove (2) we observe that if n is odd and k is even then $\delta = n - k$ is odd and so any (n, δ) -graph has a vertex of degree at least $\delta + 1$ so we can take $\Delta = \delta + 1 = n - k + 1$ in (3.1) and we obtain (2). $\square$

Corollary 3.5.

- (1) $\gamma(n, n-3) = \gamma_r(n, n-3) = 2$ if $n > 3$.
- (2) $\gamma(n, n-4) = \gamma_r(n, n-4) = 2$ if $n \geq 7$.
- (3) $\gamma(n, n-5) = \gamma_r(n, n-5) = 2$ if $n > 13$.
- (4) $\gamma(n, n-6) = \gamma_r(n, n-6) = 2$ if $n \geq 21$ or $n = 19$.
- (5) $\gamma(n, n-7) = \gamma_r(n, n-7) = 2$ if $n > 31$. $\square$

4. $\gamma(n, \delta)$ FOR $n \leq 16$.

In Table 1 we give a list of values (or bounds) for $\gamma(n, \delta)$ when $n \leq 16$. In this section we justify the entries of this table. See Table 3 in Appendix A for a summary of how entries in Table 1 are obtained. From Propositions 2.3, 2.4, 2.5, 3.2 and Corollary 3.5 we obtain immediately the exact values of $\gamma(n, \delta)$ for all values of n and δ for $n \leq 16$ except for the following 33 cases:

- (1) $n = 9, \delta = 4$.
- (2) $n = 10, \delta = 4, 5$.
- (3) $n = 11, \delta = 4, 5, 6$.
- (4) $n = 12, \delta = 4, 5, 6, 7$.
- (5) $n = 13, \delta = 4, 5, 6, 7, 8$.
- (6) $n = 14, \delta = 4, 5, 6, 7, 8$.
- (7) $n = 15, \delta = 4, 5, 6, 7, 8, 9$.
- (8) $n = 16, \delta = 4, 5, 6, 7, 8, 9, 10$.

The number of unknown $\gamma(n, \delta)$ can be reduced further by using the upper bounds given in Proposition 3.1 (taking $s = 1$) and Reed's Theorem. See Table 2 for upper bounds computed using Proposition 3.1. Let $Ub(n, \delta)$ denote $M(n, \delta, \delta+1, 1)$ if $n\delta$ is even or $M(n, \delta, \delta+2, 1)$ if $n\delta$ is odd. If the entry in the (n, δ) -th cell of Table 2 is a single number then that number is $Ub(n, \delta)$ and is, in fact, equal to $\gamma(n, \delta)$. So only an example suffices to establish $\gamma(n, \delta)$ in these cases. Tight lower bounds are given by the graphs $G(n, \delta, \gamma)$ in Appendix B. Each graph $G(n, \delta, \gamma)$ listed in Appendix B is an (n, δ) graph with domination number γ . These graphs

are regular if $n\delta$ is even and almost regular otherwise. After applying these results we have only the following 15 remaining cases:

- (1) $n = 10, \delta = 5$.
- (2) $n = 11, \delta = 4$.
- (3) $n = 12, \delta = 7$.
- (4) $n = 13, \delta = 5, 8$.
- (5) $n = 14, \delta = 4, 6$.
- (6) $n = 15, \delta = 5, 6, 9$.
- (7) $n = 16, \delta = 4, 5, 7, 9, 10$.

As indicated in Table 3 (in Appendix A) all but 6 of these cases are settled by one of the following propositions and/or the use of an exhaustive search using Brendan McKay's program `makeg` augmented with a subroutine to compute domination numbers. For example we use Proposition 4.1 below to reduce the determination of $\gamma(10, 5)$ to the determination of $\gamma_r(10, 5)$. Then we search through all 5-regular graphs of order 10 to find that $\gamma_r(10, 5) = 2$.

Table 2. Upper bounds given by Proposition 3.1 for $5 \leq n \leq 16$

$n \setminus \delta$	4	5	6	7	8	9	10	11	12	13	14	15
5	1											
6	2	1										
7	2	1	1									
8	2	2	2	1								
9	3	2	2	1	1							
10	3	2,3,7	2	2	2	1						
11	3,4,5	3	3	2	2	1	1					
12	4	3	3	2,3,8	2	2	2	1				
13	4,5,6	3,4,7	3	3	2,3,9	2	2	1	1			
14	4,5,7	4	3,4,7	3	3	2	2	2	2	1		
15	5	4,5,8	3-4*	3	3	2-3*	2	2	2	1	1	
16	5,6,5	4-5*	4	3-4*	3	2-3*	2-3*	2	2	2	2	1

An entry of the form $a, a+1, \lambda$ indicates that $\gamma(n, \delta) = a$ and $Ub(n, \delta) = a+1$. λ is the least positive integer for which $M(n, \delta, \lambda+1, 1) = a$. Thus in these cases one obtains a tight upper bound by assuming the existence of a vertex of degree λ . In cells containing $x-y^*$ the value of $\gamma(n, \delta)$ is unknown, but our current best upper bound is given by $Ub(n, \delta) = y$ and x is our current best lower bound.

Several times below we need the following trivial result.

Lemma 4.0. *Let $\mathfrak{F}$ be a set of two element subsets of a four element set X . If the following two conditions hold*

- (1) $A \cap B \neq \emptyset$ for all $A, B \in \mathfrak{F}$, and
- (2) $\bigcup_{A \in \mathfrak{F}} A = X$,

then $\bigcap_{A \in \mathfrak{F}} A \neq \emptyset$ $\square$

Proposition 4.1. $2 \leq \gamma_r(10, 5) = \gamma(10, 5) \leq 3$.

Proof. From Proposition 3.1 (see Table 2) we obtain $\gamma(10, 5) \leq 3$ so it suffices to show that if $G = (V, E)$ is a $(10, 5)$ -graph that is not regular then $\gamma(G) \leq 2$. If $\Delta \geq 7$ then by Lemma 3.3 (or Table 2), $\gamma(G) \leq 2$. So suppose that $\Delta = 6$. Let x be a vertex of degree 6. Then V is the disjoint union of $N[x]$, the closed neighborhood of x , and the 2-set $A = \{a, b\}$. If the induced graph $H = G[A] \setminus \{4\}$ has 2 edges then

it can be dominated by a single vertex. So we can assume that H has at most one edge. Thus H has at least one isolated vertex, say, c . This means that c is adjacent to at least 5 vertices in the open neighborhood $N(x)$ of x and each of the remaining two vertices a and b are adjacent to at least 4 vertices of $N(x)$. It follows that there is at least one vertex y in $N(x)$ that is adjacent to all three vertices in A . Hence $\{x, y\}$ is a dominating set for G . $\square$

Proposition 4.2. $3 \leq \gamma_r(11, 4) = \gamma(11, 4) \leq 4$

Proof. By Proposition 3.1 $\gamma(11, 4) \leq 4$. By Lemma 2.1 and the above

$$\gamma_r(11, 4) \geq \gamma_r(6, 4) + \gamma_r(5, 4) = 3.$$

So it suffices to show that if G an $(11, 4)$ graph that is not regular then $\gamma(G) \leq 3$.

But it is immediate from Proposition 3.1 that if $\Delta \geq 5$ then $\gamma \leq 3$. $\square$

Proposition 4.3. $2 \leq \gamma_r(12, 7) = \gamma(12, 7) \leq 3$

Proof. Any regular $(12, 7)$ -graph has $2 \leq \gamma$. By Proposition 3.1 if G is a $(12, 7)$ -graph with $\gamma(G) \leq 3$ and $\Delta \geq 8$, then $\gamma(G) \leq 2$, and the proposition follows. $\square$

Proposition 4.4. $\gamma_r(13, 5) = \gamma(13, 5) = 3$.

Proof. By Appendix B there is an almost regular $(13, 5)$ -graph with domination number 3 so $3 \leq \gamma_r(13, 5) \leq \gamma(13, 5)$. It therefore suffices to show that $\gamma(13, 5) \leq 3$. If $G = (V, E)$ is a $(13, 5)$ -graph with maximum degree $\Delta \geq 7$ then from Proposition 3.1 we have $\gamma(G) \leq 3$. Now if $\Delta \leq 6$ then since $n\delta$ is odd G will always have a vertex x of degree 6. Thus V is a disjoint union of the closed neighborhood $N[x]$ of x and a set A of cardinality 6. Consider the 6×12 matrix M whose rows are indexed by the elements of A and the columns are indexed by the elements of $V - \{x\}$. If $a \in A$ is adjacent to or equal to $y \in V - \{x\}$ let $M_{a,y} = 1$. Otherwise, let $M_{a,y} = 0$. Now each row of M has at least 6 ones, so M has in all at least 36 ones.

If there is a column with 4 ones this means there is a vertex $y \in V - \{x\}$ that covers all but two vertices, say, z, w in A . If the sets $N[z]$, $N[w]$ are disjoint their union contains all but one vertex t . Then $\{x, y, t\}$ is a dominating set and we are

done. But if $v \in N[z] \cap N[w]$, then v dominates both z and w so $\{x, y, v\}$ is a dominating set.

We are left with the case where each column of M contains at most three ones. This implies that each column contains exactly three ones. Hence each vertex in the subgraph $H = \langle A \rangle$ has degree exactly 2. Hence H is a $(6,2)$ -graph and by Proposition 2.4 can be dominated by two vertices. Hence G can be dominated by x together with these two vertices and we are done. $\square$

Proposition 4.5. $2 \leq \gamma_r(13, 8) = \gamma(13, 8) \leq 3$.

Proof. Any regular $(13,8)$ -graph G has domination number at least 2. By Lemma 3.3 if G has a vertex of degree greater than 8 then $\gamma(G) \leq 2$. $\square$

Proposition 4.6. $\gamma_r(14, 4) = \gamma(14, 4) = 4$.

Proof. From Appendix B there is a 4-regular graph of order 14 with domination number 4. This lower bound also follows from Lemma 2.1 and the fact that $\gamma(7, 4) = \gamma_r(7, 4) = 2$ by Corollary 3.5. Thus it suffices to show that if $G = (V, E)$ is a $(14,4)$ -graph then $\gamma(G) \leq 4$. If there is a vertex of degree 7 we obtain $\gamma(G) \leq 4$ from Proposition 3.1. Hence we may assume that Δ is at most 6.

We consider two cases:

- (1) There are vertices a and b such that $|N[a] \cup N[b]| \geq 10$.
- (2) For all vertices a and b we have $|N[a] \cup N[b]| \leq 9$.

In case (1) if $|N[a] \cup N[b]| \geq 11$ we get $\gamma(G) \leq 4$ immediately from Proposition 3.1 with $s = 2$ and $\lambda = 11$. Hence can assume that $|N[a] \cup N[b]| = 10$. We let A be the set of 4 vertices not in $N[a] \cup N[b]$ and $H = \langle A \rangle$. Let

$$B = (N[a] \cup N[b]) - \{a, b\}.$$

For each vertex $v \in B$ let A_v be the set of vertices in A that are adjacent to v . If $|A_v| \geq 3$ for some v then v covers at least three vertices from A and then clearly $\gamma(G) \leq 4$. Hence we may assume that $|A_v| \leq 2$ for each $v \in B$. Note that the sum of the $|A_v|$'s is the number of edges from A to B .

If H has 2 edges then $\gamma(H) \leq 2$ so $\gamma(G) \leq 4$. So we may assume that H has at most one edge. We consider the two subcases:

- (1a) H has no edges; and
- (1b) H has exactly one edge.

If (1a) holds then there are at least $4 \cdot 4 = 16$ edges from A to B . Thus, the sum of the $|A_v|$'s is at least 16. Since $|A_v| \leq 2$ for all $v \in B$ we must have $|A_v| = 2$ for all eight v in B . If $A_{v_1} \cap A_{v_2} = \emptyset$ then $A_{v_1} \cup A_{v_2} = A$ and so $\{a, b, v_1, v_2\}$ is a dominating set for G . So we can assume that $A_{v_1} \cap A_{v_2} \neq \emptyset$ for all v_1 and v_2 . The sets A_v must cover A as the vertices of A have degree at least 4 in G . Clearly the hypotheses of Lemma 4.0 hold for $\mathcal{F} = \{A_v \mid v \in B\}$ and hence the A_v 's contain a common element. But this common element would be a vertex of degree 8, contrary to the assumption that $\Delta \leq 6$. This settles case (1a).

Assume that (1b) holds. Let $A = \{x, y, z, w\}$ and assume that $\{z, w\}$ is the single edge in H . In this case there are at least 14 edges from A to B and there must be at least 6 of the A_v 's that have cardinality 2. If there were more than 6 the argument for case (1a) repeated would show the existence of a vertex of degree greater than 6, a situation already handled. Thus we may assume that there are exactly 6 vertices v , in B such that $|A_v| = 2$. If for some i , $A_{v_i} = \{x, y\}$ then $\{a, b, v_i, z\}$ is a domination set for G . We show this must happen. If not, since both x and y have degree at least 4 in G but degree 0 in H we can assume that A_{v_i} contains x for $i = 1, 2, 3, 4$ and that A_{v_i} contains y for $i = 5, 6, 7, 8$. This means that the two A_v 's that have one element must contain either an x or a y . It follows that there are at least three of the two element A_v 's that contain z and at least three, that contain w . So it is clear that we cannot avoid having either two A_v 's of the form $\{x, z\}$ and $\{y, w\}$ or, of the form $\{x, w\}$ and $\{y, z\}$. But this contradicts our assumption that the two element A_v 's are never disjoint. This completes the proof in case (1b) and hence the proof of case (1).

Assume that case (2) holds. Note that if G is not regular then it contains a vertex of degree at least 5 and by Proposition 2.1 there will be two vertices that

cover at least 10 vertices which puts us back in case (1). Thus we can assume that G is 4-regular. Note that this implies that $N[a] \cap N[b] \neq \emptyset$ for all vertices a and b . Hence any two vertices can be covered by a single vertex. Thus if we are able to show that we can cover 12 vertices with 3 vertices we will know that $\gamma(G) \leq 4$. By Proposition 3.1 there are two vertices a and b such that $N[a] \cup N[b] = 9$. Let

$$B = (N[a] \cup N[b]) - \{a, b\}$$

and let

$$A = V - (N[a] \cup N[b]).$$

Now again let $H = \langle A \rangle$. If H contains a vertex x of degree 2 we can cover 12 vertices with the a , b and x and we are done. So we can assume that H has only vertices of degree 1 and hence H has at most 2 edges. It follows that there must be at least 16 edges with one end in A and the other end in B . Since $|A| = 5$ and $|B| = 7$ there must be at least one vertex $x \in B$ that is incident with at least 3 vertices in A . Then a , b and x cover all but 2 vertices in A and as before we are done. $\square$

Lemma 4.7. *A graph $G = (V, E)$ with $|V| = 7$, $|E| \geq 10$, and $\Delta \leq 3$ satisfies $\gamma(G) \leq 2$.*

Proof. The graph $G = (V, E)$ must have six vertices of degree 3 and a single vertex of degree 2. Let v be a vertex of degree 3 which is adjacent to a vertex of degree 2. Let A denote the set of three vertices not adjacent to v . If $\langle A \rangle$ has two or more edges, then clearly G can be dominated by v together with a vertex of degree 2 taken from $\langle A \rangle$. So we may assume that $\langle A \rangle$ has at most one edge. The sum of the degrees (in G) of the vertices of A is 9 since the single vertex of degree 2 is not in A . Since $\langle A \rangle$ has at most one edge, there are at least $7 = 9 - 2$ edges between $N(v)$ and A . Hence at least one vertex, say x , in $N(v)$ must be incident with 3 of these edges. It follows that $\{x, v\}$ is a dominating set for G . $\square$

The restriction that $\Delta \leq 3$ in the preceding lemma is essential. A graph with

vertices. Begin with the tetrahedron K_4 . Add an additional vertex connecting it to any two vertices of the tetrahedron. Finally, add two additional vertices, connecting each to one of the remaining two vertices of the tetrahedron. This graph has 10 edges and 7 vertices, but has maximum degree 4 and minimum degree 1. The graphs described by the lemma had maximum degree 3 and minimum degree 2. There is also a graph with 7 vertices, 10 edges, minimum degree 2, maximum degree 4 with domination number 3.

Proposition 4.8. $\gamma_r(14, 6) = \gamma(14, 6) = 3$.

Proof. By Appendix B there is a regular $(14, 6)$ -graph whose domination number is 3. That $\gamma_r(14, 6) = \gamma(14, 6)$ follows from Proposition 3.1. It remains to prove that $\gamma_r(14, 6) \leq 3$.

Let $G = (V, E)$ be a regular $(14, 6)$ -graph. Since every closed neighborhood contains 7 vertices, $\gamma(G) = 2$ if there are disjoint closed neighborhoods in G . Thus we can assume that any two closed neighborhoods $N[x]$ and $N[y]$ intersect in at least one point. This implies that any two vertices can be covered by a single vertex. So if we can cover 12 vertices with just 2 vertices we will be done.

Let $v \in V$ and set $A = V - N[v]$. If $\gamma(\langle A \rangle) \leq 2$ we are clearly done. If $\langle A \rangle$ has a vertex of degree at least 4 it covers 5 vertices in addition to the 7 covered by v and we are done. So we can assume that the maximum degree of a vertex in $\langle A \rangle$ is 3. If there are at least 10 edges in $\langle A \rangle$ we are done by Lemma 4.7. Thus we may assume that $\langle A \rangle$ has at most 9 edges. An endpoint of an edge in G is either in A or $N(v)$. Since $\langle A \rangle$ has at most 9 edges and G is 6-regular there are at least $24 = |A| \cdot 6 - 2 \cdot 9$ edges from A to $N(v)$. If some vertex in $N(v)$ covers 5 or more of the vertices in A we are done. Thus each of the 6 vertices in $N(v)$ is incident with at most 4 of the vertices in A . This implies that there are exactly 24 edges between A and $N(v)$ and that each $x \in N(v)$ is incident to v and to exactly four vertices in A . It follows that $\langle N(v) \rangle$ is a 1-regular graph.

Thus we can assume that *for every* vertex $v \in V$ the graph $\langle N(v) \rangle$ is 1-regular. But we shall see that this is impossible: Let $v \in V$ and let $x, y \in N(v)$ be chosen

such that $x \sim y$. It is clear that

$$N[v] \cap N[x] \cap N[y] = N[v] \cap N[x] = N[v] \cap N[y] = \{v, x, y\}$$

as x and y are adjacent only to each other in $\langle N(v) \rangle$. Also we have

$$N[x] \cap N[y] = \{v, x, y\}.$$

For suppose, to the contrary, that $w \in N[x] \cap N[y]$ and $w \notin \{v, x, y\}$. Then $y \in N(x)$ is adjacent to both $v, w \in N(x)$ which contradicts the assumption that $\langle N(x) \rangle$ is 1-regular. But then we have

$$\begin{aligned} |N[v] \cup N[x] \cup N[y]| &= \\ &= |N[v]| + |N[x]| + |N[y]| \\ &\quad - |N[v] \cap N[x]| - |N[v] \cap N[y]| - |N[x] \cap N[y]| \\ &\quad + |N[v] \cap N[x] \cap N[y]| \\ &= 7 + 7 + 7 - 3 - 3 - 3 + 3 = 15, \end{aligned}$$

which cannot be since there are only 14 vertices in G . $\square$

Proposition 4.9. $\gamma(15, 5) = \gamma_r(15, 5) = 4$.

Proof. The almost regular graph $G(15, 5, 4)$ given in Appendix B has domination number 4, so it suffices to prove that $\gamma(15, 5) \leq 4$. Let $G = (V, E)$ be a $(15, 5)$ -graph. If G has a vertex of at least degree at least 8 then $\gamma(G) \leq 4$ by Proposition 3.1. Since the minimum degree and the order are odd, there must be a vertex, say, x of degree at least 6. Again by Proposition 3.1 there is a vertex y such that $|N[x] \cup N[y]| \geq 11$. If $|N[x] \cup N[y]| \geq 12$ then Proposition 3.1 shows that $\gamma(G) \leq 4$. So we can assume that $|N[x] \cup N[y]| = 11$. Let

$$A = \{a, b, c, d\} = V - (N[x] \cup N[y]),$$

$$B = (N[x] \cup N[y]) - \{x, y\},$$

and $H = \langle A \rangle$. If H has domination number 1 or 2 we are done. So we may assume that H has at most one edge. It follows that there are at least $4 + 5 + 2 = 11$ edges

with one end in B and the other end in A . For each vertex v in B let A_v denote the vertices in A adjacent to v . If any A_v has three or more elements we are clearly done. So we may assume that each A_v has at most 2 elements. But since there are at least 18 edges joining A to B we must have $|A_v| = 2$ for each $v \in B$. Now if $A_{v_1} \cap A_{v_2} = \emptyset$ then $A_{v_1} \cup A_{v_2} = A$ and so $\{x, y, v_1, v_2\}$ is a dominating set for G . Thus we may assume that $A_{v_1} \cap A_{v_2} \neq \emptyset$ for all v_1 and v_2 in B . By Lemma 4.0 the A_v 's have an element in common. But the common element would have degree at least 9, a case we already covered. This completes the proof. $\square$

Proposition 4.10. $\gamma_r(16, 4) = \gamma(16, 4) = 5$.

Proof. Since $\gamma_r(6, 4) = 2$ and $\gamma_r(10, 4) = 3$ there is, by Lemma 2.1, a regular graph whose domination number is 5. It remains to prove that $\gamma(16, 4) \leq 5$.

Given a $(16, 4)$ graph, we may assume that any two closed neighborhoods $N[x]$ and $N[y]$ intersect in at least one point. If they were disjoint, then Proposition 3.1 would apply with $|S| = 2$ and $\lambda = 10$ vertices covered.

Applying Proposition 3.1 directly with $|S| = 1$ and $\lambda = 5$ we obtain $g_4 = 2$. Thus, 4 vertices cover all but (at most) 2 vertices. The closed neighborhoods of the remaining two vertices are not disjoint and hence they can be covered by a single vertex yielding a dominating set of at most 5 vertices. $\square$

Proposition 4.11. $4 \leq \gamma_r(16, 5) = \gamma(16, 5) \leq 5$.

Proof. A regular $(16, 5)$ -graph whose domination number is 4 may be formed by taking the disjoint union of two regular $(8, 5)$ -graphs whose domination numbers are 2. That $\gamma(16, 5) \leq 5$ follows from Proposition 3.1. It will then suffice to prove that if a $(16, 5)$ -graph $G = (V, E)$ is not regular then $\gamma(G) = 4$.

We will show that for any $x, y \in V$, $N[x] \cap N[y] \neq \emptyset$. Given this, choose $x \in V$ of degree at least 6 and apply Proposition 3.1 with $S = \{x\}$ to conclude that $g_3 \leq 2$. This will mean that 3 vertices of G cover all but (at most) 2 vertices of G and the remaining 2 can be covered by a single vertex since their closed neighborhoods are not disjoint, completing the proof.

Let $x, y \in V$ and assume $N[x] \cap N[y] = \emptyset$. The vertices x and y both have degree 5 or we are done by Proposition 3.1 starting with $S = \{x, y\}$. Let $B = N(x) \cup N(y)$ and let $A = V - N[x] - N[y]$. Clearly $|A| = 4$ and $|B| = 10$. If A can be covered by two vertices, we are done and hence $\langle A \rangle$ has at most 1 edge. Thus, there are at least $4 \cdot 5 - 2 = 18$ edges from A to B . For each $v \in B$, let A_v denote the set of vertices in A which are adjacent to v . If there exists $v \in B$ such that $|A_v| \geq 3$, then at most one point z of A is not covered by v and $\{x, y, v, z\}$ is a dominating set. Hence we may assume that $|A_v| \leq 2$ for each $v \in B$. Let B' denote the set of vertices $v \in B$ for which $|A_v| = 2$. Since there are at least 18 edges joining B to A we must have $|B'| \geq 8$. Note that the A_v 's, $v \in B'$, must cover A for if a vertex in A is not adjacent to any vertex in B' then it has degree at most 3. If there exists $u, v \in B'$ such that A_u and A_v are disjoint then $\{x, y, u, v\}$ would be a dominating set. So it follows from Lemma 4.0 that the sets A_v , $v \in B'$, have a common vertex. This common element must be a vertex of degree at least 8. Given a vertex of degree 8, Proposition 3.1 applies to show that the domination number of G , in this case, is at most 4. $\square$

5. MISCELLANEOUS REMARKS

The following proposition tempts one to conjecture that $\gamma_r(n, 4) = \gamma(n, 4) = \lfloor \frac{n}{3} \rfloor$.

Proposition 5.1. *For $5 \leq n \leq 16$,*

$$\gamma_r(n, 4) = \gamma(n, 4) = \lfloor \frac{n}{3} \rfloor$$

and for $n \geq 17$,

$$\lfloor \frac{n}{3} \rfloor \leq \gamma_r(n, 4).$$

Proof. The first sentence follows from Appendix A. If $n \geq 16$ then we can write $n = 6\ell + s$ where $\ell \geq 1$ and $s \in \{6, 7, 8, 9, 10, 11\}$. Now by Lemma 2.1 we have

$$\lfloor \frac{n}{3} \rfloor = 2\ell + \lfloor \frac{s}{3} \rfloor = \ell\gamma_r(6, 4) + \gamma_r(s, 4) \leq \gamma_r(6\ell + s, 4) = \gamma_r(n, 4). \quad \square$$

One of the authors has conjectured that $\gamma_r(n, \delta) = \gamma(n, \delta)$ is always true. Indeed, no example to the contrary has been found in our search. Even when the value of

$\gamma(n, \delta)$ is not known, one is sometimes able to show that $\gamma_r(n, \delta) = \gamma(n, \delta)$, see, e.g., Proposition 4.11. The other author conjectures that $\gamma_r(n, \delta) = \gamma(n, \delta)$ is false. The discussion following Lemma 4.7 provides an example where a more uneven distribution of degree for the vertices of a graph leads to a higher value of γ even though the number of vertices and the number of edges remains the same.

Finally we remark that if graphs are assumed to be connected at least in the cases $\delta = 1$ or $\delta = 2$ different results may be expected. See, for example, [5] where it is proved that with a few exceptions the domination number of a connected $(n, 2)$ -graph is at most $2n/5$.

Acknowledgement. The authors are grateful to David Fisher, Teresa Haynes, Bill McCuaig, Boris Shekhtman and Stephen Suen for useful discussions and/or help with references. Special thanks to Brendan McKay for the use of his program makeg.

REFERENCES

1. I. Anderson, *Combinatorics of Finite Sets*, Clarendon Press, Oxford, New York, 1987..
2. W. E. Clark, D. Fisher, B. Shekhtman and S. Suen, *Upper bounds for the domination number of a graph*, preprint.
3. J. F. Fink, M. S. Jacobson, L. K. Kinch and J. Roberts, *On graphs having domination number half their order*, Period. Math. Hungar. **16** (1985), 287–293..
4. T. W. Haynes, *Domination in graphs: a brief overview*, preprint.
5. W. McCuaig and B. Shepherd, *Domination in graphs with minimum degree two*, J. Graph Theory **13** (1989), 749–762.
6. B. D. McKay, *Isomorph-free exhaustive generation*, preprint.
7. O. Ore, *Theory of Graphs*, Amer. Math. Soc. Colloq. Publ. 38, American Mathematical Society, Providence, 1962..
8. C. Payan and N. H. Xuong, *Domination-balanced graphs*, J. Graph Theory **6** (1982), 23–32..
9. B. Reed, *Paths, stars, and the number three*, Combin. Probab. and Comput. **5** (1996), 277–295.

APPENDIX A

Table 3. Derivation of $\gamma(n, \delta)$ for $n \leq 16$, $0 \leq \delta \leq n - 1$.

$n \setminus \delta$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	2.3															
2	2.3	2.3														
3	2.3	2.3	2.4													
4	2.3	2.3	2.4	3.1												
5	2.3	2.3	2.4	3.1	3.1											
6	2.3	2.3	2.4	3.1	3.1	3.1										
7	2.3	2.3	2.4	3.1	3.1	3.1	3.1									
8	2.3	2.3	2.4	R	3.1	3.1	3.1	3.1								
9	2.3	2.3	2.4	R	3.1a	3.1	3.1	3.1	3.1							
10	2.3	2.3	2.4	R	3.1a	4.1c	3.1	3.1	3.1	3.1						
11	2.3	2.3	2.4	R	4.2c	3.1a	3.1a	3.1	3.1	3.1	3.1					
12	2.3	2.3	2.4	R	3.1b	3.1a	3.1a	4.3c	3.1	3.1	3.1	3.1				
13	2.3	2.3	2.4	R	Rd	4.4	3.1a	3.1a	4.5c	3.1	3.1	3.1	3.1			
14	2.3	2.3	2.4	R	4.6	3.1a	4.8e	3.1a	3.1a	3.1	3.1	3.1	3.1	3.1		
15	2.3	2.3	2.4	R	3.1f	4.9	3.1a	3.1a	3.1a	3.1	3.1	3.1	3.1	3.1	3.1	
16	2.3	2.3	2.4	R	4.10	3.1b	3.1b	3.1a	3.1a	3.1	3.1	3.1	3.1	3.1	3.1	3.1

The number in the (n, δ) -th cell is the number of the proposition which justifies the corresponding value in Table 1. The letter code is as follows:

- a With lower bound from Appendix B graph $G(n, \delta, \gamma)$
- b Also used Lemma 2.1 with $\gamma(n, \delta) = 2 * \gamma(n/2, \delta)$
- c Also required computer search
- d Also used Lemma 2.1 with $\gamma(7, 4) + \gamma(6, 4) = 4$
- e Confirmed by computer search using 11 days of cpu time
- f Also used Lemma 2.1 with $\gamma(9, 4) + \gamma(6, 4) = 5$
- R Reed's Theorem

APPENDIX B

Each graph $G(n, \delta, \gamma)$ listed below is an (n, δ) -graph with domination number γ . All graphs are either regular or almost regular. The vertices are numbered $0, 1, \dots, n-1$ and the i -th row represents the neighbors of the vertex in the first entry of that row. These graphs were found by Brendan McKay's program makeg augmented by a program to check domination numbers or by *ad hoc* construction in two case.

$$G(9, 3, 3) = \begin{bmatrix} 0 & 1 & 2 & 4 \\ 1 & 0 & 2 & 3 \\ 2 & 0 & 1 & 3 \\ 3 & 1 & 2 & 4 \\ 4 & 0 & 3 & 5 \\ 5 & 4 & 6 & 7 \\ 6 & 5 & 7 & 8 \\ 7 & 5 & 6 & 8 \\ 8 & 4 & 6 & 7 \end{bmatrix} \quad G(11, 3, 4) = \begin{bmatrix} 0 & 1 & 2 & 5 \\ 1 & 0 & 2 & 4 \\ 2 & 0 & 1 & 3 \\ 3 & 2 & 4 & 7 \\ 4 & 1 & 3 & 5 \\ 5 & 0 & 4 & 6 \\ 6 & 5 & 8 & 9 \\ 7 & 3 & 8 & 9 \\ 8 & 6 & 7 & 10 \\ 9 & 6 & 7 & 10 \\ 10 & 4 & 8 & 9 \end{bmatrix} \quad 10$$

$$G(9, 4, 3) = \begin{bmatrix} 0 & 1 & 2 & 5 & 8 \\ 1 & 0 & 2 & 4 & 7 \\ 2 & 0 & 1 & 3 & 6 \\ 3 & 2 & 4 & 5 & 8 \\ 4 & 1 & 3 & 5 & 7 \\ 5 & 0 & 3 & 4 & 6 \\ 6 & 2 & 5 & 7 & 8 \\ 7 & 1 & 4 & 6 & 8 \\ 8 & 0 & 3 & 6 & 7 \end{bmatrix} \quad G(10, 4, 3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 9 \\ 1 & 0 & 2 & 3 & 8 \\ 2 & 0 & 1 & 3 & 5 \\ 3 & 0 & 1 & 2 & 4 \\ 4 & 3 & 5 & 6 & 7 \\ 5 & 2 & 4 & 6 & 7 \\ 6 & 4 & 5 & 8 & 9 \\ 7 & 4 & 5 & 8 & 9 \\ 8 & 1 & 6 & 7 & 9 \\ 9 & 0 & 6 & 7 & 8 \end{bmatrix}$$

$$G(11, 5, 3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 5 & 9 \\ 1 & 0 & 2 & 3 & 5 & 8 \\ 2 & 0 & 1 & 3 & 4 & 7 \\ 3 & 0 & 1 & 2 & 4 & 7 \\ 4 & 2 & 3 & 5 & 6 & 8 \\ 5 & 0 & 1 & 4 & 6 & 10 \\ 6 & 4 & 5 & 7 & 8 & 9 \\ 7 & 2 & 3 & 6 & 9 & 10 \\ 8 & 1 & 4 & 6 & 9 & 10 \\ 9 & 0 & 6 & 7 & 8 & 10 \\ 10 & 4 & 5 & 7 & 8 & 9 \end{bmatrix} \quad 10 \quad G(11, 6, 3) = \begin{bmatrix} 0 & 1 & 2 & 4 & 7 & 9 & 10 \\ 1 & 0 & 2 & 3 & 5 & 7 & 10 \\ 2 & 0 & 1 & 3 & 4 & 6 & 8 \\ 3 & 1 & 2 & 4 & 5 & 8 & 9 \\ 4 & 0 & 2 & 3 & 5 & 6 & 9 \\ 5 & 1 & 3 & 4 & 6 & 7 & 10 \\ 6 & 2 & 4 & 5 & 7 & 8 & 10 \\ 7 & 0 & 1 & 5 & 6 & 8 & 9 \\ 8 & 2 & 3 & 6 & 7 & 9 & 10 \\ 9 & 0 & 3 & 4 & 7 & 8 & 10 \\ 10 & 0 & 1 & 5 & 6 & 8 & 9 \end{bmatrix}$$

$$G(12, 5, 3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 9 & 10 \\ 1 & 0 & 2 & 3 & 4 & 8 \\ 2 & 0 & 1 & 3 & 4 & 5 \\ 3 & 0 & 1 & 2 & 4 & 5 \\ 4 & 1 & 2 & 3 & 5 & 11 \\ 5 & 2 & 3 & 4 & 6 & 7 \\ 6 & 5 & 7 & 8 & 10 & 11 \\ 7 & 5 & 6 & 8 & 9 & 11 \\ 8 & 1 & 6 & 7 & 9 & 10 \\ 9 & 0 & 7 & 8 & 10 & 11 \\ 10 & 0 & 6 & 8 & 9 & 11 \end{bmatrix} \quad G(12, 6, 3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 5 & 7 & 11 \\ 1 & 0 & 2 & 3 & 4 & 8 & 10 \\ 2 & 0 & 1 & 3 & 4 & 7 & 11 \\ 3 & 0 & 1 & 2 & 4 & 6 & 9 \\ 4 & 1 & 2 & 3 & 5 & 6 & 10 \\ 5 & 0 & 4 & 6 & 7 & 10 & 11 \\ 6 & 3 & 4 & 5 & 7 & 8 & 9 \\ 7 & 0 & 2 & 5 & 6 & 8 & 9 \\ 8 & 1 & 6 & 7 & 9 & 10 & 11 \\ 9 & 3 & 6 & 7 & 8 & 10 & 11 \\ 10 & 1 & 4 & 5 & 8 & 9 & 11 \end{bmatrix}$$

$$G(13,6,3) = \begin{bmatrix} 0 & 6 & 7 & 8 & 9 & 10 & 11 \\ 1 & 6 & 7 & 8 & 9 & 10 & 11 \\ 2 & 6 & 8 & 9 & 10 & 11 & 12 \\ 3 & 6 & 8 & 9 & 10 & 11 & 12 \\ 4 & 7 & 8 & 9 & 10 & 11 & 12 \\ 5 & 7 & 8 & 9 & 10 & 11 & 12 \\ 6 & 0 & 1 & 2 & 3 & 7 & 12 \\ 7 & 0 & 1 & 4 & 5 & 6 & 12 \\ 8 & 0 & 1 & 2 & 3 & 4 & 5 \\ 9 & 0 & 1 & 2 & 3 & 4 & 5 \\ 10 & 0 & 1 & 2 & 3 & 4 & 5 \\ 11 & 0 & 1 & 2 & 3 & 4 & 5 \\ 12 & 2 & 3 & 4 & 5 & 6 & 7 \end{bmatrix} \quad G(13,5,3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 4 & 6 \\ 1 & 0 & 2 & 3 & 4 & 5 \\ 2 & 0 & 1 & 3 & 4 & 5 \\ 3 & 0 & 1 & 2 & 4 & 5 \\ 4 & 0 & 1 & 2 & 3 & 5 \\ 5 & 1 & 2 & 3 & 4 & 6 \\ 6 & 0 & 5 & 7 & 8 & 9 & 12 \\ 7 & 6 & 8 & 10 & 11 & 12 \\ 8 & 6 & 7 & 9 & 10 & 11 \\ 9 & 6 & 8 & 10 & 11 & 12 \\ 10 & 7 & 8 & 9 & 11 & 12 \\ 11 & 7 & 8 & 9 & 10 & 12 \\ 12 & 6 & 7 & 9 & 10 & 11 \end{bmatrix}$$

$$G(13,4,4) = \begin{bmatrix} 0 & 1 & 2 & 3 & 12 \\ 1 & 0 & 2 & 3 & 10 \\ 2 & 0 & 1 & 3 & 5 \\ 3 & 0 & 1 & 2 & 4 \\ 4 & 3 & 6 & 7 & 11 \\ 5 & 2 & 6 & 7 & 11 \\ 6 & 4 & 5 & 8 & 9 \\ 7 & 4 & 5 & 8 & 9 \\ 8 & 6 & 7 & 10 & 12 \\ 9 & 6 & 7 & 10 & 12 \\ 10 & 1 & 8 & 9 & 11 \\ 11 & 4 & 5 & 10 & 12 \\ 12 & 0 & 8 & 9 & 11 \end{bmatrix} \quad G(13,7,3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 6 & 7 & 8 & 12 \\ 1 & 0 & 2 & 3 & 4 & 6 & 10 & 12 \\ 2 & 0 & 1 & 3 & 4 & 5 & 10 & 11 \\ 3 & 0 & 1 & 2 & 4 & 5 & 8 & 9 \\ 4 & 1 & 2 & 3 & 5 & 6 & 7 & 9 \\ 5 & 2 & 3 & 4 & 6 & 7 & 11 & 12 \\ 6 & 0 & 1 & 4 & 5 & 7 & 8 & 11 & 12 \\ 7 & 0 & 4 & 5 & 6 & 8 & 9 & 10 \\ 8 & 0 & 3 & 6 & 7 & 9 & 10 & 11 \\ 9 & 3 & 4 & 7 & 8 & 10 & 11 & 12 \\ 10 & 1 & 2 & 7 & 8 & 9 & 11 & 12 \\ 11 & 2 & 5 & 6 & 8 & 9 & 10 & 12 \\ 12 & 0 & 1 & 5 & 6 & 9 & 10 & 11 \end{bmatrix}$$

$$G(14,6,3) = \begin{bmatrix} 0 & 6 & 7 & 8 & 9 & 10 & 11 \\ 1 & 6 & 7 & 9 & 10 & 12 & 13 \\ 2 & 7 & 8 & 9 & 10 & 11 & 12 \\ 3 & 7 & 9 & 10 & 11 & 12 & 13 \\ 4 & 8 & 9 & 10 & 11 & 12 & 13 \\ 5 & 8 & 9 & 10 & 11 & 12 & 13 \\ 6 & 0 & 1 & 8 & 11 & 12 & 13 \\ 7 & 0 & 1 & 2 & 3 & 8 & 13 \\ 8 & 0 & 2 & 4 & 5 & 6 & 7 \\ 9 & 0 & 1 & 2 & 3 & 4 & 5 \\ 10 & 0 & 1 & 2 & 3 & 4 & 5 \\ 11 & 0 & 2 & 3 & 4 & 5 & 6 \\ 12 & 1 & 2 & 3 & 4 & 5 & 6 \\ 13 & 1 & 3 & 4 & 5 & 6 & 7 \end{bmatrix} \quad G(14,5,4) = \begin{bmatrix} 0 & 5 & 7 & 8 & 9 & 13 \\ 1 & 6 & 7 & 8 & 10 & 11 \\ 2 & 6 & 7 & 9 & 10 & 11 \\ 3 & 9 & 10 & 11 & 12 & 13 \\ 4 & 9 & 10 & 11 & 12 & 13 \\ 5 & 0 & 8 & 10 & 11 & 12 \\ 6 & 1 & 2 & 8 & 12 & 13 \\ 7 & 0 & 1 & 2 & 12 & 13 \\ 8 & 0 & 1 & 5 & 6 & 9 \\ 9 & 0 & 2 & 3 & 4 & 8 \\ 10 & 1 & 2 & 3 & 4 & 5 \\ 11 & 1 & 2 & 3 & 4 & 5 \\ 12 & 3 & 4 & 5 & 6 & 7 \\ 13 & 0 & 3 & 4 & 6 & 7 \end{bmatrix}$$

$$G(14,4,4) = \begin{bmatrix} 0 & 7 & 8 & 9 & 10 \\ 1 & 7 & 8 & 9 & 11 \\ 2 & 7 & 8 & 12 & 13 \\ 3 & 7 & 8 & 12 & 13 \\ 4 & 9 & 10 & 11 & 12 \\ 5 & 9 & 10 & 11 & 13 \\ 6 & 10 & 11 & 12 & 13 \\ 7 & 0 & 1 & 2 & 3 \\ 8 & 0 & 1 & 2 & 3 \\ 9 & 0 & 1 & 4 & 5 \\ 10 & 0 & 4 & 5 & 6 \\ 11 & 1 & 4 & 5 & 6 \\ 12 & 2 & 3 & 4 & 6 \end{bmatrix} \quad G(14,7,3) = \begin{bmatrix} 0 & 5 & 6 & 7 & 8 & 9 & 12 & 13 \\ 1 & 6 & 8 & 9 & 10 & 11 & 12 & 13 \\ 2 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 3 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 4 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 5 & 0 & 6 & 8 & 10 & 11 & 12 & 13 \\ 6 & 0 & 1 & 5 & 7 & 9 & 10 & 11 \\ 7 & 0 & 2 & 3 & 4 & 6 & 10 & 11 \\ 8 & 0 & 1 & 2 & 3 & 4 & 5 & 13 \\ 9 & 0 & 1 & 2 & 3 & 4 & 6 & 12 \\ 10 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 11 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 12 & 0 & 1 & 2 & 3 & 4 & 5 & 9 \end{bmatrix}$$

$$G(14, 8, 3) = \begin{bmatrix} 0 & 4 & 5 & 6 & 7 & 8 & 11 & 12 & 13 \\ 1 & 4 & 5 & 6 & 7 & 9 & 10 & 11 & 13 \\ 2 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 3 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 4 & 0 & 1 & 6 & 8 & 9 & 10 & 12 & 13 \\ 5 & 0 & 1 & 7 & 8 & 9 & 10 & 11 & 12 \\ 6 & 0 & 1 & 2 & 3 & 4 & 10 & 11 & 12 \\ 7 & 0 & 1 & 2 & 3 & 5 & 10 & 12 & 13 \\ 8 & 0 & 2 & 3 & 4 & 5 & 10 & 11 & 13 \\ 9 & 1 & 2 & 3 & 4 & 5 & 11 & 12 & 13 \\ 10 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 11 & 0 & 1 & 2 & 3 & 5 & 6 & 8 & 9 \\ 12 & 0 & 2 & 3 & 4 & 5 & 6 & 7 & 9 \\ 13 & 0 & 1 & 2 & 3 & 4 & 7 & 8 & 9 \end{bmatrix} \quad G(15, 5, 4) = \begin{bmatrix} 0 & 7 & 8 & 9 & 5 & 14 \\ 1 & 7 & 8 & 10 & 6 & 14 \\ 2 & 7 & 9 & 10 & 11 & 6 \\ 3 & 9 & 10 & 11 & 12 & 13 \\ 4 & 9 & 10 & 11 & 12 & 13 \\ 5 & 0 & 8 & 10 & 11 & 12 \\ 6 & 1 & 2 & 8 & 12 & 13 \\ 7 & 0 & 1 & 2 & 12 & 13 \\ 8 & 0 & 1 & 9 & 5 & 6 \\ 9 & 0 & 2 & 8 & 3 & 4 \\ 10 & 1 & 2 & 3 & 4 & 5 \\ 11 & 2 & 3 & 4 & 5 & 14 \\ 12 & 3 & 4 & 5 & 6 & 7 & 14 \\ 13 & 7 & 3 & 4 & 6 & 14 \\ 14 & 0 & 1 & 11 & 12 & 13 \end{bmatrix}$$

$$G(15, 6, 3) = \begin{bmatrix} 0 & 7 & 8 & 9 & 10 & 11 & 12 \\ 1 & 7 & 8 & 9 & 12 & 13 & 14 \\ 2 & 7 & 10 & 11 & 12 & 13 & 14 \\ 3 & 8 & 9 & 10 & 11 & 12 & 13 \\ 4 & 8 & 9 & 10 & 11 & 12 & 14 \\ 5 & 8 & 9 & 10 & 11 & 13 & 14 \\ 6 & 8 & 9 & 10 & 11 & 13 & 14 \\ 7 & 0 & 1 & 2 & 12 & 13 & 14 \\ 8 & 0 & 1 & 3 & 4 & 5 & 6 \\ 9 & 0 & 1 & 3 & 4 & 5 & 6 \\ 10 & 0 & 2 & 3 & 4 & 5 & 6 \\ 11 & 0 & 2 & 3 & 4 & 5 & 6 \\ 12 & 0 & 1 & 2 & 3 & 4 & 7 \\ 13 & 1 & 2 & 3 & 5 & 6 & 7 \\ 14 & 1 & 2 & 4 & 5 & 6 & 7 \end{bmatrix} \quad G(15, 7, 3) = \begin{bmatrix} 0 & 1 & 2 & 3 & 4 & 5 & 7 & 8 \\ 1 & 0 & 2 & 3 & 4 & 5 & 7 & 8 \\ 2 & 0 & 1 & 3 & 4 & 5 & 6 & 13 \\ 3 & 0 & 1 & 2 & 4 & 5 & 6 & 12 \\ 4 & 0 & 1 & 2 & 3 & 5 & 6 & 11 \\ 5 & 0 & 1 & 2 & 3 & 4 & 6 & 10 \\ 6 & 2 & 3 & 4 & 5 & 7 & 9 & 14 \\ 7 & 0 & 1 & 6 & 8 & 9 & 10 & 11 & 14 \\ 8 & 0 & 1 & 7 & 9 & 12 & 13 & 14 \\ 9 & 6 & 7 & 8 & 10 & 11 & 12 & 13 \\ 10 & 5 & 7 & 9 & 11 & 12 & 13 & 14 \\ 11 & 4 & 7 & 9 & 10 & 12 & 13 & 14 \\ 12 & 3 & 8 & 9 & 10 & 11 & 13 & 14 \\ 13 & 2 & 8 & 9 & 10 & 11 & 12 & 14 \\ 14 & 6 & 7 & 8 & 10 & 11 & 12 & 13 \end{bmatrix}$$

$$G(15, 8, 3) = \begin{bmatrix} 0 & 5 & 7 & 9 & 10 & 11 & 12 & 13 & 14 \\ 1 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 2 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 3 & 6 & 7 & 8 & 10 & 11 & 12 & 13 & 14 \\ 4 & 6 & 7 & 8 & 10 & 11 & 12 & 13 & 14 \\ 5 & 0 & 8 & 9 & 10 & 11 & 12 & 13 & 14 \\ 6 & 1 & 2 & 3 & 4 & 9 & 11 & 12 & 14 \\ 7 & 0 & 1 & 2 & 3 & 4 & 9 & 10 & 14 \\ 8 & 1 & 2 & 3 & 4 & 5 & 9 & 13 & 14 \\ 9 & 0 & 1 & 2 & 5 & 6 & 7 & 8 & 14 \\ 10 & 0 & 1 & 2 & 3 & 4 & 5 & 7 & 12 \\ 11 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 13 \\ 12 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 10 \\ 13 & 0 & 1 & 2 & 3 & 4 & 5 & 8 & 11 \\ 14 & 0 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \end{bmatrix} \quad G(16, 8, 3) = \begin{bmatrix} 0 & 5 & 7 & 9 & 10 & 11 & 12 & 13 & 14 \\ 1 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 2 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 \\ 3 & 6 & 7 & 8 & 10 & 11 & 12 & 13 & 14 \\ 4 & 6 & 7 & 8 & 10 & 11 & 12 & 13 & 14 \\ 5 & 0 & 8 & 9 & 10 & 11 & 13 & 14 & 15 \\ 6 & 1 & 2 & 3 & 4 & 9 & 12 & 14 & 15 \\ 7 & 0 & 1 & 2 & 3 & 4 & 9 & 14 & 15 \\ 8 & 1 & 2 & 3 & 4 & 5 & 13 & 14 & 15 \\ 9 & 0 & 1 & 2 & 5 & 6 & 7 & 14 & 15 \\ 10 & 0 & 1 & 2 & 3 & 4 & 5 & 12 & 15 \\ 11 & 0 & 1 & 2 & 3 & 4 & 5 & 13 & 15 \\ 12 & 0 & 1 & 2 & 3 & 4 & 6 & 10 & 15 \\ 13 & 0 & 1 & 2 & 3 & 4 & 5 & 8 & 11 \\ 14 & 0 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 15 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \end{bmatrix}$$