

Short Score Certificates for Upset Tournaments

Jeffrey L. Poet and Bryan L. Shader
Department of Mathematics
University of Wyoming
Laramie, Wyoming 82071,
jpoet@uwyo.edu, bshader@uwyo.edu.

Submitted: January 15, 1998; Accepted: May 5, 1998.

Abstract

A score certificate for a tournament, T , is a collection of arcs of T which can be uniquely completed to a tournament with the same score-list as T 's, and the score certificate number of T is the least number of arcs in a score certificate of T . Upper bounds on the score certificate number of upset tournaments are derived. The upset tournaments on n vertices are in one-to-one correspondence with the ordered partitions of $n-3$, and are “almost” transitive tournaments. For each upset tournament on n vertices a general construction of a score certificate with at most $2n-3$ arcs is given. Also, for the upset tournament, T_λ , corresponding to the ordered partition λ , a score certificate with at most $n+2k+3$ arcs is constructed, where k is the number of parts of λ of size at least 2. Lower bounds on the score certificate number of T_λ in the case that each part is sufficiently large are derived. In particular, the score certificate number of the so-called nearly transitive tournament on n vertices is shown to be $n+3$, for $n \geq 10$.

1 Introduction

Some recent research has been concerned with the problem of efficiently conveying the information contained in a binary relation [AR, KTF, R]. In this paper we continue this line of research by studying the problem for a specific

class of binary relations. Throughout we use the graph-theoretic notation and terminology in [CL].

Let $V = \{v_1, v_2, \dots, v_n\}$. A *tournament*, T , on V is a directed graph obtained by replacing each edge of the complete graph on V with a directed arc. Thus, a tournament represents a complete, asymmetric, irreflexive binary relation. Throughout we use the notation and terminology for tournaments in [M]. We denote the arc from vertex v_i to vertex v_j by the ordered pair (v_i, v_j) . The *score* of a vertex $v_j \in V$ is its outdegree. The *score-list* of T is the multiset of outdegrees of the vertices of T .

In this paper we are concerned with efficiently conveying the information contained in a tournament. More precisely, a *score certificate* for T is a subset, D , of arcs such that T is the unique tournament on V that contains each arc in D and whose score-list is the same as T 's. In other words, a score certificate for T is a collection, D , of arcs with the property that T can be unambiguously determined by knowing the arcs of D and the score-list of T . The *score certificate number* of T is the least number of arcs in a score certificate for T , and is denoted by $sc(T)$.

Score certificates were first studied in [FKT, KTF], where it is proven that if T is not a regular tournament on 3 or 5 vertices, then

$$n - 1 \leq sc(T). \quad (1)$$

Thus, to convey the results of a tournament on n vertices (via score certificates) at least $n - 1$ arcs are required in all but the two exceptional cases. Note that the score certificate number of the transitive tournament on n vertices is $n - 1$. In [AR], it is shown that there exists an $\epsilon > 0$ independent of n such that

$$sc(T) \leq (1/2 - \epsilon)n^2 \quad (2)$$

for all tournaments T of order n . Also, in [AR] it is noted that there exist tournaments on n vertices with score certificates at least $(7/24 + o(1))n^2$. In [FKT] it is shown that

$$n - 1 \leq sc(N_n) \leq 3n - 9,$$

where N_n is the tournament with vertices $\{v_1, v_2, \dots, v_n\}$, and arcs (v_1, v_n) and (v_i, v_j) for all other $\{v_i, v_j\}$ with $i > j$. The tournament N_n is called the *nearly transitive tournament* on n vertices. Other than these results, little is known about the score certificate number.

In this paper, we study the score certificate number for a special family of tournaments which generalize the nearly transitive tournament. An *upset tournament* on $n \geq 4$ vertices is a tournament whose score-list is

$$\{1, 1, 2, 3, \dots, n-4, n-3, n-2, n-2\}.$$

Properties of upset tournaments have been studied in [BL].

In Section 2, we show that

$$sc(T) \leq 2n - 3 \quad (3)$$

for each upset tournament T on n vertices. The upset tournaments on n vertices possess special structural properties. In particular, they each have a unique hamiltonian cycle and they are in one-to-one correspondence with the ordered partitions of $n - 3$. In Section 2, we also give an upper bound on the score certificate number of an upset tournament which is in terms of the corresponding ordered partition. In most cases, this bound improves that of (3). In Section 3, we derive lower bounds for a special family of upset tournaments. As a consequence we show that the score certificate number of the nearly transitive tournament on n vertices equals $n + 3$, for $n \geq 10$.

We conclude this introductory section with necessary preliminaries concerning upset tournaments. Clearly, a transitive tournament is not strongly connected and has no 3-cycles. It follows from results in [M] that each upset tournament on n vertices has exactly $n - 2$ 3-cycles, and this is the least among strongly connected tournaments on n vertices. Thus, one can view the upset tournaments as a class of strongly connected tournaments that are “almost” transitive. Since equality holds in (1) for the transitive tournaments, it is natural to examine the score certificate number of upset tournaments.

It follows from the results in [BL] that the upset tournaments have a special structure, which we now describe. An *ordered partition* of the positive integer ℓ is a tuple of positive integers whose sum is ℓ . An ordered partition $\lambda = (p_1, p_2, \dots, p_k)$ of $n - 3$ determines a tournament T_λ on n vertices as follows. Let $V = \{v_1, \dots, v_n\}$ and let v_i and v_j be vertices with $i < j$. If (v_i, v_j) is (v_1, v_2) , (v_{n-1}, v_n) or one of

$$(v_2, v_{2+p_1}), (v_{2+p_1}, v_{2+p_1+p_2}), \dots, (v_{2+p_1+\dots+p_{k-1}}, v_{2+p_1+\dots+p_{k-1}+p_k}),$$

then (v_i, v_j) is an arc of T_λ . Otherwise, (v_j, v_i) is an arc of T_λ . It is easy to verify that T_λ is an upset tournament. For example, the nearly transitive

tournament on n vertices is isomorphic to $T_{(n-3)}$. It follows from results in [BL] that the nonisomorphic upset tournaments of order n are precisely the T_λ where λ runs over all ordered partitions of $n-3$. Hence, there are exactly 2^{n-4} non-isomorphic upset tournaments of order n .

An arc (v_i, v_j) of a tournament T is an *upset arc* provided the score of v_i is less than or equal to that of v_j . The upset arcs of T_λ are precisely the arcs (v_1, v_2) , (v_{n-1}, v_n) , and $(v_{2+p_1+p_2+\dots+p_j}, v_{2+p_1+p_2+\dots+p_j+p_{j+1}})$ for $j = 0, 1, 2, \dots, k-1$. Note that upset arcs of an upset tournament form a path, which we call its *upset path*.

An upset tournament T with vertices $v_1, \dots, v_n$ is in *standard form* provided (v_i, v_j) with $i < j$ is an arc of T if and only if (v_i, v_j) is in T 's upset path. Thus, each T_λ is in standard form, and each upset tournament can be put into standard form by relabelling its vertices.

Pictorially, each upset tournament, T , in standard form can be constructed by placing the vertices $v_1, v_2, \dots, v_n$ in order from bottom to top on a vertical line, drawing an path from v_1 to v_n that goes through v_2 and v_{n-1} and consists entirely of upward-oriented arcs, and orienting all other arcs downward. Moreover, up to isomorphism, the upset tournaments are precisely the tournaments which have this type of picture. For example, the upset tournaments with 6 vertices (in standard form) are given in Figure 1. Here, as is customary when illustrating tournaments, only the upward-oriented arcs are shown.




Figure 1

2 Upper bounds

In this section we derive two linear upper bounds on the score certificate number of an arbitrary upset tournament on n vertices. The first does not depend upon the partition corresponding to the tournament, but the second does. We begin by describing certain arcs that are contained in each score certificate of an upset tournament.

Let T be an upset tournament of order n . We define a *forced arc* of T to be an arc (v_i, v_j) of T where the score of v_i is 1 more than the score of v_j . If (v_i, v_j) is a forced arc, then the tournament obtained from T by reversing this arc is also an upset tournament. Thus, every score certificate of T contains each forced arc of T .

It is easy to determine the forced arcs of the upset tournament T_λ corresponding to the ordered partition $\lambda = (p_1, p_2, \dots, p_k)$ of $n - 3$. When $n = 4$, all non-upset arcs are forced arcs. Otherwise the forced arcs of T_λ are (v_n, v_{n-2}) , (v_3, v_1) , and each of the arcs in each of the paths

$$v_{2+p_1+\dots+p_{j-1}+p_j}, v_{2+p_1+\dots+p_{j-1}+p_j-1}, v_{2+p_1+\dots+p_{j-1}+p_j-2}, \dots, v_{2+p_1+\dots+p_{j-1}}$$

where $p_j \geq 2$. For example, the forced arcs of the nearly transitive tournament on $n \geq 5$ vertices are the arcs (v_n, v_{n-2}) , (v_3, v_1) , and the arcs in the path $v_{n-1}, v_{n-2}, \dots, v_2$.

We call the two vertices of an upset tournament with outdegree 1 the *bottom vertices* and the two vertices with indegree 1 the *top vertices*. The following shows that each upset tournament can be unambiguously determined given its bottom vertices and any collection of its arcs which contains each upset arc and each forced arc.

Lemma 2.1 *Let T be an upset tournament in standard form with vertices $v_1, v_2, \dots, v_n$. Let α be a subset of arcs of T containing each forced arc and each upset arc. Let T' be an upset tournament with vertices $v_1, v_2, \dots, v_n$ such that v_1 and v_2 are the bottom vertices of T' and each arc in α is an arc of T' . Then $T' = T$.*

Proof. Let D be the digraph with vertices $v_1, v_2, \dots, v_n$ whose arcs are those in α , and let γ be the set of arcs in T that are not in T' . Note that T' is the tournament obtained from T by reversing the arcs in γ .

We claim that γ is empty, and hence $T' = T$. For suppose not. Since either (v_j, v_{j+1}) or (v_{j+1}, v_j) is in α for $j = 1, \dots, n-1$ and α contains T 's upset arcs, all arcs in γ have the form (v_j, v_i) where $j-1 > i$. Among all the arcs in γ choose one, say (v_j, v_i) , such that i is minimum. Since v_1 and v_2 are the bottom vertices of T and of T' , $i \notin \{1, 2\}$. Since neither v_{n-1} nor v_n is a terminal vertex of an arc in γ , $i \leq n-2$. Thus, the score of v_i in T is $i-1$.

We claim that no vertex, v_k , in T' has score equal to $i-1$. If $k < i$, then the choice of (v_j, v_i) implies that the scores of v_k in T and in T' are equal, and hence less than $i-1$. If $k = i$, then the score of v_i in T is less than its score in T' . Suppose that $i < k < n$. Then the score of v_k in T is $k-1$. There are at most $k-i-1$ arcs in γ of the form (v_k, v_ℓ) where $k-1 > \ell \geq i$. Thus, the score of v_k in T' is at least $k-1-(k-i-1) = i$. Finally, suppose that $k = n$. Then the score of v_k in T is $n-2$. The arcs (v_n, v_{n-1}) , and (v_n, v_{n-2}) are not in γ . It follows that the score of v_n in T' is at least $n-2-(n-i-2) = i$. Therefore, we have reached the contradiction that no vertex in T' has score $i-1$.

We conclude that γ is empty, and hence that T is the unique upset tournament which contains α and has bottom vertices v_1 and v_2 . ■

We now use Lemma 2.1 to construct a score certificate with $2n-3$ arcs for each upset tournament on n vertices.

Theorem 2.2 *Let T be an upset tournament on n vertices. Then*

$$sc(T) \leq 2n-3.$$

Proof. Without loss of generality we may assume that T is in standard form. It is easy to verify the result for $T = T_{(1)}$, $T = T_{(1,1)}$ and $T = T_{(2)}$. Hence, we may assume that $n \geq 6$.

Let α be the set consisting of the upset arcs and the forced arcs of T . Consider the digraph, D , whose vertices are those of T and whose arcs are those in α . We first determine the outdegree of each vertex in D .

The vertices v_1 , v_2 and v_n each have outdegree 1 in D . The outdegree of v_3 in D is 2. If v_j is a vertex which is not on the upset path and $4 \leq j < n$, then (v_j, v_{j-1}) is a forced arc of T and v_j has outdegree 1 in D . Otherwise, v_j is a vertex on the upset path with $4 \leq j < n$. Let (v_i, v_j) be the upset arc

whose terminal vertex is v_j . If $j - i = 1$, then the outdegree of v_j in D is 1. If $j - i \neq 1$, then (v_j, v_{j-1}) is a forced arc of T , and the outdegree of v_j in D is 2.

Since $n \geq 6$, it can be verified that at least one of v_4 or v_5 has outdegree 1 in D . Let m be the smallest index such that $m \notin \{1, 2, n\}$ and v_m has outdegree 1 in D . Note that $m \geq 4$. Let

$$\beta = \{(v_j, v_{j-2}) : m < j < n, \text{ and } v_j \text{ has outdegree 1 in } D\} \cup \{(v_n, v_{n-3})\}.$$

It is easy to verify that β is a collection of arcs in T that are not in α . Let D' be the digraph whose vertices are $v_1, v_2, \dots, v_n$ and whose arcs are those in $\alpha \cup \beta$. By construction, each vertex in D' other than v_1, v_2, v_m has outdegree 2, and each of these 3 vertices has outdegree 1. Hence $\alpha \cup \beta$ has $2n - 3$ arcs.

We show that $\alpha \cup \beta$ is a score certificate for T . Suppose that T' is an upset tournament with vertices $v_1, v_2, \dots, v_n$ which has all the arcs in $\alpha \cup \beta$. We show that T' is T .

The vertices of outdegree 1 in D' are v_1, v_2 , and v_m . Also, (v_1, v_2) is an arc of D' , the arc in D' with initial vertex v_2 does not terminate at v_m , and the arc in D' with initial vertex at v_m does not terminate at v_1 or v_2 . If the outdegree of v_m in T' is 1, then (v_1, v_m) and (v_2, v_m) are arcs of T' , and we are led to the contradiction that v_m is the only vertex in T' with outdegree 1. Hence v_1 and v_2 are the bottom vertices of T' . Lemma 2.1 now implies that $T' = T$, and thus $\alpha \cup \beta$ is a score certificate for T . ■

Later we will show that the bound in Theorem 2.2 is not sharp in general. The construction of score certificates for upset tournaments given in the proof of Theorem 2.2 makes little use of the special structure of upset tournaments. We now present a construction which exploits the following property.

Proposition 2.3 *Let T be an upset tournament on n vertices. Then T contains a unique hamiltonian cycle.*

Proof. Without loss of generality we may assume that the vertices of T are labelled so that T is in standard form. Since T is strongly connected, it is well-known (see [M]) that T has a hamiltonian cycle.

Let H denote a hamiltonian cycle in T . The upset path must be a subpath of H , since it is the unique path from v_1 to v_n . All other arcs (v_i, v_j) in T have

$i > j$. Thus, the subpath of H from v_n to v_1 must be the path which starts at v_n , traverses through the vertices not on the upset path in decreasing order, and ends at v_1 . Therefore, H is unique. ■

Let T be a tournament with a hamiltonian cycle H . For vertices u and v , we denote the subpath of H from u to v by uHv . If (v_i, v_j) and (v_k, v_ℓ) are arcs of T not in H , we say that (v_k, v_ℓ) is *nested inside* (v_i, v_j) provided v_iHv_j contains v_kHv_ℓ . The following lemma is useful in identifying the location of the bottom vertices along the hamiltonian cycle of an upset tournament.

Lemma 2.4 *Let T be an upset tournament with hamiltonian cycle H . Suppose (v_i, v_j) and (v_k, v_ℓ) are arcs of T not in H such that (v_k, v_ℓ) is nested inside (v_i, v_j) . Then neither of the bottom vertices of T is on v_iHv_k .*

Proof. Neither v_i nor v_k is a bottom vertex, since each is the initial vertex of an arc in H and an arc not in H . Similarly, neither v_j nor v_ℓ is a top vertex.

Suppose to the contrary that a bottom vertex w is on v_iHv_k . Then T 's top vertices must occur on wHv_ℓ , for otherwise, (v_k, v_ℓ) is an upset arc not in H . Hence, the scores of the vertices along $v_\ellHv_i$ are in decreasing order. In particular, the score of v_j is greater than that of v_i , which leads to the contradiction that (v_i, v_j) is an upset arc of T not in H . ■

We now note that each of the six arcs in the upset tournament $T_{(1)}$ is a forced arc or is on the hamiltonian cycle. Hence, $T_{(1)}$ has a score certificate consisting of the forced arcs and the arcs in its hamiltonian cycle. For upset tournaments on five or more vertices we have the following result.

Theorem 2.5 *Let T be an upset tournament on $n \geq 5$ vertices. There exists a score certificate for T consisting of the arcs in its unique hamiltonian cycle, its forced arcs, and one other arc.*

Proof. Let H be the hamiltonian cycle of T . Without loss of generality we may assume $T = T_\lambda$ where $\lambda = (p_1, p_2, \dots, p_k)$. Let β be the set consisting of the arcs in H and the forced arcs of T .

By Lemma 2.1, it suffices to show that there exists an arc e such that each upset tournament, T' , on the same vertices as T which contains e and the arcs in β has the same bottom vertices as T .

The choice of the arc e depends upon λ . We consider four cases.

CASE 1: Each p_i equals 1.

Since $n \geq 5$, the arc (v_{n-1}, v_1) is in T but not in β . Take e to be this arc. Note that e is nested inside the arc (v_3, v_1) of T , and that (v_3, v_1) is not in H . Applying Lemma 2.4, we conclude that none of the vertices $v_3, v_4, \dots, v_{n-1}$ is a bottom vertex of T' . Since v_n is the initial vertex of two arcs in β , v_n is not a bottom vertex of T' . Hence, T' has the same bottom vertices as T .

CASE 2: For some $j \geq 2$, $p_1 = p_2 = \dots = p_{j-1} = 1$ and $p_j \geq 2$.

Then (v_{j+2}, v_{j+1}) is a forced arc of T , $(v_{j+2}, v_1) \in H$ and there exists a unique vertex v_q such that $(v_q, v_{j+2}) \in H$. Clearly, (v_q, v_1) is an arc of T not in H . Take e to be this arc. Note that e is nested inside the arc (v_3, v_1) , and that (v_3, v_1) is not in H . Applying Lemma 2.4, we conclude that the bottom vertices of T' are contained in $\{v_1, v_2, v_{j+2}\}$. Since (v_{j+2}, v_1) and (v_{j+2}, v_{j+1}) are in β , v_{j+2} is not a bottom vertex of T' . Hence, T' has the same bottom vertices as T .

CASE 3: $k = 1$.

Then $p_1 = n - 3$, and since $n \geq 5$ the arc (v_n, v_2) is in T and not in β . Take e to be this arc. Note that (v_3, v_2) is an arc of β not in H , and is nested inside e . By Lemma 2.4, we conclude that the bottom vertices of T' are contained in $\{v_1, v_2, v_{n-1}\}$. Since (v_{n-1}, v_n) and (v_{n-1}, v_{n-2}) are arcs in β , v_{n-1} is not a bottom vertex of T' . Hence, T' has the same bottom vertices as T .

CASE 4: $k \geq 2$ and $p_1 \geq 2$.

Then (v_3, v_2) is a forced arc of T which is not in H . The arc $(v_{2+p_1+p_2}, v_2)$ is an arc of T not in β . Take e to be this arc. Then (v_3, v_2) is nested inside e . By Lemma 2.4, we conclude that the bottom vertices of T' are contained in $\{v_1, v_2, v_{2+p_1}\}$. Since $(v_{2+p_1}, v_{2+p_1+p_2})$ and (v_{2+p_1}, v_{1+p_1}) are in β , v_{2+p_1} is not a bottom vertex of T' . Hence, T' has the same bottom vertices as T .

The desired conclusion is reached in each case, and the proof is complete. ■

Examples of score certificates, constructed as in the proof of Theorem 2.5, for three upset tournaments on 11 vertices are given in Figure 2. Using Theorem 2.5 we can derive an upper bound on the score certificate number of T_λ .




Figure 2

Corollary 2.6 *Let $\lambda = (p_1, p_2, \dots, p_k)$ be an ordered partition of $n-3$, where $n \geq 5$. Then*

$$sc(T_\lambda) \leq n + 2x + y + 1$$

where x equals $|\{i : 1 \leq i \leq k \text{ and } p_i \geq 2\}|$, and y equals $|\{i : i \in \{1, k\} \text{ and } p_i = 1\}|$.

Proof. By Theorem 2.5, there is a score certificate for T_λ consisting of the arcs in its hamiltonian cycle, its forced arcs, and one additional arc. The hamiltonian cycle has n arcs, the number of forced arcs not in the hamiltonian

cycle and not incident to v_1 or v_n is $2x$, and the number of forced arcs incident to v_1 or v_n and not in the hamiltonian cycle is y . The result now follows. ■

If $\lambda = (2, 2, \dots, 2)$, then by Corollary 2.6, $sc(T_\lambda) \leq 2n - 3 + 1$, and this bound is worse than that given in Theorem 2.2. However, it can be verified that the score certificate for $T_{(2,2,\dots,2)}$ constructed in the proof of Corollary 2.6 is not minimal (with respect to set inclusion). For example, one can delete the arcs (v_6, v_1) and (v_{10}, v_{11}) from the score certificate for $T_{(2,2,2,2)}$ shown in Figure 2 and still have a score certificate.

For arbitrary λ , it is easy to see that the bound for $sc(T_\lambda)$ in Corollary 2.6 is usually better than that in Theorem 2.2. Indeed, Corollary 2.6 implies that the score certificate number of the nearly transitive tournament on n vertices is at most $n + 3$.

3 Lower bounds

In this section we derive lower bounds for the score certificate number of certain upset tournaments. In particular, we show that the score certificate number of the nearly transitive tournament on n vertices equals $n + 3$, for $n \geq 10$.

The class of upset tournaments that we study in this section is the class of T_λ where each part p_i of $\lambda = (p_1, \dots, p_k)$ is at least $k + 6$. For such tournaments, Corollary 2.6 guarantees a score certificate with $n + 2k + 1$ arcs. We begin by showing that for such a tournament, each score certificate with less than $n + 2k + 1$ arcs contains each arc in its upset path, other than (v_1, v_2) and (v_{n-1}, v_n) . For convenience, we define the *truncated upset path* of T_λ to be the path obtained from its upset path by removing the first and last arc.

Lemma 3.1 *Let $\lambda = (p_1, \dots, p_k)$ be an ordered partition of $n - 3$ such that each $p_i \geq k + 6$. Then each score certificate for T_λ of cardinality less than $n + 2k + 1$ contains each arc in T_λ 's truncated upset path.*

Proof. Suppose to the contrary that there exists a score certificate, D , for T_λ with less than $n + 2k + 1$ arcs that does not contain each arc in the truncated upset path. Note that if γ is a cycle of T_λ , then the tournament

obtained from T_λ by reversing the arcs of γ has the same score-list as T_λ . Since D is a score certificate, it follows that D contains at least one arc in each cycle of T_λ .

For $m = 1, 2, \dots, k$, let x_m be the number of non-forced arcs in D having both initial and terminal vertices in

$$\{v_r, v_{r+1}, \dots, v_t\},$$

where $r = 2 + p_1 + p_2 + \dots + p_{m-1}$ and $t = 2 + p_1 + p_2 + \dots + p_m$. First suppose that the upset arc (v_r, v_t) is not in D . For each s with $r + 1 < s < t - 1$,

$$v_t \rightarrow v_s \rightarrow v_r \rightarrow v_t$$

is a 3-cycle in T_λ , and hence D contains at least one of the non-forced arcs (v_t, v_s) and (v_s, v_r) . Therefore, $x_m \geq p_m - 3 \geq k + 3$. If (v_r, v_t) is in D , then certainly $x_m \geq 1$.

It follows that the number of non-forced arcs in D is at least

$$\sum_{m=1}^k x_m \geq (k - 1) + k + 3 = 2k + 2.$$

Since D contains the $n - 1$ forced arcs, we are led to the contradiction that D has at least $n + 2k + 1$ arcs. ■

Let $\lambda = (p_1, \dots, p_k)$ be an ordered partition of $n - 3$ with each $p_i \geq k + 6$. An immediate consequence of Lemma 3.1 is that

$$sc(T_\lambda) \geq n + k - 1. \tag{4}$$

In the remainder of this paper we show that this bound can be improved to

$$sc(T_\lambda) \geq n + k + 2.$$

The technique employed is as follows. Let δ be the set consisting of the arcs in the truncated upset path of T_λ and the forced arcs of T_λ . We exhibit a family $T_1, \dots, T_5$ of upset tournaments on the vertices $v_1, \dots, v_n$ with the following properties:

- (a) $T_i \neq T_\lambda$ for $i = 1, 2, 3, 4, 5$, and

- (b) for each pair e and f of arcs which are in T_λ and not in δ , there exists a j such that both e and f are arcs of T_j .

We claim that this implies that $sc(T_\lambda) \geq n + k + 2$. To see this, let D be a score certificate for T_λ . If D does not contain the truncated upset path of T_λ , then Lemma 3.1 implies that D has at least $n + 2k + 1$ arcs. Otherwise, D contains each arc in δ . The existence of such $T_1, \dots, T_5$ implies that no set of arcs consisting of δ and at most two other arcs is a score certificate for T_λ . Thus, D has at least 3 arcs not in δ . Therefore, D has at least $n + k + 2$ arcs.

In order to easily describe the tournaments $T_1, \dots, T_5$, we introduce some notation. Throughout the remainder of the paper we identify the vertex v_i with the index i . Let T be an upset tournament with vertices $1, 2, \dots, n$ not necessarily in standard form. Let π be the permutation so that when T is put in standard form, the vertices from bottom to top are $\pi(1), \pi(2), \dots, \pi(n)$. If we start with the n -tuple $(\pi(1), \pi(2), \dots, \pi(n))$ and boldface each entry in the n -tuple corresponding to a vertex on T 's upset path, we obtain a n -tuple which completely describes T . For example, the nearly transitive tournament on n vertices in standard form is denoted by the n -tuple

$$(\mathbf{1}, \mathbf{2}, 3, 4, \dots, n-3, n-2, \mathbf{n-1}, \mathbf{n}),$$

and the 7-tuple $(\mathbf{3}, \mathbf{4}, 1, \mathbf{7}, \mathbf{2}, \mathbf{5}, \mathbf{6})$ determines the upset tournament illustrated in Figure 3.



Figure 3

If T is the tournament described by a given n -tuple, then (i, j) is an arc of T if and only if

- (1) both i and j are boldfaced and j is the first boldfaced entry to the right of i ,
- (2) both i and j are boldfaced, j is to the left of i and there is at least one boldfaced entry between j and i , or
- (3) i or j is not boldfaced and j occurs to the left of i .

Theorem 3.2 *Let $\lambda = (p_1, \dots, p_k)$ be an ordered partition of $n - 3$ such that each $p_i \geq k + 6$. Then $sc(T_\lambda) \geq n + k + 2$.*

Proof. It suffices to exhibit tournaments $T_1, \dots, T_5$ satisfying (a) and (b) above.

Without loss of generality we may assume that T_λ has upset path 1, 2, $i_1, i_2, \dots, i_{k-1}, n - 1, n$ and hence is determined by

$$\begin{aligned}
 &(\mathbf{1}, \mathbf{2}, 3, \dots, i_1 - 1, \\
 &\mathbf{i}_1, i_1 + 1, \dots, i_2 - 1, \\
 &\mathbf{i}_2, i_2 + 1, \dots, i_3 - 1, \\
 &\quad \vdots \\
 &\mathbf{i}_{k-1}, i_{k-1} + 1, \dots, n - 2, \\
 &\mathbf{n} - \mathbf{1}, \mathbf{n}).
 \end{aligned}$$

Let T_1 be the upset tournament determined by

$$\begin{aligned}
 &(\mathbf{1}, \mathbf{4}, \mathbf{3}, \mathbf{2}, 5, \dots, i_1 - 1, \\
 &\mathbf{i}_1, i_1 + 1, \dots, i_2 - 1, \\
 &\mathbf{i}_2, i_2 + 1, \dots, i_3 - 1, \\
 &\quad \vdots \\
 &\mathbf{i}_{k-1}, i_{k-1} + 1, \dots, n - 2, \\
 &\mathbf{n} - \mathbf{1}, \mathbf{n}).
 \end{aligned}$$

It is easy to verify that T_1 contains T_λ 's truncated upset path and forced arcs. Also, the set of arcs in T_λ that are not in T_1 is

$$X = \{(4, 2), (4, 1)(1, 2)\}.$$

Analogously, we define T_2 to be the upset tournament determined by

$$\begin{aligned} & (1, 2, \\ & \quad 3, \dots, i_1 - 1, \mathbf{i}_1, \\ & \quad i_1 + 1, \dots, i_2 - 1, \mathbf{i}_2 \\ & \quad \vdots \\ & \quad i_{k-2} + 1, \dots, i_{k-1} - 1, \mathbf{i}_{k-1}, \\ & i_{k-1} + 1, \dots, n - 4, \mathbf{n} - 1, \mathbf{n} - 2, \mathbf{n} - 3, \mathbf{n}). \end{aligned}$$

and observe that the set of arcs in T_λ that are not in T_2 is

$$Y = \{(n, n - 3), (n - 1, n), (n - 1, n - 3)\}.$$

Note that X and Y are disjoint. Let e and f be arcs of T_λ that are not forced arcs and not in its truncated upset path. If it is not the case that X contains one of e or f and Y contains the other, then the arcs e and f are simultaneously in at least one of T_1 and T_2 . Thus, we assume without loss of generality that $e \in X$ and $f \in Y$.

Let T_3 be the upset tournament determined by

$$\begin{aligned} & (\mathbf{i}_1 - 1, \mathbf{i}_1 - 2, 1, \mathbf{i}_1 - 3, \mathbf{i}_1 - 4, \dots, 3, 2, \\ & \quad \mathbf{i}_1, i_1 + 1, i_1 + 2, \dots, i_2 - 1, \\ & \quad \mathbf{i}_2, i_2 + 1, \dots, i_3 - 1 \\ & \quad \vdots \\ & \quad \mathbf{i}_{k-1}, i_{k-1} + 1, \dots, n - 2, \\ & \quad \mathbf{n} - 1, \mathbf{n}), \end{aligned}$$

where we identify i_1 and $n - 1$ in the case $k = 1$. It can be verified that T_3 contains T_λ 's truncated upset path and forced arcs. Also T_3 contains the arcs in $\{(4, 1)\} \cup Y$, and hence if $e = (4, 1)$ then T_3 contains both e and f .

Analogously, we define T_4 to be the upset tournament determined by

$$\begin{aligned} & (1, 2 \\ & \quad 3, 4, \dots, i_1 - 1, \mathbf{i}_1 \\ & \quad i_1 + 1, i_1 + 2, \dots, i_2 - 1, \mathbf{i}_2 \\ & \quad \vdots \\ & \quad i_{k-2} + 1, i_{k-2} + 2, \dots, i_{k-1} - 1, \mathbf{i}_{k-1} \\ & \mathbf{n} - 1, \mathbf{n} - 2, \dots, \mathbf{i}_{k-1} + 3, n, \mathbf{i}_{k-1} + 2, \mathbf{i}_{k-1} + 1) \end{aligned},$$

where we identify vertices 2 and i_{k-1} in the case that $k = 1$, and conclude that if $f = (n, n - 3)$ then T_4 contains both e and f .

Finally, let T_5 be the upset tournament determined by

$$(\mathbf{n}, \mathbf{n} - 2, \mathbf{n} - 3, n - 1, \mathbf{n} - 4, \mathbf{n} - 5, \dots, \mathbf{5}, 2, \mathbf{4}, \mathbf{3}, \mathbf{1}),$$

where each of the entries between $n - 5$ and 5 is boldfaced. It can be verified that T_1 contains T_λ 's truncated upset path and forced arcs. Also, T_5 contains the arcs in $\{(1, 2), (4, 2), (n - 1, n), (n, n - 2)\}$. Hence, if $e \neq (4, 1)$ and $f \neq (n, n - 3)$, then T_5 contains both e and f .

Therefore, the upset tournaments $T_1, \dots, T_5$ satisfy property (b), and it is clear that they also satisfy property (a). ■

In Figure 4, we illustrate the tournaments $T_1, \dots, T_5$ which are constructed in the proof of Theorem 3.2 for the upset tournament $T_{(8,8)}$.

The following is an immediate consequence of Corollary 2.6 and Theorem 3.2.

Corollary 3.3 *Let $n \geq 10$, and let N_n be the nearly transitive tournament on n vertices. Then*

$$sc(N_n) = n + 3.$$

The following chart summarizes the results for tournaments on n vertices corresponding to the partition $\lambda = (p_1, \dots, p_k)$, where each $p_i \geq k + 6$, and is a consequence of Corollary 2.6 and Theorem 3.2.

number of parts of λ	minimum number of vertices	minimum size of each part	lower bound for $sc(T_\lambda)$	upper bound for $sc(T_\lambda)$
1	10	7	$n + 3$	$n + 3$
2	19	8	$n + 4$	$n + 5$
3	30	9	$n + 5$	$n + 7$
4	43	10	$n + 6$	$n + 9$
k	$k^2 + 6k + 3$	$k + 6$	$n + k + 2$	$n + 2k + 1$




Figure 4

References

- [AR] N. Alon and M. Ruszinkó, Short certificates for tournaments, *The Electronic Journal of Combinatorics*, (1997), 4:# R12.

- [BL] R. Brualdi and Q. Li, Upsets in round robin tournaments, *J. Combinatorial Theory B*, (1983), **35**:62-77.
- [CL] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Third edition, Chapman and Hall, London, 1996.
- [FKT] P. Fishburn, J. H. Kim, and P. Tetali, Tournament Certificates, *Technical Memorandum, AT&T Bell Laboratories*, (1994), DIMACS Technical Report No. 94-05.
- [KTF] J. H. Kim, P. Tetali, and P. Fishburn, Score certificates for tournaments, *J. Graph Theory*, (1997), **24**:117-138.
- [M] J. W. Moon, *Topics on Tournaments*, Holt, Rinehart, and Winston, Inc., New York, 1968.
- [RB] K. B. Reid and L. W. Beineke, *Selected Topics in Graph Theory* (L. W. Beineke and R. J. Wilson, Eds.), Chap. 7, 385-415, Academic Press, New York, 1978.
- [R] A. Rubinstein, Why are certain properties of binary relations relatively more common in natural language?, *Econometrica*, (1996), **64**:343-355.