

Direct bijective computation of the generating series for 2 and 3-connection coefficients of the symmetric group

Alejandro H. Morales
LaCIM
Université du Québec à Montréal
Montréal PQ H3C 3P8, Canada
ahmorales@lacim.ca

Ekaterina A. Vassilieva
LIX
Ecole Polytechnique
91128 Palaiseau Cedex, France
katya@lix.polytechnique.fr

Submitted: Jan 17, 2011; Accepted: March 20, 2013; Published: XX
Mathematics Subject Classifications: 05A15, 05A19

Abstract

We evaluate combinatorially certain connection coefficients of the symmetric group that count the number of factorizations of a long cycle as a product of three permutations. Such factorizations admit an important topological interpretation in terms of unicellular constellations on orientable surfaces. Algebraic computation of these coefficients was first done by Jackson using irreducible characters of the symmetric group. However, bijective computations of these coefficients are so far limited to very special cases. Thanks to a new bijection that refines the work of Schaeffer and Vassilieva in [17] and Vassilieva in [18], we give an explicit closed form evaluation of the generating series for these coefficients. The main ingredient in the bijection is a modified oriented tricolored tree tractable to enumerate. Finally, reducing this bijection to factorizations of a long cycle into two permutations, we get the analogue formula for the corresponding generating series.

Keywords: Connection Coefficients; Factorizations; Cacti; Symmetric Group

1 Introduction

1.1 Generating series for connection coefficients

In what follows, we denote by $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \vdash n$ an integer partition of n and $\ell(\lambda) = k$ the length or number of parts of λ . We also write $\lambda = [1^{n_1(\lambda)}, 2^{n_2(\lambda)}, \dots]$ where $n_i(\lambda)$ is the number of parts i in λ .

Let \mathfrak{S}_n be the symmetric group on n elements, and \mathcal{C}_λ be the conjugacy class in \mathfrak{S}_n of permutations with cycle type λ , where $\lambda \vdash n$. Given $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(r)}, \mu \vdash n$, let $k_{\lambda^{(1)}, \dots, \lambda^{(r)}}^\mu$ be the number of ordered factorizations in \mathfrak{S}_n of a fixed permutation $\gamma \in \mathcal{C}_\mu$ as a product $\alpha_1 \cdots \alpha_r$ of r permutations $\alpha_i \in \mathcal{C}_{\lambda^{(i)}}$. These numbers are called *connection coefficients* of the symmetric group. The problem of computing these coefficients has received significant

attention and a good account of its history and references can be found in [9]. We focus on the cases $k_{\lambda,\mu}^n$ and $k_{\lambda,\mu,\nu}^n$: i.e. when $r = 2$ and 3 , $\mu = (n)$ and γ is the long cycle $\gamma_n = (1, 2, \dots, n)$.

In addition, for $\lambda \vdash n$ we use the monomial symmetric function $m_\lambda(\mathbf{x})$ on indeterminates $\mathbf{x} = (x_1, x_2, \dots)$ which is the sum of all different monomials obtained by permuting the variables of $x_1^{\lambda_1} x_2^{\lambda_2} \dots$, and the power symmetric function $p_\lambda(\mathbf{x})$, defined multiplicatively as $p_\lambda = p_{\lambda_1} p_{\lambda_2} \dots$ where $p_n(\mathbf{x}) = m_n(\mathbf{x}) = \sum_i x_i^n$. Also, if $\lambda = [1^{n_1(\lambda)}, 2^{n_2(\lambda)}, \dots]$, let $\text{Aut}(\lambda) = \prod_i n_i(\lambda)!$.

Our combinatorial results can be stated as follows:

Theorem 1.1. *The numbers $k_{\lambda,\mu,\nu}^n$ of factorizations of the long cycle γ_n into an ordered product of three permutations of types λ, μ , and ν respectively satisfy:*

$$\sum_{\lambda,\mu,\nu \vdash n} k_{\lambda,\mu,\nu}^n p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) p_\nu(\mathbf{z}) = \sum_{\lambda,\mu,\nu \vdash n} \frac{n!^2 M_{\ell(\lambda),\ell(\mu),\ell(\nu)}^{(n-1)}}{\binom{n-1}{\ell(\lambda)-1} \binom{n-1}{\ell(\mu)-1} \binom{n-1}{\ell(\nu)-1}} m_\lambda(\mathbf{x}) m_\mu(\mathbf{y}) m_\nu(\mathbf{z}), \quad (1.2)$$

where:

$$M_{\ell(\lambda),\ell(\mu),\ell(\nu)}^{(n-1)} = \binom{n-1}{\ell(\nu)-1} \sum_{g \geq 0} \binom{n-\ell(\mu)}{\ell(\lambda)-1-g} \binom{n-\ell(\nu)}{g} \binom{n-1-g}{n-\ell(\mu)}.$$

Corollary 1.3. [15] *The numbers $k_{\lambda,\mu}^n$ of factorizations of the long cycle γ_n into an ordered product of two permutations of cycle types λ and μ respectively satisfy:*

$$\sum_{\lambda,\mu \vdash n} k_{\lambda,\mu}^n p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) = \sum_{\lambda,\mu \vdash n} \frac{n(n-\ell(\lambda))!(n-\ell(\mu))!}{(n+1-\ell(\lambda)-\ell(\mu))!} m_\lambda(\mathbf{x}) m_\mu(\mathbf{y}), \quad (1.4)$$

We will see in Section 2 that the coefficients on the right hand sides of (1.2) and (1.4) are non-negative integers.

Remark 1.5. *Equations (1.2) and (1.4) can be obtained algebraically using the irreducible characters of the symmetric group, the Murnaghan-Nakayama rule, and symmetric function identities (see [12]). Here, we derive these equations through a bijection.*

1.2 Background

In the setting of the connection coefficients $k_{\lambda^{(1)}, \dots, \lambda^{(r)}}^n$, we define the *genus* $\mathbf{g}(\lambda^{(1)}, \dots, \lambda^{(r)})$ of the partitions $\lambda^{(i)}$ by the equation

$$\ell(\lambda^{(1)}) + \dots + \ell(\lambda^{(r)}) = (r-1)n + 1 - 2\mathbf{g}(\lambda^{(1)}, \dots, \lambda^{(r)}). \quad (1.6)$$

We can take \mathbf{g} to be a non-negative integer, since otherwise it is easy to show that $k_{\lambda^{(1)}, \dots, \lambda^{(r)}}^n = 0$.

Except for special cases there are no closed formulas for the connection coefficients $k_{\lambda^{(1)}, \dots, \lambda^{(r)}}^n$. For instance, using an inductive combinatorial argument Bédard and Goupil

[1] found a formula for $k_{\lambda,\mu}^n$ in the case $\mathbf{g}(\lambda,\mu) = 0$. This was extended by Goulden and Jackson [6] to evaluate $k_{\lambda^{(1)},\dots,\lambda^{(r)}}^n$ in the case $\mathbf{g}(\lambda^{(1)},\dots,\lambda^{(r)}) = 0$ via a bijection with a set of ordered rooted r -cacti on n r -gons. Later, using characters of the symmetric group and a combinatorial development, Goupil and Schaeffer [9] derived an expression for connection coefficients of arbitrary genus as a sum of positive terms (see Biane [3] for a succinct algebraic derivation; Poulalhon and Schaeffer [16] and Irving [11] for further generalizations). As a general rule, these developments are quite intricate and the formulas obtained are rather complicated.

Interestingly, if we consider the generating series for the coefficients $k_{\lambda^{(1)},\dots,\lambda^{(r)}}^n$ as in the LHS of (1.2), the coefficients of their expansion in the basis of monomial symmetric functions, as in the RHS of (1.2), can be computed in closed form thanks to a result by Jackson [12] obtained algebraically using the theory of the irreducible characters of the symmetric group. There are direct bijections for a variant of the case of two factors (i.e. $r = 2$) like the celebrated Harer-Zagier formula [10]: see Lass [14], Goulden and Nica [8], and Bernardi [2]. In this paper we follow this approach and introduce the notion of *partitioned tricolored (bicolored) 3-cacti (maps)* of given type, refining the work of Schaeffer and Vassilieva in [17] and Vassilieva in [18], and use a purely combinatorial argument to derive the explicit generating series for $k_{\lambda,\mu,\nu}^n$ and $k_{\lambda,\mu}^n$ in Equations (1.2) and (1.4) respectively.

1.3 Outline of paper

The paper is organized as follows: in Section 2 we introduce the *partitioned 3-cacti* and the *cactus trees* (the enumeration of the latter is postponed to Section 4) and relate them via a bijection Θ described in Section 3. Finally, in Section 5 we prove Corollary 1.3.

2 Combinatorial reformulation

2.1 Cacti and partitioned cacti

Factorizations in the symmetric group counted by $k_{\lambda,\mu,\nu}^n$ admit a direct interpretation in terms of *unicellular 3-constellations* also named *3-cacti* with white, black, and grey vertices of respective degree distribution λ , μ , and ν . Within a topological point of view, 3-cacti are specific *maps* which in turn are 2-cell decompositions of an oriented surface into a finite number of vertices (0-cells), edges (1-cells) and faces (2-cells) homeomorphic to open discs (see [13] for more details about maps and their applications). Maps are defined up to a homeomorphism of the surface that preserves its orientation, the type of cells, and incidences in the graph. 3-cacti are maps with black faces and one white face (thus the term *unicellular*) such that: (i) each edge separates a black face and the white face and (ii) all the black faces are triangles each composed of exactly a white, a black, and a grey vertex following each other in clockwise order. As a consequence, the degree of the white face is a multiple of 3. Often, cacti refer to planar maps (embedded in an orientable surface of genus 0). In this paper we assume that they can be embedded in an orientable surface of any genus. Besides, we consider only *rooted* cacti, i.e. cacti with a marked black face. We assume as well that each black triangle is labeled with an index in $\{1, 2, \dots, n\}$ with the convention that the

marked triangle is labeled 1. In what follows, we define the *degree* of a vertex in a cactus as the number of triangles it belongs to, and the degree distribution of the vertices of a given color is the integer partition of n formed by the degrees of all the vertices of this color.

The next classical result (see [13]) relates rooted 3-cacti with factorizations of the long cycle $\gamma_n = (1, 2, \dots, n)$.

Proposition 2.1. *Rooted 3-cacti with n black triangles are in bijection with 3-tuples $(\alpha_1, \alpha_2, \alpha_3)$ of permutations in \mathfrak{S}_n such that $\alpha_1\alpha_2\alpha_3 = \gamma_n$. Under this bijection the white (black and grey, resp.) vertices correspond to cycles of π_1 (π_2 and π_3 , resp.).*

A sketch of the proof of this classical result can be found in [18]. Each white, black, or grey vertex of a given 3-cacti corresponds to a cycle of permutation α_1 , α_2 , or α_3 respectively, and the cycle is encoded by the local counter-clockwise order of the triangles around the vertex. The fact that $\alpha_1\alpha_2\alpha_3 = \gamma_n$ corresponds to saying that traversing the map starting on the white vertex of the triangle labeled 1 and keeping the white face on the *right* we visit, in order, the white-black edges belonging to triangles labeled $1, 2, \dots, n$. A consequence of Proposition 2.1 is that for integer partitions λ, μ, ν of n , the number of 3-cacti of degree distribution λ, μ, ν is the number $k_{\lambda, \mu, \nu}^n$ of factorizations defined in Section 1.1. Moreover, by the Euler-characteristic, the genus of the underlying surface of the 3-cacti of degree distribution λ, μ, ν is given by Equation 1.6. Figure 1 shows a 3-cactus embedded on the sphere (genus 0) and a 3-cactus embedded on a torus (genus 1).




Figure 1: Two examples of rooted 3-cacti embedded in a surface of genus 0 (left) and genus 1 (right)

Example 2.2. *The cactus on the left hand side in Figure 1 can be associated to the three permutations:*

$$\alpha_1 = (1)(24)(3)(5) \tag{2.3}$$

$$\alpha_2 = (1)(23)(45) \tag{2.4}$$

$$\alpha_3 = (15)(2)(3)(4) \tag{2.5}$$

The degree distribution of this cactus is $\lambda = [1^3, 2^1]$, $\mu = [1^1, 2^2]$, $\nu = [1^3, 2^1]$.

The cactus on the right hand side corresponds to the three permutations:

$$\alpha_1 = (1236)(4)(5) \quad (2.6)$$

$$\alpha_2 = (153)(2)(4)(6) \quad (2.7)$$

$$\alpha_3 = (134)(2)(5)(6) \quad (2.8)$$

The degree distribution of this cactus is $\lambda = [1^2, 4^1]$, $\mu = [1^3, 3^1]$, $\nu = [1^3, 3^1]$.

Partitioned 3-cacti

Cacti with a given vertex degree distribution are in general non-planar and non-recursive objects, and no direct bijection is known to compute their cardinality. However, we provide an interpretation of the formal power series $\sum_{\lambda, \mu, \nu \vdash n} k_{\lambda, \mu, \nu}^n p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) p_\nu(\mathbf{z})$ as a generating function for *partitioned cacti*. We are able to give an explicit formula for this generating function by introducing a new bijective mapping for partitioned cacti. Intuitively, partitioned cacti are rooted 3-cacti where the set of vertices of each color are partitioned into blocks. Such objects have also been widely studied, for instance by Lass [14] and Bernardi [2] under the term of colored maps; and by Goulden and Nica [8], Schaeffer and Vassilieva in [17], and Vassilieva in [18] as partitioned maps or cacti.

To define partitioned cacti we use π to denote a set partition of a set of n elements with blocks $\{\pi^1, \dots, \pi^p\}$. The type of a set partition, $\text{type}(\pi) \vdash n$, is the integer partition of n obtained by considering the cardinalities of the blocks of π . We are now ready for the definition:

Definition 2.9. (*Partitioned 3-cactus*) A partitioned 3-cactus is a 4-tuple $(\pi_1, \pi_2, \pi_3, \kappa)$ such that κ is a rooted 3-cactus with n triangles, and π_1, π_2 and π_3 are set partitions on the set of white, black, and grey vertices respectively. By abuse of notation, hereinafter we view π_1, π_2 , and π_3 as set partitions on $\{1, 2, \dots, n\}$ where a block is composed of the labels of the triangles incident to the vertices contained in the block. In what follows, the blocks of π_1, π_2 , and π_3 are denoted $\pi_1^{(i)}, \pi_2^{(j)}$, and $\pi_3^{(k)}$ with the only restriction that $1 \in \pi_1^{(\ell(\lambda))}$.

For $\lambda, \mu, \nu \vdash n$, we let $\mathcal{C}(\lambda, \mu, \nu)$ be the set of partitioned cacti $(\pi_1, \pi_2, \pi_3, \kappa)$ where the set partitions π_1, π_2 , and π_3 of $\{1, 2, \dots, n\}$ have type λ, μ , and ν respectively. Let $C(\lambda, \mu, \nu) = |\mathcal{C}(\lambda, \mu, \nu)|$.

Remark 2.10. Following Proposition 2.1, we can interpret the cactus κ as a 3-tuple of permutations $(\alpha_1, \alpha_2, \alpha_3)$ where $\alpha_3 = \alpha_2^{-1} \circ \alpha_1^{-1} \circ \gamma_n$. As a result, partitioned cacti in $\mathcal{C}(\lambda, \mu, \nu)$ can be seen as 5-tuples $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ where π_1, π_2 , and π_3 are set partitions of $\{1, 2, \dots, n\}$ of types λ, μ , and ν respectively with the property that: for $k = 1, 2, 3$, if an integer l of a given cycle c of α_k belongs to a given block of π_k then all the integers in the cycle c also belong to this block.

Example 2.11. Take the cactus on the right hand side of Figure 1 and add partitions $\pi_1 = \{\pi_1^{(1)}, \pi_1^{(2)}\}, \pi_2 = \{\pi_2^{(1)}, \pi_2^{(2)}\}, \pi_3 = \{\pi_3^{(1)}, \pi_3^{(2)}, \pi_3^{(3)}\}$, where $\pi_1^{(1)} = \{4, 5\}, \pi_1^{(2)} = \{1, 2, 3, 6\}; \pi_2^{(1)} = \{1, 3, 4, 5\}, \pi_2^{(2)} = \{2, 6\}; \pi_3^{(1)} = \{1, 3, 4, 6\}, \pi_3^{(2)} = \{2\}, \pi_3^{(3)} = \{5\}$. This gives a partitioned cactus $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}([2^1, 4^1], [2^1, 4^1], [1^2, 4^1])$ depicted in Figure 2. Similarly to [17], we associate a particular shape to each of the blocks of the partitions.




Figure 2: Illustration of the Partitioned 3-Cactus from Example 2.11.

Link between cacti and partitioned cacti

Consider the partial order on integer partitions given by refinement. That is $\lambda \preceq \mu$ if and only if the parts of μ are unions of parts of λ , and we say that μ is *coarser* than λ . If $\lambda \preceq \mu$ let $\bar{R}_{\lambda,\mu}$ be the number of ways to *coarse* λ to obtain μ . For example, if $\lambda = [1^2, 2^2]$ and $\mu = [1, 2, 3]$ then $\bar{R}_{\lambda,\mu} = 4$. It is well known that $p_\lambda = \sum_{\mu \succeq \lambda} \text{Aut}(\mu) \bar{R}_{\lambda,\mu} m_\mu$ [19, Prop. 7.7.1].

We use this partial order on integer partitions to obtain an immediate relation between $C(\lambda, \mu, \nu)$ and $k_{\lambda,\mu,\nu}^n$.

Proposition 2.12. *For partitions $\rho, \delta, \epsilon \vdash n$ we have :*

$$C(\rho, \delta, \epsilon) = \sum_{\lambda \preceq \rho, \mu \preceq \delta, \nu \preceq \epsilon} \bar{R}_{\lambda\rho} \bar{R}_{\mu\delta} \bar{R}_{\nu\epsilon} k_{\lambda,\mu,\nu}^n. \quad (2.13)$$

Proof. Let $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\rho, \delta, \epsilon)$. If $\alpha_1 \in \mathcal{C}_\lambda$, $\alpha_2 \in \mathcal{C}_\mu$, and $\alpha_3 = \alpha_2^{-1} \alpha_1^{-1} \gamma_n \in \mathcal{C}_\nu$ then by the definition of the partitioned cacti, we have that $\text{type}(\pi_1) = \rho \succeq \lambda$, $\text{type}(\pi_2) = \delta \succeq \mu$, and $\text{type}(\pi_3) = \epsilon \succeq \nu$. Thus, if we further refine $C(\rho, \delta, \epsilon)$ by the cycle types of the permutations, i.e. if

$$\mathcal{C}_{\lambda,\mu,\nu}(\rho, \delta, \epsilon) = \{(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\rho, \delta, \epsilon) \mid (\alpha_1, \alpha_2, \alpha_2^{-1} \alpha_1^{-1} \gamma_n) \in \mathcal{C}_\lambda \times \mathcal{C}_\mu \times \mathcal{C}_\nu\},$$

then $\mathcal{C}(\rho, \delta, \epsilon) = \bigcup_{\lambda \preceq \rho, \mu \preceq \delta, \nu \preceq \epsilon} \mathcal{C}_{\lambda,\mu,\nu}(\rho, \delta, \epsilon)$ where the union is disjoint. Finally, if

$$C_{\lambda,\mu,\nu}(\rho, \delta, \epsilon) = |\mathcal{C}_{\lambda,\mu,\nu}(\rho, \delta, \epsilon)|$$

then it is easy to see that $C_{\lambda,\mu,\nu}(\rho, \delta, \epsilon) = \bar{R}_{\lambda\rho} \bar{R}_{\mu\delta} \bar{R}_{\nu\epsilon} k_{\lambda,\mu,\nu}^n$. □

Using $p_\lambda = \sum_{\mu \succeq \lambda} \text{Aut}(\mu) \bar{R}_{\lambda,\mu} m_\mu$ Proposition 2.12 is equivalent to:

$$\sum_{\lambda, \mu, \nu \vdash n} k_{\lambda,\mu,\nu}^n p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) p_\nu(\mathbf{z}) = \sum_{\lambda, \mu, \nu \vdash n} \text{Aut}(\lambda) \text{Aut}(\mu) \text{Aut}(\nu) C(\lambda, \mu, \nu) m_\lambda(\mathbf{x}) m_\mu(\mathbf{y}) m_\nu(\mathbf{z}) \quad (2.14)$$

In the special case when we have partitions ρ, δ and ϵ of n where $\ell(\rho) + \ell(\delta) + \ell(\epsilon) = 2n + 1$, the following proposition holds:

Proposition 2.15 ([6]). *For partitions ρ, δ and ϵ of n where $\ell(\rho) + \ell(\delta) + \ell(\epsilon) = 2n + 1$ we have that $C(\rho, \delta, \epsilon) = k_{\rho, \delta, \epsilon}^n = n^2(\ell(\rho) - 1)!(\ell(\delta) - 1)!(\ell(\epsilon) - 1)!/Aut(\rho)Aut(\delta)Aut(\epsilon)$.*

Proof. Let $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\rho, \delta, \epsilon)$ with $\ell(\rho) + \ell(\delta) + \ell(\epsilon) = 2n + 1$, and $\alpha_3 = \alpha_2^{-1}\alpha_1^{-1}\gamma_n$. If $\alpha_1 \in \mathcal{C}_\lambda$, $\alpha_2 \in \mathcal{C}_\mu$, $\alpha_3 \in \mathcal{C}_\nu$, then $\ell(\lambda) + \ell(\mu) + \ell(\nu) = 2n + 1 - 2g(\lambda, \mu, \nu) \leq 2n + 1$. But $\ell(\lambda) \geq \ell(\rho)$, $\ell(\mu) \geq \ell(\delta)$, and $\ell(\nu) \geq \ell(\epsilon)$ therefore $\rho = \lambda$, $\delta = \mu$, and $\epsilon = \nu$; and π_1 , π_2 , and π_3 are the underlying set partitions in the cycle decompositions of α_1, α_2 , and α_3 respectively. Thus $C(\delta, \rho, \epsilon) = k_{\rho, \delta, \epsilon}^n$. But, as shown in [6, Thm. 2.2], $k_{\rho, \delta, \epsilon}^n = n^2(\ell(\rho) - 1)!(\ell(\delta) - 1)!(\ell(\epsilon) - 1)!/Aut(\rho)Aut(\delta)Aut(\epsilon)$ since the genus $g(\rho, \delta, \epsilon) = 0$. As a result, $C(\rho, \delta, \epsilon) = n^2(\ell(\rho) - 1)!(\ell(\delta) - 1)!(\ell(\epsilon) - 1)!/Aut(\rho)Aut(\delta)Aut(\epsilon)$ \square

As mentioned before, explicit computation of the right hand side of equation (2.14) is made possible thanks to a new bijective description of partitioned cacti of given type which is a refinement of a bijection in [17] and [18]. Partitioned cacti are indeed in one-to-one correspondence with particular sets of cactus trees (and three additional simple combinatorial objects), which are recursive planar objects whose number one can compute with classical methods like Lagrange inversion. Next we define such trees, in Section 4 we compute the number of such trees.

2.2 Cactus trees

Before we state the actual definition of the tree structure used as the main ingredient in the proof of Theorem 1.1, we give preparatory explanations. Ordered trees are non cyclic graphs usually defined recursively as a root vertex v and an ordered sequence (possibly the empty set) of ordered trees, called descending trees, each having its root vertex connected to v by an edge. The root of a descending vertex is called a descending vertex. The root vertices of the descending trees of a given vertex are considered as its children. Although it follows the same kind of recursive definition, the tree structure we introduce has the following differences:

- vertices are of three different colors, say white, black and grey;
- the ordered sequence of children of a given vertex is not composed of vertices connected to it through an edge. A child can be: (i) a *thorn* or half edge, i.e an edge, connecting this given vertex to no other (as a result, no descending tree is attached to this kind of child), (ii) a *full edge*, i.e. an edge connecting the given vertex to a descending one with the restriction that only a black (resp. grey, white) vertex can be connected this way to a white (resp. black, grey) one, (iii) a *triangle* connecting the given vertex to two descending ones with the restriction that only a black and grey (resp. grey and white, white and black) can be connected this way to a white (resp. black, grey) one. Triangles are made of three edges connecting the two descending vertices to the ascending one and the two descending vertices between themselves. The two descending vertices are the roots of two descending trees. The three kinds of children are illustrated on Figure 3.

We are now ready to state the definition of the cactus trees:




Figure 3: Example of three types of possible children, the ordered sequence of children attached to the white vertex is: edge, thorn, triangle, thorn, triangle

Definition 2.16 (Cactus Tree). Let $\widetilde{\mathcal{CT}}(p_1, p_2, p_3, g, w, b)$ be the set of cactus trees $\widetilde{\tau}$ with p_1 white vertices, p_2 black vertices, and p_3 grey vertices, g triangles children of grey vertices, w triangles children of white vertices, and b triangles children of black vertices such that:

- (i) the root of $\widetilde{\tau}$ is a white vertex,
- (ii) the ordered set of children of each white (resp. black, grey) vertex consists of three kinds of objects: thorns; full edges connecting this white (resp. black, grey) vertex to a black (resp. grey, white) one; triangles connecting this white (resp. black, grey) vertex to both a black and a grey (resp. grey and white, white and black) one,
- (iii) the edge connecting a white (resp. black, grey) vertex to the black (resp. grey, white) one in a descending triangle comes before the one connecting it to the grey (resp. white, black) vertex according to the ordering of the children of this white (resp. black, grey) vertex.

Within a cactus tree, the degree of a vertex v is defined by:

$$\deg(v) = c + \varepsilon, \quad (2.17)$$

where c is the number of children (that can be either thorns, edges or triangles) and ε is 1 for a non-root vertex, 0 otherwise. With this definition of degree, we write the set $\widetilde{\mathcal{CT}}(p_1, p_2, p_3, g, w, b)$ as the disjoint union

$$\widetilde{\mathcal{CT}}(p_1, p_2, p_3, g, w, b) = \bigcup_{\ell(\lambda)=p_1, \ell(\mu)=p_2, \ell(\nu)=p_3} \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b).$$

where $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ is the set of cactus trees in $\widetilde{\mathcal{CT}}(p_1, p_2, p_3, g, w, b)$ with degree distribution $\lambda, \mu, \nu \vdash n$ of the white, black, and grey vertices respectively.

In addition, in what follows we will denote by $\mathcal{CT}(p_1, p_2, p_3, g, w, b)$ the set of cactus trees similar to those in $\widetilde{\mathcal{CT}}(p_1, p_2, p_3, g, w, b)$ but without thorns. We define $\mathcal{CT}(\lambda, \mu, \nu, g, w, b)$ similarly. Moreover, we will use the expression *tricolored tree* when only full edges are allowed in the set of children of each vertex. Finally, we may use the integers $(1, 2, \dots, p_1)$ (resp. $(1, 2, \dots, p_2)$, $(1, 2, \dots, p_3)$) to label the white (resp. black, grey) vertices of a given cactus tree or tricolored tree. The resulting object is called labeled cactus tree or labeled tricolored tree respectively.

Example 2.18. The cactus tree in Figure 4 belongs to

$$\widetilde{\mathcal{CT}}([1^2, 2^1, 4^1], [1^1, 3^1, 4^1], [1^1, 2^2, 3^1], 1, 1, 1)$$




Figure 4: Example of a Cactus Tree. The white vertex in the bottom is the root of the tree. The tree has $p_1 = 3$ white vertices, $p_2 = 3$ black vertices and $p_3 = 4$ grey vertices, and three triangles each children of white, black, and grey vertices respectively.

Proposition 2.19. *The number of cactus trees is:*

$$|\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)| = n \cdot \frac{(\ell(\lambda) - 1)! (\ell(\mu) - 1)! (\ell(\nu) - 1)!}{\text{Aut}(\lambda) \text{Aut}(\mu) \text{Aut}(\nu)} \frac{(g(w - \ell(\nu)) + \ell(\mu)\ell(\nu))}{(n + 1 - \ell(\lambda) - \ell(\mu) + g)} \times \\ \times \binom{n - \ell(\lambda)}{w, \ell(\mu) - g - w} \binom{n - \ell(\mu)}{b, \ell(\nu) - w - b} \binom{n - \ell(\nu)}{g, \ell(\lambda) - 1 - g - b}. \quad (2.20)$$

The proof of this proposition is carried out using the Lagrange inversion theorem (see e.g. [7, 1.2.13]) and it is postponed to Section 4.

2.3 Reformulation of the main theorem

Let $\mathcal{OP}_r^{(m)}$ be the set of all ordered r -subsets of $[m]$. By definition $|\mathcal{OP}_r^{(m)}| = (m)_r = m(m - 1) \cdots (m - r + 1)$. We have the following proposition:

Proposition 2.21. *Theorem 1.1 is equivalent to:*

$$C(\lambda, \mu, \nu) = \sum_{g, w, b \geq 0} |\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)| |\mathfrak{S}_{n+1-\ell(\lambda)-\ell(\nu)+b}| |\mathfrak{S}_{n-\ell(\mu)-\ell(\nu)+w}| |\mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}| \quad (2.22)$$

Proof. According to Equation 2.14, Theorem 1.1 is equivalent to the equality

$$C(\lambda, \mu, \nu) = \frac{n!^2}{\text{Aut}(\lambda) \text{Aut}(\mu) \text{Aut}(\nu)} \frac{1}{\binom{n-1}{\ell(\lambda)-1} \binom{n-1}{\ell(\mu)-1} \binom{n-1}{\ell(\nu)-1}} M_{\ell(\lambda), \ell(\mu), \ell(\nu)}^{(n-1)}$$

After basic simplifications on the binomial coefficients, a *summand* in the RHS of Equation (2.22) reduces to:

$$\frac{(n-1)!^2 (\ell(\mu)\ell(\nu) + g(w - \ell(\nu)))}{\binom{n-1}{\ell(\lambda)-1} \binom{n-1}{\ell(\mu)-1} \binom{n-1}{\ell(\nu)-1} \text{Aut}(\lambda) \text{Aut}(\mu) \text{Aut}(\nu)} \binom{n}{w, g, b, \ell(\lambda) - 1 - g - b, \ell(\mu) - g - b, \ell(\nu) - w - b}$$

Then we sum over g , w , and b the terms depending on these parameters. Arranging properly the terms depending on w and b , and simplifying sums on these two parameters thanks to

the Vandermonde's convolution formula, we obtain:

$$\begin{aligned} \sum_{g,w,b} (\ell(\mu)\ell(\nu) + g(w - \ell(\nu))) \binom{n}{w, g, b, \ell(\lambda) - 1 - g - b, \ell(\mu) - g - b, \ell(\nu) - w - b} &= \\ &= n^2 \binom{n-1}{\ell(\nu)-1} \sum_g \binom{n-\ell(\mu)}{\ell(\lambda)-1-g} \binom{n-\ell(\nu)}{g} \binom{n-1-g}{n-\ell(\mu)}. \end{aligned}$$

Which leads directly to the desired result. \square

As a direct consequence of Proposition 2.21, Theorem 1.1 reduces to:

Theorem 2.23. *There is an explicit bijection $\Theta_{\lambda,\mu,\nu}^n$ between partitioned 3-cacti in $\mathcal{C}(\lambda, \mu, \nu)$ and tuples $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$, where*

$$\begin{aligned} \tilde{\tau} &\in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b), \\ \sigma_1 &\in \mathfrak{S}_{n+1-\ell(\lambda)-\ell(\nu)+b}, \\ \sigma_2 &\in \mathfrak{S}_{n-\ell(\mu)-\ell(\nu)+w}, \\ \chi &\in \mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)} \end{aligned}$$

for some $g, w, b \geq 0$.

The next section is devoted to proving this theorem by describing the bijection $\Theta_{\lambda,\mu,\nu}^n$.

3 Description of the bijection

3.1 Additional definitions

Before we get to the description of $\Theta_{\lambda,\mu,\nu}^n$, we need two additional ingredients: a linear order on the white (black and grey, resp.) vertices and their children (as defined in the beginning of Section 2.2) which we call *white reverse level traversal (RLT)* (black and grey reverse level traversal, resp.), and partial permutations.

Definition 3.1 (reverse level traversals (RLT)). *For trees $\tilde{\tau} \in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ (or for $\tau \in \mathcal{CT}(\lambda, \mu, \nu, g, w, b)$), we define the white Reverse Levels Traversal (RLT) as the following linear order in $\tilde{\tau}$ of the white vertices and their children.*

We divide the white vertices of $\tilde{\tau}$ into levels depending on their height from the root (where the height is defined as the number of edges in the shortest path with vertex sequence white-grey-black-white-... to the root). So the first level consists of the root, the second level consists of the white vertices at height 3 from the root, etc. The white RLT is a traversal of all the white vertices and their children (thorns, edges, triangles) from left to right, first at the level of maximum height, then the level of second maximum height, ... up to the root vertex. The children of each white vertex are traversed from left to right before the vertex itself¹

¹In words, the white RLT can be viewed as a *reverse breadth first traversal* of the white vertices and their children.

The black and grey RLT are defined similarly with respect to black vertices and their children, and grey vertices and their children. Figure 5 depicts the three RLT for the cactus tree in Example 2.18.




Figure 5: Examples of (a) White, (b) black, and (c) grey reverse level traversals (RLT) on the cactus tree of Example 2.18.

Definition 3.2 (Partial permutations). *Given two sets X and Y and a non negative integer m , let $\mathcal{PP}(X, Y, m)$ be the set of bijections from any m -subset of X to any m -subset of Y . These bijections are called partial permutations. Then $|\mathcal{PP}(X, Y, m)| = \binom{|X|}{m} \binom{|Y|}{m} m!$.*

3.2 Bijective mapping Θ for 3-cacti that preserves type

We proceed with the description of Θ :

$$\Theta_{\lambda, \mu, \nu}^n : \mathcal{C}(\lambda, \mu, \nu) \xrightarrow{\sim} \widetilde{\mathcal{CT}}(\lambda, \mu, \nu) \times \mathfrak{S}_{n+1-\ell(\lambda)-\ell(\nu)+b} \times \mathfrak{S}_{n-\ell(\mu)-\ell(\nu)+w} \times \mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}.$$

Within the construction, we use

$$(p_1, p_2, p_3) := (\ell(\lambda), \ell(\mu), \ell(\nu)).$$

3.2.1 The cactus tree $\tilde{\tau}$

Let $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\lambda, \mu, \nu)$. We construct a cactus tree $\tilde{\tau} \in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ following the procedure below.

(i) *Cactus tree*: The first step is to construct the cactus tree τ and relabeling permutations in the same way as in [18]. For completeness, we also include here the construction. Let $m_2'^{(j)}$ ($1 \leq j \leq p_2$) be the maximal element of $\alpha_3^{-1} \alpha_2^{-1} (\pi_2^{(j)})$ and $m_i^{(j)}$ for $i = 1, 3$ ($1 \leq j \leq p_i$) be the maximal element of the block $\pi_i^{(j)}$.

We first construct the labeled tricolored tree T with p_1 white, p_2 black, and p_3 grey vertices satisfying: the root of T is the white vertex with label p_1 and the incidence relations and order of children are given in Table 1.







Incidence relations	Order of children
for $1 \leq j \leq p_2$  if $m_2^{(j)} \in \alpha_3^{-1} \alpha_2^{-1} (\pi_1^{(i)})$	 if $\alpha_2 \alpha_3 (m_2^{(j_1)}) < \alpha_2 \alpha_3 (m_2^{(j_2)})$
for $1 \leq k \leq p_3$  if $m_3^{(k)} \in \alpha_3^{-1} (\pi_2^{(j)})$	 if $\alpha_3^{-1} \alpha_2^{-1} \alpha_3 (m_3^{(k_1)}) < \alpha_3^{-1} \alpha_2^{-1} \alpha_3 (m_3^{(k_2)})$
for $1 \leq i \leq p_1 - 1$  if $m_1^{(i)} \in \pi_3^{(k)}$	 if $\alpha_3^{-1} (m_1^{(i_1)}) < \alpha_3^{-1} (m_1^{(i_2)})$

Table 1: Incidence relations and order of children of the labeled tricolored tree T . Each row of the table shows the incidence relations and order of children of the white, black and grey vertices respectively.

Lemma 3.3 ([18]). *The procedure above defines a labeled 3-colored tree T .*

We construct the labeled cactus tree Υ from T by forming triangles children of the different vertices following the rules in Table 2.




Rules for adding triangles		
for $1 \leq i \leq p_1 - 1, 1 \leq j \leq p_2$  if $\alpha_2 \alpha_3 (m_2^{(j)}) = m_1^{(i)}$	for $1 \leq j \leq p_2, 1 \leq k \leq p_3$  if $\alpha_3^{-1} \alpha_2^{-1} \alpha_3 (m_3^{(k)}) = m_2^{(j)}$ (i.e. $\alpha_3 (m_3^{(k)}) = \alpha_2 \alpha_3 (m_2^{(j)})$)	for $1 \leq k \leq p_3, 1 \leq i \leq p_1 - 1$  if $\alpha_3^{-1} (m_1^{(i)}) = m_3^{(k)}$ (i.e. $m_1^{(i)} = \alpha_3 (m_3^{(k)})$)

Table 2: Rules for forming triangles children of white, black and grey vertices in the labeled tricolored tree T in order to obtain the labeled tricolored cactus tree Υ .

Finally, we remove the labels of Υ to obtain a cactus tree τ .

Example 3.4. *The construction of T , Υ , and τ for the partitioned cactus in Example 2.11 is depicted in Figure 6.*

(ii) *Relabeling permutations:* These permutations θ_1, θ_2 , and θ_3 are defined by considering the *reverse* labeled cactus tree Υ' resulting from the labeling of τ , based on three independent




Figure 6: tricolored tree T and cactus trees Υ and τ associated to Example 2.11. The tree T is the 3-colored labeled tree built following rules in Table 1. The labeled cactus tree Υ is obtained from T by forming triangles following the rules in Table 2. The cactus tree τ is obtained from Υ' by removing the labels of the vertices.

reverse-labeling procedure for white, black, and grey vertices. We do a white RLT and label the *white vertices only* (as they are traversed) with the labels $1, 2, \dots, p_1$. Similarly, we do a black RLT and label the *black vertices only* with labels $1, 2, \dots, p_2$, and do a grey RLT to label the *grey vertices only* with labels $1, 2, \dots, p_3$. Next, we reindex the blocks of π_1, π_2 and π_3 using the new indices from Υ' : if a white vertex is labeled i in T and i' in Υ' , we set $\pi_1^{i'} = \pi_1^{(i)}$ (and $m_1^{i'} = m_1^{(i)}$). Black and grey blocks are reindexed in a similar fashion. Let u^i, v^j, w^k be the strings obtained by writing the elements of $\pi_1^i, \pi_2^j, \pi_3^k$ respectively in increasing order. Denote $u = u^1 \dots u^{p_1}$, $v = v^1 \dots v^{p_2}$, $w = w^1 \dots w^{p_3}$ the concatenations of the strings defined above. We define $\theta_1 \in \mathfrak{S}_n$ by setting u as the first line and $[n]$ as the second line of the two-line representation of this permutation. Similarly, we define the relabeling permutations θ_2 and θ_3 from v and w respectively.

Example 3.5. *Following up on Example 3.4, we construct the relabeling permutations θ_1, θ_2 , and θ_3 . We have:*

$$\theta_1 = \left(\begin{array}{cc|cccc} 4 & 5 & 1 & 2 & 3 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array} \right), \quad \theta_2 = \left(\begin{array}{cc|cc} 1 & 3 & 2 & 6 \\ 1 & 2 & 5 & 6 \end{array} \right), \quad \theta_3 = \left(\begin{array}{c|cc|cccc} 5 & 2 & 1 & 3 & 4 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array} \right).$$

We define the multisets:

$$\begin{aligned} S_1 &= \left\{ \theta_1(m_1^i) \right\}_{i=1}^{p_1-1} \cup \left\{ \theta_1(\alpha_2 \alpha_3(m_2'^j)) \right\}_{j=1}^{p_2} \subset [n], \\ S_2 &= \left\{ \theta_2(m_2'^j) \right\}_{j=1}^{p_2-1} \cup \left\{ \theta_2(\alpha_3^{-1} \alpha_2^{-1} \alpha_3(m_3^k)) \right\}_{k=1}^{p_3} \subset [n-1], \\ S_3 &= \left\{ \theta_3(m_3^k) \right\}_{k=1}^{p_3-1} \cup \left\{ \theta_3(\alpha_3^{-1}(m_1^i)) \right\}_{i=1}^{p_1-1} \subset [n-1]. \end{aligned}$$

They are multisets since we allow some elements to be repeated once when there are triangles in Υ' . Note that the sizes of the underlying sets of S_1, S_2 , and S_3 are $p_1 + p_2 - 1 - g$, $p_2 + p_3 - 1 - w$, and $p_1 + p_3 - 2 - b$ respectively. We use these multisets to label the vertices, the edges and triangles of the cactus tree τ with three types of labels: *circle* for θ_1 , *square* for θ_2 , and *triangle* for θ_3 (represented as labels \odot , \square , and \triangle as illustrated in Figure 7). We assign $\theta_1(m_1^i)$ to the white vertices indexed i in Υ' , $\theta_2(m_2'^j)$ to the black vertices indexed

j in Υ' , and $\theta_3(m_3^k)$ to the grey vertices indexed k in Υ' . Children of a white vertex (edges and triangles) are labeled $\theta_1(\alpha_2\alpha_3(m_2^j))$ if the black vertex at the end of the edge or within the triangle is indexed by j in Υ' . The children of the black and grey vertices are labeled in a similar fashion with $\{\theta_2(\alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^k))\}_{k=1}^{p_3}$ and $\{\theta_3(\alpha_3^{-1}(m_1^i))\}_{i=1}^{p_1-1}$ respectively.

Let Υ'' be the resulting cactus tree with these new additional labels.

Let $\overline{S_i}$ (for $i = 1, 2, 3$) be the ordered multiset obtained by arranging the elements of S_i in non-decreasing order (for $i = 1, 2, 3$).

Lemma 3.6. *Let $\mathbf{d} = (d_1, \dots, d_{p_1+p_2-1})$ ($\mathbf{d}' = (d'_1, \dots, d'_{p_2+p_3})$, and $\mathbf{d}'' = (d''_1, \dots, d''_{p_1+p_3-1})$ resp.) be the ordered set of labels of the first type (second and third type resp.) obtained by traversing Υ'' up to, but not including the root vertex according to the white RLT (black and grey RLT resp.) defined in Section 3. We have:*

$$\mathbf{d} = \overline{S_1}, \quad \mathbf{d}' = \overline{S_2}, \quad \mathbf{d}'' = \overline{S_3}.$$

Proof. Let $\theta_1(m_1^0) = 0$. By construction, if $\theta_1(\alpha_2\alpha_3(m_2^j))$ in Υ'' is the label of a child of a white vertex with label $\theta_1(m_1^i)$, $1 \leq i \leq p_1$, then $\alpha_2\alpha_3(m_2^j) \in \pi_1^i$ and $\alpha_2\alpha_3(m_2^j) \leq m_1^i$. As θ_1 is increasing among the blocks and within them then:

$$\theta_1(m_1^{i-1}) < \theta_1(\alpha_2\alpha_3(m_2^j)) < \theta_1(m_1^i),$$

Now, if two children of the same white vertex have respective labels $\theta_1(\alpha_2\alpha_3(m_2^{j_1}))$ and $\theta_1(\alpha_2\alpha_3(m_2^{j_2}))$ in Υ'' , and black vertex j_1 is to the left of j_2 ; then by construction we have $\alpha_2\alpha_3(m_2^{j_1}) < \alpha_2\alpha_3(m_2^{j_2})$. Again, since θ_1 is increasing within blocks of π_1 then $\theta_1(\alpha_2\alpha_3(m_2^{j_1})) < \theta_1(\alpha_2\alpha_3(m_2^{j_2}))$. Finally, the white RLT of the circle labels in Υ'' (up to but not including the root) yields $\overline{S_1}$. Similarly, black and grey RLTs of the square and triangle labels in Υ'' yield $\overline{S_2}$ and $\overline{S_3}$ respectively. \square

(iii) *Thorns:* Recall the definition of \mathbf{d}, \mathbf{d}' , and \mathbf{d}'' in Lemma 3.6. We add thorns to the white vertices for each missing element of $[n]$ in \mathbf{d} , and add thorns to black and grey vertices for each missing element of $[n-1]$ in \mathbf{d}' and \mathbf{d}'' , respectively. More specifically, we add $n+1-p_1-p_2+g$ thorns to the white vertices, $n-p_2-p_3+w$ thorns to the black vertices, and $n+1-p_1-p_3+b$ thorns to the grey vertices of Υ'' in the following fashion:

1. If $d_1 > 1$ and d_1 is the label of a (white) vertex, we connect $d_1 - 1$ thorns to it. If a child of a white vertex has label d_1 , we connect $d_1 - 1$ thorns to the ascending white vertex on the left of child d_1 .
2. For $1 < l < p_1 + p_2 - 1$, if $d_l > d_{l-1} + 1$ we follow one of the four following cases:
 - (a) d_l and d_{l-1} are both the label of white vertices in Υ'' , white vertex d_l (short for vertex corresponding to d_l) has no child and it is the white vertex following d_{l-1} in the white RLT of Υ'' . If so, we connect $d_l - d_{l-1} - 1$ thorns to d_l .
 - (b) d_l is the first label of a child and d_{l-1} is the first label of a white one, then d_l is the leftmost child of the white vertex following d_{l-1} . If so, we connect $d_l - d_{l-1} - 1$ thorns to the ascending white vertex of d_l on its left
 - (c) d_l is the first label of a white vertex and d_{l-1} is the first label of a child, then d_{l-1}

is the rightmost child of d_l . If so, we connect $d_l - d_{l-1} - 1$ thorns to d_l on the right of d_{l-1} .

(d) Finally, if d_l and d_{l-1} are both the first label of children, they have the same white ascending vertex. We connect $d_l - d_{l-1} - 1$ thorns to the ascending white vertex between them.

3. If $d_{p_1+p_2-1} < n$, we connect $n - d_{p_1+p_2-1} - 1$ thorns to the root vertex on the right of its rightmost child.

Again, we can think of this as adding a thorn to the the white vertices for each element of $[n]$ not included in \mathbf{d} .

A similar construction is applied to add thorns to the black and grey vertices following the sequence of integers \mathbf{d}' and \mathbf{d}'' . Finally we remove all the labels to get the cactus tree $\tilde{\tau}$.

Example 3.7. Figure 7 depicts the construction of the cactus tree $\tilde{\tau}$ corresponding to the partitioned cactus in Example 2.11.




Figure 7: Construction of the cactus tree $\tilde{\tau}$ associated to Example 2.11

The next two lemmas show that $\tilde{\tau}$ preserves the type of the partitioned cacti, and that Υ'' can be recovered from $\tilde{\tau}$ via white, black, and grey RLTs.

Lemma 3.8. $\tilde{\tau}$ as defined above belongs to $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ where g, w, b are the number of triangles in $\tilde{\tau}$ rooted in grey, white, and black vertices respectively.

Proof. We check the vertex degrees of $\tilde{\tau}$. If we take two successive white vertices $i - 1$ and i according to white RLT of Υ'' with labels $\theta_1(m_1^{i-1})$ and $\theta_1(m_1^i)$, ($i < p_1$), a thorn is connected to i for any missing integer of the interval $[\theta_1(m_1^{i-1}), \theta_1(m_1^i) - 1]$ in \mathbf{d} . This number of missing integers is equal to $\theta_1(m_1^i) - 1 - \theta_1(m_1^{i-1}) - f_i$ where f_i is the number of children of i . As i is not the root vertex, there is an edge between i and its ancestor so that the resulting degree \deg for i (as defined in (2.17)) is:

$$\deg(i) = f_i + (\theta_1(m_1^i) - 1 - \theta_1(m_1^{i-1}) - f_i) + 1 = \theta_1(m_1^i) - \theta_1(m_1^{i-1}), \quad \forall i \in [p_1 - 1], \quad (3.9)$$

Furthermore, $n - \theta_1(m_1^{p_1-1}) - f_p$ thorns are connected to the root vertex (since $\varepsilon_{p_1} = 0$) so that:

$$\deg(p_1) = n - \theta_1(m_1^{p_1-1}) \quad (3.10)$$

But, according to the construction of θ ,

$$\theta_1(\pi_1^1) = [\theta_1(m_1^1)] \quad (3.11)$$

$$\theta_1(\pi_1^i) = [\theta_1(m_1^i)] \setminus [\theta_1(m_1^{i-1})], \quad (2 \leq i \leq p_1 - 1) \quad (3.12)$$

$$\theta_1(\pi_1^{p_1}) = [n] \setminus [\theta_1(m_1^{p_1-1})] \quad (3.13)$$

Subsequently:

$$\deg(i) = |\pi_1^i|, \quad \forall i \in [p_1]. \quad (3.14)$$

And $\lambda = \text{type}(\pi)$ is the white vertex degree distribution of $\tilde{\tau}$. In a similar fashion, μ and ν are the black and grey vertex degree distribution of $\tilde{\tau}$. \square

Lemma 3.15. *Assign circle labels $1, 2, \dots, n$ to the white vertices and their children (including thorns) in $\tilde{\tau}$ in increasing order according to the white RLT, add two other sets of labels $1, 2, \dots, n$ (square and triangle) to the black and grey vertices and their children in increasing order according to the black and grey RLT. The labeling of the vertices and children that are not thorns is the same as in Υ'' .*

Proof. According to the construction of $\tilde{\tau}$, we add thorns to Υ'' when integers are missing in its RLTs so that the thorns would take these missing integers as labels when traversing the cactus tree. As a result, the labels of the vertices in the RLTs of $\tilde{\tau}$ are still \mathbf{d}, \mathbf{d}' , and \mathbf{d}'' and since they still appear in the same order, we have the desired result. \square

3.2.2 The permutations σ_1 and σ_2 and the ordered set χ and the

In the previous subsection we explained how to obtain the cactus tree $\tilde{\tau}$ from the partitioned 3-cactus in $\mathcal{C}(\lambda, \mu, \nu)$. We now move on to explain how to obtain the permutations σ_1 in $\mathfrak{S}_{n+1-\ell(\lambda)-\ell(\nu)+b}$ and σ_2 in $\mathfrak{S}_{n-\ell(\lambda)-\ell(\nu)+b}$, and the ordered set χ in $\mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}$.

(i) *Permutations σ_1, σ_2 :* Let E and F be the following sets:

$$E = [n] \setminus \left(\left\{ \theta_1(m_1^i) \right\}_{i=1}^{p_1-1} \cup \left\{ \theta_1(\alpha_3(m_3^k)) \right\}_{k=1}^{p_3-1} \right),$$

$$F = [n] \setminus \left(\left\{ \theta_1(\alpha_2 \alpha_3(m_2'^j)) \right\}_{j=1}^{p_2} \cup \left\{ \theta_1(\alpha_3(m_3^k)) \right\}_{k=1}^{p_3-1} \right).$$

We define partial permutations $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ in the following way:

$$\begin{aligned} \tilde{\sigma}_1 : E &\rightarrow [n-1] \setminus S_3 \\ u &\mapsto \theta_3 \alpha_3^{-1} \theta_1^{-1}(u) \\ \tilde{\sigma}_2 : F &\rightarrow [n-1] \setminus S_2 \\ u &\mapsto \theta_2 \alpha_3^{-1} \alpha_2^{-1} \theta_1^{-1}(u). \end{aligned}$$

Let $\sigma_1 \in \mathfrak{S}_{n+1-p_1-p_3+b}$ and $\sigma_2 \in \mathfrak{S}_{n-p_2-p_3+w}$ be the order isomorphic permutations corresponding to $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ respectively.

(ii) *Ordered set χ :* We define the ordered set $\tilde{\chi} = \{ \theta_1(\alpha_3(m_3^k)) \mid \theta_1(\alpha_3(m_3^k)) \notin S_1 \}_{k=1}^{p_3}$. Then, let $\rho : [n] \setminus S_1 \vdash [n - |S_1|]$ be the indexing permutation associating to any integer




Figure 8: Summary of output $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ of the map $\Theta_{\lambda, \mu, \nu}^n$ applied to the partitioned 3-cactus from Example 2.11.

$i \in [n] \setminus S_1$ its position in $\overline{[n] \setminus S_1}$ where $\overline{[n] \setminus S_1}$ is the ordered (increasing) set of $[n] \setminus S_1$. The ordered set χ is defined as follows:

$$\chi = \rho(\tilde{\chi}) \quad (3.16)$$

As $|S_1| = n - (\ell(\lambda) - 1) - \ell(\mu) + g$ and $|\{\theta_1(\alpha_3(m_3^k)) \mid \theta_1(\alpha_3(m_3^k)) \notin S_1\}| = \ell(\nu) - w - b$, χ belongs to the set $\mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}$.

Example 3.17. Getting back to Example 2.11, computing the partial permutations leads to:

$$\tilde{\sigma}_1 = \begin{pmatrix} 1 & 3 & 5 \\ 4 & 5 & 3 \end{pmatrix}, \quad \tilde{\sigma}_2 = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix},$$

and

$$\sigma_1 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}.$$

For the ordered set we have: $\tilde{\chi} = (4)$ and $\chi = (3)$.

In summary, the map $\Theta_{\lambda, \mu, \nu}^n$ applied to $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ in $\mathcal{C}([2^1, 4^1], [2^1, 4^1], [1^2, 4^1])$ from Example 2.11 gives the 4-tuple $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ depicted in Figure 8.

3.3 Showing the map Θ is a bijection

To show that $\Theta_{\lambda, \mu, \nu}^n$ is a one-to-one correspondence we take any element $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ in

$$\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b) \times \mathfrak{S}_{n+1-\ell(\lambda)-\ell(\nu)+b} \times \mathfrak{S}_{n-\ell(\mu)-\ell(\nu)+w} \times \mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}$$

and show that there is a unique element $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ in $\mathcal{C}(\lambda, \mu, \nu)$ such that $\Theta_{\lambda, \mu, \nu}^n(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) = (\tilde{\tau}, \sigma_1, \sigma_2, \chi)$. Let $p_1 = \ell(\lambda)$, $p_2 = \ell(\mu)$, and $p_3 = \ell(\nu)$. We proceed with a two step proof:

- (i) The first step is to notice that $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ defines a unique cactus tree τ belonging to $\mathcal{CT}(p_1, p_2, p_3, g, w, b)$, unique multisets $\{S_i\}_{1 \leq i \leq 3}$, as well as a unique ordered set $\tilde{\chi}$ belonging to $\mathcal{OP}_{p_3-w-b}^{(n+1-p_1-p_2+g)}$. Labeling each vertex and children of τ with $1, 2, \dots, n$

in increasing order according to the three reverse levels traversals and removing the three sets of thorns (together with their labels) gives a labeled cactus tree Υ'' that leads to $\tilde{\tau}$ according to Θ . This labeled cactus tree is the unique one that can lead to $\tilde{\tau}$ since within Θ , $\tilde{\tau}$, and Υ'' have the same underlying cactus tree structure τ , and according to Lemma 3.15, $\tilde{\tau}$ determines the labels of Υ'' .

Then, using Lemma 3.6, the three series of labels (except the root's) in Υ'' are by construction the three sets $\{S_i\}_{1 \leq i \leq 3}$. The knowledge of S_1 and χ uniquely determines $\tilde{\chi}$. As a result, exactly one 7-tuple $(\tau, S_1, S_2, S_3, \sigma_1, \sigma_2, \tilde{\chi})$ is associated to $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ by the final steps of the mapping Θ .

- (ii) The bijection Θ_{n,p_1,p_2,p_3} in [18] is identical to the first steps (up to the construction of $\tau, S_1, S_2, S_3, \sigma_1, \sigma_2$ and $\tilde{\chi}$) of $\Theta_{\lambda,\mu,\nu}^n$. Therefore by [18, Sec. 6] there is a unique 5-tuple $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ in $\mathcal{C}(p_1, p_2, p_3, n) = \bigcup_{\ell(\lambda)=p_1, \ell(\mu)=p_2, \ell(\nu)=p_3} \mathcal{C}(\lambda, \mu, \nu)$ mapped to the 7-tuple $(\tau, S_1, S_2, S_3, \sigma_1, \sigma_2, \tilde{\chi})$ by Θ_{n,p_1,p_2,p_3} and equivalently by the first steps of $\Theta_{\lambda,\mu,\nu}^n$. According to [18], the types of π_1 , π_2 , and π_3 are directly recovered from $\{S_i\}_{1 \leq i \leq 3}$ and τ . Furthermore, using Lemma 3.8, the vertex degree distribution of $\tilde{\tau}$ is equal to the type of the partitions encoded by the elements in $\{S_i\}_{1 \leq i \leq 3}$ corresponding to the relabeling of the maximum elements of the blocks. Finally, as the vertex degree distribution in $\tilde{\tau}$ is (λ, μ, ν) , so is the type of (π_1, π_2, π_3) . Therefore, $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ belongs to $\mathcal{C}(\lambda, \mu, \nu)$ as desired.

4 Proof of Proposition 2.19: computation of the number of cactus trees

In this section we prove Proposition 2.19 where we compute the cardinality of the set $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$. To do this, we consider its generating function F :

$$F = \sum_{\lambda, \mu, \nu \vdash n} \sum_{g, w, b \geq 0} |\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)| x_1^{\ell(\lambda)} x_2^{\ell(\mu)} x_3^{\ell(\nu)} x_4^g x_5^w x_6^b \mathbf{t}^{\mathbf{n}(\lambda)} \mathbf{u}^{\mathbf{n}(\mu)} \mathbf{v}^{\mathbf{n}(\nu)}. \quad (4.1)$$

That is, the white, black, and grey vertices are marked respectively by indeterminates x_1, x_2 and x_3 . Triangles children of a grey, white, and black vertex are marked respectively by x_4, x_5 , and x_6 . Furthermore, t_i, u_j , and v_k mark respectively white vertices of degree i , black vertices of degree j and grey vertices of degree k . And $\mathbf{t} = (t_1, t_2, \dots)$, $\mathbf{u} = (u_1, u_2, \dots)$, $\mathbf{v} = (v_1, v_2, \dots)$ and $\mathbf{n}(\epsilon) = (n_1(\epsilon), n_2(\epsilon), \dots)$ for $\epsilon \vdash n$ where $n_i(\epsilon)$ is the number of i parts of ϵ .

The evaluation of F is performed thanks to the multivariate Lagrange inversion theorem (see e.g. [7, 1.2.13]). We propose a recursive decomposition of the desired set of cactus trees sharing similar ideas with [6].

In a similar fashion as in [6], we introduce W , B , and G as the generating functions of the sets \mathcal{W} , \mathcal{B} , and \mathcal{G} of non empty *planted* cactus trees with respectively white, black, and grey root vertices. Construction rules for these sets of cactus trees are identical to those of $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ with the only exception that an additional *planted* edge is connected to the root vertex on the left of the leftmost child (vertex or thorn). We take this additional edge

into account in the root's degree. Finally, let T_g , T_w , and T_b be respectively the generating functions of triangles children of a grey, white, and black vertices. Immediately:

$$T_g = x_4 \quad (4.2)$$

$$T_w = x_5 \quad (4.3)$$

$$T_b = x_6 \quad (4.4)$$

Any cactus tree in $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ can be decomposed in a tuple of planted cactus trees in \mathcal{W} , \mathcal{B} , and \mathcal{G} . The rule for the decomposition is based on the nature of the leftmost child of the white root in a given cactus tree τ of $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$:

- (i) If the leftmost child is a thorn then τ is equivalent to the cactus tree in \mathcal{W} with the planted edge instead of this leftmost thorn.
- (ii) If the leftmost child is an edge connected to black vertex v , then τ is equivalent to the pair (τ_1, τ_2) in $\mathcal{W} \times \mathcal{B}$ where τ_2 is the cactus tree descending from v with v as the root and the edge linking v to the root of τ replaced by the planted edge. τ_1 is the remaining cactus tree descending from the root of τ with the edge linking it to v as the planted edge.
- (iii) If the leftmost child is a triangle containing black vertex v_1 and grey vertex v_2 then τ is equivalent to the tuple $(\tau_1, \tau_2, \tau_3, t_w)$ in $\mathcal{W} \times \mathcal{B} \times \mathcal{G} \times TT_w$ (TT_w is the singleton set composed of the triangle child of a white vertex) where τ_2 and τ_3 are the descending trees from v_1 and v_2 with the edge linking τ 's root and v_1 and the edge linking v_1 and v_2 replaced by a planted edge. τ_1 is the remaining descending cactus tree from its root with the leftmost triangle replaced by the planted edge.

One can check easily that the numbers of triangles, white, black, and grey vertices and their degree distribution are stable by the bijective transformation described above. The complicated case above is case (iii) where the edges linking v_1 and v_2 , and the edge linking the white root of τ and v_2 are replaced by nothing in (respectively) τ_2 and τ_1 . However in Definition 2.17 of the degree of a vertex in τ , these edges were already not taken into account for the degree of respectively v_1 and the root vertex. As a consequence :

$$F = W + W \cdot B + W \cdot B \cdot G \cdot T_w \quad (4.5)$$

This decomposition is illustrated in Figure 9.

To determine F , we show that W, B, G, T_g, T_w , and T_b satisfy a system of functional equations. Namely, as shown in Figure 10 any planted cactus tree in \mathcal{W} , τ can be decomposed into:

- its white root,
- the cactus trees rooted in a black vertex descending from the root,
- a triple composed of a black rooted cactus tree, a grey rooted cactus tree, a triangle for each triangle descending from the root,
- the positions of the triangles in the list of children.

Let i denote the degree of the root vertex (of degree $i + 1$), j the number of black children not belonging to a triangle and k the number of descending triangles. The vectors \mathbf{j} and \mathbf{k} give the positions of the j black vertices and k triangles within the i children. Using the




Figure 9: Illustration of the decomposition into planted trees

decomposition above, we have:

$$W = x_1 \sum_{i \geq 0} t_{i+1} \sum_{0 \leq j+k \leq i} \sum_{\mathbf{j}, \mathbf{k}} B^j (B \cdot G \cdot T_w)^k \quad (4.6)$$

Then:

$$\begin{aligned} W &= x_1 \sum_{i \geq 0} t_{i+1} \sum_{0 \leq j+k \leq i} \binom{i}{j, k} B^j (B \cdot G \cdot T_w)^k \\ W &= x_1 \sum_{i \geq 0} t_{i+1} (1 + B + B \cdot G \cdot T_w)^i \end{aligned}$$

Similarly,

$$\begin{aligned} B &= x_2 \sum_{i \geq 0} u_{i+1} (1 + G + G \cdot W \cdot T_b)^i \\ G &= x_3 \sum_{i \geq 0} v_{i+1} (1 + W + W \cdot B \cdot T_g)^i \end{aligned}$$

Finally :

$$(W, B, G, T_g, T_w, T_b) = \mathbf{x} \Phi(W, B, G, T_g, T_w, T_b) \quad (4.7)$$




Figure 10: Decomposition of a white rooted planted cactus tree

where $\mathbf{x} = (x_1, x_2, x_3, x_4, x_5, x_6)$ and $\Phi = (\Phi_i)_{1 \leq i \leq 6}$ with:

$$\Phi_1(W, B, G, T_g, T_w, T_b) = \sum_{i \geq 0} t_{i+1} (1 + B + B \cdot G \cdot T_w)^i \quad (4.8)$$

$$\Phi_2(W, B, G, T_g, T_w, T_b) = \sum_{i \geq 0} u_{i+1} (1 + G + G \cdot W \cdot T_b)^i \quad (4.9)$$

$$\Phi_3(W, B, G, T_g, T_w, T_b) = \sum_{i \geq 0} v_{i+1} (1 + W + W \cdot B \cdot T_g)^i \quad (4.10)$$

$$\Phi_i = 1 \text{ for } 4 \leq i \leq 6 \quad (4.11)$$

Using the multivariate Lagrange inversion formula for monomials (see [7, 1.2.9]), we find :

$$k_1 k_2 k_3 k_4 k_5 k_6 [\mathbf{x}^{\mathbf{k}}] W^{r_1} B^{r_2} G^{r_3} T_w^{r_5} = \sum_{\{\mu_{ij}\}} \|\delta_{ij} k_j - \mu_{ij}\| \prod_{1 \leq i \leq 6} [W^{\mu_{i1}} B^{\mu_{i2}} G^{\mu_{i3}} T_g^{\mu_{i4}} T_w^{\mu_{i5}} T_b^{\mu_{i6}}] \Phi_i^{k_i} \quad (4.12)$$

where $\|\cdot\|$ denotes the determinant, $\mathbf{k} = (k_1, k_2, k_3, k_4, k_5, k_6)$, δ_{ij} is the Kronecker delta function and the sum is over all 6×6 integer matrices $\{\mu_{ij}\}$ such that:

- $\mu_{11} = \mu_{14} = \mu_{16} = \mu_{22} = \mu_{24} =$

- $\mu_{25} = \mu_{33} = \mu_{35} = \mu_{36} = 0$

- $\mu_{ij} = 0$ for $i \geq 4$

- $\mu_{21} + \mu_{31} = k_1 - r_1$

- $\mu_{12} + \mu_{32} = k_2 - r_2$

- $\mu_{13} + \mu_{23} = k_3 - r_3$

- $\mu_{34} = k_4$

- $\mu_{15} = k_5 - r_5$

- $\mu_{26} = k_6$

$$\text{i.e. } \mu = \begin{bmatrix} 0 & * & * & 0 & * & 0 \\ * & 0 & * & 0 & 0 & * \\ * & * & 0 & * & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Looking for zero contribution terms in expression (4.12), we notice that G and T_w have necessarily the same degree in the formal power series expansion of Φ_1 . Hence, a non zero contribution of

$$[W^{\mu_{11}} B^{\mu_{12}} G^{\mu_{13}} T_1^{\mu_{14}} T_2^{\mu_{15}} T_3^{\mu_{16}}] \Phi_1^{k_1}$$

implies $\mu_{13} = \mu_{15} = k_5 - r_5$. Similar remarks give non zero contributions only for $\mu_{21} = k_6$ and $\mu_{32} = k_4$. As a result, only that one matrix μ yields a non zero contribution.

For this particular μ ,

$$\begin{aligned} \frac{1}{k_4 k_5 k_6} \|\delta_{ij} k_j - \mu_{ij}\| &= r_1 (k_2 k_3 - (k_3 - k_5) k_4) \\ &+ r_2 (k_6 k_3 + (k_6 + r_1 - k_1)(r_3 + k_4 - r_5 - k_3)) \\ &+ r_3 (k_4 k_6 - k_2(k_6 + r_1 - k_1)) \end{aligned} \quad (4.13)$$

$$= \frac{1}{k_4 k_5 k_6} \Delta(\mathbf{k}, \mathbf{r}). \quad (4.14)$$

Let $co(k)$ denotes the set of sequences of non negative integers of total sum k . The next step is to notice that:

$$\begin{aligned} \Phi_1^{k_1} &= \sum_{\mathbf{s} \in co(k_1)} \binom{k_1}{\mathbf{s}} \prod_{i \geq 0} \left[t_{i+1} (1 + B + B \cdot G \cdot T_w)^i \right]^{s_i} \\ \Phi_1^{k_1} &= \sum_{\mathbf{s} \in co(k_1)} \binom{k_1}{\mathbf{s}} \prod_{i \geq 0} t_{i+1}^{s_i} \sum_{a_1, a_2} \binom{\sum_i i s_i}{a_1 - a_2, a_2} B_1^a (GT_w)^{a_2}. \end{aligned} \quad (4.15)$$

As a result, the coefficient in $W^{\mu_{11}} B^{\mu_{12}} G^{\mu_{13}} T_g^{\mu_{14}} T_w^{\mu_{15}} T_b^{\mu_{16}}$ is equal to

$$\sum_{\mathbf{s} \in co(k_1)} \binom{k_1}{\mathbf{s}} \prod_{i \geq 0} t_{i+1}^{s_i} \binom{\sum_i i s_i}{k_5 - r_5, k_2 - k_4 - k_5 - r_2 + r_5}. \quad (4.16)$$

Similarly, we have:

$$\begin{aligned} [W^{\mu_{21}} B^{\mu_{22}} G^{\mu_{23}} T_g^{\mu_{24}} T_w^{\mu_{25}} T_b^{\mu_{26}}] \Phi_2^{k_2} &= \\ \sum_{\mathbf{s} \in co(k_2)} \binom{k_2}{\mathbf{s}} \prod_{i \geq 0} u_{i+1}^{s_i} \binom{\sum_i i s_i}{k_6, k_3 - k_5 - k_6 - r_3 + r_5} \end{aligned} \quad (4.17)$$

$$\begin{aligned} [W^{\mu_{31}} B^{\mu_{32}} G^{\mu_{33}} T_g^{\mu_{34}} T_w^{\mu_{35}} T_b^{\mu_{36}}] \Phi_3^{k_3} &= \\ \sum_{\mathbf{s} \in co(k_3)} \binom{k_3}{\mathbf{s}} \prod_{i \geq 0} v_{i+1}^{s_i} \binom{\sum_i i s_i}{k_4, k_1 - k_4 - k_6 - r_1}. \end{aligned} \quad (4.18)$$

Putting everything together gives:

$$\begin{aligned} [x_1^{\ell(\lambda)} x_2^{\ell(\mu)} x_3^{\ell(\nu)} x_4^g x_5^w x_6^b \mathbf{t}^{\mathbf{n}(\lambda)} \mathbf{u}^{\mathbf{n}(\mu)} \mathbf{v}^{\mathbf{n}(\nu)}] W^{r_1} B^{r_2} G^{r_3} T_w^{r_5} &= \\ \frac{\Delta(\ell(\lambda), \ell(\mu), \ell(\nu), g, w, b, \mathbf{r})}{\ell(\lambda) \ell(\mu) \ell(\nu)} \binom{\ell(\lambda)}{\mathbf{n}(\lambda)} \binom{\sum_i i n_{i+1}(\lambda)}{w - r_5, \ell(\mu) - g - w - r_2 + r_5} \times \\ \times \binom{\ell(\mu)}{\mathbf{n}(\mu)} \binom{\sum_i i n_{i+1}(\mu)}{b, \ell(\nu) - w - b - r_3 + r_5} \\ \times \binom{\ell(\nu)}{\mathbf{n}(\nu)} \binom{\sum_i i n_{i+1}(\nu)}{g, \ell(\lambda) - g - b - r_1}. \end{aligned} \quad (4.19)$$

Noticing that for $\epsilon \vdash n$

$$\sum_{i \geq 0} i n_{i+1}(\epsilon) = \sum_{i \geq 0} (i+1) n_{i+1}(\epsilon) - \sum_{i \geq 0} n_{i+1}(\epsilon) = n - \ell(\epsilon). \quad (4.20)$$

And summing for $\mathbf{r} \in \{(1, 0, 0, 0), (1, 1, 0, 0), (1, 1, 1, 1)\}$ gives the desired result.

5 Proof of Corollary 1.3 and restriction of bijection Θ when $\nu = [1^n]$

We look more closely at the case when one of the partitions, say ν , is $[1^n]$. We need the following definitions:

Definition 5.1 (Partitioned bicolored map). *Given partitions $\lambda, \mu \vdash n$, let $\mathcal{C}(\lambda, \mu)$ be the set of triples (π_1, π_2, α) such that $\alpha \in \mathfrak{S}_n$, π_1, π_2 are set partitions of $[n]$ with $\text{type}(\pi_1) = \lambda$ and $\text{type}(\pi_2) = \mu$, and each block of π_1 and π_2 is a union of cycles of α and $\beta = \alpha^{-1}\gamma_n$ respectively. The elements of $\mathcal{C}(\lambda, \mu)$ are called unicellular partitioned bicolored maps of type λ and μ . Let $C(\lambda, \mu) = |\mathcal{C}(\lambda, \mu)|$.*

Definition 5.2 (Ordered rooted bicolored thorn trees). *For $\lambda, \mu \vdash n$ such that $\ell(\lambda) + \ell(\mu) \leq n + 1$, we define $\widetilde{\mathcal{BT}}(\lambda, \mu)$ as the set of ordered rooted bicolored trees with $\ell(\lambda)$ white vertices, $\ell(\mu)$ black vertices, $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to the black vertices and $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to the white vertices. The white (respectively black) vertices' degree distribution (accounting the thorns) is specified by λ (respectively μ). The root is a white vertex.*

Again, adapting the Lagrange inversion developed in [6], we get:

$$|\widetilde{\mathcal{BT}}(\lambda, \mu)| = \frac{n}{\text{Aut}(\lambda)\text{Aut}(\mu)} \frac{(n - \ell(\lambda))!(n - \ell(\mu))!}{(n + 1 - \ell(\lambda) - \ell(\mu))!^2}.$$

We now prove Corollary 1.3:

Proof. We have $\mathcal{C}(\lambda, \mu, [1^n]) = \mathcal{C}(\lambda, \mu)$, the number of unicellular partitioned bicolored maps of type λ and μ . Indeed, as the cycles of α_3 refine the blocks of π_3 , if $\nu = [1^n]$ then $\pi_3 = \{\{1\}, \{2\}, \dots, \{n\}\}$ and $\alpha_3 = \iota$, the identity permutation. Then extracting the coefficient of $m_{1^n}(\mathbf{z})$ to both sides of (2.14) we obtain

$$\begin{aligned} \sum_{\lambda, \mu \vdash n} \text{Aut}(\lambda)\text{Aut}(\mu)\text{Aut}(1^n)C(\lambda, \mu)m_\lambda(\mathbf{x})m_\mu(\mathbf{y}) &= [m_{1^n}(\mathbf{z})] \sum_{\lambda, \mu, \nu \vdash n} k_{\lambda, \mu, \nu}^n p_\lambda(\mathbf{x})p_\mu(\mathbf{y})p_\nu(\mathbf{z}) \\ &= \sum_{\nu \vdash n, \nu \preceq 1^n} \text{Aut}(1^n)\bar{R}_{\nu, 1^n} \sum_{\lambda, \mu \vdash n} k_{\lambda, \mu, \nu}^n p_\lambda(\mathbf{x})p_\mu(\mathbf{y}) \end{aligned}$$

Since $\bar{R}_{\nu, 1^n} = 1$ if $\nu = 1^n$ and zero otherwise, we obtain

$$\sum_{\lambda, \mu \vdash n} \text{Aut}(\lambda)\text{Aut}(\mu)C(\lambda, \mu)m_\lambda(\mathbf{x})m_\mu(\mathbf{y}) = \sum_{\lambda, \mu \vdash n} k_{\lambda, \mu, 1^n}^n p_\lambda(\mathbf{x})p_\mu(\mathbf{y}),$$

where $k_{\lambda, \mu, 1^n}^n = k_{\lambda, \mu}^n$. □

Next, we say what the bijection $\Theta_{\lambda, \mu, \nu}^n$ of Theorem 2.23 does in this case ($\nu = [1^n]$). This matches the bijection in [15] which in turn matches the bijection in [6] when $\mathbf{g}(\lambda, \mu) = 0$ and is a refinement of a bijection in [17].

Corollary 5.3. *There is a bijection between partitioned bicolored maps $\mathcal{C}(\lambda, \mu, n)$ and pairs (\tilde{t}, σ) where $\tilde{t} \in \widetilde{\mathcal{BT}}(\lambda, \mu)$ and $\sigma \in \mathfrak{S}_{n+1-\ell(\lambda)-\ell(\mu)}$.*




Table 3: Local rules for reducing cactus tree $\tilde{\tau}$ when $\nu = [1^n]$.

Proof. From above we have that $\mathcal{C}(\lambda, \mu, [1^n]) = \mathcal{C}(\lambda, \mu)$. Let

$$(\tilde{\tau}, \sigma_1, \sigma_2, \chi) := \Theta_{\lambda, \mu, [1^n]}^n(\pi_1, \pi_2, [1^n], \alpha_1, \alpha_1^{-1}\gamma)$$

for $(\pi_1, \pi_2, \alpha_1) \in \mathcal{C}(\lambda, \mu)$. We know that $\ell(\mu) = n$ forces $\alpha_3 = \iota$, the identity permutation. In this case $m_2^{(j)}$ is the maximal element of $\alpha_2^{-1}(\pi_2^{(j)})$. But α_2 preserves the blocks of π_2 , thus $m_2^{(j)}$ is just the maximal element of $\pi_2^{(j)}$, call this $m_2^{(j)}$. First, we show that in this case $\tilde{\tau}$ can be reduced to a tree $\tilde{t} \in \widetilde{\mathcal{BT}}(\lambda, \mu)$. Then, we show that σ_1, σ_2 are trivial permutations and that χ can be regarded as a permutation in $\mathfrak{S}_{n+1-\ell(\lambda)-\ell(\mu)}$.

From the incidence rules in Table 1, we see that each black vertex j has $|\pi_2^j|$ children (one for each element of the block). And $\ell(\lambda) - 1$ of the grey vertices have one child (one for each $m_1^{(i)}$, $1 \leq i \leq \ell(\lambda) - 1$), the other grey vertices have none. Recall w, b, g count the number of triangles in $\tilde{\tau}$ children of white, black, and grey vertices respectively. From the rules in Table (2) for adding triangles children of the different vertices, we see that $w = \ell(\mu)$. And if a grey vertex has a white child, then these two vertices are part of a triangle child of a black vertex, so $b = \ell(\lambda) - 1$. For triangles children of grey vertices, if $\alpha_2(m_2^{(j)}) = m_1^{(i)}$ for some i and j ($1 \leq i \leq \ell(\lambda) - 1$ and $1 \leq j \leq \ell(\mu)$), then $m_1^{(i)} \in \pi_2^{(j)}$ and $m_1^{(i)} \leq m_2^{(j)}$ (α_2 preserves blocks of π_2). But $\alpha_1(m_1^{(i)}) \leq m_1^{(i)}$ (α_1 preserves blocks of π_1), so $\gamma(m_2^{(j)}) \leq m_2^{(j)}$. This only happens if $m_2^{(j)} = n$ which means $1 = \gamma(n) \in \pi^{(i)}$ and $i = \ell(\lambda)$, a contradiction. Thus $g = 0$; there are no triangles children of a grey vertex.

In terms of the thorns, the cactus $\tilde{\tau}$ has $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to white vertices and since $n - \ell(\mu) - \ell(\nu) + w = 0$ and $n + 1 - \ell(\lambda) - \ell(\nu) + b = 0$, $\tilde{\tau}$ has no thorns connected to black and grey vertices.

From above we see that each grey vertex is either: (i) within a triangle child of a black vertex, (ii) a vertex of a triangle child of a white vertex, and (iii) a leaf (note that there are $n - (\ell(\lambda) - 1) - \ell(\mu)$ of these). Then depending on the case we do the following reductions: (i) and (ii) triangle to the edge linking the white and the black vertex, (iii) leaf to thorn connected to a black vertex. We summarize this reduction graphically in Table 3:

The outcome is an ordered bicolored tree \tilde{t} with $\ell(\lambda)$ white vertices and $\ell(\mu)$ black vertices. This tree \tilde{t} has $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to white vertices and $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to black vertices. Moreover, this reduction $\tilde{\tau} \rightarrow \tilde{t}$ is reversible.

In addition, since $\tilde{\tau}$ had no thorns connected to black and grey vertices ($n + 1 - \ell(\lambda) - \ell(\nu) + b = 0$ and $n - \ell(\mu) - \ell(\nu) + w = 0$), then σ_1 and σ_2 are trivial permutations. Since $\ell(\nu) - w - b = n + 1 - \ell(\lambda) - \ell(\mu) = n + 1 - \ell(\lambda) - \ell(\mu) + g$, then we see that χ is just a permutation σ in $\mathfrak{S}_{n+1-\ell(\lambda)-\ell(\mu)}$.

In summary, we have a bijection from (λ, μ, n) to the desired pair (\tilde{t}, σ) . \square

Example 5.4. Let $\alpha_1 = (189\ 10)(25)(3467)$, $\alpha_2 = (15427)(3)(6)(8)(9)(10)$, $\alpha_3 = \iota(\alpha_1\alpha_2 = \alpha_1\alpha_2\alpha_3 = \gamma_{10})$, $\pi_1 = \{\{3, 4, 6, 7\}, \{1, 2, 5, 8, 9, 10\}\}$, $\pi_2 = \{\{1, 2, 4, 5, 7, 10\}, \{3, 9\}, \{6, 8\}\}$, $\pi_3 = \{\{1\}, \{2\}, \dots, \{10\}\}$. Then $\Theta_{\lambda, \mu, [1^n]}^n(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) = (\tilde{\tau}, \emptyset, \emptyset, 251364)$ where $\tilde{\tau}$ and its reduction \tilde{t} are depicted below:




Figure 11: Example of reduction of a cactus tree to a bicolored thorn trees when $\nu = [1^{10}]$.

Acknowledgements

We thank the anonymous referee for several helpful comments and suggestions. A.H.M. visited the Combinatorial Models Group of École Polytechnique while working on this project.

References

- [1] F. Bédard and A. Goupil The poset of conjugacy classes and decomposition of products in the symmetric group, *Can. Math. Bull.*, 35(2):152–160, 1992.
- [2] O. Bernardi. An analogue of the Harer-Zagier formula for unicellular maps on general surfaces, *arXiv:1011.2311*. 2010.
- [3] P. Biane. Nombre de factorisations d’un grand cycle, *Séminaire Lotharingien de Combinatoire*, 51:4, 2004.
- [4] G. Chapuy. *Combinatoire bijective des cartes de genre supérieur*, PhD. thesis, École Polytechnique, June 2009.
- [5] G. Chapuy. A new combinatorial identity for unicellular maps, via a direct bijective approach, *DMTCS Proceedings (FPSAC)*, AK:289–300, 2009.
- [6] I. P. Goulden and D. M. Jackson. The Combinatorial Relationship Between Trees, Cacti and Certain Connection Coefficients for the Symmetric Group, *European Journal of Combinatorics*, 13:357–365, 1992.
- [7] I. P. Goulden and D. M. Jackson. *Combinatorial Enumeration*, Dover, Mineola N.Y. 2004.
- [8] I. P. Goulden and A. Nica A direct bijection for the Harer-Zagier formula, *J. Comb. Theory Ser. A*, 111(2):224–238, 2005.
- [9] A. Goupil A. and G. Schaeffer. Factoring n-cycles and counting maps of given genus, *European Journal of Combinatorics*, 19:819–834(16), 1998.
- [10] J. Harer and D. Zagier. The Euler characteristic of the moduli space of curves. *Invent. Math.*, 85(3):457–485, 1986.

- [11] J. Irving. On the number of factorizations of a full cycle, *J. Comb. Theory Ser. A*, 113(7):1549–1554, 2006.
- [12] D. M. Jackson. Some combinatorial problems associated with products of conjugacy classes of the symmetric group, *J. Comb. Theory Ser. A*, 49(2):363–369, 1988.
- [13] S. Lando and A.K. Zvonkin *Graphs on surfaces and their applications*, Springer-Verlag, 2004.
- [14] B. Lass Démonstration combinatoire de la formule de Harer-Zagier, *C.R. Acad. Sci. Paris*, 333.Série I:155–160, 2001.
- [15] A. H. Morales and E. Vassilieva, Bijective Enumeration of Bicolored Maps of Given Vertex Degree Distribution, *DMTCS Proceedings (FPSAC)*, AK:661–672, 2009.
- [16] D. Poulalhon and G. Schaeffer. Factorizations of large cycles in the symmetric group, *Discrete Math.*, 254:433–458, 2000.
- [17] G. Schaeffer and E. Vassilieva. A bijective proof of Jackson’s formula for the number of factorizations of a cycle, *J. Comb. Theory Ser. A*, 115(6):903–924, 2008.
- [18] E. Vassilieva Bijective Enumeration of 3-Factorizations of an N -Cycle *Annals of Combinatorics*, 16(2):367–387, 2012.
- [19] R. P. Stanley. *Enumerative Combinatorics*, volume 2. Cambridge University Press, Cambridge, 1999.