

# On arc-transitive metacyclic covers of graphs with order twice a prime

Zhaohong Huang\*

School of Mathematics and Statistics Science  
Ludong University  
Yantai, P. R. China  
1328897542@qq.com

Jiangmin Pan<sup>†</sup>

School of Statistics and Mathematics  
Yunnan University of Finance and Economics  
Kunming, P. R. China  
jmpan@ynu.edu.cn

Submitted: July, 8, 2017; Accepted: June 28, 2018; Published: TBD

© The authors. Released under the CC BY-ND license (International 4.0).

## Abstract

Quite a lot of attention has been paid recently to the characterization and construction of edge- or arc-transitive abelian (mostly cyclic or elementary abelian) covers of symmetric graphs, but there are rare results for nonabelian covers since the voltage graph techniques are generally not easy to be used in this case. In this paper, we will classify certain metacyclic arc-transitive covers of all non-complete symmetric graphs with prime valency and twice a prime order  $2p$  (involving the complete bipartite graph  $K_{p,p}$ , the Petersen graph, the Heawood graph, the Hadamard design on 22 points and an infinite family of prime-valent arc-regular graphs of dihedral groups). A few previous results are extended.

**Mathematics Subject Classifications:** 20B15, 20B30, 05C25

## 1 Introduction

Throughout the paper, by a graph, we mean a connected, simple and undirected graph with valency at least three.

---

\*Supported by Training foundation of Shandong Province (ZR2018PA005).

<sup>†</sup>Corresponding author; supported by National Natural Science Foundation of China (11461007, 11231008)

For a graph  $\Gamma$  and an automorphism group  $X \leq \text{Aut}\Gamma$ ,  $\Gamma$  is called  $X$ -vertex-transitive,  $X$ -edge-transitive or  $X$ -arc-transitive, if  $X$  is transitive on its vertex set, edge set or arc set, respectively. For a positive integer  $s$ , an  $s$ -arc of  $\Gamma$  is a sequence  $\alpha_0, \alpha_1, \dots, \alpha_s$  of  $s+1$  vertices such that  $\alpha_{i-1}, \alpha_i$  are adjacent for  $1 \leq i \leq s$  and  $\alpha_{i-1} \neq \alpha_{i+1}$  for  $1 \leq i \leq s-1$ . Then  $\Gamma$  is called  $(X, s)$ -arc-transitive or  $(X, s)$ -arc-regular if  $X$  is transitive or regular on the set of  $s$ -arcs of  $\Gamma$ , respectively. In particular, if  $\text{Aut}\Gamma$  is regular on the set of  $s$ -arcs of  $\Gamma$ , then  $\Gamma$  is simply called  $s$ -arc-regular.

An essential method for studying edge- or arc-transitive graphs is taking normal quotient graphs. Let  $\Gamma$  be a graph with vertex set  $V\Gamma$ , and suppose that  $X \leq \text{Aut}\Gamma$  acts edge- or arc-transitively on  $\Gamma$  and has an intransitive normal subgroup  $N$ . Denote by  $V\Gamma_N$  the set of  $N$ -orbits on  $V\Gamma$ . Then the *normal quotient graph* of  $\Gamma$  induced by  $N$ , denoted by  $\Gamma_N$ , is defined with vertex set  $V\Gamma_N$  and two vertices  $B, C \in V\Gamma_N$  are adjacent if and only if some vertex in  $B$  is adjacent in  $\Gamma$  to some vertex in  $C$ . If further, for each adjacent  $B$  and  $C$ , the induced subgraph  $[B, C] \cong nK_2$  is a complete matching, where  $n = |B| = |C|$ , then  $\Gamma$  is called a *normal cover* (or *regular cover*, or *cover* for short) of  $\Gamma_N$ . In some cases, to emphasize the groups  $X$  and  $N$ ,  $\Gamma$  is called an  $X$ -edge-transitive or  $X$ -arc-transitive  $N$ -cover of  $\Gamma_N$  respectively. If  $N$  is a cyclic, abelian or nonabelian group, then  $\Gamma$  is called a *cyclic cover*, *abelian cover* or *nonabelian cover* of  $\Gamma_N$ , respectively. We remark that ‘the cover of graph’ can also be defined by using notions of fibre-preserving group and covering transformation group, refer to [9, 25, 26].

From the above definition, an important strategy for studying transitive graphs naturally arises, involving the following two steps. Step 1 would be concerned to obtain a characterization of ‘basic’ transitive graphs (that is, the graphs which are not covers of their normal quotient graphs). Step 2 would then consists in approaching all transitive covers of basic graphs.

Characterizing covers of graphs is thus often a key step for studying edge- or arc-transitive graphs. By using voltage graph techniques (which generally are powerful for finding cyclic and elementary abelian covers, refer to [21, 25, 26, 36]), a lot of classifications of transitive cyclic or elementary abelian covers of symmetric graphs have been obtained, for example, see [16, 26, 27, 28] and references therein for edge- or arc-transitive cyclic and elementary abelian covers of certain small graphs, and see [11, 12, 13, 39, 41] for 2-arc-transitive cyclic and certain elementary abelian covers of  $K_n$  (the complete graph) and  $K_{n,n} - nK_2$  (the bipartite complete graph with a 1-factor removed), see also [40] for 2-arc-transitive metacyclic covers of  $K_n$ . Quite recently, a new approach was founded by investigating the action of universal group of basic graph and then used to find all arc-transitive abelian covers of a few small cubic symmetric graphs (see [5, 6]). However, since the voltage graph techniques are generally difficult for treating nonabelian covers, the results of nonabelian covers are rare.

The main purpose of this paper is to determine all arc-transitive  $K$ -covers of graph  $\Sigma$ , where  $\Sigma$  is a non-complete graph with prime valency  $r$  and twice a prime order  $2p$ , and  $K \cong \mathbb{Z}_m : \mathbb{Z}_q$  is a split metacyclic but not cyclic group (the arc-transitive cyclic covers of  $\Sigma$  have been determined by [32]), with  $q$  a prime less than  $r$ . We notice that certain special cases have been investigated in the literature: see [14] for  $K$  cyclic and  $\Sigma = K_{3,3}$

( $p = 3$  and  $r = 3$ ); [15] for  $K$  cyclic and  $\Sigma = \text{O}_2$  the Petersen graph ( $p = 5$  and  $r = 3$ ); [37] for  $K$  cyclic and  $\Sigma$  the Heawood graph ( $p = 7$  and  $r = 3$ ); [30] for  $K$  cyclic and  $\Sigma = \text{K}_{p,p}$  the complete bipartite graph with  $p > 3$ ; and see [42] for  $K$  a dihedral group and  $r = 3$ .

The terminology and notation used in this paper are standard. For example, for a positive integer  $n$ , we denote by  $\mathbb{Z}_n$ ,  $\text{D}_n$  with  $n$  even,  $\text{A}_n$  and  $\text{S}_n$  the cyclic group and the dihedral group of order  $n$ , the alternating and the symmetric group of degree  $n$ , respectively. For two groups  $N$  and  $H$ , we denote by  $N \times H$  the direct product of  $N$  and  $H$ , by  $N.H$  an extension of  $N$  by  $H$ , and if such an extension is split, then we write  $N : H$  instead of  $N.H$ . Also, we denote by  $\text{F}_n$ ,  $\text{F}_n\text{A}$ ,  $\text{F}_n\text{B}$  etc. with  $n$  a positive integer the corresponding cubic graphs of order  $n$  in the Foster census, see [1] or [4].

The main result of this paper is the following theorem. For convenience, the graphs appearing in the theorem which are not explained above are introduced in Section 2.

**Theorem 1.** *Let  $\Gamma$  be an  $X$ -arc-transitive  $K$ -cover of a graph  $\Sigma$ , where  $X \leq \text{Aut}\Gamma$ ,  $\Sigma$  is a non-complete graph of odd prime valency  $r$  and order  $2p$  with  $p$  a prime, and  $K \cong \mathbb{Z}_m : \mathbb{Z}_q$  is a metacyclic but not cyclic group with  $q < r$  a prime. Then one of the following statements holds.*

- (1)  $\Sigma \cong \text{K}_{3,3}$ ,  $K \cong \mathbb{Z}_{2n} \times \mathbb{Z}_2$  with  $n$  odd, and  $\Gamma$  is characterized in [5, Theorem 5.1];
- (2)  $\Sigma \cong \text{CD}(2p, 3)$  with  $p \equiv 1 \pmod{3}$ , and either
  - (i)  $K \cong \mathbb{Z}_2^2$  and  $\Gamma \cong \text{CGD}(8p, 3)$  is 1-arc-regular. or
  - (ii)  $K \cong \mathbb{Z}_m \times \mathbb{Z}_2$ , and  $\Gamma \cong \text{CGD}(2, \frac{mp}{2}, \lambda)$ , or  $\text{CGD}(2p, \frac{m}{2p}, \lambda)$  with  $p \mid m$ , where  $m = 2 \cdot 3^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t} \geq 6$ ,  $s \leq 1$ ,  $0 \leq t$  and  $p_1, p_2, \dots, p_t$  are distinct primes such that  $3 \mid (p_i - 1)$  for  $i = 1, 2, \dots, t$ .
- (3)  $\Sigma \cong \text{O}_2$ , and the tuple  $(\Gamma, K)$  is listed in the following table.

Row	$\Gamma$	$K$	$s$ – arc – regular
1	F40	$\text{D}_4$	3 – arc – regular
2	F60	$\text{D}_6$	2 – arc – regular
3	F120B	$\text{D}_{12}$	2 – arc – regular

The structure of this paper as follows. We give some preliminary results and introduce examples that appear in Theorems 1 in Section 2, and prove some technical lemmas in Section 3. Then, we classify the mentioned covers of the complete bipartite graph  $\text{K}_{p,p}$  and graphs  $\text{CD}(2p, r)$  in Sections 3 and 4 respectively, and complete the proof of Theorem 1 in Sections 5.

## 2 Preliminaries and examples

### 2.1 Preliminary results

For a group  $G$  and its subgroup  $H$ , let  $C_G(H)$  and  $N_G(H)$  denote the centralizer and normalizer of  $H$  in  $G$ , respectively.

**Lemma 2.** ([22, Ch. I, Lemma 4.5]) *Let  $G$  be a group and  $H$  a subgroup of  $G$ . Then  $N_G(H)/C_G(H) \leq \text{Aut}(H)$ .*

A graph  $\Gamma$  is called a *Cayley graph* of a group  $G$  if there is a subset  $S \subseteq G \setminus \{1\}$ , with  $S = S^{-1} := \{g^{-1} \mid g \in S\}$ , such that  $V\Gamma = G$  and two vertices  $g$  and  $h$  are adjacent if and only if  $hg^{-1} \in S$ . This Cayley graph is denoted by  $\text{Cay}(G, S)$ . It is well known that a graph  $\Sigma$  is isomorphic to a Cayley graph of a group  $R$  if and only if  $\text{Aut}\Sigma$  contains a subgroup which is isomorphic to  $R$  and acts regularly on  $V\Sigma$ , see [2, Proposition 16.3]. If this regular subgroup is normal in  $X$  with  $X \leq \text{Aut}\Sigma$ , then  $\Sigma$  is called an  *$X$ -normal Cayley graph*. In particular, if this regular subgroup is normal in  $\text{Aut}\Sigma$ , then  $\Sigma$  is called a *normal Cayley graph*.

Let  $\Gamma = \text{Cay}(G, S)$ . Let

$$\hat{G} = \{\hat{g} \mid \hat{g} : x \mapsto xg, \text{ for all } g, x \in G\},$$

$$\text{Aut}(G, S) = \{\sigma \in \text{Aut}(G) \mid S^\sigma = S\}.$$

Then both  $\hat{G}$  and  $\text{Aut}(G, S)$  are subgroups of  $\text{Aut}\Gamma$ . Further, the following nice property holds.

**Lemma 3.** ([19, Lemma 2.1]) *Let  $\Gamma = \text{Cay}(G, S)$  be a Cayley graph. Then the normalizer  $N_{\text{Aut}\Gamma}(\hat{G}) = \hat{G} : \text{Aut}(G, S)$ .*

We remark that  $\hat{G}$  is the right regular representation of  $G$ , and so isomorphic to  $G$ . For convenience, we will often write  $\hat{G}$  as  $G$ .

Let  $H$  be a group acting transitively on a set  $\Omega$ . Then  $H$  is called *primitive* on  $\Omega$  if  $H$  preserves no nontrivial partition of  $\Omega$ . A graph  $\Gamma$  is called  *$X$ -locally-primitive* with  $X \leq \text{Aut}\Gamma$ , if the vertex stabilizer  $X_\alpha := \{x \in X \mid \alpha^x = \alpha\}$  acts primitively on the neighbor set  $\Gamma(\alpha)$  for each vertex  $\alpha$ . Clearly, an edge-transitive graph with odd prime valency is locally-primitive. By Lemma 3, one easily has the following assertion.

**Lemma 4.** *Let  $\Gamma = \text{Cay}(G, S)$  be an  $X$ -normal locally-primitive Cayley graph of a group  $G$ , where  $G \triangleleft X \leq \text{Aut}\Gamma$ . Then  $X_1 \leq \text{Aut}(G, S)$  and elements in  $S$  are involutions, where  $1$  denotes the vertex of  $\Gamma$  corresponding to the identity element of  $G$ . In particular, if  $G$  is abelian, then  $G$  is an elementary abelian 2-group.*

The following theorem provides a basic method for studying vertex-transitive locally-primitive graphs (see [18] for the vertex-intransitive case), where parts (1–3) were first proved by Praeger [33, Theorem 4.1] for 2-arc-transitive graphs and slightly generalized to locally-primitive graphs by [24, Lemma 2.5]; part (4) follows easily by part (1).

**Theorem 5.** Let  $\Gamma$  be an  $X$ -vertex-transitive locally-primitive graph, and let  $N \triangleleft X$  have at least three orbits on  $V\Gamma$ . Then the following statements hold.

- (1)  $N$  is semiregular on  $V\Gamma$ ,  $X/N \leq \text{Aut}\Gamma_N$ ,  $\Gamma_N$  is  $X/N$ -locally-primitive, and  $\Gamma$  is an  $X$ -locally-primitive  $N$ -cover of  $\Gamma_N$ .
- (2)  $\Gamma$  is  $(X, s)$ -arc-transitive if and only if  $\Gamma_N$  is  $(X/N, s)$ -arc-transitive, where  $1 \leq s \leq 5$  or  $s = 7$ .
- (3)  $X_\alpha \cong (X/N)_\delta$ , where  $\alpha \in V\Gamma$  and  $\delta \in V\Gamma_N$ .
- (4) If  $X$  has a normal subgroup  $M$  which is contained in  $N$ , then  $\Gamma_M$  is an  $X/M$ -locally-primitive  $N/M$ -cover of  $\Gamma_N$ .

Given a group  $G$  and two subgroups  $L, R$  such that  $L \cap R$  is core-free in  $G$  (that is,  $L \cap R$  contains no nontrivial normal subgroup of  $G$ ). Define a bipartite graph  $\Gamma = \text{Cos}(G, L, R)$ , called *bi-coset graph*, with vertex set  $[G : L] \cup [G : R]$  and

$$\{Lx, Ry\} \text{ is an edge in } \Gamma \iff yx^{-1} \in RL.$$

The following lemma gives some basic properties of the bi-coset graphs, refer to [18, Lemmas 3.5, 3.7].

**Lemma 6.** With notation above,  $\Gamma$  satisfies the following properties:

- (1)  $\Gamma$  is connected if and only if  $\langle L, R \rangle = G$ ;
- (2)  $G \leq \text{Aut}\Gamma$ , and  $\Gamma$  is  $G$ -edge-transitive and  $G$ -vertex-intransitive;
- (3)  $\Gamma(L) = \{Rx \mid x \in L\}$  and  $\Gamma(R) = \{Lx \mid x \in R\}$ ;
- (4)  $|\Gamma(\alpha)| = |L : L \cap R|$  and  $|\Gamma(\beta)| = |R : L \cap R|$ , where  $\alpha \in [G : L]$  and  $\beta \in [G : R]$ .

Conversely, each  $G$ -vertex-intransitive and  $G$ -edge-transitive graph is isomorphic to  $\text{Cos}(G, G_\alpha, G_\beta)$ , where  $\alpha$  and  $\beta$  are adjacent vertices.

A group  $G$  is called perfect if  $G = G'$ , where  $G'$  is the commutator subgroup of  $G$ ; and an extension  $G = N.H$  is called a *central extension* if  $N$  is contained in the centre  $Z(G)$  of  $G$ . If a group  $G$  is perfect and  $G/Z(G)$  is isomorphic to a simple group  $T$ , then  $G$  is called a *covering group* of  $T$ . Schur [35] showed that a simple (and more generally, perfect) group  $T$  possesses a universal covering group  $G$  with the property that every covering group of  $T$  is a homomorphic image of  $G$ ; in this case, the center  $Z(G)$  is called the Schur multiplier of  $T$ , refer to [20, P.43]. The Schur multipliers of all simple groups are known ([20, P. 302–303]). The following lemma is known.

**Lemma 7.** Let  $G = N.T$ , where  $N$  is cyclic or isomorphic to  $\mathbb{Z}_p^2$  with  $p$  a prime, and  $T$  is a nonabelian simple group. Then  $G = N.T$  is a central extension,  $G = NG'$ , and  $G' = M.T$  is a perfect group, where  $M$  is a subgroup of both  $N$  and the Schur Multiplier of  $T$ .

The next lemma gives an observation regarding the vertex stabilizer of permutation groups, that will be used later.

**Lemma 8.** ([32, Lemma 3.4]) *Suppose that  $G = NH$  is a permutation group on a set  $\Omega$ , where  $N \triangleleft G$  and  $H \leq G$ . Then  $G_\alpha \cong N_\alpha.o$  for  $\alpha \in \Omega$ , where  $o \leq H/(H \cap N)$ . In particular, if  $N$  is transitive on  $\Omega$ , then  $o \cong H/(H \cap N)$ .*

## 2.2 Examples

We now introduce examples appearing in Theorem 1.

The first family of examples arises from Cayley graphs of dihedral groups, stated in Example 9, where the first two letters CD of the notation  $\text{CD}(2p, r)$  stand for ‘Cayley graph of a dihedral group’.

**Example 9.** Let  $G = \langle a, b \mid a^p = b^2 = 1, ab = a^{-1} \rangle \cong D_{2p}$  be a dihedral group, with  $p$  an odd prime. Let  $r$  be an odd prime and  $k$  a solution of the congruence equation

$$x^{r-1} + x^{r-2} + \cdots + x + 1 \equiv 0 \pmod{p}.$$

Define

$$\text{CD}(2p, r) = \text{Cay}(G, \{b, ab, a^{k+1}b, \dots, a^{k^{r-2}+k^{r-3}+\cdots+1}b\}).$$

We give some remarks here.

- (a) The congruence equation above has a solution if and only if  $p = r$  or  $r \mid (p - 1)$ , see [17, Lemma 3.3].
- (b) The graph  $\text{CD}(2p, r)$  is a bipartite graph, and up to isomorphism, it is independent of the choice of  $k$  by [17, Corollary 3.2]. This is the reason why the notation used does not involve  $k$ .
- (c) The graph  $\text{CD}(2p, p) \cong K_{p,p}$ ,  $\text{CD}(14, 3)$  is the Heawood graph, and  $\text{CD}(22, 5)$  is the incidence graph of valency 5 (the another one is of valency 6) of the Hadamard design on 22 points.

A group  $G$  is called a *generalized dihedral group*, if  $G = H : \langle a \rangle$  for some abelian subgroup  $H$  and an involution  $a$  such that  $h^a = h^{-1}$  for each  $h \in H$ . This group is denoted by  $\text{Dih}(H)$ . Obviously,  $\text{Dih}(\mathbb{Z}_n) \cong D_{2n}$ . The next family of examples arises from Cayley graphs of generalized dihedral groups, which was first obtained in [42].

**Example 10.** Let  $m$  and  $k$  be positive integers. Let

$$\text{Dih}(\mathbb{Z}_{mk} \times \mathbb{Z}_m) = \langle a, b, c \mid a^2 = b^{mk} = c^m = 1, b^a = b^{-1}, c^a = c^{-1}, bc = cb \rangle$$

be a generalized dihedral group. Assume that  $\lambda = 0$  for  $k = 1$ , and  $\lambda^2 + \lambda + 1 \equiv 0 \pmod{k}$  for  $k > 1$ . Define

$$\text{CGD}(m, k, \lambda) = \text{Cay}(\text{Dih}(\mathbb{Z}_{mk} \times \mathbb{Z}_m), \{a, ab, ab^\lambda c\}).$$

The first three letters CGD of the graph  $\text{CGD}(m, k, \lambda)$  stand for ‘Cayley graph of a generalized dihedral group’. By the notice before [42, Theorem 3.1], up to isomorphism, the cubic graph  $\text{CGD}(2, p, \lambda)$  with order  $8p$  is independent of the choice of  $\lambda$ , we thus denote it simply by  $\text{CGD}(8p, 3)$ .

The next lemma is quoted from [3], which classifies arc-transitive prime-valent graphs of order twice a prime.

**Lemma 11.** *Suppose that  $\Sigma$  is a non-complete arc-transitive graph of valency  $r$  and order  $2p$ , where  $r$  and  $p$  are odd primes. Then  $r \leq p$  and one of the following holds.*

- (1)  $p = 5$ ,  $r = 3$ ,  $\Sigma \cong \text{O}_2$  and  $\text{Aut} \Sigma \cong \text{S}_5$ .
- (2)  $\Sigma \cong \text{K}_{p,p}$  and  $\text{Aut} \Sigma \cong \text{S}_p \wr \text{S}_2$ .
- (3)  $\Sigma \cong \text{CD}(2p, r)$  with  $r \mid (p - 1)$ , and one of the following is true.
  - (i)  $(p, r) = (7, 3)$  and  $\text{Aut} \Sigma \cong \text{PGL}(2, 7)$ ;
  - (ii)  $(p, r) = (11, 5)$  and  $\text{Aut} \Sigma \cong \text{PGL}(2, 11)$ ;
  - (iii)  $(p, r) \neq (7, 3)$  and  $(11, 5)$ , and  $\text{Aut} \Sigma \cong \text{D}_{2p} : \mathbb{Z}_r$ .

A classification of arc-transitive cyclic covers of prime-valent graphs with order twice a prime is obtained by [32]. By [32, Theorem 1.1] (together with [30, Theorem 1.1] for a detailed description in one case), we have the following result.

**Theorem 12.** *Let  $\Gamma$  be an arc-transitive  $\mathbb{Z}_n$ -cover of a graph  $\Sigma$ , with odd prime valency  $r$  and twice a prime order  $2p$ . Then one of the following statements holds.*

- (1)  $\Sigma \cong \text{K}_4$ ,  $n = 2$  and  $\Gamma \cong \text{K}_{4,4} - 4\text{K}_2$ , or  $n = 4$  and  $\Gamma \cong \text{P}(8, 3)$  is the generalized Petersen graph.
- (2)  $\Sigma \cong \text{O}_2$ ,  $n = 2$ , and  $\Gamma$  is the Dodecahedron graph or the standard double cover of  $\text{O}_2$ .
- (3)  $\Sigma \cong \text{K}_{2p}$  with  $p \geq 3$  and  $2p - 1 = r$ ,  $n = 2$  and  $\Gamma \cong \text{K}_{2p, 2p} - 2p\text{K}_2$ .
- (4)  $\Sigma \cong \text{K}_{p,p}$  and  $n = p^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t}$ , where  $s \leq 1$ ,  $0 \leq t$  and  $p_1, p_2, \dots, p_t$  are distinct primes such that  $p \mid (p_i - 1)$  for  $i = 1, 2, \dots, t$ .
- (5)  $\Sigma \cong \text{CD}(2p, r)$  with  $r \mid (p - 1)$ , and  $n = r^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t}$ , where  $s \leq 1$ ,  $0 \leq t$  and  $p_1, p_2, \dots, p_t$  are distinct primes such that  $r \mid (p_i - 1)$  for  $i = 1, 2, \dots, t$ .

### 3 Technical lemmas

In this section, we prove some technical lemmas for the subsequent discussion.

For a group  $G$ , the socle of  $G$ , denoted by  $\text{soc}(G)$ , is the product of all minimal normal subgroups of  $G$ . Obviously,  $\text{soc}(G)$  is a characteristic subgroup of  $G$ .

**Lemma 13.** Suppose  $G \cong (\mathbb{Z}_{p^s} \times \mathbb{Z}_p) : \mathbb{Z}_3$ , where  $p \equiv 1 \pmod{3}$  is a prime and  $s \geq 1$ . Then  $G$  has a normal cyclic subgroup of order  $p^s$ .

*Proof.* By assumption, we may write  $G = (\langle a \rangle \times \langle b \rangle) : \langle c \rangle$ , where  $o(a) = p^s$ ,  $o(b) = p$  and  $o(c) = 3$ .

If  $s = 1$ , then  $G \cong \mathbb{Z}_p^2 : \mathbb{Z}_3$ . Noting that  $\mathbb{Z}_p^2$  has exactly  $p + 1$  subgroups isomorphic to  $\mathbb{Z}_p$ . Since  $3 \mid (p - 1)$ ,  $p + 1 \equiv 2 \pmod{3}$ , hence the action of  $\langle c \rangle \cong \mathbb{Z}_3$  on these  $p + 1$  subgroups is not fixed-point free. So at least one such subgroup must be invariant under this action, namely  $\langle c \rangle$  normalizes a cyclic subgroup  $\mathbb{Z}_p$ , which is clearly normal in  $G$ , the lemma is true in this case.

Assume  $s \geq 2$ . Let  $H = \langle a \rangle \times \langle b \rangle$ . Then  $\text{soc}(H) = \langle a^{p^{s-1}} \rangle \times \langle b \rangle \cong \mathbb{Z}_p^2$  is a characteristic subgroup of  $H$ , and so is normal in  $G$ . Hence  $\langle c \rangle$  normalizes  $\text{soc}(H)$ . As  $\langle c \rangle \cong \mathbb{Z}_3$  acts reducibly on  $\text{soc}(H)$  and  $(3, p) = 1$ , by Maschke's theorem (see [38, Theorem 1.4]),  $\langle c \rangle$  acts completely reducible on  $\text{soc}(H)$ , that is, we may write  $\text{soc}(H) = \langle b_1, b_2 \rangle$  such that  $c$  normalizes both  $\langle b_1 \rangle$  and  $\langle b_2 \rangle$ . Noting that at least one of  $b_1$  and  $b_2$  is not in  $\langle a \rangle$ , say  $b_1$ , then  $H = \langle a \rangle \times \langle b_1 \rangle$ . Hence, replacing  $b$  by  $b_1$  if necessary, we may assume  $b^c = b^k$  for some  $k$ . As  $o(b^k) = o(b) = p$ ,  $(k, p) = 1$ .

If  $a^c \in \langle a \rangle$ , the group  $\langle a \rangle$  is a required normal cyclic subgroup. Suppose  $a^c = a^i b^j$  with  $p \nmid j$ . If  $p \mid (k - i)$ , then  $b^i = b^k$  and  $(i, p) = 1$ , by a simple computation, we have  $a^{c^2} = (a^i b^j)^c = (a^i b^j)^i b^{kj} = a^{i^2} b^{2ij}$ , and  $a^{c^3} = (a^i b^j)^{i^2} b^{2ijk} = a^{i^3} b^{3i^2 j} \notin \langle a \rangle$  as  $p \nmid 3i^2 j$ , which is a contradiction because  $o(c) = 3$ . So  $p \nmid (k - i)$ , and hence the congruence equation

$$(k - i)x \equiv -j \pmod{p}$$

has a solution, say  $x = l$ . Now, as  $o(ab^l) = p^s$  and  $(ab^l)^c = a^i b^j b^{kl} = a^i b^{il} = (ab^l)^i$ ,  $\langle ab^l \rangle \cong \mathbb{Z}_{p^s}$  is normal in  $G$ .  $\square$

The next lemma presents some properties of the automorphism groups of metacyclic groups isomorphic to  $\mathbb{Z}_m : \mathbb{Z}_2$ .

**Lemma 14.** Let  $K = \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_m : \mathbb{Z}_2$  be a metacyclic group, where  $m = 2^d n > 1$  with  $d \geq 0$  and  $n$  odd. Then the following statements hold.

- (1) If  $m$  is odd, then  $K \cong \mathbb{Z}_s \times D_{2t}$ , where  $st = m$ , and  $(s, t) = 1$ .
- (2) If  $d = 1$ , then either  $K$  is abelian, or the center  $Z(K) \cong \mathbb{Z}_{2n_1}$  and  $K/Z(K) \cong D_{2n_2}$ , where  $n_1 n_2 = n$ ,  $n_2 > 1$  is odd and  $(n_1, n_2) = 1$ .
- (3) If  $d \geq 1$ , then  $\langle a^2 \rangle$  is a characteristic subgroup of  $K$  with index 4.
- (4) Suppose  $d \geq 2$  and  $K$  is normal in an overgroup  $X$ . Then  $X$  has a normal subgroup which is contained in  $K$  with index 2.

*Proof.* (1) Suppose that  $\langle a^e \rangle \cong \mathbb{Z}_{p^l}$  is a Sylow  $p$ -subgroup of  $\langle a \rangle$ , where  $p$  is an odd prime dividing  $m$  and  $l$  is a positive integer. Then  $\langle a^e \rangle \triangleleft K$ , so  $(a^e)^b = (a^e)^k$  for some integer  $k$ . Since  $b^2 = 1$ ,  $k^2 \equiv 1 \pmod{p^l}$ , and so  $p^l \mid (k - 1)(k + 1)$ . As  $(k - 1, k + 1)$  divides 2



and  $p^l \geq 3$ , we have  $k \equiv \pm 1 \pmod{p^l}$ . Hence either  $b$  centralizes  $\langle a^e \rangle$ , or  $\langle a^e, b \rangle \cong D_{2p^l}$ . Now, as  $\langle a \rangle \cong \mathbb{Z}_{p_1^{s_1}} \times \cdots \times \mathbb{Z}_{p_k^{s_k}}$  for some distinct odd primes  $p_1, \dots, p_k$ , and  $\langle b \rangle \cong \mathbb{Z}_2$  acts trivially or dihedrally on each Sylow  $p$ -subgroup of  $\langle a \rangle$ , collecting factors on which  $\langle b \rangle$  acts trivially gives the center  $\mathbb{Z}_s$  of  $K$  while collecting the remaining factors together with  $\langle b \rangle$  gives the dihedral group  $D_{2t}$ . Hence  $K \cong \mathbb{Z}_s \times D_{2t}$ .

(2) Since  $d = 1$ ,  $\langle a^n \rangle \in Z(K)$  and  $K = \langle a^n \rangle \times (\langle a^2 \rangle : \langle b \rangle)$ . By part (1),  $\langle a^2 \rangle : \langle b \rangle \cong \mathbb{Z}_{n_1} \times D_{2n_2}$ , where  $n_1 n_2 = n$  and  $(n_1, n_2) = 1$ . If  $n_2 = 1$ ,  $K$  is abelian; if  $n_2 > 1$ , then  $Z(K) \cong \mathbb{Z}_{2n_1}$  and  $K/Z(K) \cong D_{2n_2}$ .

(3) Suppose  $a^b = a^q$  with  $q$  an integer. Then  $q^2 \equiv 1 \pmod{m}$ , so  $q$  is odd as  $m$  is even. Let  $K^{(2)} = \langle g^2 \mid g \in K \rangle$ . Clearly,  $K^{(2)} \supseteq \langle a^2 \rangle$  is a characteristic subgroup of  $K$ . Since each element of  $K$  has a form  $a^i$  or  $a^i b$  for some integer  $i$ ,  $(a^i)^2 \in \langle a^2 \rangle$  and  $(a^i b)^2 = a^{(1+q)i} \in \langle a^2 \rangle$ , we conclude that  $\langle a^2 \rangle = K^{(2)}$  is a characteristic subgroup of  $K$  with index 4.

(4) By part (3),  $\langle a^2 \rangle$  is a characteristic subgroup of  $K$ , so is  $\langle a^4 \rangle$ , hence  $\langle a^4 \rangle \triangleleft X$  and  $K/\langle a^4 \rangle \triangleleft X/\langle a^4 \rangle$ . Since  $K/\langle a^4 \rangle \cong \mathbb{Z}_4 : \mathbb{Z}_2$  is a split extension, we have  $K/\langle a^4 \rangle \cong \mathbb{Z}_4 \times \mathbb{Z}_2$  or  $D_8$ . For the former case,  $K/\langle a^4 \rangle$  has a characteristic subgroup, say  $M/\langle a^4 \rangle$ , such that  $M/\langle a^4 \rangle \cong \mathbb{Z}_2^2$ , so  $M/\langle a^4 \rangle \triangleleft X/\langle a^4 \rangle$ , it follows that  $M$  is normal in  $X$  and is of index 2 in  $K$ . For the latter case,  $K/\langle a^4 \rangle$  has a characteristic subgroup, say  $N/\langle a^4 \rangle$ , such that  $N/\langle a^4 \rangle \cong \mathbb{Z}_4$ , it follows that  $N$  is normal in  $X$  and is of index 2 in  $K$ .  $\square$

The following lemma is regarding normal arc-transitive Cayley graphs.

**Lemma 15.** *Let  $G \cong \mathbb{Z}_2^2 \times D_{2p}$  with  $p \geq 5$  a prime. Then there is no normal arc-transitive Cayley graph of  $G$  with prime valency  $r \geq 5$ .*

*Proof.* Suppose that, on the contrary,  $\Gamma = \text{Cay}(G, S)$  is an  $X$ -normal arc-transitive Cayley graph of  $G$ , where  $G \triangleleft X \leq \text{Aut} \Gamma$  and  $S \subseteq G \setminus \{1\}$ . Set  $G = (\langle a \rangle \times \langle b \rangle) \times (\langle c \rangle : \langle d \rangle)$ , where  $\langle a \rangle \times \langle b \rangle \cong \mathbb{Z}_2^2$  and  $\langle c \rangle : \langle d \rangle \cong D_{2p}$ . By Lemma 4,  $S = s^{(\sigma)}$ , where  $s \in G$  is an involution and  $\sigma \in \text{Aut}(G)$  is of order  $r$ . Noting that  $\langle a, b \rangle \cong \mathbb{Z}_2^2$  is a characteristic subgroup of  $G$ , the restriction  $\sigma|_{\langle a, b \rangle}$  of  $\sigma$  on  $\langle a, b \rangle$  is an automorphism of  $\langle a, b \rangle$ . Clearly, the order of  $\sigma|_{\langle a, b \rangle}$  divides  $o(\sigma) = r$ , and so is 1 or  $r$ . As  $r \geq 5$  and  $\text{Aut}(\mathbb{Z}_2^2) \cong S_3$  has no element with order bigger than 3, we further conclude that  $\sigma|_{\langle a, b \rangle}$  is the identity automorphism of  $\langle a, b \rangle$ . If  $s \in \langle a, b \rangle$ , then  $\langle S \rangle = \langle s^{(\sigma)} \rangle \subseteq \langle a, b \rangle < G$ ,  $\Gamma$  is disconnected, which is a contradiction. Thus  $s \in G \setminus \langle a, b \rangle$ , and so  $s = a^i b^j c^k d$  for some integers  $i, j, k$ . Noting that there is an automorphism  $\tau$  which fixes  $a, b, c$  and maps  $s$  to  $d$ , and  $\text{Cay}(G, S) \cong \text{Cay}(G, S^\tau)$ , we may, up to isomorphism, assume  $s = d$ . Now, since  $\langle c \rangle \cong \mathbb{Z}_p$  is a characteristic subgroup of  $G$ ,  $\sigma$  has a form:

$$\sigma : a \rightarrow a, \quad b \rightarrow b, \quad c \rightarrow c^k, \quad d \rightarrow a^u b^v c^w d.$$

By computation,  $d = d^{\sigma^r} = a^{ru} b^{rv} c^{w(1+k+\cdots+k^{r-1})} d$ . So  $a^{ru} = b^{rv} = 1$ , implying  $a^u = b^v = 1$  as  $r$  is odd and  $a, b$  are involutions. It follows that  $\langle s^{(\sigma)} \rangle = \langle d^{(\sigma)} \rangle \subseteq \langle c, d \rangle < G$ , which contradicts the connectivity of  $\Gamma$ .  $\square$

## 4 Metacyclic covers of $K_{p,p}$

For convenience, we give the following Hypothesis 4.1 and Notations 4.2, which will be used throughout the rest of the paper.

**Hypothesis 4.1.** Let  $\Gamma$  be an  $X$ -arc-transitive  $K$ -cover of a graph  $\Sigma$ , where  $X \leq \text{Aut}\Gamma$ ,  $\Sigma$  is a non-complete graph of odd prime valency  $r$  and order  $2p$  with  $p$  a prime, and  $K := \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_m : \mathbb{Z}_q$  is a metacyclic but not cyclic group with  $q < r$  a prime. Then  $K \triangleleft X$  is semiregular on  $V\Gamma$ , and  $Y := X/K \leq \text{Aut}\Sigma$  acts transitively on the arc set of  $\Sigma$ .

**Notation 4.2.** If  $\Sigma$  is a bipartite graph, let  $\Delta_1$  and  $\Delta_2$  be the biparts of  $\Sigma$ , let  $Y^+ = Y_{\Delta_1} = Y_{\Delta_2}$  be the stabilizer of  $Y$  on the biparts, and let  $X^+ = K.Y^+$  (the full preimages of  $Y^+$  under the natural homomorphism of  $X$  to  $X/K$ ). Then  $|X : X^+| = |Y : Y^+| = 2$  and  $Y = \langle Y^+, y \rangle$  for some  $y \in Y \setminus Y^+$ .

By Hypothesis 4.1, the tuple  $(\Sigma, \text{Aut}\Sigma)$  is listed in Lemma 11. We will prove Theorem 1 by analysing all the candidates  $\Sigma$  in Lemma 11. Recall that, for a group  $G$ , a subgroup  $H$  is called the *Frattini subgroup* of  $G$  if  $H$  is the intersection of all maximal subgroups of  $G$ ; and for a prime  $t \mid |G|$ ,  $H$  is called a *Hall  $t'$ -subgroup* if  $(|H|, t) = 1$  and  $|G : H|$  is a  $t$ -power. It is well known that the Frattini subgroup and normal Hall  $t'$ -subgroups are characteristic subgroups. As usual, we use  $G_t$  and  $G_{t'}$  to denote a Sylow  $t$ -subgroup and a Hall  $t'$ -subgroup of  $G$ , respectively.

First, we prove an induction lemma.

**Lemma 16.** *With notation above, the following statements hold.*

- (1) *If  $q \nmid m$ , then  $q = 2$  and  $\Sigma \cong \text{O}_2$ .*
- (2) *If  $q \mid m$ , then there exists a graph  $\Omega$  being an arc-transitive  $\mathbb{Z}_q^2$ -cover of  $\Sigma$ .*

*Proof.* If  $q \nmid m$ , then  $\langle a \rangle$  is a characteristic subgroup of  $K$ , so is normal in  $X$ . Since  $K/\langle a \rangle \cong \mathbb{Z}_q$ , by Theorem 5(4), the normal quotient graph  $\Gamma_{\langle a \rangle}$  is an  $X/\langle a \rangle$ -arc-transitive  $\mathbb{Z}_q$ -cover of  $\Sigma$ . Then, as  $q < r = \text{val}(\Sigma)$  and  $\Gamma$  is not a complete graph, by Theorem 12, we have  $q = 2$  and  $\Sigma \cong \text{O}_2$ , as in part (1) of the lemma.

Assume now  $q \mid m$ . Then  $m = q^d n$  with  $d \geq 1$  and  $(q, n) = 1$ . Clearly,  $\langle a^{q^d} \rangle \cong \mathbb{Z}_n$  is a normal Hall  $q'$ -subgroup of  $K$ , so is characteristic in  $K$  and normal in  $X$ . Let  $\Gamma_1$  denote the normal quotient graph  $\Gamma_{\langle a^{q^d} \rangle}$ . By Theorem 5,  $\Gamma_1$  is an  $X/\langle a^{q^d} \rangle$ -arc-transitive  $K/\langle a^{q^d} \rangle$ -cover of  $\Sigma$ . Let  $\Phi$  be the Frattini subgroup of  $K/\langle a^{q^d} \rangle$ . Then  $\Phi$  is characteristic in  $K/\langle a^{q^d} \rangle$  and normal in  $X/\langle a^{q^d} \rangle$ . Since  $K/\langle a^{q^d} \rangle \cong \mathbb{Z}_{q^d} : \mathbb{Z}_q$ , by [34, 5.3.2],  $(K/\langle a^{q^d} \rangle)/\Phi \cong \mathbb{Z}_q^2$ . It then follows from Theorem 5 that  $\Omega := (\Gamma_1)_\Phi$  is an arc-transitive  $\mathbb{Z}_q^2$ -cover of  $\Sigma$ .  $\square$

In this section, we will determine all arc-transitive  $K$ -covers of the complete bipartite graph  $K_{p,p}$ .

The following lemma excludes the case where  $p \geq 5$ .

**Lemma 17.** *Suppose  $\Sigma \cong K_{p,p}$  with  $p \geq 5$ . Then no graph  $\Gamma$  exists.*

*Proof.* By Lemma 16, it is sufficient to show that Lemma 17 is true for the case where  $K \cong \mathbb{Z}_q^2$ . Recall that  $q < r = \text{val}(\Sigma) = p$ .

Let  $K_1$  and  $K_2$  be the kernels of  $Y^+$  acting on  $\Delta_1$  and  $\Delta_2$ , respectively. For any  $\alpha \in \Delta_1$ , since  $y \notin Y^+$ , we have  $\alpha^{y^{-1}} \in \Delta_2$  and  $\alpha^{y^{-1}K_2y} = (\alpha^{y^{-1}})^{K_2y} = (\alpha^{y^{-1}})^y = \alpha$ , so  $K_2^y = y^{-1}K_2y \subseteq K_1$ . Similarly,  $K_1^y \subseteq K_2$ . Thus  $K_1 = K_2^y$  and  $K_2 = K_1^y$ . Since  $Y$  is arc-transitive on  $\Sigma$ ,  $Y_\alpha$  is transitive on the neighbor set  $\Sigma(\alpha) = \Delta_2$ . If  $Y^+$  acts faithfully on  $\Delta_1$ , then  $Y^+$  can be viewed as a transitive permutation group of degree  $p$  on  $\Delta_1$ , so  $Y^+ \leq S_p$ . It follows that  $Y_\alpha^+ \leq S_{p-1}$ , which contradicts that  $Y_\alpha^+ = Y_\alpha$  is transitive on  $\Delta_2$  of size  $p$ .

Hence,  $Y^+$  acts unfaithfully on both  $\Delta_1$  and  $\Delta_2$ , that is,  $K_1 \cong K_2 \neq 1$ . Clearly,  $K_1 \cap K_2 = 1$ , so  $K_1K_2 = K_1 \times K_2 \triangleleft Y^+$  and  $Y^+ \cong (K_1 \times K_2).P$ , where  $P \cong Y^+/(K_1 \times K_2)$ . It follows that  $K_i.P \cong Y^+/K_{3-i} = (Y^+)^{\Delta_{3-i}}$  is a transitive permutation group of degree  $p$ , where  $i = 1, 2$ . By the classification of transitive permutation groups of prime degree (see [8, P. 99]), we have that either  $K_i.P \leq \mathbb{Z}_p : \mathbb{Z}_{p-1}$  is an affine group or  $K_i.P$  is an almost simple 2-transitive group. Let  $T_i = \text{soc}(K_i.P)$  and let  $G = K.(T_1 \times T_2)$ .

If  $T_i$  is nonabelian, then  $T_i$  is 2-transitive on  $\Delta_i$ , so  $T_1 \times T_2$  is transitive on the set of all 4-cycles of  $\Sigma \cong K_{p,p}$ . Noting that  $G \cong \mathbb{Z}_q^2.(T_1 \times T_2)$  is a central extension by Lemma 7, then with quite similar discussion as in [10, Theorem 4.2] (or [30, Lemma 5.1]), we have that  $\Gamma$  is disconnected, which is a contradiction.

Assume now that  $T_i$  is abelian. Then  $T_1 \times T_2 \cong \mathbb{Z}_p^2$  is edge-transitive and vertex-intransitive on  $\Sigma$ , and so  $G = K.(T_1 \times T_2) \cong \mathbb{Z}_q^2 : \mathbb{Z}_p^2$  is edge-transitive and vertex-intransitive on  $\Gamma$ . By Lemma 6,  $\Gamma \cong \text{Cos}(G, G_\alpha, G_\beta)$  for some adjacent vertices  $\alpha$  and  $\beta$  of  $\Gamma$ . Clearly,  $G_\alpha \cong G_\beta \cong \mathbb{Z}_p$ . If  $p \geq q + 2$ , then  $|\text{Aut}(\mathbb{Z}_q^2)| = |\text{GL}(2, q)| = q(q-1)^2(q+1)$  is not a multiple of  $p$ , by Lemma 2, the centralizer  $C_G(K) = G$ , and so  $G \cong \mathbb{Z}_q^2 \times \mathbb{Z}_p^2$  is abelian. It follows that  $\langle G_\alpha, G_\beta \rangle \leq G_p < G$ , so  $\Gamma$  is disconnected by Lemma 6, which is a contradiction. Thus  $p \leq q + 1$  and hence  $q = p - 1$  as  $q < p$ . Now, both  $p$  and  $p - 1$  are primes, the only possibility is  $p = 3$  and  $q = 2$ , which is a contradiction to the assumption  $p \geq 5$ .  $\square$

We next consider the case  $\Sigma \cong K_{3,3}$ .

**Lemma 18.** *Suppose  $\Sigma \cong K_{3,3}$ . Then  $K \cong \mathbb{Z}_{2n} \times \mathbb{Z}_2$  with  $n$  odd, and  $\Gamma$  is characterized in [5, Theorem 5.1].*

*Proof.* By assumption,  $r = \text{val}(\Sigma) = 3$ , so  $q = 2$  and  $K = \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_m : \mathbb{Z}_2$ . Then Lemma 16(1) implies that  $m$  is even. Set  $m = 2^d n$  with  $d \geq 1$  and  $n$  odd.

If  $d \geq 2$ , since  $K \triangleleft X$ , by Lemma 14(4),  $X$  has a normal subgroup, say  $M$ , such that  $M$  is contained in  $K$  with index 2. Then, by Theorem 5,  $\Gamma_M$  is an arc-transitive  $\mathbb{Z}_2$ -cover of  $\Sigma$ , so  $|V\Gamma_M| = 12$ ; however, there is no cubic arc-transitive graph of order 12 by [4], which is a contradiction.

Therefore,  $d = 1$ . Assume  $K$  is nonabelian. Then  $\langle a^n \rangle \cong \mathbb{Z}_2$  is in the center of  $K$ ,  $\langle a^2 \rangle$  is a characteristic subgroup of  $K$  and as  $K$  is nonabelian,  $b$  is not in the center of  $K$ , we conclude that  $\langle a \rangle = \langle a^2 \rangle \times \langle a^n \rangle$  is normal in  $X$ . By Theorem 5,  $\Gamma_{\langle a \rangle}$  is an arc-transitive

$\mathbb{Z}_2$ -cover of  $\Sigma$ , in particular,  $|V\Gamma| = 12$ , which is a contradiction by [4]. Thus  $K$  is abelian,  $K \cong \mathbb{Z}_{2n} \times \mathbb{Z}_2$  with  $n$  odd, and  $\Gamma$  is characterized in [5, Theorem 5.1].  $\square$

## 5 Metacyclic covers of $\mathbf{CD}(2p, r)$

Let  $\Gamma, X, \Sigma, K$  and  $Y$  be as in Hypothesis 4.1, and as  $\mathbf{CD}(2p, r)$  is a bipartite graph, we use Notation 4.2.

In this section, we will determine all arc-transitive  $K$ -covers of the bipartite graphs  $\mathbf{CD}(2p, r)$  with  $r \mid (p - 1)$ . Noting that  $K = \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_m : \mathbb{Z}_q$  and  $q < r$  are primes.

**Lemma 19.** *Assume  $\Sigma \cong \mathbf{CD}(2p, r)$  with  $r \mid (p - 1)$  and  $Y \cong D_{2p} : \mathbb{Z}_r$ . Then  $q = 2$  and  $r = 3$ .*

*Proof.* Suppose that, on the contrary,  $q \geq 3$ . Since  $Y \cong D_{2p} : \mathbb{Z}_r$ ,  $\Sigma$  is a  $Y$ -normal arc-transitive Cayley graph of the dihedral group  $D_{2p}$ , so  $\Gamma$  is an  $X$ -normal arc-transitive Cayley graph of  $R := K.D_{2p}$ . Also, by Lemma 16(1), we have  $q \mid m$ .

We first consider the case where  $m = q$ . Then  $R \cong \mathbb{Z}_q^2 : D_{2p}$  and  $X = K.Y \cong R : \mathbb{Z}_r$ . Since  $K$  is abelian, the centralizer  $C := C_R(K) \supseteq K$ . If  $C = K$ , Lemma 2 implies  $D_{2p} \cong R/K = R/C \leq \text{Aut}(K) \cong \text{Aut}(\mathbb{Z}_q^2) \cong \text{GL}(2, q)$ , as  $3 \leq q < r < p$ ,  $|\text{GL}(2, q)| = q(q - 1)^2(q + 1)$  is not a multiple of  $p$ , which is a contradiction. On the other hand, if  $C = R$ , then  $R = K \times R_{q'}$ , so  $R_{q'} \cong D_{2p}$  is normal in  $X$  and has  $q^2$  orbits on  $V\Gamma$ , by Theorem 5,  $\Gamma_{R_{q'}}$  is arc-transitive of odd valency  $r$  and odd order  $q^2$ , which is also a contradiction. Hence  $K < C < R$ , and  $C/K \neq 1$  is a proper normal subgroup of  $R/K \cong D_{2p}$ , we further conclude that  $C = R_{2'} \cong \mathbb{Z}_q^2 \times \mathbb{Z}_p$  and  $\mathbb{Z}_p \cong R_p \triangleleft X$ . Now, Theorem 5 implies that  $\Gamma_{R_p}$  is an  $X/R_p$ -normal arc-transitive Cayley graph of the group  $R/R_p$ . Clearly,  $R/R_p \cong R_{p'} = K : \mathbb{Z}_2 \cong \mathbb{Z}_q^2 : \mathbb{Z}_2$ . As  $q \geq 3$ ,  $R_{p'}$  is nonabelian by Lemma 4. Suppose that  $c \in R_{p'}$  is an involution. If  $x^c = y \notin \langle x \rangle$  for some  $x \in K$ , as  $o(c) = 2$ ,  $y^c = x$ , so  $(xy)^c = yx = xy$  and  $(xy^{-1})^c = yx^{-1} = (xy^{-1})^{-1}$ , hence  $R_{p'} = \langle xy \rangle \times \langle xy^{-1}, c \rangle \cong \mathbb{Z}_q \times D_{2q}$ . Since  $\Gamma_{R_p}$  is an  $X/R_p$ -normal Cayley graph with valency  $r$  of  $R/R_p$ , we may assume  $\Gamma_{R_p} = \text{Cay}(R/R_p, S)$ , where  $S \subseteq R/R_p \setminus \{1\}$ . By Lemma 4, elements in  $S$  are involutions, and as  $R/R_p \cong R_{p'} \cong \mathbb{Z}_q \times D_{2q}$ , we have  $\langle S \rangle \leq D_{2q} < R/R_p$ ,  $\Gamma$  is disconnected, which is a contradiction. Therefore,  $x^c \in \langle x \rangle$  for each  $x \in K$ , and  $\langle x, c \rangle \cong \mathbb{Z}_{2q}$  or  $D_{2q}$  by Lemma 14(1). Since  $R_{p'}$  is nonabelian, and as proved above,  $R_{p'} \not\cong \mathbb{Z}_q \times D_{2q}$ , we further have that  $R_{p'} \cong \text{Dih}(\mathbb{Z}_q^2)$  is a generalized dihedral group. Now, as  $X/R_p \cong (R/R_p) : \mathbb{Z}_r \cong \mathbb{Z}_q^2 : \mathbb{Z}_r : \mathbb{Z}_2$ , and  $(X/R_p)_u \cong \mathbb{Z}_r$  with  $u \in V(\Gamma_{R_p})$  is not normal in  $X/R_p$ , by Lemma 2,  $\mathbb{Z}_q^2$  is self-centralized in  $X/R_p$  and  $\mathbb{Z}_r : \mathbb{Z}_2 \leq \text{Aut}(\mathbb{Z}_q^2) \cong \text{GL}(2, q)$ . Hence  $r$  divides  $|\text{GL}(2, q)| = q(q - 1)^2(q + 1)$ . However, as  $q < r$  are primes, the only possibility is  $q = 2$  and  $r = 3$ , which contradicts the assumption that  $q \geq 3$ .

For the general case, by Lemma 16, there is a graph  $\Omega$  which is an arc-transitive  $\mathbb{Z}_q^2$ -cover of  $\Sigma$ , then above discussion also leads to a contradiction.

Therefore,  $q = 2$  and  $R \cong \mathbb{Z}_2^2.D_{2p}$ . We now prove  $r = 3$ . Since  $r \geq 3$  divides  $p - 1$ ,  $p \geq 7$ , and as  $|\text{Aut}(\mathbb{Z}_2^2)| = |\text{S}_3| = 6$ , Lemma 2 implies  $R = (\mathbb{Z}_2^2 \times \mathbb{Z}_p).\mathbb{Z}_2$  and so  $R_p \triangleleft X$ . By Theorem 5,  $\Gamma_{R_p}$  is an  $X/R_p$ -normal arc-transitive Cayley graph of the

group  $R/R_p \cong R_2 \cong \mathbb{Z}_2^2$ . If  $R_2$  is nonabelian, then  $R_2 \cong Q_8$  is a quaternion group or isomorphic to  $D_8$ . The former case is not possible by Lemma 4 as  $Q_8$  has unique involution, and the latter case is not possible by [29, Lemma 3.1]. Thus  $R_2$  is abelian. By Lemma 4,  $R_2 \cong \mathbb{Z}_2^3$  and  $R \cong \mathbb{Z}_2^3 \times D_{2p}$ . Recall that  $\Gamma$  is an  $X$ -normal arc-transitive Cayley graph of  $R$ , by Lemma 15, we have  $r < 5$ , hence  $r = 3$ , as required.  $\square$

The next lemma excludes the case where  $\Sigma \cong \text{CD}(22, 5)$ .

**Lemma 20.** *Assume  $\Sigma \cong \text{CD}(22, 5)$ . Then no graph  $\Gamma$  exists.*

*Proof.* With Lemma 16, we only need to prove the lemma for the case  $K \cong \mathbb{Z}_q^2$ .

Since  $\Sigma \cong \text{CD}(22, 5)$ ,  $\text{Aut}\Sigma \cong \text{PGL}(2, 11)$  by Lemma 11. Since  $Y \leq \text{Aut}\Sigma$  acts arc-transitively on  $\Sigma$ , 110 divides  $|Y|$ , by Atlas [7],  $Y \cong D_{22} : \mathbb{Z}_5, \text{PSL}(2, 11)$  or  $\text{PGL}(2, 11)$ . Noting that  $\text{CD}(22, 5)$  is a bipartite graph,  $Y$  has a normal subgroup  $Y^+$  with index 2, so  $Y \not\cong \text{PSL}(2, 11)$ . If  $Y \cong D_{22} : \mathbb{Z}_5$ , by Lemma 19, we have  $\text{val}(\Sigma) = 3$ , yielding a contradiction.

Assume  $Y \cong \text{PGL}(2, 11)$ . Then  $Y^+ \cong \text{PSL}(2, 11)$  and  $Y_\delta = Y_\delta^+ \cong A_5$  as  $|V\Sigma| = 22$ , where  $\delta \in V\Sigma$ . By Lemma 7,  $X^+ = K.Y^+ \cong \mathbb{Z}_q^2.\text{PSL}(2, 11)$  is a central extension,  $X^+ = K(X^+)'$ , and as the Schur multiplier of  $\text{PSL}(2, 11)$  is  $\mathbb{Z}_2$  (see [20, P. 302]),  $(X^+)' \cong \text{PSL}(2, 11)$  or  $\text{SL}(2, 11)$ . For  $\alpha \in V\Gamma$ , by Theorem 5(3),  $(X^+)'_\alpha \triangleleft X_\alpha^+ = X_\alpha \cong Y_\delta \cong A_5$ , so  $(X^+)'_\alpha = 1$  or  $A_5$ . If  $(X^+)'_\alpha = 1$ , as  $X^+ = K(X^+)'$ , Lemma 8 implies  $A_5 \cong X_\alpha^+ \leq K/(K \cap (X^+)' )$ , which is a contradiction as  $K$  is abelian. Thus  $(X^+)'_\alpha \cong A_5$ , and hence  $(X^+)' \not\cong \text{SL}(2, 11)$  since  $\text{SL}(2, s)$  with  $s$  an odd prime power has a unique involution by [23, Lemma 7.4]. Therefore,  $(X^+)' \cong \text{PSL}(2, 11)$ ,  $X^+ = K \times (X^+)'$  and each orbit of  $(X^+)'$  in  $V\Gamma$  has length  $|\text{PSL}(2, 11) : A_5| = 11$ . Since  $|V\Gamma| = 22q^2$ ,  $(X^+)' \triangleleft X$  has  $2q^2 \geq 8$  orbits on  $V\Gamma$ . By Theorem 5,  $\Gamma_{(X^+)}'$  is an  $X^+/(X^+)'$ -arc-transitive graph, which is also a contradiction as  $X^+/(X^+)' \cong \mathbb{Z}_q^2$  is abelian.  $\square$

With Lemmas 19 and 20, we may now determine all arc-transitive  $K$ -covers of  $\text{CD}(2p, r)$  with  $r \mid (p - 1)$ .

**Lemma 21.** *Assume  $\Sigma \cong \text{CD}(2p, r)$  with  $r \mid (p - 1)$ . Then  $r = 3$  and  $\Gamma$  is as in part (2) of Theorem 1.*

*Proof.* By Lemma 20,  $(p, r) \neq (11, 5)$ . If  $(p, r) = (7, 3)$ , the  $q = 2$  as  $q < r$ ; if  $(p, r) \neq (11, 5)$  and  $(7, 3)$ , by Lemma 11,  $\text{Aut}\Sigma \cong D_{2p} : \mathbb{Z}_r$  is arc-regular on  $\Sigma$ , so  $Y = \text{Aut}\Sigma$  as  $Y$  is arc-transitive on  $\Sigma$ , by Lemma 19, we also have  $q = 2$  and  $r = 3$ . Now, by Lemma 16,  $m$  is even. Suppose  $m = 2^d n$  with  $(2, n) = 1$  and  $d \geq 1$ .

Assume  $d \geq 2$ . By Lemma 14(4),  $X$  has a normal subgroup, say  $M$ , such that  $M$  is contained in  $K$  with index 2. It then follows from Theorem 5 that  $\Gamma_M$  is an arc-transitive  $\mathbb{Z}_2$ -cover of  $\text{CD}(2p, 3)$ , which is impossible by Theorem 12(5).

Hence,  $d = 1$  and  $K = \langle a \rangle : \langle b \rangle \cong \mathbb{Z}_{2n} : \mathbb{Z}_2$ . If  $n = 1$ , by [42, Theorem 3.1],  $\Gamma \cong \text{CGD}(8p, 3)$  (as  $CD_{8p}$  there), part (2)(i) of Theorem 1 holds. If  $K$  is nonabelian, by Lemma 14(2), the center  $Z(K) \cong \mathbb{Z}_{2n_1}$  and  $K/Z(K) \cong D_{2n_2}$ , where  $n_1 n_2 = n$ ,  $n_2 > 1$  and  $(n_1, n_2) = 1$ . Then, by Theorem 5,  $\Gamma_{Z(K)}$  is an arc-transitive  $D_{2n_2}$ -cover of  $\Sigma$ , as  $n_2$

is odd, which is not possible by [42, Theorem 3.1]. Hence  $K \cong \mathbb{Z}_{2n} \times \mathbb{Z}_2$  is abelian. Since  $\langle a^2 \rangle \triangleleft X$  and  $K/\langle a^2 \rangle \cong \mathbb{Z}_2^2$ , by Theorem 5,  $\Gamma_{\langle a^2 \rangle}$  is an  $X/\langle a^2 \rangle$ -arc-transitive  $\mathbb{Z}_2^2$ -cover of  $\Sigma \cong \text{CD}(2p, 3)$ , then [42, Theorem 3.1] implies  $\Gamma_{\langle a^2 \rangle} \cong \text{CGD}(8p, 3)$  is a 1-arc-regular normal Cayley graph of the generalized dihedral group  $\text{Dih}(\mathbb{Z}_{2p} \times \mathbb{Z}_2)$ , so  $X/\langle a^2 \rangle \cong \text{Dih}(\mathbb{Z}_{2p} \times \mathbb{Z}_2) : \mathbb{Z}_3$ . Noting that  $\langle a^2 \rangle \cdot \mathbb{Z}_2^2 \cong K \triangleleft X$ , we further conclude  $X \cong (\mathbb{Z}_{2n} \times \mathbb{Z}_2) \cdot \text{D}_{2p} \cdot \mathbb{Z}_3$ . Further, as  $K_2 \cong \mathbb{Z}_2^2$  is normal in  $X$  and  $K/K_2 \cong \mathbb{Z}_n$ , by Theorem 5,  $\Gamma_{K_2}$  is an arc-transitive  $\mathbb{Z}_n$ -cover of  $\text{CD}(2p, 3)$ . It then follows from Theorem 12(5) that  $n = 3^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t}$ , where  $s \leq 1$ ,  $0 \leq t$ , and  $p_1, p_2, \dots, p_t$  are distinct primes such that  $3 \mid p_i - 1$  for  $i = 1, 2, \dots, t$ .

Recall that  $X \cong (\mathbb{Z}_{2n} \times \mathbb{Z}_2) \cdot \text{D}_{2p} \cdot \mathbb{Z}_3$ . Let  $R \triangleleft X$  such that  $R \cong K \cdot \text{D}_{2p}$ , and let  $C = C_R(K_{2'})$ . Then  $K \subseteq C \triangleleft R$ . If  $C = K$ , Lemma 2 implies  $\text{D}_{2p} \cong Y/K = Y/C \leq \text{Aut}(K_{2'}) \cong \mathbb{Z}_n^*$ , yielding a contradiction. Thus  $C > K$  and  $1 \neq C/K \triangleleft Y/K \cong \text{D}_{2p}$ , implying  $C \supseteq K \cdot \mathbb{Z}_p$ . Let  $L \subseteq C$  such that  $L/K \cong \mathbb{Z}_p$ . Then  $L$  is abelian and  $L \cong \mathbb{Z}_{2np} \times \mathbb{Z}_2$ , or  $\mathbb{Z}_{2n} \times \mathbb{Z}_{2p}$  with  $p \mid n$ . Clearly,  $L \subseteq K \cdot \text{D}_{2p}$  is semiregular and has exactly two orbits on  $V\Gamma$ , by [42, Lemma 3.1],  $\Gamma$  is a Cayley graph of the generalized dihedral group  $\text{Dih}(L)$ . Then, by [42, Proposition 2.6], we have  $\Gamma \cong \text{CGD}(2, np, \lambda)$ , or  $\text{CGD}(2p, n/p, \lambda)$  with  $p \mid n$ , as in part (2)(ii) of Theorem 1.  $\square$

## 6 Proof of Theorem 1

To complete the proof of Theorem 1, we first prove that the graphs  $\Gamma$  in part (2)(ii) of Theorem 1 are really examples.

**Lemma 22.** *Suppose that  $p \equiv 1 \pmod{3}$  is a prime and  $m = 2 \cdot 3^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t}$ , where  $s \leq 1$ ,  $0 \leq t$ , and  $p_1, p_2, \dots, p_t$  are distinct primes such that  $3 \mid (p_i - 1)$  for  $i = 1, 2, \dots, t$ . Then the graphs  $\text{CGD}(2, \frac{mp}{2}, \lambda)$  and  $\text{CGD}(2p, \frac{m}{2p}, \lambda)$  with  $p \mid m$  are arc-transitive  $(\mathbb{Z}_m \times \mathbb{Z}_2)$ -covers of  $\text{CD}(2p, 3)$ .*

*Proof.* (i) Let  $\Omega = \text{CGD}(2, \frac{mp}{2}, \lambda)$ . By Example 10,  $\Omega = \text{Cay}(G, \{a, b, ab^{-\lambda}c\})$ , where

$$G = \langle a, b, c \mid a^2 = b^{mp} = c^2 = 1, b^a = b^{-1}, c^a = c^{-1}, bc = cb \rangle,$$

and  $\lambda^2 + \lambda + 1 \equiv 0 \pmod{\frac{mp}{2}}$ . By [42, Proposition 2.6(4)],  $\Omega$  is an arc-regular normal Cayley graph of  $G$ , hence  $\text{Aut}(\Omega) \cong G : \mathbb{Z}_3$ .

Clearly, the center  $Z(G) = \langle b^{\frac{mp}{2}} \rangle \times \langle c \rangle \cong \mathbb{Z}_2^2$  and  $G^{(2p)} := \langle g^{2p} \mid g \in G \rangle = \langle b^{2p} \rangle \cong \mathbb{Z}_{\frac{m}{2}}$  are characteristic subgroups of  $G$ . Set  $M = Z(G)G^{(2p)}$ . Since  $(\frac{mp}{2}, 2p) = (3^s p_1^{r_1} p_2^{r_2} \cdots p_t^{r_t} p, 2p) = p$ , we have  $M = \langle b^{\frac{mp}{2}}, b^{2p} \rangle \times \langle c \rangle = \langle b^p \rangle \times \langle c \rangle \cong \mathbb{Z}_m \times \mathbb{Z}_2$  is a characteristic subgroup of  $G$ , and hence  $M \triangleleft \text{Aut}(\Omega)$ . By Theorem 5,  $\Omega$  is an arc-transitive  $(\mathbb{Z}_m \times \mathbb{Z}_2)$ -cover of  $\Gamma_M$ , and as  $|V(\Omega_M)| = 2p$  with  $p \equiv 1 \pmod{3}$  and  $\text{val}(\Omega_M) = 3$ , by Lemma 11,  $\Omega_M \cong \text{CD}(2p, 3)$ .

(ii) Set  $\Omega' = \text{CGD}(2p, \frac{m}{2p}, \lambda)$  with  $p \mid m$ . Then  $\Omega' = \text{Cay}(G, \{a, b, ab^{-\lambda}c\})$  by Example 10, where

$$G = \langle a, b, c \mid a^2 = b^m = c^{2p} = 1, b^a = b^{-1}, c^a = c^{-1}, bc = cb \rangle,$$

and  $\lambda = 0$  if  $m = 2p$ , and  $\lambda^2 + \lambda + 1 \equiv 0 \pmod{\frac{m}{2p}}$  if  $\frac{m}{2p} > 1$ . By [42, Proposition 2.6],  $\Omega'$  is a normal Cayley graph of  $G$ , and  $\Omega'$  is 2-arc-regular if  $\frac{m}{2p} = 1$  or 3, or is 1-arc-regular if  $\frac{m}{2p} > 3$ .

Hence  $\text{Aut}(\Omega') \cong G : \mathbb{Z}_3$  or  $G : S_3$ , it follows that there exists an automorphism group  $X \leq \text{Aut}(\Omega')$  such that  $X \cong G : \mathbb{Z}_3$  acts arc-regularly on  $\Omega'$ . Let  $\langle f \rangle = X_\alpha \cong \mathbb{Z}_3$  with  $\alpha \in V\Omega'$ . Noting that  $Z(G) = \langle b^{\frac{m}{2}} \rangle \times \langle c^p \rangle \cong \mathbb{Z}_2^2$  and  $G_{2'} = \langle b^2 \rangle \times \langle c^2 \rangle \cong \mathbb{Z}_{\frac{m}{2}} \times \mathbb{Z}_p$  are characteristic subgroups of  $G$ , we have  $\langle b, c \rangle = \langle b^{\frac{m}{2}}, b^2 \rangle \times \langle c^p, c^2 \rangle = Z(G)G_{2'}$  is characteristic in  $G$  and so normal in  $X$ . Let  $H = \langle b, c, f \rangle$ . Then  $H = \langle b, c \rangle : \langle f \rangle \cong (\mathbb{Z}_m \times \mathbb{Z}_{2p}) : \mathbb{Z}_3$  and  $X = \langle H \rangle : \langle a \rangle$ .

**Claim.**  $X$  has a normal subgroup  $N \subseteq G$  such that  $N \cong \mathbb{Z}_m \times \mathbb{Z}_2$ .

Since  $p \equiv 1 \pmod{3}$  and  $p \mid m$ , we may suppose  $p = p_1$  and  $s_1 \geq 1$ . Now,  $\langle b, c \rangle_{p'} \cong \mathbb{Z}_{\frac{m}{p^{s_1}}} \times \mathbb{Z}_2$  is normal in  $H$ , with order  $\frac{2m}{p^{s_1}}$ . Consider the group  $R := (\langle b^{\frac{m}{p^{s_1}}} \rangle \times \langle c^2 \rangle) : \langle f \rangle$ . Then  $R \cong (\mathbb{Z}_{p^{s_1}} \times \mathbb{Z}_p) : \mathbb{Z}_3$ . As  $3 \mid (p-1)$ , by Lemma 13,  $R$  has a normal cyclic subgroup  $M \subseteq \langle b, c \rangle$  with order  $p^{s_1}$ . Noting that  $M$  is normal in  $H$ , then  $N := \langle b, c \rangle_{p'} M \cong \mathbb{Z}_m \times \mathbb{Z}_2$  is contained in  $\langle b, c \rangle$  and normal in  $H$ , and as  $x^a = x^{-1}$  for each  $x \in \langle b, c \rangle$ ,  $N$  is normal in  $\langle H, a \rangle = X$ , the claim is true.

Now, by Theorem 5,  $\Omega'$  is an arc-transitive  $(\mathbb{Z}_m \times \mathbb{Z}_2)$ -cover of  $\Omega'_N$ . Since  $p \equiv 1 \pmod{3}$ , and  $\Omega'_N$  is of valency 3 and order  $2p$ , by Lemma 11,  $\Omega'_M \cong \text{CD}(2p, 3)$ .  $\square$

Now, we are ready to complete the proof of Theorem 1.

**Proof of Theorem 1.** By assumption,  $\Sigma$  is  $X/K$ -arc-transitive of order  $2p$ , and so  $\Sigma$  is listed in Lemma 11. To be precise,  $\Sigma \cong \text{O}_2, \text{K}_{p,p}$  or  $\text{CD}(2p, r)$  with  $r \mid (p-1)$ .

Assume  $\Gamma \cong \text{O}_2$ . All edge-transitive metacyclic covers of  $\text{O}_2$  are determined in [31], which are arc-transitive and exactly consist of seven graphs. Noting that  $K$  is not a cyclic group and a quaternion group, by [31, Theorem 1.1], the tuple  $(\Gamma, K)$  satisfies part (3) of Theorem 1.

Assume  $\Sigma \cong \text{K}_{p,p}$ . By Lemma 17, we have  $p < 5$ , then by Lemma 18, we have  $K \cong \mathbb{Z}_{2n} \times \mathbb{Z}_2$  with  $n$  odd, and  $\Gamma$  is characterized in [5, Theorem 5.1], as in part (1) of Theorem 1.

Assume  $\Sigma \cong \text{CD}(2p, r)$  with  $r \mid (p-1)$ . By Lemma 21, we have  $r = 3$  and part (2) of Theorem 1 holds.

Finally, by Lemma 22, the graphs  $\Gamma$  in part (2)(ii) of Theorem 1 are arc-transitive  $\mathbb{Z}_m \times \mathbb{Z}_2$ -covers of  $\text{CD}(2p, r)$ , and the graph  $\text{CGD}(8p, 3)$  in part (2)(i) is arc-transitive  $\text{D}_4$ -cover of  $\text{CD}(2p, 3)$  by [42, Theorem 3.1], thus the graphs  $\Gamma$  in Theorem 1 are really examples.

This completes the proof of Theorem 1.  $\square$

## Acknowledgements

The authors would like to thank the anonymous referee for the helpful comments.

## References

- [1] I. Z. Bouwer (ed.). The Foster Census. Charles Babbage Research Center, Winnipeg, 1988.
- [2] N. Biggs. Algebraic Graph Theory, Cambridge University Press, 2nd Ed, New York, 1992.
- [3] Y. Cheng, and J. Oxley. On weakly symmetric graphs of order twice a prime. *J. Combin. Theory Ser. B*, 42:196–211, 1987.
- [4] M. Conder, and P. Dobcsányi. Trivalent symmetric graphs on up to 768 vertices. *J. Combin. Math. Comput.*, 40:41–63, 2002.
- [5] M. Conder, and J. C. Ma. Arc-transitive abelian regular covers of cubic graphs. *J. Algebra*, 387:215–242, 2013.
- [6] M. Conder, and J. C. Ma. Arc-transitive abelian regular covers of the Heawood graph. *J. Algebra*, 387:243–267, 2013.
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson. Atlas of Finite Groups, Clarendon Press, Oxford, 1985.
- [8] J. Dixon, and B. Mortimer. Permutation Groups, Springer-Verlag, New York, 1996.
- [9] D. Ž. Djoković. Automorphism of graphs and coverings. *J. Combin. Theory Ser. B*, 16:243–247, 1974.
- [10] S. F. Du, D. Malnič, and D. Marušič. Classification of 2-arc-transitive dihedrants. *J. Combin. Theory Ser. B*, 98:1349–1372, 2008.
- [11] S. F. Du, D. Marušič, and A. O. Waller. On 2-arc-transitive covers of complete graphs. *J. Combin. Theory Ser. B*, 74:276–290, 1998.
- [12] S. F. Du, J. H. Kwak, and M. Y. Xu. On 2-arc-transitive covers of complete graphs with covering transformation group  $\mathbb{Z}_p^3$ . *J. Combin. Theory Ser. B*, 93:73–93, 2005.
- [13] S. F. Du, and W. Q. Xu. 2-arc-transitive regular covers of  $K_{n,n} - nK_2$  having the covering transformation group  $\mathbb{Z}_p^3$ . *J. Aust. Math. Soc.*, 101:145–170, 2016.
- [14] Y. Q. Feng, and J. H. Kwak.  $s$ -Regular cyclic coverings of the complete bipartite graphs  $K_{3,3}$ . *J. Graph Theory*, 45:101–112, 2004.
- [15] Y. Q. Feng, and J. H. Kwak. Classifying cubic symmetric graphs of order  $10p$  or  $10p^2$ . *Sci. China*, 49:300–319, 2006.
- [16] Y. Q. Feng, and J. H. Kwak. Cubic symmetric graphs of order a small number times a prime or a prime square. *J. Combin. Theory Ser. B*, 97:627–646, 2007.
- [17] Y. Q. Feng, and Y. T. Li. One-regular graphs of square-free order of prime valency. *Europ. J. Combin.*, 32:261–275, 2011.
- [18] M. Giudici, C. H. Li, and C. E. Praeger. Analysing finite locally  $s$ -arc-transitive graphs. *Trans. Amer. Math. Soc.*, 356:291–317, 356.
- [19] C. D. Godsil. On the full automorphism group of a graph. *Combinatorica*, 1:243–256, 1981.



- [20] D. Gorenstein. Finite Simple Groups, Plenum Press, New York, 1982.
- [21] J. T. Gross, and T. W. Tucker. Generating all graph coverings by permutation voltage assignment. *Discrete Math.*, 18:273–283, 1977.
- [22] B. Huppert, *Finite Groups*, Springer-Verlag, Berlin, 1967.
- [23] I. M. Isaacs. Finite Group Theory, American Mathematics Society, 2008.
- [24] C. H. Li, and J. M. Pan. Finite 2-arc-transitive abelian Cayley graphs. *Europ. J. Combin.*, 29:148–158, 2007.
- [25] A. Malnič, R. Nedela, and M. Škovič. Lifting graph automorphisms by voltage assignments. *Europ. J. Combin.*, 21:927–947, 2000.
- [26] A. Malnič, D. Marušič, and P. Potočnik. Elementary abelian covers of graphs. *J. Algebraic Combin.*, 20:71–97, 2004.
- [27] A. Malnič, D. Marušič, and P. Potočnik. On cubic graphs admitting an edge-transitive solvable group. *J. Algebraic Combin.*, 20:99–113, 2004.
- [28] A. Malnič, and P. Potočnik. Invariant subspaces, duality, and covers of the Petersen graph. *Europ. J. Combin.*, 27:971–989, 2008.
- [29] J. M. Pan. Locally primitive Cayley graphs of dihedral groups. *Europ. J. Combin.*, 36:39–52, 2014.
- [30] J. M. Pan, and Z. H. Huang. Arc-transitive regular cyclic covers of the complete bipartite graphs  $K_{p,p}$ . *J. Algebraic Combin.*, 42:619–633, 2015.
- [31] J. M. Pan, and Z. H. Huang. Edge-transitive regular metacyclic covers of the Petersen graph. *Graphs Combin.*, 32:777–784, 2016.
- [32] J. M. Pan, Z. H. Huang, and S. Y. Ding. Arc-transitive regular cyclic covers of symmetric graphs of order twice a prime. *Discrete Math.*, 340:811–816, 2017.
- [33] C. E. Praeger. An O’Nan-Scott theorem for finite quasiprimitive permutation groups and an application to 2-arc transitive graphs. *J. London Math. Soc.*, 47:227–239, 1992.
- [34] D. J. S. Robinson. A Course in the Theory of Groups, Springer-Verlag, New York, 1982.
- [35] I. Schur. Untersuchen über die Darstellung der endlichen Gruppen durch gebrochenen linearen Substitutionen. *Crelle J.*, 132:85–137, 1907.
- [36] M. Škovič. A contribution to the theory of voltage graphs. *Discrete Math.*, 61:281–292, 1986.
- [37] C. Q. Wang, and Y. Hao. Edge-transitive regular  $\mathbb{Z}_n$ -covers of the Heawood graph. *Discrete Math.*, 310:1752–1758, 2010.
- [38] P. Webb. Finite Group Representation for the Pure Mathematician, Springer-Verlag, Berlin, 1992.
- [39] W. Q. Xu, and S. F. Du. 2-arc-transitive cyclic covers of  $K_{n,n} - nK_2$ . *J. Algebraic Combin.*, 39:883–902, 2014.

- [40] W. Q. Xu, S. F. Du, J. H. Kwak, and M. Y. Xu. 2-arc-transitive metacyclic covers of complete graphs. *J. Combin. Theory Ser. B*, 111:54–74, 2015.
- [41] W. Q. Xu, Y. H. Zhu, and S. F. Du. 2-arc-transitive regular covers of  $K_{n,n} - nK_2$  with the covering transformation group  $\mathbb{Z}_p^2$ . *Ars. Math. Contemp.*, 10(2):269–280, 2016.
- [42] J. X. Zhou, and Y. Q. Feng. Edge-transitive dihedral or cyclic covers of cubic symmetric graphs of order  $2p$ . *Combinatorica*, 34(1):115–128, 2014.