

# Small Ramsey Numbers

Stanisław P. Radziszowski  
Department of Computer Science  
Rochester Institute of Technology  
Rochester, NY 14623, spr@cs.rit.edu  
<http://www.cs.rit.edu/~spr>

version #0: August 1, 1994; revision #18: April 24, 2026  
<http://doi.org/10.37236/21>

**ABSTRACT:** Ramsey theory is often regarded as the study of how order emerges from randomness. It studies the conditions wherein a combinatorial object necessarily contains some smaller given objects. The role of Ramsey numbers is to quantify some of the general existential theorems in Ramsey theory. This dynamic survey covers results and problems related to classical Ramsey numbers where the edges of complete graphs and hypergraphs are colored, while the avoided graphs are arbitrary. Thus, classical Ramsey numbers, graph Ramsey numbers, multicolor Ramsey numbers and hypergraph Ramsey numbers are all covered.

We do not attempt a complete coverage of the asymptotic behavior of Ramsey numbers, but rather we concentrate on their specific values. Numerous variations of Ramsey numbers, such as anti-Ramsey numbers, bipartite Ramsey numbers, Gallai-Ramsey numbers, and many other variants listed in the closing section *Exclusions*, are not covered.

We present data which, to the best of our knowledge, includes all known nontrivial values and bounds for specific graph, multicolor and hypergraph Ramsey numbers. Many general results pertaining to the most studied cases are also presented. We give references to all cited bounds and values, as well as to previous overview articles, similar compilations, electronic resources, or books covering the area.

This compilation is a fast source of references not only to the best known results for specific parameters, but also to general results. This survey gives a bigger picture of the development of various subareas from a theoretical, practical and computational perspective. While it is not planned to serve as a complete source of definitions or theorems, these can be easily accessed via the references gathered here.

## **Keywords:**

Ramsey numbers, extremal graph theory, coloring of graphs and hypergraphs

## **MSC 2020, Mathematics Subject Classification:**

primary 05C55, 05D10; secondary 05C15, 05C35, 05C38

**Versions 1993-2026**

1993, February, preliminary technical report, RIT-TR-93-009 [Ra2]

1994, August 1, the first version posted on the web at the *E-JC*

**Revisions at the *E-JC***

year, month	revision#	#pages	#references
1994, August 1	original	28 (13+15)	208
1994, November 7	#1	28 (13+15)	211
1995, August 28	#2	28 (13+15)	220
1996, March 25	#3	29 (14+15)	226
1997, July 11	#4	28 (15+13)	252
1998, July 9	#5	30 (16+14)	274
1999, July 5	#6	35 (19+16)	301
2000, July 25	#7	36 (21+15)	308
2001, July 12	#8	38 (22+16)	327
2002, July 15	#9	42 (24+18)	357
2004, July 4	#10	48 (27+21)	403
2006, August 1	#11	60 (36+24)	452
2009, August 4	#12	72 (42+30)	526
2011, August 22	#13	84 (50+34)	597
2014, January 12	#14	94 (56+38)	661
2017, March 3	#15	104 (60+44)	765
2021, January 15	#16	116 (67+49)	856
2024, June 7	#17	133 (77+56)	965
2026, April 24	#18	149 (87+62)	1066

The total number of pages is broken into the number of pages in the survey itself and in the references. See Section 9.2 for more statistics about references. All revisions are available at <http://doi.org/10.37236/21> or <https://www.cs.rit.edu/~spr/EJJC/eline.html>.

<b>Table of Contents of Revision #18</b>	<b>3</b>
<b>1. Scope and Notation</b>	<b>5</b>
<b>2. Classical Two-Color Ramsey Numbers</b>	<b>6</b>
2.1 Values and bounds for $R(k, l)$ , $k \leq 10$ , $l \leq 15$	6
2.2 Bounds for $R(k, l)$ , higher parameters	9
2.3 General results on $R(k, l)$	12
2.4 Some pillars of asymptotics of $R(k, l)$	14
<b>3. Two Colors: <math>K_n - e</math>, <math>K_3</math>, <math>K_{m, n}</math></b>	<b>16</b>
3.1 Dropping one edge from complete graph	16
3.2 Triangle versus other graphs	19
3.3 Complete bipartite graphs	20
3.3.1 Numbers	20
3.3.2 General results	23
<b>4. Two Colors: Numbers Involving Cycles</b>	<b>25</b>
4.1 Cycles, cycles versus paths and stars	25
4.1.1 Cycles	25
4.1.2 Cycles versus paths	26
4.1.3 Cycles versus stars	27
4.2 Cycles versus complete graphs	27
4.3 Cycles versus wheels	29
4.4 Cycles versus books	31
4.5 Cycles versus other graphs	33
<b>5. General Graph Ramsey Numbers in Two Colors</b>	<b>34</b>
5.1 Paths	34
5.2 Wheels	34
5.3 Books	35
5.4 Trees and forests	38
5.5 Stars, stars versus other graphs	39
5.6 Paths versus other graphs	40
5.7 Fans, fans versus other graphs	41
5.8 Wheels versus other graphs	42
5.9 Books versus other graphs	44
5.10 Trees and forests versus other graphs	44
5.11 Unicyclic graphs	46
5.12 Cases for $n(G)$ , $n(H) \leq 5$	47
5.13 Miscellaneous cases	47
5.14 Multiple copies of graphs, disconnected graphs	48
5.15 Ramsey goodness of graphs	49
5.16 General results for special graphs	51
5.17 General results for sparse graphs	51
5.18 General results	53

<b>6. Multicolor Graph Ramsey Numbers</b>	<b>55</b>
6.1 Bounds for classical numbers	55
6.1.1 Diagonal cases	55
6.1.2 Off-diagonal cases	56
6.2 General results for complete graphs	58
6.3 Cycles	60
6.3.1 Three colors	60
6.3.2 More colors	62
6.3.3 Cycles versus other graphs	64
6.4 Paths, paths versus other graphs	65
6.4.1 Three color paths and path-cycle cases	65
6.4.2 More colors	67
6.5 Special cases	68
6.6 General results for special graphs	69
6.7 General results	71
<b>7. Hypergraph Ramsey Numbers</b>	<b>73</b>
7.1 Values and bounds for numbers	73
7.2 Cycles and paths	74
7.2.1 Two colors	74
7.2.2 More colors	76
7.3 Berge cycles and Berge hypergraphs	76
7.4 General results for 3-uniform hypergraphs	77
7.5 General results	78
<b>8. Cumulative Data, Books and Surveys</b>	<b>80</b>
8.1 Cumulative data for two colors	80
8.2 Cumulative data for three colors	81
8.3 Electronic resources	81
8.4 Books and surveys	82
<b>9. Concluding Remarks</b>	<b>85</b>
9.1 Exclusions	85
9.2 Journal paper counts	86
Acknowledgements	87
<b>References</b>	<b>88-149</b>

Note that this survey covers the results and problems related to the classical Ramsey numbers, where the edges of complete graphs and hypergraphs are colored, while the avoided graphs or hypergraphs can be arbitrary. Numerous variations of Ramsey numbers not covered in this compilation are listed in Section 9.1, *Exclusions*.

## 1. Scope and Notation

Ramsey Theory studies conditions when a combinatorial object contains necessarily some smaller given objects. The role of Ramsey numbers is to quantify some of the general existential theorems in Ramsey Theory. There is vast literature on Ramsey type problems starting in 1930 with the original paper of Ramsey [Ram]. Graham, Rothschild and Spencer in their book [GRS] present an exciting development of Ramsey Theory, for many more pointers to books and surveys see Section 8.4. The subject has grown amazingly, in particular with regard to asymptotic bounds for various types of Ramsey numbers, but the progress on evaluating the basic numbers themselves has been unsatisfactory for a long time. In the last few decades, however, considerable progress has been obtained in this area, often by employing computer algorithms. The few known exact values and several bounds for different numbers are scattered among many technical papers. This compilation is a fast source of references for the best results known for specific numbers, but also to general results. This survey gives a bigger picture of the development of various subareas from a theoretical, practical and computational perspective. While it is not planned to serve as a complete source of definitions or theorems, these can be easily accessed via the references gathered here.

Let  $G_1, G_2, \dots, G_m$  be graphs or  $s$ -uniform hypergraphs ( $s$  is the number of vertices in each edge).  $R(G_1, G_2, \dots, G_m; s)$  denotes the  $m$ -color **Ramsey number** for  $s$ -uniform graphs/hypergraphs, avoiding  $G_i$  in color  $i$  for  $1 \leq i \leq m$ . It is defined as the least integer  $n$  such that, in any coloring with  $m$  colors of the  $s$ -subsets of a set of  $n$  elements, for some  $i$  the  $s$ -subsets of color  $i$  contain a sub-(hyper)graph isomorphic to  $G_i$  (not necessarily induced). The value of  $R(G_1, G_2, \dots, G_m; s)$  is fixed under permutations of the first  $m$  arguments. If  $s=2$  (standard graphs) then  $s$  can be omitted. If  $G_i$  is a complete graph  $K_k$ , then we may write  $k$  instead of  $G_i$ , and if  $G_i = G$  for all  $i$  we may use the abbreviation  $R_m(G; s)$  or  $R_m(G)$ . For  $s=2$ ,  $K_k - e$  denotes a  $K_k$  without one edge, and for  $s=3$ ,  $K_k - t$  denotes a  $K_k$  without one triangle (hyperedge).

The graph  $nG$  is formed by  $n$  disjoint copies of  $G$ ,  $G \cup H$  stands for vertex disjoint union of graphs, and the **join**  $G+H$  is obtained by adding all of the edges between vertices of  $G$  and  $H$  to  $G \cup H$ .  $P_i$  is a **path** on  $i$  vertices,  $C_i$  is a **cycle** of length  $i$ , and  $W_i$  is a **wheel** with  $i-1$  spokes, i.e. a graph formed by some vertex  $x$ , connected to all vertices of the cycle  $C_{i-1}$  (thus  $W_i = K_1 + C_{i-1}$ ).  $K_{n,m}$  is a complete  $n$  by  $m$  bipartite graph, in particular  $K_{1,n}$  is a **star** graph. The **book** graph  $B_i = K_2 + \bar{K}_i = K_1 + K_{1,i}$  has  $i+2$  vertices, and can be seen as  $i$  triangular pages attached to a single edge. The **fan** graph  $F_n$  is defined by  $F_n = K_1 + nK_2$ . For a graph  $G$ ,  $n(G)$  and  $e(G)$  denote the number of vertices and edges, respectively, and  $\delta(G)$  and  $\Delta(G)$  minimum and maximum degree of  $G$ . Finally,  $\chi(G)$  denotes the chromatic number of  $G$ . In general, we follow the notation used by West [West].

Section 2 contains the data for the classical two-color Ramsey numbers  $R(k, l)$  for complete graphs, Section 3 for the much studied two-color cases of  $K_n - e$ ,  $K_3$ ,  $K_{m,n}$ , and Section 4 for numbers involving cycles. Section 5 lists other often studied two-color cases for general graphs. The multicolor and hypergraph cases are gathered in Sections 6 and 7, respectively. Section 8 gives pointers to books and other cumulative data. Finally, Section 9.1 lists most Ramsey numbers variants not covered in this survey, and Section 9.2 gives counts of papers in our references which appeared in different journals.

## 2. Classical Two-Color Ramsey Numbers

### 2.1. Values and bounds for $R(k, l)$ , $k \leq 10$ , $l \leq 15$

$k \setminus l$	3	4	5	6	7	8	9	10	11	12	13	14	15
3	6	9	14	18	23	28	36	40 41	47 50	53 59	61 68	67 77	74 87
4		18	25	36 41	49 61	59 84	73 115	92 149	102 191	128 238	139 291	148 349	159 417
5			43 46	59 87	80 143	101 216	133 316	149 442	183 633	203 848	233 1138	267 1461	275 1878
6				102 165	115 298	134 495	183 780	204 1171	262 1804	294 2566	347 3703		401 6911
7					205 540	219 1031	252 1713	292 2826	405 4553	417 6954	511 10578		22112
8						282 1870	329 3583	343 6090	457 10630		817 27485		873 63609
9							565 6588	581 12677		22325	38832	64864	
10								798 23556	45881	81123			1313

**Table Ia.** Known nontrivial values, lower bounds and upper bounds for two-color Ramsey numbers  $R(k, l) = R(k, l; 2)$ , for  $k \leq 10$  and  $k \leq l \leq 15$ . For better but unpublished upper bounds (2024) with  $k \geq 4$  see Table Ib.

$k \setminus l$	4	5	6	7	8	9	10	11	12	13	14	15
3	GG	GG	Kéry	Ka2 GrY	GR McZ	Ka2 GR	Ex5 Ang1	Ex20 GoeR1	Kol1 Les	NaRT GoeR1	Kol2 GoeR1	Kol2 GoeR1
4	GG	Ka1 MR4	Ex19 MR5	Ex3 Mac	ExT Mac	Ex16 Mac	HaKr1 Mac	ExT Spe4	SuLL Spe4	NaRT Spe4	NaRT Spe4	NaRT Spe4
5		Ex4 AnM4	Ex25 HZ1	CaET HZ1	HaKr1 Spe4	Kuz Mac	ExT Mac	Kuz HW+	Kuz HW+	Kuz HW+	Kuz HW+	2.3.h HW+
6			Ka2 Mac	ExT HZ1	ExT Mac	Kuz Mac	Kuz Mac	Tat HW+	Kuz HW+	Kuz HW+		2.3.i HW+
7				Math Mac	Tat HZ1	Kuz HZ2	Kuz Mac	XXER HW+	XSR2 HW+	XuXR HW+		HW+
8					BurR Mac	Kuz Ea1	Kuz HZ2	2.3.i HW+		XXER HW+		2.3.i HW+
9						Math ShZ1	XSR2 Ea1	HW+	HW+	HW+		
10							Math Shi2	HW+	HW+			2.3.i

**References for Table Ia.** All upper bounds for  $k \geq 4$ ,  $l \geq 6$  were improved in 2019 [AnM2] and 2023 [AnM3], see Table Ib. HW+ abbreviates HWSYZH, as enhanced in [Boza5], see 2.1.o.

$k \backslash l$	5	6	7	8	9	10	11	12	13	14	15
4	25	40	58	79	105	135	170	210	256	307	364
5	46	85	133	193	282	381	511	672	860	1081	1341
6		160	270	423	651	944	1346	1855	2499	3301	4305
7			492	832	1368	2119	3197	4665	6653	9260	12635
8				1518	2662	4402	7040	10836			
9					4956	8675	14631				
10						16064					

**Table Ib.** Upper bounds for  $R(k, l)$ ,  $l \geq k \geq 4$ ,  $l \geq 5$ . All of them were obtained by Angeltveit and McKay in 2019 [AnM2] and 2023-2024 [AnM3, AnM4], except  $R(4, 5)$  [MR4], and they improve over previously best known bounds reported in Table Ia.

We split the data into Table Ia with a separate table of references corresponding to it, and Table Ib of new upper bounds. In Table Ia, the known exact values appear as centered entries, lower bounds as top entries, and upper bounds as bottom entries. For some of the exact values two references are given when the lower and upper bound credits are different. In large computational projects from 2019 to 2023, Angeltveit and McKay ran independent computations [AnM2, AnM3], and obtained new upper bounds reported in Table Ib. These, by using the classical recursive upper bound 2.3.a, and the methods of [HW+, HYZ, LiShen], lead to improvements of other higher upper bounds listed in Table Ia but not reported in Table Ib.

- (a) The task equivalent to that of proving  $R(3, 3) \leq 6$  was the second problem in the Kürschák Mathematics Competitions in Hungary in 1947 [BaLiu]. It also was the second problem in Part I of the William Lowell Putnam Mathematical Competition held in March 1953 [Bush].
- (b) Greenwood and Gleason [GG] established the initial values  $R(3, 4) = 9$ ,  $R(3, 5) = 14$  and  $R(4, 4) = 18$  in 1955.
- (c) Kéry [Kéry] proved that  $R(3, 6) = 18$  in 1964, but only in 2007 an elementary and self-contained proof of this result appeared in English [Car].
- (d) All of the critical graphs for the numbers  $R(k, l)$  (graphs on  $R(k, l) - 1$  vertices without  $K_k$  and without  $K_l$  in the complement) are known for  $k=3$  and  $l=3, 4, 5$  [Kéry], 6 [Ka2], 7 [RaK2, McZ], 8 [BrGS] and 9 [GoeR1], and there are 1, 3, 1, 7, 191, 477142, and 1 of them, respectively. There are at least  $43 \times 10^6$   $(3, 10)$ -graphs on 39 vertices [GoeR2, Ang1]. All  $(3, k)$ -graphs, for  $k \leq 6$ , were enumerated in [RaK2], and all  $(4, 4)$ -graphs in [MR2]. There exists a unique critical graph for  $R(4, 4)$  [Ka2]. Until 2015, there

were 350904 known critical graphs for  $R(4, 5)$  [MR4], but the full set of such graphs was computed in 2016 [McK3], and there are 352366 of them.

- (e) In [MR5], strong evidence is given for the conjecture that  $R(5, 5) = 43$  and that there exist exactly 656 critical graphs on 42 vertices. The upper bound of 49 was established in 1997 [MR5]. Angelteit and McKay improved it to 48 in 2016 [AnM1], and further to 46 in 2023 [AnM3]. It is known that there does not exist any  $(5, 5)$ -critical graph which is self-complementary [Chv2].
- (f) The graphs constructed by Exoo in [Ex9-Ex25], and some others, are available electronically from <http://cs.indstate.edu/ge/RAMSEY>. Fujita [Fuj1] maintains a website with some lower bound constructions, using witness graphs obtained independently from Exoo.
- (g) The equalities  $R(3, 8) = 28$  and  $R(3, 9) = 36$  were verified using certifiable DRAT proofs with SAT+CAS technique [DuLBG]. The equality  $R(4, 5) = 25$  was verified using the HOL4 (higher order logic) interactive theorem prover system [GauB].
- (h) Formalization of statements in Ramsey theory using *Lean* 4 was presented in [NaSZ]. Other SAT-based approaches for Ramsey graphs generation and enumeration are proposed in [KirS]. In 2025, heavy computer algebra techniques were used in [Pau] to formalize in *Isabelle* the new breakthrough upper bound on  $R(k, k)$  [CamGMS] from 2023, see also item 2.4.8.
- (i) *Cyclic* (or *circulant*) graphs are often used for Ramsey graph constructions. Several cyclic graphs establishing lower bounds were given in the Ph.D. dissertation by J.G. Kalbfleisch in 1966, and many others were published in the next few decades (see [RaK1]). Harborth and Krause [HaKr1] presented all best lower bounds up to 102 from cyclic graphs avoiding complete graphs. In particular, no lower bound in Table Ia can be improved with a cyclic graph on less than 102 vertices, except possibly for  $R(3, k)$  for  $k \geq 13$ . See also items 2.3.1 and 5.18.o [HaKr1]. Larger cyclic heuristic constructions for  $R(3, k)$  were explored in [JiLTX1, JiLTX2]. Several best lower bounds from *distance colorings*, a slightly more general concept than circular graphs, are presented in [HaKr2].
- (j) The claim that  $R(5, 5) = 50$  posted on the web [Stone] is in error, and despite being shown to be incorrect more than once, this value is still being cited by some authors. The bound  $R(3, 13) \geq 60$  [XieZ] cited in the 1995 version of this survey was shown to be incorrect in [Piw1]. Another incorrect construction for  $R(3, 10) \geq 41$  was described in [DuHu].
- (k) There are really only two general upper bound inequalities useful for small parameters, namely 2.3.a and 2.3.b. Stronger upper bounds for specific parameters were difficult to obtain, and they often involved massive computations, like those for the cases of  $(3, 8)$  [McZ],  $(3, 10)$  [Goer1, Ang1],  $(4, 5)$  [MR4],  $(4, 6)$  and  $(5, 5)$  [MR5, AnM1, AnM2, AnM3]. The bound  $R(6, 6) \leq 166$ , only one more than in [Mac], is an easy consequence of a theorem in [Walk] (2.3.b) and  $R(4, 6) \leq 41$ . Since 2023, we know that  $R(6, 6) \leq 160$  [AnM3], see Table Ib.
- (l) T. Spencer [Spe4], Mackey [Mac], and Huang and Zhang [HZ2], using the bounds for minimum and maximum number of edges in  $(4, 5)$  Ramsey graphs listed in [MR3, MR5],

were able to establish new upper bounds for several higher Ramsey numbers, improving on all of the previous longstanding best results by Giraud [Gi3, Gi5, Gi6]. These were further improved as reported in Table Ib.

- (m) In Table Ia, only some of the higher bounds implied by 2.3.\* are shown, and more similar bounds could be derived. In general, we show bounds beyond the contiguous small values if they improve on results previously reported in this survey or published elsewhere. Some easy upper bounds implied by 2.3.a are marked as [Ea1].
- (n) In 2009, we recomputed the upper bounds in Table Ia marked [HZ2] using the method from the paper [HZ2], because the bounds there relied on an overly optimistic personal communication from T. Spencer. Further refinements of this method were deployed in [HZ3, ShZ1, Shi2]. The paper [Shi2] subsumes the main results of the manuscripts [ShZ1, ShZ2]. All these bounds are now improved by the bounds in Table Ib obtained in [AnM2, AnM3].
- (o) In 2013, Boza [Boza5] using the method of [HWSYZH], which is abbreviated as HW+ in Table Ia, computed the bounds marked HW+ by starting from better upper bounds for smaller parameters. Most of the currently shown bounds are thus better than those originally listed in [HWSYZH, HZ3]. All these bounds are now improved by the bounds in Table Ib obtained in [AnM2, AnM3].
- (p) In 2015, Exoo and Tatarevic obtained several lower bound improvements marked [ExT] in Tables Ia and IIa by using some modifications of general circulant constructions, but especially related to the quadratic residues Paley graph  $Q_{101}$  and the cubic residues graph  $G_{127}$ . More bounds by Tatarevic are reported in [Tat]. In 2016, Kuznetsov [Kuz] obtained several further new lower bounds building up on circulant graphs. Also in 2015 and 2016, somewhat surprisingly, Kolodyazhny [Kol1, Kol2] improved four longstanding lower bounds on  $R(3, k)$ . The newest improvements of small lower bounds were found by Exoo for  $R(5, 6)$  in 2023, and by Nagda, Raghavan and Thakurta for  $R(3, 13)$ ,  $R(3, 18)$  and  $R(4, k)$  for  $k = 13, 14, 15$  [NaRT] in 2026.
- (q) Some lower bounds in Table Ia, like for  $R(6, 8)$  or  $R(8, 8)$ , may seem rather weak, yet they are not easy to improve. For comments on  $R(8, 8)$  see [ExT].
- (r) A perspective on lower bounds for Ramsey numbers as a statistical physics problem is discussed in [WoGKSF]. It builds up on another approach using adiabatic quantum optimization [RanMCG], but also on more classical heuristic algorithm developed by Exoo in many papers [Ex-nn].

## 2.2. Bounds for $R(k, l)$ , higher parameters

- (a) The upper bounds in Tables Ia and IIa marked [GoeR1, Les, Back1] were obtained mainly by deriving lower bounds for several cases of  $e(3, k, n)$ , which denotes the minimum number of edges in  $n$ -vertex triangle-free graphs with independence number less than  $k$ . The study of  $e(3, k, n)$  was also the main tool for the results obtained in [GrY, GR, RaK2, RaK3, GoeR2].

$l$	15	16	17	18	19	20	21	22	23
$k$									
3	74 Kol2 87 GoeR1	82 Ex21 97 Back3	92 W1+ 109 Back1	100 NaRT 120 Back4	106 W1+ 132 Back3	111 Ex16 145 Les	122 W1+ 157 Back4	131 W2+ 171 Back2	139 XWCS 185 Back2
4	158 Tat	170 Tat	200 Lia+	205 2.3.e	213 2.3.h	234 Ex16	242 SLZL	314 LinCa	
5	275 2.3.h	293 ExT	388 XSR2	396 2.3.h	411 XSR2	424 XSR2	441 2.3.i	492 Ihr	521 2.3.i
6	401 2.3.i	434 SLLL	548 SLLL	614 SLLL	710 SLLL	878 SLLL	888 2.3.h	1070 SLLL	
7		629 2.3.i	729 2.3.i	797 2.3.i	908 SLLL		1214 SLLL		
8	873 2.3.i		1005 2.3.i	1049 2.2.h	1237 2.2.h		1617 2.3.i		

**Table IIa.** Known bounds for higher two-color Ramsey numbers  $R(k, l)$ , with references. Lower and upper bounds are given for  $k = 3$ , only lower bounds for  $k \geq 4$ ; Lia+, W1+ and W2+ abbreviate LiaWXS, WWY1 and WSLX2, respectively.

$l$	24	25	26	27	28	29	30	31	32
$k$									
3	143 W1+	154 W2+	161 FuLS	172 LiLi	179 FuLS	190 LiLi	197 FuLS	208 FuLS	217 LiLi

$l$	33	34	35	36	37	38	39	40	41
$k$									
3	227 LiaX	234 LiaX	248 LiaX	255 FuLS	267 LiaX	278 LiaX	290 LiaX	298 LiaX	311 FuLS

$l$	42	43	44	45	46	47	48	49	50
$k$									
3	320 FuLS	333 FuLS	339 FuLS	354 Ji+	362 FuLS	380 Ji+	384 Ji+	402 Ji+	

**Table IIb.** Known lower bounds for higher Ramsey numbers  $R(3, l)$  for  $l \geq 24$ . W1+, W2+ and Ji+ abbreviate WSLX1, WSLX2 and JiLTX2, respectively.

$k$	11	12	13	14	15	16	17
lower bound	1597	1640	2557	2989	5485	5605	8917
reference	2.2.c	Tat	2.2.c	2.2.c	2.2.c	2.2.c	LuSL
$k$	18	19	20	21	22	23	24
lower bound	11005	17885	21725	30925	39109	49421	
reference	LuSL	LuSL	Ex23	Ex23	Ex23	Ex23	

**Table IIc.** Known lower bounds for diagonal Ramsey numbers  $R(k, k)$  for  $k \geq 11$ ;  
All lower bounds for  $k \geq 13$  are from Paley graphs, see also 2.2.c below.

- (b) *Ramsey Calculus* [Back1], is an extensive manuscript by Backelin, which, among other goals, addresses the derivation of  $e(3, k, n)$  and the corresponding realisers while avoiding reliance on computer assisted results as far as possible. It achieves the derivation of several lower bounds for  $e(3, k+1, n)$  better than those in [GoeR1, RaK3, RaK4] for  $n$  close to and above  $13k/4$ . Better lower bounds on  $e(3, k, n)$  sometimes lead to better upper bounds on  $R(3, l)$ , like for  $l = 18$  and  $l = 20$  [Back4]. Further improvements to bounds on  $e(3, k, n)$  were obtained in [Krü].
- (c) The construction by Mathon (discovered later but independently by Shearer [She2], see also items 2.3.j, 6.2.k and 6.2.l), using the data obtained by Shearer [She4] for primes up to 7000, implies the lower bounds in Table IIc marked 2.2.c. The first two bounds credited in Table IIc to [LuSL] also follow similarly from the data in [She4]. The same approach does not improve on the bound  $R(12, 12) \geq 1639$  [XSR2], later increased to 1640 [Tat]. The bounds in [Ex23] were obtained by extending data for Paley graphs beyond [She4] and improving on [LiaWXCS].
- (d) The lower bounds marked [XuXR], [XXER], [XSR2], 2.3.e, 2.3.h and 2.3.i need not be cyclic. Several of the Cayley colorings from [Ex16] are also non-cyclic. All other lower bounds listed in Table IIa/b were obtained by construction of circular graphs.
- (e) The graphs establishing lower bounds marked 2.3.h can be constructed by using appropriately chosen graphs  $G$  and  $H$  with a common  $m$ -vertex induced subgraph, similarly as it was done in several cases in [XuXR].
- (f) Yu [Yu2] constructed a special class of triangle-free cyclic graphs establishing several lower bounds for  $R(3, k)$ , for  $k \geq 61$ . All of these bounds can be improved by the inequalities in 2.3.c and data from Tables Ia and IIa/b.
- (g) Unpublished bound  $R(4, 22) \geq 314$  [LiSLW] improved over 282 given in [SuL]. [LinCa] obtained the same bound, and also  $R(4, 25) \geq 458$ . Not yet published bounds  $R(3, 23) \geq 139$  [XWCS] and  $R(4, 17) \geq 200$  [LiaWXS] improve over 137 and 182 obtained in [WSLX2] and [LuSS1], respectively. The bound  $R(9, 17) \geq 1411$  is given in [XuXR]. Large cyclic heuristic constructions for  $R(3, k)$  for  $k < 50$  were explored in [JiLTX1, JiLTX2].
- (h) Two special cases,  $R(8, 18) \geq 1049$  and  $R(8, 19) \geq 1237$ , can be obtained by applying 2.3.i and 2.3.h below. In both cases we start with the 816-vertex graph  $G$ , witnessing  $R(8, 13) \geq 817$ , obtained by 2.3.i. Next, for properly chosen graphs  $H$  in the application

of 2.3.h, we have large common subgraphs of  $G$  and  $H$ , namely the 101-vertex witness of  $R(6, 6) \geq 102$  and the 204-vertex witness of  $R(7, 7) \geq 205$ , respectively.

- (i) One can expect that the lower bounds in Tables IIa/b are weaker than those in Table Ia, especially smaller ones, in the sense that some of them should not be that hard to improve, in contrast to the bounds in Table Ia.

### 2.3. General results on $R(k, l)$

- (a)  $R(k, l) \leq R(k-1, l) + R(k, l-1)$ , with strict inequality when both terms on the right hand side are even [GG]. There are obvious generalizations of this inequality for avoiding graphs other than complete.
- (b)  $R(k, k) \leq 4R(k, k-2) + 2$  [Walk].
- (c) Explicit construction for  $R(3, 3k+1) \geq 4R(3, k+1) - 3$ , for all  $k \geq 2$  [CleDa], explicit construction for  $R(3, 4k+1) \geq 6R(3, k+1) - 5$ , for all  $k \geq 1$  [ChCD].
- (d) Explicit triangle-free graphs with independence  $k$  on  $\Omega(k^{3/2})$  vertices [Alon2, CoPR]. For other constructive results in relation to  $R(3, k)$  see [BrBH1, BrBH2, Fra1, Fra2, FrLo, Goer1, Gri, Klam1, Loc, RaK2, RaK3, RaK4, Stat, Yu1]. See also 2.4.3 and 2.4.4 in the next subsection.
- (e) The study of bounds for the difference between consecutive Ramsey numbers was initiated in [BEFS], where the bound  $R(k, l) \geq R(k, l-1) + 2k - 3$ , for  $k, l \geq 3$ , was established by a construction. In 1980, Erdős and Sós (cf. [Erd2, ChGra2]) asked: If we set  $\Delta_{k, l} = R(k, l) - R(k, l-1)$ , then is it true that  $\Delta_{k, k+1}/k \rightarrow \infty$  as  $k \rightarrow \infty$ ? Only easy bounds on  $\Delta_{k, l}$  are known, in particular for  $k = 3$  we have  $3 \leq \Delta_{3, l} \leq l$ . For some discussion of the roadblocks on the latter see [XSR2, Goer2, ZhuXR]. It is also known that  $R(3, k) \geq R(3, K_{k-1} - e) + 4$  [ZhuXR].
- (f) A conjecture that  $R(k, l) \geq R(k-1, l+1)$ , for all  $3 \leq k \leq l$ , is dubbed DC for the Diagonal Conjecture. Its implications, evidence for validity, and some related problems are discussed in [LiaRX]. For the multicolor version of the DC and its consequences see item 6.2.v.
- (g) By taking a disjoint union of two critical graphs one can easily see that  $R(k, p) \geq s$  and  $R(k, q) \geq t$  imply  $R(k, p+q-1) \geq s+t-1$ . Xu and Xie [XuX1] improved this construction to yield better general lower bounds, in particular  $R(k, p+q-1) \geq s+t+k-3$ .
- (h) For  $2 \leq p \leq q$  and  $3 \leq k$ , if  $(k, p)$ -graph  $G$  and  $(k, q)$ -graph  $H$  have a common induced subgraph on  $m$  vertices without  $K_{k-1}$ , then  $R(k, p+q-1) > n(G) + n(H) + m$ . In particular, this construction implies the bounds  $R(k, p+q-1) \geq R(k, p) + R(k, q) + k - 3$  and  $R(k, p+q-1) \geq R(k, p) + R(k, q) + p - 2$  [XuX1, XuXR], with some small improvements, such as using the term  $k-2$  instead of  $k-3$  in the first bound for  $k \geq 5$  [XSR2].
- (i)  $R(2k-1, l) \geq 4R(k, l-1) - 3$  for  $l \geq 5$  and  $k \geq 2$ , and in particular for  $k=3$  we have  $R(5, l) \geq 4R(3, l-1) - 3$  [XXER].

- (j) If the quadratic residues Paley graph  $Q_p$  of prime order  $p = 4t + 1$  contains no  $K_k$ , then  $R(k, k) \geq p + 1$  and  $R(k + 1, k + 1) \geq 2p + 3$  [Math, She2]. Data for larger  $p$  was obtained in [LuSL], and further for  $p$  up to 25000 in [Ex23]. See also 3.1.f, and items 6.2.k and 6.2.l for similar multicolor results.
- (k) Study of Ramsey numbers for large disjoint unions of graphs [Bu0, Bu9], in particular  $R(nK_k, nK_l) = n(k + l - 1) + R(K_{k-1}, K_{l-1}) - 2$ , for  $n$  large enough [Bu8].
- (l)  $R(k, l) \geq L(k, l) + 1$ , where  $L(k, l)$  is the maximal order of any cyclic  $(k, l)$ -graph. A compilation of many best cyclic bounds was presented in [HaKr1].
- (m) The graphs critical for  $R(k, l)$  are  $(k - 1)$ -vertex connected and  $(2k - 4)$ -edge connected, for  $k, l \geq 3$  [BePi]. This was improved to vertex connectivity  $k$  for  $k \geq 5$  and  $l \geq 3$  in [XSR2].
- (n) All Ramsey-critical  $(k, l)$ -graphs are Hamiltonian for  $k \geq l - 1 \geq 1$  and  $k \geq 3$ , except when  $(k, l) = (3, 2)$  [XSR2].
- (o) Two-color lower bounds can be obtained by using items 6.2.m, 6.2.n and 6.2.o with  $r = 2$ . Some generalizations of these were obtained in [ZLLS].

## 2.4. Some pillars of asymptotics of $R(k, l)$

In this section we give only some selected pointers to the literature dealing with asymptotics of two-color Ramsey numbers. This survey was designed mostly for small, finite, and combinatorial results, but still we wish to give the reader some useful and representative references to more traditional papers studying the infinite. In particular, we include the main breakthroughs in this area, such as the progress on asymptotics of  $R(3, k)$ ,  $R(4, k)$  and  $R(k, k)$ .

- (1) In 1947, Erdős gave a simple probabilistic proof that  $R(k, k) > 2^{k/2}$  [Erd1]. In 1975, Spencer [Spe1] improved it to  $R(k, k) > \sqrt{2}e^{-1}k2^{k/2}(1+o(1))$ . More probabilistic asymptotic lower bounds were obtained in [Spe1, Spe2, AlPu].
- (2) The limit of  $R(k, k)^{1/k}$ , if it exists, is between  $\sqrt{2}$  and 4 [GRS, GrRö, ChGra2]. This limit is now known to be strictly below 4, as implied by the last result of item (8) below.
- (3) In 1995, Kim obtained a breakthrough result by proving that  $R(3, k) = \Theta(k^2/\log k)$  [Kim]. By 2013, the best known lower and upper bounds constants were  $1/4$  [BohK2, BohK3] and 1 (implicit in [She1]), respectively. An independent proof of the lower bound constant  $1/4$  was presented in [FizGM]. In 2025, an improvement to  $1/3$  and a conjecture that  $R(3, k) = (1/2 + o(1))(k^2/\log k)$  were posted on arXiv [CamJMS]. Soon afterwards, the proof of the lower bound  $1/2$  on the constant was announced [HeHKP].
- (4) Other asymptotic and general results on triangle-free graphs in relation to  $R(3, k)$  can be found in [Boh, AlBK, AjKS, Alon2, CleDa, ChCD, CoPR, Gri, FrLo, Loc, She1, She3].
- (5) Explicit constructions yielded the lower bounds  $R(4, k) \geq \Omega(k^{8/5})$ ,  $R(5, k) \geq \Omega(k^{5/3})$  and  $R(6, k) \geq \Omega(k^2)$  [KosPR]. For the same cases of  $k$  classical probabilistic arguments give  $\Omega((k/\log k)^{5/2})$ ,  $\Omega((k/\log k)^3)$  and  $\Omega((k/\log k)^{7/2})$ , respectively [Spe2]. These were improved to  $\Omega(k^{5/2}/(\log k)^2)$ ,  $\Omega(k^3/(\log k)^{8/3})$  and  $\Omega(k^{7/2}/(\log k)^{13/4})$ , respectively, in [Boh, BohK1], and in general to  $R(s, t) \geq \Omega(t^{(s+1)/2}/(\log t)^{(s^2-s-4)/(2s-4)})$ , for fixed  $s$  and large  $t$  [BohK1].
- (6) In a 2023 arXiv posting (journal paper appeared in 2024), Mattheus and Verstraëte [MatVer] proved that there exist constants  $c_1$  and  $c_2$  such that for all  $k \geq 3$  we have

$$c_1 k^3/\log^4 k \leq R(4, k) \leq c_2 k^3/\log^2 k.$$

A brief background and explanation of this breakthrough result, and its link to finite geometry, are presented in [Bish].

- (7) Explicit construction of a graph with clique and independence  $k$  on  $2^{c \log^2 k / \log \log k}$  vertices was presented by Frankl and Wilson [FraWi], and further constructions by Chung [Chu3] and Grolmusz [Grol1, Grol2]. In 2012, the best explicit construction for large  $k$  by Barak et al. [BarRSW] improved over [FraWi] by giving such a graph on  $2^{2^{(\log \log k)^c}}$  vertices for some  $c > 1$ , or equivalently, on  $n$  vertices, where  $\log \log n = (\log \log k)^c$ . In 2016, this was improved by Cohen [Coh] to  $\log \log n = (\log k)^d$ , for a positive constant  $d$ , and in a 2023 arXiv posting, Xin Li [LiXin] presents explicit Ramsey graphs on  $n$

vertices with no clique or independent set of order  $\log^c n$ , for some  $c > 1$ . Explicit constructions such as these are usually weaker than known probabilistic results.

- (8) In 2009, Conlon [Con1] obtained the best until then upper bound for the diagonal case

$$R(k+1, k+1) \leq \binom{2k}{k} k^{-c \log k / \log \log k}.$$

In 2020 on arXiv (journal paper appeared in 2023), Sah [Sah] improved it to

$$R(k+1, k+1) \leq \binom{2k}{k} e^{-c(\log k)^2}.$$

In 2023, Campos, Griffiths, Morris and Sahasrabudhe [CamGMS] posted a 57-page long paper on arXiv showing that there exists  $\varepsilon > 0$  such that

$$R(k, k) \leq (4 - \varepsilon)^k.$$

Thus, in (2), the upper bound on the limit  $R(k, k)^{1/k}$  is  $4 - \varepsilon$ . A concrete improved upper bound  $R(k, k) \leq 3.8^k$  was obtained in [GuNNW], and much simplified proof of this bound is in [BalB+].

- (9) A survey by Yuval Wigderson [Wig3] presents the latest developments concerning the upper bound in (8) as in [CamGMS], with particular attention to the book algorithm, the main tool used therein. A gentle introduction to several upper bound breakthroughs, in particular those in (3), (6) and (8), was presented by Morris [Mor].
- (10) In 2025, Ma, Shen and Xie [MaSX] posted a 41-page long paper on arXiv claiming that their results give an exponential improvement for the lower bound on the off-diagonal cases  $R(k, ck)$ , the first such improvement since 1947. A simpler proof and an enhancement of this bound was posted by Hunter, Milojević and Sudakov [HunMS].
- (11) Other older asymptotic bounds can be found, for example, in [Chu3, McDS, Boh, BohK1] (lower bound) and [Tho] (upper bound), and for many other bounds in the general case of  $R(k, l)$  consult [Spe2, GRS, GrRö, Chu4, ChGra2, LiRZ1, AlPu, Kriv, ConFS7]. Two nontechnical essays by Sloman on recent developments in this area appeared in the *Quanta Magazine* in 2023 [Slo].

**3. Two Colors:  $K_n - e, K_3, K_{m, n}$**

**3.1. Dropping one edge from complete graph**

This section contains known values and nontrivial bounds for the two color case when the avoided graphs are complete or have the form  $K_k - e$ , but not both are complete.

- (a) The exact values in Table IIIa involving  $K_3 - e$  are obvious, since one can easily see that  $R(K_3 - e, K_k) = R(K_3 - e, K_{k+1} - e) = 2k - 1$  for all  $k \geq 2$ .
- (b) More bounds (beyond those shown in Tables IIIa/b) can be easily obtained using Table Ia/b, an obvious generalization of the inequality  $R(k, l) \leq R(k-1, l) + R(k, l-1)$ , and by the monotonicity of Ramsey numbers, in this case  $R(K_{k-1}, G) \leq R(K_k - e, G) \leq R(K_k, G)$ .

$G \setminus H$	$K_3 - e$	$K_4 - e$	$K_5 - e$	$K_6 - e$	$K_7 - e$	$K_8 - e$	$K_9 - e$	$K_{10} - e$	$K_{11} - e$
$K_3 - e$	3	5	7	9	11	13	15	17	19
$K_3$	5	7	11	17	21	25	31	37	42 45
$K_4 - e$	5	10	13	17	28	30 32	36 45	43 57	73
$K_4$	7	11	19	30	37 49	52 71	62 102	135	170
$K_5 - e$	7	13	22	37	65	66 91	69 136	188	261
$K_5$	9	16	30	43 59	65 102	81 172	121 273	381	511
$K_6 - e$	9	17	37	45 70	66 124	83 206	334	505	757
$K_6$	11	21	43 51	58 104	202	353	615	944	1346
$K_7 - e$	11	28	65	66 124	247	432	761	1218	1964
$K_7$	13	28 29	65 79	80 178	364	702	1278	2119	3197
$K_8$	15	36 39	69 117	284	636	1277	2504	4402	7040
$K_9$	17	41 53	75 169	446	1053	2293	4667	8675	14631
$K_{10}$	19	49 68	233	660	1680	3847	8345	16064	

**Table IIIa.** Bounds on the Ramsey numbers  $R(G, H)$ , for complete or missing one edge graphs  $G$  and  $H$ , but not both complete. Known exact values appear as centered entries, lower bounds as top entries, and upper bounds as bottom entries.

$G$	$H$	$K_4-e$	$K_5-e$	$K_6-e$	$K_7-e$	$K_8-e$	$K_9-e$	$K_{10}-e$	$K_{11}-e$
$K_3$		ChH2	Clan	FRS1	GrH	Ra1	Ra1	MPR GoeR2	WWY2 GoeR2
$K_4-e$		ChH1	FRS2	McR	McR	AnM2 LidP	VO BZ2	GoeVO BZ2	BZ2
$K_4$		ChH2	EHM1	Boza6 JamKR	Ex14 LidP	VO BZ3	VO Ea1	AnM3	AnM3
$K_5-e$		FRS2	CE+	VO	VO	Ea1 BZ3	Ea1 Ea1	BZ3	BZ3
$K_5$		BoH	Ex6 Ang2	Ea1 Ea1	VO LidP	VO BZ3	VO BZ3	AnM3	AnM3
$K_6-e$		McR	VO	Ex14 HZ3	VO LidP	VO BZ3	BZ3	BZ3	BZ3
$K_6$		McN/ ShWR	VO Ea1	Ea1 BZ4	BZ3	BZ4	BZ3	AnM3	AnM3
$K_7-e$		McR	VO	VO LidP	Ea1	BZ3	BZ3	BZ3	BZ3
$K_7$		Ea1 LidP	Ea1 BZ3	Ea1 BZ3	BZ4	BZ4	BZ4	AnM3	AnM3
$K_8$		VO LidP	Ea1 BZ3	BZ3	BZ3	BZ4	BZ4	AnM3	AnM3
$K_9$		VO BZ2	Ea1 BZ3	BZ3	BZ3	BZ3	BZ4	AnM3	AnM3
$K_{10}$		VO BZ2	BZ3	BZ3	BZ3	BZ3	BZ3	AnM3	

**References for Table IIIa.**

CE+ abbreviates CIEHMS, for some details of BZ1-BZ4 see item 3.1.g, the bounds marked [AnM3] are trivially implied by entries in Table Ib.

$k$	11	12	13	14	15	16
lower bound	42 WWY2	49 VO	55 GoeR2	61 VO	69 WWY2	74 Ea1
upper bound	45 GoeR2	53 GoeR2	62 GoeR2	71 GoeR2	80 GoeR2	91 GoeR2

**Table IIIb.** Lower and upper bounds for  $R(K_3, K_k - e)$  for  $11 \leq k \leq 16$ .

- (c) Upper bounds for Ramsey numbers  $R(K_k, K_l - e)$  marked [AnM3] in references for Table IIIa are trivially implied by the bounds on  $R(K_k, K_l)$  in Table Ib. These were obtained in 2023 by Angeltveit and McKay using linear programming, and they improve over upper bounds in Table Ia. [AnM2] points to little older and weaker bounds for some higher cases. The upper bounds by Lidický and Pfender [LidP] use flag algebras.

(d) Two special exact values, and several other bounds were obtained by Van Overberghe [VO] in 2020. The surprisingly large exact values  $R(K_5-e, K_6-e) = 37$  and  $R(K_5-e, K_7-e) = 65$  exploit some previously known strongly regular graphs on 27, 36 and 64 vertices, namely the Schläfli graph,  $NO^-(6, 2)$  and  $VO^-(6, 2)$ . See the website by A. E. Brouwer [Brou] for a great collection of strongly regular graphs, including those used in [VO].

(e) Some lower bounds beyond the range of Table IIIa are as follows:

$49 \leq R(K_3, K_{12}-e)$	[GoeVO]
$61 \leq R(K_3, K_{14}-e)$	[GoeVO]
$82 \leq R(K_4-e, K_{16}-e)$	[VO]
$128 \leq R(K_4, K_{12}-e)$	[Shao]
$2987 \leq R(K_{14}-e, K_{14}-e)$	[LiShen]

The results of MS thesis [VO] are contained and extended in a journal article [GoeVO]. The bound  $R(K_4, K_{12}-e) \geq 128$  [Shao] is derived by using one color of the (4, 4, 4;127)-coloring defined in [HiIr]. For comments related to the bound in [LiShen], see the next item.

(f) If the quadratic residues Paley graph  $Q_p$  of prime order  $p = 4t + 1$  contains no  $K_k - e$ , then  $R(K_{k+1}-e, K_{k+1}-e) \geq 2p + 1$ . In particular,  $R(K_{14}-e, K_{14}-e) \geq 2987$  [LiShen]. This was generalized to  $K_k - F$  for some small graphs  $F$  instead of an edge  $e (=K_2)$  [WaLi]. See also item 2.3.j.

(g) This item follows from personal communications by Boza [Boza5, Boza8]. The upper bound  $R(K_6, K_5-e) \leq 53$  [BZ1] was obtained in 2012, while [BZ2, BZ3, BZ4] are from 2024, all by Boza. [BZ2] refers to computer-free simple proofs of some upper bounds, those marked [BZ3] use the methods and theorems in [LiShen, HYZ, HWSYZH], and [BZ4] use in addition item 5.16.n. These approaches expand on 2.1.o, or are implied by [Boza6], the previous work [Boza1, Boza3, BoPo], the method of [HZ3], and the bounds given in [GoeR2, AnM2, AnM3]. The enumeration of all  $(K_6, K_4-e)$ -graphs [ShWR] is used in [BoPo]. The upper bounds in Table IIIa marked [BZ3] were recomputed by Boza in 2026 to account for the implications of the upper bound  $R(K_5, K_5-e) \leq 30$  obtained by Angeltveit [Ang2].

(h) All  $(K_3, K_k - e)$ -graphs were enumerated for  $k \leq 6$  [Ra1] and  $k = 7$  [Fid2, GoeR2]. Full sets of  $(K_l, K_k - e)$ -graphs were posted for the parameters  $(K_3, K_k - e)$  for  $k \leq 7$ ,  $(K_4, K_k - e)$  for  $k \leq 5$ , and  $(K_5, K_k - e)$  for  $k \leq 4$  ([Fid2], available until 2014), and other full and restricted families at [BrCGM, Fuj1].

(i) The number of  $(K_3, K_k - e)$ -critical graphs for  $k = 4, 5$  and  $8$  is 4, 2 and 9, respectively [MPR]. There are 7 critical graphs for  $R(K_3, K_9-e)$ , and at least 40 such graphs for  $R(K_3, K_{10}-e)$  [GoeR2].

(j) The critical graphs are unique for:  $R(K_3, K_k - e)$  for  $k=3$  [Tr], 6 and 7 [Ra1],  $R(K_4-e, K_4-e)$  [FRS2],  $R(K_5-e, K_5-e)$  [Ra3, GoeVO] and  $R(K_4-e, K_7-e)$  [McR]. All of the critical graphs for the cases  $R(K_4-e, K_4)$  [EHM1],  $R(K_4-e, K_5)$  and  $R(K_5-e, K_4)$  [DzFi1] are known, and there are 5, 13 and 6 of them, respectively.

The unpublished value of  $R(K_4 - e, K_6)$  [McN] was confirmed in [ShWR], where in addition all 24976 critical graphs were found. All  $(K_4 - e, K_6 - e)$ -graphs were enumerated in [GoeVO].

- (k) If  $m \leq n$  then  $R(K_4 - e, K_{m+n+1}) \geq R(3, m+1) + R(3, n+1) + n$ .  
Study of the growth of  $R(K_4 - e, K_n)$  and its relationship to  $R(K_3, K_n)$  [JiLSX].
- (l)  $R(K_k - e, K_k - e) \leq 4R(K_{k-2}, K_k - e) - 2$  [LiShen].  
For a similar inequality for complete graphs see 2.3.b.
- (m) Study of the cases  $R(K_m, K_n - K_{1,s})$  and  $R(K_m - e, K_n - K_{1,s})$ , with several exact values for special parameters [ChaMR]. This study was extended to some cases involving  $R(K_m - K_3)$  [MonCR].
- (n) The upper bounds from [ShZ2] are subsumed by a later article [Shi2].
- (o) The upper bounds in [HZ3] were obtained by a reasoning generalizing the bounds for classical numbers in [HZ2]. Several other results from Section 2.3 apply, though checking in which situation they do may require looking inside the proofs whether they still hold for  $K_n - e$ . The upper bounds in the manuscript [HTHZ1] are based on [HZ3].

### 3.2. Triangle versus other graphs

- (a)  $R(3, k) = \Theta(k^2/\log k)$  [Kim].  
For more comments on asymptotics see Section 2.4.(3) and the items 3.2.p/q below.
- (b) Explicit construction for  $R(3, 3k+1) \geq 4R(3, k+1) - 3$ , for all  $k \geq 2$  [CleDa],  
explicit construction for  $R(3, 4k+1) \geq 6R(3, k+1) - 5$ , for all  $k \geq 1$  [ChCD].
- (c) Explicit triangle-free graphs with independence  $k$  on  $\Omega(k^{3/2})$  vertices [Alon2, CoPR].
- (d) Ramsey numbers for  $K_3$  versus books [RoS1], wheels [BuE3, RaJi], fans [LiR2, GuGS],  
and generalized fans [HaoLin, BudPra, WaQi]; see also the corresponding and other sub-sections of Section 5.
- (e)  $R(K_3, K_7 - 2P_2) = R(K_3, K_7 - 3P_2) = 18$  [SchSch2].
- (f)  $R(K_3, K_3 + \bar{K}_m) = R(K_3, K_3 + C_m) = 2m + 5$ , for  $m \geq 212$  [Zhou1],  
 $R(K_3, K_2 + T_n) = 2n + 3$  for  $n$ -vertex trees  $T_n$ , for  $n \geq 4$  [SonGQ],  
 $R(K_3, K_1 + nK_3) = 6n + 1$ , for  $n \geq 3$  [HaoLin].
- (g)  $R(K_3, G) = 2n(G) - 1$  for any connected  $G$  on at least 4 vertices and with at most  $(17n(G) + 1)/15$  edges, in particular for  $G = P_i$  and  $G = C_i$ , for all  $i \geq 4$  [BEFRS1].
- (h)  $R(K_3, Q_n) = 2^{n+1} - 1$  for large  $n$  [GrMFSS], where  $Q_n$  is the  $n$ -dimensional hypercube.  
For related publications on the general case of  $R(K_m, Q_n)$  see [FizGMSS, ConFLS] and item 5.17.o.
- (i) Relations between  $R(3, k)$  and graphs with large  $\chi(G)$  [BiFJ],  
further detailed study of the relation between  $R(3, k)$  and the chromatic gap [GySeT].

- (j)  $R(K_3, G) \leq 2e(G) + 1$  for any graph  $G$  without isolated vertices [Sid3, GodK, HngJL].
- (k)  $R(K_3, G) \leq n(G) + e(G)$  for all  $G$ , a conjecture [Sid2].
- (l)  $R(K_3, G)$  for all connected  $G$  up to 9 vertices [BrBH1, BrBH2].
- (m)  $R(K_3, G)$  for all graphs  $G$  on 10 vertices [BrGS], except 10 cases (three of which, including  $G = K_{10} - e$ , were solved [GoeR2]). See also several items in Section 8.1.
- (n)  $R(nK_3, nK_3) = 5n$  for  $n \geq 2$ ,  $R(mK_3, nK_3) = 3m + 2n$  for  $m \geq n \geq 2$  [BES], and  $R(c(nK_3), c(nK_3)) = 7n - 2$  for  $n \geq 2$ , where  $c(nK_3)$  is any connected graph containing  $n$  vertex disjoint triangles [GySá3].
- (o) Formulas for  $R(nK_3, mG)$  for all  $G$  of order 4 without isolates [Zeng].
- (p) For every positive constant  $c$ , and for  $\Delta$  and  $n$  large enough, there exists  $n$ -vertex graph  $G$  with  $\Delta(G) \leq \Delta$  for which  $R(K_3, G) > cn$  [Bra3].
- (q)  $R(K_3, K_{k,k}) = \Theta(k^2/\log k)$  [LinLi2].
- (r) For  $R(K_3, K_n)$  see Section 2, and for  $R(K_3, K_n - e)$  see Section 3.1.
- (s) Since  $B_1 = F_1 = C_3 = W_3 = K_3$ , other sections apply. For some other cases involving triangles see also [Boh, AjKS, BrBH1, BrBH2, FrLo, Fra1, Fra2, BiFJ, Gri, GySeT, Loc, Klam1, LiZa2, RaK2, RaK3, RaK4, She1, She3, Spe2, Stat, Yu1].

### 3.3. Complete bipartite graphs

This subsection gathers information on Ramsey numbers where specific bipartite graphs are avoided in edge colorings of  $K_n$  (as everywhere in this survey), in contrast to the often studied bipartite Ramsey numbers, which are not covered in this survey, where the edges of complete bipartite graphs  $K_{n,m}$  are colored.

#### 3.3.1. Numbers

Table IVa presents data for  $C_4 = K_{2,2}$  versus stars, while the following Tables IVb and IVc gather information about small complete bipartite graphs. Their first versions were based on the surveys by Lortz and Mengersen [LoM3, LoM4]. All cases involving  $K_{1,2} = P_3$  are solved by a formula for  $R(P_3, G)$ , which holds for all isolate-free graphs  $G$ , derived in [ChH2]. The star versus star numbers are given below in the item 3.3.2.a and in Section 5.5.

- (a) Note that for graph  $G$  to avoid  $K_{1,n}$  is equivalent to  $\delta(G) < n$ . Thus, for general monotonicity we have, for example, that rows of Table IVb are nondecreasing, but we do not know if they are strictly increasing.
- (b) See function  $f(n)$  in the item 3.3.2.d as one involving  $C_4$ . Similarly, we also have  $R(C_4, K_{1,n+1}) \leq R(C_4, K_{1,n}) + 2$  [Chen], which is the same claim as in 3.3.2.c.

$n$	$R(C_4, K_{1,n})$	reference	$n$	$R(C_4, K_{1,n})$	reference
2	4	ChH2	22	28	SunSh
3	6	ChH2	23	29	Par5
4	7	Par3	24	30	WuSZR
5	8	Par3	25	31	Par3
6	9	FRS4	26	32	Par3
7	11	FRS4	27	33	Boza8
8	12	Tse1	28	35	Boza8
9	13	Par3/Tse1	29	36	Boza8
10	14	Par3/Tse1	30	37	Boza8
11	16	Tse1	31	38	Boza8
12	17	Tse1	32	39	Boza8
13	18	DyDz2	33	40	Boza8
14	19	DyDz2	34	41	WuSR
15	20	Law2/DyDz2	35	42	WuSR
16	21	Par3/DyDz2	36	43	WuSR
17	22	Par3	37	44	Boza8
18	23	ZhaBC1	38	45	ZhaCC2
19	24	WuSR	39	46-47	WuSR/DyDz2
20	25	WuSR	40	47	ZhaCC2
21	27	Par5	41	49	Boza8

**Table IVa.** Values and bounds for Ramsey numbers  $R(C_4, K_{1,n})$  for  $n \leq 41$ ,  $C_4 = K_{2,2}$ .

(c)  $R(C_4, K_{1,n}) = R(C_4, W_{n+1})$  for  $n \geq 6$  [ZhaBC1]. See also items (b)-(f) in Section 4.3 concerning results on Ramsey numbers of wheels versus  $C_4$ . Values and bounds on several cases higher than those in Table IVa are reported in [NoBa, WuSR].

(d) For all odd prime powers  $q$  we have [Boza8]:

$$q^2 + 1 \leq R(C_4, K_{1,q^2-q+1}) \leq q^2 + 2 \leq R(C_4, K_{1,q^2-q+2}) \leq q^2 + 3.$$

A number of similar results were included in the early papers by Parsons [Par3, Par5], and many in newer papers such as [ZhaBC1, WuSR, WuSZR, ZhaCC2, DyDz2, ZhaCC3]. In 2025, an extensive summary of results involving  $C_4$ , and in particular on  $R(C_4, K_{1,n})$ , was published in [ChenZZ7].

(e) Formula for  $R(K_{1,n}, K_{k_1, k_2, \dots, k_t, m})$  for  $m$  large enough, in particular for  $t=1, k_1=2$  with  $n \leq 5, m \geq 3$  and  $n=6, m \geq 11$ , for example  $R(K_{1,5}, K_{2,7}) = 15$  [Stev].

(f) The values and bounds in larger cases of  $R(K_{2,2}, K_{2,n})$  are 20, 22, 22, 24, 25, 26, 27, 28/29, 30 and 32 for  $12 \leq n \leq 21$ , respectively. All of them were given in [HaMe4], except those for  $n = 14, 15$  and  $18$ , which were obtained in [Dyb1]. More exact values for prime powers  $\lceil \sqrt{n} \rceil$  and  $\lceil \sqrt{n} \rceil + 1$  can be found in [HaMe4].

(g) The known values of  $R(K_{2,2}, K_{3,n})$  are 15, 16, 17, 20 and 22 for  $6 \leq n \leq 10$  [Lortz], and  $R(K_{2,2}, K_{3,12}) = 24$  [Shao]. See Tables IVa and IVb for the smaller cases, and [HaMe4] for upper bounds and values for some prime powers  $\lceil \sqrt{n} \rceil$ .

(h)  $R(K_{2,n}, K_{2,n})$  is equal to 46, 50, 54, 57 and 62 for  $12 \leq n \leq 16$ , respectively.

The first open diagonal case is  $65 \leq R(K_{2,17}, K_{2,17}) \leq 66$  [EHM2].

The status of all higher cases for  $n < 30$  is listed in [LoM1].

$p, q$ $m, n$	1, 2	1, 3	1, 4	1, 5	1, 6	2, 2	2, 3	2, 4	2, 5	3, 3	3, 4
2, 2	4 ChH2	6 ChH2	7 Par3	8 Par3	9 FRS4	6 ChH1					
2, 3	5 ChH2	7 FRS4	9 Stev	10 FRS4	11 FRS4	8 HaMe4	10 Bu4				
2, 4	6 ChH2	8 HaMe3	9 Stev	11 HaMe4	13 LoM4	9 HaMe4	12 ExRe	14 EHM2			
2, 5	7 ChH2	9 HaMe3	11 Stev	13 Stev	14 LoM4	11 HaMe4	13 LoM3	16 LoM1	18 EHM2		
2, 6	8 ChH2	10 HaMe3	11 Stev	14 Stev	15* Shao	12 HaMe4	14 LoM3	17 LoM3	20 LoM1		
3, 3	7 ChH2	8 HaMe3	11 LoM4	12 LoM4	13 LoM4	11 Lortz	13 HaMe3	16 LoM4	18 LoM4	18 HaMe3	
3, 4	7 ChH2	9 HaMe3	11 LoM4	13 LoM4	14 LoM4	11 Lortz	14 LoM4	17 Sh1+	20 VO-LidP	19-20 VO-LidP	25 VO-LidP
3, 5	9 ChH2	10 HaMe3	13 Sh1+	15 Sh1+	17 LidP	14 HaMe4	17* Shao	19-20 VO-LidP	21-23 VO-LidP	21-24 VO-LidP	25-29 VO-LidP

**Table IVb.** Ramsey numbers  $R(K_{m,n}, K_{p,q})$ ; unpublished result marked with \*, Sh1+ abbreviates ShaXBP, the bound  $R(K_{3,5}, K_{2,5}) \geq 21$  is also in [ShaoWX].

$m$	2	3	4	5	6	7	8	9	10	11
6	12 HaMe4	14 LoM3	17 LoM3	20 LoM1	21 EHM2					
7	14 HaMe4	17 LoM3	19 LoM3	21 LoM3	24 LoM1	26 EHM2				
8	15 HaMe4	18 LoM3	20 LoM3	22-23 LoM3	25 VO-LoM3	28 LoM1	30 EHM2			
9	16 HaMe4	19 LoM3	22 LoM3	25* Shao	27* Shao	29* Shao	32 LoM1	33 EHM2		
10	17 HaMe4	21 LoM3	24 LoM3	27 LoM3	27-29 LoM3	29-31 VO-LoM3	33 VO-LoM3	36 LoM1	38 EHM2	
11	18 HaMe4	$\geq 22$ VO	$\geq 25$ VO	$\geq 28$ VO	$\geq 29$ VO	$\geq 33$ VO	35 VO-LoM3	37 VO-LoM3	40 LoM1	42 EHM2

**Table IVc.** Known Ramsey numbers  $R(K_{2,n}, K_{2,m})$  for  $6 \leq n \leq 11$  and  $2 \leq m \leq 11$ . Results marked [VO] are from MS thesis by Van Overberghe, and they now have appeared in a journal paper [GoeVO]. Unpublished results improving over [LoM3] are marked with a \*.

- (i) Some exact values and bounds for parameters beyond the range of Tables IVb/c are:

$$\begin{array}{ll}
 R(K_{1,4}, K_{4,4}) = R(K_{1,5}, K_{4,4}) = 13 & \text{[ShaXBP]} \\
 R(K_{1,4}, K_{1,2,3}) = R(K_{1,4}, K_{2,2,2}) = 11 & \text{[GuSL]} \\
 R(K_{1,7}, K_{2,3}) = 13 & \text{[Par4, Par6]} \\
 R(K_{2,2}, K_{4,4}) = 14 & \text{[HaMe4]} \\
 R(K_{2,2}, K_{4,5}) = 15 & \text{[Shao]} \\
 R(K_{2,2}, K_{4,6}) = 16 & \text{[Shao]} \\
 R(K_{2,2}, K_{5,5}) = R(K_{2,3}, K_{3,5}) = 17 & \text{[Shao]} \\
 R(K_{3,5}, K_{3,5}) = 33 & \text{[VO][LidP]} \\
 33 \leq R(K_{4,4}, K_{4,4}) \leq 49 & \text{[VO][LidP]}
 \end{array}$$

- (j) A number of general upper and lower bounds for  $R(K_{s,t}, K_{s,t})$ , in particular for small fixed  $s$ , and for some slightly off-diagonal cases were obtained in [LoM2]. They can be used to derive the upper bounds for the cases listed in (i) above and (k) below.
- (k) Several lower bounds of the form  $R(K_{s,t}, K_{s,t}) \geq m$  from distance colorings, a slightly more general concept than circular graphs, were presented in [HaKr2] for the following triples  $(s, t, m)$ : (3,6,38), (3,7,42), (3,8,43), (3,9,54), (4,5,42), (4,6,43), (4,7,54), (5,5,54).

### 3.3.2. General results

- (a)  $R(K_{1,n}, K_{1,m}) = n + m - \varepsilon$ , where  $\varepsilon = 1$  if both  $n$  and  $m$  are even and  $\varepsilon = 0$  otherwise [Har1]. It is also a special case of multicolor numbers for stars obtained in [BuRo1].
- (b)  $R(K_{1,3}, K_{m,n}) = m + n + 2$  for  $m, n \geq 1$  [HaMe3].
- (c)  $R(K_{1,n+1}, K_{2,2}) \leq R(K_{1,n}, K_{2,2}) + 2$  [Chen].
- (d)  $R(K_{1,n}, K_{2,2}) = f(n) \leq n + \lceil \sqrt{n} \rceil + 1$ , with  $f(q^2) = q^2 + q + 1$  and  $f(q^2 + 1) = q^2 + q + 2$  for every  $q$  which is a prime power [Par3]. Furthermore,  $f(n) \geq n + \sqrt{n} - 6n^{11/40}$  [BEFRS4]. For more bounds on  $f(n)$  see [Par5, Chen, ChenJ, MoCa, WuSR, WuSZR, ZhaBC1, BoRa]. Summary of what is known and further progress were reported in two 2017 papers [ZhaCC2, ZhaCC3]. The values of  $f(n)$  are known for all  $n \leq 38$ , see Table IVa and Section 3.3.1. In 2025, an extensive summary of results involving  $C_4$ , and in particular on  $R(C_4, K_{1,n})$ , was published in [ChenZZ7]. Compare also to items in Section 4.3 involving  $C_4$ .
- (e)  $R(K_{2,\lambda+1}, K_{1,\nu-k+1})$  is either  $\nu + 1$  or  $\nu + 2$  if there exists a  $(\nu, k, \lambda)$ -difference set. This and other related results are presented in [Par4, Par5]. See also [GoCM, GuLi].
- (f) Formulas and bounds on  $R(K_{2,2}, K_{2,n})$ , and bounds on  $R(K_{2,2}, K_{m,n})$ . In particular, we have  $R(K_{2,2}, K_{k,n}) = n + k\sqrt{n} + c$ , for  $k = 2, 3, 4$ , some prime powers  $\lceil \sqrt{n} \rceil$  and  $\lceil \sqrt{n} \rceil + 1$ , and some  $-1 \leq c \leq 3$  [HaMe4]. An improvement of the latter for some special cases of  $n$  was obtained in [Dyb1]. Asymptotics of  $R(K_{2,2}, K_{n,n})$  is discussed in [LiuLi2], where in particular the lower bound  $R(K_{2,2}, K_{n,n}) = \Omega(n^{3/2}/\log n)$  is presented. See also item 4.2.d.

- (g)  $R(K_{2,n}, K_{2,n}) \leq 4n - 2$  for all  $n \geq 2$ , and the equality holds if and only if there exists a strongly regular  $(4n - 3, 2n - 2, n - 2, n - 1)$ -graph [EHM2].
- (h) Conjecture that  $4n - 3 \leq R(K_{2,n}, K_{2,n}) \leq 4n - 2$  for all  $n \geq 2$ . Many special cases are solved and several others are discussed in [LoM1].
- (i)  $R(K_{2,n-1}, K_{2,n}) \leq 4n - 4$  for all  $n \geq 3$ , with the equality if there exists a symmetric Hadamard matrix of order  $4n - 4$ . There are only 4 cases in which the equality is still open for  $3 \leq n \leq 58$ , namely 30, 40, 44 and 48 [LoM1].
- (j)  $R(K_{2,n-s}, K_{2,n}) \leq 4n - 2s - 3$  for  $s \geq 2$  and  $n \geq s + 2$ , with the equality in many cases involving Hadamard matrices or strongly regular graphs. Asymptotics of  $R(K_{2,n}, K_{2,m})$  for  $m \gg n$  [LoM3].
- (k) Some algebraic lower and upper bounds on  $R(K_{s,n}, K_{t,m})$  for various combinations of  $n, m$  and  $1 \leq t, s \leq 3$  [BaiLi, BaLX]. A general lower bound  $R(K_{m,n}) \geq 2^m(n - n^{0.525})$  for large  $n$  [Dong]. For every fixed real  $\alpha > 0$  and integer  $m \geq 1$ , for large  $n$ , we have  $R(K_{m, \text{round}(\alpha n)}, K_{m,n}) = ((\alpha^{1/m} + 1)^m + o(1))n$  [LiuDu].
- (l) Upper bounds for  $R(K_{2,2}, K_{m,n})$  for  $m, n \geq 2$ , with several cases identified for which the equality holds. Special focus on the cases for  $m = 2$  [HaMe4].
- (m) Let  $G$  be any isolate-free graph with  $p$  vertices and  $q \geq 2$  edges. Then it holds that  $R(K_{2,2}, G) \leq 2q + 1$ , with the equality for  $G = qK_2$  or  $G = K_3$ , and  $R(K_{2,2}, G) \leq 2p + q - 2$ . Some generalizations to  $R(K_{2,k}, G)$  [JRB].
- (n) Bounds for the numbers of the form  $R(K_{k,n}, K_{k,m})$ , especially for fixed  $k$  and close to the diagonal cases. Asymptotics of  $R(K_{3,n}, K_{3,m})$  for  $m \gg n$  [LoM2]. Asymptotic upper bounds on  $R(K_{2,n}, K_{2,m})$  and on  $R(K_{3,3}, K_{m,n})$  for  $m \leq 3$  [WaLL].
- (o)  $R(nK_{1,3}, mK_{1,3}) = 4n + m - 1$  for  $n \geq m \geq 1, n \geq 2$  [BES].
- (p) Asymptotics for  $K_{2,m}$  versus  $K_n$  [CaLRZ]. Upper bound asymptotics for  $K_{k,m}$  versus  $K_n$  [LiZa1] and for some bipartite graphs  $K_n$  [JiSa].
- (q) Special two-color cases apply in the study of asymptotics for multicolor Ramsey numbers for complete bipartite graphs [ChGra1].

## 4. Two Colors: Numbers Involving Cycles

Note: The paper *Ramsey Numbers Involving Cycles* [Ra4] is based on the revision #12 of this survey. It collects and comments on the results involving cycles versus any graphs, in two or more colors. It contains some more details than this survey, but only until 2009.

### 4.1. Cycles, cycles versus paths and stars

Note that  $P_2 = K_2$ ,  $B_1 = F_1 = C_3 = W_3 = K_3$ ,  $B_2 = K_4 - e$ ,  $P_3 = K_3 - e$ ,  $W_4 = K_4$  and  $C_4 = K_{2,2}$  imply other identities not mentioned explicitly. Conversely, subsections on wheels or fans in other sections enhance and complement the contents of this section.

#### 4.1.1. Cycles

- (a)  $R(C_3, C_3) = 6$  [BaLiu, Bush, GG],  
 $R(C_4, C_4) = 6$  [ChH1].
- (b)  $R(C_3, C_n) = 2n - 1$  for  $n \geq 4$ ,  $R(C_4, C_n) = n + 1$  for  $n \geq 6$ ,  
 $R(C_5, C_n) = 2n - 1$  for  $n \geq 5$ , and  $R(C_6, C_6) = 8$  [ChaS].
- (c) Result obtained independently in [Ros1] and [FS1], a new simpler proof in [KáRos]:

$$R(C_m, C_n) = \begin{cases} 2n - 1 & \text{for } 3 \leq m \leq n, m \text{ odd, } (m, n) \neq (3, 3), \\ n - 1 + m/2 & \text{for } 4 \leq m \leq n, m \text{ and } n \text{ even, } (m, n) \neq (4, 4), \\ \max\{n - 1 + m/2, 2m - 1\} & \text{for } 4 \leq m < n, m \text{ even and } n \text{ odd.} \end{cases}$$

- (d)  $R(mC_3, nC_3) = 3n + 2m$  for  $n \geq m \geq 1$ ,  $n \geq 2$  [BES].
- (e)  $R(mC_4, nC_4) = 2n + 4m - 1$  for  $m \geq n \geq 1$ ,  $(n, m) \neq (1, 1)$  [MiSa, LiWa1].
- (f) Formulas for  $R(mC_4, nC_5)$  [LiWa2].
- (g) Formulas and bounds for  $R(mC_n, sC_t)$  [Den2, Biel1].
- (h) Characterization of all graphs critical for  $R(C_4, C_n)$  [WuSR].
- (i) Study of  $R(S_1, S_2)$ , where  $S_1$  and  $S_2$  are sets of cycles [Hans].  
 A conjecture generalizing 4.1.c stated in [Hans] was proved in [WaCh2].
- (j) Unions of cycles, formulas and bounds for various cases including diagonal, different lengths, different multiplicities [MiSa, Den2], disjoint cycles versus  $K_n$  [Fuj2], and their relation to 2-local Ramsey numbers [Biel1].
- (k) Asymptotics for powers of cycles [AllBS]. Exact values for squares of paths  $P_n$  and cycles  $C_{3n}$  for sufficiently large  $n$ , in particular we have  $R(C_{3n}^2, C_{3n}^2) = 9n - 3$  for large  $n$  [AllMRS].

**4.1.2. Cycles versus paths**

(l) Result obtained by Faudree, Lawrence, Parsons and Schelp in 1974 [FLPS]:

$$R(C_m, P_n) = \begin{cases} 2n - 1 & \text{for } 3 \leq m \leq n, \quad m \text{ odd,} \\ n - 1 + m/2 & \text{for } 4 \leq m \leq n, \quad m \text{ even,} \\ \max\{m - 1 + \lfloor n/2 \rfloor, 2n - 1\} & \text{for } 2 \leq n \leq m, \quad m \text{ odd,} \\ m - 1 + \lfloor n/2 \rfloor & \text{for } 2 \leq n \leq m, \quad m \text{ even.} \end{cases}$$

$m$	$n$	$R(P_m, P_n)$	$R(P_m, C_n)$	$R(C_m, P_n)$	$R(C_m, C_n)$
3	3	3	5	5	6
3	4	4	4	7	7
3	5	5	5	9	9
3	6	6	6	11	11
3	7	7	7	13	13
3	8	8	8	15	15
3	9	9	9	17	17
3	10	10	10	19	19
4	4	5	5	5	6
4	5	6	7	6	7
4	6	7	7	7	7
4	7	8	8	8	8
4	8	9	9	9	9
4	9	10	10	10	10
4	10	11	11	11	11
5	5	6	9	9	9
5	6	7	7	11	11
5	7	8	9	13	13
5	8	9	9	15	15
5	9	10	10	17	17
5	10	11	11	19	19
6	6	8	8	8	8
6	7	9	11	9	11
6	8	10	10	10	10
6	9	11	11	11	11
6	10	12	12	12	12
7	7	9	13	13	13
7	8	10	10	15	15
7	9	11	13	17	17
7	10	12	12	19	19
8	8	11	11	11	11
8	9	12	15	12	15
8	10	13	13	13	13
9	9	12	17	17	17
9	10	13	13	19	19
10	10	14	14	14	14

**Table Va.** Ramsey numbers  $R(P_m, P_n), R(P_m, C_n), R(C_m, P_n), R(C_m, C_n)$  for  $3 \leq m \leq n \leq 10$ . We have  $R(P_m, P_n) \leq R(P_m, C_n) \leq R(C_m, C_n)$ , for other details see [FLPS], and items 4.1.a/b/c/l/m, 5.1.a.

- (m) For all  $n$  and  $m$  it holds that  $R(P_m, P_n) \leq R(C_m, P_n) \leq R(C_m, C_n)$ . Each of the two inequalities can become an equality, and, as derived in [FLPS], all four possible combinations of  $<$  and  $=$  hold for an infinite number of pairs  $(m, n)$ . See Table Va for all the cases of  $3 \leq m \leq n \leq 10$ , also including  $R(C_n, P_m)$ . For example, if both  $m$  and  $n$  are even, and at least one of them is greater than 4, then all four Ramsey numbers for  $(P_m, P_n), (C_m, P_n), (C_n, P_m), (C_m, C_n)$  are the same, however, for no  $m$  and  $n$  all four of them are different. For some related generalizations see [BEFRS2].

### 4.1.3. Cycles versus stars

Only partial results for  $C_m$  versus stars are known. Lawrence [Law1] settled the cases for odd  $m$  and for long cycles (see also [Clark, Par6]). The case for short even cycles is open, and it is related in particular to bipartite graphs. A number of results for  $C_4 = K_{2,2}$  are collected in Sections 3.3.1 and 3.3.2. The most known general exact result [Law1] is:

$$R(C_m, K_{1,n}) = \begin{cases} 2n+1 & \text{for odd } m \leq 2n+1, \\ m & \text{for } m \geq 2n. \end{cases}$$

Some new cases for even  $m$  not too small with respect to  $n$  were settled in 2016, in particular the exact values of  $R(C_6, K_{1,n})$  for all  $n \leq 11$  were completed in [ZhaBC6]. The equality  $R(C_6, K_{1,12}) = 17$  was obtained in [SunSh]. The progress on asymptotics for large even  $m$ , and exact values for large even  $m$  and  $n$  not too large were obtained in [AllLPZ].

## 4.2. Cycles versus complete graphs

Since 1976, it was conjectured that  $R(C_n, K_m) = (n-1)(m-1) + 1$  for all  $n \geq m \geq 3$ , except  $n = m = 3$  [FS4, EFRS2]. Various parts of this conjecture were proved as follows: for  $n \geq m^2 - 2$  [BoEr], for  $n > 3 = m$  [ChaS], for  $n \geq 4 = m$  [YHZ1], for  $n \geq 5 = m$  [BolJY+], for  $n \geq 6 = m$  [Schi1], for  $n \geq m \geq 7$  with  $n \geq m(m-2)$  [Schi1], for  $n \geq 7 = m$  [ChenCZ1], and for  $n \geq 4m + 2, m \geq 3$  [Nik]. Open conjectured cases are marked in Table Vb by "conj."

In 2019, Keevash, Long and Skokan [KeeLS] proved the above conjecture for  $n \geq C \log m / \log \log m$  for some absolute constant  $C \geq 1$ , and furthermore that for any  $\epsilon > 0$  and  $n > n(\epsilon)$ , for the lower bound it holds that  $R(C_n, K_m) > m \log m \gg (n-1)(m-1) + 1$  for all  $3 \leq n \leq (1-\epsilon) \log m / \log \log m$ .

- (a) The first column in Table Vb gives data from the first row in Table I.
- (b) Joint credit [He2/JR4] in Table Vb refers to two cases in which Hendry [He2] announced the values without presenting the proofs, which later were given in [JR4]. The special cases of  $R(C_6, K_5) = 21$  [JR2] and  $R(C_7, K_5) = 25$  were solved independently in [YHZ2] and [BolJY+]. The double pointer [JaBa/ChenCZ1] refers to two independent papers, similarly as [JaAl/ZZ3], except that in the latter case [ZZ3] refers to an

unpublished manuscript. For joint credits marked in Table Vb with "-", the first reference is for the lower bound and the second for the upper bound. The cases of the conjecture for  $R(C_n, K_8)$ , for  $10 \leq n \leq 15$ , are confirmed in [Ban2].

- (c) Erdős et al. [EFRS2] asked what is the minimum value of  $R(C_n, K_m)$  for fixed  $m$ , and they suggested that it might be possible that  $R(C_n, K_m)$  first decreases monotonically, then attains a unique minimum, then increases monotonically with  $n$ . If so, then the results in [KeeLS] stated above imply that this transition of behavior happens at  $n = \Theta(\log m / \log \log m)$ .

	$C_3$	$C_4$	$C_5$	$C_6$	$C_7$	$C_8$	$C_9$	...	$C_n$ for $n \geq m$
$K_3$	6 2.1.a	7 ChaS	9 ...	11	13	15	17	...	$2n - 1$ ChaS
$K_4$	9 GG	10 ChH2	13 He4/JR4	16 JR2	19 YHZ1	22 ...	25	...	$3n - 2$ YHZ1
$K_5$	14 GG	14 Clan	17 He2/JR4	21 JR2	25 YHZ2	29 BolJY+	33 ...	...	$4n - 3$ BolJY+
$K_6$	18 Kéry	18 Ex2-RoJa1	21 JR5	26 Schi1	31 ...	36	41	...	$5n - 4$ Schi1
$K_7$	23 Ka2-GrY	22 RaT-JR1	25 Schi2	31 CheCZN	37 CheCZN	43 JaBa/Ch+	49 Ch+	...	$6n - 5$ Ch+
$K_8$	28 GR-McZ	26 RaT	29 LidP	36 ChenCX	43 ChenCZ1	50 JaAl/ZZ3	57 BatJA	...	$7n - 6$ conj.
$K_9$	36 Ka2-GR	30 RaT-LaLR	33-36 LidP	41 LidP	49 BanAA	56 BanAA	65 conj.	...	$8n - 7$ conj.
$K_{10}$	40-41 Ex5-Ang1	36 LaLR						...	$9n - 8$ conj.
$K_{11}$	47-50 Ex20-GoeR1	40-43 VO-BoRa						...	$10n - 9$ conj.
$K_{12}$	53-59 Kol1-Les	43-51 BoRa						...	$11n - 10$ conj.

**Table Vb.** Known Ramsey numbers  $R(C_n, K_m)$ ;  
Ch+ abbreviates ChenCZ1, for comments on joint credits see 4.2.b.

- (d) There exist constants  $c_1, c_2 > 0$  such that  $c_1(m^{3/2}/\log m) \leq R(C_4, K_m) \leq c_2(m/\log m)^2$ . The lower bound, obtained by Bohman and Keevash ([BohK1] in 2010, see also 4.2.e/f/g below) improved over an almost 40 years old bound  $c(m/\log m)^{3/2}$  by Spencer [Spe2], using the probabilistic method. The upper bound was reported in a paper by Caro, Li, Rousseau and Zhang [CaLRZ], who in turn give the credit to an unpublished work by Szemerédi from 1980. A refined upper bound,  $R(C_4, K_m) \leq (1 + o(1))(m/\log m)^2$ , was presented by Liu and Li [LiuLi2] in 2021.

- (e) Erdős, in 1981, in the Ramsey problems section of the paper [Erd3] formulated a challenge by asking for a proof of  $R(C_4, K_m) < m^{2-\varepsilon}$ , for some  $\varepsilon > 0$ . To date, no such proof is known.
- (f) The lower bound asymptotics  $R(C_n, K_m) = \Omega(m^{(n-1)/(n-2)}/\log m)$ , for fixed  $n$  and large  $m$ , was obtained by Bohman and Keevash [BohK1]. An improvement of the latter by a polylogarithmic factor for odd  $n$  and improved exponents to  $11/8$  and  $11/9$  for  $C_5$  and  $C_7$ , respectively [MuVer]. Further improvement to exponents  $10/7$  and  $5/4$  for  $C_5$  and  $C_7$ , with a polylogarithmic factor, in [ConMMV]. Note that for  $n = 4$  it gives the lower bound in 4.2.d above. See also [Spe2, FS4, AIRö] for previous related results.
- (g) In 2025, the lower bound in (f) above was further improved to  $m^{(n-1)/(n-2) + \varepsilon_n + o(1)}$  for fixed odd  $n > 7$  and large  $k$  [CamJMPS].
- (h) Enumeration of all  $(C_n, K_4)$ -graphs for  $n \leq 7$  [JaNR].  
Classification of all  $(C_n, K_6)$ -critical graphs for  $n \geq 15$  [JaNS].
- (i) A theta graph  $\theta_n$  is obtained from the cycle  $C_n$  by adding one edge between some of its nonadjacent vertices. Summary of what is known about  $R(\theta_n, K_k)$ , and an additional result for  $k = 6$ , are collected in [BanJBJ]. The cases for  $k = 5$  and  $n = 7, 8, 9$  are solved in [JaBBJ], and the cases for  $k = 7$  and  $(n = 7$  or  $n \geq 14)$  are solved in [Ban1].
- (j) Let  $C_{\leq n}$  be the set of cycles of length at most  $n$ , and let the *girth*  $g(G)$  be the length of the shortest cycle in graph  $G$ . Probabilistic lower bound asymptotics for  $R(C_{\leq n}, K_m)$  [Spe2] currently is the same as for  $R(C_n, K_m)$ , for fixed  $n$ . However, there are clear differences already for girth 4 and 5 and small  $m$ : Backelin [Back1, Back2] found that  $R(C_{\leq 4}, K_m) = 6, 8, 11, 15, 18$  for  $m = 3, 4, 5, 6, 7$ , and that  $R(C_{\leq 5}, K_m) = 5, 8, 10, 13, 15$ , also for  $m = 3, 4, 5, 6, 7$ , respectively.
- (k) Erdős et al. [EFRS2] proved various facts about  $R(C_{\leq n}, K_m)$ , and in particular that it is equal to  $2m - 1$  for  $n \geq 2m - 1$ , and to  $2m$  for  $m < n < 2m - 1$ . The upper asymptotics for  $R(C_{\leq n}, K_m)$  is implied in the study of independence number in graphs with odd girth  $n$  [Den1]. The following close to the diagonal exact values were obtained in [WuSL]:  $R(C_{\leq n}, K_n)$  is equal to  $2n$  and  $2n + 1$  for odd  $n$  and even  $n$ , respectively, and  $R(C_{\leq n}, K_{n+1}) = 2n + 3$  for odd  $n \geq 5$  and even  $n \geq 16$ .
- (l)  $R(C_{\geq n}, K_{m_1, \dots, m_k}) = (k - 1)(n - 1) + m_1$  for  $m_1 \leq \dots \leq m_k$ ,  $5m_{k-1} + 3m_k \leq n$  [PoSu1]. The same equality holds for  $R(C_n, K_{m_1, \dots, m_k})$  for large  $m_i$ 's and very large  $n$  [PoSu2]. See also Section 5.15.
- (m) Upper bound asymptotics [BoEr, FS4, EFRS2, CaLRZ, Sud1, LiZa2, AIRö, DoLL2].

### 4.3. Cycles versus wheels

Note: In this survey the wheel graph  $W_n = K_1 + C_{n-1}$  has  $n$  vertices, while some authors use the definition  $W_n = K_1 + C_n$  with  $n + 1$  vertices. The generalized wheel  $W_{k,n}$  is defined by  $W_{k,n} = K_k + C_n$ , so  $W_n = W_{1,n-1}$ . For the cases involving  $W_3 = C_3$  versus  $C_m$  see Sections 3.2 and 4.2.

- (a)  $R(C_3, W_n) = 2n - 1$  for  $n \geq 6$  [BuE3]. All critical graphs have been enumerated. The critical graphs are unique for  $n = 3, 5$ , and for no other  $n$  [RaJi].
- (b)  $R(C_4, W_n) = 14, 16, 17$  for  $n = 11, 12, 13$ , respectively [Tse1],  
 $R(C_4, W_n) = 18, 19, 20, 21$  for  $n = 14, 15, 16, 17$ , respectively [DyDz2], and several higher values and bounds, including 9 cases of  $n$  between 18 and 44 [WuSR, WuSZR].
- (c)  $R(C_4, W_n) \leq n + \lceil (n-1)/3 \rceil$  for  $n \geq 7$  [SuBUB], which was improved to  $R(C_4, W_n) \leq n + \sqrt{n-2} + 1$  for  $n \geq 11$  [DyDz2].
- (d)  $R(C_4, W_{q^2+1}) = q^2 + q + 1$  for prime power  $q \geq 4$  [DyDz2], exact values of  $R(C_4, W_{q^2+2})$  and  $R(C_4, W_{q^2-i})$  for special  $q$  and small  $i$  [WuSZR].
- (e)  $R(C_4, W_n) = R(C_4, K_{1,n-1})$  for  $n \geq 7$  [ZhaBC1], an alternate proof relying less on prior computational results [ChenZZ7] (in these papers the definition  $W_n = K_1 + C_n$  is used). Thus, the data in Table IVa and several items in Section 3.3.1 on  $R(C_4, K_{1,n})$  apply also to wheels.
- (f) Tight bounds on  $R(C_4, W_n)$  for  $46 \leq n \leq 93$  [NoBa].

	$C_3$	$C_4$	$C_5$	$C_6$	$C_7$	$C_8$	$C_m$	for
$W_4$	9 GG	10 ChH2	13 He4	16 JR2	19 YHZ1	22 ...	$3m-2$ ...	$m \geq 4$ YHZ1
$W_5$	11 Clan	9 Clan	9 He2	11 JR2	13 SuBB2	15 ...	$2m-1$ ...	$m \geq 5$ SuBB2
$W_6$	11 BuE3	10 JR3	13 ChvS	16 SuBB2	19 SuBB2	22 ...	$3m-2$ ...	$m \geq 4$ SuBB2
$W_7$	13 BuE3	9 Tse1	13 LuLL	11 LuLL	13 LuLL		$2m-1$ ...	$m \geq 10$ Ch1
$W_8$	15 BuE3	11 Tse1	15 LuLL	16 LuLL	19 Ch2	22 ...	$3m-2$ ...	$m \geq 6$ Ch2
$W_9$	17 BuE3	12 Tse1	17 LuLL	13 LuLL	17 LuLL		$2m-1$ ...	$m \geq 13$ Ch1
$W_{10}$	19 BuE3	13 Tse1		16 Z1	19 Z2		$3m-2$ ...	$m \geq 9$ Ch2
$W_n$ for	... $2n-1$ $n \geq 6$ BuE3		$2n-1$ $n \geq 19$ Zhou2		$2n-1$ $n \geq 29$ Zhou2			cycles  large wheels

**Table VI.** Ramsey numbers  $R(W_n, C_m)$  for  $n \leq 10, m \leq 8$ ; Ch1, Ch2, Z1, Z2 abbreviate ChenCMN, ChenCNZ, ZhaBC6, ZhaZZ, respectively.

- (g)  $R(C_7, W_n) = 2n - 1$  for  $n = 9, 10, 11$  [ZhaZZ].
- (h)  $R(W_n, C_m) = 2n - 1$  for odd  $m$  with  $n \geq 5m - 6$  [Zhou2]. The range of  $n$  was extended in [ZhaZC].

- (i)  $R(W_n, C_m) = 3m - 2$  for even  $n \geq 4$  with  $m \geq n - 1$ ,  $m \neq 3$ , was conjectured by Surahmat et al. [SuBT1, SuBT2, Sur]. Parts of this conjecture were proved in [SuBT1, ZhaCC1, Shi5, ZhaBC2, ZhaZC], and the proof was completed in [ChenCNZ].
- (j) Conjecture that  $R(W_n, C_m) = 2m - 1$  for odd  $n \geq 3$  and all  $m \geq 5$  with  $m > n$  [Sur]. It was proved for  $2m \geq 5n - 7$  [SuBT1], and improved to  $2m \geq 3n - 1$  in [ChenCMN]. For further progress see also [Shi5, ZhaBC2, Sanh, RaeZ, Alw, ZhaCh5].
- (k) In Table VI, observe four distinct situations with respect to the parity of  $m$  and  $n$ .
- (l) Cycles are Ramsey unsaturated for some wheels [AliSur], see also comments on [BaLS] in item 5.18.d.
- (m) Cycles versus multiple copies of wheels,  $tW_n$  [WaZh1, WaZh2].
- (n) Cycles versus generalized wheels  $W_{k,n}$  [Sur, SuBTB, Shi5, ZhaBC2, BieDa].

**4.4. Cycles versus books**

	$C_3$	$C_4$	$C_5$	$C_6$	$C_7$	$C_8$	$C_9$	$C_{10}$	$C_{11}$	$C_m$	for
$B_2$	7 RoS1	7 Fal7	9 Cal	11 Fal9	13 ...	15	17	19	21	$2m - 1$ ...	$m \geq 4$ Fal9
$B_3$	9 RoS1	9 Fal7	10 Fal9	11 JR2	13 Shi5	15 Fal9	17 ...	19	21	$2m - 1$ ...	$m \geq 6$ Fal9
$B_4$	11 RoS1	11 Fal7	11 Fal9	12 Sal1	13 Sal1	15 Shi5	17 Shi5	19 Fal9	21 ...	$2m - 1$ ...	$m \geq 7$ Fal9
$B_5$	13 RoS1	12 Fal7	13 Fal9	14 Sal1	15 Sal1	15 Sal2	17 Sal2	19 Shi5	21 Shi5	$2m - 1$ ...	$m \geq 8$ Fal9
$B_6$	15 RoS1	13 Fal7	15 Fal9	16 Sal2	17 Sal2	18 Sal2	18 Sal2		21 Shi5	$2m - 1$ ...	$m \geq 11$ Shi5
$B_7$	17 RoS1	16 Fal7	17 Fal9	16 Sal2	19 Sal2	20 Sal2	21 Sal2			$2m - 1$ ...	$m \geq 13$ Shi5
$B_8$	19 RoS1	17 Tse1	19 Fal9	17 Sal2	19 Sal2	22 Sal2	$\geq 23$ Sal2			$2m - 1$ ...	$m \geq 14$ Shi5
$B_9$	21 RoS1	18 Tse1	21 Fal9	18 Sal2			$\geq 25$ Sal2	$\geq 26$ Sal2		$2m - 1$ ...	$m \geq 16$ Shi5
$B_{10}$	23 RoS1	19 Tse1	23 Fal9	19 Sal2				$\geq 28$ Sal2		$2m - 1$ ...	$m \geq 17$ Shi5
$B_{11}$	25 RoS1	20 Tse1	25 Fal9							$2m - 1$ ...	$m \geq 19$ Shi5
$B_n$ for	... $2n + 3$ $n \geq 2$ RoS1	$\approx n$ some (c)(d)	... $2n + 3$ $n \geq 4$ Fal9		$2n + 3$ $n \geq 15$ Fal9		$2n + 3$ $n \geq 23$ Fal9		$2n + 3$ $n \geq 31$ Fal9		cycles  large books

**Table VII.** Ramsey numbers  $R(B_n, C_m)$  for  $n, m \leq 11$ ; et al. abbreviations: Fal7/FRS7, Fal9/FRS9, Cal/ChaRSPS, Sal1/ShaXBP, Sal2/ShaXB.

- (a) For the cases of  $B_1 = K_3$  versus  $C_m$  see Section 4.2.  
The exact values for the cases (3, 7), (4, 8), (4, 9), (5, 10), (5, 11) were obtained independently in [Sal1, Sal2]/[ShaXBP, ShaXB] using computer algorithms.
- (b)  $R(C_4, B_8) = 17$  [Tse2] (it was reported incorrectly in [FRS7] to be 16),  
 $R(C_4, B_{12}) = 21$  [Tse1],  $R(C_4, B_{13}) = 22$ ,  $R(C_4, B_{14}) = 24$  [Tse2],  
 $R(C_4, B_{16}) = 27$ ,  $R(C_4, B_{18}) = 29$  [LiLP],  
 $R(C_4, B_{17}) \leq 28$  was obtained in [BoRa].
- (c)  $R(C_4, B_n) \leq g(g(n))$ , where  $g(n) = n + \lfloor \sqrt{n-1} \rfloor + 2$ , and  
 $q^2 + q + 2 \leq R(C_4, B_{q^2-q+1}) \leq q^2 + q + 4$  for prime power  $q$  [FRS7].  
 $B_n$  is a subgraph of  $B_{n+1}$ , hence likely  $R(C_4, B_n) = n + O(\sqrt{n})$   
(compare to  $R(C_4, K_{2,n})$  in Section 3.3).
- (d)  $R(C_4, B_{(m-1)^2+t-2}) \leq m^2 + t$  for  $m \geq 4$  and  $0 \leq t \leq m-1$ , and an infinite number of exact values of  $R(C_4, B_n)$ . In particular,  $R(C_4, B_{q^2-q-2}) = q^2 + q - 1$  for all prime powers  $q \geq 4$  [LiLP].
- (e) Lower and upper bounds on  $R(C_4, B_n^{(k)})$  for  $k \geq 3$ , and exact values for infinitely many  $n$  for all  $k \geq 3$  [DouLLP], which generalize results in [LiLP] for  $k = 2$ , since  $B_n^{(2)} = B_n$ .
- (f) Upper bound on  $R(C_4, B_n)$  in terms of the upper bound on  $R(C_4, K_{1,n})$  [BoRa].
- (g)  $R(C_5, B_n) = 2n + 3$  for  $n \geq 4$ , and 9, 9, 10, for  $n = 1, 2, 3$ , respectively [FRS9].
- (h)  $R(B_n, C_m) = 2n + 3$  for odd  $m \geq 5$  with  $n \geq 4m - 13$ , and  
 $R(B_n, C_m) = 2m - 1$  for  $n \geq 1$ ,  $m \geq 2n + 2$  [FRS9]. The range of  $m$  in the latter was extended to  $m \geq 2n - 1 \geq 7$  in [ShaXB], and to  $m > (6n + 7)/4$  in [Shi5]. Further improvements for odd  $m$  [Gup].
- (i) Close to the diagonal we have  $R(B_n, C_n) \geq 3n - 2$  and  $R(B_{n-1}, C_n) \geq 3n - 4$  for  $n \geq 3$  [ShaXB], and for all  $n \geq 1000$  it holds [LinP]:

$$R(B_n, C_m) = \begin{cases} 3m - 2 & \text{for } 8n/9 + 112 \leq m \leq n, \\ 3n - 1 & \text{if } m = n + 1, \\ 3n & \text{for } n + 2 \leq m \leq (3n + 1)/2, \\ 2m - 1 & \text{if } m = \lceil 3n/2 \rceil + 1. \end{cases}$$

- (j) More exact values for some  $m$  and  $n$  beyond the above constraints were obtained in [HuL+]. More theorems on  $R(B_n, C_m)$  in [FRS7, FRS9, NiRo4, Zhou1, LinP, LiLP].
- (k) Cycles versus some generalized books  $B_n^{(k)} = nK_1 + K_k$  [Shi5]. Exact asymptotics for odd cycles versus  $B_n^{(k)}$  [LiuLi1], and for general cases close to the diagonal [LinP]. Long even cycles versus  $B_n^{(k)}$  [LinS].

#### 4.5. Cycles versus other graphs

- (a)  $C_4$  versus stars [Par3, Par4, Par5, Chen, ChenJ, DyDz2, GoMC, MoCa, WuSR, WuSZR, SunSh]. For several exact results see the cases of  $C_4 = K_{2,2}$  in the Tables IVa/b/c, and for general results see several items in Sections 3.3.1, 3.3.2, and 4.\*.
- (b)  $R(C_4, T_n) = \max\{4, n+1, R(C_4, K_{1,\Delta})\}$  for  $n$ -vertex trees  $T_n$  with maximum degree  $\Delta$  [BEFRS4]. For some related results see also [EFRS4, Bu7, Chen].
- (c)  $C_4$  versus unions of stars [HaABS, Has, HaJu]
- (d)  $C_4$  versus all graphs on six vertices [JR3]
- (e)  $C_4$  versus various types of complete bipartite graphs, see [LiuLi2] and Section 3.3.
- (f)  $R(C_4, G) \leq 2q+1$  for any isolate-free graph  $G$  with  $q \geq 2$  edges, and the equality holds for  $G = qK_2$  or  $G = K_3$  [RoJa2, JRB].
- (g)  $R(C_4, G) \leq 2p+q-2$  for any isolate-free graph  $G$  on  $p$  vertices and  $q \geq 2$  edges [JRB].
- (h)  $R(C_5, K_4-e) = 9$  [ChaRSPS],  
 $R(C_5, K_6-e) = 17$  [JR4], and  $C_5$  versus all graphs on six vertices [JR4].
- (i)  $R(C_6, K_5-e) = 17$ ,  $C_6$  versus all graphs on five vertices [JR2],  
 $C_6$  versus all stars up to  $K_{1,12}$  [ZhaBC6, SunSh].
- (j)  $R(C_{2m+1}, G) = 2n-1$  for sufficiently large sparse graphs  $G$  on  $n$  vertices, in particular  $R(C_{2m+1}, T_n) = 2n-1$  for all  $n > 1512m+756$ , for  $n$ -vertex trees  $T_n$  [BEFRS2]. The range of  $n$  for trees in the latter was extended to  $n \geq 25(2m+1)$  in [Bren1]. More on cycles versus trees long ago [FSS2] and recently [HuCh].
- (k)  $R(C_n, G) \leq 2q + \lfloor n/2 \rfloor - 1$ , for  $3 \leq n \leq 5$ , for any isolate-free graph  $G$  with  $q > 3$  edges [RoJa2]. It was conjectured that it also holds for other  $n$  [EFRS9]. In 2026, the bound  $R(C_n, G) \leq 2q + \lfloor (n-1)/2 \rfloor$  was proved for  $q$  large with respect to  $n$  [CamFMPP]. Independent progress for odd  $n$  [HngJL].
- (l) Study of conditions for graphs  $G$  for which  $R(C_n, G) = (n-1)(\chi(G)-1) + \sigma(G)$ , where  $\sigma(G)$  is chromatic surplus of  $G$  [AlIBS, HasHKL].
- (m) Cycles versus fans and other graphs [Shi5], large even cycles versus fans [YouLin1].
- (n) Exact asymptotics of odd cycles versus generalized fans [LiuLi1].
- (o) Cycles versus  $nW_4$  [Sudar5], cycles and unions of cycles versus  $mK_n$  [Sudar4].
- (p) Monotone paths and cycles [Lef2]. This 1993 work was followed by many papers on the so-called *ordered Ramsey numbers*, about which only a couple of hints are made in this survey, and they are not covered here otherwise.
- (q) Cycles versus  $K_{n,m}$  and multipartite complete graphs [BoEr, PoSu1], and Ramsey goodness of cycles [PoSu2]
- (r) Cycles versus generalized books and wheels [Shi5, Sur, SuBTB]
- (s) Cycles versus special graphs of the form  $K_n+G$  with small  $n \leq 3$  and sparse  $G$  [Shi5]
- (t) Odd cycles versus  $K_{2,m}$  [Gup]

## 5. General Graph Numbers in Two Colors

This section includes data with respect to general graph results. We tried to include all nontrivial values and identities regarding exact results, or references to them, but only those out of general bounds and other results which, in our opinion, may have a direct connection to the evaluation of specific numbers. If some small value cannot be found below, it may be covered by the cumulative data gathered in Section 8, or be a special case of a general result listed in this section. Note that  $P_2 = K_2$ ,  $B_1 = F_1 = C_3 = W_3 = K_3$ ,  $B_2 = K_4 - e$ ,  $P_3 = K_3 - e$ ,  $W_4 = K_4$  and  $C_4 = K_{2,2}$  imply other identities not mentioned explicitly.

### 5.1. Paths

- (a)  $R(P_m, P_n) = n + \lfloor m/2 \rfloor - 1$  for all  $n \geq m \geq 2$  [GeGy]  
Complete classification of  $R(P_m, P_n)$ -critical graphs [Hook]
- (b) See Table Va in Section 4.1.2 for comparison to  $R(C_m, P_n)$  and  $R(C_m, C_n)$
- (c) Stripes  $mP_2$  [CocL1, CocL2, Lor]  
Trails (paths with repeated vertices) [Osu, ConT]  
Disjoint unions of paths (also called linear forests) [BuRo2, FS2]  
Monotone paths [CaYZ], ordered path powers [Mub2], see also 4.5.p
- (d) Goodness of paths [PoSu1], see also Section 5.15.b/c/e
- (e) Asymptotics for powers of paths [AllBS]. Exact values for squares of paths  $P_n$  for sufficiently large  $n$ , in particular we have  $R(P_{3n}^2, P_{3n}^2) = R(P_{3n+1}^2, P_{3n+1}^2) = 9n - 3$ , and  $R(P_{3n+2}^2, P_{3n+2}^2) = 9n + 1$  for large  $n$  [AllMRS].
- (f) Complexity classification of  $(P_k, P_l)$ -goodness (i.e. the existence of  $(P_k, P_l)$ -colorings for input graphs, which is distinct from the *goodness* of Section 5.15) was completed in [HasHR], and extended to  $(P_k, H)$ -goodness for all 2-connected graphs  $H$  [Hass].

### 5.2. Wheels

Note: In this survey the wheel graph  $W_n = K_1 + C_{n-1}$  has  $n$  vertices, while some authors use the definition  $W_n = K_1 + C_n$  with  $n + 1$  vertices.

- (a)  $R(W_3, W_n) = 2n - 1$  for all  $n \geq 6$  [BuE3],  
all critical colorings for  $R(W_3, W_n)$  for all  $n \geq 3$  [Raji].
- (b) The graph  $3K_{m-1}$  is a witness of  $3m - 2 \leq R(W_m, W_n)$  for all even  $n$ , and the graph  $2K_{m-1}$  is a witness of  $2m - 1 \leq R(W_m, W_n)$  for all  $m$  and  $n$ . In Table VIII, the lower bounds without a credit are implied by these inequalities. The lower bound on  $R(W_{2k+1}, W_{2k+1})$  was improved to  $5k - O(1)$  [ZhaCh5].
- (c)  $R(W_n, W_n) \leq 8n - 10$  for even  $n$ , and  $R(W_n, W_n) \leq 6n - 8$  for odd  $n$  [MaoWMS]. For large  $n$ , these upper bounds were improved to  $16n/3 + O(1)$  and  $4n + O(1)$ , for even and odd  $n$ , respectively [ZhaCh5].

$m$	3	4	5	6	7	8	9	10
3	6	9 GG	11 Clan	11 BuE3	13 BuE3	15 BuE3	17 BuE3	19 BuE3
4		18 GG	17 He3	19 FM	21 V-L	22-26 LidP	$\geq 25$	
5			15 He2	17 FM	15 W-V-L	17 V-L	18 V-L	$\geq 21$ VO
6				17 FM	19 LidP	22-26 LidP	$\geq 25$	
7					19 V-L	19-21 LidP	$\geq 19$ VO	$\geq 21$ VO
8						22-25 LidP	$\geq 25$	
9							$\geq 21$ VO	
10								$\geq 28$

**Table VIII.** Ramsey numbers  $R(W_m, W_n)$  for  $m \leq n \leq 10$ .  
 Joint credits for the parameters (4,7), (5,7), (5,8), (5,9) and (7,7)  
 use abbreviations W/[We1], V/[VO],[VO1] and L/[LidP],[LidMPV].

- (d) All critical colorings (2, 1 and 2) for  $R(W_n, W_6)$ , for  $n = 4, 5, 6$  [FM], all critical colorings for  $R(W_5, W_7)$  and  $R(W_5, W_9)$  [LidMPV].
- (e)  $R(W_6, W_6) = 17$ ,  $R(4,4) = 18$  and  $\chi(W_6) = 4$  give a counterexample  $G = W_6$  to the Erdős conjecture (Erd2, see also [GRS]) that  $R(G, G) \geq R(K_{\chi(G)}, K_{\chi(G)})$ .
- (f) The value  $R(W_5, W_5) = 15$  was given in the Hendry's table [He2] without a proof. Later the proof was published in [HaMe2].

### 5.3. Books

The book graph  $B_n$  is defined by  $B_n = K_2 + nK_1$ .

The generalized book  $B_{k,n}$  is defined by  $B_{k,n} = K_k + nK_1$ , sometimes written as  $B_n^{(k)}$ .

For cycles versus books see Section 4.4, and books versus other graphs Section 5.9.

- (a)  $R(B_m, B_n) \leq R(B_{m+1}, B_n)$  and  $R(B_m, B_n) = R(B_n, B_m)$  hold for all  $m, n \geq 1$ .
- (b)  $R(B_1, B_n) = 2n + 3$  for all  $n > 1$  [RoS1].
- (c) Extending row 2 of Table IXa:  $R(B_2, B_{12}) = 28$ ,  $R(B_2, B_{13}) = 29$  [We2],  $R(B_2, B_n) \leq 2n + 6$  for all  $n > 1$  [RoS1],  $R(B_2, B_n) \leq 2n + 5$  for  $12 \leq n \leq 22$ ,  $R(B_2, B_n) \leq 2n + 4$  for  $23 \leq n \leq 37$ ,  $R(B_2, B_n) = 2n + 3$  for  $n \geq 38$  [FRS8].

$m$	$n$	1	2	3	4	5	6	7	8	9	10	11
1		6	7 ChH2	9 Clan	11 RoS1	13 RoS1	15 RoS1	17 RoS1	19 RoS1	21 RoS1	23 RoS1	25 RoS1
2			10 ChH1	11 Clan	13 Rou	16 RoS1	17 Rou	18 BILR	21 V-W-L	22 V-W-L	25 V-W-L	28 FRS8
3				14 RoS1	15 Sh1+	17 RoS1	19 W-L	20 W-L				
4					18 RoS1	19 We1	22 RoS1	$\leq 23$ L		27-28 5.3.lk		
5						21 RoS1	23 V-L	$\geq 25$ W-L		$\geq 28$ VO1		
6							26 5.3.g	27 V-L	29 V-W-L			
7								30 5.3.g	31 V-L	$\geq 33$ W-L	36 5.3.lk	37-38 5.3.lk
8									33 V-W-L	35 VO1	$\geq 37$ W-L	
9										38 5.3.g	39 V-W	41 V-W
10											42 5.3.g	43 V-W

**Table IXa.** Ramsey numbers  $R(B_m, B_n)$  for  $m \leq 10$  and  $m \leq n \leq 11$ .

Several credits to [We1, We2], [VO, VO1] and [LidP, LidMPV] use abbreviations W, V and L, respectively. Upper bounds in VO1 follow from 5.3.g/k. Sh1+ abbreviates [ShaXBP].

$m$	$n$	$R(B_m B_n)$	$v$	$k$	$\lambda$	$\mu$
11	11	46	45	22	10	11
14	17	64	63	30	13	15
23	26	100	99	48	22	24
22	37	120	119	54	21	27
29	38	136	135	64	28	32
34	37	144	143	70	33	35
47	50	196	195	96	46	48
46	58	210	209	100	45	50
56	56	226	225	112	55	56
38	82	244	243	110	37	60
62	65	256	255	126	61	63
69	71	281	280	135	70	60

**Table IXb.** Exact values of  $R(B_m, B_n)$  from strongly regular  $(v, k, \lambda, \mu)$ -graphs on up to 280 vertices, using 5.3.k/l [NiRo3]. It includes only the cases beyond the range of Table IXa, and excludes the cases of  $m = n$  for  $4n + 1$  prime power, as in 5.3.g.

- (d) There are 4 critical graphs for  $R(B_2, B_3)$ , a unique graph for  $R(B_3, B_4)$  [ShaXBP], 3 for  $R(B_2, B_6)$  and 65 for  $R(B_2, B_7)$  [BILR]. In 2025, all critical graphs for  $R(B_m, B_n)$  for many small parameters  $m$  and  $n$  were generated by Wesley [We2].
- (e) Unpublished result  $R(B_2, B_6) = 17$  [Rou] was confirmed in [BILR].  
 $R(B_{11}, B_{12}) = 47$  and  $R(B_{12}, B_{13}) = 51$  were obtained in [VO1].
- (f)  $4n + 1 \leq R(B_n, B_n) \leq 4n + 2$  for all  $n \leq 21$  [LidMPV],  $R(B_{14}, B_{14}) = 57$  [We2].
- (g)  $R(B_n, B_n) = 4n + 2$  for  $4n + 1$  a prime power.  
 If  $4n + 1$  is not the sum of two integer squares, then  $R(B_n, B_n) \leq 4n + 1$  [RoS1].  
 $R(B_n, B_n) = (4 + o(1))n$  [RoS1, NiRS].
- (h) Conjecture that  $R(B_{n-1}, B_n) = 4n - 1$  holds for all  $n \geq 2$ . It holds for the cases when  $2n - 1$  is a prime power (1 mod 4) [We2]. It also holds for  $n \leq 21$  [LidMPV, We2]. It was stated that  $R(B_{n-1}, B_n) \leq 4n - 1$  holds for  $n \geq 2$ , implicit in [RoS1].
- (i) Several best known lower bound constructions for diagonal and near-diagonal book numbers  $R(B_m, B_n)$  use 2-block polycirculant method [VO, VO1, We2, LidMPV]. Some are pointed to in Table IXa, others are beyond its range.
- (j)  $R(B_{n-2}, B_n) \leq 4n - 3$  if  $n \equiv 2 \pmod{3}$  [RoS1],  
 $R(B_{n-2}, B_n) \geq 4n - 3$  for  $n \leq 21$  [LidMPV],  
 $R(B_{12}, B_{14}) = 53$ ,  $R(B_{15}, B_{17}) = 65$  [We2], also implied by the above.
- (k)  $R(B_m, B_n) \leq 2(m + n + 1)$  if  $2(m + n) + 1 > (n - m)^2/3$ ,  
 $R(B_{k^2-2}, B_{k^2+1}) = 2k^2$  if  $k = 3^m 2^{m+n-1}$  for some non-negative integers  $m$  and  $n$  [RoS1].
- (l) Strongly regular graphs often provide good lower bounds. If there exists a strongly regular graph with the parameters  $(v, k, \lambda, \mu)$ , then  $R(B_{\lambda+1}, B_{v-2k+\mu-1}) \geq v + 1$ . The lower bounds for a number of specific larger cases, like  $R(B_{62}, B_{65}) = 256$  [RoS1] or  $254 \leq R(B_{37}, B_{88}) \leq 255$  [Par6], are implied by the existence of a strongly regular graph with suitable parameters. 12 exact values of  $R(B_m, B_n)$ , beyond Table IXa, where this lower bound meets the upper bound in 5.3.k were collected by Nikiforov and Rousseau [NiRo3], and they are presented in Table IXb. For a great collection of strongly regular graphs see the website by A. E. Brouwer [Brou].
- (m)  $R(B_m, B_n) = 2n + 3$  for all  $n \geq cm$  for some  $c < 10^6$  [NiRo2, NiRo3]. A strengthening of this result implying that  $R(B_{\lceil n/4 \rceil}, B_n) = (9n/4 + o(1))n$  [ChenL], and more lower and upper bounds for off-diagonal cases are in [ChenLY].
- (n) For generalized books  $B_n^{(k)} = nK_1 + K_k$ , Conlon proved that  $R(B_n^{(k)}, B_n^{(k)}) = 2^k n + o_k(n)$  [Con4]. A simplified proof, better control of the error term, and a proof that all extremal colorings for this Ramsey problem are quasirandom are in the follow-up papers: for the diagonal case [ConFW1], and off-diagonal [ConFW2]. All this more than answered some old questions by Erdős and others. The lower bound  $R(B_n^{(3)}, B_n^{(3)}) \geq 8n + 5$ , for prime powers  $4n + 1$ , was established in [LinLi3].

- (o) Using combinations of items 5.3.a-l leads to more bounds not listed in Table IXa. Other general claims involving  $R(B_m, B_n)$  can be found in [RoS1, FRS8, Par6, NiRo2, NiRo3, NiRS, LiRZ2, LidMPV, We2].

#### 5.4. Trees and forests

In this subsection  $T_n$  and  $F_n$  denote an  $n$ -vertex tree and forest, respectively.

- (a)  $R(T_n, T_n) \leq 4n + 1$  [ErdG]. Note that if  $T_n$  were a set of all  $n$ -vertex trees, then one might say that  $R(T_n, T_n) = n$ , since for every graph  $G$  at least one of  $G$  or  $\bar{G}$  is connected, and thus it contains an  $n$ -vertex spanning tree.
- (b)  $R(T_n, T_n) \geq \lfloor (4n - 1)/3 \rfloor$  [BuE2], see also Section 5.17.
- (c) In 1976, Burr and Erdős conjectured that  $R(T_n, T_n) \leq 2n - 2$  for even  $n$ , and that it is at most  $2n - 3$  for odd  $n$  [BuE2]. Note that this is the same as asking if  $R(T_n, T_n) \leq R(K_{1,n-1}, K_{1,n-1})$ . In 2011, Zhao [Zhao] proved that  $R(T_n, T_n) \leq 2n - 2$  and thus confirmed the conjecture for even  $n$ . Independently, in 2013 Ajtai et al. [AjKSS] announced a full proof for large  $n$ . In 2025, Montgomery et al. posted an arXiv paper [MonPY2] with a proof that trees with linearly bounded maximum degree have  $R(T, T) = \max\{t_1 + 2t_2, 2t_1\} - 1$ , where  $t_1 \geq t_2$  are the sizes of the bipartite classes of  $T$ . This recent progress subsumes some of the results in items (d)-(m) below.
- (d) For general discussion of related problems see [Bu7, FSS2, ChGra2], in particular of the conjecture that  $R(T_m, T_n) \leq n + m - 2$  holds for all trees [FSS2].
- (e) If  $\Delta(T_m) = m - 2$  and  $\Delta(T_n) = n - 2$  then the exact values of  $R(T_m, T_n)$  are known, and they are between  $n + m - 5$  and  $n + m - 3$  depending on  $n$  and  $m$ . In particular, we have  $R(T_n, T_n) = 2n - 5$  for even  $n$  and  $R(T_n, T_n) = 2n - 4$  for odd  $n$  [GuoV].
- (f) Examples of families  $T_m$  and  $T_n$  (including  $P_n$ ) for which  $R(T_m, T_n) = n + m - c$ ,  $c = 3, 4, 5$  [SunZ1], extending the results in [GuoV].
- (g) View the tree  $T$  as a bipartite graph with parts  $t_1$  and  $t_2$ ,  $t_2 \geq t_1$ , then define  $b(T) = \max\{2t_1 + t_2 - 1, 2t_2 - 1\}$ . Then the bound  $R(T, T) \geq b(T)$  holds always,  $R(T, T) = b(T)$  holds for many classes of trees [EFRS3, GeGy], and asymptotically [HaLT], but cases for inequality have been found [GrHK]. Lower bounds in the asymmetric cases of the form  $R(S, T) \geq b(S, T)$  based on bipartition types of  $S$  and  $T$  [Yan].
- (h) Comments in [BallS] about some conjectures on Ramsey saturation of non-star trees, which would imply that  $R(T_n, T_n) \leq 2n - 2$  holds for sufficiently large  $n$ .
- (i) Formulas for  $R(T_m, T_n)$  for some subcases of when  $T_m$  and  $T_n$  satisfy  $\Delta(T_m) = m - 3$  and  $\Delta(T_n) \geq n - 3$  [SunWW].
- (j)  $R(T_m, K_{1,n}) \leq m + n - 1$ , with equality for  $(m - 1) \mid (n - 1)$ , and for sufficiently large  $n$  for almost all trees  $T_m$  [Bu0]. Many cases were identified for which we have  $R(T_m, K_{1,n}) = m + n - 2$  [Coc, ZhZ1], see also [Bu0].
- (k)  $R(T_m, K_{1,n}) \leq m + n$  if  $T_m$  is not a star and  $(m - 1) \nmid (n - 1)$ , and some classes of trees and stars for which the equality holds [GuoV]. Further classes of trees and stars for which we have  $R(T_m, K_{1,n}) \leq m + n - c$ , and for which the equality holds with  $c = 3$  or

$c = 2$  [YanP].

- (l) In a sequence of papers [SunZ1, SunZ2, SunW, SunWW], Zhi-Hong Sun et al. obtain several exact results for  $R(S, T)$ , where the trees  $S$  and  $T$  have high maximum degree  $\Delta \geq n - 3$ , or one of them has high maximum degree and the other is a path.
- (m) Formulas for some cases of brooms [EFRS3], where broom is a star with a path attached to its center. These results were extended to all diagonal cases for brooms [YuLi]. Note that a tree  $T_n$  with  $\Delta(T_n) = n - 2$  is a broom, and this case is listed in 5.4.e.
- (n)  $R(F_n, F_n) > n + \log_2 n - O(\log \log n)$  [BuE2], forests are tight for this bound [CsKo].
- (o) Forests, linear forests (unions of paths) [BuRo2, FS3, CsKo].
- (p) Extensive tables of  $R(T_m, T_n)$  for  $6 \leq m, n \leq 8$ , for many concrete pairs of trees, which were obtained through an adiabatic quantum optimization algorithm [RanMCG].
- (q) Tristars and fountains [BroNN].
- (r) Paths versus trees [FSS2], see also other parts of this survey involving special graphs, in particular Sections 5.5, 5.6, 5.10, 5.13, 5.15 and 5.17.

### 5.5. Stars, stars versus other graphs

$R(K_{1,n}, K_{1,m}) = n + m - \varepsilon$ , where  $\varepsilon = 1$  for even  $n$  and  $m$ , and  $\varepsilon = 0$  otherwise [Har1]. This is also a special case of multicolor numbers for stars 6.6.e obtained in [BuRo1].

$R(K_{1,n}, K_m) = n(m - 1) + 1$  by Chvátal's theorem [Chv1].

Stars versus  $C_4$  [Par3, Par4, Par5, BEFRS4, Chen, ChenJ], until 2002

Stars versus  $C_4$  [GoMC, MoCa, ZhaBC1, ZhaCC2, ZhaCC3, ChenZZ7], since 2004

Stars versus  $K_{2,n}$  [Par4, GoMC]

Stars versus  $K_{n,m}$  [Stev, Par3, Par4]

See also Section 3.3.1

$R(K_{1,4}, B_4) = 11$ , some general cases for books versus stars [RoS2]

$R(K_{1,4}, K_{1,2,3}) = R(K_{1,4}, K_{2,2,2}) = 11$  [GuSL]

Stars versus paths [Par2, BEFRS2, AbdMT]

Stars versus cycles [Law1, Clark, ZhaBC6, SunSh], see also [Par6] and Section 4.1

Stars versus  $2K_2$  [MeO]

Stars versus stripes  $mP_2$  [CocL1, CocL2, Lor]

Stars versus bistars [AlmHS]

Stars versus kipsas  $K_1 + P_n$  [LiZB]

Values and bounds on  $R(S_n, K_6 - 3K_2)$  [LoM10], see also [LoM9]

Stars versus  $W_5$  and  $W_6$  [SuBa1]

$nK_{1,m}$  versus  $W_5$  [BaHA]

Stars versus  $W_9$  [Zhang2, ZhaCZ1]

Stars versus wheels [HaBA1, ChenZZ2, Kor, LiSch, HagMa]

Stars versus books [ChaRSPS, RoS2]

Stars versus fans [ZhaBC3, ZhaoW, HuP2]  
 Stars versus trees [Bu0, Cheng, Coc, GuoV, SunZ1, SunZ2, SunWW, ZhZ1, YanP]  
 Stars versus  $K_n - tK_2$  [Hua1, Hua2]  
 Stars versus almost all connected graphs on 6 vertices [LoM7], see also item 8.1.p  
 For even  $n$ ,  $K_{1,n}$  is  $H$ -good, where  $H$  is the 6-vertex Hajós graph [HeZZ]  
 Tight bounds on  $R(G, K_{1,n})$  for special graphs  $G$  [HuZC3]

Union of two stars [Gros2]  
 Double stars versus  $K_{2,q}$  and  $sK_2$  versus  $K_s + C_n$  [SuAUB]  
 Odd-linked double stars [KarK], even-linked double stars [LiuCh]  
 Asymptotics for double stars\* [NoSZ, FloS]  
 Unions of stars versus  $C_4$  and  $W_5$  [HaABS, Has, HaJu]  
 Unions of stars versus wheels [BaHA, HaBA2, SuBAU1]  
 See item 6.6.i for work on multicolor cases for double stars

## 5.6. Paths versus other graphs

Note: for cycles versus  $P_n$  see Section 4.1.2.

$P_3$  versus all isolate-free graphs [ChH2]  
 Paths versus stars [Par2, BEFRS2, AbdMT]  
 Paths versus trees [FS4, FSS2, SunZ1, SunZ2, SunWW]  
 Paths versus books [RoS2]  
 Paths versus  $K_n$  [Par1]  
 Paths versus  $2K_n$  [SuAM, SuAAM]  
 Paths versus  $K_{n,m}$  [Häg]  
 Paths versus some balanced complete multipartite graphs [Pokr]  
 Paths versus  $W_5$  and  $W_6$  [SuBa1]  
 Paths versus  $W_7$  and  $W_8$  [Bas]  
 Paths versus wheels [BaSu, ChenZZ1, SaBr3, Zhang1]  
 Paths versus wheels, the last piece completed [LiNing2]  
 $R(P_n, mW_4) = 2n + m - 2$  [Sudar1]  
 Paths versus beaded wheels [AliBT2]  
 Paths versus sunflower graphs [AliTJ]  
 Paths versus powers of paths [Pokr, AllBS, AllMRS]  
 Paths versus fans [SaBr2]  
 Paths versus kivas  $K_1 + P_m$  [SaBr1, SaBr4, LiZBBH]  
 Paths versus  $K_1 + F$ , where  $F$  is a linear forest [LiNing1]  
 Paths versus Jahangir graphs [SuTo]  
 Paths and cycles versus trees [FSS2]

---

\* double star is a union of two stars with their centers joined by an edge

- Powers of paths [AllBS, AllMRS], see also Section 5.1.e
- Unions of paths [BuRo2]
- Paths and unions of paths versus  $tK_n$  [Sudar2]
- Paths and unions of paths versus Jahangir graphs [AliBas, AliBT1, AliSur]
- Paths and unions of paths versus  $K_{2m} - mK_2$  [AliBB]
- Goodness of paths for  $tK_n$  [Sudar3]
- Goodness of paths, study of graphs  $H$  for which  $P_n$  is  $H$ -good [PoSu1, BotMds]
- Sparse graphs versus paths and cycles [BEFRS2, FanLin2]
- Graphs with long tails [Bu2, BuG]
- Long paths versus other good graphs [PeiLi, PeiCLY]
- Paths versus generalized wheels [BieDa]
- Monotone paths [Lef2, CaYZ, MuSuk5], and monotone cycles [Lef2], see also 4.5.p

**5.7. Fans, fans versus other graphs**

The fan graph  $F_n$  is defined by  $F_n = K_1 + nK_2$ ,  
 and generalized fan  $F_{k,n}$  is defined by  $F_{k,n} = K_1 + nK_k$ .  
 The fan graph  $F_n$  has  $2n + 1$  vertices, and  $F_{k,n}$  has  $kn + 1$  vertices.

- (a)  $R(F_1, F_n) = R(K_3, F_n) = 4n + 1$  for  $n \geq 2$ , and  
 bounds for  $R(F_m, F_n)$  [LiR2, GuGS, ChenYZ].
- (b)  $R(F_2, F_n) = 4n + 1$  for  $n \geq 2$ , and  
 $R(F_m, F_n) \leq 4n + 2m$  for  $n \geq m \geq 2$  [GuGS, LinLi1].
- (c)  $R(F_m, F_n) = 4n + 1$  for  $n$  sufficiently larger than  $m$  [LinLD, ZhaBC4].
- (d)  $9n/2 - 5 \leq R(F_n, F_n) \leq 11n/2 + 6$  for all  $n \geq 1$  [ChenYZ],  
 upper bound growth improved to  $R(F_n, F_n) \leq 31n/6 + 15$  for all  $n \geq 1$  [DvoMe],  
 lower bounds for  $R(F_m, F_n)$  close to the diagonal [ZhaCh5].

$m$	$n$	1	2	3	4
1		6 2.1.a	9 LiR2	13 LiR2	17 LiR2
2			9 Bu4	13 LinLi1	17 LinLi1
3				14 ZhaoW	
4					$\leq 28$ 5.7.d

**Table X.** Ramsey numbers  $R(F_m, F_n)$ , for  $1 \leq m \leq n \leq 4$ . Note that  $F_1 = K_3$ .

- (e)  $R(K_4, F_n) = 6n + 1$  for  $n \geq 3$  [SuBB3],  
 $R(K_5, F_n) = 8n + 1$  for  $n \geq 5$  [ZhaCh],  
 $R(K_6, F_n) = 10n + 1$  for  $n \geq 6$  [KaOS],  
 A conjecture that  $R(K_m, F_n) = 2mn - 2n + 1$  for  $n \geq m \geq 4$  [SuBB3].
- (f) Fans versus paths, formulas for a number of cases including  $R(P_6, F_n)$  [SaBr2],  
 missing case  $R(P_6, F_4) = 12$  solved in [Shao].  
 $R(F_m, K_n) \leq (1 + o(1))n^2/\log n$  [LiR2].  
 Lower bounds on  $R(F_2, K_n)$  from cyclic graphs for  $n \leq 9$  [Shao].
- (g) Fans versus wheels [ZhaBC5, MengZZ, ZeZZ]  
 Fans versus stars [ZhaoW, HuP2]  
 Fans versus trees and stars [ZhaBC3, Bren2]  
 Fans versus trees [SherSBO, SherBSO]  
 Fans versus generalized books  $K_k + nK_1$  [LiuLi3]  
 Fans versus cycles [Shi5]  
 Fans versus large even cycles [YouLin1]  
 Fans versus unicyclic graphs [Bren2]
- (h) Ramsey goodness of fans in general case [ZhaCh4], for sparse graphs [HuZC1], and for  
 the 6-vertex Hajós graph [HeZZ]. Ramsey goodness of fans, and more general  $K_1 + nH$   
 graphs, for complete  $p$ -partite graphs [ChuLin]. See also Section 5.15.
- (i)  $R(K_3, F_{3,n}) = 6n + 1$ , for  $n \geq 3$  [HaoLin], extended to the case  $n = 2$  [BudPra]  
 $R(K_3, F_{4,n}) = 8n + 1$ , for  $n \geq 4$  [WaQi]
- (j)  $K_m$  and other graphs versus generalized fans  $F_{k,n}$  [LiR2, WaQi]  
 $nK_m$  versus  $F_{G,n} = K_1 + nG$ , where  $n \geq 2$  and  $G$  is a graph [HaHT]  
 Asymptotics of odd cycles versus generalized fans  $F_{G,n}$  [LiuLi1]  
 Generalized fans versus special chromatic graphs  $G$  [JiHou]

### 5.8. Wheels versus other graphs

Notes: In this survey the wheel graph  $W_n = K_1 + C_{n-1}$  has  $n$  vertices, while some authors use the definition  $W_n = K_1 + C_n$  with  $n + 1$  vertices. For cycles versus  $W_n$  see Section 4.3. Consider also similarity of wheels to other graphs, like fans, kipas [LiZBBH], sunflower [AliTJ], and Jahangir graphs [SuTo].

$$R(W_5, K_5 - e) = 17 \text{ [He2][YH]}$$

$$R(W_5, K_5) = 27 \text{ [He2][RaST]}$$

$$35 \leq R(W_5, K_6) \leq 36 \text{ [ZhaCh5, LidP]}$$

$$45 \leq R(W_5, K_7) \leq 50 \text{ [VO, LidP]}$$

$$35 \leq R(W_6, K_6) \leq 40 \text{ [ZhaCh5, LidP]}$$

$$45 \leq R(W_6, K_7) \leq 55 \text{ [GoeVO, LidP]}$$

Using the blow-up technique, it was shown that for all  $n \geq 1$  we have:

$$\begin{aligned} R(W_5, K_n) &\geq 2R(K_3, K_n) - 1, \\ R(W_6, K_n) &\geq 2R(K_3, K_n) - 1, \text{ and} \\ R(W_7, K_n) &\geq 2R(K_4 - e, K_n) - 1, \end{aligned}$$

and it was conjectured that these lower bounds for  $W_5$  and  $W_7$  are tight [ZhaCh5].

$W_5$  and  $W_6$  versus stars and paths [SuBa1]

$W_5$  versus  $nK_{1,m}$  [BaHA]

$W_5$  versus unions of stars [Has]

$W_5$  versus theta graphs  $\theta_n$  [JaBVR]

$W_5$  and  $W_6$  versus trees [BaSNM]

$W_7$  and  $W_8$  versus paths [Bas]

$W_7$  versus trees  $T_n$  with  $\Delta(T_n) \geq n - 3$ , other special trees  $T$ , and  $T_n$  for  $n \leq 8$  [ChenZZ3, ChenZZ5, ChenZZ6]

$W_7$  and  $W_8$  versus trees [ChenZZ4, ChenZZ5]

$W_8$  versus trees  $T_n$  with  $\Delta(T_n) \geq n - 5$  [ChngBTW]

$W_9$  versus stars [Zhang2, ZhaCZ1, ZhaCC4, ZhaCC7]

$W_9$  versus trees of high maximum degree [ZhaCZ2]

$W_{2n}$  versus trees of high maximum degree [HafBa]

$R(W_n, C_4) = R(K_{1,n-1}, C_4)$  for  $n \geq 7$  [ZhaBC1], an alternate proof relying less on prior computational results [ChenZZ7] (in these papers the definition  $W_n = K_1 + C_n$  is used). Thus, the data in Table IVa and several items in Section 3.3.1 on  $R(C_4, K_{1,n})$  apply to wheels and stars.

Wheels versus stars [HaBA1, ChenZZ2, Kor, LiSch, HagMa]

Wheels  $W_n$ , for even  $n$ , versus star-like trees [SuBB1]

Wheels versus paths [BaSu, ChenZZ1, SaBr3, Zhang1]

Wheels versus paths, the last piece completed [LiNing2]

Wheels versus fans and wheels [ZhaBC5, MengZZ, ZeZZ]

Wheels versus trees [RaeZ, ZhuZL, ZhaChZ2]

Wheels versus books [Zhou3]

Wheels versus  $K_{2,n}$  [GupM]

Wheels versus unions of stars [BaHA, HaBA2, SuBAU1]

Wheels versus linear forests (disjoint unions of paths) [SuBa2]

Some cases of wheels versus  $K_n - K_{1,s}$  [ChaMR]

Generalized wheels versus cycles [Shi5, BieDa]

Generalized wheels  $W_{k,6}$  and  $W_{k,7}$  versus trees [Wang2, ChngTW]

Generalized wheels versus trees [WaCh, ZhaChZ1]

Upper asymptotics for  $R(W_n, K_m)$  [Song5, SonBL]

Upper asymptotics for generalized wheels versus  $K_n$  [Song9]

### 5.9. Books versus other graphs

The book graph is defined by  $B_n = K_2 + nK_1$ .

The generalized book  $B_{k,n}$  is defined by  $B_{k,n} = K_k + nK_1$ , it is also written as  $B_n^{(k)}$ .

For cycles versus books first see Section 4.4.

$$R(B_3, K_4) = 14 \text{ [He3]}$$

$$R(B_3, K_5) = 20 \text{ [He2][BaRT]}$$

$$R(B_4, K_{1,4}) = 11, \text{ some general cases for books versus stars [RoS2]}$$

Cyclic lower bounds for  $R(B_m, K_n)$  for  $m \leq 7, n \leq 9$

and for  $R(B_3, K_n - e)$  for  $n \leq 7$  [Shao, ShaoWX]

$$R(T_n, B_m) = 2n - 1 \text{ for all } n \geq 3m - 3 \text{ [EFRS7]}$$

$B_4$  versus trees [LoM8]

Trees  $T_n$  are  $2B_2$ -good for  $n \geq 5$  [GuoHP]

Books versus  $K_n$  [LiR1, Sud2]

Books versus paths [RoS2]

Books versus cycles [FRS7, FRS9]

Books versus stars [ChaRSPS, RoS2]

Books versus trees [EFRS7, ZhaCZ]

Books versus wheels [Zhou3]

Books versus  $K_2 + C_n$  [Zhou3]

Books versus multipartite graphs [BFRS, EFRS7]

Books and  $(K_1 + \text{tree})$  versus  $K_n$  [LiR1]

$B_{3,n}$  versus cycles [Shi5]

$B_{k,n}$  versus  $K_m$  [NiRo1, NiRo4]

$B_{k,n}$  versus  $K_1 + C_4$  [LinLiu]

$B_{k,n}$  versus paths and trees [RoS2, FSR]

$B_{k,n}$  versus fans [LiuLi3]

$B_{k,n}$  versus graphs with small bandwidth [YouLC]

Asymptotics of  $B_{k,n}$  versus  $C_n$  [LinP]

$C_{(2m+1)}$ -goodness of  $B_{k,n}$  and  $K_k + nH$  [LiuLi5]

$G$ -goodness of  $B_{k,n}$  for  $s(G) = 1$  and large  $n$  [LiuLi7]

A 2023 paper *Ramsey Goodness of Books Revisited* [FoxHW] reviews the subject of the title and proves a result that  $R(G, B_{k,n}) = (p-1)(n-1) + 1$  for generalized books  $B_{k,n}$  (also called  $k$ -books), where  $G$  is a complete  $p$ -partite graph with some constraints on the sizes of its parts. Some exact values for  $p = 3$  [FanHL]. See also Section 5.15.

### 5.10. Trees and forests versus other graphs

In this subsection  $T_n$  and  $F_n$  denote  $n$ -vertex tree and forest, respectively.

$$R(T_n, K_m) = (n-1)(m-1) + 1 \text{ [Chv1]}$$

$$R(T_n, K_m - e) = n + 1 \text{ if } (m=3 \text{ and } n \text{ is even and } T_n = K_{1,n-1}), \text{ otherwise}$$

$$R(T_n, K_m - e) = (n-1)(m-2) + 1 \text{ [ChaGP].}$$

$R(T_n, C_4) = \max\{4, n+1, R(C_4, K_{1,\Delta})\}$  for  $T_n$  with maximum degree  $\Delta$  [BEFRS4].  
For some related results see also [EFRS4, Bu7, Chen].

$R(T_n, C_{2m+1}) = 2n - 1$  for all  $n > 1512m + 756$  [BEFRS2].  
The range of  $n$  was extended to  $n \geq 25(2m + 1)$  in [Bren1].

$R(T_n, B_m) = 2n - 1$  for all  $n \geq 3m - 3$  [EFRS7]

$R(F_{nk}, K_m) = (n-1)(m-2) + nk$  for all forests  $F_{nk}$  consisting of  $k$  trees with  $n$  vertices each, also exact formula for all other cases of forests versus  $K_m$  [Stahl]

Exact results for almost all small ( $n(G) \leq 5$ ) connected graphs  $G$  versus all trees [FRS4]

Stripes versus trees and unicyclic graphs [HuP2]

Trees and unicyclic graphs versus fans [Bre2]

Trees versus  $K_1 + P_m$  and fans [SherSBO, SherBSO]

Large trees are  $tK_m$ -good for  $t, m \geq 3$  [LuoP2]

All Ramsey-critical graphs for large trees versus  $tK_m$  [CheLC]

Trees versus stars [Bu0, Cheng, Coc, GuoV, ZhZ1, YanP]

Trees versus unions of complete graphs of different sizes [HuLuo]

Trees versus paths [FS4, FSS2, BEFRS2]

Trees versus cycles [FSS2, EFRS6]

Trees versus odd cycles [BEFRS2, Bren1]

Trees versus odd cycles sharing one vertex [HuCh]

Trees versus books [EFRS7, ZhaCZ]

Trees versus  $B_4$  [LoM8]

Trees  $T_n$  are  $2B_2$ -good for  $n \geq 5$  [GuoHP]

Trees versus fans [ZhaBC3]

Trees versus  $W_5$  and  $W_6$  [BaSNM]

Trees versus  $W_7$  and  $W_8$  [ChenZZ4, ChenZZ5]

Some trees versus wheels [RaeZ, ZhuZL]

Trees versus wheels [ZhaBC5, ZhaChZ2]

Trees  $T_n$  with  $\Delta(T_n) \geq n - 3$ , other special trees  $T$ ,  
and  $T_n$  for  $n \leq 8$  versus  $W_7$  [ChenZZ3, ChenZZ5, ChenZZ6]

Trees  $T_n$  with  $\Delta(T_n) \geq n - 5$  versus  $W_8$  [ChngBTW]

Trees  $T_n$  with  $\Delta(T_n) \geq n - 4$  versus  $W_9$  [ZhaCZ2]

Trees  $T_n$  with large  $\Delta(T_n)$  versus  $W_{2m}$  [HafBa]

Star-like trees versus odd wheels [SuBB1, ChenZZ3]

Trees versus  $K_n + \bar{K}_m$  [RoS2, FSR]

Trees versus generalized wheels  $W_{k,6}$  and  $W_{k,7}$  [Wang2, ChngTW]

Trees versus generalized wheels [WaCh, ZhaChZ1]

Trees versus generalized books [RoS2, FSR]

Trees versus bipartite graphs [BEFRS4, EFRS6]

Trees versus almost complete graphs [GoJa2]

Trees versus multipartite complete graphs [EFRS8, BEFRSGJ]

$R(T, G)$  for most non-star trees  $T$  and  $n(G) \leq 6$  [LoM9], see item 8.1.p

Linear forests versus  $3K_3$  and  $2K_4$  [SuBAU2]

Linear forests versus  $2K_m$  [SuAAM]

Linear forests versus  $tK_n$  [Sudar2, Sudar3]

Linear forests versus wheels [SuBa2]

Forests versus almost complete graphs [ChaGP]

Forests versus complete graphs [BuE1, Stahl, BaHA]

Forests versus disjoint union of complete graphs [HuP1]

Study of graphs  $G$  for which all or almost all trees are  $G$ -good [BuF, BEFRSGJ], see also Section 5.15, 5.16 and 5.17, for the definitions and more pointers.

Ramsey goodness of bounded degree trees [BalPS], an extension of the results and confirmation of a conjecture therein [MonPY1]. See also Section 5.15.

See also various parts of this survey for special trees, and Section 5.4.

### 5.11. Unicyclic graphs

*Unicyclic* graphs have exactly one cycle. Note that the deletion of any edge from the cycle in a (connected) unicyclic graph creates a (tree)/forest. In 1979, Grossman [Gros1] conjectured that for any connected unicyclic graph  $G$  of odd girth with  $n(G) \geq 4$ , we have  $R(G, G) = 2n(G) - 1$ . Grossman's conjecture is known to hold for the following graphs  $G$ :

- (a)  $C_3$  with pendant edges emanating from its three vertices, or pendant edges emanating from two vertices and a path emanating from the third vertex [Gros1],
- (b)  $C_{2k+1}$  with pendant edges at a single vertex [Köh],
- (c)  $C_{2k+1}$  with a pendant edge at two adjacent vertices [KrRod],
- (d) Let  $C_{k*1}H$  be the graph obtained by identifying any vertex  $u$  of  $C_k$  with any degree-one vertex of  $H$ . If  $H$  is disconnected, then a degree-one vertex in each component of  $H$  is identified with  $u$ . Then for odd  $k$ , we can take  $G = C_{k*1}K_{1,n}$  for  $n \geq 1$ , or  $G = C_{k*1}(nK_2 \cup K_{1,m})$  for  $n \geq 1$  and  $m \geq 2$  [LowKap1].

Other results for Ramsey numbers involving unicyclic graphs can be found at:

- (e) Fans versus unicyclic graphs [Bren2],
- (f) Stripes versus trees and unicyclic graphs [HuP2],
- (g) Some work on the diagonal graph Ramsey numbers of even cycles with pendant edges (at a single vertex) can be found in [LowKap1, LowKap2].

For some multicolor cases for unicyclic graphs see item 6.6.k.

### 5.12. Cases for $n(G), n(H) \leq 5$

Clancy [Clan], in 1977, presented a table of  $R(G, H)$  for all isolate-free graphs  $G$  with  $n(G) = 5$  and  $H$  with  $n(H) \leq 4$ , except 5 entries. All five of the open entries have been solved as follows:

$R(B_3, K_4) = 14$	[He3]
$R(K_5, K_4 - e) = 16$	[BoH]
$R(W_5, K_4) = 17$	[He2]
$R(K_5 - e, K_4) = 19$	[EHM1]
$R(K_5, K_4) = R(4, 5) = 25$	[MR4]

An interesting case in [Clan] is:

$$R(K_4, K_5 - P_3) = R(K_4, K_4 + e) = R(4, 4) = 18$$

Hendry [He2], in 1989, presented a table of  $R(G, H)$  for all graphs  $G$  and  $H$  on 5 vertices without isolates, except 7 entries. Six of the open entries have been solved:

$R(K_5, K_4 + e) = R(4, 5) = 25$	[Ka1][MR4]
$R(K_5, K_5 - P_3) = 25$	[Ka1][Boza2, CalSR]
$R(K_5, B_3) = 20$	[He2][BaRT]
$R(K_5, W_5) = 27$	[He2][RaST]
$R(W_5, K_5 - e) = 17$	[He2][YH]
$R(K_5, K_5 - e) = 30$	[Ex6][Ang2]

The still open case for  $K_5$  versus  $K_5$  is:

$$43 \leq R(K_5, K_5) \leq 46 \quad [\text{Ex4}][\text{AnM4}]$$

All critical colorings for the case  $R(C_5 + e, K_5) = 17$  were found by Hendry [He5].

### 5.13. Miscellaneous cases

$R(P, P) \geq 19$ , where  $P$  is the 10-vertex Petersen graph [HaKr2]

$R(Q_3, Q_3) = 13$ , where  $Q_3$  is the 8-vertex cube graph [LidP], see also items 5.17.m/n/o

$30 \leq R(K_{2,2,2}, K_{2,2,2}) \leq 31$ , where  $K_{2,2,2}$  is the octahedron [HaKr2, LidP]

$K_{2,m}$  and  $C_{2m}$  versus  $K_n$  [CaLRZ]

$K_{2,n}$  versus any isolate-free graph [RoJa2, JRB]

Union of two stars [Gros2]

Double stars [GrHK, BahS, NoSZ, FloS]

Odd-linked double stars [KarK], even-linked double stars [LiuCh]

Graphs with bridge versus  $K_n$  [Li1]

Graphs with long tails [Bu2, BuG]

Formulas for some cases of brooms<sup>1)</sup> [EFRS3], extended to all diagonal cases [YuLi]  
 Lower and upper bounds for kipas<sup>2)</sup> [ZhaCh5]  
 Kipas versus linear forests [LiMSY]  
 Values, lower and upper bounds for combs<sup>3)</sup>, suns<sup>4)</sup>, and combs versus suns [XuWLM]  
 Multipartite complete graphs [BFRS, FRS3, Stev]  
 Multipartite complete graphs versus trees [EFRS8, BEFRSGJ]  
 Multipartite complete graphs versus sparse graphs [EFRS4]

**5.14. Multiple copies of graphs, disconnected graphs**

- (a)  $2K_2$  versus isolate-free graphs [ChH2],  
 $nK_2$  versus isolate-free graphs [FSS1].
- (b) Note that  $P_2 = K_2$ . In several papers graphs  $mP_2$  are called stripes.  
 $nK_2$  versus  $mK_2$ , in particular  $R(nK_2, nK_2) = 3n - 1$  for  $n \geq 1$  [CocL1, CocL2, Lor],  
 $R(mK_2 \cup nK_3, kK_2 \cup lK_3) = 2m + 3n + k + 2l - 1$  for  $m + n \geq k + l \geq 1$  [YaoHMZ].
- (c)  $R(nK_3, nK_3) = 5n$  for  $n \geq 2$ ,  $R(mK_3, nK_3) = 3m + 2n$  for  $m \geq n \geq 2$  [BES],  
 formulas for  $R(nK_3, mG)$  for all isolate-free graphs  $G$  on 4 vertices [Zeng].
- (d) Lorimer and Mullins [LorMu] proved that

$$R(nK_3, mK_4) = \begin{cases} 9 & \text{for } m = n = 1, \\ 3n + 3m + 1 & \text{for } n \geq m > 1, \\ 3n + 5 & \text{for } n \geq 2 \text{ and } m = 1, \\ 2n + 4m + 1 & \text{for } m \geq n \text{ and } m \geq 2. \end{cases}$$

- (e)  $R(nK_4, nK_4) = 7n + 4$  and  $R(nK_5, nK_5) = 9n + 16$  for large  $n$  [Bu8], see (h) below
- (f)  $R(mK_{1,3}, nK_{1,3}) = 4m + n - 1$  for  $2 \leq n \leq m$  [BES],  
 general cases for  $mK_{1,n}$  versus  $sK_{1,t}$  [LuoP1].
- (g)  $m(K_{1,3} + e)$  versus  $n(K_{1,3} + e)$  and  $m(K_4 - e)$  versus  $n(K_4 - e)$  [HuPZ]
- (h)  $R(mK_n, mK_n) \leq (2n - 1)m + c_n$  for  $n \geq 4$  and some constant  $c_n$  [BES],  
 $R(mK_s, mK_t) = m(s + t - 1) + R(K_{s-1}, K_{t-1}) - 2$  for sufficiently large  $m$  [Bu8].
- (i)  $R(nH) = (2k - \alpha)n + c$  holds for all sufficiently large  $n$ , where  $H$  is a  $k$ -vertex isolate-free graph with independence number  $\alpha = \alpha(H)$ , and  $c = c(H)$  is a constant dependent on  $H$  [BES]. Now, it is known that there exists  $C > 0$  such that this equality holds for all  $n \geq 2^{Ck}$ . Further, in the case of complete graphs we have the equality

---

1) broom is a star with a path attached to its center  
 2) kipas graph is  $K_1 + P_n$   
 3) comb is a path with pendant edges attached to all its vertices  
 4) sun is a cycle with pendant edges attached to all its vertices

$R(nK_k) = (2k - 1)n + R(K_{k-1}) - 2$  [BucSu]. Significant improvements were obtained for sparse graphs, in particular those with bounded maximum degree [SulTr].

- (j)  $R(mG, nH) \leq (m - 1)n(G) + (n - 1)n(H) + R(G, H)$  [BES]
- (k) Variety of results for numbers of the form  $R(nG, mH)$  [Bu0, BES, HaBA2, SuBAU1, SuBAU4, SuAUB, Sudar2, Sudar3].
- (l)  $R(F, G \cup H) \leq \max\{R(F, G) + n(H), R(F, G)\}$  [Par6]
- (m)  $R(G, H)$  for all disconnected isolate-free graphs  $H$  on at most 6 vertices versus all  $G$  on at most 5 vertices, except 3 cases [LoM5]. Missing cases were completed in [KroMe].
- (n) Cycles and unions of cycles versus  $mK_n$  [Sudar4],  $nK_{1,m}$  versus  $W_5$  [BaHA], cycles versus  $nW_4$  [Sudar5].
- (o) Cycles versus multiple copies of wheels,  $tW_n$  [WaZh1, WaZh2].
- (p) Results on  $R(G, G)$  for  $G = P_k \cup K_{1,n}$ : exact formulas for  $k$  equal to 2 or 3, and lower and upper bounds for  $k \geq 4$  [ZhouLMW], where these results are used as tools to derive some Gallai-Ramsey numbers.
- (q) Trees and forests versus  $K_m \cup K_n$  [HuP1],  
Trees versus unions of complete graphs of different sizes [HuLuo],  
Trees  $T_n$  are  $2B_2$ -good for  $n \geq 5$  [GuoHP],  
All Ramsey-critical graphs for large trees versus  $tK_m$  [CheLC],  
Disjoint unions of paths (linear forests) [BuRo2, FS2],  
Linear forests versus  $3K_3 \cup 2K_4$  [SuBAU2],  
Linear forests versus kipas [LiMSY].
- (r) Forests versus  $K_n$  [Stahl, BaHA] and  $W_n$  [BaHA]. Generalizations to forests versus other graphs  $G$  in terms of  $\chi(G)$  and the chromatic surplus of  $G$  [Biel4], and for linear forests versus  $2K_n$  [SuAM].
- (s) Disconnected graphs versus other graphs,  $R(nH, G)$  [BuE1, GoJa1, BucSu],  $R(nB_k, G)$  for sparse  $G$  [HuZC2],  $nK_k$ -goodness for sparse graphs [ZhaCh3].
- (t) Let  $c(nK_k)$  denote the set of connected graphs containing  $n$  vertex-disjoint  $K_k$ 's. Then  $R(c(nK_3), c(nK_3)) = 7n - 2$  for  $n \geq 2$  [GySá3], and for  $n \geq R(k, k)$  and  $k \geq 4$  we have  $R(c(nK_k), c(nK_k)) = (k^2 - k + 1)n - k + 1$  [Rob]. These are just samples of results on the so called connected Ramsey numbers, otherwise not covered in this survey.
- (u) See Section 4.1 for cases involving unions of cycles.  
See also [Bu9, BuE1, LorMu, MiSa, Den2, Biel1, Biel2].

### 5.15. Ramsey goodness of graphs

For graphs  $G$  and  $H$ , consider the equality  $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$ , where  $s(G)$  is the chromatic surplus of  $G$ , defined as the minimum number of vertices in some color class under all proper vertex colorings using  $\chi(G)$  colors [Bu2]. By canonical construction we have the lower bound. Graph  $H$  is called  $G$ -good if the equality holds. In

the special case of  $G = K_n$ , a graph  $H$  satisfying  $R(K_n, H) = (n-1)(n(H)-1)+1$  is called Ramsey  $n$ -good [BuE3]. An early summary of related results can be found in [FRS5]. In 2009, an extensive survey and study of  $n$ -goodness was written by Nikiforov and Rousseau [NiRo4], and another one in 2013 by Allen, Brightwell and Skokan [AllBS].

- (a) Graph  $H$  on  $n$  vertices is 3-good if  $R(K_3, H) = 2n - 1$ , several types of such graphs have been identified, see items 3.2.d/f/g/h.
- (b) Sparse graphs versus paths and cycles [BEFRS2], study of graphs  $G$  for which all or almost all trees are  $G$ -good [BuF, BEFRSGJ].
- (c) Goodness of paths and cycles:  $P_n$  is  $G$ -good for  $n \geq 4e(G)$  [PoSu1],  $C_n$  is  $G$ -good for  $n \geq 10^{60}e(G)$  and  $s(G) \geq \chi(G)^{22}$  [PoSu2]. These results prove a conjecture in [AllBS]. More on goodness of paths [BotMdS].
- (d) Goodness for graphs with bounded maximum degree and bounded bandwidth [AllSK].
- (e) Goodness of paths for  $tK_n$  [Sudar3], goodness of cycles for  $tK_n$  [Sudar4], long paths versus other good graphs [PeiLi, PeiCLY],
- (f) Ramsey goodness of bounded degree trees [BalPS]. An extension of the results and confirmation of a conjecture in the latter [MonPY1].
- (g) *Ramsey Goodness of Books Revisited* [FoxHW] reviews the subject and proves that  $R(G, B_{k,n}) = (p-1)(n-1)+1$  for generalized books  $B_{k,n}$  (also called  $k$ -books), where  $G$  is a complete  $p$ -partite graph with some constraints on the sizes of its parts. A conjecture on near Ramsey goodness posed in [FoxHW] is disproved in [FanLin1].
- (h) For a fixed graph  $G$  with  $\chi(G) = k+1$ ,  $s(G) = 1$ , for all  $p \geq 2$ , we have that large  $K_p + nH$  is  $G$ -good if and only if  $G$  is a subgraph of  $mK_2 + K_{k-1}(m)$ , where  $K_{k-1}(m)$  is the complete  $k-1$ -partite graph with all parts of size  $m$  [LiuLi6].
- (i) Trees  $T_n$  are  $2B_2$ -good for  $n \geq 5$  [GuoHP], large trees are  $tK_m$ -good for  $t, m \geq 3$  [LuoP2], all critical graphs [CheLC],  $C_{2m+1}$ -goodness of  $B_{k,n}$  and  $K_k + nH$  [LiuLi5],  $G$ -goodness of  $B_{k,n}$  for  $s(G)=1$  and large  $n$  [LiuLi7],  $W_m$ -goodness of  $K_{2,n}$  [GupM],  $nK_k$ -goodness of sparse graphs [ZhaCh3],  $R(nB_k, G)$  for sparse  $G$  [HuZC2].
- (j) Ramsey goodness of fans in general case [ZhaCh4], for some sparse graphs [HuZC1], and in particular for the 6-vertex Hajós graph [HeZZ]. Ramsey goodness of fans, and more general  $K_1 + nH$  graphs, for complete  $p$ -partite graphs [ChuLin].
- (k) During the last five decades, many papers on Ramsey goodness, often related to  $R(G, H)$  for large and sparse graphs  $H$ , were written. Several of them are cited in other sections of this survey: older ones ( $< 2010$ ) [Bu0, Bu5, BuG, BEFRS3, FaSi, EFRS4, EFRS5, FRS3, LiR4, Biel2, SuBAU3, Song6, LiBie, LinLiu, FRS7, LiR4, NiRo1, Song6] and newer ( $\geq 2010$ ) [Biel4, LiBie, LinLD, Sudar1, Sudar2, SuBAU3]. See also surveys and overviews listed in Section 8.4.
- (l) For some results on hypergraph concept of goodness see item 7.5.p.

**5.16. General results for special graphs**

- (a)  $R(K_m^p, K_n^q) = R(K_m, K_n)$  for  $m, n \geq 3$ ,  $m + n \geq 8$ ,  $p \leq m/(n-1)$  and  $q \leq n/(m-1)$ , where  $K_s^t$  is a  $K_s$  with an additional vertex connected to it by  $t$  edges [BEFS]. Some applications can be found in [BILR].
- (b)  $R(K_{2,k}, G) \leq kq + 1$  for  $k \geq 3$ , for isolate-free graphs  $G$  with  $q \geq 2$  edges [RoJa2, JRB].
- (c)  $R(W_6, W_6) = 17$  and  $\chi(W_6) = 4$  [FM]. This gives a counterexample  $G = W_6$  to the Erdős conjecture (see [GRS])  $R(G, G) \geq R(K_{\chi(G)}, K_{\chi(G)})$ , since  $R(4, 4) = 18$ .
- (d)  $R(G + K_1, H) \leq R(K_{1, R(G, H)}, H)$  [BuE1].
- (e)  $R(\bar{K}_2 + G, \bar{K}_2 + G) \leq 4R(G, \bar{K}_2 + G) - 2$  [LiShen].
- (f) For arbitrary fixed graphs  $G$  and  $H$ , if  $n$  is sufficiently large then we have  $R(K_2 + G, K_1 + nH) = (k + 1)mn + 1$ , where  $k = \chi(G)$  and  $m = |V(H)|$  [LiR2].
- (g) Study of  $nK_m$  versus  $F_{G,n} = K_1 + nG$ , where  $n \geq 2$  and  $G$  is a graph [HaHT]. Study of  $R(G + K_1, nH + K_1)$  [LinLD]. Further lower bounds based on the Paley graphs, in particular for  $R(K_3 + \bar{K}_n, K_3 + \bar{K}_n)$  [LinLS].
- (h)  $R(K_{p+1}, B_q^r) = p(q + r - 1) + 1$  for generalized books  $B_q^r = K_r + qK_1$ , for sufficiently large  $q$  [NiRo1]. Formula for  $R(K_1 + C_4, B_q^r)$  for sufficiently large  $q$  [LinLiu].
- (i) Study of the cases  $R(K_m, K_n - K_{1,s})$  and  $R(K_m - e, K_n - K_{1,s})$ , with several exact values for special parameters [ChaMR]. This study was extended to some cases involving  $R(K_m - K_3)$  [MonCR].
- (j) Study of  $R(T + K_1, K_n)$  for trees  $T$  [LiR1]. Asymptotic upper bounds for  $R(T + K_2, K_n)$  [Song7], see also [SonGQ].
- (k) Bounds on  $R(H + \bar{K}_n, K_n)$  for general  $H$  [LiR3]. Also, for fixed  $k$  and  $m$ , as  $n \rightarrow \infty$ ,  $R(K_k + \bar{K}_m, K_n) \leq (m + o(1))n^k / (\log n)^{k-1}$  [LiRZ1].
- (l) Asymptotics of  $R(H + \bar{K}_n, K_n)$ . In particular, the order of magnitude of  $R(K_{m,n}, K_n)$  is  $n^{m+1} / (\log n)^m$  [LiTZ]. Upper asymptotics for  $R(K_s + K_{m,n}, K_k)$  [Song9].
- (m) Study of the largest  $k$  such that if the star  $K_{1,k}$  is removed from  $K_r$ ,  $r = R(G, H)$ , any edge 2-coloring of the remaining part still contains monochromatic  $G$  or  $H$ , as for  $K_r$ , for various special  $G$  and  $H$  [HoIs].
- (n) Let  $G''$  be a graph obtained from  $G$  by deleting two vertices with adjacent edges. Then  $R(G, H) \leq A + B + 2 + 2\sqrt{(A^2 + AB + B^2)/3}$ , where  $A = R(G'', H)$  and  $B = R(G, H'')$  [LiRZ2].

**5.17. General results for sparse graphs**

- (a)  $R(K_n, T_m) = (n-1)(m-1) + 1$  for any tree  $T_m$  on  $m$  vertices [Chv1].
- (b) Graphs yielding  $R(K_n, G) = (n-1)(n(G)-1) + 1$ , called Ramsey  $n$ -good [BuE3], and related results [EFRS5]. Extensive surveys and further study of  $n$ -goodness appeared in [NiRo4, AllBS]. See Section 5.15 for more comments and pointers.

- (c)  $R(C_{2m+1}, G) = 2n - 1$  for sufficiently large sparse graphs  $G$  on  $n$  vertices, little more complicated formulas for  $P_{2m+1}$  instead of  $C_{2m+1}$  [BEFRS2].
- (d) Conjecture that  $R(G, G) \leq 12n(G)$  for all planar  $G$ , for sufficiently large  $n$  [AllBS].
- (e)  $R(G, G) \leq c_d n(G)$  for all  $G$ , where constant  $c_d$  depends only on the maximum degree  $d$  in  $G$  [ChRST]. The constant was improved in [GRR1, FoxSu1]. Tight lower and upper bounds for bipartite  $G$  [GRR2, Con2, ConFS7, ConFS8]. Further improvements of the constant  $c_d$  in general were obtained in [ConFS4], and for graphs with bounded bandwidth in [AllBS].
- (f)  $R(G, G) \leq c_d n$  for all  $d$ -arrangeable graphs  $G$  on  $n$  vertices, in particular with the same constant for all planar graphs [ChenS]. The constant  $c_d$  was improved in [Eaton]. An extension to graphs not containing a subdivision of  $K_d$  [RöTh].
- (g) Ramsey numbers grow linearly for degenerate graphs versus some sparser graphs, arrangeable graphs, crowns, graphs with bounded maximum degree, planar graphs, and graphs without any topological minor of a fixed clique [Shi3].
- (h) Study of  $L$ -sets, which are sets of graphs or pairs of graphs whose Ramsey numbers  $R(G_i, H_i)$  are linear in the number of vertices. Conjecture that Ramsey numbers grow linearly for  $d$ -degenerate graphs (graph is  $d$ -degenerate if all its subgraphs have minimum degree at most  $d$ ) [BuE1], 1973. Progress towards this conjecture was obtained by several authors, including [KoRö1, KoRö2, KoSu, FoxSu1, FoxSu2]. Further progress, and the proof of the conjecture were obtained in 2017, using its relation to the chromatic number [Lee].
- (i) Ramsey number is *linear* in a class of graphs  $X$  if  $R_X(p, q) \leq c(p+q)$  for some constant  $c$  and all  $p, q$ , where we color the edges of graphs in  $X$ . A conjecture that this linearity holds for  $X$  if and only if the co-chromatic number is bounded in  $X$  [AtLZ], and for certain hereditary classes  $X$  [AleALZ]. Discussion of various old and new classes of Ramsey linear graphs [NeOs]. A similar concept of classes of graphs for other constraints on  $R(p, q)$  are studied in [Loz].
- (j) Study of graphs  $G$ , called *Ramsey size linear*, for which there exists a constant  $c_G$  such that for all  $H$  with no isolates  $R(G, H) \leq c_G e(H)$  [EFRS9], 1993. An overview and further results were given in [BalSS], 2002, and more recently in [BraGS], 2024.
- (k)  $R(G, G) < 6n$  for all  $n$ -vertex graphs  $G$ , in which no two vertices of degree at least 3 are adjacent [LiRS]. This improves the result  $R(G, G) \leq 12n$  in [Alon1]. In an early paper by Burr and Erdős [BuE1] it was proved that if any two points of degree at least 3 are at distance at least 3 then  $R(G, G) \leq 18n$ .
- (l)  $R(G_{a,b}, G_{a,b}) = (3/2 + o(1))ab$ , where  $G_{a,b}$  is the rectangular  $a \times b$  grid graph. Other similar results follow for bipartite planar graphs with bounded degree and grids of higher dimension [MoSST].
- (m)  $R(Q_n, Q_n) \leq 2^{2.62n + o(n)}$ , for the  $n$ -dimensional hypercube  $Q_n$  with  $2^n$  vertices [Shi1]. This bound can also be derived from a theorem in [KoRö1]. An improvement was obtained in [Shi4], a further one to  $R(Q_n, Q_n) \leq 2^{2n+5n}$  in [FoxSu1], then another decrease of the upper bound to  $2^{2n+6}$  in [ConFS8], and the latest one to  $2^{2n-cn+1} + 2$ ,

for some positive  $c$  and sufficiently large  $n$  [Tikh].

- (n) A lower bound construction for  $12 \leq R(Q_3, Q_3)$  was presented in [HaKr2], and  $R(Q_3, Q_3) = 13$  was shown in [LidP]. This is not to be confused with the poset Ramsey numbers for  $Q_n$  (studied in [BohP], for example, but otherwise not covered in this survey), where the vertices of  $Q_m$  are colored and monochromatic copies of  $Q_n$  are being avoided.
- (o)  $R(K_m, Q_n) = (m-1)(2^n - 1) + 1$  for every fixed  $m$  and sufficiently large  $n$  [FizGMSS]. This improves on the results in [ConFLS] and [GrMFSS]. The apparent contradiction with publication years is due to the timing of publication processes.
- (p)  $R(nB_k, G)$  for sparse  $G$  [HuZC2],  $nK_k$ -goodness for sparse graphs [ZhaCh3].
- (q) Using SAT to compute  $R(G, G)$  for several small sparse graphs  $G$  [LowKKB].
- (r) See also earlier subsections 5.\* for various specific sparse graphs.

### 5.18. General results

- (a)  $R(G, H) \geq (\chi(G) - 1)(c(H) - 1) + 1$ , where  $\chi(G)$  is the chromatic number of  $G$ , and  $c(H)$  is the size of the largest connected component of  $H$ . [ChH2].
- (b)  $R(G, G) > (s 2^{e(G)-1})^{1/n(G)}$ , where  $s$  is the number of automorphisms of  $G$ . Hence  $R(K_{n,n}, K_{n,n}) > 2^n$ , see also item 6.7.m [ChH3].
- (c)  $R(G, G) \geq \lfloor (4n(G) - 1)/3 \rfloor$  for any connected  $G$ , and  $R(G, G) \geq 2n - 1$  for any connected nonbipartite  $G$ . These bounds can be achieved for all  $n \geq 4$  [BuE2].
- (d) Graph  $G$  is Ramsey saturated if  $R(G + e, G + e) > R(G, G)$  for every edge  $e$  in  $\bar{G}$ . The paper [BallS] contains several theorems involving cycles, cycles with chords and trees on Ramsey saturated and unsaturated graphs, and also seven conjectures including one stating that almost all graphs are Ramsey unsaturated. Some classes of graphs were proved to be Ramsey unsaturated [Ho]. Special cases involving cycles and Jahangir graphs were studied in [AliSur]. Discussion of how much  $R(G, G)$  can change if one vertex from  $G$  is removed [Wig2].
- (e) Relations between  $R(3, k)$  and graphs with large  $\chi(G)$  [BiFJ]. Further detailed study of the relation between  $R(3, k)$  and the chromatic gap [GySeT].
- (f)  $R(G, H) > h(G, d)n(H)$  for all nonbipartite  $G$  and almost every  $d$ -regular  $H$ , for some  $h$  unbounded in  $d$  [Bra3].
- (g) Lower asymptotics of  $R(G, H)$  depending on the average degree of  $G$  and the size of  $H$  [DoLL1]. This continued the study initiated in [EFRS5], later much enhanced for both lower and upper bounds in [Sud3].
- (h) Lower bound asymptotics of  $R(G, H)$  for large dense  $H$  [LiZa1].
- (i) Upper bounds on  $R(G, K_n)$  for dense graphs  $G$  [Con3].
- (j) A conjecture posed by Erdős in 1983 that there exists a constant  $c$  such that  $R(G, G) \leq 2^{c\sqrt{e(G)}}$  for all isolate-free graphs  $G$  [Erd4]. Discussion of this conjecture and partial results, proof for bipartite graphs and progress in other cases are included in

[AIKS]. In 2011, Sudakov [Sud5] completed the proof of this conjecture. An extension of the latter to some off-diagonal cases is presented in [MaOm1], and an improvement of the constant for bipartite graphs is given in [JoPe]. For the multicolor case see item 6.7.1.

- (k) Lower bound on  $R(G, K_n)$  depending on the density of subgraphs of  $G$  [Kriv]. This construction for  $G = K_m$  produces a bound similar to the best known probabilistic lower bound by Spencer [Spe2]. Further lower and upper bounds on  $R(G, K_n)$  in terms of  $n$  and  $e(G)$  can be found in [Sud3, HngJL].
- (l) The graphs  $K_n$  and  $K_n + K_{n-1}$  are Ramsey equivalent for  $n \geq 4$ , i.e. every graph arrows both of them or neither of them. This equivalence does not hold for  $n = 3$ , and every graph witnessing such nonequivalence contains  $K_6$  [BILi]. See references therein for history and further results on Ramsey equivalent and nonequivalent pairs of graphs.
- (m) Relations between the cases of  $G$  or  $G + K_1$  versus  $H$  or  $H + K_1$  [BuE1].
- (n) Study of cyclic graphs yielding lower bounds for Ramsey numbers. Exact formulas for paths and cycles, and values for small complete graphs and for graphs with up to five vertices [HaKr1].
- (o) Relations between some Ramsey graphs and block designs [Par3, Par4].
- (p) Ramsey numbers for line graphs and perfect graphs [BeHHS], with generalization and extensions to other graph classes, in particular to circular interval graphs [BeHHRS].
- (q) Lidický and Pfender used flag algebras to constrain the space of feasible Ramsey colorings of various types. This was implemented, and then led to a number of new upper bounds listed throughout this survey [LidP].
- (r) Relations between the Shannon capacity of noisy communication channels and graph Ramsey numbers [Li2]. See also Section 6 in [Ros2], and [XuR3].
- (s) Given integer  $m$  and graphs  $G$  and  $H$ , determining whether  $R(G, H) \leq m$  holds is NP-hard [Bu6]. Further early complexity results related to Ramsey theory were presented by Burr in [Bu10].
- (t) Complexity classification of  $(P_k, P_l)$ -goodness (i.e. the existence question of  $(P_k, P_l)$ -colorings for input graphs, which is distinct from the *goodness* of Section 5.15) was completed in [HasHR], and then extended to  $(P_k, H)$ -goodness for all 2-connected graphs  $H$  [Hass].
- (u) Ramsey arrowing is  $\Pi_2^P$ -complete, a rare natural example of a problem higher than NP in the polynomial hierarchy of computational complexity theory [Scha]. Superpolynomial lower bound on the length of resolution proofs that a graph is Ramsey is obtained in [LauPRT].
- (v) Graph sequences for Ramsey numbers with prescribed rate of growth [PavPS]. A conjecture in the latter involving in addition connectivity was proved in [AhS].
- (w) Special cases of multicolor results listed in Section 6. See also surveys listed in Section 8.

### 6. Multicolor Ramsey Numbers

Until 2016, the only known value of a multicolor classical Ramsey number was:

$$R_3(3) = R(3, 3, 3) = R(3, 3, 3; 2) = 17 \quad \text{[GG]}$$

2 critical colorings (on 16 vertices) [KaSt, LayMa]  
 2 colorings on 15 vertices [Hein]  
 115 colorings on 14 vertices [PR1]

Now, we know one more case, namely  $R(3, 3, 4) = 30$ , for some details see 6.1.b.

#### 6.1. Bounds for classical numbers

General upper bound, implicit in [GG]:

$$R(k_1, \dots, k_r) \leq 2 - r + \sum_{i=1}^r R(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_r) \quad (a)$$

The inequality in (a) is strict if the right hand side is even and at least one of the terms in the summation is even. It is suspected that this upper bound is never tight for  $r \geq 3$  and  $k_i \geq 3$ , except for  $r = k_1 = k_2 = k_3 = 3$ . However, only some cases are known to improve over (a), namely:  $R_4(3) \leq 62$  [FeKR],  $R(3, 3, 4) \leq 31$  [PR1, PR2],  $R(3, 3, 4) \leq 30$  [CodFIM], and more recently  $R(3, 3, 6) \leq 91$ ,  $R(3, 4, 5) \leq 157$  and  $R_3(4) \leq 229$  using a version of (a) which is based on mod 3 instead of parity [Boza9].

#### 6.1.1. Diagonal Cases

$m$	3	4	5	6	7	8	9	10
$r$								
3	17 GG	128 HiIr	454 Ex23	1106 Row3	3214 XuR1	7174 Row5	15041 Row5	23094 Row5
4	51 Chu1	634 XXER	4073 Row3	23502 Row5	94874 Row5	182002 Row5	719204 Row5	
5	162 Ex10	4176 Row1	41626 Row5	258506 Row5				
6	538 FreSw	32006 Row1	441606 Row5					
7	1698 Row4	160024 Row1						
8	5288 Row3							
9	17805 AgCP+							

**Table XIa.** Known lower bounds for small parameter diagonal multicolor Ramsey numbers  $R_r(m)$ , with references.

A general construction of linear Ramsey graphs as described by Rowley [Row2, Row3] in 2020 leads to lower bounds in higher cases, such as  $R_6(6) \geq 4515702$ . The templates by Rowley were generalized, which yielded many new lower bound constructions for Schur numbers, which in turn give lower bounds for Ramsey numbers  $R_k(3)$  for  $10 \leq k \leq 15$  [AgCP+]. Other lower bounds, implied by general constructions such as those in Section 6.2, are not listed.

The most studied and intriguing open case is

$$[\text{Chu1}] \quad 51 \leq R_4(3) = R(3, 3, 3, 3) \leq 62 \quad [\text{FeKR}]$$

The construction for  $51 \leq R_4(3)$  as described in [Chu1] is correct, but be warned of a typo found by Christopher Frederick in 2003 (there is a triangle (31, 7, 28) in color 1 in the displayed matrix). It was shown that the bound 51 cannot be improved by using group partitioning into disjoint union of symmetric product-free sets [Ana1], neither by some other group partition means [Ana2]. The inequality 6.1.a implies  $R_4(3) \leq 66$ , Folkman [Fol] in 1974 improved this bound to 65, and Sánchez-Flores [Sán] in 1995 proved  $R_4(3) \leq 64$ .

The upper bounds in  $162 \leq R_5(3) \leq 307$ ,  $538 \leq R_6(3) \leq 1838$ ,  $1698 \leq R_7(3) \leq 12861$ ,  $128 \leq R_3(4) \leq 230$  and  $634 \leq R_4(4) \leq 6306$  are implied by 6.1.a (we repeat lower bounds from Table XIa just to see easily the ranges). All the latter and other upper bounds obtainable from known smaller bounds and 6.1.a can be computed with the help of a LISP program written by Kerber and Rowat [KerRo]. In 2026, Boza improved the upper bound on  $R_3(4)$  by 1 to 229 ([Boza9], see also 6.1.a). In addition, the bound  $R_4(4) \leq 6301$  is implicit in [Boza9].

In general, we have:

$$c^{rk} \leq R_r(k) \leq e^{-\delta k} r^{rk}$$

for some positive constant  $\delta = \delta(r)$ ,  $c > 1$ , and large  $k$ . The lower bound follows from an early 1947 paper by Erdős [Erd1] by applying 6.2.p [Lef1], while the upper bound is a recent improvement from 2024 [BalB+] to classics. See also other items in Section 6.2.

### 6.1.2. Off-diagonal Cases

Three colors:

$m$	4	5	6	7	8	9	10	11	12	13	14	15	16
$k$													
3	30 Ka2	45 Ex2	61 ExT	85 Ex18	103 Ex18	129 Ex18	150 ExT	174 ExT	194 ExT	217 ExT	242 ExT	269 ExT	291 ExT
4	55 KrLR	89 Ex17	117 Ex17	152 ExT	193 6.2.g	242 ExT							
5	89 Ex17	139 Ex17	181 Ex17	241 6.2.g									

**Table XIb.** Known nontrivial lower bounds for 3-color Ramsey numbers of the form  $R(3, k, m)$ , with references. See also 6.1.c/d/e/f below.

- (b) In several past revisions of this survey we wrote: "The other most studied, and perhaps the only open case of a classical multicolor Ramsey number, for which we can anticipate exact evaluation in the not-too-distance future is

$$[\text{Ka2}] \quad 30 \leq R(3, 3, 4) \leq 31 \quad [\text{PR1}, \text{PR2}]$$

In [PR1] it was conjectured that  $R(3, 3, 4) = 30$ , and the results in [PR2] eliminate some cases which could give  $R(3, 3, 4) = 31$ . Since 2016, we can write that  $R(3, 3, 4) = 30$  due to the computations completed by Codish, Frank, Itzhakov and Miller [CodFIM].

- (c) In addition to Table XIb, the bounds  $303 \leq R(3, 6, 6)$ ,  $609 \leq R(3, 7, 7)$  and  $1689 \leq R(3, 9, 9)$  were derived in [XXER] (used there for building other lower bounds for some diagonal cases). These three bounds were improved to 338, 674 and 1844, respectively, by Rowley [Row5].
- (d) The upper bounds in the inequalities  $45 \leq R(3, 3, 5) \leq 57$ ,  $55 \leq R(3, 4, 4) \leq 77$  and  $89 \leq R(3, 4, 5) \leq 158$  are implied by 6.1.a. We repeat lower bounds from Table XIb to show explicitly the current ranges. In 2026, Boza obtained the upper bounds  $R(3, 3, 6) \leq 91$ ,  $R(3, 4, 5) \leq 157$  and  $R_3(4) \leq 229$  using a version of 6.1.a, which is based on mod 3 instead of parity of its parts.
- (e) In 2015, Exoo and Tatarevic obtained several lower bounds improvements which are marked as [ExT] in Table XIb. The same paper improves also on several classical two-color cases in Table Ia, see also comments 2.1.p and 2.1.q.
- (f) For three colors, Rowley [Row5] gives also the lower bounds:

$$\begin{array}{ll} 338 \leq R(3, 6, 6) & 674 \leq R(3, 7, 7) \\ 941 \leq R(3, 8, 8) & 1844 \leq R(3, 9, 9) \\ 2841 \leq R(3, 10, 10) & 10769 \leq R(8, 8, 9) \end{array}$$

- (g) The lower bound  $3214 \leq R(7, 7, 7)$  [XuR1] is based on Mathon's cyclotomic colorings of Paley graphs. The idea of Mathon's colorings can be applied to the  $k$ -th power Paley digraphs, which leads to many new lower bounds on multicolor directed Ramsey numbers [McCM].

- (h) Four colors:

$$\begin{array}{ll} 97 \leq R(3, 3, 3, 4) \leq 149 & [\text{Ex17}], 6.1.a \\ 174 \leq R(3, 3, 4, 4) \leq 450 & [\text{Row2}], 6.1.a \\ 381 \leq R(3, 4, 4, 4) \leq 1576 & 6.2.j, \text{ implicit in } [\text{Boza9}] \\ 162 \leq R(3, 3, 3, 5) & [\text{XXER}] \\ 513 \leq R(3, 3, 3, 10) & 6.2.g \\ 597 \leq R(3, 3, 3, 11) & 6.2.g \\ 729 \leq R(3, 4, 5, 5) & [\text{Row2}] \\ 1430 \leq R(3, 5, 5, 5) & [\text{Row5}] \end{array}$$

Lower bounds for higher numbers can be obtained by using general constructive results from Section 6.2 below. For example, the bounds  $261 \leq R(3, 3, 15)$  and  $247 \leq R(3, 3, 3, 7)$  were not published explicitly but are implied by 6.2.g and 6.2.h, respectively.

## 6.2. General results for complete graphs

- (a)  $R(k_1, \dots, k_r) \leq 2 - r + \sum_{i=1}^r R(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_r)$  [GG].
- (b)  $R_r(3) \geq 3R_{r-1}(3) + R_{r-3}(3) - 3$  [Chu1].
- (c)  $R_r(m) \geq c_m(2m-3)^r$ , and some slight improvements of this bound for small values of  $m$  were described in [AbbH, Gi1, Gi2, Song2]. For  $m = 3$ , the best known lower bound is  $R_r(3) \geq (3.28\dots)^r$  [AgCP+].
- (d)  $R_r(3) \leq r!(e - e^{-1} + 3)/2 \approx 2.67r!$  [Wan] improved over the classical upper bound  $3r!$  in [GG, GRS]. This was further improved to  $R_r(3) \leq r!(e - 1/6) + 1 \approx 2.55r!$  for all  $r \geq 4$  [XuXC]. Drawing from the latter, further conditional upper bounds depending on the value of  $R_4(3)$  were obtained in [Eli]. In particular, assuming that  $R_4(3) = 51$ , we have  $R_r(3) \leq r!(e - 5/8) + 1 \approx 2.09r!$  for all  $r \geq 4$ .
- (e) The limit  $L = \lim_{r \rightarrow \infty} R_r(3)^{1/r}$  exists, though it can be infinite [ChGri]. It is known that  $3.28\dots < L$ , as implied by (c) above. The lower bounds on the limits  $\lim_{r \rightarrow \infty} R_r(k)^{1/r}$  for small fixed  $k$  are gathered in [Row1, Row3, Row5], see also 6.2.v. The best lower bounds for  $R_r(k)$  from the  $k$ -th residue Paley graphs for  $k=3$  and  $k=4$  are described in [DawMc], though they are much weaker than those in Table XIa. Detailed analysis of upper bounds on  $R_r(3)$  and  $R_r(4)$  for small  $r$ , but also in general for  $R_r(k)$ , is interesting, but it does not improve on previous bounds. For some older related results, mostly on the asymptotics of  $R_r(3)$ , see [AbbH, Fre, Chu2, GRS, GrRö].
- (f) In 2020, the limit  $L$  (as in the previous item e) was studied by Fox, Pach and Suk [FoxPS1] assuming a conjecture for multicolorings with bounded VC-dimension, and further for  $\lim_{r \rightarrow \infty} R_r(k)^{1/r}$  when restricted to the so-called semi-algebraic colorings [FoxPS2]. In 2024, Suk and Zeng [SukZ] established bounds on  $L$  in terms of hypergraph numbers  $R(P_{n+2}, I_n; 3)$ , where  $I_n$  is a set of ordered 3-uniform monotone paths with  $n$  jumps, see item 7.2.1.1.
- (g)  $R(3, k, l) \geq 4R(k, l-1) - 3$  for  $k \geq 3, l \geq 5$ , and in general for  $r \geq 2$  and  $k_i \geq 2$  it holds  $R(3, k_1, \dots, k_r) \geq 4R(k_1 - 1, k_2, \dots, k_r) - 3$  for  $k_1 \geq 5$ , and  $R(k_1, 2k_2 - 1, k_3, \dots, k_r) \geq 4R(k_1 - 1, k_2, \dots, k_r) - 3$  for  $k_1 \geq 5$  [XuX2, XXER].
- (h)  $R(3, 3, 3, k_1, \dots, k_r) \geq 3R(3, 3, k_1, \dots, k_r) + R(k_1, \dots, k_r) - 3$  [Rob2].  
For  $r+1$  colors, avoiding  $K_3$  in the first  $r$  colors and avoiding  $K_m$  in the last color,  $R(3, \dots, 3, m) \leq r!m^{r+1}$  [Sár1].
- (i)  $R(k_1, \dots, k_r) \geq S(k_1, \dots, k_r) + 2$ , where  $S(k_1, \dots, k_r)$  is the generalized Schur number [AbbH, Gi1, Gi2]. In particular, the special case  $k_1 = \dots = k_r = 3$  has been widely studied [Fre, FreSw, Ex10, Rob3, Row4, AgCP+].
- (j)  $R(k_1, \dots, k_r) \geq L(k_1, \dots, k_r) + 1$ , where  $L(k_1, \dots, k_r)$  is the maximal order of any cyclic  $(k_1, \dots, k_r)$ -coloring, which can be considered a special case of Schur partitions defining

(symmetric) Schur numbers. Many lower bounds for Ramsey numbers were established by cyclic colorings. The following recurrence can be used to derive lower bounds for higher parameters. In 1968, Giraud [Gi2] showed that for  $k_i \geq 3$  we have

$$L(k_1, \dots, k_r, k_{r+1}) \geq (2k_{r+1} - 3)L(k_1, \dots, k_r) - k_{r+1} + 2.$$

In 2017, this was superseded by a result obtained by Rowley [Row1]:

$$L(k_1, \dots, k_{p+q}) \geq ((2L(k_1, \dots, k_p) - 1)(2L(k_1, \dots, k_q) - 1) + 1)/2.$$

It applies not only to cyclic but also to general linear graphs. In many cases it can be surpassed by constructions involving special templates [Row3], which in turn are further enhanced for Schur numbers [AgCP+].

- (k)  $R_r(m) \geq p + 1$  and  $R_r(m + 1) \geq r(p + 1) + 1$  if there exists a  $K_m$ -free cyclotomic  $r$ -class association scheme of order  $p$  [Math].
- (l) If the quadratic residues Paley graph  $Q_p$  of prime order  $p = 4t + 1$  contains no  $K_k$ , then  $R(s, k + 1, k + 1) \geq 4ps - 6p + 3$  [XXER].
- (m)  $R_r(pq + 1) > (R_r(p + 1) - 1)(R_r(q + 1) - 1)$  [Abb1]
- (n)  $R_r(pq + 1) > R_r(p + 1)(R_r(q + 1) - 1)$  for  $p \geq q$  [XXER]
- (o)  $R(p_1q_1 + 1, \dots, p_rq_r + 1) > (R(p_1 + 1, \dots, p_r + 1) - 1)(R(q_1 + 1, \dots, q_r + 1) - 1)$  [Song3]
- (p)  $R_{r+s}(m) > (R_r(m) - 1)(R_s(m) - 1)$  [Lef1, Song2]
- (q)  $R(k_1, k_2, \dots, k_r) > (R(k_1, \dots, k_i) - 1)(R(k_{i+1}, \dots, k_r) - 1)$  in [Song1], see [XXER].
- (r)  $R(k_1, k_2, \dots, k_r) > (k_1 + 1)(R(k_2 - k_1 + 1, k_3, \dots, k_r) - 1)$  [Rob4]
- (s) Further lower bound constructions, though with more complicated assumptions, were presented in [XuX2, XXER].
- (t) Grolmusz [Grol1] generalized the classical constructive lower bound by Frankl and Wilson [FraWi] (item 2.4.7) to more colors and to hypergraphs [Grol3] (item 7.5.m).

All lower bounds in (b) through (t) above are constructive. Item (h) generalizes (b), (o) generalizes both (m) and (q), and (q) generalizes (p). (n) is stronger than (m). Finally, we note that the construction in (o) with  $q_1 = \dots = q_i = 1 = p_{i+1} = \dots = p_r$  is the same as (q).

- (u)  $R(n, n, n) \leq R(n - 2, n, n) + 8R(n - 1, n - 1, n) - 6$  for  $n \geq 3$  [HTHZ2].
- (v) A conjecture that  $R(k_1, k_2, \dots, k_r) \geq R(k_1, k_2, \dots, k_{r-2}, k_{r-1} - 1, k_r + 1)$  holds for all  $k_r \geq k_{r-1} \geq 3$  (called DC), its implications, evidence for validity, and related problems [LiaRX]. For two-color case see also item 2.3.f. If we set  $L_k = \lim_{r \rightarrow \infty} R_r(k)^{1/r}$ , then the limit  $L_k$  exists, finite or infinite, for every  $k \geq 3$  [ChGri]. If DC holds, then all  $L_k$ 's are finite or all of them are infinite, and if  $L_3$  is finite, then  $L_k < L_{k+1}$  for all  $k \geq 3$  [LiaRX]. See also item 6.2.e.
- (w) In 2020, Conlon and Ferber [ConFer] showed constructively that  $R_3(k) > 2^{7k/8 + o(k)}$  and  $R_4(k) > 2^{k/2} 3^{3k/8 + o(k)}$ , and they discussed more general best known lower and upper bounds on  $R_r(k)$ . An improvement to their construction by Wigderson [Wig1] yields  $R_r(k) \geq (2^{3r/8 - 1/4})^{k - o(k)}$ , for any fixed  $r \geq 2$ , and a further improvement by Sawin

[Saw] to 0.383796 in the exponent in place of  $3/8$  ( $= 0.375$ ). In 2026, two more improvements were posted on arXiv [CamP, AttLM].

- (x) Exact asymptotics of a very special but important case is known, namely we have  $R(3, 3, n) = \Theta(n^3 \text{poly}\text{-}\log n)$  [AlRö]. Generalizations to other parameters and more colors are studied in [HeWi].
- (y) Mattheus and Verstraëte [MatVer] established the lower bound, and He and Wigderson [HeWi] the upper bound, as in

$$c_1 k^5 / \log^8 k \leq R(4, 4, k) \leq c_2 k^5 / \log^4 k,$$

see also 6.6.o/p. Generalizations to other parameters and more colors are discussed in [HeWi]. For earlier results on general upper bounds and more asymptotics see [Chu4, ChGra2, ChGri, GRS, GrRö].

### 6.3. Cycles

Note: The paper *Ramsey Numbers Involving Cycles* [Ra4] is based on the revision #12 of this survey. It collects and comments on the results involving cycles versus any graphs, in two or more colors. It contains some more details than this survey, but only until 2009.

#### 6.3.1. Three colors

- (a) One long cycle.

The first larger paper in this area by Erdős, Faudree, Rousseau and Schelp [EFRS1] appeared in 1976. It gives several formulas and bounds for  $R(C_m, C_n, C_k)$  and  $R(C_m, C_n, C_k, C_l)$  for large  $m$ . For three colors [EFRS1] includes:

$$\begin{aligned} R(C_m, C_{2p+1}, C_{2q+1}) &= 4m - 3 \quad \text{for } p \geq 2, q \geq 1, \\ R(C_m, C_{2p}, C_{2q+1}) &= 2(m+p) - 3 \quad \text{and} \\ R(C_m, C_{2p}, C_{2q}) &= m + p + q - 2 \quad \text{for } p, q \geq 1 \quad \text{and large } m. \end{aligned}$$

- (b) Triple even cycles.

$R_3(C_{2m}) \geq 4m$  for all  $m \geq 2$  [DzNS], see also 6.3.2.d/e/f. It was proven that  $R(C_n, C_n, C_n) = (2 + o(1))n$  for even  $n$  [FiŁu1, GyRSS], which was improved to exactly  $2n$ , for large  $n$ , by Benevides and Skokan [BenSk]. In 2005, Dzido [Dzi1] conjectured that  $R_3(C_{2m}) = 4m$  for all  $m \geq 3$ . The first open case is for  $R_3(C_{10})$ , known to be at least 20. A more general result holds for some off-diagonal cases [FiŁu1]:

$$\begin{aligned} R(C_{2\lfloor \alpha_1 n \rfloor}, C_{2\lfloor \alpha_2 n \rfloor}, C_{2\lfloor \alpha_3 n \rfloor}) = \\ (\alpha_1 + \alpha_2 + \alpha_3 + \max\{\alpha_1, \alpha_2, \alpha_3\} + o(1))n, \quad \text{for all } \alpha_1, \alpha_2, \alpha_3 > 0. \end{aligned}$$

The conjectured equality  $R_3(C_{2m}) = 4m$ , whenever true, implies  $R_3(P_{2m+1}) = 4m + 1$  [DyDR] (see also Section 6.4). For general mixed-parity case see 6.3.1.d/e below.

Triple odd cycles.

Bondy and Erdős conjectured that  $R(C_n, C_n, C_n) \leq 4n - 3$  for all  $n \geq 4$  (see for example [Erd2]). If true, then for all odd  $n \geq 5$  we have  $R(C_n, C_n, C_n) = 4n - 3$ . The first open case is for  $R_3(C_9)$ , known to be at least 33. Erdős [Erd3] and other authors credit this conjecture to Bondy and Erdős, often pointing to a 1973 paper [BoEr]. Interestingly, however, the conjecture is not mentioned in this paper.

$m n k$	$R(C_m, C_n, C_k)$	references	general results
<b>3 3 3</b>	<b>17</b>	GG	2 critical colorings [KaSt, LayMa]
3 3 4	17	ExRe	
3 3 5	21	Sun1+/Tse3	$5k - 4$ for $k \geq 5$ , $m = n = 3$ [Sun1+]
3 3 6	26	Sun1+	
3 3 7	31	Sun1+	
3 4 4	12	Schu	
3 4 5	13	Sun1+/Rao/Tse3	
3 4 6	13	Sun1+/Tse3	
3 4 7	15	Sun1+/Tse3	
3 5 5	17	Tse3/LidP	
3 5 6	21	Sun1+	
3 5 7	25	Sun1+	
3 6 6	15	LidP/We3	
3 6 7	21	Sun1+	
3 7 7			
<b>4 4 4</b>	<b>11</b>	BiaS	1000 critical colorings [Ra4]
4 4 5	12	Sun2+/Tse3	
4 4 6	12	Sun2+/Tse3	$k + 2$ for $k \geq 11$ , $m = n = 4$ [Sun2+]
4 4 7	12	Sun2+/Tse3	values for $k = 8, 9, 10$ are 12, 13, 13 [Sun2+]
4 5 5	13	Tse3	
4 5 6	13	Sun1+	
4 5 7	15	Sun1+	
4 6 6	11	Tse3	
4 6 7	13	Sun1+/Tse3	
4 7 7			
<b>5 5 5</b>	<b>17</b>	YR1	1701746176 critical colorings [Nar]
5 5 6	21	Sun1+	
5 5 7	25	Sun1+	
5 6 6	15	LidP/We3	
5 6 7	21	Sun1+	
5 7 7			
<b>6 6 6</b>	<b>12</b>	YR2	$R_3(C_{2q}) \geq 4q$ for $q \geq 2$ [DzNS]
6 6 7	15	Sun1+	see 6.3.1.a for larger parameters
6 7 7			see 6.3.1.a for larger parameters
<b>7 7 7</b>	<b>25</b>	FSS3	$R_3(C_{2q+1}) = 8q + 1$ for large $q$ [KoSS1, KoSS2]
<b>8 8 8</b>	<b>16</b>	Sun/SunY	$R_3(C_{2q}) = 4q$ for large $q$ [BenSk]

**Table XII.** Ramsey numbers  $R(C_m, C_n, C_k)$  for  $m, n, k \leq 7$  and  $m = n = k = 8$ ; Sun1+ abbreviates SunYWLX, Sun2+ abbreviates SunYLZ2, the work in [SunYWLX] and [SunYLZ2] is independent from [Tse3].

Łuczak proved that  $R(C_n, C_n, C_n) \leq (4+o(1))n$ , with equality for odd  $n$  [Łuc]. The result  $R_3(C_{2m+1}) = 8m+1$  for all sufficiently large  $m$ , or equivalently  $R(C_n, C_n, C_n) = 4n-3$  for large odd  $n$ , was announced with an outline of the proof by Kohayakawa, Simonovits and Skokan [KoSS1], followed by the full proof in [KoSS2].

(d) Three mixed-parity cycles.

Ferguson [Ferg] shows that  $R(C_m, C_n, C_k) = \max\{2m+n, 2n+m, (n+m)/2+k-2\}$ , for all  $m, n, k$  sufficiently large, which generalizes and improves on all even case in [FiŁu1]. The reference [Ferg] consists of a Ph.D. thesis and three long arXiv preprints.

(e) Asymptotics for triples of cycles of mixed parity similar in form to (b) [FiŁu2].

(f)  $R(C_3, C_3, C_k) = 5k-4$  for  $k \geq 5$  [SunYWLX], and  $R(C_4, C_4, C_k) = k+2$  for  $k \geq 11$  [SunYLZ2]. All exceptions to these formulas for small  $k$  are listed in Table XII.

(g) Almost all of the off-diagonal cases in Table XII required the use of computers.

### 6.3.2. More colors

$m$	3	4	5	6	7	8
3	17	11	17	12	25	16
4	51 62	18	33 77	18 20	49	20
5	162 307	27 29	65	26	97	28
6	538 1838	34 38	129		193	

**Table XIII.** Known values and bounds for  $R_k(C_m)$  for small  $k, m$ ;

(a) For the entries in the row  $k = 3$  and in the column  $m = 3$  in Table XIII, more details and all corresponding references are in Sections 6.3.1 and 6.1, respectively. The lower bounds for  $m = 5, 7$  are implied by 6.3.2.1. The bound  $R_4(C_5) \leq 158$  follows from 6.3.2.k. Using a reasoning as in [Li4] and the equality  $R_3(C_5) = 17$  one can obtain  $R_4(C_5) \leq 137$ . The bound  $R_4(C_5) \leq 77$  was obtained in 2020 with the help of flag algebras [LidP]. The references to other cases with  $k, m \geq 4$  can be found below in this section.

$R_4(C_4) = 18$	[Ex2] [SunYLZ1]
$33 \leq R_4(C_5) \leq 77$	[6.3.2.1] [LidP]
$18 \leq R_4(C_6) \leq 20$	[SunYJLS] [ZhaSW]
$27 \leq R_5(C_4) \leq 29$	[LaWo1]
$R_5(C_6) = 26$	[SunYJLS] [SunYW]
$34 \leq R_6(C_4) \leq 38$	[Ex22] [Boza8]
$24 \leq R(C_3, C_4, C_4, C_4) \leq 27$	[DyDz1] [XuR2]
$30 \leq R(C_3, C_3, C_4, C_4) \leq 36$	[DyDz1] [XuR2]
$49 \leq R(C_3, C_3, C_3, C_4) \leq 57$	[6.7.h] [BoRa]
$18 \leq R(C_4, C_6, C_6, C_6) \leq 20$	[ZhaSW]
$18 \leq R(C_4, C_4, C_6, C_6) \leq 20$	[ZhaSW]
$R(C_4, C_4, C_4, C_6) = 19$	[ZhaSW]

- (b)  $R_k(C_4) \leq k^2 + k + 1$  for all  $k \geq 1$ ,  $R_k(C_4) \geq k^2 - k + 2$  for all  $k - 1$  which is a prime power [Ir, Chu2, ChGra1], and  $R_k(C_4) \geq k^2 + 2$  for odd prime power  $k$  [LaWo1]. The latter was extended to any prime power  $k$  in [Ling, LaMu]. For even number of colors,  $k$ , it was shown that  $R_k(C_4) \leq k^2 + k$  [Boza8]. In 2025, the latter was improved by 1 to  $R_k(C_4) \leq k^2 + k - 1$  [LiLin1].
- (c) Formulas for  $R(C_m, C_n, C_k, C_l)$  for large  $m$  [EFRS1].

Bounds in (d)-(j) below cover different situations and each is interesting in some respect.

- (d)  $R_k(C_{2m}) \geq (k + 1)m$  for odd  $k$  and  $m \geq 2$ , and  $R_k(C_{2m}) \geq (k + 1)m - 1$  for even  $k$  and  $m \geq 2$  [DzNS].
- (e)  $R_k(C_{2m}) \geq 2(k - 1)(m - 1) + 2$  [SunYXL].
- (f)  $R_k(C_{2m}) \geq k^2 + 2m - k$  for  $2m \geq k + 1$  and prime power  $k$  [SunYJLS].
- (g)  $R_k(C_{2m}) = \Theta(k^{m/(m-1)})$  for fixed  $m = 2, 3$  and  $5$  [LiLih].
- (h)  $R_k(C_{2m}) \leq 201km$  for  $k \leq 10^m/201m$  [ErdG].
- (i)  $R_k(C_{2m}) \leq 2km + o(m)$  for all fixed  $k \geq 2$  [ŁucSS].
- (j)  $R_k(C_{2m}) \leq 2(k - c_k)m + o(m)$  for some small  $c_k > 0$ , for all fixed  $k \geq 2$  [Sár2]. This was improved to an absolute constant  $c = c_k = 1/4$  in [DavJR], and further to  $c = 1/2$  in [KniSu]. See also 6.4.2.d.
- (k)  $R_k(C_5) < (18^k k!)^{1/2}/10$  [Li4].
- (l)  $2^k m < R_k(C_{2m+1}) \leq (k + 2)!(2m + 1)$  [BoEr].  
 Better upper bound  $R_k(C_{2m+1}) < 2(k + 2)!m$  was obtained in [ErdG].  
 Still better upper bound  $R_k(C_{2m+1}) \leq (c^k k!)^{1/m}$ , for some positive constant  $c$ , if all Ramsey-critical colorings for  $C_{2m+1}$  are not far from regular, was obtained in [Li4].
- (m) For each fixed  $m \geq 3$ , there exists a positive constant  $c$  such that for every  $k \geq 3$ ,  $R_k(C_{2m+1}) < c^{k-1}(k!)^{1/2+\delta}$ , where  $\delta$  is approaching 0 for large  $m$  [LinCh].

- (n)  $R_k(C_{2m+1}) \leq k2^k(2m+1) + o(m)$  for all fixed  $k \geq 4$  [ŁucSS].
- (o) Conjecture that  $R_k(C_{2m+1}) = 2^k m + 1$  for all  $m \geq 2$  was credited by several authors to Bondy and Erdős [BoEr], though only lower bound, not the conjecture, is in this paper. After more than 40 years, in 2016, Jenssen and Skokan [JenSk] posted a preprint on arXiv (which appeared in *Advances in Mathematics* in 2021) containing a proof of the conjecture for each fixed  $k$  with sufficiently large  $m$ . On the other hand, the work by Day and Johnson [DayJ] shows that the lower bound of the conjecture does not hold for each  $m$  and sufficiently large  $k$ .
- (p)  $R_k(C_{2m+1}) \leq (4m-2)^k k^{k/m} + 1$  for all  $k, m \geq 1$  [AxCJMR].
- (q)  $R(C_n, C_{l_1}, \dots, C_{l_k}) = 2^k(n-1) + 1$  for all  $l_i$ 's odd with  $l_i > 2^i$ , and sufficiently large  $n$ , and support for the conjecture that  $R_k(C_n) = 2^{k-1}(n-1) + 1$  for large odd  $n$  [AllBS].
- (r)  $R_l(C_{\leq l+1}) = 2l+3$  for all odd  $l \geq 3$ . For even  $l$  we have:  $R_4(C_{\leq 5}) = 12$ ,  $R_6(C_{\leq 7}) = 12$ , and  $R_l(C_{\leq l+1}) = 2l+3$  for  $l = 8, 10$  and  $12$  [ZhuSWZ].
- (s) Other asymptotic bounds for  $R_k(C_n)$  [Bu0, GRS, ChGra2, Li4, LiLih, ŁucSS].
- (t) Survey of multicolor cycle cases [Li3].

### 6.3.3. Cycles versus other graphs

- (a) Some cases involving  $C_4$ :

$20 \leq R(C_4, C_4, K_4) \leq 21$	[DyDz1] [LidP]
$27 \leq R(C_3, C_4, K_4) \leq 29$	[DyDz1] [BoRa]
$52 \leq R(C_4, K_4, K_4) \leq 66$	[XSR1] [BoRa]
$34 \leq R(C_4, C_4, C_4, K_4) \leq 48$	[DyDz1] [LidP]
$49 \leq R(C_3, C_4, C_4, K_4) \leq 75$	[We3] [BoRa]
$87 \leq R(C_4, C_4, K_4, K_4) \leq 177$	[XSR1] [BoRa]
$R(K_{1,3}, C_4, K_4) = 16$	[KlaM2]
$R(C_4, C_4, K_4 - e) = 16$	[DyDz1]
$R(C_4, C_4, C_4, T) = 16$ for $T = P_4$ and $T = K_{1,3}$	[ExRe]

- (b)  $R(C_4, C_4, K_{1,n}) = 6, 8, 9, 11, 13, 14$  for  $n = 1, 2, 3$  [ArKM], and  $n = 4, 5, 6$  [ZhaCC4].
- (c) Study of the general upper bound on  $R(C_4, \dots, C_4, K_{1,n})$ , which for 3 colors implies  $R(C_4, C_4, K_{1,n}) \leq n+3 + \lceil \sqrt{4n+5} \rceil$  [ZhaCC4]. This was extended to hold for wheel graphs as in  $R(C_4, C_4, W_{n+1}) \leq n+3 + \lceil \sqrt{4n+5} \rceil$  for all  $n \geq 56$  or  $n$  equal to 42, 48, 49, 50, 51 or 52. Furthermore,  $R(C_4, C_4, W_{k^2-k+1}) \leq k^2 + k + 2$  for  $k \geq 9$ , with equality for prime powers  $k$  [ZhaCC5].
- (d) Study of  $R(C_4, K_{1,m}, P_n)$ , bounds and exact values for several special  $m$  and  $n$ , in particular when  $R(C_4, K_{1,m}, P_n) = R(C_4, K_{1,q^2-s})$ , for some small  $s$  [ZhZC, SunSh].
- (e) Study of  $R(H, H, C_n)$ , for sufficiently large odd  $n$ , and for suitably defined balanced bipartite graphs  $H$  with small bandwidth [YouLin2].

- (f) Study of asymptotics for  $R(C_{2m}, C_{2m}, K_n)$  for fixed  $m$  [AIRö, ShiuLL], in particular  $R(C_4, C_4, K_n) = \Theta(n^2 \text{poly-log } n)$  [AIRö].
- (g) For fixed  $m$  and large  $n$ ,  $\Omega(n^{1+2/(2m-1)}/(\log n)^{4/(2m-1)}) \leq R(C_{2m+1}, C_{2m+1}, K_n) \leq O(n^{(1+1/m)^2}/(\log n)^{(2m+1)/m^2})$  [LiuLi4].
- (h) Study of  $R(C_n, K_{t_1}, \dots, K_{t_k})$  and  $R(C_n, K_{t_1, s_1}, \dots, K_{t_k, s_k})$  for large  $n$  [EFRS1].
- (i)  $R(C_n, K_{t_1}, \dots, K_{t_k}) = (n-1)(r-1) + 1$  for  $n \geq 4r+2$ , where  $r = R(K_{t_1}, \dots, K_{t_k})$ . This equality was obtained as a special case of more general results in [OmRa2]. Similar proof was presented later in [Mad]. Further, see items 6.6.f and 6.7.f.
- (j) Study of asymptotics of  $R(C_m, \dots, C_m, K_n)$ , in particular for any fixed number of colors  $k \geq 4$  we have  $R(C_4, \dots, C_4, K_n) = \Theta(n^2/\log^2 n)$  [AIRö]. Improved lower bounds for odd  $m$ , in particular for  $R(C_5, \dots, C_5, K_n)$  and for  $R(C_7, \dots, C_7, K_n)$  [HuLin].
- (k) Lower and upper bounds on  $R(C_4, \dots, C_4, K_{1,n})$  [ZhaCC4, ZhaCC6, Boza8]. The equality  $R(C_4, \dots, C_4, K_{1,n}) = R(C_4, \dots, C_4, W_{n+1})$  for large  $n$  was obtained in [ZhaCC6], note that it generalizes item 4.3.e.
- (l) Upper bounds for  $R(C_4, \dots, C_4, G_1, \dots, G_k)$ , when certain constraints on graphs  $G_i$  are known, applicable in particular to stars [BoRa].
- (m) Monotone paths and cycles [Lef2], see also 4.5.p.
- (n) Upper bound on  $R(C_3, \dots, C_3, K_n)$  [HngJL].  
For other combinations of  $C_3$  and  $K_n$  see Sections 2.2, 3.2, 4.2, 6.1 and 6.2.

#### 6.4. Paths, paths versus other graphs

In 2007, Gyárfás, Ruszinkó, Sárközy and Szemerédi [GyRSS] established that for all  $n$  large enough we have

$$R(P_n, P_n, P_n) = 2n - 2 + (n \bmod 2).$$

Faudree and Schelp [FS2] conjectured that the latter holds for all  $n \geq 1$ . It is true for  $n \leq 9$  (see (c) below), and the first open case is that for  $P_{10}$ . The conjectured equality  $R(C_{2m}, C_{2m}, C_{2m}) = 4m$  (see 6.3.1.b), whenever true, implies the above for three paths  $P_{2m+1}$  case [DyDR].

##### 6.4.1. Three-color path and path-cycle cases

- (a)  $R(P_m, P_n, P_k) = m + \lfloor n/2 \rfloor + \lfloor k/2 \rfloor - 2$  for  $m \geq 6(n+k)^2$  [FS2],  
the equality holds asymptotically for  $m \geq n \geq k$  with an extra term  $o(m)$  [FiLu1],  
extensions of the range of  $m, n, k$  for which (a) holds were obtained in [Biel3].
- (b)  $R(P_3, P_m, P_n) = m + \lfloor n/2 \rfloor - 1$  for  $m \geq n$  and  $(m, n) \neq (3, 3), (4, 3)$  [MaORS2].
- (c)  $R_3(P_3) = 5$  [Ea1],  $R_3(P_4) = 6$  [Ir],  
 $R(P_m, P_n, P_k) = 5$  for other  $m-n-k$  combinations with  $3 \leq m, n, k \leq 4$  [ArKM],  
 $R_3(P_5) = 9$  [YR1],  $R_3(P_6) = 10$  [YR1], and  $R_3(P_7) = 13$  [YY],  
 $R_3(P_8) = 14, R_3(P_9) = 17$  [DyDR].

- (d)  $R(P_4, P_4, P_{2n}) = 2n + 2$  for  $n \geq 2$ ,  
 $R(P_5, P_5, P_5) = R(P_5, P_5, P_6) = 9$ ,  
 $R(P_5, P_5, P_n) = n + 2$  for  $n \geq 7$ ,  
 $R(P_5, P_6, P_n) = R(P_4, P_6, P_n) = n + 3$  for  $n \geq 6$ ,  
 $R(P_6, P_6, P_{2n}) = R(P_4, P_8, P_{2n}) = 2n + 4$  for  $n \geq 14$  [OmRa1].
- (e)  $R(P_m, P_n, C_k) = 2n + 2 \lfloor m/2 \rfloor - 3$  for large  $n$  and odd  $k \geq 3$  [DzFi2],  
improvements on the range of  $m, n, k$  [Biel3, Fid1].
- (f)  $R(P_3, P_3, C_m) = 5, 6, 6$ , for  $m = 3, 4$  [ArKM], 5,  
 $R(P_3, P_3, C_m) = m$  for  $m \geq 6$  [Dzi2].  
 $R(P_3, P_4, C_m) = 7$  for  $m = 3, 4$  [ArKM] and 5,  
 $R(P_3, P_4, C_m) = m + 1$  for  $m \geq 6$  [Dzi2].  
 $R(P_4, P_4, C_m) = 9, 7, 9$  for  $m = 3, 4$  [ArKM] and 5 [Dzi2],  
 $R(P_4, P_4, C_m) = m + 2$  for  $m \geq 6$  [DzKP].
- (g)  $R(P_3, P_5, C_m) = 9, 7, 9, 7, 9$  for  $m = 3, 4, 5, 6, 7$  [Dzi2, DzFi2],  
 $R(P_3, P_5, C_m) = m + 1$  for  $m \geq 8$  [DzKP].  
A table of  $R(P_3, P_k, C_m)$  for all  $3 \leq k \leq 8$  and  $3 \leq m \leq 9$  [DzFi2].
- (h)  $R(P_4, P_5, C_m) = 11, 7, 11, 11$ , and  $m + 2$  for  $m = 3, 4, 5, 7$  and  $m \geq 23$ ,  
 $R(P_4, P_6, C_m) = 13, 8, 13, 13$ , and  $m + 3$  for  $m = 3, 4, 5, 7$  and  $m \geq 18$  [ShaXSP].
- (i)  $R(P_3, P_n, C_4) = n + 1$  for  $n \geq 6$  [DzFi2],  
 $R(P_3, P_n, C_6) = n + 2$  for  $n \geq 6$ ,  
 $R(P_3, P_n, C_8) = n + 3$  for  $n \geq 7$  [Fid1],  
 $R(P_3, P_n, C_k) = 2n - 1$ , and  
 $R(P_4, P_n, C_k) = 2n + 1$  for odd  $k \geq 3$  and  $n \geq k$  [DzFi2].
- (j)  $R(P_3, P_6, C_m) = m + 2$  for  $m \geq 23$ ,  
 $R(P_6, P_6, C_m) = R(P_4, P_8, C_m) = m + 4$  for  $m \geq 27$ ,  
 $R(P_6, P_7, C_m) = m + 4$  for  $m \geq 57$ ,  
 $R(P_4, P_n, C_4) = R(P_5, P_n, C_4) = n + 2$  for  $n \geq 5$  [OmRa1].
- (k)  $R(P_3, C_3, C_3) = 11$  [BuE3],  $R(P_3, C_4, C_4) = 8$  [ArKM],  $R(P_3, C_6, C_6) = 9$  [Dzi2],  
 $R(P_3, C_m, C_m) = R(C_m, C_m) = 2m - 1$  for odd  $m \geq 5$  [DzKP] (for  $m = 5, 7$  [Dzi2]),
- (l)  $R(P_3, C_n, C_m) = R(C_n, C_m)$  for  $n \geq 7$  and odd  $m, 5 \leq m \leq n$ , and  
some values and bounds on  $R(P_3, C_n, C_m)$  in other cases [Fid1].
- (m)  $R(P_3, C_3, C_4) = 8$  [ArKM],  $R(P_3, C_3, C_5) = 9$ ,  $R(P_3, C_3, C_6) = 11$ ,  
 $R(P_3, C_3, C_7) = 13$ ,  $R(P_3, C_4, C_5) = 8$ ,  $R(P_3, C_4, C_6) = 8$ ,  
 $R(P_3, C_4, C_7) = 8$ ,  $R(P_3, C_5, C_6) = 11$ ,  $R(P_3, C_5, C_7) = 13$  and  
 $R(P_3, C_6, C_7) = 11$  [Dzi2].
- (n)  $R(P_4, C_3, C_5) = 13$ ,  $R(P_4, C_4, C_5) = 10$ ,  $R(P_4, C_4, C_6) = 9$ ,  
 $R(P_4, C_5, C_5) = 13$ ,  $R(P_4, C_6, C_6) = 10$  [SunSh].

- (o) A formula for  $R(P_m, P_n, C_k)$  for  $k$  large enough and  $m, n$  satisfying some constraints. In addition, some cases involving  $tK_2$  instead of  $C_k$  are derived as side results [KhoDz].
- (p) Study of  $R(P_n, C_4, K_{1,m})$  [ZhZC, SunSh].
- (q) Formulas for  $R(pP_3, qP_3, rP_3)$  and  $R(pP_4, qP_4, rP_4)$  [Scob].
- (r) Lower and upper bounds on  $R_3(P_m \cup K_{1,n})$  [ZhouLMW].
- (s)  $R(P_3, K_4-e, K_4-e) = 11$  [Ex7]. All colorings which can form any color neighborhood for the case  $R_3(K_4-e)$  (see Section 6.5) were found in [Piw2].

### 6.4.2. More colors

- (a)  $R_k(P_3) = k + 1 + (k \bmod 2)$ ,  $R_k(2P_2) = k + 3$  for all  $k \geq 1$  [Ir].
- (b)  $R_k(P_4) = 2k + c_k$  for all  $k$  and some  $0 \leq c_k \leq 2$ . If  $k$  is not divisible by 3 then  $c_k = 3 - k \bmod 3$  [Ir]. Wallis [Wall] showed  $R_6(P_4) = 13$ , which already implied  $R_{3t}(P_4) = 6t + 1$ , for all  $t \geq 2$ . Independently, the case  $R_k(P_4)$  for  $k \neq 3^m$  was completed by Lindström in [Lind], and later Bierbrauer proved  $R_{3^m}(P_4) = 2(3^m) + 1$  for all  $m > 1$ .  $R_3(P_4) = 6$  [Ir].
- (c)  $R_3(P_5) = 9$  [YR1] and  $R_4(P_5) = 11$  [LiuMS]. Furthermore, for all  $k > 4$  we have  $R_k(P_5) = 3k + c$ , where  $c = 0$  for  $k$  equal to 2 or 3 mod 4,  $c = 1$  for  $k = 0 \bmod 4$ , and  $c = 2$  for  $k = 1 \bmod 4$  [LiuMS].
- (d)  $R_k(P_n) \leq (k - c_k)n + o(n)$  for some small  $c_k > 0$ , for all fixed  $k \geq 2$  [Sár2]. This was improved to an absolute constant  $c = c_k = 1/4$  in [DavJR], and further to  $c = 1/2$  in [KniSu]. See also 6.3.2.j/o.
- (e) Formula for  $R(P_{n_1}, \dots, P_{n_k})$  for large  $n_1$  [FS2], and some extensions [Biel3]. Conjectures about  $R(P_{n_1}, \dots, P_{n_k})$  when all or all but one of  $n_i$ 's are even [OmRa1].
- (f) Formulas for  $R(P_{n_1}, \dots, P_{n_k}, C_m)$  for some cases, for large  $m$  [OmRa1].
- (g) Formula for  $R(n_1P_2, \dots, n_kP_2)$ , in particular  $R(nP_2, nP_2, nP_2) = 4n - 2$  [CocL1]. New proof with characterization of all critical graphs [XuYZ]. Note how close the latter is to  $R(C_{2n}, C_{2n}, C_{2n}) = 4n$ , and see an earlier item 6.3.1.b.
- (h) Cockayne and Lorimer [CocL1] found the exact formula

$$R(n_1P_2, \dots, n_kP_2) = n_1 + 1 + \sum_{i=1}^k (n_i - 1), \text{ where } n_1 = \max\{n_1, \dots, n_k\}.$$

Later, Lorimer [Lor] extended it to a more general case of  $R(K_m, n_1P_2, \dots, n_kP_2)$ . More general cases of the latter, with multiple copies of the complete graph, paths, stars and forests, were studied in [Stahl, LorSe, LorSo, GyRSS]. A special 3-color case  $R(P_3, mP_2, nP_2) = 2m + n - 1$  for  $m \geq n \geq 3$  is given in [MaORS2], and some other cases in [KhoDz]. The general case of multicolor combinations of stars and stripes is completed in [OmRR]. Ramsey numbers for path-matchings and covering designs, generalizing  $R(n_1P_2, \dots, n_kP_2)$ , are studied in [DeBGS].

- (i) Multicolor cases for one large path or cycle involving small paths, cycles, complete and complete bipartite graphs [EFRS1].
- (j) See Sections 6.5 and 8.2, especially [ArKM, BoDD], for a number of cases for triples of small graphs.

### 6.5. Special cases

- (a) Denote  $K_3+e = K_4 - P_3$ .

$$R_3(K_3+e) = R_3(K_3) [= 17] \quad [\text{YR3, ArKM}]$$

$$R(K_3+e, K_3+e, K_4-e) = R(K_3, K_3, K_4-e) = 17 \quad [\text{ShWR}]$$

$$R(K_3+e, K_3+e, K_5-P_3) = R(K_3, K_3, K_4) [= 30] \quad [\text{ShWR}]$$

$$\text{If } R_4(K_3) = 51 \text{ then } R_4(K_3+e) = 52, \text{ and}$$

$$\text{if } R_4(K_3) > 51 \text{ then } R_4(K_3+e) = R_4(K_3) \quad [\text{ShWR}]$$

- (b)

$$R_3(K_4-e) = 28 \quad [\text{Ex7}] [\text{LidP}]$$

$$R(P_3, K_4-e, K_4-e) = 11 \quad [\text{Ex7}], \text{ all colorings } [\text{Piw2}]$$

$$R(P_3, K_4-e, K_4) = 17 \quad [\text{ArKM}]$$

$$R(P_3, K_4, K_4) = 35 \quad [\text{BuE3}], \text{ special case of 6.7.e}$$

$$472 \leq R_3(K_6-e) \quad [\text{HeLD}]$$

$$1102 \leq R_3(K_7-e) \quad [\text{HeLD}], \text{ superseded by}$$

$$1106 \leq R_3(6) [\text{Row3}]$$

- (c)

$$21 \leq R(K_3, K_4-e, K_4-e) \leq 22 \quad [\text{ShWR}] [\text{LidP}]$$

$$31 \leq R(K_3, K_4, K_4-e) \leq 40 \quad [\text{VO}] [\text{LidP}]$$

$$35 \leq R(K_4, K_4-e, K_4-e) \leq 47 \quad [\text{We3}] [\text{LidP}]$$

$$55 \leq R(K_4, K_4, K_4-e) \leq 94 \quad [\text{Ea1}] [\text{LidP}]$$

- (d)

$$R(C_4, P_4, K_4-e) = 11 \quad [\text{ArKM}]$$

$$R(C_4, P_4, K_4) = 14 \quad [\text{BoDD}]$$

$$R(C_4, C_4, K_4-e) = 16 \quad [\text{DyDz1}]$$

$$R(C_4, K_3, K_4-e) = 17 \quad [\text{BoDD}]$$

$$R(C_4, K_4-e, K_4-e) = 19 \quad [\text{BoDD}]$$

$$29 \leq R(C_4, K_4, K_4-e) \leq 36 \quad [\text{VO}] [\text{BoDD}]$$

$$52 \leq R(C_4, K_4, K_4) \leq 66 \quad [\text{XSR1}] [\text{BoRa}]$$

- (e) For prime  $p = 3q + 1$ , if the cubic residues Paley graph  $Q_p$  contains no  $K_k - e$ , then  $R_3(K_{k+1} - e) > 3p$  [HeLD]. The cases  $k=5$  and  $k=6$  give two bounds listed in (b). Also based on Paley graphs, several new lower bounds for  $R_3(K_1+G)$ , and in particular

for  $R_3(B_n)$ , were derived in [LinLS].

- (f) If  $T_n$  is the set of all  $n$ -vertex trees (and all monochromatic  $n$ -vertex trees are avoided), then  $R_3(T_n) = 2n - 2$  for even  $n$ , and  $R_3(T_n) = 2n - 1$  for odd  $n$  [GeGy].
- (g)  $R_3(G_{a,b}) = (2 + o(1))ab$ , where  $G_{a,b}$  is the rectangular  $a \times b$  grid graph. Lower and upper bounds on  $R_3(G)$  for graphs  $G$  with small bandwidth and bounded  $\Delta(G)$  [MoSST].
- (h)  $R(K_3, K_3, F_n) = 10n + 1$  for  $n \geq 6$ , where  $F_n$  is the fan graph [BudPri].
- (i) See also Section 8.2 for pointers to cumulative data for three colors.

### 6.6. General results for special graphs

Several authors use  $R_k(G; H)$  to denote the  $k$ -color graph Ramsey number  $R(G, \dots, G, H)$ , where monochromatic  $G$  is avoided in the first  $k - 1$  colors, and  $H$  in the  $k$ -th color. This is different from our notation, where  $r$  in  $R_k(G; r)$  and  $R(G; r)$  in Section 7 stands for the uniformity of considered hypergraphs.

- (a) Formulas for  $R_k(G)$ , where  $G$  is one of the graphs  $P_3$ ,  $2K_2$  and  $K_{1,3}$ , for all  $k$ , and for  $P_4$  if  $k$  is not divisible by 3 [Ir]. For some details see Section 6.4.2.b.
- (b)  $tk^2 + 1 \leq R_k(K_{2,t+1}) \leq tk^2 + k + 2$ , where the upper bound is general, and the lower bound holds when both  $t$  and  $k$  are powers of the same prime  $p$  [ChGra1, LaMu].
- (c)  $(m - 1)\lfloor (k + 1)/2 \rfloor < R_k(T_m) \leq 2km + 1$  for any tree  $T_m$  with  $m$  edges [ErdG], see also [GRS]. The lower bound can be improved for special large  $k$  [ErdG, GRS]. The upper bound was improved to  $R_k(T_m) < (m - 1)(k + \sqrt{k(k - 1)}) + 2$  in [GyTu].
- (d)  $k(\sqrt{m} - 1)/2 < R_k(F_m) < 4km$  for any forest  $F_m$  with  $m$  edges [ErdG], see [GRS]. See also pointers in other items below which involve special forests (like trees and stars).
- (e)  $R(S_1, \dots, S_k) = n + \varepsilon$ , where  $S_i$ 's are arbitrary stars,  $n = n(S_1) + \dots + n(S_k) - 2k$ , and we set  $\varepsilon = 1$  if  $n$  is even and some  $n(S_i)$  is odd, and  $\varepsilon = 2$  otherwise [BuRo1]. See also [GauST, Par6]. Note that for graph  $G$  (here the set of edges in a given color), to avoid star  $S = K_{1,n}$  is equivalent to having  $\delta(G) < n$ .
- (f) Formula for  $R(S_1, \dots, S_k, K_n)$ , where  $S_i$ 's are arbitrary stars [Jaco]. It was generalized to a formula for  $R(S_1, \dots, S_k, K_{k_1}, \dots, K_{k_r})$  expressed in terms of  $R(k_1, \dots, k_r)$  and star orders [BoCGR]. A much shorter proof of the latter was presented in [OmRa2]. Special cases for bistars [AlmHS], and bounds for stars and trees instead of stars [Bai].
- (g) Formula for  $R(S_1, \dots, S_k, nK_2)$ , where  $S_i$ 's are arbitrary stars [CocL2], and a formula for  $R(n_1K_2, \dots, n_kK_2)$  [CocL1]. A new proof with characterization of all critical graphs [XuYZ]. See also cases involving  $P_2$  in Section 6.4.2.
- (h) Formula for  $R(S_1, \dots, S_k, G)$ , where  $S_i$ 's are stars and  $G$  is a tree [ZhZ1], or  $G$  is a cycle or wheel [RaeZ], for  $G$  of some orders depending on stars. Extension of these results to larger ranges of orders of  $G$ , and for  $G$  being a path [Wang1, ZhaHou].

Special cases when  $S_i$ 's are trees and  $G$  is a wheel [RaeZ].

- (i) Formulas for  $R(S_1, \dots, S_k)$ , where each  $S_i$ 's is a star or  $m_i K_2$  [ZhZ2, ErdG, OmRR], formula for the case  $R(S, mK_2, nK_2)$  [GySá2]. Several cases of formulas and asymptotics for  $R_k(S(n, m))$ , for sufficiently large  $n$ , where  $S(m, n)$  is a double star [RuoS, Sár3]. The relation of  $R_k(S(n, m))$  to the corresponding list Ramsey numbers [RuoS]. More bounds on  $R_k(S(n, n))$  and their relation to  $R_k(P_{2n+2})$  [BalDO]. For work on two-color cases for double stars see Sections 5.5 and 5.13.
- (j) Formula for  $R(F, K_{k_1}, \dots, K_{k_r})$  in terms of  $R(K_{k_1}, \dots, K_{k_r})$  and the size and structure of any forest  $F$  [KamRa2]. This corrects a claim in an earlier version of [AlmBCL]. The latter studies the concept of multicolor  $p$ -goodness. For 2-color concept of goodness see Section 5.15.
- (k) Bounds on  $R_k(G)$  for unicyclic graphs  $G$  of odd girth. Some exact values for special graphs  $G$ , for  $k = 3$  and  $k = 4$  [KrRod]. See Section 5.11 for two-color numbers involving unicyclic graphs.
- (l)  $R_k(K_{3,3}) = (1 + o(1))k^3$  [AIRóS].
- (m)  $R_k(\theta_{3,4}) = \Theta(k^{4/3})$  for large  $k$ , where  $\theta_{3,4}$  is the graph of three internally disjoint paths  $P_4$  sharing their endpoints [LiuW].
- (n) Bounds on  $R_k(K_{s,t})$ , in particular for  $K_{2,2} = C_4$  and  $K_{2,t}$  [ChGra1, AxFM]. Asymptotics of  $R_k(K_{s,t})$  for fixed  $k$  and  $s$  [DoLi, LiTZ]. Upper bounds on  $R_k(K_{s,t})$  [SunLi]. Upper bounds on multicolor numbers where all but the last color involve  $K_{s,t}$  while the last color avoids a star or another bipartite graph [WaLL]. Upper bounds in the case when the last color is avoiding generalized books [LiLW]. Upper bounds in the case when the last color is avoiding fans [LiZZ].
- (o) Asymptotics of  $R(k, \dots, k, n)$  in  $k, n$  and the number of colors [HeWi]. Exact asymptotics  $R(K_{t,s}, K_{t,s}, K_{t,s}, K_m) = \Theta(m^t / \log^t m)$ , for any fixed  $t > 1$  and large  $s \geq (t-1)! + 1$  [AIRö].
- (p) Variety of asymptotic results on  $R(K_{2,s}, \dots, K_{2,s}, K_m)$  [LeMu]. Asymptotic upper bounds on  $R(G, \dots, G, H)$  in two cases:  $G = K_{m,n}$ ,  $H$  is any nonempty graph, and  $G = B_n$ ,  $H = K_{m,n}$  [LiW].
- (q) Bounds on  $R_k(G)$  for trees, forests, stars and cycles [Bu0]. Bounds for trees  $R_k(T)$  and forests  $R_k(F)$  [ErdG, GRS, BierB, GyTu, Bra1, Bra2, SwPr].
- (r) Study of the case  $R(K_m, n_1 P_2, \dots, n_k P_2)$  [Lor]. Other similar results include  $R(P_3, mK_2, nK_2) = 2m + n - 1$  for  $m \geq n \geq 3$  [MaORS2] and  $R(S_n, nK_2, nK_2) = 3n - 1$  [GySá2]. More general cases, with multiple copies of the complete graph, stars and forests, were investigated in [Stahl, LorSe, LorSo, GyRSS, OmRR]. See also Section 6.4.2.
- (s) See Section 8.2, especially [ArKM, BoDD], for a number of cases for other small graphs, similar to those listed in Sections 6.3 and 6.4.

### 6.7. General results

- (a) In 2001, Nešetřil and Rosenfeld [NeRo] in an overview paper with numerous historical comments discuss the connections between the Ramsey numbers  $R_r(3)$ , Schur partitions and Shannon capacity of graphs.
- (b) In 2020, the limit  $\lim_{r \rightarrow \infty} R_r(3)^{1/r}$  was studied by Fox, Pach and Suk [FoxPS1] assuming a conjecture for multicolorings with bounded VC-dimension, and further for  $\lim_{r \rightarrow \infty} R_r(k)^{1/r}$  when restricted to the so-called semi-algebraic colorings [FoxPS2]. For other related work see items 6.2.a-f.
- (c) Szemerédi's Regularity Lemma [Szem] states that the vertices of every large graph can be partitioned into similar size parts so that the edges between these parts behave almost randomly. This lemma has been used extensively in various forms to prove the upper bounds, including those studied in [BenSk, GyRSS, GySS1, HaŁP1+, HaŁP2+, KoSS1, KoSS2].
- (d)  $R(m_1 G_1, \dots, m_k G_k) \leq R(G_1, \dots, G_k) + \sum_{i=1}^k n(G_i)(m_i - 1)$ , exercise 8.3.28 in [West].
- (e) If  $G$  is connected and  $R(K_k, G) = (k-1)(n(G)-1)+1$ , in particular if  $G$  is any  $n$ -vertex tree, then  $R(K_{k_1}, \dots, K_{k_r}, G) = (R(k_1, \dots, k_r) - 1)(n - 1) + 1$  [BuE3]. A generalization for connected  $G_1, \dots, G_n$  in place of  $G$  appeared in [Jaco].
- (f) Conjecture that  $R_3(H) \leq 2^{\Delta^{1+o(1)}} n$ , where  $\Delta = \Delta(H)$  [ConFS7].
- (g) For connected graphs  $G_1, \dots, G_n$  with  $s = R(G_1, \dots, G_n)$  and  $t = R(K_{k_1}, \dots, K_{k_r})$ , if  $m \geq 2$  and  $R(G_1, \dots, G_n, K_m) = (s-1)(m-1)+1$ , then  $R(G_1, \dots, G_n, K_{k_1}, \dots, K_{k_r}) = (s-1)(t-1) + 1$  [OmRa2]. The latter implies that  $R(T_k, K_{k_1}, \dots, K_{k_r}) = (k-1)(t-1)+1$  for any  $k$ -vertex tree  $T_k$ . The same result as in [OmRa2] was presented much later in [Mad]. It also generalizes a result in [BoCGR].
- (h) If  $F, G, H$  are connected graphs then  $R(F, G, H) \geq (R(F, G)-1)(\chi(H)-1) + \min\{R(F, G), s(H)\}$ , where  $s(G)$  is the chromatic surplus of  $G$  (see Section 5.15). This leads to several formulas and bounds for  $F$  and  $G$  being stars and/or trees when  $H = K_n$  [ShiuLL].
- (i)  $R(K_{k_1}, \dots, K_{k_r}, G_1, \dots, G_s) \geq (R(k_1, \dots, k_r)-1)(R(G_1, \dots, G_s)-1) + 1$  for arbitrary graphs  $G_1, \dots, G_s$  [Bev]. This generalizes 6.2.q, but is a special case of 6.7.g.
- (j) Constructive bound  $R(G_1, \dots, G_{t^{n-1}}) \geq t^n + 1$  for decompositions of  $K_{t^n}$  [LaWo1, LaWo2].
- (k)  $R(G_1, \dots, G_k) \leq 32\Delta k^\Delta n$ , where  $n \geq n(G_i)$  and  $\Delta \geq \Delta(G_i)$  for all  $1 \leq i \leq k$ ,  $R(G_1, \dots, G_k) \leq k^{2k\Delta q} n$ , where  $q \geq \chi(G_i)$  for all  $1 \leq i \leq k$  [FoxSu1].
- (l)  $R_k(G) \leq k^{6e(G)^{2/3}k}$  for all isolate-free graphs  $G$  and  $k \geq 3$  [JoPe].  
For the original two-color conjecture, now a theorem, see item 5.18.j [Erd4].

- (m)  $R_k(G) \succ (sk^{e(G)-1})^{1/n(G)}$ , where  $s$  is the number of automorphisms of  $G$  [ChH3]. Other general bounds for  $R_k(G)$  [ChH3, Par6].
- (n) Study of  $R(G_1, \dots, G_k, G)$  for large sparse  $G$  [EFRS1, Bu3].
- (o) Study of asymptotics for  $R(H, \dots, H, K_m)$ , in particular when  $H$  is a fixed bipartite graph, and for  $R(C_n, \dots, C_n, K_m)$  [AlRö]. An improvement of the lower bound asymptotics for  $C_5$  and  $C_7$  was obtained in [XuGe], and further improvement for the same in [HuLin]. See also Sections 6.3.3.f/g.
- (p) Relations between the Shannon capacity of noisy communication channels and graph Ramsey numbers. A lower bound construction for  $R_k(m)$  implying that supremum of the Shannon capacity over all graphs with bounded independence cannot be achieved by any finite graph power [XuR3]. For some other links between Shannon capacity and Ramsey numbers see Section 6 in [Ros2], and [Li2].
- (q) See surveys listed in Section 8.

## 7. Hypergraph Numbers

### 7.1. Values and bounds for numbers

The only known value of a nontrivial classical Ramsey number for hypergraphs:

$$R(4, 4; 3) = 13 \quad \text{[MR1]}$$

There are exactly 434714 critical colorings on 12 points, none of which extend to a 2-coloring of all triples in  $K_{13-t}$  without monochromatic  $K_4$  [McK2]

The computer evaluation of  $R(4, 4; 3)$  in 1991 consisted of an improvement of the upper bound from 15 to 13. This result followed an extensive theoretical study of this number by several authors [Gi4, Isb1, Sid1].

- (a)  $35 \leq R(4, 5; 3)$  [Dyb2]
- $63 \leq R(4, 6; 3)$  [Dyb3]
- $88 \leq R(5, 5; 3)$  [Dyb3]
- $79 \leq R(4, 4, 4; 3)$  [Dyb3]
- $35 \leq R(5, 5; 4)$  [Ex24]
- $163 \leq R(5, 5, 5; 3)$  [BudHR1]

The last bound can be much improved to  $7570 \leq R(5, 5, 5; 3)$  by using  $88 \leq R(5, 5; 3)$  and a general constructive result in [BrBH], which yields  $R_k(5; 3) \geq 87^{2^{k-2}}$ .

- (b)  $R(K_4-t, K_4-t; 3) = 7$  [Ea2]
- $R(K_4-t, K_4; 3) = 8$  [Sob, Ex1, MR1]
- $R(K_4-t, K_5-t; 3) = 12$  [LidP]
- $R(K_4-t, K_4-t, K_4-t; 3) = 13$  [Ex1] [We3]
- $14 \leq R(K_4-t, K_5; 3) \leq 16$  [Ex1] [LidP]

(c) The first bound on  $R(4, 5; 3) \geq 24$  was obtained by Isbell [Isb2]. Shastri [Shas] gave a weak bound  $R(5, 5; 4) \geq 19$  (after a few improvements, now 35 in [Ex24]), nevertheless his lemmas, the stepping-up lemmas by Erdős and Hajnal (see [GRS, GrRö], also item 7.5.a), and others in [Ka3, Abb2, GRS, GrRö, HuSo, SonYL] can be used to derive better lower bounds for higher numbers.

(d) Several lower bound constructions for 3-uniform hypergraphs were presented in [HuSo]. Study of lower bounds on  $R(p, q; 4)$  can be found in [Song3] and [SonYL, Song4] (the latter two papers are almost the same in contents). Most of the concrete lower bounds in these papers can be easily improved by using the same techniques, but starting with better constructions for small parameters as listed above.

- (e)  $R(p, q; 4) \geq 2R(p-1, q; 4) - 1$  for  $p, q > 4$ , and  
 $R(p, q; 4) \geq (p-1)R(p-1, q; 4) - p + 2$  for  $p \geq 5, q \geq 7$  [SonYL].  
 Lower bound asymptotics for  $R(p, q; 4)$  [SonLi].

- (f) Recurrence relations in the form  $R(p, q; r) \geq d(R(p-1, q; r)-1)+1$ , where  $d$  depends on  $p, q$  and  $r$ , including the following: There exists  $c \geq 25$ , such that for  $k, 5 \leq k \leq c$ , and any  $p \geq k+2$  and  $q \geq k+1$ , we have  $R(p, q; r) \geq (p-1)(R(p-1, q; r)-1)+1$  [Liu]. Such relations lead to the following bounds:

$$\begin{array}{lll} R(5, 6; 4) \geq 67, & R(6, 6; 4) \geq 133, & R(7, 6; 4) \geq 661, \\ R(7, 7; 4) \geq 3961, & R(8, 8; 4) \geq 194041, & R(13, 6; 4) \geq 50689, \\ R(6, 6; 5) \geq 72. \end{array}$$

- (g)  $R(K_{1,1,c}, K_{1,1,c}; 3) = c+2$  for  $2 \leq c \leq 4$ , and a conjecture that this equality holds for all  $c \geq 5$ . The lower and upper bounds are closely related to the existence of appropriate BIBDs (balanced incomplete block designs) with block size 3. The conference and journal versions of this work [MiPal] differ on some results. See also item 7.4.h.
- (h) Lower bound asymptotics for  $R(4, n; 3)$  [ConFS2], lower bound asymptotics for  $R(5, n; 4)$  [MuSuk2, MuSuk3], and lower bound asymptotics for  $R(6, n; 4)$  [MuSuk3].
- (i) Lower and upper bounds on  $R(K_4-t, K_n; 3)$  [ErdH, MuSuk2]. Extensions to  $r$ -half-graph  $B^r$ , where  $B^3 = K_4-t$  [MuSuk2], and to  $B^3$  versus stars [MuSp].
- (j) Several constructive lower bounds for hypergraph numbers, including constructions which introduce a new color. In particular, they imply that  $R_k(5; 3)$  is equal to at least 82, 163, 131073, 262145 or 524289, for  $k = 2, 3, 4, 5$  and 6 colors, respectively [BudHR1]. Using 7.1.e and other known concrete lower bounds,  $R(5, 6; 4) \geq 67$  and  $R(4, 4, 5, 5, 5, 5; 3) \geq 17179869185$  are noted in [BudHP].

## 7.2. Cycles and paths

**Definitions.**  $P_n^{r,s}$  is called an  $s$ -path in an  $r$ -uniform hypergraph  $H$ , if it consists of  $n$  hyperedges  $\{e_1, \dots, e_n\}$  in  $E(H)$ , such that  $|e_i \cap e_{i+1}| = s$  for all  $1 \leq i < n$ , and all other vertices in  $e_j$ 's are distinct [Peng]. An  $s$ -cycle  $C_n^{r,s}$  is defined analogously. Several authors use the terms of *loose* paths and *loose* cycles, which are 1-path and 1-cycles, and *tight* paths and *tight* cycles, the latter most often for 3-uniform hypergraphs when they are 2-paths and 2-cycles, respectively.

In the following items (b) to (i), when  $r = 3$  or  $r$  is implied by the context, we write  $C_n$  and  $P_n$  for the  $r$ -uniform loose cycles and paths,  $C_n^{r,1}$  and  $P_n^{r,1}$ , respectively. In other cases special comments are added.

### 7.2.1. Two colors

- (a) Tetrahedron is formed by four triples on the set of four points. The Ramsey number of tetrahedron is  $R(4, 4; 3) = 13$  [MR1].

- (b) For loose cycles and paths,  $R(C_3, C_3; 3) = 7$ ,  $R(C_4, C_4; 3) = 9$ , and for the  $r$ -uniform case we have in general  $R(P_3, P_3; r) = R(P_3, C_3; r) = R(C_3, C_3; r) + 1 = 3r - 1$  and  $R(P_4, P_4; r) = R(P_4, C_4; r) = R(C_4, C_4; r) + 1 = 4r - 2$ , for  $r \geq 3$ . These results and discussion of several related cases were presented in [GyRa].
- (c)  $R(P_m, P_n; 3) = R(C_m, C_n; 3) + 1 = R(P_m, C_n; 3) = 2m + \lfloor (n+1)/2 \rfloor$ , for all  $m \geq n$ , and  $R(C_m, P_n; 3) = 2m + \lfloor (n-1)/2 \rfloor$ , for  $m > n$  [MaORS1, OmSh1]. The formulas for 3-uniform two-color cases of some loose paths versus stars are derived in [ZhCh].
- (d) For loose cycles,  $R(C_{2n}, C_{2n}; 3) > 5n - 2$  and  $R(C_{2n+1}, C_{2n+1}; 3) > 5n + 1$ , and asymptotically these lower bounds are tight [HaLP1+]. Generalizations to  $r$ -uniform hypergraphs and graphs other than cycles appeared in [GySS1].
- (e) For loose cycles,  $R(C_n, C_n; r) = (r-1)n + \lfloor (n-1)/2 \rfloor$  for  $n \geq 2$ ,  $r \geq 8$  [OmSh2], and it also holds for  $r=4$  [OmSh3]. Further extensions to off-diagonal cases as in (c) are obtained in [OmSh4]. Based on these results, it was conjectured that for  $n \geq m \geq 3$  and  $r \geq 3$ , we have  $R(C_n, C_m; r) = (r-1)n + \lfloor (m-1)/2 \rfloor$ . In [Shah], the known cases of this conjecture are discussed, and it is shown that it holds for  $r=5$  with large  $n$ .
- (f) For tight cycles,  $R(C_{3n}, C_{3n}; 3) \approx 4n$  and  $R(C_{3n+i}, C_{3n+i}; 3) \approx 6n$  for  $i=1$  or  $2$ , and for tight paths  $R(P_n, P_n; 3) \approx 4n/3$  [HaLP2+]. Some related results are discussed in [PoRRS]. Also for tight paths and cycles, but 4-uniform, we have  $R(P_n, P_n; 4) \approx 5n/4$  and  $R(C_{4n}, C_{4n}; 4) \approx 5n$  [LoPfe]. For tight cycles further generalization to all uniformities was obtained in [Pfe].
- (g) Exact values for Ramsey numbers involving  $s$ -paths for even  $r$  and  $s = r/2$ , in particular for  $P_n^{r,s}$  versus  $P_3^{r,s}$  and  $P_4^{r,s}$ , when this value is  $(n+1)s + 1$  [Peng].
- (h) Lower and upper asymptotic bounds for  $R(C_3^{3,1}, K_m; 3)$  and  $R(C_3^{r,1}, K_m; r)$  [KosMV2].
- (i) Lower and upper asymptotic bounds for  $R(C_s, K_m; 3)$  for tight cycles  $C_s$  [MuR]. An improvement of the upper bound from the latter was obtained in [Mub1].
- (j) Upper bounds on asymptotics of  $R(C_n^{r,1}, K_m; r)$  for even and odd  $n$  [ColGJ]. Improvements of the results from the latter, in particular for the case of  $n=5$  and  $r=3$ , and for general  $n$  [Mér].
- (k) Summary of known values and ranges for hypergraph numbers for loose paths (and some other trees) versus complete hypergraphs,  $R(P_m, K_n; 3)$ , for  $n \leq 10$  and odd  $m$  [BudPe].
- (l) Study of the growth rate of  $R(P_m, K_n; r)$  for tight paths  $P_m$  with  $m \geq r+3$ , and links between the growth of  $R(P_{r+1}, K_n; r)$  and  $R(n, n; r)$  [MuSuk1]. The correct tower growth rate for ordered tight paths versus cliques [Mub2]. Growth rate of  $R(P_{n+2}, I_n; 3)$ , where  $I_n$  are ordered 3-uniform monotone paths with  $n$  jumps, and its relation to the classical problem of growth of  $R_n(3)$  [SukZ], see items 6.2.b/c/d/e/f.
- (m) Study of  $R(G, nH; r)$  and  $R(mG, nH; r)$  for  $G$  and  $H$  being loose/tight path, cycles and stars, including several exact results for large  $m$  or  $n$  [OmRa3]. The case of loose  $t$ -tight paths versus stars and some tripartite hypergraphs is explored in [BudHR2].
- (n) Let  $F$  be the Fano plane, seen as a 3-uniform hypergraph of 7 hyperedges. If  $P_n$  and  $C_n$  are tight path and cycle on  $n$  vertices, respectively, then for sufficiently large  $n$  we have

$$R(P_n, F; 3) = 2n - 1 \text{ and } R(C_n, F; 3) = 2n - 1 \text{ [BalCSW].}$$

### 7.2.2. More colors

- (a) For loose cycles,  $R_3(C_3; 3) = R(C_3, C_3, C_3; 3) = 8$ , and in general for  $k \geq 4$  colors Gyárfás and Raeisi established the bounds  $k + 5 \leq R_k(C_3; 3) \leq 3k$  [GyRa].
- (b) For loose paths, we have  $R_3(P_3; 3) = 9$  and  $10 \leq R_4(P_3; 3) \leq 12$  [Jack]. This was improved to  $R_k(P_3; 3) = k + 6$  for all  $2 \leq k \leq 9$  [JacPR, PoRu], and extended to  $k = 10$  [Pol]. The general upper bound  $R_k(P_3; 3) \leq 2k + \sqrt{18k + 1} + 2$  was obtained in [ŁuPo1], then improved to  $R_k(P_3; 3) \leq 1.975k + 7\sqrt{k} + 2$  [ŁuPo2], and then further improved to  $R_k(P_3; 3) \leq 1.546k$  for large  $k$  [BohZ]. For the *messy path*  $M_3 = \{abc, bcd, def\}$ , we have  $R_k(M_3; 3) \leq 1.6k$  for large  $k$  [BohZ].
- (c) The case of  $R_k(P_3; r)$  for loose path was asymptotically solved in [ŁuPR]. A general upper bound  $R_k(P_n; r) \leq (r - 1)kn$ , for all  $k, r \geq 2$  and  $n \geq 3$ , was obtained in [DuRu]. Do not be confused by notation in some of the papers when their bounds are expressed in terms of  $r$  colors and  $k$ -uniform paths (in contrast to  $k$ -colors and  $r$ -uniform used in this survey).
- (d) For tight paths  $P_{m_i}$ , study of the growth rate of  $R(P_{m_1}, \dots, P_{m_k}, K_m; r)$  [MuSuk1].
- (e) Study of Turán and Ramsey numbers of sets of minimal 3-uniform paths of length 4 for up to 4 colors [HanPR]. Minimality of path here means that there are no redundant edge intersections, in particular no vertex belongs to more than two edges.

### 7.3. Berge cycles and Berge hypergraphs

A 3-uniform *Berge cycle* is formed by  $n$  distinct vertices, such that all consecutive pairs of vertices are in an edge of the cycle, and all of the cycle edges are distinct. 3-uniform Berge cycles are not determined uniquely.

- (a) For 3-uniform Berge cycles and two colors, we have  $R(C_n, C_n; 3) = n$  for  $n \geq 5$  [GyLSS]. Some results for Berge-cycle versus Berge-complete numbers are obtained in [MahS], and the asymptotics of Berge-cycles versus complete hypergraph is derived in [NieVer].
- (b) Gyárfás, Sárközy and Szemerédi proved that, for sufficiently large  $n$ , every 2-coloring of the edges of the complete 4-uniform hypergraph  $K_n$  contains a monochromatic 3-tight Berge cycle  $C_n$  [GySS2]. Exact formulas and bounds for Berge- $K_n$  hypergraphs, including higher uniformity  $r$  [SaTWZ].
- (c) For 3-uniform Berge cycles, we have  $R_3(C_n; 3) = (1 + o(1))5n/4$  [GySá1], and also  $R(C_n, C_n, C_3; 3) = n + 1$  [MahS]. Some special cases for  $r$ -uniform hypergraphs with respect to Berge cycles were studied in [GyLSS], and for small uniformity  $r$  in [DeMST]. See also the next two items.

- (d) Study of 3-uniform Berge- $G$  graphs in  $r$  colors: asymptotic lower and upper bounds, and several exact values for small  $r$  with  $G = K_3$  or  $G = K_4$ . Some asymptotics in the nonuniform case [AxGy]. This extends the results in 7.3.a [GyLSS] and 7.3.c.
- (e) Exact formulas and bounds for Berge- $K_n$  hypergraphs, including multiple colors [AxGy] and higher uniformity  $r$  [AxGy, SaTWZ]. Progress on the conjecture that every  $(r-1)$ -coloring of  $K_n^r$ , for fixed  $r$  and large  $n$ , contains a monochromatic Hamiltonian Berge cycle [MaOm2]. Determination of some cases of uniformity  $r$ , number of colors and  $G$ , for which the Ramsey number of Berge- $G$  is superlinear [Gerbn]. Further study of multicolor Ramsey numbers for such Berge- $G$  hypergraphs, with some equalities and improved asymptotics, were obtained in [GerMOV, Pálv].
- (f) Exact order of growth of the multicolor numbers for Berge cycles of length 4, 5, 6, 7, 10 and 11 for small uniformities, and for multicolor cases of Berge- $K_{a,b}$  for certain  $a, b$  and  $r$ , were derived in [DeMST].
- (g) The growth rate of  $R(F, K_n; 3)$  for  $F = \{123, 124, 345\}$  [MatMNV].

#### 7.4. General results for 3-uniform hypergraphs

- (a)  $2^{cn^2} < R(n, n; 3) < 2^{2^n}$  is credited to Erdős, Hajnal and Rado (see [ChGra2] p. 30).
- (b) For some  $a, b$  the numbers  $R(m, a, b; 3)$  are at least exponential in  $m$  [AbbS].
- (c) Improved lower and upper asymptotics for  $R(s, n; 3)$  for fixed  $s$  and large  $n$ , proof of related Erdős and Hajnal conjecture on the growth of  $R(4, n; 3)$ , and the lower bound  $2^{n^{c \ln n}} < R(n, n, n; 3)$  [ConFS2].
- (d) Let  $S_n$  denote the 3-uniform star on  $n+1$  vertices consisting of all  $n(n-1)/2$  triangles sharing one vertex. Then we have  $2^{c \log^2 n} \leq R(K_4, S_n; 3) \leq 2^{c' n^{2/3} \log n}$ , for some positive constants  $c, c'$  [ConFH+]. Study of the rate of growth of  $R(K_n, H; 3)$  for fixed hypergraph  $H$  [ConFG1+]. Further progress in [ConFG2+].
- (e) The hedgehog  $H_t$  is a 3-uniform hypergraph with  $t+t(t-1)/2$  vertices such that for every  $(i, j)$  with  $1 \leq i < j \leq t$  there exists a unique vertex  $k > t$  such that  $ijk$  is an edge, and  $H_t$  has no other edges. Conlon, Fox and Rödl studied the bounds on  $R_k(H_t; 3)$  for  $2 \leq k \leq 4$  and large  $t$  [ConFR]. The hypergraphs  $H_t$  constitute the first family of hypergraphs whose Ramsey numbers show a strong dependence on the number of colors: their 2-color Ramsey numbers grow polynomially in  $t$ , while in the 4-color case they grow exponentially.  $R_k(H_t; 3) = O(t^2 \ln t)$  was obtained in [FoxLi].
- (f) For two colors and some 3-uniform hypergraphs related to hedgehogs: there exists a 1-degenerate hypergraph  $G$  with  $R(G, G; 3) = \Omega(n^{3/2}/\log n)$ , and for generalized hedgehogs  $H$  we have  $R(H, H; 3) = O(n^{3/2})$  [AllBP].
- (g)  $R(G, G; 3) \leq cn(H)$  for some constant  $c$  depending only on the maximum degree of a 3-uniform hypergraph  $H$  [CooFKO1, NaORS]. Similar results were proved for  $r$ -uniform hypergraphs in [KüCFO, Ishi, CooFKO2, ConFS1], see also item 7.5.g.

- (h) Asymptotic lower bounds for  $R(K_{a,b,c}, K_{a,b,c}; 3)$ , where  $K_{a,b,c}$  is formed by all  $abc$  triples on sets of orders  $a, b, c$ . The conference and journal versions of this work [MiPal] differ on some results. See also item 7.1.g.
- (i) Asymptotic or exact values of  $R_k(H; 3)$  when  $H$  is a bow  $\{abc, ade\}$ , kite  $\{abc, abd\}$ , tight path  $P_3^{3,2} = \{abc, bcd, cde\}$ , or windmill  $\{abc, bde, cef, bce\}$ , and a special concrete case for 6-color kite,  $R_6(\text{kite}; 3) = 8$ .
- (j)  $R_k(K_3; 2) \leq R_{4k}(K_{4-t}; 3) \leq R_{4k}(K_3; 2) + 1$  [AxGLM].
- (k) Variety of general lower bound constructions for 3-uniform complete or complete missing one hyperedge hypergraphs from liftings of graphs, for two and more colors. For example, we have  $R(K_{2s_1-1-t}, K_{2s_2-1}, K_{2s_3-1}; 3) \geq R(s_1, s_2, s_3)$  [BudHMP] and  $R(K_5, K_{43-t}, K_{43-t}, K_{43-t}; 3) > 1257480$  [BudHLS].
- (l) If  $G$  is a 3-uniform  $H$ -free hypergraph, then  $G$  contains a complete or empty tripartite subgraph with parts of order  $(\log n(H))^{c+1/2}$ , where  $c > 0$  depends only on  $H$ . Furthermore, for  $k \geq 4$  no analogue of it can hold for  $k$ -uniform hypergraphs [ConFS5].
- (m) Upper bounds on  $R_k(H; 3)$  for complete multipartite 3-uniform hypergraphs  $H$ , a 4-color case, and some other general and special cases [ConFS1, ConFS2, ConFS3].  $R_k(H; 3)$  ranges from  $\sqrt{6k}(1+o(1))$  to double exponential in  $k$  [AxGLM]. Determination when  $R_k(H; 3)$  behaves polynomially, exponentially or double-exponentially in  $k$  [BraFS1].

### 7.5. General results

- (a) If  $R(n, n; r) > m$ , then we have  $R(2n+r-4, 2n+r-4; r+1) > 2^m$  for all  $n > r \geq 3$  (see [GRS] p. 106). This is the so-called stepping-up lemma, usually credited to Erdős and Hajnal. An improvement of the stepping-up lemma implying better lower bounds for a few types of hypergraph Ramsey numbers was obtained by Conlon, Fox and Sudakov [ConFS6].
- (b) For  $k$  colors, lower bounds on  $R_k(n; r)$  are discussed in [AbbW, DuLR, ConFS6, AxGLM, JaKSY]. In particular, if  $R_k(n; r) > m$ , then we have  $R_{k+3}(n+1; r+1) > 2^m$  for  $r \geq 3$  [JaKSY].
- (c) General lower bounds for large number of colors were given in an early paper by Hirschfeld [Hir], and some of them were later improved in [AbbL].
- (d) Lower and upper asymptotics of  $R(s, n; k)$  for fixed  $s$  [ConFS2, MuSuk2, MuSuk3]. Lower bounds for  $R(k+1, k+1; k)$ ,  $R(k+1, k+2; k)$ ,  $R(k+1, 2k+1; k)$  and  $R(k+2, k+2; k)$  in terms of the exp-tower function  $\text{tw}_k$  [PuRW].
- (e) Exact and asymptotic results generalizing 7.2.1.d/e to  $r$ -uniform case for cycles, and 2- and 3-color cases for all  $r$ -uniform diamond matchings [GySS1]. Connections between the existence of large sets of combinatorial  $t$ -designs to some bounds on Ramsey-type problems [Gy].
- (f) Extension of results from 7.4.d to higher uniformities [HeNWY].

- (g) Study of  $R(G, nH; r)$  and  $R(mG, nH; r)$  for loose/tight path and cycles (possibly with some additions), stars,  $r$ -partite hypergraphs, including several exact results for large  $m$  or  $n$  [OmRa3].
- (h)  $R(H, H; r) \leq cn(H)^{1+\varepsilon}$ , for some constant  $c = c(\Delta, r, \varepsilon)$  depending only on the maximum degree of  $H$ ,  $r$  and  $\varepsilon > 0$  [KoRö3]. The proofs of the linear bound  $cn(H)$  were obtained independently in [KüCFO] and [Ishi], the latter including the multicolor case, and then without regularity lemma in [ConFS1]. More discussion of lower and upper bounds for various cases can be found in [ConFS1, ConFS2, ConFS3, CooFKO2].
- (i) Let  $T_r$  be an  $r$ -uniform hypergraph with  $r$  edges containing a fixed  $(r-1)$ -vertex set  $S$  and the  $(r+1)$ -st edge intersecting all former edges in one vertex outside  $S$ . Then  $R(T_r, K_r; r) = O(t^r/\log t)$  [KosMV1].
- (j) Study of tree-star and tree-complete cases of Ramsey numbers for  $r$ -uniform hypergraphs [Nie]. Several bounds and equalities for special cases [BudHR1, BudCli]. This was posed and explored as a problem of which trees are Ramsey  $n$ -good hypergraphs [BudPe]. Further results towards the conjecture that all  $r$ -uniform trees are  $n$ -good [BudCli]. Study of the Ramsey numbers of disjoint union of  $H$ -good hypergraphs [KamRa1].
- (k) Let  $H^r(s, t)$  be the complete  $r$ -partite  $r$ -uniform hypergraph with  $r-2$  parts of size 1, one part of size  $s$ , and one part of size  $t$  (for example, for  $r=2$  it is the same as  $K_{s, t}$ ). For the multicolor numbers, Lazebnik and Mubayi [LaMu] proved that

$$tk^2 - k + 1 \leq R_k(H^r(2, t+1); r) \leq tk^2 + k + r,$$

where the lower bound holds when both  $t$  and  $k$  are powers of the same prime  $p$ . For the general case of  $H^r(s, t)$ , more bounds are presented in [LaMu].

- (l)  $R_k(H; r)$  is polynomial in  $k$  when a fixed  $r$ -uniform  $H$  is  $r$ -partite, otherwise it is at least exponential in  $k$  [AxGLM].
- (m) Grolmusz [Grol1] generalized the classical constructive lower bound by Frankl and Wilson [FraWi] (item 2.4.7) to more colors and to hypergraphs [Grol3].
- (n) Lower and upper asymptotics, and other theoretical results on hypergraph numbers, are discussed in [GrRö, GRS, ConFS2, ConFS3, ConFS7, Song8, MuSuk1, MuSuk2, MuSuk3]. An extensive overview of progress and open problems in hypergraph Ramsey theory by Mubayi and Suk was compiled in 2018 [MuSuk4].
- (o) For fixed uniformity  $r$  and  $k$ -colors, the Ramsey number of any  $r$ -uniform hypergraph with  $m$  edges is at most  $\text{tw}_r(O(\sqrt{m}))$ , where  $\text{tw}$  is the exp-tower function [BraFS2]. See also [ConFS1, ConFS4].
- (p) Ramsey goodness of  $r$ -uniform paths, or the lack thereof; or the contrast with and similarities to the concept of Ramsey goodness for graphs [BoLo].
- (q) Lower bounds for Ramsey numbers of bounded degree hypergraphs [BraHS].
- (r) Bounds on the Ramsey numbers of daisies [PuRS], enhanced in [Sal].

## 8. Cumulative Data, Books and Surveys

### 8.1. Cumulative data for two colors

- (a)  $R(G, G)$  for all graphs  $G$  without isolates on at most 4 vertices [ChH1].
- (b)  $R(G, H)$  for all graphs  $G$  and  $H$  without isolates on at most 4 vertices [ChH2].
- (c)  $R(G, H)$  for all graphs  $G$  on at most 4 vertices and  $H$  on 5 vertices, except five entries [Clan], now all solved, see Section 5.12. All critical colorings for the isolate-free graphs  $G$  and  $H$  studied in [Clan] were found in [He4].
- (d)  $R(G, G)$  for all graphs  $G$  without isolates and with at most 6 edges [Bu4],  
 $R(G, G)$  for all graphs  $G$  without isolates and with at most 7 edges [He1],  
 $R(G, G)$  for all graphs  $G$  on 5 vertices and with 7 or 8 edges [HaMe2].
- (e)  $R(G, H)$  for all graphs  $G$  and  $H$  on 5 vertices without isolates, except 7 entries [He2]. Only 2 cases are still open, see Section 5.12 and the paragraph at the end of this section.
- (f) Tables of  $R(G, H)$  for most connected graphs on up to 5 vertices and  $R(G, G)$  for all isolate-free graphs with up to 7 edges [ReWi].
- (g)  $R(G, H)$  for all disconnected isolate-free graphs  $H$  on at most 6 vertices versus all  $G$  on at most 5 vertices, except 3 cases [LoM5]. Missing cases were completed in [KroMe].
- (h)  $R(G, H)$  for some  $G$  on 5 vertices versus all connected graphs on 6 vertices [LoM6].
- (i)  $R(G, H)$  for  $G = K_{1,3} + e$  and  $G = K_4 - e$  versus all connected graphs  $H$  on 6 vertices, except  $R(K_4 - e, K_6)$  [HoMe]. The result  $R(K_4 - e, K_6) = 21$  was claimed by McNamara [McN, unpublished], now confirmed in [ShWR].
- (j)  $R(G, H)$  for some graphs  $G$  with 4 vertices versus all graphs  $H$  with 7 vertices [Boza4].
- (k)  $R(G, T)$  for all connected graphs  $G$  with  $n(G) \leq 5$ , and almost all trees  $T$  [FRS4].
- (l)  $R(T_m, T_n)$  for  $6 \leq m, n \leq 8$ , for  $k$ -vertex trees  $T_k$  [RanMCG].
- (m)  $R(K_3, G)$  for all connected graphs  $G$  on 6 vertices [FRS1].  
 $R(K_3, G)$  for all connected graphs  $G$  on 7 vertices [Jin],  
some errors in the latter were found [SchSch1].
- (n)  $R(K_3, G)$  for all connected graphs  $G$  on at most 8 vertices [Brin]. The numbers for  $K_3$  versus sets of graphs with fixed number of edges, on at most 8 vertices, were presented in [KlaM1].  $R(K_3, G)$  for all connected graphs  $G$  on 9 vertices [BrBH1, BrBH2].
- (o)  $R(K_3, G)$  for all graphs  $G$  on 10 vertices, except 10 cases [BrGS].  
Three of the open cases, including  $G = K_{10} - e$ , were solved [GoR2].
- (p)  $R(S, G)$  for stars  $S$  versus almost all connected graphs  $G$  on 6 vertices [LoM7]. This was extended to  $R(T, G)$  for most non-star trees  $T$ , in particular for all trees on at most 5 vertices versus all connected graphs  $G$  on at most 6 vertices [LoM9]. The values and bounds for still missing cases are presented in [LoM8, LoM10].
- (q) Formulas for  $R(nK_3, mG)$  for all  $G$  of order 4 without isolates [Zeng].

- (r)  $R(C_4, G)$  for all graphs  $G$  on at most 6 vertices [JR3],  
this work was followed by two errata listed in the references.  
 $R(C_5, G)$  for all graphs  $G$  on at most 6 vertices [JR4],  
 $R(C_6, G)$  for all graphs  $G$  on at most 5 vertices [JR2].
- (s)  $R(K_{2,n}, K_{2,m})$  for all  $2 \leq n, m \leq 10$  except 8 cases, for which lower and upper bounds are given [LoM3]. Further data for other complete bipartite graphs are gathered in Section 3.3 and [LoMe4].
- (t) All best lower bounds up to 102 from cyclic graphs. Formulas for best cyclic lower bounds for paths and cycles, and values for small complete graphs and for graphs with up to five vertices [HaKr1].

Chvátal and Harary [ChH1, ChH2] formulated several simple but very useful observations on how to discover values of some numbers. All five missing entries in the tables of Clancy [Clan] have been solved (Section 5.12). Out of 7 open cases in [He2] 6 have been solved, including  $R(4, 5) = R(G_{19}, G_{23}) = 25$  and other cases listed in Section 5.12. The only still open case is for  $K_5$  versus  $K_5$  (Section 2.1). Many extremal and other Ramsey graphs for various parameters are available at [BrCGM, McK1, Ex18, Fuj1], see Section 8.3 below.

## 8.2. Cumulative data for three colors

- (a)  $R_3(G)$  for all graphs  $G$  with at most 4 edges and no isolates [YR3].
- (b)  $R_3(G)$  for all graphs  $G$  with 5 edges and no isolates, except  $K_4 - e$  [YR1].  
This last open case is now solved, namely, we have  $R_3(K_4 - e) = 28$  [Ex7][LidP].
- (c)  $R_3(G)$  for all graphs  $G$  with 6 edges and no isolates, except 10 cases [YY].
- (d)  $R(F, G, H)$  for many triples of isolate-free graphs with at most 4 vertices [ArKM].  
Some of the missing cases completed in [KlaM2].
- (e) Extension of [ArKM] to most triples of graphs with at most 4 vertices [BoDD].
- (f)  $R(P_3, P_k, C_m)$  for all  $3 \leq k \leq 8$  and  $3 \leq m \leq 9$  [DzFi2].

## 8.3. Electronic resources

- (a) W. Gasarch [Gas] maintains a website gathering over 60 pointers to literature on applications of Ramsey theory in computer science, and in particular logic, complexity theory and algorithms, <http://www.cs.umd.edu/~gasarch/TOPICS/ramsey/ramsey.html>.
- (b) Many of the Ramsey graph constructions found by G. Exoo [Ex1-Ex25] are posted at <http://cs.indstate.edu/ge/RAMSEY>.
- (c) G. Brinkmann, K. Coolsaet, J. Goedgebeur and H. Mélot, *House of Graphs: A database of interesting graphs* [BrCGM], <http://hog.grinvin.org>.

- (d) B.D. McKay, presents some graphs related to classical Ramsey numbers [McK1], <http://cs.anu.edu.au/people/bdm/data/ramsey.html>.
- (e) Set of Ramsey problems with comments and references by a team of students of Fan Chung, University of California San Diego [UCSD], *Erdős' Problems on Graphs, Ramsey Theory*, <http://www.math.ucsd.edu/~erdosproblems/RamseyTheory.html> (2010-2012).
- (f) H. Fujita, collection of Ramsey graph constructions [Fuj1], <http://opal.inf.kyushu-u.ac.jp/~fujita/ramsey.html>, (2014-2017).
- (g) M. Rubey, an electronic GUI resource for values of some small Ramsey numbers [Rub], <http://www.findstat.org/StatisticsDatabase/St000479>.
- (h) S. Van Overberghe, Ramsey graph constructions associated with MS Thesis, Ghent University, Belgium, 2020 [VO], <https://github.com/Steven-VO/circulant-Ramsey>. Extended by many newer lower bound constructions involving wheels and books [VO1, LidMPV] pointed to in Sections 5.2 and 5.3.
- (i) A.E. Brouwer, Parameters of Strongly Regular Graphs [Brou], used mainly in 3.1.d and 5.3.1, <https://www.win.tue.nl/~aeb/graphs/srg/srgtab.html>.

#### 8.4. Books and surveys

##### Books and special works

- (1980) *Ramsey Theory* by R.L. Graham, B.L. Rothschild and J.H. Spencer [GRS], first edition 1980, second edition 1990, paperback of the second edition 2013.
- (1983) Special volume of the *Journal of Graph Theory* [JGT].
- (1996) A chapter in *Handbook of Combinatorics* by J. Nešetřil [Neš].
- (1997) Among 114 open problems and conjectures of Paul Erdős, presented and commented by F.R.K. Chung, 31 are concerned directly with Ramsey numbers [Chu4]. 216 references are given. An extended version of this work was prepared jointly with R.L. Graham [ChGra2] in 1998.
- (2001) An extensive chapter on Ramsey theory in a widely used student textbook and researcher's guide of graph theory by D. West [West].
- (2002) *Ramsey Theory and Paul Erdős* by R.L. Graham and J. Nešetřil [GrNe].
- (2003) Special issue of *Combinatorics, Probability and Computing* [CoPC].
- (2004) *Ramsey Theory on the Integers* by B. Landman and A. Robertson [LaRo], first edition 2004, second edition 2014.
- (2009) History, results and people of Ramsey theory. *The mathematical coloring book, mathematics of coloring and the colorful life of its creators* by A. Soifer [Soi1].
- (2011) *Ramsey Theory. Yesterday, Today and Tomorrow*, a special volume in the series *Progress in Mathematics* [Soi2]. A survey of Ramsey numbers involving cycles by this author is included there [Ra4].

- (2015) *Recent Developments in Graph Ramsey Theory* by D. Conlon, J. Fox and B. Sudakov [ConFS7].
- (2015) *Rudiments of Ramsey Theory*, a new edition of the classics by R.L. Graham and S. Butler [GrBu].
- (2018) *Ramsey Theory, Unsolved Problems and Results* by Xiaodong Xu, Meilian Liang and Haipeng Luo [XuLL].
- (2021) *The Discrete Mathematical Charms of Paul Erdős: A Simple Introduction* by Vašek Chvátal [Chv2].
- (2022) *Elementary Methods of Graph Ramsey Theory*, an extensive presentation of the area by Yusheng Li and Qizhong Lin [LiLin2].
- (2023) *Basics of Graph Ramsey Theory* by Veselin Jungić [Jun].
- (2024) *The New Mathematical Coloring Book, Mathematics of Coloring and the Colorful Life of Its Creators*, greatly extended and revised first edition of the 2009 coloring book [Soi1] by Alexander Soifer [Soi3].

### Surveys and overviews

- (1974) A general survey of results in Ramsey graph theory by S.A. Burr [Bu0]
- (1978) A general survey of results in Ramsey graph theory by T.D. Parsons [Par6]
- (1979) A survey of noncomplete Ramsey graph theory by S.A. Burr [Bu1]
- (1980) Survey of results and new problems on multiplicities and Ramsey multiplicities by S.A. Burr and V. Rosta [BuRo3]
- (1981) Summary of progress by Frank Harary [Har2]
- (1983) A survey of bounds and values by F.R.K. Chung and C.M. Grinstead [ChGri]
- (1984) A review of Ramsey graph theory for newcomers by F.S. Roberts [Rob1]
- (1987) An overview of progress so far and plans for the future, *What Can We Hope to Accomplish in Generalized Ramsey Theory?* by S. Burr [Bu7]
- (1987) Survey of asymptotic problems by R.L. Graham and V. Rödl [GrRö]
- (1991) Survey by R.J. Faudree, C.C. Rousseau and R.H. Schelp of graph goodness results, i.e. conditions for the formula  $R(G, H) = (\chi(G) - 1)(n(H) - 1) + s(G)$  [FRS5]
- (1996) Survey of zero-sum Ramsey theory by Y. Caro [Caro]
- (2004) Dynamic survey of Ramsey theory applications by V. Rosta [Ros2]. A website maintained by W. Gasarch [Gas] gathers over 60 pointers to literature on applications of Ramsey theory in computer science.
- (2009) *Ramsey Goodness and Beyond* by V. Nikiforov and C.C. Rousseau [NiRo4]
- (2010) *Recent Developments in Extremal Combinatorics* by B. Sudakov [Sud4]
- (2010) *Hypergraph Ramsey Numbers* by D. Conlon, J. Fox and B. Sudakov [ConFS2].
- (2013) *Ramsey-Goodness - and Otherwise* by P. Allen, G. Brightwell and J. Skokan [AllBS]

- (2013) *Problems in Graph Theory from Memphis*, "a summary of problems and results coming out of the 20 year collaboration between Paul Erdős and the authors", by R.J. Faudree, C.C. Rousseau and R.H. Schelp [FRS6].
- (2016) *On Some Open Questions for Ramsey and Folkman Numbers* [XuR4].
- (2018) *A Survey of Hypergraph Ramsey Problems* by D. Mubayi and A. Suk [MuSuk4].
- (2020) *An Introduction to Ramsey's Theorem* by A. Tripathi [Tri].
- (2021) *New Directions in Ramsey Theory* by Gary Chartrand and Ping Zhang [ChaZ].
- (2024) *Upper Bounds on Diagonal Ramsey Numbers* [after Campos, Griffiths, Morris, and Sahasrabudhe], by Yuval Wigderson [Wig3].
- (2025) Survey of two-color and multicolor Ramsey numbers involving  $C_4$  by Chen Yaojun, Zhang Xuemei and Zhang Yanbo [ChenZZ7].
- (2025) *Numbers in Ramsey Theory* by Fan Chung, a nice survey of open problems involving Ramsey numbers, in the first issue of a new journal *Journal of Open Mathematical Problems* [Chu5].
- (2025) *Recent Progress in Ramsey Theory* by Jacques Verstraëte [Ver].
- (2026) A gentle introduction to recent upper bound breakthroughs by Robert Morris [Mor].
- (2026) *Ramsey Theory*, an extensive chapter in *Basic Graph Theory* book by Béla Bollobás and Robert Morris [BolM].

The surveys by S.A. Burr [Bu0, Bu1] and T.D. Parsons [Par6] contain extensive chapters on general exact results in graph Ramsey theory. F. Harary presented the state of the theory in 1981 in [Har2], where he also gathered many references including seven to other early surveys of this area. More than three decades ago, Chung and Grinstead in their survey paper [ChGri] gave much less data than in this work, but they included a broad discussion of different methods used in Ramsey computations in the classical case. S. A. Burr, one of the most experienced researchers in Ramsey graph theory, formulated in [Bu7] seven conjectures on Ramsey numbers for sufficiently large and sparse graphs, and reviewed the evidence for them found in the literature. Three of them have been refuted in [Bra3].

For newer extensive presentations see [GRS, GrRö, FRS5, Neš, Chu4, ChGra2, ConFS7], though these focus on asymptotic theory not on the numbers themselves. A very welcome addition is the 2004 compilation of applications of Ramsey theory by V. Rosta [Ros2]. This survey could not be complete without recommending special volumes of the *Journal of Graph Theory* [JGT, 1983] and *Combinatorics, Probability and Computing* [CoPC, 2003], which, besides a number of research papers, include historical notes and present to us Frank P. Ramsey (1903-1930) as a person. Read a colorful book by A. Soifer [Soi1, 2009] on history and results in Ramsey theory, followed by a collection of essays and technical papers based on presentations from the 2009 Ramsey theory workshop at DIMACS [Soi2, 2011]. A 70-page long paper from 2015, entitled *Recent Developments in Graph Ramsey Theory*, by D. Conlon, J. Fox and B. Sudakov [ConFS7] documents in details what the title says. Finally, since the previous revision #16 of this survey, some new and revised books appeared: in 2021 by V. Chvátal [Chv2], in 2022 by Li and Lin [LiLin2], and in 2024 an extended and revised book by Soifer [Soi3].

The historical perspective and, in particular, the timeline of progress on prior best bounds, can be obtained by comparing all the previous versions of this survey since 1994 at the author's website <http://www.cs.rit.edu/~spr/EIJC/eline.html> or at the portal of the *E-JC* journal <http://www.combinatorics.org/ojs/index.php/eljc/issue/view/Surveys>.

## 9. Concluding Remarks

### 9.1. Exclusions

This compilation does not include much information on numerous variations of Ramsey numbers, nor related topics, like

anti-Ramsey numbers,	oriented Ramsey numbers,
ascending Ramsey index,	oriented size Ramsey numbers,
avoiding sets of graphs,	pattern Ramsey numbers,
barycentric Ramsey numbers,	planar Ramsey numbers,
bipartite Ramsey numbers,	poset Ramsey numbers,
blowup Ramsey numbers,	potential Ramsey numbers,
canonical Ramsey numbers,	proper Ramsey numbers,
chromatic Ramsey numbers,	quantum Ramsey numbers,
class Ramsey numbers,	quasi-Ramsey numbers,
complementary Ramsey numbers,	rainbow Ramsey numbers,
connected Ramsey numbers,	Ramsey equivalence,
connected size Ramsey numbers,	Ramsey game numbers,
cover Ramsey numbers,	Ramsey games,
defective Ramsey numbers,	Ramsey-minimal graphs,
degree Ramsey numbers,	Ramsey multiplicities,
directed Ramsey numbers,	Ramsey-Turán numbers,
distance Ramsey numbers,	Ramsey sequences of graphs,
edge-chromatic Ramsey numbers,	regular Ramsey numbers,
edge-disjoint Ramsey numbers,	restricted online Ramsey numbers,
edge-ordered Ramsey numbers,	restricted size Ramsey numbers,
Folkman numbers,	Schur numbers,
Gallai-Ramsey numbers,	semi-algebraic Ramsey numbers,
generalized Ramsey numbers,	set-coloring Ramsey numbers,
induced Ramsey numbers,	Sidon-Ramsey numbers,
inverse Ramsey numbers,	signed Ramsey numbers,
irredundant Ramsey numbers,	singular Ramsey numbers,
list Ramsey numbers,	size multipartite Ramsey numbers,
local Ramsey numbers,	size Ramsey numbers,
$k$ -Ramsey numbers,	star-critical Ramsey numbers,
mixed Ramsey numbers,	sub-Ramsey numbers,
multipartite Ramsey numbers,	Van der Waerden numbers,
odd-Ramsey numbers,	vertex size Ramsey numbers,
online Ramsey numbers,	weakened Ramsey numbers,
ordered Ramsey numbers,	zero-sum Ramsey numbers,
ordered size Ramsey numbers,	or coloring graphs other than complete.

Interested readers can find such information in some of the surveys listed in Section 8 here, or just enter it into the search box of a web browser for a quick summary and further pointers.

Readers may be also interested in knowing that the US patent 6965854 B2 issued on November 15, 2005 claims a method of using Ramsey numbers in "Methods, Systems and Computer Program Products for Screening Simulated Traffic for Randomness." Check the original document at <http://www.uspto.gov/patft> if you wish to find out whether your usage of Ramsey numbers is covered by this patent.

## 9.2. Journal paper counts

Out of 1066 references gathered in this survey, most are papers which appeared in more than 100 different periodicals (in addition to books, conference proceedings, arXiv postings, and personal communications). The most popular periodicals were:

<i>Discrete Mathematics</i>	110
<i>Journal of Combinatorial Theory</i> (old, Series A and B)	65
<i>Journal of Graph Theory</i>	65
<i>Electronic Journal of Combinatorics</i>	57
<i>Graphs and Combinatorics</i>	48
<i>Ars Combinatoria</i>	33
<i>European Journal of Combinatorics</i>	32
<i>Journal of Combinatorial Mathematics and Combinatorial Computing</i>	32
<i>Discrete Applied Mathematics</i>	28
<i>Australasian Journal of Combinatorics</i>	25
<i>Combinatorica</i>	22
<i>Utilitas Mathematica</i>	19
<i>SIAM Journal on Discrete Mathematics</i>	18
<i>Combinatorics, Probability and Computing</i>	17
<i>Discussiones Mathematicae Graph Theory</i>	12
<i>Congressus Numerantium</i>	12
<i>Random Structures and Algorithms</i>	10
<i>Applied Mathematics Letters</i>	8
<i>Mathematica Applicata</i>	7
arXiv preprints	57

Some of the original arXiv pointers remain listed together with their later published versions, if we think that this gives additional information. Such arXiv postings are not included in the count 57 above. Please also remember that arXiv preprints are not peer-reviewed.

The results of 204 references depend on computer algorithms.

## Acknowledgements

I would like to thank many individuals, collaborators and casual readers, who helped me to improve and make more complete the consecutive versions of this survey. In particular, I am very grateful for their continuous encouragement and suggestions to add a new section or enhance parts related to specific areas.

The author would like to specially thank his Ramsey-collaborator Xiaodong Xu. He has seen more revisions of this survey than one would wish. The author would also like to extend his thanks to Brendan McKay, Geoffrey Exoo and Heiko Harborth for their help in gathering data throughout the years. The help of all those who contributed to the development and improvement of new revisions over the years is greatly appreciated. Special thanks go to Yanbo Zhang who recommended several changes to the latest revision #18.

The author apologizes for any omissions or other errors in reporting results belonging to the scope of this work. Suggestions for any kind of corrections or additions will be greatly appreciated and considered for inclusion in the next revision of this survey. Keep in mind, however, that this survey is already quite large, and the priority is given to the classical Ramsey numbers, in contrast to many other types listed in Section 9.1, *Exclusions*.

## References

The references are ordered alphabetically by the last name of the first author, and where multiple papers have the same first author they are ordered by the last name of the second author, etc. We preferred that all work by the same author be in consecutive positions. Unfortunately, this causes that some of the abbreviations are not in alphabetical order. For example, [BaRT] is earlier on the list than [BaLS]. We also wish to explain a possible confusion with respect to the order of parts and spelling of Chinese names. We put them without any abbreviations, often with the last name written first as is customary in original. Sometimes this is different from the citations in other sources. One can obtain all variations of writing any specific name by consulting the authors database of *Mathematical Reviews* at <http://www.ams.org/mathscinet/search>, or *zbMATH* (formerly *Zentralblatt für Mathematik*) at <http://www.zbmath.org/authors>.

Papers containing results obtained with the help of computer algorithms have been marked with stars. We identify two such categories of papers: those marked with \* involving some use of computers where the results are easily verifiable with some computations, and those marked with \*\* where cpu intensive algorithms have to be implemented to replicate or verify the results. The first category contains mostly constructions done by algorithms, while the second mostly nonexistence results or claims of complete enumerations of special classes of graphs.

A, Ba, Bl, Bu	page 88
Ca, Cl, D, E	page 97
Fa, Fi, Ga, Gu, Ha, He, I	page 105
J, K, La, Li, Lia, Lo	page 115
M, N, O, P, Q, Ra, Ro	page 125
Sa, Sh, Si, Su, Sun	page 134
T, U, V, W, X, Y, Z	page 141 - page 149

## A

- [Abb1] H.L. Abbott, Some Problems in Combinatorial Analysis, *Ph.D. thesis*, Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, 1965.
- [Abb2] H.L. Abbott, A Theorem Concerning Higher Ramsey Numbers, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) Vol. 1, 25-28, Colloq. Math. Soc. János Bolyai, Vol. 10, North-Holland, Amsterdam, 1975.
- [AbbH] H.L. Abbott and D. Hanson, A Problem of Schur and Its Generalizations, *Acta Arithmetica*, **20** (1972) 175-187.
- [AbbL] H.L. Abbott and Andy Liu, Remarks on a Paper of Hirschfeld Concerning Ramsey Numbers, *Discrete Mathematics*, **39** (1982) 327-328.
- [AbbS] H.L. Abbott and M.J. Smuga-Otto, Lower Bounds for Hypergraph Ramsey Numbers, *Discrete Applied Mathematics*, **61** (1995) 177-180.
- [AbbW] H.L. Abbott and E.R. Williams, Lower Bounds for Some Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **16** (1974) 12-17.
- [-] Adiwijaya, see [SuAM, SuAAM].
- [AbdMT] Kh.Kh. Abdullin, D.B. Mokeev and D.S. Taletskii, On the Ramsey Number  $R(K_{1,s}, P_t)$  (in Russian), *Matematicheskie Zametki*, **119**(1) (2026) 3-8.

- [AgCP+]\* R. Ageron, P. Casteras, T. Pellerin, Y. Portella, A. Rimmel and J. Tomasik, New Lower Bounds for Schur and Weak Schur Numbers, *preprint*, <http://arxiv.org/abs/2112.03175> (2022).
- [AjKS] M. Ajtai, J. Komlós and E. Szemerédi, A Note on Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **29** (1980) 354-360.
- [AjKSS] M. Ajtai, J. Komlós, M. Simonovits and E. Szemerédi, Erdős-Sós Conjecture, *in preparation* (2013).
- [AhS] I. Ahme and A. Scott, Graphs with Arbitrary Ramsey Number and Connectivity, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.76, **31**(4) (2024), 7 pages.
- [AleALZ] B. Alecu, A. Atminas, V. Lozin and V. Zamaraev, Graph Classes with Linear Ramsey Numbers, *Discrete Mathematics*, **344** (2021) 112307, 14 pages.
- [AliBB] K. Ali, A.Q. Baig and E.T. Baskoro, On the Ramsey Number for a Linear Forest Versus a Cocktail Party Graph, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **71** (2009) 173-177.
- [AliBas] K. Ali and E.T. Baskoro, On the Ramsey Numbers for a Combination of Paths and Jahangirs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **65** (2008) 113-119.
- [AliBT1] K. Ali, E.T. Baskoro and I. Tomescu, On the Ramsey Numbers for Paths and Generalized Jahangir Graphs  $J_{s,m}$ , *Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.)*, **51**(99) (2008) 177-182.
- [AliBT2] K. Ali, E.T. Baskoro and I. Tomescu, On the Ramsey Number for Paths and Beaded Wheels, *Journal of Prime Research in Mathematics*, **5** (2009) 133-138.
- [AliSur] K. Ali and Surahmat, A Cycle or Jahangir Ramsey Unsaturated Graphs, *Journal of Prime Research in Mathematics*, **2** (2006) 187-193.
- [AliTJ] K. Ali, I. Tomescu and I. Javaid, On Path-Sunflower Ramsey Numbers, *Mathematical Reports, Bucharest*, **17** (2015) 385-390.
- [AllBP] P. Allen, S. Boyadzhyska and M. Pavez-Signé, Ramsey Numbers for 1-Degenerate 3-Graphs, *preprint*, <http://arxiv.org/abs/2507.23623> (2025).
- [AllBS] P. Allen, G. Brightwell and J. Skokan, Ramsey-Goodness - and Otherwise, *Combinatorica*, **33** (2013) 125-160.
- [AllLPZ] P. Allen, T. Łuczak, J. Polcyn and Yanbo Zhang, The Ramsey Number of a Long Even Cycle Versus a Star, *Journal of Combinatorial Theory, Series B*, **162** (2023) 144-153.
- [AllMRS] P. Allen, D. Mergoni Cecchelli, B. Roberts and J. Skokan, The Ramsey Numbers of Squares of Paths and Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P2.11, **31**(2) (2024), 7 pages.
- [AlmBCL] J. Alm, P. Bahls, K. Coffey and C. Langhoff, Generalizing  $p$ -Goodness to Ordered Graphs, *preprint*, <http://arxiv.org/abs/1412.3071> (2020).
- [AlmHS] J.F. Alm, N. Hommowun and A. Schneider, Mixed, Multi-color, and Bipartite Ramsey Numbers Involving Trees of Small Diameter, *preprint*, <http://arxiv.org/abs/1403.0273> (2014).
- [Alon1] N. Alon, Subdivided Graphs Have Linear Ramsey Numbers, *Journal of Graph Theory*, **18** (1994) 343-347.
- [Alon2] N. Alon, Explicit Ramsey Graphs and Orthonormal Labelings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R12, **1** (1994), 8 pages.
- [AlBK] N. Alon, S. Ben-Shimon and M. Krivelevich, A Note on Regular Ramsey Graphs, *Journal of Graph Theory*, **64** (2010) 244-249.
- [AlKS] N. Alon, M. Krivelevich and B. Sudakov, Turán Numbers of Bipartite Graphs and Related Ramsey-Type Questions, *Combinatorics, Probability and Computing*, **12** (2003) 477-494.
- [AlPu] N. Alon and P. Pudlák, Constructive Lower Bounds for Off-Diagonal Ramsey Numbers, *Israel Journal of Mathematics*, **122** (2001) 243-251.
- [AlRö] N. Alon and V. Rödl, Sharp Bounds for Some Multicolor Ramsey Numbers, *Combinatorica*, **25** (2005) 125-141.

- [AlR6S] N. Alon, L. Rónyai and T. Szabó, Norm-Graphs: Variations and Applications, *Journal of Combinatorial Theory, Series B*, **76** (1999) 280-290.
- [-] A.A. Al-Rhayyel, see [BanAA].
- [-] A. Alrifai, see [BanAA].
- [Alw] R. Alweiss, Ramsey Numbers of Odd Cycles Versus Larger Even Wheels, *Discrete Mathematics*, **341** (2018) 981-989.
- [-] B.M.N. Alzaleq, see [BatJA, JaAl].
- [Ana1]\* C.S. Anabanti, A Counterexample on a Group Partitioning Problem, *Birkbeck Mathematics Preprint Series*, #37, Birkbeck University of London, (2017), 7 pages.
- [Ana2]\* C.S. Anabanti, The Ramsey Number  $R_4(3)$  Is Not Solvable by Group Partition Means, *Quasigroups and Related Systems*, **31** (2023) 165-174.
- [Ang1]\*\* V. Angelteit,  $R(3, 10) \leq 41$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.30, **32**(4) (2025), 11 pages.
- [Ang2]\*\* V. Angelteit, An Exact Value for the Ramsey Number  $R(K_5, K_5 - e)$ , *preprint*, <http://arxiv.org/abs/2602.11459> (2026).
- [AnM1]\*\* V. Angelteit and B.D. McKay,  $R(5, 5) \leq 48$ , *Journal of Graph Theory*, **89** (2018) 5-13.
- [AnM2]\*\* V. Angelteit and B.D. McKay, *personal communication* (2020).
- [AnM3]\*\* V. Angelteit and B.D. McKay, *personal communication* (2024).
- [AnM4]\*\* V. Angelteit and B.D. McKay,  $R(5, 5) \leq 46$ , *Journal of Graph Theory*, **11x** (2026) 1-11.
- [ArKM] J. Arste, K. Klamroth and I. Mengersen, Three Color Ramsey Numbers for Small Graphs, *Utilitas Mathematica*, **49** (1996) 85-96.
- [-] H. Assiyatun, see [HaABS, HaBA1, HaBA2, BaHA, SuAAM, SuAUB, SuBAU1, SuBAU2, SuBAU3, SuBAU4].
- [AtLZ] A. Atminas, V. Lozin and V. Zamaraev, Linear Ramsey Numbers, *Proceedings of IWOCA 2018*, Singapore, LNCS 10979, Springer, (2018) 26-38.
- [-] A. Atminas, see also [AleALZ].
- [AttLM] Yamaan Attwa, Albert López Vidal and Patrick Morris, A Note on Multicolour Ramsey Numbers and Random Sphere Graphs, *preprint*, <http://arxiv.org/abs/2602.02155> (2026).
- [AxCJMR] M. Axenovich, W. Cames van Batenburg, O. Janzer, L. Michel and M. Rundström, An Improved Upper Bound for the Multicolour Ramsey Number of Odd Cycles, *preprint*, <http://arxiv.org/abs/2510.17981> (2025).
- [AxFM] M. Axenovich, Z. Füredi and D. Mubayi, On Generalized Ramsey Theory: the Bipartite Case, *Journal of Combinatorial Theory, Series B*, **79** (2000) 66-86.
- [AxGy] M. Axenovich and A. Gyárfás, A Note on Ramsey Numbers for Berge- $G$  Hypergraphs, *Discrete Mathematics*, **342** (2019) 1245-1252.
- [AxGLM] M. Axenovich, A. Gyárfás, Hong Liu and D. Mubayi, Multicolor Ramsey Numbers for Triple Systems, *Discrete Mathematics*, **322** (2014) 69-77.

## Ba - Bi

- [BaRT]\* A. Babak, S.P. Radziszowski and Kung-Kuen Tse, Computation of the Ramsey Number  $R(B_3, K_5)$ , *Bulletin of the Institute of Combinatorics and its Applications*, **41** (2004) 71-76.
- [Back1] J. Backelin, Contributions to a Ramsey Calculus, *manuscript* 2000-2012.
- [Back2] J. Backelin, *personal communication* (2013).
- [Back3]\* J. Backelin, Edge number report 1: state of the art estimates for  $n \leq 43$ , *preprint*, <http://arxiv.org/abs/1410.1843> (2014).

- [Back4] J. Backelin, *personal communication* (2017).
- [BahS] P. Bahls and T.S. Spencer, On the Ramsey Numbers of Trees with Small Diameter, *Graphs and Combinatorics*, **29** (2013) 39-44.
- [-] P. Bahls, see also [AlmBCL].
- [Bai] Bai Lufeng, Multi-color Ramsey Numbers for Trees Versus Complete Graphs (in Chinese), *Mathematics in Practice and Theory*, **43** (2013) 252-254.
- [BaiLi] Bai Lufeng and Li Yusheng, Algebraic Constructions and Applications in Ramsey Theory, *Advances in Mathematics, China*, **35** (2006) 167-170.
- [BaLX] Bai Lufeng, Li Yusheng and Xu Zhiqiang, Algebraic Constructions and Applications in Ramsey Theory, *Journal of Mathematical Study (China)*, **37** (2004) 245-249.
- [-] Bai Lufeng, see also [SonBL].
- [-] A.Q. Baig, see [AliBB].
- [BalDO] Deepak Bal, Louis DeBiasio and Ella Oren-Dahan, On the Multicolor Ramsey Numbers of Balanced Double Stars, *Discrete Mathematics*, **349** (2026) 114748, 11 pages.
- [BaLL+]\* V. Balaji, P. Lamb, A. Lott, D. Patel, A. Rice, S. Singh and C.R. Ward, The Pigeonhole Principle and Multicolor Ramsey Numbers, *Involve: A Journal of Mathematics*, **15**(5) (2022) 857-884.
- [BalB+] P. Balister, B. Bollobás, M. Campos, S. Griffiths, E. Hurley, R. Morris, J. Sahasrabudhe and M. Tiba, Upper Bounds for Multicolour Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2410.17197> (2025).
- [BalLS] P.N. Balister, J. Lehel and R.H. Schelp, Ramsey Unsaturated and Saturated Graphs, *Journal of Graph Theory*, **51** (2006) 22-32.
- [BalSS] P.N. Balister, R.H. Schelp and M. Simonovits, A Note on Ramsey Size-Linear Graphs, *Journal of Graph Theory*, **39** (2002) 1-5.
- [BalPS] I. Balla, A. Pokrovskiy and B. Sudakov, Ramsey Goodness of Bounded Degree Trees, *Combinatorics, Probability and Computing*, **27**(3) (2018) 289-309.
- [BalCSW] J. Balogh, F.C. Clemen, J. Skokan and A.Z. Wagner, The Ramsey Number of Fano Plane Versus Tight Path, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.60, **27**(1) (2020), 16 pages.
- [Ban1] A. Baniabedruhman, The Theta-Complete Graph Ramsey Number  $R(\theta_n, K_7)$ ;  $n = 7$ ;  $n \geq 14$ , *Jordan Journal of Mathematics and Statistics*, **14**(3) (2021) 517-526.
- [Ban2] A. Baniabedruhman, The Cycle-Complete Graph Ramsey Numbers  $R(C_n, K_8)$  for  $10 \leq n \leq 15$ , *Jordan Journal of Mathematics and Statistics*, **16**(4) (2023) 703-718.
- [BanAA] A. Baniabedruhman, A. Alrifai and A.A. Al-Rhayyel, The Cycle-Complete Graph Ramsey Numbers  $R(C_7, K_9)$  and  $R(C_8, K_9)$ , *Italian Journal of Pure and Applied Mathematics*, **48** (2022) 325-335.
- [BanJBJ] A. Baniabedruhman, M.M.M. Jaradat, M.S. Bataineh and A.M.M. Jaradat, On the Ramsey Number for Theta Graphs Versus the Complete Graph of order Six, *Journal of Mathematics*, (2024), article ID 2416730, 6 pages.
- [-] A. Baniabedruhman, see also [JaBa, JaBBJ].
- [-] Qiquan Bao, see [ShaXB, ShaXBP].
- [BarRSW] B. Barak, A. Rao, R. Shaltiel and A. Wigderson, 2-Source Dispersers for  $n^{o(1)}$  Entropy, and Ramsey Graphs Beating the Frankl-Wilson Construction, *Annals of Mathematics* (2), **176** (2012) 1483-1544.
- [BaLiu] *Hungarian Problem Book IV*, translated and edited by Robert Barrington Leigh and Andy Liu, The Mathematical Association of America, 2011.
- [Bas] E.T. Baskoro, The Ramsey Number of Paths and Small Wheels, *Majalah Ilmiah Himpunan Matematika Indonesia*, MIHMI, **8** (2002) 13-16.
- [BaHA] E.T. Baskoro, Hasmawati and H. Assiyatun, The Ramsey Numbers for Disjoint Unions of Trees, *Discrete Mathematics*, **306** (2006) 3297-3301.

- [BaSu] E.T. Baskoro and Surahmat, The Ramsey Number of Paths with respect to Wheels, *Discrete Mathematics*, **294** (2005) 275-277.
- [BaSNM] E.T. Baskoro, Surahmat, S.M. Nababan and M. Miller, On Ramsey Graph Numbers for Trees Versus Wheels of Five or Six Vertices, *Graphs and Combinatorics*, **18** (2002) 717-721.
- [-] E.T. Baskoro, see also [AliBB, AliBas, AliBT1, AliBT2, HafBa, HaABS, HaBA1, HaBA2, NoBa, SherBSO, SherSBO, SuAUB, SuBa1, SuBa2, SuBAU1, SuBAU2, SuBAU3, SuBAU4, SuBB1, SuBB2, SuBB3, SuBB4, SuBT1, SuBT2, SuBTB, SuBUB].
- [BatJA] M.S.A. Bataineh, M.M.M. Jaradat and L.M.N. Al-Zaleq, The Cycle-Complete Graph Ramsey Number  $r(C_9, K_8)$ , *International Scholarly Research Network - Algebra*, Article ID 926191, (2011), 10 pages.
- [-] M.S.A. Bataineh, see also [BanJBJ, JaBBJ, JaBVR].
- [BeHHRs] R. Belmonte, P. Heggenes, P. van 't Hof, A. Rafiey and R. Saei, Graph Classes and Ramsey Numbers, *Discrete Applied Mathematics*, **173** (2014) 16-27.
- [BeHHS] R. Belmonte, P. Heggenes, P. van 't Hof and R. Saei, Ramsey Numbers for Line Graphs and Perfect Graphs, *Proceedings of COCOON 2012*, Sydney, Australia, LNCS 7434, Springer, (2012) 204-215.
- [BenSk] F.S. Benevides and J. Skokan, The 3-Colored Ramsey Number of Even Cycles, *Journal of Combinatorial Theory, Series B*, **99** (2009) 690-708.
- [-] S. Ben-Shimon, see [AlBK].
- [-] S. Bereg, see [LowKKB].
- [Bev] D. Bevan, *personal communication* (2002).
- [BePi] A. Beveridge and O. Pikhurko, On the Connectivity of Extremal Ramsey Graphs, *Australasian Journal of Combinatorics*, **41** (2008) 57-61.
- [BiaS] A. Bialostocki and J. Schönheim, On Some Turán and Ramsey Numbers for  $C_4$ , in *Graph Theory and Combinatorics* (ed. B. Bollobás), Academic Press, London, (1984) 29-33.
- [Biel1] H. Bielak, Ramsey and 2-local Ramsey Numbers for Disjoint Unions of Cycles, *Discrete Mathematics*, **307** (2007) 319-330.
- [Biel2] H. Bielak, Ramsey Numbers for a Disjoint Union of Some Graphs, *Applied Mathematics Letters*, **22** (2009) 475-477.
- [Biel3] H. Bielak, Multicolor Ramsey Numbers for Some Paths and Cycles, *Discussiones Mathematicae Graph Theory*, **29** (2009) 209-218.
- [Biel4] H. Bielak, Ramsey Numbers for a Disjoint Union of Good Graphs, *Discrete Mathematics*, **310** (2010) 1501-1505.
- [BieDa] H. Bielak and K. Dabrowska, The Ramsey Numbers for Some Subgraphs of Generalized Wheels Versus Cycles and Paths, *Annales Universitatis Mariae Curie-Skłodowska Lublin-Polonia, Sectio A*, **LXIX** (2015) 1-7.
- [-] H. Bielak, see also [LiBie, LiZBBH].
- [Bier] J. Bierbrauer, Ramsey Numbers for the Path with Three Edges, *European Journal of Combinatorics*, **7** (1986) 205-206.
- [BierB] J. Bierbrauer and A. Brandis, On Generalized Ramsey Numbers for Trees, *Combinatorica*, **5** (1985) 95-107.
- [BiFJ] C. Biró, Z. Füredi and S. Jahanbekam, Large Chromatic Number and Ramsey Graphs, *Graphs and Combinatorics*, **29** (2013) 1183-1191.
- [Bish] A. Bishnoi, Finite geometry paves the way for breakthroughs in Ramsey theory, *Nieuw Archief voor Wiskunde*, NAW 5/24 (2023) 201-203.

**Bl - Br**

- [BILR]\* K. Black, D. Leven and S.P. Radziszowski, New Bounds on Some Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **78** (2011) 213-222.

- [BLi] T. Bloom and A. Liebenau, Ramsey Equivalence of  $K_n$  and  $K_n + K_{n-1}$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.4, **25**(3) (2018), 17 pages.
- [Boh] T. Bohman, The Triangle-Free Process, *Advances in Mathematics*, **221** (2009) 1653-1677.
- [BohK1] T. Bohman and P. Keevash, The Early Evolution of the  $H$ -Free Process, *Inventiones Mathematicae*, **181** (2010) 291-336.
- [BohK2] T. Bohman and P. Keevash, Dynamic Concentration of the Triangle-Free Process, *Seventh European Conference on Combinatorics, Graph Theory and Applications*, 489-495, CRM Series, 16, Pisa, 2013.
- [BohK3] T. Bohman and P. Keevash, Dynamic Concentration of the Triangle-Free Process, *preprint*, <http://arxiv.org/abs/1302.5963> (2013), 52 pages. Revised version (2019), 75 pages. *Random Structures and Algorithms*, **58**(2) (2021) 221-293.
- [BohP] Tom Bohman and Fei Peng, A Construction for Boolean Cube Ramsey Numbers, *Order*, **40** (2023) 327-333.
- [BohZ] Tom Bohman and Emily Zhu, On Multicolor Ramsey Numbers of Triple System Paths of Length 3, *SIAM Journal on Discrete Mathematics*, **37**(3) (2023) 1419-1435.
- [BolJY+] B. Bollobás, C.J. Jayawardene, Yang Jian Sheng, Huang Yi Ru, C.C. Rousseau and Zhang Ke Min, On a Conjecture Involving Cycle-Complete Graph Ramsey Numbers, *Australasian Journal of Combinatorics*, **22** (2000) 63-71.
- [BolM] B. Bollobás and R. Morris, *Basic Graph Theory*, chapter 4: *Ramsey Theory*, 132-173, Cambridge University Press, 2026.
- [-] B. Bollobás, see also [BalB+, JRB].
- [BoH] R. Bolze and H. Harborth, The Ramsey Number  $r(K_4-x, K_5)$ , in *The Theory and Applications of Graphs*, (Kalamazoo, MI, 1980), John Wiley & Sons, New York, (1981) 109-116.
- [BoEr] J.A. Bondy and P. Erdős, Ramsey Numbers for Cycles in Graphs, *Journal of Combinatorial Theory, Series B*, **14** (1973) 46-54.
- [BotMds] F. Botler, L. Moreira and J.P. de Souza, Ramsey Goodness of Paths Versus Unbalanced Graphs, *Discrete Mathematics*, **349** (2026) 114831, 6 pages.
- [BoLo] Simona Boyadzhyska and Allan Lo, Ramsey Goodness of  $k$ -Uniform Paths, or the Lack Thereof, *European Journal of Combinatorics*, **129** (2025) 104021, 24 pages.
- [-] Simona Boyadzhyska, see also [AllBP].
- [Boza1] L. Boza, Nuevas Cotas Superiores de Algunos Números de Ramsey del Tipo  $r(K_m, K_n - e)$ , in proceedings of the *VII Jornada de Matemática Discreta y Algoritmica*, JMDA 2010, Castro Urdiales, Spain, July 2010.
- [Boza2] L. Boza, The Ramsey Number  $r(K_5 - P_3, K_5)$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P90, **18**(1) (2011), 10 pages.
- [Boza3]\* L. Boza, Upper Bounds for Some Ramsey Numbers of  $K_n - e$  Versus  $K_m$ , *manuscript* (2012).
- [Boza4]\* L. Boza, Números de Ramsey de Algunos Grafos de 4 Vértices y Todos los Grafos de 7 Vértices, in proceedings of the *VIII Jornada de Matemática Discreta y Algoritmica*, JMDA 2012, Almeria, Spain, July 2012.
- [Boza5]\* L. Boza, University of Seville, *personal communication* (2013).
- [Boza6]\* L. Boza, Sobre el Número de Ramsey  $R(K_4, K_6 - e)$ , *VIII Encuentro Andaluz de Matemática Discreta*, Sevilla, Spain, October 2013.
- [Boza7]\* L. Boza, Sobre los Números de Ramsey  $R(K_5 - e, K_5)$  y  $R(K_6 - e, K_4)$ , *IX Jornada de Matemática Discreta y Algoritmica*, JMDA 2014, Tarragona, Spain, July 2014.
- [Boza8]\* L. Boza, Exact Values and Bounds for Ramsey Numbers of  $C_4$  Versus a Star Graph, *preprint*, <http://arxiv.org/abs/2409.12770> (2024).
- [Boza9] L. Boza, New Upper Bounds for the Classical Ramsey Numbers  $R(4, 4, 4)$ ,  $R(3, 4, 5)$  and  $R(3, 3, 6)$ , *preprint*, <http://arxiv.org/abs/2603.10851> (2026).

- [BoCGR] L. Boza, M. Cera, P. Garcia-Vázquez and M.P. Revuelta, On the Ramsey Numbers for Stars Versus Complete Graphs, *European Journal of Combinatorics*, **31** (2010) 1680-1688.
- [BoDD]\* L. Boza, J. Dybizbański and T. Dzido, Three Color Ramsey Numbers for Graphs With at Most 4 Vertices, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P47, **19**(4) (2012), 16 pages.
- [BoPo]\* L. Boza and J.R. Portillo, Sobre el Número de Ramsey  $R(K_4 - e, K_7)$ , in proceedings of the *VIII Jornada de Matemática Discreta y Algoritmica*, JMDA 2012, Almeria, Spain, July 2012.
- [BoRa] L. Boza and S. Radziszowski, Some Upper Bounds on Ramsey Numbers Involving  $C_4$ , *Discussiones Mathematicae Graph Theory*, **45** (2025) 845-856.
- [BraFS1] D. Bradač, J. Fox and B. Sudakov, The Growth Rate of Multicolor Ramsey Numbers of 3-Graphs, *Research in the Mathematical Sciences*, (2024), 11:52, 16 pages.
- [BraFS2] D. Bradač, J. Fox and B. Sudakov, Ramsey Numbers of Hypergraphs with a Given Size, *Mathematical Proceedings of the Cambridge Philosophical Society*, **178** (2025) 31-44.
- [BraGS] D. Bradač, L. Gishboliner and B. Sudakov, On Ramsey Size-Linear Graphs and Related Questions, *SIAM Journal on Discrete Mathematics*, **38**(1) (2024) 10.1137/22M1481713.
- [BraHS] D. Bradač, Z. Hunter and B. Sudakov, Lower Bounds for Ramsey Numbers of Bounded Degree Hypergraphs, *preprint*, <http://arxiv.org/abs/2502.20863> (2025).
- [-] A. Brandis, see [BierB].
- [Bra1] S. Brandt, Subtrees and Subforests in Graphs, *Journal of Combinatorial Theory, Series B*, **61** (1994) 63-70.
- [Bra2] S. Brandt, Sufficient Conditions for Graphs to Contain All Subgraphs of a Given Type, *Ph.D. thesis*, Freie Universität Berlin, 1994.
- [Bra3] S. Brandt, Expanding Graphs and Ramsey Numbers, *preprint No. A 96-24*, <ftp://ftp.math.fu-berlin.de/pub/math/publ/pre/1996> (1996).
- [BrBH1]\*\* S. Brandt, G. Brinkmann and T. Harmuth, All Ramsey Numbers  $r(K_3, G)$  for Connected Graphs of Order 9, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R7, **5** (1998), 20 pages.
- [BrBH2]\*\* S. Brandt, G. Brinkmann and T. Harmuth, The Generation of Maximal Triangle-Free Graphs, *Graphs and Combinatorics*, **16** (2000) 149-157.
- [Bren1] M. Brennan, Ramsey Numbers of Trees Versus Odd Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.2, **23**(3) (2016), 12 pages.
- [Bren2] M. Brennan, Ramsey Numbers of Trees and Unicyclic Graphs Versus Fans, *Discrete Mathematics*, **340** (2017) 969-983.
- [-] C. Bright, see [DuLBG].
- [-] G. Brightwell, see [AllBS].
- [Brin]\*\* G. Brinkmann, All Ramsey Numbers  $r(K_3, G)$  for Connected Graphs of Order 7 and 8, *Combinatorics, Probability and Computing*, **7** (1998) 129-140.
- [BrCGM]\* G. Brinkmann, K. Coolsaet, J. Goedgebeur and H. Mélot, House of Graphs: A database of interesting graphs, *Discrete Applied Mathematics*, **161** (2013) 311-314.
- [BrGS]\*\* G. Brinkmann, J. Goedgebeur and J.C. Schläge-Puchta, Ramsey Numbers  $R(K_3, G)$  for Graphs of Order 10, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P36, **19**(4) (2012), 23 pages.
- [-] G. Brinkmann, see also [BrBH1, BrBH2].
- [-] T. Britz, see [ChngBTW].
- [-] H.J. Broersma, see [LiZBBH, LiZB, SaBr1, SaBr2, SaBr3, SaBr4, SuBB1, SuBB2, SuBB3, SuBB4, SuBTB, SuBUB, ZhaBC1, ZhaBC2, ZhaBC3, ZhaBC4, ZhaBC5, ZhaBC6].
- [BroNN] S. Brooks, T. Nguyen and E. Nystrom, The Ramsey Number and Saturation of the Tristar, *manuscript* (2016).

- [Brou] A.E. Brouwer, *Parameters of Strongly Regular Graphs*, status of feasible parameters up to 1300 vertices and of several SRG family parameters, <https://www.win.tue.nl/~aeb/graphs/srg/srgtab.html>.
- [-] Chad E. Brown, see [GauB].
- [BrBH] M. Bruce, M. Budden and J. Hiller, Lexicographic Products of  $r$ -Uniform Hypergraphs and Some Applications to Hypergraph Ramsey Theory, *Australasian Journal of Combinatorics*, **70** (2018) 390-401.

## Bu

- [BucSu] M. Bucić and B. Sudakov, Tight Ramsey Bounds for Multiple Copies of a Graph, *Advances in Combinatorics*, <https://doi.org/10.19086/aic.2023.1> (2023):1, 22 pages.
- [BudCli] M. Budden and J. Clifton, Hypergraph Ramsey Numbers Involving Trees, Stars, and Complete Hypergraphs, *Integers*, <http://math.colgate.edu/~integers>, **22** (2022) #A92, 14 pages.
- [BudHLS] M. Budden, J. Hiller, J. Lambert and C. Sanford, The Lifting of Graphs to 3-Uniform Hypergraphs and Some Applications to Hypergraph Ramsey Theory, *Involve: A Journal of Mathematics*, **10** (2017) 65-76.
- [BudHMP] M. Budden, J. Hiller, T. Meek and A. Penland, Algebraic Properties of a Hypergraph Lifting Map, *Integers*, <http://math.colgate.edu/~integers>, **21** (2021) #A77, 12 pages.
- [BudHP] M. Budden, J. Hiller and A. Penland, Constructive Methods in Gallai-Ramsey Theory for Hypergraphs, *Integers*, <http://math.colgate.edu/~integers>, **20A** (2020) #A4, 14 pages.
- [BudHR1] M. Budden, J. Hiller and A. Rapp, Generalized Ramsey Theorems for  $r$ -Uniform Hypergraphs, *Australasian Journal of Combinatorics*, **63** (2015) 142-152.
- [BudHR2] M. Budden, J. Hiller and A. Rapp, Hypergraph Ramsey Numbers Involving Paths, *Acta Universitatis Apulensis*, **48** (2016) 75-87.
- [BudPe] M. Budden and A. Penland, Trees and  $n$ -Good Hypergraphs, *Australasian Journal of Combinatorics*, **72**(2) (2018) 329-349. Corrigendum in **75**(1) (2019) 171-173.
- [BudPra] M. Budden and R. Prange, Ramsey Theory for a Generalized Fan Versus Triangles, *Utilitas Mathematica*, **123** (2025) 165-178.
- [BudPri] M. Budden and H. Privette, Multicolor Ramsey Theory for a Fan Versus Complete Graphs, *Discrete Mathematics Letters*, **13** (2024) 143-149.
- [-] M. Budden, see also [BrBH].
- [BurR]\* J.P. Burling and S.W. Reyner, Some Lower Bounds of the Ramsey Numbers  $n(k, k)$ , *Journal of Combinatorial Theory, Series B*, **13** (1972) 168-169.
- [Bu0] S.A. Burr, Generalized Ramsey Theory for Graphs - a Survey, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 52-75.
- [Bu1] S.A. Burr, A Survey of Noncomplete Ramsey Theory for Graphs, *Annals of the New York Academy of Sciences*, **328**(1) (1979) 58-75.
- [Bu2] S.A. Burr, Ramsey Numbers Involving Graphs with Long Suspended Paths, *Journal of the London Mathematical Society* (2), **24** (1981) 405-413.
- [Bu3] S.A. Burr, Multicolor Ramsey Numbers Involving Graphs with Long Suspended Path, *Discrete Mathematics*, **40** (1982) 11-20.
- [Bu4] S.A. Burr, Diagonal Ramsey Numbers for Small Graphs, *Journal of Graph Theory*, **7** (1983) 57-69.
- [Bu5] S.A. Burr, Ramsey Numbers Involving Powers of Sparse Graphs, *Ars Combinatoria*, **15** (1983) 163-168.
- [Bu6] S.A. Burr, Determining Generalized Ramsey Numbers is NP-Hard, *Ars Combinatoria*, **17** (1984) 21-25.
- [Bu7] S.A. Burr, What Can We Hope to Accomplish in Generalized Ramsey Theory?, *Discrete Mathematics*, **67** (1987) 215-225.

- [Bu8] S.A. Burr, On the Ramsey Numbers  $r(G, nH)$  and  $r(nG, nH)$  When  $n$  Is Large, *Discrete Mathematics*, **65** (1987) 215-229.
- [Bu9] S.A. Burr, On Ramsey Numbers for Large Disjoint Unions of Graphs, *Discrete Mathematics*, **70** (1988) 277-293.
- [Bu10] S.A. Burr, On the Computational Complexity of Ramsey-type Problems, Mathematics of Ramsey Theory, *Algorithms and Combinatorics*, **5**, Springer, Berlin, 1990, 46-52.
- [BuE1] S.A. Burr and P. Erdős, On the Magnitude of Generalized Ramsey Numbers for Graphs, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds., Keszthely 1973) Vol. 1, 215-240, Colloq. Math. Soc. János Bolyai, Vol. 10, North-Holland, Amsterdam, 1975.
- [BuE2] S.A. Burr and P. Erdős, Extremal Ramsey Theory for Graphs, *Utilitas Mathematica*, **9** (1976) 247-258.
- [BuE3] S.A. Burr and P. Erdős, Generalizations of a Ramsey-Theoretic Result of Chvátal, *Journal of Graph Theory*, **7** (1983) 39-51.
- [BEFRS1] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, An Extremal Problem in Generalized Ramsey Theory, *Ars Combinatoria*, **10** (1980) 193-203.
- [BEFRS2] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for the Pair Sparse Graph-Path or Cycle, *Transactions of the American Mathematical Society*, **269** (1982) 501-512.
- [BEFRS3] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Ramsey Number for the Pair Complete Bipartite Graph-Graph of Limited Degree, in *Graph Theory with Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley & Sons, New York, (1985) 163-174.
- [BEFRS4] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Some Complete Bipartite Graph-Tree Ramsey Numbers, *Annals of Discrete Mathematics*, **41** (1989) 79-89.
- [BEFRSGJ] S.A. Burr, P. Erdős, R.J. Faudree, C.C. Rousseau, R.H. Schelp, R.J. Gould and M.S. Jacobson, Goodness of Trees for Generalized Books, *Graphs and Combinatorics*, **3** (1987) 1-6.
- [BEFS] S.A. Burr, P. Erdős, R.J. Faudree and R.H. Schelp, On the Difference between Consecutive Ramsey Numbers, *Utilitas Mathematica*, **35** (1989) 115-118.
- [BES] S.A. Burr, P. Erdős and J.H. Spencer, Ramsey Theorems for Multiple Copies of Graphs, *Transactions of the American Mathematical Society*, **209** (1975) 87-99.
- [BuF] S.A. Burr and R.J. Faudree, On Graphs  $G$  for Which All Large Trees Are  $G$ -good, *Graphs and Combinatorics*, **9** (1993) 305-313.
- [BFRS] S.A. Burr, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Ramsey Numbers Involving Starlike Multipartite Graphs, *Journal of Graph Theory*, **7** (1983) 395-409.
- [BuG] S.A. Burr and J.W. Grossman, Ramsey Numbers of Graphs with Long Tails, *Discrete Mathematics*, **41** (1982) 223-227.
- [BuRo1] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Stars, *Utilitas Mathematica*, **4** (1973) 217-220.
- [BuRo2] S.A. Burr and J.A. Roberts, On Ramsey Numbers for Linear Forests, *Discrete Mathematics*, **8** (1974) 245-250.
- [BuRo3] S.A. Burr and V. Rosta, On the Ramsey Multiplicities of Graphs - Problems and Recent Results, *Journal of Graph Theory*, **4** (1980) 347-361.
- [Bush] L.E. Bush, The William Lowell Putnam Mathematical Competition (question #2 in Part I asks for the proof of  $R(3,3) \leq 6$ ), *American Mathematical Monthly*, **60** (1953) 539-542.
- [-] S. Butler, see [GrBu].

## Ca - Ch

- [-] J.W. Cain, see [LinCa].
- [CaET]\* N.J. Calkin, P. Erdős and C.A. Tovey, New Ramsey Bounds from Cyclic Graphs of Prime Order, *SIAM Journal on Discrete Mathematics*, **10** (1997) 381-387.
- [CalSR]\* J.A. Calvert, M.J. Schuster and S.P. Radziszowski, Computing the Ramsey Number  $R(K_5 - P_3, K_5)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **82** (2012) 131-140.
- [CamFMPP] S. Cambie, A. Freschi, P. Morawski, K. Petrova and A. Pokrovskiy, Ramsey Number of a Cycle Versus a Graph of a Given Size, *preprint*, <http://arxiv.org/abs/2601.10238> (2026).
- [-] Wouter Cames van Batenburg, see [AxCJMR].
- [CamGMS] M. Campos, S. Griffiths, R. Morris and J. Sahasrabudhe, An Exponential Improvement for Diagonal Ramsey, *preprint*, <http://arxiv.org/abs/2303.09521> (2023), 57 pages. To appear in *Annals of Mathematics*, Princeton University.
- [CamJMPS] M. Campos, M. Jenssen, M. Michelen, F. Pfender and J. Sahasrabudhe, A Polynomial Improvement for the Odd Cycle-Complete Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2511.10641> (2025).
- [CamJMS] M. Campos, M. Jenssen, M. Michelen and J. Sahasrabudhe, A New Lower Bound for the Ramsey Numbers  $R(3, k)$ , *preprint*, <http://arxiv.org/abs/2505.13371> (2025), 52 pages.
- [CamP] M. Campos and C. Pohoata, An Update on Multicolor Ramsey Lower Bounds, *preprint*, <http://arxiv.org/abs/2601.15183> (2026), 9 pages.
- [-] M. Campos, see also [BalB+].
- [Car] D. Cariolaro, On the Ramsey Number  $R(3, 6)$ , *Australasian Journal of Combinatorics*, **37** (2007) 301-304.
- [Caro] Y. Caro, Zero-Sum Problems - A Survey, *Discrete Mathematics*, **152** (1996) 93-113.
- [CaLRZ] Y. Caro, Li Yusheng, C.C. Rousseau and Zhang Yuming, Asymptotic Bounds for Some Bipartite Graph - Complete Graph Ramsey Numbers, *Discrete Mathematics*, **220** (2000) 51-56.
- [CaYZ] Y. Caro, R. Yuster and C. Zarb, Ramsey Numbers for Degree Monotone Paths, *Discrete Mathematics*, **340** (2017) 124-131.
- [-] M. Cera, see [BoCGR].
- [-] P. Casteras, see [AgCP+].
- [ChaMR] J. Chappelon, L.P. Montejano and J.L. Ramirez Alfonsin, On Ramsey Numbers of Complete Graphs with Dropped Stars, *Discrete Applied Mathematics*, **210** (2016) 200-206.
- [-] J. Chappelon, see also [MonCR].
- [ChaGP] G. Chartrand, R.J. Gould and A.D. Polimeni, On Ramsey Numbers of Forests Versus Nearly Complete Graphs, *Journal of Graph Theory*, **4** (1980) 233-239.
- [ChaRSPS] G. Chartrand, C.C. Rousseau, M.J. Stewart, A.D. Polimeni and J. Sheehan, On Star-Book Ramsey Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 203-214.
- [ChaS] G. Chartrand and S. Schuster, On the Existence of Specified Cycles in Complementary Graphs, *Bulletin of the American Mathematical Society*, **77** (1971) 995-998.
- [ChaZ] Gary Chartrand and Ping Zhang, New Directions in Ramsey Theory, *Discrete Mathematics Letters*, **6** (2021) 84-96.
- [-] Gang Chen, see [LiuCh].
- [Chen] Chen Guantao, A Result on  $C_4$ -Star Ramsey Numbers, *Discrete Mathematics*, **163** (1997) 243-246.
- [ChenS] Chen Guantao and R.H. Schelp, Graphs with Linearly Bounded Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **57** (1993) 138-149.
- [ChenYZ] Guantao Chen, Xiaowei Yu and Yi Zhao, Improved Bounds on the Ramsey Number of Fans, *European Journal of Combinatorics*, **96** (2021) 103347, 7 pages.

- [-] Chen Hong, see also [LiaWXCS, XWCS].
- [ChenJ] Chen Jie, The Lower Bound of Some Ramsey Numbers (in Chinese), *Journal of Liaoning Normal University, Natural Science*, **25** (2002) 244-246.
- [-] Kun Chen, see [ZhaCZ].
- [-] Ming Chen, see [PeiCLY].
- [-] Pingge Chen, see [CheLC].
- [-] Weiji Chen, see [LinCh].
- [ChenL] Xun Chen and Qizhong Lin, New Upper Bounds for Ramsey Numbers of Books, *European Journal of Combinatorics*, **115** (2024) 103785, 9 pages.
- [ChenLY] Xun Chen, Qizhong Lin and Chunlin You, Ramsey Numbers of Large Books, *Journal of Graph Theory*, **101** (2022) 124-133.
- [-] Xun Chen, see also [YouLC].
- [ChenCMN] Yaojun Chen, T.C. Edwin Cheng, Zhengke Miao and C.T. Ng, The Ramsey Numbers for Cycles Versus Wheels of Odd Order, *Applied Mathematics Letters*, **22** (2009) 1875-1876.
- [ChenCNZ] Yaojun Chen, T.C. Edwin Cheng, C.T. Ng and Yunqing Zhang, A Theorem on Cycle-Wheel Ramsey Number, *Discrete Mathematics*, **312** (2012) 1059-1061.
- [ChenCX] Yaojun Chen, T.C. Edwin Cheng and Ran Xu, The Ramsey Number for a Cycle of Length Six Versus a Clique of Order Eight, *Discrete Applied Mathematics*, **157** (2009) 8-12.
- [ChenCZ1] Yaojun Chen, T.C. Edwin Cheng and Yunqing Zhang, The Ramsey Numbers  $R(C_m, K_7)$  and  $R(C_7, K_8)$ , *European Journal of Combinatorics*, **29** (2008) 1337-1352.
- [ChenZZ1] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Paths Versus Wheels, *Discrete Mathematics*, **290** (2005) 85-87.
- [ChenZZ2] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Stars Versus Wheels, *European Journal of Combinatorics*, **25** (2004) 1067-1075.
- [ChenZZ3] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers  $R(T_n, W_6)$  for  $\Delta(T_n) \geq n-3$ , *Applied Mathematics Letters*, **17** (2004) 281-285.
- [ChenZZ4] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers of Trees Versus  $W_6$  or  $W_7$ , *European Journal of Combinatorics*, **27** (2006) 558-564.
- [ChenZZ5] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers  $R(T_n, W_6)$  for Small  $n$ , *Utilitas Mathematica*, **67** (2005) 269-284.
- [ChenZZ6] Chen Yaojun, Zhang Yunqing and Zhang Ke Min, The Ramsey Numbers  $R(T_n, W_6)$  for  $T_n$  without Certain Deletable Sets, *Journal of Systems Science and Complexity*, **18** (2005) 95-101.
- [ChenZZ7] Chen Yaojun, Zhang Xuemei and Zhang Yanbo, Star-Quadrilateral Ramsey Number and Beyond, *Advances in Mathematics, China*, **54**(2) (2025) 292-314.
- [-] Chen Yaojun, see also [CheCZN, HuCh, HuZC1, HuZC2, HuZC3, WaCh1, WaCh2, ZhCh, ZhZC, ZhaBC1, ZhaBC2, ZhaBC3, ZhaBC4, ZhaBC5, ZhaBC6, ZhaCC1, ZhaCC2, ZhaCC3, ZhaCC4, ZhaCC5, ZhaCC6, ZhaCh1, ZhaCh3, ZhaCh4, ZhaCh5, ZhaCC7, ZhaCZ1, ZhaCZ2, ZhaZC, ZhaChZ1, ZhaChZ2].
- [-] Chen Zhi, see [XuXC].
- [Cheng] Cheng Ying, On Graphs Which Do Not Contain Certain Trees, *Ars Combinatoria*, **19** (1985) 119-151.
- [CheCZN] T.C. Edwin Cheng, Yaojun Chen, Yunqing Zhang and C.T. Ng, The Ramsey Numbers for a Cycle of Length Six or Seven Versus a Clique of Order Seven, *Discrete Mathematics*, **307** (2007) 1047-1053.
- [-] T.C. Edwin Cheng, see also [ChenCMN, ChenCNZ, ChenCX, ChenCZ1, ZhaCC1, ZhaCC2, ZhaCC3, ZhaCC4, ZhaCC5, ZhaCC6, ZhaCC7].

- [CheLC] Zhiyu Cheng, Zhidan Luo and Pingge Chen, All Ramsey Critical Graphs for a Large Tree Versus  $tK_m$ , *preprint*, <http://arxiv.org/abs/2506.18235> (2025).
- [ChngBTW] Zhi Yee Chng, Thomas Britz, Ta Sheng Tan and Kok Bin Wong, The Ramsey Numbers for Trees of Large Maximum Degree Versus the Wheel Graph  $W_8$ , *Bulletin of the Malaysian Mathematical Sciences Society*, **47** (2024) 134, 44 pages.
- [ChngTW] Zhi Yee Chng, Ta Sheng Tan and Kok Bin Wong, On the Ramsey Numbers for the Tree Graphs Versus Certain Generalised Wheel Graphs, *Discrete Mathematics*, **344** (2021) 112440, 12 pages.
- [Chu1] F.R.K. Chung, On the Ramsey Numbers  $N(3, 3, \dots, 3; 2)$ , *Discrete Mathematics*, **5** (1973) 317-321.
- [Chu2] F.R.K. Chung, On Triangular and Cyclic Ramsey Numbers with  $k$  Colors, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 236-241.
- [Chu3] F.R.K. Chung, A Note on Constructive Methods for Ramsey Numbers, *Journal of Graph Theory*, **5** (1981) 109-113.
- [Chu4] F.R.K. Chung, Open problems of Paul Erdős in Graph Theory, *Journal of Graph Theory*, **25** (1997) 3-36.
- [Chu5] Fan Chung, Numbers in Ramsey Theory, *Journal of Open Mathematical Problems*, **1**(1) (2025) 1-15.
- [ChCD] F.R.K. Chung, R. Cleve and P. Dagum, A Note on Constructive Lower Bounds for the Ramsey Numbers  $R(3, t)$ , *Journal of Combinatorial Theory, Series B*, **57** (1993) 150-155.
- [ChGra1] F.R.K. Chung and R.L. Graham, On Multicolor Ramsey Numbers for Complete Bipartite Graphs, *Journal of Combinatorial Theory, Series B*, **18** (1975) 164-169.
- [ChGra2] F.R.K. Chung and R.L. Graham, *Erdős on Graphs, His Legacy of Unsolved Problems*, A K Peters, Wellesley, Massachusetts (1998).
- [ChGri] F.R.K. Chung and C.M. Grinstead, A Survey of Bounds for Classical Ramsey Numbers, *Journal of Graph Theory*, **7** (1983) 25-37.
- [ChuLin] Fan Chung and Qizhong Lin, Fan-Complete Ramsey Numbers, *Advances in Applied Mathematics*, **171** (2025) 102939, 19 pages.
- [-] F.R.K. Chung, see also [UCSD].
- [Chv1] V. Chvátal, Tree-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 93.
- [Chv2] V. Chvátal, *The Discrete Mathematical Charms of Paul Erdős: A Simple Introduction*, Cambridge University Press, first edition, 2021.
- [ChH1] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, II. Small Diagonal Numbers, *Proceedings of the American Mathematical Society*, **32** (1972) 389-394.
- [ChH2] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, III. Small Off-Diagonal Numbers, *Pacific Journal of Mathematics*, **41** (1972) 335-345.
- [ChH3] V. Chvátal and F. Harary, Generalized Ramsey Theory for Graphs, I. Diagonal Numbers, *Periodica Mathematica Hungarica*, **3** (1973) 115-124.
- [ChRST] V. Chvátal, V. Rödl, E. Szemerédi and W.T. Trotter Jr., The Ramsey Number of a Graph with Bounded Maximum Degree, *Journal of Combinatorial Theory, Series B*, **34** (1983) 239-243.
- [ChvS] V. Chvátal and A. Schwenk, On the Ramsey Number of the Five-Spoked Wheel, in *Graphs and Combinatorics* (R. Bari and F. Harary eds.), Springer LNM **406**, Berlin, (1974) 247-261.

### CI - Cs

- [Clan] M. Clancy, Some Small Ramsey Numbers, *Journal of Graph Theory*, **1** (1977) 89-91.
- [Clap] C. Clapham, The Ramsey Number  $r(C_4, C_4, C_4)$ , *Periodica Mathematica Hungarica*, **18** (1987) 317-318.
- [CIEHMS] C. Clapham, G. Exoo, H. Harborth, I. Mengersen and J. Sheehan, The Ramsey Number of  $K_5 - e$ , *Journal of Graph Theory*, **13** (1989) 7-15.

- [Clark] L. Clark, On Cycle-Star Graph Ramsey Numbers, *Congressus Numerantium*, **50** (1985) 187-192.
- [-] L. Clark, see also [RanMCG].
- [-] F.C. Clemen, see [BalCSW].
- [CleDa] R. Cleve and P. Dagum, A Constructive  $\Omega(t^{1.26})$  Lower Bound for the Ramsey Number  $R(3, t)$ , *International Computer Science Institute*, TR-89-009, Berkeley, CA, 1989.
- [-] R. Cleve, see also [ChCD].
- [-] J. Clifton, see [BudCli].
- [Coc] E.J. Cockayne, Some Tree-Star Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **17** (1974) 183-187.
- [CocL1] E.J. Cockayne and P.J. Lorimer, The Ramsey Number for Stripes, *Journal of the Australian Mathematical Society, Series A*, **19** (1975) 252-256.
- [CocL2] E.J. Cockayne and P.J. Lorimer, On Ramsey Graph Numbers for Stars and Stripes, *Canadian Mathematical Bulletin*, **18** (1975) 31-34.
- [CoPR] B. Codenotti, P. Pudlák and G. Resta, Some Structural Properties of Low-Rank Matrices Related to Computational Complexity, *Theoretical Computer Science*, **235** (2000) 89-107.
- [CodFIM]\* M. Codish, M. Frank, A. Itzhakov and A. Miller, Computing the Ramsey Number  $R(4, 3, 3)$  Using Abstraction and Symmetry Breaking, *Constraints*, **21** (2016) 375-393.
- [-] K. Coffey, see [AlmBCL].
- [Coh] G. Cohen, Two-Source Dispersers for Polylogarithmic Entropy and Improved Ramsey Graphs, in *Proceedings of the 48-th Annual ACM Symposium on Theory of Computing, STOC'16*, Cambridge MA, 278-284. Extended version on arXiv, <http://arxiv.org/abs/1506.04428> (2015). *SIAM Journal on Computing*, **50**(3) (2021) 10.1137/16M1096219.
- [ColGJ] C. Collier-Cartaino, N. Graber and Tao Jiang, Linear Turán Numbers of Linear Cycles and Cycle-Complete Ramsey Numbers, *Combinatorics, Probability and Computing*, **27**(3) (2018) 358-386.
- [CoPC] Special issue on Ramsey theory of *Combinatorics, Probability and Computing*, **12** (2003), Numbers 5 and 6.
- [Con1] D. Conlon, A New Upper Bound for Diagonal Ramsey Numbers, *Annals of Mathematics*, **170** (2009) 941-960.
- [Con2] D. Conlon, Hypergraph Packing and Sparse Bipartite Ramsey Numbers, *Combinatorics, Probability and Computing*, **18** (2009) 913-923.
- [Con3] D. Conlon, The Ramsey Number of Dense Graphs, *Bulletin of the London Mathematical Society*, **45** (2013) 483-496.
- [Con4] D. Conlon, The Ramsey Number of Books, *Advances in Combinatorics*, **3** (2019), 12 pages, <https://doi.org/10.19086/aic.10808>.
- [ConFer] D. Conlon and A. Ferber, Lower Bounds for Multicolor Ramsey Numbers, *Advances in Mathematics*, **378** (2021) 107528, 5 pages.
- [ConFG1+] D. Conlon, J. Fox, B. Gunby, Xiaoyu He, D. Mubayi, A. Suk and J. Verstraëte, On Off-Diagonal Hypergraph Ramsey Numbers, *International Mathematics Research Notices*, rnaf122 (2025), Issue 11, 22 pages.
- [ConFG2+] David Conlon, Jacob Fox, Benjamin Gunby, Xiaoyu He, Dhruv Mubayi, Andrew Suk, Jacques Verstraëte and Hung-Hsun Hans Yu, When Are Off-Diagonal Hypergraph Ramsey Numbers Polynomial?, *Proceedings of the American Mathematical Society*, **153** (2025) 4605-4617.
- [ConFH+] D. Conlon, J. Fox, Xiaoyu He, D. Mubayi, A. Suk and J. Verstraëte, Hypergraph Ramsey Numbers of Cliques Versus Stars, *Random Structures and Algorithms*, **63** (2023) 610-623.
- [ConFLS] D. Conlon, J. Fox, C. Lee and B. Sudakov, Ramsey Numbers of Cubes Versus Cliques, *Combinatorica*, **36** (2016) 37-70.

- [ConFR] D. Conlon, J. Fox and V. Rödl, Hedgehogs Are Not Color Blind, *Journal of Combinatorics*, **8** (2017) 475-485.
- [ConFS1] D. Conlon, J. Fox and B. Sudakov, Ramsey Numbers of Sparse Hypergraphs, *Random Structures and Algorithms*, **35** (2009) 1-14.
- [ConFS2] D. Conlon, J. Fox and B. Sudakov, Hypergraph Ramsey Numbers, *Journal of the American Mathematical Society*, **23** (2010) 247-266.
- [ConFS3] D. Conlon, J. Fox and B. Sudakov, Large Almost Monochromatic Subsets in Hypergraphs, *Israel Journal of Mathematics*, **181** (2011) 423-432.
- [ConFS4] D. Conlon, J. Fox and B. Sudakov, On Two Problems in Graph Ramsey Theory, *Combinatorica*, **32** (2012) 513-535.
- [ConFS5] D. Conlon, J. Fox and B. Sudakov, Erdős-Hajnal-type Theorems in Hypergraphs, *Journal of Combinatorial Theory, Series B*, **102** (2012) 1142-1154.
- [ConFS6] D. Conlon, J. Fox and B. Sudakov, An Improved Bound for the Stepping-Up Lemma, *Discrete Applied Mathematics*, **161** (2013) 1191-1196.
- [ConFS7] D. Conlon, J. Fox and B. Sudakov, Recent Developments in Graph Ramsey Theory, *Surveys in Combinatorics*, London Mathematical Society Lecture Note Series, (2015) 49-118.
- [ConFS8] D. Conlon, J. Fox and B. Sudakov, Short Proofs of Some Extremal Results II, *Journal of Combinatorial Theory, Series B*, **121** (2016) 173-196.
- [ConFW1] D. Conlon, J. Fox and Y. Wigderson, Ramsey Numbers of Books and Quasirandomness, *Combinatorica*, **42**(3) (2022) 309-363.
- [ConFW2] D. Conlon, J. Fox and Y. Wigderson, Off-Diagonal Book Ramsey Numbers, *Combinatorics, Probability and Computing*, **32** (2023) 516-545.
- [ConMMV] D. Conlon, S. Mattheus, D. Mubayi and J. Verstraëte, Ramsey Numbers and the Zarankiewicz Problem, *Bulletin of the London Mathematical Society*, **56** (2024) 2014-2023.
- [ConT] D. Conlon and M. Tyomkyn, Ramsey Numbers of Trails and Circuits, *Journal of Graph Theory*, **102** (2023) 194-196.
- [CooFKO1] O. Cooley, N. Fountoulakis, D. Kühn and D. Osthus, 3-Uniform Hypergraphs of Bounded Degree Have Linear Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **98** (2008) 484-505.
- [CooFKO2] O. Cooley, N. Fountoulakis, D. Kühn and D. Osthus, Embeddings and Ramsey Numbers of Sparse  $k$ -uniform Hypergraphs, *Combinatorica*, **29** (2009) 263-297.
- [-] O. Cooley, see also [KüCFO].
- [-] K. Coolsaet, see [BrCGM].
- [CsKo] R. Csákány and J. Komlós, The Smallest Ramsey Numbers, *Discrete Mathematics*, **199** (1999) 193-199.

## D

- [-] K. Dabrowska, see [BieDa].
- [-] P. Dagum, see [ChCD, CleDa].
- [DavJR] E. Davies, M. Jenssen and B. Roberts, Multicolour Ramsey Numbers of Paths and Even Cycles, *European Journal of Combinatorics*, **63** (2017) 124-133.
- [DawMc]\* M.L. Dawsey and D. McCarthy, Generalized Paley Graphs and Their Complete Subgraphs of Orders Three and Four, *Research in the Mathematical Sciences*, **8** 18, Springer, (2021).
- [DayJ] A.N. Day and J.R. Johnson, Multicolour Ramsey Numbers of Odd Cycles, *Journal of Combinatorial Theory, Series B*, **124** (2017) 56-63.
- [DeBGS] L. DeBiasio, A. Gyárfás and G.N. Sárközy, Ramsey Numbers of Path-Matchings, Covering Designs and 1-Cores, *Journal of Combinatorial Theory, Series B*, **146** (2021) 124-140.

- [–] L. DeBiasio, see also [BalDO].
- [Den1] T. Denley, The Independence Number of Graphs with Large Odd Girth, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R9, **1** (1994), 12 pages.
- [Den2] T. Denley, The Ramsey Numbers for Disjoint Unions of Cycles, *Discrete Mathematics*, **149** (1996) 31-44.
- [DeMST] Z. DeStefano, H. Mahon, F. Simutis and M. Tait, Multicolor Ramsey Numbers for Berge Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.26, **29**(4) (2022), 10 pages.
- [Dong] Dong Lin, A Note on a Lower Bound for  $r(K_{m,n})$ , *Journal of Tongji University (Natural Science)*, **38** (2010) 776,778.
- [DoLi] Lin Dong and Yusheng Li, A Construction for Ramsey Numbers for  $K_{m,n}$ , *European Journal of Combinatorics*, **31** (2010) 1667-1670.
- [DoLL1] Lin Dong, Yusheng Li and Qizhong Lin, Ramsey Numbers Involving Graphs with Large Degrees, *Applied Mathematics Letters*, **22** (2009) 1577-1580.
- [DoLL2] Dong Lin, Li Yusheng and Lin Qizhong, Ramsey Numbers of Cycles vs. Large Complete Graph, *Advances in Mathematics, China*, **39** (2010) 700-702.
- [–] Dong Lin, see also [HeLD, LinLD].
- [DouLLP] Chunyang Dou, Tianyu Li, Qizhong Lin and Xing Peng, The Ramsey Number of the 4-Cycle Versus a Book Graph, *preprint*, <http://arxiv.org/abs/2506.10477> (2025).
- [–] Bangwei Du, see [LiuDu].
- [DuHu] Duan Chanlun and Huang Wenke, Lower Bound of Ramsey Number  $r(3, 10)$  (in Chinese), *Acta Scientiarum Naturalium Universitatis Nei Mongol*, **31** (2000) 468-470.
- [DuRu] A. Dudek and A. Ruciński, Monochromatic Loose Paths in Multicolored  $k$ -Uniform Cliques, *Discrete Mathematics and Theoretical Computer Science, Graph Theory*, **21**(4) (2019) #7, 15 pages.
- [DuLR] D. Duffus, H. Lefmann and V. Rödl, Shift Graphs and Lower Bounds on Ramsey Numbers  $r_k(l; r)$ , *Discrete Mathematics*, **137** (1995) 177-187.
- [DuLBG]\*\* Conor Duggan, Zhengyu Li, Curtis Bright and Vijay Ganesh, A SAT + Computer Algebra System Verification of the Ramsey Problem  $R(3, 8)$ , *Proceedings of the AAAI Conference on Artificial Intelligence 2024, Student Abstracts*, <https://doi.org/10.1609/aaai.v38i21.30437>. Full paper posted on arXiv, Verified Certificates via SAT and Computer Algebra Systems for the Ramsey  $R(3, 8)$  and  $R(3, 9)$  Problems (order of the first two authors swapped), <http://arxiv.org/abs/2502.06055> (2025).
- [DvoMe] V. Dvořák and H. Metrebian, A New Upper Bound for the Ramsey Number of Fans, *European Journal of Combinatorics*, **110** (2023) 103680, 14 pages.
- [Dyb1]\* J. Dybizbański, On Some Ramsey Numbers of  $C_4$  Versus  $K_{2,n}$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **87** (2013) 137-145.
- [Dyb2]\* J. Dybizbański, A Lower Bound on the Hypergraph Ramsey Number  $R(4, 5; 3)$ , *Contributions to Discrete Mathematics*, **13**(2) (2018) 112-115.
- [Dyb3]\* J. Dybizbański, University of Gdańsk, *personal communication* (2018). Addendum to [Dyb2] at <https://inf.ug.edu.pl/ramsey>.
- [DyDz1]\* J. Dybizbański and T. Dzido, On Some Ramsey Numbers for Quadrilaterals, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P154, **18**(1) (2011), 12 pages.
- [DyDz2] J. Dybizbański and T. Dzido, On Some Ramsey Numbers for Quadrilaterals Versus Wheels, *Graphs and Combinatorics*, **30** (2014) 573-579.
- [DyDR] J. Dybizbański, T. Dzido and S.P. Radziszowski, On Some Three-Color Ramsey Numbers for Paths, *Discrete Applied Mathematics*, **204** (2016) 133-141.
- [–] J. Dybizbański, see also [BoDD].
- [Dzi1]\* T. Dzido, Ramsey Numbers for Various Graph Classes (in Polish), *Ph.D. thesis*, University of Gdańsk, Poland, November 2005.

- [Dzi2]\* T. Džido, Multicolor Ramsey Numbers for Paths and Cycles, *Discussiones Mathematicae Graph Theory*, **25** (2005) 57-65.
- [DzFi1]\* T. Džido and R. Fidytek, The Number of Critical Colorings for Some Ramsey Numbers, *International Journal of Pure and Applied Mathematics*, ISSN 1311-8080, **38** (2007) 433-444.
- [DzFi2]\* T. Džido and R. Fidytek, On Some Three Color Ramsey Numbers for Paths and Cycles, *Discrete Mathematics*, **309** (2009) 4955-4958.
- [DzKP] T. Džido, M. Kubale and K. Piwakowski, On Some Ramsey and Turán-type Numbers for Paths and Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R55, **13** (2006), 9 pages.
- [DzNS] T. Džido, A. Nowik and P. Szuca, New Lower Bound for Multicolor Ramsey Numbers for Even Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #N13, **12** (2005), 5 pages.
- [ - ] T. Džido, see also [BoDD, DyDz1, DyDz2, DyDR, KhoDz].

## E

- [Ea1] Easy to obtain by simple combinatorics from other results, in particular by using graphs establishing lower bounds with smaller parameters.
- [Ea2] Unique 2-(6,3,2) design gives lower bound 7, upper bound is easy.
- [Ea3] Every edge (3, 3, 3;2)-coloring of  $K_{15}$  has 35 edges in each color [Hein], and since the number of triangles in  $K_{16}$  is not divisible by 3, hence no required triangle-coloring of  $K_{16}$  exists.
- [Eaton] N. Eaton, Ramsey Numbers for Sparse Graphs, *Discrete Mathematics*, **185** (1998) 63-75.
- [Eli] S. Eliahou, An Adaptive Upper Bound on the Ramsey Numbers  $R(3, \dots, 3)$ , *Integers*, <http://math.colgate.edu/~integers>, **20** (2020) #A54, 7 pages.
- [Erd1] P. Erdős, Some Remarks on the Theory of Graphs, *Bulletin of the American Mathematical Society*, **53** (1947) 292-294.
- [Erd2] P. Erdős, Some New Problems and Results in Graph Theory and Other Branches of Combinatorial Mathematics, *Combinatorics and Graph Theory* (Calcutta 1980), Berlin-NY Springer, LNM **885** (1981) 9-17.
- [Erd3] P. Erdős, On the Combinatorial Problems Which I Would Most Like to See Solved, *Combinatorica*, **1** (1981) 25-42.
- [Erd4] P. Erdős, On Some Problems in Graph Theory, Combinatorial Analysis and Combinatorial Number Theory, *Graph Theory and Combinatorics*, (Cambridge 1983), 1-17, Academic Press, London-New York, 1984.
- [EFRS1] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalized Ramsey Theory for Multiple Colors, *Journal of Combinatorial Theory, Series B*, **20** (1976) 250-264.
- [EFRS2] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, On Cycle-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **2** (1978) 53-64.
- [EFRS3] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Numbers for Brooms, *Congressus Numerantium*, **35** (1982) 283-293.
- [EFRS4] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Sparse Graph Ramsey Numbers, *Combinatorica*, **5** (1985) 311-318.
- [EFRS5] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Ramsey Problem of Harary on Graphs with Prescribed Size, *Discrete Mathematics*, **67** (1987) 227-233.
- [EFRS6] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Extremal Theory and Bipartite Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 103-112.
- [EFRS7] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, The Book-Tree Ramsey Numbers, *Scientia*, Series A: Mathematical Sciences, Valparaíso, Chile, **1** (1988) 111-117.
- [EFRS8] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Multipartite Graph-Tree Graph Ramsey Numbers, in *Graph Theory and Its Applications: East and West, Proceedings of the First China-USA*

- International Graph Theory Conference*, Annals of the New York Academy of Sciences, **576** (1989) 146-154.
- [EFRS9] P. Erdős, R.J. Faudree, C.C. Rousseau and R.H. Schelp, Ramsey Size Linear Graphs, *Combinatorics, Probability and Computing*, **2** (1993) 389-399.
- [ErdG] P. Erdős and R.L. Graham, On Partition Theorems for Finite Sets, in *Infinite and Finite Sets*, (A. Hajnal, R. Rado and V.T. Sós eds.) Vol. 1, 515-527, Colloq. Math. Soc. János Bolyai, Vol. 10, North Holland, 1975.
- [ErdH] P. Erdős and A. Hajnal, On Ramsey Like Theorems, Problems and Results, *Combinatorics*, Conference on Combinatorial Mathematics, Math. Institute, Oxford, (1972) 123-140.
- [-] P. Erdős, see also [BoEr, BuE1, BuE2, BuE3, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRSGJ, BEFS, BES, CaET].
- [Ex1]\* G. Exoo, Ramsey Numbers of Hypergraphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **2** (1987) 5-11.
- [Ex2]\* G. Exoo, Constructing Ramsey Graphs with a Computer, *Congressus Numerantium*, **59** (1987) 31-36.
- [Ex3]\* G. Exoo, Applying Optimization Algorithm to Ramsey Problems, in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi ed.), SIAM Philadelphia, (1989) 175-179.
- [Ex4]\* G. Exoo, A Lower Bound for  $R(5, 5)$ , *Journal of Graph Theory*, **13** (1989) 97-98.
- [Ex5]\* G. Exoo, On Two Classical Ramsey Numbers of the Form  $R(3, n)$ , *SIAM Journal on Discrete Mathematics*, **2** (1989) 488-490.
- [Ex6]\* G. Exoo, A Lower Bound for  $r(K_5 - e, K_5)$ , *Utilitas Mathematica*, **38** (1990) 187-188.
- [Ex7]\* G. Exoo, Three Color Ramsey Number of  $K_4 - e$ , *Discrete Mathematics*, **89** (1991) 301-305.
- [Ex8]\* G. Exoo, Indiana State University, *personal communication* (1992).
- [Ex9]\* G. Exoo, Announcement: On the Ramsey Numbers  $R(4, 6)$ ,  $R(5, 6)$  and  $R(3, 12)$ , *Ars Combinatoria*, **35** (1993) 85. The construction of a graph proving  $R(4, 6) \geq 35$  is presented in detail at <http://cs.indstate.edu/ge/RAMSEY> (2001).
- [Ex10]\* G. Exoo, A Lower Bound for Schur Numbers and Multicolor Ramsey Numbers of  $K_3$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R8, **1** (1994), 3 pages.
- [Ex12]\* G. Exoo, Some New Ramsey Colorings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R29, **5** (1998), 5 pages. The constructions are available electronically at <http://cs.indstate.edu/ge/RAMSEY>. The lower bounds presented in this paper have been improved.
- [Ex14]\* G. Exoo, Indiana State University, *New Lower Bounds for Table III*, (2000). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>.
- [Ex16]\* G. Exoo, Indiana State University, *personal communication* (2005-2006). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>.
- [Ex17]\* G. Exoo, Indiana State University, *personal communication* (2010-2011). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>.
- [Ex18]\* G. Exoo, Indiana State University, *personal communication* (2012-2013). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>.
- [Ex19]\* G. Exoo, On the Ramsey Number  $R(4, 6)$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P66, **19**(1) (2012), 5 pages.
- [Ex20]\* G. Exoo, On Some Small Classical Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P68, **20**(1) (2013), 6 pages.
- [Ex21]\* G. Exoo, Ramsey Colorings from  $p$ -Groups, *in preparation*, (2013).
- [Ex22]\* G. Exoo, Indiana State University, *personal communication* (2015). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>.
- [Ex23]\* G. Exoo, Indiana State University, *personal communication* (2017-2019). Constructions available at <http://cs.indstate.edu/ge/RAMSEY>, data on Paley graphs at <http://cs.indstate.edu/ge/Paley>.

- [Ex24]\* G. Exoo, Indiana State University, *personal communication* (2021).
- [Ex25]\* G. Exoo, Indiana State University, A Lower Bound for  $R(5, 6)$ , *manuscript* (2023).
- [EHM1] G. Exoo, H. Harborth and I. Mengersen, The Ramsey Number of  $K_4$  Versus  $K_5 - e$ , *Ars Combinatoria*, **25A** (1988) 277-286.
- [EHM2] G. Exoo, H. Harborth and I. Mengersen, On Ramsey Number of  $K_{2,n}$ , in *Graph Theory, Combinatorics, Algorithms, and Applications* (Y. Alavi, F.R.K. Chung, R.L. Graham and D.F. Hsu eds.), SIAM Philadelphia, (1989) 207-211.
- [ExRe]\* G. Exoo and D.F. Reynolds, Ramsey Numbers Based on  $C_5$ -Decompositions, *Discrete Mathematics*, **71** (1988) 119-127.
- [ExT]\* G. Exoo and M. Tatarovic, New Lower Bounds for 28 Classical Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.11, **22**(3) (2015), 12 pages. Graphs available at the journal site and at <http://cs.indstate.edu/ge/RAMSEY/ExTa>.
- [-] G. Exoo, see also [CIEHMS, XXER].

### Fa - Fe

- [FanHL] Chunchao Fan, Junqiang Huang and Qizhong Lin, Ramsey Numbers of Large Books Versus Multipartite Graphs, *Graphs and Combinatorics*, **40**:125 (2024), 14 pages.
- [FanLin1] Chunchao Fan and Qizhong Lin, Ramsey Non-Goodness Involving Books, *Journal of Combinatorial Theory, Series A*, **199** (2023) 105780, 19 pages.
- [FanLin2] Chunchao Fan and Qizhong Lin, Ramsey Numbers for Sparse Graphs Versus Path or Cycle, *preprint*, <http://arxiv.org/abs/2507.11835> (2025).
- [FLPS] R.J. Faudree, S.L. Lawrence, T.D. Parsons and R.H. Schelp, Path-Cycle Ramsey Numbers, *Discrete Mathematics*, **10** (1974) 269-277.
- [FM]\*\* R.J. Faudree and B.D. McKay, A Conjecture of Erdős and the Ramsey Number  $r(W_6)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **13** (1993) 23-31.
- [FRS1] R.J. Faudree, C.C. Rousseau and R.H. Schelp, All Triangle-Graph Ramsey Numbers for Connected Graphs of Order Six, *Journal of Graph Theory*, **4** (1980) 293-300.
- [FRS2] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Studies Related to the Ramsey Number  $r(K_5 - e)$ , in *Graph Theory and Its Applications to Algorithms and Computer Science*, (Y. Alavi et al. eds.), John Wiley and Sons, New York, (1985) 251-271.
- [FRS3] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Generalizations of the Tree-Complete Graph Ramsey Number, in *Graphs and Applications*, (F. Harary and J.S. Maybee eds.), John Wiley and Sons, New York, (1985) 117-126.
- [FRS4] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Small Order Graph-Tree Ramsey Numbers, *Discrete Mathematics*, **72** (1988) 119-127.
- [FRS5] R.J. Faudree, C.C. Rousseau and R.H. Schelp, A Good Idea in Ramsey Theory, in *Graph Theory, Combinatorics, Algorithms, and Applications* (San Francisco, CA 1989), SIAM Philadelphia, PA (1991) 180-189.
- [FRS6] R.J. Faudree, C.C. Rousseau and R.H. Schelp, Problems in Graph Theory from Memphis, in *The Mathematics of Paul Erdős II*, R.L. Graham et al. (eds.), Springer, New York, (2013) 95-118.
- [FRS7] R.J. Faudree, C.C. Rousseau and J. Sheehan, More from the Good Book, in *Proceedings of the Ninth Southeastern Conference on Combinatorics, Graph Theory, and Computing*, Utilitas Mathematica Publ., *Congressus Numerantium*, **XXI** (1978) 289-299.
- [FRS8] R.J. Faudree, C.C. Rousseau and J. Sheehan, Strongly Regular Graphs and Finite Ramsey Theory, *Linear Algebra and its Applications*, **46** (1982) 221-241.
- [FRS9] R.J. Faudree, C.C. Rousseau and J. Sheehan, Cycle-Book Ramsey Numbers, *Ars Combinatoria*, **31** (1991) 239-248.

- [FS1] R.J. Faudree and R.H. Schelp, All Ramsey Numbers for Cycles in Graphs, *Discrete Mathematics*, **8** (1974) 313-329.
- [FS2] R.J. Faudree and R.H. Schelp, Path Ramsey Numbers in Multicolorings, *Journal of Combinatorial Theory, Series B*, **19** (1975) 150-160.
- [FS3] R.J. Faudree and R.H. Schelp, Ramsey Numbers for All Linear Forests, *Discrete Mathematics*, **16** (1976) 149-155.
- [FS4] R.J. Faudree and R.H. Schelp, Some Problems in Ramsey Theory, in *Theory and Applications of Graphs*, (conference proceedings, Kalamazoo, MI 1976), Lecture Notes in Mathematics **642**, Springer, Berlin, (1978) 500-515.
- [FSR] R.J. Faudree, R.H. Schelp and C.C. Rousseau, Generalizations of a Ramsey Result of Chvátal, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 351-361.
- [FSS1] R.J. Faudree, R.H. Schelp and J. Sheehan, Ramsey Numbers for Matchings, *Discrete Mathematics*, **32** (1980) 105-123.
- [FSS2] R.J. Faudree, R.H. Schelp and M. Simonovits, On Some Ramsey Type Problems Connected with Paths, Cycles and Trees, *Ars Combinatoria*, **29A** (1990) 97-106.
- [FSS3] R.J. Faudree, A. Schelten and I. Schiermeyer, The Ramsey Number  $r(C_7, C_7, C_7)$ , *Discussiones Mathematicae Graph Theory*, **23** (2003) 141-158.
- [FaSi] R.J. Faudree and M. Simonovits, Ramsey Problems and Their Connection to Turán-Type Extremal Problems, *Journal of Graph Theory*, **16** (1992) 25-50.
- [-] R.J. Faudree, see also [BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRSGJ, BEFS, BuF, BFRS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9].
- [FeKR]\*\* S. Fettes, R.L. Kramer and S.P. Radziszowski, An Upper Bound of 62 on the Classical Ramsey Number  $R(3, 3, 3, 3)$ , *Ars Combinatoria*, **72** (2004) 41-63.
- [-] A. Ferber, see [ConFer].
- [Ferg] D.G. Ferguson, Topics in Graph Colouring and Graph Structures, *Ph.D. thesis*, Department of Mathematics, London School of Economics and Political Science, London, 2013. The problems on Ramsey theory are presented also in three arXiv preprints "The Ramsey Number of Mixed-Parity Cycles I, II and III", <http://arxiv.org/abs/1508.07154>, [1508.07171](http://arxiv.org/abs/1508.07171) and [1508.07176](http://arxiv.org/abs/1508.07176) (2015).

## Fi - Fu

- [Fid1]\* R. Fidytek, Two- and Three-Color Ramsey Numbers for Paths and Cycles, *manuscript* (2010).
- [Fid2]\* R. Fidytek, *personal communication*, Ramsey Graphs  $R(K_n, K_m - e)$ , <http://fidytek.inf.ug.edu.pl/ramsey> (2010), available until 2014.
- [-] R. Fidytek, see also [DzFi1, DzFi2].
- [FiŁu1] A. Figaj and T. Łuczak, The Ramsey Number for a Triple of Long Even Cycles, *Journal of Combinatorial Theory, Series B*, **97** (2007) 584-596.
- [FiŁu2] A. Figaj and T. Łuczak, The Ramsey Numbers for a Triple of Long Cycles, *Combinatorica*, **38**(4) (2018) 827-845.
- [FizGM] G. Fiz Pontiveros, S. Griffiths and R. Morris, The Triangle-Free Process and  $R(3, k)$ , *Memoirs of the American Mathematical Society*, Vol. **263**, Number 1274 (2020), 125 pages, first version on arXiv, <http://arxiv.org/abs/1302.6279> (2013).
- [FizGMSS] G. Fiz Pontiveros, S. Griffiths, R. Morris, D. Saxton and J. Skokan, The Ramsey Number of the Clique and the Hypercube, *Journal of the London Mathematical Society*, **89** (2014) 680-702.
- [-] G. Fiz Pontiveros, see also [GrMFSS].

- [FloS] Freddy Flores Dubó and Maya Stein, On the Ramsey Number of the Double Star, *Discrete Mathematics*, **348** (2025) 114227, 4 pages.
- [Fol] J. Folkman, Notes on the Ramsey Number  $N(3, 3, 3, 3)$ , *Journal of Combinatorial Theory, Series A*, **16** (1974) 371-379.
- [-] N. Fountoulakis, see [CooFKO1, CooFKO2, KüCFO].
- [FoxHW] Jacob Fox, Xiaoyu He and Yuval Wigderson, Ramsey Goodness of Books Revisited, *Advances in Combinatorics*, <https://doi.org/10.19086/aic.2023.4> (2023):4, 21 pages.
- [FoxLi] Jacob Fox and Ray Li, On Ramsey Numbers of Hedgehogs, *Combinatorics, Probability and Computing*, **29** (2020) 101-112.
- [FoxPS1] J. Fox, J. Pach and A. Suk, Bounded VC-Dimension Implies the Schur-Erdős Conjecture, *36th International Symposium on Computational Geometry*, Dagstuhl Publishing, Germany, 46:1-46:8, (2020). *Combinatorica*, **41**(6) (2021) 803-813.
- [FoxPS2] J. Fox, J. Pach and A. Suk, The Schur-Erdős Problem for Semi-Algebraic Colorings, *Israel Journal of Mathematics*, **239** (2020) 39-57.
- [FoxSu1] J. Fox and B. Sudakov, Density Theorems for Bipartite Graphs and Related Ramsey-type Results, *Combinatorica*, **29** (2009) 153-196.
- [FoxSu2] J. Fox and B. Sudakov, Two Remarks on the Burr-Erdős Conjecture, *European Journal of Combinatorics*, **30** (2009) 1630-1645.
- [-] J. Fox, see also [BraFS1, BraFS2, ConFG1+, ConFG2+, ConFH+, ConFLS, ConFR, ConFS1, ConFS2, ConFS3, ConFS4, ConFS5, ConFS6, ConFS7, ConFS8, ConFW1, ConFW2].
- [-] M. Frank, see [CodFIM].
- [FraWi] P. Frankl and R.M. Wilson, Intersection Theorems with Geometric Consequences, *Combinatorica*, **1** (1981) 357-368.
- [Fra1] K. Fraughnaugh Jones, Independence in Graphs with Maximum Degree Four, *Journal of Combinatorial Theory, Series B*, **37** (1984) 254-269.
- [Fra2] K. Fraughnaugh Jones, Size and Independence in Triangle-Free Graphs with Maximum Degree Three, *Journal of Graph Theory*, **14** (1990) 525-535.
- [FrLo] K. Fraughnaugh and S.C. Locke, Finding Independent Sets in Triangle-Free Graphs, *SIAM Journal on Discrete Mathematics*, **9** (1996) 674-681.
- [Fre] H. Fredricksen, Schur Numbers and the Ramsey Numbers  $N(3, 3, \dots, 3; 2)$ , *Journal of Combinatorial Theory, Series A*, **27** (1979) 376-377.
- [FreSw]\* H. Fredricksen and M.M. Sweet, Symmetric Sum-Free Partitions and Lower Bounds for Schur Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R32, **7** (2000), 9 pages.
- [-] A. Freschi, see [CamFMPP].
- [-] Lars Fritz, see [WoGKSF].
- [FuLS]\*\* F. Furini, I. Ljubić and P. San Segundo, New IP-based Lower Bounds for Small Ramsey Numbers Using Circulant Graphs, *preprint*, <https://www.researchgate.net/publication/354381524>, 2021, 29 pages.
- [Fuj1]\* H. Fujita, Ramsey Numbers and Ramsey Graphs, <http://opal.inf.kyushu-u.ac.jp/~fujita/ramsey.html>, 2014-2017.
- [Fuj2] S. Fujita, Generalized Ramsey Numbers for Graphs with Three Disjoint Cycles Versus a Complete Graph, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P14, **19**(2) (2012), 11 pages.
- [-] Z. Füredi, see [AxFM, BiFJ].

## Ga - Gr

- [ - ] F. Gaitan, see [RanMCG].
- [ - ] Vijay Ganesh, see [DuLBG].
- [ - ] P. Garcia-Vázquez, see [BoCGR].
- [Gas] W. Gasarch, Applications of Ramsey Theory to Computer Science, collection of pointers to papers, <http://www.cs.umd.edu/~gasarch/TOPICS/ramsey/ramsey.html> (2009, 2011, 2017).
- [GauST] S. Gautam, A.K. Srivastava and A. Tripathi, On Multicolour Noncomplete Ramsey Graphs of Star Graphs, *Discrete Applied Mathematics*, **156** (2008) 2423-2428.
- [GauB]\* Thibault Gauthier and Chad E. Brown, A Formal Proof of  $R(4, 5) = 25$ , *15th International Conference on Interactive Theorem Proving*, ITP 2024, Dagstuhl Publishing, article #16, 18 pages.
- [ - ] Gennian Ge, see [XuGe].
- [Gerbi]\* R. Gerbicz, New Lower Bounds for Two Color and Multicolor Ramsey Numbers, *preprint*, <http://arxiv.org/abs/1004.4374> (2010), pointed to in revision #14. Since 2015, better bounds in all cases were obtained by others.
- [Gerbn] D. Gerbner, On Berge-Ramsey Problems, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P2.39, **27**(2) (2020), 8 pages.
- [GerMOV] D. Gerbner, A. Methuku, G. Omid and M. Vizer, Ramsey Problems for Berge Hypergraphs, *SIAM Journal on Discrete Mathematics*, **34** (2020) 351-369.
- [GeGy] L. Gerencsér and A. Gyárfás, On Ramsey-Type Problems, *Annales Universitatis Scientiarum Budapestinensis, Eötvös Sect. Math.*, **10** (1967) 167-170.
- [ - ] Aris Giotis, see [WoGKSF].
- [Gi1] G. Giraud, Une généralisation des nombres et de l'inégalité de Schur, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A437-A440.
- [Gi2] G. Giraud, Minoration de certains nombres de Ramsey binaires par les nombres de Schur généralisés, *C.R. Acad. Sc. Paris, Séries A-B*, **266** (1968) A481-A483.
- [Gi3] G. Giraud, Nouvelles majorations des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **268** (1969) A5-A7.
- [Gi4] G. Giraud, Majoration du nombre de Ramsey ternaire-bicolore en (4,4), *C.R. Acad. Sc. Paris, Séries A-B*, **269** (1969) A620-A622.
- [Gi5] G. Giraud, Une minoration du nombre de quadrangles unicolores et son application à la majoration des nombres de Ramsey binaires-bicolores, *C.R. Acad. Sc. Paris, Séries A-B*, **276** (1973) A1173-A1175.
- [Gi6] G. Giraud, Sur le problème de Goodman pour les quadrangles et la majoration des nombres de Ramsey, *Journal of Combinatorial Theory, Series B*, **27** (1979) 237-253.
- [ - ] L. Gishboliner, see [BraGS].
- [ - ] A.M. Gleason, see [GG].
- [GodK] W. Goddard and D.J. Kleitman, An Upper Bound for the Ramsey Numbers  $r(K_3, G)$ , *Discrete Mathematics*, **125** (1994) 177-182.
- [GoeR1]\*\* J. Goedgebeur and S.P. Radziszowski, New Computational Upper Bounds for Ramsey Numbers  $R(3, k)$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P30, **20**(1) (2013), 28 pages.
- [GoeR2]\*\* J. Goedgebeur and S.P. Radziszowski, The Ramsey Number  $R(3, K_{10} - e)$  and Computational Bounds for  $R(3, G)$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P19, **20**(4) (2013), 25 pages.
- [GoeVO]\*\* J. Goedgebeur and S. Van Overberghe, New Bounds for Ramsey Numbers  $R(K_k - e, K_l - e)$ , *Discrete Applied Mathematics*, **307** (2022) 212-221.

- [-] J. Goedgebeur, see also [BrCGM, BrGS].
- [GoMC] A. Gonçalves and E.L. Monte Carmelo, Some Geometric Structures and Bounds for Ramsey Numbers, *Discrete Mathematics*, **280** (2004) 29-38.
- [GoJa1] R.J. Gould and M.S. Jacobson, Bounds for the Ramsey Number of a Disconnected Graph Versus Any Graph, *Journal of Graph Theory*, **6** (1982) 413-417.
- [GoJa2] R.J. Gould and M.S. Jacobson, On the Ramsey Number of Trees Versus Graphs with Large Clique Number, *Journal of Graph Theory*, **7** (1983) 71-78.
- [-] R.J. Gould, see also [BEFRSGJ, ChaGP].
- [-] N. Graber, see [ColGJ].
- [GrBu] R.L. Graham and S. Butler, Rudiments of Ramsey Theory, second edition, *CBMS Regional Conference Series in Mathematics*, 123, American Mathematical Society 2015.
- [GrNe] R.L. Graham and J. Nešetřil, Ramsey Theory and Paul Erdős (Recent Results from a Historical Perspective), *Bolyai Society Mathematical Studies*, **11**, Budapest (2002) 339-365.
- [GrRö] R.L. Graham and V. Rödl, Numbers in Ramsey Theory, in *Surveys in Combinatorics*, (ed. C. Whitehead), Cambridge University Press, 1987, 111-153.
- [GRR1] R.L. Graham, V. Rödl and A. Ruciński, On Graphs with Linear Ramsey Numbers, *Journal of Graph Theory*, **35** (2000) 176-192.
- [GRR2] R.L. Graham, V. Rödl and A. Ruciński, On Bipartite Graphs with Linear Ramsey Numbers, Paul Erdős and his mathematics, *Combinatorica*, **21** (2001) 199-209.
- [GRS] R.L. Graham, B.L. Rothschild and J.H. Spencer, *Ramsey Theory*, John Wiley & Sons, first edition 1980, second edition 1990, paperback of the second edition 2013.
- [-] R.L. Graham, see also [ChGra1, ChGra2, ErdG].
- [GrY] J.E. Graver and J. Yackel, Some Graph Theoretic Results Associated with Ramsey's Theorem, *Journal of Combinatorial Theory*, **4** (1968) 125-175.
- [GG] R.E. Greenwood and A.M. Gleason, Combinatorial Relations and Chromatic Graphs, *Canadian Journal of Mathematics*, **7** (1955) 1-7.
- [GrH] U. Grenda and H. Harborth, The Ramsey Number  $r(K_3, K_7 - e)$ , *Journal of Combinatorics, Information & System Sciences*, **7** (1982) 166-169.
- [GrMFSS] S. Griffiths, R. Morris, G. Fiz Pontiveros, D. Saxton and J. Skokan, On the Ramsey Number of the Triangle and the Cube, *Combinatorica*, **36** (2016) 71-89.
- [-] S. Griffiths, see also [BalB+, CamGMS, FizGM, FizGMSS].
- [Gri] J.R. Griggs, An Upper Bound on the Ramsey Numbers  $R(3, k)$ , *Journal of Combinatorial Theory, Series A*, **35** (1983) 145-153.
- [GR]\*\* C. Grinstead and S. Roberts, On the Ramsey Numbers  $R(3, 8)$  and  $R(3, 9)$ , *Journal of Combinatorial Theory, Series B*, **33** (1982) 27-51.
- [-] C. Grinstead, see also [ChGri].
- [Grol1] V. Grolmusz, Superpolynomial Size Set-Systems with Restricted Intersections mod 6 and Explicit Ramsey Graphs, *Combinatorica*, **20** (2000) 73-88.
- [Grol2] V. Grolmusz, Low Rank Co-Diagonal Matrices and Ramsey Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R15, **7** (2000), 7 pages.
- [Grol3] V. Grolmusz, Set-Systems with Restricted Multiple Intersections, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R8, **9** (2002), 10 pages.
- [Gros1] J.W. Grossman, Some Ramsey Numbers of Unicyclic Graphs, *Ars Combinatoria*, **8** (1979) 59-63.
- [Gros2] J.W. Grossman, The Ramsey Numbers of the Union of Two Stars, *Utilitas Mathematica*, **16** (1979) 271-279.

- [GrHK] J.W. Grossman, F. Harary and M. Klawe, Generalized Ramsey Theory for Graphs, X: Double Stars, *Discrete Mathematics*, **28** (1979) 247-254.
- [-] J.W. Grossman, see also [BuG].

### Gu - Gy

- [GuLi] Gu Hua and Li Yusheng, On Ramsey Number of  $K_{2,t+1}$  vs  $K_{1,n}$ , *Journal of Nanjing University Mathematical Biquarterly*, **19** (2002) 150-153.
- [GuSL] Gu Hua, Song Hongxue and Liu Xiangyang, Ramsey Numbers  $r(K_{1,4}, G)$  for All Three-Partite Graphs  $G$  of Order Six, *Journal of Southeast University*, (English Edition), **20** (2004) 378-380.
- [-] Gu Hua, see also [SonGQ].
- [-] B. Gunby, see [ConFG1+, ConFG2+].
- [GuoHP] Xiao-bing Guo, Si-nan Hu and Yue-jian Peng, Ramsey Numbers of Trees Versus Multiple Copies of Books, *Acta Mathematicae Applicatae Sinica, English Series*, **40** (2024) 600-612.
- [GuoV] Guo Yubao and L. Volkmann, Tree-Ramsey Numbers, *Australasian Journal of Combinatorics*, **11** (1995) 169-175.
- [-] L. Gupta, see [GuGS].
- [GuNNW] P. Gupta, N. Ndiaye, S. Norin and L. Wei, Optimizing the CGMS Upper Bound on Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2407.19026> (2024).
- [Gup] Sayan Gupta, A Study of Two Ramsey Numbers Involving Odd Cycles, *preprint*, <http://arxiv.org/abs/2504.15693> (2025).
- [GupM] Sayan Gupta and Kaushik Majumder, An Exact Ramsey Number of Large Bipartite Graphs Versus Odd Wheel, *preprint*, <http://arxiv.org/abs/2511.14867> (2025).
- [GuGS] S.K. Gupta, L. Gupta and A. Sudan, On Ramsey Numbers for Fan-Fan Graphs, *Journal of Combinatorics, Information & System Sciences*, **22** (1997) 85-93.
- [Gy] A. Gyárfás, Large Sets of  $t$ -Designs and a Ramsey-Type Problem, *Integers*, **18** (2018) A53, 3 pages, <http://math.colgate.edu/~integers>.
- [GyLSS] A. Gyárfás, J. Lehel, G.N. Sárközy and R.H. Schelp, Monochromatic Hamiltonian Berge-Cycles in Colored Complete Uniform Hypergraphs, *Journal of Combinatorial Theory, Series B*, **98** (2008) 342-358.
- [GyRa] A. Gyárfás and G. Raeisi, The Ramsey Number of Loose Triangles and Quadrangles in Hypergraphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P30, **19**(2) (2012), 9 pages.
- [GyRSS] A. Gyárfás, M. Ruszinkó, G.N. Sárközy and E. Szemerédi, Three-color Ramsey Numbers for Paths, *Combinatorica*, **27** (2007) 35-69. *Corrigendum* in **28** (2008) 499-502.
- [GySá1] A. Gyárfás and G.N. Sárközy, The 3-Colour Ramsey Number of a 3-Uniform Berge Cycle, *Combinatorics, Probability and Computing*, **20** (2011) 53-71.
- [GySá2] A. Gyárfás and G.N. Sárközy, Star Versus Two Stripes Ramsey Numbers and a Conjecture of Schelp, *Combinatorics, Probability and Computing*, **21** (2012) 179-186.
- [GySá3] A. Gyárfás and G.N. Sárközy, Ramsey Number of a Connected Triangle Matching, *Journal of Graph Theory*, **83** (2016) 109-119.
- [GySS1] A. Gyárfás, G.N. Sárközy and E. Szemerédi, The Ramsey Number of Diamond-Matchings and Loose Cycles in Hypergraphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R126, **15** (2008), 14 pages.
- [GySS2] A. Gyárfás, G.N. Sárközy and E. Szemerédi, Monochromatic Hamiltonian 3-Tight Berge Cycles in 2-Colored 4-Uniform Hypergraphs, *Journal of Graph Theory*, **63** (2010) 288-299.
- [GySeT] A. Gyárfás, A. Sebő and N. Trotignon, The Chromatic Gap and Its Extremes, *Journal of Combinatorial Theory, Series B*, **102** (2012) 1155-1178.

- [GyTu] A. Gyárfás and Z. Tuza, An Upper Bound on the Ramsey Number of Trees, *Discrete Mathematics*, **66** (1987) 309-310.
- [-] A. Gyárfás, see also [AxGy, AxGLM, DeBGS, GeGy].

## Ha

- [HafBa] Y. Hafidh and E.T. Baskoro, The Ramsey Number for Tree Versus Wheel with Odd Order, *Bulletin of the Malaysian Mathematical Sciences Society*, **44**(4) (2021) 2151-2160.
- [Häg] R. Häggkvist, On the Path-Complete Bipartite Ramsey Number, *Discrete Mathematics*, **75** (1989) 243-245.
- [HagMa] Sh. Haghi and H.R. Maimani, A Note on the Ramsey Number of Even Wheels Versus Stars, *Discusiones Mathematicae Graph Theory*, **38** (2018) 397-404.
- [HaHT] A. Hamm, P. Hazelton and S. Thompson, On Ramsey and Star-Critical Ramsey Numbers for Generalized Fans Versus  $nK_m$ , *Discrete Applied Mathematics*, **305** (2021) 64-70. Corrigendum in **376** (2025) 462-464.
- [-] A. Hajnal, see [ErdH].
- [HanPR] Jie Han, J. Polcyn and A. Ruciński, Turán and Ramsey Numbers for 3-Uniform Minimal Paths of Length 4, *Journal of Graph Theory*, **98**(3) (2021) 460-498.
- [Han]\* D. Hanson, Sum-Free Sets and Ramsey Numbers, *Discrete Mathematics*, **14** (1976) 57-61.
- [-] D. Hanson, see also [AbbH].
- [Hans] M. Hansson, On Generalized Ramsey Numbers for Two Sets of Cycles, *Discrete Applied Mathematics*, **238** (2018) 86-94.
- [HaoLin] Yiyuan Hao and Qizhong Lin, Ramsey Number of  $K_3$  Versus  $F_{3,n}$ , *Discrete Applied Mathematics*, **251** (2018) 345-348.
- [Har1] F. Harary, Recent Results on Generalized Ramsey Theory for Graphs, in *Graph Theory and Applications*, (Y. Alavi et al. eds.) Springer, Berlin (1972) 125-138.
- [Har2] F. Harary, Generalized Ramsey Theory I to XIII: Achievement and Avoidance Numbers, in *Proceedings of the Fourth International Conference on the Theory and Applications of Graphs*, (Kalamazoo, MI 1980), John Wiley & Sons, (1981) 373-390.
- [-] F. Harary, see also [ChH1, ChH2, ChH3, GrHK].
- [HaKr1]\*\* H. Harborth and S. Krause, Ramsey Numbers for Circulant Colorings, *Congressus Numerantium*, **161** (2003) 139-150.
- [HaKr2]\*\* H. Harborth and S. Krause, Distance Ramsey Numbers, *Utilitas Mathematica*, **70** (2006) 197-200.
- [HaMe1] H. Harborth and I. Mengersen, An Upper Bound for the Ramsey Number  $r(K_5 - e)$ , *Journal of Graph Theory*, **9** (1985) 483-485.
- [HaMe2] H. Harborth and I. Mengersen, All Ramsey Numbers for Five Vertices and Seven or Eight Edges, *Discrete Mathematics*, **73** (1988/89) 91-98.
- [HaMe3] H. Harborth and I. Mengersen, The Ramsey Number of  $K_{3,3}$ , in *Combinatorics, Graph Theory, and Applications*, Vol. 2 (Y. Alavi, G. Chartrand, O.R. Oellermann and J. Schwenk eds.), John Wiley & Sons, (1991) 639-644.
- [-] H. Harborth, see also [BoH, CIEHMS, EHM1, EHM2, GrH].
- [HaMe4] M. Harborth and I. Mengersen, Some Ramsey Numbers for Complete Bipartite Graphs, *Australasian Journal of Combinatorics*, **13** (1996) 119-128.
- [-] T. Harmuth, see [BrBH1, BrBH2].
- [HasHKL] John Haslegrave, Joseph Hyde, Jaehoon Kim and Hong Liu, Ramsey Numbers of Cycles Versus General Graphs, *Forum of Mathematics, Sigma*, **11**:e10 (2023) 1-18.

- [Has] Hasmawati, The Ramsey Numbers for Disjoint Union of Stars, *Journal of the Indonesian Mathematical Society*, **16** (2010) 133-138.
- [HaABS] Hasmawati, H. Assiyatun, E.T. Baskoro and A.N.M. Salman, Ramsey Numbers on a Union of Identical Stars Versus a Small Cycle, in *Computational Geometry and Graph Theory, Kyoto CGGT 2007, LNCS 4535*, Springer, Berlin (2008) 85-89.
- [HaBA1] Hasmawati, E.T. Baskoro and H. Assiyatun, Star-Wheel Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **55** (2005) 123-128.
- [HaBA2] Hasmawati, E.T. Baskoro and H. Assiyatun, The Ramsey Numbers for Disjoint Unions of Graphs, *Discrete Mathematics*, **308** (2008) 2046-2049.
- [HaJu] Hasmawati and Jusmawati, Ramsey Number on a Union of Stars Versus a Small Cycle, *Jurnal Matematika Statistika dan Komputasi*, Universitas Hasanuddin, **13** (2017) 152-157.
- [-] Hasmawati, see also [BaHA].
- [HasHR] Z.R. Hassan, E. Hemaspaandra and S. Radziszowski, The Complexity of  $(P_k, P_l)$ -Arrowing, *24th International Symposium on Fundamentals of Computation Theory*, FCT 2023, Trier, Germany, September 18-21, LNCS, Springer, 2023.
- [Hass] Z.R. Hassan, The Complexity of  $(P_3, H)$ -Arrowing and Beyond, *49th International Symposium on Mathematical Foundations of Computer Science MFCS 2024*, Bratislava, Slovakia, August 26-30, Leibniz International Proceedings in Informatics, article #59, 16 pages.
- [HaLP1+] P.E. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, M. Simonovits and J. Skokan, The Ramsey Number for Hypergraph Cycles I, *Journal of Combinatorial Theory, Series A*, **113** (2006) 67-83.
- [HaLP2+] P.E. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński and J. Skokan, The Ramsey Number for 3-Uniform Tight Hypergraph Cycles, *Combinatorics, Probability and Computing*, **18** (2009) 165-203.
- [HaLT] P.E. Haxell, T. Łuczak and P.W. Tingley, Ramsey Numbers for Trees of Small Maximum Degree, *Combinatorica*, **22** (2002) 287-320.
- [-] P. Hazelton, see [HaHT].

## He - Hu

- [HeLD]\* Changxiang He, Yusheng Li and Lin Dong, Three-Color Ramsey Numbers of  $K_n$  Dropping an Edge, *Graphs and Combinatorics*, **28** (2012) 663-669.
- [HeZZ] Jiafu He, Haiyu Zeng and Yanbo Zhang, Ramsey Goodness of Stars and Fans for the Hajós Graph, *Discrete Mathematics and Theoretical Computer Science*, **27**(3) (2025), #11, 11 pages.
- [HeNWY] Xiaoyu He, Jiayi Nie, Yuval Wigderson and Hung-Hsun Hans Yu, Off-Diagonal Ramsey Numbers for Linear Hypergraphs, *preprint*, <http://arxiv.org/abs/2507.05641> (2025).
- [HeWi] Xiaoyu He and Yuval Wigderson, Multicolor Ramsey Numbers via Pseudorandom Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.32, **27**(1) (2020), 8 pages.
- [-] Xiaoyu He, see also [ConFG1+, ConFG2+, ConFH+, FoxHW].
- [HeHKP] Zion Hefty, Paul Horn, Dylan King and Florian Pfender, Improving  $R(3, k)$  in Just Two Bites, *preprint*, <http://arxiv.org/abs/2510.19718> (2025).
- [-] P. Heggernes, see [BeHHRS, BeHHS].
- [Hein] K. Heinrich, Proper Colourings of  $K_{15}$ , *Journal of the Australian Mathematical Society, Series A*, **24** (1977) 465-495.
- [-] E. Hemaspaandra, see [HasHR].
- [He1] G.R.T. Hendry, Diagonal Ramsey Numbers for Graphs with Seven Edges, *Utilitas Mathematica*, **32** (1987) 11-34.
- [He2] G.R.T. Hendry, Ramsey Numbers for Graphs with Five Vertices, *Journal of Graph Theory*, **13** (1989) 245-248.

- [He3] G.R.T. Hendry, The Ramsey Numbers  $r(K_2 + \bar{K}_3, K_4)$  and  $r(K_1 + C_4, K_4)$ , *Utilitas Mathematica*, **35** (1989) 40-54, addendum in **36** (1989) 25-32.
- [He4] G.R.T. Hendry, Critical Colorings for Clancy's Ramsey Numbers, *Utilitas Mathematica*, **41** (1992) 181-203.
- [He5] G.R.T. Hendry, Small Ramsey Numbers II. Critical Colorings for  $r(C_5 + e, K_5)$ , *Quaestiones Mathematica*, **17** (1994) 249-258.
- [-] G.R.T. Hendry, see also [YH].
- [HiIr]\* R. Hill and R.W. Irving, On Group Partitions Associated with Lower Bounds for Symmetric Ramsey Numbers, *European Journal of Combinatorics*, **3** (1982) 35-50.
- [-] J. Hiller, see [BrBH, BudHLS, BudHMP, BudHP, BudHR1, BudHR2].
- [Hir] J. Hirschfeld, A Lower Bound for Ramsey's Theorem, *Discrete Mathematics*, **32** (1980) 89-91.
- [HngJL] Eng Keat Hng, Meng Ji and Ander Lamaison, Ramsey Size Linear and Generalization, *preprint*, <http://arxiv.org/abs/2603.25453> (2026).
- [Ho] Pak Tung Ho, On Ramsey Unsaturated and Saturated Graphs, *Australasian Journal of Combinatorics*, **46** (2010) 13-18.
- [HoMe] M. Hoeth and I. Mengersen, Ramsey Numbers for Graphs of Order Four Versus Connected Graphs of Order Six, *Utilitas Mathematica*, **57** (2000) 3-19.
- [-] Pim van 't Hof, see [BeHHRS, BeHHS].
- [-] P. Holub, see [LiZBBH].
- [-] N. Hommowun, see [AlmHS].
- [Hook] J. Hook, Critical Graphs for  $R(P_n, P_m)$  and the Star-Critical Ramsey Number for Paths, *Discussiones Mathematicae Graph Theory*, **35** (2015) 689-701.
- [HoIs] J. Hook and G. Isaak, Star-Critical Ramsey Numbers, *Discrete Applied Mathematics*, **159** (2011) 328-334.
- [-] Paul Horn, see [HeHKP].
- [-] Xinmin Hou, see [JiHou, ZhaHou].
- [HuL+] Fu-Tao Hu, Qizhong Lin, Tomasz Łuczak, Bo Ning and Xing Peng, Ramsey Numbers of Books Versus Long Cycles, *SIAM Journal on Discrete Mathematics*, **39**(1) (2025) 550-561.
- [HuLuo] Sinan Hu and Zhidan Luo, Ramsey Numbers for a Large Tree Versus Multiple Copies of Complete Graphs of Different Sizes, *Discussiones Mathematicae Graph Theory*, **45** (2025) 419-429.
- [HuP1] Sinan Hu and Yuejian Peng, The Ramsey Number for a Forest Versus Disjoint Union of Complete Graphs, *Graphs and Combinatorics*, **39**:26 (2023), 16 pages.
- [HuP2] Si-nan Hu and Yue-jian Peng, Ramsey Numbers of Stripes Versus Trees and Unicyclic Graphs, *Journal of the Operations Research Society of China*, (2023), <https://doi.org/10.1007/s40305-023-00494-0>.
- [-] Si-nan Hu, see also [GuoHP].
- [HuLin] Xinyu Hu and Qizhong Lin, Ramsey Numbers and a General Erdős-Rogers Function, *Discrete Mathematics*, **347** (2024) 114203, 10 pages.
- [HuPZ] Caixia Huang, Yuejian Peng and Yiran Zhang, Ramsey Numbers for Multiple Copies of Graphs in a Component, *Graphs and Combinatorics*, **40**:94 (2024), 23 pages.
- [HuSo] Huang Da Ming and Song En Min, Properties and Lower Bounds of the Third Order Ramsey Numbers (in Chinese), *Mathematica Applicata*, **9** (1996) 105-107.
- [Hua1] Huang Guotai, Some Generalized Ramsey Numbers (in Chinese), *Mathematica Applicata*, **1** (1988) 97-101.
- [Hua2] Huang Guotai, An Unsolved Problem of Gould and Jacobson (in Chinese), *Mathematica Applicata*, **9** (1996) 234-236.
- [-] Huang Jian, see [HTHZ1, HTHZ2, HWSYZH].

- [-] Junqiang Huang, see [FanHL].
- [HuCh] Ting Huang and Yaojun Chen, Ramsey Numbers of Trees Versus Odd Cycles Sharing One Vertex, *Discrete Mathematics*, **349** (2026) 115050, 8 pages.
- [HuZC1] Ting Huang, Yanbo Zhang and Yaojun Chen, Fan-Goodness of Sparse Graphs, *preprint*, <http://arxiv.org/abs/2507.09832> (2025).
- [HuZC2] Ting Huang, Yanbo Zhang and Yaojun Chen, Ramsey Numbers of Sparse Graphs Versus Disjoint Books, *preprint*, <http://arxiv.org/abs/2507.09827> (2025).
- [HuZC3] Ting Huang, Yanbo Zhang and Yaojun Chen, Minimum Degree and Sparse Connected Spanning Subgraphs, *preprint*, <http://arxiv.org/abs/2507.03264> (2025).
- [-] Huang Wenke, see [DuHu].
- [HTHZ1] (also abbreviated as HT+) Yiru Huang, Fuping Tan, Jian Huang and Kemin Zhang, New Upper Bounds for Ramsey Number  $R(K_m - e, K_n - e)$ , *manuscript* (2016).
- [HTHZ2] Huang Yiru, Tan Fuping, Huang Jian and Zhang Chaohui, On the Upper Bounds Formulas of Multicolored Ramsey Number (in Chinese), *Journal of Jishou University*, Natural Science Edition, **38** (2017) 1-6.
- [HWSYZH] (also abbreviated as HW+) Huang Yi Ru, Wang Yuandi, Sheng Wancheng, Yang Jiansheng, Zhang Ke Min and Huang Jian, New Upper Bound Formulas with Parameters for Ramsey Numbers, *Discrete Mathematics*, **307** (2007) 760-763.
- [HYZ] Yi Ru Huang, Jian Sheng Yang and Ke Min Zhang, A Note on Ramsey Numbers with Two Parameters, *European Journal of Combinatorics*, **27** (2006) 574-576.
- [HZ1] Huang Yi Ru and Zhang Ke Min, A New Upper Bound Formula on Ramsey Numbers, *Journal of Shanghai University, Natural Science*, **7** (1993) 1-3.
- [HZ2] Huang Yi Ru and Zhang Ke Min, An New Upper Bound Formula for Two Color Classical Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **28** (1998) 347-350.
- [HZ3] Huang Yi Ru and Zhang Ke Min, New Upper Bounds for Ramsey Numbers, *European Journal of Combinatorics*, **19** (1998) 391-394.
- [-] Huang Yi Ru, see also [BoJY+, YHZ1, YHZ2].
- [-] Zhong Huang, see [YaoHMZ].
- [HunMS] Zach Hunter, Aleksa Milojević and Benny Sudakov, Gaussian Random Graphs and Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2512.17718> (2025).
- [-] Zach Hunter, see also [BraHS].
- [-] Eoin Hurley, see [BalB+].
- [-] Joseph Hyde, see [HasHKL].

## I

- [Ihr]\* F. Ihringer, Two Small Improvements on Ramsey Numbers, *unpublished report*, <http://math.ihringer.org/publications.php> (2020).
- [Ir] R.W. Irving, Generalised Ramsey Numbers for Small Graphs, *Discrete Mathematics*, **9** (1974) 251-264.
- [-] R.W. Irving, see also [HiIr].
- [-] G. Isaak, see [HoIs].
- [Isb1] J.R. Isbell,  $N(4, 4; 3) \geq 13$ , *Journal of Combinatorial Theory*, **6** (1969) 210.
- [Isb2] J.R. Isbell,  $N(5, 4; 3) \geq 24$ , *Journal of Combinatorial Theory*, Series A, **34** (1983) 379-380.
- [Ishi] Y. Ishigami, Linear Ramsey Numbers for Bounded-Degree Hypergraphs, *Electronic Notes in Discrete Mathematics*, **29** (2007) 47-51.

[-] A. Itzhakov, see [CodFIM].

## J

- [Jack] E. Jackowska, The 3-Color Ramsey Number for a 3-Uniform Loose Path of Length 3, *Australasian Journal of Combinatorics*, **63** (2015) 314-320.
- [JacPR] E. Jackowska, J. Polcyn and A. Ruciński, Multicolor Ramsey Numbers and Restricted Turán Numbers for the Loose 3-Uniform Path of Length Three, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.5, **24** (2017), 21 pages.
- [Jaco] M.S. Jacobson, On the Ramsey Number for Stars and a Complete Graph, *Ars Combinatoria*, **17** (1984) 167-172.
- [-] M.S. Jacobson, see also [BEFRSGJ, GoJa1, GoJa2].
- [-] S. Jahanbekam, see [BiFJ].
- [JamKR]\*\* D. James, E. Kahan and E. Rauer,  $R(K_6 - e, K_4) = 30$ , *Graphs and Combinatorics*, **41:27** (2025), 17 pages.
- [-] O. Janzer, see [AxCJMR].
- [JaBBJ] A.M.M. Jaradat, A. Baniabedalruhman, M.S. Bataineh and M.M.M. Jaradat, The Theta-Complete Graph Ramsey Number  $r(\theta_k, K_5)$ ;  $k=7, 8, 9$ , *Italian Journal of Pure and Applied Mathematics*, **46** (2021) 614-620.
- [-] A.M.M. Jaradat, see also [BanJBJ].
- [JaAl] M.M.M. Jaradat and B.M.N. Alzaleq, The Cycle-Complete Graph Ramsey Number  $r(C_8, K_8)$ , *SUT Journal of Mathematics*, **43** (2007) 85-98.
- [JaBa] M.M.M. Jaradat and A.M.M. Baniabedalruhman, The Cycle-Complete Graph Ramsey Number  $r(C_8, K_7)$ , *International Journal of Pure and Applied Mathematics*, **41** (2007) 667-677.
- [JaBVR] M.M.M. Jaradat, M.S. Bataineh, T. Vetric and A.M.M. Rabaiah, A Note on the Ramsey Numbers for Theta Graphs Versus the Wheel of Order 5, *AKCE International Journal of Graphs and Combinatorics*, **15** (2018) 187-189.
- [-] M.M.M. Jaradat, see also [BanJBJ, BatJA, JaBBJ].
- [JaKSY] B. Jartoux, C. Keller, S. Smorodinsky and Y. Yuditsky, On Multicolour Ramsey Numbers and Subset Coloring of Hypergraphs, *SIAM Journal on Discrete Mathematics*, **36(3)** (2022) 1848-1860.
- [-] I. Javaid, see [AliTJ].
- [JaNR]\* C.J. Jayawardene, D. Narváez and S. Radziszowski, Star-Critical Ramsey Numbers for Cycles Versus  $K_4$ , *Discussiones Mathematicae Graph Theory*, **41** (2021) 381-390.
- [JaNS] C.J. Jayawardene, W.C.W. Navaratna and J.N. Senadheera, All Ramsey Critical Graphs for Large Cycles vs a Complete Graph of Order Six, *Journal of the National Science Foundation of Sri Lanka*, **52(1)** (2024) 113-123.
- [JR1] C.J. Jayawardene and C.C. Rousseau, An Upper Bound for the Ramsey Number of a Quadrilateral Versus a Complete Graph on Seven Vertices, *Congressus Numerantium*, **130** (1998) 175-188.
- [JR2] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers  $r(C_6, G)$  for All Graphs  $G$  of Order Less than Six, *Congressus Numerantium*, **136** (1999) 147-159.
- [JR3] C.J. Jayawardene and C.C. Rousseau, The Ramsey Numbers for a Quadrilateral vs. All Graphs on Six Vertices, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **35** (2000) 71-87. Erratum in **51** (2004) 221. Erratum by L. Boza in **89** (2014) 155-156.
- [JR4] C.J. Jayawardene and C.C. Rousseau, Ramsey Numbers  $r(C_5, G)$  for All Graphs  $G$  of Order Six, *Ars Combinatoria*, **57** (2000) 163-173.
- [JR5] C.J. Jayawardene and C.C. Rousseau, The Ramsey Number for a Cycle of Length Five vs. a Complete Graph of Order Six, *Journal of Graph Theory*, **35** (2000) 99-108.

- [JRB} C. Jayawardene, C.C. Rousseau and B. Bollobás, How Ramsey Theory Can Be Used to Solve Harary's Problem for  $K_{2,k}$ , *preprint*, <http://arxiv.org/abs/1901.01552> (2019).
- [-] C.J. Jayawardene, see also [BolJY+, RoJa1, RoJa2].
- [JenSk] M. Jenssen and J. Skokan, Exact Ramsey Numbers of Odd Cycles via Nonlinear Optimisation, *Advances in Mathematics*, **376** (2021), paper no. 107444, 46 pages.
- [-] M. Jenssen, see also [DavJR, CamJMPS, CamJMS].
- [-] Meng Ji, see [HngJL].
- [-] Jiang Baoqi, see [SunYJLS].
- [JiHou] Taiping Jiang and Xinmin Hou, Ramsey Numbers of Color Critical Graphs Versus Large Generalized Fans, *preprint*, <http://arxiv.org/abs/2308.10546> (2023).
- [JiSa] Tao Jiang and M. Salerno, Ramsey Numbers of Some Bipartite Graphs Versus Complete Graphs, *Graphs and Combinatorics*, **27** (2011) 121-128.
- [-] Tao Jiang, see also [ColGJ].
- [JiLSX] Yu Jiang, Meilian Liang, Yongqi Sun and Xiaodong Xu, On Ramsey Numbers  $R(K_4 - e, K_t)$ , *Graphs and Combinatorics*, **37** (2021) 651-661.
- [JiLTX1]\* Yu Jiang, Meilian Liang, Yanmei Teng and Xiaodong Xu, The Cyclic Triangle-Free Process, *Symmetry*, **11** (2019) 955.
- [JiLTX2]\* Yu Jiang, Meilian Liang, Yanmei Teng and Xiaodong Xu, Random Cyclic Triangle-Free Graphs of Prime Order, *Journal of Mathematics*, Hindawi, volume 2021, article ID 5595919, 6 pages.
- [Jin]\*\* Jin Xia, Ramsey Numbers Involving a Triangle: Theory & Applications, *Technical Report RIT-TR-93-019*, MS thesis, Department of Computer Science, Rochester Institute of Technology, 1993.
- [-] Jin Xia, see also [RaJi].
- [-] J.R. Johnson, see [DayJ].
- [JoPe] K. Johst and Y. Person, On the Multicolor Ramsey Number of a Graph with  $m$  Edges, *Discrete Mathematics*, **339** (2016) 2857-2860.
- [Jun] V. Jungić, *Basics of Ramsey Theory*, Routledge Taylor & Francis, 2023.
- [-] Jusmawati, see [HaJu].
- [JGT] *Journal of Graph Theory*, special volume on Ramsey theory, **7**, Number 1, (1983).

## K

- [KaOS] S. Kadota, T. Onozuka and Y. Suzuki, The Graph Ramsey Number  $R(F_1, K_6)$ , *Discrete Mathematics*, **342** (2019) 1028-1037.
- [-] E. Kahan, see [JamKR].
- [Ka1] J.G. Kalbfleisch, Construction of Special Edge-Chromatic Graphs, *Canadian Mathematical Bulletin*, **8** (1965) 575-584.
- [Ka2]\* J.G. Kalbfleisch, Chromatic Graphs and Ramsey's Theorem, *Ph.D. thesis*, University of Waterloo, January 1966.
- [Ka3] J.G. Kalbfleisch, On Robillard's Bounds for Ramsey Numbers, *Canadian Mathematical Bulletin*, **14** (1971) 437-440.
- [KaSt] J.G. Kalbfleisch and R.G. Stanton, On the Maximal Triangle-Free Edge-Chromatic Graphs in Three Colors, *Journal of Combinatorial Theory*, **5** (1968) 9-20.
- [KamRa1] A. Kamranian and G. Raeisi, Ramsey Number of Disjoint Union of Good Hypergraphs, *Iranian Journal of Science and Technology, Transactions Science*, **44** (2020) 1649-1652.
- [KamRa2] A. Kamranian and G. Raeisi, On the Star-Critical Ramsey Number of a Forest Versus Complete Graphs, *Iranian Journal of Science and Technology, Transaction Science*, **46** (2022) 499-505.

- [-] Ross Kang, see [WoGKSF].
- [-] Ardak Kapbasov, see [LowKap1, LowKap2, LowKKB].
- [-] Arman Kapbasov, see [LowKKB].
- [KarK] C.D. Karamchedu and M.M. Klawe, On the Ramsey Numbers of Odd-Linked Double Stars, *Discrete Mathematics*, **345** (2022) 113001, 12 pages.
- [KáRos] G. Károlyi and V. Rosta, Generalized and Geometric Ramsey Numbers for Cycles, *Theoretical Computer Science*, **263** (2001) 87-98.
- [KeeLS] P. Keevash, E. Long and J. Skokan, Cycle-Complete Ramsey Numbers, *International Mathematics Research Notices*, rnz119 (2019) 1-26, Volume (2021), Issue 1, 275-300.
- [-] P. Keevash, see also [BohK1, BohK2, BohK3].
- [-] C. Keller, see [JaKSY].
- [KerRo] M. Kerber and C. Rowan, CommonLisp program for computing upper bounds on classical Ramsey numbers, <http://www.cs.bham.ac.uk/~mmk/demos/ramsey-upper-limit.lisp> (2009).
- [Kéry] G. Kéry, On a Theorem of Ramsey (in Hungarian), *Matematikai Lapok*, **15** (1964) 204-224.
- [KhoDz] F. Khoeini and T. Dzido, On Some Three Color Ramsey Numbers for Paths, Cycles, Stripes and Stars, *Graphs and Combinatorics*, **35** (2019) 559-567.
- [-] Jaehoon Kim, see [HasHKL].
- [Kim] Jeong Han Kim, The Ramsey Number  $R(3, t)$  Has Order of Magnitude  $t^2/\log t$ , *Random Structures and Algorithms*, **7** (1995) 173-207.
- [-] Dylan King, see [HeHKP].
- [KirS]\* M. Kirchweger and S. Szeider, SAT Modulo Symmetries for Graph Generation and Enumeration, *ACM Transactions on Computational Logic*, **25**(3) (2024), article #18, 29 pages.
- [KlaM1] K. Klamroth and I. Mengersen, Ramsey Numbers of  $K_3$  Versus  $(p, q)$ -Graphs, *Ars Combinatoria*, **43** (1996) 107-120.
- [KlaM2] K. Klamroth and I. Mengersen, The Ramsey Number of  $r(K_{1,3}, C_4, K_4)$ , *Utilitas Mathematica*, **52** (1997) 65-81.
- [-] K. Klamroth, see also [ArKM].
- [-] M. Klawe, see [GrHK, KarK].
- [-] D.J. Kleitman, see [GoK].
- [KniSu] C. Knierim and P. Su, Improved Bounds on the Multicolor Ramsey Numbers of Paths and Even Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.26, **26**(1) (2019), 17 pages.
- [KoSS1] Y. Kohayakawa, M. Simonovits and J. Skokan, The 3-colored Ramsey Number of Odd Cycles, *Electronic Notes in Discrete Mathematics*, **19** (2005) 397-402.
- [KoSS2] Y. Kohayakawa, M. Simonovits and J. Skokan, The 3-colored Ramsey Number of Odd Cycles, to appear in *Journal of Combinatorial Theory, Series B* (2013).
- [Köh] W. Köhler, On a Conjecture by Grossman, *Ars Combinatoria*, **23** (1987) 103-106.
- [-] J. Komlós, see [CsKo, AjKS, AjKSS].
- [Kol1]\* M. Kolodyazhny, Novye Nizhnie Granitsy Chisel Ramseyeya  $R(3, 12)$  i  $R(3, 13)$  (in Russian), *Matematicheskoye i Informacionnoye Modelirovanie*, Tyumen, **14** (2015) 126-130.
- [Kol2]\* M. Kolodyazhny, graphs available at <http://aluarium.net/forum/wiki-article-17.html>, *personal communication* (2016).
- [Kor] A. Korolova, Ramsey Numbers of Stars Versus Wheels of Similar Sizes, *Discrete Mathematics*, **292** (2005) 107-117.
- [KosMV1] A. Kostochka, D. Mubayi and J. Verstraëte, On Independent Sets in Hypergraphs, *Random Structures and Algorithms*, **44** (2014) 224-239.

- [KosMV2] A. Kostochka, D. Mubayi and J. Verstraëte, Hypergraph Ramsey Numbers: Triangles Versus Cliques, *Journal of Combinatorial Theory, Series A*, **120** (2013) 1491-1507.
- [KosPR] A. Kostochka, P. Pudlák and V. Rödl, Some Constructive Bounds on Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **100** (2010) 439-445.
- [KoRö1] A.V. Kostochka and V. Rödl, On Graphs with Small Ramsey Numbers, *Journal of Graph Theory*, **37** (2001) 198-204.
- [KoRö2] A.V. Kostochka and V. Rödl, On Graphs with Small Ramsey Numbers, II, *Combinatorica*, **24** (2004) 389-401.
- [KoRö3] A.V. Kostochka and V. Rödl, On Ramsey Numbers of Uniform Hypergraphs with Given Maximum Degree, *Journal of Combinatorial Theory, Series A*, **113** (2006) 1555-1564.
- [KoSu] A.V. Kostochka and B. Sudakov, On Ramsey Numbers of Sparse Graphs, *Combinatorics, Probability and Computing*, **12** (2003) 627-641.
- [-] R.L. Kramer, see [FeKR].
- [KrRod] I. Krasikov and Y. Roditty, On Some Ramsey Numbers of Unicyclic Graphs, *Bulletin of the Institute of Combinatorics and its Applications*, **33** (2001) 29-34.
- [-] S. Krause, see [HaKr1, HaKr2].
- [KrLR]\* D.L. Kreher, Wei Li and S. Radziszowski, Lower Bounds for Multicolor Ramsey Numbers from Group Orbits, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 87-95.
- [-] D.L. Kreher, see also [RaK1, RaK2, RaK3, RaK4].
- [Kriv] M. Krivelevich, Bounding Ramsey Numbers through Large Deviation Inequalities, *Random Structures and Algorithms*, **7** (1995) 145-155.
- [-] M. Krivelevich, see also [AIBK, AIKS].
- [KroMe] M. Krone and I. Mengersen, The Ramsey Numbers  $r(K_5 - 2K_2, 2K_3)$ ,  $r(K_5 - e, 2K_3)$  and  $r(K_5, 2K_3)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **81** (2012) 257-260.
- [Krü] O. Krüger, An Invariant for Minimum Triangle-Free Graphs, *Australasian Journal of Combinatorics*, **74** (2019) 371-388.
- [-] M. Kubale, see [DzKP].
- [KüCFO] D. Kühn, O. Cooley, N. Fountoulakis and D. Osthus, Ramsey Numbers of Sparse Hypergraphs, *Electronic Notes in Discrete Mathematics*, **29** (2007) 29-33.
- [-] D. Kühn, see also [CooFKO1, CooFKO2].
- [Kuz]\* E. Kuznetsov, Computational Lower Limits on Small Ramsey Numbers, *preprint*, <http://arxiv.org/abs/1505.07186> (2016).

## La - Le

- [-] Hong-Jian Lai, see [XuWLM].
- [-] P.C.B. Lam, see [ShiuLL].
- [-] Ander Lamaison, see [HngJL].
- [-] Powers Lamb, see [BaLL+].
- [-] J. Lambert, see [BudHLS].
- [LaRo] B. Landman and A. Robertson, *Ramsey Theory on the Integers*, Student Mathematical Library, American Mathematical Society, first edition **24** (2004), second edition **73** (2014).
- [LaLR]\*\* A. Lange, I. Livinsky and S.P. Radziszowski, Computation of the Ramsey Numbers  $R(C_4, K_9)$  and  $R(C_4, K_{10})$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **97** (2016) 139-154.
- [-] C. Langhoff, see [AlmBCL].

- [LauPRT] M. Lauria, P. Pudlák, V. Rödl and N. Thapen, The Complexity of Proving That a Graph Is Ramsey, *ICALP 2013, Part I, LNCS 7965* (2013) 684-695. *Combinatorica*, **37**(2) (2017) 253-268.
- [Law1] S.L. Lawrence, Cycle-Star Ramsey Numbers, *Notices of the American Mathematical Society*, **20** (1973) Abstract A-420.
- [Law2] S.L. Lawrence, Bipartite Ramsey Theory, *Notices of the American Mathematical Society*, **20** (1973) Abstract A-562.
- [-] S.L. Lawrence, see also [FLPS].
- [LayMa] C. Laywine and J.P. Mayberry, A Simple Construction Giving the Two Non-isomorphic Triangle-Free 3-Colored  $K_{16}$ 's, *Journal of Combinatorial Theory, Series B*, **45** (1988) 120-124.
- [LaMu] F. Lazebnik and D. Mubayi, New Lower Bounds for Ramsey Numbers of Graphs and Hypergraphs, *Advances in Applied Mathematics*, **28** (2002) 544-559.
- [LaWo1] F. Lazebnik and A. Woldar, New Lower Bounds on the Multicolor Ramsey Numbers  $r_k(C_4)$ , *Journal of Combinatorial Theory, Series B*, **79** (2000) 172-176.
- [LaWo2] F. Lazebnik and A. Woldar, General Properties of Some Families of Graphs Defined by Systems of Equations, *Journal of Graph Theory*, **38** (2001) 65-86.
- [Lee] Choongbum Lee, Ramsey Numbers of Degenerate Graphs, *Annals of Mathematics*, **185** (2017) 791-829.
- [-] Choongbum Lee, see also [ConFLS].
- [Lef1] H. Lefmann, A Note on Ramsey Numbers, *Studia Scientiarum Mathematicarum Hungarica*, **22** (1987) 445-446.
- [Lef2] H. Lefmann, Ramsey Numbers for Monotone Paths and Cycles, *Ars Combinatoria*, **35** (1993) 271-279.
- [-] H. Lefmann, see also [DuLR].
- [-] J. Lehel, see [BaLS, GyLSS].
- [LeMu] J. Lenz and D. Mubayi, Multicolor Ramsey Numbers for Complete Bipartite Versus Complete Graphs, *Journal of Graph Theory*, **77** (2014) 19-38.
- [Les]\* A. Lesser, Theoretical and Computational Aspects of Ramsey Theory, *Examensarbeten i Matematik*, Matematiska Institutionen, Stockholms Universitet, **3**, <http://www2.math.su.se/gemensamt/grund/exjobb/matte/2001> (2001).
- [-] D. Leven, see [BILR].

## Li

- [-] Li Bingxi, see [SunYWLX, SunYXL].
- [LiBie] Binlong Li and H. Bielak, On the Ramsey-Goodness of Paths, *Graphs and Combinatorics*, **32** (2016) 2541-2549.
- [LiNing1] Binlong Li and Bo Ning, On Path-Quasar Ramsey Numbers, *Annales Universitatis Mariae Curie-Skłodowska Lublin-Polonia, Sectio A*, **LXVIII** (2014) 11-17.
- [LiNing2] Binlong Li and Bo Ning, The Ramsey Numbers of Paths Versus Wheels: a Complete Solution, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.41, **21**(4) (2014), 30 pages.
- [LiSch] Binlong Li and I. Schiermeyer, On Star-Wheel Ramsey Numbers, *Graphs and Combinatorics*, **32** (2016) 733-739.
- [LiZBBH] Binlong Li, Yanbo Zhang, H. Bielak, H. Broersma and P. Holub, Closing the Gap on Path-Kipas Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.21, **22**(3) (2015), 7 pages.
- [LiZB] Binlong Li, Yanbo Zhang and H. Broersma, An Exact Formula for All Star-Kipas Ramsey Numbers, *Graphs and Combinatorics*, **33** (2017) 141-148.

- [LiWa1] Li Da Yong and Wang Zhi Jian, The Ramsey Number  $r(mC_4, nC_4)$  (in Chinese), *Journal of Shanghai Tiedao University*, **20** (1999) 66-70, 83.
- [LiWa2] Li Da Yong and Wang Zhi Jian, The Ramsey Numbers  $r(mC_4, nC_5)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **45** (2003) 245-252.
- [-] Dongxin Li, see [ZhuZL].
- [-] Li Guiqing, see [SLLL, SLZL].
- [LiLi]\* Li Mingbo and Li Yusheng, Ramsey Numbers and Triangle-Free Cayley Graphs (in Chinese), *Journal of Tongji University (Natural Science)*, Shanghai, **43**(11) (2015) 1750-1752.
- [-] Li Jinwen, see [ZLLS].
- [LiMSY] Ping Li, Ya-ping Mao, Ingo Schiermeyer and Yi-fan Yao, Ramsey Numbers for Linear Forest-Kipas and Its Applications in Gallai-Ramsey Numbers, *Acta Mathematicae Applicatae Sinica, English Series*, November 2025, 15 pages, doi.org/10.1007/s10255-025-0052-3.
- [LiSLW]\* Li Qiao, Su Wenlong, Luo Haipeng and Wu Kang, Lower Bounds for Some Two-Color Ramsey Numbers, *manuscript* (2011). By 2016, the results there have been improved by others.
- [-] Li Qiao, see also [SuLL, SLLL].
- [-] Ray Li, see [FoxLi].
- [LiLin1] Tia-nyu Li and Qizhong Lin, Upper Bounds on the Multicolor Ramsey Numbers  $r_k(C_4)$ , *Acta Mathematicae Applicatae Sinica, English Series*, **41**(1) (2025) 286-294.
- [LiLP] Tianyu Li, Qizhong Lin and Xing Peng, Ramsey Numbers of the Quadrilateral Versus Books, *Journal of Graph Theory*, **103**(2) (2023) 309-322.
- [-] Tianyu Li, see also [DouLLP].
- [-] Wei Li, see [KrLR].
- [LiXin] Xin Li, Two Source Extractors for Asymptotically Optimal Entropy, and (Many) More, *IEEE 64-th Annual Symposium on Foundations of Computer Science (FOCS)*, Santa Cruz, CA, USA, 2023.
- [LiLW] Yan Li, Yusheng Li and Ye Wang, Multicolor Ramsey Numbers of Bipartite Graphs and Large Books, *Graphs and Combinatorics*, **39**:21 (2023), 7 pages.
- [LiW] Yan Li and Ye Wang, Two Multicolor Ramsey Numbers Involving Bipartite Graphs, *COCOA 2023*, LNCS 14462 (2024) 147-151.
- [LiZZ] Yan Li, Yahui Zhang and Ping Zhang, Multicolor Ramsey Numbers and Star-Critical Ramsey Numbers Involving Fans, *Discussiones Mathematicae Graph Theory*, **45** (2025) 395-409.
- [-] Yan Li, see also [WaLL].
- [Li1] Li Yusheng, Some Ramsey Numbers of Graphs with Bridge, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **25** (1997) 225-229.
- [Li2] Li Yusheng, The Shannon Capacity of a Communication Channel, Graph Ramsey Number and a Conjecture of Erdős, *Chinese Science Bulletin*, **46** (2001) 2025-2028.
- [Li3] Yusheng Li, Ramsey Numbers of a Cycle, *Taiwanese Journal of Mathematics*, **12** (2008) 1007-1013.
- [Li4] Yusheng Li, The Multi-Color Ramsey Number of an Odd Cycle, *Journal of Graph Theory*, **62** (2009) 324-328.
- [LiLih] Yusheng Li and Ko-Wei Lih, Multi-Color Ramsey Numbers of Even Cycles, *European Journal of Combinatorics*, **30** (2009) 114-118.
- [LiLin2] Yusheng Li and Qizhong Lin, *Elementary Methods of Graph Ramsey Theory*, Springer, Applied Mathematical Sciences 211, 2022.
- [LiR1] Li Yusheng and C.C. Rousseau, On Book-Complete Graph Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **68** (1996) 36-44.
- [LiR2] Li Yusheng and C.C. Rousseau, Fan-Complete Graph Ramsey Numbers, *Journal of Graph Theory*, **23** (1996) 413-420.

- [LiR3] Li Yusheng and C.C. Rousseau, On the Ramsey Number  $r(H + \bar{K}_n, K_n)$ , *Discrete Mathematics*, **170** (1997) 265-267.
- [LiR4] Li Yusheng and C.C. Rousseau, A Ramsey Goodness Result for Graphs with Many Pendant Edges, *Ars Combinatoria*, **49** (1998) 315-318.
- [LiRS] Li Yusheng, C.C. Rousseau and L. Soltés, Ramsey Linear Families and Generalized Subdivided Graphs, *Discrete Mathematics*, **170** (1997) 269-275.
- [LiRZ1] Li Yusheng, C.C. Rousseau and Zang Wenan, Asymptotic Upper Bounds for Ramsey Functions, *Graphs and Combinatorics*, **17** (2001) 123-128.
- [LiRZ2] Li Yusheng, C.C. Rousseau and Zang Wenan, An Upper Bound on Ramsey Numbers, *Applied Mathematics Letters*, **17** (2004) 663-665.
- [LiShen] Yusheng Li and Jian Shen, Bounds for Ramsey Numbers of Complete Graphs Dropping an Edge, *European Journal of Combinatorics*, **29** (2008) 88-94.
- [LiTZ] Li Yusheng, Tang Xueqing and Zang Wenan, Ramsey Functions Involving  $K_{m,n}$  with  $n$  Large, *Discrete Mathematics*, **300** (2005) 120-128.
- [LiZa1] Li Yusheng and Zang Wenan, Ramsey Numbers Involving Large Dense Graphs and Bipartite Turán Numbers, *Journal of Combinatorial Theory, Series B*, **87** (2003) 280-288.
- [LiZa2] Li Yusheng and Zang Wenan, The Independence Number of Graphs with a Forbidden Cycle and Ramsey Numbers, *Journal of Combinatorial Optimization*, **7** (2003) 353-359.
- [-] Li Yusheng, see also [BaiLi, BaLX, CaLRZ, Doli, DoLL1, DoLL2, GuLi, HeLD, LiLi, LinLi1, LinLi2, LinLi3, LinLD, LinLS, LiLW, LiuLi1, LiuLi2, LiuLi3, LiuLi4, LiuLi5, LiuLi6, LiuLi7, PeiLi, PeiCLY, ShiuLL, SonLi, SunLi, WaLi, WaLL, YuLi].
- [-] Li Zhenchong, see [LuSL, LuLL].
- [-] Zhengyu Li, see [DuLBG].
- [-] Zhihui Li, see [ZhouLMW].

### Lia - Lj

- [LiaRX] Meilian Liang, S. Radziszowski and Xiaodong Xu, On a Diagonal Conjecture for Classical Ramsey Numbers, *Discrete Applied Mathematics*, **267** (2019) 195-200.
- [LiaX]\*\* Meilian Liang and Xiaodong Xu, Some New Lower Bounds for Ramsey Numbers  $R(3, k)$ , *in preparation*, (2024).
- [-] Meilian Liang, see also [JiLSX, JiLTX1, JiLTX2, LuLL, XuLL].
- [LiaWXS]\* Wenzhong Liang, Kang Wu, Xiaodong Xu and Wenlong Su, New Lower Bounds for Seven Classical Ramsey Numbers, *in preparation*, (2011).
- [LiaWXCS]\* Liang Wenzhong, Wu Kang, Xu Chengzhang, Chen Hong and Su Wenlong, Using the Two Stage Automorphism of Paley to Calculate the Lower Bound of Ramsey, *Journal of Inner Mongolia Normal University*, **41** (2012) 591-596.
- [LidP]\*\* B. Lidický and F. Pfender, Semidefinite Programming and Ramsey Numbers, *preprint*, <http://arxiv.org/abs/1704.03592> (2017), revised version (2020). *SIAM Journal on Discrete Mathematics*, **35**(4) (2021) 10.1137/18M1169473.
- [LidMPV]\* B. Lidický, G. McKinley, F. Pfender and S. Van Overberghe, Small Ramsey Numbers for Books, Wheels, and Generalizations, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.64, **32**(4) (2025), 20 pages.
- [-] A. Liebenau, see [BILi].
- [-] Ko-Wei Lih, see [LiLih].
- [LinCh] Qizhong Lin and Weiji Chen, New Upper Bound for Multicolor Ramsey Number of Odd Cycles, *Discrete Mathematics*, **342** (2019) 217-220.

- [LinLi1] Qizhong Lin and Yusheng Li, On Ramsey Numbers of Fans, *Discrete Applied Mathematics*, **157** (2009) 191-194.
- [LinLi2] Qizhong Lin and Yusheng Li, Ramsey Numbers of  $K_3$  and  $K_{n,n}$ , *Applied Mathematics Letters*, **25** (2012) 380-384.
- [LinLi3] Qizhong Lin and Yusheng Li, Quasi-Random Graphs of Given Density and Ramsey Numbers, *Pure and Applied Mathematics Quarterly*, **18**(6) (2022) 2537-2549.
- [LinLD] Qizhong Lin, Yusheng Li and Lin Dong, Ramsey Goodness and Generalized Stars, *European Journal of Combinatorics*, **31** (2010) 1228-1234.
- [LinLS]\* Qizhong Lin, Yusheng Li and Jian Shen, Lower Bounds for  $r_2(K_1+G)$  and  $r_3(K_1+G)$  from Paley Graph and Generalization, *European Journal of Combinatorics*, **40** (2014) 65-72.
- [LinLiu] Qizhong Lin and Xiudi Liu, Ramsey Numbers Involving Large Books, *SIAM Journal on Discrete Mathematics*, **35** (2021) 23-34.
- [LinP] Qizhong Lin and Xing Peng, Large Book-Cycle Ramsey Numbers, *SIAM Journal on Discrete Mathematics*, **35** (2021) 532-545.
- [LinS] Qizhong Lin and Shixi Song, Ramsey Numbers of Long Even Cycles Versus Books, *preprint*, <http://arxiv.org/abs/2509.26323> (2025).
- [-] Qizhong Lin, see also [ChenL, ChenLY, ChuLin, DoLL1, DoLL2, DouLLP, FanHL, FanLin1, FanLin2, HaoLin, HuLin, HuL+, LiLin1, LiLP, LiLin2, YouLin1, YouLin2, YouLC].
- [-] Lin Xiaohui, see [SunYJLS, SunYLZ1, SunYLZ2].
- [LinCa]\* M. Lindsay and J.W. Cain, Improved Lower Bounds on the Classical Ramsey Numbers  $R(4, 22)$  and  $R(4, 25)$ , *preprint*, <http://arxiv.org/abs/1510.06102> (2015).
- [Lind] B. Lindström, Undecided Ramsey-Numbers for Paths, *Discrete Mathematics*, **43** (1983) 111-112.
- [Ling] A.C.H. Ling, Some Applications of Combinatorial Designs to Extremal Graph Theory, *Ars Combinatoria*, **67** (2003) 221-229.
- [-] Andy Liu, see [AbbL, BaLiu].
- [LiuMS] Henry Liu, Bojan Mohar and Yongtang Shi, Multicoloured Ramsey Numbers of the Path of Length Four, *preprint*, <http://arxiv.org/abs/2108.06477> (2021).
- [-] Hong Liu, see [AxGLM, HasHKL].
- [-] Liu Linzhong, see [ZLLS].
- [LiuDu] Meng Liu and Bangwei Du, Ramsey Numbers of Complete Bipartite Graphs, *Graphs and Combinatorics*, **41**:30 (2025), 4 pages.
- [LiuLi1] Meng Liu and Yusheng Li, Ramsey Numbers of a Fixed Odd-Cycle and Generalized Books and Fans, *Discrete Mathematics*, **339** (2016) 2481-2489.
- [LiuLi2] Meng Liu and Yusheng Li, Ramsey Numbers and Bipartite Ramsey Numbers via Quasi-Random Graphs, *Discrete Mathematics*, **344** (2021), 112162, 5 pages.
- [LiuLi3] Meng Liu and Yusheng Li, Ramsey Numbers of Fans and Large Books, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.52, **29**(1) (2022), 9 pages.
- [LiuLi4] Meng Liu and Yusheng Li, Ramsey Numbers Involving an Odd Cycle and Large Complete Graphs in Three Colors, *Graphs and Combinatorics*, **38**:182 (2022), 6 pages.
- [LiuLi5] Meng Liu and Yusheng Li, A Note on Ramsey Numbers Involving Large Books, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.7, **31**(1) (2024), 6 pages.
- [LiuLi6] Meng Liu and Yusheng Li, On Graphs for Which Large Books Are Ramsey Good, *Journal of Graph Theory*, **108**(3) (2025) 543-559.
- [LiuLi7] Meng Liu and Yusheng Li, Ramsey Goodness of Large Books Revisited, *Discrete Mathematics*, **349** (2026), 114937, 7 pages.
- [LiuW] Meng Liu and Ye Wang, Polynomial Resultants and Ramsey Numbers of a Theta Graph, *Advances in Applied Mathematics*, **167** (2025) 102881, 9 pages.

- [-] Liu Shu Yan, see [SonBL].
- [Liu]\* Sixue Cliff Liu, Lower Bounds for Small Ramsey Numbers on Hypergraphs, in *Proceedings of COCOON 2019*, Xi'an, China, LNCS 11653, Springer, (2019) 412-424.
- [LiuCh] Siyue Liu and Gang Chen, On the Ramsey Numbers of Even-Linked Double Stars, *Australasian Journal of Combinatorics*, **94**(3) (2026) 385-402.
- [-] Liu Xiangyang, see [GuSL].
- [-] Xiudi Liu, see [LinLiu].
- [-] Liu Yanwu, see [SonYL].
- [-] Zhiguo Liu, see [WuSL].
- [-] I. Livinsky, see [LaLR].
- [-] I. Ljubić, see [FuLS].

### Lo - Lu

- [LoPfe] Allan Lo and Vincent Pfenninger, The Ramsey Number for 4-Uniform Tight Cycles, *Extended Abstracts EuroComb 2021*, Springer, TM 14, (2021) 442-445. Full paper in *SIAM Journal on Discrete Mathematics*, **39**(1) (2025) 361-387.
- [-] Allan Lo, see also [BoLo].
- [Loc] S.C. Locke, Bipartite Density and the Independence Ratio, *Journal of Graph Theory*, **10** (1986) 47-53.
- [-] S.C. Locke, see also [FrLo].
- [-] E. Long, see [KeeLS].
- [-] Albert López Vidal, see [AttLM].
- [Lor] P.J. Lorimer, The Ramsey Numbers for Stripes and One Complete Graph, *Journal of Graph Theory*, **8** (1984) 177-184.
- [LorMu] P.J. Lorimer and P.R. Mullins, Ramsey Numbers for Quadrangles and Triangles, *Journal of Combinatorial Theory, Series B*, **23** (1977) 262-265.
- [LorSe] P.J. Lorimer and R.J. Segegin, Ramsey Numbers for Multiple Copies of Complete Graphs, *Journal of Graph Theory*, **2** (1978) 89-91.
- [LorSo] P.J. Lorimer and W. Solomon, The Ramsey Numbers for Stripes and Complete Graphs 1, *Discrete Mathematics*, **104** (1992) 91-97. Corrigendum in **131** (1994) 395.
- [-] P.J. Lorimer, see also [CocL1, CocL2].
- [Lortz] R. Lortz, A Note on the Ramsey Number of  $K_{2,2}$  Versus  $K_{3,n}$ , *Discrete Mathematics*, **306** (2006) 2976-2982.
- [LoM1] R. Lortz and I. Mengersen, On the Ramsey Numbers  $r(K_{2,n-1}, K_{2,n})$  and  $r(K_{2,n}, K_{2,n})$ , *Utilitas Mathematica*, **61** (2002) 87-95.
- [LoM2] R. Lortz and I. Mengersen, Bounds on Ramsey Numbers of Certain Complete Bipartite Graphs, *Results in Mathematics*, **41** (2002) 140-149.
- [LoM3]\* R. Lortz and I. Mengersen, Off-Diagonal and Asymptotic Results on the Ramsey Number  $r(K_{2,m}, K_{2,n})$ , *Journal of Graph Theory*, **43** (2003) 252-268.
- [LoM4]\* R. Lortz and I. Mengersen, Further Ramsey Numbers for Small Complete Bipartite Graphs, *Ars Combinatoria*, **79** (2006) 195-203.
- [LoM5] R. Lortz and I. Mengersen, Ramsey Numbers for Small Graphs Versus Small Disconnected Graphs, *Australasian Journal of Combinatorics*, **51** (2011) 89-108.
- [LoM6] R. Lortz and I. Mengersen, On the Ramsey Numbers of Certain Graphs of Order Five Versus All Connected Graphs of Order Six, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **90** (2014) 197-222.

- [LoM7] R. Lortz and I. Mengersen, On the Ramsey Numbers for Stars Versus Connected Graphs of Order Six, *Australasian Journal of Combinatorics*, **73** (2019) 1-24.
- [LoM8] R. Lortz and I. Mengersen, All Missing Ramsey Numbers for Trees Versus the Four-Page Book, *Electronic Journal of Graph Theory and Applications*, **9**(2) (2021) 377-385.
- [LoM9] R. Lortz and I. Mengersen, On the Ramsey Numbers of Non-Star Trees Versus Connected Graphs of Order Six, *Discussiones Mathematicae Graph Theory*, **43** (2023) 331-349.
- [LoM10] R. Lortz and I. Mengersen, On the Ramsey Number  $r(S_n, K_6 - 3K_2)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **113** (2020) 119-126, and **124** (2025) 229-234.
- [-] Andrew Lott, see [BaLL+].
- [LowKap1]\* R. Low and A. Kapbasov, New Diagonal Graph Ramsey Numbers of Unicyclic Graphs, *Theory and Applications of Graphs*, **10**(1) (2023), article 9, 12 pages.
- [LowKap2]\* R. Low and A. Kapbasov, Diagonal Graph Ramsey Numbers of Even Cycles with Pendant Edges, *Australasian Journal of Combinatorics*, **94**(1) (2026) 221-234.
- [LowKKB]\* R. Low, A. Kapbasov, A. Kapbasov and S. Bereg, Computation of New Diagonal Graph Ramsey Numbers, *Electronic Journal of Graph Theory and Applications*, **10**(2) (2022) 575-588.
- [Loz] V. Lozin, Ramsey Numbers and Graph Parameters, *Graphs and Combinatorics*, **40**:29 (2024), 13 pp.
- [-] V. Lozin, see also [AleALZ, AtLZ].
- [Łuc] T. Łuczak,  $R(C_n, C_n, C_n) \leq (4 + o(1))n$ , *Journal of Combinatorial Theory, Series B*, **75** (1999) 174-187.
- [ŁuPo1] T. Łuczak and J. Polcyn, On the Multicolor Ramsey Number for 3-Paths of Length Three, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.27, **24** (2017), 4 pages.
- [ŁuPo2] T. Łuczak and J. Polcyn, The Multipartite Ramsey Number for the 3-Path of Length Three, *Discrete Mathematics*, **341** (2018) 1270-1274.
- [ŁuPR] T. Łuczak, J. Polcyn and A. Ruciński, On Multicolor Ramsey Numbers for Loose  $k$ -Paths of Length Three, *European Journal of Combinatorics*, **71** (2018) 43-50.
- [ŁucSS] T. Łuczak, M. Simonovits and J. Skokan, On the Multi-Colored Ramsey Numbers of Cycles, *Journal of Graph Theory*, **69** (2012) 169-175.
- [-] T. Łuczak, see also [AllŁPZ, FiŁu1, FiŁu2, HaŁP1+, HaŁP2+, HaŁT, HuL+].
- [LuSL]\* Luo Haipeng, Su Wenlong and Li Zhenchong, The Properties of Self-Complementary Graphs and New Lower Bounds for Diagonal Ramsey Numbers, *Australasian Journal of Combinatorics*, **25** (2002) 103-116.
- [LuSS1]\* Luo Haipeng, Su Wenlong and Shen Yun-Qiu, New Lower Bounds of Ten Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **24** (2001) 81-90.
- [LuSS2]\* Luo Haipeng, Su Wenlong and Shen Yun-Qiu, New Lower Bounds for Two Multicolor Classical Ramsey Numbers, *Radovi Matematički*, **13** (2004) 15-21, pointed to in past revisions. Since 2015, better bounds were obtained by others.
- [-] Luo Haipeng, see also [LiSLW, SuL, SuLL, SLLL, SLZL, WSLX1, WSLX2, XuLL].
- [LuLL]\* Liang Luo, Meilian Liang and Zhenchong Li, Computation of Ramsey Numbers  $R(C_m, W_n)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **81** (2012) 145-149.
- [LuoP1] Zhidan Luo and Yuejian Peng, Ramsey Numbers of Multiple Copies of Stars, *Discrete Mathematics*, **345** (2022) 112801, 8 pages.
- [LuoP2] Zhidan Luo and Yuejian Peng, A Large Tree Is  $tK_m$ -Good, *Discrete Mathematics*, **346** (2023) 113502, 5 pages.
- [-] Zhidan Luo, see also [CheLC, HuLuo].

## M

- [MaSX] Jie Ma, Wujie Shen and Shengjie Xie, An Exponential Improvement for Ramsey Lower Bounds, *preprint*, <http://arxiv.org/abs/2507.12926> (2025).
- [Mac]\* J. Mackey, Combinatorial Remedies, *Ph.D. thesis*, Department of Mathematics, University of Hawaii, 1994.
- [-] W. Macready, see [RanMCG].
- [Mad] P. Madarasi, The Ramsey Number of a Long Cycle and Complete Graphs, *preprint*, <http://arxiv.org/abs/2003.12691> (2020).
- [-] C. Magnant, see [MaoWMS].
- [MaOm1] L. Maherani and G.R. Omid, Around a Conjecture of Erdős on Graph Ramsey Numbers, *preprint*, <http://arxiv.org/abs/1211.6287> (2012).
- [MaOm2] L. Maherani and G.R. Omid, Monochromatic Hamiltonian Berge-Cycles in Colored Hypergraphs, *Discrete Mathematics*, **340** (2017) 2043-2052.
- [MaORS1] L. Maherani, G.R. Omid, G. Raeisi and M. Shahsiah, The Ramsey Number of Loose Paths in 3-Uniform Hypergraphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P12, **20**(1) (2013), 8 pages.
- [MaORS2] L. Maherani, G.R. Omid, G. Raeisi and M. Shahsiah, On Three-Color Ramsey Number of Paths, *Graphs and Combinatorics*, **31** (2015) 2299-2308.
- [MahS] L. Maherani and M. Shahsiah, On Ramsey Numbers of 3-Uniform Berge Cycles, *Discrete Mathematics*, **347** (2024) 113877, 10 pages.
- [-] H. Mahon, see [DeMST].
- [-] H.R. Maimani, see [HagMa].
- [-] Kaushik Majumder, see [GupM].
- [MaoWMS] Yaping Mao, Zhao Wang, Colton Magnant and Ingo Schiermeyer, Ramsey and Gallai-Ramsey Number for Wheels, *Graphs and Combinatorics*, **38**:42 (2022), 17 pages.
- [-] Yaping Mao, see also [LiMSY, XuWLM, YaoHMZ, ZhouLMW].
- [Math]\* R. Matheron, Lower Bounds for Ramsey Numbers and Association Schemes, *Journal of Combinatorial Theory*, Series B, **42** (1987) 122-127.
- [MatMNV] Sam Mattheus, Dhruv Mubayi, Jiayi Nie and Jacques Verstraëte, Off-Diagonal Ramsey Numbers for Slowly Growing Hypergraphs, *Random Structures and Algorithms*, **66**:e21284 (2025), 8 pages.
- [MatVer] S. Mattheus and J. Verstraëte, The Asymptotics of  $r(4, t)$ , *Annals of Mathematics*, **199**:2 (2024) 919-949.
- [-] S. Mattheus, see also [ConMMV].
- [-] J.P. Mayberry, see [LayMa].
- [McCM]\* D. McCarthy and C. Monico, A Matheron-Type Construction for Digraphs and Improved Lower Bounds for Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P2.42, **32**(2) (2025), 10 pages.
- [-] D. McCarthy, see also [DawMc].
- [McDS] C. McDiarmid and A. Steger, Tidier Examples for Lower Bounds on Diagonal Ramsey Numbers, *Journal of Combinatorial Theory*, Series A, **74** (1996) 147-152.
- [McK1]\*\* B.D. McKay, Australian National University, *personal communication* (2003+). Graphs available at <http://cs.anu.edu.au/people/bdm/data/ramsey.html>.
- [McK2]\*\* B.D. McKay, A Class of Ramsey-Extremal Hypergraphs, *Transactions on Combinatorics*, **6**(3) (2017) 37-43.
- [McK3]\*\* B.D. McKay, Australian National University, *personal communication* (2016).

- [MPR]\*\* B.D. McKay, K. Piwakowski and S.P. Radziszowski, Ramsey Numbers for Triangles Versus Almost-Complete Graphs, *Ars Combinatoria*, **73** (2004) 205-214.
- [MR1]\*\* B.D. McKay and S.P. Radziszowski, The First Classical Ramsey Number for Hypergraphs is Computed, *Proceedings of the Second Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA'91, San Francisco, (1991) 304-308.
- [MR2]\* B.D. McKay and S.P. Radziszowski, A New Upper Bound for the Ramsey Number  $R(5, 5)$ , *Australian Journal of Combinatorics*, **5** (1992) 13-20.
- [MR3]\*\* B.D. McKay and S.P. Radziszowski, Linear Programming in Some Ramsey Problems, *Journal of Combinatorial Theory, Series B*, **61** (1994) 125-132.
- [MR4]\*\* B.D. McKay and S.P. Radziszowski,  $R(4, 5) = 25$ , *Journal of Graph Theory*, **19** (1995) 309-322.
- [MR5]\*\* B.D. McKay and S.P. Radziszowski, Subgraph Counting Identities and Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **69** (1997) 193-209.
- [McZ]\*\* B.D. McKay and Zhang Ke Min, The Value of the Ramsey Number  $R(3, 8)$ , *Journal of Graph Theory*, **16** (1992) 99-105.
- [-] B.D. McKay, see also [AnM1, AnM2, AnM3, AnM4, FM].
- [-] G. McKinley, see [LidMPV].
- [McN]\*\* J. McNamara, SUNY Brockport, *personal communication* (1995).
- [McR]\*\* J. McNamara and S.P. Radziszowski, The Ramsey Numbers  $R(K_4 - e, K_6 - e)$  and  $R(K_4 - e, K_7 - e)$ , *Congressus Numerantium*, **81** (1991) 89-96.
- [-] T. Meek, see [BudHMP].
- [-] H. Mélot, see [BrCGM].
- [MengZZ] Yanjun Meng, Yanbo Zhang and Yunqing Zhang, The Ramsey Number of Wheels Versus a Fan of Order Five, *Ars Combinatoria*, **137** (2018) 365-372.
- [MeO] I. Mengersen and J. Oeckermann, Matching-Star Ramsey Sets, *Discrete Applied Mathematics*, **95** (1999) 417-424.
- [-] I. Mengersen, see also [ArKM, CIEHMS, EHM1, EHM2, HoMe, HaMe1, HaMe2, HaMe3, HaMe4, KlaM1, KlaM2, KroMe, LoM1, LoM2, LoM3, LoM4, LoM5, LoM6, LoM7, LoM8, LoM9, LoM10].
- [-] D. Mergoni Cecchelli, see [AllMRS].
- [Mér] A. Méroueh, The Ramsey Number of Loose Cycles Versus Cliques, *Journal of Graph Theory*, **90** (2019) 172-188.
- [-] A. Methuku, see [GerMOV].
- [-] H. Metrebian, see [DvoMe].
- [-] Zhengke Miao, see [ChenCMN].
- [-] L. Michel, see [AxCJMR].
- [-] M. Michelen, see [CamJMPS, CamJMS].
- [-] A. Miller, see [CodFIM].
- [-] M. Miller, see [BaSNM].
- [-] A. Milojević, see [HunMS].
- [MiPal] Tapas Kumar Mishra and Sudebkumar Prasant Pal, Lower Bounds for Ramsey Numbers for Complete Bipartite and 3-Uniform Tripartite Subgraphs, WALCOM 2013, LNCS 7748, Springer, Berlin (2013) 257-264. *Journal of Graph Algorithms and Applications*, **17**(6) (2013) 671-688.
- [MiSa] H. Mizuno and I. Sato, Ramsey Numbers for Unions of Some Cycles, *Discrete Mathematics*, **69** (1988) 283-294.
- [-] B. Mohar, see [LiuMS].
- [-] D.B. Mokeev, see [AbdMT].

- [-] C. Monico, see [McCM].
- [MoCa] E.L. Monte Carmelo, Configurations in Projective Planes and Quadrilateral-Star Ramsey Numbers, *Discrete Mathematics*, **308** (2008) 3986-3991.
- [-] E.L. Monte Carmelo, see also [GoMC].
- [MonCR] L.P. Montejano, J. Chappelon and J.L. Ramirez Alfonsin, Ramsey for Complete Graphs with a Dropped Edge or a Triangle, *Electronic Notes in Discrete Mathematics*, **62** (2017) 21-25.
- [-] L.P. Montejano, see also [ChaMR].
- [MonPY1] R. Montgomery, M. Pavez-Signé and Jun Yan, Ramsey Numbers of Bounded Degree Trees Versus General Graphs, *Journal of Combinatorial Theory, Series B*, **173** (2025) 102-145.
- [MonPY2] R. Montgomery, M. Pavez-Signé and Jun Yan, Ramsey Numbers of Trees, *preprint*, <http://arxiv.org/abs/2509.07934> (2025), 59 pages.
- [-] P. Morawski, see [CamFMPP].
- [-] L. Moreira, see [BotMdS].
- [-] P. Morris, see [AttLM].
- [Mor] R. Morris, Some Recent Results in Ramsey Theory, *preprint*, <http://arxiv.org/abs/2601.05221> (2026).
- [-] R. Morris, see also [BalB+, BolM, CamGMS, FizGM, FizGMSS, GrMFSS].
- [MoSST] G.O. Mota, G.N. Sárközy, M. Schacht and A. Taraz, Ramsey Numbers for Bipartite Graphs with Small Bandwidth, *European Journal of Combinatorics*, **48** (2015) 165-176.
- [Mub1] D. Mubayi, Improved Bounds for the Ramsey Number of Tight Cycles Versus Cliques, *Combinatorics, Probability and Computing*, **25** (2016) 791-796.
- [Mub2] D. Mubayi, Variants of the Erdős-Szekeres and Erdős-Hajnal Ramsey Problems, *European Journal of Combinatorics*, **62** (2017) 197-205.
- [MuR] D. Mubayi and V. Rödl, Hypergraph Ramsey Numbers: Tight Cycles Versus Cliques, *Bulletin of the London Mathematical Society*, **48** (2016) 127-134.
- [MuSp] D. Mubayi and N. Spanier,  $K_4^-$ -free Triple Systems Without Large Stars in the Complement, *preprint*, <http://arxiv.org/abs/2504.06076> (2025).
- [MuSuk1] D. Mubayi and A. Suk, Off-Diagonal Hypergraph Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **125** (2017) 168-177.
- [MuSuk2] D. Mubayi and A. Suk, Constructions in Ramsey Theory, *Journal of the London Mathematical Society*, **97** (2018) 247-257.
- [MuSuk3] D. Mubayi and A. Suk, New Lower Bounds for Hypergraph Ramsey Numbers, *Bulletin of the London Mathematical Society*, **50** (2018) 189-201.
- [MuSuk4] D. Mubayi and A. Suk, A Survey of Hypergraph Ramsey Problems, in *Discrete Mathematics and Applications*, Springer Optimization and Its Applications, Cham, vol. 165 (2020).
- [MuSuk5] D. Mubayi and A. Suk, Ramsey Numbers of Cliques Versus Monotone Paths, *European Journal of Combinatorics*, **118** (2024) 103922, 7 pages.
- [MuVer] D. Mubayi and J. Verstraëte, A Note on Pseudorandom Ramsey Graphs, *Journal of the European Mathematical Society*, **26**(1) (2024) 153-161.
- [-] D. Mubayi, see also [AxFM, AxGLM, ConFG1+, ConFG2+, ConFH+, ConMMV, KosMV1, KosMV2, LaMu, LeMu, MatMNV].
- [-] P.R. Mullins, see [LorMu].
- [-] S. Musdalifah, see [SuAM, SuAAM].

## N

- [-] S.M. Nababan, see [BaSNM].

- [NaRT]\*\* Ansh Nagda, Prabhakar Raghavan and Abhradeep Thakurta, Reinforced Generation of Combinatorial Structures: Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2603.09172> (2026).
- [NaORS] B. Nagle, S. Olsen, V. Rödl and M. Schacht, On the Ramsey Number of Sparse 3-Graphs, *Graphs and Combinatorics*, **24** (2008) 205-228.
- [Nar]\* D. Narváez, Some Multicolor Ramsey Numbers Involving Cycles, *MS thesis*, Department of Computer Science, Rochester Institute of Technology, 2015.
- [NaSZ]\* David E. Narváez, Cruise Song and Ningxin Zhang, Formalizing Finite Ramsey Theory in Lean 4, *Proceedings of the 17th International Conference on Intelligent Computer Mathematics*, CICM 2024, Montréal, QC, LNAI 14960, (2024) 91-108.
- [-] D. Narváez, see also [JaNR].
- [-] W.C.W. Navaratna, see [JaNS].
- [-] Ndiamé Ndiaye, see [GuNNW].
- [Neš] J. Nešetřil, Ramsey Theory, chapter 25 in *Handbook of Combinatorics*, ed. R.L. Graham, M. Grötschel and L. Lovász, The MIT-Press, Vol. II, 1996, 1331-1403.
- [NeOs] J. Nešetřil and P. Ossona de Mendez, Fraternal Augmentations, Arrangeability and Linear Ramsey Numbers, *European Journal of Combinatorics*, **30** (2009) 1696-1703.
- [NeRo] J. Nešetřil and M. Rosenfeld, I. Schur, C.E. Shannon and Ramsey Numbers, A Short Story, *Discrete Mathematics*, **229** (2001) 185-195.
- [-] J. Nešetřil, see also [GrNe].
- [-] C.T. Ng, see [ChenCMN, ChenCNZ, CheCZN].
- [-] T. Nguyen, see [BroNN].
- [Nie] Jiayi Nie, On Tight Tree-Complete Hypergraph Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2412.19461> (2024).
- [NieVer] Jiayi Nie and Jacques Verstraëte, Ramsey Numbers for Nontrivial Berge Cycles, *SIAM Journal on Discrete Mathematics*, **36**(1) (2022) 10.1137/21M1396770.
- [-] Jiayi Nie, see also [HeNWY, MatMNV].
- [Nik] V. Nikiforov, The Cycle-Complete Graph Ramsey Numbers, *Combinatorics, Probability and Computing*, **14** (2005) 349-370.
- [NiRo1] V. Nikiforov and C.C. Rousseau, Large Generalized Books Are  $p$ -Good, *Journal of Combinatorial Theory, Series B*, **92** (2004) 85-97.
- [NiRo2] V. Nikiforov and C.C. Rousseau, Book Ramsey Numbers I, *Random Structures and Algorithms*, **27** (2005) 379-400.
- [NiRo3] V. Nikiforov and C.C. Rousseau, A Note on Ramsey Numbers for Books, *Journal of Graph Theory*, **49** (2005) 168-176.
- [NiRo4] V. Nikiforov and C.C. Rousseau, Ramsey Goodness and Beyond, *Combinatorica*, **29** (2009) 227-262.
- [NiRS] V. Nikiforov, C.C. Rousseau and R.H. Schelp, Book Ramsey Numbers and Quasi-Randomness, *Combinatorics, Probability and Computing*, **14** (2005) 851-860.
- [-] Bo Ning, see [HuL+, LiNing1, LiNing2].
- [NoSZ] S. Norin, Yue Ru Sun and Yi Zhao, Asymptotics of Ramsey Numbers of Double Stars, *preprint*, <http://arxiv.org/abs/1605.03612> (2016).
- [-] S. Norin, see also [GuNNW].
- [NoBa]\* E. Noviani and E.T. Baskoro, On the Ramsey Number of 4-Cycle Versus Wheel, *Indonesian Journal of Combinatorics*, **1** (2016) 9-21.
- [-] A. Nowik, see [DzNS].
- [-] E. Nystrom, see [BroNN].

## O

- [-] J. Oeckermann, see [MeO].
- [-] F. Oktariani, see [SherBSO, SherSBO].
- [-] S. Olsen, see [NaORS].
- [OmRa1] G.R. Omid and G. Raeisi, On Multicolor Ramsey Number of Paths Versus Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P24, **18**(1) (2011), 16 pages.
- [OmRa2] G.R. Omid and G. Raeisi, A Note on the Ramsey Number of Stars - Complete Graphs, *European Journal of Combinatorics*, **32** (2011) 598-599.
- [OmRa3] G.R. Omid and G. Raeisi, Ramsey Numbers for Multiple Copies of Hypergraphs, *Graphs and Combinatorics*, **40**:133 (2024), 12 pages.
- [OmRR] G.R. Omid, G. Raeisi and Z. Rahimi, Star Versus Stripes Ramsey Numbers, *European Journal of Combinatorics*, **67** (2018) 268-274.
- [OmSh1] G.R. Omid and M. Shahsiah, Ramsey Numbers of 3-Uniform Loose Paths and Loose Cycles, *Journal of Combinatorial Theory, Series A*, **121** (2014) 64-73.
- [OmSh2] G.R. Omid and M. Shahsiah, Diagonal Ramsey Numbers of Loose Cycles in Uniform Hypergraphs, *SIAM Journal on Discrete Mathematics*, **31** (2017) 1634-1669.
- [OmSh3] G.R. Omid and M. Shahsiah, Ramsey Numbers of 4-Uniform Loose Cycles, *Discrete Applied Mathematics*, **230** (2017) 112-120.
- [OmSh4] G.R. Omid and M. Shahsiah, Ramsey Numbers of Uniform Loose Paths and Cycles, *Discrete Mathematics*, **340** (2017) 1426-1434.
- [-] G.R. Omid, see also [GerMOV, MaOm1, MaOm2, MaORS1, MaORS2].
- [-] T. Onozuka, see [KaOS].
- [-] E. Oren-Dahan, see [BalDO].
- [-] P. Ossona de Mendez, see [NeOs].
- [-] D. Osthus, see [CooFKO1, CooFKO2, KüCFO].
- [Osu] M. Osumi, Ramsey Numbers of Trails, *IEICE Transactions on Fundamentals of Electronics, Communications and Computers*, **E105-A**(9) (2022) 1235-1240.

## P

- [-] J. Pach, see [FoxPS1, FoxPS2].
- [-] S.P. Pal, see [MiPal].
- [Pálv] D. Pálvölgyi, Exponential Lower Bound for Berge-Ramsey Problems, *Graphs and Combinatorics*, **37** (2021) 1433-1435.
- [-] Linqiang Pan, see [ShaXBP, ShaXSP].
- [Par1] T.D. Parsons, The Ramsey Numbers  $r(P_m, K_n)$ , *Discrete Mathematics*, **6** (1973) 159-162.
- [Par2] T.D. Parsons, Path-Star Ramsey Numbers, *Journal of Combinatorial Theory, Series B*, **17** (1974) 51-58.
- [Par3] T.D. Parsons, Ramsey Graphs and Block Designs, I, *Transactions of the American Mathematical Society*, **209** (1975) 33-44.
- [Par4] T.D. Parsons, Ramsey Graphs and Block Designs, *Journal of Combinatorial Theory, Series A*, **20** (1976) 12-19.
- [Par5] T.D. Parsons, Graphs from Projective Planes, *Aequationes Mathematicae*, **14** (1976) 167-189.
- [Par6] T.D. Parsons, Ramsey Graph Theory, in *Selected Topics in Graph Theory*, (L.W. Beineke and R.J. Wilson eds.), Academic Press, (1978) 361-384.

- [-] T.D. Parsons, see also [FLPS].
- [-] Dhruv Patel, see [BaLL+].
- [Pau]\* Lawrence C. Paulson, Formalising New Mathematics in Isabelle: Diagonal Ramsey, *16th International Conference on Interactive Theorem Proving, ITP 2025*, LIPICS Dagstuhl Publishing, article #18, 18 pages.
- [PavPS] M. Pavez-Signé, S. Piga and N. Sanhueza-Matamala, Ramsey Numbers with Prescribed Rate of Growth, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.24, **30**(3) (2023), 11 pages.
- [-] M. Pavez-Signé, see also [AllBP, MonPY1, MonPY2].
- [PeiLi] Chaoping Pei and Yusheng Li, Ramsey Numbers Involving a Long Path, *Discrete Mathematics*, **339** (2016) 564-570.
- [PeiCLY] Chaoping Pei, Ming Chen, Yusheng Li and Pei Yu, Ramsey Good Graphs with Long Suspended Paths, *Graphs and Combinatorics*, **34** (2018) 759-767.
- [-] T. Pellerin, see [AgCP+].
- [-] Fei Peng, see [BohP].
- [Peng] Xing Peng, The Ramsey Number of Generalized Loose Paths in Hypergraphs, *Discrete Mathematics*, **339** (2016) 539-546.
- [-] Xing Peng, see also [DouLLP, HuL+, LiLP, LinP].
- [-] Yuejian Peng, see [GuoHP, HaLP1+, HaLP2+, HuPZ, HuP1, HuP2, LuoP1, LuoP2, YanP].
- [-] A. Penland, see [BudHMP, BudHP, BudPe].
- [-] Y. Person, see [JoPe].
- [-] K. Petrova, see [CamFMPP].
- [-] F. Pfender, see [CamJMPS, HeHKP, LidP, LidMPV].
- [Pfe] V. Pfenninger, On  $k$ -uniform Tight Cycles: the Ramsey Number for  $C_{kn}^{(k)}$  and an Approximate Lehel's Conjecture, *Combinatorica*, **45** (2025), article #47, 22 pages.
- [-] V. Pfenninger, see also [LoPfe].
- [-] S. Piga, see [PavPS].
- [-] O. Pikhurko, see [BePi].
- [Piw1]\* K. Piwakowski, Applying Tabu Search to Determine New Ramsey Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R6, **3**(1) (1996), 4 pages. The lower bounds presented in this paper have been improved.
- [Piw2]\*\* K. Piwakowski, A New Upper Bound for  $R_3(K_4 - e)$ , *Congressus Numerantium*, **128** (1997) 135-141.
- [PR1]\*\* K. Piwakowski and S.P. Radziszowski,  $30 \leq R(3, 3, 4) \leq 31$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **27** (1998) 135-141.
- [PR2]\*\* K. Piwakowski and S.P. Radziszowski, Towards the Exact Value of the Ramsey Number  $R(3, 3, 4)$ , *Congressus Numerantium*, **148** (2001) 161-167.
- [-] K. Piwakowski, see also [MPR, DzKP].
- [-] C. Pohoata, see [CamP].
- [Pokr] A. Pokrovskiy, Calculating Ramsey Numbers by Partitioning Coloured Graphs, *Journal of Graph Theory*, **84** (2017) 477-500.
- [PoSu1] A. Pokrovskiy and B. Sudakov, Ramsey Goodness of Paths, *Journal of Combinatorial Theory, Series B*, **122** (2017) 384-390.
- [PoSu2] A. Pokrovskiy and B. Sudakov, Ramsey Goodness of Cycles, *SIAM Journal on Discrete Mathematics*, **34** (2020) 1884-1908.
- [-] A. Pokrovskiy, see also [BalPS, CamFMPP].

- [Pol] J. Polcyn, One More Turán Number and Ramsey Number for the Loose 3-Uniform Path of Length Three, *Discussiones Mathematicae Graph Theory*, **37** (2017) 443-464.
- [PoRRS] J. Polcyn, V. Rödl, A. Ruciński and E. Szemerédi, Short Paths in Quasi-Random Triple Systems with Sparse Underlying Graphs, *Journal of Combinatorial Theory, Series B*, **96** (2006) 584-607.
- [PoRu] J. Polcyn and A. Ruciński, Refined Turán Numbers and Ramsey Numbers for the Loose 3-Uniform Path of Length Three, *Discrete Mathematics*, **340** (2017) 107-118.
- [-] J. Polcyn, see also [AllŁPZ, HanPR, JacPR, ŁuPo1, ŁuPo2, ŁuPR].
- [-] A.D. Polimeni, see [ChaGP, ChaRSPS].
- [-] Y. Portella, see [AgCP+].
- [-] J.R. Portillo, see [BoPo].
- [-] R. Prange, see [BudPra].
- [-] H. Privette, see [BudPri].
- [-] L.M. Pretorius, see [SwPr].
- [PuRS] P. Pudlák, V. Rödl and M. Sales, On the Ramsey Numbers of Daisies I, *Combinatorics, Probability and Computing*, **33** (2024) 795-806.
- [PuRW] P. Pudlák, V. Rödl and W. Wesley, A Lower Bound on the Ramsey Number  $R_k(k+1, k+1)$ , *preprint*, <http://arxiv.org/abs/2412.16637> (2024).
- [-] P. Pudlák, see also [AlPu, CoPR, KosPR, LauPRT].

## Q

- [-] Jianguo Qian, see [WaQi].
- [-] Qian Xinjin, see [SonGQ].

## Ra - Re

- [-] A.M.M. Rabaiah, see [JaBVR].
- [Ra1]\*\* S.P. Radziszowski, The Ramsey Numbers  $R(K_3, K_8 - e)$  and  $R(K_3, K_9 - e)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **8** (1990) 137-145.
- [Ra2] S.P. Radziszowski, Small Ramsey Numbers, *Technical Report RIT-TR-93-009*, Department of Computer Science, Rochester Institute of Technology, 1993.
- [Ra3]\*\* S.P. Radziszowski, On the Ramsey Number  $R(K_5 - e, K_5 - e)$ , *Ars Combinatoria*, **36** (1993) 225-232.
- [Ra4] S.P. Radziszowski, Ramsey Numbers Involving Cycles, in *Ramsey Theory: Yesterday, Today and Tomorrow* (ed. A. Soifer), Progress in Mathematics 285, Springer-Birkhauser 2011, 41-62.
- [RaJi] S.P. Radziszowski and Jin Xia, Paths, Cycles and Wheels in Graphs without Antitriangles, *Australasian Journal of Combinatorics*, **9** (1994) 221-232.
- [RaK1]\* S.P. Radziszowski and D.L. Kreher, Search Algorithm for Ramsey Graphs by Union of Group Orbits, *Journal of Graph Theory*, **12** (1988) 59-72.
- [RaK2]\*\* S.P. Radziszowski and D.L. Kreher, On  $R(3, k)$  Ramsey Graphs: Theoretical and Computational Results, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 37-52.
- [RaK3]\*\* S.P. Radziszowski and D.L. Kreher, Upper Bounds for Some Ramsey Numbers  $R(3, k)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **4** (1988) 207-212.
- [RaK4] S.P. Radziszowski and D.L. Kreher, Minimum Triangle-Free Graphs, *Ars Combinatoria*, **31** (1991) 65-92.
- [RaT]\* S.P. Radziszowski and Kung-Kuen Tse, A Computational Approach for the Ramsey Numbers  $R(C_4, K_n)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **42** (2002) 195-207.

- [RaST]\* S.P. Radziszowski, J. Stinehour and Kung-Kuen Tse, Computation of the Ramsey Number  $R(W_5, K_5)$ , *Bulletin of the Institute of Combinatorics and its Applications*, **47** (2006) 53-57.
- [-] S.P. Radziszowski, see also [BaRT, BILR, BoRa, CalSR, DyDR, FeKR, GoeR1, GoeR2, HasHR, JaNR, KrLR, LaLR, LiaRX, MPR, MR1, MR2, MR3, MR4, MR5, McR, PR1, PR2, ShWR, WuSR, WuSZR, XuR1, XuR2, XuR3, XuR4, XSR1, XSR2, XXER, XuXR, ZhuXR].
- [RaeZ] G. Raeisi and A. Zaghian, Ramsey Number of Wheels Versus Cycles and Trees, *Canadian Mathematical Bulletin*, **59** (2016) 190-196.
- [-] G. Raeisi, see also [GyRa, KamRa1, KamRa2, MaORS1, MaORS2, OmRa1, OmRa2, OmRa3, OmRR].
- [-] Arash Rafiey, see [BeHHRs].
- [-] Prabhakar Raghavan, see [NaRT].
- [-] Z. Rahimi, see [OmRR].
- [-] J.L. Ramirez Alfonsin, see [ChaMR, MonCR].
- [Ram] F.P. Ramsey, On a Problem of Formal Logic, *Proceedings of the London Mathematical Society*, **30** (1930) 264-286.
- [RanMCG]\* M. Ranjbar, W. Macready, L. Clark and F. Gaitan, Generalized Ramsey Numbers through Adiabatic Quantum Optimization, *Quantum Information Processing*, **15** (2016) 3519-3542.
- [-] A. Rao, see [BarRSW].
- [Rao]\* S. Rao, Applying a Genetic Algorithm to Improve the Lower Bounds of Multi-Color Ramsey Numbers, *MS thesis*, Department of Computer Science, Rochester Institute of Technology, 1997.
- [-] A. Rapp, see [BudHR1, BudHR2].
- [-] E. Rauer, see [JamKR].
- [ReWi] R.C. Read and R.J. Wilson, *An Atlas of Graphs*, Clarendon Press, Oxford, 1998.
- [-] G. Resta, see [CoPR].
- [-] M.P. Revuelta, see [BoCGR].
- [-] S.W. Reyner, see [BurR].
- [-] D.F. Reynolds, see [ExRe].
- [-] Alex Rice, see [BaLL+].
- [-] A. Rimmel, see [AgCP+].

## Ro - Ru

- [Rob] B. Roberts, Ramsey Numbers of Connected Clique Matchings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P1.36, **24** (2017), 7 pages.
- [-] B. Roberts, see also [AllMRS, DavJR].
- [Rob1] F.S. Roberts, *Applied Combinatorics*, Prentice-Hall, Englewood Cliffs, 1984.
- [-] J.A. Roberts, see [BuRo1, BuRo2].
- [-] S. Roberts, see [GR].
- [Rob2]\* A. Robertson, New Lower Bounds for Some Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R12, **6** (1999), 6 pages.
- [Rob3]\* A. Robertson, Difference Ramsey Numbers and Issai Numbers, *Advances in Applied Mathematics*, **25** (2000) 153-162.
- [Rob4] A. Robertson, New Lower Bounds Formulas for Multicolored Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R13, **9** (2002), 6 pages.
- [-] A. Robertson, see also [LaRo].

- [-] Y. Roditty, see [KrRod].
- [RöTh] V. Rödl and R. Thomas, Arrangeability and Clique Subdivisions, in *The Mathematics of Paul Erdős II*, 236-239, Algorithms and Combinatorics **14**, Springer, Berlin, 1997.
- [-] V. Rödl, see also [AlRö, ChRST, ConFR, DuLR, GrRö, GRR1, GRR2, HaLP1+, HaLP2+, KosPR, KoRö1, KoRö2, KoRö3, LauPRT, MuR, NaORS, PoRRS, PuRS, PuRW].
- [-] L. Rónyai, see [AlRóS].
- [-] M. Rosenfeld, see [NeRo].
- [Ros1] V. Rosta, On a Ramsey Type Problem of J.A. Bondy and P. Erdős, I & II, *Journal of Combinatorial Theory*, Series B, **15** (1973) 94-120.
- [Ros2] V. Rosta, Ramsey Theory Applications, Dynamic Survey in *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #DS13, (2004), 43 pages.
- [-] V. Rosta, see also [BuRo3, KáRos].
- [-] B.L. Rothschild, see [GRS].
- [Rou] C.C. Rousseau, *personal communication* (2006).
- [RoJa1] C.C. Rousseau and C.J. Jayawardene, The Ramsey Number for a Quadrilateral vs. a Complete Graph on Six Vertices, *Congressus Numerantium*, **123** (1997) 97-108.
- [RoJa2] C.C. Rousseau and C.J. Jayawardene, Harary's Problem for  $K_{2,k}$ , *manuscript* (1999).
- [RoS1] C.C. Rousseau and J. Sheehan, On Ramsey Numbers for Books, *Journal of Graph Theory*, **2** (1978) 77-87.
- [RoS2] C.C. Rousseau and J. Sheehan, A Class of Ramsey Problems Involving Trees, *Journal of the London Mathematical Society* (2), **18** (1978) 392-396.
- [-] C.C. Rousseau, see also [BolJY+, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRSGJ, BFRS, CaLRZ, ChaRSPS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FRS1, FRS2, FRS3, FRS4, FRS5, FRS6, FRS7, FRS8, FRS9, FSR, JR1, JR2, JR3, JR4, JR5, JRB, LiR1, LiR2, LiR3, LiR4, LiRS, LiRZ1, LiRZ2, NiRo1, NiRo2, NiRo3, NiRo4, NiRS].
- [-] C. Rowan, see [KerRo].
- [Row1]\* F. Rowley, Constructive Lower Bounds for Ramsey Numbers from Linear Graphs, *Australasian Journal of Combinatorics*, **68** (2017) 385-395.
- [Row2]\* F. Rowley, Some Further Results in Ramsey Graph Construction, *Australasian Journal of Combinatorics*, **78** (2020) 1-10.
- [Row3]\* F. Rowley, A Generalised Linear Ramsey Graph Construction, *Australasian Journal of Combinatorics*, **81**(2) (2021) 245-256.
- [Row4]\* F. Rowley, An Improved Lower Bound for  $S(7)$  and Some Interesting Templates, *preprint*, <http://arxiv.org/abs/2107.03560> (2021).
- [Row5]\* F. Rowley, Improved Lower Bounds for Multicolor Ramsey Numbers Using SAT-Solvers, *preprint*, <http://arxiv.org/abs/2203.13476> (2022).
- [-] P. Rowlinson, see [YR1, YR2, YR3].
- [Rub] M. Rubey, Technische Universität Wien, an electronic resource for values of small Ramsey numbers, <http://www.findstat.org/StatisticsDatabase/St000479>, 2016.
- [-] A. Ruciński, see [DuRu, HanPR, JacPR, GRR1, GRR2, HaLP1+, HaLP2+, ŁuPR, PoRRS, PoRu].
- [-] Mathieu Rundström, see [AxCJMR].
- [RuoS] Jake Ruotolo and Zi-Xia Song, Multicolor Ramsey Number for Double Stars, *Discrete Mathematics*, **347** (2024) 114034, 9 pages.
- [-] M. Ruzinkó, see [GyRSS].

## Sa - Se

- [-] Reza Saei, see [BeHHRS, BeHHS].
- [Sah] Ashwin Sah, Diagonal Ramsey via Effective Quasirandomness, *Duke Mathematical Journal*, **172**(3) (2023) 545-567.
- [-] J. Sahasrabudhe, see [BalB+, CamGMS, CamJMPS, CamJMS].
- [-] M. Salerno, see [JiSa].
- [Sal] M. Sales, On the Ramsey Number of Daisies II, *preprint*, <http://arxiv.org/abs/2211.10385> (2024).
- [-] M. Sales, see also [PuRS].
- [SaTWZ] N. Salia, C. Tompkins, Zhiyu Wang and O. Zamora, Ramsey Numbers of Berge-Hypergraphs and Related Structures, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P4.40, **26**(4) (2019), 25 pages.
- [SaBr1] A.N.M. Salman and H.J. Broersma, The Ramsey Numbers of Paths Versus Kipases, *Electronic Notes in Discrete Mathematics*, **17** (2004) 251-255.
- [SaBr2] A.N.M. Salman and H.J. Broersma, Paths-Fan Ramsey Numbers, *Discrete Applied Mathematics*, **154** (2006) 1429-1436.
- [SaBr3] A.N.M. Salman and H.J. Broersma, The Ramsey Numbers for Paths Versus Wheels, *Discrete Mathematics*, **307** (2007) 975-982.
- [SaBr4] A.N.M. Salman and H.J. Broersma, Path-Kipas Ramsey Numbers, *Discrete Applied Mathematics*, **155** (2007) 1878-1884.
- [-] A.N.M. Salman, see also [HaABS].
- [-] P. San Segundo, see [FuLS].
- [Sán] A. Sánchez-Flores, An Improved Bound for Ramsey Number  $N(3, 3, 3, 3; 2)$ , *Discrete Mathematics*, **140** (1995) 281-286.
- [-] C. Sanford, see [BudHLS].
- [Sanh] N. Sanhueza-Matamala, Stability and Ramsey Numbers for Cycles and Wheels, *Discrete Mathematics*, **339** (2016) 1557-1565.
- [-] N. Sanhueza-Matamala, see also [PavPS].
- [-] Suhadi Wido Saputro, see [SherBSO, SherSBO].
- [Sár1] G.N. Sárközy, Monochromatic Cycle Partitions of Edge-Colored Graphs, *Journal of Graph Theory*, **66** (2011) 57-64.
- [Sár2] G.N. Sárközy, On the Multi-Colored Ramsey Numbers of Paths and Even Cycles, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.53, **23**(3) (2016), 9 pages.
- [Sár3] G.N. Sárközy, Improved Monochromatic Double Stars in Edge Colorings, *Graphs and Combinatorics*, **38**:78 (2022), 10 pages.
- [-] G.N. Sárközy, see also [DeBGS, GyLSS, GyRSS, GySá1, GySá2, GySá3, GySS1, GySS2, MoSST].
- [-] I. Sato, see [MiSa].
- [Saw] Will Sawin, An Improved Lower Bound for Multicolor Ramsey Numbers and A Problem of Erdős, *Journal of Combinatorial Theory, Series A*, **188** (2022) 105579, 11 pages.
- [-] D. Saxton, see [FizGMSS, GrMFSS].
- [-] M. Schacht, see [MoSST, NaORS].
- [Scha] M. Schaefer, Graph Ramsey Theory and the Polynomial Hierarchy, *Journal of Computer and System Sciences*, **62** (2001) 290-322.
- [-] R.H. Schelp, see [BalLS, BalSS, BEFRS1, BEFRS2, BEFRS3, BEFRS4, BEFRSGJ, BEFS, BFRS, ChenS, EFRS1, EFRS2, EFRS3, EFRS4, EFRS5, EFRS6, EFRS7, EFRS8, EFRS9, FLPS, FRS1, FRS2, FRS3, FRS4, FRS5, FRS6, FS1, FS2, FS3, FS4, FSR, FSS1, FSS2, GyLSS, NiRS].

- [SchSch1]\* A. Schelten and I. Schiermeyer, Ramsey Numbers  $r(K_3, G)$  for Connected Graphs  $G$  of Order Seven, *Discrete Applied Mathematics*, **79** (1997) 189-200.
- [SchSch2] A. Schelten and I. Schiermeyer, Ramsey Numbers  $r(K_3, G)$  for  $G \cong K_7 - 2P_2$  and  $G \cong K_7 - 3P_2$ , *Discrete Mathematics*, **191** (1998) 191-196.
- [-] A. Schelten, see also [FSS3].
- [Schi1] I. Schiermeyer, All Cycle-Complete Graph Ramsey Numbers  $r(C_m, K_6)$ , *Journal of Graph Theory*, **44** (2003) 251-260.
- [Schi2] I. Schiermeyer, The Cycle-Complete Graph Ramsey Number  $r(C_5, K_7)$ , *Discussiones Mathematicae Graph Theory*, **25** (2005) 129-139.
- [-] I. Schiermeyer, see also [FSS3, LiSch, LiMSY, MaoWMS, SchSch1, SchSch2].
- [-] J.C. Schlage-Puchta, see [BrGS].
- [-] A. Schneider, see [AlmHS].
- [-] J. Schönheim, see [BiaS].
- [Schu] C.-U. Schulte, Ramsey-Zahlen für Bäume und Kreise, *Ph.D. thesis*, Heinrich-Heine-Universität Düsseldorf, (1992).
- [-] D. Schuricht, see [WoGKSF].
- [-] M.J. Schuster, see [CalSR].
- [-] S. Schuster, see [ChaS].
- [-] A. Schwenk, see [ChvS].
- [Scob] M.W. Scobee, On the Ramsey Number  $R(m_1P_3, m_2P_3, m_3P_3)$  and Related Results, ..., *MA thesis*, University of Louisville (1993).
- [-] Alex Scott, see [AhS].
- [-] A. Sebő, see [GySeT].
- [-] R.J. Segegin, see [LorSe].
- [-] J.N. Senadheera, see [JaNS].

## Sh

- [Shah] M. Shahsiah, Ramsey Numbers of 5-Uniform Loose Cycles, *Graphs and Combinatorics*, **38:5** (2022), 23 pages.
- [-] M. Shahsiah, see also [MaORS1, MaORS2, MahS, OmSh1, OmSh2, OmSh3, OmSh4].
- [-] R. Shaltiel, see [BarRSW].
- [Shao]\* Zehui Shao, *personal communication* (2008).
- [ShaoWX]\* Shao Zehui, Wang Zicheng and Xiao Jianhua, Lower Bounds for Ramsey Numbers Based on Simulated Annealing Algorithm (in Chinese), *Computer Engineering and Applications*, **45** (2009) 70-71.
- [ShaXBP]\* Zehui Shao, Jin Xu, Qiquan Bao and Linqiang Pan, Computation of Some Generalized Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **75** (2010) 217-228.
- [ShaXB]\* Zehui Shao, Xiaodong Xu and Qiquan Bao, On the Ramsey Numbers  $R(C_m, B_n)$ , *Ars Combinatoria*, **94** (2010) 265-271.
- [ShaXSP]\* Zehui Shao, Xiaodong Xu, Xiaolong Shi and Linqiang Pan, Some Three-Color Ramsey Numbers  $R(P_4, P_5, C_k)$  and  $R(P_4, P_6, C_k)$ , *European Journal of Combinatorics*, **30** (2009) 396-403.
- [-] Zehui Shao, see also [SunSh, XSR1, XSR2].
- [Shas] A. Shastri, Lower Bounds for Bi-Colored Quaternary Ramsey Numbers, *Discrete Mathematics*, **84** (1990) 213-216.
- [She1] J.B. Shearer, A Note on the Independence Number of Triangle-Free Graphs, *Discrete Mathematics*, **46** (1983) 83-87.

- [She2]\* J.B. Shearer, Lower Bounds for Small Diagonal Ramsey Numbers, *Journal of Combinatorial Theory, Series A*, **42** (1986) 302-304.
- [She3] J.B. Shearer, A Note on the Independence Number of Triangle-Free Graphs II, *Journal of Combinatorial Theory, Series B*, **53** (1991) 300-307.
- [She4]\* J.B. Shearer, Independence Numbers of Paley Graphs (data for primes 1 mod 4 up to 7000), <http://www.research.ibm.com/people/s/shearer/indpal.html> (1996).
- [-] J. Sheehan, see [ChaRSPS, CIEHMS, FRS7, FRS8, FRS9, FSS1, RoS1, RoS2].
- [-] Jian Shen, see [LiShen, LinLS].
- [-] Wujie Shen, see [MaSX].
- [-] Shen Yun-Qiu, see [LuSS1, LuSS2].
- [-] Sheng Wancheng, see [HWSYZH].
- [SherBSO] I. Sherlin, E.T. Baskoro, S.W. Saputro and F. Oktariani, On the Ramsey Numbers for Trees Versus Fans, *Journal of Information Processing*, **33** (2025) 1025-1032.
- [SherSBO] I. Sherlin, S.W. Saputro, E.T. Baskoro and F. Oktariani, On Ramsey Numbers for Trees Versus Fans of Even Order, *Indonesian Journal of Combinatorics*, **8**(1) (2024) 13-25.
- [ShWR]\* D. Shetler, M. Wurtz and S.P. Radziszowski, On Some Multicolor Ramsey Numbers Involving  $K_3+e$  and  $K_4-e$ , *SIAM Journal on Discrete Mathematics*, **26** (2012) 1256-1264.
- [-] Shi Lei, see [SunYJLS].
- [Shi1] Lingsheng Shi, Cube Ramsey Numbers Are Polynomial, *Random Structures and Algorithms*, **19** (2001) 99-101.
- [Shi2] Lingsheng Shi, Upper Bounds for Ramsey Numbers, *Discrete Mathematics*, **270** (2003) 251-265.
- [Shi3] Lingsheng Shi, Linear Ramsey Numbers of Sparse Graphs, *Journal of Graph Theory*, **50** (2005) 175-185.
- [Shi4] Lingsheng Shi, The Tail Is Cut for Ramsey Numbers of Cubes, *Discrete Mathematics*, **307** (2007) 290-292.
- [Shi5] Lingsheng Shi, Ramsey Numbers of Long Cycles Versus Books or Wheels, *European Journal of Combinatorics*, **31** (2010) 828-838.
- [ShZ1] Shi Ling Sheng and Zhang Ke Min, An Upper Bound Formula for Ramsey Numbers, *manuscript* (2001).
- [ShZ2] Shi Ling Sheng and Zhang Ke Min, A Sequence of Formulas for Ramsey Numbers, *manuscript* (2001).
- [-] Xiaolong Shi, see [ShaXSP].
- [-] Yongtang Shi, see [LiuMS].
- [ShiuLL] Shiu Wai Chee, Peter Che Bor Lam and Li Yusheng, On Some Three-Color Ramsey Numbers, *Graphs and Combinatorics*, **19** (2003) 249-258.

### Si - St

- [Sid1] A.F. Sidorenko, On Turán Numbers  $T(n, 5, 4)$  and Number of Monochromatic 4-cliques in 2-colored 3-graphs (in Russian), *Voprosy Kibernetiki*, **64** (1980) 117-124.
- [Sid2] A.F. Sidorenko, An Upper Bound on the Ramsey Number  $R(K_3, G)$  Depending Only on the Size of the Graph  $G$ , *Journal of Graph Theory*, **15** (1991) 15-17.
- [Sid3] A.F. Sidorenko, The Ramsey Number of an  $N$ -Edge Graph Versus Triangle Is at Most  $2N+1$ , *Journal of Combinatorial Theory, Series B*, **58** (1993) 185-196.
- [-] M. Simonovits, see [AjKSS, BalSS, FSS2, FaSi, HaLP1+, KoSS1, KoSS2, LucSS].
- [-] F. Simutis, see [DeMST].

- [-] Sakshi Singh, see [BaLL+].
- [-] J. Skokan, see [AllBS, AllMRS, BalCSW, BenSk, FizGMSS, GrMFSS, HaLP1+, HaLP2+, JenSk, KeeLS, KoSS1, KoSS2, ŁucSS].
- [Slo] Leila Sloman, A Very Big Small Leap Forward in Graph Theory (May), The Lawlessness of Large Numbers (July), essays in *Quanta Magazine*, May/July 2023.
- [-] S. Smorodinsky, see [JaKSY].
- [-] M.J. Smuga-Otto, see [AbbS].
- [Sob] A. Sobczyk, Euclidian Simplices and the Ramsey Number  $R(4, 4; 3)$ , *Technical Report #10, Clemson University* (1967).
- [Soi1] A. Soifer, *The Mathematical Coloring Book, Mathematics of Coloring and the Colorful Life of Its Creators*, Springer 2009.
- [Soi2] A. Soifer, *Ramsey Theory: Yesterday, Today and Tomorrow*, Progress in Mathematics 285, Springer-Birkhauser 2011.
- [Soi3] A. Soifer, *The New Mathematical Coloring Book, Mathematics of Coloring and the Colorful Life of Its Creators*, Springer 2024.
- [-] W. Solomon, see [LorSo].
- [-] L. Soltés, see [LiRS].
- [-] Cruise Song, see [NaSZ].
- [Song1] Song En Min, Study of Some Ramsey Numbers (in Chinese), a note (announcement of results without proofs), *Mathematica Applicata*, **4**(2) (1991) 6.
- [Song2] Song En Min, New Lower Bound Formulas for the Ramsey Numbers  $N(k, k, \dots, k; 2)$  (in Chinese), *Mathematica Applicata*, **6** (1993) suppl., 113-116.
- [Song3] Song En Min, An Investigation of Properties of Ramsey Numbers (in Chinese), *Mathematica Applicata*, **7** (1994) 216-221.
- [Song4] Song En Min, Properties and New Lower Bounds of the Ramsey Numbers  $R(p, q; 4)$  (in Chinese), *Journal of Huazhong University of Science and Technology*, **23** (1995) suppl. II, 1-4.
- [SonYL] Song En Min, Ye Weiguo and Liu Yanwu, New Lower Bounds for Ramsey Number  $R(p, q; 4)$ , *Discrete Mathematics*, **145** (1995) 343-346.
- [-] Song En Min, see also [HuSo, ZLLS].
- [Song5] Song Hongxue, Asymptotic Upper Bounds for Wheel-Complete Graph Ramsey Numbers, *Journal of Southeast University* (English Edition), ISSN 1003-7985, **20** (2004) 126-129.
- [Song6] Song Hongxue, A Ramsey Goodness Result for Graphs with Large Pendent Trees, *Journal of Mathematical Study (China)*, **42** (2009) 36-39.
- [Song7] Song Hong-xue, Asymptotic Upper Bounds for  $K_2 + T_m$  : Complete Graph Ramsey Numbers, *Journal of Mathematics (China)*, **30** (2010) 797-802.
- [Song8] Song Hongxue, Asymptotic Lower Bounds of Ramsey Numbers for  $r$ -Uniform Hypergraphs, *Advances in Mathematics, China*, **40** (2011) 179-186.
- [Song9] Hongxue Song, Asymptotic Upper Bounds for  $K_{1,m,k}$  : Complete Graph Ramsey Numbers, *Ars Combinatoria*, **111** (2013) 137-144.
- [SonBL] Song Hong Xue, Bai Lu Feng and Liu Shu Yan, Asymptotic Upper Bounds for the Wheel-Complete Graph Ramsey Numbers (in Chinese), *Acta Mathematica Scientia, Series A*, ISSN 1003-3998, **26** (2006) 741-746.
- [SonGQ] Song Hongxue, Gu Hua and Qian Xinjin, On the Ramsey Number of  $K_3$  Versus  $K_2 + T_n$  (in Chinese), *Journal of Liaoning Normal University, Natural Science Edition*, ISSN 1000-1735, **27** (2004) 142-145.

- [SonLi] Song Hongxue and Li Yusheng, Asymptotic Lower Bounds of Ramsey Numbers for 4-Uniform Hypergraphs (in Chinese), *Journal of Nanjing University Mathematical Biquarterly*, **26** (2009) 216-224.
- [-] Song Hongxue, see also [GuSL].
- [-] Shixi Song, see [LinS].
- [-] Zi-Xia Song, see [RuoS].
- [-] N. Spanier, see [MuSp].
- [-] J.P. de Souza, see [BotMds].
- [Spe1] J.H. Spencer, Ramsey's Theorem - A New Lower Bound, *Journal of Combinatorial Theory, Series A*, **18** (1975) 108-115.
- [Spe2] J.H. Spencer, Asymptotic Lower Bounds for Ramsey Functions, *Discrete Mathematics*, **20** (1977) 69-76.
- [Spe3] J.H. Spencer, Eighty Years of Ramsey  $R(3, k)$  ... and Counting! in *Ramsey Theory: Yesterday, Today and Tomorrow* (ed. A. Soifer), Progress in Mathematics 285, Springer-Birkhauser 2011, 27-39.
- [-] J.H. Spencer, see also [BES, GRS].
- [-] T.S. Spencer, see [BahS].
- [Spe4]\* T. Spencer, University of Nebraska at Omaha, *personal communication* (1993), and, Upper Bounds for Ramsey Numbers via Linear Programming, *manuscript* (1994).
- [-] A.K. Srivastava, see [GauST].
- [Stahl] S. Stahl, On the Ramsey Number  $R(F, K_m)$  where  $F$  is a Forest, *Canadian Journal of Mathematics*, **27** (1975) 585-589.
- [-] R.G. Stanton, see [KaSt].
- [Stat] W. Staton, Some Ramsey-type Numbers and the Independence Ratio, *Transactions of the American Mathematical Society*, **256** (1979) 353-370.
- [-] A. Steger, see [McDS].
- [-] Maya Stein, see [FloS].
- [-] J. Stinehour, see [RaST].
- [Stev] S. Stevens, Ramsey Numbers for Stars Versus Complete Multipartite Graphs, *Congressus Numerantium*, **73** (1990) 63-71.
- [-] M.J. Stewart, see [ChaRSPS].
- [Stone] J.C. Stone, Utilizing a Cancellation Algorithm to Improve the Bounds of  $R(5, 5)$ , (1996). This paper claimed incorrectly that  $R(5, 5) = 50$ .

## Su - Sul

- [-] Pascal Su, see [KniSu].
- [SuL]\* Su Wenlong and Luo Haipeng, Prime Order Cyclic Graphs and New Lower Bounds for Three Classical Ramsey Numbers  $R(4, n)$  (in Chinese), *Journal of Mathematical Study*, **31**, 4 (1998) 442-446.
- [SuLL]\* Su Wenlong, Luo Haipeng and Li Qiao, New Lower Bounds of Classical Ramsey Numbers  $R(4, 12)$ ,  $R(5, 11)$  and  $R(5, 12)$ , *Chinese Science Bulletin*, **43**, 6 (1998) 528.
- [SLLL]\* Su Wenlong, Luo Haipeng, Li Guiqing and Li Qiao, Lower Bounds of Ramsey Numbers Based on Cubic Residues, *Discrete Mathematics*, **250** (2002) 197-209.
- [SLZL]\* Su Wenlong, Luo Haipeng, Zhang Zhengyou and Li Guiqing, New Lower Bounds of Fifteen Classical Ramsey Numbers, *Australasian Journal of Combinatorics*, **19** (1999) 91-99.
- [-] Su Wenlong, see also [LiaWXCS, LiaWXS, LuSL, LiSLW, LuSS1, LuSS2, WSLX1, WSLX2, XWCS].

- [Sud1] B. Sudakov, A Note on Odd Cycle-Complete Graph Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #N1, **9** (2002), 4 pages.
- [Sud2] B. Sudakov, Large  $K_r$ -Free Subgraphs in  $K_s$ -Free Graphs and Some Other Ramsey-Type Problems, *Random Structures and Algorithms*, **26** (2005) 253-265.
- [Sud3] B. Sudakov, Ramsey Numbers and the Size of Graphs, *SIAM Journal on Discrete Mathematics*, **21** (2007) 980-986.
- [Sud4] B. Sudakov, Recent Developments in Extremal Combinatorics: Ramsey and Turán Type Problems, *Proceedings of the International Congress of Mathematicians*, Hyderabad, India, 2010, 2579-2606.
- [Sud5] B. Sudakov, A Conjecture of Erdős on Graph Ramsey Numbers, *Advances in Mathematics*, **227** (2011) 601-609
- [-] B. Sudakov, see also [AIKS, BalPS, BraFS1, BraFS2, BraGS, BraHS, BucSu, ConFLS, ConFS1, ConFS2, ConFS3, ConFS4, ConFS5, ConFS6, ConFS7, ConFS8, FoxSu1, FoxSu2, HunMS, KoSu, PoSu1, PoSu2].
- [-] A. Sudan, see [GuGS].
- [Sudar1] I.W. Sudarsana, The Goodness of Long Path with Respect to Multiple Copies of Small Wheel, *Far East Journal of Mathematical Sciences*, **59** (2011) 47-55.
- [Sudar2] I.W. Sudarsana, The Goodness of Long Path with Respect to Multiple Copies of Complete Graphs, *Journal of the Indonesian Mathematical Society*, **20** (2014) 31-35.
- [Sudar3] I.W. Sudarsana, The Goodness of Path or Cycle with Respect to Multiple Copies of Complete Graphs of Order Three, *Ars Combinatoria*, **126** (2016) 359-367.
- [Sudar4] I.W. Sudarsana, On the Ramsey Number for Cycle with Respect to Identical Copies of Complete Graphs, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **99** (2016) 61-68.
- [Sudar5] I.W. Sudarsana, A Note on the Ramsey Number for a Cycle with Respect to a Disjoint Union of Wheels, *Electronic Journal of Graph Theory and Applications*, **9**(2) (2021) 561-566.
- [SuAM] I.W. Sudarsana, Adiwijaya and S. Musdalifah, The Ramsey Number for a Linear Forest Versus Two Identical Copies of Complete Graphs, COCOON 2010, LNCS 6196, Springer, Berlin (2010) 209-215.
- [SuAAM] I.W. Sudarsana, H. Assiyatun, Adiwijaya and S. Musdalifah, The Ramsey Number for a Linear Forest Versus Two Identical Copies of Complete Graph, *Discrete Mathematics, Algorithms and Applications*, **2** (2010) 437-444.
- [SuAUB] I.W. Sudarsana, H. Assiyatun, S. Uttunggadewa and E.T. Baskoro, On the Ramsey Numbers  $R(S_{2,m}, K_{2,q})$  and  $R(sK_2, K_s + C_n)$ , *Ars Combinatoria*, **119** (2015) 235-246.
- [SuBAU1] I.W. Sudarsana, E.T. Baskoro, H. Assiyatun and S. Uttunggadewa, The Ramsey Number of a Certain Forest with Respect to a Small Wheel, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **71** (2009) 257-264.
- [SuBAU2] I.W. Sudarsana, E.T. Baskoro, H. Assiyatun and S. Uttunggadewa, The Ramsey Numbers of Linear Forest Versus  $3K_3 \cup 2K_4$ , *Journal of the Indonesian Mathematical Society*, **15** (2009) 61-67.
- [SuBAU3] I.W. Sudarsana, E.T. Baskoro, H. Assiyatun and S. Uttunggadewa, The Ramsey Numbers for the Union Graph with  $H$ -Good Components, *Far East Journal of Mathematical Sciences*, **39** (2010) 29-40.
- [SuBAU4] I.W. Sudarsana, E.T. Baskoro, H. Assiyatun and S. Uttunggadewa, On the Union of Graph Ramsey Numbers, *Applied Mathematical Sciences*, **8**(16) (2014) 767-773.
- [SukZ] Andrew Suk and Ji Zeng, 3-Uniform Monotone Paths and Multicolor Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2411.15649> (2024).
- [-] Andrew Suk, see also [ConFG1+, ConFG2+, ConFH+, FoxPS1, FoxPS2, MuSuk1, MuSuk2, MuSuk3, MuSuk4, MuSuk5].
- [SulTr] A. Sulser and M. Trujić, Ramsey Numbers for Multiple Copies of Sparse Graphs, *Journal of Graph Theory*, **106** (2024) 843-870.

## Sun - Sz

- [SunSh]\*\* Minhong Sun and Zehui Shao, Exact Values of Some Generalized Ramsey Numbers, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **107** (2018) 277-283.
- [Sun]\* Sun Yongqi, Research on Ramsey Numbers of Some Graphs (in Chinese), *Ph.D. thesis*, Dalian University of Technology, China, July 2006.
- [SunY]\* Sun Yongqi and Yang Yuansheng, Study of the Three Color Ramsey Number  $R_3(C_8)$  (in Chinese), *Journal of Beijing Jiaotong University*, **35** (2011) 14-17.
- [SunYJLS] Sun Yongqi, Yang Yuansheng, Jiang Baoqi, Lin Xiaohui and Shi Lei, On Multicolor Ramsey Numbers for Even Cycles in Graphs, *Ars Combinatoria*, **84** (2007) 333-343.
- [SunYLZ1]\* Sun Yongqi, Yang Yuansheng, Lin Xiaohui and Zheng Wenping, The Value of the Ramsey Number  $R_4(C_4)$ , *Utilitas Mathematica*, **73** (2007) 33-44.
- [SunYLZ2]\* Sun Yongqi, Yang Yuansheng, Lin Xiaohui and Zheng Wenping, On the Three Color Ramsey Numbers  $R(C_m, C_4, C_4)$ , *Ars Combinatoria*, **84** (2007) 3-11.
- [SunYW]\* Sun Yongqi, Yang Yuansheng and Wang Zhihai, The Value of the Ramsey Number  $R_5(C_6)$ , *Utilitas Mathematica*, **76** (2008) 25-31.
- [SunYWLX]\* Sun Yongqi, Yang Yuansheng, Wang Wei, Li Bingxi and Xu Feng, Study of Three Color Ramsey numbers  $R(C_{m_1}, C_{m_2}, C_{m_3})$  (in Chinese), *Journal of Dalian University of Technology*, ISSN 1000-8608, **46** (2006) 428-433.
- [SunYXL] Sun Yongqi, Yang Yuansheng, Xu Feng and Li Bingxi, New Lower Bounds on the Multicolor Ramsey Numbers  $R_r(C_{2m})$ , *Graphs and Combinatorics*, **22** (2006) 283-288.
- [-] Sun Yongqi, see also [JiLSX, WuSL, WuSR, WuSZR, ZhaSW, ZhuSWZ].
- [-] Yue Ru Sun, see [NoSZ].
- [SunLi] Sun Yuqin and Li Yusheng, On an Upper Bound of Ramsey Number  $r_k(K_{m,n})$  with Large  $n$ , *Heilongjiang Daxue Ziran Kexue Xuebao*, ISSN 1001-7011, **23** (2006) 668-670.
- [SunZ1] Zhi-Hong Sun, Ramsey Numbers for Trees, *Bulletin of the Australian Mathematical Society*, **86** (2012) 164-176.
- [SunZ2] Zhi-Hong Sun, Ramsey Numbers for Trees II, *preprint*, <http://arxiv.org/abs/1410.7637> (2014). Revised version (2016). *Czechoslovak Mathematical Journal*, **71** (2021), no. 146, 351-372.
- [SunW] Zhi-Hong Sun and Lin-Lin Wang, Turán's Problem for Trees, *Journal of Combinatorics and Number Theory*, **3** (2011) 51-69.
- [SunWW] Zhi-Hong Sun, Lin-Lin Wang and Yi-Li Wu, Turán's Problem and Ramsey Numbers for Trees, *Colloquium Mathematicum*, **139** (2015) 273-298.
- [Sur] Surahmat, Cycle-Wheel Ramsey Numbers. Some results, open problems and conjectures. *Math Track*, ISSN 1817-3462, 1818-5495, **2** (2006) 56-64.
- [SuBa1] Surahmat and E.T. Baskoro, On the Ramsey Number of a Path or a Star Versus  $W_4$  or  $W_5$ , *Proceedings of the 12-th Australasian Workshop on Combinatorial Algorithms*, Bandung, Indonesia, July 14-17 (2001) 174-179.
- [SuBa2] Surahmat and E.T. Baskoro, The Ramsey Number of Linear Forest Versus Wheel, paper presented at the *13-th Australasian Workshop on Combinatorial Algorithms (AWOCA)*, Fraser Island, Queensland, Australia, July 7-10, 2002.
- [SuBB1] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Star-like Trees Versus Large Odd Wheels, *Technical Report #1621*, Faculty of Mathematical Sciences, University of Twente, The Netherlands, (2002).
- [SuBB2] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Cycles Versus Small Wheels, *Integers: Electronic Journal of Combinatorial Number Theory*, <http://www.integers-ejcnt.org/vol4.html>, #A10, **4** (2004), 9 pages.

- [SuBB3] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Fans Versus  $K_4$ , *Bulletin of the Institute of Combinatorics and its Applications*, **43** (2005) 96-102.
- [SuBB4] Surahmat, E.T. Baskoro and H.J. Broersma, The Ramsey Numbers of Large Star and Large Star-Like Trees Versus Odd Wheels, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **65** (2008) 153-162.
- [SuBT1] Surahmat, E.T. Baskoro and I. Tomescu, The Ramsey Numbers of Large Cycles Versus Wheels, *Discrete Mathematics*, **306** (2006), 3334-3337.
- [SuBT2] Surahmat, E.T. Baskoro and I. Tomescu, The Ramsey Numbers of Large Cycles Versus Odd Wheels, *Graphs and Combinatorics*, **24** (2008), 53-58.
- [SuBTB] Surahmat, E.T. Baskoro, I. Tomescu and H.J. Broersma, On Ramsey Numbers of Cycles with Respect to Generalized Even Wheels, *manuscript* (2006).
- [SuBUB] Surahmat, E.T. Baskoro, S. Uttunggadewa and H.J. Broersma, An Upper Bound for the Ramsey Number of a Cycle of Length Four Versus Wheels, in *LNCS 3330*, Springer, Berlin (2005) 181-184.
- [SuTo] Surahmat and I. Tomescu, On Path-Jahangir Ramsey Numbers, *Applied Mathematical Sciences*, **8**(99) (2014) 4899-4904.
- [-] Surahmat, see also [AliSur, BaSu, BaSNM].
- [-] Y. Suzuki, see [KaOS].
- [SwPr] C.J. Swanepoel and L.M. Pretorius, Upper Bounds for a Ramsey Theorem for Trees, *Graphs and Combinatorics*, **10** (1994) 377-382.
- [-] M.M. Sweet, see [FreSw].
- [-] T. Szabó, see [AIRóS].
- [-] S. Szeider, see [KirS].
- [Szem] E. Szemerédi, Regular Partitions of Graphs, *Problèmes Combinatoires et Théorie des Graphes* (Orsay, 1976), *Colloques Internationaux du Centre National de la Recherche Scientifique, CNRS Paris*, **260** (1978) 399-401.
- [-] E. Szemerédi, see also [AjKS, AjKSS, ChRST, GyRSS, GySS1, GySS2, PoRRS].
- [-] P. Szuca, see [DzNS].

## T

- [-] M. Tait, see [DeMST].
- [-] D.S. Taletskii, see [AbdMT].
- [-] Fuping Tan, see [HTHZ1, HTHZ2].
- [-] Ta Sheng Tan, see [ChngBTW, ChngTW].
- [-] Tang Xueqing, see [LiTZ].
- [-] A. Taraz, see [MoSST].
- [Tat]\* M. Tatarevic, *personal communication*, graph constructions for lower bounds on Ramsey numbers at <http://github.com/milostatarevic/ramsey-numbers/tree/master/graphs> (2020).
- [-] M. Tatarevic, see also [ExT].
- [-] Yanmei Teng, see [JiLTX1, JiLTX2].
- [-] Abhradeep Thakurta, see [NaRT].
- [-] N. Thapen, see [LauPRT].
- [-] R. Thomas, see [RöTh].
- [Tho] A. Thomason, An Upper Bound for Some Ramsey Numbers, *Journal of Graph Theory*, **12** (1988) 509-517.

- [-] S. Thompson, see [HaHT].
- [-] M. Tiba, see [BalB+].
- [Tikh] K. Tikhomirov, A Remark on the Ramsey Number of the Hypercube, *European Journal of Combinatorics*, **120** (2024) 103954, 24 pages.
- [-] P.W. Tingley, see [HaLT].
- [-] J. Tomasik, see [AgCP+].
- [-] I. Tomescu, see [AliBT1, AliBT2, AliTJ, SuBT1, SuBT2, SuBTB, SuTo].
- [-] C. Tompkins, see [SaTWZ].
- [-] C.A. Tovey, see [CaET].
- [Tri] A. Tripathi, An Introduction to Ramsey's Theorem, *Proceedings of Telangana Academy of Sciences*, **1** (2020) 202-213.
- [-] A. Tripathi, see also [GauST].
- [Tr] Trivial results.
- [-] N. Trotignon, see [GySeT].
- [-] W.T. Trotter Jr., see [ChRST].
- [-] M. Trujić, see [SulTr].
- [Tse1]\* Kung-Kuen Tse, On the Ramsey Number of the Quadrilateral Versus the Book and the Wheel, *Australasian Journal of Combinatorics*, **27** (2003) 163-167.
- [Tse2]\* Kung-Kuen Tse, A Note on the Ramsey Numbers  $R(C_4, B_n)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **58** (2006) 97-100.
- [Tse3]\* Kung-Kuen Tse, A Note on Some Ramsey Numbers  $R(C_p, C_q, C_r)$ , *Journal of Combinatorial Mathematics and Combinatorial Computing*, **62** (2007) 189-192.
- [-] Kung-Kuen Tse, see also [BaRT, RaST, RaT].
- [-] Z. Tuza, see [GyTu].
- [-] M. Tyomkyn, see [ConT].

## U

- [UCSD] University of California San Diego, Fan Chung and a team of students, *Erdős' Problems on Graphs, Ramsey Theory*, <http://www.math.ucsd.edu/~erdosproblems/RamseyTheory.html> (2010-2012).
- [-] S. Uttungadewa, see [SuAUB, SuBAU1, SuBAU2, SuBAU3, SuBAU4, SuBUB].

## V

- [VO]\*\* Steven Van Overberghe, Algorithms for Computing Ramsey Numbers, *MS Thesis in Mathematics*, Ghent University, Belgium, 2020. Constructions at <https://github.com/Steven-VO/circulant-Ramsey>.
- [VO1]\*\* Steven Van Overberghe, Ghent University, *personal communication* (2024). Constructions available at <https://github.com/Steven-VO/circulant-Ramsey>.
- [-] Steven Van Overberghe, see also [GoeVO, LidMPV].
- [Ver] J. Verstraëte, Recent Progress in Ramsey Theory, *preprint*, <http://arxiv.org/abs/2503.22094> (2025).
- [-] J. Verstraëte, see [ConFG1+, ConFG2+, ConFH+, ConMMV, KosMV1, KosMV2, MatMNV, MatVer, MuVer, NieVer].
- [-] T. Vetric, see [JaBVR].
- [-] M. Vizer, see [GerMOV].
- [-] L. Volkmann, see [GuoV].

## W

- [-] A.Z. Wagner, see [BalCSW].
- [Walk] K. Walker, Dichromatic Graphs and Ramsey Numbers, *Journal of Combinatorial Theory*, **5** (1968) 238-243.
- [Wall] W.D. Wallis, On a Ramsey Number for Paths, *Journal of Combinatorics, Information & System Sciences*, **6** (1981) 295-296.
- [Wan] Wan Honghui, Upper Bounds for Ramsey Numbers  $R(3, 3, \dots, 3)$  and Schur Numbers, *Journal of Graph Theory*, **26** (1997) 119-122.
- [-] Wang Gongben, see [WW, WWY1, WWY2].
- [-] Lin-Lin Wang, see [SunW, SunWW].
- [Wang1] Longqin Wang, Some Multi-Color Ramsey Numbers on Stars Versus Path, Cycle or Wheel, *Graphs and Combinatorics*, **36** (2020) 515-524.
- [Wang2] Longqin Wang, The Ramsey Numbers of Trees Versus Generalized 6-Wheels or Generalized 7-Wheels, *Graphs and Combinatorics*, **38**:153 (2022), 9 pages.
- [WaCh1] Longqin Wang and Yaojun Chen, The Ramsey Numbers of Trees Versus Generalized Wheels, *Graphs and Combinatorics*, **35** (2019) 189-193.
- [WaCh2] Longqin Wang and Yaojun Chen, The Ramsey Numbers of Two Sets of Cycles, *Journal of Graph Theory*, **96**(1) (2021) 129-136.
- [WaQi] Maoqun Wang and Jianguo Qian, Ramsey Numbers for Complete Graphs Versus Generalized Fans, *Graphs and Combinatorics*, **38**:186 (2022), 8 pages.
- [WW]\* Wang Qingxian and Wang Gongben, New Lower Bounds of Ramsey Numbers  $r(3, q)$  (in Chinese), *Acta Scientiarum Naturalium, Universitatis Pekinensis*, **25** (1989) 117-121. The lower bounds presented in this paper have been improved.
- [WWY1]\* Wang Qingxian, Wang Gongben and Yan Shuda, A Search Algorithm And New Lower Bounds for Ramsey Numbers  $r(3, q)$ , *manuscript* (1994).
- [WWY2]\* Wang Qingxian, Wang Gongben and Yan Shuda, The Ramsey Numbers  $R(K_3, K_q - e)$  (in Chinese), *Beijing Daxue Xuebao Ziran Kexue Ban*, **34** (1998) 15-20.
- [-] Wang Wei, see [SunYWLX, SunYXL].
- [WaLi] Ye Wang and Yusheng Li, Lower Bounds for Ramsey Numbers of  $K_n$  with a Small Subgraph Removed, *Discrete Applied Mathematics*, **160** (2012) 2063-2068.
- [WaLL] Ye Wang, Yusheng Li and Yan Li, Ramsey Numbers of Several  $K_{t,s}$  and A Large  $K_{m,n}$ , *Discrete Mathematics*, **345** (2022) 112987, 9 pages.
- [-] Ye Wang, see also [LiLW, LiW, LiuW].
- [-] Wang Yuandi, see [HWSYZH].
- [-] Zhao Wang, see [MaoWMS].
- [WaZh1] Zhaofa Wang and Yanbo Zhang, Ramsey Numbers of Cycles Versus Multiple Wheels, *Filomat*, **39**(33) (2025) 11961-11967.
- [WaZh2] Zhaofa Wang and Yanbo Zhang, Ramsey Numbers of Cycles Versus Multiple Odd Wheels, *Bulletin of the Australian Mathematical Society*, DOI:10.1017/S0004972725100476 (2025), 9 pages.
- [-] Wang Zhi Jian, see [LiWa1, LiWa2].
- [-] Wang Zhihai, see [SunYW].
- [-] Zhiyu Wang, see [SaTWZ].
- [-] Wang Zicheng, see [ShaoWX].
- [-] Christine Rose Ward, see [BaLL+].

- [-] Bing Wei, see [ZhaoW].
- [-] Louis Wei, see [GuNNW].
- [-] Meiqin Wei, see [XuWLM, ZhouLMW].
- [We1] William J. Wesley, Algebraic and Boolean Methods for Computation and Certification of Ramsey-Type Numbers, *PhD dissertation*, University of California, Davis, 2023.
- [We2]\* William J. Wesley, Lower Bounds for Book Ramsey Numbers, *Discrete Mathematics*, **349** (2026) 114913, 13 pages.
- [We3]\* William J. Wesley, New Bounds for Some Small Multicolor Ramsey Numbers, *preprint*, <http://arxiv.org/abs/2509.03784> (2025).
- [-] William J. Wesley, see also [PuRW].
- [West] D. West, *Introduction to Graph Theory*, second edition, Prentice Hall, 2001.
- [Wh] E.G. Whitehead, The Ramsey Number  $N(3, 3, 3, 3; 2)$ , *Discrete Mathematics*, **4** (1973) 389-396.
- [-] Avi Wigderson, see [BarRSW].
- [Wig1] Yuval Wigderson, An Improved Lower Bound on Multicolor Ramsey Numbers, *Proceedings of the American Mathematical Society*, **149**(6) (2021) 2371-2374.
- [Wig2] Yuval Wigderson, Ramsey Numbers Upon Vertex Deletion, *Journal of Graph Theory*, **106** (2024) 663-675.
- [Wig3] Yuval Wigderson, Upper Bounds on Diagonal Ramsey Numbers [after Campos, Griffiths, Morris, and Sahasrabudhe], *preprint*, <http://arxiv.org/abs/2411.09321> (2024). Also in *Séminaire BOURBAKI*, **77** (2024-2025), no. 1230, 48 pages.
- [-] Yuval Wigderson, see also [FoxHW, HeNWY, HeWi, ConFW1, ConFW2].
- [-] E.R. Williams, see [AbbW].
- [-] R.J. Wilson, see [ReWi].
- [-] R.M. Wilson, see [FraWi].
- [-] A. Woldar, see [LaWo1, LaWo2].
- [-] Kok Bin Wong, see [ChngBTW, ChngTW].
- [WoGKSF]\* Jurriaan Wouters, Aris Giotis, Ross Kang, Dirk Schuricht and Lars Fritz, Lower Bounds for Ramsey Numbers As a Statistical Physics Problem, *Journal of Statistical Mechanics: Theory and Experiment*, (2022) 033211, 14 pages.
- [WSLX1]\* Kang Wu, Wenlong Su, Haipeng Luo and Xiaodong Xu, New Lower Bound for Seven Classical Ramsey Numbers  $R(3, q)$ , *Applied Mathematics Letters*, **22** (2009) 365-368.
- [WSLX2]\* Kang Wu, Wenlong Su, Haipeng Luo and Xiaodong Xu, A Generalization of Generalized Paley Graphs and New Lower Bounds for  $R(3, q)$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #N25, **17** (2010), 10 pages.
- [-] Wu Kang, see also [LiaWXCS, LiaWXS, LiSLW, XWCS].
- [WuSL] Yali Wu, Yongqi Sun and Zhiguo Liu, The Ramsey Numbers  $R(C_{\leq n}, K_m)$ , *Ars Combinatoria*, **131** (2017) 227-237.
- [WuSR] Wu Yali, Sun Yongqi and S.P. Radziszowski, Wheel and Star-Critical Ramsey Numbers for Quadrilateral, *Discrete Applied Mathematics*, **186** (2015) 260-271.
- [WuSZR] Wu Yali, Sun Yongqi, Zhang Rui and S.P. Radziszowski, Ramsey Numbers of  $C_4$  Versus Wheels and Stars, *Graphs and Combinatorics*, **31** (2015) 2437-2446.
- [-] Wu Yali, see also [ZhaSW, ZhuSWZ].
- [-] Yi-Li Wu, see [SunWW].
- [-] M. Wurtz, see [ShWR].

## X

- [-] Xiao Jianhua, see [ShaoWX].
- [XieZ]\* Xie Jiguo and Zhang Xiaoxian, A New Lower Bound for Ramsey Number  $r(3, 13)$  (in Chinese), *Journal of Lanzhou Railway Institute*, **12** (1993) 87-89.
- [-] Shengjie Xie, see [MaSX].
- [-] Xie Zheng, see [XuX1, XuX2, XuXC, XXER, XuXR].
- [XWCS]\* Chengzhang Xu, Kang Wu, Hong Chen and Wenlong Su, New Lower Bounds for Some Ramsey Numbers Based on Cyclic Graphs, *in preparation*, (2011).
- [-] Xu Chengzhang, see also [LiaWXCS].
- [XuYZ] Chuandong Xu, Hongna Yang and Shenggui Zhang, On Characterizing the Critical Graphs for Matching Ramsey Numbers, *Discrete Applied Mathematics*, **287** (2020) 15-20.
- [-] Jin Xu, see [ShaXBP].
- [-] Xu Feng, see [SunYWLX, SunYXL].
- [-] Ran Xu, see [ChenCX].
- [XuWLM] Xiao Xu, Meiqin Wei, Hong-Jian Lai and Yaping Mao, Ramsey and Gallai-Ramsey Numbers for Comb and Sun Graphs, *Discrete Applied Mathematics*, **362** (2025) 131-145.
- [XuLL] Xiaodong Xu, Meilian Liang and Haipeng Luo, *Ramsey Theory. Unsolved Problems and Results*. De Gruyter, Berlin; University of Science and Technology of China Press, 2018.
- [XuR1] Xiaodong Xu and S.P. Radziszowski, An Improvement to MATHON's Cyclotomic Ramsey Colorings, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #N1, **16**(1) (2009), 5 pages.
- [XuR2] Xiaodong Xu and S.P. Radziszowski,  $28 \leq R(C_4, C_4, C_3, C_3) \leq 36$ , *Utilitas Mathematica*, **79** (2009) 253-257.
- [XuR3] Xiaodong Xu and S.P. Radziszowski, Bounds on Shannon Capacity and Ramsey Numbers from Product of Graphs, *IEEE Transactions on Information Theory*, **59** (2013) 4767-4770.
- [XuR4] Xiaodong Xu and S.P. Radziszowski, On Some Open Questions for Ramsey and Folkman Numbers, in *Graph Theory, Favorite Conjectures and Open Problems*, Vol. 1, edited by R. Gera, S. Hedetniemi and C. Larson, Problem Books in Mathematics, Springer 2016, 43-62.
- [XSR1]\* Xiaodong Xu, Zehui Shao and S.P. Radziszowski, Bounds on Some Ramsey Numbers Involving Quadrilateral, *Ars Combinatoria*, **90** (2009) 337-344.
- [XSR2]\* Xiaodong Xu, Zehui Shao and S.P. Radziszowski, More Constructive Lower Bounds on Classical Ramsey Numbers, *SIAM Journal on Discrete Mathematics*, **25** (2011) 394-400.
- [XuX1]\* Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on the Ramsey Numbers  $r(k, l)$ , *manuscript* (2002).
- [XuX2] Xu Xiaodong and Xie Zheng, A Constructive Approach for the Lower Bounds on Multicolor Ramsey Numbers, *manuscript* (2002).
- [XuXC] Xu Xiaodong, Xie Zheng and Chen Zhi, Upper Bounds for Ramsey Numbers  $R_n(3)$  and Schur Numbers (in Chinese), *Mathematics in Economics*, **19**(1) (2002) 81-84.
- [XXER]\* Xu Xiaodong, Xie Zheng, G. Exoo and S.P. Radziszowski, Constructive Lower Bounds on Classical Multicolor Ramsey Numbers, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #R35, **11**(1) (2004), 24 pages.
- [XuXR] Xu Xiaodong, Xie Zheng and S.P. Radziszowski, A Constructive Approach for the Lower Bounds on the Ramsey Numbers  $R(s, t)$ , *Journal of Graph Theory*, **47** (2004) 231-239.
- [-] Xu Xiaodong, see also [JiLSX, JiLTX1, JiLTX2, LiaRX, LiaWXS, LiaX, ShaXB, ShaXSP, WSLX1, WSLX2, ZhuXR].
- [-] Xu Zhiqiang, see [BaLX].

- [XuGe] Zixiang Xu and Gennian Ge, A Note on Multicolor Ramsey Number of Small Odd Cycles Versus a Large Clique, *Discrete Mathematics*, **345** (2022) 112823, 4 pages..

## Y

- [-] J. Yackel, see [GrY].
- [Yan] Jun Yan, Asymmetric Ramsey Numbers of Trees, *preprint*, <http://arxiv.org/abs/2511.15673> (2025).
- [-] Jun Yan, see also [MonPY1, MonPY2].
- [-] Yan Shuda, see [WWY1, WWY2].
- [YanP] Zilong Yan and Yuejian Peng, Tree Embeddings and Tree-Star Ramsey Numbers, *Journal of Graph Theory*, **100**(4) (2022) 733-745.
- [-] Hongna Yang, see [XuYZ].
- [YHZ1] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min, The Value of the Ramsey Number  $R(C_n, K_4)$  is  $3(n-1)+1$  ( $n \geq 4$ ), *Australasian Journal of Combinatorics*, **20** (1999) 205-206.
- [YHZ2] Yang Jian Sheng, Huang Yi Ru and Zhang Ke Min,  $R(C_6, K_5) = 21$  and  $R(C_7, K_5) = 25$ , *European Journal of Combinatorics*, **22** (2001) 561-567.
- [-] Yang Jian Sheng, see also [BoJY+, HWSYZH, HYZ].
- [YY]\*\* Yang Yuansheng, On the Third Ramsey Numbers of Graphs with Six Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **17** (1995) 199-208.
- [YH]\* Yang Yuansheng and G.R.T. Hendry, The Ramsey Number  $r(K_1+C_4, K_5-e)$ , *Journal of Graph Theory*, **19** (1995) 13-15.
- [YR1]\*\* Yang Yuansheng and P. Rowlinson, On the Third Ramsey Numbers of Graphs with Five Edges, *Journal of Combinatorial Mathematics and Combinatorial Computing*, **11** (1992) 213-222.
- [YR2]\* Yang Yuansheng and P. Rowlinson, On Graphs without 6-Cycles and Related Ramsey Numbers, *Utilitas Mathematica*, **44** (1993) 192-196.
- [YR3]\* Yang Yuansheng and P. Rowlinson, The Third Ramsey Numbers for Graphs with at Most Four Edges, *Discrete Mathematics*, **125** (1994) 399-406.
- [-] Yang Yuansheng, see also [SunY, SunYJLS, SunYLZ1, SunYLZ2, SunYW, SunYWLX, SunYXL].
- [YaoHMZ] Yifan Yao, Zhong Huang, Yaping Mao and Jiannan Zhou, Ramsey and Gallai-Ramsey Numbers for Multiple Triangles of Graphs and Their Multiplicities, *Discrete Applied Mathematics*, **377** (2025) 195-203.
- [-] Yifan Yao, see also [LiMSY].
- [-] Ye Weiguo, see [SonYL].
- [YouLin1] Chunlin You and Qizhong Lin, Ramsey Numbers of Large Even Cycles and Fans, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.14, **30**(3) (2023), 17 pages.
- [YouLin2] Chunlin You and Qizhong Lin, Three-Color Ramsey Numbers of an Odd Cycle Versus Bipartite Graphs with Small Bandwidth, *Graphs and Combinatorics*, **39**:46 (2023), 18 pages.
- [YouLC] Chunlin You, Qizhong Lin and Xun Chen, Ramsey Numbers of Large Books and Bipartite Graphs with Small Bandwidth, *Discrete Mathematics*, **344** (2021) 112427, 9 pages.
- [-] Chunlin You, see also [ChenLY].
- [-] Hung-Hsun Hans Yu, see [ConFG2+, HeNWY].
- [YuLi] Pei Yu and Yusheng Li, All Ramsey Numbers for Brooms in Graphs, *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P3.29, **23**(3) (2016), 8 pages.
- [-] Pei Yu, see also [PeiCLY].
- [Yu1]\* Yu Song Nian, A Computer Assisted Number Theoretical Construction of  $(3, k)$ -Ramsey Graphs, *Annales Universitatis Scientiarum Budapestinensis, Sect. Comput.*, **10** (1989) 35-44.

- [Yu2]\* Yu Song Nian, Maximal Triangle-Free Circulant Graphs and the Function  $K(c)$  (in Chinese), *Journal of Shanghai University, Natural Science*, **2** (1996) 678-682.
- [-] Xiaowei Yu, see [ChenYZ].
- [-] Y. Yuditsky, see [JaKSY].
- [-] R. Yuster, see [CaYZ].

## Z

- [-] A. Zaghian, see [RaeZ].
- [-] V. Zamaraev, see [AleALZ, AtLZ].
- [-] O. Zamora, see [SaTWZ].
- [-] Zang Wenan, see [LiRZ1, LiRZ2, LiTZ, LiZa1, LiZa2].
- [-] C. Zarb, see [CaYZ].
- [ZeZZ] Haiyu Zeng, Yanbo Zhang and Feng Zhao, Ramsey Numbers of Small Wheels Versus Fans, *Bulletin of the Malaysian Mathematical Sciences Society*, **49** (2026), article #1, 13 pages.
- [-] Haiyu Zeng, see also [HeZZ].
- [-] Ji Zeng, see [SukZ].
- [Zeng] Zeng Wei Bin, Ramsey Numbers for Triangles and Graphs of Order Four with No Isolated Vertex, *Journal of Mathematical Research & Exposition*, **6** (1986) 27-32.
- [-] Zhang Chaohui, see [HTHZ2].
- [ZhCh] Fangfang Zhang and Yaojun Chen, The Ramsey Numbers of 3-Uniform Loose Path Versus Star, *Graphs and Combinatorics*, **38**:20 (2022), 11 pages.
- [ZhZC] Fangfang Zhang, Yunqing Zhang and Yaojun Chen, On Three-color Ramsey Numbers  $R(C_4, K_{1,m}, P_n)$ , *Finite Fields and Their Applications*, **55** (2019) 97-108.
- [-] Hanshuo Zhang, see [ZhuSWZ].
- [ZhZ1] Zhang Ke Min and Zhang Shu Sheng, Some Tree-Stars Ramsey Numbers, *Proceedings of the Second Asian Mathematical Conference 1995*, 287-291, World Sci. Publishing, River Edge, NJ, 1998.
- [ZhZ2] Zhang Ke Min and Zhang Shu Sheng, The Ramsey Numbers for Stars and Stripes, *Acta Mathematica Scientia*, **25A** (2005) 1067-1072.
- [-] Zhang Ke Min, see also [BolJY+, ChenZZ1, ChenZZ2, ChenZZ3, ChenZZ4, ChenZZ5, ChenZZ6, HTHZ1, HWSYZH, HYZ, HZ1, HZ2, HZ3, McZ, ShZ1, ShZ2, YHZ1, YHZ2, ZhaCZ1, ZhaCZ2, ZZ3].
- [ZhaCC1] Lianmin Zhang, Yaojun Chen and T.C. Edwin Cheng, The Ramsey Numbers for Cycles Versus Wheels of Even Order, *European Journal of Combinatorics*, **31** (2010) 254-259.
- [ZhaCZ] Lianmin Zhang, Kun Chen and Dongmei Zhu, Some Tree-Book Ramsey Numbers, *Ars Combinatoria*, **130** (2017) 97-102.
- [-] Lianmin Zhang, see also [ZhuZL].
- [-] Ningxin Zhang, see also [NaSZ].
- [-] Ping Zhang, see [ChaZ].
- [-] Ping Zhang, see [LiZZ].
- [ZhaSW]\* Zhang Rui, Sun Yongqi and Wu Yali, On the Four Color Ramsey Numbers for Hexagons, *Ars Combinatoria*, **111** (2013) 515-522.
- [-] Zhang Rui, see also [WuSZR].
- [-] Shenggui Zhang, see [XuYZ].
- [-] Zhang Shu Sheng, see [ZhZ1, ZhZ2].

- [-] Zhang Xiaoxian, see [XieZ].
- [ZhaHou] Xuejun Zhang and Xinmin Hou, Multicolor Ramsey Numbers of Stars Versus a Path, *Journal of Mathematical Research with Applications*, **44**(4) (2024) 465-469.
- [ZhaCC2] Xuemei Zhang, Yaojun Chen and T.C. Edwin Cheng, Some Values of Ramsey Numbers for  $C_4$  Versus Stars, *Finite Fields and Their Applications*, **45** (2017) 73-85.
- [ZhaCC3] Xuemei Zhang, Yaojun Chen and T.C. Edwin Cheng, Polarity Graphs and Ramsey Numbers for  $C_4$  Versus Stars, *Discrete Mathematics*, **340** (2017) 655-660.
- [ZhaCC4] Xuemei Zhang, Yaojun Chen and T.C. Edwin Cheng, On Three Color Ramsey Numbers for  $R(C_4, C_4, K_{1,n})$ , *Discrete Mathematics*, **342** (2019) 285-291.
- [ZhaCC5] Xuemei Zhang, Yaojun Chen and T.C. Edwin Cheng, On the 3-Color Ramsey Numbers  $R(C_4, C_4, W_n)$ , *Graphs and Combinatorics*, **38**:103 (2022), 9 pages.
- [ZhaCC6] Xuemei Zhang, Yaojun Chen and T.C. Edwin Cheng, Bounds for Two Multicolor Ramsey Numbers Concerning Quadrilaterals, *Finite Fields and Their Applications*, **79** (2022) 101999, 16 pages.
- [-] Zhang Xuemei, see also [ChenZZ7].
- [-] Yahui Zhang, see [LiZZ].
- [ZhaBC1] Yanbo Zhang, Hajo Broersma and Yaojun Chen, A Remark on Star- $C_4$  and Wheel- $C_4$  Ramsey Numbers, *Electronic Journal of Graph Theory and Applications*, **2** (2014) 110-114.
- [ZhaBC2] Yanbo Zhang, Hajo Broersma and Yaojun Chen, Three Results on Cycle-Wheel Ramsey Numbers, *Graphs and Combinatorics*, **31** (2015) 2467-2479.
- [ZhaBC3] Yanbo Zhang, Hajo Broersma and Yaojun Chen, Ramsey Numbers of Trees Versus Fans, *Discrete Mathematics*, **338** (2015) 994-999.
- [ZhaBC4] Yanbo Zhang, Hajo Broersma and Yaojun Chen, A Note on Ramsey Numbers for Fans, *Bulletin of the Australian Mathematical Society*, **92** (2015) 19-23.
- [ZhaBC5] Yanbo Zhang, Hajo Broersma and Yaojun Chen, On Fan-Wheel and Tree-Wheel Ramsey Numbers, *Discrete Mathematics*, **339** (2016) 2284-2287.
- [ZhaBC6] Yanbo Zhang, Hajo Broersma and Yaojun Chen, Narrowing Down the Gap on Cycle-Star Ramsey Numbers, *Journal of Combinatorics*, **7** (2016) 481-493.
- [ZhaCh1] Yanbo Zhang and Yaojun Chen, The Ramsey Numbers of Fans Versus a Complete Graph of Order Five, *Electronic Journal of Graph Theory and Applications*, **2** (2014) 66-69.
- [ZhaCh3] Yanbo Zhang and Yaojun Chen, Trichotomy and  $tK_m$ -goodness of Sparse Graphs, *preprint*, <http://arxiv.org/abs/2505.04142> (2025).
- [ZhaCh4] Yanbo Zhang and Yaojun Chen, Ramsey Goodness of Fans, *Discrete Mathematics*, **349**(2) (2026) 114771, 8 pages.
- [ZhaCh5] Yanbo Zhang and Yaojun Chen, New Bounds for Ramsey Numbers Involving Graphs with a Center, *submitted for publication* (2025).
- [ZhaChZ1] Yanbo Zhang, Yaojun Chen and Yunqing Zhang, Ramsey Numbers of Trees Versus Generalized Wheels, *Discrete Mathematics*, **349** (2026) 114938, 7 pages.
- [ZhaChZ2] Yanbo Zhang, Yaojun Chen and Yunqing Zhang, On Tree-Wheel Ramsey Numbers, *Discrete Applied Mathematics*, **383** (2026) 80-84.
- [ZhaZC] Yanbo Zhang, Yunqing Zhang and Yaojun Chen, The Ramsey Numbers of Wheels Versus Odd Cycles, *Discrete Mathematics*, **323** (2014) 76-80.
- [ZhaZZ] Zhang Yanbo, Zhu Shiping and Zhang Yunqing, Ramsey Numbers for 7-Cycle Versus Wheels with Small Order (in Chinese), *Journal of Nanjing University, Mathematical Biquarterly*, **30** (2013) 48-55.
- [-] Yanbo Zhang, see also [AllLPZ, ChenZZ7, HeZZ, HuZC1, HuZC2, HuZC3, LiZBBH, LiZB, MengZZ, WaZh1, WaZh2, ZeZZ].
- [-] Yiran Zhang, see [HuPZ].

- [-] Zhang Yuming, see [CaLRZ].
- [Zhang1] Zhang Yunqing, On Ramsey Numbers of Short Paths Versus Large Wheels, *Ars Combinatoria*, **89** (2008) 11-20.
- [Zhang2] Zhang Yunqing, The Ramsey Numbers for Stars of Odd Small Order Versus a Wheel of Order Nine, *Nanjing Daxue Xuebao Shuxue Bannian Kan*, ISSN 0469-5097, **25** (2008) 35-40.
- [ZhaCC7] Yunqing Zhang, T.C. Edwin Cheng and Yaojun Chen, The Ramsey Numbers for Stars of Odd Order Versus a Wheel of Order Nine, *Discrete Mathematics, Algorithms and Applications*, **1** (2009) 413-436.
- [ZhaCZ1] Yunqing Zhang, Yaojun Chen and Kemin Zhang, The Ramsey Numbers for Stars of Even Order Versus a Wheel of Order Nine, *European Journal of Combinatorics*, **29** (2008) 1744-1754.
- [ZhaCZ2] Yunqing Zhang, Yaojun Chen and Kemin Zhang, The Ramsey Numbers for Trees of High Degree Versus a Wheel of Order Nine, *manuscript* (2009).
- [ZZ3] Yunqing Zhang and Ke Min Zhang, The Ramsey Number  $R(C_8, K_8)$ , *Discrete Mathematics*, **309** (2009) 1084-1090.
- [-] Zhang Yunqing, see also [ChenCNZ, ChenCZ1, ChenZZ1, ChenZZ2, ChenZZ3, ChenZZ4, ChenZZ5, ChenZZ6, CheCZN, MengZZ, ZhZC, ZhaZC, ZhaChZ1, ZhaChZ2, ZhaZZ].
- [-] Zhang Zhengyou, see [SLZL].
- [ZLLS] Zhang Zhongfu, Liu Linzhong, Li Jinwen and Song En Min, Some Properties of Ramsey Numbers, *Applied Mathematics Letters*, **16** (2003) 1187-1193.
- [-] Feng Zhao, see [ZeZZ].
- [ZhaoW] Qinghong Zhao and Bing Wei, Some Exact Values on Ramsey Numbers Related to Fans, *preprint*, <http://arxiv.org/abs/2211.02338> (2022).
- [Zhao] Yi Zhao, Proof of the  $(n/2 - n/2 - n/2)$  Conjecture for Large  $n$ , *Electronic Journal of Combinatorics*, <http://www.combinatorics.org>, #P27, **18**(1) (2011), 61 pages.
- [-] Yi Zhao, see also [ChenYZ, NoSZ].
- [-] Zheng Wenping, see [SunYLZ1, SunYLZ2].
- [Zhou1] Zhou Huai Lu, Some Ramsey Numbers for Graphs with Cycles (in Chinese), *Mathematica Applicata*, **6** (1993) 218.
- [Zhou2] Zhou Huai Lu, The Ramsey Number of an Odd Cycle with Respect to a Wheel (in Chinese), *Journal of Mathematics, Shuxue Zazhi* (Wuhan), **15** (1995) 119-120.
- [Zhou3] Zhou Huai Lu, On Book-Wheel Ramsey Number, *Discrete Mathematics*, **224** (2000) 239-249.
- [ZhouLMW] Jiannan Zhou, Zhihui Li, Yaping Mao and Meiqin Wei, Ramsey and Gallai-Ramsey Numbers for the Union of Paths and Stars, *Discrete Applied Mathematics*, **325** (2023) 297-308.
- [ZhuZL] Dongmei Zhu, Lianmin Zhang and Dongxin Li, The Ramsey Numbers of Large Trees Versus Wheels, *Bulletin of the Iranian Mathematical Society*, **42**(4) (2016) 879-880.
- [-] Dongmei Zhu, see also [ZhaCZ].
- [-] Emily Zhu, see [BohZ].
- [ZhuXR] Rujie Zhu, Xiaodong Xu and S.P. Radziszowski, A Small Step Forwards on the Erdős-Sós Problem Concerning the Ramsey Numbers  $R(3, k)$ , *Discrete Applied Mathematics*, **214** (2016) 216-221.
- [-] Zhu Shiping, see [ZhaZZ].
- [ZhuSWZ] Weiguo Zhu, Yongqi Sun, Yali Wu and Hanshuo Zhang, Exact Values of Multicolor Ramsey Numbers  $R_l(C_{\leq l+1})$ , *Graphs and Combinatorics*, **36** (2020) 839-852.